

ON SIMULTANEOUS RATIONALITY OF TWO AHMES SERIES

VJEKOSLAV KOVAČ

ABSTRACT. Paul Erdős asked how rapidly a sequence of positive integers (n_k) can grow if both series $\sum_k 1/n_k$ and $\sum_k 1/(n_k - 1)$ have rational sums. In this note we show that there exists an exponentially growing sequence (n_k) with this property. Previous records had polynomial growth, even for easier variants of the problem, regarding the series $\sum_k 1/n_k$ and $\sum_k 1/(n_k - d)$ for any concrete nonzero integer d .

1. INTRODUCTION

A series of unit fractions

$$\sum_{k=1}^{\infty} \frac{1}{n_k} \quad \text{for some positive integers } n_1 < n_2 < n_3 < \dots \quad (1.1)$$

was named an *Ahmes series* by Erdős and Straus [12], but the term has since been seldom used and always in relation with rationality/irrationality problems [22, 14]. A folklore result is that

$$\lim_{k \rightarrow \infty} n_k^{1/2^k} = \infty$$

is a sufficient condition guaranteeing that the sum (1.1) is an irrational number [12, 8]. Conversely, shifted Sylvester's sequence [21, A129871],

$$n_1 = 2, \quad n_{k+1} = n_k^2 - n_k + 1 \text{ for } k \geq 1,$$

has asymptotics $n_k \sim c_0^{2^k}$ for a particular constant $c_0 = 1.2640847\dots$ (see [13, p. 109], [23]) and the sum of its reciprocals equals 1. By shifting Sylvester's sequence further, one then immediately obtains sequences satisfying $n_k \sim c^{2^k}$ for arbitrarily large constants c , the reciprocals of which still sum to a rational number. We conclude that the aforementioned irrationality result is sharp; this observation is borrowed from [8, p. 2].

Erdős came up with the following related problem, which has been posed on several occasions in the 1980s [11, p. 64], [9, p. 334], [10, p. 104], and recently also on Thomas Bloom's website *Erdős problems* [3, Problem #265]. We choose the formulation stated at the *Symposium on Transcendental Number Theory* held in Durham in 1986 and published in its proceedings [10, p. 104–105]:

Once I asked: Assume that $\sum \frac{1}{n_k}$ and $\sum \frac{1}{n_k - 1}$ are both rational. How fast can n_k tend to infinity? I was (and am) sure that $n_k^{1/k} \rightarrow \infty$ is possible but $n_k^{1/2^k}$ must tend to 1. Unfortunately almost nothing is known. David Cantor observed that

$$\sum_{k=3}^{\infty} \frac{1}{\binom{k}{2}} \quad \text{and} \quad \sum_{k=3}^{\infty} \frac{1}{\binom{k}{2} - 1}$$

2020 *Mathematics Subject Classification*. Primary 11J72; Secondary 11D68.

Key words and phrases. irrationality condition, series of fractions, sequence growth.

are both rational and we do not know any sequence with this property which tends to infinity faster than polynomially.

In [9, p. 334] Erdős specifically mentioned the exponential growth as already being an interesting open problem:

(...) and we could never decide if n_k can increase exponentially or even faster.

On a different occasion, in Erdős and Graham's 1980 book on open problems in combinatorial number theory, a weaker property is also mentioned:

If 1 is replaced by a larger constant then higher degree polynomials can be used. For example, if $p(x) = x^3 + 6x^2 + 5x$ then both $\sum_{n \geq 1} \frac{1}{p(n)}$ and $\sum_{n \geq 1} \frac{1}{p(n)+8}$ are rational (since both $p(n)$ and $p(n) + 8$ completely split over the integers). Similar examples are known using polynomials with degrees as large as 10 (see [15]).

There is a harmless typo in the last quote that the author deciphered as: $p(n) + 8$ should be replaced with $p(n) - 12$ and the two series should sum over $n \geq 2$. Also note that studying rationality of the two series $\sum_k 1/n_k$ and $\sum_k 1/(n_k - d)$, for a fixed positive integer d , is indeed a simpler problem. Namely, it is solved by multiplying by d any sequence (n_k) that makes $\sum_k 1/n_k$ and $\sum_k 1/(n_k - 1)$ rational.

Our goal is to answer the question partially, by showing that the exponential growth of (n_k) is indeed possible, just as Erdős suspected. The result can be formulated rigorously as Theorem 1 below. Super-exponential growth still remains out of reach and it seems to require more sophisticated tools.

Theorem 1. *There exists a strictly increasing sequence of positive integers $(n_k)_{k=1}^{\infty}$ such that*

$$n_k > 1.01^k \text{ for every index } k, \quad (1.2)$$

while both

$$\sum_{k=1}^{\infty} \frac{1}{n_k} \text{ and } \sum_{k=1}^{\infty} \frac{1}{n_k - 1} \quad (1.3)$$

are rational numbers.

Our proof of Theorem 1 is not explicitly constructive, in the sense that the sequence (n_k) will be defined implicitly, but this might be viewed as an advantage. The main idea is very simple and it shares some similarity with the proof of a different conjecture of Erdős and Straus, recently given in [19]. We will initially arrange $\sum_k 1/n_k$ to be the sum of several, carefully chosen, geometric series with ratio $1/2$, which clearly evaluates to a rational number. Then we will describe a procedure of changing the sequence terms in a way that $\sum_k 1/n_k$ remains the same, but $\sum_k 1/(n_k - 1)$ attains all values from a non-degenerate interval. The proof will be completed by merely choosing a rational number from that interval. After the proof we will use a computer to convince ourselves additionally: we will generate the first 1000 terms of one such sequence and calculate the approximation errors. This also justifies that the proof of Theorem 1 is a bit more concrete and more on the computational side, since otherwise the asymptotic reasoning could have been used.

Further motivation for Theorem 1 comes from the fact that many obvious guesses for exponentially decaying series (1.3) are known to have at least one irrational sum. For instance, Chowla [6] conjectured and Erdős [7] proved that

$$\sum_{k=1}^{\infty} \frac{1}{q^k - 1}$$

is an irrational number for every integer $q \geq 2$. More generally, Borwein [4, 5] solved an open problem of Erdős by showing that

$$\sum_{k=1}^{\infty} \frac{1}{q^k + r}$$

is irrational whenever $q \geq 2$ is an integer and r is a rational number different from 0 and any of $-q^k$; an alternative proof appeared in [1]. Various sums of reciprocals of Fibonacci or Lucas numbers are also known to be irrational [20].

2. PROOF OF THEOREM 1

All 19 numbers from the interval $(2^6, 2^7)$ that are relatively prime to 2, 3, and 7 are listed in the increasing order as

$$\begin{aligned} s_0 &= 65, & s_1 &= 67, & s_2 &= 71, & s_3 &= 73, & s_4 &= 79, & s_5 &= 83, & s_6 &= 85, \\ s_7 &= 89, & s_8 &= 95, & s_9 &= 97, & s_{10} &= 101, & s_{11} &= 103, & s_{12} &= 107, \\ s_{13} &= 109, & s_{14} &= 113, & s_{15} &= 115, & s_{16} &= 121, & s_{17} &= 125, & s_{18} &= 127. \end{aligned}$$

A sequence $\epsilon = (\epsilon_m)_{m=0}^{\infty}$ of zeroes and ones will be chosen later, so let it be arbitrary for now. The set of all numbers appearing in the desired sequence $(n_k)_{k=1}^{\infty}$ will be of the form

$$A_{\epsilon} := \left(\bigcup_{\substack{i \geq 0, 0 \leq j \leq 18 \\ \text{such that } \epsilon_{19i+j}=0}} \{9 \cdot 2^i s_j, 21 \cdot 2^i s_j\} \right) \cup \left(\bigcup_{\substack{i \geq 0, 0 \leq j \leq 18 \\ \text{such that } \epsilon_{19i+j}=1}} \{7 \cdot 2^i s_j, 63 \cdot 2^i s_j\} \right).$$

Since

$$\frac{1}{9} + \frac{1}{21} = \frac{1}{7} + \frac{1}{63} = \frac{10}{63}, \quad (2.1)$$

the sum

$$\sum_{n \in A_{\epsilon}} \frac{1}{n} = \frac{10}{63} \underbrace{\left(\sum_{i=0}^{\infty} \frac{1}{2^i} \right)}_{=2} \left(\sum_{j=0}^{18} \frac{1}{s_j} \right)$$

is equal to a rational number independent of ϵ . Also,

$$\sum_{n \in A_{\epsilon}} \frac{1}{n-1} = y + \sum_{m=0}^{\infty} \epsilon_m x_m \quad (2.2)$$

where we have denoted

$$y := \sum_{j=0}^{18} \sum_{i=0}^{\infty} \left(\frac{1}{9 \cdot 2^i s_j - 1} + \frac{1}{21 \cdot 2^i s_j - 1} \right)$$

and

$$x_{19i+j} := \frac{1}{7 \cdot 2^i s_j - 1} + \frac{1}{63 \cdot 2^i s_j - 1} - \frac{1}{9 \cdot 2^i s_j - 1} - \frac{1}{21 \cdot 2^i s_j - 1}.$$

It remains to choose the coefficients ϵ in a way that (2.2) is a rational number too. This will certainly be possible if the set

$$\left\{ \sum_{m=0}^{\infty} \epsilon_m x_m : (\epsilon_m)_{m=0}^{\infty} \in \{0, 1\}^{\mathbb{N}_0} \right\} \quad (2.3)$$

happens to be a non-degenerate interval. It is an easy exercise, which has already been known to Kakeya [17, 18] (also see the survey paper [2]), that

$$x_m > 0 \text{ for } m = 0, 1, 2, 3, \dots, \quad (2.4a)$$

$$x_0 \geq x_1 \geq x_2 \geq x_3 \geq \dots, \quad (2.4b)$$

$$\sum_{m=0}^{\infty} x_m < \infty, \quad (2.4c)$$

$$\sum_{l=m+1}^{\infty} x_l \geq x_m \text{ for } m = 0, 1, 2, 3, \dots \quad (2.4d)$$

are sufficient conditions for (2.3) to be a closed bounded interval, namely $[0, \sum_{m=0}^{\infty} x_m] \subset \mathbb{R}$.

In the verification of all four conditions (2.4a)–(2.4d) we need the estimate

$$\frac{1}{n} + \frac{1}{n^2} + \frac{1}{n^3} \leq \frac{1}{n-1} \leq \frac{1}{n} + \frac{1}{n^2} + \frac{2}{n^3} \quad (2.5)$$

valid for $n \geq 2$; it is an easy consequence of

$$n^3 \left(\frac{1}{n-1} - \frac{1}{n} - \frac{1}{n^2} \right) = \frac{n}{n-1} \in [1, 2].$$

From (2.5) and (2.1), denoting

$$\Delta := \frac{1}{7^2} + \frac{1}{63^2} - \frac{1}{9^2} - \frac{1}{21^2} = \frac{8}{1323} > 0,$$

and observing

$$0 < \frac{2}{7^3} + \frac{2}{63^3} - \frac{1}{9^3} - \frac{1}{21^3} < \frac{1}{200}, \quad -\frac{1}{25000} < \frac{1}{7^3} + \frac{1}{63^3} - \frac{2}{9^3} - \frac{2}{21^3} < 0,$$

we can write

$$\frac{\Delta}{2^{2i} s_j^2} - \frac{1}{25000 \cdot 2^{3i} s_j^3} < x_{19i+j} < \frac{\Delta}{2^{2i} s_j^2} + \frac{1}{200 \cdot 2^{3i} s_j^3}.$$

Finally, recalling $s_j > 2^6$, we obtain

$$\left(1 - \frac{3}{20000}\right) \cdot \frac{\Delta}{2^{2i} s_j^2} < x_{19i+j} < \left(1 + \frac{3}{200}\right) \cdot \frac{\Delta}{2^{2i} s_j^2} \quad (2.6)$$

for all $i \geq 0$ and $0 \leq j \leq 18$.

Condition (2.4a) is immediate from the lower bound in (2.6), while Condition (2.4c) is clear from the corresponding upper bound. Next, for $0 \leq j \leq 17$ we have

$$\frac{x_{19i+j+1}}{x_{19i+j}} < \frac{1 + 3/200}{1 - 3/20000} \left(\max_{0 \leq j \leq 17} \frac{s_j}{s_{j+1}} \right)^2 < 1,$$

while

$$\frac{x_{19(i+1)}}{x_{19i+18}} < \frac{1 + 3/200}{1 - 3/20000} \left(\frac{s_{18}}{2s_0} \right)^2 < 1,$$

both thanks to (2.6) again. This verifies Condition (2.4b). Finally, to check Condition (2.4d), we take $m = 19i + j$, $i \geq 0$, $0 \leq j \leq 7$, apply (2.6), and recall $2^6 < s_j < 2^7$:

$$\frac{1}{x_{19i+j}} \sum_{l=19i+j+1}^{\infty} x_l > \frac{1}{x_{19i+j}} \sum_{l=19(i+1)}^{19(i+1)+18} x_l > \frac{1 - 3/20000}{1 + 3/200} \cdot \frac{19 \cdot (1/4) \cdot 2^{-14}}{2^{-12}} > 1.$$

Now we finally know that (2.2) attain values from a whole interval, so there really exists $\epsilon \in \{0, 1\}^{\mathbb{N}_0}$ such that

$$\sum_{n \in A_\epsilon} \frac{1}{n} \in \mathbb{Q} \quad \text{and} \quad \sum_{n \in A_\epsilon} \frac{1}{n-1} \in \mathbb{Q}.$$

Let $(n_k)_{k=1}^{\infty}$ be the strictly increasing sequence that enumerates A_{ϵ} . It remains to check the claim (1.2) about its growth. Clearly, for any integer $m \geq 6$ only the elements

$$9 \cdot 2^i s_j, 21 \cdot 2^i s_j \text{ or } 7 \cdot 2^i s_j, 63 \cdot 2^i s_j$$

of A_{ϵ} corresponding to the indices $0 \leq i \leq m-7$ and an arbitrary $0 \leq j \leq 18$ can be less than 2^m and there are at most $38(m-6)$ such terms. Consequently, for every $k \in \mathbb{N}$ and the unique integer $m \geq 6$ such that $38(m-6) < k \leq 38(m-5)$, we have

$$n_k \geq 2^m > 2^{k/38} > 1.01^k$$

and we are done.

3. INITIAL TERMS OF A POSSIBLE SEQUENCE

The proof from the previous section shows that there exists a sequence $(n_k)_{k=1}^{\infty}$ satisfying (1.2), such that

$$\sum_{k=1}^{\infty} \frac{1}{n_k} = \frac{20}{63} \sum_{j=0}^{18} \frac{1}{s_j} = 0.0655394679\dots, \quad (3.1)$$

while

$$\sum_{k=1}^{\infty} \frac{1}{n_k - 1} \text{ can be any number from } [0.0655851987\dots, 0.0656041482\dots].$$

We can, for instance, target

$$\sum_{k=1}^{\infty} \frac{1}{n_k - 1} = \frac{41}{625} = 0.0656, \quad (3.2)$$

and achieve this by a recursive greedy construction:

$$\epsilon_m := \begin{cases} 0 & \text{if } x_m + \sum_{l=0}^{m-1} \epsilon_l x_l > 41/625 - y, \\ 1 & \text{if } x_m + \sum_{l=0}^{m-1} \epsilon_l x_l \leq 41/625 - y \end{cases}$$

for $m = 0, 1, 2, \dots$. The empty sum for $m = 0$ is understood to be 0. We can use Mathematica [16] to generate $(n_k)_{k=1}^{1000}$ as the smallest 1000 elements of the obtained set A_{ϵ} ; the sequence begins:

$$\begin{aligned} n_1 &= 455, & n_2 &= 469, & n_3 &= 497, & n_4 &= 511, & n_5 &= 553, \\ n_6 &= 581, & n_7 &= 595, & n_8 &= 623, & n_9 &= 665, & n_{10} &= 679, \dots \end{aligned}$$

The finite sums $\sum_{k=1}^{1000} 1/n_k$ and $\sum_{k=1}^{1000} 1/(n_k - 1)$ respectively differ from (3.1) and (3.2) by less than 10^{-9} .

ACKNOWLEDGMENT

This work was supported in part by the Croatian Science Foundation under the project HRZZ-IP-2022-10-5116 (FANAP).

REFERENCES

- [1] Tewodros Amdeberhan and Doron Zeilberger. q -Apéry irrationality proofs by q -WZ pairs. *Adv. in Appl. Math.*, 20(2):275–283, 1998. doi:10.1006/aama.1997.0565.
- [2] Artur Bartoszewicz, Małgorzata Filipczak, and Franciszek Prus-Wiśniowski. Topological and algebraic aspects of subsums of series. In *Traditional and present-day topics in real analysis*, pages 345–366. Faculty of Mathematics and Computer Science. University of Łódź, Łódź, 2013.
- [3] Thomas Bloom. Erdős problems. <https://www.erdosproblems.com/>. Accessed: June 24, 2024.
- [4] Peter B. Borwein. On the irrationality of $\sum(1/(q^n + r))$. *J. Number Theory*, 37(3):253–259, 1991. doi:10.1016/S0022-314X(05)80041-1.

- [5] Peter B. Borwein. On the irrationality of certain series. *Math. Proc. Cambridge Philos. Soc.*, 112(1):141–146, 1992. doi:10.1017/S030500410007081X.
- [6] Sarvadaman Chowla. On series of the Lambert type which assume irrational values for rational values of the argument. *Proc. Nat. Inst. Sci. India*, 13:171–173, 1947.
- [7] Paul Erdős. On arithmetical properties of Lambert series. *J. Indian Math. Soc. (N.S.)*, 12:63–66, 1948.
- [8] Paul Erdős. Some problems and results on the irrationality of the sum of infinite series. *J. Math. Sci.*, 10:1–7, 1975.
- [9] Paul Erdős. E. Straus (1921–1983). *Rocky Mountain J. Math.*, 15(2):331–341, 1985. Number theory (Winnipeg, Man., 1983). doi:10.1216/RMJ-1985-15-2-331.
- [10] Paul Erdős. On the irrationality of certain series: problems and results. In *New advances in transcendence theory (Durham, 1986)*, Alan Baker (Ed.), pages 102–109. Cambridge Univ. Press, Cambridge, 1988.
- [11] Paul Erdős and Ronald L. Graham. *Old and new problems and results in combinatorial number theory*, volume 28 of *Monographies de L’Enseignement Mathématique*. Université de Genève, L’Enseignement Mathématique, Geneva, 1980.
- [12] Paul Erdős and Ernst G. Straus. On the irrationality of certain Ahmes series. *J. Indian Math. Soc. (N.S.)*, 27:129–133, 1964.
- [13] Ronald L. Graham, Donald E. Knuth, and Oren Patashnik. *Concrete mathematics*. Addison-Wesley Publishing Company, Advanced Book Program, Reading, 1989.
- [14] Jaroslav Hančl and Robert Tijdeman. On the irrationality of Cantor and Ahmes series. *Publ. Math. Debrecen*, 65(3-4):371–380, 2004. doi:10.5486/pmd.2004.3254.
- [15] Godfrey H. Hardy and Edward M. Wright. *An introduction to the theory of numbers*. Oxford, at the Clarendon Press, 1954. 3rd ed.
- [16] Wolfram Research, Inc. Mathematica, Version 13.0.0. Champaign, IL, 2021. URL: <https://www.wolfram.com/mathematica>.
- [17] Sōichi Kakeya. On the partial sums of an infinite series. *Tōhoku Sci. Rep.*, 3:159–164, 1914.
- [18] Sōichi Kakeya. On the set of partial sums of an infinite series. *Proc. Tokyo Math.-Phys. Soc., 2nd ser.*, 7:250–251, 1914.
- [19] Vjekoslav Kovač. On the set of points represented by harmonic subseries. preprint, 2024. URL: <https://arxiv.org/abs/2405.07681>.
- [20] Tapani Matala-Aho and Marc Prévost. Quantitative irrationality for sums of reciprocals of Fibonacci and Lucas numbers. *Ramanujan J.*, 11(2):249–261, 2006. doi:10.1007/s11139-006-6511-4.
- [21] Neil J. A. Sloane and The OEIS Foundation Inc. The on-line encyclopedia of integer sequences, 2024. URL: <http://oeis.org>.
- [22] Robert Tijdeman and Pingzhi Yuan. On the rationality of Cantor and Ahmes series. *Indag. Math. (N.S.)*, 13(3):407–418, 2002. doi:10.1016/S0019-3577(02)80018-0.
- [23] Ilan Vardi. *Computational recreations in Mathematica*. Addison-Wesley Publishing Company, Advanced Book Program, Redwood City, CA, 1991.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, UNIVERSITY OF ZAGREB, BIJENIČKA CESTA 30,
10000 ZAGREB, CROATIA

Email address: vjekovac@math.hr