

OPTIMAL PARTIAL TRANSPORT FOR METRIC PAIRS

MAURICIO CHE

ABSTRACT. In this article we study Figalli and Gigli’s formulation of optimal transport between non-negative Radon measures in the setting of metric pairs. We carry over classical characterisations of optimal plans to this setting and prove that the resulting spaces of measures, $\mathcal{M}_p(X, A)$, are complete, separable and geodesic whenever the underlying space, X , is so. We also prove that, for $p > 1$, $\mathcal{M}_p(X, A)$ preserves the property of being non-branching, and for $p = 2$ it preserves non-negative curvature in the Alexandrov sense. Finally, we prove isometric embeddings of generalised spaces of persistence diagrams $\mathcal{D}_p(X, A)$ into the corresponding spaces $\mathcal{M}_p(X, A)$, generalising a result by Divol and Lacombe. As an application of this framework, we show that several known geometric properties of spaces of persistence diagrams follow from those of $\mathcal{M}_p(X, A)$, including the fact that $\mathcal{D}_2(X, A)$ is an Alexandrov space of non-negative curvature whenever X is a proper non-negatively curved Alexandrov space.

1. INTRODUCTION

Optimal transport provides a geometric perspective on the study of spaces of probability measures. It can be formulated as the problem of minimising the total cost of transferring a given amount of mass between two given distributions, provided we know the cost of delivering mass between any two locations. Remarkably simple though this framework might seem, it has found applications to several different areas and has become a prominent body of research in the intersection of analysis, geometry, and probability (see the encyclopedic presentation in [28] for a detailed account of the history of this subject).

An important restriction for classic optimal transport is that it only makes sense for measures with the same total mass, and it is therefore interesting to explore ways to define optimal transport between unbalanced measures. Different approaches have been proposed (see for example [9, 12, 16, 17, 19, 22] and references therein), and very recently, Savaré and Sodini, in [26], presented a general formulation that includes previous approaches in the case of finite Radon measures.

In [17], Figalli and Gigli introduced a notion of optimal transport between non-negative Radon measures defined on bounded domains in Euclidean space, motivated by finding solutions to evolution equations with Dirichlet boundary conditions, in the spirit of Jordan–Kinderlehrer–Otto scheme [20]. The idea here is to use the boundary of the domain as an unlimited supply and storage of mass, such that it can be used to compensate the difference in mass between the given measures, as long as one pays the cost of transporting it to the boundary (see Section 3 for rigorous definitions). For the sake of simplicity, refer to this notion as *optimal partial transport*, although we acknowledge that this terminology has been used in the past to denote the related work by Figalli [16].

More recently, in [14], Divol and Lacombe revisited and applied this theory to study spaces of persistence diagrams, which arise in topological data analysis (see [13, 15, 29] for foundational results on this area). They extended definitions in [17] to Radon measures on unbounded domains in \mathbb{R}^n and proved several useful properties of the resulting metric spaces (e.g. completeness, separability, characterisation of convergence, existence of Fréchet means). They also proved that the space of

Date: 26th June 2024.

Supported by CONAHCYT (Mexico) Doctoral Scholarship No. 769708.

persistence diagrams endowed with the Wasserstein distances can be isometrically embedded into these spaces of measures, by regarding persistence diagrams as discrete measures on a half-plane in \mathbb{R}^2 . Also based on the framework of optimal partial transport, Bate and Garcia Pulido proved the embeddability of finite atomic measures, endowed with the optimal partial transport metrics, into Hilbert spaces [4].

In this article, we extend the theory of optimal partial transport to the setting of *metric pairs*. Namely, we consider ordered pairs (X, A) , where X is a proper (i.e. closed and bounded subsets are compact), complete and separable metric space, and A is a closed, non-empty subset of X (cf. [1, 10, 11]), and non-negative Radon measures μ on $X \setminus A$ such that the function $\text{dist}(\cdot, A)$ is in $L^p(\mu)$. As a result, we obtain one-parameter families of metric spaces $\{(\mathcal{M}_p(X, A), \text{Wb}_p)\}_{p \in [1, \infty)}$ associated to each metric pair (X, A) . This is a natural extension of Figalli and Gigli's definition, which is recovered when $X = \overline{\Omega}$ and $A = \partial\Omega$, where Ω is a bounded open set in \mathbb{R}^n . We also recover Divol and Lacombe's setting when $X = \{(x, y) \in \mathbb{R}^2 : x \leq y\}$ and $A = \{(x, y) \in \mathbb{R}^2 : x = y\}$.

The structure of the paper is as follows. We start by proving the existence of optimal partial transport plans and the fact that Wb_p defines a metric on $\mathcal{M}_p(X, A)$ (Theorems 3.5 and 3.6), generalising results in [14, 17]. Along the way, we correct an oversight in the proof of [14, Proposition 3.2] (see discussion in the proof of Theorem 3.5). We then prove that the spaces $\mathcal{M}_p(X, A)$, endowed with the optimal partial transport distance Wb_p , are complete, separable and geodesic, provided that X has the same properties (Propositions 5.1 and 5.2). Moreover, when $p > 1$, we prove that $\mathcal{M}_p(X, A)$ is non-branching (Proposition 5.4), and for $p = 2$, we prove that $\mathcal{M}_2(X, A)$ is non-negatively curved in the Alexandrov sense, whenever X is so (Theorem 5.5). Finally, in section 6, we prove that the generalised spaces of persistence diagrams $\mathcal{D}_p(X, A)$ introduced in [10] can be isometrically embedded into $\mathcal{M}_p(X, A)$, generalising [14, Proposition 3.5]. It is important to note that, combined with Theorem 5.5, this gives an alternative proof of the fact that $\mathcal{D}_2(X, A)$ is a non-negatively curved Alexandrov space whenever X is proper and non-negatively curved [10, Proposition 7.3] (see also [7] for related results), which in the case $X = \{(x, y) \in \mathbb{R}^2 : x \leq y\}$ and $A = \{(x, y) \in \mathbb{R}^2 : x = y\}$ yields the known result, due to Turner, Mileyko, Mukherjee and Harer, that the usual space of persistence diagrams endowed with the L^2 -Wasserstein distance is non-negatively curved [27, Theorem 2.5].

Acknowledgements. I would like to express my gratitude to Fernando Galaz-García for all his support and advise, as this work is part of my doctoral thesis under his supervision. I would also like to thank Jaime Santos-Rodríguez, Martin Kerin, Kohei Suzuki, Alpár Mészáros, Amit Einav, Mo Dick Wong, Norbert Peyerimhoff and Mohammad Al Attar for all the valuable comments and fruitful discussions during several sessions of our reading seminar in Durham University, where the contents of this article were first discussed. Finally, I also thank Javier Casado and Manuel Mellado Cuerno for the careful reading of a preliminary version of this paper. I have been financially supported by CONAHCYT (Mexico), through the Doctoral Scholarship No. 769708.

2. PRELIMINARIES

2.1. Metric geometry. We briefly recall well-known definitions and results about metric spaces, geodesics, and lower curvature bounds in the Alexandrov sense (see [8] for a more detailed discussion).

Definition 2.1. Let X be a metric space. We denote by $\mathcal{C}([a, b], X)$ the space of continuous curves $\xi: [a, b] \rightarrow X$, endowed with the uniform metric. For any $t \in [a, b]$, $e_t: \mathcal{C}([a, b], X) \rightarrow X$ is the evaluation map given by $e_t(\xi) = \xi_t = \xi(t)$.

A *constant speed geodesic*, or simply a *geodesic*, is a continuous curve $\xi \in \mathcal{C}([0, 1], X)$ such that

$$d(\xi_s, \xi_t) = d(\xi_0, \xi_1)|s - t|$$

for any $s, t \in [0, 1]$. We denote by $\text{Geo}(X)$ the space of geodesics in X , endowed with the uniform metric. We say that X is a *geodesic space* if for any $x, y \in X$ there exists $\xi \in \text{Geo}(X)$ such that $\xi_0 = x$ and $\xi_1 = y$.

It is known that if X is a complete, separable and geodesic space, then $\text{Geo}(X)$, endowed with the uniform metric, is complete and separable. Moreover, if X is a proper space then $\text{Geo}(X)$ is proper as well, by the Arzelà-Ascoli theorem.

Definition 2.2. We say that a geodesic space X is *non-branching* if for any $t \in (0, 1)$ the map $(e_0, e_t): \text{Geo}(X) \rightarrow X \times X$ is injective.

Alexandrov spaces are synthetic generalisations of Riemannian manifolds with sectional curvature bounded from below. This generalisation comes from the classical Toponogov's comparison theorem in Riemannian geometry (see [18, 23]).

More precisely, the n -dimensional *model space* with constant sectional curvature κ is given by

$$\mathbb{M}_\kappa^n = \begin{cases} \mathbb{S}_\kappa^n, & \text{if } \kappa > 0, \\ \mathbb{R}^n, & \text{if } \kappa = 0, \\ \mathbb{H}_\kappa^n, & \text{if } \kappa < 0, \end{cases}$$

where \mathbb{S}_κ^n and \mathbb{H}_κ^n are the sphere and the hyperbolic space with their canonical metrics re-scaled by $1/\sqrt{|\kappa|}$. A *geodesic triangle* $\triangle pqr$ in X consists of three points $p, q, r \in X$ and three minimising geodesics $[pq]$, $[qr]$, $[rp]$ between those points. A *comparison triangle* in \mathbb{M}_κ^2 for $\triangle pqr$ is a geodesic triangle $\tilde{\triangle}_\kappa pqr = \triangle \tilde{p}\tilde{q}\tilde{r}$ in \mathbb{M}_κ^2 such that

$$d(\tilde{p}, \tilde{q}) = d(p, q), \quad d(\tilde{q}, \tilde{r}) = d(q, r), \quad d(\tilde{r}, \tilde{p}) = d(r, p).$$

Definition 2.3. We say that X is an *Alexandrov space with curvature bounded below by κ* , and denote it by $\text{curv}(X) \geq \kappa$, if X is complete, geodesic and satisfies the following condition:

(T $_\kappa$) For any geodesic triangle $\triangle pqr$, any comparison triangle $\tilde{\triangle}_\kappa pqr$ in \mathbb{M}_κ^2 and any point $x \in [qr]$, the corresponding point $\tilde{x} \in [\tilde{q}\tilde{r}]$ such that $d(\tilde{q}, \tilde{x}) = d(q, x)$ satisfies

$$d(p, x) \geq d(\tilde{p}, \tilde{x}).$$

Remark 2.4. Observe that condition (T $_0$) can be formulated as follows: for any geodesic triangle $p, q, r \in X$, any geodesic $\xi \in \text{Geo}(X)$ with $\xi_0 = q$ and $\xi_1 = r$,

$$(2.1) \quad d(p, \xi_t)^2 \geq (1-t)d(p, q)^2 + td(p, r)^2 - (1-t)td(q, r)^2$$

holds for any $t \in [0, 1]$. This is due to the right hand side of the inequality above being the square of $|\tilde{p} - \tilde{\xi}_t|$ in the comparison triangle $\tilde{\triangle}_0 pqr$.

Remark 2.5. Condition (T $_\kappa$) is equivalent to the following:

(A $_\kappa$) For any $p \in X$ and any $\xi^1, \xi^2 \in \text{Geo}(X)$ such that $\xi_0^1 = \xi_0^2 = p$, the function $(s, t) \mapsto \tilde{\angle}_\kappa \xi_s^1 p \xi_t^2$ is non-increasing in both s and t , where $\tilde{\angle}_\kappa \xi_s^1 p \xi_t^2$ denotes the angle at \tilde{p} in the comparison triangle $\tilde{\triangle}_\kappa \xi_s^1 p \xi_t^2$.

Condition (A_κ) implies that the *angle* between $\xi^1, \xi^2 \in \text{Geo}(X)$ with $\xi_0^1 = \xi_0^2$, given by

$$\angle(\xi^1, \xi^2) = \lim_{s,t \rightarrow 0} \widetilde{\angle}_\kappa \xi_s^1 p \xi_t^2$$

is well-defined. Geodesics that make an angle zero determine an equivalence class called *geodesic direction*. The set of geodesic directions at a point $p \in X$ is denoted by Σ'_p . When equipped with the angle metric \angle , the set Σ'_p is a metric space. The completion of (Σ'_p, \angle) is called the *space of directions of X at p* , and is denoted by Σ_p . Note that in a closed Riemannian manifold the space of directions at any point is isometric to the unit sphere in the tangent space to the manifold at the given point.

2.2. Optimal transport. We now recall definitions and classical results from optimal transport (see [2, 25, 28] for detailed expositions).

Let X be a metric space and $\mathcal{P}(X)$ be the set of Borel probability measures on X . The *support* of a Borel measure μ on X , denoted by $\text{supp}(\mu)$, is the smallest closed set $E \subset X$ such that $\mu(X \setminus E) = 0$. For any Borel map $T: X \rightarrow Y$ between metric spaces, the *push-forward map* $T_\# : \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$ is given by

$$T_\# \mu(E) = \mu(T^{-1}(E))$$

for any Borel set $E \subset Y$.

Definition 2.6. Given $\mu, \nu \in \mathcal{P}(X)$, we say that $\gamma \in \mathcal{P}(X \times X)$ is a *transport plan* between μ and ν if $\pi_\#^1 \gamma = \mu$ and $\pi_\#^2 \gamma = \nu$, where $\pi^1, \pi^2: X \times X \rightarrow X$ are the coordinate maps. The set of transport plans between μ and ν is denoted by $\text{Adm}(\mu, \nu)$. For any $p \in [1, \infty)$, the *L^p -Wasserstein metric* between $\mu, \nu \in \mathcal{P}(X)$ is defined as

$$(2.2) \quad W_p(\mu, \nu) = \inf_{\gamma \in \text{Adm}(\mu, \nu)} \left(\int d(x, y)^p d\gamma(x, y) \right)^{1/p}.$$

Any minimiser γ for (2.2) is an *optimal plan* between μ and ν . The set of optimal plans between μ and ν is denoted by $\text{Opt}(\mu, \nu)$.

Theorem 2.7. *Let X be a complete and separable metric space, and $\mu, \nu \in \mathcal{P}(X)$ such that*

$$(2.3) \quad \int_X d(x, x_0)^p d\mu(x), \int_X d(y, x_0)^p d\nu(y) < \infty$$

for some (and therefore any) $x_0 \in X$. Then $\text{Opt}(\mu, \nu) \neq \emptyset$. Moreover, W_p defines a metric in $\mathcal{P}_p(X)$, the set of measures in $\mathcal{P}(X)$ satisfying (2.3).

The following result, commonly known in the optimal transport jargon as the *gluing lemma*, plays a role in the proof of Theorem 2.7, and will be useful later on.

Theorem 2.8. *Let $\mu^1, \mu^2, \mu^3 \in \mathcal{P}(X)$, $\gamma^{12} \in \text{Adm}(\mu^1, \mu^2)$, and $\gamma^{23} \in \text{Adm}(\mu^2, \mu^3)$. Then there exists $\gamma^{123} \in \mathcal{P}(X \times X \times X)$ such that*

$$\begin{aligned} \pi_\#^{12}(\gamma^{123}) &= \gamma^{12}, \\ \pi_\#^{23}(\gamma^{123}) &= \gamma^{23}. \end{aligned}$$

More generally, if $\Gamma^1 \in \mathcal{P}(\mathcal{X}^1)$, $\Gamma^2 \in \mathcal{P}(\mathcal{X}^2)$, and $F^i: \mathcal{X}^i \rightarrow \mathcal{X}$, $i = 1, 2$, are measurable maps such that $F_\#^1 \Gamma^1 = F_\#^2 \Gamma^2$, then there exists

$$\hat{\Gamma} \in \mathcal{P}(\{(x_1, x_2) \in \mathcal{X}^1 \times \mathcal{X}^2 : F^1(x_1) = F^2(x_2)\})$$

such that $\pi_\#^i \hat{\Gamma} = \Gamma^i$, $i = 1, 2$.

It is also possible to characterise optimal plans in terms of cyclical monotonicity and the existence of Kantorovich potentials.

Definition 2.9. We say that a set $\Gamma \subset X \times X$ is *c-cyclically monotone* if, for any $n \in \mathbb{N}$, any $\{(x_i, y_i)\}_{i=1}^n \subset \Gamma$ and any permutation σ of $\{1, \dots, n\}$,

$$\sum_{i=1}^n d(x_i, y_i)^p \leq \sum_{i=1}^n d(x_i, y_{\sigma(i)})^p$$

holds. The *c-transform* of a function $\phi: X \rightarrow \mathbb{R} \cup \{-\infty\}$ is given by

$$\phi^c(y) = \inf_{x \in X} \{d(x, y)^p - \phi(x)\}.$$

We say that a function $\phi: X \rightarrow \mathbb{R} \cup \{-\infty\}$ is *c-concave* if there exists a function $\psi: X \rightarrow \mathbb{R} \cup \{-\infty\}$ such that $\phi(x) = \psi^c(x)$.

The *c-superdifferential* of a c-concave function ϕ is the set

$$\partial_+^c \phi = \{(x, y) \in X \times X : c(x, y) = \phi(x) + \phi^c(y)\}.$$

Theorem 2.10. *Let X be a complete and separable metric space, $\mu, \nu \in \mathcal{P}_p(X)$, and $\gamma \in \text{Adm}(\mu, \nu)$. Then the following conditions are equivalent:*

- (1) $\gamma \in \text{Opt}(\mu, \nu)$;
- (2) $\text{supp}(\gamma)$ is c-cyclically monotone;
- (3) There is a c-concave function ϕ (known as a Kantorovich potential of γ) with $\max\{0, \phi\} \in L^1(\mu)$ and $\text{supp}(\gamma) \subset \partial_+^c \phi$.

A sequence $\{\mu\}_{n \in \mathbb{N}}$ of non-negative Borel measures in a metric space X is *weakly convergent* to μ if

$$\int_X f d\mu_n \rightarrow \int_X f d\mu \quad \text{for any } f \in \mathcal{C}_b(X),$$

where $\mathcal{C}_b(X)$ is the set of continuous and bounded functions on X . On the other hand, $\{\mu_n\}_{n \in \mathbb{N}}$ is *vaguely convergent* to μ , and we write $\mu_n \xrightarrow{v} \mu$, whenever

$$\int_X f d\mu_n \rightarrow \int_X f d\mu \quad \text{for any } f \in \mathcal{C}_c(X),$$

where $\mathcal{C}_c(X)$ is the set of compactly supported continuous functions on X .

Lemma 2.11 below is a particular case of the classical Prokhorov's theorem. Lemmas 2.12 and 2.13 below are analogous results about the vague topology (see [21] for details and more general formulations).

Lemma 2.11. *Let \mathcal{F} be a set of non-negative Borel measures with finite mass (i.e. $\mu_n(X) < \infty$ for all $n \in \mathbb{N}$) on a complete and separable metric space X . Then \mathcal{F} is weakly precompact (i.e. every sequence in \mathcal{F} has a weakly convergent subsequence) if and only if the following conditions hold:*

- (1) \mathcal{F} has uniformly bounded total variation, i.e. $\sup_{\mu \in \mathcal{F}} \mu(X) < \infty$,
- (2) \mathcal{F} is tight, i.e. for any $\varepsilon > 0$ there exist a compact set $K \subset X$ such that $\sup_{\mu \in \mathcal{F}} \mu(X \setminus K) < \varepsilon$.

Lemma 2.12. *Let \mathcal{F} be a set of non-negative Radon measures on a complete, separable and locally compact metric space X . Then \mathcal{F} is vaguely precompact if and only if*

$$\sup\{\gamma(K) : \gamma \in \mathcal{F}\} < \infty$$

for any compact $K \subset X$.

Lemma 2.13. *Let $\{\gamma_n\}_{n \in \mathbb{N}}$ be a sequence of non-negative Radon measures on a complete, separable and locally compact metric space X . Then the following conditions are equivalent:*

(1) $\gamma_n \xrightarrow{v} \gamma$ for some $\gamma \in \mathcal{M}(X)$.

(2) For any bounded open $U \subset X$ and any bounded closed $F \subset X$,

$$\gamma(U) \leq \liminf_{n \rightarrow \infty} \gamma_n(U)$$

and

$$\gamma(F) \geq \limsup_{n \rightarrow \infty} \gamma_n(F).$$

3. OPTIMAL PARTIAL TRANSPORT FOR METRIC PAIRS

A *metric pair* is an ordered pair (X, A) where X is a metric space and $A \subseteq X$ is closed and non-empty. Moreover, we will only consider metric pairs where X is complete, separable and proper, the latter meaning that closed and bounded sets in X are compact.

We denote by $\mathcal{M}(X)$ the set of non-negative Radon measures on X , i.e. inner regular Borel measures that are finite on compact sets. We endow $\mathcal{M}(X)$ with the vague topology.

Given a metric pair (X, A) , let $\mathcal{M}(X, A)$ be the set of non-negative Radon measures on X supported in $\Omega = X \setminus A$. Let $p \in [1, \infty)$ and define

$$\mathcal{M}_p(X, A) = \left\{ \mu \in \mathcal{M}(X, A) : \int_X d(x, A)^p d\mu(x) < \infty \right\}.$$

Given $\mu, \nu \in \mathcal{M}_p(X, A)$, we define

$$(3.1) \quad \text{Wb}_p(\mu, \nu) = \inf_{\gamma \in \text{Adm}(\mu, \nu)} \left(\int_{X \times X} d(x, y)^p d\gamma(x, y) \right)^{1/p}$$

where $\text{Adm}(\mu, \nu)$ is the set of measures $\gamma \in \mathcal{M}(X \times X)$ whose marginals restricted to Ω are μ and ν , i.e. the following equations hold

$$(3.2) \quad \pi_{\#}^1(\gamma)|_{\Omega} = \mu, \quad \pi_{\#}^2(\gamma)|_{\Omega} = \nu,$$

where $\pi^1, \pi^2: X \times X \rightarrow X$ are the projections onto the first and second factor.

Remark 3.1. Given $\gamma \in \mathcal{M}(X \times X)$, we denote

$$C(\gamma) = \int_{X \times X} d(x, y)^p d\gamma(x, y)$$

and $\gamma_R^S = \gamma|_{R \times S}$ for any $R, S \subset X$. In particular,

$$\gamma = \gamma_{\Omega}^{\Omega} + \gamma_{\Omega}^A + \gamma_A^{\Omega} + \gamma_A^A.$$

Moreover, it is easy to check that if γ satisfies (3.2) then $\gamma - \gamma_A^A$ does as well, and clearly

$$C(\gamma - \gamma_A^A) \leq C(\gamma).$$

Therefore, any minimiser γ of (3.1) satisfies that γ_A^A is supported in the diagonal of $A \times A$, i.e. the set $\Delta(A \times A) = \{(a, a) : a \in A\}$.

Remark 3.2. Regarding the terminology of partial transport plans, this comes from the fact that, whenever $\gamma \in \mathcal{M}(X \times X)$ satisfies (3.2), the measure γ_Ω^Ω can be regarded as a transport plan between measures $\tilde{\mu} \leq \mu$ and $\tilde{\nu} \leq \nu$ with the same total mass, therefore effectively transporting part of the mass of μ into part of the mass of ν .

It is clear that the zero measure belongs to $\mathcal{M}_p(X, A)$, and for any $\mu \in \mathcal{M}_p(X, A)$ we have

$$(3.3) \quad \text{Wb}_p^p(\mu, 0) = \int_X d(x, A)^p d\mu(x).$$

This is a consequence of the following lemma.

Lemma 3.3. *Let (X, A) be a metric pair, and assume that X is complete, separable and proper. Then there exists a measurable map $\text{proj}_A: X \rightarrow A$ such that $d(x, \text{proj}_A(x)) = d(x, A)$ for all $x \in X$.*

Proof. Since A is closed and X is complete and separable, it follows that A is complete and separable endowed with the restricted metric, and since X is proper, the set

$$E = \{(x, y) \in X \times A : d(x, y) = d(x, A)\}$$

is σ -compact. Moreover, by the continuity of d , it follows that E is closed. Therefore, by [3, Theorem 1], the claim follows. \square

We can consider now the partial transport plan $\gamma = (\text{id}, \text{proj}_A)_\# \mu \in \text{Adm}(\mu, 0)$, which satisfies

$$C(\gamma) = \int_X d(x, \text{proj}_A(x))^p d\mu(x) = \int_X d(x, A)^p d\mu(x).$$

On the other hand, for any $\tilde{\gamma} \in \text{Adm}(\mu, 0)$ we have that $\pi_{\#}^2(\tilde{\gamma})|_\Omega = 0$, which implies that $\tilde{\gamma}(X \times \Omega) = 0$. Therefore, we have

$$\begin{aligned} C(\tilde{\gamma}) &= \int_{X \times X} d(x, y)^p d\tilde{\gamma}(x, y) \\ &= \int_{X \times A} d(x, y)^p d\tilde{\gamma}(x, y) \\ &\geq \int_{X \times A} d(x, A)^p d\tilde{\gamma}(x, y) \\ &= \int_X d(x, A)^p d\mu(x). \end{aligned}$$

This proves equation (3.3).

The following lemma is a natural generalisation of the well-known gluing lemma for probability measures.

Lemma 3.4. *Let $\mu^1, \mu^2, \mu^3 \in \mathcal{M}_p(X, A)$, $\gamma^{12} \in \text{Adm}(\mu^1, \mu^2)$, and $\gamma^{23} \in \text{Adm}(\mu^2, \mu^3)$. Then there exists $\gamma^{123} \in \mathcal{M}(X \times X \times X)$ such that*

$$\begin{aligned} \pi_{\#}^{12}(\gamma^{123}) &= \gamma^{12} - (\gamma^{12})_A^A + \sigma^{12}, \\ \pi_{\#}^{23}(\gamma^{123}) &= \gamma^{23} - (\gamma^{23})_A^A + \sigma^{23}, \end{aligned}$$

where σ^{12} and σ^{23} are both concentrated in $\Delta(A \times A)$.

Proof. From the hypothesis, we can see that

$$\mu^2 = \pi_{\#}^2(\gamma^{12})|_{\Omega} = \pi_{\#}^2((\gamma^{12})_X^{\Omega} + (\gamma^{12})_{\Omega}^A)|_{\Omega} = \pi_{\#}^2((\gamma^{12})_X^{\Omega}).$$

Analogously, $\mu^2 = \pi_{\#}^1((\gamma^{23})_{\Omega}^X)$. Therefore, by applying the classical gluing lemma, since μ^2 is a Radon measure, we can find $\tilde{\gamma}^{123} \in \mathcal{M}(X \times \Omega \times X)$ such that

$$\begin{aligned}\pi_{\#}^{12}(\tilde{\gamma}^{123}) &= (\gamma^{12})_X^{\Omega}, \\ \pi_{\#}^{23}(\tilde{\gamma}^{123}) &= (\gamma^{23})_{\Omega}^X.\end{aligned}$$

We define

$$\begin{aligned}\tilde{\sigma}^{12} &= (\pi^1, \pi^1, \pi^2)_{\#}((\gamma^{23})_A^{\Omega}) \in \mathcal{M}(A \times A \times X), \\ \tilde{\sigma}^{23} &= (\pi^1, \pi^2, \pi^2)_{\#}((\gamma^{12})_{\Omega}^A) \in \mathcal{M}(X \times A \times A), \\ \gamma^{123} &= \tilde{\gamma}^{123} + \tilde{\sigma}^{12} + \tilde{\sigma}^{23} \in \mathcal{M}(X \times X \times X), \\ \sigma^{12} &= (\pi^1, \pi^1)_{\#}((\gamma^{23})_A^{\Omega}) \in \mathcal{M}(\Delta(A \times A)), \\ \sigma^{23} &= (\pi^2, \pi^2)_{\#}((\gamma^{12})_{\Omega}^A) \in \mathcal{M}(\Delta(A \times A)).\end{aligned}$$

We can check that these measures work. Indeed,

$$\begin{aligned}\pi_{\#}^{12}(\gamma^{123}) &= \pi_{\#}^{12}(\tilde{\gamma}^{123}) + \pi_{\#}^{12}(\tilde{\sigma}^{12}) + \pi_{\#}^{12}(\tilde{\sigma}^{23}) \\ &= (\gamma^{12})_X^{\Omega} + \pi_{\#}^{12}(\pi^1, \pi^1, \pi^2)_{\#}((\gamma^{23})_A^{\Omega}) + \pi_{\#}^{12}(\pi^1, \pi^2, \pi^2)_{\#}((\gamma^{12})_{\Omega}^A) \\ &= (\gamma^{12})_X^{\Omega} + (\pi^1, \pi^1)_{\#}((\gamma^{23})_A^{\Omega}) + (\pi^1, \pi^2)_{\#}((\gamma^{12})_{\Omega}^A) \\ &= (\gamma^{12})_X^{\Omega} + \sigma^{12} + (\gamma^{12})_{\Omega}^A \\ &= \gamma^{12} - (\gamma^{12})_A^A + \sigma^{12}.\end{aligned}$$

Similarly with $\pi_{\#}^{23}(\gamma^{123})$. □

Theorem 3.5. *Let (X, A) be a metric pair, where X is complete, separable and proper. Then, for any $\mu, \nu \in \mathcal{M}_p(X, A)$, the set*

$$\text{Opt}(\mu, \nu) = \{\gamma \in \text{Adm}(\mu, \nu) : C(\gamma) = \text{Wb}_p^p(\mu, \nu)\}$$

is non-empty. Moreover, the set

$$\{\gamma \in \text{Opt}(\mu, \nu) : \gamma_A^A = 0\}$$

is vaguely compact.

Proof. Let $\mu, \nu \in \mathcal{M}_p(X, A)$ and, for every $k \in \mathbb{N}$, let $\gamma_k \in \text{Adm}(\mu, \nu)$ be such that

$$(3.4) \quad \text{Wb}_p(\mu, \nu) \leq C(\gamma_k)^{1/p} < \text{Wb}_p(\mu, \nu) + 1/k.$$

Moreover, we can assume that $(\gamma_k)_A^A = 0$, due to Remark 3.1. In other words, we will regard each γ_k as a measure in $\mathcal{M}(X \times X \setminus A \times A)$. It is also important to remark that, since $X \times X$ is complete and separable, then $X \times X \setminus A \times A$, being an open subset, admits a complete and separable metric [6], which is sufficient to apply Lemmas 2.12 and 2.13.

Now observe that $\{\gamma_k\}_{k \in \mathbb{N}}$ is a vaguely precompact subset of $\mathcal{M}(X \times X \setminus A \times A)$. Indeed, for any compact $K \subset X \times X \setminus A \times A$ there are compact sets $K', K'' \subset \Omega$ such that $K \subset K' \times X \cup X \times K''$, therefore

$$\gamma_k(K) \leq \gamma_k(K' \times X) + \gamma_k(X \times K'') = \mu(K') + \nu(K''),$$

and the claim follows due to Lemma 2.12.

As a consequence of the previous observation, up to passing to a subsequence, we can assume that there exists $\gamma \in \mathcal{M}(X \times X \setminus A \times A)$ such that $\gamma_k \xrightarrow{v} \gamma$. Moreover, we can see that $\gamma \in \text{Adm}(\mu, \nu)$. Indeed, if $f \in \mathcal{C}_c(\Omega)$ then

$$(3.5) \quad \int_{\Omega} f \, d\pi_{\#}^1 \gamma = \int_{\Omega \times X} f \circ \pi^1 \, d\gamma = \lim_{k \rightarrow \infty} \int_{\Omega \times X} f \circ \pi^1 \, d\gamma_k = \int_{\Omega} f \, d\mu.$$

There is a slight technical issue in the previous chain of equations (which is the same as in the proof of [14, Proposition 3.2]), when X is unbounded. Namely, $f \circ \pi^1$ is not compactly supported, and therefore vague convergence alone is not sufficient to get the second equation above, whenever X is unbounded. However, this can be fixed by proving that, for any compact $C \subset \Omega$, the sequence $\{(\gamma_k)_C^X\}_{k \in \mathbb{N}}$ is weakly convergent. Indeed, if X is unbounded, take $p_0 \in X$ and $0 < r < R$ such that $C \subset \overline{B}_r(p_0) \subset \overline{B}_R(p_0)$. By Hölder's inequality, we get

$$(3.6) \quad \left| \int_{C \times (X \setminus \overline{B}_R(p_0))} d(y, p_0) - d(x, p_0) \, d\gamma_k(x, y) \right|^p \leq \gamma_k(C \times (X \setminus \overline{B}_R(p_0)))^{p-1} \int_{C \times (X \setminus \overline{B}_R(p_0))} d(x, y)^p \, d\gamma_k(x, y),$$

whereas

$$(3.7) \quad \left| \int_{C \times (X \setminus \overline{B}_R(p_0))} d(y, p_0) - d(x, p_0) \, d\gamma_k(x, y) \right|^p \geq (R - r)^p \gamma_k(C \times (X \setminus \overline{B}_R(p_0)))^p$$

since $\text{dist}(\overline{B}_r(p_0), X \setminus \overline{B}_R(p_0)) \geq R - r$. Combining (3.6), (3.7) and (3.4), we get

$$\gamma_k(C \times (X \setminus \overline{B}_R(p_0))) \leq \frac{\text{Wb}_p(\mu, \nu) + 1}{(R - r)^p},$$

which can be made arbitrarily small, uniformly over $k \in \mathbb{N}$, by fixing r and letting R tend to infinity. This proves that $\{(\gamma_k)_C^X\}_{k \in \mathbb{N}}$ is tight. Moreover, since $\gamma_k \in \text{Adm}(\mu, \nu)$, each $(\gamma_k)_C^X$ has total mass $\mu(C) < \infty$, therefore $\{(\gamma_k)_C^X\}_{k \in \mathbb{N}}$ has uniformly bounded total variation. By Lemma 2.11, the claim follows.

By applying the previous claim to $C = \text{supp}(f)$, whenever $f \in \mathcal{C}_c(\Omega)$, we get that $\{(\gamma_k)_{\text{supp}(f)}^X\}_{k \in \mathbb{N}}$ weakly converges to $\gamma_{\text{supp}(f)}^X$, therefore justifying equation (3.5). We thus get that $\pi_{\#}^1 \gamma|_{\Omega} = \mu$, and analogously $\pi_{\#}^2 \gamma|_{\Omega} = \nu$.

Finally, by Lemma 2.13 applied to the sequence $\{d(\cdot, \cdot)^p \gamma_k\}_{k \in \mathbb{N}}$, which is vaguely convergent to $d(\cdot, \cdot)^p \gamma$, and any bounded open set $U \subset X \times X$, we get

$$C(\gamma_U^U) \leq \liminf_{k \rightarrow \infty} C((\gamma_k)_U^U) \leq \text{Wb}_p^p(\mu, \nu).$$

By the monotone convergence theorem,

$$C(\gamma) \leq \text{Wb}_p^p(\mu, \nu),$$

which implies that $\gamma \in \text{Opt}(\mu, \nu)$.

Regarding the second part of the theorem, observe that if $\{\gamma_k\}_{k \in \mathbb{N}} \subset \text{Opt}(\mu, \nu)$ and $\gamma_k(A \times A) = 0$ for all $k \in \mathbb{N}$ then, by the previous arguments, up to passing to a subsequence, $\gamma_k \xrightarrow{v} \gamma$ for some $\gamma \in \mathcal{M}(X \times X \setminus A \times A)$. The fact that $\gamma \in \text{Opt}(\mu, \nu)$ also follows from the arguments above. \square

Theorem 3.6. *Let (X, A) be a metric pair, where X is complete, separable and proper. Then the function Wb_p is a metric on $\mathcal{M}_p(X, A)$. Moreover, Wb_p is lower semi-continuous with respect to the vague topology.*

Proof. Let $\mu, \nu \in \mathcal{M}_p(X, A)$. Then the measure

$$\gamma = (\text{id}, \text{proj}_A)_\#(\mu) + (\text{proj}_A, \text{id})_\#(\nu)$$

is in $\text{Adm}(\mu, \nu)$, and it satisfies

$$C(\gamma) = \int_X d(x, A)^p d\mu(x) + \int_X d(y, A)^p d\nu(y) < \infty.$$

It is also clear that Wb_p is non-negative and symmetric. Moreover, due to Proposition 3.5, $\text{Wb}_p(\mu, \nu) = 0$ if and only if $C(\gamma) = 0$ for some $\gamma \in \text{Adm}(\mu, \nu)$, which is equivalent to

$$\int_{X \times X} d(x, y)^p d\gamma(x, y) = 0.$$

However, this is the same as $\text{supp}(\gamma) \subset \Delta(X \times X)$, which in turn is equivalent to $\pi_{\#}^1 \gamma = \pi_{\#}^2 \gamma$. The latter implies that $\mu = \nu$. Conversely, if $\mu = \nu$ then we can take $\gamma = (\text{id}, \text{id})_\# \mu \in \text{Adm}(\mu, \nu)$, which satisfies $C(\gamma) = 0$.

For the triangle inequality, we need Lemma 3.4. Indeed, let $\mu^1, \mu^2, \mu^3 \in \mathcal{M}_p(X, A)$ and choose $\gamma^{12} \in \text{Opt}(\mu^1, \mu^2)$ and $\gamma^{23} \in \text{Opt}(\mu^2, \mu^3)$ (which we can do thanks to Proposition 3.5). By Lemma 3.4, there is $\gamma^{123} \in \mathcal{M}(X \times X \times X)$ such that

$$\begin{aligned} \pi_{\#}^{12} \gamma^{123} &= \gamma^{12} + \sigma^{12}, \\ \pi_{\#}^{23} \gamma^{123} &= \gamma^{23} + \sigma^{23}, \end{aligned}$$

with $\text{supp}(\sigma^{12}), \text{supp}(\sigma^{23}) \subset \Delta(A \times A)$. In particular, $\pi_{\#}^1(\gamma^{123})|_{\Omega} = \mu^1$ and $\pi_{\#}^3(\gamma^{123})|_{\Omega} = \mu^3$, therefore $\pi_{\#}^{13}(\gamma^{123}) \in \text{Adm}(\mu^1, \mu^3)$. This implies

$$\begin{aligned} \text{Wb}_p(\mu^1, \mu^3) &\leq C(\pi_{\#}^{13}(\gamma^{123}))^{1/p} \\ &= \|d \circ \pi^{1,3}\|_{L^p(\gamma^{123})} \\ &\leq \|d \circ \pi^{1,2} + d \circ \pi^{2,3}\|_{L^p(\gamma^{123})} \\ &\leq \|d \circ \pi^{1,2}\|_{L^p(\gamma^{123})} + \|d \circ \pi^{2,3}\|_{L^p(\gamma^{123})} \\ &= \|d\|_{L^p(\gamma^{12} + \sigma^{12})} + \|d\|_{L^p(\gamma^{23} + \sigma^{23})} \\ &= C(\gamma^{12}(x, y))^{1/p} + C(\gamma^{23}(y, z))^{1/p} \\ &= \text{Wb}_p(\mu^1, \mu^2) + \text{Wb}_p(\mu^2, \mu^3). \end{aligned}$$

To prove that Wb_p is lower semi-continuous with respect to the vague topology, let $\mu_n \xrightarrow{v} \mu$ and $\nu_n \xrightarrow{v} \nu$. If $\liminf_{n \rightarrow \infty} \text{Wb}_p(\mu_n, \nu_n) = \infty$, the result follows trivially. Otherwise, up to passing to a subsequence, we can assume that

$$\liminf_{n \rightarrow \infty} \text{Wb}_p(\mu_n, \nu_n) = \lim_{n \rightarrow \infty} \text{Wb}_p(\mu_n, \nu_n).$$

Take $\gamma_n \in \text{Opt}(\mu_n, \nu_n)$, and without loss of generality, we can assume that $(\gamma_n)_A^A = 0$. By similar arguments to those in the proof of Proposition 3.5, $\{\gamma_n\}_{n \in \mathbb{N}}$ is vaguely precompact, therefore, up to passing to another subsequence, we can assume that $\gamma_n \xrightarrow{v} \gamma$ for some $\gamma \in \mathcal{M}(X \times X \setminus A \times A)$. Moreover, we can also prove that for any compact $C \subset \Omega$, $\{(\gamma_n)_C^X\}_{n \in \mathbb{N}}$ is tight and each $(\gamma_n)_C^X$ has total mass bounded above by $\mu(C)$ due to Lemma 2.13. Therefore, $\{(\gamma_n)_C^X\}_{n \in \mathbb{N}}$ is weakly convergent

to γ_C^X . It follows that $\pi_{\#}^1 \gamma|_{\Omega} = \mu$, and analogously $\pi_{\#}^2 \gamma|_{\Omega} = \nu$. Therefore $\gamma \in \text{Adm}(\mu, \nu)$ and due to Lemma 2.13 applied to the measures $\{d(\cdot, \cdot)^p \gamma_n\}_{n \in \mathbb{N}}$, which vaguely converge to $d(\cdot, \cdot)^p \gamma$, and bounded open sets $U \subset X \times X$, we get

$$C(\gamma_U^U) \leq \liminf_{n \rightarrow \infty} C((\gamma_n)_U^U) \leq \lim_{n \rightarrow \infty} \text{Wb}_p^p(\mu_n, \nu_n)$$

and by the monotone convergence theorem, the claim follows. \square

4. CHARACTERISATION OF OPTIMAL PARTIAL TRANSPORT

We now move on to study a characterisation of optimal partial transport plans analogous to that of optimal plans in the classic sense. Let us define $c(x, y) = d(x, y)^p$ and $\tilde{c}(x, y) = \min\{d(x, y)^p, d(x, A)^p + d(y, A)^p\}$, and

$$\mathcal{S} = \{(x, y) \in X \times X : \tilde{c}(x, y) = c(x, y)\}.$$

Proposition 4.1. *Let $\gamma \in \mathcal{M}(X \times X)$ be a measure satisfying*

$$\int d(x, A)^p + d(y, A)^p d\gamma(x, y) < \infty.$$

Then the following are equivalent:

- (1) $\gamma \in \text{Opt}(\pi_{\#}^1(\gamma)|_{\Omega}, \pi_{\#}^2(\gamma)|_{\Omega})$.
- (2) γ is concentrated on \mathcal{S} and the set $\text{supp}(\gamma) \cup A \times A$ is \tilde{c} -cyclically monotone.
- (3) there is a c -concave function ϕ such that both ϕ and ϕ^c are identically 0 on A and $\text{supp}(\gamma) \subset \partial_+^c \phi$.

Moreover, $d(x, y) = d(x, A)$ for γ_{Ω}^A -a.e. (x, y) and $d(x, y) = d(y, A)$ for γ_A^{Ω} -a.e. (x, y) , whenever γ is optimal.

Proof. We start proving that (1) implies (2). We denote $\bar{\mu} = \pi_{\#}^1 \gamma$, $\bar{\nu} = \pi_{\#}^2 \gamma$, $\mu = \bar{\mu}|_{\Omega}$ and $\nu = \bar{\nu}|_{\Omega}$, and define

$$\tilde{\gamma} = \gamma|_{\mathcal{S}} + (\pi^1, P \circ \pi^1)_{\#} \gamma|_{X \times X \setminus \mathcal{S}} + (P \circ \pi^2, \pi^2)_{\#} \gamma|_{X \times X \setminus \mathcal{S}}.$$

It is clear that $\tilde{\gamma} \in \text{Adm}(\mu, \nu)$ and

$$C(\tilde{\gamma}) = \int_{\mathcal{S}} d(x, y)^p d\gamma(x, y) + \int_{X \times X \setminus \mathcal{S}} d(x, A)^p + d(y, A)^p d\gamma(x, y) \leq C(\gamma)$$

with strict inequality if and only if $\gamma(X \times X \setminus \mathcal{S}) > 0$. Since $\gamma \in \text{Opt}(\mu, \nu)$, we get that $C(\tilde{\gamma}) = C(\gamma)$, which implies that $\gamma(X \times X \setminus \mathcal{S}) = 0$, i.e. γ is concentrated on \mathcal{S} . In particular,

$$C(\gamma) = \int_{X \times X} \tilde{c}(x, y) d\gamma(x, y).$$

Now suppose that $\eta \in \mathcal{M}(X \times X)$ satisfies $\pi_{\#}^1(\eta) = \bar{\mu}$ and $\pi_{\#}^2(\eta) = \bar{\nu}$. Then we can define $\tilde{\eta}$ in analogy to how we defined $\tilde{\gamma}$. Clearly $\tilde{\eta} \in \text{Adm}(\mu, \nu)$, and

$$C(\tilde{\eta}) = \int_{X \times X} \tilde{c}(x, y) d\eta(x, y).$$

In particular,

$$\int_{X \times X} \tilde{c}(x, y) d\gamma(x, y) = C(\gamma) \leq C(\tilde{\eta}) = \int_{X \times X} \tilde{c}(x, y) d\eta(x, y)$$

for any η with the same marginals as γ . In other words, γ is an optimal plan between $\bar{\mu}$ and $\bar{\nu}$ with respect to the cost function \tilde{c} in the usual sense. The classical theory implies that $\text{supp}(\gamma)$ is \tilde{c} -cyclically monotone. Moreover, given $\{(x_i, y_i)\}_{i=1}^n \subset \text{supp}(\gamma) \cup A \times A$ and $\sigma \in \Sigma_n$, we have

$$\begin{aligned} \sum_{i=1}^n \tilde{c}(x_i, y_{\sigma(i)}) &= \sum_{\substack{(x_i, y_i) \in \text{supp}(\gamma) \\ (x_{\sigma(i)}, y_{\sigma(i)}) \in \text{supp}(\gamma)}} \tilde{c}(x_i, y_{\sigma(i)}) + \sum_{\substack{(x_i, y_i) \in \text{supp}(\gamma) \\ (x_{\sigma(i)}, y_{\sigma(i)}) \in A \times A \setminus \text{supp}(\gamma)}} \tilde{c}(x_i, y_{\sigma(i)}) \\ &+ \sum_{\substack{(x_i, y_i) \in A \times A \setminus \text{supp}(\gamma) \\ (x_{\sigma(i)}, y_{\sigma(i)}) \in \text{supp}(\gamma)}} \tilde{c}(x_i, y_{\sigma(i)}) + \sum_{\substack{(x_i, y_i) \in A \times A \setminus \text{supp}(\gamma) \\ (x_{\sigma(i)}, y_{\sigma(i)}) \in A \times A \setminus \text{supp}(\gamma)}} \tilde{c}(x_i, y_{\sigma(i)}) \end{aligned}$$

where it is clear that the number of summands in the second and the third sums are the same, whereas the fourth sum vanishes since $\tilde{c}(x, y) = 0$ for any $(x, y) \in A \times A$. Let $\{j_i\}_{i=1}^p$, $\{k_i\}_{i=1}^q$ and $\{l_i\}_{i=1}^q$ be the sets of indices for the first, second and third sums, respectively, and define a permutation $\tilde{\sigma}$ of the indices $\{j_1, \dots, j_p, k_1, \dots, k_q\} = \{i : (x_i, y_i) \in \text{supp}(\gamma)\}$ by $\tilde{\sigma}(j_r) = \sigma(j_r)$ for $r = 1, \dots, p$ and $\tilde{\sigma}(k_s) = \sigma(l_s)$ for $s = 1, \dots, q$. Then

$$\begin{aligned} \sum_{i=1}^n \tilde{c}(x_i, y_{\sigma(i)}) &= \sum_{i=1}^p \tilde{c}(x_{j_i}, y_{\sigma(j_i)}) + \sum_{i=1}^q \tilde{c}(x_{k_i}, y_{\sigma(k_i)}) + \tilde{c}(x_{l_i}, y_{\sigma(l_i)}) \\ &= \sum_{i=1}^p \tilde{c}(x_{j_i}, y_{\sigma(j_i)}) + \sum_{i=1}^q \tilde{c}(x_{k_i}, y_{\sigma(k_i)}) + \tilde{c}(y_{\sigma(k_i)}, x_{l_i}) + \tilde{c}(x_{l_i}, y_{\sigma(l_i)}) \\ &\geq \sum_{i=1}^p \tilde{c}(x_{j_i}, y_{\sigma(j_i)}) + \sum_{i=1}^q \tilde{c}(x_{k_i}, y_{\sigma(l_i)}) \\ &= \sum_{(x_i, y_i) \in \text{supp}(\gamma)} \tilde{c}(x_i, y_{\tilde{\sigma}(i)}) \\ &\geq \sum_{(x_i, y_i) \in \text{supp}(\gamma)} \tilde{c}(x_i, y_i) \\ &= \sum_{i=1}^n \tilde{c}(x_i, y_i) \end{aligned}$$

where in the second and last equalities we have used the fact that $\tilde{c}(x, y) = 0$ for any $(x, y) \in A \times A$, whereas for the first and second inequalities we have used the easily checked inequality $\tilde{c}(x, y) + \tilde{c}(z, y) \geq \tilde{c}(x, z)$ that holds for any $x, z \in X$ and $y \in A$, and the fact that $\text{supp}(\gamma)$ is \tilde{c} -cyclically monotone. This shows (2).

Now, to prove that (2) implies (3), observe that, by the classical theory, $\text{supp}(\gamma) \cup A \times A$ being \tilde{c} -cyclically monotone implies that there exists a \tilde{c} -concave function, say ϕ , such that $\text{supp}(\gamma) \cup A \times A \subset \partial_+^{\tilde{c}} \phi$. In particular,

$$\phi(x) + \phi^{\tilde{c}}(y) = \tilde{c}(x, y) = 0$$

for any $(x, y) \in A \times A$, which implies that both ϕ and $\phi^{\tilde{c}}$ are constant on A . Since the \tilde{c} -concavity is invariant under addition of constants, we can assume that $\phi = \phi^{\tilde{c}} = 0$ on A .

On the other hand, we can see that ϕ is c -concave. Indeed, one can prove that, for any $y \in X$, the map $x \mapsto \tilde{c}(x, y)$ is c -concave. Indeed, if we define $\psi_y: X \rightarrow \mathbb{R} \cup \{-\infty\}$ by

$$\psi_y(z) = \begin{cases} -\infty & \text{if } z \in X \setminus (A \cup \{y\}) \\ 0 & \text{if } z = y \\ -d(y, A)^p & \text{if } z \in A, \end{cases}$$

then it is clear that

$$\inf_{z \in X} c(x, z) - \psi_y(z) = \min \left\{ d(x, y)^p, \inf_{z \in A} d(x, z)^p + d(y, A)^p \right\} = \tilde{c}(x, y),$$

and since ϕ is \tilde{c} -concave, there exists a function $\psi: X \rightarrow \mathbb{R} \cup \{-\infty\}$ such that $\phi = \psi^{\tilde{c}}$, that is,

$$\begin{aligned} \phi(x) &= \inf_{y \in X} \tilde{c}(x, y) - \psi(y) \\ &= \inf_{y \in X} \inf_{z \in X} c(x, z) - \psi_y(z) - \psi(y) \\ &= \inf_{z \in X} \inf_{y \in X} c(x, z) - \psi_y(z) - \psi(y) \\ &= \inf_{z \in X} c(x, z) - \sup_{y \in X} \psi_y(z) + \psi(y) \\ &= \inf_{z \in X} c(x, z) - \eta(z), \end{aligned}$$

where $\eta(z) = \sup_{y \in X} \psi_y(z) + \psi(y)$ (observe this is a well defined function $X \rightarrow \mathbb{R} \cup \{-\infty\}$ since, if $z \in A$ then

$$\sup_{y \in X} \psi_y(z) + \psi(y) = \sup_{y \in X} -d(y, A)^p + \psi(y) = - \inf_{y \in X} \tilde{c}(z, y) - \psi(y) = -\phi(z) = 0$$

and if $z \notin A$, then

$$\psi_y(z) + \psi(y) = \begin{cases} -\infty & \text{if } y \neq z \\ \psi(z) & \text{if } y = z \end{cases}$$

which implies that $\sup_{y \in X} \psi_y(z) + \psi(y) = \psi(z)$.

Moreover, if $(x, y) \in \partial_+^{\tilde{c}} \cap \mathcal{S}$ then

$$\phi^{\tilde{c}}(y) = \tilde{c}(x, y) - \phi(x) = c(x, y) - \phi(x)$$

whereas

$$\phi^{\tilde{c}}(y) \leq \tilde{c}(x', y) - \phi(x') \leq c(x', y) - \phi(x')$$

for any $x' \in X$, which means that $\phi^{\tilde{c}}(y) = \phi^c(y)$, therefore $(x, y) \in \partial_+^c \phi$. In particular, since γ is concentrated on $\partial_+^{\tilde{c}} \cap \mathcal{S}$, we get that it is also concentrated on $\partial_+^c \phi$, which implies that $\text{supp}(\gamma) \subset \partial_+^c \phi$.

On the other hand, if $x \in A$ then, since $\phi = \phi^{\tilde{c}} = 0$ on A , we get

$$\phi(x) + \phi^{\tilde{c}}(x) = \tilde{c}(x, x) = c(x, x) = 0$$

which implies that $(x, x) \in \partial_+^{\tilde{c}} \phi \cap \mathcal{S}$, therefore $(x, x) \in \partial_+^c \phi$. In particular,

$$\phi^c(x) = \phi(x) + \phi^c(x) = c(x, x) = 0$$

therefore $\phi^c = 0$ on A . This proves (3).

Finally, to prove that (3) implies (1), consider ϕ a c -concave function such that $\text{supp}(\gamma) \subset \partial_+^c \phi$ and $\phi = \phi^c = 0$ on A , and choose $\tilde{\gamma} \in \text{Adm}(\mu, \nu)$. Then, since $\pi_{\#}^1(\gamma)|_{\Omega} = \pi_{\#}^1(\tilde{\gamma})|_{\Omega}$, we get

$$\int_X \phi \, d\pi_{\#}^1(\gamma) = \int_{\Omega} \phi \, d\pi_{\#}^1(\gamma) = \int_{\Omega} \phi \, d\pi_{\#}^1(\tilde{\gamma}) = \int_X \phi \, d\pi_{\#}^1(\tilde{\gamma}).$$

Analogously with ϕ^c . Then, we get

$$\begin{aligned}
\int_{X \times X} d(x, y)^p d\gamma(x, y) &= \int_{X \times X} \phi(x) + \phi^c(y) d\gamma(x, y) \\
&= \int_X \phi(x) d\pi_{\#}^1(\gamma)(x) + \int_X \phi^c(y) d\pi_{\#}^2(\gamma)(y) \\
&= \int_X \phi(x) d\pi_{\#}^1(\tilde{\gamma})(x) + \int_X \phi^c(y) d\pi_{\#}^2(\tilde{\gamma})(y) \\
&= \int_{X \times X} \phi(x) + \phi^c(y) d\tilde{\gamma}(x, y) \\
&\leq \int_{X \times X} d(x, y)^p d\tilde{\gamma}(x, y)
\end{aligned}$$

where the last inequality is due to the inequality $\phi(x) + \phi^c(y) \leq c(x, y)$ that holds for general $(x, y) \in X \times X$. This argument implies that $\gamma \in \text{Opt}(\mu, \nu)$.

To prove (4) we only need to observe that, whenever $\gamma \in \text{Opt}(\mu, \nu)$ for some $\mu, \nu \in \mathcal{M}_p(X, A)$, then by (2) we have

$$\text{supp}(\gamma_{\Omega}^A) \subset \text{supp}(\gamma) \cap \Omega \times A \subset \mathcal{S} \cap \Omega \times A$$

and for any $(x, y) \in \mathcal{S} \cap \Omega \times A$ we have

$$d(x, A)^p \leq d(x, y)^p = \min\{d(x, y)^p, d(x, A)^p\} \leq d(x, A)^p$$

which implies $d(x, y) = d(x, A)$ for any $(x, y) \in \text{supp}(\gamma_{\Omega}^A)$. Analogously with γ_A^{Ω} . \square

5. PROPERTIES OF $\mathcal{M}_p(X, A)$

In this section we prove that $\mathcal{M}_p(X, A)$ inherits different properties from the space X . Namely, we prove that $\mathcal{M}_p(X, A)$ is complete, separable, geodesic and non-negatively curved in the Alexandrov sense whenever the underlying space X is so. Proofs are similar to the corresponding results for the classical Wasserstein spaces of probability measures, and we include them for the sake of completeness.

5.1. Completeness and separability.

Proposition 5.1. *The space $\mathcal{M}_p(X, A)$ is complete and separable.*

Proof. For the separability of $\mathcal{M}_p(X, A)$, if we choose a countable dense set $S \subset X$ and define,

$$F = \left\{ \sum_{i \in I} q_i \delta_{x_i} : x_i \in S \cap \Omega, q_i \in \mathbb{Q}_+, I \subset \mathbb{N} \text{ is finite} \right\},$$

it is easy to check that F is countable and dense in $\mathcal{M}_p(X, A)$.

To prove that $\mathcal{M}_p(X, A)$ is complete, we consider a Cauchy sequence $\{\mu_n\}_{n \in \mathbb{N}} \subset \mathcal{M}_p(X, A)$. Since $\{\text{Wb}_p(\mu_n, 0)\}_{n \in \mathbb{N}}$ is a bounded sequence in \mathbb{R} , say by some $C > 0$, then for any compact $K \subset \Omega$ we have

$$\mu_n(K) \leq \frac{1}{r^p} \int_{\Omega} d(x, A)^p d\mu_n(x) = \frac{1}{r^p} \text{Wb}_p^p(\mu_n, 0) \leq \frac{C^p}{r^p}$$

where $d(x, A) > r$ for any $x \in K$. In particular

$$\sup\{\mu_n(K) : n \in \mathbb{N}\} < \infty$$

for any compact $K \subset \Omega$, which due to Lemma 2.12, implies that $\{\mu_n : n \in \mathbb{N}\}$ is vaguely precompact. Therefore it has a subsequence $\{\mu_{n_k}\}_{k \in \mathbb{N}}$ vaguely convergent to some μ . By the lower semi continuity of Wb_p , we get that

$$\text{Wb}_p^p(\mu, 0) \leq \liminf_{k \rightarrow \infty} \text{Wb}_p^p(\mu_{n_k}, 0) < \infty,$$

therefore $\mu \in \mathcal{M}_p(X, A)$. Moreover, for any $n \in \mathbb{N}$,

$$\text{Wb}_p(\mu_n, \mu) \leq \liminf_{k \rightarrow \infty} \text{Wb}_p^p(\mu_n, \mu_{n_k}),$$

which implies

$$\lim_{n \rightarrow \infty} \text{Wb}_p(\mu_n, \mu) \leq \lim_{n \rightarrow \infty} \liminf_{k \rightarrow \infty} \text{Wb}_p^p(\mu_n, \mu_{n_k}) = 0$$

where the last inequality comes from the fact that $\{\mu_n\}_{n \in \mathbb{N}}$ is Cauchy. Therefore $\mu_n \rightarrow \mu$ in $\mathcal{M}_p(X, A)$, which implies the completeness. \square

5.2. Geodesics.

Proposition 5.2. *Let (X, A) be a complete, separable, proper, and geodesic metric pair. Then $\mathcal{M}_p(X, A)$ is geodesic as well. Furthermore, if $(\mu_t)_{t \in [0,1]}$ is a constant speed geodesic in $\mathcal{M}_p(X, A)$, then there exists a measure $\gamma \in \mathcal{M}(\text{Geo}(X))$ such that $(e_0, e_1)_\#(\gamma) \in \text{Opt}(\mu_0, \mu_1)$ and*

$$\mu_t = (e_t)_\#(\gamma^m)|_\Omega$$

for any $t = i/2^m$ with $i \in \{0, \dots, m\}$.

For the proof of Theorem 5.2 we need the following technical observation.

Lemma 5.3. *There is a measurable map $\text{GeoSel}: X \times X \rightarrow \text{Geo}(X)$ such that*

$$(e_0, e_1) \circ \text{GeoSel} = \text{id}_{X \times X}.$$

Proof. Since X and $\text{Geo}(X)$ are complete, separable and proper, the set

$$E = \{(x, y, \xi) \in X \times X \times \text{Geo}(X) : \xi_0 = x, \xi_1 = y, d(x, y) = \mathcal{L}(\xi)\}$$

is σ -compact. Moreover, by the continuity of d , the lower semi-continuity of \mathcal{L} , and the continuity of the evaluation maps e_0, e_1 , it follows that E is closed. Therefore, by [3, Theorem 1], the claim follows. \square

Proof of Proposition 5.2. Let $\mu^0, \mu^1 \in \mathcal{M}_p(X, A)$ and choose $\gamma \in \text{Opt}(\mu^0, \mu^1)$. By Lemma 5.3 there is a measurable map $\text{GeoSel}: X \times X \rightarrow \text{Geo}(X)$ such that $\text{GeoSel}(x, y)$ is a constant speed geodesic joining x and y . We define $\gamma \in \mathcal{M}(\text{Geo}(X))$ by

$$\gamma = \text{GeoSel}_\#(\gamma).$$

Observe that for any $(x, y) \in \mathcal{S}$, $t \in (0, 1)$ and $\xi \in \text{Geo}(X)$ such that $(e_0, e_1)(\xi) = (x, y)$, we have $\xi_t \in \Omega$. Indeed, if this is not the case, then we have that for some choice of (x, y) , t and ξ satisfying said conditions, $\xi_t \in A$. Therefore,

$$d(x, A)^p + d(y, A)^p \leq d(x, \xi_t)^p + d(y, \xi_t)^p = ((1-t)^p + t^p)d(x, y)^p < d(x, y)^p$$

which contradicts the fact that $(x, y) \in \mathcal{S}$. In particular, since γ is supported in the set of geodesics joining points in \mathcal{S} , we get that $\text{supp}((e_t)_\# \gamma) \subset \Omega$ for any $t \in (0, 1)$. Let $\mu_t \in \mathcal{M}(X, A)$ be given by

$$\mu_t = (e_t)_\#(\gamma)|_\Omega.$$

We claim that the curve $(\mu_t)_{t \in [0,1]}$ is a constant speed geodesic joining μ^0 and μ^1 in $\mathcal{M}_p(X, A)$. Indeed, since $(e_0, e_t) \circ \text{GeoSel} = \text{id}$ then

$$\mu^0 = \pi_{\#}^1 \gamma|_{\Omega} = \pi_{\#}^1 (e_0, e_1)_{\#} \text{GeoSel}_{\#}(\gamma)|_{\Omega} = (e_0)_{\#}(\gamma)|_{\Omega} = \mu_0$$

and, analogously, $\mu^1 = \mu_1$. Moreover,

$$\begin{aligned} \text{Wb}_p^p(\mu_t, \mu_s) &= \text{Wb}_p^p((e_t)_{\#}(\gamma)|_{\Omega}, (e_s)_{\#}(\gamma)|_{\Omega}) \\ &\leq C((e_t, e_s)_{\#}(\gamma)) \\ &= \int_{X \times X} d(x, y)^p d(e_t, e_s)_{\#}(\gamma)(x, y) \\ &= \int_{X \times X} d(\text{GeoSel}(x, y)_t, \text{GeoSel}(x, y)_s)^p d\gamma(x, y) \\ &= |t - s|^p \int_{X \times X} d(x, y)^p d\gamma(x, y) \\ &= |t - s|^p \text{Wb}_p^p(\mu_0, \mu_1). \end{aligned}$$

This argument implies both that $\mu_t \in \mathcal{M}_p(X, A)$ for any $t \in [0, 1]$, by the triangle inequality, and that $(\mu_t)_{t \in [0,1]} \in \text{Geo}(\mathcal{M}_p(X, A))$.

Now, for the second claim, let us assume that $(\mu_t)_{t \in [0,1]}$ is a geodesic in $\mathcal{M}_p(X, A)$. We can find a measure $\gamma^m \in \mathcal{M}(\mathcal{C}([0, 1], X))$ such that

$$(e_{i/2^m}, e_{(i+1)/2^m})_{\#} \gamma^m \in \text{Opt}(\mu_{i/2^m}, \mu_{(i+1)/2^m}), \quad i \in \{0, \dots, 2^m - 1\}$$

and γ^m -a.e. ξ is a geodesic when restricted to each interval $[i/2^m, (i+1)/2^m]$. Indeed, for each $m \in \mathbb{N}$ and each $i \in \{0, \dots, 2^m - 1\}$, choose $\gamma^{i,m} \in \text{Opt}(\mu_{i/2^m}, \mu_{(i+1)/2^m})$ and apply Lemma 3.4 iteratively to get a measure $\gamma^m \in \mathcal{M}(X^{2^m+1})$ such that

$$\pi_{\#}^{i,i+1} \gamma^m = \gamma^{i,m} + \sigma^{i,m}$$

where $\sigma^{i,m}$ is supported in $\Delta(A \times A)$. Then, by considering the measurable map $G^m: X^{2^m+1} \rightarrow \mathcal{C}([0, 1], X)$ given by

$$(x_0, \dots, x_{2^m}) \mapsto \text{GeoSel}(x_0, x_1) * \dots * \text{GeoSel}(x_{2^m}, x_{2^m+1}),$$

where $*$ denotes concatenation of paths, we can define

$$\gamma^m = G_{\#}^m(\gamma^m).$$

We now claim that γ^m is concentrated in $\text{Geo}(X)$. Indeed, for any $j, k \in \{0, \dots, 2^m\}$, we have

$$\begin{aligned} \|d(e_{j/2^m}, e_{k/2^m})\|_{L^p(\gamma^m)} &\leq \left\| \sum_{i=j}^{k-1} d(e_{i/2^m}, e_{(i+1)/2^m}) \right\|_{L^p(\gamma^m)} \\ &\leq \sum_{i=j}^{k-1} \|d(e_{i/2^m}, e_{(i+1)/2^m})\|_{L^p(\gamma^m)} \\ &= \sum_{i=j}^{k-1} \text{Wb}_p(\mu_{i/2^m}, \mu_{(i+1)/2^m}) \\ &= \text{Wb}_p(\mu_{j/2^m}, \mu_{k/2^m}) \end{aligned}$$

where the first inequality is given by the monotonicity of the integral; the second one is the triangle inequality in $L^p(\gamma^m)$, and the last two lines are consequences of the definition of γ^m and the fact

that $(\mu_t)_{t \in [0,1]}$ is a geodesic. As a consequence, we get that

$$(e_{j/2^m}, e_{k/2^m})_{\#} \gamma^m \in \text{Opt}(\mu_{j/2^m}, \mu_{k/2^m})$$

and

$$d(e_{j/2^m}, e_{k/2^m}) = \sum_{i=j}^{k-1} d(e_{i/2^m}, e_{(i+1)/2^m})$$

in the support of γ^m , which in turn implies that $\text{supp}(\gamma^m) \subset \text{Geo}(X)$.

We now proceed to prove that $\{\gamma^m\}_{m \in \mathbb{N}}$ is relatively compact with respect to the vague topology. Indeed, since $(e_0, e_1)_{\#}(\gamma^m) \in \text{Opt}(\mu_0, \mu_1)$, then by Proposition 4.1 we know that $(e_0, e_1)_{\#}(\gamma^m)$ is concentrated in $\mathcal{S} \cap E_{\Omega}$, which in turn implies that γ^m is concentrated in $(e_0, e_1)^{-1}(\mathcal{S} \cap E_{\Omega})$. Therefore, if $\mathcal{K} \subset \text{Geo}(X)$ is a compact set, then

$$\gamma^m(\mathcal{K}) \leq \gamma^m((e_{1/2})^{-1}(e_{1/2}(\mathcal{K}))) = (e_{1/2})_{\#}(\gamma^m)(e_{1/2}(\mathcal{K})) = \mu_{1/2}(e_{1/2}(\mathcal{K})),$$

and since $\mu_{1/2}$ is a Radon measure and $e_{1/2}(\mathcal{K})$ is compact, we get that

$$\sup_{m \in \mathbb{N}} \gamma^m(\mathcal{K}) < \infty,$$

which proves the claim.

We can also see that, for any compact $C \subset X$ and any dyadic rational $t \in [0, 1]$, $\{\gamma^m|_{e_t^{-1}(C)}\}_{m \in \mathbb{N}}$ is weakly precompact. Indeed, if $p_0 \in X$ and $R > r > 0$ are such that $C \subset \overline{B}_r(p_0) \subset \overline{B}_R(p_0)$, then an argument analogous to that in the proof of Theorem 3.5 shows that

$$\gamma^m(e_t^{-1}(C) \setminus (e_0, e_1)^{-1}(\overline{B}_R(p_0) \times \overline{B}_R(p_0))) \leq \frac{\text{Wb}_p^p(\mu_0, \mu_1)}{(R-r)^p}$$

where $(e_0, e_1)^{-1}(\overline{B}_R(p_0) \times \overline{B}_R(p_0))$ is closed and, being the set of geodesics with endpoints in $\overline{B}_R(p_0)$, is contained in $\text{Geo}(\overline{B}_{2R}(X))$, which is compact due to Arzelà–Ascoli and the fact that X is proper. By fixing $r > 0$ and letting R tend to infinity, the right hand side of the inequality above can be made arbitrarily small, uniformly over $m \in \mathbb{N}$, which implies that $\{\gamma^m|_{e_t^{-1}(C)}\}_{m \in \mathbb{N}}$ is tight. Moreover, for sufficiently large $m \in \mathbb{N}$, each $\gamma^m|_{e_t^{-1}(C)}$ has total mass $\mu_t(C) < \infty$, therefore we have uniformly bounded total variation, and the claim follows due to Theorem 2.11.

As a consequence of the previous observations, we can assume that, up to passing to a subsequence, $\{\gamma^m\}_{m \in \mathbb{N}}$ is vaguely convergent to some $\gamma \in \mathcal{M}(\text{Geo}(X))$, and for any compact $C \subset \Omega$ and any dyadic rational $t \in [0, 1]$, $\{\gamma^m|_{e_t^{-1}(C)}\}_{m \in \mathbb{N}}$ is weakly convergent to $\gamma|_{e_t^{-1}(C)}$. Thus, if $f \in \mathcal{C}_c(\Omega)$, $n \in \mathbb{N}$ and $i \in \{0, \dots, 2^n\}$, then

$$\begin{aligned} \int_{\Omega} f d(e_{i/2^n})_{\#} \gamma &= \int_{e_{i/2^n}^{-1}(\text{supp}(f))} f \circ e_{i/2^n} d\gamma \\ &= \lim_{m \rightarrow \infty} \int_{e_{i/2^n}^{-1}(\text{supp}(f))} f \circ e_{i/2^n} d\gamma^m \\ &= \lim_{m \rightarrow \infty} \int_{\text{supp}(f)} f d(e_{i/2^n})_{\#} \gamma^m \\ &= \int_{\Omega} f d\mu_{i/2^n}, \end{aligned}$$

which implies $(e_{i/2^n})_{\#}\gamma = \mu_{i/2^n}$. Finally, for any other $t \in [0, 1]$, let $\{t_k\}_{k \in \mathbb{N}}$ and $\{t^l\}_{l \in \mathbb{N}}$ be two sequences of dyadic rational numbers converging to t , with $t_k \leq t \leq t^l$ for any $k, l \in \mathbb{N}$, and observe that $(e_{t_k}, e_t)_{\#}\gamma \in \text{Adm}(\mu_{t_k}, (e_t)_{\#}\gamma|_{\Omega})$. Therefore,

$$\begin{aligned} \text{Wb}_p^p(\mu_{t_k}, (e_t)_{\#}\gamma|_{\Omega}) &\leq C((e_{t_k}, e_t)_{\#}\gamma) \\ &\leq C((e_{t_k}, e_{t^l})_{\#}\gamma) \\ &= \text{Wb}_p^p(\mu_{t_k}, \mu_{t^l}). \end{aligned}$$

By letting $k, l \rightarrow \infty$, we get that $(e_t)_{\#}\gamma|_{\Omega} = \mu_t$ as claimed. \square

Proposition 5.4. *Let (X, A) be a complete, separable, proper, geodesic, and non-branching metric pair, and $p \in (1, \infty)$. Then $\mathcal{M}_p(X, A)$ is non branching as well. Furthermore, if $(\mu_t)_{t \in [0, 1]} \subset \mathcal{M}_p(X, A)$ is a constant speed geodesic, then for any $t \in (0, 1)$ and any $\gamma \in \text{Opt}(\mu_0, \mu_t)$, γ_X^{Ω} is unique and it is induced by a map.*

Proof. Let $(\mu_t)_{t \in [0, 1]} \in \text{Geo}(\mathcal{M}_p(X, A))$, $t \in (0, 1)$ and consider $\gamma^1 \in \text{Opt}(\mu_0, \mu_t)$ and $\gamma^2 \in \text{Opt}(\mu_t, \mu_1)$. By the proof of Theorem 5.2, there is measure $\gamma \in \mathcal{M}(X^3)$ such that

$$\pi_{\#}^{1,2}\gamma = \gamma^1 + \sigma^1 \quad \text{and} \quad \pi_{\#}^{2,3}\gamma = \gamma^2 + \sigma^2$$

for some σ^1, σ^2 supported in $\Delta(A \times A)$, and such that $\pi_{\#}^{1,3}\gamma \in \text{Opt}(\mu_0, \mu_1)$. Moreover,

$$d(x, y) = td(x, z), \quad d(y, z) = (1 - t)d(x, z)$$

which implies that x, y, z lie in a geodesic, for any $(x, y, z) \in \text{supp}(\gamma)$,

Now consider $(x, y, z), (x', y, z') \in \text{supp}(\gamma)$. By Proposition 4.1, we know that $\text{supp}(\pi_{\#}^{1,3}\gamma)$ is in the superdifferential of a c -concave function, which implies it is c -cyclically monotone. Therefore,

$$\begin{aligned} d(x, z)^p + d(x', z')^p &\leq d(x, z')^p + d(x', z)^p \\ &\leq (d(x, y) + d(y, z'))^p + (d(x', y) + d(y, z))^p \\ &= (td(x, z) + (1 - t)d(x', z'))^p + (td(x', z') + (1 - t)d(x, z))^p \\ &\leq td(x, z)^p + (1 - t)d(x', z')^p + td(x', z')^p + (1 - t)d(x, z)^p \\ &= d(x, z)^p + d(x', z')^p \end{aligned}$$

where the last inequality is due to the convexity of $t \mapsto t^p$ for $p > 1$. Moreover, the strong convexity of the same function, and the fact the all the inequalities above are equations, imply that $d(x, z) = d(x', z')$ and

$$d(x, y) + d(y, z') = d(x, z').$$

In particular, x, y, z' lie in a geodesic. Since X is non-branching, we get that $z = z'$, and analogously we get that $x = x'$. In other words, the map $\pi^2: (x, y, z) \mapsto y$ is injective in $\text{supp}(\gamma)$. In particular, if T is the inverse of $\pi^2|_{\text{supp}(\gamma)}$, we get that

$$(\pi^1 \circ T, \text{id})_{\#}\mu_t = (\gamma^1)_{\Omega}^X \quad \text{and} \quad (\text{id}, \pi^3 \circ T)_{\#}\mu_t = (\gamma^2)_{\Omega}^X.$$

Therefore, $(\gamma^1)_{\Omega}^X$ and $(\gamma^2)_{\Omega}^X$ are induced by maps. Moreover, this also implies that $(\gamma^1)_{\Omega}^X$ is unique, because otherwise we could construct $\gamma \in \text{Opt}(\mu_0, \mu_t)$ such that

$$\gamma_X^{\Omega} = \frac{1}{2} \left((\pi^1 \circ T, \text{id})_{\#}\mu_t + (\pi^1 \circ T', \text{id})_{\#}\mu_t \right),$$

which would not be induced by a map.

Finally, to prove that $\mathcal{M}_p(X, A)$ is non-branching, consider geodesics $(\mu_t)_{t \in [0,1]}$, $(\mu'_t)_{t \in [0,1]}$ and $t_0 \in (0, 1)$ such that $\mu_0 = \mu'_0$ and $\mu_{t_0} = \mu'_{t_0}$. Let $\gamma, \gamma' \in \mathcal{M}(\text{Geo}(X))$ be such that $\mu_t = (e_t)_\# \gamma|_\Omega$ and $\mu'_t = (e_t)_\# \gamma'|_\Omega$ for all $t \in [0, 1]$. Then we have

$$(e_0, e_{t_0})_\# \gamma, (e_0, e_{t_0})_\# \gamma' \in \text{Opt}(\mu_0, \mu_{t_0})$$

which, due to our previous arguments, implies

$$(e_0, e_{t_0})_\# \gamma = (e_0, e_{t_0})_\# \gamma|_{X \times \Omega} = (e_0, e_{t_0})_\# \gamma'|_{X \times \Omega} = (e_0, e_{t_0})_\# \gamma',$$

where we have used that $\text{supp}((e_{t_0})_\# \gamma) \subset \Omega$ for $t_0 \in (0, 1)$ (see Proposition 5.2). Since X is non-branching, the map $(e_0, e_{t_0}): \text{Geo}(X) \rightarrow X \times X$ is injective, which implies

$$\gamma = \gamma',$$

therefore

$$\mu_1 = (e_1)_\# \gamma|_\Omega = (e_1)_\# \gamma'|_\Omega = \mu'_1,$$

and the proposition follows. \square

5.3. Non-negative curvature.

Theorem 5.5. *Assume that (X, A) is a complete, separable, proper, geodesic and non-negatively curved Alexandrov space. Then $\mathcal{M}_2(X, A)$ is a non-negatively curved Alexandrov space too.*

Proof. Let $(\mu_t)_{t \in [0,1]}$ be a constant speed geodesic in $\mathcal{M}_2(X, A)$, which we know exists because X is geodesic. Let also $\nu \in \mathcal{M}_2(X, A)$ be some measure. By Proposition 5.2, we know there exists $\gamma \in \mathcal{M}(\text{Geo}(X))$ such that $(e_0, e_1)_\# \gamma \in \text{Opt}(\mu_0, \mu_1)$ and $(e_{1/2})_\# \gamma|_\Omega = \mu_{1/2}$. Consider also $\gamma \in \text{Opt}(\mu_{1/2}, \nu)$. By observing that $(e_{1/2})_\# \gamma|_\Omega = \pi_\#^1 \gamma|_\Omega = \mu_{1/2}$ and applying the disintegration theorem, we get a measure $\alpha \in \mathcal{M}(\text{Geo}(X) \times X)$ such that

$$\begin{aligned} \pi_\#^{\text{Geo}(X)} \alpha &= \gamma \\ (e_{1/2}, \pi^X)_\# \alpha &= \gamma \end{aligned}$$

In particular, $(e_0, \pi^X)_\# \alpha \in \text{Adm}(\mu_0, \nu)$ and $(e_1, \pi^X)_\# \alpha \in \text{Adm}(\mu_1, \nu)$, therefore

$$\begin{aligned} 2\text{Wb}_2^2(\mu_{1/2}, \nu) &= \int_{X \times X} 2d(x, z)^2 d\gamma(x, z) \\ &= \int_{\text{Geo}(X) \times X} 2d(\xi_{1/2}, z) d\alpha(\xi, z) \\ &\geq \int_{\text{Geo}(X) \times X} d(\xi_0, z)^2 + d(\xi_1, z)^2 - 2d(\xi_0, \xi_1)^2 d\alpha(\xi, z) \\ &= \int_{X \times X} d(x, z)^2 d(e_0, \pi^X)_\# \alpha(x, z) + \int_{X \times X} d(x, z)^2 d(e_1, \pi^X)_\# \alpha(x, z) \\ &\quad - 2 \int d(x, y)^2 d((e_0, e_1) \circ \pi^{\text{Geo}(X)})_\# \alpha(x, y) \\ &= \int_{X \times X} d(x, z)^2 d(e_0, \pi^X)_\# \alpha(x, z) + \int_{X \times X} d(x, z)^2 d(e_1, \pi^X)_\# \alpha(x, z) \\ &\quad - 2 \int d(x, y)^2 d(e_0, e_1)_\# \gamma(x, y) \\ &\geq \text{Wb}_2^2(\mu_0, \nu) + \text{Wb}_2^2(\mu_1, \nu) - 2\text{Wb}_2^2(\mu_0, \mu_1), \end{aligned}$$

which proves the claim. \square

An interesting fact about the geometric structure of $\mathcal{M}_2(X, A)$ when X is a non-negatively curved Alexandrov space is that the zero measure is always an *extremal point*, i.e. a point at which the space of directions has diameter bounded above by $\pi/2$ (see [24] for a more general exposition about extremal points and extremal sets in Alexandrov spaces).

Proposition 5.6. *The space of directions at the zero measure, $\Sigma_0(\mathcal{M}_2(X, A))$, has diameter no greater than $\pi/2$.*

Proof. Let $\mu, \nu \in \mathcal{M}_2(X, A)$. Then, we know that

$$\text{Wb}_2(\mu, 0)^2 + \text{Wb}_2(\nu, 0)^2 \geq \text{Wb}_2(\mu, \nu)^2$$

since the transport plan $(\text{id}, \text{proj}_A)_\# \mu + (\text{proj}_A, \text{id})_\# \nu \in \text{Adm}(\mu, \nu)$ is suboptimal. Therefore, if $\xi_1, \xi_2 \in \text{Geo}(\mathcal{M}_2(X, A))$ are geodesics with $\xi_1(0) = \xi_2(0) = 0$, then

$$\cos \angle(\xi_1, \xi_2) = \lim_{s, t \rightarrow 0} \frac{\text{Wb}_2(\xi_1(s), 0)^2 + \text{Wb}_2(\xi_2(t), 0)^2 - \text{Wb}_2(\xi_1(s), \xi_2(t))^2}{2\text{Wb}_2(\xi_1(s), 0)\text{Wb}_2(\xi_2(t), 0)} \geq 0$$

which implies that $\angle(\xi_1, \xi_2) \leq \pi/2$. \square

6. EMBEDDING OF $\mathcal{D}_p(X, A)$ INTO $\mathcal{M}_p(X, A)$

One of the motivations for considering $\mathcal{M}_p(X, A)$ is that it admits a natural inclusion of the space of generalised persistence diagrams $\mathcal{D}_p(X, A)$, as defined in [10]. Indeed, we have the natural map

$$\begin{aligned} \mathcal{D}_p(X, A) &\longrightarrow \mathcal{M}_p(X, A) \\ \sigma &\longmapsto \sum_{x \in \sigma} \delta_x. \end{aligned}$$

Proposition 6.1. *Let $\mu \in \mathcal{M}_p(X, A)$, $r > 0$ and $A_r = \{x \in X : d(x, A) \leq r\}$. Let $\mu^r = \mu|_{X \setminus A_r}$. Then $\text{Wb}_p(\mu^r, \mu) \rightarrow 0$ when $r \rightarrow 0$. Similarly, if $\sigma \in \mathcal{D}_p(X, A)$, we have $d_p(\sigma^r, \sigma) \rightarrow 0$ as $r \rightarrow 0$.*

Proof. Let $\gamma \in \text{Adm}(\mu, \mu^r)$ be the transport plan given by

$$\gamma = (\text{id}, \text{id})_\# \mu|_{X \setminus A_r} + (\text{id}, \text{proj}_A)_\# \mu|_{A_r}.$$

Therefore,

$$\text{Wb}_p^p(\mu, \mu^r) \leq \int_{A_r} d(x, A)^p d\mu(x).$$

By the monotone convergence theorem applied to μ with the functions $f_r(x) = d(x, A)^p \cdot 1_{X \setminus A_r}(x)$, we conclude that $\text{Wb}_p(\mu, \mu^r) \rightarrow 0$ as $r \rightarrow 0$. Similar arguments show that $d_p(\sigma, \sigma^r) \rightarrow 0$ as $r \rightarrow 0$. \square

Theorem 6.2. *For $\sigma, \tau \in \mathcal{D}_p(X, A)$, $\text{Wb}_p(\sigma, \tau) = d_p(\sigma, \tau)$.*

Proof. First we consider the case when both $\sigma \setminus A$ and $\tau \setminus A$ have finite cardinality (counting multiplicity), that is, $\sigma \setminus A = \{x_1, \dots, x_m\}$ and $\tau \setminus A = \{y_1, \dots, y_n\}$ for some $x_i, y_j \in \Omega$, $i = 1, \dots, m, j = 1, \dots, n$. Let us define

$$\begin{aligned} \tilde{\sigma} &= \{x_1, \dots, x_m, \text{proj}_A(y_1), \dots, \text{proj}_A(y_n)\}, \\ \tilde{\tau} &= \{y_1, \dots, y_n, \text{proj}_A(x_1), \dots, \text{proj}_A(x_m)\}. \end{aligned}$$

Then, it is clear that

$$d_p^p(\sigma, \tau) = d_p^p(\tilde{\sigma}, \tilde{\tau}) = \min\langle P, C \rangle_{\text{HS}},$$

where P runs over all permutation matrices of size $(m+n) \times (m+n)$, $\langle \cdot, \cdot \rangle_{\text{HS}}$ denotes the Hilbert–Schmidt inner product of square matrices, and

$$C_{ij} = \begin{cases} d(x_i, y_j)^p & \text{if } 1 \leq i \leq m, 1 \leq j \leq n \\ d(x_i, p_A(x_{j-n}))^p & \text{if } 1 \leq i \leq m, n < j \leq m+n \\ d(y_j, p_A(y_{i-m}))^p & \text{if } m < i \leq m+n, 1 \leq j \leq n \\ 0 & \text{if } m < i \leq m+n, n < j \leq m+n \end{cases}.$$

Similarly, it is clear that

$$(6.1) \quad \text{Wb}_p^p(\sigma, \tau) = \min_M \langle M, C \rangle_{\text{HS}}$$

where M runs over all matrices of size $(m+n) \times (m+n)$ such that $M_{ij} \geq 0$ and $\sum_{i=1}^{m+n} M_{ij} = \sum_{j=1}^{m+n} M_{ij} = 1$.

However, it is known that minimisers in equation (6.1) are permutation matrices (see [25, 5]). This proves the finite case.

For arbitrary $\sigma, \tau \in \mathcal{D}_p(X, A)$, consider $r > 0$ and observe that both σ^r and τ^r contain finitely many points in Ω . Then, due to Proposition 6.1, we get that

$$\text{Wb}_p(\sigma, \tau) = \lim_{r \rightarrow 0} \text{Wb}_p(\sigma^r, \tau^r) = \lim_{r \rightarrow 0} d_p(\sigma^r, \tau^r) = d_p(\sigma, \tau). \quad \square$$

REFERENCES

1. Andrés Ahumada Gómez and Mauricio Che, *Gromov-Hausdorff convergence of metric pairs and metric tuples*, Differential Geom. Appl. **94** (2024), Paper No. 102135, 29. MR 4727066
2. Luigi Ambrosio and Nicola Gigli, *A user’s guide to optimal transport*, Modelling and Optimisation of Flows on Networks, Lecture Notes in Mathematics, vol. 2062, Springer, Berlin, Heidelberg, 2013.
3. Edward A. Azoff, *Borel measurability in linear algebra*, Proc. Amer. Math. Soc. **42** (1974), 346–350. MR 327799
4. David Bate and Ana Lucía García Pulido, *Bi-lipschitz embeddings of the space of unordered m -tuples with a partial transportation metric*, Mathematische Annalen (2024).
5. Garrett Birkhoff, *Three observations on linear algebra*, Univ. Nac. Tucumán. Revista A. **5** (1946), 147–151. MR 20547
6. Nicolas Bourbaki, *Elements of mathematics. General topology. Part 2*, Hermann, Paris; Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1966. MR 205211
7. Peter Bubenik and Irina Hartsock, *Topological and metric properties of spaces of generalized persistence diagrams*, J. Appl. Comput. Topol. (2024).
8. Dmitri Burago, Yuri Burago, and Sergei Ivanov, *A course in metric geometry*, Graduate Studies in Mathematics, vol. 33, American Mathematical Society, Providence, RI, 2001. MR 1835418
9. Luis A. Caffarelli and Robert J. McCann, *Free boundaries in optimal transport and Monge-Ampère obstacle problems*, Ann. of Math. (2) **171** (2010), no. 2, 673–730. MR 2630054
10. Mauricio Che, Fernando Galaz-García, Luis Guijarro, and Ingrid Membrillo Solis, *Metric geometry of space of persistence diagrams*, arXiv:2109.14697, 2022.
11. Mauricio Che, Fernando Galaz-García, Luis Guijarro, Ingrid Membrillo Solis, and Motiejus Valiunas, *Basic metric geometry of the bottleneck distance*, Proceedings of the American Mathematical Society (2024).
12. Lénaïc Chizat, Gabriel Peyré, Bernhard Schmitzer, and François-Xavier Vialard, *Unbalanced optimal transport: dynamic and Kantorovich formulations*, J. Funct. Anal. **274** (2018), no. 11, 3090–3123. MR 3782988
13. David Cohen-Steiner, Herbert Edelsbrunner, and John Harer, *Stability of persistence diagrams*, Discrete Comput. Geom. **37** (2007), no. 1, 103–120. MR 2279866
14. Vincent Divol and Théo Lacombe, *Understanding the topology and the geometry of the space of persistence diagrams via optimal partial transport*, Journal of Applied and Computational Topology **5** (2021), no. 1, 1–53. MR 4224153
15. Herbert Edelsbrunner, David Letscher, and Afra Zomorodian, *Topological persistence and simplification*, 41st Annual Symposium on Foundations of Computer Science (Redondo Beach, CA, 2000), IEEE Comput. Soc. Press, Los Alamitos, CA, 2000, pp. 454–463. MR 1931842
16. Alessio Figalli, *The optimal partial transport problem*, Arch. Ration. Mech. Anal. **195** (2010), no. 2, 533–560. MR 2592287

17. Alessio Figalli and Nicola Gigli, *A new transportation distance between non-negative measures, with applications to gradient flows with Dirichlet boundary conditions*, Journal de Mathématiques Pures et Appliquées. Neuvième Série **94** (2010), no. 2, 107–130. MR 2665414
18. Detlef Gromoll, Wilhelm Klingenberg, and Wolfgang Meyer, *Riemannsche Geometrie im Grossen*, Lecture Notes in Mathematics, vol. No. 55, Springer-Verlag, Berlin-New York, 1968. MR 229177
19. Leonid G. Hanin, *Kantorovich-Rubinstein norm and its application in the theory of Lipschitz spaces*, Proc. Amer. Math. Soc. **115** (1992), no. 2, 345–352. MR 1097344
20. Richard Jordan, David Kinderlehrer, and Felix Otto, *The variational formulation of the Fokker-Planck equation*, SIAM Journal on Mathematical Analysis **29** (1998), no. 1, 1–17 (English).
21. Olav Kallenberg, *Random measures*, fourth ed., Akademie-Verlag, Berlin; Academic Press, Inc., London, 1986. MR 854102
22. Matthias Liero, Alexander Mielke, and Giuseppe Savaré, *Optimal entropy-transport problems and a new Hellinger-Kantorovich distance between positive measures*, Invent. Math. **211** (2018), no. 3, 969–1117. MR 3763404
23. Wolfgang Meyer, *Toponogov's Theorem and Applications*, Lecture Notes, College on Differential Geometry, Trieste, 1989.
24. Anton Petrunin, *Semiconcave functions in Alexandrov's geometry*, Surveys in differential geometry. Vol. XI, Surv. Differ. Geom., vol. 11, Int. Press, Somerville, MA, 2007, pp. 137–201. MR 2408266
25. Filippo Santambrogio, *Optimal transport for applied mathematicians*, Progress in Nonlinear Differential Equations and their Applications, vol. 87, Birkhäuser/Springer, Cham, 2015, Calculus of variations, PDEs, and modeling. MR 3409718
26. Giuseppe Savaré and Giacomo Enrico Sodini, *A relaxation viewpoint to Unbalanced Optimal Transport: Duality, optimality and Monge formulation*, J. Math. Pures Appl. (9) **188** (2024), 114–178. MR 4756504
27. Katharine Turner, Yuriy Mileyko, Sayan Mukherjee, and John Harer, *Fréchet means for distributions of persistence diagrams*, Discrete Comput. Geom. **52** (2014), no. 1, 44–70. MR 3231030
28. Cédric Villani, *Optimal transport*, Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 338, Springer-Verlag, Berlin, 2009, Old and new. MR 2459454
29. Afra Zomorodian and Gunnar Carlsson, *Computing persistent homology*, Discrete Comput. Geom. **33** (2005), no. 2, 249–274. MR 2121296

(Che) DEPARTMENT OF MATHEMATICAL SCIENCES, DURHAM UNIVERSITY, UNITED KINGDOM.

Email address: mauricio.a.che-moguel@durham.ac.uk