

# Efficient classical algorithm for simulating boson sampling with inhomogeneous partial distinguishability

S.N. van den Hoven,<sup>1</sup> E. Kanis,<sup>1</sup> and J. J. Renema<sup>1</sup>

<sup>1</sup>*MESA+ Institute for Nanotechnology, University of Twente,  
P. O. box 217, 7500 AE Enschede, The Netherlands*

(Dated: June 26, 2024)

Boson sampling is one of the leading protocols for demonstrating a quantum advantage, but the theory of how this protocol responds to noise is still incomplete. We extend the theory of classical simulation of boson sampling with partial distinguishability to the case where the degree of indistinguishability between photon pairs is different between different pairs.

## I. INTRODUCTION

Quantum computers are expected to outperform classical computers in certain well-defined tasks such as the hidden subgroup problems for abelian finite groups, which includes prime factorization [1], simulations of quantum systems [2], and unstructured search [3]. However, building a universal fault-tolerant quantum computer is no easy task, due to the extreme degree of control over a large number of quantum particles required. As an intermediate step, experimental research has focused on the demonstration of a quantum advantage [4], i.e. a computational task where quantum hardware outperforms all classical hardware in wall-clock time, on a well-defined computational problem not necessarily of any practical utility. Such demonstrations have been claimed in superconducting circuits [5–7], and photons [8–10].

These quantum advantage claims caused substantial debate, with several later being outperformed by classical simulations [11–15]. This was possible despite strong guarantees of computational complexity because experimental hardware suffers from various forms of noise, which introduce decoherence, and reduce the degree to which the task which the device is performing is truly quantum mechanical, thereby opening up loopholes for classical simulation strategies to exploit. Similar to the situation in Bell tests, these simulation strategies demarcate the regime where a classical explanation for the observed data cannot be ruled out. They therefore serve a vital function in assessing the success or failure of a quantum advantage demonstration.

One protocol for a quantum advantage demonstration is boson sampling [16]. In boson sampling, single photons are sent through a large-scale linear interferometer. The computational task is to provide samples from the output state measured in the Fock basis (see Fig. 1). Complexity arises ultimately from quantum interference between the exponentially many ways in which the photons can traverse the interferometer to produce a single outcome. The main sources of noise in boson sampling are photon loss [17], where some of the photons do not emerge from the output of the interferometer, and photon

distinguishability [18], where the particles carry which-path information in their internal quantum states.

Several strategies exist to classically simulate imperfect boson sampling, including ones based on approximating the quantum state using tensor networks [14, 19–21], ones aimed at reproducing the marginal photon distributions behind some number of optical modes [22, 23], and ones based on phase-space methods [24–27]. Some methods are specifically aimed at spoofing certain benchmarks which have themselves been put forward as proxies for computational complexity [28].

The classical simulation technique that we focus on here makes use of the fact that imperfections dampen quantum interference more strongly between paths through the interferometer that exhibit a higher degree of classical dissimilarity, i.e. which would be more different if the particles were fully classical [29, 30]. This allows us to establish a notion of distance between the paths, with the attenuation of quantum interference between two paths depending exponentially on the distance. Since it can be shown that there are only polynomially many paths shorter than a given distance, truncating the quantum interference at a fixed distance produces an approximation to the output probability, which is both efficiently computable and maintains its accuracy as the system size is scaled up.

Interestingly, this bosonic algorithm has a direct counterpart in the simulation of qubit-based systems [31], as do some of the other algorithms. It is an open question whether this is a coincidence or a symptom of some deeper structure of the problem of demonstrating a quantum advantage, with Kalai and Kindler conjecturing [32] that the susceptibility of boson sampling to noise is an intrinsic feature of a non-error corrected approach to demonstrating a quantum advantage.

However, this algorithm suffers from some restrictions. In particular, it assumes that the degree of indistinguishability between all photons is equal. In the case of varying indistinguishability among pairs of photons, the only solution available to the algorithm is to approximate the degree of indistinguishability between all photons as that of the highest pairwise indistinguishability. This substantially reduces the applicability of the algo-

rithm to a real experiment, where such fluctuations inevitably occur. In the most pathological case, an experiment with two fully indistinguishable photons and otherwise all distinguishable photons could not be classically simulated, even though only two-photon quantum interference occurs in this case.

In this work, we eliminate the dependency on this assumption, demonstrating a classical simulation of noisy boson sampling that is efficient for realistic models of dissimilar photon indistinguishability. We focus on two experimentally relevant cases: first the case of identical and independent fluctuations in the state of the photons, and secondly on the situation where there are two species of photons, one with partial distinguishability and one with full indistinguishability. For these cases, we show extensions of the algorithm of [29] which achieve better performance than the original, extending the area of the parameter space which is susceptible to classical simulation. For the case of identical and independently distributed partial distinguishabilities, we find that the complexity is entirely governed by the mean of the distribution of Hong-Ou-Mandel [33] visibilities. For the case of the two-species model, we find that the sampling problem is still classically simulable, but at an additional cost exponential in the number of fully indistinguishable particles.

We achieve these results by reworking and simplifying the derivation of [29], to more easily accommodate more complex partial distinguishability distributions. We therefore show that the sensitivity of the hardness of boson sampling to imperfections is not a result of the specific assumptions made in the classical simulation techniques. Moreover, these results provide evidence for the idea that the sensitivity of quantum advantage demonstrations to noise is intrinsic rather than dependent on the specific model of noise chosen.

We focus specifically on partial distinguishability as a source of error, motivated by the idea that both optical loss and indistinguishability, as well as other errors, all affect the computational complexity of the boson sampling problem in similar ways [30, 34], meaning that any is paradigmatic for the others. We leave full extension of our results to optical loss and other imperfections to future work.

## II. CLASSICAL ALGORITHM FOR BOSON SAMPLING WITH PARTIALLY DISTINGUISHABLE PHOTONS

In this section, we will revisit the algorithm for efficiently classically simulating boson sampling with partial distinguishable particles as described in [29][35]. We demonstrate a simplified proof for the algorithm, that is heavily inspired by the proof given in [29], but allows us to extend the algorithm to other cases more easily.




FIG. 1. Schematic representation of a boson sampling device. The photon states enter the linear optical network on the left. The photons interfere and create an output state. Upon measuring where the photons leave the interferometer, a sample from the created output state has been drawn

This section will be structured as follows. First we will revisit the theory needed to describe interference experiments with partial distinguishable photons. Then, we will show that we can efficiently approximate transition probabilities by neglecting contributions of high-order multiphoton interference. Lastly we will show that the error induced by such an approximation on the total probability distribution is independent of the number of photons. We start by considering boson sampling with partially distinguishable photons. Previous research has demonstrated a method to compute the probability of detecting a certain output configuration  $\mathbf{s}$  in the Fock-basis[36, 37]. This expression allows for arbitrary multiphoton input states and arbitrary number-resolving photon detectors.

Under the assumption of pure input states and lossless detectors which are insensitive to the internal state of the photon, it has been shown that the results of [36, 37] can be rewritten in the a compact form [18]. The probability of measuring a particular detection outcome  $\mathbf{s}$  is given by:

$$P = \frac{1}{\prod_i r_i! s_i!} \sum_{\sigma \in S_n} \left[ \prod_{j=1}^n S_{j, \sigma(j)} \right] \text{perm}(M \circ M_\sigma^*), \quad (1)$$

where  $M$  is a submatrix of the unitary representation of the interferometer  $U$ , constructed by selecting rows and columns of  $U$  corresponding to the input modes and output modes of interest ( $M = U_{d(\mathbf{r}), d(\mathbf{s})}$ ), where  $d(\mathbf{r})$  and  $d(\mathbf{s})$  represent the mode assignment lists of the input and output states respectively. Note that the size of  $M$  is determined by the number of photons  $n$  considered in the sampling task.  $r_i$  and  $s_i$  denote the  $i^{\text{th}}$  element of the mode occupation lists of the input state and the output state respectively.  $S_n$  denotes the symmetric group,  $\mathcal{S}$  denotes the distinguishability matrix where  $S_{i,j} = \langle \psi_i | \psi_j \rangle$ , the overlap between photon  $i$  and  $j$  represented by their wave-functions  $\psi_i$  and  $\psi_j$  respectively.  $\circ$  represents the Hadamard product,  $*$  represents the element wise conjugation and  $M_\sigma$  represents  $M$  where its

rows are permuted according to  $\sigma$ . Lastly the permanent of matrix  $M$  with shape  $n \times n$  is defined as

$$\text{Perm}(M) = \sum_{\sigma \in S_n} \prod_{i=1}^n M_{i,\sigma(i)}. \quad (2)$$

For the moment, we will continue by assuming that all particles are equally distinguishable, i.e. that:

$$\mathcal{S}_{ij} = x + (1-x)\delta_{ij}. \quad (3)$$

Note that the assumption made in Eq. (3) is in the literature often referred to as the orthogonal bad bit model [38, 39], and is supported by experimental evidence.

Using the assumption of Eq. (3), we note that the quantity  $\prod_{j=1}^n \mathcal{S}_{j,\sigma(j)}$  will only depend on the number of fixed points of  $\sigma$ . (A fixed point is a point in  $\sigma$  that maps to itself after the permutation, or  $\sigma(j) = j$ .) We can therefore rewrite Eq. (1) as:

$$P(s) = \frac{1}{\prod_i r_i! s_i!} \sum_{j=0}^n \sum_{\tau \in \sigma_j} x^j \text{Perm}(M \circ M_\tau^*). \quad (4)$$

Here,  $\sigma_j$  denotes the set of all permutations with  $n-j$  fixed points.

The term  $x^j$  introduces exponential dampening in  $j$ . For this reason, it is natural to truncate the series at some value  $k < n$ :

$$P_k(s) = \frac{1}{\prod_i r_i! s_i!} \sum_{j=0}^k \sum_{\tau \in \sigma_j} x^j \text{Perm}(M \circ M_\tau^*), \quad (5)$$

leaving an expression for the error:

$$Q_k = \frac{1}{\prod_i r_i! s_i!} \sum_{j=k+1}^n \sum_{\tau \in \sigma_j} x^j \text{Perm}(M \circ M_\tau^*). \quad (6)$$

Note that all terms in Eq. (4) for a given  $j$  correspond to all contributions to the probability where  $j$  photons interfere with each other and  $n-j$  photons undergo classical transmission. Hence, by truncating Eq. (4), we only consider those contributions to the probability where at most  $k$  photons interfere with each other.

In the remainder of this section, we will focus on non-collisional input and output states, hence  $\prod_i (r_i!)(s_i!) = 1$ . We will show two things. The first is that Eq. (5) can be computed efficiently on a classical computer for arbitrary system sizes  $n$ . The second is that, under the assumption that  $M$  is filled with elements that are i.i.d. complex Gaussian, the error term in Eq. (6) decreases exponentially as  $n$  increases. It decreases such that, the expectation value of the  $L_1$ -distance between the approximate distribution and the real distribution is upper bounded by the following expression:

$$\mathbb{E} \left( \sum_s |P(s) - P_k(s)| \right) < \sqrt{\frac{x^{2k+2}}{1-x^2}}. \quad (7)$$

This upper bound holds regardless of the system size. Note that if the number of modes is much larger than the number of photons, any  $n \times n$  submatrix of a Haar-random matrix will be close in variation distance to a matrix filled with complex i.i.d. Gaussians [16]. We can use the first result to efficiently draw samples from the approximate distribution via a Metropolis sampler, and we use the second result to show that the distribution that is sampled from is close in variation distance to the real distribution, thus resulting in an efficient classical algorithm for imperfect boson sampling.

To demonstrate that we can efficiently compute  $P_k$  from Eq. (5), we use the fact that an algorithm exists for approximating the permanent of a matrix with real non-negative elements [40]. Additionally, we use the Laplace expansion to split up the permanents from Eq. (6) into sums of the product of two permanents of smaller matrices. One of these two permanents is filled with non-negative elements. We rewrite each term in Eq. (5) by Laplace expanding about the rows that correspond to the fixed points of  $\tau$ :

$$\text{Perm}(M \circ M_\tau^*) = \sum_{\rho \in \binom{n}{j}} \text{Perm}(M_{I_p, \rho} \circ M_{\tau_p, \rho}^*) \text{Perm}(|M_{\tau_u, \bar{\rho}}|^2). \quad (8)$$

Here  $\tau_u$  and  $\tau_p$  are the unpermuted and permuted parts of  $\tau$  respectively, i.e. those parts that correspond to fixed points (cycles of length 1) and longer cycles, respectively. Given that  $\tau$  has  $n-j$  fixed points,  $\rho$  is a  $j$ -combination of  $n$ ,  $\bar{\rho}$  is its complementary set and  $I_p$  is the identity permutation for the elements of  $\rho$ . Eq. (8) now contains two permanents. The second permanent contains a matrix with only real non-negative elements and can be efficiently approximated via the JSV algorithm [40]. The other permanent contains a matrix with complex elements. The size of these matrices is however determined by  $j$ , which due to our truncation has a maximum value of  $k$ . To compute Eq. (5) we thus need to compute permanents of complex matrices of size  $j$  and permanents of real, non-negative matrices of size  $n-j$ . We need to do both of these calculations  $\binom{n}{j}$  times for each  $\tau \in \sigma_j$  for all  $j \leq k$ , which results in a polynomial scaling of computational costs with  $n$  to evaluate  $P_k$ .

We now continue with a derivation for Eq. (7). In the main text we will give a sketch of this derivation, in appendix A we will give a full derivation. It is important to note that the derivation follows the same ideas as presented in [29], but differs in some key details. These differences allow us to find similar upper bounds for adjacent boson sampling experiments as will be elaborated on in the following sections.

The derivation consists of the following steps:

1. Using Jensen's inequality, note that  $\mathbb{E}(|Q_k|) \leq \sqrt{\text{Var}(Q_k)}$

2. Using the definition of the permanent, note that  $Q_k$  is a large sum where each term is described by a product containing elements of  $M$
3. Using Bienamaymé's identity, note that the variance of a large sum is equal to the covariance between all pairs of terms in this large sum
4.  $M$  is assumed to be filled with i.i.d. complex Gaussian elements, and hence  $E(M_{ij}) = 0$ ,  $E((M_{ij})^2) = 0$ ,  $E((M_{ij}^*)^2) = 0$ ,  $E(|M_{ij}|^2) = \frac{1}{m}$  and  $E(|M_{ij}|^4) = \frac{2}{m^2}$  for all  $i, j$ .
5. We use these properties of the elements of  $M$  to find that almost all of these correlations are equal to zero.
6. We use simple combinatorics to count the number of covariances that contribute the same non-zero amount to  $\text{Var}(Q_k)$
7. We find that the error term approximates a truncated geometric series
8. We have now found an approximate expression for  $\text{Var}(Q_k) \approx \frac{(n!)^2}{m^n} \frac{x^{2k+2} - x^{2n+2}}{1-x^2}$  and as a result an upper bound for  $E(|Q_k|)$  for a typical non-collisional output configuration
9. By counting the number of non-collisional output configurations we find the upper bound on  $L_1$ -distance of interest as presented in inequality 7

To conclude this section, we have revisited a known algorithm as described in [29][35]. The algorithm approximates transition probabilities in a boson sampling experiment by neglecting high order interference contributions to the transition probability. We have demonstrated that these approximated transition probabilities are efficiently computable. Moreover we have demonstrated that, under the assumption that all photons are equally distinguishable, the  $L_1$ -distance over all non-collisional outputs between the approximated distribution and the real distribution is upper bounded as demonstrated in inequality 7. Notably, this upper bound is independent of the system size of the boson sampling experiment. In the following section we will relax this assumption.

### III. GENERAL PARTIAL DISTINGUISHABILITY

In the previous section, we have sketched an efficient classical algorithm for boson sampling with partially distinguishable photons, with full details given in Appendix A. To show that the approximate distribution we can efficiently sample from is close to the real distribution it was assumed that all particles are equally distinguishable, see Eq. (3). However, realistic single-photon sources do not

adhere to the assumption that all photons are equally imperfect. The algorithm as proposed in [29] circumvents this problem by computing the upper bound in Eq. (7) as if all photons are equally imperfect and as good as the best photon pair present. This way an upper bound can be found in general, but depending on the variations in the quality of the particles, this bound may be very loose. Here, we relax the assumption made in Eq. (3). Although we are not able to tighten this upper bound in general, we show that for two experimentally relevant generalizations, we are able to tighten this bound.

We will start by noting that without the assumption made in Eq. (3), we can still efficiently evaluate our truncated probability. For general, pure partially distinguishable photons, Eq. (5) reads:

$$P_k = \frac{1}{\prod_i r_i! s_i!} \sum_{j=0}^k \sum_{\tau \in \sigma_j} \left( \prod_{i=1}^n \mathcal{S}_{i, \tau(i)} \right) \text{Perm}(M \circ M_\tau^*). \quad (9)$$

If we compare Eq. (9) with Eq. (5), we notice that the only difference is that  $x^j$  is substituted with  $\prod_{i=1}^n \mathcal{S}_{i, \tau(i)}$ . It takes a multiplication of  $j$  factors to compute  $\prod_{i=1}^n \mathcal{S}_{i, \tau(i)}$  and hence Eq. (9) is still efficiently computable on a classical computer.

We will continue to derive an upper bound for the  $L_1$ -distance between the approximate and the real distribution. For general, pure partially distinguishable photons, Eq. (6) becomes

$$Q_k = \frac{1}{\prod_i r_i! s_i!} \sum_{j=k+1}^n \sum_{\tau \in \sigma_j} \left( \prod_{i=1}^n \mathcal{S}_{i, \tau(i)} \right) \text{Perm}(M \circ M_\tau^*). \quad (10)$$

We note that again the only difference between Eq. (6) and Eq. (10) is that  $x^j$  is substituted with  $\prod_{i=1}^n \mathcal{S}_{i, \tau(i)}$ . We also note that  $\prod_{i=1}^n \mathcal{S}_{i, \tau(i)}$  is completely independent of all elements in  $M$ . After all, the choice for the interferometer is completely independent of the quality of the photons used in the experiment. We are considering the expectation value of  $|Q_k|$  over the ensemble of Haar-unitaries, we thus note that  $E(\prod_{i=1}^n \mathcal{S}_{i, \tau(i)}) = \prod_{i=1}^n \mathcal{S}_{i, \tau(i)}$ . As a result, all steps in appendix A are valid up until Eq. (A14). We continue by directly substituting  $\prod_{i=1}^n \mathcal{S}_{i, \tau(i)}$  for  $x^j$  in Eq. (A14) to find

$$\begin{aligned} \text{Var}(Q_k) = & \\ & \sum_{j=k+1}^n \sum_{\tau \in \sigma_j} \left( \prod_{i=1}^n |\mathcal{S}_{i, \tau(i)}|^2 \right) \sum_{\rho \in \mathcal{S}_n} \sum_{p=0}^{n-j} R_{n-j, p} 2^p \left( \frac{1}{m^2} \right)^n. \end{aligned} \quad (11)$$

We note that  $\sum_{p=0}^{n-j} R_{n-j, p} 2^p \left( \frac{1}{m^2} \right)^n$  is independent of  $\tau$ . We would like to find an expression (or an upper bound) for  $\prod_{i=1}^n |\mathcal{S}_{i, \tau(i)}|^2$  that is independent of  $\tau$ , because that could allow us to simplify Eq. (11) further

by recognizing a truncated geometric series. We note that for a general overlap matrix  $\mathcal{S}$ , we can simplify Eq. (11) by realizing that  $\prod_{i=1}^n |\mathcal{S}_{i,\tau(i)}|^2 \leq \max(|\mathcal{S}_{ij}|)^{2j}$ , but this approach has already been mentioned in [29]. In the remainder of this section, we will discuss two different experimentally relevant cases for which we are able to tighten the upper bound for the  $L_1$ -distance.

### A. Independent and identically distributed orthogonal bad bits

First, we will consider the internal modes of the  $i^{\text{th}}$  photon to be:

$$|\psi\rangle = \sqrt{x_i}|\Psi_0\rangle + \sqrt{1-x_i}|\Psi_i\rangle, \quad (12)$$

where all  $x_i$  are independent and identically distributed variables and  $\langle\Psi_i|\Psi_j\rangle = \delta_{ij}$ . This model is a variation on the orthogonal bad bit model[38, 41] where we allow for deviations in the quality of the photons. We argue that the first case we consider is experimentally relevant, because upon manufacturing single-photon sources, the target indistinguishability is most likely the same for all sources but deviations in the manufacturing process result in fluctuations in the quality of the individual sources. Hence, we can reasonably expect that the quality of the photons are independent and randomly sampled from the same distribution. With this assumption, we can simplify Eq. (11).

The overlap matrix  $\mathcal{S}$  then becomes:

$$\mathcal{S}_{ij} = \begin{cases} 1 & \text{for } i = j \\ \sqrt{x_i}\sqrt{x_j}^* & \text{for } i \neq j \end{cases} \quad (13)$$

Every permutation  $\tau$  can uniquely be described with its cycle notation, from the cycle notation it becomes clear that

$$\mathbb{E}\left(\prod_{r=1}^n |\mathcal{S}_{r,\tau(r)}|^2\right) = \prod_{i \in \text{nonfix}(\tau)} \mathbb{E}(|x_i|^2). \quad (14)$$

Since all  $x_i$  follow the same distribution, the expression in Eq. (14) is independent of  $\tau \in \sigma_j$ , and only depends on the number of fixed points of  $\tau$ . Hence we can simplify Eq. (11), following the same arguments as presented in Eqs. (A14), (A15) and (A16):

$$\begin{aligned} \text{Var}(Q_k) &= \\ \sum_{j=k+1}^n \mathbb{E}(|x_i|^2)^j R_{n,n-j} n! \sum_{p=0}^{n-j} R_{n-j,p} 2^p \left(\frac{1}{m^2}\right)^n \\ &\approx \sum_{j=k+1}^n \mathbb{E}(|x_i|^2)^j \frac{(n!)^2}{m^{2n}}. \end{aligned} \quad (15)$$

Where we note the Gaussian distribution of the elements of  $M$ , and in particular the fact that  $E(M_{ij}) = 0$ , implies that all cross terms between the variance of functions of  $S$  and functions of  $M$  cancel (see Eq. (A8)).

Since  $\mathbb{E}(|x_i|^2) \leq 1$  for  $|\psi_i\rangle$  to be normalized, we can thus recognize a truncated geometric series again to find an upper bound on the trace distance between our approximated distribution and the real distribution:

$$\mathbb{E}\left(\sum_s |P(s) - P_k(s)|\right) = \mathbb{E}\left(\sum_s |Q_k(s)|\right) \quad (16)$$

$$< \sqrt{\frac{\mathbb{E}(|x_i|^2)^{k+1}}{1 - \mathbb{E}(|x_i|^2)}} \quad (17)$$

We find that Eq. (17) is equal to 7, with the substitution of  $\mathbb{E}(|x_i|^2)$  for  $x^2$ . Conveniently,  $|x|^2$  is the visibility of a Hong-Ou-Mandel interference experiment [33]. For independent sources, the complexity of boson sampling is therefore governed by the average of the HOM visibilities. This result improves on our earlier work, which could only upper bound the complexity of boson sampling on independent sources by the maximum of their visibilities. For the specific case of Gaussian i.i.d. distinguishabilities, we note that we can evaluate  $\mathbb{E}(x_i^2)$  as the expectation value of a non-central chi-squared distribution with one degree of freedom and  $\mathbb{E}(x_i^2) = \mu^2 + \sigma^2$ .

Figure 2 shows the effect of this tighter bound on the regime of the parameter space which can be efficiently simulated. We assume  $\max(x_i) = \mu + 2\sigma$ , motivated by the observation that the probability is then about  $\frac{1}{2}$  that  $\max(x_i) < \mu + 2\sigma$  in the case of 30 photons already, and decreasing further if  $n$  increases. The areas of the parameter space in Fig 2 which are in between the solid and dotted lines are the areas of the parameter space which could not be simulated before.

### B. Few indistinguishable photons

For the second case, we consider the internal modes of the  $i^{\text{th}}$  photon to be:

$$|\psi_i\rangle = \begin{cases} |\Psi_0\rangle & \text{for } i \leq p \\ \sqrt{x_i}|\Psi_0\rangle + \sqrt{1-x_i}|\Psi_i\rangle & \text{for } i > p \end{cases} \quad (18)$$

Here, again all  $x_i$  are independent variables according to the same distribution and  $\langle\Psi_i|\Psi_j\rangle = \delta_{ij}$ . The overlap matrix  $\mathcal{S}$  then becomes:

$$\mathcal{S}_{ij} = \begin{cases} 1 & \text{for } i = j \text{ and } i, j \leq p \\ \sqrt{x_i}\sqrt{x_j}^* & \text{for } (i \neq j \text{ and } i, j > p) \\ \sqrt{x_i} & \text{for } (i > p \text{ and } j \leq p) \\ \sqrt{x_j}^* & \text{for } (i \leq p \text{ and } j > p) \end{cases} \quad (19)$$




FIG. 2. Solutions to Eqs. (7) (solid) and (17) (dashed) for two different upper bounds on the  $L_1$ -distance (1% and 5%). We consider an i.i.d. orthogonal bad bit model where the standard deviation is given by  $\sigma = 0.02$ . Eq. (7) assumes  $\max(x_i) = \mu + 2\sigma$ . The lines indicate pairs of  $\mu$  and  $k$  for which an upper bound on the  $L_1$ -distance is given by 1% (blue) or 5% (orange).

We argue that this second case is experimentally relevant, because according to the algorithm as presented in section II it was a seemingly good strategy to spend a lot of resources to make a subset of all your single photons as good as possible, while neglecting the others.

With this overlap matrix in which  $p$  photons are indistinguishable and the others are independently sampled from the same distribution, we can again simplify Eq. (11). We do this by realizing that

$$\mathbb{E} \left( \left( \prod_{r=1}^n |\mathcal{S}_{r,\tau(r)}|^2 \right) \right) \leq \mathbb{E}(|x_i|^2)^{\max(j-p,0)}. \quad (20)$$

We can then simplify Eq. (11), following the same arguments as presented in Eqs. (A14), (A15) and (A16):

$$\begin{aligned} \text{Var}(Q_k) &< \\ \sum_{j=k+1}^n \mathbb{E}(|x_i|^2)^{\max(j-p,0)} R_{n,n-j} n! \sum_{p=0}^{n-j} R_{n-j,p} 2^p \left( \frac{1}{m^2} \right)^n \\ &\approx \sum_{j=k+1}^n \mathbb{E}(|x_i|^2)^{\max(j-p,0)} \frac{(n!)^2}{m^{2n}}. \end{aligned} \quad (21)$$

If we now consider  $k+1 > p$  and if we again realize that this is now a truncated geometric series, we can find an upper bound for the expectation value of the  $L_1$ -

distance.

$$\begin{aligned} \mathbb{E} \left( \sum_s |P(s) - P_k(s)| \right) &\leq \\ \binom{m}{n} \sqrt{\sum_{j=k+1}^n \mathbb{E}(|x_i|^2)^{j-p} \frac{n!^2}{m^{2n}}} &< \\ \sqrt{\sum_{j=k-p+1}^n \mathbb{E}(|x_i|^2)^j} &< \\ \sqrt{\frac{\mathbb{E}(|x_i|^2)^{k-p+1}}{1 - \mathbb{E}(|x_i|^2)}} & \end{aligned} \quad (22)$$

Let's look at Eq. (22). Adding  $p$  perfectly indistinguishable photons can be negated by truncating at  $k' = k + p$  instead of  $k$ .




FIG. 3. Solutions to Eq. (22) for an  $L_1$ -distance of 1%. We consider an i.i.d. orthogonal bad bit model where the standard deviation is given by  $\sigma = 0.02$  and in addition,  $p$  photons are perfectly indistinguishable. The lines indicate pairs of  $\mu$  and  $k$  for which an upper bound on the  $L_1$ -distance is given by 1%.

#### IV. DISCUSSION AND CONCLUSION

We have extended classical simulation techniques for noisy boson sampling. These new classical simulation techniques push the boundaries for the required qualities of the resources needed for an optical quantum computer. Our results strengthen the intuition that the fragility of computational complexity to noise is itself a robust phenomenon, that does not depend on the particular details of how that noise is modeled. Future work will focus on including other inhomogeneous noise sources into this model, such as unbalanced optical loss.

### Appendix A: Formal derivation for the upper bound on the $L_1$ -distance

In this section we will give a formal derivation of the upper bound as presented in Eq. (7). We will follow the steps as presented in the main text.

We start by realizing that  $Q_k$  from Eq. (6) is real, we know  $|Q_k| = \sqrt{Q_k^2}$ . Jensen's inequality for a concave function [42] yields

$$\mathbb{E}(|Q_k|) = \mathbb{E}(\sqrt{Q_k^2}) \leq \sqrt{\mathbb{E}(Q_k^2)} \quad (\text{A1})$$

We note

$$\begin{aligned} & \mathbb{E}(\text{Perm}(M \circ M_\tau^*)) = \\ & \mathbb{E}\left(\sum_{\rho \in S_n} \prod_{i=1}^n (M \circ M_\tau^*)_{i,\rho(i)}\right) = \\ & \sum_{\rho \in S_n} \mathbb{E}\left(\prod_{i=1}^n (M \circ M_\tau^*)_{i,\rho(i)}\right), \end{aligned} \quad (\text{A2})$$

where we used the definition of the permanent and the linearity of the expectation value. We are interested in the situation where our truncation parameter  $k$  is larger than zero, and as a result, all permutations  $\tau$  that we consider have a nonzero amount of points which are not fixed. In other words:

$$\exists q \text{ s.t. } \tau(q) \neq q \forall \tau \quad (\text{A3})$$

and Eq. (A2) can be written as

$$\begin{aligned} & \sum_{\rho \in S_n} \mathbb{E}\left(\prod_{i=1}^n (M \circ M_\tau^*)_{i,\rho(i)}\right) = \\ & \sum_{\rho \in S_n} \mathbb{E}\left(\prod_{\substack{i=1 \\ i \neq q}}^n (M \circ M_\tau^*)_{i,\rho(i)}\right) \mathbb{E}(M_{q,\rho(q)}) \mathbb{E}(M_{\tau(q),\rho(q)}^*) \\ & = 0. \end{aligned} \quad (\text{A4})$$

Here we used the fact that  $\mathbb{E}(XY) = \mathbb{E}(X)\mathbb{E}(Y)$  if  $X$  and  $Y$  are independent variables and we note that  $M_{q,\rho(q)}$  and  $M_{\tau(q),\rho(q)}^*$  are independent of all other factors. We further use that the expectation of our i.i.d. complex Gaussian elements is zero,  $\mathbb{E}(M_{ij}) = 0$  for all  $ij$ . If we again use the linearity of the expectation value, we find that

$$\mathbb{E}(Q_k) = \mathbb{E}\left(\sum_{j=k+1}^n \sum_{\tau \in \sigma_j} x^j \text{Perm}(M \circ M_\tau^*)\right) = 0, \quad (\text{A5})$$

hence

$$\mathbb{E}(|Q_k|) \leq \sqrt{\mathbb{E}(Q_k^2) - \mathbb{E}(Q_k)^2} = \sqrt{\text{Var}(Q_k)}. \quad (\text{A6})$$

The square root of the variance thus provides an upper bound for  $Q_k$ . We derive an expression for  $\text{Var}(Q_k)$  in the remainder of this section.

Using Bienaymé's identity [43] we find

$$\begin{aligned} & \text{Var}(Q_k) \\ &= \text{Var}\left(\sum_{j=k+1}^n \sum_{\tau \in \sigma_j} x^j \text{Perm}(M \circ M_\tau^*)\right) \\ &= \text{Var}\left(\sum_{j=k+1}^n \sum_{\tau \in \sigma_j} x^j \sum_{\rho \in S_n} \prod_{r=1}^n (M \circ M_\tau^*)_{r,\rho(r)}\right) \\ &= \text{Var}\left(\sum_{j=k+1}^n \sum_{\tau \in \sigma_j} x^j \sum_{\rho \in S_n} \prod_{r=1}^n (M_{r,\rho(r)} M_{\tau(r),\rho(r)}^*)\right) \\ &= \sum_{j=k+1}^n \sum_{j'=k+1}^n \sum_{\tau \in \sigma_j} \sum_{\tau' \in \sigma_{j'}} x^j x^{j'} \sum_{\rho \in S_n} \sum_{\rho' \in S_n} \dots \\ & \text{Cov}\left(\prod_{r=1}^n (M_{r,\rho(r)} M_{\tau(r),\rho(r)}^*), \prod_{r=1}^n (M_{r,\rho'(r)} M_{\tau'(r),\rho'(r)}^*)\right). \end{aligned} \quad (\text{A7})$$

We use the definition of the covariance between two complex random variables to find

$$\begin{aligned} & \text{Cov}\left(\prod_{r=1}^n M_{r,\rho(r)} M_{\tau(r),\rho(r)}^*, \prod_{r=1}^n M_{r,\rho'(r)} M_{\tau'(r),\rho'(r)}^*\right) \\ &= \mathbb{E}\left(\prod_{r=1}^n M_{r,\rho(r)} M_{\tau(r),\rho(r)}^* M_{r,\rho'(r)}^* M_{\tau'(r),\rho'(r)}\right) \dots \\ & \quad - \mathbb{E}\left(\prod_{r=1}^n M_{r,\rho(r)} M_{\tau(r),\rho(r)}^*\right) \times \\ & \quad \mathbb{E}\left(\prod_{r=1}^n M_{r,\rho'(r)}^* M_{\tau'(r),\rho'(r)}\right) \end{aligned} \quad (\text{A8})$$

If we focus on the second term in Eq. (A8), we realize that again, for all  $\tau$  that we will consider, this term evaluates to zero for the same reasons as given in Eqs. (A3) and (A4). Then Eq. (A8) reduces to

$$\begin{aligned} & \text{Cov}\left(\prod_{r=1}^n M_{r,\rho(r)} M_{\tau(r),\rho(r)}^*, \prod_{r=1}^n M_{r,\rho'(r)} M_{\tau'(r),\rho'(r)}^*\right) \\ &= \mathbb{E}\left(\prod_{r=1}^n M_{r,\rho(r)} M_{\tau(r),\rho(r)}^* M_{r,\rho'(r)}^* M_{\tau'(r),\rho'(r)}\right). \end{aligned} \quad (\text{A9})$$

We will continue to show that the expression in Eq. (A9) is equal to zero for almost all of the combinations of  $\tau$ ,  $\rho$ ,  $\tau'$  and  $\rho'$ . In this demonstration, it is crucial to assume that all elements of our submatrix  $M$  are i.i.d. complex

Gaussian, or:

$$M_{ij} \sim \mathcal{CN}(0, \frac{1}{m}) \forall i, j \quad (\text{A10})$$

From Eq. (A10) it follows that

1.  $E(M_{ij}) = 0$
2.  $E((M_{ij})^2) = E((M_{ij}^*)^2) = 0$
3.  $E(|M_{ij}|^2) = \frac{E(\chi_2^2)}{2m} = \frac{1}{m}$
4.  $E(|M_{ij}|^4) = \frac{2}{m^2}$

If we inspect the equations listed above, we note that Eq. (A9) will only evaluate to a nonzero amount when for all  $r$  one of the following conditions is true

1.  $r = \tau(r)$  and  $r = \tau'(r)$
2.  $\tau(r) = \tau'(r)$  and  $\rho(r) = \rho'(r)$

We note that the first condition is true for all  $r \in \text{fix}(\tau)$ , if  $\tau$  and  $\tau'$  share the same fixed points. For all non-fixed points, condition 2 must thus be true and we conclude that only if  $\tau = \tau'$  and  $\rho(q) = \rho'(q) \forall q \in \text{nonfix}(\tau)$ , Eq. (A9) will evaluate to a non-zero value. Here  $\text{fix}(\tau)$  and  $\text{nonfix}(\tau)$  denote the set of all fixed points of the permutation  $\tau$  and its complementary set respectively. Then, Eq.(A7) reduces to:

$$\begin{aligned} \text{Var}(Q_k) &= \sum_{j=k+1}^n \sum_{\tau \in \sigma_j} x^{2j} \sum_{\rho \in S_n} \sum_{\substack{\rho' \in S_n \\ \rho(q) = \rho'(q) \forall q \in \text{nonfix}(\tau)}} \dots \\ &E \left( \prod_{r=1}^n (M_{r, \rho(r)} M_{\tau(r), \rho(r)}^*) (M_{r, \rho'(r)}^* M_{\tau(r), \rho'(r)}) \right). \end{aligned} \quad (\text{A11})$$

We now split up the product.

$$\begin{aligned} \text{Var}(Q_k) &= \sum_{j=k+1}^n \sum_{\tau \in \sigma_j} x^{2j} \sum_{\rho \in S_n} \sum_{\substack{\rho' \in S_n \\ \rho(q) = \rho'(q) \forall q \in \text{nonfix}(\tau)}} \dots \\ &E \left( \prod_{r \in \text{fix}(\tau)} M_{r, \rho(r)} M_{\tau(r), \rho(r)}^* M_{r, \rho'(r)}^* M_{\tau(r), \rho'(r)} \right) \times \\ &E \left( \prod_{q \in \text{nonfix}(\tau)} M_{q, \rho(q)} M_{\tau(q), \rho(q)}^* M_{q, \rho'(q)}^* M_{\tau(q), \rho'(q)} \right) \end{aligned} \quad (\text{A12})$$

which, if we use that  $\tau(r) = r \forall r \in \text{fix}(\tau)$  and  $\rho(q) =$

$\rho'(q) \forall q \in \text{nonfix}(\tau)$ , reduces to

$$\begin{aligned} \text{Var}(Q_k) &= \sum_{j=k+1}^n \sum_{\tau \in \sigma_j} x^{2j} \sum_{\rho \in S_n} \sum_{\substack{\rho' \in S_n \\ \rho(q) = \rho'(q) \forall q \in \text{nonfix}(\tau)}} \dots \\ &E \left( \prod_{r \in \text{fix}(\tau)} |M_{r, \rho(r)}|^2 |M_{r, \rho'(r)}|^2 \right) \times \\ &E \left( \prod_{q \in \text{nonfix}(\tau)} |M_{q, \rho(q)}|^2 |M_{\tau(q), \rho(q)}|^2 \right). \end{aligned} \quad (\text{A13})$$

If we now realize that  $M_{q, \rho(q)}$  is independent of  $M_{\tau(q), \rho(q)}$  for all  $q \in \text{nonfix}(\tau)$ ,  $M_{r, \rho(r)}$  is equal to  $M_{r, \rho'(r)}$  if  $\rho'(r) = \rho(r)$  and independent otherwise,  $E(|M_{ij}|^2) = \frac{1}{m}$  and  $E(|M_{ij}|^4) = \frac{2}{m^2}$ , Eq. (A13) reduces to

$$\begin{aligned} \text{Var}(Q_k) &= \sum_{j=k+1}^n \sum_{\tau \in \sigma_j} x^{2j} \sum_{\rho \in S_n} \sum_{p=0}^{n-j} \dots \\ &R_{n-j, p} \left( \frac{1}{m^2} \right)^j \left( \frac{2}{m^2} \right)^p \left( \frac{1}{m^2} \right)^{n-j-p} \\ &= \sum_{j=k+1}^n x^{2j} R_{n, n-j} n! \sum_{p=0}^{n-j} \dots \\ &R_{n-j, p} \left( \frac{1}{m^2} \right)^j \left( \frac{2}{m^2} \right)^p \left( \frac{1}{m^2} \right)^{n-j-p} \\ &= \sum_{j=k+1}^n x^{2j} R_{n, n-j} n! \sum_{p=0}^{n-j} R_{n-j, p} 2^p \left( \frac{1}{m^2} \right)^n. \end{aligned} \quad (\text{A14})$$

Here  $R_{n, k}$  is Rencontre's number that counts the number of ways one can permute the set  $\{1, \dots, n\}$  with  $k$  fixed points.  $R_{n, k} = \frac{n!}{k!} \sum_{q=0}^{n-k} \frac{(-1)^q}{q!}$ , and Eq. (A14) becomes:

$$\begin{aligned} \text{Var}(Q_k) &= \dots \\ &\sum_{j=k+1}^n x^{2j} \frac{(n!)^2}{m^{2n}} \sum_{q=0}^j \frac{(-1)^q}{q!} \sum_{p=0}^{n-j} \frac{2^p}{p!} \sum_{r=0}^{n-j-p} \frac{(-1)^r}{r!}. \end{aligned} \quad (\text{A15})$$

Now  $\sum_{q=0}^j \frac{(-1)^q}{q!} \approx \frac{1}{e}$ ,  $\sum_{p=0}^{n-j} \frac{2^p}{p!} \approx e^2$  and  $\sum_{r=0}^{n-j-p} \frac{(-1)^r}{r!} \approx \frac{1}{e}$  when  $j$ ,  $n - j - p$  and  $n - j$  are large respectively. And thus  $\sum_{q=0}^j \frac{(-1)^q}{q!} \sum_{p=0}^{n-j} \frac{2^p}{p!} \sum_{r=0}^{n-j-p} \frac{(-1)^r}{r!} \approx 1$ . Note that for  $j = 4$ ,  $n - j = 5$  and  $n - j - p = 4$  these approximations already have errors below 2%. Hence,

$$\text{Var}(Q_k) \approx \sum_{j=k+1}^n x^{2j} \frac{(n!)^2}{m^{2n}}. \quad (\text{A16})$$



We realize that Eq. (A16) describes a truncated geometric series.

$$\begin{aligned} \text{Var}(Q_k) &\approx \frac{(n!)^2 x^{2k+2} - x^{2n+2}}{m^{2n} (1-x^2)} \\ &< \frac{(n!)^2 x^{2k+2}}{m^{2n} (1-x^2)} \end{aligned} \quad (\text{A17})$$

Finally, we use Eq. (A17) to find an upper bound for the expectation value of the  $L_1$ -distance between the approximate distribution and the real distribution over the Haar-unitaries. In the following expression the sum with

index  $s$ , runs over all non-collisional outputs.

$$\begin{aligned} \mathbb{E} \left( \sum_s |P(s) - P_k(s)| \right) &= \mathbb{E} \left( \sum_s |Q_k(s)| \right) \\ &= \sum_s \mathbb{E} (|Q_k(s)|) \\ &< \binom{m}{n} \frac{n!}{m^n} \sqrt{\frac{x^{2k+2}}{1-x^2}} \\ &< \sqrt{\frac{x^{2k+2}}{1-x^2}} \end{aligned} \quad (\text{A18})$$

- 
- [1] P. W. Shor, *SIAM Journal on Computing* **26**, 1484 (1997).  
[2] S. Lloyd, *Science* **273**, 1073 (1996).  
[3] L. K. Grover, in *Proceedings of the Twenty-Eighth Annual ACM Symposium on Theory of Computing* (Association for Computing Machinery, 1996), STOC '96, p. 212–219, ISBN 0897917855.  
[4] A. W. Harrow and A. Montanaro, *Nature* **549**, 203–209 (2017).  
[5] F. Arute, K. Arya, R. Babbush, D. Bacon, J. C. Bardin, R. Barends, R. Biswas, S. Boixo, F. G. Brandao, D. A. Buell, et al., *Nature* **574**, 505 (2019).  
[6] Y. Wu, W.-S. Bao, S. Cao, F. Chen, M.-C. Chen, X. Chen, T.-H. Chung, H. Deng, Y. Du, D. Fan, et al., *Phys. Rev. Lett.* **127**, 180501 (2021).  
[7] Q. Zhu, S. Cao, F. Chen, M.-C. Chen, X. Chen, T.-H. Chung, H. Deng, Y. Du, D. Fan, M. Gong, et al., *Science Bulletin* **67**, 240–245 (2022), ISSN 2095-9273.  
[8] H.-S. Zhong, H. Wang, Y.-H. Deng, M.-C. Chen, L.-C. Peng, Y.-H. Luo, J. Qin, D. Wu, X. Ding, Y. Hu, et al., *Science* **370**, 1460 (2020).  
[9] H. Wang, J. Qin, X. Ding, M.-C. Chen, S. Chen, X. You, Y.-M. He, X. Jiang, L. You, Z. Wang, et al., *Phys. Rev. Lett.* **123**, 250503 (2019).  
[10] H.-S. Zhong, Y.-H. Deng, J. Qin, H. Wang, M.-C. Chen, L.-C. Peng, Y.-H. Luo, D. Wu, S.-Q. Gong, H. Su, et al., *Phys. Rev. Lett.* **127**, 180502 (2021).  
[11] E. Pednault, J. A. Gunnels, G. Nannicini, L. Horesh, and R. Wisnieff, *Leveraging secondary storage to simulate deep 54-qubit sycamore circuits* (2019), arXiv:1910.09534.  
[12] F. Pan and P. Zhang, *Simulating the sycamore quantum supremacy circuits* (2021), arXiv:2103.03074.  
[13] J. Tindall, M. Fishman, E. M. Stoudenmire, and D. Sels, *PRX Quantum* **5**, 010308 (2024).  
[14] C. Oh, M. Liu, Y. Alexeev, B. Fefferman, and L. Jiang, *Classical algorithm for simulating experimental gaussian boson sampling* (2023), arXiv:2306.03709.  
[15] Anonymous, *Quantum disadvantage* (2024), Proc. SIGBOVIK 2024.  
[16] S. Aaronson and A. Arkhipov, in *Proceedings of the Forty-Third Annual ACM Symposium on Theory of Computing* (Association for Computing Machinery, New York, NY, USA, 2011), STOC '11, p. 333–342, ISBN 9781450306911.  
[17] D. J. Brod and M. Oszmaniec, *Quantum* **4**, 267 (2020).  
[18] M. C. Tichy, *Phys. Rev. A* **91**, 022316 (2015).  
[19] M. Lubasch, A. A. Valido, J. J. Renema, W. S. Kolthammer, D. Jaksch, M. S. Kim, I. Walmsley, and R. García-Patrón, *Phys. Rev. A* **97**, 062304 (2018).  
[20] D. Cilluffo, N. Lorenzoni, and M. B. Plenio, *Simulating gaussian boson sampling with tensor networks in the heisenberg picture* (2023), arXiv:2305.11215.  
[21] M. Liu, C. Oh, J. Liu, L. Jiang, and Y. Alexeev, *Phys. Rev. A* **108**, 052604 (2023).  
[22] B. Villalonga, M. Y. Niu, L. Li, H. Neven, J. C. Platt, V. N. Smelyanskiy, and S. Boixo, *Efficient approximation of experimental gaussian boson sampling* (2021), arXiv:2109.11525.  
[23] J. F. F. Bulmer, B. A. Bell, R. S. Chadwick, A. E. Jones, D. Moise, A. Rigazzi, J. Thorbecke, U.-U. Haus, T. V. Vaerenbergh, R. B. Patel, et al., *Sci. Adv.* **8** (2022).  
[24] S. Rahimi-Keshari, T. C. Ralph, and C. M. Caves, *Phys. Rev. X* **6**, 021039 (2016).  
[25] H. Qi, D. J. Brod, N. Quesada, and R. García-Patrón, *Phys. Rev. Lett.* **124**, 100502 (2020).  
[26] P. D. Drummond, B. Opanchuk, A. Delliios, and M. D. Reid, *Phys. Rev. A* **105**, 012427 (2022).  
[27] A. S. Delliios, M. D. Reid, and P. D. Drummond, *AAPPS Bulletin* **33** (2023), ISSN 2309-4710.  
[28] C. Oh, L. Jiang, and B. Fefferman, *Phys. Rev. Lett.* **131**, 010401 (2023).  
[29] J. J. Renema, A. Menssen, W. R. Clements, G. Triginer, W. S. Kolthammer, and I. A. Walmsley, *Phys. Rev. Lett.* **120**, 220502 (2018).  
[30] J. Renema, V. Shchesnovich, and R. Garcia-Patron, *Classical simulability of noisy boson sampling* (2018), arXiv:1809.01953.  
[31] D. Aharonov, X. Gao, Z. Landau, Y. Liu, and U. Vazirani, in *Proceedings of the 55th Annual ACM Symposium on Theory of Computing* (Association for Computing Machinery, New York, NY, USA, 2023), STOC 2023, p. 945–957.  
[32] G. Kalai and G. Kindler, *Gaussian noise sensitivity and bosonsampling* (2014), arXiv:1409.3093.  
[33] C.-K. Hong, Z.-Y. Ou, and L. Mandel, *Phys. Rev. Lett.* **59**, 2044 (1987).

- [34] J. J. Renema, H. Wang, J. Qin, X. You, C. Lu, and J. Pan, *Phys. Rev. A* **103**, 023722 (2021).
- [35] R. García-Patrón, J. J. Renema, and V. Shchesnovich, *Quantum* **3**, 169 (2019).
- [36] V. Shchesnovich, *Phys. Rev. A* **89**, 022333 (2014).
- [37] V. Shchesnovich, *Phys. Rev. A* **91**, 013844 (2015).
- [38] C. Sparrow, Ph.D. thesis, Imperial College London (2018).
- [39] R. D. Shaw, A. E. Jones, P. Yard, and A. Laing, *Errors in heralded circuits for linear optical entanglement generation* (2023), 2305.08452.
- [40] M. Jerrum, A. Sinclair, and E. Vigoda, *J. ACM* **51**, 671–697 (2004).
- [41] J. Marshall, *Phys. Rev. Lett.* **129**, 213601 (2022).
- [42] F. M. Dekking, C. Kraaikamp, H. P. Lopuhaä, and L. E. Meester, in *A Modern Introduction to Probability and Statistics: Understanding Why and How*, edited by F. M. Dekking, C. Kraaikamp, H. P. Lopuhaä, and L. E. Meester (Springer, London, 2005), Springer Texts in Statistics, pp. 103–114, ISBN 978-1-84628-168-6.
- [43] A. Klenke, in *Probability Theory: A Comprehensive Course*, edited by A. Klenke (Springer, London, 2014), Universitext, pp. 101–130, ISBN 978-1-4471-5361-0.