A reduction of the "cycles plus K_4 's" problem

Aseem Dalal^{*} Jessica McDonald[†] Songling Shan[‡]

Abstract

Let H be a 2-regular graph and let G be obtained from H by gluing in vertexdisjoint copies of K_4 . The "cycles plus K_4 's" problem is to show that G is 4colourable; this is a special case of the *Strong Colouring Conjecture*. In this paper we reduce the "cycles plus K_4 's" problem to a specific 3-colourability problem. In the 3-colourability problem, vertex-disjoint triangles are glued (in a limited way) onto a disjoint union of triangles and paths of length at most 12, and we ask for 3-colourability of the resulting graph.

1 Introduction

In this paper all graphs are assumed to be simple, unless explicitly stated otherwise. The reader is referred to [20] for standard terminology.

Given vertex-disjoint graphs G_1, \ldots, G_q , H with $|\bigcup_{1 \leq i \leq q} V(G_i)| \leq |V(H)|$, we glue G_1, \ldots, G_q onto H by defining an injective function $f : \bigcup_{1 \leq i \leq q} V(G_i) \to V(H)$, and then forming a new graph G with V(G) = V(H) and $E(G) = E(H) \cup \bigcup_{1 \leq i \leq q} E_i$, where $E_i = \{f(a)f(b) : ab \in E(G_i), f(a)f(b) \notin E(H)\}$. The graph G is said to have been obtained from H by gluing in G_1, \ldots, G_q . Consider the following question:

Question 1.1. Suppose that H is a graph with $\Delta(H) \leq 2$, and suppose that G is obtained from H by gluing on vertex-disjoint triangles. When is $\chi(G) \leq 3$?

For Question 1.1, if H contains a C_4 , then certainly a K_4 may be created in G which would make $\chi(G) \not\leq 3$. Having C_4 's in H is not the only thing that could go wrong however: Fleischner and Stiebitz [7] found an infinite family of examples where H does not contain any C_4 components, but where $\chi(G) \not\leq 3$, answering a question of Erdős [5]. The smallest of Fleishner and Stiebitz's examples has $H = C_5 \cup C_{10}$; other known graphs H with $\Delta(H) = 2$ but which can yield negative answers to Question 1.1 (i.e. there is a way to glue on vertex-disjoint triangles that gives a 4-chromatic graph) include $H = C_3 \cup C_6$ (Öhman [18]) and $H = C_7 \cup K_1 \cup K_1$ (Sachs, see [7]); for much more, see

^{*}Indian Institute of Technology Delhi, Department of Mathematics, Delhi India. Email: aseem.dalal@gmail.com.

[†]Auburn University, Department of Mathematics and Statistics, Auburn U.S.A. Email: mcdonald@auburn.edu. Supported in part by Simons Foundation Grant #845698

[‡]Auburn University, Department of Mathematics and Statistics, Auburn U.S.A. Email: szs0398@auburn.edu. Supported in part by NSF grant DMS-2345869.

Öhman. On the other hand, Fleishner and Stiebitz [8] (and later Sachs [19]) famously provided a positive answer to Question 1.1: they proved that if H is a single cycle then the G obtained is always 3-colourable. This "cycle plus triangles" theorem answered another question of Erdős, which also had origins in the work of Du, Hsu, and Hwang [4] (see [8] for more history).

If we hope for a positive answer to Question 1.1 for some H which has multiple components, it seems wise to avoid cycles of length at least four in H, and instead consider H to be a disjoint union of paths and triangles. Here there are two extremes, both of which always yield positive answers to Question 1.1. If H is a disjoint union of paths, then by joining all these paths into a cycle, we form a "cycle plus triangles" graph G which is 3-colourable (by Felishner and Stiebitz [8]). On the other hand if H is a disjoint union of triangles, then we can form a 3-regular auxiliary bipartite graph which describes the interaction between triangles of H and added triangles, and the 3edge-colourability of this graph implies the 3-colourability of G. Things become more difficult when glued-in triangles join both path and triangle components in H. For G, Has in Question 1.1, we say that in G, two triangles T, T' in H are path-linked if there exists at least one glued-in triangle Y with $T \cap Y, T' \cap Y \neq \emptyset$, and the third vertex of Y is on some path in G of length at least two. If such gluings are strictly limited - so that in G, every H-triangle is path-linked to at most one other H-triangle, then we conjecture that G is 3-colourable. In fact, we also restrict our path components in H to length at most 12 to eliminate other potential problems; note that if all H-paths have length at most 2 then our auxiliary bipartite argument above again works to show 3-colourability.

Conjecture 1.2. Let H be a graph which is a disjoint union of triangles and paths of length at most 12, and let G be obtained from H by gluing on vertex-disjoint triangles. Moreover, suppose that in G, every H-triangle is path-linked to at most one other H-triangle. Then G is 3-colourable.

Given all its restrictions, we hope that Conjecture 1.2 may be approachable. Our main result in this paper is that the truth of Conjecture 1.2 would imply a seemingly very difficult conjecture, the "cycles plus K_4 's problem" that we state here now.

Conjecture 1.3. Let H be a graph with $\Delta(H) \leq 2$ and let G be obtained from H by gluing on vertex-disjoint copies of K_4 . Then $\chi(G) \leq 4$.

Note that Conjecture 1.3, as stated above, is not just "cycles plus K_4 's", but rather it allows paths in the base graph as well. However it is easy to see that it suffices to prove Conjecture 1.3 for 2-regular H (other H being subgraphs of these); hence the nickname. Our main result of this paper is concretely the following.

Theorem 1.4. Conjecture 1.2 implies Conjecture 1.3.

While we reduce Conjecture 1.3 to Conjecture 1.2 in this paper, we do not show that they are equivalent. However, an equivalence cannot be so far away from the truth. If Conjecture 1.3 holds, then by deleting any one of the four colour classes we get a 3-colourable graph G', and G' is obtained by gluing vertex-disjoint triangles onto a graph H' which is a disjoint union of paths and cycles.

To give some context for Conjecture 1.3, we must mention the strong chromatic number of a graph H, denoted $s\chi(H)$, which was introduced independently by Alon [2] and Fellows [6] in the late 1980's. Skipping this definition, let us just say that the *Strong Colouring Conjecture* posits that $s\chi(H) \leq 2\Delta(H)$; exact attribution of the conjecture is tricky, but the 1995 book of Jensen and Toft [13] (Section 4.14) has more on the early history of strong colouring. Support for the Strong Colouring Conjecture has been given by numerous papers, eg. Haxell [11][12], by Aharoni, Berger and Ziv [1], Axenovich and Martin [3], Johansson, Johansson and Markström [14], and Lo and Sanhueza-Matamala [15]. On the other hand, while the truth of the Strong Colouring Conjecture is trivial for $\Delta(H) = 1$, (where it asks essentially for the union of two matchings to be bipartite), the $\Delta(H) = c$ case is open for all constants $c \geq 2$. In the most glaring open case of $\Delta(H) = 2$, a result of Haxell [11] says that $s\chi(H) \leq 5$, but the conjecture is precisely Conjecture 1.3 (see e.g. [17]). The only cases of Conjecture 1.3 known to be true is when H has at most one odd cycle of length exceeding 3, or H has at most 3 triangles (McDonald and Puelo [17]). It seems that a new approach is needed to make a breakthrough on this problem, and our hope is that the reduction to Conjecture 1.2 may help.

This paper proceeds as follows. The following section discusses so-called *independent sets of representatives* (ISRs), and states two results of Haxell [9][10] that will be needed for our main proof. We also prove a result about combining two ISRs into one, which generalizes a prior theorem of the second author and Puleo [17], and may be of independent interest. The third section of the paper contains our proof of Theorem 1.4.

2 Independent Sets of Representatives

If H is a graph and let V_1, \ldots, V_n be disjoint subsets of V(H). An independent set of representatives (ISR) of $\{V_1, \ldots, V_n\}$ in G is set $R \subseteq V(G)$ such that R is independent in G and R contains exactly one vertex from each set V_i . If $R \subseteq V(G)$ is independent in G and contains at most one vertex from each set V_i , then R is said to be a partial ISR; R is said to hit those V_i for which it contains a representative. Note that if R is a partial ISR of $\{V_1, \ldots, V_n\}$ in G then it is an ISR of $\{V_i : R \text{ hits } V_i\}$.

Haxell has proved the following.

Theorem 2.1 (Haxell [10]). If H is a graph with $\Delta(H) \leq 2$ and V_1, \ldots, V_n are disjoint subsets of V(H) with each $|V_i| \geq 4$, then (V_1, \ldots, V_n) has an ISR.

To state a second result of Haxell, we need a few additional definitions. To this end, a *total dominating set* in a graph G is a set of vertices X such that every vertex in G is adjacent to a vertex in X. In particular, every vertex of X must also have a neighbor in X. The *total domination number* of G, written $\bar{\gamma}(G)$, is the size of a smallest total dominating set; if G has isolated vertices, then by convention we set $\bar{\gamma}(G) = \infty$. Note that by this definition, every (nonempty) graph has total domination number at least two. Given a graph H and disjoint subsets $V_1, \ldots, V_n \subset V(H)$, for each $S \subset [n]$ we define a subgraph H_S by taking the subgraph induced by the vertex set $\bigcup_{i \in S} V_i$ and deleting all edges in $H[V_i]$ for every $i \in S$.

Theorem 2.2 (Haxell [9]). Let H be a graph and let V_1, \ldots, V_n be disjoint subsets of V(H). If, for all $S \subset [n]$, we have $\bar{\gamma}(H_S) \geq 2|S| - 1$, then (V_1, \ldots, V_n) has an ISR.

It is worth noting we have stated Theorem 2.2 as it appeared in [17] – the original formulation was in terms of hypergraphs.

Haxell's Theorems 2.1 and 2.2 are both about the existence of a single ISR. We now prove a theorem about combining two ISRs into one. To this end, let $\mathcal{X} = \{X_1, \ldots, X_p\}$ and $\mathcal{Y} = \{Y_1, \ldots, Y_q\}$ be two collections of pairwise disjoint subsets of V(G), and suppose that $R_{\mathcal{X}}, R_{\mathcal{Y}}$ are ISRs of \mathcal{X}, \mathcal{Y} , respectively, in G. If $G[R_{\mathcal{X}} \cup R_{\mathcal{Y}}]$ is an independent set of vertices, then we view this union $R_{\mathcal{X}} \cup R_{\mathcal{Y}}$ as the successful combination of two ISRs into one, since it is simultaneously an ISR of both \mathcal{X} and \mathcal{Y} . In fact, we would be just as happy if we could find $R \subseteq R_{\mathcal{X}} \cup R_{\mathcal{Y}}$ that was simultaneously an ISR of both \mathcal{X} and \mathcal{Y} . This won't always be possible, but we will look to delete vertices from $R_{\mathcal{X}} \cup R_{\mathcal{Y}}$ so that the resulting R is independent and is somehow close to hitting all the sets in both \mathcal{X} and \mathcal{Y} . We say that an edge $e \in G[R_{\mathcal{X}} \cup R_{\mathcal{Y}}]$ is a \mathcal{X} -edge $(\mathcal{Y}$ -edge) if both endpoints of e are in X_i (Y_i) for some $X_i \in \mathcal{X}$ $(Y_i \in \mathcal{Y})$; note that it is possible for an edge to be both a \mathcal{X} -edge and a \mathcal{Y} -edge. We define $E_{\mathcal{X}\mathcal{Y}}$ to be all those edges of $G[R_{\mathcal{X}} \cup R_{\mathcal{Y}}]$ that are neither \mathcal{X} -edges nor \mathcal{Y} -edges.

Theorem 2.3. Let G be a graph. Suppose that $R_{\mathcal{X}}$ is an ISR of \mathcal{X} in G and $R_{\mathcal{Y}}$ is an ISR of \mathcal{Y} in G. For all $X \in \mathcal{X}$, denote by v_X the representative of X in $R_{\mathcal{X}}$. Then G has an independent set $R \subseteq R_{\mathcal{X}} \cup R_{\mathcal{Y}}$ that is an ISR of \mathcal{Y} and such that for every $X \in \mathcal{X}$:

- (a) if v_X is not incident to any E_{XY} -edges, then R hits X, and;
- (b) if v_X is incident to at least one E_{XY} -edge, then R hits $X \cup \{w\}$ for some E_{XY} -edge wv_X .

Before we proceed to the proof of Theorem 2.3, it may be remarked that if $E_{\mathcal{X}\mathcal{Y}} = \emptyset$, then Theorem 2.3 provides an independent set R which is an ISR of both \mathcal{X} and \mathcal{Y} . In this way Theorem 2.3 is a generalization of Lemma 3.3 of the second author and Puleo in [17] (their hypothesis that $(\mathcal{X}, \mathcal{Y})$ is an "admissible-pair" in G implies that $E_{\mathcal{X}\mathcal{Y}} = \emptyset$). Let us now prove Theorem 2.3.

Proof of Theorem 2.3. Initially, let $R_0 = R_{\mathcal{X}} \cup R_{\mathcal{Y}}$. The set R_0 clearly hits every $X_i \in \mathcal{X}$ and $Y_j \in \mathcal{Y}$ in G, but as there may be edges between $R_{\mathcal{X}}$ and $R_{\mathcal{Y}}$, the set R_0 may not be independent. We will describe an algorithm for iteratively deleting vertices from R_0 in order to obtain an independent subset of R_0 which still hits every $Y \in \mathcal{Y}$ and also meets conditions (a) and (b) for each $X \in \mathcal{X}$. First however, let us prove the following claim.

Claim 1. Every vertex of $R_{\mathcal{X}}$ is incident to at most one \mathcal{Y} -edge, and every vertex of $R_{\mathcal{Y}}$ is incident to at most one \mathcal{X} -edge.

Proof of Claim. Suppose that $u \in R_{\mathcal{X}}$ and that uv_1 , uv_2 are two different \mathcal{Y} -edges incident to u. Then $v_1, v_2 \in R_{\mathcal{Y}}$ with $\{u, v_1\} \subseteq Y_1$ and $\{u, v_2\} \subseteq Y_2$ for some $Y_1, Y_2 \in \mathcal{Y}$. Since the sets in \mathcal{Y} are pairwise vertex-disjoint, this implies that $Y_1 = Y_2$. But since $v_1, v_2 \in R_{\mathcal{Y}}$, we must have $Y_1 \neq Y_2$, contradiction. The same argument, interchanging the roles of \mathcal{X} and \mathcal{Y} , proves the claim about \mathcal{X} -edges.

Our algorithm defines a sequence of vertex sets R_0, R_1, \ldots starting with $R_0 = R_{\mathcal{X}} \cup R_{\mathcal{Y}}$. Given some set R_i , we either produce a new set R_{i+1} and proceed to the next round, or produce the final set R, via the following algorithm.

Step 1. If there is a vertex $v \in R_{\mathcal{Y}}$ which is isolated in $G[R_i] \setminus E_{\mathcal{X}\mathcal{Y}}$ but is incident to least one $E_{\mathcal{X}\mathcal{Y}}$ -edge in $G[R_i]$, then: form R_{i+1} from R_i by deleting all vertices that are adjacent to v in $G[R_i]$, and then go back to the start of **Step 1**.

Step 2. If there is a vertex $v \in R_{\mathcal{Y}}$ which has degree 1 in $G[R_i] \setminus E_{\mathcal{XY}}$, and this one edge is a \mathcal{X} -edge, then: form R_{i+1} from R_i by deleting all vertices that are adjacent to v in $G[R_i]$, and then go back to the start of **Step 1**.

Step 3. If there is a vertex $v \in R_{\mathcal{X}}$ which has degree 1 in $G[R_i]$, and this one edge is a \mathcal{Y} -edge, then: form R_{i+1} from R_i by deleting the one vertex in R_i that is adjacent to v in $G[R_i]$, and then go back to the start of **Step 1**.

Step 4. Otherwise, obtain R from R_i by deleting every vertex of $R_{\mathcal{Y}} \cap R_i$ that has positive degree in $G[R_i]$, and then terminate.

We call any vertex v found in **Steps** 1–3 above a *dangerous* vertex (for R_i), and only reach Step 4 when R_i has no dangerous vertices. Note that vertices which were not initially dangerous for R_0 may become dangerous for some later R_i as their neighbors are deleted. However, once a dangerous vertex v is found in one of **Steps** 1–3, all of v's neighbours in $G[R_i]$ are deleted, and hence v will be isolated in $G[R_j]$ for all $j \ge i$, and will remain until the end of the algorithm and be a member of our terminal R.

The algorithm always terminates, since $|R_{i+1}| < |R_i|$ whenever R_i has a dangerous vertex. Moreover, the terminal set R is always independent due to **Step 4**. It remains to show that R hits every set $Y \in \mathcal{Y}$ and also meets conditions (a) and (b) for each $X \in \mathcal{X}$.

Consider any set $Y \in \mathcal{Y}$. Let w be the representative of Y in $R_{\mathcal{Y}}$. If $w \in R$, then R hits Y and we are done. Therefore, we suppose that w was deleted by the algorithm. If $w \in R_{\mathcal{Y}} \cap R_{\mathcal{X}}$ then w is isolated in $G[R_0]$ and never gets deleted. So $w \in R_{\mathcal{Y}} \setminus R_{\mathcal{X}}$ and hence must have been deleted in **Step 3** or **Step 4**.

First suppose that w was deleted in **Step 3** due to v being a dangerous vertex for R_i . Then, based on our earlier comments, $v \in R$. Since the deletion of w happened in **Step 3**, we know that vw is a \mathcal{Y} -edge, so $v, w \in Y'$ for some $Y' \in \mathcal{Y}$. But the sets in \mathcal{Y} are disjoint and $w \in Y$, so in fact Y' = Y, and v ensures that R hits Y.

Now assume that w was deleted in **Step 4** after determining that R_i has no dangerous vertices. We know that w is not isolated in $G[R_i]$, since otherwise it would not have been deleted in **Step 4**. So since w is not dangerous for R_i according to **Step 1**, it must be incident to at least one \mathcal{X} -edge or \mathcal{Y} -edge in $G[R_i]$. In fact, since w is not dangerous for R_i according to **Step 2**, and given Claim 1, w must have either degree one or two in $G[R_i]$; in the former case its incident edge is a \mathcal{Y} -edge and in the latter case it is incident to both a \mathcal{X} -edge and a \mathcal{Y} -edge (which are distinct). In either case we know that there is a \mathcal{Y} -edge incident to w in $G[R_i]$, say wu. Since $w \in Y$ we get that $u \in Y$ as well. Since w is the lone representative for Y in $R_{\mathcal{Y}}$ (or since $u \sim w$ and $R_{\mathcal{Y}}$ is independent) we know that $u \notin R_{\mathcal{Y}}$, so u is not deleted in this **Step 4**. But since this is the very last step of our algorithm, no subsequent step could have deleted u either, so $u \in R$ at the end. Hence R hits Y.

Now consider any $X \in \mathcal{X}$. If $v_X \in R$, then R hits X (and thus hits $X \cup \{u\}$ for every $E_{\mathcal{X}\mathcal{Y}}$ -edge uv_X in G) and we are done. If $v_X \in R_{\mathcal{X}} \cap R_{\mathcal{Y}}$ then v_X is isolated in $G[R_0]$ and never gets deleted. Therefore, we suppose v_X was deleted by the algorithm and $v_X \in R_{\mathcal{X}} \setminus R_{\mathcal{Y}}$. Hence, v_X must have been deleted in **Step 1** or **Step 2**. First assume that v_X was deleted in **Step 1** due to w being a dangerous vertex for R_i . Then, based on our earlier comments, $w \in R$. But then v_X, w are joined by an E_{XY} -edge and R hits $X \cup \{w\}$.

Now we may assume that v_X was deleted in **Step 2** due to w being a dangerous vertex for R_i . Then, again based on our earlier comments, $w \in R$. We know that either $v_X w$ is an \mathcal{X} -edge or $v_X w$ is an $E_{\mathcal{X}\mathcal{Y}}$ -edge. In the latter case, R hits $X \cup \{w\}$. In the former case, $w \in X$ and so R hits X. Either way, conditions (a) and (b) are satisfied. \Box

3 Proof of Theorem 1.4

Within our proof of Theorem 1.4, we will find occasion to use the following classic result of Lovász [16] (see also Theorem 20 in [21]).

Theorem 3.1 (Lovász [16]). Let d, D be a non-negative integers and let $k = \lfloor \frac{D}{d+1} \rfloor + 1$. If G is a graph with maximum degree D, then V(G) can be partitioned onto k sets V_1, \ldots, V_k such that $G[V_i]$ has maximum degree at most d for all $i \in \{1, 2, \ldots, k\}$.

Let us now proceed with our main proof.

Proof. (Theorem 1.4) Let H be a graph with $\Delta(H) \leq 2$, and let G be obtained from H by gluing on vertex-disjoint copies of K_4 . We show that, under the assumption that Conjecture 1.2 is true, $\chi(G) \leq 4$. As previously discussed, we may assume that H is 2-regular.

Let X_1, \ldots, X_p be the vertex sets of the components of H and let Y_1, \ldots, Y_q be the vertex sets of the added copies of K_4 . Let $\mathcal{X} = \{X_1, \ldots, X_p\}$ and let $\mathcal{Y} = \{Y_1, \ldots, Y_q\}$. Let $\mathcal{T} \subseteq \mathcal{X}$ correspond to those components of H which are triangles, with $t = |\mathcal{T}|$. For any $\mathcal{W} \subseteq \mathcal{X}$ define $G^{\mathcal{W}}$ to be the the graph with vertex set \mathcal{T} and where two vertices are joined by an edge if the corresponding X_i, X_j have the property: there exists (at least one) $Y \in \mathcal{Y}$ with $Y \cap X_i, Y \cap X_j, Y \cap X_k \neq \emptyset$ for some $X_k \in \mathcal{X}$ corresponding to a cycle of length at least four.

Choose \mathcal{M} such that:

- (M1) $\mathcal{M} \subseteq \mathcal{T};$
- (M2) $\Delta(G^{\mathcal{M}}) \leq 1;$
- (M3) $|\mathcal{M}|$ is maximum, subject to (M1) and (M2), and;
- (M4) $|E(G^{\mathcal{M}})|$ is minimum, subject to (M1), (M2), (M3).

We can prove the following about the size of \mathcal{M} .

Claim 2. $|\mathcal{M}| \geq \frac{t}{4}$.

Proof of Claim. Let $T \in \mathcal{T}$ and consider its degree in $G^{\mathcal{X}}$. Suppose $v \in T$ and $v \in Y \in \mathcal{Y}$. If Y contributes to the degree of T in $G^{\mathcal{X}}$, then at least one of the four vertices in Y must be a member of a non-triangle cycle in H. But in this case, Y contains vertices from at most two different members of \mathcal{T} , aside from T, and hence Y contributes at most two to the degree of T in $G^{\mathcal{X}}$. Since T has size three, this means that T has degree at most 6 in $G^{\mathcal{X}}$. Applying Theorem 3.1 to the graph $G^{\mathcal{X}}$ with D = 6 and d = 1 gives a partition of $V(G^{\mathcal{X}})$ into four parts, each of which induces a subgraph of maximum degree at most one. If we take the subset of \mathcal{T} corresponding to the largest of these parts, then it has at least t/4 elements.

We consider now the long cycles in H, which we intend to break into short pieces by deleting some vertices. In particular, we let $L = \{i : X_i \in \mathcal{X}, |X_i| \ge 15\}$. For each $i \in L$, if X_i consists of the vertex set of the cycle $(x_1, x_2, \ldots, x_{l_i})$, we let $p_i = \lfloor \frac{l_i}{5} \rfloor$ and define $X_i^* = \{x_{5(j)-4} : 1 \le j \le p_i\}$. We then let $X_i^1, \ldots, X_i^{p_i}$ be the vertex sets of the p_i cycle-segments created by deleting X_i^* , namely $X_i^j = \{x_{5j-3}, \ldots, x_{5j}\}$ for $1 \le j \le p_i - 1$ and $X_i^{p_i} = \{x_{5p_i-3}, \ldots, x_{l_i}\}$. Note that $|X_i^j| = 4$ for $1 \le j \le p_i - 1$ and $4 \le |X_i^{p_i}| \le 8$.

Let G' be the graph formed from G by deleting all the vertices in all the sets X_i^* , that is, $G' = G \setminus (\bigcup_{i \in L} X_i^*)$. We obtain \mathcal{X}' from \mathcal{X} by replacing each X_i , $i \in L$, with the p_i sets $X_i^1, \ldots X_i^{p_i}$. Let $\mathcal{X}' = \{X'_1, X'_2, \ldots, X'_{p'}\}$, noting that $p' \ge p = |\mathcal{X}|$. Note also that we still have $\mathcal{M} \subseteq \mathcal{X}'$, since triangles are not affected by the deletion process.

The goal in this next section of our proof will be to find an ISR of $\mathcal{X}' \setminus \mathcal{M}$ in the graph G'. Since we are unconcerned with hitting the triangle parts, it will serve us to form an auxiliary graph G'' by taking the disjoint union of G' along with $m = |\mathcal{M}|$ copies of K_m . For all $X'_i \notin \mathcal{M}$, define $X''_i = X'_i$ and otherwise define X''_i to be X'_i together with one vertex from each copy of K_m , chosen so that $X''_1, \ldots, X''_{p'}$ are disjoint, and let \mathcal{X}'' be this last collection of sets. We will aim, through the next two claims, to get an ISR of \mathcal{X}'' in G'' via Theorem 2.2. When restricted to G', such an ISR would contain a representative from each set in \mathcal{X}' except possibly those in \mathcal{M} . So we would indeed be able to get an ISR of $\mathcal{X}' \setminus \mathcal{M}$ in the graph G', as we have said we want.

In order to apply Theorem 2.2 (to get an ISR of \mathcal{X}'' in G''), we start by letting $S \subset [p']$, with \mathcal{S} the set of $X''_i \in \mathcal{X}''$ corresponding to S. Let $t_s = |\mathcal{T} \cap \mathcal{S}|$. We consider the graph G''_S defined with respect to $X''_1, \ldots, X''_{p'}$ (as discussed prior to the statement of Theorem 2.2). Note that the set of edges removed from G'' to make G''_S are exactly the same as the set of edges removed from G' to make G'_S , since each X''_i is obtained from X'_i by adding either an independent set or nothing. So G''_S is just the disjoint union of G'_S and m copies of $K_{|\mathcal{S} \cap \mathcal{M}|}$; in particular note that $G''_S = G'_S$ when $\mathcal{S} \cap \mathcal{M} = \emptyset$.

Claim 3. Let $\operatorname{comp}(G'_S)$ be the number of components of G'_S . Then in general, $\operatorname{comp}(G'_S) \geq \frac{1}{4}|V(G'_S)|$. Moreover, if $\mathcal{S} \cap \mathcal{M} = \emptyset$, then $\operatorname{comp}(G'_S) \geq \frac{1}{4}(|V(G'_S)| + t_s)$.

Proof of Claim. Since all edges of H are removed when forming G'_S , each component of G'_S has size at most four, and we immediately get the $\frac{1}{4}|V(G'_S)|$ bound. Now suppose that $S \cap \mathcal{M} = \emptyset$.

Let $T \in \mathcal{T} \cap S$. Then $T \notin \mathcal{M}$, by assumption. We claim that in $G^{\mathcal{X}}$ (note that this graph is formed prior to any deletions or breaking of long cycles), there are at least two triangles in \mathcal{M} that are each in some common glued K_4 with T. Certainly we cannot have no edges, since then $\tilde{\mathcal{M}} = \mathcal{M} \cup \{T\}$ has $|\tilde{\mathcal{M}}| > |\mathcal{M}|$, violating (M3). So suppose for a contradiction that there is exactly one such edge in $G^{\mathcal{X}}$, say from T to $\tilde{T} \in \mathcal{M}$. We know that \tilde{T} has at most one edge in $G^{\mathcal{X}}$ joining it to other triangles in \mathcal{M} . If it actually has no such edges, then again $\tilde{\mathcal{M}} = \mathcal{M} \cup \{T\}$ satisfies (M1) and (M2) but $|\tilde{\mathcal{M}}| > |\mathcal{M}|$ violating (M3). So \tilde{T} must have exactly one edge in $G^{\mathcal{X}}$ joining it to other triangles in \mathcal{M} ; now $\tilde{\mathcal{M}} = (\mathcal{M} \setminus {\tilde{T}}) \cup {T}$ satisfies (M1), (M2), (M3), but $|E(G^{\tilde{\mathcal{M}}})| < |E(G^{\mathcal{M}})|$, violating (M4). So indeed, there are at least two edges from T to \mathcal{M} in $G^{\mathcal{X}}$. Let E_T be such a pair of edges.

Either E_T comes from one $Y \in \mathcal{Y}$ containing vertices from both T and two different triangles in \mathcal{M} , or it comes from two different $Y^1, Y^2 \in \mathcal{Y}$, both of which contain a vertex from T and a triangle in \mathcal{M} . In all cases, by the definition of $G^{\mathcal{X}}$, each of Y, Y^1, Y^2 must contain at least one vertex from a cycle of H of size at least four; this cycle may or may not be in \mathcal{S} . Partition $\mathcal{S} \cap \mathcal{T}$ into $\mathcal{S}_1, \mathcal{S}_2$ so that for a given $T \in \mathcal{S} \cap \mathcal{T}, T \in \mathcal{S}_1$ if E_T comes from one $Y \in \mathcal{Y}$, and $T \in \mathcal{S}_2$ if E_T comes from two $Y^1, Y^2 \in \mathcal{Y}$.

If $T \in S_1$, the vertex in $T \cap Y$ is in a component of size at most 2 in G_S (since $\mathcal{M} \cap \mathcal{S} = \emptyset$). If $T \in S_2$, the two vertices in $T \cap Y', T \cap Y''$ are both in components of size at most 3 in G_S (since $\mathcal{M} \cap \mathcal{S} = \emptyset$). However it could be that the third vertex in such a component (other than the vertex in T and the vertex in the longer cycle) is another triangle in S_2 . Overall, for every triangle in S_1 we may count one component of G_S of size at most three (in fact size at most two), and for every triangle in S_2 we may count one component of size at most three. In fact, all these vertices in triangles still exist in G'_S (since we only delete vertices from long cycles), so the previous sentence remains true if we replace G_S with G'_S . Since we already know that all components in G'_S have size at most 4, we get:

$$\operatorname{comp}(G'_{S}) \geq \frac{1}{4}(|V(G'_{S})| - 2|S_{1}| - 3|S_{2}|) + |S_{1}| + |S_{2}|$$

$$= \frac{1}{4}|V(G'_{S})| + \frac{1}{2}|S_{1}| + \frac{1}{4}|S_{2}|$$

$$= \frac{1}{4}|V(G'_{S})| + \frac{1}{2}|S_{1}| + \frac{1}{4}|S_{2}|$$

$$\geq \frac{1}{4}|V(G'_{S})| + \frac{1}{4}(|S_{1}| + |S_{2}|) = \frac{1}{4}(|V(G'_{S})| + t_{s}).$$

We will now use Claim 3 as part of our proof of the following.

Claim 4. $\bar{\gamma}(G''_S) \geq 2|S|$.

Proof of Claim. We proceed by cases according to $|S \cap \mathcal{M}|$. If this value is one, then G''_S contains isolates, and the result follows trivially.

Suppose first that $|S \cap \mathcal{M}| \geq 2$. Then $\bar{\gamma}(K_{|S \cap \mathcal{M}|}) = 2$, and using the result of Claim 2, we get

$$\bar{\gamma}(G_S'') = \bar{\gamma}(G_S') + m \cdot \bar{\gamma}(K_{|\mathcal{S} \cap \mathcal{M}|}) = \bar{\gamma}(G_S') + 2m \ge \bar{\gamma}(G_S') + \frac{t}{2} \tag{1}$$

Since every component of G'_S has total domination number at least 2, by the first bound in Claim 3, we get

$$\bar{\gamma}(G'_S) \ge 2(|V(G'_S)|/4) = \frac{1}{2}|V(G'_S)|.$$
 (2)

On the other hand, all $X'_i \in \mathcal{X}'$ have at least three vertices, with the only parts of size three corresponding to triangles in H (recall that our broken cycle pieces have at least four vertices). It is possible that all t of the triangle parts are included in \mathcal{S} , but even still,

$$|V(G'_S)| \ge 4|S| - t.$$
 (3)

Combining (1), (2), and (3), we get

$$\bar{\gamma}(G_S'') \ge \frac{1}{2}(4|S| - t) + \frac{t}{2} = 2|S|$$

We may now assume that $S \cap \mathcal{M} = \emptyset$. Then $\bar{\gamma}(G''_S) = \bar{\gamma}(G'_S)$. Since every component of G'_S has total domination number at least 2, and by the second bound in Claim 3, we get

$$\bar{\gamma}(G'_S) \ge 2(\frac{1}{4}(|V(G'_S)| + t_s)) = \frac{1}{2}|V(G'_S)| + \frac{t_s}{2}.$$
(4)

We know that exactly t_s triangle parts are included in \mathcal{S} , so we get

$$|V(G'_S)| \ge 4|S| - t_s.$$
 (5)

Combining (4) and (5), we get

$$\bar{\gamma}(G''_S) = \bar{\gamma}(G'_S) \ge \frac{1}{2}(4|S| - t_s) + \frac{t_s}{2} = 2|S|.$$

By Claim 4, we know that $\bar{\gamma}(G_S'') \geq 2|S| - 1$, so we may finally apply Theorem 2.2 to get an ISR of \mathcal{X}'' in G''. As previously discussed, this in turn allows us to get an ISR of $\mathcal{X}' \setminus \mathcal{M}$ in G', which we have said was our goal for this section of the proof. Let $\tilde{\mathcal{X}} = \mathcal{X}' \setminus \mathcal{M}$, and call this last ISR $R_{\tilde{\mathcal{X}}}$. In fact, $R_{\tilde{\mathcal{X}}}$ is also an ISR of $\tilde{\mathcal{X}}$ in G, since no edges are added between vertices of $R_{\tilde{\mathcal{X}}}$ when moving back to G from G'.

Since $\Delta(H) \leq 2$ and $|Y_i| \geq 4$ for all *i*, Theorem 2.1 guarantees that \mathcal{Y} has an ISR in H, say $R_{\mathcal{Y}}$.

Since $R_{\mathcal{Y}}$ is an ISR of \mathcal{Y} in G, and the $R_{\tilde{\mathcal{X}}}$ is an ISR of $\tilde{\mathcal{X}}$ in G, we can now apply Theorem 2.3. The result is a set R that is independent in G, that is an ISR of \mathcal{Y} , and that hits $\bigcup_{v \in V(C)} X_v$ for each component C of the graph $\mathcal{G}_{\tilde{\mathcal{XY}}}$. Since R hits every set in $\mathcal{Y}, G - R$ is obtained from some graph $H' \subseteq H$ by gluing on triangles.

Claim 5. H' is a disjoint union of triangles and paths of length at most 12.

Proof of Claim. Consider a cycle in H of length at most 14 not belonging to \mathcal{M} , say represented by $X \in \mathcal{X}$. Such a cycle is unaltered when moving from G, \mathcal{X} to G', \mathcal{X}' so $X \in \mathcal{X}'$ and since $X \notin \mathcal{M}, X \in \tilde{\mathcal{X}}$. Suppose v is the representative of X in $R_{\tilde{\mathcal{X}}}$. Consider $\mathcal{G}_{\tilde{\mathcal{X}}\mathcal{Y}}$. Since v is the representative of X in $\tilde{\mathcal{X}}, X_v = X$. Now any vertex $w \in R_{\mathcal{Y}}$ adjacent to v in G either belongs to X (implying that vw is an $\tilde{\mathcal{X}}$ -edge) or both $w, v \in Y$ for some $Y \in \mathcal{Y}$ (implying that vw is an \mathcal{Y} -edge). Therefore, by definition, there is no $E_{\tilde{\mathcal{X}}\mathcal{Y}}$ edge incident to v. So by Theorem 2.3, R hits X. This means that in G - R, all that remains of the cycle represented by X in H (i.e. its contribution to H') is a subgraph of a path with at most 13 vertices (i.e. a path of length at most 12).

Consider now a cycle $(x_1, x_2, \ldots, x_{\ell_i})$ in H where $\ell_i \geq 15$, say represented by $X_i \in \mathcal{X}$. Suppose, for a contradiction, that there exists a path segment P of $\ell \geq 14$ vertices (so with length at least 13) which is a subgraph of this cycle and such that $V(P) \cap R = \emptyset$; without loss of generality suppose that $P = (x_1, \ldots, x_\ell)$

We know that $|X_i^j| = 4$ for $1 \le j \le p_i - 1$ and $4 \le |X_i^{p_i}| \le 8$. Moreover, any two X_i^j s are separated (along the path on cycle X_i) by exactly one vertex and that belongs to X_i^* . Now since $\ell > 8 + 1 + 4 = 13$, P contains a path segment $aX_i^j b$ such that $a, b \in X_i^*$ and $X_i^j \in \mathcal{X}'$. We know that $R_{\tilde{\mathcal{X}}}$ uniquely hits X_i^j . Let w_j be this representative for X_i^j , that is, $w_j = X_i^j \cap R_{\tilde{\mathcal{X}}}$.

If there is no $E_{\tilde{\chi}\mathcal{Y}}$ -edge incident to w_j , then by Theorem 2.3 R hits X_i^j , contradicting our assumption that $V(P) \cap R = \emptyset$. Therefore, there must be at least one $E_{\tilde{\chi}\mathcal{Y}}$ -edge incident to w_j ; let v be a second endpoint of such an edge. Note that $v \in \{a, b\}$ because if not then vw_j is either an $\tilde{\mathcal{X}}$ -edge or a \mathcal{Y} -edge. So, either v = a and w_j is the lowestindexed member of X_i^j , or v = b and w_j is the highest-indexed member of X_i^j . By symmetry, we can assume that v = a. There cannot be any other $E_{\tilde{\mathcal{X}}\mathcal{Y}}$ -edges incident to w_j , since $|X_i^j| \ge 4$ means that w_j is certainly not followed by b on P. Then by Theorem 2.3, R hits $X_i^j \cup \{a\} \subseteq V(P)$, contradiction.

Consider now the graph H' in the context of Claim 5. Note that every triangle in H' is from \mathcal{M} . By condition (M2), we know that $\Delta(G^{\mathcal{M}}) \leq 1$. In H' this means that if T_1 is a triangle in H', then there is at most one other triangle T_2 in H' such that they are both joined in G - R by some added K_3 , say Y', whose third vertex is from some cycle of length at least four in H.

In G-R, a pair of triangles in H' are path-linked iff there is some added K_3 containing vertices from both triangles as well as from a path in H' of length at least two. Since paths in H' of length at least two were all part of a cycle of length at least four in H (paths in H' coming from triangles in H have length one), the above paragraph tells us that in G - R every H'-triangle is path-linked to at most one other H'-triangle. So the truth of Conjecture 1.2 would give a 3-colouring of G - R, and hence a 4-colouring of G.

References

- Ron Aharoni, Eli Berger, and Ran Ziv, Independent systems of representatives in weighted graphs, Combinatorica 27 (2007), no. 3, 253–267.
- [2] N. Alon, The linear arboricity of graphs, Israel J. Math. 62 (1988), no. 3, 311–325.
- [3] Maria Axenovich and Ryan Martin, On the strong chromatic number of graphs, SIAM J. Discrete Math. 20 (2006), no. 3, 741–747.
- [4] D. Z. Du, D. F. Hsu, and F. K. Hwang, The Hamiltonian property of consecutive-d digraphs, vol. 17, 1993, Graph-theoretic models in computer science, II (Las Cruces, NM, 1988–1990), pp. 61–63.
- [5] Paul Erdős, On some of my favourite problems in graph theory and block designs, vol. 45, 1990, Graphs, designs and combinatorial geometries (Catania, 1989), pp. 61– 73 (1991).
- [6] Michael R. Fellows, Transversals of vertex partitions in graphs, SIAM J. Discrete Math. 3 (1990), no. 2, 206–215.
- [7] H. Fleischner and M. Stiebitz, Some remarks on the cycle plus triangles problem, The mathematics of Paul Erdős, II, Algorithms Combin., vol. 14, Springer, Berlin, 1997, pp. 136–142.
- [8] Herbert Fleischner and Michael Stiebitz, A solution to a colouring problem of P. Erdős, Discrete Math. 101 (1992), no. 1-3, 39–48, Special volume to mark the centennial of Julius Petersen's "Die Theorie der regulären Graphs", Part II.

- [9] P. E. Haxell, A condition for matchability in hypergraphs, Graphs Combin. 11 (1995), no. 3, 245–248.
- [10] _____, A note on vertex list colouring, Combinatorics, Probability & Computing 10 (2001), no. 4, 345.
- [11] _____, On the strong chromatic number, Combin. Probab. Comput. 13 (2004), no. 6, 857–865.
- [12] _____, An improved bound for the strong chromatic number, J. Graph Theory 58 (2008), no. 2, 148–158.
- [13] Tommy R. Jensen and Bjarne Toft, Graph coloring problems, Wiley-Interscience Series in Discrete Mathematics and Optimization, John Wiley & Sons, Inc., New York, 1995, A Wiley-Interscience Publication.
- [14] Anders Johansson, Robert Johansson, and Klas Markström, Factors of r-partite graphs and bounds for the strong chromatic number, Ars Combin. 95 (2010), 277– 287.
- [15] Allan Lo and Nicolás Sanhueza-Matamala, An asymptotic bound for the strong chromatic number, Combinatorics, Probability and Computing (2017).
- [16] L. Lovász, On decomposition of graphs, Studia Sci. Math. Hungar. 1 (1966), 237–238.
- [17] Jessica McDonald and Gregory J. Puleo, Strong coloring 2-regular graphs: cycle restrictions and partial colorings, J. Graph Theory 100 (2022), no. 4, 653–670.
- [18] Lars-Daniel Ohman, Strong chromatic numbers of graphs with maximum degree 2, unpublished manuscript, available at https://lars-daniel.se/papers/index.htm.
- [19] H. Sachs, Elementary proof of the cycle-plus-triangles theorem, Combinatorics, Paul Erdős is eighty, Vol. 1, Bolyai Soc. Math. Stud., János Bolyai Math. Soc., Budapest, 1993, pp. 347–359.
- [20] Douglas B. West, Introduction to graph theory, 2 ed., Prentice Hall, September 2000.
- [21] David R. Wood, Defective and clustered graph colouring, Electron. J. Combin. DS23 (2018), 71.