Spectral conditions implying the existence of doubly chorded cycles without or with constraints

Leyou Xu^{*}, Bo Zhou[†]

School of Mathematical Sciences, South China Normal University Guangzhou 510631, P.R. China

Abstract

What spectral conditions imply a graph contains a chorded cycle? This question was asked by R.J. Gould in 2022. We answer three modified versions of Gould's question by giving tight spectral conditions that imply the existence of doubly chorded cycle, a doubly chorded cycle with two chords incident to a vertex, and a doubly chorded cycle on five vertices with two chords incident to one vertex, in a graph, respectively.

Keywords: doubly chorded cycle, spectral condition, spectral radius

Mathematics Subject Classification: 05C50, 05C38

1 Introduction

One of the classical questions in combinatorics is as follows: What bounds for the size imply an *n*-vertex graph contains a subgraph with a certain prescribed structure? Its spectral version is: What spectral conditions imply an *n*-vertex graph contains a subgraph with a certain prescribed structure?

We consider only simple graphs. Let G be a graph with vertex set V(G) and edge set E(G). Let $\delta(G)$ denote the minimum degree of G. The adjacency matrix A(G) of G is the matrix $(a_{uv})_{u,v\in V(G)}$, where $a_{uv} = 1$ if u and v are adjacent and 0 otherwise. The largest eigenvalue of A(G) is called the spectral radius of G, denoted by $\rho(G)$. The above question may be rephrased via the spectral radius as: Among n-vertex graphs that do not contain a subgraph with a certain prescribed structure, what is the maximum spectral radius and which graphs achieve this value? Such questions, originated from [5] and called Brualdi–Solheid–Turán type problems in [20], received much attention in recent years, see, e.g. [8, 18, 20, 23, 27, 28].

^{*}Email: leyouxu@m.scnu.edu.cn

[†]Email: zhoubo@m.scnu.edu.cn

Let K_s denote the complete graph on s vertices. For positive integers n_1, \ldots, n_k with $k \ge 2$, let K_{n_1,\ldots,n_k} denote the complete k-partite graph with part sizes n_1, \ldots, n_k .

Let C be a cycle of a graph. We say an edge that joins two vertices of a cycle C is a chord of C if the edge is not itself an edge of C. If C has at least one chord, then it is called a chorded cycle, and if C has at least two chords, then it is called a doubly chorded cycle (DCC for short). The study of cycles in graphs is well established. The existence of cycles with additional properties such as containing at least s chords for some number $s \ge 1$ received much attention, see [7, 10, 12, 15] and references therein. For s = 1, 2, the cycles are just chorded cycles and doubly chorded cycles.

Given a graph G, Pósa [21] suggested the problem of finding degree conditions that imply the existence of a chorded cycle in G. Czipser proved (see problem 10.2, p. 65 in [19]) that $\delta(G) \ge 3$ suffices. Gould asked in [15] the following question: What spectral conditions imply a graph contains a chorded cycle? Zheng et al. [29] answered Gould's question by showing that any graph G of order $n \ge 6$ with $\rho(G) \ge \rho(K_{2,n-2})$ contains a chorded cycle unless $G \cong K_{2,n-2}$. Different answers are provided via the signless Laplacian spectral radius [24]. Gould et al. [16] observed that it is only slightly more difficult to show that $\delta(G) \ge 3$ implies G contains a DCC. So it is more natural to consider conditions implying the existence of DCCs [16, 22]. We try to answer the following three questions, which may be viewed as modified versions of Gould's original question.

Question 1.1. What spectral conditions imply a graph on n vertices contains a DCC?

The problem to determine the maximum size of an *n*-vertex graph that does not contain a cycle with many chords incident to a vertex received much attention. Through the work of Erdős, Lewin (see [1, p. 398, no. 12]) and Bollobás [1, p. 398, no. 12], Jiang [17] showed that for any *n*-vertex graph with $n \ge 9$ that does not contain a DCC with two chords incident to a vertex, $|E(G)| \le 3n - 9$, and it is attained if $G \cong K_{3,n-3}$.

We call a DCC with two chords incident to a vertex a DCC_1 .

Question 1.2. What spectral conditions imply a graph on n vertices contains a DCC_1 ?

Let $G \cup H$ be the disjoint union of graphs G and H. The disjoint union of k copies of a graph G is denoted by kG. The join of disjoint graphs G and H, denoted by $G \vee H$, is the graph obtained from $G \cup H$ by adding all possible edges between vertices in G and vertices in H. Note that $K_1 \vee P_4$ is the unique 5-vertex DCC₁. There are spectral conditions for a graph with fixed size which imply the existence of a copy of $K_1 \vee P_4$ [25]. There is also a paper considering the spectral extremal graphs of graphs containing no copies of $K_1 \vee P_{2k}$ with $k \ge 3$ for sufficiently large order [26]. Zhao et al. [28] determined the *n*-vertex graphs that maximize the spectral radius among graphs that do not contain any wheels $W_r := P_1 \vee C_r$ for $r \ge 3$. A graph that does not contain copies of $K_1 \vee P_4$ does not contain any wheels W_r for $r \ge 4$ but it may contain W_3 (i.e., K_4). A graph that does not contains any wheels may contain $K_1 \vee P_4$.

Question 1.3. What spectral conditions imply a graph on n vertices contains a copy of $K_1 \vee P_4$?

Questions 1.1, 1.2 and 1.3 are answered by the following Theorems 1.1, 1.2 and 1.3, respectively.

Theorem 1.1. Suppose that G is an n-vertex graph containing no DCCs, where $n \ge 3$. Then $\rho(G) \le \frac{1}{2} + \sqrt{2n - \frac{15}{4}}$ with equality if and only if $G \cong K_{1,1,n-2}$.

Let F_1 be the graph obtained from K_2 and K_4 by identifying a vertex.

Theorem 1.2. Suppose that G is an n-vertex graph containing no DCC_1s . Then

$$\rho(G) \leqslant \begin{cases}
n-1 & \text{if } n \leqslant 4 \\
3.0861 & \text{if } n = 5 \\
\frac{1}{2} + \sqrt{2n - \frac{15}{4}} & \text{if } 6 \leqslant n \leqslant 9 \\
\sqrt{3(n-3)} & \text{if } n \geqslant 10
\end{cases}$$

with equality if and only if

$$G \cong \begin{cases} K_n & \text{if } n \leq 4, \\ F_1 & \text{if } n = 5, \\ K_{1,1,n-2} & \text{if } 6 \leq n \leq 9, \\ K_{3,n-3} & \text{if } n \geqslant 10. \end{cases}$$
(1.1)

By Theorem 1.2, if G is a 5-vertex graph containing no copies of $K_1 \vee P_4$, then $\rho(G) \leq \rho(F_1)$ with equality if and only if $G \cong F_1$.

Theorem 1.3. Suppose that G is an n-vertex graph containing no copies of $K_1 \vee P_4$ with $n \ge 6$. Then

$$\rho(G) \leqslant \begin{cases} \frac{n+1}{2} & \text{if } n \equiv 1 \pmod{2} \\ \frac{1+\sqrt{n^2+1}}{2} & \text{if } n \equiv 0 \pmod{4} \\ \eta(n) & \text{if } n \equiv 2 \pmod{4} \end{cases}$$

with equality if and only if $G \cong H_n$, where

$$H_n = \begin{cases} \frac{n+1}{2}K_1 \vee \frac{n-1}{4}K_2 & \text{if } n \equiv 1 \pmod{4}, \\ \frac{n-1}{2}K_1 \vee \frac{n+1}{4}K_2 & \text{if } n \equiv 3 \pmod{4}, \\ \frac{n}{2}K_1 \vee \frac{n}{4}K_2 & \text{if } n \equiv 0 \pmod{4}, \\ \frac{n}{2}K_1 \vee (\frac{n-2}{4}K_2 \cup K_1) & \text{if } n \equiv 2 \pmod{4}, \end{cases}$$

and $\eta(n)$ is the largest root of $x^3 - x^2 - \frac{n^2}{4}x + \frac{n}{2} = 0$.

2 Preliminaries

Let G be a graph. For $v \in V(G)$, we denote by $N_G(v)$ the neighborhood of v in G, and $d_G(v)$ the degree of v in G. Let $N_G[v] := \{v\} \cup N_G(v)$ be the closed neighborhood of v. For simplicity, we use d_u for $d_G(u)$ if there is no ambiguity. For $S \subseteq V(G)$, let $N_S(v) = N_G(v) \cap S$ and $d_S(v) = |N_S(v)|$ whether $v \in S$ or not. For $S, T \subset V(G)$, let e(S, T) be the number of edges between S and T. Particularly, if S = T, then e(S) denotes the number of edges in G[S] (the subgraph of G induced by S). The set S is independent if G[S] has no edges.

For a nontrivial graph G with $v \in V(G)$, G - v denotes the graph obtained from G by removing v and its incident edges.

A vertex v in a connected graph G is a cut vertex if G - v is disconnected, equivalently, there exist vertices u and w distinct from v such that there is at least one path from u to w in G and v lies on every such path in G.

For a graph G with two nonadjacent vertices u and v, denote by G + uv the graph obtained from G by adding the edge uv. If wz is an edge of G, then G - wz is the graph obtained from G be removing the edge wz.

The following lemma is an immediate consequence of the Perron-Frobenius theorem.

Lemma 2.1. Let G be a graph and u and v two nonadjacent vertices of G. If G + uv is connected, then $\rho(G + uv) > \rho(G)$.

If G is a connected graph, then A(G) is irreducible, so by Perron-Frobenius theorem, there exists a unique unit positive eigenvector of A(G) corresponding to $\rho(G)$, which we call the Perron vector of G.

Lemma 2.2. [4] Let G be a connected graph with $\{u, v, w\} \subseteq V(G)$ such that $uw \notin E(G)$ and $vw \in E(G)$. If **x** is the Perron vector of G with $x_u \ge x_v$, then $\rho(G - vw + uw) > \rho(G)$.

If \mathbf{x}' is the Perron vector of G - vw + uw in Lemma 2.2, then $x'_u > x'_v$ by Lemma 2.2. So Lemma 2.2 has the equivalent form: Let G be a connected graph with $\{u, v\} \subset V(G)$ and $N_G(v) \setminus N_G[u] \neq \emptyset$. If \mathbf{x} is the Perron vector of G with $x_u \ge x_v$, then for any nonempty $N \subseteq N_G(v) \setminus N_G[u]$, $\rho(G - \{vw : w \in N\} + \{uw : w \in N\}) > \rho(G)$.

For a graph G with $u, v \in V(G)$, the (u, v)-entry of $A(G)^2$ is equal to the number of walks of length two from u to v. If **x** is the Perron vector of G, then **x** is an eigenvector of $A(G)^2$ corresponding to the eigenvalue $\rho(G)^2$, so

$$\rho(G)^2 x_u = d_G(u) x_u + \sum_{v \in N_G(u)} d_{N_G(u)}(v) x_v + \sum_{v \in V(G) \setminus N_G[u]} d_{N_G(u)}(v) x_v.$$
(2.1)

Suppose that V(G) is partitioned as $V_1 \cup \cdots \cup V_m$. For $1 \leq i < j \leq m$, set A_{ij} to be the submatrix of A(G) with rows corresponding to vertices in V_i and columns corresponding to vertices in V_j . The quotient matrix of A(G) with respect to the partition $V_1 \cup \cdots \cup V_m$ is denoted by $B = (b_{ij})$, where $b_{ij} = \frac{1}{|V_i|} \sum_{u \in V_i} \sum_{v \in V_j} a_{uv}$. If A_{ij} has constant row sum, then we say B is an equitable quotient matrix. The following lemma is an immediate consequence of [3, Lemma 2.3.1].

Lemma 2.3. For a connected graph G, if B is an equitable quotient matrix of A(G), then the eigenvalues of B are also eigenvalues of A(G) and $\rho(G)$ is equal to the largest eigenvalue of B.

Let G_1 and G_2 be two graphs with $u \in V(G_1)$ and $v \in V(G_2)$. Denote by $G_1 uvG_2$ the graph obtained by identifying the vertices u and v, which we call this graph the coalescence of G_1 and G_2 at u and v. If u = v, then we denote by $G_1 uG_2$ for simplicity.

Lemma 2.4. [9] Let G_1 be a graph with $u \in V(G_1)$ and G_2 a connected graph with $v, w \in V(G_2)$. If $\rho(G_2 - v) < \rho(G_2 - w)$, then $\rho(G_1 uvG_2) > \rho(G_1 uwG_2)$.

Lemma 2.5. [9] Let G be a connected graph with $u \in V(G)$, H_1 and H_2 be two graphs of the same order with $v_i \in V(H_i)$ for i = 1, 2. If $\rho(H_1) > \rho(H_2)$ and $\rho(H_1 - v_1) \leq \rho(H_2 - v_2)$, then $\rho(Guv_1H_1) > \rho(Guv_2H_2)$.

Lemma 2.6. [13] We have

$$\rho(K_{1,1,n-2}) = \frac{1}{2} + \sqrt{2n - \frac{15}{4}} \text{ and } \rho(K_{3,n-3}) = \sqrt{3(n-3)}$$

Moreover, $\rho(K_{3,n-3}) > \rho(K_{1,1,n-2})$ *if and only if* $n \ge 10$.

Denote by $d_G(u, v)$ the distance between vertices u and v in a graph G. Let u and v be two vertices of G and P be a path from u to v in G. Moreover, if w and z are two vertices on P with $d_P(u, w) < d_P(u, z)$, then P[w, z] denotes the segment of P from w to z.

If a component of a graph is K_1 , K_2 , or $K_{1,s}$ with $s \ge 2$, then we call it an isolated vertex, an isolated edge, or a star (in which the vertex of degree s is its center). As usual, a copy of K_3 is called a triangle. A vertex of degree one is also called a pendant vertex.

We use I_n to denote the identity matrix of order n.

3 Proof of Theorem 1.1

Lemma 3.1. [6] If G is an n-vertex graph containing no DCCs, then $e(G) \leq 2n - 3$.

Proof of Theorem 1.1. The case for n = 3, 4 can be checked easily. Suppose that $n \ge 5$. Suppose that G is a graph that maximizes the spectral radius among all graphs on n vertices containing no DCCs. It suffices to show that $G \cong K_{1,1,n}$.

If G is not connected with components G_1, \ldots, G_r , then we obtain a graph G' by choosing $v_i \in V(G_i)$ for $i = 1, \ldots, r$ and adding edges v_1v_j for all $j = 2, \ldots, j$, which obviously does not contain a DCC, and we have by Lemma 2.1 that $\rho(G') > \rho(G)$, which is a contradiction. Thus, G is connected.

Let $\rho = \rho(G)$. Denote by **x** the Perron vector of G. Let u be a vertex with maximum entry in **x**. Let $X = N_G(u)$ and $Y = V(G) \setminus N_G[u]$. As G does not contain a DCC, a component of G[X] is an isolated vertex K_1 , an isolated edge K_2 , or a star $K_{1,r}$ for some $r \ge 2$. Let X_0 be the set of isolated vertices in G[X] and X_1 be the set of vertices of isolated edges in G[X]. Let $X_2 = X \setminus (X_0 \cup X_1)$. Let X'_2 be the set of vertices of degree at least two in G[X] (i.e., the centers of the stars in G[X]). Note that $\rho x_u = \sum_{v \in X} x_v$. From (2.1), we have

$$(\rho^{2} - \rho)x_{u} = d_{u}x_{u} + \sum_{v \in X} (d_{X}(v) - 1)x_{v} + \sum_{v \in Y} d_{X}(v)x_{v}$$

$$\leqslant \left(d_{u} + \sum_{v \in X'_{2}} (d_{X}(v) - 1) + \sum_{v \in Y} d_{X}(v) - \sum_{v \in X_{0}} \frac{x_{v}}{x_{u}} \right) x_{u}$$

$$= \left(|X| + |X_{2}| - 2|X'_{2}| + e(X,Y) - \sum_{v \in X_{0}} \frac{x_{v}}{x_{u}} \right) x_{u},$$

 \mathbf{SO}

$$\rho^2 - \rho \leqslant |X| + |X_2| - 2|X_2'| + e(X, Y) - \sum_{v \in X_0} \frac{x_v}{x_u}$$

As $K_{1,1,n-2}$ does not contain a DCC, we have by Lemma 2.6 that $\rho \ge \rho(K_{1,1,n-2}) = \frac{1}{2} + \sqrt{2n - \frac{15}{4}}$, so $\rho^2 - \rho \ge 2n - 4$. Therefore, we have

$$2n - 4 \leq |X| + |X_2| - 2|X_2'| + e(X, Y) - \sum_{v \in X_0} \frac{x_v}{x_u}.$$
(3.1)

Claim 3.1. There is no cut vertex in $V(G) \setminus \{u\}$.

Proof. Suppose to the contrary that there is a cut vertex $v \in V(G) \setminus \{u\}$. Then G - v has at least two components, one of which does not contain u, which we denote by H. Let

$$G' = G - \{vz : z \in N_H(v)\} + \{uz : z \in N_H(v)\}.$$

Then u is a cut vertex of G' and $G'[\{u\} \cup V(H)] \cong G[\{v\} \cup V(H)]$, so G' does not contain a DCC. However, we have by Lemma 2.2 that $\rho(G') > \rho$, a contradiction.

Claim 3.2. For each $vv' \in E(G[X_1])$, if $vw \in Y$ for some $w \in Y$, then either $N_X(w) \setminus \{v\} \subseteq X_0$ or $N_X(w) = \{v, v'\}$.

Proof. We first claim that $N_X(w) \setminus \{v, v'\} \subseteq X_0$. Otherwise, there is a vertex $v_1 \in X \setminus (X_0 \cup \{v, v'\})$ with $v_1 w \in E(G)$. Then $v_1 \in (X_1 \cup X_2) \setminus \{v, v'\}$. Denote by v'_1 one neighbor of v_1 in X. Then $uv'vwv_1v'_1u$ is a cycle with chords uv and uv_1 , a contradiction.

If $v_0w \in E(G)$ for some $v_0 \in X_0$ and $v'w \in E(G)$, then $uvv'wv_0u$ is a cycle with chords uv' and vw, a contradiction. So $v'w \notin E(G)$ or $v_0w \notin E(G)$.

Let Y_1 be the set of those vertices in Y with at least one neighbor in X_1 . By Claim 3.2, $d_{X_1}(w) = 1, 2$ for any $w \in Y_1$. Let $Y_1^i = \{w \in Y_1 : d_{X_1}(w) = i\}$ for i = 1, 2. Then $Y_1 = Y_1^1 \cup Y_1^2$ and

$$e(X_1, Y) = |Y_1^1| + 2|Y_1^2|$$
 and $e(X_0, Y_1^2) = 0.$ (3.2)

Claim 3.3. For each $v \in X_2 \setminus X'_2$, $N_Y(v) = \emptyset$.

Proof. Suppose that $N_Y(v) = \emptyset$, say $w \in N_Y(v)$ for some $v \in X_2 \setminus X'_2$. By Claim 3.1, there is a path P between w and u which does not contain v. Let v_0 be the unique neighbor of v in X and v' be a neighbor of v_0 different from v' in X.

If P does not contain v_0 or v', then $uv'v_0vwPu$ is a cycle with chords uv_0 and uv, a contradiction. If P contains v_0 but not v' or P contains both v_0 and v' with $d_P(w, v_0) < d_P(w, v')$, then $uvwP[w, v_0]v_0v'u$ is a cycle with chords uv_0 and vv_0 , a contradiction. Similarly, if P contains v' but not v_0 or P contains both v' and v_0 with $d_P(w, v') < d_P(w, v_0)$, then $uvwP[w, v']v'v_0u$ is a cycle with chords uv' and vv_0 , also a contradiction.

Claim 3.4. For each $v \in X'_2$, if $vw \in E(G)$ for some $w \in Y$, then $N_X(w) \setminus \{v\} \subseteq X_0$.

Proof. Suppose that there exists a vertex $v_0 \in X_1 \cup X_2 \setminus \{v\}$ such that $v_0 w \in E(G)$. Let $v' \in N_X(v) \setminus \{v_0\}$. By Claim 3.3, $vv_0 \notin E(G)$. Let v'_0 be a neighbor of v_0 in X. Then $uv'vwv_0v'_0u$ is a cycle with chords uv and uv_0 , a contradiction.

Let Y_2 be the set of those vertices in Y with at least one neighbor in X_2 . Then by Claims 3.3 and 3.4,

$$e(X_2, Y) = e(X'_2, Y_2) = |Y_2|.$$
(3.3)

By Claims 3.2 and 3.4, we have $Y_1 \cap Y_2 = \emptyset$. For $v \in X_1 \cup X'_2$, let

$$X_v = \{ z \in X_0 : N_Y(z) \cap N_Y(v) \neq \emptyset \}.$$

Claim 3.5. For each $v_0 \in X_v$ with $v \in X_1 \cup X'_2$ and $w \in N_Y(v) \cap N_Y(v_0)$, if $|N_Y(v_0)| \ge 2$, then $N_X(z) \subseteq \{v, v_0\}$ for each $z \in N_Y(v_0) \setminus \{w\}$. Moreover, if $|N_Y(v)| \ge 2$, then $|N_Y(v) \cap N_Y(v_0)| = 1$ or $N_X(z) \subseteq \{v, v_0\}$ for each $z \in N_Y(v_0)$.

Proof. Suppose that $v_1 \in N_X(z) \setminus \{v, v_0\}$. Denote by v' one neighbor of v in X. Then $uv'vwv_0zv_1u$ is a cycle with chords uv and uv_0 , a contradiction. So $N_X(z) \setminus \{v, v_0\} = \emptyset$. The second result follows immediately.

By Claim 3.5, we have $X_v \cap X_{v'} = \emptyset$ for any pair $\{v, v'\} \subseteq X_1 \cup X'_2$. By Claims 3.2 and 3.4 again, $e(X_v, Y_1 \cup Y_2) = e(X_v, N_Y(v))$.

Let $v \in X_1 \cup X'_2$. Note that

$$e(X_v, Y_1 \cup Y_2) \leqslant |X_v| + |N_Y(v)|.$$

If $|N_Y(v)| = 1$, this is obvious as $e(X_v, Y_1 \cup Y_2) = |X_v|$. Suppose that $|N_Y(v)| \ge 2$. Let $X_v^1 = \{v_0 \in X_v : |N_Y(v) \cap N_Y(v_0)| = 1\}$ and $X_v^2 = X_v \setminus X_v^1$. Then by Claim 3.5, $e(X_v^1, Y_1 \cup Y_2) = |X_v^1|$ and $N_Y(v_0) \cap N_Y(v'_0) = \emptyset$ for any pair $\{v_0, v'_0\} \subseteq X_v^2$. So $e(X_v^2, Y_1 \cup Y_2) = |\{w \in N_Y(v) : v_0w \in E(G), v_0 \in X_v^2\}| \le |N_Y(v)|$. Thus

$$e(X_v, Y_1 \cup Y_2) = e(X_v^1, Y_1 \cup Y_2) + e(X_v^2, Y_1 \cup Y_2) \leqslant |X_v^1| + |N_Y(v)| \leqslant |X_v| + |N_Y(v)|,$$

as desired.

Let $X'_0 = \bigcup_{v \in X_1 \cup X'_2} X_v$. Then

$$e(X'_0, Y_1 \cup Y_2) = \sum_{v \in X_1 \cup X_2} e(X_v, Y_1 \cup Y_2) \leq \sum_{v \in X_1 \cup X_2} (|X_v| + |N_Y(v)|) = |X'_0| + |Y'_1| + |Y_2|.$$

Let

$$Y_v = \{ z \in Y \setminus (Y_1 \cup Y_2) : N_{X_v}(z) \neq \emptyset \}.$$

By Claim 3.5, we have $e(X, Y_v) = e(X_v, Y_v) = |Y_v|$ and $Y_v \cap Y_{v'} = \emptyset$ for any $v' \in X_1 \cup X'_2 \setminus \{v\}$. Let $Y'_0 = \bigcup_{v \in X_1 \cup X'_2} Y_v$. Then $e(X_v, Y'_0) = e(X_v, Y_v) = |Y_v|$ and

$$e(X'_0, Y'_0) = \sum_{v \in X_1 \cup X'_2} e(X_v, Y'_0) = \sum_{v \in X_1 \cup X'_2} |Y_v| = |Y'_0|.$$

Note that $e(X'_0, Y) = e(X'_0, Y'_0) + e(X'_0, Y_1 \cup Y_2)$. So

$$e(X'_0, Y) \leqslant |X'_0| + |Y'_0| + |Y_1^1| + |Y_2|.$$
(3.4)

Let $X_0^* = X_0 \setminus X_0'$ and $Y_0 = Y \setminus (Y_1 \cup Y_2 \cup Y_0')$. Then $e(X_0^*, Y) = e(X_0^*, Y_0)$ and $e(X_0', Y_0) = 0$. It then follows from (3.2)–(3.4) that

$$e(X,Y) = e(X_0^*,Y) + e(X_0',Y) + e(X_1,Y) + e(X_2',Y)$$

$$\leq e(X_0^*,Y) + |X_0'| + |Y_0'| + |Y_1^1| + |Y_2| + |Y_1^1| + 2|Y_1^2| + |Y_2|$$

$$= e(X_0^*,Y_0) + |X_0'| + |Y_0'| + 2|Y_1| + 2|Y_2|.$$

Note that n = 1 + |X| + |Y|, $X = |X_0^*| + |X_0'| + |X_1| + |X_2|$ and $|Y| = |Y_0| + |Y_0'| + |Y_1| + |Y_2|$. So, from (3.1), we have

$$|X_0^*| + |X_1| + 2|X_2'| + 2|Y_0| + |Y_0'| - 2 \le e(X_0^*, Y_0) - \sum_{v \in X_0} \frac{x_v}{x_u}.$$
(3.5)

Suppose that $X_0^* \neq \emptyset$. Then $\sum_{v \in X_0} \frac{x_v}{x_u} > 0$. From (3.5), we have

$$e(X_0^*, Y_0) \ge |X_0^*| + 2|Y_0| - 1.$$
(3.6)

Let $G_1 = G[\{u\} \cup X_0^* \cup Y_0]$. As G_1 does not contain a DCC, we have $e(G_1) \leq 2|X_0^*| + 2|Y_0| - 1$ by Lemma 3.1. So, from (3.6), we have

$$2|X_0^*| + 2|Y_0| - 1 \le e(G_1) = |X_0^*| + e(X_0^*, Y_0) + e(Y_0) \ge 2|X_0^*| + 2|Y_0| - 1 + e(Y_0),$$

implying that $e(Y_0) = 0$. Thus (3.6) is an equality, so from (3.5) again, we have $X_1 = X'_2 = \emptyset$, so $G = G_1$. As G is bipartite, we have $\rho(G) \leq \sqrt{e(G)} = \sqrt{2n-3}$, so $\frac{1}{2} + \sqrt{2n-\frac{15}{4}} \leq \sqrt{2n-3}$, a contradiction. This shows that $X_0^* = \emptyset$. So (3.5) becomes

$$|X_1| + 2|X_2'| + 2|Y_0| + |Y_0'| \le 2 - \sum_{v \in X_0} \frac{x_v}{x_u}.$$
(3.7)

If $X_1 \cup X'_2 = \emptyset$, then $X'_0 = \emptyset$, so $X_0 = \emptyset$, which shows that G is trivial or disconnected, a contradiction. So $X_1 \cup X'_2 \neq \emptyset$. From (3.7) we have $(|X_1|, |X'_2|, |Y_0|, |Y'_0|) = (2, 0, 0, 0)$, (0, 1, 0, 0) and $X_0 = \emptyset$.

If $(|X_1|, |X_2'|, |Y_0|, |Y_0'|) = (0, 1, 0, 0)$, then by Claims 3.1 and 3.3, $Y_2 = \emptyset$. So $G \cong K_{1,1,n-2}$, as desired.

If $(|X_1|, |X'_2|, |Y_0|, |Y'_0|) = (2, 0, 0, 0)$, then we have from (3.1) and (3.2) that $Y_1^1 = \emptyset$ and $V(G) = \{u\} \cup X_1 \cup Y_1^2$. We claim that Y_1^2 is independent, as otherwise, there is an edge w_1w_2 with $w_1, w_2 \in Y$, which leads to a cycle $uv_1w_1w_2v_2u$ with chords v_1v_2 and v_1w_2 , where $v_1, v_2 \in X_1$, a contradiction. Therefore, $G \cong K_{1,1,n-2}$.

Combining the above cases, we conclude that $G \cong K_{1,1,n-2}$. This completes the proof. \Box

4 Proof of Theorem 1.2

Let G be an n-vertex graph that does not contain a DCC₁. From a result in [17], we have $e(G) \leq 3n - 9$ for $n \geq 9$. Note that F_1 is a graph on five vertices with no DCC₁s, and $e(F_1) = 7 > 6 = 3n - 9$. In the following lemma, we show that $e(G) \leq 3n - 9$ for $n \geq 6$. As in [17], we need a result due to Bondy [2] (see also [1, Theorem 4.11]): Let C be a longest cycle (with length c) in an n-vertex graph G, then there are at most $\lfloor \frac{1}{2}c(n-c) \rfloor$ edges with at most one end vertex on C. We also need a result of Czipszer [11] (see also [1, p. 386]) stating that if $\delta(G) \geq 4$ then G contains a DCC₁.

Lemma 4.1. Let $n \ge 6$. If G is an n-vertex graph that does not contain a DCC_1 , then $e(G) \le 3n - 9$ with equality when G is bipartite if and only if $G \cong K_{3,n-3}$.

Proof. We prove the result by induction on n.

Suppose that n = 6. Let C be a longest cycle of G and c the length of C. By Bondy's result, $e(G - E(G[V(C)])) \leq \lfloor \frac{1}{2}c(6-c) \rfloor$. As G does not contain a DCC₁, each vertex in G[V(C)] has degree at most three. Then $e(G[V(C)]) \leq \lfloor \frac{3}{2}c \rfloor$. Then $e(G) = e(G - E(G[V(C)])) + e(G[V(C)]) \leq \lfloor \frac{1}{2}c(6-c) \rfloor + \lfloor \frac{3}{2}c \rfloor$. So, if c = 3, 5, 6, then $e(G) \leq 9$. Suppose that c = 4. If $G[V(C)] \cong K_4$, then each vertex outside C is adjacent to at most one vertex on C, and it is possible that the two vertices outside C are adjacent to a vertex on C, so they may be adjacent, implying that $e(G) \leq 6+3=9$. If $G[V(C)] \ncong K_4$, then $e(G[V(C)]) \leq 5$, so $e(G) \leq \lfloor \frac{1}{2}c(6-c) \rfloor + 5=9$. This shows that $e(G) \leq 3n-9$ for n = 6.

Moreover, suppose that G is a bipartite graph with e(G) = 3n - 9 = 9. Then c = 4, 6. If c = 4, then $e(G) \leq \lfloor \frac{1}{2}c(6-c) \rfloor + 4 = 8 < 9$. If c = 6, then $e(G) = \lfloor \frac{3}{2}c \rfloor = 9$ if and only if G is regular of degree three, equivalently, $G \cong K_{3,3}$.

Suppose that $n \ge 7$. Let v be a vertex of minimum degree in G. By induction assumption, $e(G - v) \le 3(n - 1) - 9$. By Czipser's result, $\delta(G) \le 3$. So $e(G) = e(G - v) + \delta(G) \le 3(n - 1) - 9 + 3 = 3n - 9$.

Next, suppose that G is a bipartite graph with e(G) = 3n - 9. By the above argument, $d_G(v) = 3$ and e(G - v) = 3(n - 1) - 9. By induction assumption, $G - v \cong K_{3,n-4}$. If n = 7, then as G is bipartite, we have $G \cong K_{3,4}$. Suppose that $n \ge 8$. Let (V_1, V_2) be the bipartition of G - v with $|V_1| = 3$ and $|V_2| = n - 4$. Suppose that $G \ncong K_{3,n}$. As

G is bipartite, G does not contain a triangle, so $N_G(v) \subset V_2$. Let $V_1 = \{u_1, u_2, u_3\}$ and $N_G(v) = \{v_1, v_2, v_3\} \subset \{v_1, \dots, v_4\} \subseteq V_2$. Then $vv_1u_1v_4u_3v_3u_2v_1v$ is a cycle with chords v_1u_2 and v_1u_3 , a contradiction.

For positive integers n and r with $n \ge 3r + 4$ and $r \ge 1$, let u be the vertex of degree 3r in $K_1 \vee rK_3$ (any vertex if r = 1), and vw an edge of $K_{3,n-3r-3}$, where the degree of v is n - 3r - 3 and the degree of w is three, and let $H_{n,r} = (K_1 \vee rK_3)uvK_{3,n-3r-3}$ and $H'_{n,r} = (K_1 \lor rK_3) uw K_{3,n-3r-3}.$

Lemma 4.2. For $n \ge 3r + 6$ and $r \ge 1$, $\rho(H_{n,r}) \ge \rho(H'_{n,r})$ with equality if and only if n = 3r + 6.

Proof. If n = 3r + 6, then $H_{n,r} \cong H'_{n,r}$. Suppose that $n \ge 3r + 7$. Then n - 3r - 3 > 3, so $\rho(K_{3,n-3r-3} - v) = \rho(K_{2,n-3r-3}) = \sqrt{2(n-3r-3)} < \sqrt{3(n-3r-4)} = \rho(K_{3,n-3r-4}) = \rho(K_{3,n \rho(K_{3,n-3r-3}-w)$. Thus the result follows by Lemma 2.4.

Lemma 4.3. For positive integers $n \ge 3r + 4$, $\rho(K_{3,n-3}) > \rho(H_{n,r})$.

Proof. Let $U_1 = V(rK_3)$, V_1 be the set of vertices of degree three in $K_{3,n-3r-3}$ and $V_2 =$ $V(G) \setminus (U_1 \cup V_1 \cup \{u\})$. With respect to the vertex partition $V(H_{n,r}) = \{u\} \cup U_1 \cup V_1 \cup V_2$, $A(H_{n,r})$ has an equitable quotient matrix

$$B = \begin{pmatrix} 0 & 3r & n - 3r - 3 & 0 \\ 1 & 2 & 0 & 0 \\ 1 & 0 & 0 & 2 \\ 0 & 0 & n - 3r - 3 & 0 \end{pmatrix}.$$

By Lemma 2.3, $\rho(H_{n,r})$ is the largest root of f(x) = 0, where

 $f(x) = \det(xI_4 - B) = x^4 - 2x^3 + (6r - 3n + 9)x^2 + (6n - 18r - 18)x + 6nr - 18r - 18r^2.$ It is evident that $f''(x) \ge f''(\sqrt{3n-9}) > 0$ if $x \ge \sqrt{3n-9}$, so it may be checked that $f'(x) \ge f'(\sqrt{3n-9}) > 0$ if $x \ge \sqrt{3n-9}$. As $n \ge 3r+4$, we have $4n - 12 - 3r > 3\sqrt{3n-9}$, \mathbf{SO}

$$f(\sqrt{3(n-3)}) = 24rn - 72r - 18r^2 - 18r\sqrt{3n-9} = 6r(4n-12-3r-3\sqrt{3n-9}) > 0,$$

thich shows that $\rho(H_{n,r}) < \rho(K_{3,n-3}).$

which shows that $\rho(H_{n,r}) < \rho(K_{3,n-3})$.

The following result follows from Lemmas 4.2 and 4.3.

Corollary 4.1. For $n \ge 3r + 6$ and $r \ge 1$, $\rho(K_{3,n-3}) > \rho(H'_{n,r})$.

Lemma 4.4. The following results hold.

(i) For integer $n \ge 7$ with $n \equiv 1 \pmod{3}$, $\rho(K_1 \lor \frac{n-1}{3}K_3) < \sqrt{3(n-3)}$; (ii) For integer $n \ge 6$ with $n \equiv 0 \pmod{3}$,

$$\rho(K_1 \lor (K_2 \cup \frac{n-3}{3}K_3)) < \begin{cases} \sqrt{3(n-3)} & \text{if } n \ge 9, \\ \frac{1}{2} + \sqrt{2n - \frac{15}{4}} & \text{if } n = 6; \end{cases}$$

(iii) For integers $n \ge 8$ with $n \equiv 2 \pmod{3}$, $\rho(K_1 \lor (K_1 \cup \frac{n-2}{3}K_3)) < \sqrt{3(n-3)}$.

Proof. From [3, p. 19, Ex. 1.12], we know that $\rho(K_1 \vee \frac{n-1}{3}K_3) = 1 + \sqrt{n}$, so (i) follows.

Let $H = K_1 \vee (K_2 \cup \frac{n-3}{3}K_3)$. Let u be the vertex of degree n-1 in H, V_1 be the set of vertices of degree two and $V_2 = V(H) \setminus (V_1 \cup \{u\})$. With respect to the vertex partition $V(H) = \{u\} \cup V_1 \cup V_2, A(H)$ has an equitable quotient matrix

$$B' = \begin{pmatrix} 0 & 2 & n-3 \\ 1 & 1 & 0 \\ 1 & 0 & 2 \end{pmatrix}.$$

By Lemma 2.3, $\rho(H)$ is equal to the largest root of g(x) = 0, where

$$g(x) = \det(xI_3 - B') = x^3 - 3x^2 - (n-3)x + n + 1.$$

As g(x) is increasing in $[\sqrt{3n-9}, \infty)$ and $g(\sqrt{3(n-3)}) = 2(n-3)\sqrt{3n-9} - 2(4n-13) > 0$ if $n \ge 9$, we have $\rho(H) < \sqrt{3(n-3)}$ if $n \ge 9$. The result for n = 6 follows by an easy calculation. This proves (ii).

Let $G = K_1 \vee (K_1 \cup \frac{n-2}{3}K_3)$. Let u be the vertex of degree n-1 and v be the pendant vertex in G. Let $U_1 = V(G) \setminus \{u, v\}$. With vertex partition $V(G) = \{u\} \cup \{v\} \cup V_1$, A(G) has an equitable quotient matrix

$$B = \begin{pmatrix} 0 & 1 & n-2 \\ 1 & 0 & 0 \\ 1 & 0 & 2 \end{pmatrix}.$$

By Lemma 2.3, $\rho(G)$ is equal to the largest root of f(x) = 0, where

$$f(x) = \det(xI_3 - B) = x^3 - 2x^2 - (n-1)x + 2.$$

As f(x) is increasing in $[\sqrt{3n-9}, \infty]$ and $f(\sqrt{3n-9}) = (2n-8)\sqrt{3n-9} - 2(3n-10) > 0$. This proves (iii).

Lemma 4.5. For $n \ge 5 + t$ and $t \ge 2$ with $n - t \equiv 2 \pmod{3}$, $\rho(K_1 \lor (K_{1,t} \cup \frac{n-t-2}{3}K_3)) < \rho(K_{1,1,n-2})$.

Proof. Let $G = K_1 \vee (K_{1,t} \cup \frac{n-t-2}{3}K_3)$, $\rho = \rho(G)$ and **x** be the Perron vector of G. Let u and v be the vertices of degree n-1 and t+1 in G, respectively. Let V_1 be the set of vertices of degree four in G and V_2 be the set of remaining vertices. By $A(G)\mathbf{x} = \rho\mathbf{x}$, for i = 1, 2, each vertex in V_i has the same entry, denoted by x_i . By $A(G)\mathbf{x} = \rho\mathbf{x}$ at $v, w \in V_1$ and $z \in V_2$, we have

$$x_1 = \frac{1}{\rho - 2} x_u$$
 and $x_v = \frac{\rho + t}{\rho^2 - t} x_u$.

As $\frac{1}{\rho-2} < \frac{\rho+t}{\rho^2-t}$, $x_1 < x_v$. Let

$$G' = G - \{wz \in E(G) : w, z \in V_1\} + \{vw : w \in V_1\}$$

Then $G' \cong K_{1,1,n-2}$. By Rayleigh's principle,

$$\rho(G') \geqslant \mathbf{x}^{\top} A(G') \mathbf{x} = \mathbf{x}^{\top} A(G) \mathbf{x} - 2 \sum_{wz \in E(G[V_1])} x_w x_z + 2 \sum_{w \in V_1} x_v x_w > \rho,$$

as desired.

Proof of Theorem 1.2. It is trivial for n = 3, 4. Suppose in the following that $n \ge 5$. Suppose that G is a graph that maximizes the spectral radius among all graphs of order n containing no DCC₁s. Similarly as in the proof of Theorem 1.1, G is connected. Let $\rho = \rho(G)$ and let **x** be the Perron vector of G and u a vertex with maximum entry in **x**. Let $X = N_G(u)$ and $Y = V(G) \setminus N_G[u]$. From (2.1), we have

$$\rho^2 x_u = d_u x_u + \sum_{v \in X} d_X(v) x_v + \sum_{v \in Y} d_X(v) x_v \leqslant (|X| + 2e(X) + e(X, Y)) x_u,$$

 \mathbf{SO}

 $\rho^2 \leqslant |X| + 2e(X) + e(X, Y).$

As $K_{3,n-3}$ has no DCC₁s, we have $\rho \ge \rho(K_{3,n-3}) = \sqrt{3(n-3)}$, so we have

$$3(n-3) \leqslant |X| + 2e(X) + e(X,Y).$$
(4.1)

By the same argument as in the proof of Claim 3.1, we have

Claim 4.1. There is no cut vertex in $V(G) \setminus \{u\}$.

As G does not contain a DCC₁, G[X] is P_4 -free, any component G[X] is an isolated vertex K_1 , an isolated edge K_2 , a star $K_{1,r}$ for some $r \ge 2$, or a triangle K_3 . Let X_0 be the set of isolated vertices in G[X], X_1 the set of vertices of isolated edges in G[X], X_2 the set of vertices of stars in G[X], and X_3 the set of vertices of triangles in G[X]. Then $X = X_0 \cup X_1 \cup X_2 \cup X_3$.

Claim 4.2. For each $v \in X_3$, $N_Y(v) = \emptyset$.

Proof. Suppose that $N_Y(v) \neq \emptyset$ for some $v \in X_3$. Assume that $w \in N_Y(v)$. By Claim 4.1, v is not a cut vertex of G, so there is a path P connecting w and u which does not contain v. Denote by v' the vertex on P adjacent to u. Let v_1 and v_2 be two other vertices of the triangle of G[X] containing v. If both v_1 and v_2 lie outside P, then uv_1v_2vwPu is a cycle with chords uv and uv_2 , a contradiction. If one of v_1 or v_2 , say v_1 , lies on P, where we assume that $d_P(v_1, w) < d_P(v_2, w)$ if both v_1 and v_2 lie one P, then $uv_2vwP[w, v_1]v_1u$ is a cycle with chords vu and vv_1 , also a contradiction.

Let X'_2 be the set of the centers of the stars in $G[X_2]$. Let X^*_2 be the set of pendant vertices of the components $K_{1,2}$ of $G[X_2]$

Claim 4.3. For each $v \in X_2^*$, if $vw \in Y$ for some $w \in Y$, then $N_X(w) \subseteq \{v, v'\}$, where v' is the other vertex in the component $K_{1,2}$ of $G[X_2]$ containing v.

Proof. Suppose that $wv_1 \in E(G)$ for some $v_1 \in X \setminus \{v, v'\}$. If $vv_1 \in E(G)$, then $v'v_1 \in E(G)$, so $uv'v_1wvu$ is a cycle with chords uv_1 and vv_1 , a contradiction. So $vv_1 \notin E(G)$. Let v_0 be the unique neighbor of v in G[X]. Then $uv'v_0vwv_1u$ is a cycle with chords uv_0 and uv, also a contradiction.

Claim 4.4. For each $v \in X_2 \setminus (X'_2 \cup X^*_2), N_Y(v) = \emptyset$.

Proof. Suppose that $N_Y(v) \neq \emptyset$ for some $v \in X_2 \setminus (X'_2 \cup X^*_2)$. Assume that $w \in N_Y(v)$. By Claim 4.1, v is not a cut vertex of G, so there is a path P from w to u which does not contain v. Let v_0 be the neighbor of v in X and v_1, \ldots, v_t all the neighbors of v_0 in X with $v = v_t$. If all $v_0, v_1, \ldots, v_{t-1}$ lie outside P, then uv_1v_0vwPu is a cycle with chords uv_0 and uv_2 , a contradiction. In the following, if one of v_1, \ldots, v_{t-1} lies one P, then we assume that v_1 lies on P. If v_0 lies on P and v_1 lies outside P, or both v_0 and v_1 lie on P with $d_P(w, v_0) < d_P(w, v_1)$, then $uvwP[w, v_0]v_0v_1u$ is a cycle with chords uv_0 and vv_0 , a contradiction. If both v_0 and one of If v_1 lies on P and v_0 lies outside P, or both v_1 and v_0 lie on P with $d_P(w, v_1) < d_P(w, v_0)$, then $uv_2v_0vwP[w, v_1]v_1u$ is a cycle with chords is a cycle with chords v_0u and v_0v_1 , also a contradiction.

By Claim 4.3, for any $v \in X_2^*$ and any $w \in Y$, if $vw \in E(G)$, then $d_X(w) \leq 2$. Let Y_2 be the set of those vertices in Y with at least one neighbor in X_2^* . Then $e(X_2^*, Y_2) \leq 2|Y_2|$. By Claim 4.4, $e(X_2 \setminus (X_2' \cup X_2^*), Y) = 0$, so

$$e(X_2 \setminus X'_2, Y) = e(X_2^*, Y_2) \leqslant 2|Y_2|.$$
(4.2)

By a similar argument as in the proof of Claim 3.4, we have

Claim 4.5. For each $v \in X'_2$, if $vw \in Y$ for some $w \in Y$, then $N_X(w) \setminus \{v\} \subseteq X_0$.

Denote by $Y'_2 \subseteq Y$ the set of vertices with at least one neighbor in X'_2 . Then by Claim 4.5,

$$e(X'_2, Y) = e(X'_2, Y'_2) = |Y'_2|.$$
(4.3)

Claim 4.6. For each $vv' \in E(G[X_1])$, if $vw \in Y$ for some $w \in Y$, then $N_X(w) \setminus \{v, v'\} \subseteq X_0$.

Proof. Suppose that $wv_1 \in E(G)$ for some $v_1 \in X \setminus (X_0 \cup \{v, v'\})$. Denote by v'_1 a neighbor of v_1 in X. Then $uv'vwv_1v'_1u$ is a cycle with chords uv and uv_1 , a contradiction.

Let $Y_1 \subseteq Y$ be the set of those vertices with at least one neighbors in X_1 . By Claims 4.3, 4.5 and 4.6, $S \cap T = \emptyset$ for each pair of $\{S, T\} \subset \{Y_1, Y_2, Y_2'\}$.

For $v \in X_1 \cup X'_2$, let $X_v = \{z \in X_0 : N_Y(z) \cap N_Y(v) \neq \emptyset\}.$

By the same argument as in the proof of Claim 3.5, we have

Claim 4.7. For each $v_0 \in X_v$, if $|N_Y(v_0)| \ge 2$, then $N_X(z) \subseteq \{v, v_0\}$ for each $z \in N_Y(v_0) \setminus \{w\}$, where w is a common neighbor of v and v_0 . Moreover, if $|N_Y(v)| \ge 2$, then $|N_Y(v) \cap N_Y(v_0)| = 1$ or $N_X(z) \subseteq \{v, v_0\}$ for each $z \in N_Y(v_0)$.

By Claim 4.7, $X_v \cap X_{v'} = \emptyset$ for any pair $\{v, v'\} \subseteq X_1 \cup X'_2$ with $vv' \notin E(G)$ if $v, v' \in X_1$. Then for each $v \in X'_2$, by Claim 4.5, $e(X_v, Y_1 \cup Y'_2) = e(X_v, N_Y(v))$ and so by the same argument as in the previous section,

$$e(X_v, Y_1 \cup Y_2') \leqslant |X_v| + |N_Y(v)|.$$
(4.4)

Claim 4.8. If $vv' \in E(X_1)$, then $X_v \subseteq X_{v'}$ or $X_{v'} \subseteq X_v$.

Proof. Suppose to the contrary that $v_0 \in X_v \setminus X'_v$ and $v_1 \in X'_v \setminus X_v$. By the definition of X_v , there exist vertices w and w' such that $vw, v_0w, v'w', v_1w' \in E(G)$. Then $uv_0wvv'w'v_1u$ is a cycle with chords uv and uv', a contradiction. So $X_v \setminus X'_v = \emptyset$ or $X'_v \setminus X_v = \emptyset$.

For an edge $vv' \in G[X_1]$, assume that $X_{v'} \subseteq X_v$ by Claim 4.8. For $vv' \in E(X_1)$, let $Y_{vv'} = \{w \in Y_1 : vw, v'w \in E(G)\}.$

Claim 4.9. For $vv' \in G[X_1]$, $e(\{v, v'\} \cup X_v \cup X_{v'}, Y_1) \leq 2|N_Y(v) \cup N_Y(v')| + |X_v|$.

Proof. If $X_v = \emptyset$, then the result follows by Claim 4.6. Suppose next that $X_v \neq \emptyset$, say $v_0 \in X_v$. Let $w \in N_Y(v_0)$. Assume that $vw \in E(G)$.

We first claim that $Y_{vv'} \setminus \{w\} = \emptyset$. Otherwise, there is some vertex $z \in Y_{vv'}$ with $z \neq w$. Then $uv_0wvzv'u$ is a cycle with chords uv and vv', a contradiction. So either $Y'_{vv'} = \{w\}$ or $Y_{vv'} = \emptyset$.

If $Y_{vv'} = \{w\}$, then $e(\{v, v'\}, Y_1) = |N_Y(v) \cup N_Y(v')| + 1 \leq 2|N_Y(v) \cup N_Y(v')|$ and for each $z \in X_v$, z is adjacent to exactly one vertex w in Y, so $e(X_v, Y_1) = |X_v|$. Therefore, $e(\{v, v'\} \cup X_v \cup X_{v'}, Y_1) \leq 2|N_Y(v) \cup N_Y(v')| + |X_v \cup X'_v|$.

If $Y_{vv'} = \emptyset$, then $e(\{v, v'\}, Y_1) = |N_Y(v) \cup N_Y(v')|$. By Eq. (4.4), $e(X_v, Y_1) \leq |X_v| + |N_Y \cup N_Y(v')|$. Therefore, $e(\{v, v'\} \cup X_v \cup X_{v'}, Y_1) \leq 2|N_Y(v) \cup N_Y(v')| + |X_v \cup X_v'|$.

Let $X'_0 \subseteq X_0$ be the set of those vertices with at least one neighbor in $Y_1 \cup Y'_2$. Then $X'_0 = \bigcup_{v \in X_1 \cup X'_2} X_v$. For $v \in X_1 \cup X'_2$, let $Y_v \subseteq Y \setminus (Y_1 \cup Y_2 \cup Y'_2)$ be the set of those vertices with at least one neighbor in X_v . Let $Y'_0 = \bigcup_{v \in X_1 \cup X'_2} Y_v$. By Claim 4.7, $e(X_v, Y_v) = |Y_v|$ and $e(X_v, Y'_0) = e(X_v, Y_v) = |Y_v|$. By (4.3) and (4.4) and Claim 4.9,

$$e(X'_{0} \cup X_{1} \cup X'_{2}, Y)$$

$$= \sum_{vv' \in E(X_{1})} (e(\{v, v'\} \cup X_{v} \cup X_{v'}, Y_{1}) + e(X_{v} \cup X_{v'}, Y'_{0})) + \sum_{v \in X'_{2}} (e(v, Y) + e(X_{v}, Y))$$

$$\leqslant \sum_{vv' \in E(X_{1})} (2|N_{Y}(v) \cup N_{Y}(v')| + |X_{v}| + |Y_{v}|) + |Y'_{2}| + \sum_{v \in X'_{2}} (|X_{v}| + |N_{Y}(v)| + |Y_{v}|)$$

$$= 2|Y_{1}| + \sum_{vv' \in E(X_{1})} (|X_{v}| + |Y_{v}|) + 2|Y'_{2}| + \sum_{v \in X'_{2}} (|X_{v}| + |Y_{v}|),$$

i.e.,

$$e(X'_0 \cup X_1 \cup X'_2, Y) \leqslant |X'_0| + |Y'_0| + 2|Y_1| + 2|Y'_2|.$$
(4.5)

Let $X_0^* = X_0 \setminus X_0'$ and $Y_0 = Y \setminus (Y_1 \cup Y_2' \cup Y_2 \cup Y_0')$. Then $e(X_0^*, Y) = e(X_0^*, Y_0)$. By (4.2) and (4.5),

$$e(X,Y) = e(X_0^*,Y) + e(X_0' \cup X_1 \cup X_2',Y) + e(X_2 \setminus X_2',Y)$$

$$\leqslant e(X_0^*,Y_0) + |X_0'| + |Y_0'| + 2|Y_1| + 2|Y_2'| + 2|Y_2|.$$

Note that

$$e(X) = \frac{1}{2}|X_1| + |X_2| - |X_2'| + |X_3|$$

and

$$n = 1 + |X_0^*| + |X_0'| + |X_1| + |X_2| + |X_3| + |Y_0| + |Y_0'| + |Y_1| + |Y_2'| + |Y_2|.$$

From (4.1), we have

$$e(X_0^*, Y_0) \ge 2|X_0^*| + |X_0'| + |X_1| + 2|X_2'| + 3|Y_0| + 2|Y_0'| + |Y_1| + |Y_2'| + |Y_2| - 6,$$
(4.6)

so

$$e(X_0^*, Y_0) \ge 2|X_0^*| + 3|Y_0| - 6.$$
(4.7)

Case 1. $|X_0^*| + |Y_0| \ge 5$.

Let $G_1 = G[\{u\} \cup X_0^* \cup Y_0]$. From (4.7), we have

$$e(G_1) \ge |X_0^*| + e(X_0^*) + e(X_0^*, Y_0) + e(Y_0) \ge 3|X_0^*| + 3|Y_0| - 6 + e(Y_0).$$

As G_1 does not contain a DCC₁, we have $e(G_1) \leq 3|X_0^*| + 3|Y_0| - 6$, so

$$3|X_0^*| + 3|Y_0| - 6 + e(Y_0) \leqslant e(G_1) \leqslant 3|X_0^*| + 3|Y_0| - 6,$$

implying that $e(Y_0) = 0$ and $e(G_1) = 3|X_0^*| + 3|Y_0| - 6$.

As $e(Y_0) = 0$, G_1 is a bipartite graph with bipartition $(X_0^*, \{u\} \cup Y_0)$. As $e(G_1) = 3|X_0^*| + 3|Y_0| - 6$, we have by Lemma 4.1 that $G_1 \cong K_{3,|X_0^*| + |Y_0| - 2}$. So $|X_0| = 3$ or $|Y_0| = 2$.

As $e(G_1) = 3|X_0^*| + 3|Y_0| - 6$, we know from the above argument that (4.7) is an inequality. Then, from (4.6), we have $X'_0 = X_1 = X'_2 = Y'_0 = Y'_2 = Y_2 = \emptyset$, so $X = X_0^* \cup X_3$ and $Y = Y_0$.

Suppose $X_3 \neq \emptyset$. Let r be the number of triangles in G[X]. Then $|X_3| = 3r$. If $|Y_0| = 2$, then $G \cong H_{n,r}$, and if $|X_0| = 3$, then $G \cong H'_{n,r}$. By Lemma 4.3 and Corollary 4.1, we have $\rho < \rho(K_{3,n-3})$, a contradiction. It follows that $X_3 = \emptyset$, so $n \ge 6$ and $G \cong K_{3,n-3}$. Case 2. $|X_0^*| + |Y_0| \le 4$ and $X_0^* \neq \emptyset$.

As $e(X_0^*, Y_0) \leq |X_0^*||Y_0|$, we have from (4.7) that $(|X_0^*| - 3)(|Y_0| - 2) \geq 0$, so $|X_0^*| \leq 3$ and $|Y_0| \leq 2$. If $|X_0^*| = 1$ and $|Y_0| \geq 1$, then we have from (4.6) that $|X_0'| + |X_1| + 2|X_2| + 2|Y_0'| + |Y_1| + |Y_2| + |Y_2'| \leq 4 - 2|Y_0| \leq 2$, so $Y_0' = Y_1 = Y_2 = Y_2' = \emptyset$, implying u and any vertex in Y_0 belong to different components of $G - X_0^*$, that is, the vertex of X_0^* is a cut vertex, which contradicts Claim 4.1. So if $|X_0^*| = 1$, then $Y_0 = \emptyset$ and there are three possibilities as below:

- (a) $|X_0^*| = 2$ and $|Y_0| = 2$;
- (b) $|X_0^*| = 2, 3 \text{ and } |Y_0| = 1;$

(c) $|X_0^*| = 1, 2, 3 \text{ and } Y_0 = \emptyset.$

Suppose first that (a) holds. From (4.7), we have $e(X_0^*, Y_0) = 4$. Now, from (4.6), we have $X'_0 = X_1 = X'_2 = Y'_0 = Y_1 = Y_2 = \emptyset$, so $X = X_0^*$ and $Y = Y_0$. If $X_3 \neq \emptyset$, then $G \cong H_{n,\frac{|X_3|}{3}}$, so we have by Lemma 4.3 that $\rho < \rho(K_{3,n-3})$, a contradiction. Therefore $X_3 = \emptyset$, so n = 5 and $G \cong K_{3,n-3}$.

Suppose next that (b) holds. From (4.6), we have $|X'_0| + |X_1| + 2|X'_2| + |Y_1| + |Y'_2| + |Y_2| \leq 3 - |X^*_0| \leq 1$, so $X_1 = X'_2 = \emptyset$. It follows that $Y = Y_0$, say $Y_0 = \{w\}$, so G' := G + uw does not contain a DCC₁. By Lemma 2.1, $\rho(G') > \rho$, a contradiction.

Suppose finally that (c) holds. If $|X_0^*| \ge 2$, then as $Y_0 = \emptyset$, each vertex in X_0^* is a pendant vertex of G, so adding an edge between two vertices in X_0^* leads to a graph that does not contain a DCC₁, which, by Lemma 2.1, is a contradiction. So $|X_0^*| = 1$.

Claim 4.10. $X_1 \cup X_2 = \emptyset$.

Proof. Suppose to the contrary that $X_1 \cup X_2 \neq \emptyset$.

If $Y = \emptyset$, then adding an edge between the vertex of X_0^* and a vertex in $X_1 \cup X'_2$ produces a graph that does not contain a DCC₁, which, by Lemma 2.1, is a contradiction. So $Y \neq \emptyset$. Moreover, we have $Y'_2 = \emptyset$; otherwise we have by Claim 4.1 and (4.6) that $|X'_0| = |X'_2| = |Y'_2| = 1$, $X = X_0^* \cup X'_0 \cup X_2 \cup X_3$ and $Y = Y'_2$, so e(X, Y) = 2, which contradicts (4.1). This shows that $Y_1 \cup Y_2 \neq \emptyset$.

Suppose that $Y_1 \neq \emptyset$. If $X'_0 = \emptyset$, then the graph obtained from G by adding an edge between the vertex of X^*_0 and a vertex in Y_1 does not contain a DCC₁, which, by Lemma 2.1, is a contradiction. So $X'_0 \neq \emptyset$. From (4.6), $|Y_1| = 1$, so the graph obtained from G adding an edge between the vertex of X^*_0 and the vertex of Y_1 does not contain a DCC₁, which, by Lemma 2.1, is a contradiction.

Suppose that $Y_2 \neq \emptyset$. Then $X'_2 \neq \emptyset$. We have by Eq. (4.6) that $|X'_2| = 1$, $|Y_2| = 1, 2$ and $e(X_2, Y_2) = 2|Y_2|$. Let $F_0 = G[\{u\} \cup X_0^* \cup X_2 \cup Y_2]$. Then F_0 is obtained from $K_{2,2+|Y_2|}$ with bipartition $(X_2^*, \{u\} \cup X'_2 \cup Y_2)$ by adding an edge between u and the vertex of X'_2 and a pendant vertex to u. If $n = 5 + |Y_2|$, then $G = F_0$ and if $n > 5 + |Y_2|$, then G consists of F_0 and $G[V(G) \setminus V(F_0) \cup \{u\}]$ with a common vertex u. Let $F'_0 \cong K_{1,1,|Y_2|+3}$. By direct calculation, we have

$$\rho(F_0) = \begin{cases} 2.9439 & \text{if } |Y_2| = 1, \\ 3.2054 & \text{if } |Y_2| = 2, \end{cases} \text{ and } \rho(F_0 - u) = \begin{cases} 2 & \text{if } |Y_2| = 1, \\ 2.4495 & \text{if } |Y_2| = 2, \end{cases}$$

From Lemma 2.4,

$$\rho(F'_0) = \begin{cases} 3.3723 & \text{if } |Y_2| = 1, \\ 3.7016 & \text{if } |Y_2| = 2. \end{cases}$$

As $F'_0 - u \cong K_{1,|Y_2|+3}$,

$$\rho(F'_0 - u) = \begin{cases} 2 & \text{if } |Y_2| = 1, \\ \sqrt{5} & \text{if } |Y_2| = 2. \end{cases}$$

Thus,

$$\rho(F_0) < \rho(F'_0) \text{ and } \rho(F_0 - u) \ge \rho(F'_0 - u).$$

If $n = 5 + |Y_2|$, then $\rho(F_0) < \rho(F'_0)$, a contradiction, and if $n > 5 + |Y_2|$, then we have by Lemma 2.5 that $\rho(F'_0 uG[V(G) \setminus V(F_0) \cup \{u\}]) > \rho$, also a contradiction.

This shows that $X_1 \cup X_2 = \emptyset$, as desired.

By Claim 4.10, we have $X = X_0^* \cup X_3$ and $Y = \emptyset$, so $G \cong K_1 \vee (K_1 \cup \frac{n-2}{3}K_3)$, which, by Lemma 4.4, is a contradiction if $n \ge 8$. Thus n = 5 and $G \cong F_1$. **Case 3.** $X_0^* = \emptyset$.

Case 3.1. $X'_0 = \emptyset$.

In this case, $X_0 = \emptyset$. If $X_1 \cup X_2 = \emptyset$, then $G \cong K_1 \vee \frac{n-3}{3}K_3$, which, by Lemma 4.4, is a contradiction. So $X_1 \cup X_2 \neq \emptyset$.

Next, we show that $Y_1 \cup Y_2 \cup Y'_2 = \emptyset$. Suppose to the contrary that $Y_1 \cup Y_2 \cup Y'_2 \neq \emptyset$.

Claim 4.11. For each $w \in Y_1 \cup Y_2 \cup Y'_2$ and one neighbor v of w in $X_1 \cup X_2$, any path from w to u passes through some vertex of the component H of G[X] containing v.

Proof. Suppose that this is not true. let P be a path from w to u that does not pass through any vertex of H. Let v_1 be a neighbor of u on P. As $X_0 = \emptyset$, $N_X(v_1) \neq \emptyset$. Let $v'_1 \in N_X(v_1)$. Assume that $v'_1 \notin V(P)$. Then with $v' \in N_X(v)$, $uv'vwP[w, v_1]v_1v'_1u$ is a cycle with chords uv and uv_1 , a contradiction.

Claim 4.12. $Y'_2 = \emptyset$.

Proof. Otherwise, assume that $w \in Y'_2$. Let v be the neighbor of w in X'_2 . By Claims 4.1 and 4.11, there is a path from w to u containing some vertex $v^* \in N_X(v)$. Then, with $v' \in N_X(v) \setminus \{v^*\}, uv'vwP[w, v^*]v^*u$ is a cycle with chords uv and vv^* , a contradiction. \Box

By Claim 4.12, we assume that $w \in Y_1 \cup Y_2$. Let v_1 be one neighbor of w in $X_1 \cup X_2$ and H be the component of G[X] containing v_1 , which is a copy of K_2 or $K_{1,2}$. Denote by v_2 the pendant vertex in H different from v_1 .

Let $Y_H = N_Y(v_1) \cup N_Y(v_2)$.

Claim 4.13. For any $z \in Y_H$, $d_G(z) = 2$.

Proof. By Claims 4.5 and 4.6, z has one or two neighbors in H.

Suppose first that z has two neighbors in H, that is, the two neighbors are v_1 and v_2 . If $N_Y(z) \neq \emptyset$, say $z' \in N_Y(z)$, then by Claim 4.1, there is a path from z' to u which does not pass through z. By Claims 4.11 and 4.12, a shortest such path P passes through one of v_1 and v_2 , say v_1 . Then v_2 does not lie on P. So $uv_2zz'P[z', v_1]v_1u$ is a cycle with chords v_1v_2 and v_1z if $H \cong K_2$ and $uv_0v_2zz'P[z, v_1]v_1u$ is a cycle with chords v_0v_1 and v_1z if $H \cong K_{1,2}$ with center v_0 , a contradiction. It follows that $N_Y(z) = \emptyset$, so $d_G(z) = 2$.

Suppose next that z has exactly one neighbor, say v_1 , in H. Suppose that $d_Y(z) \ge 2$. Next, we show that $N_Y(z) \subseteq Y_H$. Suppose that this is not true. Then z has a neighbor in Y_0 , so $|Y_0| \ge 1$. Note that $X_0 = \emptyset$. By Claim 4.3, $e(X^*, Y_2) \le 2|Y_2|$. By Claim 4.6, $e(X_1, Y_1) \le 2|Y_1|$. Now (4.1) becomes

$$3(|X_1| + |X_2| + |Y_1| + |Y_2| + |Y_0|) - 6 \leq |X_1| + |X_2| + 2|X_2| - 2|X_2'| + 2|Y_1| + 2|Y_2|$$

i.e., $|X_1| + 2|X'_2| + 3|Y_0| + |Y_1| + |Y_2| \leq 6$, so $|Y_0| = 1$. If $X_1 \neq \emptyset$, then $z \in Y_H \subseteq Y_1$, so $Y_H = \{z\}$. If $X'_2 \neq \emptyset$, then $z \in Y_H \subset Y_2$, so $Y_H = \{z\}$. In either case, z is a cut vertex of G, a contradicting Claim 4.1. So $N_Y(z) \subseteq Y_H$. By Claims 4.1 and 4.11, there is a path P from z to u containing v_2 but not v_1 . Let $z' \in N_Y(z)$ outside P. Assume that $z'v_1 \in E(G)$. Then $uv_1z'zP[z,v_2]v_2u$ is a cycle with chords v_1v_2 and v_1z if $H \cong K_2$ and $uv_1z'zP[z,v_2]v_2v_0u$ is a cycle with chords v_0v_1 and v_1z if $H \cong K_{1,2}$ with center v_0 , a contradiction. So $d_Y(z) \leq 1$. By Claim 4.1, $d_Y(z) = 1$, so $d_G(z) = 2$.

By Claims 4.5, 4.6 and 4.13, $e(Y_H) \leq \frac{|Y_H|}{2}$, so $\sum_{z \in Y_H} x_u x_z > \sum_{zz' \in E(G[Y_H])} x_z x_{z'}$. Assume that $x_{v_1} \geq x_{v_2}$.

Suppose that $H \cong K_2$. Let

$$G' = G - \{v_2 z : z \in N_Y(v_2) \setminus N_Y(v_1)\} - \{z z' \in E(G) : z, z' \in Y_H\} + \{v_1 z : z \in N_Y(v_2) \setminus N_Y(v_1)\} + \{u z : z \in Y_H\}.$$

It is easy to see that G' does not contain a DCC₁. But we have by Rayleigh's principle that

$$\rho(G') \ge \rho + 2\sum_{z \in N_Y(v_2) \setminus N_Y(v_1)} (x_{v_1} - x_{v_2}) x_z + 2\left(\sum_{z \in Y_H} x_u x_z - \sum_{zz' \in E(G[Y_H])} x_z x_{z'}\right) > \rho,$$

a contradiction.

Suppose that $H \cong K_{1,2}$. Let v_0 be the center of H. Let

$$G' = \begin{cases} G - \{v_i z_i : z_i \in N_Y(v_i), i = 1, 2\} - \{z z' \in E(G) : z, z' \in Y_H\} \\ + \{u z, v_0 z : z \in Y_H\} \\ G - v_0 v_2 - \{v_2 z : z \in N_Y(v_2)\} - \{z z' \in E(G) : z, z' \in Y_H\} \\ + v_1 v_2 + \{u z : z \in Y_H\} + \{v_1 z : z \in N_Y(v_2) \setminus N_Y(v_1)\} \end{cases}$$
 otherwise.

Note that G' does not contain a DCC₁. Let $U_0 = N_Y(v_1) \cap N_Y(v_2)$, $U_1 = N_Y(v_1) \setminus U_0$ and $U_2 = N_Y(v_2) \setminus U_0$. If $x_{v_0} \ge x_{v_1}$, as one of U_0, U_1, U_2 is not empty, so

$$\rho(G') \ge \rho(G) + 2 \sum_{z \in U_0} (x_u + x_{v_0} - x_{v_1} - x_{v_2}) x_z$$

+ 2 $\sum_{z \in U_1} (x_{v_0} - x_{v_1}) x_z + 2 \sum_{z \in U_2} (x_{v_0} - x_{v_2}) x_z$
+ 2 $\left(\sum_{z \in U_1 \cup U_2} x_u x_z - \sum_{zz' \in E(G[U_1 \cup U_2])} x_z x_{z'} \right)$

 $> \rho$,

and otherwise, we have $x_{v_1} > x_{v_0}$, so

$$\rho(G') \ge \rho + 2(x_{v_1} - x_{v_0})x_{v_2} + 2\sum_{z \in U_2} (x_{v_1} - x_{v_2})x_z + 2\sum_{z \in U_0} (x_u - x_{v_2})x_z + 2\left(\sum_{z \in U_1 \cup U_2} x_u x_z - \sum_{zz' \in E(G[U_1 \cup U_2])} x_z x_{z'}\right) > \rho.$$

Thus we reach a contradiction in either case. This shows that $Y_1 \cup Y_2 \cup Y'_2 = \emptyset$. Thus, $Y = \emptyset$.

Suppose that there are two components in $G[X_1 \cup X_2]$, say H_1 and H_2 . Let v_i be the center of H_i for i = 1, 2 (if $H_i \cong K_2$, then v_i is arbitrary). Assume that $x_{v_1} \ge x_{v_2}$. Let

$$G' = G - \{v_2 z : z \in N_X(v_2)\} \cup \{v_1 z : z \in N_X(v_1)\}.$$

Obviously, G' does not contain a DCC₁. By Lemma 2.2, $\rho(G') > \rho$, a contradiction. So $G[X_1 \cup X_2]$ is connected, i.e., either $X_1 \neq \emptyset$ or $X_2 \neq \emptyset$.

If $X_1 \neq \emptyset$, then $|X_1| = 2$, so $G \cong K_1 \vee (K_2 \cup \frac{n-3}{3}K_3)$, which, by Lemma 4.4, is a contradiction.

If $X_2 \neq \emptyset$, then $|X'_2| = 1$, $G \cong K_1 \lor (K_{1,t} \cup \frac{n-t-2}{3}K_3)$ for some $2 \leqslant t \leqslant n-2$, so we have by Lemma 4.5 that t = n-2, i.e., $G \cong K_{1,1,n-2}$. **Case 3.2.** $X'_0 \neq \emptyset$.

In this case, $X_1 \cup X'_2 \neq \emptyset$. As $X'_0 \neq \emptyset$, we have $v_0 w \in E(G)$ for some $v_0 \in X'_0$ and some $w \in Y_1 \cup Y'_2$.

Case 3.2.1. $w \in Y_1$.

From (4.6), $Y_2 = Y'_2 = \emptyset$.

Let v_1 be a neighbor of w in X_1 and v_2 the neighbor of v_1 in X. Let $X'_1 = \{v_1, v_2\}, X''_0 = \bigcup_{v \in X'_1} X_v, Y^* = N_Y(v_1) \cup N_Y(v_2)$ and $Y' = N_Y(v_1) \cap N_Y(v_2)$.

By similar argument as in the proof of Claim 4.9, $Y' = \{w\}$ or $Y' = \emptyset$.

Suppose that $Y' = \{w\}$. Then $e(X''_0, Y_1) = |X''_0|$. As $e(X'_1, Y_1) = |Y^*| + 1$, we have by (4.1) that $|Y^*| = 1$, $e(X'_1, Y^*) = 2$ and $1 \leq |X''_0| \leq 3$. So $N_G(w) = X'_1 \cup X''_0$. Let $F_2 = G[\{u\} \cup N_G[w]]$. Then F_2 is the graph obtained from $K_{2,|X''_0|+2}$ with bipartition $(\{u, w\}, N_G(w))$ by adding an edge v_1v_2 . Thus, if $n = 4 + |X''_0|$, then $G = F_2$, and if $n > 4 + |X''_0|$, then G consists of F_2 and $G[V(G) \setminus N_G[w]]$ with a common vertex u. Let F'_2 be a copy of $K_{1,1,|X''_0|+2}$ obtained from $K_{2,|X''_0|+2}$ as above by adding an edge uw. By a direct calculation, we have

$$\rho(F_2) = \begin{cases}
2.8858 & \text{if } |X_0''| = 1, \\
3.1413 & \text{if } |X_0''| = 2, \text{ and } \rho(F_2 - u) = \begin{cases}
2.1701 & \text{if } |X_0''| = 1, \\
2.3429 & \text{if } |X_0''| = 2, \\
2.5141 & \text{if } |X_0''| = 3.
\end{cases}$$

From Lemma 2.6,

$$\rho(F_2') = \begin{cases} 3 & \text{if } |X_0''| = 1, \\ 3.3723 & \text{if } |X_0''| = 2, \\ 3.7016 & \text{if } |X_0''| = 3. \end{cases}$$

Evidently, $F'_2 - u \cong K_{1,|X'_0|+2}$, so

$$\rho(F'_2 - u) = \begin{cases} \sqrt{3} & \text{if } |X''_0| = 1, \\ 2 & \text{if } |X''_0| = 2, \\ \sqrt{5} & \text{if } |X''_0| = 3. \end{cases}$$

Thus

$$\rho(F_2) < \rho(F'_2) \text{ and } \rho(F_2 - u) > \rho(F'_2 - u).$$

If $n = 4 + |X_0''|$, then $\rho(F_2) < \rho(F_2')$, a contradiction, and if $n > 4 + |X_0''|$, then we have by Lemma 2.5 that $\rho(F_2' uG[V(G) \setminus N_G[w]]) > \rho$, also a contradiction.

Suppose that $Y' = \emptyset$. Then $e(X_1, Y_1) = |Y^*|$ and $e(X_0'', Y_1) \leq |X_0''| + |Y^*| - 1$, so we have by (4.1) that $|Y^*| = 1$ and $|X_0''| \leq 2$. As $G + v_2w$ also does not contain a DCC₁, we have by Lemma 2.1 that $\rho(G + v_2w) > \rho$, a contradiction.

Case 3.2.2. $w \in Y'_2$.

Let $v \in N_X(w)$. As G does not contain a DCC₁, $N_Y(w) = \emptyset$. So $G' = G - \{vz : z \in N_Y(v)\} + \{uz : z \in N_Y(v)\}$ also does not contain a DDC₁. By Lemma 2.2, $\rho(G') > \rho$, a contradiction.

Combining the above cases, we have $G \cong F_1, K_{1,1,n-2}, K_{3,n-3}$. Note that $\rho(F_1) > \rho(K_{1,1,3}) > \rho(K_{3,2})$. By Lemma 2.6, we have (1.1).

5 Proof of Theorem 1.3

Lemma 5.1. [28] Let H_n be defined as in Theorem 1.3. Then

$$\rho(H_n) = \begin{cases} \frac{n+1}{2} & \text{if } n \equiv 1 \pmod{2}, \\ \frac{1+\sqrt{n^2+1}}{2} & \text{if } n \equiv 0 \pmod{4}, \\ \eta(n) & \text{if } n \equiv 2 \pmod{4}, \end{cases}$$

where $\eta(n)$ is the largest root of $x^3 - x^2 - \frac{n^2}{4}x + \frac{n}{2} = 0$.

Proof of Theorem 1.3. Suppose that G is an n-vertex $K_1 \vee P_4$ -free graph that maximizes the spectral radius. By similar argument as in the proof of Theorem 1.1, G is connected. It suffices to show that $G \cong H_n$.

Let $\rho = \rho(G)$ and let **x** be the Perron vector of G and u a vertex with maximum entry in **x**. Let $X = N_G(u)$ and $Y = V(G) \setminus N_G[u]$. From (2.1), we have

$$\rho^2 \leqslant |X| + 2e(X) + e(X, Y).$$

Note that H_n is $K_1 \vee P_4$ -free, $\rho(G) \ge \rho(H_n)$. By Lemma 5.1, $\rho(G)^2 \ge \rho(H_n)^2 \ge \frac{(n+1)^2-1}{4}$, and moreover, $\rho(G)^2 \ge 16$ if n = 7 and $\rho(G)^2 \ge 21$ if n = 8. Thus

$$|X| + 2e(X) + e(X,Y) \ge \frac{(n+1)^2 - 1}{4}$$
(5.1)

and

$$|X| + 2e(X) + e(X, Y) \ge \begin{cases} 16 & \text{if } n = 7, \\ 21 & \text{if } n = 8. \end{cases}$$
(5.2)

As G is $K_1 \vee P_4$ -free, G[X] is P_4 -free, so each component of G[X] is a triangle K_3 , isolated vertex K_1 , isolated edge K_2 , or a star $K_{1,r}$ with some $r \ge 2$. In particular, $e(X) \le |X| = d_u$.

Claim 5.1. For any $w \in Y$, w is adjacent to at most one vertex of a triangle in G[X].

Proof. Suppose that there is a vertex $w \in Y$ such that it is adjacent to two vertices, say v_1, v_2 , of a triangle with vertices v_1, v_2, v_3 in G[X]. Suppose that G[X] contains a component isomorphic to K_3 with vertices v_1, v_2, v_3 . Assume that $v_1w, v_2w \in E(G)$. Then wv_1uv_3 is a copy of P_4 in G and v_2 is adjacent to all vertices of this path, so G contains $K_1 \vee P_4$ as a subgraph, a contradiction.

Let $d_2(u)$ be the number of vertices of distance two from u in G.

Claim 5.2. $d_2(u) = n - 1 - d_u$.

Proof. If G[X] contains a triangle, then $e(X,Y) \leq (d_u-2)d_2(u)$ by Claim 5.1, so

$$d_u + 2e(X) + e(X, Y) \leq d_u + 2d_u + d_u d_2(u) - 2d_2(u)$$

= $d_u(d_2(u) + 3) - 2d_2(u)$
 $\leq \frac{(d_u + d_2(u) + 3)^2}{4} - 2,$

and if G[X] does not contain any triangle, then $e(X) \leq |X| - 1 = d_u - 1$, so

$$d_u + 2e(X) + e(X,Y) \le d_u + 2(d_u - 1) + d_u d_2(u) \le \frac{(d_u + d_2(u) + 3)^2}{4} - 2.$$

In either case, we have by (5.1) that

$$\frac{(n+1)^2 - 1}{4} \leqslant d_u + 2e(X) + e(X,Y) \leqslant \frac{(d_u + d_2(u) + 3)^2}{4} - 2,$$

implying that $d_u + d_2(u) + 3 > n+1$, i.e., $d_u + d_2(u) \ge n-1$, which shows $d_u + d_2(u) = n-1$. Claim 5.3. G[X] does not contain a triangle.

Proof. Suppose that G[X] contains at least a triangle. Let t be the number of triangles in G[X].

Suppose that there is a component in G[X] that is not a triangle. By Claims 5.1 and 5.2,

$$d_u + 2e(X) + e(X, Y) \leq d_u + 2(d_u - 1) + (d_u - 2t)(n - 1 - d_u)$$

$$\leq d_u + 2(d_u - 1) + (d_u - 2)(n - 1 - d_u)$$

$$= -(d_u^2 - (n + 4)d_u + 2n),$$

so from (5.1), we have

$$d_u^2 - (n+4)d_u + 2n + \frac{(n+1)^2 - 1}{4} \leqslant 0,$$

which is impossible as the discriminant of the quadratic polynomial in d_u in the left hand side of the above inequality is $(n + 4)^2 - 8n - (n + 1)^2 + 1 < 0$ when $n \ge 9$. If n = 8, then from (5.2), we have $d_u^2 - 12d_u + 37 \le 0$, which is also impossible. Suppose that n = 7. From (5.2), we have $d_u^2 - 11d_u + 30 \le 0$, so $d_u = 5, 6$, implying that $G \cong F_3, K_1 \lor (K_3 \cup K_{1,2})$, where F_3 is a graph obtained from $K_1 \lor (K_3 \cup K_2)$ by adding a vertex adjacent to one vertex of K_3 and one vertex of K_2 . By a direct calculation, $\rho(F_3), \rho(K_1 \lor (K_3 \cup K_{1,2})) < \rho(H_n)$, a contradiction. Suppose that n = 6. Then $d_u^2 - 10d_u + 24 \le 0$, so $d_u = 4, 5$, implying that $G \cong F_4, K_1 \lor (K_3 \cup K_2)$, where F_4 is a graph obtained from $K_1 \lor (K_3 \cup K_1)$ by adding a vertex adjacent to one vertex of K_3 and the vertex of degree one. By a direct calculation, $\rho(G) < \rho(H_6)$, a contradiction. It now follows that every component in G[X] is a triangle. So $G[X] \cong tK_3$ with $d_u = 3t$ for some $t \ge 1$. By (5.1), Claims 5.1 and 5.2, we have

$$\frac{(n+1)^2 - 1}{4} \leqslant 3t + 6t + t(n-1-3t) = -3t^2 + (n+8)t,$$

i.e.,

$$3t^2 - (n+8)t + \frac{(n+1)^2 - 1}{4} \leqslant 0.$$

which is also impossible as $(n+8)^2 - 3(n+1)^2 + 3 = -2(n^2 - 5n + 32) < 0.$

By Claim 5.3, any component of G[X] with at least three vertices is a star.

Claim 5.4. Let H be a component of G[X] with at least three vertices and v be its center. For any $w \in Y$, if $vw \in E(G)$, then v is the unique neighbor of w in H.

Proof. Suppose that there is a vertex $v_1 \in V(H) \setminus \{v\}$ such that $v_1w \in E(G)$. Let v_2 be another vertex in H different from v and v_1 . Then v_2uv_1w is a P_4 in G and v is adjacent to all such vertices, which shows that G contains $K_1 \vee P_4$ as a subgraph, a contradiction. \Box

From Claim 5.4, we have $d_X(w) \leq d_u - r$ for any $w \in Y$, where r denotes the number of components of G[X] with at least three vertices.

Claim 5.5. There is at most one component with at least three vertices in G[X]. Moreover, if such a component exists, then G[X] has exactly one component and $e(X,Y) \ge (d_u-1)(n-1-d_u)-1$.

Proof. Denote by r the number of components of G[X] with at least three vertices. If $r \ge 2$, then we have by (5.1) and Claim 5.4 that

$$\frac{(n+1)^2 - 1}{4} \leqslant d_u + 2(d_u - r) + (d_u - r)(n - 1 - d_u) \leqslant d_u + 2(d_u - 2) + (d_u - 2)(n - 1 - d_u),$$

i.e.,

$$d_u^2 - (n+4)d_u + 2(n+1) + \frac{(n+1)^2 - 1}{4} \le 0,$$

which is impossible. So $r \leq 1$.

Suppose that r = 1. If G[X] has at least two components, then $e(X) \leq d_u - 2$, so we have by (5.1) that

$$\frac{(n+1)^2 - 1}{4} \leqslant d_u + 2(d_u - 2) + (d_u - 1)(n - 1 - d_u),$$

i.e.,

$$d_u^2 - (n+3)d_u + (n+3) + \frac{(n+1)^2 - 1}{4} \le 0,$$

which is impossible as $(n+3)^2 - 4(n+3) - (n+1)^2 + 1 = -3 < 0$. So G[X] contains exactly one component. If $e(X, Y) \leq (d_u - 1)(n - 1 - d_u) - 2$, then by (5.1),

$$\frac{(n+1)^2 - 1}{4} \leq d_u + 2(d_u - 1) + (d_u - 1)(n - 1 - d_u) - 2,$$

so $d_u^2 - (n+3)d_u + (n+3) + \frac{(n+1)^2 - 1}{4} \leq 0$, which is impossible. Thus $e(X, Y) \ge (d_u - 1)(n - 1 - d_u) - 1$.

Now we consider two cases.

Case 1. There is exactly one component with at least three vertices in G[X].

By Claim 5.5, $G[X] \cong K_{1,d_u-1}$. It follows from (5.1) that

$$\frac{(n+1)^2 - 1}{4} \leq |X| + 2e(X) + e(X,Y) \leq d_u + 2(d_u - 1) + (d_u - 1)(n - 1 - d_u),$$

implying that $\frac{n+3-\sqrt{5}}{2} \leq d_u \leq \frac{n+3+\sqrt{5}}{2}$. That is,

$$d_u = \begin{cases} \frac{n+1}{2}, \frac{n+3}{2}, \frac{n+5}{2} & \text{if } n \text{ is odd,} \\ \frac{n+2}{2}, \frac{n+4}{2} & \text{if } n \text{ is even.} \end{cases}$$

If n = 6, then $d_u = 4, 5$, so $G \cong H_6, K_{1,1,4}$, and a direct calculation shows that $\rho(K_{1,1,4}) = 3.3723 < 3.3923 = \rho(H_6)$, so $G \cong H_6$. If n = 7, then $d_u = 4, 5, 6$, so $G \cong H_n, 4K_1 \lor (K_2 \cup K_1), K_{1,1,6}$, and a direct calculation shows that $\rho(4K_1 \lor (K_2 \cup K_1)) = 3.3273 < 3.7016 = \rho(K_{1,1,5}) < 4 = \rho(H_n)$, so $G \cong H_7$. Suppose that $n \ge 8$. Then $d_u \ge 5$. Let v be the center of G[X]. If $vw \in E(G)$ for some $w \in Y$, then by Claim 5.4, we have $e(X, Y) \le (d_u - 1)(n - 2 - d_u) + 1 < (d_u - 1)(n - 1 - d_u) - 2$, contradicting Claim 5.5, so $vw \notin E(G)$ for any $w \in Y$. By Claim 5.5, there are at least two vertices $v_1, v_2 \in X$ that are adjacent to each vertex in Y. If G[Y] contains a path P_3 , say $w_1w_2w_3$, then $G[\{w_1, v_1, w_3, v_2\}]$ contains a path P_4 and w_2 is adjacent to the vertices of the path, so G contains $K_1 \lor P_4$, a contradiction. Thus G[Y] is does not contain any P_3 , i.e., G[Y] consists of isolated vertices or edges. As adding edges between isolated vertices in G[Y] does not lead to a copy of

 $K_1 \vee P_4$, we have by Lemma 2.1 that G[Y] contains at most one isolated vertex. Therefore, $G \cong aK_1 \vee (bK_2 \cup cK_1)$, where $a = d_u - 1$, $c \leq 1$ and a + 2b + c = n. Case 1.1 c = 0.

In this case, $G \cong aK_1 \vee \frac{n-a}{2}K_2$, and it is easy to see $\rho(G) = \frac{1}{2}(1 + \sqrt{1 + 4a(n-a)})$, so

$$a = \begin{cases} \frac{n+1}{2} & \text{if } n \equiv 1 \pmod{4}, \\ \frac{n-1}{2} & \text{if } n \equiv 3 \pmod{4}, \\ \frac{n}{2} & \text{if } n \equiv 0 \pmod{4}, \\ \frac{n+2}{2} & \text{if } n \equiv 2 \pmod{4}. \end{cases}$$

Therefore, $G \cong H_n$ if $n \equiv 0, 1, 3 \pmod{4}$. If $n \equiv 2 \pmod{4}$, then $G \cong \frac{n+2}{2}K_1 \vee \frac{n-2}{4}K_2$, and a direct calculation and Lemma 5.1 show that $\rho(G) = \frac{1+\sqrt{n^2-3}}{2} < \rho(H_n)$, a contradiction. **Case 1.2.** c = 1.

In this case, $G \cong aK_1 \vee (\frac{n-a-1}{2}K_2 \cup K_1)$ and A(G) has an equitable quotient matrix

$$B' = \begin{pmatrix} 0 & n-a-1 & 1\\ a & 1 & 0\\ a & 0 & 0 \end{pmatrix}$$

with respect to vertex partition $V(G) = V(aK_1) \cup V(\frac{n-a-1}{2}K_2) \cup V(K_1)$, so by Lemma 2.3, $\rho(G)$ is equal to the largest root of g(x, a) = 0 with $g(x, a) := \det(xI_3 - B') = x^3 - x^2 - a(n-a)x + a$. Note that

$$g(x,a) - g(x,a+1) = (n - 2a - 1)x - 1 < 0$$

as $a = d_u - 1 \ge \frac{n-1}{2}$. So

$$a = \begin{cases} \frac{n-1}{2} & \text{if } n \equiv 1 \pmod{4}, \\ \frac{n+1}{2} & \text{if } n \equiv 3 \pmod{4}, \\ \frac{n+2}{2} & \text{if } n \equiv 0 \pmod{4}, \\ \frac{n}{2} & \text{if } n \equiv 2 \pmod{4}. \end{cases}$$

Therefore, $G \cong H_n$ if $n \equiv 2 \pmod{4}$. Otherwise, $G \cong \frac{n-1}{2}K_1 \vee (\frac{n-1}{4}K_2 \cup K_1)$ if $n \equiv 1 \pmod{4}$, $G \cong \frac{n+1}{2}K_1 \vee (\frac{n-3}{4}K_2 \cup K_1)$ if $n \equiv 3 \pmod{4}$ and $G \cong \frac{n+2}{2}K_1 \vee (\frac{n-4}{4}K_2 \cup K_1)$ if $n \equiv 0 \pmod{4}$. However, by an easy calculation,

$$g(\rho(H_n), a) = \begin{cases} \frac{n-1}{2} & \text{if } n \equiv 1 \pmod{4} \\ \frac{n+1}{2} & \text{if } n \equiv 3 \pmod{4} \\ \frac{(n+2)(n+4)}{4} & \text{if } n \equiv 0 \pmod{4} \\ > 0 \end{cases}$$

so $\rho(G) < \rho(H_n)$ if $n \equiv 0, 1, 3 \pmod{4}$, a contradiction.

Combining Cases 1.1 and 1.2, we have $G \cong H_n$.

Case 2. G[X] does not contain a component with at least three vertices.

In this case, G[X] consists of isolated vertices and isolated edges, and $e(X) \leq \lfloor \frac{d_u}{2} \rfloor$. We claim that G[X] contains at most one isolated vertex, as otherwise, $e(X) \leq \frac{d_u-2}{2}$, so we have by (5.1) that

$$\frac{(n+1)^2 - 1}{4} \leqslant d_u + d_u - 2 + d_u(n-1-d_u),$$

i.e.,

$$d_u^2 - (n+1)d_u + \frac{(n+1)^2 - 1}{4} + 2 \leq 0.$$

which is impossible as $(n+1)^2 - (n+1)^2 + 1 - 8 = -7 < 0$. Similarly, $e(X, Y) \ge d_u(n-1-d_u) - 1$. By (5.1),

$$\frac{(n+1)^2 - 1}{4} \leqslant |X| + 2e(X) + e(X,Y) \leqslant d_u + 2\left\lfloor \frac{d_u}{2} \right\rfloor + d_u(n-1-d_u),$$

so d_u is even with $\frac{n}{2} \leq d_u \leq \frac{n+2}{2}$.

If n = 6, then $d_u = 4$, so the inequality above is actually an equality, so $G \cong 2K_1 \vee 2K_2$ with $\rho(2K_1 \vee 2K_2) = 3.3723 < \rho(H_6)$, a contradiction. If n = 7, then we have from (5.2) that $16 \leq d_u + d_u + d_u(6 - d_u)$, so $d_u = 4$ and $G \cong H_n$, as desired. If n = 8, then $21 \leq d_u + d_u + d_u(7 - d_u)$, which is impossible.

Suppose that $n \ge 9$. Then $d_u \ge 5$, so there are at least two adjacent vertices $v_1, v_2 \in X$ that are adjacent to each vertex in Y. If G[Y] contains an edge w_1w_2 , then $G[\{u, v_2, w_1, w_2\}]$ contains a path P_4 as a subgraph and v_1 is adjacent to the vertices of the path, so G contains $K_1 \lor P_4$ as a subgraph, a contradiction. Thus Y is independent. As adding possible edges between vertices in X and Y does not lead to a copy of $K_1 \lor P_4$, we have by Lemma 2.1 that each vertex in X is adjacent to all vertices in Y. Therefore, $G \cong (n - d_u)K_1 \lor (bK_2 \cup cK_1)$, where $c \le 1$ and $2b + c = d_u$. If $n \equiv 0 \pmod{4}$, then $d_u = \frac{n}{2}$, so $G \cong H_n$, as desired. If $n \equiv 2 \pmod{4}$, then $d_u = \frac{n+2}{2}$, so $G \cong \frac{n-2}{2}K_1 \lor \frac{n+2}{4}K_2$, for which, we have by a direct calculation that $\rho(G) = \frac{1+\sqrt{n^2-3}}{2} < \rho(H_n)$, a contradiction. If $n \equiv 1 \pmod{4}$, then $d_u = \frac{n+1}{2}$, which contradicts that d_u is even. If $n \equiv 3 \pmod{4}$, then $d_u = \frac{n+1}{2}$ and $G \cong H_n$, as desired. \Box

Acknowledgements

This work was supported by National Natural Science Foundation of China (No. 12071158).

References

- [1] B. Bollobás, Extremal Graph Theory, Academic Press, London, 1978.
- [2] J.A. Bondy, Large cycles in graphs, Discrete Math. 1 (2) (1971) 121–132.
- [3] A. Brouwer, W. Haemers, Spectra of Graphs, Springer, New York, 2012.
- [4] R.A. Brualdi, E.S. Solheid, On the spectral radius of connected graphs, Publ. Inst. Math. (Beograd) (N.S.) 39 (53) (1986) 45–54.

- [5] R.A. Brualdi, E.S. Solheid, On the spectral radius of complementary acyclic matrices of zeros and ones, SIAM J. Algebraic Discrete Methods 7 (2) (1986) 265–272.
- [6] J. Cai, L. Xu, B. Zhou, Spectral conditions for the existence of (doubly) chorded cycles in graphs with fixed size, ArXiv: 2405.03229v1.
- [7] S. Chiba, T. Yamashita, Degree conditions for the existence of vertex-disjoint cycles and paths: a survey, Graphs Combin. 34 (1) (2018) 1–83.
- [8] S. Cioabă, D. Desai, M. Tait, A spectral Erdős–Sós, theorem, SIAM J. Discrete Math. 37 (2023) 2228–2239.
- [9] C.M. Conde, E. Dratman, L.N. Grippo, On the spectral radius of block graphs having all their blocks of the same size, Linear Algebra Appl., 634 (2022) 137–148.
- [10] M. Cream, R.J. Faudree, R.J. Gould, K. Hirohata, Chorded cycles, Graphs Combin. 32 (6) (2016) 2295–2313.
- [11] J. Czipszer, Solution to Problem 127 (in Hungarian), Mat. Lapok 14 (1963) 373–374.
- [12] N. Draganić, A. Methuku, D. Munhá Correia, B. Sudakov, Cycles with many chords, Random Struct. Alg. 65 (2024) 3–16.
- [13] F. Esser, F. Harary, On the spectrum of a complete multipartite graph, Eur. J. Comb. 1 (1980) 2110–218.
- [14] Y. Gao, X. Lin, H. Wang, Vertex-disjoint double chorded cycles in bipartite graphs, Discrete Math. 342 (9) (2019) 2482–2492.
- [15] R.J. Gould, Results and problems on chorded cycles: a survey, Graphs Combin. 38 (2022) 189, 27 pp.
- [16] R.J. Gould, K. Hirohata, P. Horn, On independent doubly chorded cycles, Discrete Math. 338 (11) (2015) 2051–2071.
- [17] T. Jiang, A note on a conjecture about cycles with many incident chords, J. Graph Theory 46 (3) (2004) 180–182.
- [18] Y. Li, Y. Peng, The spectral radius of graphs with no intersecting odd cycles, Discrete Math. 345 (2022) 112907.
- [19] L. Lovász, Combinatorial Problems and Exercises, North-Holland, 1979.
- [20] V. Nikiforov, The spectral radius of graphs without paths and cycles of specified length, Linear Algebra Appl. 432 (2010) 2243–2256.
- [21] L. Pósa, Problem 127 (in Hungarian), Mat. Lapok 12 (1961) 254.

- [22] M. Santana, M. Van Bonn, Sharp minimum degree conditions for disjoint doubly chorded cycles, J. Comb. 15 (2) (2024) 217–282.
- [23] H. Wang, X. Hou, Y. Ma, Spectral extrema of graphs with bounded clique number and matching number, Linear Algebra Appl. 669 (2023) 125–135.
- [24] L. Xu, B. Zhou, Answers to Gould's question concerning the existence of chorded cycles, ArXiv: 2312.1679.
- [25] L. Yu, Y. Li, Y. Peng, Spectral extremal graphs for an graphs, ArXiv: 2404.03423.
- [26] Q. Yuan, R. Liu, J. Yuan, The spectral radius of H_{2k} -free graphs, Appl. Math. Comput. 469 (2024) 128560.
- [27] M. Zhai, R. Liu, J. Xue, A unique characterization of spectral extrema for friendship graphs, Electron. J. Combin. 29 (3) (2022) 3.32.
- [28] Y. Zhao, X. Huang, H. Lin, The maximum spectral radius of wheel-free graphs, Discrete Math. 344 (2021) 112341.
- [29] J. Zheng, X. Huang, J. Wang, Spectral radius for the existence of a chorded cycle, ArXiv: 2311.13323v2.