

# A HILTON-MILNER THEOREM FOR EXTERIOR ALGEBRAS

DENYS BULAVKA, FRANCESCA GANDINI, AND RUSS WOODROOFE

ABSTRACT. Recent work of Scott and Wilmer and of Woodroffe extends the Erdős-Ko-Rado theorem from set systems to subspaces of  $k$ -forms in an exterior algebra. We prove an extension of the Hilton-Milner theorem to the exterior algebra setting, answering in a strong way a question asked by these authors.

## 1. INTRODUCTION

A family of sets  $\mathcal{F}$  is *pairwise-intersecting* if every pair of sets in  $\mathcal{F}$  have nonempty intersection. Erdős, Ko, and Rado [6] gave an upper bound on the size of a pairwise-intersecting family of small sets, and characterized the families that achieve the upper bound.

**Theorem 1.1** (Erdős-Ko-Rado). *Let  $k \leq n/2$ . If  $\mathcal{F} \subseteq \binom{[n]}{k}$  is a pairwise-intersecting family of sets, then  $|\mathcal{F}| \leq \binom{n-1}{k-1}$ . Moreover, if  $k < n/2$  and  $|\mathcal{F}|$  achieves the upper bound, then  $\mathcal{F}$  consists of all the  $k$ -subsets containing some fixed element.*

There are many generalizations and analogues of Theorem 1.1. In this article, we focus on the extension to exterior algebras that was considered by Scott and Wilmer [17] and by the third author [18]. We first fix some notation:

*Notation 1.2.* Through the paper, let  $\mathbb{F}$  be a field. We assume for expository purposes that the characteristic is not 2, although the fundamental techniques are independent of characteristic. Let  $V$  be a vector space of dimension  $n$  over  $\mathbb{F}$ . Let  $\mathbf{e} = \{e_1, \dots, e_n\}$  be a given basis for  $V$ . If  $I$  is a subset of  $[n]$ , then let  $V^{(I)}$  be the subspace spanned by  $\{e_j : j \notin I\}$ , and write  $V^{(i)}$  for  $V^{\{i\}}$ .

A subset  $L$  of the exterior algebra is *self-annihilating* if  $L \wedge L = 0$ ; that is, if for every  $x, y \in L$ , it holds that  $x \wedge y = 0$ .

**Theorem 1.3** (Scott and Wilmer). *Let  $k \leq n/2$ . If  $L$  is a self-annihilating subspace of  $\wedge^k V$ , then  $\dim L \leq \binom{n-1}{k-1}$ .*

---

Work started while the first author was at the Department of Applied Mathematics, Faculty of Mathematics and Physics, Charles University, Prague, Czech Republic. Work of the first author is partially supported by the GAČR grant no. 22-19073S and by the Israel Science Foundation grant ISF-2480/20. Work of the second author is supported in part by the Slovenian Research Agency (research projects N1-0160, J1-3003). Work of the third author is supported in part by the Slovenian Research Agency research program P1-0285 and research projects J1-9108, N1-0160, J1-2451, J1-3003, and J1-50000.

The bound of Theorem 1.3 is an extension of the bound of Theorem 1.1, as follows. Given a set  $F = \{i_1 < i_2 < \dots < i_k\}$  in  $\mathcal{F}$ , we represent it with the exterior monomial  $e_F = e_{i_1} \wedge \dots \wedge e_{i_k}$  in  $\wedge^k V$ . Now taking  $\{e_F : F \in \mathcal{F}\}$  as a basis, using that  $e_F \wedge e_G = 0$  if and only if  $F \cap G \neq \emptyset$ , and applying the distributive axiom, we obtain the bound of Theorem 1.1 from Theorem 1.3. With this tight connection, Theorem 1.3 may indeed be viewed as a categorification of the Erdős-Ko-Rado bound.

Scott and Wilmer asked in [17, Section 2.2] whether the characterization of the maximal families in Theorem 1.1 extends to the exterior algebra setting. The third author of the current article asked in [18, Question 5.1] the even stronger question of whether the Hilton-Milner upper bound extends. We will answer both questions in the affirmative.

We first recall the theorem of Hilton and Milner [11]. We say that a pairwise-intersecting set family is *nontrivial* if  $\bigcap_{F \in \mathcal{F}} F$  is empty.

**Theorem 1.4** (Hilton-Milner). *Let  $k \leq n/2$ . If  $\mathcal{F} \subseteq \binom{[n]}{k}$  is a nontrivial pairwise-intersecting family of sets, then  $|\mathcal{F}| \leq \binom{n-1}{k-1} - \binom{n-k-1}{k-1} + 1$ .*

Our main theorem will extend Theorem 1.4 to exterior algebras. We say that a self-annihilating subset  $L$  of  $\wedge V$  is *nontrivial* if there is no 1-form in  $V = \wedge^1 V$  that annihilates  $L$ .

**Theorem 1.5.** *Let  $k \leq n/2$ . If  $L$  is a nontrivial self-annihilating subspace of  $\wedge^k V$ , then  $\dim L \leq \binom{n-1}{k-1} - \binom{n-k-1}{k-1} + 1$ .*

Theorem 1.5 categorifies Theorem 1.4 in the same way that Theorem 1.3 categorifies the bound of Theorem 1.1. Since  $\binom{n-k-1}{k-1} > 1$  when  $k < n/2$ , we recover the exterior extension of the characterization of maximal families in Theorem 1.1.

**Corollary 1.6.** *Let  $k < n/2$ . If  $L$  is a self-annihilating subspace of  $\wedge^k V$  such that  $\dim L$  achieves the upper bound of  $\binom{n-1}{k-1}$ , then  $L$  is annihilated by a 1-form.*

Theorem 1.1 was earlier shown in [13] for the restrictive case where  $k = 2$ . Draisma, Kraft, and Kuttler [3] considered bounds on nontrivially nilpotent subspaces in a somewhat different setting.

Our strategy is to adapt proofs of Theorem 1.4, variously due to Frankl and/or Füredi [7, 8, 9] and to Hurlbert and Kamat [12], to the exterior algebra setting. These proofs start with a nontrivial pairwise-intersecting set family, and apply combinatorial shifting operations. If the family becomes trivial at some step, then this guarantees strong structural properties at the previous step. Wise choices of further combinatorial shifting operations results in a nontrivial pairwise-intersecting family that is of the same size and shifted.

In order to implement this strategy in the exterior algebra setting, we introduce a parameterized family of linear maps. The limit of the action of these maps extends combinatorial shifting to exterior algebras, and preserves more structure than those used by the third author in [18, Section 4]. We act with these maps on self-annihilating subspaces and study the resulting (limit) subspaces. Our techniques are characteristic

independent; although they are based on ideas from algebraic geometry, our presentation here is self-contained and relatively elementary.

In addition to an analogue of Theorem 1.4, we also prove an extension of another theorem of Hilton and Milner. Set families  $\mathcal{F}$  and  $\mathcal{G}$  are *cross-intersecting* if for every  $F \in \mathcal{F}$  and  $G \in \mathcal{G}$ , the intersection  $F \cap G$  is nonempty. Hilton and Milner gave bounds on the size of  $|\mathcal{F}| + |\mathcal{G}|$  under the conditions that  $\mathcal{F}, \mathcal{G}$  are nonempty and cross-intersecting (as a special case of a highly technical more general result [11, Theorem 2]).

The exterior algebra analogue is as follows. Subsets  $K$  and  $L$  of  $\wedge V$  are *cross-annihilating* if  $K \wedge L = 0$ , that is, if for every  $x \in K$  and  $y \in L$  it holds that  $x \wedge y = 0$ . We show:

**Theorem 1.7.** *Let  $k \leq n/2$ . If  $K$  and  $L$  are nonzero cross-annihilating subspaces of  $\wedge^k V$ , then  $\dim K + \dim L \leq \binom{n}{k} - \binom{n-k}{k} + 1$ .*

Scott and Wilmer also consider a cross-intersecting theorem in [17, Theorem 2.4], where they bounded the product of  $\dim K$  and  $\dim L$ .

The rest of this article is organized as follows. In Section 2 we give various preliminaries and background results. In Section 3 we introduce the linear maps whose limits will allow us to replace a given self-annihilating subspace with one that is easier to analyze. In Section 4 we prove Theorem 1.7, and in Section 5 we prove Theorem 1.5.

## ACKNOWLEDGEMENTS

We thank Jake Levinson for clarifying our doubts regarding limits on the Grassmannian, and Allen Knutson for answering our questions on the relationship to limits in algebraic geometry.

## 2. PRELIMINARIES

**2.1. Combinatorial shifting and shifted families of sets.** Let  $\mathcal{F} \subseteq \binom{[n]}{k}$  be a family of sets, let  $F \in \mathcal{F}$ , and let  $1 \leq i < j \leq n$ . The *combinatorial shift* operation  $\text{shift}_{j \rightarrow i}$  is defined as follows.

$$\text{shift}_{j \rightarrow i}(F, \mathcal{F}) = \begin{cases} (F \setminus \{j\}) \cup \{i\} & \text{if } j \in F, i \notin F \text{ and } (F \setminus \{j\}) \cup \{i\} \notin \mathcal{F}, \\ F & \text{otherwise.} \end{cases}$$

$$\text{shift}_{j \rightarrow i} \mathcal{F} = \{\text{shift}_{j \rightarrow i}(F, \mathcal{F}) : F \in \mathcal{F}\}.$$

For  $I \subseteq [n]$ , we say that a family  $\mathcal{F}$  of subsets of  $[n]$  is *shifted with respect to  $I$*  if  $\text{shift}_{j \rightarrow i} \mathcal{F} = \mathcal{F}$  for each pair  $i < j$  with  $i, j \in I$ . If  $I = [n]$ , we simply say that  $\mathcal{F}$  is *shifted*. It is well-known that iteratively applying combinatorial shift operations results in a shifted family [7, Section 2]; see also Theorem 3.9 below.

The technique of combinatorial shifting was introduced by Erdős, Ko, and Rado in [6] to prove Theorem 1.1. It has since become a standard tool in combinatorial set theory [7].

**2.2. Exterior algebras.** The *exterior algebra* over the vector space  $V$ , denoted  $\wedge V$ , is a graded algebra that is described as follows. For each  $0 \leq k \leq n$ , we define the  $k$ th graded component  $\wedge^k V$  as the span of elements  $\{e_S : S \in \binom{[n]}{k}\}$ . Here, we identify  $\wedge^1 V$  with  $V$  by identifying  $e_{\{i\}}$  with  $e_i$ ; we identify  $\wedge^0 V$  with  $\mathbb{F}$ . The elements of  $\wedge^k V$  are called *k-forms*.

Now the product is defined to satisfy the following axioms:  $e_\emptyset \wedge e_S = 1e_S = e_S$ ,  $e_i \wedge e_j = -e_j \wedge e_i$ , and  $e_S = e_{s_1} \wedge e_{s_2} \wedge \cdots \wedge e_{s_k}$  for  $S = \{s_1 < s_2 < \cdots < s_k\}$ . The elements  $e_S$  are *monomials with respect to the basis e*.

The exterior algebra is an antisymmetric analogue of the polynomial algebra, and more background may be found in most advanced algebra textbooks, such as [4]. It has long been used as a tool for proving results in combinatorial set theory [2].

Given a  $k$ -form  $x$  and an understood basis, the *support* of  $x$  is the set of monomials with nonzero coefficient in expression for  $x$  as a linear combination of monomials. The monomial  $e_S$  has *variable set*  $\{e_i : i \in S\}$ . When it causes no confusion, we may identify a variable set  $\{e_i : i \in S\}$  with the underlying set of indices  $S$ .

Given a linear operator  $M$  on  $V$ , we extend the linear operator to  $\wedge V$  by sending  $e_S = e_{s_1} \wedge e_{s_2} \wedge \cdots \wedge e_{s_k}$  to  $Me_{s_1} \wedge Me_{s_2} \wedge \cdots \wedge Me_{s_k}$  and extending linearly. We notice that in particular, for any  $x, y$  in  $\wedge V$ , we have that  $M(x \wedge y) = Mx \wedge My$ .

The following result on annihilation by 1-forms is well-known; see e.g. [10, Lecture 6].

**Lemma 2.1.** *For  $v \in \wedge^1 V$ ,  $x \in \wedge^k V$ , we have that  $v \wedge x = 0$  if and only if  $x = v \wedge x'$  for some  $x' \in \wedge^{k-1} V$ .*

**2.3. Limit actions on the Grassmannian.** Write  $\mathbb{P}V$  for the projective space defined on the vector space  $V$ , consisting of (equivalence classes of) vectors of  $V \setminus \{0\}$ , considered up to scalar multiplication. If  $v_1, v_2, \dots, v_k$  are linearly independent vectors in  $V$ , then we may identify the subspace spanned by these vectors with the point  $[v_1 \wedge v_2 \wedge \cdots \wedge v_k]$  in  $\mathbb{P}(\wedge^k V)$ . We get the *Grassmannian*  $\text{Gr}(k, V)$ , consisting of all points in  $\mathbb{P}(\wedge^k V)$  that may be written in this form. Thus, the Grassmannian is a geometric object whose points correspond to  $k$ -dimensional subspaces of  $V$ .

We will need to take limits as  $t \rightarrow 0$  of actions of families of matrices parameterized by  $t$ . The limits we will take are basically of an algebraic geometry nature, but we did not find accessible literature on the topic. For completeness, we give an elementary description, which we hope will be accessible to any reader who has had a solid graduate course in (linear) algebra.

We start by describing the action on an arbitrary projective space  $\mathbb{P}V$ . Next, extend our field  $\mathbb{F}$  to  $\mathbb{F}(t)$  by adjoining a transcendental  $t$ . Thus,  $\mathbb{F}(t)$  consists of ratios of polynomials in  $t$  with coefficients in  $\mathbb{F}$ .

We now extend coefficients in our vector space  $V$  over  $\mathbb{F}$  to a vector space  $V(t) \cong V \otimes_{\mathbb{F}} \mathbb{F}(t)$  over  $\mathbb{F}(t)$ . Thus, an element of  $V(t)$  is a linear combination of  $e_1, \dots, e_n$  with coefficients in  $\mathbb{F}(t)$ . By clearing denominators and/or dividing by  $t$  in  $\mathbb{P}V(t)$ , we may write an arbitrary projective element  $v(t)$  as a combination of  $e_1, \dots, e_n$ , where each coefficient is a polynomial in  $t$ , and where the coefficients do not have a common divisor polynomial of positive  $t$ -degree. This form is the *canonical representative of  $v(t)$* . In

particular, the (polynomial) coefficient of at least one  $e_i$  has a nonzero coefficient of the  $t$ -degree zero term. Now the map  $\text{eval}_0 : \mathbb{P}V(t) \rightarrow \mathbb{P}V$  obtained by evaluating the polynomials in the canonical representative at  $t = 0$  is a well-defined map.

Now let  $N(t)$  be a nonsingular linear map over  $\mathbb{F}(t)$ . We define the *limit* of the action of  $N(t)$  on  $\mathbb{P}V$  to be the composition

$$\mathbb{P}V \xrightarrow{\iota} \mathbb{P}V(t) \xrightarrow{N(t)} \mathbb{P}V(t) \xrightarrow{\text{eval}_0} \mathbb{P}V,$$

where  $\iota$  is the standard inclusion map. We write  $N$  for the resulting map  $\mathbb{P}V \rightarrow \mathbb{P}V$ .

**Example 2.2.** For the limit of  $N(t)$  acting on any fixed vector  $v$  of  $\mathbb{P}\mathbb{R}^2$ , we can write  $N(t)v = w$  in coordinates as  $[w_1(t), w_2(t)]$ . This point corresponds to the formal ratio  $w_1(t)/w_2(t)$ . Our description of the limit map resembles closely the usual procedure for evaluating  $\lim_{t \rightarrow 0} w_1(t)/w_2(t)$  from a first calculus course. Note that here the limit point  $[\alpha, 1]$  corresponds to the real number  $\alpha$ , while the (single) limit point  $[1, 0]$  corresponds to the calculus limits  $\pm\infty$ .

The action on  $\mathbb{P}(\wedge^k V)$  is now a special case: the basis for  $V$  gives a basis of monomials for  $\wedge^k V$ , and a linear map  $N(t)$  on  $V(t)$  induces a linear map on  $\wedge^k V(t)$ . Similarly for  $\mathbb{P}(\wedge^r(\wedge^k V))$ , given a nonnegative integer  $r$ .

We are ready to discuss the limit of the action on  $\text{Gr}(r, V)$  and  $\text{Gr}(r, \wedge^k V)$ .

**Proposition 2.3.** *Let  $N(t)$  be a nonsingular linear map  $V(t) \rightarrow V(t)$ . If  $L = v_1 \wedge \cdots \wedge v_r$  is a point on  $\text{Gr}(r, V) \subseteq \mathbb{P}(\wedge^r V)$ , then the limit  $NL$  is also in  $\text{Gr}(r, V)$ , and is the vector subspace spanned by  $\{Nw : w \in L\}$ .*

*Proof.* We first notice that a vector  $w$  is in  $L$  if and only if  $w \wedge L = 0$ . Now  $N(t)(w \wedge L) = N(t)w \wedge N(t)L = 0$ , and so the same happens in  $t$ -degree zero in the canonical representative. It follows that if  $w \wedge L = 0$ , then  $Nw \wedge NL = 0$ .

Conversely, we need to show that  $NL$  is annihilated by  $r$  linearly independent vectors in the image of  $N$ . Suppose that  $e_{i_1} \wedge \cdots \wedge e_{i_r}$  is in the support of  $NL$ . Then  $e_{i_1} \wedge \cdots \wedge e_{i_r}$  has a  $t$ -degree zero component in the canonical representative of  $N(t)L$ . The coefficient of this monomial is an  $r \times r$  minor of the the matrix with columns  $N(t)v_1, \dots, N(t)v_r$ . By Gaussian elimination, we may find vectors  $N(t)w_1, \dots, N(t)w_r$  in  $N(t)L$  so that the corresponding minor of  $N(t)w_1, \dots, N(t)w_r$  is diagonal and has a  $t$ -degree zero component. It follows that  $Nw_1, \dots, Nw_r$  are linearly independent, as desired.  $\square$

*Remark 2.4.* Equivalently, if the matrix  $W$  with columns  $N(t)v_1, \dots, N(t)v_r$  has an  $r \times r$  minor with a  $t$ -degree zero component, then let  $G$  be the corresponding  $r \times r$  submatrix. Now the columns of  $WG^{-1}$  are vectors in  $N(t)L$  whose limits are linearly independent.

**Corollary 2.5.** *Let  $N(t)$  be a nonsingular linear map  $V(t) \rightarrow V(t)$ . If  $L_0 \subseteq L$  are vector subspaces of  $V$ , then also  $NL_0 \subseteq NL$ .*

As before, the limit action  $N$  on  $\text{Gr}(r, \wedge^k V)$  is a special case.

For the purpose of this paper, we need that the limit preserves the self-annihilating and cross-annihilating properties. Experts in algebraic geometry will recognize this as following from the fact that these properties correspond to closed sets in the Zariski topology. An elementary proof is also easy.

**Lemma 2.6.** *Let  $N(t)$  be a nonsingular linear map  $V(t) \rightarrow V(t)$ . If  $L$  and  $L'$  are cross-annihilating subspaces of  $\wedge L$ , then  $NL$  and  $NL'$  are also cross-annihilating.*

*Proof.* Let  $x$  be in  $L$  and  $x'$  in  $L'$ . The  $t$ -degree zero term of the exterior product of the canonical representatives of  $N(t)x$  and  $N(t)x'$  is the product of the  $t$ -degree zero terms of  $N(t)x$  and  $N(t)x'$ . Now if  $x \wedge x' = 0$ , so that also  $N(t)x \wedge N(t)x' = 0$ , then we recover that  $Nx \wedge Nx' = 0$ .  $\square$

For further reading on projective space and the Grassmannian, we refer the reader to [1, 10], for example. Accessible literature on limits in algebraic geometry (also known as *degenerations* or *specializations*) seems to be a bit difficult to find. Artin [1, Chapter 5] discusses limits of curves in algebraic geometry briefly, mainly over the field of complex numbers. The description we have given is known (in a somewhat more abstract setting) as the *valuative criterion*, as discussed by Newstead in [16, Preliminaries]. Eisenbud and Harris [5, II.3] also give a somewhat similar description to ours in a more abstract and general situation.

### 3. SLOWER SHIFTING

In this section, we introduce a new family of linear operators and their limits, similar to those given by the third author in [18, Section 4.1], and earlier by Knutson in [15]. In the case where a subspace of the exterior algebra does not admit a basis of monomials, our matrices will preserve more structure in comparison to those of [18], yielding a slower and gentler shifting operation.

Let  $i$  and  $j$  be between 1 and  $n$ . As before, let  $\mathbb{F}(t)$  be the field extension with a new transcendental element  $t$ , and let  $V(t) \cong V \otimes \mathbb{F}(t)$  be the extension of  $V$  to coefficients in  $\mathbb{F}(t)$ .

**Definition 3.1.** Let  $N_{j \rightarrow i}(t)$  be the linear map sending  $e_j \mapsto e_i + te_j$ , and fixing all other basis elements. The limit of this linear map is the *slow shifting* operation.

Also of some interest is the linear map  $M_i(t)$ , sending  $e_i \mapsto te_i$  and fixing all other  $e_j$ . Obviously, both  $N_{j \rightarrow i}(t)$  and  $M_i(t)$  depend on the choice of (ordered) basis  $\mathbf{e}$ . In particular, as matrices acting on the left, we have

$$N_{j \rightarrow i}(t) = \begin{matrix} & & i & & j & & \\ & & & & & & \\ & i & & & & & \\ & & 1 & & 1 & & \\ & & & \ddots & & & \\ & j & & & t & & \\ & & & & & & 1 \end{matrix}, \quad M_i(t) = \begin{matrix} & & & & i & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & t & & \\ & & & & & \ddots & \\ & & & & & & 1 \end{matrix}.$$

As in Section 2.3, we may extend an action on  $V(t)$  to the Grassmannian, and consider the limit action obtained by evaluating the canonical representative at  $t = 0$ . Denote the limit of  $N_{j \rightarrow i}(t)$  as  $N_{j \rightarrow i}$ , and that of  $M_i(t)$  as  $M_i$ .

**3.1. The action of  $N_{j \rightarrow i}$ .** We first describe the action of  $N_{j \rightarrow i}$  on (the projectivization of) elements of  $V$ . As in Notation 1.2, we denote by  $V^{(j)}$  the vector subspace of  $V$  spanned by  $\mathbf{e} \setminus \{e_j\}$ . Let  $v = x + e_j$  be in  $\mathbb{P}V$ , where  $x$  is in  $V^{(j)}$ . Then  $N_{j \rightarrow i}(t)v = x + e_i + te_j$ . This yields as canonical representative

$$N_{j \rightarrow i}(t)v = \begin{cases} x + e_i + te_j & \text{if } e_i + x \neq 0, \\ e_j & \text{otherwise.} \end{cases} \quad \text{Thus, } N_{j \rightarrow i}v = \begin{cases} x + e_i & \text{if } e_i + x \neq 0, \\ e_j & \text{otherwise.} \end{cases}$$

We now extend this action to  $\mathbb{P}(\wedge^k V)$ . Here, if  $m = x + e_j \wedge y$ , where the supports of  $x$  and  $y$  do not contain  $e_j$  in their variable sets, then  $N_{j \rightarrow i}(t)m = x + e_i \wedge y + te_j \wedge y$ . Similarly to the above, this yields as canonical representative

$$(3.1) \quad N_{j \rightarrow i}(t)m = \begin{cases} x + e_i \wedge y + te_j \wedge y & \text{if } e_i \wedge y + x \neq 0, \\ e_j \wedge y & \text{otherwise; and} \end{cases}$$

$$N_{j \rightarrow i}m = \begin{cases} x + e_i \wedge y & \text{if } e_i \wedge y + x \neq 0, \\ e_j \wedge y & \text{otherwise.} \end{cases}$$

We observe that when  $m$  is a monomial with variable set  $F$ , then  $N_{j \rightarrow i}m$  is the monomial with variable set  $\text{shift}_{j \rightarrow i}F$ . More generally, if  $L$  is generated by monomials whose variable sets form the set system  $\mathcal{F}$ , then  $N_{j \rightarrow i}$  is generated by the monomials with variable sets  $\text{shift}_{j \rightarrow i}\mathcal{F}$ . The latter statement follows quickly from Lemma 3.5 below, or a direct proof with bases is also not difficult.

**3.2. Comparison with other operations.** Our operation  $N_{j \rightarrow i}$  slowly changes  $e_j$ 's to  $e_i$ 's in an exterior  $k$ -form, while leaving alone monomials whose variable sets avoid both. In contrast, the shifting operations considered in [15, 18] may quickly replace non-trivial systems with trivial systems. Algebraic shifting [14] or the related techniques with initial monomials in [17] have similar defects.

Indeed, one essential ingredient of our proof is Lemma 5.4 below, which says that if the 1-form  $\ell$  annihilates a subspace of  $k$ -forms after applying  $N_{j \rightarrow i}$ , then the 2-form  $\ell \wedge (e_i - e_j)$  annihilates before shifting. This is an exterior algebra analogue of the fact that if  $\text{shift}_{j \rightarrow i}\mathcal{F}$  becomes trivial, then before shifting, every set in  $\mathcal{F}$  contains at least one of  $i$  or  $j$ . The analogue fails for the operations of [14, 15, 17, 18].

To illustrate, we rephrase the operation of [15, 18] in the language of the current paper: it is the limit  $O_{j \rightarrow i}$  as  $t \rightarrow 0$  of the linear map  $O_{j \rightarrow i}(t)$  sending  $e_j \mapsto e_i + te_j$  and  $e_h \mapsto te_h$  for  $h \neq j$ . We consider the  $k$ -form  $z = e_1 \wedge x + e_2 \wedge y + w$ , where  $x, y$ , and  $w$  are in  $\wedge V^{\{\{1,2\}\}}$ . For  $n$  large enough relative to  $k$ , one can choose  $x, y, w$  so that  $z$  is not annihilated by any 2-form. Now  $N_{2 \rightarrow 1}z$  is  $e_1 \wedge x + e_1 \wedge y + w$ . Depending on the choice of  $w$ , this may not even be annihilated by any 2-form, let alone a 1-form. In contrast, it is straightforward to see that  $O_{2 \rightarrow 1}z$  is  $e_1 \wedge y$ , which is annihilated by  $e_1$ . We see that the analogue of Lemma 5.4 for  $O_{j \rightarrow i}$  does not hold, even for 1-dimensional subspaces!

Similar drawbacks hold for initial monomial techniques, which replace  $z$  with a monomial in a single step.

**3.3. The action of  $M_i$ .** Although it will be of less importance to us, it is not difficult to describe the action of  $M_i$ . If  $v = x + e_i$  is in  $\mathbb{P}V$  for  $x \in V^{(i)}$ , then

$$M_i(t)v = \begin{cases} x + te_i & \text{if } x \neq 0, \\ e_i & \text{otherwise.} \end{cases} \quad \text{Thus, } M_i v = \begin{cases} x & \text{if } x \neq 0, \\ e_i & \text{otherwise.} \end{cases}$$

Extending this action to  $\mathbb{P}(\wedge^k V)$ , we get for  $m = x + e_i \wedge y$  (where the supports of  $x$  and  $y$  do not contain  $e_i$  in their variable sets) the following.

$$M_i(t)m = \begin{cases} x + te_i \wedge y & \text{if } x \neq 0, \\ e_i \wedge y & \text{otherwise.} \end{cases} \quad M_i m = \begin{cases} x & \text{if } x \neq 0, \\ e_i \wedge y & \text{otherwise.} \end{cases}$$

**3.4. Stabilization of slow shifting.** We now show that a subspace  $L$  of  $\wedge^k V$  stabilizes under repeated actions of our slow shifting  $N_{j \rightarrow i}$  operations.

A limit action of  $N_{j \rightarrow i}$  on  $L$  is said to be *fixing* if  $N_{j \rightarrow i} L = L$ ; we will mainly be interested in non-fixing actions. Consider the following procedure.

**Algorithm 3.2** (Slow shifting procedure). *Given  $L$  a subspace of  $\wedge^k V$ , and  $I \subseteq [n]$ :*

*While there exists a pair  $i < j$  chosen from  $I$  so that  $N_{j \rightarrow i} L \neq L$ ,*

*set  $L := N_{j \rightarrow i} L$ .*

*Return  $L$ .*

We will require some lemmas. We consider  $\wedge V^{(j)}$  as a subalgebra of  $\wedge V$ . We say that  $L$  is *monomial with respect to  $e_j$*  if

$$L = \left( L \cap \wedge^k V^{(j)} \right) \oplus \left( L \cap \left( e_j \wedge \wedge^{k-1} V^{(j)} \right) \right).$$

That is,  $L$  is monomial with respect to  $e_j$  if we can find a basis of elements that are either multiples of  $e_j$ , or else do not have any monomials with  $e_j$  in the support. We also call such a basis *monomial with respect to  $e_j$* .

**Lemma 3.3.** *If  $1 \leq i < j \leq n$ , then  $N_{j \rightarrow i} L$  is monomial with respect to  $e_j$ .*

*Proof.* Immediate by (3.1). □

If  $L$  is monomial with respect to  $e_h$ , then the slow shift  $N_{j \rightarrow i}$  distributes over this monomiality for  $h \neq i, j$ , as follows.

**Lemma 3.4.** *Let  $L$  be a subspace of  $\wedge^k V$ , and let  $h \in [n]$ . If  $L$  is monomial with respect to  $e_h$ , and  $1 \leq i < j \leq n$  are distinct from  $h$ , then*

$$\begin{aligned} N_{j \rightarrow i} L &= N_{j \rightarrow i} \left( L \cap \wedge^k V^{(h)} \right) \oplus N_{j \rightarrow i} \left( L \cap \left( e_h \wedge \wedge^{k-1} V^{(h)} \right) \right) \\ &= \left( N_{j \rightarrow i} L \cap \wedge^k V^{(h)} \right) \oplus \left( N_{j \rightarrow i} L \cap \left( e_h \wedge \wedge^{k-1} V^{(h)} \right) \right). \end{aligned}$$

*Proof.* It is immediate by (3.1) that

$$N_{j \rightarrow i} \left( L \cap \wedge^k V^{(h)} \right) \subseteq \wedge^k V^{(h)} \quad \text{and} \quad N_{j \rightarrow i} \left( L \cap \left( e_h \wedge \wedge^{k-1} V^{(h)} \right) \right) \subseteq e_h \wedge \wedge^{k-1} V^{(h)}.$$



The proof now follows by counting dimensions.  $\square$

If  $L$  is monomial with respect to  $e_j$ , then the slow shift  $N_{j \rightarrow i}$  respects the monomiality in a weaker manner.

**Lemma 3.5.** *Let  $L$  be a subspace of  $\Lambda^k V$  and  $j \leq n$ . Let  $K$  be the largest subspace of  $\Lambda^{k-1} V^{(j)}$  such that  $e_j \wedge K \subseteq L$ . If  $L$  is monomial with respect to  $e_j$ , and if  $1 \leq i < j$ , then*

$$N_{j \rightarrow i} L = \left( \left( L \cap \bigwedge^k V^{(j)} \right) + (e_i \wedge K) \right) \oplus (e_j \wedge K'),$$

where  $K'$  is the largest subspace of  $K$  such that  $e_i \wedge K' \subseteq L$ .

*Proof.* It is immediate that  $L \cap \Lambda^k V^{(j)}$  is fixed under  $N_{j \rightarrow i}$ , while from definition

$$N_{j \rightarrow i}(e_j \wedge K) = (e_i \wedge K) \oplus (e_j \wedge \langle x \in K : e_i \wedge x = 0 \rangle).$$

More broadly, for each  $y \in K$  so that  $e_i \wedge y \in L$ , we have  $N_{j \rightarrow i}(e_j \wedge y - e_i \wedge y) = e_j \wedge y$ . The proof now follows by counting dimensions.  $\square$

**Corollary 3.6.** *If  $L$  is monomial with respect to  $e_j$ , then for each  $e_j \wedge x$  in  $N_{j \rightarrow i} L$ , it also holds that  $e_i \wedge x$  is in  $N_{j \rightarrow i} L$ .*

The following will be useful as a base case for an inductive argument that Algorithm 3.2 terminates.

**Corollary 3.7.** *For any  $1 \leq i < j \leq n$  and subspace  $L$  of  $\Lambda^k V$ , we have*

$$N_{j \rightarrow i} N_{j \rightarrow i} N_{j \rightarrow i} L = N_{j \rightarrow i} N_{j \rightarrow i} L.$$

Moreover, if  $L$  is monomial with respect to  $e_j$ , then  $N_{j \rightarrow i} N_{j \rightarrow i} L = N_{j \rightarrow i} L$ .

We caution that there are (non-monomial) examples where  $N_{j \rightarrow i} N_{j \rightarrow i} L \neq N_{j \rightarrow i} L$ . Consider the following:

**Example 3.8.** The span of  $e_1 \wedge e_2 \wedge e_3 - e_1 \wedge e_2 \wedge e_4$  requires two applications of  $N_{4 \rightarrow 3}$  to stabilize. The first application yields the span of  $e_1 \wedge e_2 \wedge e_4$ . Although this is monomial, it is not stable under  $N_{4 \rightarrow 3}$ . Performing  $N_{4 \rightarrow 3}$  again yields  $e_1 \wedge e_2 \wedge e_3$ , which is stable under further slow shifts.

Although it is clear that the shifting algorithm for set systems terminates, the analogue for slow shifting of exterior subspaces requires a slightly more careful analysis.

**Theorem 3.9.** *Algorithm 3.2 terminates, for any choice of a sequence of non-fixing slow shifts.*

*Proof.* We work by induction on the size of the set of permissible indices  $I$  given as input to the algorithm. If  $|I| = 1$ , then the result is trivial, and if  $|I| = 2$ , then the result follows from Corollary 3.7.

If  $|I| > 2$ , then let  $b$  be the greatest index in  $I$ . If there is no  $N_{b \rightarrow i}$  operation in the sequence, then by induction (examining  $I \setminus \{b\}$ ) the algorithm terminates. The first  $N_{b \rightarrow i}$  operation makes  $L$  monomial with respect to  $e_b$ . By Lemma 3.5, each additional non-fixing  $N_{b \rightarrow i}$  operation reduces the dimension of the subspace consisting of multiples of  $e_b$ . By Lemma 3.4, each other slow shift operation preserves the dimension of this

subspace. Thus, there are finitely many non-fixing  $N_{b \rightarrow i}$  operations. The result now follows by induction on  $I \setminus \{b\}$ .  $\square$

As in the set system case [7, Proposition 2.2], it is efficient to shift first from the greatest indices. Here we base our lexicographic order on the usual order on  $[n]$ .

**Proposition 3.10.** *If we choose at each step of Algorithm 3.2 the lexicographically last ordered pair  $(j, i)$  where  $N_{j \rightarrow i}$  is non-fixing, then the algorithm terminates in at most  $|I| - 1 + \binom{|I|}{2}$  iterations.*

*Proof.* Let  $b$  be the last element of  $I$ . We first apply  $N_{b \rightarrow i}$  over all  $i < b$ , in decreasing order, performing the first such shift twice if possible (and so necessary). After the first slow shift, we have a subspace  $L$  that is monomial with respect to  $e_b$ . Additional slow shifts preserve monomiality with respect to  $e_b$  by Lemma 3.5. Moreover, since  $L \cap \wedge^k V^{(b)} \subseteq N_{b \rightarrow i} \left( L \cap \wedge^k V^{(b)} \right)$ , each  $N_{b \rightarrow i}$  is non-fixing for at most one application.

Suppose that we have completed all slow shifts  $N_{b \rightarrow i}$  on  $L$ , and let  $e_b \wedge K$  be as in Lemma 3.5. Repeated application of Lemma 3.5 then gives that  $e_i \wedge K \subseteq L$  for each  $i < b$  in  $I$ , hence that  $\bigoplus_{i \in I} e_i \wedge K \subseteq L$ . Now by another application of Lemma 3.5, the subspace  $\bigoplus_{i \in I} e_i \wedge K$  is preserved under all further  $N_{j \rightarrow i}$  operations with  $i, j \in I$ . In particular,  $N_{b \rightarrow i}$  will fix  $L$  for all  $i$  throughout the remainder of the slow shifting procedure.

That the process terminates now follows by induction on  $I \setminus \{b\}$ . Since in the worst case we shift at each pair  $i < j$ , and one extra time at each element of  $I$  except the least in order to guarantee monomiality, we have at most  $|I| - 1 + \binom{|I|}{2}$  slow shifts.  $\square$

*Remark 3.11.* It is instructive to compare and contrast with [7, Proposition 2.2], which applies a similar sequence of combinatorial shifting operations to sets in the case where  $I = [n]$ . The set situation requires only  $\binom{n}{2}$  operations, since it is not required to first make the system monomial with respect to each  $i$ .

By Theorem 3.9 and Proposition 3.10, given a subspace  $L$  of  $\wedge^k V$ , we may find a subspace of the same dimension that is stable under  $N_{j \rightarrow i}$  over all  $i < j$  both in  $I$ . We now describe the resulting subspaces.

**Proposition 3.12.** *Let  $L$  be a subspace of  $\wedge^k V$ , and let  $I \subseteq [n]$ . If  $L$  is stable under slow shifting  $N_{j \rightarrow i}$  over all  $i < j$  in  $I$ , then for each  $x \in \wedge V^{(I)}$ , the variable sets of the monomials  $y$  in  $\wedge V^{([n] \setminus I)}$  so that  $x \wedge y \in L$  form a shifted set system.*

*Proof.* Let  $x \wedge y$  be in the hypothesis. By definition,  $N_{j \rightarrow i} x \wedge y = x \wedge N_{j \rightarrow i} y$ . As we observed in Section 3.1 that  $N_{j \rightarrow i}$  acts on monomials as combinatorial shifting on their variable sets, the result follows.  $\square$

**Theorem 3.13.** *Let  $L$  be a subspace of  $\wedge^k V$ , let  $I \subseteq [n]$ , and let  $a$  be  $\min I$ . If  $L$  is stable under slow shifting  $N_{j \rightarrow i}$  over all  $i < j$  in  $I$ , then  $L$  has a basis consisting of elements of the form  $x \wedge y$ , where  $x$  is a homogeneous form in  $\wedge V^{(I \setminus a)}$  and  $y$  is a monomial whose variable set is a subset of  $\{e_h : h \in I\}$ .*

*Proof.* Immediate from repeated application of Lemmas 3.3 and 3.4, together with basic facts about direct sums.  $\square$

Thus, Theorem 3.13 says that if  $L$  is stable under  $N_{j \rightarrow i}$  over  $I$ , then  $L$  is simultaneously monomial with respect to all variables with index in  $I$  except for the least indexed. In certain circumstances, we can also get monomiality with respect to the least index of  $I$ .

**Corollary 3.14.** *Let  $L$  be a subspace of  $\bigwedge^k V$ . If  $L$  is stable under slow shifting  $N_{j \rightarrow i}$  over all  $i, j \in [n]$ , then  $L$  has a monomial basis.*

*Proof.* Up to constant multiplication, the only homogeneous forms in  $\bigwedge V^{\{2, \dots, n\}}$  are 1 and  $e_1$ .  $\square$

We remark that if  $L$  is stable under  $N_{j \rightarrow i}$  over all  $i < j$  in  $I$ , but is not monomial with respect to  $a = \min I$ , then it is easy to see that applying  $M_a$  (as in Section 3.3) will result in a system that is monomial with respect to all indices in  $I$ . Indeed, one might see  $M_a$  as a “degenerate” slow shifting operation, in the sense that we send  $e_a$  to zero (as in Section 2.3), but do not have a lesser indexed variable available with which to replace it.

An essential step of our main proof will use shifting over  $I = \{3, \dots, n\}$ .

**Corollary 3.15.** *Let  $L$  be a subspace of  $\bigwedge^k V$  that contains  $e_1 \wedge e_2 \wedge \bigwedge^{k-2} V$  and that is annihilated by  $e_1 \wedge e_2$ . If  $L$  is stable under slow shifting  $N_{j \rightarrow i}$  over all  $i, j \geq 3$ , then  $L$  has a basis consisting of elements that are monomial with respect to all  $e_j$  with  $j \geq 3$ .*

*Proof.* In the basis yielded by Theorem 3.13, the form  $x$  has exterior degree 0, 1, 2, or 3. Without loss of generality, the basis contains the monomials of  $e_1 \wedge e_2 \wedge \bigwedge^{k-2} V$ . It is immediate that for degree 0 or 3, we have the desired. For degree 1, the annihilation condition gives that  $x = \lambda_1 e_1 + \lambda_2 e_2$ . For degree 2, we eliminate  $e_1 \wedge e_2$  using the containment condition, leaving  $x = \lambda_1 e_1 \wedge e_3 + \lambda_2 e_2 \wedge e_3$ .  $\square$

It seems worth noting that, in the case  $I = \{3, \dots, n\}$ , a subspace stable under all  $N_{j \rightarrow i}$ 's decomposes in a pleasing manner.

**Corollary 3.16.** *Let  $L$  be a subspace of  $\bigwedge^k V$  that contains  $e_1 \wedge e_2 \wedge \bigwedge^{k-2} V$  and that is annihilated by  $e_1 \wedge e_2$ . If  $L$  is stable under slow shifting  $N_{j \rightarrow i}$  over all  $i, j \geq 3$ , then there are pairwise linearly independent vectors  $w_1, \dots, w_m$  in the span of  $e_1, e_2$  and subspaces  $K_1, \dots, K_m$  of  $\bigwedge^{k-1} V^{\{1,2\}}$  so that*

$$L = e_1 \wedge e_2 \wedge \bigwedge^{k-2} V + \sum w_i \wedge K_i,$$

and with the following additional properties:

- (1) For every distinct  $i, j \in [m]$ , we have  $K_i \cap K_j = \bigcap_h K_h$ .
- (2) Each  $K_i$  as well as  $\bigcap_h K_h$  has a basis of monomials whose variable sets form a shifted set system.

*Proof.* Take  $K_i$  to be maximal under the condition that  $w_i \wedge K_i \subseteq L$  and apply Corollary 3.15. Suppose  $x \in K_i \cap K_j$ . Then  $w_i \wedge x$  and  $w_j \wedge x$  are in  $L$ . Thus, arbitrary combinations of  $w_i \wedge x$  and  $w_j \wedge x$  are in  $L$ . In particular, as the span of  $w_i$  and  $w_j$  is the same as the span of  $e_1$  and  $e_2$ , it holds that  $w_h \wedge x$  is in  $L$ . Monomiality and shiftedness follows from Corollary 3.15 and Proposition 3.12.  $\square$

#### 4. CROSS-ANNIHILATING SUBSPACES

In this section, we prove Theorem 1.7. With the framework that we have developed, the proof is easy. By Lemma 2.6, slow shifting preserves the cross-annihilating property. Applying the slow shifting procedure over  $I = [n]$ , by Theorem 3.9, Corollary 3.14 and Proposition 3.12 (with  $x = 1$ ), we can reduce to the case where  $K$  and  $L$  have a monomial basis. The variable sets of the monomials in the bases form shifted, nonempty, cross-intersecting set systems. The result now follows by [11, Theorem 2].

#### 5. NONTRIVIAL SELF-ANNIHILATING SUBSPACES

In this section, we prove Theorem 1.5.

We use a similar proof strategy as in the proof of the set system result. We briefly sketch this proof: one reduces to the case where either 1 or 2 is in every set in the set system, then apply a cross-intersecting bound. A difficulty in extending this approach to the exterior algebra case is that an element of  $L$  may have both  $e_1$  and  $e_2$  occurring in its support, but not have  $e_1 \wedge e_2$  as a factor.

**5.1. Main lemma.** The following technical lemma says that Theorem 1.5 holds under the extra conditions of annihilation by a 2-form and partial annihilation by a 1-form factor.

**Lemma 5.1.** *Let  $k \leq n/2$ , and let  $f, g \in V$ . If  $L$  is a nontrivial self-annihilating subspace of  $\Lambda^k V$ , such that  $L$  is annihilated by  $f \wedge g$ , and where some element of  $L$  is annihilated by  $f$  but not by  $g$ , then  $\dim L \leq \binom{n-1}{k-1} - \binom{n-k-1}{k-1} + 1$ .*

*Proof.* Let  $V_0 \subseteq V$  be a complementary subspace to the span of  $\{f, g\}$ . Without loss of generality, it holds that  $A = f \wedge g \wedge \Lambda^{k-2} V$  is contained in  $L$ , as otherwise  $A + L$  satisfies the hypothesis.

We now split  $L$  into the direct sum of three subspaces: the subspace  $A$  as before, the subspace  $B$  of elements that are annihilated by  $f$  but not by  $g$ , and a complementary subspace  $C$ . Moreover,  $B$  consists of elements of the form  $f \wedge x$ , and we can choose  $C$  to consist of elements of the form  $f \wedge x + g \wedge y$ , where  $x, y \in \Lambda^{k-1} V_0$  and  $y \neq 0$ .

Now we take  $B' \subseteq \Lambda^{k-1} V_0$  to be  $\{x : f \wedge x \in B\}$ , and  $C' \subseteq \Lambda^{k-1} V_0$  to be

$$\{y : f \wedge x + g \wedge y \in C \text{ for some } x\}.$$

Since no element of  $C$  is annihilated by  $f$ , we get that  $\dim C' = \dim C$ . Since  $B$  and  $C$  are cross-annihilating, and by the definition of exterior multiplication, we obtain that  $B'$  and  $C'$  are cross-annihilating.

But now by Theorem 1.7, we see that

$$\dim L = \dim A + \dim B' + \dim C' \leq \binom{n-2}{k-2} + \binom{n-2}{k-1} - \binom{n-k-1}{k-1} + 1.$$

The result now follows by the Pascal's triangle identity. □

**5.2. Other lemmas.** We also need several lemmas that parallel results in combinatorial set theory.

The first is immediate by Lemma 2.6, and parallels that if 1 or 2 is in every set  $F \in \mathcal{F}$ , then the same holds after combinatorial shifting.

**Lemma 5.2.** *If  $L \subseteq \wedge^k V$  is a subspace so that  $e_1 \wedge e_2 \wedge L = 0$ , then also  $e_1 \wedge e_2 \wedge (N_{j \rightarrow i} L) = 0$  for every  $3 \leq i < j \leq n$ .*

The second is immediate by computation.

**Lemma 5.3.** *Let  $f, g, \ell$  be 1-forms of  $\wedge V$ , such that  $f$  and  $g$  are linearly independent. If  $L$  is a subspace of  $\wedge^k V$  so that  $f \wedge g \wedge \wedge^{k-2} V \subseteq L$ , and so that  $\ell$  annihilates  $L$ , then  $\ell$  is in the span of  $\{f, g\}$ .*

The third is non-trivial, and requires analysis of the slow shifting operation.

**Lemma 5.4.** *Let  $\ell$  be a 1-form of  $\wedge V$ . If  $L$  is a subspace of  $\wedge^k V$  such that  $\ell \wedge N_{j \rightarrow i} L = 0$ , then  $\ell \wedge (e_i - e_j) \wedge L = 0$ .*

*Moreover, if  $L$  is nontrivial, then  $\ell \wedge (e_i - e_j) \neq 0$ .*

Broadly speaking, Lemma 5.4 is an extension of the fact in combinatorial set theory that if  $\text{shift}_{j \rightarrow i} \mathcal{F}$  becomes trivial, then every set in  $\mathcal{F}$  contains at least one of  $i, j$ .

*Proof (of Lemma 5.4).* By (3.1), whenever  $x + e_j \wedge y$  is in  $L$  (where the variable sets of the supports of  $x$  and  $y$  do not contain  $e_j$ ), we have  $\ell \wedge (x + e_i \wedge y) = 0$ . Thus,

$$\begin{aligned} \ell \wedge (e_i - e_j) \wedge (x + e_j \wedge y) &= \ell \wedge e_i \wedge x - \ell \wedge e_j \wedge (x + e_i \wedge y) \\ &= \ell \wedge e_i \wedge x. \end{aligned}$$

Since  $\ell \wedge (x + e_i \wedge y) = 0$ , we have also  $\ell \wedge e_i \wedge x = \ell \wedge e_i \wedge (x + e_i \wedge y) = 0$ .

Finally, if  $L$  is nontrivial, then there is an  $m = x + e_j \wedge y$  that is not annihilated by  $e_i - e_j$ . Thus,  $e_i \wedge x - e_j \wedge (x + e_i \wedge y) \neq 0$ , and in particular  $x \neq -e_i \wedge y$ . But then (3.1) gives that  $N_{j \rightarrow i} m = x + e_i \wedge y$ . It now follows by computation that  $(e_i - e_j) \wedge m = (e_i - e_j) \wedge N_{j \rightarrow i} m$ , so that the latter is nonzero. This yields that  $\ell$  and  $e_i - e_j$  are linearly independent, as desired.  $\square$

**5.3. Proof of Theorem 1.5.** Let  $L$  be a nontrivial self-annihilating subspace of maximal dimension. As the theorem is trivial for  $k = 1$ , assume that  $k \geq 2$ . Apply slow shifting operations over all  $i, j \in [n]$  and with respect to the standard basis  $\mathbf{e}$  until either  $L$  stabilizes or is annihilated by a 1-form.

If  $L$  stabilizes to a nontrivial self-annihilating subspace under these slow shifting operations, then by Corollary 3.14, the resulting  $L$  is monomial with respect to  $\mathbf{e}$ , and the desired bound follows from (the shifted version of) Hilton-Milner for set systems, Theorem 1.4.

If  $N_{j \rightarrow i} L$  is annihilated by a 1-form  $\ell$ , then we change to the ordered basis  $\mathbf{f}$  with  $f_1 = \ell$ ,  $f_2 = e_j - e_i$ , and  $f_3, \dots, f_n$  chosen arbitrarily to fill out the basis. It follows from Lemma 5.4 that  $f_1$  and  $f_2$  are indeed linearly independent, and that  $L$  is annihilated by  $f_1 \wedge f_2$ . Apply slow shifting operations with respect to the basis  $\mathbf{f}$  for  $i, j \in \{3, \dots, n\}$  until either  $L$  stabilizes or is annihilated by a 1-form. We may assume by maximality

that  $f_1 \wedge f_2 \wedge \wedge^{k-2} V \subseteq L$ , and notice that Lemma 5.2 gives this subspace to be preserved by the slow shifting.

If  $L$  stabilizes, then by Corollary 3.15, we may find a basis consisting of elements of the form  $f_1 \wedge f_2 \wedge x$  and  $(\lambda_1 f_1 + \lambda_2 f_2) \wedge y$ . In particular, the resulting  $L$  satisfies the conditions of Lemma 5.1 for some  $f, g$  in the span of  $\{f_1, f_2\}$ . The desired bound follows.

Finally, if  $N_{j \rightarrow i} L$  is annihilated by a 1-form  $\ell'$  for some  $3 \leq i < j$ , then Lemma 5.4 gives that  $\ell' \wedge (f_j - f_i)$  annihilates  $L$ . Lemma 5.3 gives that  $\ell'$  is in  $\text{span}\{f_1, f_2\}$ ; take  $g$  to be such that  $\ell' \wedge g = f_1 \wedge f_2$ . By maximality,  $\ell' \wedge (f_j - f_i) \wedge \wedge^{k-2} V$  is contained in  $L$ .

Now  $f_1 \wedge f_2 \wedge \wedge^{k-2} V = \ell' \wedge g \wedge \wedge^{k-2} V \subseteq L$ , and (since  $g$  is in the span of  $\{f_1, f_2\}$ ) there are elements of  $\ell' \wedge (f_j - f_i) \wedge \wedge^{k-2} V$  that are annihilated by  $\ell'$  but not by  $g$ . Thus, we may apply Lemma 5.1 to finish the proof.

## REFERENCES

- [1] Michael Artin, *Algebraic geometry—notes on a course*, Graduate Studies in Mathematics, vol. 222, American Mathematical Society, Providence, RI, 2022.
- [2] László Babai and Peter Frankl, *Linear algebra methods in combinatorics*, unpublished manuscript, 1992.
- [3] Jan Draisma, Hanspeter Kraft, and Jochen Kuttler, *Nilpotent subspaces of maximal dimension in semi-simple Lie algebras*, *Compos. Math.* **142** (2006), no. 2, 464–476.
- [4] David S. Dummit and Richard M. Foote, *Abstract algebra*, 3rd ed., John Wiley & Sons, Inc., Hoboken, NJ, 2004.
- [5] David Eisenbud and Joe Harris, *The geometry of schemes*, Graduate Texts in Mathematics, vol. 197, Springer-Verlag, New York, 2000.
- [6] Paul Erdős, Chao Ko, and Richard Rado, *Intersection theorems for systems of finite sets*, *Quart. J. Math. Oxford Ser. (2)* **12** (1961), 313–320.
- [7] Peter Frankl, *The shifting technique in extremal set theory*, *Surveys in combinatorics 1987* (New Cross, 1987), London Math. Soc. Lecture Note Ser., vol. 123, Cambridge Univ. Press, Cambridge, 1987, pp. 81–110.
- [8] ———, *A simple proof of the Hilton-Milner theorem*, *Mosc. J. Comb. Number Theory* **8** (2019), no. 2, 97–101.
- [9] Peter Frankl and Zoltán Füredi, *Nontrivial intersecting families*, *J. Combin. Theory Ser. A* **41** (1986), no. 1, 150–153.
- [10] Joe Harris, *Algebraic geometry: a first course*, Graduate Texts in Mathematics, vol. 133, Springer-Verlag, New York, 1995.
- [11] A. J. W. Hilton and E. C. Milner, *Some intersection theorems for systems of finite sets*, *Quart. J. Math. Oxford Ser. (2)* **18** (1967), 369–384.
- [12] Glenn Hurlbert and Vikram Kamat, *New injective proofs of the Erdős-Ko-Rado and Hilton-Milner theorems*, *Discrete Math.* **341** (2018), no. 6, 1749–1754, arXiv:1609.04714.
- [13] Grigory Ivanov and Seyda Köse, *Erdős-Ko-Rado and Hilton-Milner theorems for two-forms*, *Discrete Math.* **346** (2023), no. 6, Paper No. 113363, 2, arXiv:2201.10892.
- [14] Gil Kalai, *Algebraic shifting*, *Computational commutative algebra and combinatorics* (Osaka, 1999), *Adv. Stud. Pure Math.*, vol. 33, Math. Soc. Japan, Tokyo, 2002, pp. 121–163.
- [15] Allen Knutson, *Schubert calculus and shifting of interval positroid varieties*, unpublished preprint, 2014, arXiv:1408.1261.
- [16] P. E. Newstead, *Introduction to moduli problems and orbit spaces*, *Tata Institute of Fundamental Research Lectures on Mathematics and Physics*, vol. 51, Tata Institute of Fundamental Research, Bombay; Narosa Publishing House, New Delhi, 1978.

- [17] Alex Scott and Elizabeth Wilmer, *Combinatorics in the exterior algebra and the Bollobás two families theorem*, J. Lond. Math. Soc. (2) **104** (2021), no. 4, 1812–1839, arXiv:1907.06019.
- [18] Russ Woodroffe, *An algebraic groups perspective on Erdős-Ko-Rado*, Linear Multilinear Algebra **70** (2022), no. 22, 7825–7835, arXiv:2007.03707.

EINSTEIN INSTITUTE OF MATHEMATICS, HEBREW UNIVERSITY, JERUSALEM 91904, ISRAEL

*Email address:* `Denys.Bulavka@mail.huji.ac.il`

*URL:* <https://kam.mff.cuni.cz/~dbulavka/>

DEPARTMENT OF MATHEMATICS, STATISTICS, AND COMPUTER SCIENCE, ST. OLAF COLLEGE,  
NORTHFIELD MN, USA

*Email address:* `fra.gandi.phd@gmail.com`

*URL:* <https://sites.google.com/a/umich.edu/gandini/>

UNIVERZA NA PRIMORSKEM, GLAGOLJAŠKA 8, 6000 KOPER, SLOVENIA

*Email address:* `russ.woodrooffe@famnit.upr.si`

*URL:* <https://osebje.famnit.upr.si/~russ.woodrooffe/>