

On induced L-infinity action of diffeomorphisms on Cochains

Andrey Losev¹, Dmitrii Sheptunov², and Xin Geng³

¹Shanghai Institute for Mathematics and Interdisciplinary Sciences

²Moscow Institute of Physics and Technology

³University of Science and Technology of China

June 27, 2024

Abstract

One of the approaches to quantum gravity is to formulate it in terms of De Rham algebra, choose a triangulation of space-time, and replace differential forms by cochains (that form a finite dimensional vector space). The key issue of general relativity is the action of diffeomorphisms of space-time on fields. In this paper, we induce the action of diffeomorphisms on cochains by homotopy transfer (or, equivalently, BV integral) that leads to a L_∞ action. We explicitly compute this action for the space-time being an interval, a circle, and a square.

1 Introduction

The quantum theory of gravity is one of the main goals of theoretical physics. The main problem is in ultraviolet divergences that are present due to the infinite dimension of the space of differential forms on the smooth manifold. Therefore, to quantize gravity in the spirit of Feynman integral, we should modify the space of differential forms. In ([1]) we proposed the replacement of the differential forms by A-infinity algebra, that was called Feynman geometry.

The particular class of Feynman geometries may be obtained in the following construction. Consider triangulation of a manifold. Decompose the space of differential forms as complexes as a direct sum of an infinite-dimensional complex of differential forms that have zero integrals against all chains of the triangulation, and a finite dimensional complex of cochains dual to such chains of the triangulation. The latter would be a replacement of the algebra of differential forms. In works of Mnev ([2, 3]) and Sullivan ([4]) it was shown how wedge product on differential forms leads to higher products of cochains in the procedure known as homotopic transfer, which in mathematical physics is just a BV integral against a Lagrangian submanifold determined by the homotopy.

Since we are planning to study gravity the important issue is the general covariance. In particular, we should know how diffeomorphisms are acting on cochains. The hint is that we know that diffeomorphisms are acting on differential forms by Lie derivative. Thus, in this paper, we will apply the BV integral (homotopic transfer) to get the action of vector fields on manifold on cochains. It is not surprising that we will get not the representation of the Lie algebra but L-infinity representation. This construction in the context of transferring the action of supersymmetry to cohomology was earlier done in ([5]).

We believe that explicit formulas for such representation would be an important stepping stone in understanding of quantum gravity in this approach.

The paper is organized as follows. In section 2 we review relation between L_∞ representations and solutions to classical master equations (CME) of the special form. We also recall in this section that BV integral leads to induction of L_∞ representation (also known as homotopic transfer). These results are not new, but we present them for completeness.

In section 3 we proceed to explicit calculation of effective BV-action. To begin with, we consider an interval as a manifold. Further, in sections 4 and 5, explicit formulas for the action in the case of a circle and a square are obtained.

2 BV formalism, L_∞ module structure, BV integral and homotopic transfer

2.1 How to understand L_∞ module

The concept of L_∞ module ([6]) could be easily understood by analogy with the concept of a module over a Lie algebra. The latter may be explained as follows. Take a Lie algebra L , a vector space M (considered as an abelian Lie algebra) and study $L \oplus M$. Now deform the Lie algebra structure on $L \oplus M$ considered as a vector space keeping the structure constants of L . This would lead to structure constants

$$T : L \otimes M \rightarrow M$$

and condition that T gives a structure of the Lie algebra module is equivalent to Jacobi identity for the deformed Lie algebra.

If we apply the same procedure to L_∞ algebra, namely, extend it by a complex M and deform, we will get a set of maps

$$T_p : \bigwedge^p L \rightarrow End(M)$$

subject to quadratic relations that follow from the L_∞ relations.

2.2 BV formalism

A convenient way to write down these relations and to operate with them is provided by BV formalism (see [7]).

In this formalism we study a supermanifold equipped with the odd symplectic structure and a function S_{BV} on it, that is called BV action. S_{BV} should satisfy quantum master equation, but in this work we will restrict ourselves by its classical limit, that we will write explicitly below.

For our purpose it is enough to consider as a supermanifold $T^*[1]W$, where W is a supervector space and symplectic structure is a canonical one. In BV language linear coordinates on W are called fields (and we will denote them as η^a) and dual linear coordinates on the fiber are called antifields and we will denote them as η_a^* .

The classical master equation (CME) takes the form

$$\sum_{a=1}^{dim V} (-1)^{|\eta^a|} \left(\frac{\partial S_{BV}}{\partial \eta^a} \frac{\partial S_{BV}}{\partial \eta_a^*} + \frac{\partial S_{BV}}{\partial \eta_a^*} \frac{\partial S_{BV}}{\partial \eta^a} \right) = 0 \quad (1)$$

where $|\eta^a|$ is parity of η^a .

Using BV formalism is useful because one can reformulate the concept of L_∞ algebra and the concept of L_∞ module as a particular solutions to CME. Moreover, the BV formalism contains the procedure of BV integral allowing to get new solutions of CME from the old ones.

Namely, if $W = W_1 \oplus W_2$, and \mathcal{L} is the Lagrangian submanifold in $T^*[1]W_1$, then the integral

$$\int_{\mathcal{L}} \exp(S_{BV}/\hbar) = \exp(S_{BV}^{ind}(\eta_2, \eta_2^*, \hbar)/\hbar) \quad (2)$$

produces the induced BV action $S_{BV}^{ind}(\hbar)$ that satisfies quantum master equation. For our purpose it is enough to consider the value of S^{ind} at $\hbar = 0$ that gives a solution to CME on the space $T^*[1]W_2$. Note, that

$$S_{BV}^{ind}(\eta_2, \eta_2^*, 0) = \text{extr}_{\mathcal{L}} S_{BV}(\eta, \eta^*) \quad (3)$$

2.3 L_∞ algebra and L_∞ module from the BV point of view

Consider the Lie algebra L with a basis $\{ t_a \}$ and structure constants f_{ab}^e :

$$[t_a, t_b] = f_{ab}^d t_d \quad (4)$$

Consider $L[1]$, it means that $L[1]$ is an odd space with odd coordinates c^a .

Correspondingly, dual coordinates on the fiber $T^*[1](L[1])$ are even and according to our conventions we denote them as c_a^*

It is easy to show that CME for the action

$$S_L = f_{ab}^d c^a c^b c_d^* \quad (5)$$

is equivalent to Jacobi identity for the structure constants f_{ab}^d .

Consider an (even) vector space M with the basis e_i and linear coordinates φ^i . The fiber $T^*[1]M$ has odd dual coordinates φ_i^* . It is equally easy to check that M is a module over Lie algebra iff the map

$$L \rightarrow \text{End}(M), t_a \mapsto T_{aj}^i \quad (6)$$

is such that

$$S_M = c^a T_{aj}^i \varphi_i^* \varphi^j + \frac{1}{2} f_{ab}^d c^a c^b c_d^* \quad (7)$$

solves CME.

Note, that the latter case may be considered as a particular case of the former one for the superalgebra obtained by extension of L by the odd module M .

It is instructive to notice that the L_∞ algebra structure, i.e. the collection of maps

$$\bigwedge L \rightarrow L, \quad t_{a_1} \wedge \dots \wedge t_{a_p} \mapsto f_{a_1 \dots a_p}^b t_b \quad (8)$$

satisfying quadratic relations can be packed into a solution to CME for

$$S_{L_\infty} = \sum_{p=1}^n f_{a_1 \dots a_p}^b c^{a_1} \dots c^{a_p} c_b^* \quad (9)$$

Now we are ready to explain the concept of L_∞ module in BV formalism.

Let us replace a supervector space M by a complex with the differential Q and introduce the matrix Q_i^j as follows:

$$Q(e_i) = Q_i^j e_j, \quad Q^2 = 0 \quad (10)$$

Define a map

$$\bigwedge L \rightarrow \text{End}(M), \quad t_{a_1} \wedge \dots \wedge t_{a_p} \mapsto (T_{a_1 \dots a_p})_i^j \quad (11)$$

Then one can show that this map gives a structure of L_∞ module iff the following action S_{M_∞} solves CME:

$$S_{M_\infty} = \varphi_i^* Q_j^i \varphi^j + \sum_{p=1}^n \frac{1}{p!} c^{a_1} \dots c^{a_p} (T_{a_1 \dots a_p})_j^i \varphi_i^* \varphi^j + \frac{1}{2} f_{ab}^d c^a c^b c_d^* \quad (12)$$

Like in the case of ordinary module over a Lie algebra, the structure of L_∞ module may be considered as a special case of L_∞ algebra on the space $L \oplus M$.

2.4 Induction of the L_∞ module by contraction of the acyclic subcomplex in the BV formalism

Kadeishvili-like theorems ([8]) state that by contraction of the acyclic subcomplex we can pass from the Lie algebra to L_∞ algebra. Here we will describe how in works in the particular case of L modules and L_∞ modules.

Let us apply the BV induction procedure for the Lie algebra acting on a complex M with the action commuting with the differential Q in the complex:

$$[T_a, Q] = 0 \quad (13)$$

Let M be a direct sum of two subcomplexes:

$$M = M_Z \oplus M_A, \quad (14)$$

such that M_A is acyclic, and h is the contracting homotopy:

$$hQ + Qh = Proj_{M_A}, \quad h^2 = 0, \quad hM_Z = 0 \quad (15)$$

Take

$$W_1 = M_A, \quad W_2 = L[1] \oplus M_Z \quad (16)$$

and consider linear Lagrangian submanifold given by equations

$$hw = 0; \quad h^*w^* = 0 \quad (17)$$

where h^* is a conjugated homotopy operator on the dual space.

Since S_M is quadratic polynomial in $w_A \in M_A$ and $w_A^* \in M_A^*[1]$, it is easy to compute its critical value on the Lagrangian submanifold.

It is given by

$$S^{ind} = \langle w_Z^*, Qw_Z \rangle + \frac{1}{2} f_{ab}^d c^a c^b c_d^* + \sum_{k=1}^{\infty} c^{a_1} \dots c^{a_k} \langle w_Z^*, T_{a_1} h \dots h T_{a_k} w_Z \rangle \quad (18)$$

where $w_Z \in M_Z$ and $w_Z^* \in M_Z^*[1]$. If we introduce an universal action operator

$$T_c = c^a T_a$$

the formula above takes the simple form

$$S^{ind} = \langle w_Z^*, Qw_Z \rangle + \frac{1}{2} f_{ab}^d c^a c^b c_d^* + \sum_{k=1}^{\infty} \langle w_Z^*, \underbrace{T_c h \dots h T_c}_{k \text{ times}} w_Z \rangle \quad (19)$$

Thus we have an induced structure of L_∞ module also known as homotopy transfer, see for example ([9]).

2.5 Application to the action of diffeomorphisms on cochains

In our approach to discretized gravity, we take as a module M the space Ω_X of differential forms on a manifold X , as a differential - De Rham operator d on these forms, as a Lie algebra - the algebra of diffeomorphisms on X .

Now we proceed to discretization. We consider a triangulation of X , and take as M_A the subspace Ω_A of differential forms ω having zero integrals against all the chains z of the triangulation:

$$\int_z \omega = 0 \quad (20)$$

As a second summand we take differential forms Ω_Z dual to chains of the triangulation. Such choice is not unique. Whitney considered polynomial forms. However, in this paper we allow more general choices, restricted by condition that

$$d\Omega_Z \in \Omega_Z$$

Applying the construction described above, we get L_∞ module structure on Ω_Z . In the following sections we will study explicit examples.

3 Interval

In this section we consider the manifold $X = [0, 1]$ with the coordinate t , $0 \leq t \leq 1$. The corresponding Lie algebra $L = Vect([0, 1])$ is spanned by

$$v_k = t^k \frac{d}{dt}, k \in \mathbb{Z}, k \geq 0 \quad (21)$$

The structure constants can be written in this basis as

$$[v_k, v_l] = (l - k)t^{k+l-1} \frac{d}{dt} = (l - k)v_{k+l-1} \quad (22)$$

$$f_{kl}^m = (l - k)\delta_{k+l-1}^m \quad (23)$$

We will also use the notation like in (2.4)

$$v_c = \sum_k c^k v_k = \sum_k c^k t^k \frac{d}{dt} = \nu_c(t) \frac{d}{dt} \quad (24)$$

The action of $v_c \in L[1]$ on Ω_X is given by the Lie derivative which we denote \mathcal{L}_{v_c} . So the action (12) takes the form

$$S = \langle \varphi^*, d\varphi \rangle + (-1)^{|\varphi^*|} \langle \varphi^*, \mathcal{L}_{v_c} \varphi \rangle + \frac{1}{2} f_{ab}^d c^a c^b c_d^* \quad (25)$$

We construct the triangulation of $[0, 1]$ as follows. Choose $n + 1$ points $t_0 < t_1 < \dots < t_n$ such that $t_0 = 0, t_n = 1$. Using this we set the space of chains

$$Z = Span\langle \alpha^0 = [t_0], \dots, \alpha^n = [t_n], \beta^0 = [t_0, t_1], \dots, \beta^{n-1} = [t_{n-1}, t_n] \rangle \quad (26)$$

Further consider n functions $\alpha_1, \dots, \alpha_n$ with the condition $\alpha_j(t_i) = \delta_j^i$ and define $\alpha_0 = 1 - \sum_{k=1}^n \alpha_k$. 1-forms are introduced as

$$\beta_j = d\left(\sum_{k=j+1}^n \alpha_k\right), 0 \leq j \leq n-1 \quad (27)$$

Note that:

$$\int_{t_i}^{t_{i+1}} \beta_j = \int_{t_i}^{t_{i+1}} d\left(\sum_{k=j+1}^n \alpha_k\right) = \sum_{k=j+1}^n \alpha_k(t_{i+1}) - \alpha_k(t_i) = \sum_{k=j+1}^n \delta_k^{i+1} - \delta_k^i = \delta_j^i \quad (28)$$

Hence we get $\langle \alpha^i, \alpha_j \rangle = \langle \beta^i, \beta_j \rangle = \delta_j^i$, $\langle \alpha^i, \beta_j \rangle = \langle \beta^i, \alpha_j \rangle = 0$, where \langle, \rangle is the pairing between chains and differential forms given by integration, and the space of forms dual to chains can be written as:

$$\Omega_Z = Span(\alpha_0, \dots, \alpha_n, \beta_0, \dots, \beta_{n-1}) \quad (29)$$

Therefore, fields and antifields are

$$\varphi_Z = \sum_{i=0}^n \alpha_i \varphi^i + \sum_{j=0}^{n-1} \beta_j \psi^j \quad (30)$$

$$\varphi_Z^* = \sum_{i=0}^n \varphi_i^* \alpha^i + \sum_{j=0}^{n-1} \psi_j^* \beta^j \quad (31)$$

To construct the L_∞ module structure on Ω_Z , we need to provide a homotopy h satisfying (15). We write this condition in the form $dh + hd = id - P_Z$, where the projector $P_Z : \Omega_X \rightarrow \Omega_Z$ is defined as

$$P_Z(g(t)) = \sum_j g(t_j) \alpha_j \quad (32)$$

for a 0-form $g(t)$ and

$$P_Z(f(t)dt) = \sum_j \left(\int_{t_j}^{t_{j+1}} f(\tau) d\tau \right) \beta_j \quad (33)$$

for a 1-form $f(t)dt$.

Let us set now

$$h(f(t)dt) = \int_0^t f(\tau) d\tau - \sum_j \left(\int_{t_j}^{t_{j+1}} f(\tau) d\tau \right) \int_0^t \beta_j \quad (34)$$

and check the relation $dh + hd = id - P_Z$ for 0-forms:

$$(dh + hd)g(t) = h(g'(t)dt) = \int_0^t g'(\tau) d\tau - \sum_j \left(\int_{t_j}^{t_{j+1}} g'(\tau) d\tau \right) \int_0^t \beta_j = \quad (35)$$

$$= g(t) - g(0) - \sum_j \left(g(t_{j+1}) - g(t_j) \right) \int_0^t d \left(\sum_{k=j+1}^n \alpha_k \right) = \quad (36)$$

$$= g(t) - g(0) - \sum_j \left(g(t_{j+1}) - g(t_j) \right) \sum_{k=j+1}^n \alpha_k(t) = \quad (37)$$

$$= g(t) - g(0) - \sum_{k=1}^n \alpha_k(t) \sum_{j=0}^{k-1} \left(g(t_{j+1}) - g(t_j) \right) = \quad (38)$$

$$= g(t) - g(0) - \sum_{k=1}^n \alpha_k(t) \left(g(t_k) - g(t_0) \right) = \quad (39)$$

$$= g(t) - g(t_0) \left(1 - \sum_{k=1}^n \alpha_k \right) - \sum_{k=1}^n g(t_k) \alpha_k = (id - P_Z)g(t) \quad (40)$$

and for 1-forms:

$$(dh + hd)f(t)dt = d \left[\int_0^t f(\tau)d\tau - \sum_j \left(\int_{t_j}^{t_{j+1}} f(\tau)d\tau \right) \int_0^t \beta_j \right] = \quad (41)$$

$$= f(t)dt - \sum_j \left(\int_{t_j}^{t_{j+1}} f(\tau)d\tau \right) \beta_j = (id - P_Z)f(t)dt \quad (42)$$

Finally, the induced action (18) becomes

$$S^{ind} = \langle \varphi_Z^*, d\varphi_Z \rangle + (-1)^{|\varphi_Z^*|} \langle \varphi_Z^*, \mathcal{L}_{v_c} \varphi_Z \rangle + \langle \varphi_Z^*, \mathcal{L}_{v_c} h \mathcal{L}_{v_c} \varphi_Z \rangle + \frac{1}{2} f_{ab}^d c^a c^b c_d^* \quad (43)$$

All the terms containing more than one operator h vanish since h lowers form's degree by 1.

To compute the induced action we introduce notations

$$\beta_j^i = \sum_{k=j+1}^n \alpha'_k(t_i) \quad (44)$$

$$\alpha^j \nu_c = \nu_c(t_j) = \nu_c^j \quad (45)$$

$$\alpha^j \nu'_c = \nu'_c(t_j) = \nu_c'^j \quad (46)$$

What remains is to calculate the distinct components of the expression for the induced action

$$\langle \beta^j, d\alpha_i \rangle = \int_{t_j}^{t_{j+1}} d\alpha_i = \alpha_i(t_{j+1}) - \alpha_i(t_j) = \delta_i^{j+1} - \delta_i^j \quad (47)$$

$$\langle \alpha^i, \mathcal{L}_{v_c} \alpha_j \rangle = \alpha^i \nu_c(t) \frac{d\alpha_j}{dt} = \nu_c(t_i) \alpha'_j(t_i) = \nu_c^i (\beta_{j-1}^i - \beta_j^i) \quad (48)$$

$$\begin{aligned} \langle \beta^i, \mathcal{L}_{v_c} \beta_j \rangle &= \beta^i d \left(\nu_c(t) \sum_{k=j+1}^n \alpha'_k \right) = \int_{t_i}^{t_{i+1}} d \left(\nu_c(t) \sum_{k=j+1}^n \alpha'_k \right) \\ &= \sum_{k=j+1}^n \nu_c(t_{i+1}) \alpha'_k(t_{i+1}) - \nu_c(t_i) \alpha'_k(t_i) = \nu_c^{i+1} \beta_j^{i+1} - \nu_c^i \beta_j^i \end{aligned} \quad (49)$$

$$\begin{aligned} \langle \alpha^i, \mathcal{L}_{v_c} h \mathcal{L}_{v_c} \beta_j \rangle &= \alpha^i \mathcal{L}_{v_c} h d \left(\nu_c(t) \sum_{k=j+1}^n \alpha'_k \right) = \alpha^i \mathcal{L}_{v_c} (id - P_Z) \left(\nu_c(t) \sum_{k=j+1}^n \alpha'_k \right) = \\ &= \alpha^i \mathcal{L}_{v_c} \left(\nu_c(t) \sum_{k=j+1}^n \alpha'_k - \sum_m \nu_c^m \beta_j^m \alpha_m \right) = \\ &= \nu_c^i \nu_c'^i \beta_j^i - \nu_c^i \sum_m \nu_c^m \beta_j^m (\beta_{m-1}^i - \beta_m^i) \end{aligned} \quad (50)$$

Thus, the resulting L_∞ module structure is

$$\begin{aligned} S^{ind} = & \psi_i^*(\delta_j^{i+1} - \delta_j^i)\varphi^j - \varphi_i^*\nu_c^i(\beta_{j-1}^i - \beta_j^i)\varphi^j + \psi_i^*(\nu_c^{i+1}\beta_j^{i+1} - \nu_c^i\beta_j^i)\psi^j + \\ & + \varphi_i^*\left(\nu_c^i\nu_c^i\beta_j^i - \nu_c^i\sum_m \nu_c^m\beta_j^m(\beta_{m-1}^i - \beta_m^i)\right)\psi^j + \frac{1}{2}f_{ab}^dc^ac^bc_d^* \end{aligned} \quad (51)$$

4 Circle

In this section we consider the manifold $X = S^1$ with the coordinate $0 \leq t < 2\pi$ and the Lie algebra $L = Vect(S^1)$. Let

$$v_k = e^{ikt} \frac{d}{dt}, k \in \mathbb{Z} \quad (52)$$

be a basis in $L = Vect(S^1)$. Then, the corresponding structure constants are

$$[v_k, v_l] = i(l-k)e^{i(m+n)t} \frac{d}{dt} = i(l-k)v_{k+l} \quad (53)$$

$$f_{kl}^m = i(l-k)\delta_{k+l}^m \quad (54)$$

Let us use the following notation for the subsequent computations

$$v_c = \sum_k c^k v_k = \sum_k c^k e^{ikt} \frac{d}{dt} = \nu_c(t) \frac{d}{dt} \quad (55)$$

The action (12) takes the form analogous to the interval case

$$S = \langle \varphi^*, d\varphi \rangle + (-1)^{|\varphi^*|} \langle \varphi^*, \mathcal{L}_{v_c}\varphi \rangle + \frac{1}{2}f_{ab}^dc^ac^bc_d^* \quad (56)$$

The triangulation of S^1 is constructed as follows. Choose n points $t_0 = 0, \dots, t_{n-1}$ and 1-forms $\beta_0, \dots, \beta_{n-1}$ such that $\int_{t_i}^{t_{i+1}} \beta_j = \delta_j^i$. Denote $\beta_j^i := \vartheta_j(t_i)$, where $\beta_j = \vartheta_j(t)dt$. Define the 0-forms:

$$\alpha_0(t) = 1 + \int_0^t (\beta_{n-1} - \beta_0) \quad (57)$$

$$\alpha_j(t) = \int_0^t (\beta_{j-1} - \beta_j), j \neq 0 \quad (58)$$

One can check that $\alpha_j(t_i) = \delta_j^i$.

Then, the spaces of chains and cochains are

$$Z = \text{Span}\langle \alpha^0 = [t_0], \dots, \alpha^{n-1} = [t_{n-1}], \beta^0 = [t_0, t_1], \dots, \beta^{n-1} = [t_{n-1}, t_0] \rangle \quad (59)$$

$$\Omega_Z^\bullet = \text{Span}\langle \alpha_0, \dots, \alpha_{n-1}, \beta_0, \dots, \beta_{n-1} \rangle \quad (60)$$

The corresponding fields and antifields are

$$\varphi_Z^* = \sum_{i=0}^{n-1} \varphi_i^* \alpha^i + \sum_{j=0}^{n-1} \psi_j^* \beta^j \quad (61)$$

$$\varphi_Z = \sum_{i=0}^{n-1} \alpha_i \varphi^i + \sum_{j=0}^{n-1} \beta_j \psi^j \quad (62)$$

As mentioned earlier in section 3, it is necessary to provide h satisfying the condition $dh + hd = id - P_Z$, where the projector $P_Z : \Omega_X \rightarrow \Omega_Z$ is defined in this case as

$$P_Z(g(t)) = \sum_j g(t_j) \alpha_j \quad (63)$$

for a 0-form $g(t)$ and

$$P_Z(f(t)dt) = \sum_j \left(\int_{t_j}^{t_{j+1}} f(\tau) d\tau \right) \beta_j \quad (64)$$

for a 1-form $f(t)dt$.

Let us set the homotopy h

$$h(f(t)dt) = \int_0^t f(\tau) d\tau - \sum_j \left(\int_{t_j}^{t_{j+1}} f(\tau) d\tau \right) \int_0^t \beta_j \quad (65)$$

and check the the relation $dh + hd = id - P_Z$ for 1-forms:

$$\begin{aligned} (dh + hd)(f(t)dt) &= d \left[\int_0^t f(\tau) d\tau - \sum_j \left(\int_{t_j}^{t_{j+1}} f(\tau) d\tau \right) \int_0^t \beta_j \right] = \\ &= f(t)dt - \sum_j \left(\int_{t_j}^{t_{j+1}} f(\tau) d\tau \right) \beta_j = (id - P_Z)f(t)dt \end{aligned} \quad (66)$$

and for 0-forms:

$$\begin{aligned}
(dh + hd)(g(t)) &= \int_0^t g'(\tau) d\tau - \sum_j \left(\int_{t_j}^{t_{j+1}} g'(\tau) d\tau \right) \int_0^t \beta_j = \\
&= g(t) - g(0) - \sum_j (g(t_{j+1}) - g(t_j)) \int_0^t \beta_j = \\
&= g(t) - g(0) - g(0) \left(\int_0^t (\beta_{n-1} - \beta_0) \right) - \\
&\quad - \sum_{j \neq 0} g(t_j) \left(\int_0^t (\beta_{j-1} - \beta_j) \right) = \\
&= g(t) - \sum_j g(t_j) \alpha_j = (id - P_Z)g(t)
\end{aligned} \tag{67}$$

The induced action (18) is analogous to the case of the interval

$$S^{ind} = \langle \varphi_Z^*, d\varphi_Z \rangle + (-1)^{|\varphi_Z^*|} \langle \varphi_Z^*, \mathcal{L}_{v_c} \varphi_Z \rangle + \langle \varphi_Z^*, \mathcal{L}_{v_c} h \mathcal{L}_{v_c} \varphi_Z \rangle + \frac{1}{2} f_{ab}^d c^a c^b c_d^* \tag{68}$$

We introduce notations

$$\nu_c(t_j) = \nu_c^j \tag{69}$$

$$\nu_c'(t_j) = \nu_c'^j \tag{70}$$

Finally, let us compute the action

$$d\alpha_j = \beta_{j-1} - \beta_j \tag{71}$$

$$\langle \beta^i, d\alpha_j \rangle = \delta_{j-1}^i - \delta_j^i \tag{72}$$

$$\langle \alpha^i, \mathcal{L}_{v_c} \alpha_j \rangle = \nu_c^i (\beta_{j-1}^i - \beta_j^i) \tag{73}$$

$$\langle \beta^i, \mathcal{L}_{v_c} \beta_j \rangle = \int_{t_i}^{t_{i+1}} d(\nu_c(t) \vartheta_j(t)) = \nu_c^{i+1} \beta_j^{i+1} - \nu_c^i \beta_j^i \tag{74}$$

$$\begin{aligned}
h\mathcal{L}_{v_c} \beta_j &= \int_0^t d(\nu_c \vartheta_j) - \sum_m \left(\int_{t_m}^{t_{m+1}} d(\nu_c \vartheta_j) \right) \int_0^t \beta_m = \\
&= \nu_c(t) \vartheta_j(t) - \nu_c^0 \beta_j^0 - \sum_m \left(\nu_c^{m+1} \beta_j^{m+1} - \nu_c^m \beta_j^m \right) \int_0^t \beta_m = \\
&= \nu_c(t) \vartheta_j(t) - \sum_m \nu_c^m \beta_j^m \alpha_m(t)
\end{aligned} \tag{75}$$

$$\begin{aligned}
\mathcal{L}_{v_c} h \mathcal{L}_{v_c} \beta_j &= \nu_c(t) \frac{\partial}{\partial t} \left(\nu_c(t) \vartheta_j(t) - \sum_m \nu_c^m \beta_j^m \alpha_m(t) \right) = \\
&= \nu_c(t) \nu_c'(t) \vartheta_j(t) + \nu_c(t) \nu_c(t) \vartheta_j'(t) - \\
&\quad - \nu_c(t) \sum_m \nu_c^m \beta_j^m (\vartheta_{m-1}(t) - \vartheta_m(t))
\end{aligned} \tag{76}$$

$$\langle \alpha^i, \mathcal{L}_{v_c} h \mathcal{L}_{v_c} \beta_j \rangle = \nu_c^i \nu_c^i \beta_j^i - \nu_c^i \sum_m \nu_c^m \beta_j^m (\beta_{m-1}^i - \beta_m^i) \tag{77}$$

Thus, we get the L_∞ module structure

$$\begin{aligned}
S^{ind} &= \psi_i^* (\delta_{j-1}^i - \delta_j^i) \varphi^j - \varphi_i^* \nu_c^i (\beta_{j-1}^i - \beta_j^i) \varphi^j + \psi_i^* (\nu_c^{i+1} \beta_j^{i+1} - \nu_c^i \beta_j^i) \psi^j + \\
&\quad + \varphi_i^* \left(\nu_c^i \nu_c^i \beta_j^i - \nu_c^i \sum_m \nu_c^m \beta_j^m (\beta_{m-1}^i - \beta_m^i) \right) \psi^j + \frac{1}{2} f_{ab}^d c^a c^b c_d^*
\end{aligned} \tag{78}$$

5 Square

In this section we consider the manifold $X = [0, 1] \times [0, 1]$ with the coordinates x, y . The vector field v_c takes the form

$$v_c(x, y) = \nu_c^x(x, y) \frac{\partial}{\partial x} + \nu_c^y(x, y) \frac{\partial}{\partial y}, \tag{79}$$

$$\nu_c^x = \sum_{n,m} c^{x,nm} x^n y^m, \tag{80}$$

$$\nu_c^y = \sum_{n,m} c^{y,nm} x^n y^m \tag{81}$$

The triangulation is given by

$$\gamma^* = [0, 1] \times [0, 1] \tag{82}$$

$$\beta^0 = [(0, 0), (1, 0)], \beta^1 = [(1, 0), (1, 1)], \beta^2 = [(1, 1), (0, 1)], \beta^3 = [(0, 1), (0, 0)] \tag{83}$$

$$\alpha^0 = (0, 0), \alpha^1 = (1, 0), \alpha^2 = (1, 1), \alpha^3 = (0, 1) \tag{84}$$

$$Z = \text{Span}(\alpha^0, \dots, \alpha^3, \beta^0, \dots, \beta^3, \gamma^*) \tag{85}$$

Let us define the constructions that will be used in further calculations:
2-form $\gamma = dx dy$; 1-forms:

$$\beta_0 = \left(\int_y^1 d\tilde{y} \right) dx = (1 - y) dx \tag{86}$$

$$\beta_1 = \left(\int_0^x d\tilde{x} \right) dy = x dy \tag{87}$$

$$\beta_2 = -\left(\int_0^y d\tilde{y}\right)dx = -ydx \quad (88)$$

$$\beta_3 = -\left(\int_x^1 d\tilde{x}\right)dy = -(1-x)dy; \quad (89)$$

and functions:

$$\alpha_0 = 1 + \int (\beta_3 - \beta_0) = (1-x)(1-y) \quad (90)$$

$$\alpha_1 = \int (\beta_0 - \beta_1) = x(1-y) \quad (91)$$

$$\alpha_2 = \int (\beta_1 - \beta_2) = xy \quad (92)$$

$$\alpha_3 = \int (\beta_2 - \beta_3) = y(1-x). \quad (93)$$

Since $d\beta_j = \gamma$, forms $\beta_i - \beta_j$ are closed, and the functions α_i are defined correctly. Note that $\langle \gamma^*, \gamma \rangle = 1$, $\langle \beta^i, \beta_j \rangle = \delta_j^i$, $\langle \alpha^i, \alpha_j \rangle = \delta_j^i$. Then the space dual to the space of chains is

$$\Omega_Z = \text{Span}(\alpha_0, \dots, \alpha^3, \beta_0, \dots, \beta^3, \gamma) \quad (94)$$

The fields under consideration are

$$\varphi_Z = \sum_{i=0}^3 \alpha_i \varphi^i + \sum_{j=0}^3 \beta_j \psi^j + \gamma \omega \quad (95)$$

$$\varphi_Z^* = \sum_{i=0}^3 \varphi_i^* \alpha^i + \sum_{j=0}^3 \psi_j^* \beta^j + \omega^* \gamma^* \quad (96)$$

We introduce the projector P_Z (g , f and e are 0, 1 and 2-forms respectively) analogously to the previous sections

$$P_Z(g) = \sum_j \langle \alpha^j, g \rangle \alpha_j \quad (97)$$

$$P_Z(f) = \sum_j \langle \beta^j, f \rangle \beta_j \quad (98)$$

$$P_Z(e) = \langle \gamma^*, e \rangle \gamma \quad (99)$$

To define the homotopy consider the operator $I : \Omega_X \rightarrow \Omega_X$; for 2-forms:

$$I(e(x, y)dx dy) = \frac{1}{2} \left(\int_0^x e(\tilde{x}, y) d\tilde{x} \right) dy - \frac{1}{2} \left(\int_0^y e(x, \tilde{y}) d\tilde{y} \right) dx \quad (100)$$

for 1-forms:

$$I(f_x(x, y)dx + f_y(x, y)dy) = \frac{1}{2} \left(\int_0^x f_x(\tilde{x}, 0)d\tilde{x} + \int_0^y f_y(x, \tilde{y})d\tilde{y} + \int_0^y f_y(0, \tilde{y})d\tilde{y} + \int_0^x f_x(\tilde{x}, y)d\tilde{x} \right) \quad (101)$$

The last expression is simply the sum of the integrals of an 1-form f along the edges of the rectangle with vertices $(0, 0), (0, x), (0, y), (x, y)$. For 0-forms $g(t)$ $I(g) = 0$. Using I , we can now define the homotopy (e and f are 2- and 1- forms respectively):

$$h(e) = I(e) - \sum_i \langle \beta^i, I(e) \rangle \beta_i \quad (102)$$

$$h(f) = I\left(f - \sum_i \langle \beta^i, f \rangle \beta_i\right) \quad (103)$$

Let us check the condition (15) in the form $dh + hd = id - P_Z$:

$$\begin{aligned} (dh + hd)g(x, y) &= h(dg) = I\left(dg - \sum_i \langle \beta^i, dg \rangle \beta_i\right) = \\ &= g(x, y) - g(0, 0) - I\left(\sum_i (\langle \alpha^{i+1}, g \rangle - \langle \alpha^i, g \rangle) \beta_i\right) = \\ &= g(x, y) - g(0, 0) - \sum_i \langle \alpha^i, g \rangle I(\beta_{i-1} - \beta_i) = \\ &= g(x, y) - \sum_i \langle \alpha^i, g \rangle \alpha_i = (id - P_Z)g \end{aligned} \quad (104)$$

$$\begin{aligned} (dh + hd)e &= d(h(e)) = d\left(I(e) - \sum_i \langle \beta^i, I(e) \rangle \beta_i\right) = \\ &= e - \left(\sum_i \langle \beta^i, I(e) \rangle\right) \gamma = e - \langle \gamma^*, e \rangle \gamma = (id - P_Z)e \end{aligned} \quad (105)$$

$$hdf = hd(f_x dx + f_y dy) = h(df) = I(df) - \sum_i \langle \beta^i, I(df) \rangle \beta_i \quad (106)$$

$$\begin{aligned} 2I(df) &= \left(f_x(x, y) - f_x(x, 0) - \int_0^y \frac{\partial f_y}{\partial x} d\tilde{y}\right) dx + \\ &+ \left(f_y(x, y) - f_y(0, y) - \int_0^x \frac{\partial f_x}{\partial y} d\tilde{x}\right) dy \end{aligned} \quad (107)$$

$$\sum_i \langle \beta^i, I(df) \rangle \beta_i = \left(\sum_i \langle \beta^i, f \rangle\right) \frac{\beta_1 + \beta_2}{2} = \langle \gamma^*, df \rangle I(\gamma) \quad (108)$$

$$2dI(f) = f_x(x, y)dx + f_y(x, y)dy + f_x(x, 0)dx + f_y(0, y)dy + \left(\int_0^y \frac{\partial f_y}{\partial x} d\tilde{y} \right) dx + \left(\int_0^x \frac{\partial f_x}{\partial y} d\tilde{x} \right) dy \quad (109)$$

$$I(df) + dI(f) = f \quad (110)$$

$$\begin{aligned} (hd + dh)f &= I(df) + dI(f) - \sum_i \langle \beta^i, I(df) \rangle \beta_i - \sum_i \langle \beta^i, f \rangle dI(\beta_i) = \\ &= f - \sum_i \langle \beta^i, f \rangle \beta_i - \langle \gamma^*, df \rangle I(\gamma) + \sum_i \langle \beta^i, f \rangle I(d\beta_i) = \\ &= (1 - P_Z)f + \sum_i \langle \beta^i, f \rangle I(\gamma) - \langle \gamma^*, df \rangle I(\gamma) = (1 - P_Z)f \end{aligned} \quad (111)$$

Let us compute the components of action

$$d\alpha_j = \beta_{j-1} - \beta_j \quad (112)$$

$$\langle \beta^i, d\alpha_j \rangle = \delta_{j-1}^i - \delta_j^i \quad (113)$$

$$d\beta_j = \gamma \quad (114)$$

$$\langle \gamma^*, d\beta_j \rangle = 1 \quad (115)$$

$$\begin{aligned} \langle \alpha^i, \mathcal{L}_{v_c} \alpha_j \rangle &= \langle \alpha^i, \nu_c^x \frac{\partial}{\partial x} \int (\beta_{j-1} - \beta_j) + \nu_c^y \frac{\partial}{\partial y} \int (\beta_{j-1} - \beta_j) \rangle \\ &= \nu_i (\beta_{j-1}^i - \beta_j^i) \end{aligned} \quad (116)$$

$$\begin{aligned} \langle \beta^i, \mathcal{L}_{v_c} \beta_j \rangle &= \langle \beta^i, d(v_c \beta_j) + i_{v_c} d\beta_j \rangle = \nu_{i+1} \beta_j^{i+1} - \nu_i \beta_j^i + \\ &+ \int_{\beta^i} \nu_c^x dy - \nu_c^y dx \end{aligned} \quad (117)$$

$$\langle \gamma^*, \mathcal{L}_{v_c} \gamma \rangle = \langle \gamma^*, d(\nu_c^x dy - \nu_c^y dx) \rangle = \int_{[0,1]^2} \mathbf{div}(v_c) dx dy \quad (118)$$

$$\begin{aligned}
\langle \alpha^i, \mathcal{L}_{v_c} h \mathcal{L}_{v_c} \beta_j \rangle &= \langle \alpha^i, \mathcal{L}_{v_c} h(d(v_c \beta_j) + i_{v_c} d\beta_j) \rangle = \langle \alpha^i, \mathcal{L}_{v_c} (hd + dh)(v_c \beta_j) \rangle + \\
&+ \langle \alpha^i, \mathcal{L}_{v_c} h(\nu_c^x dy - \nu_c^y dx) \rangle = \langle \alpha^i, \mathcal{L}_{v_c} (v_c \beta_j - \sum_m \nu_m \beta_j^m \alpha_m) \rangle + \\
&+ \langle \alpha^i, \mathcal{L}_{v_c} h(\nu_c^x dy - \nu_c^y dx) \rangle \\
&= \nu_{x,i} \frac{\partial(\nu_i \beta_j)}{\partial x} + \nu_{y,i} \frac{\partial(\nu_i \beta_j)}{\partial y} - \sum_m \nu_i (\beta_{m-1}^i - \beta_m^i) \nu_m \beta_j^m + \\
&+ \frac{1}{2} \langle \alpha^i, \nu_c^x(x, y) \nu_c^x(x, 0) - \nu_c^x \int_0^y \frac{\partial \nu_c^y}{\partial x}(x, \tilde{y}) dy \rangle + \\
&+ \frac{1}{2} \langle \alpha^i, -\nu_c^y(x, y) \nu_c^y(0, y) + \nu_c^y \int_0^y \frac{\partial \nu_c^x}{\partial x}(\tilde{x}, y) dx \rangle - \\
&- \langle \alpha^i, \sum_k \left(\nu_c^x \frac{\partial I(\beta_k)}{\partial x} + \nu_c^y \frac{\partial I(\beta_k)}{\partial y} \right) \int_{\beta^k} \nu_c^x dy - \nu_c^y dx \rangle
\end{aligned} \tag{119}$$

Denote

$$\begin{aligned}
\xi := \mathcal{L}_{v_c} h \mathcal{L}_{v_c} \gamma &= \mathcal{L}_{v_c} h d(\nu_c^x dy - \nu_c^y dx) = \mathcal{L}_{v_c} h(\mathbf{div}(v_c) dx dy) \\
&= \frac{1}{2} \left(d \left(\nu_c^y (\nu_c^x(x, y) - \nu_c^x(0, y)) + \int_0^x \frac{\partial \nu_c^y}{\partial y} d\tilde{x} \right) - \right. \\
&- \nu_c^x (\nu_c^y(x, y) - \nu_c^y(x, 0)) + \left. \int_0^y \frac{\partial \nu_c^x}{\partial x} dy \right) + \mathbf{div}(v_c) (\nu_c^x dy - \nu_c^y dx) - \\
&- \left(\int_{[0,1]^2} \mathbf{div}(v_c) dx dy \right) \left(d(v_c(\beta_1 + \beta_2)) + 2(\nu_c^x dy - \nu_c^y dx) \right)
\end{aligned} \tag{120}$$

The most complicated term (cubic in c^a) has the form

$$\langle \alpha^i, \mathcal{L}_{v_c} h \mathcal{L}_{v_c} h \mathcal{L}_{v_c} \gamma \rangle = \langle \alpha^i, \mathcal{L}_{v_c} h \xi \rangle = \langle \alpha^i, \mathcal{L}_{v_c} \left(I(\xi) - \sum_i I(\beta_i) \int_{\beta^i} \xi \right) \rangle \tag{121}$$

Thus, from (112)-(121) we get the L_∞ module structure

$$\begin{aligned}
S^{ind} &= \psi_i^* \langle \beta^i, d\alpha_j \rangle \varphi^j + \omega^* \langle \gamma^*, d\beta_j \rangle \psi^j - \varphi_i^* \langle \alpha^i, \mathcal{L}_{v_c} \alpha_j \rangle \varphi^j + \psi_i^* \langle \beta^i, \mathcal{L}_{v_c} \beta_j \rangle \psi^j - \\
&- \omega^* \langle \gamma^*, \mathcal{L}_{v_c} \gamma \rangle \omega + \varphi_i^* \langle \alpha^i, \mathcal{L}_{v_c} h \mathcal{L}_{v_c} \beta_j \rangle \psi^j + \psi_i^* \langle \beta^i, \mathcal{L}_{v_c} h \mathcal{L}_{v_c} \gamma \rangle \omega + \\
&+ \varphi_i^* \langle \alpha^i, \mathcal{L}_{v_c} h \mathcal{L}_{v_c} h \mathcal{L}_{v_c} \gamma \rangle \omega + \frac{1}{2} f_{ab}^d c^a c^b c_d^*
\end{aligned} \tag{122}$$

References

- [1] Sen Hu and Andrey Losev. “Feynman geometries”. In: *60 Years of Yang–Mills Gauge Field Theories: CN Yang’s Contributions to Physics*. World Scientific, 2016, pp. 453–471.
- [2] Pavel Nikolaevich Mnev. “On the simplicial super-BF model”. In: *Zapiski Nauchnykh Seminarov POMI* 331 (2006), pp. 84–90.
- [3] Pavel Mnev. “Discrete BF theory”. In: *arXiv preprint arXiv:0809.1160* (2008).
- [4] Ruth Lawrence and Dennis Sullivan. “A free differential Lie algebra for the interval”. In: *arXiv preprint math.AT/0610949* (2006).
- [5] Victor Alexandrov et al. “On pure spinor superfield formalism”. In: *Journal of High Energy Physics* 2007.10 (2007), p. 074.
- [6] Maxim Kontsevich. “Deformation quantization of Poisson manifolds”. In: *Letters in Mathematical Physics* 66 (2003), pp. 157–216.
- [7] Albert Schwarz. “Semiclassical approximation in Batalin-Vilkovisky formalism”. In: *Communications in Mathematical Physics* 158 (1993), pp. 373–396.
- [8] Tornike V Kadeishvili. “On the theory of homology of fiber spaces”. In: *Uspekhi Mat. Nauk* 35.3 (1980), p. 213.
- [9] Alex S Arvanitakis et al. “Homotopy Transfer and Effective Field Theory I: Tree-level”. In: *Fortschritte der Physik* 70.2-3 (2022), p. 2200003.