Dynamics of multidimensional fundamental and vortex solitons in random media

Volodymyr M. Lashkin*

¹Institute for Nuclear Research, Pr. Nauki 47, Kyiv 03028, Ukraine and ²Space Research Institute, Pr. Glushkova 40 k.4/1, Kyiv 03187, Ukraine

We study the dynamics of fundamental and vortex solitons in the framework of the nonlinear Schrödinger equation with the spatial dimension $D \ge 2$ with a multiplicative random term depending on the time and space coordinates. To this end, we develop a new technique for calculating the even moments of the Nth order. The proposed formalism does not use closure procedures for the nonlinear term, as well as the smallness of the random term and the use of perturbation theory. The essential point is the quadratic form of the autocorrelation function of the random field and the special stochastic change of variables. Using variational analysis to determine the field of structures in the deterministic case, we analytically calculate a number of statistical characteristics describing the dynamics of fundamental and vortex solitons in random medium, such as the mean intensities, the variance of the intensity, the centroid and spread of the structures, the spatial mutual coherence function etc. In particular, we show that, under the irreversible action of fluctuations, the solitons spread out, i.e., no collapse occurs.

I. INTRODUCTION

The propagation of nonlinear waves in random media is an important problem in nonlinear physics and has been intensively studied over the past few decades, i. e. almost immediately with the widespread use of the concept of soliton in physics [1–4]. This is due both to general theoretical interest, where the problems of fundamental nonlinear (primarily soliton) physics and statistical physics are intertwined, but also from a practical point of view of applied problems of optics, acoustics, plasma physics, physics of disordered systems, etc., where fluctuating parameters and randomness appears in a natural way. Depending on the problems under study, the source of fluctuations can be the refractive index of the medium, which changes randomly and continuously in time and space, the presence of turbulence in laboratory and space plasmas and in the ionosphere and troposphere, particles randomly distributed in space (e. g. rain, fog, hail), random interaction potential in disordered systems, etc. [5-7]. Note that, as is well known, the wave equation in a random medium even in the linear case cannot be solved explicitly in the sense that it is impossible to determine the probability distribution of the wave field as a function of the probability distribution of the medium [5, 6]. This applies even more to the propagation of nonlinear waves in random media. The fundamental difficulty faced by any statistical description of nonlinear waves is the statistical closure problem, in which the time evolution of the moments of order n for the wave field is coupled by the nonlinearity to moments of order n + 1. To obtain closed nonlinear equations for statistical characteristics (for example, moments of the *n*th order), various approximate statistical closure procedures are used. This provides a way of expressing a moment of some order in terms of lower-order moments. Nevertheless, a fairly

extensive literature is devoted to nonlinear waves in random media. In the vast majority of cases, this concerns the one-dimensional (1D) case that is soliton propagation in random media. In particular, the propagation of solitons in the presence of randomness in models of the nonlinear Schrödinger (NLS) equation, the Kortewegde Vries (KdV) equation and the sine-Gordon equation, as well as shock waves of the stochastic Burgers equation, has been studied in sufficient detail [8–19] (see also the reviews and books [1-4, 20] and references therein). The phenomenon of Anderson localization in disordered nonlinear systems with a random potential within the framework of the discrete NLS equation model was studied in Refs. [21–23], where the unlimited spreading of an initially localized wave packet was shown. Attempts were also made to directly numerically simulate solitons within the framework of the stochastic KdV and NLS equations [24, 25]. Examples of the 1D solitons in random media with a nonlocal nonlinearity were considered in Refs. [26, 27]. Alfvén and ion-cyclotron solitons in plasma with fluctuating parameters in the models of the derivative NLS equation and the Fokas-Lenells equation were studied in Refs. [28, 29] and [30], respectively. Solitons of the modified NLS equation in the presence of a random addition to the refractive index were considered in Refs. [31, 32]. Two-dimensional (2D) solitons of the Kadomtsev-Petviashvili equation with a noise term were explored in Ref. [33]. Much less attention has been devoted to multidimensional solitons and nonlinear structures in random media. The point is that, as is known, stationary solutions (if any) of nonlinear evolution equations in the multidimensional case often turn out to be unstable and subject to the phenomenon of collapse or self-focusing [34, 35, 37]. The collapse induced by the cubic nonlinearity is critical in the 2D space, which means that it sets in when the norm of the underlying wave field exceeds a certain finite critical value, and supercritical in the three-dimensional (3D) space, where an initial state with an arbitrarily small norm may blow up due to the collapse [36–40]. Note that to deal with realistic mod-

^{*}Electronic address: vlashkin62@gmail.com

els, saturation had been suggested as a way to stabilize such a catastrophic collapse or self-focusing and produce stable soliton structures of higher dimensions.

Already in early studies [41–43] on the self-focusing of optical beams in random media, it was shown that a random addition to the refractive index can significantly weaken or completely prevent the effect of self-focusing of the beam. In this case, different approximations were used, for example, an incoherent self-interacting light beam was studied in the random phase approximation in Ref. [41], the Rytov method for wave propagation in random media generalized to include self-focusing effect was considered in Ref. [42], and in Ref. [43] a purely Markov approximation was used. Using the so-called method of polynomial representation of the moments, it was shown in Ref. [44] that, under fairly general assumptions, collapse does not occur on average. In Ref. [45], the results of Ref. [44] were confirmed (in particular, that collapse on average is impossible) and a sufficient criterion for arresting the collapse of a pulse in a nonlinear waveguide with a randomly fluctuating refractive index was obtained. Note that in all the above-mentioned works, in the absence of randomness (that is, in the deterministic case), the considered models were reduced to the 2D NLS equation. The stochastic 2D NLS equation with multiplicative noise was also considered in Ref. [46], where the perturbation theory was developed in the framework of the so-called ε -expansion, and in Ref. [47] using the socalled δ -expansion. In both cases, the authors argued that the developed perturbation theory makes it possible to effectively reduce the stochastic 2D NLS equation to the linear stochastic equation. The effects of randomness have been discussed in recent papers [48–50], and it has also been noted that stochasticity in multidimensional NLS equation leads to the prevention of the collapse phenomenon. As indicated above, the collapse in a multidimensional NLS with a random term may not arrest in the general case, but be significantly weakened. First of all, this concerns 3D models where there is no critical collapse regime. In addition, the possibility of preventing collapse apparently depends on the type of random perturbation and its statistical properties. For example, in Ref. [51] the possibility of collapse in the 3D NLS model with a random term in the form of noise depending only on time against the background of a trapping parabolic potential was demonstrated both by an analytical variational method and by numerical simulation. It was shown that the expected collapse time is inversely proportional to the integrated covariance of the time autocorrelation function.

In most cases, when calculating statistical characteristics the random term is small and then a perturbation theory along with the appropriate closure procedures can be applied. One of the few exceptions is the work of Besieris [52] (a brief synopsis of this work can be found, for example, in Refs. [1, 3]), where a novel statistical technique is proposed that allows one to calculate the moments of even orders associated with the 1D NLS stochastic equation without the need to derive a closed equation for the moments. A necessary condition for this was the quadratic form of the noise autocorrelation function. In this work, some statistical characteristics describing the soliton propagation in the presence of noise were also calculated.

In the present paper, following the ideas of Ref. [52], we consider the NLS equation with multiplicative random term depending on time and space variables in the spatial dimension $D \ge 2$. In this case, no procedure of statistical closure for the nonlinear term. The main thrust of the research will be concerned with the analysis of two specific stochastic nonlinear equations for even moments employing the quadratic form of the autocorrelation function. After introducing a stochastic change of variables and a fundamental ansatz, we will show explicitly how each of the aforementioned stochastic nonlinear equations can be mapped into an explicit deterministic equation for a new dependent variable and a system of stochastic nonlinear coupled ordinary differential equations governing the remaining elements of the transformation. The ensuing analysis proceeds two-fold. At one level, we must be able to solve a deterministic nonlinear partial differential equation governing the new dependent variable. At the other level, we must proceed to derive the Fokker-Planck equation and to find the joint probability density function for the stochastic system. Combining the two solutions and inverting the mapping, we are able to find explicit expressions for any even moment of the wave field. As an application of the developed formalism, we consider the influence of noise on the 2D and 3D fundamental (ground state) and vortex solitons, which are unstable in the absence of randomness. The vortex soliton (spinning soliton) is the localized nonlinear structure with embedded vorticity and ringlike in the 2D or toroidal in 3D case field intensity distribution, with the dark hole at the center where the phase dislocation takes place: a phase circulation around the azimuthal axis is equal to $2m\pi$. An integer m is referred to as topological charge. We calculate some averaged observables for these structures and show that both fundamental and vortex solitons do not collapse.

The paper is organized as follows. In Sec. II, using Donsker-Furutsu-Novikov functional formalism, a stochastic equation for even moments is obtained. The effective nonlinear stochastic equation and a stochastic change of variables are proposed in Sec. III. In Sec. IV., the Fokker-Planck equation for the joint probability density is derived and solved. Using the variational method, analytical expressions for the fundamental and vortex solitons are found Sec. V. In Sec. VI, the statistical characteristics for the fundamental and vortex solitons are analytically calculated. The conclusion is made in Sec. VII.

II. BASIC EQUATIONS

We start with the normalized NLS equation in the spatial dimension $D \ge 2$ with a multiplicative random term,

$$i\frac{\partial\varphi}{\partial t} + \Delta\varphi + |\varphi|^2\varphi + \varepsilon(\mathbf{r}, t)\varphi = 0, \qquad (1)$$

where $\varepsilon(\mathbf{r}, t)$ is a zero-mean real homogeneous in space and stationary in time gaussian random field with the autocorrelation function

$$\langle \varepsilon(\mathbf{r}, t)\varepsilon(\mathbf{r}', t')\rangle = A(\mathbf{r} - \mathbf{r}') C(t - t')$$
(2)

where the angular brackets $\langle \dots \rangle$ means ensemble averaging. Without loss of generality, we consider both 2D and 3D cases (the 1D case was considered in Ref. [52]). This model equation arises in a number of important physical applications. In particular, these are the following.

- 1. The propagation of a continuous wave beam (the 2D case) or the transmission of a spatiotemporal light signal (the 3D case) in a bulk self-focusing optical medium with the Kerr nonlinearity and a random addition to the refractive index. Here t is the propagation distance z, and the x- and y-axes are two transverse directions for the 2D case. In the 3D case, the third coordinate z replaced by the local time $\tau = t z/v_0$ (v_0 is the group velocity of the carrier wave) [54].
- 2. The propagation of strong electromagnetic waves in the turbulent ionosphere [55], where $\varepsilon(\mathbf{r}, t)$ stands for the turbulent part of the refractive index.
- 3. The nonlinear Langmuir waves in the turbulent unmagnetized plasma. Here $\varepsilon(\mathbf{r}, t)$ corresponds to the turbulent part of the plasma density perturbation [56].
- 4. Nonlinear matter waves in the Bose-Einstein condensates (BEC) with random potentials (BEC with disordering) [57, 58].

Our ultimate goal in connection with Eq. (1) is to compute the Nth order moments $\langle \varphi^{(N)}(\mathbf{x},t) \rangle$, where

$$\varphi^{(N)}(\mathbf{x},t) = \prod_{m=1}^{N/2} \varphi(\mathbf{r}_{2m-1},t)\varphi^*(\mathbf{r}_{2m},t), \qquad (3)$$

and $\mathbf{x} = (\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N)$. An evolution equation for $\varphi^{(N)}(\mathbf{x}, t)$ follows readily from the definition (3) and the basic equation (1) and has the form

$$i\frac{\partial\varphi^{(N)}}{\partial t} + \sum_{m=1}^{N} \xi_m \Delta_m \varphi^{(N)} + \sum_{m=1}^{N} \xi_m |\varphi(\mathbf{r}_m, t)|^2 \varphi^{(N)} + \sum_{m=1}^{N} \xi_m \varepsilon(\mathbf{r}_m, t) \varphi^{(N)} = 0, \qquad (4)$$

where

$$\Delta_m = \frac{\partial^2}{\partial x_m^2} + \frac{\partial^2}{\partial y_m^2},\tag{5}$$

for the two-dimensional case, and

$$\Delta_m = \frac{\partial^2}{\partial x_m^2} + \frac{\partial^2}{\partial y_m^2} + \frac{\partial^2}{\partial z_m^2} \tag{6}$$

for the three-dimensional one. The quantity ξ_m entering into Eq. (4) is assumed to be $\xi_m = 1$ if *m* is odd, and $\xi_m = -1$ if *m* is even. Ensemble averaging of the equation (4) results in the expression

$$i\frac{\partial\langle\varphi^{(N)}\rangle}{\partial t} + \sum_{m=1}^{N} \xi_m \Delta_m \langle\varphi^{(N)}\rangle + \sum_{m=1}^{N} \xi_m \langle|\varphi(\mathbf{r}_m, t)|^2 \varphi^{(N)}\rangle + \sum_{m=1}^{N} \xi_m \langle\varepsilon(\mathbf{r}_m, t)\varphi^{(N)}\rangle = 0$$
(7)

with the initial condition $\langle \varphi^{(N)}(\mathbf{x}, 0) \rangle = \varphi_0^{(N)}(\mathbf{x})$. Equation (7) is not a closed equation for $\langle \varphi^{(N)} \rangle$ due to the third and fourth terms on the left-hand side. To calculate an average of the product of the gaussian random function $\eta(s)$ with the functional $R[\eta]$ depending on it, we will use the Donsker-Furutsu-Novikov functional formalism [59–61], where *s* denotes the set of all arguments, both continuous and discrete, on which the random function depends. The corresponding expression is

$$\langle \eta(s)R[\eta] \rangle = \int \langle \eta(s)\eta(s') \rangle \left\langle \frac{\delta R[\eta]}{\delta \eta(s')} \right\rangle ds', \qquad (8)$$

where $\delta/\delta\eta$ denotes the functional derivative, and the integral extends over the region in which the function is defined and, as usual, summation over the repeated indices of the discrete variables is assumed. A partial closure in Eq. (7) can be performed for a special class of random fields $\varepsilon(\mathbf{r}, t)$ when $C(t - t') = \delta(t - t')$ in Eq. (2), where $\delta(\tau)$ is the Dirac delta function (the white noise in time). Then, using Eq. (8) we have for the average in the last term of Eq. (7),

$$\left\langle \varepsilon(\mathbf{r}_{m},t)\varphi^{(N)}(\mathbf{x},t)\right\rangle = \int \left\langle \varepsilon(\mathbf{r}_{m},t)\varepsilon(\mathbf{r}_{m}',t')\right\rangle$$
$$\times \left\langle \frac{\delta\varphi^{(N)}(\mathbf{x},t)}{\delta\varepsilon(\mathbf{r}_{m}',t')}\right\rangle d\mathbf{r}_{m}' dt' = \int A(\mathbf{r}_{m}-\mathbf{r}_{m}')$$
$$\times \left\langle \frac{\delta\varphi^{(N)}(\mathbf{x},t)}{\delta\varepsilon(\mathbf{r}_{m}',t)}\right\rangle d\mathbf{r}_{m}'. \tag{9}$$

The functional derivative $\delta \varphi^{(N)} / \delta \varepsilon$ required in Eq. (9) can be found from the original equation (4). To this end we rewrite Eq. (4) in the integral form

$$\varphi^{(N)}(\mathbf{x},t) = \varphi_0^{(N)}(\mathbf{x}) + i \int_0^t \sum_{m=1}^N \xi_m [\Delta_m + |\varphi(\mathbf{r}_m,t')|^2 + \varepsilon(\mathbf{r}_m,t')] \varphi^{(N)}(\mathbf{x},t') dt'.$$
(10)

Then, taking the functional derivative in (10) and using

$$\frac{\delta\varepsilon(\mathbf{r},t)}{\delta\varepsilon(\mathbf{r}',t')} = \delta(\mathbf{r} - \mathbf{r}')\delta(t - t'), \qquad (11)$$

(see, e. g. [3]), we find for 0 < t' < t

$$\frac{\delta\varphi^{(N)}(\mathbf{x},t)}{\delta\varepsilon(\mathbf{r}'_{m},t')} = i\sum_{n=1}^{N} \xi_{n} \int_{t'}^{t} \left\{ \left[\Delta_{n} + \varepsilon(\mathbf{r}_{n},t'') \right] \frac{\delta\varphi^{(N)}(\mathbf{x},t'')}{\delta\varepsilon(\mathbf{r}'_{m},t')} + \frac{\delta[|\varphi^{(N)}(\mathbf{r}_{n},t'')|^{2}\varphi^{(N)}(\mathbf{x},t'')]}{\delta\varepsilon(\mathbf{r}'_{m},t')} \right\} + \frac{i}{2}\sum_{n=1}^{N} \xi_{n}\varphi^{(N)}(\mathbf{x},t')\delta(\mathbf{r}_{n}-\mathbf{r}'_{m}), \quad (12)$$

where we have used the causality principle $\delta \varphi^{(N)}(\mathbf{x}, t'') / \delta \varepsilon(\mathbf{r}', t') = 0$ for t'' < t', (the factor 1/2 appears due to $\int_0^\infty \delta(t) dt$). In the limit $t' \to t$ we get

$$\frac{\delta\varphi^{(N)}(\mathbf{x},t)}{\delta\varepsilon(\mathbf{r}'_m,t)} = \frac{i}{2} \sum_{n=1}^{N} \xi_n \varphi^{(N)}(\mathbf{x},t) \delta(\mathbf{r}_n - \mathbf{r}'_m) \qquad (13)$$

Substituting Eq. (13) into Eq. (9), we obtain

$$\langle \varepsilon(x_m, t)\varphi^{(N)} \rangle = \frac{i}{2} \sum_{n=1}^{N} \xi_n A(\mathbf{r}_m - \mathbf{r}_n) \langle \varphi^{(N)} \rangle.$$
(14)

Finally substituting this into Eq. (7) we have

$$i\frac{\partial\langle\varphi^{(N)}\rangle}{\partial t} + \sum_{m=1}^{N} \xi_m \Delta_m \langle\varphi^{(N)}\rangle + \sum_{m=1}^{N} \xi_m \langle|\varphi(\mathbf{r}_m, t)|^2 \varphi^{(N)}\rangle + \frac{i}{2} \sum_{m=1}^{N} \sum_{n=1}^{N} \xi_m \xi_n A(\mathbf{r}_m - \mathbf{r}_n) \langle\varphi^{(N)}\rangle = 0.$$
(15)

Equation (15) is not a closed equation for $\langle \varphi^{(N)} \rangle$ due to the presence of nonlinearities. An attempt to effect a closure of the third term on the left-hand side of Eq. (15) will result in the infinite hierarchy of coupled moment equations mentioned in the introduction.

In many physical applications, the correlation function $A(\boldsymbol{\rho})$ is isotropic with the characteristic scale ρ_c and of power-law type for $|\boldsymbol{\rho}| < \rho_c$, so that for the structure function $A(0) - A(|\boldsymbol{\rho}|)$ one can write as

$$A(0) - A(|\boldsymbol{\rho}|) = A(0)(\rho/\rho_c)^{\nu}, \qquad (16)$$

where $\nu \leq 2$ (see, e. g. discussion in Ref. [62]). The case $\nu = 5/3$, in particular, corresponds to the well-known Kolmogorov spectrum $\sim \kappa^{-11/3}$, where κ is the wave number. In what follows, as in Ref. [52], we restrict ourselves to the case $\nu = 2$ which corresponds to a simplified or quadratic Kolmogorov spectrum. Such a quadratic approximation of the autocorrelation function occurs in many physically interesting situations [6, 63, 64]. In particular, the quadratic spectrum appears in the turbulence

(e. g. in turbulent atmosphere and ionosphere) on scales smaller than the Kolmogorov length scale [6]. In addition, strictly speaking, the use of such a quadratic form is justified only if the functionals of $A(0) - A(|\boldsymbol{\rho}|)$ is not sensitive at too large distances, which takes into account the exponential localization of the structures considered below. Then we have

$$A(\mathbf{r} - \mathbf{r}') = A(0) \left[1 - \frac{(\mathbf{r} - \mathbf{r}')^2}{\rho_c^2} \right], \qquad (17)$$

and further we denote $B = 3A(0)/(8\rho_c^2)$ (the factor 3/8 is introduced to simplify the appearance of expressions in Sec. VI). Then inserting Eq. (17) into Eq. (15) one can obtain

$$i\frac{\partial\langle\varphi^{(N)}\rangle}{\partial t} + \sum_{m=1}^{N} \xi_m \Delta_m \langle\varphi^{(N)}\rangle + \sum_{m=1}^{N} \xi_m \langle|\varphi(\mathbf{r}_m, t)|^2 \varphi^{(N)}\rangle - \frac{3i}{8}B \sum_{m(18)$$

This equation, like Eq. (15), is of course also not a closed equation for $\langle \varphi^{(N)} \rangle$.

III. EFFECTIVE NONLINEAR STOCHASTIC EQUATION

Let us consider the auxiliary nonlinear stochastic equation for the function φ_{e} ,

$$i\frac{\partial\varphi_{\rm e}}{\partial t} + \Delta\varphi_{\rm e} + |\varphi_{\rm e}|^2\varphi_{\rm e} + \frac{1}{2}\mathbf{r}\cdot\mathbf{f}(t)\varphi_{\rm e} = 0, \qquad (19)$$

where $\mathbf{f}(t)$ is a zero-mean, stationary, δ -correlated in time, vector real Gaussian random process characterized by the correlation matrix

$$\langle f_i(t)f_j(t')\rangle = \frac{3}{2}B\delta_{ij}\,\delta(t-t'),\tag{20}$$

and *B* is defined in Eq. (17). In the one-dimensional case, Eq. (19) was previously proposed in Ref. [52] (see also Eq. (2.3) in Ref. [1], and Eq. (3.2.4) in Ref. [3]). We define the function $\varphi_{\rm e}^{(N)}$ in the same way as in Eq. (3) by replacing φ with $\varphi_{\rm e}$,

$$\varphi_{\mathbf{e}}^{(N)}(\mathbf{x},t) = \prod_{m=1}^{N/2} \varphi_{\mathbf{e}}(\mathbf{r}_{2m-1},t) \varphi_{\mathbf{e}}^*(\mathbf{r}_{2m},t).$$
(21)

Then, by analogy with Eq. (4), we get

$$i\frac{\partial\varphi_{\mathrm{e}}^{(N)}}{\partial t} + \sum_{m=1}^{N}\xi_{m}\Delta_{m}\varphi_{\mathrm{e}}^{(N)} + \sum_{m=1}^{N}\xi_{m}|\varphi_{\mathrm{e}}(\mathbf{r}_{m},t)|^{2}\varphi_{\mathrm{e}}^{(N)} + \frac{1}{2}\sum_{m=1}^{N}\xi_{m}\mathbf{r}_{m}\cdot\mathbf{f}(t)\varphi_{\mathrm{e}}^{(N)} = 0.$$
(22)

Averaging Eq. (22) yields

$$i\frac{\partial\langle\varphi_{\rm e}^{(N)}\rangle}{\partial t} + \sum_{m=1}^{N} \xi_m \Delta_m \langle\varphi_{\rm e}^{(N)}\rangle + \sum_{m=1}^{N} \xi_m \langle|\varphi_{\rm e}(\mathbf{r}_m, t)|^2 \varphi_{\rm e}^{(N)}\rangle + \frac{1}{2} \sum_{m=1}^{N} \xi_m \mathbf{r}_m \cdot \langle \mathbf{f}(t) \varphi_{\rm e}^{(N)}\rangle = 0, \qquad (23)$$

The statistical averaging in the fourth term on the lefthand side of Eq. (23) can be handled by the Donsker-Furutsu-Novikov formula (8). Then, using Eq. (20), we have

$$\left\langle \mathbf{f}(t)\varphi_{\mathrm{e}}^{(N)}\right\rangle = \frac{3}{2}B\left\langle \frac{\delta\varphi_{\mathrm{e}}^{(N)}}{\delta\mathbf{f}(t)}\right\rangle$$
 (24)

The functional derivative can be found from Eq. (22),

$$\frac{\delta\varphi_{\rm e}^{(N)}(\mathbf{x},t)}{\delta\mathbf{f}(t)} = \frac{i}{4} \sum_{n=1}^{N} \xi_n \mathbf{r}_n \varphi_{\rm e}^{(N)}(\mathbf{x},t), \qquad (25)$$

and then we have

$$\left\langle \mathbf{f}(t)\varphi_{\mathrm{e}}^{(N)}\right\rangle = \frac{3i}{8}B\sum_{n=1}^{N}\xi_{n}\mathbf{r}_{n}\langle\varphi_{\mathrm{e}}^{(N)}(\mathbf{x},t)\rangle.$$
 (26)

Substituting Eq. (26) into Eq. (23) we have

$$i\frac{\partial\langle\varphi_{\rm e}^{(N)}\rangle}{\partial t} + \sum_{m=1}^{N}\xi_{m}\Delta_{m}\langle\varphi_{\rm e}^{(N)}\rangle + \sum_{m=1}^{N}\xi_{m}\langle|\varphi_{\rm e}(\mathbf{r}_{m},t)|^{2}\varphi_{\rm e}^{(N)}\rangle + \frac{3i}{8}B\sum_{m=1}^{N}\sum_{n=1}^{N}\xi_{m}\xi_{n}\mathbf{r}_{m}\cdot\mathbf{r}_{n}\langle\varphi_{\rm e}^{(N)}\rangle = 0.$$
(27)

Using identity (it can be proved, for example, by the method of mathematical induction)

$$-\sum_{m$$

and comparing Eq. (18) and Eq. (27) we can see that the equations for $\langle \varphi^{(N)} \rangle$ and $\langle \varphi_{\rm e}^{(N)} \rangle$ are identical, that is $\langle \varphi^{(N)} \rangle = \langle \varphi_{\rm e}^{(N)} \rangle$. We emphasize that the equivalence of equations (27) and (18) is valid only under the assumption of the quadratic Kolmogorov spectrum of the random process $\varepsilon(\mathbf{r}, t)$. Equation (27), being the same with Eq. (18), is not a closed equation for $\langle \varphi_{\rm e}^{(N)} \rangle$, but further we will show that this equation, in contrast to Eq. (18), admits a global statistical analysis. We use the following ansatz consisting in the change of variables

$$\mathbf{R}(t) = \mathbf{r} - \mathbf{u}(t) \tag{29}$$

and introducing of a new function $\psi(\mathbf{R}, t)$ by

$$\varphi_{\mathbf{e}}(\mathbf{r},t) = \psi(\mathbf{R}(t),t) \exp\{i[\mathbf{u}_t(t) \cdot \mathbf{R}(t)/2 + \gamma(t)]\}, \quad (30)$$

where the subscript denotes differentiation in t. By direct substitution Eq. (30) into Eq. (19) one can see that ψ satisfy the equation

$$i\frac{\partial\psi}{\partial t} + \Delta_{\mathbf{R}}\psi + |\psi|^2\psi = 0, \qquad (31)$$

where $\Delta_{\mathbf{R}} = \partial^2 / \partial R_1^2 + \partial^2 / \partial R_2^2 + + \partial^2 / \partial R_3^2$ (and the corresponding expression for the 2D case), provided that the functions $\mathbf{u}(t)$ and $\gamma(t)$ satisfy equations

$$\mathbf{u}_{tt} = \mathbf{f}(t), \ \mathbf{u}(0) = 0, \ \mathbf{u}_t(0) = 0,$$
 (32)

and

$$\gamma_t = \frac{1}{4}\mathbf{u}_t^2(t) + \frac{1}{2}\mathbf{u}(t) \cdot \mathbf{f}(t), \ \gamma(0) = 0$$
(33)

for $t \ge 0$. The choice of zero initial data provides a one-to-one mapping of the dependent variable ψ onto φ_e . Thus, the ansatz (30) has transformed the stochastic nonlinear equations (19) into an explicitly deterministic nonlinear equations for $\psi(\mathbf{R}, t)$. It should be noted, however, that ψ is random function by virtue of its dependence on $\mathbf{u}(t)$ through $\mathbf{R}(t)$ in Eq. (29). The solution ψ of Eq. (31) is used in the expression (30) for φ_e . This function, in turn, is a functional of \mathbf{u} , \mathbf{u}_t and γ because of its dependence on ψ and the presence of the exponential factor in Eq. (30). The computation of statistical moments of φ_e require knowledge of the joint probability density functional of the random processes $\mathbf{u},\,\mathbf{u}_t$ and γ . Here, however, we are only interested in even Nth order moments $\langle \varphi_{\mathbf{e}}^{(N)} \rangle$ or, equivalently, moments of Eq. (3), and then $\varphi_{\rm e}$ is independent of γ . Thus, our statistical analysis is limited to finding only the joint density functional of **u** and \mathbf{u}_t .

In the conclusion of this section we would like to make a remark regarding Eq. (19) and its 1D version. It should be noted that the replacements Eqs. (29) and (30) reducing the non-autonomous equation (19) with an explicit dependence on t to the autonomous equation (31) is rather nontrivial. In the 1D case, such a replacement was proposed by Besieris in Ref. [52]. On the other hand, in the 1D case, as is known, Eq. (31) is completely integrable, has N-soliton solutions and can be solved by the inverse scattering transform (IST) method [65]. Indeed, by generalizing the spectral Zakharov-Shabat problem for the NLS equation, it was shown that the onedimensional version of Eq. (19) can be solved using the IST [66, 67]

IV. FOKKER-PLANCK EQUATION

Under the assumptions made about the vector process \mathbf{f} , equation (32) with the external random force \mathbf{f} determined by the autocorrelation matrix (20) is the equation for one Brownian particle and it can be written as a system

$$\mathbf{u}_t = \mathbf{v},\tag{34}$$

$$\mathbf{v}_t = \mathbf{f}(t),\tag{35}$$

with the initial conditions $\mathbf{v}(0) = 0$ and $\mathbf{u}(0) = 0$. Then, the Fokker-Planck equation [68] for the joint probability density function $p(\mathbf{u}, \mathbf{v}, t)$ has the form

$$\frac{\partial p}{\partial t} + \mathbf{v} \frac{\partial p}{\partial \mathbf{u}} = \frac{3B}{4} \frac{\partial^2 p}{\partial \mathbf{v}^2} \tag{36}$$

with the initial condition $p(\mathbf{u}, \mathbf{v}, 0) = \delta(\mathbf{u})\delta(\mathbf{v})$. In the Appendix, we derived a more general Fokker-Planck equation for a joint probability density function $p(\mathbf{u}, \mathbf{v}, \gamma, t)$ that also depends on γ . Equation (36) follows from (94) when the dependence on γ is neglected. It can readily be solved and the solution has the form of the six-dimensional gaussian distribution with the moments

$$\langle u_i u_j \rangle = \delta_{ij} \sigma_{11}^2, \ \langle u_i v_j \rangle = \delta_{ij} \sigma_{12}^2, \ \langle v_i v_j \rangle = \delta_{ij} \sigma_{22}^2,$$
(37)

where i, j = x, y, z, and

$$\sigma_{11}^2 = \frac{1}{2}Bt^3, \ \sigma_{12}^2 = \frac{3}{4}Bt^2, \ \sigma_{22}^2 = \frac{3}{2}Bt.$$
 (38)

Separate components of \mathbf{u} and \mathbf{v} are statistically independent due to Eqs. (20), (34) and (35), so that the sixdimensional gaussian distribution is the product of the two-dimensional distributions for the each component,

$$p(\mathbf{u}, \mathbf{v}, t) = \prod_{i=1}^{D} p(u_i, v_i, t).$$
(39)

For $p(u_x, v_x, t)$, for example, one obtains [68]

$$p(u_x, v_x, t) = \frac{1}{2\pi\sqrt{\det\Sigma}} \times \exp\left[-\frac{1}{2\det\Sigma} \left(\sigma_{22}^2 u_x^2 + \sigma_{11}^2 v_x^2 - 2\sigma_{12}^2 u_x v_x\right)\right], \quad (40)$$

where Σ is the 2 × 2 covariance matrix with the elements σ_{mn}^2 , m, n = 1, 2 and det $\Sigma = 3B^2t^4/16$. Then the joint probability density function of **u** and **v** is

$$p(\mathbf{u}, \mathbf{v}, t) = \frac{1}{(2\pi)^D (\det \Sigma)^{D/2}}$$
$$\times \exp\left[-\frac{1}{2 \det \Sigma} \left(\sigma_{22}^2 \mathbf{u}^2 + \sigma_{11}^2 \mathbf{v}^2 - 2\sigma_{12}^2 \mathbf{u} \cdot \mathbf{v}\right)\right]. \quad (41)$$

Note that in the limit $t \to \infty$ the joint probability density function $p(\mathbf{u}, \mathbf{v}, t)$ tends to zero as $\sim 1/t^{2D}$. From Eq. (41) it follows that

$$p(\mathbf{u},t) = \int p(\mathbf{u},\mathbf{v},t) d^D \mathbf{v} = \frac{1}{(2\pi)^{D/2} \sigma_{11}^D} \exp\left(-\frac{\mathbf{u}^2}{2\sigma_{11}^2}\right),$$
(42)

$$p(\mathbf{v},t) = \int p(\mathbf{u},\mathbf{v},t) \, d^D \mathbf{u} = \frac{1}{(2\pi)^{D/2} \sigma_{22}^D} \exp\left(-\frac{\mathbf{v}^2}{2\sigma_{22}^2}\right).$$
(43)

In the limit $t \to \infty$, the probability density functions $p(\mathbf{u},t)$ and $p(\mathbf{v},t)$ behave as $\sim 1/t^{3D/2}$ and $\sim 1/t^{D/2}$,

respectively. Given the known joint probability density function $p(\mathbf{u}, \mathbf{v}, t)$, we can write

$$\langle \varphi^{(N)} \rangle = \langle \varphi_{\mathbf{e}}^{(N)} \rangle = \int \varphi_{\mathbf{e}}^{(N)} p(\mathbf{u}, \mathbf{v}, t) d^{D} \mathbf{u} d^{D} \mathbf{v}, \quad (44)$$

where $\varphi_{e}^{(N)}$ is constructed from solutions of Eq. (31) using Eqs. (29) and (30).

V. DETERMINISTIC STATIONARY SOLUTIONS

In this section we will consider stationary solutions of Eq. (31) in the form of multidimensional coherent structures, namely the fundamental soliton and the vortex soliton. Since corresponding exact analytical solutions do not exist (and numerical solutions are not adequate for further analysis), we use the variational method. Note that these solutions are purely formal in the sense that here we do not study the problem of the stability of such solutions. Equation (31) can be written in Hamiltonian form

$$i\frac{\partial\psi}{\partial t} = \frac{\delta H}{\delta\psi^*},\tag{45}$$

where the Hamiltonian is

$$H = \int \left(|\nabla \psi|^2 - \frac{|\psi|^4}{2} \right) d^D \mathbf{R}.$$
 (46)

Other integrals of motion are the norm (also called energy, number of particles, or in optics wave power)

$$N = \int |\psi|^2 d^D \mathbf{R},\tag{47}$$

the momentum

$$\mathbf{P} = \frac{i}{2} \int (\psi \nabla \psi^* - \psi^* \nabla \psi) d^D \mathbf{R}$$
(48)

and the angular momentum

$$\mathbf{M} = i \int \mathbf{R} \times (\psi^* \nabla \psi - \psi \nabla \psi^*) d^D \mathbf{R}.$$
 (49)

An extensive literature is devoted to the NLS equation (31) with the dimension of space D (see, for example, [69] and numerous references therein). It is well known that the NLS equation with D = 1 is completely integrable and admits N-soliton solutions [65], while for $D \ge 2$ the initial field distribution undergoes collapse (or self-focusing if t has the meaning of a spatial variable along the beam) or spreads out [38–40]. The collapse is critical if D = 2, which means that it sets in when the norm of the underlying wave field exceeds a certain finite critical value, and supercritical in 3D, where an initial state with an arbitrarily small norm may blow up. Under this, the initial perturbation in any case never reaches a stationary

state (in other words, stationary solutions are unstable). We are still looking for formal stationary solutions of Eq. (31) of the form

$$\psi(\mathbf{R},t) = \phi(\mathbf{R}) \mathrm{e}^{i\lambda t}.$$
 (50)

These solutions satisfy the equation

$$-\lambda\phi + \Delta_{\mathbf{R}}\phi + |\phi|^2\phi = 0, \qquad (51)$$

and resolve the variational problem $\delta S=0$ for the functional

$$S = H + \lambda N. \tag{52}$$

Note that in this paper we restrict ourselves to motionless and non-rotating solutions. Solutions moving with the constant velocity \mathbf{v} correspond to the additional term $\mathbf{v} \cdot \mathbf{P}$ in the functional S [53]. Due to the Galilean invariance of Eq. (31), moving solutions are easily obtained from motionless solutions by the appropriate Galilean transformation. For solutions rotating with the constant angular velocity ω , i.e., which are stationary in the reference frame rotating with constant angular velocity ω , the functional S would also contain term $\boldsymbol{\omega} \cdot \mathbf{M}$. The procedure for finding such solutions (both by variational and direct numerical methods) is described in [70] and these solutions are vortex clusters (azimuthon solitons). In this paper, we do not consider such rotating stationary structures, although the developed formalism is fully applicable to them as well.

The solution of Eq. (51) in the form of a fundamental soliton (ground state) corresponds to the radially symmetric case with boundary conditions $\partial \phi / \partial R = 0$ at zero R = 0 and $\phi \to 0$ at infinity $R \to \infty$. The vortex soliton corresponds to ansatz $\phi = \phi(R) \exp(im\theta)$, where θ is the polar angle and m is an integer (topological charge), with boundary conditions $\phi = 0$ at zero R = 0 and $\phi \to 0$ at infinity $R \to \infty$. The important integral of motion associated with this type of solitary wave is the angular momentum. One can show that for the vortex soliton $M_z = mN$. Solutions of Eq. (51) in the form of fundamental and vortex solitons can be found numerically in both 2D and 3D cases [71], and the corresponding solutions turn out to be unstable.

For an analytical treatment, following Ref. [53], we consider the trial function of the form $\phi(\mathbf{R}) = A\phi_a(\mathbf{R}/a)$, where A and a are unknown parameters to be determined by the variational procedure from the system of equations $\partial S/\partial A = 0$ and $\partial S/\partial a = 0$. For not too large values of λ , or equivalently N, the variational analysis is in good agreement with the solutions obtained by exact numerical simulation. We denote $\boldsymbol{\xi} = \mathbf{R}/a$ and for an arbitrary form of the $\phi_a(\boldsymbol{\xi})$, provided ϕ_a vanishes at infinity fast enough so that the corresponding integrals are convergent, one can get

$$A^{2} = \frac{2\lambda\delta}{\beta(4-D)}, \quad a^{2} = \frac{\alpha(4-D)}{D\lambda\delta}, \quad (53)$$

where

$$\alpha = \int |\nabla \varphi_a|^2 d^D \boldsymbol{\xi}, \quad \beta = \int \frac{|\varphi_a|^4}{2} d^D \boldsymbol{\xi}, \quad (54)$$

$$\delta = \int |\varphi_a|^2 d^D \boldsymbol{\xi}.$$
 (55)

We choose the test functions in the form:

(i) The ground state (fundamental soliton)

$$\phi_a(\boldsymbol{\xi}) = \exp(-\xi^2/2). \tag{56}$$

(ii) The vortex soliton

$$\phi_a(\boldsymbol{\xi}) = \xi^{|m|} \exp(-\xi^2/2) \mathrm{e}^{im\theta}, \qquad (57)$$

and further we restrict ourselves to the case m = 1. Substituting Eqs. (56) and (57) into Eqs. (54) and (55), and calculating the corresponding integrals one can obtain, following Ref. [53], for the fundamental soliton in the 2D case $\alpha = \delta = \pi$, $\beta = \pi/4$, and in the 3D case $\alpha = 3\pi^{3/2}/2$, $\beta = \sqrt{2\pi^{3/2}}/8$, $\delta = \pi^{3/2}$. For the vortex soliton we have $\alpha = 2\pi, \beta = \pi/8$, $\delta = \pi$ in the 2D case, and $\alpha = 5\pi^{3/2}/2, \beta = \sqrt{2\pi^{3/2}}/16$, $\delta = \pi^{3/2}$ in the 3D case. Then, for the parameters A (the amplitude) and a (characteristic size of the structures), from Eq. (53) we can obtain for the fundamental soliton,

$$A^{2} = \begin{cases} 4\lambda & \text{for 2D,} \\ 8\sqrt{2}\lambda & \text{for 3D} \end{cases}, \quad a^{2} = \begin{cases} 1/\lambda & \text{for 2D,} \\ 1/(2\lambda) & \text{for 3D} \end{cases},$$
(58)

and for the vortex soliton,

$$A^{2} = \begin{cases} 8\lambda & \text{for 2D,} \\ 16\sqrt{2}\lambda & \text{for 3D} \end{cases}, \quad a^{2} = \begin{cases} 2/\lambda & \text{for 2D,} \\ 5/(6\lambda) & \text{for 3D} \end{cases}.$$
(59)

Since $N = A^2 a^D \delta$, it follows from Eq. (58) that for the 3D fundamental soliton $\partial N/\partial \lambda < 0$, that is, according to the Vakhitov-Kolokolov criterion, the soliton is unstable. For a two-dimensional soliton, we have $\partial N/\partial \lambda = 0$, which corresponds to the marginal instability (critical collapse).

VI. AVERAGED OBSERVABLES FOR FUNDAMENTAL AND VORTEX SOLITONS

Due to the parametric dependence of \mathbf{R} on time in Eq. (29), solutions (50), (56) and (57) of Eq. (31) presented in the previous section with replacement Eq. (30) are not stationary at all and, moreover, are random functions. In this case they have the form

$$\varphi_{\rm e}(\mathbf{r},t) = A \exp\left[-\frac{(\mathbf{r}-\mathbf{u})^2}{2a^2} + i\frac{\mathbf{v}\cdot(\mathbf{r}-\mathbf{u})}{2a^2} + i\gamma + i\lambda t\right]$$
(60)



FIG. 1: The mean 2D vortex soliton intensity $\langle I(\mathbf{r}, t) \rangle$ in the presence of random fluctuations according to the analytical prediction Eq. (64) for the parameters $\lambda = 1$ and B = 0.05.

for the fundamental soliton, and

$$\varphi_{\mathbf{e}}(\mathbf{r},t) = A \frac{\left[(x-u_x)+i(y-u_y)\right]}{a}$$
$$\times \exp\left[-\frac{(\mathbf{r}-\mathbf{u})^2}{2a^2} + i\frac{\mathbf{v}\cdot(\mathbf{r}-\mathbf{u})}{2a^2} + i\gamma + i\lambda t\right] \qquad (61)$$

for the vortex soliton, respectively. Our first task is to compute the mean intensities $\langle |\varphi(\mathbf{r},t)|^2 \rangle = \langle |\varphi_e(\mathbf{r},t)|^2 \rangle$ of the fundamental and vortex solitons. Next we denote $I(\mathbf{r},t) = |\varphi(\mathbf{r},t)|^2$. In this case the function $I(\mathbf{r},t)$ is a function of only of the random process $\mathbf{u}(t)$. Its ensemble average, therefore, given by

$$\langle I(\mathbf{r},t)\rangle = \int p(\mathbf{u},t)I(\mathbf{r},t)d^{D}\mathbf{u},$$
 (62)

where $p(\mathbf{u}, t)$ is determined by Eq. (42) and $I(\mathbf{r}, t)$ is determined from Eqs. (60) and (61) for the fundamental and vortex soliton respectively. Calculating the integrals in Eq. (62), we obtain for the fundamental soliton,

$$\langle I(\mathbf{r},t)\rangle = \frac{A^2 a^D}{(a^2 + Bt^3)^{D/2}} \exp\left[-\frac{r^2}{(a^2 + Bt^3)}\right],$$
 (63)

where A and a are determined by Eq. (58), and for the vortex soliton

$$\langle I(\mathbf{r},t) \rangle = \frac{A^2 a^D [r_{\perp}^2 a^2 + Bt^3 (a^2 + Bt^3)]}{(a^2 + Bt^3)^{D/2+2}} \\ \times \exp\left[-\frac{r^2}{(a^2 + Bt^3)}\right], \tag{64}$$

where A and a are determined by Eq. (59). Note that in the presence of randomness, as can be seen from (64), the zero at the center of the vortex soliton disappears. In both cases the mean intensity spreads out monotonically due to irreversible effects of randomness, and no collapse occurs. The mean 2D vortex soliton intensity $\langle I(\mathbf{r},t)\rangle$ with the parameters $\lambda = 1$, B=0.05 for different times is shown in Fig. 1. As can be seen, the conservative equation (1) conserves the norm $\int |\varphi|^2 d^D \mathbf{r}$ for any random realization of $\varepsilon(\mathbf{r},t)$ and then the averaged norm

$$\langle N \rangle = \int \langle I(\mathbf{r}, t) \rangle d^D \mathbf{r}$$
 (65)

is also conserved. For analytical expressions obtained by the variational method in the previous section, we have $\langle N \rangle = A^2 a^D \pi^{D/2}$. Next, we calculate the mean momentum density $\mathbf{p} = i(\psi \nabla \psi^* - \psi^* \nabla \psi)/2$ and the mean angular momentum density $\mathbf{m} = [\mathbf{r} \times \mathbf{p}]$. These quantities are nonzero only for a vortex soliton, since here we consider motionless structures. Then we have

$$\langle \mathbf{p}(\mathbf{r},t) \rangle = \int p(\mathbf{u},t)\mathbf{p}(\mathbf{r},t)d^{D}\mathbf{r},$$
 (66)

and calculating integrals we find for the mean momentum density,

$$\langle \mathbf{p}(\mathbf{r},t) \rangle = \frac{A^2 a^D[\hat{\mathbf{z}} \times \mathbf{r}]}{(a^2 + Bt^3)^{D/2+1}} \exp\left[-\frac{r^2}{(a^2 + Bt^3)}\right], \quad (67)$$

and for the mean angular momentum density we have $\langle \mathbf{m}(\mathbf{r},t) \rangle = [\mathbf{r} \times \langle \mathbf{p}(\mathbf{r},t) \rangle].$

The mean centroid of the structure, defined by

$$\bar{\mathbf{r}}(t) = \frac{1}{\langle N \rangle} \int \mathbf{r} \langle I(\mathbf{r}, t) \rangle d^D \mathbf{r}, \qquad (68)$$

as follows from Eqs. (63) and (64), is zero for all t. A measure of the spreading of the structure can be found by using the definition

$$\mathbf{r}_{s}^{2}(t) = \frac{1}{\langle N \rangle} \int [\mathbf{r} - \bar{\mathbf{r}}(t)]^{2} \langle I(\mathbf{r}, t) \rangle d^{D} \mathbf{r}.$$
 (69)

Calculating integrals and taking into account that $\bar{\mathbf{r}}(t) = 0$ we find that the standard deviation evolves in t as

$$[\mathbf{r}_s^2(t)]^{1/2} = \left[\frac{D}{2}(a^2 + Bt^3)\right]^{1/2} \tag{70}$$

for the fundamental soliton, and

$$[\mathbf{r}_s^2(t)]^{1/2} = \left[\frac{(D+2)a^2 + DBt^3}{2}\right]^{1/2}$$
(71)

for the vortex soliton. Dependence of $(\mathbf{r}_s^2)^{1/2}$ on t (recall that the evolutionary variable t for 2D beams plays the role of a spatial coordinate along the beam propagation) of the form $\sim (\text{const} + t^3)^{1/2}$ is in agreement with the result of Ref. [45] obtained with the aid of the Donsker-Furutsu-Novikov formalism and virial theorem for the 2D NLS equation. It is interesting to note that the same *t*-dependence of the 2D beam position standard deviation was found experimentally with high accuracy in an stochastic nonlocal focusing Kerr medium such as liquid crystals [72].

Consider then the spatial mutual coherence function $\Gamma(\mathbf{r}_1, \mathbf{r}_2, t) = \langle \varphi^{(2)}(\mathbf{r}_1, \mathbf{r}_2, t) \rangle$ that characterizes the angular spread and coherence properties of the wave field and is defined as

$$\Gamma(\mathbf{r}_1, \mathbf{r}_2, t) = \int p(\mathbf{u}, \mathbf{v}, t) \varphi(\mathbf{r}_1, t) \varphi^*(\mathbf{r}_2, t) d^D \mathbf{u} d^D \mathbf{v}.$$
 (72)

This function is important and is especially often used when studying wave beams in optics in the presence of a random addition to the refractive index. One can see that the function $\varphi(\mathbf{r}_1, t)\varphi^*(\mathbf{r}_2, t)$ depends on both **u** and **v**. For the fundamental soliton we have

$$\varphi(\mathbf{r}_1, t)\varphi^*(\mathbf{r}_2, t) = A^2 \exp\left[-\frac{(\mathbf{r}_1 - \mathbf{u})^2 + (\mathbf{r}_2 - \mathbf{u})^2}{2a^2} + i\frac{\mathbf{v} \cdot (\mathbf{r}_1 - \mathbf{r}_2)}{2a^2}\right],$$
(73)

and for the vortex soliton

$$\varphi(\mathbf{r}_{1},t)\varphi^{*}(\mathbf{r}_{2},t) = \frac{A^{2}}{a^{2}}[x_{1} - u_{x} + i(y_{1} - u_{y})][x_{2} - u_{x} + i(y_{2} - u_{y})]\exp\left[-\frac{(\mathbf{r}_{1} - \mathbf{u})^{2} + (\mathbf{r}_{2} - \mathbf{u})^{2}}{2a^{2}} + i\frac{\mathbf{v}\cdot(\mathbf{r}_{1} - \mathbf{r}_{2})}{2a^{2}}\right].$$
(74)

To avoid a rather cumbersome expression, we present the corresponding expression for the spatial mutual coherence function only for the fundamental soliton. Using Eqs. (41) and (72) we get

$$\Gamma(\mathbf{r}_{1},\mathbf{r}_{2},t) = \frac{A^{2}a^{D}}{(a^{2}+Bt^{3})^{D/2}} \exp\left\{-\frac{Bt(3a^{2}+4a^{2}t^{2}+3Bt^{3}/4)(\mathbf{r}_{1}-\mathbf{r}_{2})^{2}+8a^{4}(\mathbf{r}_{1}^{2}+\mathbf{r}_{2}^{2})-6iBa^{2}t^{2}(\mathbf{r}_{1}^{2}-\mathbf{r}_{2}^{2})}{16a^{4}(a^{2}+Bt^{3})}\right\}.$$
 (75)

For $\mathbf{r}_1 = \mathbf{r}_2$ we recover the result Eq. (63). From the known $\Gamma(\mathbf{r}_1, \mathbf{r}_2, t)$, one can find the mean Wigner distribution density function which is defined as follows, [73]

$$W(\mathbf{p}, \mathbf{r}, t) = \frac{1}{(2\pi)^D} \int e^{i\mathbf{p}\cdot\mathbf{y}} \Gamma\left(\mathbf{r} + \frac{\mathbf{y}}{2}, \mathbf{r} - \frac{\mathbf{y}}{2}, t\right) d^D \mathbf{y},$$
(76)

and has numerous applications not only in statistical mechanics, but also in quantum and classical optics, signal processing theory etc. This quantity, as is well known, is real, but not necessarily positive everywhere. Provided that $W(\mathbf{p}, \mathbf{r}, t)$ is normalized to unity, this function is different from zero in a region of which the volume in phase space is at least equal to $(2\pi)^D$ and $W(\mathbf{p}, \mathbf{r}, t)$ can never be sharply localized in \mathbf{r} and \mathbf{p} . Inserting Eq. (75) into Eq. (76) and calculating integrals we have

$$W(\mathbf{p}, \mathbf{r}, t) = \frac{2^{D} A^{2} a^{3D}}{\pi^{D+1/2} [Bt(3a^{2} + 4a^{2}t^{2} + 3Bt^{3}/4) + 4a^{4}]^{D/2}} \times \exp\left\{-\frac{\mathbf{r}^{2}}{a^{2} + Bt^{3}} - \frac{4a^{4}(a^{2} + Bt^{3})}{[Bt(3a^{2} + 4a^{2}t^{2} + 3Bt^{3}/4) + 4a^{4}]} \left[\frac{3Bt^{2}}{4a^{2}(a^{2} + Bt^{3})}\mathbf{r} + \mathbf{p}\right]^{2}\right\}.$$
(77)

Let us consider the higher moment orders. We define the centroid of the structure by

$$\mathbf{r}_{c}(t) = \frac{1}{\langle N \rangle} \int \mathbf{r} I(\mathbf{r}, t) d^{D} \mathbf{r}.$$
 (78)

As we have seen from Eq. (68), the mean centroid $\mathbf{\bar{r}}(t) = \langle \mathbf{r}_c(t) \rangle$ have found to be equal to zero for both

fundamental and vortex soliton. It is interesting, however, to consider also the variance of the random function



FIG. 2: Time dependence of the variance of intensity at the center σ^2 of the fundamental soliton for $\lambda = 1$ and different values of *B* (noise intensity).

$$\mathbf{r}_{c}(t)$$

$$\left\langle [\mathbf{r}_{c}(t) - \bar{\mathbf{r}}(t)]^{2} \right\rangle = \int p(\mathbf{u}, t) [\mathbf{r}_{c}(t) - \bar{\mathbf{r}}(t)]^{2} d\mathbf{u}$$
$$= \frac{1}{\langle N \rangle^{2}} \int \int \mathbf{r}_{1} \cdot \mathbf{r}_{2} p(\mathbf{u}, t) I(\mathbf{r}_{1}, t) I(\mathbf{r}_{2}, t) d^{D} \mathbf{u} d^{D} \mathbf{r}_{1} d^{D} \mathbf{r}_{2}.$$
(79)

Integration in Eq. (79) taking into account Eqs. (42), (63) and (64) yields

$$\left\langle [\mathbf{r}_c(t) - \bar{\mathbf{r}}(t)]^2 \right\rangle = CBt^3,$$
 (80)

where C = 1 and C = 3/2 for the fundamental soliton in the 2D and 3D cases, respectively. For the vortex soliton, we have C = 2 and C = 27/8, respectively. The quantity (80) becomes unbounded as $t \to \infty$. This indicates the stochastic instability of these structures due to the presence of random fluctuations. Since most experiments measure the intensity and not the field itself, intensity fluctuations are of particular interest. An important characteristic in this case is the variance of the intensity

$$\sigma_I^2(\mathbf{r},t) \equiv \left\langle [I(\mathbf{r},t) - \langle I(\mathbf{r},t) \rangle]^2 \right\rangle = \left\langle I^2(\mathbf{r},t) \right\rangle - \left\langle I(\mathbf{r},t) \right\rangle^2.$$
(81)

The normalized variance $\sigma_I^2/\langle I \rangle^2$ is commonly referred to as the scintillation (flicker) index [5, 6]. The calculations can be carried out without difficulty and we have for the fundamental soliton

$$\sigma_I^2(\mathbf{r}, t) = A^4 \left\{ \frac{a^D \exp\left[-2r^2/(a^2 + 2Bt^3)\right]}{(a^2 + 2Bt^3)^{D/2}} - \frac{a^{2D} \exp\left[-2r^2/(a^2 + Bt^3)\right]}{(a^2 + Bt^3)^D} \right\},$$
(82)

and for the vortex soliton,

$$\sigma_{I}^{2}(\mathbf{r},t) = A^{4} \left\{ a^{D+4} \exp\left[-\frac{2r^{2}}{(a^{2}+2Bt^{3})}\right] \right\}$$

$$\times \frac{[a^{2}r_{\perp}^{2}+2Bt^{3}(a^{2}+2Bt^{3})]^{2}-2B^{2}t^{6}(a^{2}+2Bt^{3})^{2}}{(a^{2}+2Bt^{3})^{D/2+4}}$$

$$-a^{2(D+2)} \exp\left[-\frac{2r^{2}}{(a^{2}+Bt^{3})}\right] \frac{[a^{2}r_{\perp}^{2}+Bt^{3}(a^{2}+Bt^{3})]^{2}}{(a^{2}+Bt^{3})^{D+4}} \right\}$$
(83)

The variance of the intensity at the center of the fundamental and vortex solitons can be easily found from Eqs. (82) and (83), and have the form

$$\sigma_I^2(0,t) = A^4 \left\{ \frac{a^D}{(a^2 + 2Bt^3)^{D/2}} - \frac{a^{2D}}{(a^2 + Bt^3)^D} \right\}.$$
 (84)

and

$$\sigma_I^2(0,t) = A^4 \left\{ \frac{2a^{D+4}B^2t^6}{(a^2+2Bt^3)^{D/2+2}} - \frac{a^{2(D+2)}B^2t^6}{(a^2+Bt^3)^{D+2}} \right\}.$$
(85)

respectively. The function $\sigma_I^2(0, t)$ in both cases is a nonmonotonic function of time and reaches a maximum at some fixed moment of time, which increases with decreasing noise intensity B. In the limit $t \to \infty$, the variance of the intensity at the center tends to zero. Time dependence of the variance of intensity at the center of the fundamental soliton for $\lambda = 1$ and different values of Bis shown in Fig. 2.

VII. CONCLUSION

We have presented a new technique for calculating the even moments of the Nth order for the NLS equation with the spatial dimension $D \ge 2$ with a multiplicative random term depending on the time and space coordinates. The proposed formalism does not use closure procedures for the nonlinear term, as well as the smallness of the random term and the use of perturbation theory. Along with the original Eq. (1), we have considered an auxiliary stochastic nonlinear equation. We have derived equations (nonclosed) for the even moments of the Nth order both for the original and for the auxiliary (effective) equation. Assuming that the autocorrelation function of the random field is quadratic, we have shown that the even moments of the original equation and the effective one coincide. By a stochastic change of variables and a fundamental ansatz, we have demonstrated how the effective stochastic nonlinear equation can be mapped into an explicit deterministic equation for a new dependent variable and a system of stochastic nonlinear ordinary differential equations. From this system of Langevin-type equations, we have derived the corresponding Fokker-Planck equation for the joint probability density function and obtained its Gaussian-type solution. The even moments of the field are then expressed in terms of the

integrals of the product of the deterministic fields and the corresponding probability density function. As a specific example, we have used analytical expressions for the fundamental and vortex solitons, obtained using variational analysis. We have analytically calculated a number of statistical characteristics describing the dynamics of fundamental and vortex solitons in random medium, such as the mean intensities, the variance of the intensity, the centroid and spread of the structures, the spatial mutual coherence function etc. In particular, we have shown that, under the irreversible action of fluctuations, the solitons spread out, i.e., no collapse (beam self-focusing) occurs.

It is important to note that for specific realizations (there are an infinite number of them) of noise, the multidimensional fundamental and vortex solitons are unstable and collapse. In this regard, it is necessary to emphasize that the arrest of the collapse of the considered coherent structures is understood only in the sense of average values (more precisely, even moments) when, for example, analytical expressions for the average width of the structure clearly indicate spreading, and not collapse (beam self-focusing).

Note that the developed method is also applicable to odd moments, and of particular interest is the calculation of the first moment, that is, the mean field $\langle \varphi \rangle$. This requires a joint probability density function $p(\mathbf{u}, \mathbf{v}, \gamma)$. Then, in particular, we have for the mean field

$$\langle \varphi(\mathbf{r},t) \rangle = \int \varphi_{\mathbf{e}}(\mathbf{r},t) p(\mathbf{u},\mathbf{v},\gamma,t) d^{D} \mathbf{u} d^{D} \mathbf{v} d\gamma,$$
 (86)

where $\varphi_{\mathbf{e}}$ is determined by Eqs. (60) or (61). The equation for $p(\mathbf{u}, \mathbf{v}, \gamma)$ can readily be obtained (see Appendix). However, the analytical solution of the resulting equation seems to encounter significant difficulties.

The formalism developed in this paper can also be applied to other problems that can be formulated in terms of the model of the stochastic NLS equation (1). We emphasize, however, that this approach is applicable only for a very specific type (quadratic spatial dependence) of spatial noise correlator.

Appendix: derivation of the extended Fokker-Planck equation

The microscopic ("fine-grained") distribution function associated to the system (33)-(35), is

$$F(\mathbf{u}, \mathbf{v}, \gamma, t) = \delta[\mathbf{u} - \mathbf{u}(t)]\delta[\mathbf{v} - \mathbf{v}(t)]\delta[\gamma - \gamma(t)], \quad (87)$$

$$F(\mathbf{u}, \mathbf{v}, \gamma, 0) = \delta(\mathbf{u})\delta(\mathbf{v})\delta(\gamma).$$
(88)

The joint probability density function is defined as

 $p(\mathbf{u}, \mathbf{v}, \gamma, t) = \langle F(\mathbf{u}, \mathbf{v}, \gamma, t) \rangle.$ (89) Differentiating Eq. (87) with respect to t, we have the Liouville equation for F,

with the initial condition (88). Averaging of Eq. (90) gives

$$\left(\frac{\partial}{\partial t} + \mathbf{v}\frac{\partial}{\partial \mathbf{u}} + \frac{\mathbf{v}^2}{4}\frac{\partial}{\partial\gamma}\right)\langle F\rangle + \frac{1}{2}\left(\frac{\partial}{\partial\mathbf{v}} + \mathbf{u}\frac{\partial}{\partial\gamma}\right)\langle\mathbf{f}F\rangle = 0.$$
(91)

To determine $\langle \mathbf{f} F \rangle$, we use Donsker-Furutsu-Novikov formula (8),

$$\langle \mathbf{f}(t)F(\mathbf{u},\mathbf{v},\gamma,t)\rangle = \int_0^\infty \langle f_i(t)f_j(t')\rangle \left\langle \frac{\delta F(\mathbf{u},\mathbf{v},\gamma,t)}{\delta f_j(t')} \right\rangle dt'$$
(92)

The functional derivative $\langle \delta F / \delta f_j \rangle$ can be found from Eq. (90), and then using Eq. (20) one can obtain

$$\langle \mathbf{f}(t)F(\mathbf{u},\mathbf{v},\gamma,t)\rangle = -\frac{3B}{2} \left(\frac{\partial}{\partial \mathbf{v}} + \mathbf{u}\frac{\partial}{\partial \gamma}\right) \langle F(\mathbf{u},\mathbf{v},\gamma,t)\rangle.$$
(93)

Inserting this into Eq. (91), we finally get the Fokker-Planck equation,

$$\begin{bmatrix} \frac{\partial}{\partial t} + \mathbf{v} \frac{\partial}{\partial \mathbf{u}} + \frac{\mathbf{v}^2}{4} \frac{\partial}{\partial \gamma} - \frac{3B}{4} \left(\frac{\partial^2}{\partial \mathbf{v}^2} + 2\mathbf{u} \frac{\partial^2}{\partial \mathbf{v} \partial \gamma} + \mathbf{u}^2 \frac{\partial^2}{\partial \gamma^2} \right) \right] p(\mathbf{u}, \mathbf{v}, \gamma, t) = 0, \qquad (94)$$

with the initial condition $p(\mathbf{u}, \mathbf{v}, \gamma, 0) = \delta(\mathbf{u})\delta(\mathbf{v})\delta(\gamma)$.

 F. G. Bass, Y. S. Kivshar, V. V. Konotop, and Y. A. Sinitsyn, Dynamics of solitons under random perturbations, Phys. Rep. 157, 63-181 (1988).

[2] S. A. Gredeskul and Y. S. Kivshar, Propagation and scat-

- [3] V. V.Konotop and L. Vázquez, Nonlinear random waves (World Scientific Publishing, Singapore, 1994).
- [4] F. Abdullaev and J. Garnier, Optical solitons in random media, in *Progress in Optics Vol. 48*, edited by E. Wolf (Elsevier, New York, 2005) pp. 35-106.
- [5] V. Tatarskii, Wave propagation in a Turbulent Medium (Dover, New York, 1967).
- [6] A. Ishimaru, Wave propagation and Scattering in Random Media (Academic Press, New York, 1978).
- [7] I. M. Lifshits, S. A. Gredeskul, and L. A. Pastur, *Intro*duction to the Theory of Disorder Systems (JohnWiley & Sons, New York, 1988).
- [8] F. Kh. Abdullaev, Soliton collective excitations in stochastic molecular crystals, Theor. Math. Phys. 51, 607-610 (1982).
- [9] J. N. Elgin, Inverse scattering theory with stochastic initial potentials, Phys. Lett. A 110, 441-443 (1985).
- [10] J. P. Gordon and H. A. Haus, Random walk of coherently amplified solitons in optical fiber transmission, Opt. Lett. 11, 665-667 (1986).
- [11] B. A. Malomed, Emission from, quasi-classical quantization, and stochastic decay of sine-Gordon solitons in external fields, Physica D 27, 113-157 (1987).
- [12] F. G. Bass, Yu. S. Kivshar, V. V. Konotop and S. A. Puzenko, Propagation of nonlinear incoherent pulses in single-mode optical fibres, Opt. Commun. 68, 385-390 (1988).
- [13] Yu. S. Kivshar, S. A. Gredeskul, A. Sánchez and L. Vázquez, Localization decay induced by strong nonlinearity in disordered systems, Phys. Rev. Lett. 64, 1693-1696 (1990).
- [14] J. Garnier, Solitons in random media with long-range correlation, Waves Random Media 11, 149-162 (2001).
- [15] F. Kh. Abdullaev, D. V. Navotny, B. B. Baizakov, Optical pulse propagation in fibers with random dispersion, Physica D **192**, 83-94 (2004).
- [16] C. Conti, Solitonization of the Anderson localization, Phys. Rev. A 86, 061801(R) (2012).
- [17] E. Michaely and S. Fishman, Effective noise theory for the nonlinear Schrödinger equation with disorder, Phys. Rev. E 85, 046218 (2012).
- [18] G. Schwiete and A. M. Finkel'stein, Effective theory for the propagation of a wave packet in a disordered and nonlinear medium, Phys. Rev. A 87, 043636 (2013).
- [19] C. Weiss, S. L.Cornish, S. A.Gardiner, and H.-P.Breuer, Superballistic center-of-mass motion in one-dimensional attractive bose gases: Decoherence-induced gaussian random walks in velocity space, Phys. Rev. A 93, 013605 (2016).
- [20] Yu. S. Kivshar and B. A. Malomed, Dynamics of solitons in nearly integrable systems, Rev. Mod. Phys. 61, 763-915 (1989).
- [21] A. S. Pikovsky and D. L. Shepelyansky, Destruction of Anderson localization by a weak nonlinearity, Phys. Rev. Lett. 100, 094101 (2008).
- [22] S. Flach, D. O. Krimer, and Ch. Skokos, Universal spreading of wave packets in disordered nonlinear systems, Phys. Rev. Lett. **102**, 024101 (2009).
- [23] A. V. Milovanov and A. Iomin, Localizationdelocalization transition on a separatrix system of nonlinear Schrödinger equation with disorder, Europhys. Lett. 100, 10006 (2012).

- [24] A. Debussche and J. Printems, Numerical simulations of the stochastic Korteweg de Vries equation, Physica D 134, 200-226 (1999).
- [25] A. Debussche and L. D. Menza, Numerical simulation of focusing stochastic nonlinear Schrödinger equations, Physica D 162, 131-154 (2002).
- [26] V. Folli and C. Conti, Frustrated brownian motion of nonlocal solitary waves, Phys. Rev. Lett. 104, 193901 (2010).
- [27] F. Maucher, W. Krolikowski, and S. Skupin, Stability of solitary waves in random nonlocal nonlinear media, Phys. Rev. A 85, 063803 (2012).
- [28] M. S. Ruderman, Propagation of solitons of the derivative nonlinear Schrödinger equation in a plasma with fluctuating density, Phys. Plasmas 9, 2940-2945 (2002).
- [29] V. M. Lashkin, Alfvén soliton and emitted radiation in the presence of perturbations, Phys. Rev. E 74, 016603 (2006).
- [30] V. M. Lashkin, Perturbation theory for solitons of the Fokas-Lenells equation: Inverse scattering transform approach, Phys. Rev. E 103, 042203 (2021).
- [31] E V. Doktorov and I. S. Kuten, The Gordon-Haus effect for modified NLS solitons, Europhys. Lett. 53, 22 (2001).
- [32] V. M. Lashkin, Soliton of modified nonlinear Schrödinger equation with random perturbations, Phys. Rev. E 69, 016611 (2004).
- [33] V. A. Gorodtsov, Stochastic Kadomtsev–Petviashvili equation, JETP 90, 1105-1113 (2000).
- [34] R. Y. Chiao, E. Garmire, and C. H. Townes, Self-trapping of optical beams, Phys. Rev. Lett. 13, 479–482 (1964).
- [35] P. L. Kelley, Self-focusing of optical beams, Phys. Rev. Lett. 15, 1005-1008 (1965).
- [36] E. A. Kuznetsov, A. M. Rubenchik, and V. E. Zakharov, Soliton stability in plasmas and hydrodynamics, Phys. Rep. 142, 103-165 (1986).
- [37] V. E. Zakharov, Collapse of Langmuir waves, Sov. Phys. JETP 35, 908-914 (1972).
- [38] V. E. Zakharov and E. A. Kuznetsov, Solitons and collapses: two evolution scenarios of nonlinear wave systems, Physics-Uspekhi 55, 535-556 (2012).
- [39] G. Fibich, The Nonlinear Schrödinger Equation: Singular Solutions and Optical Collapse (Springer, Heidelberg, 2015).
- [40] L. Bergé, Wave collapse in physics: principles and applications to light and plasma waves, Phys. Rep. 303, 259-370 (1998).
- [41] G. A. Pasmanik, Self-interaction of incoherent light beams, Sov. Phys. JETP 39, 234-238 (1974).
- [42] J. R. Jokipii and J. Marburger, Homogeneity requirements for minimizing self-focusing damage by strong electromagnetic waves, Appl. Phys. Lett. 23, 696-698 (1973).
- [43] V. V. Vorob'ev, The broadening of a light beam in a nonlinear medium having random inhomogeneities of the refractive index, Izv. Vuz. Radiofiz. 13, 1053-1057 (1970).
- [44] S. N. Vlasov, V. A. Petrishchev, and V. I. Talanov, Averaged description of wave beams in linear and nonlinear medis (the method of moments), Radiophys. Quantum Electron. 14, 1062-1070 (1974).
- [45] Yu. B. Gaididei and P. L. Christiansen, Spatiotemporal collapse in a nonlinear waveguide with a randomly fluctuating refractive index, Opt. Lett. 23, 1090-1092 (1998).
- [46] B. K. Shivamoggi, L. C. Andrews, and R. L. Phillips, Nonlinear wave propagation in a random medium, Physica A 275, 86-98 (2000).

- [47] R. A. Van Gorder, δ -expansion method for nonlinear stochastic differential equations describing wave propagation in a random medium, Phys. Rev. E **82**, 056712 (2010).
- [48] B. Hafizi, J. R. Peñano, J. P. Palastro, R. P. Fischer, and G. DiComo, Laser beam self-focusing in turbulent dissipative media, Opt. Lett. 42, 298-301 (2017).
- [49] J. Peñano, J. P. Palastro, B. Hafizi, M. H. Helle, and G. P. DiComo, Self-channeling of high-power laser pulses through strong atmospheric turbulence, Phys. Rev. A 96, 013829 (2017).
- [50] Y. Deng, H. Wang, X. Ji, X. Li, H. Yu, and L. Chen, Characteristics of high-power partially coherent laser beams propagating upwards in the turbulent atmosphere, Opt. Express 28, 27927-27939 (2020).
- [51] J. Garnier, F. Kh. Abdullaev, and B. B. Baizakov, Collapse of a Bose-Einstein condensate induced by fluctuations of the laser intensity, Phys. Rev. A 69, 053607 (2004).
- [52] I. M. Besieris, Solitons in randomly inhomogeneous media, in *Nonlinear Electromagnetics*, edited by P. L. E. Uslenghi (Academic Press, New York, 1980) pp. 87-116.
- [53] T. J. Alexander and L. Bergé, Ground states and vortices of matter-wave condensates and optical guided waves, Phys. Rev. E 65, 026611 (2002).
- [54] Y. S. Kivshar and G. P. Agrawal, Optical Solitons: From Fibers to Photonic Crystals (Academic Press, San Diego, 2003).
- [55] A. V. Gurevich, Nonlinear Phenomena in the Ionosphere (Springer-Verlag, Berlin, 1978).
- [56] M. V. Goldman, Strong turbulence of plasma waves, Rev. Mod. Phys. 56, 709-735 (1984).
- [57] V. I. Yukalov, E. P. Yukalova, K. V. Krutitsky, and R. Graham, Bose-Einstein-condensed gases in arbitrarily strong random potentials, Phys. Rev. A 76, 053623 (2007).
- [58] Z.-Y. Sun and X. Yu, Transport of nonautonomous solitons in two-dimensional disordered media, Ann. Phys., 1600323 (2017).
- [59] K. Furutsu, On the statistical theory of electromagnetic waves in a fluctuating medium (i), J. Res. NBS 67D, 303-323 (1963).

- [60] M. D. Donsker, On function space integrals, in Analysis in Function Space, edited by W. T. Martin and I. Segal (M.I.T. Press, Cambridge, 1964) pp. 17-30.
- [61] E. A. Novikov, Functionals and the random-force method in turbulence theory, Sov. Phys. JETP 20, 1290-1294 (1965).
- [62] L. C. Lee and J. R. Jokipii, Strong scintillations in astrophysics. II. A theory of temporal broadening of pulses, Astrophys. J. 201, 532-543 (1975).
- [63] C. M. Rose and I. M. Besieris, Nth-order multifrequency coherence functions: A functional path integral approach, J. Math. Phys. 20, 1530-1538 (1979).
- [64] V. Bezák, Theory of randomly inhomogeneous waveguides with slight material dispersion: I. Single-mode waveguides, Waves Random Media 8, 351-367 (1998).
- [65] S. P. Novikov, S. V. Manakov, L. P. Pitaevski, and V. E. Zakharov, *Theory of Solitons: The Inverse Scattering Method* (Consultants Bureau, New York, 1984).
- [66] V. N. Serkin and T. L. Belyaeva, High-energy optical Schrödinger solitons, JETP Lett. 74, 573-577. (2001).
- [67] V. N. Serkin and A. Hasegawa, Exactly integrable nonlinear Schrödinger equation models with varying dispersion, nonlinearity and gain: Application for soliton dispersion managements, IEEE J. Sel. Top. Quantum Electron. 8, 418-431 (2002).
- [68] N. G. van Kampen, Stochastic Processes in Physics and Chemistry (North-Holland, Amsterdam, 1992).
- [69] C. Sulem and P.-L. Sulem, The Nonlinear Schrödinger Equation (Springer, New York, 1999).
- [70] V. M. Lashkin, A. S. Desyatnikov, E. A. Ostrovskaya, and Yu. S. Kivshar, Azimuthal vortex clusters in Bose-Einstein condensates, Phys. Rev. A 85, 013620 (2012).
- [71] B. A. Malomed, Vortex solitons: Old results and new perspectives, Physica D 399, 108-137 (2019).
- [72] H. Louis, M. Tlidi, and E. Louvergneaux, Experimental evidence of dynamical propagation for solitary waves in ultra slow stochastic non-local Kerr medium, Opt. Express 24, 16206-16211 (2016).
- [73] E. Wigner, On the quantum correction for thermodynamic equilibrium, Phys. Rev. 40, 749-759 (1932).