

# Spectral Determinants of Almost Equilateral Quantum Graphs

Jonathan Harrison<sup>1</sup> and Tracy Weyand<sup>2</sup>

<sup>1</sup>Department of Mathematics, Baylor University, 1410 S.4th Street,  
Waco, 76706, Texas, United States.

<sup>2</sup>Department of Mathematics, Rose-Hulman Institute of Technology,  
5500 Wabash Avenue, Terre Haute, 47803, Indiana, United States.

## Abstract

Kirchoff's matrix tree theorem of 1847 connects the number of spanning trees of a graph to the spectral determinant of the discrete Laplacian [6]. Recently an analogue was obtained for quantum graphs relating the number of spanning trees to the spectral determinant of a Laplacian acting on functions on a metric graph with standard (Neumann-like) vertex conditions [11]. This result holds for quantum graphs where the edge lengths are close together. A quantum graph where the edge lengths are all equal is called equilateral. Here we consider equilateral graphs where we perturb the length of a single edge (almost equilateral graphs). We analyze the spectral determinant of almost equilateral complete graphs, complete bipartite graphs, and circulant graphs. This provides a measure of how fast the spectral determinant changes with respect to changes in an edge length. We apply these results to estimate the width of a window of edge lengths where the connection between the number of spanning trees and the spectral determinant can be observed. The results suggest the connection holds for a much wider window of edge lengths than is required in [11].

**Keywords:** spectral graph theory, quantum graphs, spectral determinant, spanning trees

**MSC Classification:** 35A01 , 65L10 , 65L12 , 65L20 , 65L70

# 1 Introduction

A classical result in spectral graph theory is Kirchhoff's matrix tree theorem [6]. The number of spanning trees of a graph is given by the spectral determinant; the product of the non-zero eigenvalues of the graph's discrete Laplacian.

**Theorem 1** (Kirchhoff's Matrix Tree Theorem). *For a connected graph  $G$  with  $V$  vertices,*

$$\# \text{ spanning trees} = \frac{1}{V} \det'(\mathbf{L}) = \det(\mathbf{L}[i]) , \quad (1)$$

where  $i = 1, \dots, V$ . The  $V \times V$  matrix  $\mathbf{L} = \mathbf{D} - \mathbf{A}$  is the discrete Laplacian with  $\mathbf{D}$  the diagonal matrix of vertex degrees and  $\mathbf{A}$  the adjacency matrix of the graph. The  $(V - 1) \times (V - 1)$  matrix  $\mathbf{L}[i]$  is produced by deleting the  $i$ th row and column from  $\mathbf{L}$ .

In [11] the authors obtained a similar result for quantum graphs. A quantum graph is a metric graph along with a self-adjoint operator acting on functions defined on a set of intervals associated with the edges of the graph. Functions on these intervals are related by local boundary conditions at the vertices; see e.g. [2, 13] for an introduction to quantum graphs. The spectral determinant of the quantum graph Laplacian with standard (Neumann-like) vertex conditions determines the number of spanning trees.

**Theorem 2.** *Let  $\Gamma$  be a connected metric graph with  $V$  vertices and  $E$  edges whose edge lengths are in the interval  $[\ell, \ell + \delta)$ . If*

$$\delta < \frac{\ell}{V^V 2^{E+V} \sqrt{2EV}} , \quad (2)$$

then the number of spanning trees is the closest integer to

$$T_\Gamma = \frac{\prod_{v \in \mathcal{V}} d_v}{E 2^E \ell^{\beta+1}} \det'(\mathcal{L}) , \quad (3)$$

where  $\mathcal{V}$  is the set of vertices,  $d_v$  is the degree of vertex  $v$ ,  $\mathcal{L}$  is the Laplacian with standard vertex conditions, and  $\beta = E - V + 1$  is the first Betti number of  $\Gamma$ , the number of independent cycles.

To obtain the result, the edge lengths of the metric graph are tightly constrained so they are all close to  $\ell$ . We call a metric graph where the edge lengths are all equal an *equilateral graph*. Hence the result holds for metric graphs that are close to an equilateral graph. However, heuristically, the spectral determinant of a quantum graph should determine the number of spanning trees for graphs where the edge lengths are less tightly constrained. In this article we approach this relationship from a different direction. We consider three classes of equilateral graphs where we can compute the spectral determinant analytically; fully connected graphs, fully connected bipartite graphs, and some circulant graphs. We then consider perturbing the length of a single edge to obtain an almost equilateral graph, and we ascertain how the spectral determinant and the number of spanning trees changes under the perturbation. This provides a measure of how fast  $T_\Gamma$  varies as one changes an edge length. In each case, we see that  $T_\Gamma$  can be expected to provide an accurate determination of the number of spanning trees for graphs where the edge lengths vary much more widely than is implied by the constraint (2) on  $\delta$  imposed in theorem 2. For example, the bound

$\delta < \ell/V^V$  can be expected to provide a sufficiently narrow window of edge lengths for  $T_\Gamma$  to determine the number of spanning trees for these classes of graphs. The spectral determinant was formulated for Laplace and Schrödinger operators on metric graphs in [4, 5, 7], via periodic orbits in [1], and using the zeta function in [8, 9, 12].

The article is laid out as follows. In section 2 we introduce notation for graphs and quantum graphs and define the corresponding spectral determinants. Sections 3, 4 and 5 analyze the spectral determinants of almost equilateral complete graphs, complete bipartite graphs, and certain connected circulant graphs respectively. Finally, in section 6 we apply the results for the spectral determinant to see how the function  $T_\Gamma$  behaves under perturbation of an edge length.

## 2 Background

A *graph*  $G$  consists of a set of vertices  $\mathcal{V}$  and a set of edges  $\mathcal{E}$  such that each edge  $e = (u, v) \in \mathcal{E}$  connects a pair of vertices  $u, v \in \mathcal{V}$ ; see figures 1 and 2 for examples. Two vertices are *adjacent* if there is an edge  $(u, v) \in \mathcal{E}$ , denoted  $u \sim v$ . The number of vertices is  $V = |\mathcal{V}| < \infty$ , and  $E = |\mathcal{E}| < \infty$  is the number of edges. The *degree* of vertex  $v$  is the number of vertices adjacent to  $v$ , denoted  $d_v$ . The graph is *k-regular* if  $d_v = k$  for all  $v \in \mathcal{V}$ . Throughout the paper, we assume that graphs are *connected*, so there is a path between every pair of vertices. We also assume the graphs are *simple*, with no loops or multiple edges. The *first Betti number* is  $\beta = E - V + 1$ , the number of independent cycles of  $G$ . A *tree* is a connected graph with no cycles ( $\beta = 0$ ). A *spanning tree* of  $G$  is a subgraph that is a tree and contains all vertices of  $G$ . Additionally, when we reference sets of vectors and specific elements of vectors, we use the convention that  $\mathbf{v}_j$  denotes the  $j$ th vector in the set  $\{\mathbf{v}_j\}_{j=1}^n$  whereas  $[\mathbf{v}]_j$  denotes the  $j$ th element of the vector  $\mathbf{v}$ .

We now define some classes of graphs that will be used in the article. The *fully connected graph* on  $n$  vertices  $K_n$  is the graph where every pair of vertices is connected by an edge so  $(u, v) \in \mathcal{E}$  for all  $u, v \in \mathcal{V}$  with  $u \neq v$ . The *complete bipartite graph*  $K_{m,n}$  is a graph where the set of vertices is divided into two disjoint sets,  $\mathcal{V} = \mathcal{A} \cup \mathcal{B}$  with  $\mathcal{A} \cap \mathcal{B} = \emptyset$ . Every vertex of  $\mathcal{A}$  is adjacent to every vertex of  $\mathcal{B}$  and there are no edges connecting pairs of vertices in  $\mathcal{A}$  or pairs of vertices in  $\mathcal{B}$ . The number of vertices in  $\mathcal{A}$  is  $m = |\mathcal{A}|$  and  $n = |\mathcal{B}|$ . Figure 1 shows the complete bipartite graph  $K_{3,5}$ . A *circulant graph* on  $n$  vertices is denoted  $C_n(\mathbf{a})$  where  $\mathbf{a} = (a_1, \dots, a_p)$  with  $a_1 < a_2 < \dots < a_p \leq n/2$ . Then two vertices  $i, j \in \{0, \dots, n-1\}$  are connected if  $|i - j| = a_k \pmod{n}$  for some  $k \in \{1, \dots, p\}$ . If  $a_p < n/2$ , then the circulant graph is  $2p$ -regular. A circulant graph is connected if and only if  $\gcd(n, a_1, \dots, a_p) = 1$ . Figure 2 shows the connected circulant graph  $C_{17}(2, 5)$ .

### 2.1 Quantum Graphs

A *metric graph*  $\Gamma$  is a graph where each edge  $e \in \mathcal{E}$  is assigned a length  $\ell_e > 0$ . The *total length* of a metric graph is  $\ell_{tot} = \sum_{e \in \mathcal{E}} \ell_e$ . We view the edge  $e = (u, v) \in \mathcal{E}$  as the interval  $[0, \ell_e]$  with  $x_e = 0$  at  $u$  and  $x_e = \ell_e$  at  $v$ ; the choice of orientation will not effect our results. This allows us to define a function  $f$  on  $\Gamma$  as a collection of functions,  $\{f_e\}_{e \in \mathcal{E}}$ , such that the function  $f_e$  is defined on the interval  $[0, \ell_e]$ . A metric graph is




Fig. 1 The complete bipartite graph  $K_{3,5}$ .




Fig. 2 The circulant graph  $C_{17}(2, 5)$ .

*equilateral* if all the edges have the same length. We use  $\Gamma$  to denote an equilateral graph. If  $\Gamma$  is equilateral with edge lengths  $\ell$ , then we will say that  $\Gamma'$  is an *almost equilateral graph* if the length of one edge  $e$  of  $\Gamma$  is perturbed so  $\ell_e = \ell + \epsilon$ .

A *quantum graph* is a metric graph  $\Gamma$  along with a self-adjoint differential operator. Our focus in this paper is on the *Laplace operator*, which acts as  $-\frac{d^2}{dx_e^2}$  on functions that are defined on the interval  $[0, \ell_e]$ . We consider standard Neumann-Kirchhoff conditions at the vertices:

$$\begin{cases} f \text{ is continuous on } \Gamma \text{ and} \\ \sum_{e \in \mathcal{E}_v} \frac{df_e}{dx_e}(v) = 0 \text{ at each vertex } v, \end{cases} \quad (4)$$

where  $\mathcal{E}_v$  denotes the edges containing the vertex  $v$  and the derivative  $\frac{df_e}{dx_e}(v)$  is taken in the outgoing direction at  $v$ . The second Sobolev space on  $\Gamma$  is the direct sum of second Sobolev spaces on the set of intervals:

$$H^2(\Gamma) = \bigoplus_{e \in \mathcal{E}} H^2([0, \ell_e]) . \quad (5)$$

The domain of the Laplace operator is the set of functions  $f \in H^2(\Gamma)$  that satisfy (4). With these conditions, the Laplace operator is self-adjoint with infinitely many

non-negative eigenvalues. We use the notation  $\mathcal{L}$  and  $\mathcal{L}'$  for the Laplace operator on an equilateral graph  $\Gamma$  and a corresponding almost equilateral graph  $\Gamma'$  respectively.

Our first goal is to see how the spectral determinant of an almost equilateral graph,  $\det'(\mathcal{L}')$ , is related to the spectral determinant of an equilateral graph,  $\det'(\mathcal{L})$ . To do this, we employ the following result from [7].

**Theorem 3.** *Given a connect metric graph,*

$$\det'(\mathcal{L}) = \frac{2^E \ell_{tot} \prod_{e \in \mathcal{E}} \ell_e}{\prod_{v \in \mathcal{V}} d_v} \det'(\mathbf{R}) \quad (6)$$

where  $\mathbf{R}$  is the  $V \times V$  matrix,

$$[\mathbf{R}]_{u,v} = \begin{cases} \sum_{w \sim v} \ell_{(w,v)}^{-1}, & \text{if } u = v \\ -\ell_{(u,v)}^{-1}, & \text{if } u \sim v \\ 0, & \text{otherwise.} \end{cases} \quad (7)$$

Let  $\Gamma$  be an equilateral metric graph. In this case, the matrix  $\mathbf{R} = \ell^{-1}\mathbf{L}$  in theorem 3 is

$$[\mathbf{R}]_{u,v} = \begin{cases} \ell^{-1}d_v, & \text{if } u = v \\ -\ell^{-1}, & \text{if } u \sim v \\ 0, & \text{otherwise,} \end{cases} \quad (8)$$

where  $\mathbf{L}$  is the *combinatorial Laplacian matrix*;  $\mathbf{L} = \mathbf{D} - \mathbf{A}$  where  $\mathbf{D}$  is the diagonal matrix of vertex degrees with  $[\mathbf{D}]_{v,v} = d_v$  and  $\mathbf{A}$  is the adjacency matrix with  $[\mathbf{A}]_{u,v} = 1$  if  $u \sim v$  and 0 otherwise. Note that, as the edges are not directed, the adjacency matrix is by definition symmetric. We now form a perturbed graph,  $\Gamma'$ , by changing the length of one edge  $e = (a, b) \in \mathcal{E}$  to  $\ell + \epsilon$ . Then,

$$[\mathbf{R}']_{u,v} = \begin{cases} \ell^{-1}d_v, & \text{if } u = v \neq a \text{ or } b \\ -\ell^{-1}, & \text{if } (u, v) \in \mathcal{E} - (a, b) \\ \ell^{-1}d_v - \ell^{-2}\epsilon + \mathcal{O}(\epsilon^2), & \text{if } u = v = a \text{ or } b \\ -\ell^{-1} + \ell^{-2}\epsilon + \mathcal{O}(\epsilon^2), & \text{if } (u, v) = (a, b) \\ 0, & \text{otherwise.} \end{cases} \quad (9)$$

Equivalently,

$$\mathbf{R}' = \mathbf{R} + \mathbf{M} + \mathcal{O}(\mathbf{M}^2) = \mathbf{R} - \frac{\epsilon}{\ell^2} \mathbf{Q} + \mathcal{O}(\epsilon^2), \quad (10)$$

where  $\mathbf{M} = -\frac{\epsilon}{\ell^2} \mathbf{Q}$  and

$$[\mathbf{Q}]_{u,v} = \begin{cases} 1, & \text{if } u = v = a \text{ or } b \\ -1, & \text{if } (u, v) = (a, b) \\ 0, & \text{otherwise.} \end{cases} \quad (11)$$

## 2.2 Perturbation Theory

We use the following propositions to relate the eigenvalues of  $\mathbf{R}$  and  $\mathbf{R}'$ ; see e.g. [14].

**Proposition 4.** *If the matrix  $\mathbf{R}$  has a non-degenerate eigenvalue  $\lambda$  with normalized eigenvector  $\mathbf{v}$  then  $\mathbf{R}'$  has a non-degenerate eigenvalue*

$$\lambda' = \lambda - \frac{\epsilon}{\ell^2} \mathbf{v} \cdot \mathbf{Q} \mathbf{v} + \mathcal{O}(\epsilon^2) . \quad (12)$$

For example, assume that all the eigenvalues of the matrix  $\mathbf{R}$  are simple. Let  $\{\mathbf{v}_k\}_{k=1}^V$  be the set of orthonormal eigenvectors corresponding to the eigenvalues  $\{\lambda_k\}_{k=1}^V$ . Using (11), one can see that

$$\mathbf{v}_k \cdot \mathbf{Q} \mathbf{v}_k = ([\mathbf{v}_k]_a - [\mathbf{v}_k]_b)^2, \quad (13)$$

and hence, by proposition 4, the eigenvalues of  $\mathbf{R}'$  are

$$\lambda'_k = \lambda_k - \frac{\epsilon}{\ell^2} ([\mathbf{v}_k]_a - [\mathbf{v}_k]_b)^2 + \mathcal{O}(\epsilon^2) . \quad (14)$$

We use the following proposition if an eigenvalue  $\lambda$  is degenerate.

**Proposition 5.** *Suppose that  $\lambda$  is an eigenvalue of  $\mathbf{R}$  with multiplicity  $n$  and corresponding orthonormal eigenvectors  $\{\mathbf{u}_k\}_{k=1}^n$ . Let  $\mathbf{Q}'$  be the  $n \times n$  matrix defined by,*

$$[\mathbf{Q}']_{j,k} = \mathbf{u}_j \cdot \mathbf{Q} \mathbf{u}_k . \quad (15)$$

*Then  $n$  of the eigenvalues of  $\mathbf{R}'$  are*

$$\lambda'_k = \lambda - \frac{\epsilon}{\ell^2} \lambda_k + \mathcal{O}(\epsilon^2) , \quad (16)$$

*where  $\lambda_k$  is the  $k$ th eigenvalue of  $\mathbf{Q}'$ .*

## 3 Almost Equilateral Complete Graphs

Consider the complete equilateral graph  $K_n$ . The matrix  $\mathbf{R}$  has eigenvalues 0 and  $n/\ell$  (with multiplicity  $n - 1$ ). Without loss of generality, suppose that the length of the edge connecting vertices 1 and  $n$  is increased from  $\ell$  to  $\ell + \epsilon$ . Notice that  $\{\mathbf{u}_k\}_{k=1}^{n-1}$  with,

$$[\mathbf{u}_k]_j = \frac{1}{\sqrt{k(k+1)}} \begin{cases} 1, & \text{if } j \leq k \\ -k, & \text{if } j = k + 1 \\ 0, & \text{otherwise} \end{cases} \quad (17)$$

is a set of  $n - 1$  orthonormal eigenvectors of  $\mathbf{R}$  corresponding to the eigenvalue  $n/\ell$ . Then, from equation (15), the  $(n - 1) \times (n - 1)$  matrix  $\mathbf{Q}'$  has entries

$$[\mathbf{Q}']_{j,k} = \mathbf{u}_j \cdot \mathbf{Q}\mathbf{u}_k = \begin{cases} \frac{1}{\sqrt{j(j+1)k(k+1)}}, & \text{if } j, k \neq n - 1 \\ \frac{n}{\sqrt{j(j+1)k(k+1)}}, & \text{if either } j \text{ or } k = n - 1 \\ \frac{n}{n-1}, & \text{if } j = k = n - 1. \end{cases} \quad (18)$$

Define  $n - 3$  vectors  $\{\mathbf{v}_k\}_{k=2}^{n-2}$  via,

$$[\mathbf{v}_k]_j = \begin{cases} -\frac{1}{\sqrt{k(k+1)/2}}, & \text{if } j = 1 \\ 1, & \text{if } j = k \\ 0, & \text{otherwise,} \end{cases} \quad (19)$$

and a vector  $\mathbf{v}_{n-1}$  with,

$$[\mathbf{v}_{n-1}]_j = \begin{cases} -\frac{1}{\sqrt{n(n-1)/2}}, & \text{if } j = 1 \\ \frac{1}{n}, & \text{if } j = n - 1 \\ 0, & \text{otherwise.} \end{cases} \quad (20)$$

One can verify that  $\{\mathbf{v}_k\}_{k=2}^{n-1}$  are  $n - 2$  linearly independent eigenvectors of  $\mathbf{Q}'$  with eigenvalue 0. Similarly, define  $\mathbf{v}$  via,

$$[\mathbf{v}]_j = \begin{cases} \frac{1}{\sqrt{j(j+1)/2}}, & \text{if } j \neq n - 1 \\ \frac{n}{\sqrt{n(n-1)/2}}, & \text{if } j = n - 1. \end{cases} \quad (21)$$

Then  $\mathbf{v}$  is an eigenvector of  $\mathbf{Q}'$  with eigenvalue 2. Hence, applying proposition 5, the eigenvalue  $n/\ell$  has degeneracy  $n - 2$  (up to first-order in  $\epsilon$ ) and there is a new simple eigenvalue,

$$\frac{n}{\ell} - \frac{2\epsilon}{\ell^2} + \mathcal{O}(\epsilon^2).$$

The remaining eigenvalue of  $\mathbf{R}$  is  $\lambda = 0$  with normalized eigenvector  $\mathbf{u}$  where,

$$[\mathbf{u}]_j = \frac{1}{\sqrt{n}} \quad \text{for } j = 1, 2, \dots, n. \quad (22)$$

Hence  $[\mathbf{Q}\mathbf{u}]_j = 0$  for all  $j = 1, \dots, n$  and  $\lambda' = 0 + \mathcal{O}(\epsilon^2)$ .

Combining these results,

$$\det'(\mathbf{R}') = \prod_{j=2}^n \lambda'_j = \left(\frac{n}{\ell}\right)^{n-1} - \epsilon \frac{2n^{n-2}}{\ell^n} + \mathcal{O}(\epsilon^2). \quad (23)$$

Therefore, from Theorem 3, the spectral determinant of the almost equilateral complete graph  $K_n$  is,

$$\det'(\mathcal{L}') = \frac{E2^E n^{n-2} \ell^{\beta+1}}{(n-1)^n} \left( 1 + \frac{\epsilon}{\ell} \left( 1 - \frac{2(n-2)}{n(n-1)} \right) \right) + \mathcal{O}(\epsilon^2), \quad (24)$$

where  $E = n(n-1)/2$  and  $\beta = E - n + 1$ .

## 4 Almost Equilateral Complete Bipartite Graphs

Consider a complete bipartite quantum graph  $K_{m,n}$  where each edge has length  $\ell$ . Let  $A = \{1, 2, \dots, m\}$  and  $B = \{m+1, m+2, \dots, m+n\}$  where all vertices of  $A$  are adjacent to all vertices of  $B$ ; figure 1 shows the complete bipartite graph  $K_{3,5}$ . For such a graph, from equation (8), we have,

$$[\mathbf{R}]_{u,v} = \begin{cases} \ell^{-1}n, & u = v \in A \\ \ell^{-1}m, & u = v \in B \\ -\ell^{-1}, & u \sim v \text{ (so } u \in A \text{ and } v \in B \text{ or vice versa)} \\ 0, & \text{otherwise.} \end{cases} \quad (25)$$

The eigenvalues of  $\mathbf{R} = \ell^{-1}\mathbf{L}$  are  $\ell^{-1}(m+n)$ ,  $\ell^{-1}n$  (with multiplicity  $m-1$ ),  $\ell^{-1}m$  (with multiplicity  $n-1$ ) and 0. Therefore

$$\det'(\mathbf{R}) = \frac{(m+n)m^{n-1}n^{m-1}}{\ell^{m+n-1}}. \quad (26)$$

Without loss of generality, we perturb the length of the edge  $(1, m+1)$  and apply propositions 4 and 5 to obtain the following proposition for the spectrum of  $\mathbf{R}'$ , which is proven in the following two subsections.

**Proposition 6.** *Let  $\Gamma = K_{m,n}$  be an equilateral complete bipartite graph with edge length  $\ell$ . Suppose a new graph  $\Gamma'$  is formed by increasing the length of one edge to  $\ell + \epsilon$ . Up to first-order in  $\epsilon$ , the eigenvalues of the matrix  $\mathbf{R}'$  are 0,  $\ell^{-1}(m+n) - \epsilon\ell^{-2} \left( \frac{m+n}{mn} \right)$ ,  $\ell^{-1}n$  (with multiplicity  $m-2$ ),  $\ell^{-1}m$  (with multiplicity  $n-2$ ),  $\ell^{-1}n - \epsilon\ell^{-2} \left( \frac{m-1}{m} \right)$ , and  $\ell^{-1}m - \epsilon\ell^{-2} \left( \frac{n-1}{n} \right)$ .*

### 4.1 Non-degenerate Eigenvalues

First, consider the eigenvalue  $\lambda = \ell^{-1}(m+n)$  with normalized eigenvector  $\mathbf{v}$  where,

$$[\mathbf{v}]_j = \frac{1}{\sqrt{mn(m+n)}} \begin{cases} n, & \text{if } 1 \leq j \leq m \\ -m, & \text{if } m+1 \leq j \leq m+n. \end{cases} \quad (27)$$

Applying proposition 4,

$$\lambda' = \frac{m+n}{\ell} - \frac{\epsilon}{\ell^2} \mathbf{v} \cdot \mathbf{Q}\mathbf{v} + \mathcal{O}(\epsilon^2) \quad (28)$$



$$= \frac{m+n}{\ell} - \frac{\epsilon}{\ell^2} \frac{m+n}{mn} + \mathcal{O}(\epsilon^2). \quad (29)$$

Similarly, the eigenvalue  $\lambda = 0$  has normalized eigenvector  $\mathbf{u}$  with,

$$[\mathbf{u}]_j = \frac{1}{\sqrt{m+n}} \quad \text{for } j = 1, 2, \dots, m+n. \quad (30)$$

Since  $[\mathbf{Q}\mathbf{u}]_j = 0$ , we see that  $\lambda' = 0 + \mathcal{O}(\epsilon^2)$  using proposition 4.

## 4.2 Degenerate Eigenvalues

Consider the eigenvalue  $\ell^{-1}n$  with  $m-1$  orthonormal eigenvectors  $\{\mathbf{v}_k\}_{k=1}^{m-1}$  where,

$$[\mathbf{v}_k]_j = \frac{1}{\sqrt{k(k+1)}} \begin{cases} 1, & \text{for } j \leq k \\ -k, & \text{for } j = k+1 \\ 0, & \text{for } j > k+1. \end{cases} \quad (31)$$

Then,

$$[\mathbf{Q}']_{j,k} = \mathbf{v}_j \cdot \mathbf{Q}\mathbf{v}_k = \frac{1}{\sqrt{j(j+1)k(k+1)}}. \quad (32)$$

**Lemma 7.** *The eigenvalues of  $\mathbf{Q}'$  are 0 (with multiplicity  $m-2$ ) and  $\frac{m-1}{m}$ .*

*Proof.* Consider the vectors  $\{\mathbf{u}_k\}_{k=2}^{m-1}$  of length  $m-1$  such that

$$[\mathbf{u}_k]_j = \begin{cases} -\frac{1}{\sqrt{k(k+1)/2}}, & j = 1 \\ 1, & j = k \\ 0, & \text{otherwise.} \end{cases} \quad (33)$$

Multiplying  $\mathbf{u}_k$  by row  $j$  of  $\mathbf{Q}'$ , we have

$$[\mathbf{Q}'\mathbf{u}_k]_j = -\frac{1}{\sqrt{j(j+1)k(k+1)}} + \frac{1}{\sqrt{j(j+1)k(k+1)}} = 0. \quad (34)$$

Hence zero is an eigenvalue of  $\mathbf{Q}'$  with multiplicity  $m-2$ .

Similarly, define  $\mathbf{w}$  such that

$$[\mathbf{w}]_j = \frac{1}{\sqrt{j(j+1)/2}} \quad \text{for } j = 1, \dots, m-1. \quad (35)$$

Then,

$$[\mathbf{Q}'\mathbf{w}]_j = \frac{1}{\sqrt{j(j+1)/2}} \sum_{k=1}^{m-1} \frac{1}{k(k+1)} = \left(1 - \frac{1}{m}\right) [\mathbf{w}]_j. \quad (36)$$

Hence,  $1 - 1/m$  is the remaining eigenvalue of  $\mathbf{Q}'$ .  $\square$

Similarly,  $\ell^{-1}m$  is an eigenvalue of  $\mathbf{R}$  with multiplicity  $n - 1$  and the eigenvalues of a corresponding matrix  $\mathbf{Q}'$  are 0 (with multiplicity  $n - 2$ ) and  $1 - 1/n$ . Combining the results in subsections 4.1 and 4.2 with proposition 5 produces proposition 6.

### 4.3 Spectral Determinant

The spectral determinant of  $\mathbf{R}'$  is the product of its non-zero eigenvalues. From proposition 6, for an almost equilateral complete bipartite graph,

$$\det'(\mathbf{R}') = \det'(\mathbf{R}) - \frac{\epsilon}{\ell^{m+n}} n^{m-2} m^{n-2} (m+n)(m+n-1) + \mathcal{O}(\epsilon^2). \quad (37)$$

Therefore, using equation (26) and theorem 3 with  $E = mn$  and  $V = m + n$ , the spectral determinant of the almost equilateral complete bipartite graph  $K_{m,n}$  is,

$$\det'(\mathcal{L}') = 2^E \ell^{\beta+1} \left( 1 + \frac{\epsilon}{\ell} \left( 1 - \frac{V-2}{E} \right) \right) + \mathcal{O}(\epsilon^2). \quad (38)$$

#### 4.3.1 Star Graph

For example, a *star graph* with  $n + 1$  vertices is the complete bipartite graph  $K_{n,1}$  and  $E = V - 1$ . Suppose this is an equilateral graph with edge length  $\ell$ . The eigenvalues of  $\mathbf{R}$  are  $\ell^{-1}$  (with multiplicity  $n - 1$ ), 0 and  $\ell^{-1}(n + 1)$ . Then, perturbing one edge,

$$\det'(\mathbf{R}') = \frac{V}{\ell^{V-1}} - \epsilon \frac{V}{\ell^V} + \mathcal{O}(\epsilon^2). \quad (39)$$

From (38),

$$\det'(\mathcal{L}') = 2^E \ell + \frac{\epsilon}{\ell E} + \mathcal{O}(\epsilon^2). \quad (40)$$

## 5 Almost Equilateral Circulant Graphs

Consider a  $2p$ -regular circulant graph  $C_n(\mathbf{a})$  with  $\mathbf{a} = (a_1, a_2, \dots, a_p)$  and  $a_1 < a_2 < \dots < a_p < n/2$ ; see e.g. figure 2. For an equilateral circulant graph, the matrix  $\mathbf{R}$  is a circulant matrix with entries

$$[\mathbf{R}]_{u,v} = \begin{cases} 2p\ell^{-1}, & \text{if } u = v \\ -\ell^{-1}, & \text{if } |u - v| = a_k \pmod{n} \\ 0, & \text{otherwise.} \end{cases} \quad (41)$$

Eigenvalues of an  $n \times n$  circulant matrix with first row,

$$[c_0 \ c_1 \ c_2 \ \dots \ c_{n-1}] \quad (42)$$

are

$$\lambda_j = c_0 + c_1 \omega^j + c_2 \omega^{2j} + \dots + c_{n-1} \omega^{(n-1)j}, \quad (43)$$

with corresponding eigenvectors,

$$\mathbf{v}_j = \frac{1}{\sqrt{n}} [1, \omega^j, \omega^{2j}, \dots, \omega^{(n-1)j}]^T \quad (44)$$

for  $j = 0, 1, \dots, n-1$ , where  $\omega = e^{\frac{2\pi i}{n}}$  is a primitive  $n$ th root of unity [3]. We take  $n$  to be prime which guarantees that the graph is connected and  $\omega^j$  is a primitive root of unity for  $j = 1, \dots, n-1$ . Therefore the eigenvalues of  $\mathbf{R}$  are 0 (with multiplicity 1) and  $\lambda_j = 2p\ell^{-1} - 2\ell^{-1} \sum_{k=1}^p \cos\left(\frac{2\pi j a_k}{n}\right)$ . We notice that  $\lambda_j = \lambda_{n-j}$  for  $0 < j < (n-1)/2$ , so these eigenvalue have a multiplicity of at least two. Hence,

$$\det'(\mathbf{R}) = \left(\frac{2}{\ell}\right)^{n-1} \prod_{j=1}^{(n-1)/2} \left(p - \sum_{k=1}^p \cos\left(\frac{2\pi j a_k}{n}\right)\right)^2. \quad (45)$$

In the following, we will assume that the vector  $\mathbf{a}$  is chosen so the multiplicity of the non-zero eigenvalues is exactly two. This is the generic situation and is the case, for example, when  $p = 2$  and  $n \nmid (a_1^2 + a_2^2)$  [10]. We now consider how the eigenvalues change in an almost equilateral graph.

### 5.1 The Zero Eigenvalue

Zero is the only non-degenerate eigenvalue of  $\mathbf{R}$  with a normalized eigenvector,

$$\mathbf{v}_0 = \frac{1}{\sqrt{n}} [1, \dots, 1]^T. \quad (46)$$

Then, from proposition 4, the corresponding eigenvalue of  $\mathbf{R}'$  is,

$$\lambda'_0 = 0 - \frac{\epsilon}{\ell^2} \mathbf{v}_0 \cdot \mathbf{Q} \mathbf{v}_0 + \mathcal{O}(\epsilon^2) = 0 + \mathcal{O}(\epsilon^2) \quad (47)$$

since  $\mathbf{Q} \mathbf{v}_0 = \mathbf{0}$ .

### 5.2 Eigenvalues with Multiplicity Two

Suppose  $j = 1, \dots, (n-1)/2$  so  $\lambda_j = \lambda_{n-j}$  with corresponding eigenvectors,

$$\mathbf{v}_j = \frac{1}{\sqrt{n}} [1, \omega^j, \omega^{2j}, \dots, \omega^{(n-1)j}]^T, \quad (48)$$

$$\mathbf{v}_{n-j} = \frac{1}{\sqrt{n}} [1, \omega^{-j}, \omega^{-2j}, \dots, \omega^{-(n-1)j}]^T. \quad (49)$$

Without loss of generality, suppose edge  $e = (0, a_m)$  has its length perturbed. Then,

$$[\mathbf{Q} \mathbf{v}_j]_k = \frac{1}{\sqrt{n}} \begin{cases} 1 - \omega^{a_m j}, & \text{if } k = 0 \\ \omega^{a_m j} - 1, & \text{if } k = a_m \\ 0, & \text{otherwise.} \end{cases} \quad (50)$$

Therefore,

$$\mathbf{v}_j \cdot \mathbf{Q}\mathbf{v}_j = \frac{2}{n} \omega^{a_m j} (\cos \alpha_{m,j} - 1) , \quad (51)$$

$$\mathbf{v}_{n-j} \cdot \mathbf{Q}\mathbf{v}_j = -\frac{2}{n} (\cos \alpha_{m,j} - 1) \quad (52)$$

where

$$\alpha_{m,j} = \frac{2\pi a_m j}{n} . \quad (53)$$

Similarly,

$$\mathbf{v}_{n-j} \cdot \mathbf{Q}\mathbf{v}_{n-j} = \frac{2}{n} \omega^{-a_m j} (\cos \alpha_{m,j} - 1) , \quad (54)$$

$$\mathbf{v}_j \cdot \mathbf{Q}\mathbf{v}_{n-j} = -\frac{2}{n} (\cos \alpha_{m,j} - 1) \quad (55)$$

since  $\cos \alpha_{m,n-j} = \cos \alpha_{m,j}$ . Hence,

$$\mathbf{Q}' = \frac{2(\cos \alpha_{m,j} - 1)}{n} \begin{bmatrix} \omega^{a_m j} & -1 \\ -1 & \omega^{-a_m j} \end{bmatrix} . \quad (56)$$

The eigenvalues of the matrix in square brackets are 0 and  $2 \cos \alpha_{m,j}$ . So, by proposition 5, the eigenvalues of  $\mathbf{R}'$  are

$$\lambda'_j = \lambda_j + \mathcal{O}(\epsilon^2) , \quad (57)$$

$$\lambda'_{n-j} = \lambda_j + \frac{\epsilon}{\ell^2} \frac{4 \cos \alpha_{m,j} (\cos \alpha_{m,j} - 1)}{n} + \mathcal{O}(\epsilon^2) \quad (58)$$

for  $j = 1, \dots, (n-1)/2$ .

### 5.3 Spectral Determinant

With  $n$  prime and  $\mathbf{a}$  chosen so that the eigenvalues of  $\mathbf{R}$  have at most multiplicity two, the spectral determinant of  $\mathbf{R}'$  is,

$$\det'(\mathbf{R}') = \det'(\mathbf{R}) + \frac{\epsilon}{\ell^2} \sum_{j=1}^{(n-1)/2} \frac{4 \cos \alpha_{m,j} (\cos \alpha_{m,j} - 1)}{n} \frac{\det'(\mathbf{R})}{\lambda_j} + \mathcal{O}(\epsilon^2) \quad (59)$$

$$= \det'(\mathbf{R}) \left( 1 + \frac{\epsilon}{\ell} \sum_{j=1}^{(n-1)/2} \frac{2 \cos \alpha_{m,j} (\cos \alpha_{m,j} - 1)}{n(p - \sum_{k=1}^p \cos \alpha_{k,j})} \right) + \mathcal{O}(\epsilon^2) . \quad (60)$$

Hence, applying theorem 3, the almost equilateral spectral determinant is,

$$\det'(\mathcal{L}') = \det'(\mathcal{L}) \left( 1 + \frac{\epsilon}{\ell} \left( 1 + \frac{1}{np} + \sum_{j=1}^{(n-1)/2} \frac{2 \cos \alpha_{m,j} (\cos \alpha_{m,j} - 1)}{n(p - \sum_{k=1}^p \cos \alpha_{k,j})} \right) \right) + \mathcal{O}(\epsilon^2), \quad (61)$$

where  $(0, a_m)$  is the perturbed edge and

$$\det'(\mathcal{L}) = \frac{2^{np-1} \ell^{\beta+1}}{p^{n-1}} \prod_{j=1}^{(n-1)/2} \left( p - \sum_{k=1}^p \cos \alpha_{k,j} \right)^2. \quad (62)$$

## 6 Spanning Trees

The relation between the spectral determinant and number of spanning trees was studied in [11].

**Theorem 8.** *Given an equilateral graph  $\Gamma$  with  $E$  edges of length  $\ell$  and  $V$  vertices,*

$$\# \text{ spanning trees} = T_\Gamma = \frac{\prod_{v \in \mathcal{V}} d_v}{E 2^E \ell^{\beta+1}} \det'(\mathcal{L}), \quad (63)$$

where  $\beta = E - V + 1$  is the first Betti number of the graph.

For a generic quantum graph  $\Gamma$  with edge lengths in  $[\ell, \ell + \epsilon]$ , it was shown that the number of spanning trees is the closest integer to  $T_\Gamma$  if  $\epsilon$  is sufficiently small; see theorem 2. We now apply the results for the spectral determinants of almost equilateral graphs to estimate how fast  $T_\Gamma$  deviates from the number of spanning trees.

Given an almost equilateral quantum graph  $\Gamma'$  with  $E - 1$  edges of length  $\ell$  and one edge of length  $\ell + \epsilon$ , from equations (3) and (6),

$$T_{\Gamma'} = \frac{\ell_{tot} \prod_{e \in \mathcal{E}} \ell_e}{E V \ell^{\beta+1}} \det'(\mathbf{R}'), \quad (64)$$

and

$$\ell_{tot} \prod_{e \in \mathcal{E}} \ell_e = E \ell^{E+1} + \epsilon \ell^E (E + 1) + \mathcal{O}(\epsilon^2). \quad (65)$$

### 6.1 Complete Graphs

Using equation (23), we can see that for an almost equilateral complete graph  $K_n$ ,

$$T_{\Gamma'} = \frac{\ell^{E-\beta}}{V} \det'(\mathbf{R}) + \epsilon \left( \frac{E+1}{EV} \ell^{V-2} \det'(\mathbf{R}) - \frac{2V^{V-3}}{\ell} \right) + \mathcal{O}(\epsilon^2) \quad (66)$$

$$= \# \text{ spanning trees} + \frac{\epsilon n^{n-3} (n^2 - 3n + 4)}{\ell (n-1)} + \mathcal{O}(\epsilon^2) \quad (67)$$

since  $\det'(\mathbf{R}) = (n/\ell)^{n-1}$  for a complete equilateral graph.

For a large graph,  $T_{\Gamma'} \approx \# \text{ spanning trees} + (\epsilon/\ell)n^{n-2}$ . If we use this to estimate the window of edge lengths for which  $T_{\Gamma}$  can be used to evaluate the number of spanning trees, we would expect  $[T_{\Gamma}]$ , the nearest integer to  $T_{\Gamma}$ , gives the number of spanning trees provided the edge lengths all lie in an interval  $[\ell, \ell + \delta)$  for  $\delta \lesssim \ell n^{2-n}$ .

## 6.2 Complete Bipartite Graphs

We now consider an almost equilateral complete bipartite graph,  $K_{m,n}$ . Substituting (37) into equation (64) and using (26),

$$T_{\Gamma'} = \# \text{ spanning trees} + \frac{\epsilon}{\ell} m^{n-2} n^{m-2} (mn - m - n + 2) + \mathcal{O}(\epsilon^2) \quad (68)$$

$$= \# \text{ spanning trees} + \frac{\epsilon(\beta+1) \prod_{v \in \mathcal{V}} d_v}{\ell E^2} + \mathcal{O}(\epsilon^2), \quad (69)$$

because for a complete bipartite graph  $V = m + n$  and  $E = mn$ .

To use this result to estimate a window of edge lengths for which  $[T_{\Gamma}]$  could be expected to count the spanning trees, consider a large graph  $K_{n,n}$ . Then  $T_{\Gamma'} \approx \# \text{ spanning trees} + (\epsilon/\ell)n^{2n-2}$ . Consequently  $[T_{\Gamma}]$  might be expected to provide the number of spanning trees for a metric graph with edge lengths in a window  $[\ell, \ell + \delta)$  for  $\delta \lesssim \ell n^{2-2n}$ .

### 6.2.1 Star Graphs

Since a star graph is a tree, it has only one spanning tree. Suppose that  $\Gamma'$  is an almost equilateral star graph with  $n + 1$  vertices. For a star graph, using (39) and (64),

$$T_{\Gamma'} = 1 + \frac{\epsilon}{\ell n} + \mathcal{O}(\epsilon^2), \quad (70)$$

which agrees (68) with  $m = 1$ .

## 6.3 Circulant Graphs

Consider an almost equilateral circulant graph  $C_n(\mathbf{a})$  with a prime number of vertices and whose maximum degeneracy of eigenvalues of  $\mathbf{R}$  is two. Substituting (59) in (64) and using (45),

$$T_{\Gamma'} = \# \text{ spanning trees} + \epsilon \frac{\ell^{n-2}}{n} \det'(\mathbf{R}) \left( \frac{np+1}{np} + C \right) + \mathcal{O}(\epsilon^2) \quad (71)$$

$$= \# \text{ spanning trees} + \frac{\epsilon 2^{n-1}}{\ell n} \left( \frac{np+1}{np} + C \right) \prod_{j=1}^{(n-1)/2} \left( p - \sum_{k=1}^p \cos \alpha_{k,j} \right)^2 + \mathcal{O}(\epsilon^2), \quad (72)$$

where the number of edges is  $E = np$ , the perturbed edge is  $(0, a_m)$ , and

$$C = \sum_{j=1}^{(n-1)/2} \frac{2 \cos \alpha_{m,j} (\cos \alpha_{m,j} - 1)}{n (p - \sum_{k=1}^p \cos \alpha_{k,j})}. \quad (73)$$

To estimate the size of a window of edge lengths for which  $[T_\Gamma]$  can be expected to count the number of spanning trees, it is convenient to make some simplifying assumptions. For most graphs and most angles  $\theta \in (0, \pi)$  we can expect,

$$\sum_{k=1}^p \cos(\theta a_k) \approx 0. \quad (74)$$

Then,

$$C \approx \sum_{j=1}^{(n-1)/2} \frac{2 \cos \alpha_{m,j} (\cos \alpha_{m,j} - 1)}{np} \approx \frac{1}{2p}, \quad (75)$$

where we replace the sum with averages assuming  $n$  is large. Then, for a typical large  $2p$ -regular circulant graph,

$$T_{\Gamma'} \approx \# \text{ spanning trees} + \left(\frac{\epsilon}{\ell}\right) \frac{1}{n} \left(1 + \frac{1}{2p}\right) (2p)^{n-1}. \quad (76)$$

Therefore we might expect  $[T_\Gamma]$  to evaluate the number of spanning trees for graphs where the edge lengths fall in a window  $[\ell, \ell + \delta)$  for  $\delta \lesssim \ell n (2p)^{1-n} (1 + 1/2p)^{-1}$ .

## 6.4 Summary

In theorem 2, the integer  $[T_\Gamma]$  is the number of spanning trees of a metric graph provided the edge lengths lie in a window  $[\ell, \ell + \delta)$  with

$$\delta < \frac{\ell}{V^V 2^{E+V} \sqrt{2EV}}. \quad (77)$$

Considering the families of almost equilateral graphs analyzed in this section, this window appears unnecessarily narrow. For complete graphs  $K_n$  a window of size  $\delta < \ell/n^{n-2}$  could be expected to be sufficient, while for complete bipartite graphs  $K_{n,n}$  a window of size  $\delta < \ell/n^{2n-2}$  should suffice. For typical  $2p$ -regular circulant graphs  $C_n(\mathbf{a})$ , edge lengths constrained to a window of size  $\delta < \ell n/2(2p)^{n-1}$  could be expected to be sufficient for  $[T_\Gamma]$  to measure the number of spanning trees. All of these results would agree, for example, with a window of size of  $\ell/V^V$ . This is still a tight constraint which requires all of the edge lengths to be close together on the scale of  $\ell$ . However, this argument suggests an analog of Kirchoff's matrix tree theorem for quantum graphs applies to graphs with edge lengths that fall in a window that is much wider than that required in [11].

**Acknowledgments.** TW would like to thank the Rose-Hulman leaves program and the hospitality of the Baylor mathematics department during her sabbatical, when this work was carried out.

## References

- [1] Akkermans, E., Comet, A., Desbois, J., Montambaux, G. and Texier, C.: Spectral determinant on quantum graphs. *Ann. Phys.* (2000) <https://doi.org/10.1006/aphy.2000.6056>
- [2] Berkolaiko, G. and Kuchment, P.: Introduction to quantum graphs, vol. 186 of *Mathematical Surveys and Monographs*. Amer. Math. Soc., Providence, RI (2013)
- [3] Davis, P. J.: *Circulant Matrices*. Wiley, New York, NY (1979)
- [4] Desbois, J.: Spectral determinant of Schrödinger operators on graphs. *J. Phys. A: Math. Gen.* (2000) <https://doi.org/10.1088/0305-4470/33/7/103>
- [5] Desbois, J.: Spectral determinant on graphs with generalized boundary conditions. *Eur. Phys. J. B* (2001) <https://doi.org/10.1007/s100510170013>
- [6] Kirchhoff, G.: Über die Auflösung der Gleichungen, auf welche man bei der Untersuchung der linearen Vertheilung galvanischer Ströme geführt wird. *Ann. Phys. Chem.* (1847) 72:497-508
- [7] Friedlander, L.: Determinant of the Schrödinger operator on a metric graph. In: Berkolaiko, G., Carson, R., Fulling, S. and Kuchment, P. (eds.) *Quantum Graphs and their applications*, p. 151-160 American Mathematical Society, Providence, RI (2006)
- [8] Harrison, J. M. and Kirsten K.: Zeta functions of quantum graphs. *J. Phys. A: Math. Theor.* (2011) <https://doi.org/10.1088/1751-8113/44/23/235301>
- [9] Harrison, J. M., Kirsten, K. and Texier, C.: Spectral determinants and zeta functions of Schrödinger operators on metric graphs. *J. Phys. A: Math. Theor.* (2012) <https://doi.org/10.1088/1751-8113/45/12/125206>
- [10] Harrison, J. M. and Pruss, C.: Discrete quantum unique ergodicity on circulant graphs. (In preparation.)
- [11] Harrison, J. M. and Weyand, T.: Can one hear the spanning trees of a quantum graph? *Lett. Math. Phys.* (2023) <https://doi.org/10.1007/s11005-023-01656-0>
- [12] Harrison, J. M., Weyand, T. and Kirsten, K.: Zeta functions of the Dirac operator on quantum graphs. *J. Math. Phys.* (2016) <https://doi.org/10.1063/1.4964260>



- [13] Kuchment, P.: Quantum graphs I. Some basic structures, *Waves Random Media* (2003) <https://doi.org/10.1088/0959-7174/14/1/014>
- [14] Mathews, J. and Walker, R. L.: *Mathematical Methods of Physics*, Second Edition. Addison-Wesley Publishing Company Inc., Redwood City, CA (1970)