

# On the estimation of varextropy under complete data

F. Goodarzi<sup>a</sup>, R. Zamini<sup>b1</sup>

<sup>a</sup>Faculty of Mathematical Sciences, Department of Statistics, University of Kashan,  
Kashan, Iran.

<sup>b</sup>Department of Mathematics, Faculty of Mathematical Sciences and Computer,  
Kharazmi University, Tehran, Iran.

**Abstract:** In this paper, we propose nonparametric estimators for varextropy function of an absolutely continuous random variable. Consistency of the estimators is established under suitable regularity conditions. Moreover, a simulation study is performed to compare the performance of the proposed estimators based on mean squared error (MSE) and bias. Furthermore, by using the proposed estimators some tests are constructed for uniformity. It is shown that the varextropy-based test proposed here performs well compared to the power of the other uniformity hypothesis tests.

**Keywords:** Varextropy estimator, Consistency, Goodness-of-fit test, Testing uniformity, Monte Carlo simulation.

**Mathematics Subject Classification (2010):** 62G86; 62G05.

## 1 Introduction

Suppose that  $X$  be a non-negative and an absolutely continuous random variable with the probability density function (pdf)  $f$ , distribution function  $F$  and survival function  $\bar{F}$ . A prevalent uncertainty measure is the expectation of  $-\log f(X)$ , as defined by [Shannon \(1948\)](#), and given by

$$H(X) = - \int_0^{+\infty} f(x) \log f(x) dx, \quad (1.1)$$

Many researchers have estimated the entropy of continuous random variables in their studies. Some of these researchers are [Vasicek \(1976\)](#), [van Es \(1992\)](#), [Ebrahimi et al. \(1994\)](#), [Correa \(1995\)](#)

---

<sup>1</sup>Raheleh Zamini:zamini@khu.ac.ir

and [Zamanzade \(2015\)](#). The varentropy of a random variable  $X$  is defined as

$$\begin{aligned} VE(X) &= Var[-\log f(X)] \\ &= \int_0^{+\infty} f(x)[\log f(x)]^2 dx - \left[ \int_0^{+\infty} f(x) \log f(x) dx \right]^2. \end{aligned} \quad (1.2)$$

Hence the variability in the information content of  $X$  is measured by the varentropy. The relevance of this measure to physics, computer and mathematics sciences has been pointed out in various studies, including [Madiman and Barron \(2007\)](#), [Jiang et al. \(2011\)](#), [Fradelizi et al. \(2016\)](#) and [Li et al. \(2016\)](#). Recently [Alizadeh Noughabi and Shafaei Noughabi \(2023\)](#) proposed some estimators for varentropy of a continuous random variable.

The notion of entropy is recently intertwined with a complementary dual measure, designated as extropy, by [Lad et al. \(2015\)](#). The extropy of the random variable  $X$  is defined as:

$$J(X) = -\frac{1}{2} \int_0^{+\infty} f^2(x) dx. \quad (1.3)$$

It is clear that  $J(X) \leq 0$ . Scoring the forecasting distributions using the total log-scoring rule is one of the statistical applications of extropy (see [Gneiting and Raftery \(2007\)](#)). Furthermore, the extropy has been universally investigated in commercial or scientific areas such as astronomical measurements of heat distributions in galaxies, (see [Furuichi and Mitroi \(2012\)](#) and [Vontobel \(2013\)](#)). The concept of extropy is useful in automatic speech recognition (Refer to [Becerra et al. \(2018\)](#)). Moreover, extropy is a measure better than entropy in some scenarios in statistical mechanics and thermodynamics. For more studies on extropy, see [Qiu and Jia \(2018b\)](#), [Yang et al. \(2019\)](#). Recently, the problem of estimating  $J(X)$  has been noted by [Qiu and Jia \(2018a\)](#), [Alizadeh Noughabi and Jarrahiferiz \(2019\)](#) and [Al-Labadi and Berry \(2022\)](#).

[Vaselabadi et al. \(2021\)](#) defined the varextropy for the absolutely continuous random variable  $X$  as:

$$VJ(X) = Var \left[ -\frac{1}{2} f(X) \right] = \frac{1}{4} E[f^2(X)] - J^2(X). \quad (1.4)$$

Moreover, [Goodarzi \(2024\)](#) obtained lower bounds for varextropy. The varextropy measure can be used as an alternative measure to Shannon entropy and a measure of the amount of uncertainty.

The rest of the work is structured as follows. In Section 2, we propose nonparametric estimators for varextropy function and investigate consistency of the estimators under suitable conditions. We present the results of the Monte Carlo studies on biases and mean squared errors (MSEs) of the proposed estimators in Section 3. In Section 4, we introduce some goodness-of-fit tests of uniformity by using the our proposed estimators and compare the powers of our proposed test statistics with some known test statistics. We apply the newly introduced test procedures to real data sets for illustration in Section 5.

## 2 Varextropy estimation for complete data

In this section, we propose five estimators for varextropy function under complete data and investigate some of their asymptotic properties.

### 2.1 The first estimator

Before constructing the first estimator for  $VJ(X)$ , we first review some required concepts and notations. Let  $X$  be a continuous random variable with density function  $f$  and distribution function  $F$ . Consider the class of functions based on  $f(x)$  as

$$T(f) = \int_{-\infty}^{+\infty} f(x)\Phi(f(x))w(x)dx, \quad (2.1)$$

where  $\Phi$  is a real valued differentiable function on  $[0, \infty)$  and  $w$  is a real valued function on  $[0, \infty)$ .

Assuming that  $f$  is bounded, van Es (1992) proposed the following estimator for  $T(f)$ :

$$T_{m,n} = \frac{1}{2(n-m)} \sum_{j=1}^{n-m} \Phi \left( \frac{m}{n+1} (X_{(j+m)} - X_{(j)})^{-1} \right) (w(X_{(j)}) + w(X_{(j+m)})), \quad (2.2)$$

where  $X_{(1)} \leq \dots \leq X_{(n)}$  are the order statistics corresponding to the sample  $X_1, \dots, X_n$  and  $m$  is an integer such that  $1 \leq m \leq n$ . The estimator given in (2.2) is constructed according to the fact that  $f(X_{(j)})$ , the density at the point  $X_{(j)}$ , is replaced by  $m((n+1)(X_{(j+m)} - X_{(j)})^{-1})$ , where is the local histogram estimate of  $f(X_{(j)})$ .

Under some conditions, van Es (1992) showed that  $T_{m,n}$  is a.s. consistent, i.e.,

$$T_{m,n} \rightarrow T(f), \quad a.s. \quad as \quad m, n \rightarrow \infty. \quad (2.3)$$

(See Theorem 1 of van Es (1992) for some details.) Two special cases of  $T(f)$  are

$$T'(f) = \int_0^{+\infty} f^3(x)dx, \quad (2.4)$$

and

$$T''(f) = \int_0^{+\infty} f^2(x)dx, \quad (2.5)$$

obtained by taking  $\Phi(x)$  equal to  $x^2$  and  $x$ , respectively, and  $w$  is identically equal to  $I_{[0,\infty)}(x)$ . According to  $T_{m,n}$  defined in (2.2), estimators for  $T'(f)$  and  $T''(f)$  can be proposed as

$$T'_{m,n} = \frac{1}{n-m} \sum_{j=1}^{n-m} \left( \frac{m}{n+1} (X_{(j+m)} - X_{(j)})^{-1} \right)^2, \quad (2.6)$$

and

$$T''_{m,n} = \frac{1}{n-m} \sum_{j=1}^{n-m} \left( \frac{m}{n+1} (X_{(j+m)} - X_{(j)})^{-1} \right), \quad (2.7)$$

respectively.

Now, the expression given in (1.4) for  $VJ(X)$ , motivates us to construct the first estimator of varextropy based on  $T'_{m,n}$  and  $T''_{m,n}$  as follows:

$$VJV_{m,n} = \frac{1}{4}\{T'_{m,n} - (T''_{m,n})^2\}. \quad (2.8)$$

In the following theorem, we establish the a.s. consistency of  $VJV_{m,n}$ .

**Theorem 2.1.** *Let the density function  $f$  be bounded, then  $VJV_{m,n} \rightarrow VJ(X)$  a.s., provided  $m, n \rightarrow \infty$ ,  $\frac{m}{\log n} \rightarrow \infty$  and  $\frac{m}{n} \rightarrow 0$ .*

*Proof.*

$$\begin{aligned} VJV_{m,n} - VJ(X) &= \frac{1}{4}\{T'_{m,n} - (T''_{m,n})^2\} - \frac{1}{4}\{T'(f) - (T''(f))^2\} \\ &= \frac{1}{4}\{T'_{m,n} - T'(f)\} - \frac{1}{4}\{(T''_{m,n})^2 - (T''(f))^2\}. \end{aligned} \quad (2.9)$$

Theorem 1 of van Es (1992) ensures that

$$T'_{m,n} \rightarrow T'(f) \text{ a.s. as } m, n \rightarrow \infty, \quad (2.10)$$

and

$$(T''_{m,n})^2 \rightarrow (T''(f))^2 \text{ a.s. as } m, n \rightarrow \infty. \quad (2.11)$$

(2.9), (2.10) and (2.11) complete the proof.  $\square$

## 2.2 The second estimator

Let  $\hat{f}(x)$  be the kernel density function of  $f$  defined by

$$\hat{f}(x) = \frac{1}{nh_n} \sum_{i=1}^n K\left(\frac{x - X_i}{h_n}\right), \quad (2.12)$$

where  $K$  is the kernel density function and  $h_n$  is the bandwidth parameter. The second estimator for  $VJ(X)$  can be proposed based on  $\hat{f}(x)$  as:

$$VJD = \frac{1}{4} \left[ \int \hat{f}^3(x) dx - \left( \int \hat{f}^2(x) dx \right)^2 \right]. \quad (2.13)$$

The following assumptions on  $K$  are used in the Theorem 2.2 of this section.

(A<sub>1</sub>) :  $K$  is uniformly continuous and of bounded variation  $V(K)$ .

(A<sub>2</sub>) :  $K(x) \rightarrow 0$  as  $|x| \rightarrow \infty$ .

(A<sub>3</sub>) :  $\int |x \log |x|^{\frac{1}{2}}| dK(x) < \infty$ .

**Theorem 2.2.** *Suppose  $K$  satisfies Assumptions A<sub>1</sub>–A<sub>3</sub> and  $f$  is uniformly continuous. If  $h_n \rightarrow 0$  and  $(nh_n)^{-1} \log n \rightarrow 0$  as  $n \rightarrow \infty$ , then*

$$VJD \rightarrow VJ(X) \text{ a.s. as } n \rightarrow \infty.$$

*Proof.* From (1.4) and (2.13) we have

$$VJD - VJ(X) = I_n + II_n, \quad (2.14)$$

where

$$I_n = \frac{1}{4} \left( \int_0^{+\infty} \hat{f}^3(x) dx - \int_0^{+\infty} f^3(x) dx \right), \quad (2.15)$$

and

$$II_n = \left( -\frac{1}{2} \int_0^{+\infty} f^2(x) dx \right)^2 - \left( -\frac{1}{2} \int_0^{+\infty} \hat{f}^2(x) dx \right)^2. \quad (2.16)$$

Observe that

$$\begin{aligned} |I_n| &= \frac{1}{4} \left| \int_0^{+\infty} \hat{f}^3(x) dx - \int_0^{+\infty} f^3(x) dx \right| \\ &\leq \frac{1}{4} \left| \int_0^{+\infty} \hat{f}^2(x) (\hat{f}(x) - f(x)) dx \right| \\ &\quad + \frac{1}{4} \left| \int_0^{+\infty} (\hat{f}^2(x) - f^2(x)) f(x) dx \right|. \end{aligned} \quad (2.17)$$

Under Assumptions  $A_1 - A_3$ , from Theorem A of Silverman (1978), one can see that

$$\lim_{n \rightarrow \infty} \sup_x |\hat{f}(x) - f(x)| = 0, \quad a.s. \quad (2.18)$$

On the other hand, a simple algebra calculation shows that

$$\hat{f}^2(x) - f^2(x) = \sum_{i=1}^2 (f(x))^{i-1} (\hat{f}(x))^{2-i} (\hat{f}(x) - f(x)). \quad (2.19)$$

(2.18) and (2.19) ensure that

$$\lim_{n \rightarrow \infty} \sup_x |\hat{f}^2(x) - f^2(x)| = 0, \quad a.s. \quad (2.20)$$

From (2.17), (2.18) and (2.20), one can observe that

$$\lim_{n \rightarrow \infty} I_n = 0, \quad a.s. \quad (2.21)$$

Next, to deal with  $II_n$ , observe that

$$\begin{aligned} \left| \int_0^{+\infty} \hat{f}^2(x) dx - \int_0^{+\infty} f^2(x) dx \right| &\leq \left| \int_0^{+\infty} \hat{f}^2(x) dx - \int_0^{+\infty} \hat{f}(x) dF(x) \right| \\ &\quad + \left| \int_0^{+\infty} \hat{f}(x) dF(x) - \int_0^{+\infty} f^2(x) dx \right| \\ &= \left| \int_0^{+\infty} \hat{f}(x) (\hat{f}(x) - f(x)) dx \right| \\ &\quad + \left| \int_0^{+\infty} (\hat{f}(x) - f(x)) dF(x) \right|. \end{aligned} \quad (2.22)$$

Since from (2.18),  $\hat{f}(x)$  is a.s. consistent, we can write

$$\lim_{n \rightarrow \infty} \int_0^{+\infty} \hat{f}^2(x) dx = \int_0^{+\infty} f^2(x) dx, \quad a.s. \quad (2.23)$$

As a result

$$\lim_{n \rightarrow \infty} II_n = 0, \quad a.s. \quad (2.24)$$

(2.14), (2.21) and (2.24) complete the proof.  $\square$

### 2.3 The third estimator

The third estimator is proposed by:

$$\frac{1}{4} \left( \frac{1}{n} \sum_{i=1}^n f^2(X_i) - \left( \frac{1}{n} \sum_{i=1}^n f(X_i) \right)^2 \right), \quad (2.25)$$

using the empirical distribution function  $F_n(x) = \frac{1}{n} \sum_{i=1}^n I(X_i \leq x)$ .

Since  $f$  is unknown, a natural estimator for  $VJ(X)$  can be obtained by substituting  $f$  by  $\hat{f}$  defined in (2.12), and is given by

$$\begin{aligned} VJB_n &= \frac{1}{4} \left( \int_0^{+\infty} \hat{f}^2(x) dF_n(x) - \left( \int_0^{+\infty} \hat{f}(x) dF_n(x) \right)^2 \right) \\ &= \frac{1}{4} \left( \frac{1}{n} \sum_{i=1}^n (\tilde{f}_n(X_i))^2 - \left( \frac{1}{n} \sum_{i=1}^n \tilde{f}_n(X_i) \right)^2 \right), \end{aligned} \quad (2.26)$$

where

$$\tilde{f}_n(X_i) = \frac{1}{(n-1)h_{n-1}} \sum_{j=1, j \neq i}^n K \left( \frac{X_i - X_j}{h_{n-1}} \right). \quad (2.27)$$

In the following theorem, we investigate a.s. consistency of  $VJ_n(X)$ .

**Theorem 2.3.** *Suppose that  $\lim_{x \rightarrow \infty} f(x)$  exists and is finite. If  $f$  is bounded, then under the conditions of Theorem 2.2, we can write:*

$$\lim_{n \rightarrow \infty} |VJB_n - VJ(X)| = 0, \quad a.s.$$

*Proof.* By using (1.4) and (2.26) we have

$$|VJB_n - VJ(X)| \leq |I'_n| + |I''_n|, \quad (2.28)$$

where

$$I'_n = \frac{1}{4n} \sum_{i=1}^n (\tilde{f}_n(X_i))^2 - \frac{1}{4} \int_0^{+\infty} f^2(x) dF(x), \quad (2.29)$$

and

$$I_n'' = \left( -\frac{1}{2} \int_0^{+\infty} f^2(x) dx \right)^2 - \left( -\frac{1}{2n} \sum_{i=1}^n (\tilde{f}_n(X_i)) \right)^2. \quad (2.30)$$

Using the definition of step function  $F_n(x)$  and adding and subtracting  $\frac{1}{4} \int_0^{+\infty} f^2(x) dF_n(x)$ ,  $I_n'$  can be rewritten as

$$\begin{aligned} I_n' &= \frac{1}{4} \int_0^{+\infty} \hat{f}^2(x) dF_n(x) - \frac{1}{4} \int_0^{+\infty} f^2(x) dF(x) \\ &= \frac{1}{4} \int_0^{+\infty} (\hat{f}^2(x) - f^2(x)) dF_n(x) + \frac{1}{4} \int_0^{+\infty} f^2(x) d(F_n(x) - F(x)). \end{aligned} \quad (2.31)$$

On the other hand, since  $f$  is bounded and  $\lim_{x \rightarrow \infty} f(x)$  exists and is finite, by using the integration by parts we have:

$$\frac{1}{4} \int_0^{+\infty} f^2(x) d(F_n(x) - F(x)) = -\frac{1}{2} \int_0^{+\infty} (F_n(x) - F(x)) f(x) f'(x) dx. \quad (2.32)$$

From (2.31) and (2.32), we can write:

$$I_n' = \frac{1}{4} \int_0^{+\infty} (\hat{f}^2(x) - f^2(x)) dF_n(x) - \frac{1}{2} \int_0^{+\infty} (F_n(x) - F(x)) f(x) f'(x) dx. \quad (2.33)$$

Hence

$$|I_n'| \leq \frac{1}{4} \sup_x |\hat{f}^2(x) - f^2(x)| + \frac{1}{4} \sup_x |F_n(x) - F(x)| \left( \lim_{x \rightarrow \infty} f^2(x) + (\sup_x f(x))^2 \right). \quad (2.34)$$

The a.s. convergence of  $\hat{f}^2(x)$  given in (2.20) and a.s. convergence of  $F_n(x)$  given in the Glivenko-Cantelli theorem ensure that

$$\lim_{n \rightarrow \infty} |I_n'| = 0. \quad (2.35)$$

In a similar way, using the triangle inequality and adding and subtracting  $\int_0^{+\infty} f(x) dF_n(x)$ , we can see that

$$\begin{aligned} \left| \int_0^{+\infty} \hat{f}(x) dF_n(x) - \int_0^{+\infty} f(x) dF(x) \right| &\leq \left| \int_0^{+\infty} (\hat{f}(x) - f(x)) dF_n(x) \right| \\ &\quad + \left| \int_0^{+\infty} f(x) d(F_n(x) - F(x)) \right|. \end{aligned} \quad (2.36)$$

Since  $f$  is bounded and  $\lim_{x \rightarrow \infty} f(x)$  exists and is finite, using the integration by parts we can write:

$$\int_0^{+\infty} f(x) d(F_n(x) - F(x)) = - \int_0^{+\infty} (F_n(x) - F(x)) f'(x) dx. \quad (2.37)$$

(2.36) and (2.37) imply that

$$\begin{aligned} \left| \int_0^{+\infty} \hat{f}(x) dF_n(x) - \int_0^{+\infty} f(x) dF(x) \right| &\leq \sup_x |\hat{f}(x) - f(x)| \\ &\quad + \sup_x |F_n(x) - F(x)| \left( \lim_{x \rightarrow \infty} f(x) + \sup_x f(x) \right). \end{aligned} \quad (2.38)$$

It can be concluded from the a.s. convergence of  $\hat{f}$  given in (2.18), the Glivenko-Cantelli theorem and the finiteness of  $\lim_{x \rightarrow \infty} f(x)$  and  $\sup_x f(x)$  that

$$\lim_{n \rightarrow \infty} |I_n''| = 0. \quad (2.39)$$

(2.28), (2.35) and (2.39) complete the proof.  $\square$

## 2.4 The fourth estimator

Let  $Q(u) = \inf\{x : F(x) \geq u\}$ ,  $0 \leq u \leq 1$ , be the quantile function corresponding to the distribution function  $F(x)$  and  $q(u) = Q'(u) = \frac{1}{f(Q(u))}$  be its quantile density function. Soni et al. (2012) proposed a smooth estimator for  $q(u)$  given by

$$\begin{aligned} \tilde{q}_n(u) &= \frac{1}{h_n} \int_0^1 \frac{K\left(\frac{t-u}{h_n}\right)}{f_n(Q_n(t))} dt \\ &= \frac{1}{nh_n} \sum_{i=1}^n \frac{K\left(\frac{S_i-u}{h_n}\right)}{f_n(X_{(i)})}, \end{aligned} \quad (2.40)$$

where  $S_i$  is the proportion of observations less than or equal to  $X_{(i)}$ , the  $i$ th order statistic,  $f_n(u)$  is the same  $\hat{f}(u)$  given in (2.12), and  $Q_n(u) = \inf\{x : F_n(x) \geq u\}$ ,  $0 \leq u \leq 1$ , is the empirical estimator of  $Q(u)$ .

The fourth estimator for varextropy function  $VJ(X)$  is obtained by considering the fact that  $VJ(X)$  can be rewritten in terms of the quantile function as follows:

$$\begin{aligned} VJ(X) &= \frac{1}{4} \left[ \int_0^1 f^2(Q(u)) du - \left( \int_0^1 f(Q(u)) du \right)^2 \right] \\ &= \frac{1}{4} \left[ \int_0^1 \frac{du}{(q(u))^2} - \left( \int_0^1 \frac{du}{q(u)} \right)^2 \right]. \end{aligned} \quad (2.41)$$

We consider the following plug-in estimator for quantile-based varextropy, given by

$$VJS_n = \frac{1}{4} \left[ \int_0^1 \frac{du}{(\tilde{q}_n(u))^2} - \left( \int_0^1 \frac{du}{\tilde{q}_n(u)} \right)^2 \right]. \quad (2.42)$$

In the following theorem, we prove the a.s. consistency of  $VJS_n$ .

**Theorem 2.4.** *Let  $f$  be bounded, then*

$$VJS_n \rightarrow VJ(X), \quad a.s. \quad n \rightarrow \infty.$$

*Proof.* We have

$$\begin{aligned} VJS_n - VJ(X) &= \frac{1}{4} \left[ \int_0^1 \frac{du}{(\tilde{q}_n(u))^2} - \int_0^1 \frac{du}{(q(u))^2} \right] \\ &\quad - \frac{1}{4} \left[ \left( \int_0^1 \frac{du}{\tilde{q}_n(u)} \right)^2 - \left( \int_0^1 \frac{du}{q(u)} \right)^2 \right] \\ &:= J_1 + J_2. \end{aligned} \quad (2.43)$$



Now

$$\int_0^1 \left( \frac{1}{(\tilde{q}_n(u))^2} - \frac{1}{(q(u))^2} \right) du = \int_0^1 (q(u) - \tilde{q}_n(u)) \cdot \frac{q(u) + \tilde{q}_n(u)}{(\tilde{q}_n(u))^2 q^2(u)} du. \quad (2.44)$$

Using proof of Theorem 3.4 of [Subhash et al. \(2023\)](#),  $\sup_t |\tilde{q}_n(t) - q(t)| \rightarrow 0$  as  $n \rightarrow \infty$ , hence the above right expression asymptotically reduces to

$$\int_0^1 (q(u) - \tilde{q}_n(u)) \frac{2}{q^3(u)} du. \quad (2.45)$$

Since  $f$  is bounded, there exist  $M > 0$ , such that  $|f| \leq M$  and hence

$$\int_0^1 (q(u) - \tilde{q}_n(u)) \frac{2}{q^3(u)} du \leq 2 \sup_u |q(u) - \tilde{q}_n(u)| \int_0^1 f^3(Q(u)) du \leq 2M^3 \sup_u |q(u) - \tilde{q}_n(u)| \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (2.46)$$

As a result

$$J_1 \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (2.47)$$

In a similar way, we can show that

$$J_2 \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (2.48)$$

(2.43), (2.47) and (2.48), ensure that

$$VJS_n \rightarrow VJ(X) \text{ as } n \rightarrow \infty.$$

□

## 2.5 The fifth estimator

By analogy to [Qiu and Jia \(2018a\)](#) and using the (2.41), we propose the following estimator for varextropy function:

$$VJQ_{m,n} = \frac{1}{4} \left[ \frac{1}{n} \sum_{i=1}^n \left( \frac{c_i \frac{m}{n}}{X_{(i+m)} - X_{(i-m)}} \right)^2 - \left( \frac{1}{n} \sum_{i=1}^n \frac{c_i \frac{m}{n}}{X_{(i+m)} - X_{(i-m)}} \right)^2 \right], \quad (2.49)$$

where the window size  $m$  is a positive integer smaller than  $n/2$ ,  $X_{(i)} = X_{(1)}$  if  $i < 1$ ,  $X_{(i)} = X_{(n)}$  if  $i > n$  and

$$c_i = \begin{cases} 1 + \frac{i-1}{m}, & 1 \leq i \leq m, \\ 2, & m+1 \leq i \leq n-m, \\ 1 + \frac{n-i}{m}, & n-m+1 \leq i \leq n. \end{cases}$$

The consistency of the proposed estimator is shown in the following theorem.

**Theorem 2.5.** *Let  $X_1, \dots, X_n$  be a random sample from distribution function  $F$  with bounded probability density function  $f$  and finite variance. Then we can write:*

$$VJQ_{m,n} \xrightarrow{P} VJ(X), \text{ as } n \rightarrow \infty, \quad m \rightarrow \infty, \quad \frac{m}{n} \rightarrow 0. \quad (2.50)$$

*Proof.* A simple algebra shows that:

$$VJQ_{m,n} - VJ(X) := I_1 + I_2, \quad (2.51)$$

where

$$I_1 = \frac{1}{4} \left[ \frac{1}{n} \sum_{i=1}^n \left( \frac{c_i \frac{m}{n}}{X_{(i+m)} - X_{(i-m)}} \right)^2 - \int_0^{+\infty} f^3(x) dx \right], \quad (2.52)$$

and

$$I_2 = \frac{1}{4} \left[ \left( \frac{1}{n} \sum_{i=1}^n \frac{c_i \frac{m}{n}}{X_{(i+m)} - X_{(i-m)}} \right)^2 - \left( \int_0^{+\infty} f^2(x) dx \right)^2 \right]. \quad (2.53)$$

Theorem 2.1 of Qiu and Jia (2018a), clearly shows that  $I_2 \xrightarrow{P} 0$ . For  $I_1$ , first we write  $\frac{1}{n} \sum_{i=1}^n \left( \frac{c_i \frac{m}{n}}{X_{(i+m)} - X_{(i-m)}} \right)^2$  as follows

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n (c_i \frac{m}{n})^2 \left( \frac{1}{X_{(i+m)} - X_{(i-m)}} \right)^2 &= \frac{1}{n} \sum_{i=1}^n \left( \frac{\frac{2m}{n}}{X_{(i+m)} - X_{(i-m)}} \right)^2 \\ &\quad - \frac{2}{n} \sum_{i=1}^n \xi_{im} \frac{2m}{n} \frac{1}{X_{(i+m)} - X_{(i-m)}} \left( 1 - \frac{c_i}{2} \right) := J_1 - J_2. \end{aligned} \quad (2.54)$$

where  $\xi_{im}$  is a random point between the points  $\frac{c_i \frac{m}{n}}{X_{(i+m)} - X_{(i-m)}}$  and  $\frac{\frac{2m}{n}}{X_{(i+m)} - X_{(i-m)}}$ .

Next to deal with  $J_1$ , observe that

$$\begin{aligned} J_1 &= \frac{1}{n} \sum_{i=1}^n \left( \frac{\frac{2m}{n}}{X_{(i+m)} - X_{(i-m)}} \right)^2 = \frac{1}{n} \sum_{i=1}^n \left( \frac{F(X_{(i+m)}) - F(X_{(i-m)})}{X_{(i+m)} - X_{(i-m)}} \right)^2 \left[ \left( \frac{\frac{2m}{n}}{F(X_{(i+m)}) - F(X_{(i-m)})} \right)^2 - 1 \right] \\ &\quad + \frac{1}{n} \sum_{i=1}^n \left( \frac{F(X_{(i+m)}) - F(X_{(i-m)})}{X_{(i+m)} - X_{(i-m)}} \right)^2 \\ &:= J_{11} + J_{12}. \end{aligned} \quad (2.55)$$

Now, since  $f(x)$  is positive and continuous, there exists a value  $X_i \in (X_{(i-m)}, X_{(i+m)})$  such that

$$f(X_i) = \frac{F(X_{(i+m)}) - F(X_{(i-m)})}{X_{(i+m)} - X_{(i-m)}}, \quad (2.56)$$

hence

$$J_{11} = \frac{1}{n} \sum_{i=1}^n (f(X_i))^2 \left[ \left( \frac{\frac{2m}{n}}{F(X_{(i+m)}) - F(X_{(i-m)})} \right)^2 - 1 \right]. \quad (2.57)$$

On the other hand, since  $F_n(X_{(i+m)}) - F_n(X_{(i-m)}) = \frac{2m}{n}$ , Chung's law of the iterated logarithm (LIL) implies that:

$$\begin{aligned} \frac{F(X_{(i+m)}) - F(X_{(i-m)})}{F_n(X_{(i+m)}) - F_n(X_{(i-m)})} &\leq \frac{|F_n(X_{(i+m)}) - F(X_{(i+m)})|}{F_n(X_{(i+m)}) - F_n(X_{(i-m)})} + \frac{|F_n(X_{(i-m)}) - F(X_{(i-m)})|}{F_n(X_{(i+m)}) - F_n(X_{(i-m)})} \\ &\quad + \frac{F_n(X_{(i+m)}) - F_n(X_{(i-m)})}{F_n(X_{(i+m)}) - F_n(X_{(i-m)})} \leq \frac{n}{2m} \sup_x |F_n(x) - F(x)| \\ &\quad + \frac{n}{2m} \sup_x |F_n(x) - F(x)| + 1 \\ &\leq \frac{n}{m} O \left( \sqrt{\frac{\log \log n}{n}} \right) + 1, \end{aligned} \quad (2.58)$$

Now, since  $\lim_{n \rightarrow \infty} \frac{\sqrt{n \log \log n}}{m} = 0$ , we can see that

$$\lim_{n \rightarrow \infty} \frac{\frac{2m}{n}}{F(X_{(i+m)}) - F(X_{(i-m)})} = \lim_{n \rightarrow \infty} \frac{F_n(X_{(i+m)}) - F_n(X_{(i-m)})}{F(X_{(i+m)}) - F(X_{(i-m)})} \geq 1. \quad (2.59)$$

(2.57), (2.59) and condition of  $\sup_x f(x) < M < \infty$  result that for enough large  $n$  we can write

$$J_{11} \leq \frac{M^2}{n} \sum_{i=1}^n \left[ \frac{\left(\frac{2m}{n}\right)^2}{(F(X_{(i+m)}) - F(X_{(i-m)}))^2} - 1 \right]. \quad (2.60)$$

Now, we prove that when  $m \rightarrow \infty$ ,

$$\lim_{n \rightarrow \infty} E \left( \frac{M^2}{n} \sum_{i=1}^n \left[ \frac{\left(\frac{2m}{n}\right)^2}{(F(X_{(i+m)}) - F(X_{(i-m)}))^2} - 1 \right] \right) = 0. \quad (2.61)$$

Note that

$$\begin{aligned} \lim_{n \rightarrow \infty} E \left( \frac{M^2}{n} \sum_{i=1}^n \left[ \frac{\left(\frac{2m}{n}\right)^2}{(F(X_{(i+m)}) - F(X_{(i-m)}))^2} - 1 \right] \right) &= \lim_{n \rightarrow \infty} E \left( \frac{M^2}{n} \sum_{i=1}^m \left[ \frac{\left(\frac{2m}{n}\right)^2}{(F(X_{(i+m)}) - F(X_{(1)}))^2} - 1 \right] \right) \\ &+ \lim_{n \rightarrow \infty} E \left( \frac{M^2}{n} \sum_{i=m+1}^{n-m} \left[ \frac{\left(\frac{2m}{n}\right)^2}{(F(X_{(i+m)}) - F(X_{(i-m)}))^2} - 1 \right] \right) \\ &+ \lim_{n \rightarrow \infty} E \left( \frac{M^2}{n} \sum_{i=n-m+1}^n \left[ \frac{\left(\frac{2m}{n}\right)^2}{(F(X_{(n)}) - F(X_{(i-m)}))^2} - 1 \right] \right). \end{aligned} \quad (2.62)$$

Next notice that  $F(X_{(1)}), \dots, F(X_{(n)})$  is an orderd sample from uniform distribution and the random variable  $F(X_{(i+j)}) - F(X_{(i)})$  has the beta distribution with parameters  $j$  and  $n - j + 1$ , so we have

$$\begin{aligned} \lim_{n \rightarrow \infty} E \left( \frac{M^2}{n} \sum_{i=1}^n \left[ \frac{\left(\frac{2m}{n}\right)^2}{(F(X_{(i+m)}) - F(X_{(i-m)}))^2} - 1 \right] \right) \\ = \lim_{n \rightarrow \infty} M^2 \left( \frac{2m^2(n-1)(2m^2 + mn + 2m - 2n)}{n^2(m-1)(m-2)(2m-1)} - 1 \right) = 0. \end{aligned} \quad (2.63)$$

Since convergence in expectation to 0 implies convergence in probability to 0 for nonnegative random variables, thus

$$J_{11} \xrightarrow{P} 0. \quad (2.64)$$

Note that by (2.56) and the law of large numbers (LLN), we can write:

$$J_{12} = \frac{1}{n} \sum_{i=1}^n f^2(X_i) = \int f^2(x) dF_n(x) \xrightarrow{P} \int f^3(x) dx. \quad (2.65)$$

(2.55), (2.64) and (2.65) conclude that

$$J_1 \xrightarrow{P} \int f^3(x) dx. \quad (2.66)$$

To deal with  $J_2$ , we notice that since  $\xi_{im} \leq \frac{\frac{2m}{n}}{X_{(i+m)} - X_{(i-m)}}$ , we can write

$$\begin{aligned} |J_2| &\leq \frac{2}{n} \sum_{i=1}^n \frac{\left(\frac{2m}{n}\right)^2}{(X_{(i+m)} - X_{(i-m)})^2} \left(1 - \frac{c_i}{2}\right) \\ &= \frac{2}{n} \sum_{i=1}^n \left(\frac{2m}{n}\right)^2 \left(\frac{F(X_{(i+m)}) - F(X_{(i-m)})}{X_{(i+m)} - X_{(i-m)}}\right)^2 \frac{1 - \frac{c_i}{2}}{(F(X_{(i+m)}) - F(X_{(i-m)}))^2}. \end{aligned}$$

From (2.56) and boundedness of  $f(X_i)$ , we have

$$\begin{aligned} |J_2| &\leq \frac{2M^2}{n} \sum_{i=1}^n \frac{4m^2}{n^2} \frac{1 - \frac{c_i}{2}}{(F(X_{(i+m)}) - F(X_{(i-m)}))^2} \\ &= \frac{4M^2m}{n^3} \left[ \sum_{i=1}^m \frac{m-i+1}{(F(X_{(i+m)}) - F(X_{(1)}))^2} + \sum_{i=n-m+1}^n \frac{m-n+i}{(F(X_{(n)}) - F(X_{(i-m)}))^2} \right], \end{aligned} \quad (2.67)$$

thus

$$\begin{aligned} E(J_2) &\leq \frac{8M^2m}{n^2} \sum_{i=1}^m \frac{(n-1)(m-i+1)}{(i+m-2)(i+m-3)} \\ &= \frac{8M^2m(n-1)}{n^2} \left\{ \frac{m(2m-1)}{2(m-1)(m-2)} + \psi(m-2) - \psi(2m-2) \right\}, \end{aligned} \quad (2.68)$$

where  $\psi(x) = \frac{\Gamma'(x)}{\Gamma(x)}$  and converges to zero with  $n \rightarrow \infty$ ,  $m \rightarrow \infty$  and  $m/n \rightarrow 0$ . Thus,  $J_2$  form a series of non-positive variables with expectations approaching zero, and consequently

$$J_2 \xrightarrow{P} 0. \quad (2.69)$$

(2.51), (2.66) and (2.69) complete the proof.  $\square$

**Remark 2.6.** In Theorem 2.5, many  $m$ 's can be found that satisfy the conditions of theorem, for example  $m = O((\log \log n)^\alpha n^\beta)$  for every  $\alpha \in \mathbb{R}$  and  $\frac{1}{2} < \beta < 1$ .

### 3 Simulation study

In this section, the results of the Monte Carlo studies on mean squared errors (MSEs) and biases of our introduced estimators are presented. We consider the gamma, uniform and exponential distributions, which are considered in many references. For each sample size, 10,000 samples were generated and MSEs and biases of estimators were computed. We used the  $m = \lceil \sqrt{n} + 0.5 \rceil$  formula to estimate the varextropy suggested by Grzegorzewski and Wieczorkowski (1999). They used this heuristic formula for entropy estimation. In addition, we choose the standard normal density as the kernel and its corresponding optimal value of  $h_n = 1.06sn^{-\frac{1}{5}}$ , where  $s$  is the sample standard deviation. Tables 1-3 give simulated biases and MSEs of  $VJV_{m,n}$ ,  $VJD$ ,  $VJB_n$ ,  $VJS_n$  and  $VJQ_{m,n}$  for gamma, uniform and exponential distributed samples, respectively. It is apparent from the tables that both the absolute bias and MSE decrease as the sample size increases. The bold type in these tables indicates the varextropy estimator achieves the minimal MSE (absolute bias).

In the case of gamma distribution, Table 1, it is observed that, the estimator  $VJD$  performs best in terms of bias and MSE. Under the uniform distribution, Table 2, it can be seen that, the MSEs and biases of  $VJS_n$  are always smaller than other proposed estimators except in  $n = 10$ . According to Table 3, under exponential distribution, estimator  $VJD$  performs better

Table 1: MSE and Bias of estimators to estimate the varextropy  $VJ(X)$  of the gamma distribution  $(G(2, 1))$ .

$n$	MSE(Bias)				
	$VJV_{m,n}$	$VJD$	$VJB_n$	$VJS_n$	$VJQ_{m,n}$
10	0.0120008(0.0198195)	<b>0.0000030(-0.0004866)</b>	0.0000048(-0.0011958)	0.0000072(-0.0023191)	0.0016533(0.0065913)
20	0.0016044(0.0116757)	<b>0.0000013(-0.0005814)</b>	0.0000026(-0.0013097)	0.0000064(-0.0024858)	0.0000979(0.0030967)
30	0.0002339(0.0074458)	<b>0.0000010(-0.0005538)</b>	0.0000022(-0.0012534)	0.0000062(-0.0024651)	0.0000228(0.0020731)
40	0.0000904(0.0052193)	<b>0.0000008(-0.0005351)</b>	0.0000019(-0.0012115)	0.0000060(-0.0024254)	0.0000115(0.0016434)
50	0.0000483(0.0039833)	<b>0.0000007(-0.0005132)</b>	0.0000017(-0.0011682)	0.00000577(-0.0023838)	0.0000072(0.0013517)
100	0.0000125(0.0024089)	<b>0.0000004(-0.0004324)</b>	0.0000013(-0.0010154)	0.0000048(-0.0021808)	0.0000030(0.0010021)

Table 2: MSE and Bias of estimators to estimate the varextropy  $VJ(X)$  of the uniform distribution  $(U(0, 1))$

$n$	MSE(Bias)				
	$VJV_{m,n}$	$VJD$	$VJB_n$	$VJS_n$	$VJQ_{m,n}$
10	5.634314(0.3022407)	0.0019133(0.0405239)	<b>0.0007618(0.0191408)</b>	0.0007793(0.0197080)	0.5714468(0.1247980)
20	0.2218065(0.1563245)	0.0013200(0.0349457)	0.0003397(0.0141829)	<b>0.0001894(0.0109698)</b>	0.0163038(0.0553850)
30	0.0295767(0.0993167)	0.0010754(0.0320101)	0.0002256(0.0119855)	<b>0.0001103(0.0086761)</b>	0.0040972(0.0372555)
40	0.0113412(0.0710308)	0.0009404(0.0301090)	0.0001722(0.0107598)	<b>0.0000784(0.0074447)</b>	0.0023016(0.0282432)
50	0.0058115(0.0547022)	0.0008532(0.0287799)	0.0001409(0.0099317)	<b>0.0000610(0.0066770)</b>	0.0010221(0.0222844)
100	0.0015858(0.0336733)	0.0006416(0.0251320)	0.0000811(0.0079505)	<b>0.0000300(0.0048685)</b>	0.0003130(0.0142730)

than other proposed estimators in terms of MSE, and in terms of absolute bias, estimator  $VJD$  performs better in small samples and estimator  $VJQ_{m,n}$  in large samples.

Finally, based on the results of our simulation study, we can conclude that the estimators  $VJD$  and  $VJS_n$  have better performance.

## 4 Some tests of uniformity

In this section, we introduce a few goodness-of-fit tests of uniformity. These tests are based on our proposed varextropy estimators and their percentage points and power values are obtained by Monte Carlo simulation.

### 4.1 Test statistic

Consider the class of continuous distribution functions  $F$  with density function  $f(x)$  defined on interval  $[0, 1]$ . Due to the fact that the variance is always nonnegative, we have  $VJ(X) = \text{Var}[-\frac{1}{2}f(X)] \geq 0$ . If  $f$  is a standard uniform density function, it is easy to see that  $VJ(X) = 0$ . On the other hand, if  $VJ(X) = \text{Var}[-\frac{1}{2}f(X)] = 0$ , then we have  $f(x) = c$  on  $[0, 1]$ , where  $c$

Table 3: MSE and Bias of estimators to estimate the varextropy  $VJ(X)$  of the exponential distribution with mean equal to one.

$n$	MSE(Bias)				
	$VJV_{m,n}$	$VJD$	$VJB_n$	$VJS_n$	$VJQ_{m,n}$
10	0.4596275(0.0899478)	<b>0.0002119(-0.0125190)</b>	0.0003007(-0.0159281)	0.0003360(-0.0167951)	0.0745941(0.0470953)
20	0.0252052(0.0485333)	<b>0.0001861(-0.0129253)</b>	0.0002770(-0.0163934)	0.0003233(-0.0175661)	0.0074666(0.0248586)
30	0.0089156(0.0322723)	<b>0.0001754(-0.0127999)</b>	0.0002678(-0.0162028)	0.0003118(-0.0173596)	0.0023550(0.0179326)
40	0.0034209(0.0234392)	<b>0.0001673(-0.0125957)</b>	0.0002598(-0.0159925)	0.0002999(-0.0170549)	0.0012288(0.0140575)
50	0.0015958(0.0177894)	<b>0.0001614(-0.0124132)</b>	0.0002522(-0.0157660)	0.0002883(-0.0167369)	0.0006660( <b>0.0110643</b> )
100	0.0004385(0.0106603)	<b>0.0001415(-0.0117113)</b>	0.0002277(-0.0150125)	0.0002375(-0.0151743)	0.0002418( <b>0.0074678</b> )

is a constant and from the property of density function, it is obvious that  $f(x) = 1$ . Therefore, we can claim that  $VJ(X)$  is always nonnegative and its minimum value is uniquely obtained by the standard uniform distribution. The above discussion gives us the idea to use the proposed varextropy estimators as the test statistics of goodness-of-fit tests of uniformity.

Let  $X_1, \dots, X_n$  be a random sample from a continuous distribution function  $F(x)$  on  $[0, 1]$ . The null hypothesis is  $H_0 : F(x) = x$  and the alternative hypothesis (denoted  $H_1$ ) is the opposite of  $H_0$ . Given any significance level  $\alpha$ , our hypothesis-testing procedure can be defined by the critical region:

$$G_n = \hat{V}J(X) \geq C_{1-\alpha}, \quad (4.1)$$

where  $\hat{V}J(X)$  is one of the proposed estimators and  $C_{1-\alpha}$  is the critical value for the test with level  $\alpha$ . Notice when  $\hat{V}J(X) \xrightarrow{P} VJ(X)$ , under  $H_0$ ,  $G_n \xrightarrow{P} 0$  and  $H_1$ ,  $G_n$  converges to a number larger than zero in probability.

Due to the complexity of the calculations, it is not easy to determine the distribution of the test statistics under  $H_0$ , so the critical values are calculated using the Monte Carlo method. By utilizing five different varextropy estimators, we introduce the following test statistics for testing the uniformity:

$$GV_n = VJV_{m,n},$$

$$GD_n = VJD,$$

$$GB_n = VJB_n,$$

$$GS_n = VJS_n,$$

$$GQ_n = VJQ_{m,n}.$$

It is necessary to mention here that, also [Alizadeh Noughabi and Shafaei Noughabi \(2023\)](#) used varentropy for testing uniformity. They showed that  $VE(X) = 0$  if and only if  $f(x)$  is the standard

uniform density function and used their proposed varextropy estimators as the test statistics for goodness-of-fit tests of uniformity.

## 4.2 Power comparisons

In this subsection, we compare the powers of our proposed test statistics with Kolmogorov-Smirnov statistic (Kolmogorov (1933), Smirnov (1939)), and the test statistics proposed by Alizadeh Noughabi and Shafaei Noughabi (2023). The test statistic of Kolmogorov-Smirnov is given as follows:

$$KS = \max \left( \max_{1 \leq i \leq n} \left\{ \frac{i}{n} - X_{(i)} \right\}, \max_{1 \leq i \leq n} \left\{ X_{(i)} - \frac{i-1}{n} \right\} \right), \quad (4.2)$$

where  $X_{(1)}, \dots, X_{(n)}$  are order statistics. Moreover, the Alizadeh Noughabi-Shafaei Noughabi Statistics are given as:

$$TV_n = \frac{1}{n} \sum_{i=1}^n \log^2 (X_{(i+m)} - X_{(i-m)}) - \left[ \frac{1}{n} \sum_{i=1}^n \log (X_{(i+m)} - X_{(i-m)}) \right]^2,$$

$$TE_n = \frac{1}{n} \sum_{i=1}^n \log^2 \left( \frac{c_i m/n}{X_{(i+m)} - X_{(i-m)}} \right) - \left[ \frac{1}{n} \sum_{i=1}^n \log \left( \frac{c_i m/n}{X_{(i+m)} - X_{(i-m)}} \right) \right]^2,$$

where  $X_{(i)} = X_{(1)}$  if  $i < 1$  and  $X_{(i)} = X_{(n)}$  if  $i > n$ ,

$$TD_n = \int_{-\infty}^{+\infty} \hat{f}(x) [\log \hat{f}(x)]^2 dx - \left[ \int_{-\infty}^{+\infty} \hat{f}(x) \log \hat{f}(x) dx \right]^2,$$

where  $\hat{f}$  is the kernel density function estimation of  $f$  and is defined by

$$\hat{f}(x) = \frac{1}{nh} \sum_{j=1}^n k \left( \frac{x - X_j}{h} \right),$$

$$TB_n = \frac{1}{n} \sum_{i=1}^n (\log \hat{f}(X_{(i)}))^2 - \left[ \frac{1}{n} \sum_{i=1}^n \log \hat{f}(X_{(i)}) \right]^2,$$

where  $\hat{f}(X_{(i)}) = \frac{1}{n} \sum_{j=1}^n k \left( \frac{X_{(i)} - X_j}{h} \right)$ ,

$$TC_n = \frac{1}{n} \sum_{i=1}^n \log^2 \left( \frac{\sum_{j=i-m}^{i+m} (X_{(j)} - \bar{X}_{(i)})(j-i)}{n \sum_{j=i-m}^{i+m} (X_{(j)} - \bar{X}_{(i)})^2} \right) - \left[ \frac{1}{n} \sum_{i=1}^n \log \left( \frac{\sum_{j=i-m}^{i+m} (X_{(j)} - \bar{X}_{(i)})(j-i)}{n \sum_{j=i-m}^{i+m} (X_{(j)} - \bar{X}_{(i)})^2} \right) \right]^2,$$

where  $\bar{X}_{(i)} = \frac{1}{2m+1} \sum_{j=i-m}^{i+m} X_{(j)}$ , and

$$TA_n = \frac{1}{n} \sum_{i=1}^n \log^2 \{ \hat{f}(X_{(i+m)}) + \hat{f}(X_{(i-m)}) \} - \left[ \frac{1}{n} \sum_{i=1}^n \log \{ \hat{f}(X_{(i+m)}) + \hat{f}(X_{(i-m)}) \} \right]^2. \quad (4.3)$$

In the above statistics,  $m$  is the window size as defined in Subsection 2.5. For power comparisons,

Table 4: Percentage points of the proposed test statistics at the level  $\alpha = 0.05$ .

n	$GV_n$	$GD_n$	$GB_n$	$GS_n$	$GQ_n$
10	0.9102	0.0665	0.0574	0.0558	0.4024
20	0.4937	0.0485	0.0374	0.0277	0.1813
30	0.2845	0.0413	0.0297	0.0205	0.1365
40	0.1933	0.0371	0.0256	0.0169	0.0767
50	0.1389	0.0343	0.0226	0.0148	0.0583
75	0.0872	0.0296	0.0183	0.0116	0.0385
100	0.0715	0.0270	0.0158	0.0098	0.0320

we compute the powers of our proposed tests and the powers of tests mentioned above under the following alternative distributions:

$$\begin{aligned}
 A_k : F(x) &= 1 - (1 - x)^k, \quad 0 \leq x \leq 1 \quad (\text{for } k = 1.5, 2); \\
 B_k : F(x) &= \begin{cases} 2^{k-1}x^k, & 0 \leq x \leq 0.5 \\ 1 - 2^{k-1}(1 - x)^k, & 0.5 \leq x \leq 1 \end{cases} \quad (\text{for } k = 1.5, 2, 3); \\
 C_k : F(x) &= \begin{cases} 0.5 - 2^{k-1}(0.5 - x)^k, & 0 \leq x \leq 0.5 \\ 0.5 + 2^{k-1}(x - 0.5)^k, & 0.5 \leq x \leq 1 \end{cases} \quad (\text{for } k = 1.5, 2).
 \end{aligned}$$

[Stephens \(1974\)](#) uses these alternative distributions in his study on power comparisons of some uniformity tests.

Based on 100,000 repetitions, the critical values are estimated as shown in Table 4. Tables 5 and 6 contain the results of simulations of our proposed test statistics and competitive test statistics with 100,000 repetitions for  $n = 10, 20, 30$  and  $\alpha = 0.05$ . The bold type in these tables indicates the statistic achieving the maximal power. According to Tables 5 and 6, the performance of tests depends on alternative distributions. Against alternative  $A$ , our proposed test  $GD_n$  and KS-based test give competitive powers to different samples. For sample size ( $n = 10$ ), the test based on  $GD_n$  is the best. For moderate sample size ( $n = 20$ ), if alternative distribution is  $A_{1.5}$ ,  $GD_n$  is better and otherwise, for  $n = 30$ , KS test statistic performs better than the others. Alizadeh Noughabi-Shafaei Noughabi test statistics have very low power in this case. For alternative distribution  $B$ , our proposed test  $GD_n$  and  $GB_n$  are the best tests and other tests have low powers in this case, especially the performance of  $TV$ ,  $TE$ ,  $TC$ ,  $TA$  and  $KS$  is poor. If the alternative is  $C$ , the test based on  $TC$  is the best test, while the behavior of our proposed statistic  $GQ_n$  is better than the performance of KS.



Table 5: Power comparisons of the proposed tests at a significance level of 0.05.

n	Alternative	$GV_n$	$GD_n$	$GB_n$	$GS_n$	$GQ_n$
10	$A_{1.5}$	0.0739	<b>0.1767</b>	0.1072	0.1663	0.0764
	$A_2$	0.1150	<b>0.3959</b>	0.2033	0.3423	0.1327
	$B_{1.5}$	0.0719	0.1589	<b>0.1629</b>	0.1109	0.0456
	$B_2$	0.1126	<b>0.3494</b>	0.3292	0.2167	0.0624
	$B_3$	0.2156	<b>0.7326</b>	0.6421	0.5038	0.1294
	$C_{1.5}$	0.0808	0.0415	0.0296	0.0573	<b>0.1182</b>
	$C_2$	0.1295	0.0406	0.0300	0.0660	<b>0.2160</b>
20	$A_{1.5}$	0.0884	<b>0.2889</b>	0.1575	0.2497	0.1088
	$A_2$	0.1684	<b>0.6680</b>	0.3648	0.5377	0.2489
	$B_{1.5}$	0.0879	0.2424	<b>0.2543</b>	0.0977	0.0444
	$B_2$	0.1640	<b>0.5707</b>	0.5493	0.2611	0.1061
	$B_3$	0.3689	<b>0.9459</b>	0.8948	0.7059	0.3549
	$C_{1.5}$	0.09628	0.0373	0.0251	0.1010	<b>0.1812</b>
	$C_2$	0.1900	0.0440	0.0281	0.1339	<b>0.3924</b>
30	$A_{1.5}$	0.1114	<b>0.3933</b>	0.2092	0.3206	0.1580
	$A_2$	0.2425	<b>0.8319</b>	0.5078	0.6832	0.3994
	$B_{1.5}$	0.1100	0.3160	<b>0.3416</b>	0.1080	0.0690
	$B_2$	0.2333	<b>0.7251</b>	0.7132	0.3674	0.2224
	$B_3$	0.5500	<b>0.9907</b>	0.9736	0.8706	0.6455
	$C_{1.5}$	0.1236	0.0385	0.0228	0.1409	<b>0.2450</b>
	$C_2$	0.2827	0.0498	0.0263	0.2118	<b>0.5757</b>

## 5 Real data

**Example 5.1.** *Illowsky and Dean (2018)* used 55 data set of babies' smiling time measured in seconds. The data follows a  $U(0, 23)$  distribution. We transformed this data to  $U(0, 1)$  and obtained test statistics as:

$$GV_n = 0.01735047, GD_n = 0.02292441, GB_n = 0.01267519, GS_n = 0.002416893, GQ_n = 0.008132604.$$

The values belong to the acceptance region. Therefore, Assumption  $H_0$  is accepted, that is, the data follows the standard uniform distribution.

**Example 5.2.** We consider the data of 20 wild lizards collected by the zoologists in the southwestern United States. The total length (mm) of each given as follows:

179, 157, 169, 146, 143, 131, 159, 142, 141, 130, 142, 116, 130, 140, 138, 137, 134, 114, 90, 114.

Using common tests such as Kolmogorov Smirnov, Shapiro-Wilk and Anderson-Darling, it can be

Table 6: Power comparisons of the Alizadeh Noughabi-Shafaei Noughabi tests and KS test at a significance level of 0.05.

n	Alternative	TV	TE	TD	TB	TC	TA	KS
10	$A_{1.5}$	0.0526	0.0625	0.0811	0.0814	0.0642	0.0832	<b>0.1542</b>
	$A_2$	0.0828	0.1014	0.1311	0.1244	0.0883	0.1427	<b>0.3871</b>
	$B_{1.5}$	0.0295	0.0213	<b>0.0826</b>	0.0810	0.0298	0.0414	0.0367
	$B_2$	0.0195	0.0213	0.1110	<b>0.1226</b>	0.0169	0.0422	0.0447
	$B_3$	0.0185	0.0319	<b>0.1873</b>	0.1810	0.0264	0.0609	0.0873
	$C_{1.5}$	0.1627	0.1409	0.0530	0.0519	<b>0.1779</b>	0.0736	0.1148
	$C_2$	0.3329	0.2909	0.0733	0.0620	<b>0.3530</b>	0.0924	0.2001
20	$A_{1.5}$	0.0731	0.1185	0.1421	0.1429	0.0804	0.1502	<b>0.2796</b>
	$A_2$	0.1523	0.2684	0.2924	0.2992	0.1772	0.3322	<b>0.6979</b>
	$B_{1.5}$	0.0082	0.0335	0.1526	<b>0.1582</b>	0.0196	0.0585	0.0579
	$B_2$	0.0053	0.0593	0.3020	<b>0.3067</b>	0.0126	0.0848	0.1246
	$B_3$	0.0093	0.1317	0.5122	<b>0.5166</b>	0.0385	0.1746	0.4133
	$C_{1.5}$	0.2930	0.2316	0.0324	0.0334	<b>0.2933</b>	0.1027	0.1501
	$C_2$	0.6430	0.5415	0.0425	0.0384	<b>0.6673</b>	0.1326	0.3159
30	$A_{1.5}$	0.0942	0.1835	0.2001	0.2039	0.1155	0.2326	<b>0.4066</b>
	$A_2$	0.2342	0.4228	0.4429	0.4532	0.2913	0.5028	<b>0.8751</b>
	$B_{1.5}$	0.0053	0.0582	0.2433	<b>0.2441</b>	0.0192	0.1215	0.0837
	$B_2$	0.0083	0.1622	0.4851	<b>0.4946</b>	0.0339	0.2716	0.2411
	$B_3$	0.0299	0.3916	0.7583	<b>0.7629</b>	0.1242	0.5309	0.7524
	$C_{1.5}$	0.4023	0.3224	0.0259	0.0232	<b>0.4029</b>	0.0832	0.1892
	$C_2$	0.8224	0.7425	0.0353	0.0244	<b>0.8450</b>	0.0942	0.4383

seen that the data follows a normal distribution. Now we want to transform the data into standard uniform distribution using the integral probability transformation. Therefore if  $U_i = F_0(X_i)$  for  $i = 1, \dots, n$ , where  $F_0(X_i)$  is distribution function of the normal distribution with estimated mean and variance of the data. The transform data is obtained as follows:

0.9804, 0.8326, 0.9408, 0.6620, 0.6056, 0.3715, 0.8562, 0.5864, 0.5670, 0.3530, 0.5864, 0.1419, 0.3530, 0.5475, 0.5081, 0.4884, 0.4289, 0.1205, 0.0091, 0.1205.

Next, the values of the proposed test statistics based on transformed sample are computed as:  $GV_n = 0.1453019$ ,  $GD_n = 0.03378436$ ,  $GB_n = 0.02535139$ ,  $GS_n = 0.002548617$ ,  $GQ_n = 0.03390633$ . By comparing with the percentage points of the test statistics at the 0.05 level given in Table 4, which are equal to 0.4937, 0.0485, 0.0374, 0.0277 and 0.1813, respectively, since the values of the test statistics are smaller than the corresponding critical values it can be conclude that the

assumption of normality of the data cannot be rejected at the 0.05 significance level.

**Example 5.3.** In this example, the data set symbolizes the simulated strengths of glass fibers presented by [Mahmoud and Mandouh \(2000\)](#)

1.014, 1.081, 1.082, 1.185, 1.223, 1.248, 1.267, 1.271, 1.272, 1.275, 1.276, 1.278, 1.286, 1.288, 1.292, 1.304, 1.306, 1.355, 1.361, 1.364, 1.379, 1.409, 1.426, 1.459, 1.460, 1.476, 1.481, 1.484, 1.501, 1.506, 1.524, 1.526, 1.535, 1.541, 1.568, 1.579, 1.581, 1.591, 1.593, 1.602, 1.666, 1.670, 1.684, 1.691, 1.704, 1.731, 1.735, 1.747, 1.748, 1.757, 1.800, 1.806, 1.867, 1.876, 1.878, 1.910, 1.916, 1.972, 2.012, 2.456, 2.592, 3.197, 4.121.

The values of the proposed test statistics after obtaining transformed sample are  $GV_n = 0.4381156$ ,  $GD_n = 0.05200372$ ,  $GB_n = 0.03980463$ ,  $GS_n = 0.03055442$ ,  $GQ_n = 0.05855942$ . By looking at the Table 4, we can see that because the test statistics are larger than the percentage points obtained for  $n=63$ , therefore, the assumption of normality will be rejected at the significance level of  $\alpha = 0.05$ .

**Example 5.4.** The following data had presented in [Fuller et al. \(1994\)](#) and represents the strength of glass for aircraft window.

18.83, 20.8, 21.657, 23.03, 23.23, 24.05, 24.321, 25.5, 25.52, 25.8, 26.69, 26.77, 26.78, 27.05, 27.67, 29.9, 31.11, 33.2, 33.73, 33.76, 33.89, 34.76, 35.75, 35.91, 36.98, 37.08, 37.09, 39.58, 44.045, 45.29, 45.381.

By using Kolmogorov-Smirnov (K-S) distance test statistic, [Alshenawy \(2020\)](#) showed that, among the distributions fitted to the data, including the exponential and A distributions, A distribution have the best fitted for data. Now by calculating the proposed test statistics after obtaining transformed sample, we show that A distribution, unlike the exponential distribution, is a suitable distribution to fit the data. Since, for A distribution, test statistics are lower than critical values

Table 7: The CDF, MLE and the proposed test statistics.

Model	CDF	MLE	$GV_n$	$GD_n$	$GB_n$	$GS_n$	$GQ_n$
Exponential	$1 - \exp(-\beta x)$	0.032	0.7052102	0.2151081	0.1541674	0.08482313	0.07347814
A	$\exp(\frac{1}{\beta}(1 - \exp(\frac{\beta}{x})))$	125.662	0.1404678	0.02569515	0.0144915	0.01595046	0.07580088

0.1875, 0.04088, 0.028615, 0.020166 and 0.07852028, thus it appears to fit the data very well.

## 6 Conclusion

In this paper, we proposed estimators of varextropy. The consistency of the proposed estimators is proved. We employed Monte-Carlo simulation to derive the mean squared error (MSE) and bias of the estimators across various distributions. In addition, we assessed the suggested estimators based on their bias and MSE characteristics. We introduced some goodness-of-fit tests of

uniformity using the proposed estimators. Then, we obtained the critical and power values of the proposed tests by Monte Carlo simulation and compared the powers with the competitive powers in Alizadeh Noughabi and Shafaei Noughabi (2023) and Kolmogorov-Smirnov. Our simulation findings demonstrated that the suggested tests perform well in comparison with the other tests of uniformity.

## 7 Declarations

**Conflict of interest** The authors declare no conflict of interest.

## References

- Alizadeh Noughabi, H., and Jarrahiferiz, J. (2019), ‘On the estimation of extropy’, *Journal of Nonparametric Statistics*, 31(1), 88-99.
- Alizadeh Noughabi, H., and Shafaei Noughabi, M. (2023), ‘Varentropy estimators with applications in testing uniformity’, *Journal of Statistical Computation and Simulation*, Doi: 10.1080/00949655.2023.2196627
- Al-Labadi, L., and Berry, S. (2022), ‘Bayesian estimation of extropy and goodness of fit tests’, *Journal of Applied Statistics*, 49, 357–370.
- Alshenawy, R. (2020), ‘A new one parameter distribution: properties and estimation with applications to complete and type II censored data’, *Journal of Taibah University for Science*, 14(1), 11–18.
- Becerra, A., de la Rosa, J. I., González, E., Pedroza, A. D., and Escalante, N. I. (2018), ‘Training deep neural networks with non-uniform frame-level cost function for automatic speech recognition’, *Multimedia Tools and Applications*, 77, 27231–27267.
- Correa, J.C. (1995), ‘A new estimator of entropy’, *Communication in Statistics-Theory and Methods*, 24(10), 2439–2449.
- Ebrahimi, N., Pflughoeft, K., and Soofi, E. (1994), ‘Two measures of sample entropy’, *Statistics and Probability Letters*, 20(3), 225–234.
- Fradelizi, M., Madiman, M., and Wang, L. (2016), ‘Optimal concentration of information content for logconcave densities’. In C. Houdré, D. Mason, P. Reynaud-Bouret & J. Rosiński (eds.), *High Dimensional Probability VII. Progress in Probability*, vol. 71, Cham, Springer, pp. 45-60.

- Fuller, E.J., Frieman, S., Quinn, J., Quinn, G., Carter, W. (1994), ‘Fracture mechanics approach to the design of glass aircraft windows’, *A case study. International Society for Optics and Photonics*, 419–430.
- Furuichi, S., and Mitroi, F.C. (2012), ‘Mathematical inequalities for some divergences’, *Physica A*, 391, 388–400.
- Goodarzi, F. (2024), ‘Results on conditional variance in parallel system and lower bounds for varextropy’, *Communication in Statistics-Theory and Methods*, 53, 1590-1610.
- Gneiting, T., and Raftery, A.E. (2007), ‘Strictly proper scoring rules, prediction, and estimation’, *Journal of the American statistical Association*, 102(477), 359–378.
- Grzegorzewski, P., and Wieczorkowski, R. (1999), ‘Entropy-based goodness-of-fit test for exponentiality’, *Communications in Statistics-Theory and Methods*, 28, 1183–1202.
- Illovsky, B., and Dean, S. (2018), ‘Introductory statistics’. Houston: OpenStax.
- Jiang, J., Wang, R., Pezeril, M., and Wang, Q.A. (2011), ‘Application of varentropy as a measure of probabilistic uncertainty for complex networks’, *Science Bulletin*, 56, 3677–3682.
- Kolmogorov, A.N. (1933), ‘Sulla determinazione empirica di una legge didistribuziane’, *Giornale Z. dell’IstitutaItaliano degli Attuari*, 4, 83–91.
- Lad, F., Sanfilippo, G., and Agrò, G. (2015), ‘Extropy: Complementary Dual of Entropy’, *Statistical Science*, 30, 40-58.
- Li, J., Fradelizi, M., and Madiman, M. (2016), ‘Information concentration for convex measures’, *IEEE International Symposium on Information Theory*, Barcelona, 1128-1132.
- Madiman, M., and Barron, A. (2007), ‘Generalized entropy power inequalities and monotonicity properties of information’, *IEEE Transactions on Information Theory*, 53, 2317-2329.
- Mahmoud, M.R., and Mandouh, R.M. (2013), ‘On the transmuted frechet distribution’, *Journal of Applied Sciences Research*, 9(10), 5553-5561.
- Shannon, C.E. (1948), ‘A mathematical theory of communication’, *The Bell System Technical Journal*, 27 (1948), 379-432.
- Silverman, B.W. (1978), ‘Weak and strong uniform consistency of the kernel estimate of a density and its derivatives’, *The annals of statistics*, 6, 177-184.
- Smirnov, N. (1939), ‘Estimate of derivation between empirical distribution functions in two independent samples’ (in Russian). *Bulletin Moscow University*, 2, 3–16.

- Soni, P., Dewan, I., and Jain, K. (2012), ‘Nonparametric estimation of quantile density function’, *Computational Statistics and Data Analysis*, 56(12), 3876–3886.
- Stephens, M.A. (1974), ‘EDF statistics for goodness of fit and some comparisons’, *Journal of the American Statistical Association*, 69(347), 730–737.
- Subhash, S., Sunoj, S.M., Sankaran, P.G., and Rajesh, G. (2023), ‘Nonparametric estimation of quantile-based entropy function’, *Communications in Statistics - Simulation and Computation*, 52, 1805-1821.
- Qiu, G., and Jia, K. (2018a), ‘Extropy estimators with applications in testing uniformity’, *Journal of Nonparametric Statistics*, 30(1), 182–196.
- Qiu, G., and Jia, K. (2018b), ‘The residual extropy of order statistics’, *Statistics and Probability Letters*, 133, 15-22.
- van Es, B. (1992), ‘Estimating functionals related to a density by a class of statistics based on spacings’, *Scandinavian Journal of Statistics*, 19(1), 61–72.
- Vaselabadi, N.M., Tahmasebi, S., Kazemi, M.R., and Buono, F. (2021), ‘Results on varextropy measure of random variables’, *Entropy*, 23, 356.
- Vasicek, O. (1976), ‘A test for normality based on sample entropy’, *Journal of the Royal Statistical Society: Series B*, 38, 54–59.
- Vontobel, P.O. (2013), ‘The Bethe permanent of a nonnegative matrix’, *IEEE Transactions on Information Theory*, 59, 1866–1901.
- Yang, J., Xia, W., and Hu, T. (2019), ‘Bounds on extropy with variational distance constraint’, *Probability in the Engineering and Informational Sciences*, 33(2), 186–204.
- Zamanzade, E. (2015), ‘Testing uniformity based on new entropy estimators’, *Journal of Statistical Computation and Simulation*, 85(16), 3191-3205.