# THERMODYNAMICS OF VISCOUS FLOWS ON SURFACES

A. DUYUNOVA, V. LYCHAGIN, AND S. TYCHKOV

Abstract. In this paper, thermodynamics of a moving medium is considered. Main goal of this paper is to describe thermodynamics state equations and equations of co-existence manifolds of media on two- and three-dimensional manifolds. To this end the phase space of a medium is extended with the deformation tensor and the stress tensor.

# 1. INTRODUCTION

This paper we continues our study [\[2\]](#page-4-0) of thermodynamics of a moving medium. We consider a two- and three-dimensional flows of a medium and introduce an extended thermodynamic phase space that includes tensor quantities, namely, deformation  $\Delta$  and a stress  $\sigma$  together with the ordinary quantities: mass density  $\rho$ , entropy density s, energy density e, temperature  $T$  and chemical potential  $\eta$ . To analyze this phase space we use the methods, which are usually applied (cf. [\[3\]](#page-4-1)) to the classical case only. Note that by a moving medium we mean a medium that flows with a non-zero acceleration.

This paper illustrates the difference between still and moving media from the thermodynamics standpoint.

Recall [\[1\]](#page-4-2) that the generalized Navier–Stokes system describing motion of a fluid on an oriented *n*-dimensional Riemannian manifold  $(M, g)$  has the form

<span id="page-0-0"></span>(1.1)  
\n
$$
\begin{cases}\n\rho \left( \frac{\partial X}{\partial t} + \nabla_X X \right) = \text{div} \,\sigma, \\
\frac{\partial \rho}{\partial t} + X(\rho) + \rho \,\text{div} \, X = 0, \\
c_p \rho \left( \frac{\partial T}{\partial t} + X(T) \right) = \langle \sigma, \Delta \rangle + \text{div}(\varkappa \,\text{grad}\, T)\n\end{cases}
$$

where  $\rho$  and e are the mass and the energy densities,  $\sigma \in T^*M \otimes TM$  and  $\Delta \in$  $TM \otimes T^*M$  are the stress and the deformation tensor fields, X is the flow velocity field,  $\varkappa$  is a thermal conductivity of the medium,  $c_p$  is a heat capacity at constant pressure.

The deformation tensor  $\Delta$  is defined as  $\Delta = d_{\nabla} X \in TM$ , where TM is the tangent space at a point of M, and  $\nabla$  is the Levi-Civita connection associated with g, and  $d\sigma$  is the covariant differential.

Note that if a medium flows with a constant velocity, which means  $\Delta = 0$ , the usual thermodynamics is applied.

The paper is organized as follows. In Section [2](#page-1-0) we recall main results regarding the geometrical approach to thermodynamics. The notion of a Newton medium

<sup>2010</sup> *Mathematics Subject Classification.* Primary 76S05, 58J37.

*Key words and phrases.* thermodynamics, Navier–Stokes equation, co-existence curve.

is used to determine how the thermodynamic state depends on the deformation tensor. In Section [3](#page-2-0) these constructions are applied to two- and three-dimensional flows. We calculate a co-existence curve for phases of moving medium. In the Sections [4](#page-2-1) and [5,](#page-3-0) we study a special case of flows on a two-dimensional manifold and give an example of the Navier–Stokes equations for flows on a plane.

# 2. Thermodynamics of moving media

<span id="page-1-0"></span>In this section, the necessary for the paper thermodynamics constructions are recalled.

The thermodynamic state of the fluid in a small neighborhood of a point is given by the extensive quantities: mass density  $\rho$ , internal energy density  $e$ , entropy density s, deformation tensor  $\Delta$ , and the intensive ones: chemical potential  $\eta$ , temperature T and stress tensor  $\sigma$ .

Since we consider flows on an oriented Riemannian  $(M, q)$ , its thermodynamic phase space is a bundle  $\Phi \to M$  with a fiber

$$
\Phi_a = \mathbb{R}^5 \times \operatorname{End} \mathrm{T}_a^* M \times \operatorname{End} \mathrm{T}_a M
$$

at a point  $a, \Phi_a = (s, \eta, \rho, T, e, \sigma, \Delta)$ . This space is equipped with a contact structure

$$
\alpha = de - T dS - \sum_{i,j} \sigma_{ij} d\Delta_{ij} - \eta d\rho.
$$

The Levi-Civita connection  $\nabla$  allows to compare thermodynamic states at different points of the manifold  $M$  with contact transformations generated by the parallel transports. This means that the thermodynamics states of the moving medium is invariant with respect to the Riemannian holonomy group of the manifold M.

The thermodynamic state can be given with a single function  $e = e(s, \rho, \Delta)$ . But it is more convenient to introduce the density of Helmholtz free energy  $h =$  $h(\rho, T, \Delta) = e(s, \rho, \Delta) - Ts$ . In terms of h, we have  $\sigma = h_{\Delta}$ ,  $\eta = h_{\rho}$ ,  $e = h - Th_T$ . In terms of the Helmholtz free energy, the pseudo-Riemannian structure  $\kappa$  has the form

<span id="page-1-1"></span>(2.1)  

$$
\kappa = \frac{1}{T} \left( \frac{\partial^2 h}{\partial T^2} dT^2 - \frac{\partial^2 h}{\partial \rho^2} d\rho^2 - \sum_{i,j=1}^n \sum_{k,l=1}^n \frac{\partial^2 h}{\partial \Delta_{ij} \partial \Delta_{kl}} d\Delta_{ij} \cdot d\Delta_{kl}
$$

$$
- 2 \sum_{i,j=1}^n \frac{\partial^2 h}{\partial \Delta_{ij} \partial \rho} d\Delta_{ij} \cdot d\rho).
$$

We consider Newtonian media, i. e. media that satisfy 'Hooke's law', i. e. the stress tensor  $\sigma$  depends on the deformation  $\Delta$  linearly. Also, since  $\sigma_{ij} = h_{\Delta_{ij}}$ , we get that the Helmholtz free energy density is a quadratic function in  $\Delta$ .

As we mentioned above, it is required that the thermodynamic state is  $SO(g)$ -invariant [\[1\]](#page-4-2). Thus free Helmholtz energy is a function of  $SO(g)$ -invariants of the deformation  $\Delta$ ,

$$
h = \frac{1}{2} \left( a(\rho, T) \operatorname{Tr} \Delta^2 + b(\rho, T) \operatorname{Tr} \Delta \Delta' + c(\rho, T) \operatorname{Tr}^2 \Delta \right) + d(\rho, T) \operatorname{Tr} \Delta + e(\rho, T),
$$

where  $\Delta'$ ,  $\Delta'_{ij} = \sum$  $_{k,l}$  $g_{ik}^{-1}\Delta_{lk}g_{lj}$ , is the operator adjoint to  $\Delta$ , and a, b, c, d, e are some functions. The stress tensor takes the form

$$
\sigma = a(\rho, T)\Delta' + b(\rho, T)\Delta + (c(\rho, T) \operatorname{Tr} \Delta + d(\rho, T))\mathbf{1}.
$$

It is an established assumption [\[6\]](#page-4-3) that the stress is equal to the negative thermodynamic pressure p when the deformation vanishes, i. e.,  $p = -d$ .

Recall  $[1]$  that thermodynamic pressure  $p$  is expressed in terms of the Helmholtz free energy density  $h_0(\rho, T)$  for still media as follows,

$$
p = \rho \frac{\partial h_0(\rho, T)}{\partial \rho} - h_0(\rho, T).
$$

### 3. Thermodynamics of two- and three-dimensional flows

<span id="page-2-0"></span>In this section, the general picture we discussed in Section [2](#page-1-0) is applied to twoand three-dimensional flows.

Since we consider an isotropic Newtonian medium, the Helmholtz free energy density is a quadratic function in  $\Delta$ . Thus we have

$$
h = \frac{1}{2} \left( (\mu + \tau) \operatorname{Tr} \Delta^2 + (\mu - \tau) \operatorname{Tr} \Delta \Delta' + \left( \zeta - \frac{2}{n} \mu \right) \operatorname{Tr}^2 \Delta \right) - p \operatorname{Tr} \Delta + h_0.
$$

Here  $n = 2, 3$  is the dimension of the manifold.

Calculating the stress tensor  $\sigma = h_{\Delta}$ , we get a well-known linear function in  $\Delta$ , that is,

$$
\sigma = -p\mathbf{1} + \mu(\Delta + \Delta') + \tau(\Delta - \Delta') + \left(\zeta - \frac{2}{n}\mu\right) \operatorname{Tr} \Delta \mathbf{1}.
$$

Here  $\mu$  is the viscosity,  $\zeta$  is the second, or volume, viscosity, and  $\tau$  is the rotational viscosity. These quantities are considered as functions of the density  $\rho$  and the temperature T.

For brevity, we introduce the following notation:

$$
d_1 = \text{Tr}\,\Delta, \quad d_2 = \text{Tr}\,\Delta^2, \quad d_3 = \text{Tr}\,\Delta\Delta'.
$$

Substituting this function into [\(2.1\)](#page-1-1), we obtain a formula for the pseudo-Riemannian, which is quite long, so we do not give it here.

Calculating the condition for degeneracy of this form, we get the following lemma.

Lemma 3.1. Co-existence manifold of a moving Newton medium is defined by the equation:

$$
\frac{\partial^2 h}{\partial T^2} \left( \left( d_2 + d_3 - \frac{2}{n} d_1^2 \right) \mu^2 \frac{\partial}{\partial \rho} \left( \frac{1}{\mu} \frac{\partial}{\partial \rho} \ln \mu \right) + (d_2 - d_3) \tau^2 \frac{\partial}{\partial \rho} \left( \frac{1}{\tau} \frac{\partial}{\partial \rho} \ln \tau \right) + d_1^2 \zeta^2 \frac{\partial}{\partial \rho} \left( \frac{1}{\zeta} \frac{\partial}{\partial \rho} \ln \zeta \right) - 2d_1 \left( \frac{\partial^2 p}{\partial \rho^2} + \frac{\partial p}{\partial \rho} \frac{\partial}{\partial \rho} \ln \zeta \right) + \left( \frac{\partial p}{\partial \rho} \right)^2 + 2 \frac{\partial^2 h_0}{\partial \rho^2} \right) = 0.
$$

#### 4. Thermodynamics of flows on surfaces

<span id="page-2-1"></span>In this section, we study a special case of flows on a two-dimensional manifold with a metric g and the associated volume form  $\Omega<sub>g</sub>$ .

Denote the Riemannian holonomy group at a point  $a \in M$  as  $H(M, a) \in SO(g)$ . Then the thermodynamic state given by the Helmholtz free energy has to be a function quadratic in  $H(M, a)$ -invariants.

Consider the first non-trivial case of the Berger classification [\[5\]](#page-4-4) of irreducible Riemannian holonomy groups,  $H(M, a) = U(1)$ , dim  $M = 2$ , and M is a Kähler manifold.

In this case, there is a complex structure on  $M, J \in \text{End }TM$ , i.e.,  $J^2 = -1, J$ is an isometry of the metric  $g$ , and  $J$  is preserved by the parallel transport.

Lemma 4.1. All first and second degree polynomial invariants are generated by the invariants

$$
t_1 = \text{Tr }\Delta
$$
,  $t_2 = \text{Tr }J\Delta$ ,  $t_3 = \text{Tr }\Delta(\Delta + \Delta')$ .

Proof. Let us enumerate all possible first and second order invariants of the operators J,  $\Delta$ ,  $\Delta'$ . According to the Procesi theorem [\[7\]](#page-4-5), the following invariants generate all polynomial invariants of order equal or less than two.

$$
t_1 = \text{Tr }\Delta, \quad t_2 = \text{Tr } J\Delta, \quad t_3 = \text{Tr }\Delta(\Delta + \Delta'), \quad t_4 = \text{Tr }\Delta(\Delta - \Delta'),
$$
  
 $t_5 = \text{Tr } J\Delta^2, \quad t_6 = \text{Tr } J\Delta' J\Delta, \quad t_7 = \text{Tr } (J\Delta)^2.$ 

In this list, invariants that vanish and those which coincide up to a sign are omitted. Straightforward computations show that

$$
t_4 = t_2^2
$$
,  $t_5 = t_1 t_2$ ,  $t_6 = \frac{t_3}{2} - \frac{t_2^2}{2} - t_1^2$ ,  $t_7 = \frac{t_3}{2} + \frac{t_2^2}{2} - t_1^2$ .

Note that these relations between invariants  $t_1, \ldots, t_7$  are true only for flows on surfaces.  $\Box$ 

Thus, the density of the free Helmholtz energy for Newtonian media has the form

$$
h = \frac{1}{2} \left( \mu \operatorname{Tr} \Delta(\Delta + \Delta') + \tau \operatorname{Tr}^2 J \Delta + \alpha (\operatorname{Tr} \Delta)(\operatorname{Tr} J \Delta) + (\zeta - \mu) \operatorname{Tr}^2 \Delta \right) - (p \operatorname{Tr} \Delta + q \operatorname{Tr} J \Delta) + h_0.
$$

Note that, in addition to the three viscosities  $\mu$ ,  $\tau$ ,  $\zeta$  (see Section [3\)](#page-2-0) and the pressure p, we have another viscosity-like quantity  $\alpha$  and a 'rotational' pressure q.

Substituting this quadratic equation of state into  $(2.1)$  we get the following lemma.

Lemma 4.2. Co-existence curve of a moving Newton medium on a surface is defined by the equation:

$$
h_{TT} \left( \left( \left( \left( t_1^2 - t_4 \right) \mu_{\rho \rho} - \eta_{\rho \rho} t_1^2 - \alpha_{\rho \rho} t_1 t_2 - \tau_{\rho \rho} t_2^2 + 2 p_{\rho \rho} t_1 + 2 q_{\rho \rho} t_2 - 2 h_{0, \rho \rho} \right) \alpha^2 \right. \\
\left. + 8 \left( \tau_{\rho} t_2 + \frac{\alpha_{\rho} t_1}{2} - q_{\rho} \right) \left( \eta_{\rho} t_1 + \frac{\alpha_{\rho} t_2}{2} - p_{\rho} \right) \alpha + \left( \left( 4 \left( t_4 - t_1^2 \right) \mu_{\rho \rho} + 4 \eta_{\rho \rho} t_1^2 \right) \right. \\
\left. + 4 \alpha_{\rho \rho} t_1 t_2 + 4 \tau_{\rho \rho} t_2^2 - 8 p_{\rho \rho} t_1 - 8 q_{\rho \rho} t_2 + 8 h_{0, \rho \rho} \right) \tau - 8 \left( \tau_{\rho} t_2 + \frac{\alpha_{\rho} t_1}{2} - q_{\rho} \right)^2 \right) \eta \\
\left. - 8 \tau \left( \eta_{\rho} t_1 + \frac{\alpha_{\rho} t_2}{2} - p_{\rho} \right)^2 \right) \mu + 8 \left( t_1^2 - t_4 \right) \left( \tau \eta - \frac{\alpha^2}{4} \right) \mu_{\rho}^2 \right) = 0.
$$

### 5. Navier–Stokes equations on a plane

<span id="page-3-0"></span>As an example, we consider Navier–Stokes equations on a Euclidean plane  $(g =$  $dx^2+dy^2$ , given that the thermodynamic state equation  $h = h(\rho, T, \Delta)$  is quadratic in  $\Delta$ , and viscosities  $\mu$ ,  $\tau$ ,  $\zeta$  and  $\alpha$  are constants.

Let x, y be coordianates on the plane,  $u(t, x, y)\partial_x + v(t, x, y)\partial_y$  be the flow velocity field.

Then the stress tensor  $\sigma = h_{\Delta}$  is given by

$$
\sigma = (-p + (\mu + \zeta)u_x + (\zeta - \mu)v_y + \alpha(u_y - v_x))\partial_x \otimes dx
$$
  
+
$$
(-q + (\mu + \tau)u_y + (\mu - \tau)v_x + \alpha(u_x + v_y))\partial_x \otimes dy
$$
  
-
$$
(-q - (\mu - \tau)u_y - (\mu + \tau)v_x + \alpha(u_x + v_y))\partial_y \otimes dx
$$
  
+
$$
(-p + (\mu + \zeta)v_y + (\zeta - \mu)u_x + \alpha(u_y - v_x))\partial_y \otimes dy
$$

Substituting the stress tensor into the first equation of  $(1.1)$ , we get

$$
\begin{cases}\n\rho(u_t + uu_x + vu_y) + p_x - q_y - (\mu + \zeta)u_{xx} + (\tau - \mu)u_{yy} - (\tau + \zeta)v_{xy} + \alpha(v_{xx} + v_{yy}) = 0, \\
\rho(v_t + uv_x + vv_y) + p_y + q_x - (\mu + \zeta)v_{yy} + (\tau - \mu)v_{xx} - (\tau + \zeta)u_{xy} - \alpha(u_{xx} + u_{yy}) = 0.\n\end{cases}
$$

Acknowledgments. This research was partially supported by Russian Science Foundation grant number 21-71-20034.

### **REFERENCES**

- <span id="page-4-2"></span>[1] A. Duyunova, V. Lychagin, and S. Tychkov, "On Dynamics and Thermodynamics of Moving Media," Lobachevskii J. Math. 45, no. 2, 663–676 (2024).
- <span id="page-4-0"></span>[2] A. Duyunova, V. Lychagin, and S. Tychkov, "Thermodynamics of One-Dimensional Flows," Lobachevskii J. Math. 44, no. 9, 3913–3917, (2023).
- <span id="page-4-1"></span>[3] V. Lychagin and M. Roop, "Shock waves in Euler flows of gases," Lobachevskii J. Math. 41 (12), 2473–2481 (2020).
- [4] V. Lychagin, "Thermodynamics as a Theory of Measurement," J. of Geometry and Physics 172, 104430 (2022).
- <span id="page-4-4"></span>[5] M. Berger, "Sur les groupes d'holonomie homogènes de variétés à connexion affine et des variétés riemanniennes," Bull. Soc. Math. Fr. 83, 279-330 (1955).
- <span id="page-4-3"></span>[6] L. D. Landau and E. M. Lifshitz, *Fluid Mechanics, 2nd ed.* (Pergamon Press, Oxford, England; New York, 1987).
- <span id="page-4-5"></span>[7] C. Procesi, *Lie Groups: an Approach Through Invariants and Representations* (Springer, New York, 2007).

*Email address*: anna.duyunova@yahoo.com

V. A. Trapeznikov Institute of Control Sciences of Russian Academy of Sciences 65 PROFSOYUZNAYA STREET, MOSCOW 117997, RUSSIA

*Email address*: valentin.lychagin@uit.no

V. A. Trapeznikov Institute of Control Sciences of Russian Academy of Sciences 65 PROFSOYUZNAYA STREET, MOSCOW 117997, RUSSIA

#### *Email address*: tychkovsn@ipu.ru

V. A. Trapeznikov Institute of Control Sciences of Russian Academy of Sciences 65 PROFSOYUZNAYA STREET, MOSCOW 117997, RUSSIA