

# BERGMAN PROJECTION INDUCED BY RADIAL WEIGHT ACTING ON GROWTH SPACES

ÁLVARO MIGUEL MORENO, JOSÉ ÁNGEL PELÁEZ, AND JARI TASKINEN

ABSTRACT. Let  $\omega$  be a radial weight on the unit disc of the complex plane  $\mathbb{D}$  and denote  $\omega_x = \int_0^1 s^x \omega(s) ds$ ,  $x \geq 0$ , for the moments of  $\omega$  and  $\hat{\omega}(r) = \int_r^1 \omega(s) ds$  for the tail integrals. A radial weight  $\omega$  belongs to the class  $\hat{\mathcal{D}}$  if satisfies the upper doubling condition

$$\sup_{0 < r < 1} \frac{\hat{\omega}(r)}{\hat{\omega}\left(\frac{1+r}{2}\right)} < \infty.$$

If  $\nu$  or  $\omega$  belongs to  $\hat{\mathcal{D}}$ , it is described the boundedness of the Bergman projection  $P_\omega$  induced by  $\omega$  on the growth space  $L_{\hat{\nu}}^\infty = \{f : \|f\|_{\infty, \nu} = \text{ess sup}_{z \in \mathbb{D}} |f(z)| \hat{\nu}(z) < \infty\}$  in terms of neat conditions on the moments and/or the tail integrals of  $\omega$  and  $\nu$ . Moreover, it is solved the analogous problem for  $P_\omega$  from  $L_{\hat{\nu}}^\infty$  to the Bloch type space  $B_{\hat{\nu}}^\infty = \{f \text{ analytic in } \mathbb{D} : \|f\|_{B_{\hat{\nu}}^\infty} = \sup_{z \in \mathbb{D}} (1 - |z|) \hat{\nu}(z) |f'(z)| < \infty\}$ . We also study similar questions for exponentially decreasing radial weights.

## 1. INTRODUCTION AND MAIN RESULTS

The question of when the Bergman projection  $P_\omega$  induced by a radial weight  $\omega$  on the unit disc  $\mathbb{D}$  of the complex plane, is a bounded operator in given function spaces is fundamental in the theory of spaces of analytic functions on  $\mathbb{D}$ . This is not only due to the mathematical difficulties the question raises, but also to its numerous applications in operator theory. Indeed, bounded analytic projections can be used to establish duality relations and to obtain useful equivalent norms in spaces of analytic functions [5, 13, 14, 20]. It has been recently described the radial weights  $\omega$  such that  $P_\omega$  is bounded from  $L^\infty$  to the classical Bloch space  $\mathcal{B}$  [14]. In this paper we are also interested in the natural limit case  $p = \infty$ , and our main results provide complete characterizations of the radial weights  $\omega$  (respectively,  $\nu$ ) such that  $P_\omega$  is bounded on the growth space induced by  $\nu$ , when  $\nu$  (resp.  $\omega$ ) satisfies an upper doubling condition. In addition, we will consider bounded projections onto weighted Bloch spaces in the setting of doubling weights, and we also obtain new results on the boundedness of  $P_\omega$ , when  $\omega$  and  $\nu$  belong to certain classes of exponentially decreasing weights. In order to present the precise statements of our results some definitions are needed.

Let  $\mathcal{H}(\mathbb{D})$  denote the space of analytic functions in  $\mathbb{D}$ . For a nonnegative function  $\omega \in L^1([0, 1))$ , the extension to  $\mathbb{D}$ , defined by  $\omega(z) = \omega(|z|)$  for all  $z \in \mathbb{D}$ , is called a radial weight. For  $0 < p < \infty$  and such an  $\omega$ , the weighted Lebesgue space  $L_\omega^p$  consists of complex-valued measurable functions  $f$  on  $\mathbb{D}$  such that

$$\|f\|_{L_\omega^p}^p = \int_{\mathbb{D}} |f(z)|^p \omega(z) dA(z) < \infty,$$

where  $dA(z) = \frac{dx dy}{\pi}$  is the normalized Lebesgue area measure on  $\mathbb{D}$ . The corresponding weighted Bergman space is  $A_\omega^p = L_\omega^p \cap \mathcal{H}(\mathbb{D})$ . Throughout this paper we assume  $\hat{\omega}(z) = \int_{|z|}^1 \omega(s) ds > 0$  for all  $z \in \mathbb{D}$ , for otherwise  $A_\omega^p = \mathcal{H}(\mathbb{D})$ . We also consider the Lebesgue space

---

*Key words and phrases.* Bergman projection; boundedness; weighted sup-norm; radial weight; doubling weight; exponential weight; weighted Hardy space; Szegő projection .

The first two authors are supported in part by Ministerio de Ciencia e Innovación, Spain, project PID2022-136619NB-I00 and La Junta de Andalucía, project FQM210. The third author was supported in part by the HORIZON-MSCA-2022-PF-01 project 101109510 LARGE BERGMAN as well as by the Magnus Ehrnrooth Foundation and the Väisälä Foundation of the Finnish Academy of Sciences and Letters.

$L^\infty$  of complex-valued measurable functions  $f$  on  $\mathbb{D}$  such that  $\|f\|_\infty = \text{ess sup}_{z \in \mathbb{D}} |f(z)| < \infty$  and the Hardy space is  $H^\infty = L^\infty \cap \mathcal{H}(\mathbb{D})$ .

For a radial weight  $\omega$ , the orthogonal Bergman projection  $P_\omega$  from  $L_\omega^2$  to  $A_\omega^2$  is

$$P_\omega(f)(z) = \int_{\mathbb{D}} f(\zeta) \overline{B_z^\omega(\zeta)} \omega(\zeta) dA(\zeta),$$

where  $B_z^\omega$  are the reproducing kernels of  $A_\omega^2$ . As usual,  $A_\alpha^p$  stands for the classical weighted Bergman space induced by the standard radial weight  $\omega(z) = (\alpha + 1)(1 - |z|^2)^\alpha$ ,  $B_z^\alpha$  are the kernels of  $A_\alpha^2$ , and  $P_\alpha$  denotes the corresponding Bergman projection.

One of the main obstacles throughout this work is the lack of explicit expressions for the Bergman reproducing kernel  $B_z^\omega$ . For a radial weight  $\omega$ , the kernel has the representation  $B_z^\omega(\zeta) = \sum e_n(\overline{z}) e_n(\zeta)$  for each orthonormal basis  $\{e_n\}$  of  $A_\omega^2$ , and therefore we are basically forced to work with the formula  $B_z^\omega(\zeta) = \sum_{n=0}^{\infty} \frac{(\overline{z}\zeta)^n}{2\omega_{2n+1}}$  induced by the normalized monomials.

Here  $\omega_{2n+1}$  are the odd moments of  $\omega$ , and in general from now on we write  $\omega_x = \int_0^1 r^x \omega(r) dr$  for all  $x \geq 0$ . Therefore the influence of the weight to the kernel is transmitted by its moments through this infinite sum, and nothing more than that can be said in general. This is in stark contrast with the neat expression  $(1 - \overline{z}\zeta)^{-(2+\alpha)}$  of the standard Bergman kernel  $B_z^\alpha$  which is easy to work with. It is well known that for  $1 < p < \infty$  and  $\alpha > -1$ , the Bergman projection  $P_\alpha$  acts as a bounded operator from  $L_\alpha^p$  to  $A_\alpha^p$  [20, Section 4]. However,  $P_\alpha$  is never bounded from  $L^\infty$  to  $H^\infty$ . In fact, this is a general phenomenon (known e.g. in the isomorphic theory of Banach spaces) rather than a particular case.

**Theorem A.** *There does not exist any bounded projection from  $L^\infty$  to  $H^\infty$ . In particular,  $P_\omega$  is not bounded from  $L^\infty$  to  $H^\infty$  for any radial weight.*

In view of the previous result it is natural to look for a substitute of  $H^\infty$  in the target space when the Bergman projection  $P_\omega$  acts on  $L^\infty$ . As for this question, it is known that the standard Bergman projection  $P_\alpha$  is bounded and onto from  $L^\infty$  to the Bloch space  $\mathcal{B}$ , which consists of  $f \in \mathcal{H}(\mathbb{D})$  such that  $\|f\|_{\mathcal{B}} = \sup_{z \in \mathbb{D}} |f'(z)|(1 - |z|^2) + |f(0)| < \infty$  [20, Theorem 5.2]. Furthermore, it has been recently proved that the Bergman projection  $P_\omega$  induced by a radial weight  $\omega$  acts as a bounded operator from the space  $L^\infty$  to  $\mathcal{B}$  if and only if  $\omega \in \widehat{\mathcal{D}}$ , and  $P_\omega : L^\infty \rightarrow \mathcal{B}$  is bounded and onto if and only if  $\omega \in \mathcal{D} = \widehat{\mathcal{D}} \cap \check{\mathcal{D}}$  [14, Theorems 1-2-3]. Let us recall the reader that a radial weight  $\omega$  belongs to  $\widehat{\mathcal{D}}$ , and it is called upper doubling, if there exists  $C = C(\omega) > 0$  such that

$$\widehat{\omega}(r) \leq C \widehat{\omega} \left( \frac{1+r}{2} \right), \quad r \rightarrow 1^-,$$

and  $\omega \in \check{\mathcal{D}}$  if there exists  $K = K(\omega) > 1$  and  $C = C(\omega) > 1$  such that

$$\widehat{\omega}(r) \geq C \widehat{\omega} \left( 1 - \frac{1-r}{K} \right), \quad r \rightarrow 1^-.$$

We say that  $\omega$  is a doubling weight if  $\omega \in \mathcal{D}$ . Standard weights belong to  $\widehat{\mathcal{D}}$  but exponentially decreasing weights do not belong to  $\widehat{\mathcal{D}}$ . Moreover, the containment in  $\widehat{\mathcal{D}}$  or  $\check{\mathcal{D}}$  does not require differentiability, continuity or strict positivity. In fact, weights in these classes may vanish on a relatively large part of each outer annulus  $\{z : r \leq |z| < 1\}$  of  $\mathbb{D}$ . For basic properties of the aforementioned classes, concrete nontrivial examples and more, see [11, 12, 14] and the relevant references therein.

A natural next step within this theory and the main objective of the paper is to study the properties of a radial weight  $\omega$  which are determinative so that  $P_\omega$  is bounded on the growth space  $L_v^\infty = \{f \text{ measurable} : \|f\|_{\infty, v} = \text{ess sup}_{z \in \mathbb{D}} |f(z)|v(z) < \infty\}$ , where  $v$  is a radial, continuous and decreasing weight with  $\lim_{r \rightarrow 1^-} v(r) = 0$ . We denote  $H_v^\infty = L_v^\infty \cap \mathcal{H}(\mathbb{D})$ . Sections 2–4 deal with cases of doubling weights, and in the previous definitions of spaces,  $v$  will be equal to  $\widehat{v}$  for a radial weight  $\nu$ . In Section 4.1 we will study the boundedness of  $P_\omega$

from  $L_{\hat{\nu}}^{\infty}$  to  $H_{\hat{\nu}}^{\infty}$  under some hypothesis on  $\nu$ . In particular, assuming  $\nu \in \hat{\mathcal{D}}$  we will prove the following characterization of the pairs of radial weights  $(\omega, \nu)$  such that this fact happens.

**Theorem 1.** *Let  $\omega$  be a radial weight and  $\nu \in \hat{\mathcal{D}}$ . Then, the following conditions are equivalent:*

- (i)  $P_{\omega} : L_{\hat{\nu}}^{\infty} \rightarrow H_{\hat{\nu}}^{\infty}$  is bounded;
- (ii)  $\nu \in \check{\mathcal{D}}$  and

$$\sup_{x \geq 0} \frac{\nu_x}{\omega_{2x}} \left( \frac{\omega}{\hat{\nu}} \right)_x < \infty; \quad (1.1)$$

- (iii)  $\omega \in \hat{\mathcal{D}}$ ,  $\nu \in \check{\mathcal{D}}$  and

$$\sup_{0 \leq r < 1} \frac{\hat{\nu}(r)}{\hat{\omega}(r)} \int_r^1 \frac{\omega(s)}{\hat{\nu}(s)} ds < \infty; \quad (1.2)$$

- (iv)  $\frac{\omega}{\hat{\nu}} \in \mathcal{D}$  and  $\nu \in \check{\mathcal{D}}$ .

It is worth mentioning that  $\int_r^1 \frac{\omega(s)}{\hat{\nu}(s)} ds \geq \frac{\hat{\omega}(r)}{\hat{\nu}(r)}$  holds for all radial weights and that (1.2) means nothing else but that the function  $\frac{1}{\hat{\nu}}$  satisfies a Bekollé-Bonami type condition  $B_{1,\omega}$ , see [16, Theorem 3] for related results. As for the proof of Theorem 1, the condition (1.1) is deduced from (i) using the the reformulation

$$\sup_{a \in \mathbb{D}} \hat{\nu}(a) \int_{\mathbb{D}} |B_a^{\omega}(\zeta)| \frac{\omega(\zeta)}{\hat{\nu}(\zeta)} dA(\zeta) < \infty, \quad (1.3)$$

together with a Hausdorff-Young inequality [4, Theorem 6.1]. Then it is proved that condition (1.1) implies that  $\omega \in \hat{\mathcal{D}}$  which joined to [13, Theorem 1] allows to express the integral in (1.3) in terms of  $\hat{\omega}$  and  $\hat{\nu}$ . Finally, some technical calculations imply that  $\nu \in \check{\mathcal{D}}$  (see Proposition 13 below) and we get (ii). The proofs of (ii) $\Leftrightarrow$ (iii) $\Leftrightarrow$ (iv) involve several descriptions of the classes of radial weight  $\hat{\mathcal{D}}$  and  $\check{\mathcal{D}}$ . In order to prove (iii) $\Rightarrow$ (i), [13, Theorem 1] is employed again.

In the case of standard weights  $\omega(r) = (1-r^2)^{\beta}$ ,  $\nu(r) = (1-r^2)^{\alpha}$ ,  $\alpha, \beta > -1$ , the projection  $P_{\omega}$  is bounded on  $L_{\hat{\nu}}^{\infty}$ , if and only if

$$\beta > \alpha. \quad (1.4)$$

This follows from the Forelli-Rudin estimates, see [20], Lemma 3.10 and the proof of Theorem 3.12. Since both  $\omega$  and  $\nu$  belong to  $\mathcal{D}$ , we conclude that (1.4) is an equivalent formulation of (1.1) and (1.2) in this simple case.

As a byproduct of Theorem 1 we deduce the next result which might be expected in view of Theorem A, because roughly speaking the spaces  $L_{\hat{\nu}}^{\infty}$  are "small" growth spaces for  $\nu \in \hat{\mathcal{D}} \setminus \check{\mathcal{D}}$ .

**Corollary 2.** *Let  $\nu \in \hat{\mathcal{D}}$ . Then, there is a radial weight  $\omega$  such that  $P_{\omega} : L_{\hat{\nu}}^{\infty} \rightarrow H_{\hat{\nu}}^{\infty}$  is bounded if and only if  $\nu \in \check{\mathcal{D}}$ . In particular, if  $\nu \in \hat{\mathcal{D}} \setminus \check{\mathcal{D}}$  and a  $\omega$  is a radial weight  $P_{\omega}$  is not bounded from  $L_{\hat{\nu}}^{\infty}$  to  $H_{\hat{\nu}}^{\infty}$ .*

Theorem 1 is also related with the boundedness of the Szegő projection. In Section 3 we recall some literature references on the proof of Theorem A and its relations to the boundedness of the Szegő projection, since these can be used to give an alternative proof for one of our results in Corollary 2.

In Section 4.2 we will study the boundedness of  $P_{\omega}$  from  $L_{\hat{\nu}}^{\infty}$  to  $H_{\hat{\nu}}^{\infty}$  assuming some hypothesis on  $\omega$ . The following main results will be proved using similar techniques to those employed in the proof of Theorem 1.

**Theorem 3.** *Let  $\nu$  be a radial weight and  $\omega \in \hat{\mathcal{D}}$ . Then, the following conditions are equivalent:*

- (i)  $P_{\omega} : L_{\hat{\nu}}^{\infty} \rightarrow H_{\hat{\nu}}^{\infty}$  is bounded;
- (ii)  $\nu \in \check{\mathcal{D}}$  and (1.1) holds;
- (iii)  $\nu \in \check{\mathcal{D}}$  and (1.2) holds;
- (iv)  $\frac{\omega}{\hat{\nu}} \in \mathcal{D}$  and  $\nu \in \mathcal{D}$ .

**Corollary 4.** *Let  $\omega \in \widehat{\mathcal{D}}$ . Then there is a radial weight  $\nu$  such that  $P_\omega : L_\nu^\infty \rightarrow H_\nu^\infty$  is bounded if and only if  $\omega \in \check{\mathcal{D}}$ . In particular, if  $\omega \in \widehat{\mathcal{D}} \setminus \check{\mathcal{D}}$  and a  $\nu$  is a radial weight  $P_\omega$  is not bounded from  $L_\nu^\infty$  to  $H_\nu^\infty$ .*

Bearing in mind Theorem A, [14, Theorem 1] and our previous results it is also interesting to study the boundedness from  $L_\nu^\infty$  to the Bloch type space

$$B_\nu^\infty = \left\{ f \in \mathcal{H}(\mathbb{D}) : \|f\|_{B_\nu^\infty} = \sup_{z \in \mathbb{D}} (1 - |z|)\nu(z)|f'(z)| < \infty \right\},$$

where  $\nu$  is a continuous and decreasing weight with  $\lim_{r \rightarrow 1^-} \nu(r) = 0$ . In Section 4.3 we will prove the following results for  $\nu = \widehat{\nu}$ .

**Theorem 5.** *Let  $\omega$  be a radial weight and  $\nu \in \widehat{\mathcal{D}}$ . Then, the following conditions are equivalent:*

- (i)  $P_\omega : L_\nu^\infty \rightarrow \mathcal{B}_\nu^\infty$  is bounded;
- (ii) (1.1) holds;
- (iii)  $\omega \in \widehat{\mathcal{D}}$  and (1.2) holds.

**Theorem 6.** *Let  $\nu$  be a radial weight and  $\omega \in \widehat{\mathcal{D}}$ . Then, the following conditions are equivalent:*

- (i)  $P_\omega : L_\nu^\infty \rightarrow \mathcal{B}_\nu^\infty$  is bounded;
- (ii) (1.1) holds;
- (iii) (1.2) holds.

We point out that on the contrary to the boundedness of  $P_\omega : L_\nu^\infty \rightarrow H_\nu^\infty$ , the boundedness of  $P_\omega : L_\nu^\infty \rightarrow \mathcal{B}_\nu^\infty$ ,  $\omega, \nu \in \widehat{\mathcal{D}}$ , does not require that  $\omega, \nu \in \check{\mathcal{D}}$ .

Section 5 is devoted to the study of the cases with exponentially decreasing weights, where we use techniques different from those for Theorems 1–6. We postpone there the detailed definitions and the formulation of the results.

Finally, concerning the notation in this paper, we denote by  $C$  (respectively  $C(\omega)$  etc.) generic positive constants independent of the variables involved (resp. depending only on  $\omega$ ), the values of which may vary from place to place. As usual,  $A \lesssim B$  ( $B \gtrsim A$ ) for nonnegative functions  $A, B$  means that  $A \leq CB$ , for some constant  $C$ . Furthermore, we write  $A \asymp B$  when  $A \lesssim B$  and  $A \gtrsim B$ .

## 2. PRELIMINARY RESULTS ON RADIAL WEIGHTS

Throughout the paper we will employ different descriptions of the weight classes  $\widehat{\mathcal{D}}$ ,  $\check{\mathcal{D}}$ . The next result gathers several characterizations of  $\widehat{\mathcal{D}}$  proved in [11, Lemma 2.1].

**Lemma B.** *Let  $\omega$  be a radial weight. Then, the following statements are equivalent:*

- (i)  $\omega \in \widehat{\mathcal{D}}$ ;
- (ii) There exist  $C = C(\omega) \geq 1$  and  $\alpha_0 = \alpha_0(\omega) > 0$  such that

$$\widehat{\omega}(s) \leq C \left( \frac{1-s}{1-t} \right)^\alpha \widehat{\omega}(t), \quad 0 \leq s \leq t < 1,$$

for all  $\alpha \geq \alpha_0$ ;

- (iii)

$$\omega_x = \int_0^1 s^x \omega(s) ds \asymp \widehat{\omega} \left( 1 - \frac{1}{x} \right), \quad x \in [1, \infty);$$

- (iv) There exists  $C(\omega) > 0$  such that  $\omega_x \leq C\omega_{2x}$ , for any  $x \geq 1$ .

In particular, we will repeatedly use Lemma B(iii) and the fact that  $\sup_{x \geq 1} \frac{\widehat{\omega}(1-\frac{1}{x})}{\omega_x} < \infty$  for any radial weight  $\omega$ .

We will also use the following descriptions of the class  $\check{\mathcal{D}}$ , see [10, Lemma 4] and [15, Lemma B].

**Lemma C.** *Let  $\omega$  be a radial weight. Then, the following statements are equivalent:*

- (i)  $\omega \in \check{\mathcal{D}}$ ;  
(ii) There exist  $C = C(\omega) > 0$  and  $\beta = \beta(\omega) > 0$  such that

$$\hat{\omega}(s) \leq C \left( \frac{1-s}{1-t} \right)^\beta \hat{\omega}(t), \quad 0 \leq t \leq s < 1;$$

- (iii) For each (or some)  $\gamma > 0$  there exists  $C = C(\gamma, \omega) > 0$  such that

$$\int_0^r \frac{ds}{\hat{\omega}(s)^\gamma (1-s)} \leq \frac{C}{\hat{\omega}(r)^\gamma}, \quad 0 \leq r < 1;$$

- (iv) There exist  $K(\omega) > 1$  and  $C = C(\omega) > 0$  such that

$$\int_r^{1-\frac{1-r}{K}} \omega(s) ds \geq C \int_r^1 \omega(s) ds, \quad 0 \leq r < 1.$$

We will also use the following results.

**Lemma 7.** Let  $\omega$  be a radial weight and let  $\varphi$  be a mapping  $\varphi : [0, 1) \rightarrow (0, \infty)$ .

- (i) If  $\omega \in \hat{\mathcal{D}}$  and  $\varphi$  is a non-decreasing function, such that  $\omega\varphi$  is a weight, then  $\omega\varphi \in \hat{\mathcal{D}}$ .  
(ii) If  $\omega \in \check{\mathcal{D}}$  and  $\varphi$  is a non-increasing function, then  $\omega\varphi \in \check{\mathcal{D}}$ .

*Proof.* (i). Observe that  $\omega \in \hat{\mathcal{D}}$  if and only if there exists  $C(\omega) > 0$  such that

$$\int_r^{\frac{1+r}{2}} \omega(s) ds \leq C \int_{\frac{1+r}{2}}^1 \omega(s) ds, \quad 0 \leq r < 1.$$

Applying this, and the fact that  $\varphi$  is non-decreasing, we obtain that

$$\begin{aligned} \frac{1}{\varphi\left(\frac{1+r}{2}\right)} \int_r^{\frac{1+r}{2}} \omega(s)\varphi(s) ds &\leq \int_r^{\frac{1+r}{2}} \omega(s) ds \\ &\leq C \int_{\frac{1+r}{2}}^1 \omega(s) ds \leq C \frac{1}{\varphi\left(\frac{1+r}{2}\right)} \int_{\frac{1+r}{2}}^1 \omega(s)\varphi(s) ds, \quad 0 \leq r < 1, \end{aligned}$$

then

$$\int_r^{\frac{1+r}{2}} \omega(s)\varphi(s) ds \leq C \int_{\frac{1+r}{2}}^1 \omega(s)\varphi(s) ds, \quad 0 \leq r < 1,$$

so  $\omega\varphi \in \hat{\mathcal{D}}$ .

(ii). By Lemma C(iv), there holds  $\omega \in \check{\mathcal{D}}$  if and only if there are  $K > 1$  and  $C(\omega) > 0$  such that

$$\int_r^{1-\frac{1-r}{K}} \omega(s) ds \geq C \int_{1-\frac{1-r}{K}}^1 \omega(s) ds, \quad 0 \leq r < 1.$$

So, since  $\varphi$  is a non-increasing function,

$$\begin{aligned} \int_r^{1-\frac{1-r}{K}} \omega(s)\varphi(s) ds &\geq \varphi\left(1 - \frac{1-r}{K}\right) \int_r^{1-\frac{1-r}{K}} \omega(s) ds \\ &\geq C\varphi\left(1 - \frac{1-r}{K}\right) \int_{1-\frac{1-r}{K}}^1 \omega(s) ds \geq \int_{1-\frac{1-r}{K}}^1 \omega(s)\varphi(s) ds, \quad 0 \leq r < 1. \end{aligned}$$

Hence  $\omega\varphi \in \check{\mathcal{D}}$ . This finishes the proof.  $\square$

**Lemma 8.** Let  $\omega \in \mathcal{D}$  and  $\nu \in \mathcal{D}$ . Then  $\omega\hat{\nu} \in \mathcal{D}$  and

$$\int_r^1 \omega(s)\hat{\nu}(s) ds = \hat{\omega}(r)\hat{\nu}(r), \quad 0 \leq r < 1. \quad (2.1)$$

*Proof.* By Lemma 7(ii), we have  $\omega\hat{\nu} \in \check{\mathcal{D}}$ . On the other hand, by Lemma C(iv) there are  $K > 1$  and  $C(\omega) > 0$  such that

$$\int_r^{1-\frac{1-r}{K}} \omega(s)ds \geq C \int_r^1 \omega(s)ds, \quad 0 \leq r < 1.$$

So, Lemma B(ii) together with the above inequality yield

$$\int_r^1 \omega(s)\hat{\nu}(s)ds \geq \hat{\nu} \left(1 - \frac{1-r}{K}\right) \int_{1-\frac{1-r}{K}}^1 \omega(s)ds \gtrsim \hat{\nu}(r)\hat{\omega}(r), \quad 0 \leq r < 1.$$

Therefore, using that  $\omega, \nu \in \hat{\mathcal{D}}$

$$\int_r^1 \omega(s)\hat{\nu}(s)ds \leq \hat{\omega}(r)\hat{\nu}(r) \lesssim \hat{\omega} \left(\frac{1+r}{2}\right) \hat{\nu} \left(\frac{1+r}{2}\right) \lesssim \int_{\frac{1+r}{2}}^1 \omega(s)\hat{\nu}(s)ds,$$

so  $\omega\hat{\nu} \in \hat{\mathcal{D}}$  and (2.1) holds. This finishes the proof.  $\square$

### 3. PRELIMINARY RESULTS ON THE BOUNDEDNESS OF $P_\omega : L_\nu^\infty \rightarrow H_\nu^\infty$

**3.1. On the boundedness of the Szegő projection in growth spaces.** Some known results on the boundedness of the Szegő projection  $R$  (also known as the Riesz projection) will be relevant for our studies on Bergman projections. Recall that if  $f$  is a complex valued harmonic function on  $\mathbb{D} \ni z = re^{i\theta}$  with a series representation  $f(re^{i\theta}) = \sum_{m=-\infty}^{\infty} f_m r^{|m|} e^{im\theta}$  converging at least uniformly on every compact subset of  $\mathbb{D}$ , then the Szegő projection  $R$  by definition maps  $f$  to the analytic function  $\sum_{m=0}^{\infty} f_m r^m e^{im\theta}$ , where the series also converges at least uniformly on compact subsets. See [20, Section 9.1].

As for the known proofs of Theorem A, if  $P$  were a bounded projection from  $L^\infty$  onto  $H^\infty$ , then one could consider the restriction of  $P$  as a bounded projection from  $L^\infty(\partial\mathbb{D})$  onto  $H^\infty$ . But it is well-known that such a projection does not exist. Indeed, by techniques explained in [18, Theorem 5.18, Example 5.19], see also [20, Theorem 9.7], one can show that the existence of such a bounded projection would imply the untrue fact that the Szegő projection  $R : L^\infty(\partial\mathbb{D}) \rightarrow H^\infty$  is bounded (e.g.[20], p. 258).

Another way to see Theorem A follows from the fact that  $H^\infty$  is known not to be isomorphic to the Banach space  $\ell^\infty$  of all bounded sequences (e.g. [8, Theorem 1.1]). Recall that a closed subspace of a Banach space is called *complemented*, if there exists a bounded projection onto it. Now, by [17], [6, p.111], the space  $L^\infty$  is isomorphic to the Banach space  $\ell^\infty$ , which is a *prime* Banach space ([6, Theorem 2.a.7]), i.e. all of its complemented subspaces are again isomorphic to  $\ell^\infty$ . We thus find that  $H^\infty$  is not complemented in  $L^\infty$ .

The next result is probably also known to experts, but we indicate the simple proof for the convenience of the reader.

**Proposition 9.** *Let  $\omega$  and  $\nu$  be radial weights such that  $P_\omega : L_\nu^\infty \rightarrow H_\nu^\infty$  is bounded. Then, the restriction of  $P_\omega$  onto the closed subspace  $h_\nu^\infty$  of  $L_\nu^\infty$ , consisting of harmonic functions, coincides with the Szegő projection and consequently,  $R : h_\nu^\infty \rightarrow H_\nu^\infty$  is bounded.*

*Proof.* Assume that  $P_\omega : L_\nu^\infty \rightarrow H_\nu^\infty$  is bounded and  $f(re^{i\theta}) = \sum_{m=-\infty}^{\infty} f_m r^{|m|} e^{im\theta} \in h_\nu^\infty \subset L_\nu^\infty$  and  $z = re^{i\theta} \in \mathbb{D}$ . Bearing in mind that  $B_z^\omega(\zeta) = \sum_{n=0}^{\infty} \frac{(\bar{z}\zeta)^n}{2\omega_{2n+1}}$  we obtain

$$\begin{aligned} P_\omega f(z) &= \int_{\mathbb{D}} \sum_{n=0}^{\infty} \frac{(z\bar{\zeta})^n}{2\omega_{2n+1}} f(\zeta)\omega(\zeta)dA(\zeta) \\ &= \frac{1}{\pi} \int_0^1 \int_0^{2\pi} \sum_{n=0}^{\infty} (2\omega_{2n+1})^{-1} r^n e^{in\theta} \sum_{m=-\infty}^{\infty} f_m s^{n+|m|+1} e^{i\varphi(m-n)} \omega(s)d\varphi ds \\ &= \sum_{n=0}^{\infty} (\omega_{2n+1})^{-1} f_n r^n e^{in\theta} \int_0^1 s^{2n+1} \omega(s)ds = \sum_{n=0}^{\infty} f_n r^n e^{in\theta} = Rf(z). \end{aligned} \quad (3.1)$$

This finishes the proof.  $\square$

Now, let us observe that the "only if part" of Corollary 2 can be deduced from Proposition 9 and some results in [8] (see also [7]). In fact, some calculations show that  $\nu \in \widehat{\mathcal{D}}$  if and only if

$$\sup_{n \in \mathbb{N}} \frac{\widehat{\nu}(1 - 2^{-n})}{\widehat{\nu}(1 - 2^{-n-1})} < \infty$$

and  $\nu \in \check{\mathcal{D}}$  if and only if

$$\inf_{k \in \mathbb{N}} \limsup_{n \in \mathbb{N}} \frac{\widehat{\nu}(1 - 2^{-n-k})}{\widehat{\nu}(1 - 2^{-n})} < \infty.$$

Moreover, Proposition 6.4 and the first lines of p. 26 of [8] show that if  $\nu \in \widehat{\mathcal{D}}$ , then the Szegő projection  $R$  is bounded from  $h_{\widehat{\nu}}^{\infty}$  to  $H_{\widehat{\nu}}^{\infty}$  if and only if  $\nu \in \check{\mathcal{D}}$ , which together with Proposition 9 proves that if  $\omega$  is a radial weight and  $\nu \in \widehat{\mathcal{D}} \setminus \check{\mathcal{D}}$  then  $P_{\omega}$  is not bounded from  $L_{\widehat{\nu}}^{\infty}$  to  $H_{\widehat{\nu}}^{\infty}$ . Later on we will prove Corollary 2 using Theorem 1.

**3.2. Preliminary results on the boundedness of  $P_{\omega} : L_{\widehat{\nu}}^{\infty} \rightarrow H_{\widehat{\nu}}^{\infty}$ .** In this section we will prove some preliminary results of their own interest, which will be used in the proof of Theorem 1.

**Proposition 10.** *Let  $\omega$  and  $\nu$  be radial weights. Then,  $P_{\omega} : L_{\widehat{\nu}}^{\infty} \rightarrow H_{\widehat{\nu}}^{\infty}$  is bounded if and only if*

$$\sup_{a \in \mathbb{D}} \widehat{\nu}(a) \int_{\mathbb{D}} |B_a^{\omega}(\zeta)| \frac{\omega(\zeta)}{\widehat{\nu}(\zeta)} dA(\zeta) < \infty. \quad (3.2)$$

*Proof.* It is clear that  $P_{\omega} : L_{\widehat{\nu}}^{\infty} \rightarrow H_{\widehat{\nu}}^{\infty}$  is bounded if (3.2) holds. Reciprocally, assume that  $P_{\omega} : L_{\widehat{\nu}}^{\infty} \rightarrow H_{\widehat{\nu}}^{\infty}$  is bounded and for each  $a \in \mathbb{D}$ , consider the function

$$f_a(\zeta) = \begin{cases} \frac{B_a^{\omega}(\zeta)}{\widehat{\nu}(\zeta)|B_a^{\omega}(\zeta)|} & \text{if } B_a^{\omega}(\zeta) \neq 0 \\ 0 & \text{if } B_a^{\omega}(\zeta) = 0 \end{cases}.$$

Then, for each  $a \in \mathbb{D}$ ,  $\|f_a\|_{L_{\widehat{\nu}}^{\infty}} = 1$  and

$$\begin{aligned} \widehat{\nu}(|a|) \int_{\mathbb{D}} |B_a^{\omega}(\zeta)| \frac{\omega(\zeta)}{\widehat{\nu}(\zeta)} dA(\zeta) &= \widehat{\nu}(|a|) \int_{\mathbb{D}} f_a(\zeta) \overline{B_a^{\omega}(\zeta)} \omega(\zeta) dA(\zeta) \\ &\leq \sup_{z \in \mathbb{D}} \widehat{\nu}(|z|) \left| \int_{\mathbb{D}} f_a(\zeta) \overline{B_z^{\omega}(\zeta)} \omega(\zeta) dA(\zeta) \right| \leq \|P_{\omega}\| \|f_a\|_{L_{\widehat{\nu}}^{\infty}} = \|P_{\omega}\|. \end{aligned}$$

Therefore, (3.2) holds. This finishes the proof.  $\square$

It is worth noticing that an argument similar to that of Proposition 10 allows to give an alternative proof of the fact that  $P_{\omega}$  is not bounded from  $L^{\infty}$  to  $H^{\infty}$  for any radial weight  $\omega$ . In fact, arguing as in the the proof of Proposition 10 it follows that  $P_{\omega}$  is bounded from  $L^{\infty}$  to  $H^{\infty}$  if and only if

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |B_a^{\omega}(\zeta)| \omega(\zeta) dA(\zeta) < \infty.$$

Moreover, using Hardy's inequality [4, Theorem 5.1], for each  $a \in \mathbb{D} \setminus \{0\}$

$$\begin{aligned} \int_{\mathbb{D}} |B_a^{\omega}(\zeta)| \omega(\zeta) dA(\zeta) &= 2 \int_0^1 M_1(B_a^{\omega}, s) s \omega(s) ds \\ &\geq 2\pi \int_0^1 \left( \sum_{n=0}^{\infty} \frac{s^n |a|^n}{2(n+1)\omega_{2n+1}} \right) s \omega(s) ds \\ &= \pi \sum_{n=0}^{\infty} \frac{|a|^n}{(n+1)\omega_{2n+1}} \int_0^1 s^{n+1} \omega(s) ds \\ &= \pi \sum_{n=0}^{\infty} \frac{|a|^n \omega_{n+1}}{(n+1)\omega_{2n+1}} \geq \pi \sum_{n=0}^{\infty} \frac{|a|^n}{n+1} = \frac{\pi}{|a|} \log \left( \frac{1}{1-|a|} \right), \end{aligned}$$

Therefore,  $\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |B_a^\omega(\zeta)| \omega(\zeta) dA(\zeta) = \infty$  and consequently  $P_\omega$  is not bounded from  $L^\infty$  to  $H^\infty$ .

We will use Proposition 10 to obtain a necessary condition for the boundedness of  $P_\omega : L_{\hat{\nu}}^\infty \rightarrow H_{\hat{\nu}}^\infty$  in terms of moments and tails of  $\omega$  and  $\nu$ .

**Proposition 11.** *Let  $\omega$  and  $\nu$  radial weights such that  $P_\omega : L_{\hat{\nu}}^\infty \rightarrow H_{\hat{\nu}}^\infty$  is bounded. Then, there exists  $C(\omega, \nu) > 0$  such that*

$$\left(\frac{\omega}{\hat{\nu}}\right)_x \leq C \frac{\omega_{2x}}{\hat{\nu} \left(1 - \frac{1}{x}\right)}, \quad x \geq 1. \quad (3.3)$$

*Proof.* By Proposition 10 there exists  $C(\omega, \nu) > 0$ , such that

$$\sup_{z \in \mathbb{D}} \hat{\nu}(z) \int_{\mathbb{D}} |B_z^\omega(\zeta)| \frac{\omega(\zeta)}{\hat{\nu}(\zeta)} dA(\zeta) \leq C,$$

and by applying [4, Theorem 6.1],

$$\begin{aligned} C &\geq \sup_{z \in \mathbb{D}} \hat{\nu}(z) \int_{\mathbb{D}} |B_z^\omega(\zeta)| \frac{\omega(\zeta)}{\hat{\nu}(\zeta)} dA(\zeta) = \sup_{z \in \mathbb{D}} \hat{\nu}(z) \int_0^1 \frac{\omega(r)}{\hat{\nu}(r)} M_1(B_z^\omega, r) dr \\ &\gtrsim \sup_{z \in \mathbb{D}, n \in \mathbb{N}} \hat{\nu}(z) \int_0^1 \frac{\omega(r)}{\hat{\nu}(r)} \frac{|z|^n r^n}{\omega_{2n+1}} dr \geq \hat{\nu} \left(1 - \frac{1}{n}\right) \frac{\left(\frac{\omega}{\hat{\nu}}\right)_n}{\omega_{2n+1}} \left(1 - \frac{1}{n}\right)^n \\ &\gtrsim \hat{\nu} \left(1 - \frac{1}{n}\right) \frac{\left(\frac{\omega}{\hat{\nu}}\right)_n}{\omega_{2n}}, \quad n \in \mathbb{N}, \quad n \geq 2. \end{aligned} \quad (3.4)$$

For  $x \geq 1$ , take  $N \in \mathbb{N}$  such that  $N \leq x < N + 1$ , then the previous inequality yields

$$\left(\frac{\omega}{\hat{\nu}}\right)_x \leq \left(\frac{\omega}{\hat{\nu}}\right)_N \leq C \frac{\omega_{2N}}{\hat{\nu} \left(1 - \frac{1}{N}\right)} \leq C \frac{\omega_{2(N+1)}}{\hat{\nu} \left(1 - \frac{1}{N}\right)} \leq C \frac{\omega_{2x}}{\hat{\nu} \left(1 - \frac{1}{x}\right)}.$$

This finishes the proof.  $\square$

#### 4. BOUNDEDNESS OF BERGMAN PROJECTION ON GROWTH SPACE INDUCED BY RADIAL DOUBLING WEIGHT.

**4.1. Boundedness of  $P_\omega : L_{\hat{\nu}}^\infty \rightarrow H_{\hat{\nu}}^\infty$  assuming conditions on  $\nu$ .** Theorem 1 and Corollary 2 will be proved in this section. Prior to presenting their proofs, two preliminary results are needed.

**Proposition 12.** *Let  $\omega$  be a radial weight and  $\nu \in \hat{\mathcal{D}}$  such that (1.1) holds. Then,  $\omega \in \hat{\mathcal{D}}$  and  $\omega \hat{\nu} \in \hat{\mathcal{D}}$ .*

*Proof.* Let  $x \geq 1$ . By applying Lemma B (iv) and (1.1)

$$\begin{aligned} \omega_x &\lesssim \frac{1}{\nu_x} \int_0^1 \omega(s) \hat{\nu}(s) s^{2x} ds = \frac{1}{\nu_x} \int_0^1 \omega(s) s^{2x} \int_s^1 \nu(t) dt ds \\ &\leq \frac{1}{\nu_x} \int_0^1 \omega(s) s^{\frac{3x}{2}} \int_s^1 t^{\frac{x}{2}} \nu(t) dt ds \leq \omega_{\frac{3x}{2}} \frac{\nu_{\frac{x}{2}}}{\nu_x} \lesssim \omega_{\frac{3x}{2}}, \quad x \geq 1. \end{aligned} \quad (4.1)$$

Hence,  $\omega_x \lesssim \omega_{\frac{3x}{2}}$ . So, iteration of this inequality yields  $\omega_x \lesssim \omega_{\frac{3x}{2}} \lesssim \omega_{\frac{9x}{4}} \leq \omega_{2x}$ . Therefore, by Lemma B (iv),  $\omega \in \hat{\mathcal{D}}$ .

Now, a similar argument, the fact that  $\omega, \nu \in \hat{\mathcal{D}}$  and (1.1) imply

$$(\omega \hat{\nu})_x \leq \omega_{\frac{x}{2}} \nu_{\frac{x}{2}} \lesssim \omega_x \nu_x \lesssim (\omega \hat{\nu})_{2x}, \quad x \geq 1,$$

so  $\omega \hat{\nu} \in \hat{\mathcal{D}}$  by Lemma B (iv).  $\square$

**Proposition 13.** *Let  $\nu$  be a radial weight and  $\omega \in \hat{\mathcal{D}}$  such that  $P_\omega : L_{\hat{\nu}}^\infty \rightarrow H_{\hat{\nu}}^\infty$  is bounded. Then,  $\nu \in \check{\mathcal{D}}$ .*



*Proof.* By Proposition 10, there exists a constant  $C > 0$  such that

$$\int_{\mathbb{D}} |B_z^\omega(\zeta)| \frac{\omega(\zeta)}{\widehat{\nu}(\zeta)} dA(\zeta) \leq \frac{C}{\widehat{\nu}(z)}, \quad z \in \mathbb{D}.$$

Now, by [13, Theorem 1] and Fubini's theorem, for  $\frac{1}{2} \leq |z| < 1$

$$\int_{\mathbb{D}} |B_z^\omega(\zeta)| \frac{\omega(\zeta)}{\widehat{\nu}(\zeta)} dA(\zeta) = \int_0^1 \frac{\omega(s)}{\widehat{\nu}(s)} \int_0^{|z|^s} \frac{dt}{\widehat{\omega}(t)(1-t)} ds = \int_0^{|z|} \frac{1}{\widehat{\omega}(t)(1-t)} \int_{\frac{t}{|z|}}^1 \frac{\omega(s)}{\widehat{\nu}(s)} ds dt,$$

Applying Lemma B (ii), we obtain that

$$\begin{aligned} \int_{\mathbb{D}} |B_z^\omega(\zeta)| \frac{\omega(\zeta)}{\widehat{\nu}(\zeta)} dA(\zeta) &\gtrsim \int_0^{|z|} \frac{\widehat{\omega}(\frac{t}{|z|})}{\widehat{\omega}(t)} \cdot \frac{dt}{\widehat{\nu}(\frac{t}{|z|})(1-t)} \geq \int_0^{2|z|-1} \frac{\widehat{\omega}(\frac{t}{|z|})}{\widehat{\omega}(t)} \cdot \frac{dt}{\widehat{\nu}(t)(1-t)} \\ &\gtrsim \int_0^{2|z|-1} \left( \frac{1 - \frac{t}{|z|}}{1-t} \right)^\beta \cdot \frac{dt}{\widehat{\nu}(t)(1-t)} \\ &\geq \int_0^{2|z|-1} \left( \frac{1 - \frac{2t}{1+t}}{1-t} \right)^\beta \cdot \frac{dt}{\widehat{\nu}(t)(1-t)} \\ &= \int_0^{2|z|-1} \frac{dt}{\widehat{\nu}(t)(1-t)}, \quad |z| \geq \frac{1}{2}. \end{aligned}$$

Then, we take  $|z| = \frac{1+r}{2}$  and observe that there is  $C = C(\omega, \nu) > 0$  such that

$$\int_0^r \frac{dt}{\widehat{\nu}(t)(1-t)} \leq \frac{C}{\widehat{\nu}(\frac{1+r}{2})}, \quad 0 \leq r < 1.$$

Therefore if  $0 \leq r \leq s < 1$ , then

$$\frac{1}{\widehat{\nu}(r)} \log \frac{1-r}{1-s} = \frac{1}{\widehat{\nu}(r)} \int_r^s \frac{dt}{1-t} \leq \int_r^s \frac{dt}{\widehat{\nu}(t)(1-t)} \leq \frac{C}{\widehat{\nu}(\frac{1+s}{2})}.$$

Finally, take  $K > 2$  such that  $\frac{\log \frac{K}{2}}{C} > 1$  and choose  $s = 1 - \frac{2}{K}(1-r) > r$ . Then,  $\frac{1+s}{2} = 1 - \frac{1-r}{K}$ , and

$$\widehat{\nu}(r) \geq \frac{\log \frac{K}{2}}{C} \widehat{\nu} \left( 1 - \frac{1-r}{K} \right), \quad 0 \leq r < 1.$$

This shows that  $\nu \in \check{\mathcal{D}}$  and finishes the proof. □

*Proof of Theorem 1.* We will prove (i) $\Rightarrow$ (ii) $\Leftrightarrow$ (iii) $\Leftrightarrow$ (iv) and finally (iii) $\Rightarrow$ (i).

Assume that  $P_\omega : L_\nu^\infty \rightarrow H_\nu^\infty$  is bounded. Then, (1.1) holds by Proposition 11 and Lemma B (iii). Next, Proposition 12 implies that  $\omega \in \widehat{\mathcal{D}}$  and therefore  $\nu \in \check{\mathcal{D}}$  by Proposition 13, so we get (ii). Now, if (ii) holds, then Proposition 12 implies that  $\omega \in \widehat{\mathcal{D}}$ . So, (1.2) follows from (1.1) and Lemma B (iii). That is, (iii) is satisfied. Reciprocally, if (iii) holds, then  $\frac{\omega}{\nu} \in \widehat{\mathcal{D}}$  by Lemma 7, and (1.1) follows from (1.2) and Lemma B (iii). Therefore we get (ii). Now let us prove (iii) $\Rightarrow$ (iv). Firstly,  $\frac{\omega}{\nu} \in \widehat{\mathcal{D}}$  by Lemma 7. In order to prove that  $\frac{\omega}{\nu} \in \check{\mathcal{D}}$  and to make the notation easier, we denote  $\mu = \frac{\omega}{\nu}$ . Then, (1.2) is equivalent to the existence of  $C > 1$  such that

$$\widehat{\mu}(r) \leq C \frac{\widehat{\omega}(r)}{\widehat{\nu}(r)}, \quad 0 \leq r < 1. \quad (4.2)$$

On the other hand, an integration by parts yields

$$\widehat{\omega}(r) = \widehat{\nu}(r) \widehat{\mu}(r) - \int_r^1 \nu(s) \widehat{\mu}(s) ds, \quad 0 \leq r < 1,$$

so (4.2) is equivalent to the existence of  $C \in (0, 1)$  such that

$$\frac{\int_r^1 \nu(s) \hat{\mu}(s) ds}{\hat{\nu}(r)} \leq C \hat{\mu}(r), \quad 0 \leq r < 1. \quad (4.3)$$

By Lemma C (ii) there exists  $C_2 > 0$  and  $\alpha > 0$  such that for every  $K > 1$

$$\frac{\hat{\nu}\left(1 - \frac{1-r}{K}\right)}{\hat{\nu}(r)} \leq \frac{C_2}{K^\alpha}, \quad 0 \leq r < 1.$$

Therefore, taking  $K$  such that  $\frac{C_2}{K^\alpha} < 1 - C$ , we obtain that

$$\begin{aligned} C \hat{\mu}(r) &\geq \frac{\int_r^1 \nu(s) \hat{\mu}(s) ds}{\hat{\nu}(r)} \geq \frac{\int_r^{1-\frac{1-r}{K}} \nu(s) \hat{\mu}(s) ds}{\hat{\nu}(r)} \geq \hat{\mu}\left(1 - \frac{1-r}{K}\right) \frac{\hat{\nu}(r) - \hat{\nu}\left(1 - \frac{1-r}{K}\right)}{\hat{\nu}(r)} \\ &= \hat{\mu}\left(1 - \frac{1-r}{K}\right) \left(1 - \frac{\hat{\nu}\left(1 - \frac{1-r}{K}\right)}{\hat{\nu}(r)}\right) \geq \hat{\mu}\left(1 - \frac{1-r}{K}\right) \left(1 - \frac{C_2}{K^\alpha}\right), \quad 0 \leq r < 1. \end{aligned}$$

Consequently,  $\mu \in \check{\mathcal{D}}$  and (iv) holds. Conversely, if (iv) holds, we get  $\omega \in \hat{\mathcal{D}}$  by Lemma 8. Moreover, since  $\frac{\omega}{\hat{\nu}} \in \check{\mathcal{D}}$ , Lemma C(iv) implies that there are  $K > 1$  and  $C > 0$  such that

$$\int_r^1 \frac{\omega(s)}{\hat{\nu}(s)} ds \leq C \int_r^{1-\frac{1-r}{K}} \frac{\omega(s)}{\hat{\nu}(s)} ds, \quad 0 \leq r < 1.$$

This together with Lemma B(ii) give

$$\begin{aligned} \hat{\omega}(r) &= \int_r^1 \omega(s) ds = \int_r^1 \frac{\omega(s)}{\hat{\nu}(s)} \hat{\nu}(s) ds \geq \int_r^{1-\frac{1-r}{K}} \frac{\omega(s)}{\hat{\nu}(s)} \hat{\nu}(s) ds \\ &\geq \hat{\nu}\left(1 - \frac{1-r}{K}\right) \int_r^{1-\frac{1-r}{K}} \frac{\omega(s)}{\hat{\nu}(s)} ds \gtrsim \hat{\nu}(r) \int_r^1 \frac{\omega(s)}{\hat{\nu}(s)} ds, \quad 0 \leq r < 1, \end{aligned}$$

which yields (iii).

Finally assume that (iii) holds and let us prove that  $P_\omega : L_\nu^\infty \rightarrow H_\nu^\infty$  is bounded. By Proposition 10, it is sufficient to prove that

$$\sup_{a \in \mathbb{D}, |a| \geq \frac{1}{2}} \hat{\nu}(a) \int_{\mathbb{D}} |B_a^\omega(z)| \frac{\omega(z)}{\hat{\nu}(z)} dA(z) < \infty.$$

By [13, Theorem 1] and Fubini's Theorem, this is equivalent with proving that there is  $C > 0$  such that

$$\hat{\nu}(a) \int_0^{|a|} \frac{1}{\hat{\omega}(t)(1-t)} \int_{\frac{t}{|a|}}^1 \frac{\omega(r)}{\hat{\nu}(r)} dr dt \leq C, \quad |a| \geq \frac{1}{2}.$$

By applying (1.2) and Lemma C (iii)

$$\begin{aligned} \hat{\nu}(a) \int_0^{|a|} \frac{1}{\hat{\omega}(t)(1-t)} \int_{\frac{t}{|a|}}^1 \frac{\omega(r)}{\hat{\nu}(r)} dr dt &\leq \hat{\nu}(a) \int_0^{|a|} \frac{1}{\hat{\omega}(t)(1-t)} \int_t^1 \frac{\omega(r)}{\hat{\nu}(r)} dr dt \\ &\lesssim \hat{\nu}(a) \int_0^{|a|} \frac{dt}{\hat{\nu}(t)(1-t)} dt \lesssim 1, \end{aligned}$$

so  $P_\omega : L_\nu^\infty \rightarrow B_\nu^\infty$  is bounded. This finishes the proof.  $\square$

*Proof of Corollary 2.* If  $P_\omega : L_\nu^\infty \rightarrow H_\nu^\infty$  is bounded then  $\nu \in \check{\mathcal{D}}$  by Theorem 1. Reciprocally, if  $\nu \in \mathcal{D}$  by Lemma B(ii), there exist  $C, \alpha > 0$  such that

$$\frac{1}{\hat{\nu}(s)} \leq C \left(\frac{1-r}{1-s}\right)^\alpha \frac{1}{\hat{\nu}(r)}, \quad 0 \leq r \leq s < 1.$$

Therefore, if  $\omega(z) = (1 - |z|)^\gamma$  with  $\gamma > \alpha - 1$ , then

$$\int_r^1 \frac{\omega(s)}{\widehat{\nu}(s)} ds \lesssim \frac{(1-r)^\alpha}{\widehat{\nu}(r)} \int_r^1 (1-s)^{\gamma-\alpha} ds \asymp \frac{(1-r)^{\gamma+1}}{\widehat{\nu}(r)} \asymp \frac{\widehat{\omega}(r)}{\widehat{\nu}(r)},$$

that is (1.2) holds. Then,  $P_\omega$  is bounded from  $L_\nu^\infty$  to  $H_\nu^\infty$  by Theorem 1. This finishes the proof.  $\square$

**4.2. Boundedness of  $P_\omega : L_\nu^\infty \rightarrow H_\nu^\infty$  assuming conditions on  $\omega$ .** This section is dedicated to the proof of Theorem 3 and Corollary 4 among other results.

*Proof of Theorem 3.* We will prove (i) $\Leftrightarrow$ (iii) $\Leftrightarrow$ (ii) and (iii) $\Leftrightarrow$ (iv).

If (i) holds, then  $\nu \in \widetilde{\mathcal{D}}$  by Proposition 13 and (3.3) holds by Proposition 11. So, using that  $\omega \in \widehat{\mathcal{D}}$  and Lemma B, we get (1.2). Hence, (iii) holds. Reciprocally, if (iii) holds, using (1.2) and the fact that  $\omega \in \widehat{\mathcal{D}}$

$$\widehat{\nu}(r) \lesssim \frac{\widehat{\omega}(r)}{\int_r^1 \frac{\omega(s)}{\widehat{\nu}(s)} ds} \lesssim \frac{\widehat{\omega}\left(\frac{1+r}{2}\right)}{\int_{\frac{1+r}{2}}^1 \frac{\omega(s)}{\widehat{\nu}(s)} ds} \leq \widehat{\nu}\left(\frac{1+r}{2}\right), \quad 0 < r < 1.$$

That is,  $\nu \in \widehat{\mathcal{D}}$ . So, (i) follows from Theorem 1. Next, assuming again that (iii) holds we have  $\nu \in \widehat{\mathcal{D}}$ . Moreover,  $\frac{\omega}{\nu} \in \widehat{\mathcal{D}}$  by Lemma 7(i). Therefore (1.1) follows from (1.2) and Lemma B. So (ii) is satisfied. Reciprocally, if (ii) holds, using that  $\omega \in \widehat{\mathcal{D}}$ , Lemma B and (1.1), we get (1.2) and thus also (iii). Finally, if (iii) holds, arguing as above yields  $\nu \in \widehat{\mathcal{D}}$ , and so  $\frac{\omega}{\nu} \in \mathcal{D}$  by Theorem 1. Hence, (iv) holds. On the other hand, (iv) $\Rightarrow$ (iii) follows from Theorem 1. This finishes the proof.  $\square$

*Proof of Corollary 4.* Let  $\omega \in \widehat{\mathcal{D}}$ . If  $P_\omega : L_\nu^\infty \rightarrow H_\nu^\infty$  is bounded, then Theorem 3 yields  $\frac{\omega}{\nu} \in \mathcal{D}$  and  $\nu \in \mathcal{D}$ , hence, by Lemma 8,  $\omega \in \mathcal{D}$ . On the other hand, if  $\omega \in \mathcal{D}$ , then, by Lemma B (ii), there exists  $0 < \beta$  such that

$$\left(\frac{1-r}{1-s}\right)^\beta \widehat{\omega}(s) \lesssim \widehat{\omega}(r). \quad 0 \leq r \leq s < 1.$$

So, in particular  $\widehat{\omega}(s) \lesssim (1-s)^\beta$ ,  $0 \leq s < 1$ . Now, take  $0 < \gamma < \beta$  and consider the standard weight  $\nu(r) = (1-r)^{\gamma-1}$ . We have  $\widehat{\nu}(r) \asymp (1-r)^\gamma$  and

$$\begin{aligned} \int_r^1 \frac{\omega(s)}{(1-s)^\gamma} ds &\lesssim \frac{\widehat{\omega}(r)}{(1-r)^\gamma} + \int_r^1 \frac{\gamma \widehat{\omega}(s)}{(1-s)^{\gamma+1}} ds \\ &\lesssim \frac{\widehat{\omega}(r)}{(1-r)^\gamma} + \frac{\widehat{\omega}(r)}{(1-r)^\beta} \int_0^1 (1-s)^{\beta-\gamma-1} ds \lesssim \frac{\widehat{\omega}(r)}{(1-r)^\gamma}, \quad 0 \leq r < 1. \end{aligned}$$

Consequently, by Theorem 3,  $P_\omega$  is bounded from  $L_\nu^\infty$  to  $H_\nu^\infty$ .  $\square$

We finish this section proving that both weights,  $\omega$  and  $\nu$ , have to be doubling if the operator  $P_\omega : L_\nu^\infty \rightarrow H_\nu^\infty$  is bounded and one of them is assumed to be upper doubling.

**Corollary 14.** *Let  $\omega$  and  $\nu$  be radial weights such that  $P_\omega : L_\nu^\infty \rightarrow H_\nu^\infty$  is bounded and  $\omega \in \widehat{\mathcal{D}}$  or  $\nu \in \widehat{\mathcal{D}}$ . Then, both weights  $\omega, \nu \in \mathcal{D}$ .*

*Proof.* Assume that  $P_\omega : L_\nu^\infty \rightarrow H_\nu^\infty$  is bounded and  $\omega \in \widehat{\mathcal{D}}$ , then  $\omega \in \mathcal{D}$  by Corollary 4 and  $\nu \in \mathcal{D}$  by Theorem 3. On the other hand, if  $P_\omega : L_\nu^\infty \rightarrow H_\nu^\infty$  is bounded and  $\nu \in \widehat{\mathcal{D}}$ , then  $\nu \in \mathcal{D}$  and  $\frac{\omega}{\nu} \in \mathcal{D}$  by Theorem 1, so by Lemma 8  $\omega \in \mathcal{D}$ . This finishes the proof.  $\square$

**4.3. Boundedness of  $P_\omega : L_{\hat{\nu}}^\infty \rightarrow \mathcal{B}_{\hat{\nu}}^\infty$ .** In this section, we will prove Theorem 5 and Theorem 6 among other results.

Firstly, by mimicking the proof of Proposition 10 we get the analogous result for the boundedness of  $P_\omega : L_{\hat{\nu}}^\infty \rightarrow \mathcal{B}_{\hat{\nu}}^\infty$ .

**Proposition 15.** *Let  $\omega$  and  $\nu$  be radial weights. Then,  $P_\omega : L_{\hat{\nu}}^\infty \rightarrow \mathcal{B}_{\hat{\nu}}^\infty$  is bounded if and only if*

$$\sup_{z \in \mathbb{D}} (1 - |z|) \hat{\nu}(z) \int_{\mathbb{D}} |(B_\zeta^\omega)'(z)| \frac{\omega(\zeta)}{\hat{\nu}(\zeta)} dA(\zeta) < \infty.$$

As a byproduct of Proposition 15 we get the following.

**Proposition 16.** *Let  $\omega$  and  $\nu$  radial weights. If  $P_\omega : L_{\hat{\nu}}^\infty \rightarrow \mathcal{B}_{\hat{\nu}}^\infty$  is bounded, then (3.3) holds.*

*Proof.* The proof follows the ideas from the proof of Proposition 11. We include the details for the sake of completeness. If  $P_\omega : L_{\hat{\nu}}^\infty \rightarrow \mathcal{B}_{\hat{\nu}}^\infty$  is bounded, then Proposition 15 and [4, Theorem 6.1] yield

$$\begin{aligned} \infty &> \sup_{z \in \mathbb{D}} (1 - |z|) \hat{\nu}(z) \int_{\mathbb{D}} |(B_\zeta^\omega)'(z)| \frac{\omega(\zeta)}{\hat{\nu}(\zeta)} dA(\zeta) \\ &\gtrsim \sup_{z \in \mathbb{D}, n \in \mathbb{N}} (1 - |z|) \hat{\nu}(z) \frac{n \int_0^1 |z|^{n-1} r^{n+1} \frac{\omega(r)}{\hat{\nu}(r)} dr}{\omega_{2n+1}} \\ &\gtrsim \sup_{n \in \mathbb{N}} \left(1 - \frac{1}{n}\right)^{n-1} \hat{\nu} \left(1 - \frac{1}{n}\right) \frac{\left(\frac{\omega}{\hat{\nu}}\right)_{n+1}}{\omega_{2n+1}} \\ &\gtrsim \hat{\nu} \left(1 - \frac{1}{n}\right) \frac{\left(\frac{\omega}{\hat{\nu}}\right)_{n+1}}{\omega_{2n+1}}, \quad n \geq 2. \end{aligned}$$

So arguing as in the proof of Proposition 11 one gets (3.3). This finishes the proof.  $\square$

*Proof of Theorem 5.* We will prove (i)  $\Rightarrow$  (ii)  $\Rightarrow$  (iii)  $\Rightarrow$  (i).

Assume that  $P_\omega : L_{\hat{\nu}}^\infty \rightarrow \mathcal{B}_{\hat{\nu}}^\infty$  is bounded. Then, Proposition 16 together with Lemma B imply (1.1). Hence (ii) is satisfied. Next assuming that (ii) holds, Proposition 12 implies that  $\omega \in \widehat{\mathcal{D}}$  and consequently (1.2) follows from (1.1) and Lemma B, so we get (iii). Finally, if (iii) holds, then using that  $z(B_\zeta^\omega)'(z) = \overline{\zeta(B_\zeta^\omega)'(\zeta)}$ , and [13, Theorem 1], we obtain

$$\begin{aligned} \int_{\mathbb{D}} |(B_\zeta^\omega)'(z)| \frac{\omega(\zeta)}{\hat{\nu}(\zeta)} dA(\zeta) &\lesssim \int_0^{|z|} \frac{1}{\widehat{\omega}(t)(1-t)^2} \int_t^1 \frac{\omega(s)}{\widehat{\nu}(s)} ds dt \\ &\lesssim \int_0^{|z|} \frac{dt}{\widehat{\nu}(t)(1-t)^2} \lesssim \frac{1}{\widehat{\nu}(z)(1-|z|)}, \quad |z| \geq \frac{1}{2}. \end{aligned}$$

Consequently, (i) holds by Proposition 15 and thus the proof is complete.  $\square$

*Proof of Theorem 6.* We will prove (i)  $\Rightarrow$  (iii)  $\Leftrightarrow$  (ii) and (iii)  $\Rightarrow$  (i).

Assuming that  $P_\omega : L_{\hat{\nu}}^\infty \rightarrow \mathcal{B}_{\hat{\nu}}^\infty$  is bounded, Proposition 16 together with Lemma B imply (1.2). Next, if (iii) holds  $\frac{\omega}{\hat{\nu}} \in \widehat{\mathcal{D}}$  by Lemma 7. Moreover, using (1.2) we get

$$\widehat{\nu}(r) \lesssim \frac{\widehat{\omega}(r)}{\int_r^1 \frac{\omega(s) ds}{\widehat{\nu}(s)}} \lesssim \frac{\widehat{\omega}\left(\frac{1+r}{2}\right)}{\int_{\frac{1+r}{2}}^1 \frac{\omega(s) ds}{\widehat{\nu}(s)}} \leq \widehat{\nu}\left(\frac{1+r}{2}\right), \quad 0 < r < 1.$$

That is,  $\nu \in \widehat{\mathcal{D}}$ . Now, bearing in mind Lemma B we conclude that (1.1) is satisfied. Reciprocally, if (ii) holds, (1.2) follows from Lemma B. Finally, if (iii) holds and  $\omega \in \widehat{\mathcal{D}}$ , arguing as above we get  $\nu \in \widehat{\mathcal{D}}$ , so (i) follows from Theorem 5. This finishes the proof.  $\square$

Next, we will obtain some consequences from Theorem 5 and Theorem 6.

**Corollary 17.** *Let  $\omega$  and  $\nu$  be radial weights such that  $P_\omega : L_{\hat{\nu}}^\infty \rightarrow \mathcal{B}_{\hat{\nu}}^\infty$  is bounded. Then  $\omega \in \widehat{\mathcal{D}}$  if and only if  $\nu \in \widehat{\mathcal{D}}$ .*

*Proof.* Assume that  $P_\omega : L_\nu^\infty \rightarrow \mathcal{B}_\nu^\infty$  is bounded. If  $\nu \in \widehat{\mathcal{D}}$ , then  $\omega \in \widehat{\mathcal{D}}$  by Theorem 5. On the other hand, if  $\omega \in \widehat{\mathcal{D}}$ , the proof of (iii) $\Rightarrow$ (ii) in Theorem 6 gives that  $\nu \in \widehat{\mathcal{D}}$ . This finishes the proof.  $\square$

Let us observe that Theorem 1 together with Theorem 5 imply that for any radial weight  $\omega$  and  $\nu \in \mathcal{D}$ , the operators  $P_\omega : L_\nu^\infty \rightarrow \mathcal{H}_\nu^\infty$  and  $P_\omega : L_\nu^\infty \rightarrow \mathcal{B}_\nu^\infty$  are simultaneously bounded. This fact can be also deduced from the following result [?, Theorems 3.2, 3.3].

**Theorem D.** *Let  $\nu$  be a radial weight. Then, the following statements holds:*

- (i)  $H_\nu^\infty$  is continuously embedded in  $\mathcal{B}_\nu^\infty$  if and only if  $\nu \in \widehat{\mathcal{D}}$ ;
- (ii)  $H_\nu^\infty = \mathcal{B}_\nu^\infty$  with equivalence of norms if and only if  $\nu \in \mathcal{D}$ .

However, we are able to prove that the aforementioned result on the boundedness of  $P_\omega$  does not remain true for  $\nu \in \widehat{\mathcal{D}} \setminus \check{\mathcal{D}}$ .

**Corollary 18.** *Let  $\nu \in \widehat{\mathcal{D}} \setminus \check{\mathcal{D}}$ . Then, for any radial weight  $\omega$ ,  $P_\omega$  is not bounded from  $L_\nu^\infty$  to  $H_\nu^\infty$ . However, there exists a radial weight  $\omega_\nu$  such that  $P_{\omega_\nu} : L_\nu^\infty \rightarrow \mathcal{B}_\nu^\infty$  is bounded.*

*Proof.* Since  $\nu \in \widehat{\mathcal{D}} \setminus \check{\mathcal{D}}$ , Corollary 14 implies that  $P_\omega$  is not bounded from  $L_\nu^\infty$  to  $H_\nu^\infty$  for any radial weight  $\omega$ . Now, consider  $\omega_\nu = \nu \widehat{\nu}$ , then  $\widehat{\omega}_\nu(r) = (\widehat{\nu}(r))^2$ ,  $0 \leq r < 1$ , and  $\omega_\nu \in \widehat{\mathcal{D}}$ . So,  $\int_r^1 \frac{\omega_\nu(s)}{\widehat{\nu}(s)} ds = \widehat{\nu}(r) = \frac{\widehat{\omega}_\nu(r)}{\widehat{\nu}(r)}$ ,  $0 \leq r < 1$ . Therefore,  $P_{\omega_\nu} : L_\nu^\infty \rightarrow \mathcal{B}_\nu^\infty$  is bounded, by Theorem 5. This finishes the proof.  $\square$

It is also worth mentioning that there exist  $\omega \in \mathcal{D}$  such that  $P_\omega$  is not bounded from  $L_\nu^\infty$  to  $H_\nu^\infty$  but  $P_\omega : L_{\nu_\omega}^\infty \rightarrow \mathcal{B}_{\nu_\omega}^\infty$  is bounded. Take  $\omega = 1$  and  $\nu(r) = (1-r)^{-1} \left( \log \frac{e}{1-r} \right)^{-2} \in \widehat{\mathcal{D}} \setminus \check{\mathcal{D}}$ , and apply Corollary 14 and Theorem 5

We are also able to prove the following result.

**Corollary 19.** *Let  $\omega \in \widehat{\mathcal{D}} \setminus \check{\mathcal{D}}$ . Then for any radial weight  $\nu$ ,  $P_\omega$  is not bounded from  $L_\nu^\infty$  to  $H_\nu^\infty$ . However there exists a radial weight  $\nu_\omega$  such that  $P_\omega : L_{\nu_\omega}^\infty \rightarrow \mathcal{B}_{\nu_\omega}^\infty$  is bounded.*

*Proof.* Since  $\omega \in \widehat{\mathcal{D}} \setminus \check{\mathcal{D}}$ , Corollary 14 yields that  $P_\omega$  is not bounded from  $L_\nu^\infty$  to  $H_\nu^\infty$  for any radial weight  $\nu$ . Now, for  $\nu_\omega(r) = \frac{d}{dr} \left( (\widehat{\omega}(r))^{\frac{1}{2}} \right)$  we have  $\widehat{\nu}_\omega(r) = (\widehat{\omega}(r))^{\frac{1}{2}}$ ,  $0 \leq r < 1$ , and  $\nu_\omega \in \widehat{\mathcal{D}}$ . Moreover,  $\int_r^1 \frac{\omega(s)}{\widehat{\nu}_\omega(s)} ds = (\widehat{\omega}(r))^{\frac{1}{2}} = \frac{\widehat{\omega}(r)}{\widehat{\nu}_\omega(r)}$ ,  $0 \leq r < 1$ . Therefore,  $P_\omega : L_{\nu_\omega}^\infty \rightarrow \mathcal{B}_{\nu_\omega}^\infty$  is bounded by Theorem 5. This finishes the proof.  $\square$

## 5. STUDY OF THE BOUNDEDNESS OF THE BERGMAN PROJECTIONS FOR EXPONENTIALLY DECREASING WEIGHTS

**5.1. Class of exponentially decreasing weights.** In this section we consider the boundedness of Bergman projections in the space  $L_v^\infty$  for a class of exponentially decreasing weights  $v$ . In order to formulate the results we need some more definitions. As in [5], we say that a continuous function  $\rho : \mathbb{D} \rightarrow \mathbb{R}$  belongs to the class  $\mathcal{L}_0$ , if  $\lim_{|z| \rightarrow 1} \rho(z) = 0$ ,

$$\sup_{z, \zeta \in \mathbb{D}, z \neq \zeta} \frac{|\rho(z) - \rho(\zeta)|}{|z - \zeta|} < \infty \quad (5.1)$$

and for every  $\varepsilon > 0$  there exists a compact  $E \subset \mathbb{D}$  such that  $|\rho(z) - \rho(\zeta)| \leq \varepsilon |z - \zeta|$  whenever  $z, \zeta \in \mathbb{D} \setminus E$ . Furthermore, according to [5], a twice continuously differentiable, real valued subharmonic function  $\varphi$  on  $\mathbb{D}$  is said to belong to the class  $\mathcal{W}_0$ , if  $\Delta \varphi > 0$  and there exists  $\rho \in \mathcal{L}_0$  such that

$$\frac{1}{\sqrt{\Delta \varphi}} \asymp \rho. \quad (5.2)$$

on  $\mathbb{D}$ .

We say that a weight  $v$  on  $\mathbb{D}$  belongs to the class  $\mathcal{E}$ , if it is of the form  $v = e^{-\varphi}$  for some radial function  $\varphi \in \mathcal{W}_0$ . As examples of weights in this class, we mention

$$v(z) = \exp\left(-\alpha/(1-|z|^\ell)^\beta\right) \quad (5.3)$$

where  $\alpha, \beta, \ell > 0$  are constants. For these weights one can take  $\rho(z) = (1-|z|)^{1+\beta/2}$ .

For the weight  $v$  as in (5.3) with  $\ell, \beta = 1$  and  $\omega = v^2$ , the boundedness of  $P_\omega$  on  $L_v^\infty$  was proved in [2]. In the setting of exponentially decreasing weights satisfying the general condition (B) of [8], [9], examples of bounded projections from  $L_v^\infty$  onto  $H_v^\infty$  were constructed in [9]. (These projections are not necessarily of the Bergman type. All weights (5.3) satisfy condition (B), but there are many others, see the mentioned references.) We also mention the following result of [5].

**Theorem E.** *If  $v = e^{-\varphi}$  with  $\varphi \in \mathcal{W}_0$  and  $\omega = v^2$ , then the projection  $P_\omega$  is bounded from  $L_v^\infty$  onto  $H_v^\infty$ .*

We will prove in Theorem 20 a generalization of this result for the class  $\mathcal{E}$  but it is first worthwhile to recall known, related negative results. Indeed, in [3] it was shown that given a weight  $\omega$  as in (5.3) with  $\ell = 2$ , the projection  $P_\omega$  is bounded in  $L_\omega^p$ ,  $1 \leq p \leq \infty$ , if and only if  $p = 2$ . The result was extended in [19]. Furthermore, in [1, Theorems 1 and 2] it was shown that given  $v$  as in (5.3) and another weight  $\omega = \exp(-\tilde{\alpha}/(1-|z|^\ell)^{\tilde{\beta}})$ , then  $P_\omega$  is unbounded in  $L_v^\infty$ , if  $\tilde{\alpha} \neq 2\alpha$  or  $\tilde{\beta} \neq \beta$ .

**5.2. Study of the boundedness  $P_\omega : L_v^\infty \rightarrow H_v^\infty$  for perturbed exponentially decreasing weights.** In view of Theorem E and the above mentioned negative results, it is of interest to know if the projection  $P_\omega$  is bounded in  $L_v^\infty$  for some other, “smaller” perturbations  $v$  of the weight  $\omega^{1/2}$ . We will prove the following generalization, which is our main result concerning exponentially decreasing weights.

**Theorem 20.** *Assume the weight  $w = e^{-\varphi}$  belongs to  $\mathcal{E}$  with  $\varphi, \rho$  satisfying (5.2). Moreover, let  $v \in \widehat{\mathcal{D}}$  and define the weights  $\omega = w^2$  and  $v = w\widehat{v}^t\rho^\sigma$ , where  $t \in \{-1, 0, 1\}$  and  $\sigma$  is any real number. Then, the projection  $P_\omega$  is bounded from  $L_v^\infty$  onto  $H_v^\infty$ .*

Here, the factor  $\widehat{v}^t\rho^\sigma$  can be considered as a modest multiplicative perturbation of the exponential weight  $w$ . For example, for a weight  $w$  as in (5.3), the perturbation could be any negative or positive power of the boundary distance.

*Proof.* We present the proof for  $t = 1$ , similar proofs work for  $t \in \{-1, 0\}$ . We will need some results of [5]. The Bergman reproducing kernel  $B_z^\omega$  in

$$P_\omega f(z) = \int_{\mathbb{D}} f(\zeta) \overline{B_z^\omega(\zeta)} \omega(\zeta) dA(\zeta) \quad (5.4)$$

has by [5, Theorem 3.2] the upper bound

$$|B_z^\omega(\zeta)| \leq C \frac{e^{\varphi(z)+\varphi(\zeta)}}{\rho(z)\rho(\zeta)} e^{-\alpha d_\rho(z,\zeta)} \quad (5.5)$$

for some  $\alpha > 0$ , and the last factor also has for any  $M > 0$  the upper bound

$$e^{-\alpha d_\rho(z,\zeta)} \leq C(M) \left( \frac{\min(\rho(z), \rho(\zeta))}{|z-\zeta|} \right)^M, \quad (5.6)$$

see formula (23) of the reference. We will choose a large enough  $M$  later. In (5.6),  $d_\rho$  is defined according to [5] as the distance function

$$d_\rho(z, \zeta) = \inf_\gamma \int_0^1 |\gamma'(t)| \frac{dt}{\rho(\gamma(t))}, \quad (5.7)$$

where  $z, \zeta \in \mathbb{D}$  and the infimum is taken over all piecewise  $C^1$  curves  $\gamma : [0, 1] \rightarrow \mathbb{D}$  with  $\gamma(0) = z, \gamma(1) = \zeta$ .

Let  $\nu \in \widehat{\mathcal{D}}$  and  $\nu = w\widehat{\nu}\rho^\sigma = e^{-\varphi}\widehat{\nu}\rho^\sigma$  be as in the assumptions of the theorem, and let  $f \in L_v^\infty$  be such that  $\|f\|_{\infty, \nu} \leq 1$ . We can estimate for every  $z \in \mathbb{D}$ , by using (5.5) and  $|f(\zeta)| \leq \rho(\zeta)^{-\sigma} e^{\varphi(\zeta)} / \widehat{\nu}(\zeta)$ ,

$$\begin{aligned} v(z)|P_\omega f(z)| &\leq w(z)\rho(z)^\sigma \widehat{\nu}(z) \int_{\mathbb{D}} |B_z^\omega(\zeta)| |f(\zeta)| e^{-2\varphi(\zeta)} dA(\zeta) \\ &\leq w(z)\widehat{\nu}(z)\rho(z)^\sigma e^{\varphi(z)} \int_{\mathbb{D}} \frac{e^{\varphi(\zeta)}}{\rho(z)\rho(\zeta)} e^{-\alpha d_\rho(z, \zeta)} \frac{1}{e^{\varphi(\zeta)} \widehat{\nu}(\zeta) \rho(\zeta)^\sigma} dA(\zeta) \\ &\leq \widehat{\nu}(z)\rho(z)^{\sigma-1} \int_{\mathbb{D}} \frac{e^{-\alpha d_\rho(z, \zeta)}}{\widehat{\nu}(\zeta)\rho(\zeta)^{\sigma+1}} dA(\zeta). \end{aligned} \quad (5.8)$$

We continue by dividing the integration domain  $\mathbb{D}$  into two subsets,

$$\mathbb{A}_z = \left\{ \zeta \in \mathbb{D} : |\zeta| \geq \frac{1}{2}(1 + |z|) \right\} \quad \text{and} \quad \mathbb{B}_z = \mathbb{D} \setminus \mathbb{A}_z. \quad (5.9)$$

Then, there holds, for  $\zeta \in \mathbb{B}_z$ ,

$$\widehat{\nu}(\zeta) \geq \widehat{\nu}\left(\frac{1}{2}(1 + |z|)\right) \geq C\widehat{\nu}(z), \quad (5.10)$$

since  $\widehat{\nu}$  is decreasing with respect to the radius and  $\nu \in \widehat{\mathcal{D}}$ . This yields

$$\begin{aligned} \widehat{\nu}(z)\rho(z)^{\sigma-1} \int_{\mathbb{B}_z} \frac{e^{-\alpha d_\rho(z, \zeta)}}{\widehat{\nu}(\zeta)\rho(\zeta)^{\sigma+1}} dA(\zeta) &\leq C\rho(z)^{\sigma-1} \int_{\mathbb{B}_z} \frac{e^{-\alpha d_\rho(z, \zeta)}}{\rho(\zeta)^{\sigma+1}} dA(\zeta) \\ &\leq C\rho(z)^{\sigma-1} \int_{\mathbb{D}} \frac{e^{-\alpha d_\rho(z, \zeta)}}{\rho(\zeta)^{\sigma+1}} dA(\zeta). \end{aligned} \quad (5.11)$$

By Corollary 3.1 of [5], this expression is bounded by a constant.

To estimate the integral over  $\mathbb{A}_z$  we first note that since  $\nu \in \widehat{\mathcal{D}}$ , Lemma B(ii) shows that there are constants  $C, \beta > 0$  such that

$$\frac{\widehat{\nu}(z)}{\widehat{\nu}(\zeta)} \leq C \frac{(1 - |z|)^\beta}{(1 - |\zeta|)^\beta}$$

for all  $z, \zeta$  with  $|z| \leq |\zeta|$ . We use this and (5.6) to write

$$\begin{aligned} \widehat{\nu}(z)\rho(z)^{\sigma-1} \int_{\mathbb{A}_z} \frac{e^{-\alpha d_\rho(z, \zeta)}}{\widehat{\nu}(\zeta)\rho(\zeta)^{\sigma+1}} dA(\zeta) &\leq C(1 - |z|)^\beta \rho(z)^{\sigma-1} \int_{\mathbb{A}_z} \frac{e^{-\alpha d_\rho(z, \zeta)}}{\rho(\zeta)^{\sigma+1}(1 - |\zeta|)^\beta} dA(\zeta) \\ &\leq C(M)(1 - |z|)^\beta \rho(z)^{\sigma-1} \int_{\mathbb{A}_z} \frac{1}{(1 - |\zeta|)^\beta \rho(\zeta)^{\sigma+1}} \left( \frac{\min(\rho(z), \rho(\zeta))}{|z - \zeta|} \right)^M dA(\zeta) \end{aligned} \quad (5.12)$$

We now choose the number  $\gamma > 0$  such that  $\sigma + \gamma - 1 > 0$  and then  $M$  such that  $M > \sigma + \beta + \gamma + 3$ . Then, we use

$$\min(\rho(z), \rho(\zeta))^M \leq \rho(z)^\gamma \rho(\zeta)^{M-\gamma}$$

to bound (5.12) by

$$C_M(1 - |z|)^\beta \rho(z)^{\sigma+\gamma-1} \int_{\mathbb{A}_z} \frac{\rho(\zeta)^{M-(\gamma+\sigma+1)}}{(1 - |\zeta|)^\beta |z - \zeta|^M} dA(\zeta). \quad (5.13)$$

Note that by (5.1), the function  $\rho$  has the bound  $\rho(\zeta) \leq C(1 - |\zeta|)$  for all  $\zeta \in \mathbb{D}$ . Thus, (5.13) is not larger than

$$C_M(1 - |z|)^{\sigma+\beta+\gamma-1} \int_{\mathbb{A}_z} \frac{(1 - |\zeta|)^{M-(\sigma+\beta+\gamma+1)}}{|z - \zeta|^M} dA(\zeta). \quad (5.14)$$

Since  $|z - \zeta| \geq C|1 - z\bar{\zeta}|$  for  $\zeta \in \mathbb{A}$ , the Forelli-Rudin estimates [20, Lemma 3.10] and the choice of the number  $M$  show that (5.14) is bounded by a constant.

Combining (5.8) and (5.11)–(5.14) yield  $\sup_{z \in \mathbb{D}} v(z)|P_\omega f(z)| \leq C$  for all  $f \in L_v^\infty$  with  $\|f\|_{\infty, v} \leq 1$ , which proves the theorem.  $\square$

Finally, we will apply Theorem E in order to get the following result for the special exponential weights and set  $w = e^{-\varphi}$ ,  $\varphi(z) = \alpha(1 - |z|^2)^{-\beta}$ , and

$$v(z) = (1 - |z|^2)^\gamma w(z), \quad \omega(z) = (1 - |z|^2)^{2\sigma} w(z)^2, \quad (5.15)$$

where the parameters satisfy  $\alpha, \beta > 0$ ,  $\sigma, \gamma \in \mathbb{R}$ .

**Proposition 21.** *Let the weights  $v$  and  $\omega$  be as in (5.15). Then, the projection  $P_\omega$  is bounded from  $L_v^\infty$  onto  $H_v^\infty$ .*

*Proof.* We define

$$\psi(z) = \frac{\alpha}{(1 - |z|^2)^\beta} - \sigma \log(1 - |z|^2)$$

and observe that  $\psi \in \mathcal{W}_0$ , since there holds  $1/\sqrt{\Delta\psi(z)} \asymp R(z)$  for the function  $R(z) = (1 - |z|^2)^{1+\beta/2} \in \mathcal{L}_0$ . We define the weight  $W = e^{-\psi} \in \mathcal{E}$ . Then, obviously,  $\omega = W^2$  and  $v(z) = R(z)^{\frac{\gamma-\sigma}{1+\beta/2}} W(z)$  so that the result follows from Theorem 20 by choosing  $W \rightarrow w$  and  $R \rightarrow \rho$  and setting  $t = 0$ .  $\square$

## REFERENCES

- [1] J. Bonet, W. Lusky and J. Taskinen, Unbounded Bergman projections on weighted spaces with respect to exponential weights, *Integr. Equ. Oper. Theory* (2021) 61–93.
- [2] O. Constantin and J. A. Peláez, Boundedness of the Bergman projection on  $L^p$ -spaces with exponential weights, *Bull. Sci. Math.* 139 (2015) 245–268.
- [3] M. Dostanic, Unboundedness of the Bergman projections on  $L^p$ -spaces with exponential weights, *Proc. Edinb. Math. Soc.* 47 (2004) 111–117.
- [4] P. Duren, *Theory of  $H^p$  spaces*. Pure and Applied Mathematics, Vol. 38 Academic Press, New York-London 1970 xii+258 pp.
- [5] Z. Hu, X. Lv and A. P. Schuster, Bergman spaces with exponential weights, *J. Funct. Anal.* 276 (2019), no.5, 1402–1429.
- [6] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces 1*. *Ergebnisse der Mathematik und ihrer Grenzgebiete* 92, Springer (1977).
- [7] W. Lusky, On weighted spaces of harmonic and holomorphic functions, *J. London. Math. Soc.* (2) 51 (1995), 309–320.
- [8] W. Lusky, On the isomorphism classes of weighted spaces of harmonic and holomorphic functions, *Studia Math.* 175 (2006), no. 1, 19–45.
- [9] W. Lusky and J. Taskinen, Bounded holomorphic projections for exponentially decreasing weights, *J. Function Spaces Appl.* 6 (2008), no. 1, 59–70.
- [10] A. M. Moreno, J. A. Peláez and E. de la Rosa, Fractional derivative description of the Bloch space, *Potential Anal.* <https://doi.org/10.1007/s11118-023-10119-z>.
- [11] J. A. Peláez, Small weighted Bergman spaces, *Proceedings of the summer school in Complex and Harmonic analysis and related topics*, (2016).
- [12] J. A. Peláez and J. Rättyä, Weighted Bergman spaces induced by rapidly increasing weights, *Mem. Amer. Math. Soc.* 227 (2014), no. 1066.
- [13] J. A. Peláez and J. Rättyä, Two weight inequality for Bergman projection, *J. Math. Pures Appl.* 105 (2016), 102–130.
- [14] J. A. Peláez and J. Rättyä, Bergman projection induced by radial weight. *Adv. Math.* 391 (2021), Paper No. 107950, 70 pp.
- [15] J. A. Peláez and E. de la Rosa, Littlewood-Paley inequalities for fractional derivative on Bergman spaces. *Ann. Fenn. Math.* 47 (2022), no. 2, 1109–1130.



- [16] J. A. Peláez, J. Rättyä and B. D. Wick, Bergman projection induced by kernel with integral representation, *J. Anal. Math.* 138 (2019), no.1, 325–360.
- [17] A. Pełczyński, On the isomorphism of the spaces  $M$  and  $m$ , *Bull. Acad. Pol. Sci.* 6 (1958), 695–696.
- [18] W. Rudin, *Functional analysis*, McGraw-Hill (1973).
- [19] Y. E. Zeytuncu,  $L^p$ -regularity of weighted Bergman projections, *Trans. Am. Math. Soc.* 365 (2013) 2959–2976.
- [20] K. Zhu, *Operator theory in function spaces*, 2nd. *Mathematical Surveys and Monographs*, vol. 138, 2nd edn. American Mathematical Society, Providence (2007)

DEPARTAMENTO DE ANALISIS MATEMÁTICO, UNIVERSIDAD DE MÁLAGA, CAMPUS DE TEATINOS, 29071 MALAGA, SPAIN

*Email address:* `alvarommorenolopez@uma.es`

DEPARTAMENTO DE ANALISIS MATEMÁTICO, UNIVERSIDAD DE MÁLAGA, CAMPUS DE TEATINOS, 29071 MALAGA, SPAIN

*Email address:* `japelaez@uma.es`

UNIVERSITY OF HELSINKI, DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O.Box 68, 00014 HELSINKI, FINLAND

*Email address:* `jari.taskinen@helsinki.fi`