

CONVERGENCE OF TSIRELSON CONVOLUTION SYSTEMS OF PROBABILITY SPACES

REMUS FLORICEL AND PATRICK MELANSON

ABSTRACT. We associate two specific projective systems of probability spaces with any Tsirelson convolution system. If the projective limits of these systems exist, then we call the convolution system convergent and K -convergent, respectively. It is shown that convergent convolution systems give rise to continuous products of probability spaces, while K -convergent convolution systems lead to flow systems. We investigate the relationship between convergence and K -convergence, as well as their connections to two-parameter product systems of Hilbert spaces.

1. INTRODUCTION

Motivated by Arveson's theory of product systems [1, 2], B. Tsirelson initiated the study of two-parameter product systems of Hilbert spaces in [12, 13] (see also [11]) through probabilistic methods, including the analysis of stochastic processes and flows. Considering that convolution semigroups of measures can naturally be used to construct stochastic processes via Kolmogorov's extension theorem, which in turn lead to product systems, one of Tsirelson's initial steps in his investigation was to introduce the concept of a convolution system of standard probability spaces as a generalization of convolution semigroups. However, unlike convolution semigroups of measures, convolution systems may not always give rise to a stochastic flow. This limitation primarily stems from the fact that the category of probability spaces generally does not admit projective limits, making any attempt to generalize Kolmogorov's extension theorem quite challenging [3, 4, 10]. When a convolution system leads to a stochastic flow, or a flow system, as it is called in [13, Sec. 3b], the flow system can then be directly used to construct a continuous product of probability spaces (CPPS), largely due to the specific properties of the standard spaces used in this construction (see [13, Sec. 3c]). This, in turn, leads to a two-parameter product system of Hilbert spaces by transitioning to the corresponding L^2 -spaces.

The purpose of this note is to show that continuous products of (not necessarily standard) probability spaces over a linearly ordered set can be obtained from some convolution systems, even if they may not admit flow

2020 *Mathematics Subject Classification.* Primary 28A33, 60A10, 60B15, 46L53, 46L55.

Key words and phrases. convolution systems, continuous product of probability spaces, flow systems, subproduct and product system of Hilbert spaces.

systems. It also explores the interplay between convolution systems that admit CPPS and those that admit flow systems. This is achieved by associating two projective systems of probability spaces with a given convolution system (see Proposition 3.1 and Proposition 4.1). If the projective limits of these systems exist, they will be referred to as convergent and K -convergent, respectively. In these cases, they give rise to a continuous product of probability spaces and a flow system, respectively

We show in Theorem 3.3 that the product system of L^2 -Hilbert spaces of the projective CPPS associated with a convergent convolution system can be described in terms of the subproduct system of L^2 -Hilbert spaces of the convolution system through an inductive limit construction. We also show in Proposition 4.2 that any K -convergent system is convergent and find in Theorem 4.5 the necessary and sufficient conditions for a convergent system to be K -convergent.

At this end, we note that some of the techniques employed in this paper can be regarded as the projective counterparts to the inductive techniques used in [5] within the context of C^* -algebras.

Notation 1.1. All the notation used in this article is standard. The category **Prob** of probability spaces consists of probability spaces $(\Omega, \mathcal{F}, \mu)$, where \mathcal{F} is a σ -field of subsets of Ω and μ is a probability measure on \mathcal{F} , with morphisms $T : (\Omega, \mathcal{F}, \mu) \rightarrow (\Omega', \mathcal{F}', \mu')$ given by measurable and measure-preserving transformations $T : \Omega \rightarrow \Omega'$. Any such morphism T induces the Koopman isometry $U_T : L^2(\mu') \rightarrow L^2(\mu)$ by $U_T(f) = f \circ T$ for $f \in L^2(\mu')$. As usual, relations between sets and functions are to be interpreted up to a set of measure zero.

2. BACKGROUND

2.1. Convolution systems of probability spaces and subproduct systems of Hilbert spaces. In this subsection, we present the main objects of study in this article, all of which were formally introduced by B. Tsirelson in [12, 13]. It is important to note that Tsirelson's work assumes all probabilistic spaces to be standard. To maintain the highest level of generality in this note, and in line with the development of an "uncountable" measure theory as proposed in [7], we entirely drop this assumption. Additionally, we will work with convolution systems over arbitrary linearly ordered sets instead of \mathbb{R} , as suggested in [13]. Similarly, for the same reasons, we do not automatically assume the separability of Hilbert spaces, which is an important requirement in Arveson's theory of product systems.

Definition 2.1. (i) A convolution system $\mathcal{S} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{S}}$ over a linearly ordered set (\mathbb{S}, \leq) consists of a family $\{(\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}) \mid s, t \in \mathbb{S}, s < t\}$ of probability spaces and a family $\{T_{r,s,t} \mid r, s, t \in \mathbb{S}, r < s < t\}$ of morphisms $T_{r,s,t} : (\Omega_{r,s}, \mathcal{F}_{r,s}, \mu_{r,s}) \times (\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}) \rightarrow (\Omega_{r,t}, \mathcal{F}_{r,t}, \mu_{r,t})$, called the multiplication of the system, which is associative in the sense that the

diagram

$$(1) \quad \begin{array}{ccc} \Omega_{r,s} \times \Omega_{s,t} \times \Omega_{t,u} & \xrightarrow{T_{r,s,t} \times \text{id}_{\Omega_{t,u}}} & \Omega_{r,t} \times \Omega_{t,u} \\ \text{id}_{\Omega_{r,s}} \times T_{s,t,u} \downarrow & & \downarrow T_{r,t,u} \\ \Omega_{r,s} \times \Omega_{s,u} & \xrightarrow{T_{r,s,u}} & \Omega_{r,u}, \end{array}$$

commutes for all $r < s < t < u$ in \mathbb{S} .

(ii) A continuous product of probability spaces, abbreviated as CPPS, is a convolution system whose multiplication consists of isomorphisms of probability spaces.

(iii) A morphism of convolution systems, from $\mathcal{S} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{S}}$ to $\mathcal{S}' = \{\Omega'_{s,t}, \mathcal{F}'_{s,t}, \mu'_{s,t}, T'_{r,s,t}\}_{\mathbb{S}}$, is a family $\theta = \{\theta_{s,t}\}_{s < t}$ of morphisms of probability spaces $\theta_{s,t} : (\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}) \rightarrow (\mathcal{F}'_{s,t}, \Sigma'_{s,t}, \mu'_{s,t})$, which makes the diagram

$$(2) \quad \begin{array}{ccc} \Omega_{r,s} \times \Omega_{s,t} & \xrightarrow{\theta_{r,s} \times \theta_{s,t}} & \Omega'_{r,s} \times \Omega'_{s,t} \\ T_{r,s,t} \downarrow & & \downarrow T'_{r,s,t} \\ \Omega_{r,t} & \xrightarrow{\theta_{r,t}} & \Omega'_{r,t} \end{array}$$

commutative, for all $r < s < t$ in \mathbb{S}

(iv) A flow system $((\Omega, \mathcal{F}, P), \{X_{s,t}\})_{\mathcal{S}}$ over a convolution system $\mathcal{S} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{S}}$ consists of a probability space (Ω, \mathcal{F}, P) and a family $\{X_{s,t} \mid s, t \in \mathbb{S}, s < t\}$ of morphisms of probability spaces $X_{s,t} : (\Omega, \mathcal{F}, P) \rightarrow (\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t})$ that satisfies the following conditions:

- (1) \mathcal{F} is the joint of the σ -fields $\mathcal{F}_{s,t}^* = X_{s,t}^{-1}(\mathcal{F}_{s,t})$, i.e., the σ -field generated by the field $\mathcal{F}_0 = \bigcup_{s < t} \mathcal{F}_{s,t}^*$;
- (2) $X_{t_1, t_2}, X_{t_1, t_2, \dots}, X_{t_1, t_2}$ are independent, for all $t_1 < t_2 < \dots < t_n$ in \mathbb{S} ;
- (3) $X_{r,t}(\omega) = T_{r,s,t}(X_{r,s}(\omega), X_{s,t}(\omega))$, $\omega \in \Omega$, for all $r < s < t$ in \mathbb{S} .

Example 2.1. i) Let (S, \mathcal{G}) be a measurable semigroup, consisting of a semigroup S and a σ -field \mathcal{G} on S such that the semigroup operation $(s, t) \mapsto st$ is \mathcal{G} -measurable. The convolution $\mu * \nu$ of two probability measures μ and ν on \mathcal{G} is defined as the push-forward of the product measure $\mu \times \nu$ by the semigroup operation. Suppose that μ is an idempotent probability measure on \mathcal{G} , i.e., $\mu * \mu = \mu$. Then it gives rise to a trivial convolution system $\mathcal{S}_\mu = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t} = \mu, T_{r,s,t}\}_{\mathbb{S}}$ over any linearly ordered set (\mathbb{S}, \leq) , where $\Omega_{s,t} = S$, $\mathcal{F}_{s,t} = \mathcal{G}$ and $T_{r,s,t}$ is given by the semigroup operation, for all $r < s < t$ in \mathbb{S} . We recall that the structure of idempotent regular Borel probability measures on locally compact Hausdorff second-countable topological semigroups is well-understood (see e.g. [8, Th. 2.8]).

ii) More generally, let $\{\mu_t\}_{t \in \mathbb{R}}$ be a one-parameter convolution semigroup of probability measures over a measurable semigroup (S, \mathcal{G}) . Then $\mathcal{S}_{\{\mu_t\}} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{R}}$ is a convolution system over the set of all

real numbers, where $\Omega_{s,t} = S$, $\mathcal{F}_{s,t} = \mathcal{G}$, $\mu_{s,t} = \mu_{s-t}$, and $T_{r,s,t}$ is given by the semigroup operation, for all real numbers $r < s < t$.

Observation 2.1. Tsirelson’s method for constructing CPPSs from convolution systems of standard probability spaces [13, Sect. 3c] does not directly involve the given convolution system, but rather the flow system associated with it, if one exists. Also, Tsirelson showed in [13] that a convolution system over $\mathbb{S} = (\mathbb{R}, \leq)$ admits a flow system if and only if it is separable, in the sense of Definition 3b4 in [13]. For example, taking $S = (\mathbb{Z}/m\mathbb{Z}, +)$ and μ being the uniform distribution on $\mathbb{Z}/m\mathbb{Z}$, the convolution system constructed as in Example 2.1 (i), based on this data, is not separable. On the other hand, taking $S = (\mathbb{R}, +)$, the system constructed in Example 2.1 (ii) is separable.

Convolution systems and CPPSs can be regarded as the probabilistic counterparts to the notions of two-parameter subproduct systems and product systems of Hilbert spaces, respectively. However, the former appear to be more versatile than the latter. The concept of two-parameter product systems of Hilbert spaces was first introduced by Tsirelson in [12] under the name “local continuous product of Hilbert spaces”, obtained by excluding unbounded intervals from the definition of a continuous product of Hilbert spaces. Meanwhile, two-parameter subproduct systems were formally introduced in [5] under the name “Tsirelson subproduct system”, although the concept was well-known long before their formal introduction.

Definition 2.2. A two-parameter subproduct system of Hilbert spaces, or subproduct system for short, $\mathcal{H} = \{H_{s,t}, U_{r,s,t}\}_{\mathbb{S}}$ over a linearly ordered set (\mathbb{S}, \leq) consists of a family $\{H_{s,t} \mid s, t \in \mathbb{S}, s < t\}$ of Hilbert spaces $H_{s,t}$ and a family $\{U_{r,s,t} \mid r, s, t \in \mathbb{S}, r < s < t\}$ of isometries $U_{r,s,t} : H_{r,t} \rightarrow H_{r,s} \otimes H_{s,t}$ that satisfy the co-associativity law

$$(3) \quad (1_{H_{r,s}} \otimes U_{s,t,u}) U_{r,s,u} = (U_{r,s,t} \otimes 1_{H_{t,u}}) U_{r,t,u},$$

for all $r < s < t < u$ in \mathbb{S} .

If the operators $U_{r,s,t}$ are all unitary operators, then $\mathcal{H} = \{H_{s,t}, U_{r,s,t}\}_{\mathbb{S}}$ will be referred to as a two-parameter product system of Hilbert spaces, or product system for short.

The concept of morphism/isomorphism of subproduct systems is naturally defined, similarly to that of morphism/isomorphism of convolution systems.

Observation 2.2. Any convolution system $\mathcal{S} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{S}}$ gives rise to the subproduct system

$$L^2(\mathcal{S}) = \{L^2(\mu_{s,t}), U_{T_{r,s,t}}\}_{\mathbb{S}},$$

where $U_{T_{r,s,t}}$ is the Koopman isometry induced by $T_{r,s,t}$. If \mathcal{S} is a CPPS, then $L^2(\mathcal{S})$ is a product system. Moreover, any morphism $\theta = \{\theta_{s,t}\}_{0 < s < t}$ of convolution systems, from \mathcal{S} to \mathcal{S}' , induces the morphism $U_\theta = \{U_{\theta_{s,t}}\}_{0 < s < t}$ of subproduct systems, from $L^2(\mathcal{S}')$ to $L^2(\mathcal{S})$.

2.2. Projective systems of probability spaces. In this sub-section, we briefly recall the notion of a projective system and projective limit of probability spaces, which is central to the purpose of this paper. For a comprehensive discussion of the subject, we refer the reader to [3, 4, 10].

A projective system of probability spaces over a directed set (I, \leq) is a pair $\{(\Omega_i, \mathcal{F}_i, \mu_i)\}, T_{i,j}$ consisting of a family $\{(\Omega_i, \mathcal{F}_i, \mu_i)\}_{i \in I}$ of probability spaces and a family $\{T_{i,j}\}_{i, j \in I, i \leq j}$ of morphisms of probability spaces $T_{i,j} : (\Omega_j, \mathcal{F}_j, \mu_j) \rightarrow (\Omega_i, \mathcal{F}_i, \mu_i)$, $i \leq j$, satisfying the compatibility condition $T_{i,j}T_{j,k} = T_{i,k}$, for all $i \leq j \leq k$, and $T_{i,i} = \text{id}_{\Omega_i}$.

Let $\Omega = \varprojlim \{\Omega_i, T_{i,j}\}$ be the projective limit of the projective system $\{\Omega_i, T_{i,j}\}$ over (I, \leq) in the category **Set** of sets, i.e., Ω is the subset of the cartesian product $\Omega^I = \times_{i \in I} \Omega_i$ consisting of those elements $\omega = \{\omega_i\}_{i \in I}$, called threads, such that for each $i \leq j$ in I , $\omega_i = T_{i,j}(\omega_j)$. We notice that the set Ω can be empty, even in cases where the mappings $T_{i,j}$ are surjective [9, 14]. If Ω is non-empty and the coordinate projections

$$T_i : \Omega \ni \{\omega_j\}_{j \in I} \mapsto \omega_i \in \Omega_i$$

are all surjective, then, using terminology from [3], we say that the **Set**-projective system $\{\Omega_i, T_{i,j}\}$ is *simply maximal*. In this in case, the mappings $T_{i,j}$ must also be all surjective.

Assuming further that the system $\{\Omega_i, T_{i,j}\}$ is simply maximal, we consider the field $\mathcal{F}_0 = \bigcup_{i \in I} T_i^{-1}(\mathcal{F}_i)$ and the joint $\mathcal{F} = \bigvee_{i \in I} T_i^{-1}(\mathcal{F}_i)$ of the σ -fields $T_i^{-1}(\mathcal{F}_i)$. Let $\mu : \mathcal{F}_0 \rightarrow [0, 1]$ be the finitely additive set function, defined uniquely by the equation

$$(4) \quad \mu(T_i^{-1}(A)) = \mu_i(A), \quad A \in \mathcal{F}_i.$$

Generally, μ may fail to be σ -additive and thus cannot be extended to \mathcal{F} .

Definition 2.3. A simply maximal projective system of probability spaces $\{(\Omega_i, \mathcal{F}_i, \mu_i), T_{i,j}\}$ is said to be convergent if μ is σ -additive on \mathcal{F}_0 .

If this is the case, then we shall use the same notation, μ , to denote the (unique) σ -additive extension of μ to \mathcal{F} . The resulting probability space

$$(\Omega, \mathcal{F}, \mu) := \varprojlim_I \{(\Omega_i, \mathcal{F}_i, \mu_i), T_{i,j}\}$$

is called the projective limit of $\{(\Omega_i, \mathcal{F}_i, \mu_i), T_{i,j}\}$.

Numerous criteria for the existence of the projective limit have been identified (see [4, 6, 10] and the references therein); in this paper, we employ a simple yet effective one for our purposes. If there exists an ‘‘upper bound’’, consisting on a probability space (N, \mathcal{N}, ν) and a family $\{S_i\}_{i \in I}$ of epimorphisms $S_i : (N, \mathcal{N}, \nu) \rightarrow (\Omega_i, \mathcal{F}_i, \mu_i)$ such that $\mathcal{N} = \bigvee_{i \in I} S_i^{-1}(\mathcal{F}_i)$ and $T_{i,j}S_j = S_i$, for all $i, j \in I$ with $i \leq j$ then the set-function μ in (4) must be σ -additive, and thus the projective limit exists. From this perspective, the projective limit is the least upper bound (N, \mathcal{N}, ν) with the aforementioned properties. Consequently, it is unique (up to isomorphism) and possesses

the property that $(\Omega, \mathcal{F}, \mu) = \varprojlim_{I_0} \{(\Omega_i, \mathcal{F}_i, \mu_i), T_{i,j}\}$, for any cofinal subset I_0 of (I, \leq) .

2.3. Finite Partitions. Let (\mathbb{S}, \leq) be a linearly ordered set. For any two elements $s, t \in \mathbb{S}$, $s < t$, consider the partially ordered set $(\mathcal{K}_{s,t}, \subseteq)$ of all finite partitions of the interval $[s, t]$, ordered by inclusion. We notice that if $I = \{s = \iota_0 < \iota_1 < \iota_2 < \cdots < \iota_m < \iota_{m+1} = t\} \in \mathcal{K}_{s,t}$, then any refinement $J \in \mathcal{K}_{s,t}$ of I , i.e., $I \subseteq J$, can be written as

$$(5) \quad J = I_0 \cup I_1 \cup \cdots \cup I_m,$$

where $I_i = \{j \in J, \iota_i \leq j \leq \iota_{i+1}\} = \{\iota_i = \iota_{i_0} < \iota_{i_1} < \cdots < \iota_{i_{n_{I_i}}} < \iota_{i+1}\} \in \mathcal{K}_{\iota_i, \iota_{i+1}}$, for some $n_{I_i} \in \mathbb{N}$ depending on the partition I_i , for all $0 \leq i \leq m$.

We also consider the set $\mathcal{K} = \bigcup_{s < t} \mathcal{K}_{s,t}$ of all finite subsets $I \subset \mathbb{S}$, $|I| \geq 2$, ordered by inclusion as well. As above, if $I \in \mathcal{K}_{s,t}$ and $J \in \mathcal{K}$ is a refinement of I , then J can be written as

$$(6) \quad J = I_L \cup \tilde{I} \cup I_R,$$

where $I_L = \{j \in J \mid j \leq \iota_0\}$, $I_R = \{j \in J \mid j \geq \iota_{m+1}\}$ and $\tilde{I} = I_0 \cup \cdots \cup I_m \in \mathcal{K}_{s,t}$, with I_k defined as above. Note that \tilde{I} is a refinement of I .

3. CONVERGENT CONVOLUTION SYSTEMS AND ASSOCIATED CPPSs

Let $\mathcal{S} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{S}}$ be a convolution system. For any two elements $s, t \in \mathbb{S}$, $s < t$, and any partition $I \in \mathcal{K}_{s,t}$, $I = \{s = \iota_0 < \iota_1 < \iota_2 < \cdots < \iota_m < \iota_{m+1} = t\}$, we consider the product measure space $(\Omega_I, \mathcal{F}_I, \mu_I)$ over I , where $\Omega_I = \Omega_{\iota_0, \iota_1} \times \Omega_{\iota_1, \iota_2} \times \cdots \times \Omega_{\iota_m, \iota_{m+1}}$, $\mathcal{F}_I = \mathcal{F}_{\iota_0, \iota_1} \otimes \mathcal{F}_{\iota_1, \iota_2} \otimes \cdots \otimes \mathcal{F}_{\iota_m, \iota_{m+1}}$, and $\mu_I = \mu_{\iota_0, \iota_1} \times \mu_{\iota_1, \iota_2} \times \cdots \times \mu_{\iota_m, \iota_{m+1}}$.

For any refinement $J \in \mathcal{K}_{s,t}$ of I , we also consider the mapping $T_{I,J} : \Omega_J \rightarrow \Omega_I$ defined as follows:

(i) If I is the trivial partition, $I = \{s, t\}$, and $J = \{s = j_0 < j_1 < j_2 < \cdots < j_n < j_{n+1} = t\}$, then $T_{I,J} : \Omega_J \rightarrow \Omega_{s,t}$ is defined iteratively as

$$T_{I,J} = \begin{cases} T_{j_0, j_1, j_2} & \text{if } n = 1 \\ T_{j_0, j_n, j_{n+1}} (T_{\{j_0, j_n\}, J \setminus \{j_{n+1}\}} \times \text{id}_{\Omega_{j_n, j_{n+1}}}) & \text{if } n \geq 2 \end{cases}$$

where $J \setminus \{j_{n+1}\} \in \mathcal{K}_{s, j_n}$ is the partition obtained by removing the endpoint $t = j_{n+1}$ from J .

(ii) If $I = \{s = \iota_0 < \iota_1 < \iota_2 < \cdots < \iota_m < \iota_{m+1} = t\}$, $I \subseteq J$, are arbitrary partitions, then by writing J it as in (5), we define

$$T_{I,J} = T_{\{\iota_0, \iota_1\}, I_0} \times T_{\{\iota_1, \iota_2\}, I_1} \times \cdots \times T_{\{\iota_m, \iota_{m+1}\}, I_m},$$

where $T_{\{\iota_k, \iota_{k+1}\}, I_k} : \Omega_{I_k} \rightarrow \Omega_{\iota_k, \iota_{k+1}}$ are as in (i).

(iii) If $I = J$, we set $T_{I,I} = \text{id}_{\Omega_I}$.

The construction carried out above ensures that $T_{I,J} : (\Omega_J, \mathcal{F}_J, \mu_J) \rightarrow (\Omega_I, \mathcal{F}_I, \mu_I)$ is a morphism of probability spaces, for all $I, J \in \mathcal{K}_{s,t}$, $I \subseteq J$. The newly created system is projective, as shown below.

Proposition 3.1. *Let $\mathcal{S} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{S}}$ be a convolution system. Then the system $(\{(\Omega_I, \mathcal{F}_I, \mu_I)\}, \{T_{I,J}\})$ is a projective system of probability spaces over the directed set $(\mathcal{K}_{s,t}, \subseteq)$, for all $s < t$ in \mathbb{S} .*

Proof. Let $s < t$ be two elements of \mathbb{S} . We show that $T_{I,K} = T_{I,J}T_{J,K}$, for all partitions $I, J, K \in \mathcal{K}_{s,t}$, $I \subseteq J \subseteq K$. If $I = \{s, t\}$ is the trivial partition, then the identity can be deduced easily. For a general partition $I = \{s = \iota_0 < \iota_1 < \dots < \iota_m < \iota_{m+1} = t\}$, we use (5) to decompose J in terms of I as $J = I_0 \cup I_1 \cup \dots \cup I_m$, where $I_i \in \mathcal{K}_{\iota_i, \iota_{i+1}}$, for all $0 \leq i \leq m$. Similarly, we can decompose K in terms of J , thus $K = J_0 \cup J_1 \cup \dots \cup J_\ell$, where $J_0 \in \mathcal{K}_{\iota_0, \iota_1}$, $J_1 \in \mathcal{K}_{\iota_0, \iota_2}$, \dots , $J_\ell \in \mathcal{K}_{\iota_{m+1}, \iota_{m+1}}$. However, we can also decompose K

in terms of I . Explicitly, we get $K = I'_0 \cup I'_1 \cup \dots \cup I'_m$, where $I'_0 = \bigcup_{i=0}^{n_{I_0}} J_i$

and $I'_i = \bigcup_{j=n_{I_{i-1}+1}}^{n_{I_i}} J_j$ for all $1 \leq i \leq m$. Via this new decomposition,

we have that $T_{I_0, I'_0} = T_{\{\iota_0, \iota_1\}, J_0} \times T_{\{\iota_0, \iota_2\}, J_1} \times \dots \times T_{\{\iota_0, \iota_{n_{I_0}}, \iota_1\}, J_{n_{I_0}}}$, and $T_{I_k, I'_k} = T_{\{\iota_{k_0}, \iota_{k_1}\}, J_{n_{I_{k-1}+1}}} \times T_{\{\iota_{k_1}, \iota_{k_2}\}, J_{n_{I_{k-1}+2}}} \times \dots \times T_{\{\iota_{k, n_{I_k}}, \iota_{k+1}\}, J_{n_{I_k}}}$, for all $1 \leq k \leq m$, so $T_{J,K} = \times_{k=0}^m T_{I_k, I'_k}$. Consequently,

$$T_{I,K} = \times_{k=0}^m T_{\{\iota_k, \iota_{k+1}\}, I'_k} = \times_{k=0}^m T_{\{\iota_k, \iota_{k+1}\}, I_k} T_{I_k, I'_k} = T_{I,J} T_{J,K}$$

as required. \square

Definition 3.1. A convolution system $\mathcal{S} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{S}}$ is said to be convergent if for any two elements $s < t$ of \mathbb{S} , the projective system $(\{(\Omega_I, \mathcal{F}_I, \mu_I)\}, \{T_{I,J}\})$ over $(\mathcal{K}_{s,t}, \subseteq)$ is convergent.

We denote by $(\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b)$ the projective limit,

$$(\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b) = \varprojlim_{\mathcal{K}_{s,t}} \{(\Omega_I, \mathcal{F}_I, \mu_I), T_{I,J}\},$$

of the convergent projective system $\{(\Omega_I, \mathcal{F}_I, \mu_I), T_{I,J}\}$ over $(\mathcal{K}_{s,t}, \subseteq)$, and by $T_I^b : (\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b) \rightarrow (\Omega_I, \mathcal{F}_I, \mu_I)$ the canonical epimorphism (coordinate projection), for all $I \in \mathcal{K}_{s,t}$; they satisfy the compatibility relation $T_I^b = T_{I,J} T_J^b$, for all $I, J \in \mathcal{K}_{s,t}$, $I \subseteq J$.

The following result allows the construction of a CPPS from a convergent convolution system.

Theorem 3.2. *Let $\mathcal{S} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{S}}$ be a convergent convolution system. For any $r < s < t$ in \mathbb{S} , there exists an isomorphism of probability spaces*

$$T_{r,s,t}^b : (\Omega_{r,s}^b, \mathcal{F}_{r,s}^b, \mu_{r,s}^b) \times (\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b) \rightarrow (\Omega_{r,t}^b, \mathcal{F}_{r,t}^b, \mu_{r,t}^b)$$

that makes the system $\mathcal{S}^b = \{(\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b), T_{r,s,t}^b\}_{\mathbb{S}}$ into a CPPS.

Proof. Let $r, t \in \mathbb{S}$ with $r < t$. For any $s \in \mathbb{S}$ satisfying $r < s < t$, let $\mathcal{K}_{r,s,t}$ be the set of all partitions $I \in \mathcal{K}_{r,t}$ such that $s \in I$. Since $\mathcal{K}_{r,s,t}$ is a cofinal subset of $(\mathcal{K}_{r,t}, \subseteq)$, we have that $(\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b)$ is the projective limit of the convergent projective system $\{(\Omega_I, \mathcal{F}_I, \mu_I), T_{I,J}\}$ over $(\mathcal{K}_{r,s,t}, \subseteq)$. We freely identify the poset $(\mathcal{K}_{r,s,t}, \subseteq)$ with $\mathcal{K}_{r,s} \times \mathcal{K}_{s,t}$, considered with the product order, via the order isomorphism $\mathcal{K}_{r,s,t} \ni I = I_s \cup_s I \mapsto I_s \times_s I \in \mathcal{K}_{r,s} \times \mathcal{K}_{s,t}$. Given that $(\Omega_I, \mathcal{F}_I, \mu_I) = (\Omega_{I_s}, \mathcal{F}_{I_s}, \mu_{I_s}) \times (\Omega_{I_t}, \mathcal{F}_{I_t}, \mu_{I_t})$, for all $I \in \mathcal{K}_{r,s,t}$, and $T_{I,J} = T_{I_s, J_s} \times T_{I_t, J_t}$, for all $I \subseteq J$ in $\mathcal{K}_{r,t}$, we infer that the product probability space $(\Omega_{r,s}^b, \mathcal{F}_{r,s}^b, \mu_{r,s}^b) \times (\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b)$ is the least upper bound of the convergent projective system $\{(\Omega_I, \mathcal{F}_I, \mu_I), T_{I,J}\}$ over $(\mathcal{K}_{r,s,t}, \subseteq)$. Consequently, there exists an isomorphism of probability spaces $T_{r,s,t}^b : (\Omega_{r,s}^b, \mathcal{F}_{r,s}^b, \mu_{r,s}^b) \times (\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b) \rightarrow (\Omega_{r,t}^b, \mathcal{F}_{r,t}^b, \mu_{r,t}^b)$ such that

$$(7) \quad T_I^b T_{r,s,t}^b = T_{I_s}^b \times T_{I_t}^b$$

for all $I = I_s \cup_s I \in \mathcal{K}_{r,s,t}$.

It remains to show that the family $\{T_{r,s,t}^b\}$ thus constructed is associative. Let then $r < s < t < u$ be some elements of \mathbb{S} . Identifying, as above, the poset $(\mathcal{K}_{r,s,t,u}, \subseteq)$ of all partitions $I \in \mathcal{K}_{r,u}$ such that $s, t \in I$ with the product poset $\mathcal{K}_{r,s} \times \mathcal{K}_{s,t} \times \mathcal{K}_{t,u}$ via $I = I_s \cup_s I_t \cup_t I \mapsto I_s \times_s I_t \times_t I$, we have

$$(8) \quad \begin{aligned} T_I^b T_{r,s,u}^b (\text{id}_{\Omega_{r,s}^b} \times T_{s,t,u}^b) &= (T_{I_s}^b \times T_{s I_t \cup_t I}^b) (\text{id}_{\Omega_{r,s}^b} \times T_{s,t,u}^b) \\ &= T_{I_s}^b \times T_{s I_t}^b \times T_{I_t}^b. \end{aligned}$$

Similarly, $T_I^b T_{r,t,u}^b (T_{r,s,t}^b \times \text{id}_{\Omega_{t,u}^b}) = T_{I_s}^b \times T_{s I_t}^b \times T_{I_t}^b$, and the conclusion follows. \square

Definition 3.2. The system $\mathcal{S}^b = \{\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b, T_{r,s,t}^b\}_{\mathbb{S}}$ will be referred as the projective CPPS associate with \mathcal{S} .

If $T_{s,t}^b : (\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b) \rightarrow (\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b)$ is the canonical epimorphism corresponding to the trivial partition $I = \{s, t\}$, then the resulting family $\tau_{\mathcal{S}} = \{T_{s,t}^b\}_{s < t}$ is an epimorphism of convolution systems from \mathcal{S}^b to \mathcal{S} . The following result follows immediately from Theorem 3.2 and from the properties of the projective limit.

Corollary 3.2.1. *If $\theta : \mathcal{S}_1 \rightarrow \mathcal{S}_2$ is an isomorphism of convergent convolution systems, then there exists an isomorphism of CPPSs $\theta^b : \mathcal{S}_1^b \rightarrow \mathcal{S}_2^b$ such that $\theta \tau_{\mathcal{S}_1} = \tau_{\mathcal{S}_2} \theta^b$.*

We end this section by noticing that the product system of Hilbert spaces $L^2(\mathcal{S}^b)$ of the projective CPPS \mathcal{S}^b associated with a convergent convolution system $\mathcal{S} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{S}}$ can be described in terms of the subproduct system of Hilbert spaces $L^2(\mathcal{S})$ through an inductive limit construction. Indeed, for all $s < t$ in \mathbb{S} , let

$$H_{s,t} = \varinjlim_{I \in \mathcal{K}_{s,t}} L^2(\mu_I)$$

be the inductive limit of the inductive system of Hilbert spaces $(L^2(\mu_I), U_{T_{I,J}})$ over $(\mathcal{K}_{s,t}, \subseteq)$ with connecting isometries $V_I : L^2(\mu_I) \rightarrow H_{s,t}$, $I \in \mathcal{K}_{s,t}$ satisfying the compatibility relation $V_J U_{T_{I,J}} = V_I$ for all $I \subseteq J$ in $\mathcal{K}_{s,t}$. Arguing as in Theorem 3.2, one can find a unique unitary operator $U_{r,s,t} : H_{r,t} \rightarrow H_{r,s} \otimes H_{s,t}$ that satisfies the identity $U_{r,s,t} V_I = V_{I_s} \otimes V_{sI}$, for all $I = I_s \cup_s I \in \mathcal{K}_{r,s,t}$. The system

$$\mathcal{H} = \{H_{s,t}, U_{r,s,t}\}_{\mathbb{S}}$$

is a product system of Hilbert spaces. We then have:

Theorem 3.3. *If $\mathcal{S} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{S}}$ is a convergent convolution system, then the product systems of Hilbert spaces $\mathcal{H} = \{H_{s,t}, U_{r,s,t}\}_{\mathbb{S}}$ and $L^2(\mathcal{S}^b)$ are isomorphic.*

Proof. Because $U_{T_J} U_{T_{I,J}} = U_{T_I^b}$, for all $I \subseteq J$, $I, J \in \mathcal{K}_{s,t}$, and the set $\bigcup_{I \in \mathcal{K}_{s,t}} U_{T_I^b}(L^2(\mu_I))$ is everywhere dense in $L^2(\mu_{s,t}^b)$, it follows from the universal property of the inductive limit that there exists a unique unitary operator $\theta_{s,t} : H_{s,t} \rightarrow L^2(\mu_{s,t}^b)$ such that $\theta_{s,t} V_I = U_{T_I^b}$, for all $I \in \mathcal{K}_{s,t}$. Moreover, since

$$\begin{aligned} (\theta_{r,s} \otimes \theta_{s,t}) U_{r,s,t} V_I &= \theta_{r,s} V_{I_s} \otimes \theta_{s,t} V_{sI} = U_{T_{I_s}^b} \otimes U_{T_{sI}^b} \\ &= U_{T_{r,s,t}^b} U_{T_I^b} = U_{T_{r,s,t}^b} \theta_{r,t} V_I, \end{aligned}$$

for all $I = I_s \cup_s I \in \mathcal{K}_{r,s,t}$, and $r < s < t$ in \mathbb{S} , we deduce that $(\theta_{r,s} \otimes \theta_{s,t}) U_{r,s,t} = U_{T_{r,s,t}^b} \theta_{r,t}$, i.e., $\{\theta_{s,t}\}$ is an isomorphism of product systems. \square

4. K -CONVERGENT CONVOLUTION SYSTEMS AND ASSOCIATED FLOW SYSTEMS

Let $\mathcal{S} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{S}}$ be a convolution system. Consider two partitions $I, J \in \mathcal{K}$, $I \subseteq J$, and write $J = I_L \cup \tilde{I} \cup I_R$ as in (6). We set

$$X_{I,J} = \begin{cases} T_{I,J} & \text{if } I, J \in \mathcal{K}_{s,t}, \text{ for some } s < t \text{ in } \mathbb{S} \\ (T_{I,\tilde{I}}) \circ (\pi_{\tilde{I},J}) & \text{otherwise} \end{cases}$$

where $\pi_{\tilde{I},J} : \Omega_J \rightarrow \Omega_{\tilde{I}}$ is the coordinate projection. Then $X_{I,J} : (\Omega_J, \mathcal{F}_J, \mu_J) \rightarrow (\Omega_I, \mathcal{F}_I, \mu_I)$ is a morphism of probability spaces. Moreover:

Proposition 4.1. *The system $\{(\Omega_I, \mathcal{F}_I, \mu_I), X_{I,J}\}$ is a projective system of probability spaces over the directed set (\mathcal{K}, \subseteq) .*

Proof. The compatibility relation $X_{I,J} X_{J,K} = X_{I,K}$, for all $I, J, K \in \mathcal{K}$ satisfying $I \subseteq J \subseteq K$, is verified using the same approach as in Proposition 3.1. For this, suppose $I \in \mathcal{K}_{q,r}$, $J \in \mathcal{K}_{s,t}$, and $K \in \mathcal{K}_{u,v}$ where $u < s < q < r < t < v$. Following (6), we decompose J with respect to I as $J = I_L \cup \tilde{I} \cup I_R$, K with respect to J as $K = J_L \cup \tilde{J} \cup J_R$, and K with respect to I as $K = I'_L \cup \tilde{I}' \cup I'_R$. Note that the partitions $I, \tilde{I}, \tilde{I}' \in \mathcal{K}_{q,r}$ satisfy $I \subseteq \tilde{I} \subseteq \tilde{I}'$, and the partitions $J, \tilde{J} \in \mathcal{K}_{s,t}$ satisfy $J \subseteq \tilde{J}$. It then

follows that $\tilde{I}' \subseteq \tilde{J}$, and decomposition (6) of \tilde{J} with respect to \tilde{I}' is given by $\tilde{J} = (\tilde{J} \cap [u, s]) \cup \tilde{I}' \cup (\tilde{J} \cap [t, v])$. Using this, we have

$$\begin{aligned} \pi_{\tilde{I}, J} \circ (T_{J, \tilde{J}}) &= \pi_{\tilde{I}, J} \circ (T_{J \cap [u, s], \tilde{J} \cap [u, s]} \times T_{J \cap [s, t], \tilde{J} \cap [s, t]} \times T_{J \cap [t, v], \tilde{J} \cap [t, v]}) \\ &= \pi_{\tilde{I}, J} \circ (T_{I_L, \tilde{J} \cap [u, s]} \times T_{\tilde{I}, \tilde{I}'} \times T_{I_R, \tilde{J} \cap [t, v]}) \\ &= (T_{\tilde{I}, \tilde{I}'}) \circ \pi_{\tilde{I}', \tilde{J}}. \end{aligned}$$

Therefore

$$\begin{aligned} X_{I, J} X_{J, K} &= (T_{I, \tilde{I}}) \circ (\pi_{\tilde{I}, J}) \circ (T_{J, \tilde{J}}) \circ (\pi_{\tilde{J}, K}) = (T_{I, \tilde{I}}) \circ (T_{\tilde{I}, \tilde{I}'}) \circ (\pi_{\tilde{I}', \tilde{J}}) \circ (\pi_{\tilde{J}, K}) \\ &= X_{I, K}, \end{aligned}$$

as needed. \square

Definition 4.1. A convolution system $\mathcal{S} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{S}}$ is said to be K -convergent if the projective system $\{(\Omega_I, \mathcal{F}_I, \mu_I), X_{I,J}\}$ over (\mathcal{K}, \subseteq) is convergent.

If $\mathcal{S} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{S}}$ is K -convergent, then we denote by $(\Omega^b, \mathcal{F}^b, P^b)$ the projective limit of this projective system,

$$(\Omega^b, \mathcal{F}^b, P^b) = \varprojlim_{\mathcal{K}} \{(\Omega_I, \mathcal{F}_I, \mu_I), X_{I,J}\},$$

and by $X_I : (\Omega^b, \mathcal{F}^b, P^b) \rightarrow (\Omega_I, \mathcal{F}_I, \mu_I)$ the canonical epimorphism; it satisfies the compatibility relation $X_I = X_{I,J} X_J$, for all $I, J \in \mathcal{K}$, $I \subseteq J$. For a trivial partition $I = \{s < t\}$, we simply write $X_{s,t}$ instead of $X_{\{s,t\}}$. Using the properties of the projective limit, one can readily see that

$$((\Omega^b, \mathcal{F}^b, P^b), \{X_{s,t}\})_{\mathcal{S}}$$

is a flow system over \mathcal{S} . In addition, we have the following.

Proposition 4.2. *Any K -convergent convolution system is convergent.*

Proof. Let $\mathcal{S} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{S}}$ be a K -convergent convolution system, and $s < t$ be two fixed elements in \mathbb{S} . It is clear that the **Set**-projective system $\{\Omega_I, T_{I,J}\}$ over $(\mathcal{K}_{s,t}, \subseteq)$ is simply maximal. Furthermore, if $\pi_{s,t} : \Omega^b \rightarrow \Omega_{s,t}^b$ is the coordinate projection onto Ω^b , then $T_I^b \pi_{s,t} = X_I$, for all $I \in \mathcal{K}_{s,t}$, so $(\Omega^b, \mathcal{F}^b, P^b)$ is an upper bound of $\{(\Omega_I, \mathcal{F}_I, \mu_I), T_{I,J}\}$. Therefore the the projective system $\{(\Omega_I, \mathcal{F}_I, \mu_I), T_{I,J}\}$ over $(\mathcal{K}_{s,t}, \subseteq)$ must be convergent. \square

Next, we focus on finding a necessary and sufficient condition for a convergent convolution system to be K -convergent. For this purpose, we start with a convergent convolution system $\mathcal{S} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{S}}$, and let $\mathcal{S}^b = \{\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b, T_{r,s,t}^b\}_{\mathbb{S}}$ be the projective CPPS associate with \mathcal{S} . For any elements $u < s < t < v$ in \mathbb{S} , let $T_{(s,t),(u,v)}^b : \Omega_{u,v}^b \rightarrow \Omega_{s,t}^b$ be the mapping

$$T_{(s,t),(u,v)}^b = \pi_{(s,t)} \circ (\text{id}_{\Omega_{u,s}^b} \times T_{s,t,v}^b)^{-1} \circ (T_{u,s,v}^b)^{-1},$$

where $\pi_{(s,t)} : \Omega_{u,s}^b \times \Omega_{s,t}^b \times \Omega_{t,v}^b \rightarrow \Omega_{s,t}^b$ is the coordinate projection onto $\Omega_{s,t}^b$. If $u = s$, and/or $t = v$, then $T_{(s,t),(u,v)}^b$ is defined accordingly. It is clear that $T_{(s,t),(u,v)}^b : (\Omega_{u,v}^b, \mathcal{F}_{u,v}^b, \mu_{u,v}^b) \rightarrow (\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b)$ is an epimorphism of probability spaces. Furthermore, we have:

Lemma 4.3. *For any elements $u < s < t < v$ in \mathbb{S} and partitions $I \in \mathcal{K}_{s,t}$ and $J \in \mathcal{K}_{u,v}$ we have $T_I^b T_{(s,t),(u,v)}^b = X_{I,J} T_J^b$.*

Proof. By decomposing $J = I_L \cup \tilde{I} \cup I_R$ as in (6), we have

$$\begin{aligned}
T_I^b T_{(s,t),(u,v)}^b &= T_I^b \pi_{(s,t)} (\text{id}_{\Omega_{u,s}^b} \times T_{s,t,v}^b)^{-1} (T_{u,s,v}^b)^{-1} \\
&= T_{I,\tilde{I}}^b T_{\tilde{I}}^b \pi_{(s,t)} (\text{id}_{\Omega_{u,s}^b} \times T_{s,t,v}^b)^{-1} (T_{u,s,v}^b)^{-1} \\
&= T_{I,\tilde{I}}^b \pi_{\tilde{I},J} (T_{I_R}^b \times T_{\tilde{I}}^b \times T_{I_L}^b) (\text{id}_{\Omega_{u,s}^b} \times T_{s,t,v}^b)^{-1} (T_{u,s,v}^b)^{-1} \\
&= X_{I,J} (T_{I_R}^b \times T_{\tilde{I}}^b \times T_{I_L}^b) (\text{id}_{\Omega_{u,s}^b} \times T_{s,t,v}^b)^{-1} (T_{u,s,v}^b)^{-1} \\
&\stackrel{(8)}{=} X_{I,J} T_J^b,
\end{aligned}$$

as needed. \square

Proposition 4.4. *Let $\mathcal{S} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{S}}$ be a convergent convolution system. The system $(\{(\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b)\}, T_{(s,t),(u,v)})$ is a projective system of probability spaces over the partially ordered set of ordered pairs $\mathcal{C} = \{(s,t) \mid s, t \in \mathbb{S}, s \leq t\}$ with partial order $(s,t) \subseteq (u,v)$ if $u \leq s \leq t \leq v$.*

Proof. The compatibility relation $T_{(q,r),(s,t)}^b T_{(s,t),(u,v)}^b = T_{(q,r),(u,v)}^b$, for all $(q,r) \subseteq (s,t) \subseteq (u,v)$, can be checked by using the laws of associativity. The process is, however, lengthy and convoluted. We will highlight the key intermediate steps and leave the detailed completion to the reader.

$$\begin{aligned}
T_{(q,r),(s,t)}^b T_{(s,t),(u,v)}^b &= \pi_{(q,r)} \circ (T_{s,q,r}^b \times \text{id}_{\Omega_{r,t}^b})^{-1} \circ (T_{s,r,t}^b)^{-1} \circ \\
&\quad \pi_{(s,t)} \circ (\text{id}_{\Omega_{u,s}^b} \times T_{s,t,v}^b)^{-1} \circ (T_{u,s,v}^b)^{-1} \\
&= \pi_{(q,r)} \circ (T_{s,q,r}^b \times \text{id}_{\Omega_{r,t}^b})^{-1} \circ (T_{s,r,t}^b)^{-1} \circ \\
&\quad T_{s,r,t}^b \circ \pi_{(s,r) \times (r,t)} \circ (\text{id}_{\Omega_{u,s}^b} \times \text{id}_{\Omega_{s,r}^b} \times T_{r,t,v}^b)^{-1} \circ \\
&\quad (\text{id}_{\Omega_{u,s}^b} \times T_{s,r,v}^b)^{-1} \circ (T_{u,s,v}^b)^{-1} \\
&= \pi_{(q,r)} \circ (\text{id}_{\Omega_{u,s}^b} \times \text{id}_{\Omega_{s,q}^b} \times \text{id}_{\Omega_{q,r}^b} \times T_{r,t,v}^b)^{-1} \circ \\
&\quad (\text{id}_{\Omega_{u,s}^b} \times T_{s,q,r}^b \times \text{id}_{\Omega_{r,v}^b})^{-1} \circ (\text{id}_{\Omega_{u,s}^b} \times T_{s,r,v}^b)^{-1} \circ (T_{u,s,v}^b)^{-1} \\
&= \pi_{(q,r)} \circ (T_{u,s,q}^b \times \text{id}_{\Omega_{q,r}^b} \times \text{id}_{\Omega_{r,v}^b}) \circ (\text{id}_{\Omega_{u,s}^b} \times T_{s,q,r}^b \times \text{id}_{\Omega_{r,v}^b})^{-1} \circ \\
&\quad (\text{id}_{\Omega_{u,s}^b} \times T_{s,r,v}^b)^{-1} \circ (T_{u,s,v}^b)^{-1}
\end{aligned}$$

where $\pi_{(s,r) \times (r,t)} : \Omega_{u,s}^b \times \Omega_{s,r}^b \times \Omega_{r,t}^b \times \Omega_{t,v}^b \rightarrow \Omega_{s,r}^b \times \Omega_{r,t}^b$ is the coordinate projection. Noticing that $(\text{id}_{\Omega_{u,q}^b} \times \circ T_{q,r,v}^b)^{-1} \circ T_{u,q,v}^{-b} = (T_{u,s,q}^b \times$

$\text{id}_{\Omega_{q,r}^b} \times \text{id}_{\Omega_{r,v}^b}) \circ (\text{id}_{\Omega_{u,s}^b} \times T_{s,q,r}^b \times \text{id}_{\Omega_{r,v}^b})^{-1} \circ (\text{id}_{\Omega_{u,s}^b} \times T_{s,r,v}^b)^{-1} \circ T_{u,s,v}^{-b}$, it follows that $T_{(q,r),[s,t]}^b T_{(s,t),(u,v)}^b = T_{(q,r),(u,v)}^b$, as required. \square

We conclude with the following general description of the K -convergence of a convergent convolution system.

Theorem 4.5. *Let $\mathcal{S} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{S}}$ be a convergent convolution system such that the **Set**-projective system $\{\Omega_I, X_{I,J}\}$ over \mathcal{K} is simply maximal. The following conditions are equivalent:*

- (1) *The convolution system \mathcal{S} is K -convergent;*
- (2) *The projective system $\{(\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b), T_{(s,t),(u,v)}\}$ is convergent.*

Proof. (1) \Rightarrow (2). Suppose that \mathcal{S} is K -convergent. Because $T_{I,J}X_J = X_I$, for any two elements $s < t$ in \mathbb{S} and any two partitions $I, J \in \mathcal{K}_{s,t}$, $I \subseteq J$, it follows that there exists an epimorphism $X_{s,t}^b : (\Omega_{s,t}^b, \mathcal{F}^b, P^b) \rightarrow (\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b)$ such that

$$T_I^b X_{s,t}^b = X_I,$$

for all $I \in \mathcal{K}_{s,t}$ and all $s < t$ in \mathbb{S} . Using Lemma 4.3, one can readily see that $T_{(s,t),(u,v)}^b X_{u,v}^b = X_{s,t}^b$, for all $u < s < t < v$ in \mathbb{S} . Indeed,

$$T_I^b T_{(s,t),(u,v)}^b X_{u,v}^b = X_{I,J} T_J^b X_{u,v}^b = X_{I,J} X_J = X_I,$$

for all $I \in \mathcal{K}_{s,t}$, $J \in \mathcal{K}_{u,v}$, $I \subseteq J$, and the conclusion follows. Consequently $(\Omega^b, \mathcal{F}^b, P^b)$ is an upper limit for the projective system $\{(\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b), T_{(s,t),(u,v)}\}$, so this must be convergent.

(2) \Rightarrow (1). Suppose that the projective system $\{(\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b), T_{(s,t),(u,v)}\}$ is convergent, and let

$$(9) \quad ({}^b\Omega, {}^b\mathcal{F}, {}^bP) = \varprojlim_{\mathcal{C}} \{(\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b), T_{(s,t),(u,v)}\}$$

be its projective limit with canonical epimorphisms $T'_{s,t} : ({}^b\Omega, {}^b\mathcal{F}, {}^bP) \rightarrow (\Omega_{s,t}^b, \mathcal{F}_{s,t}^b, \mu_{s,t}^b)$, for all $s < t$ in \mathbb{S} . For any $I \in \mathcal{K}$, say $I \in \mathcal{K}_{s,t}$, for some $s < t$ in \mathbb{S} , consider the composite epimorphism $T_I^b T'_{s,t} : ({}^b\Omega, {}^b\mathcal{F}, {}^bP) \rightarrow (\Omega_I, \mathcal{F}_I, \mu_I)$. We deduce from Lemma 4.3 that $T_I^b T'_{s,t} = X_{I,J} T_J^b T'_{u,v}$ for all $I \subseteq J$, $I \in \mathcal{K}_{s,t}$, $J \in \mathcal{K}_{u,v}$, $u \leq s < t \leq v$. Thus $({}^b\Omega, {}^b\mathcal{F}, {}^bP)$ is an upper bound for $\{(\Omega_I, \mathcal{F}_I, \mu_I), X_{I,J}\}$ over (\mathcal{K}, \subseteq) , which leads to the convergence of the projective system $\{(\Omega_I, \mathcal{F}_I, \mu_I), X_{I,J}\}$. \square

It is worth mentioning that if $\mathcal{S} = \{\Omega_{s,t}, \mathcal{F}_{s,t}, \mu_{s,t}, T_{r,s,t}\}_{\mathbb{S}}$ is a K -convergent convolution system, then the probability space $({}^b\Omega, {}^b\mathcal{F}, {}^bP)$ defined in (9) is the least upper bound for the projective system $\{(\Omega_I, \mathcal{F}_I, \mu_I), X_{I,J}\}$ over (\mathcal{K}, \subseteq) . We postpone the proof of this fact, as well as the discussion of some relevant examples, for another occasion.

REFERENCES

- [1] William Arveson, *Continuous analogues of Fock space*, Mem. Amer. Math. Soc. **80** (1989), no. 409, iv+66.
- [2] ———, *Noncommutative dynamics and E-semigroups*, Springer Monographs in Mathematics, Springer-Verlag, New York, 2003.
- [3] Salomon Bochner, *Harmonic analysis and the theory of probability*, University of California Press, Berkeley-Los Angeles, Calif., 1955.
- [4] J. R. Choksi, *Inverse limits of measure spaces*, Proc. London Math. Soc. (3) **8** (1958), 321–342.
- [5] Remus Floricel and Brian Ketelboeter, *C*-subproduct and product systems*, J. Math. Anal. Appl. **528** (2023), no. 1, Paper No. 127485, 31.
- [6] Zdeněk Frolík, *Projective limits of measure spaces*, Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability (Univ. California, Berkeley, Calif., 1970/1971), Univ. California Press, Berkeley, CA, 1972, pp. 67–80.
- [7] Asgar Jamneshan and Terence Tao, *Foundational aspects of uncountable measure theory: Gelfand duality, Riesz representation, canonical models, and canonical disintegration*, arXiv:2010.00681v3.
- [8] Göran Högnäs and Arunava Mukherjea, *Probability measures on semigroups*, 2nd ed., Probability and its Applications (New York), Springer, New York, 2011. Convolution products, random walks, and random matrices.
- [9] Leon Henkin, *A problem on inverse mapping systems*, Proc. Amer. Math. Soc. **1** (1950), 224–225.
- [10] M. M. Rao, *Projective limits of probability spaces*, J. Multivariate Anal. **1** (1971), no. 1, 28–57.
- [11] B. S. Tsirelson and A. M. Vershik, *Examples of nonlinear continuous tensor products of measure spaces and non-Fock factorizations*, Rev. Math. Phys. **10** (1998), no. 1, 81–145.
- [12] Boris Tsirelson, *Non-isomorphic product systems*, Advances in quantum dynamics (South Hadley, MA, 2002), Contemp. Math., vol. 335, Amer. Math. Soc., Providence, RI, 2003, pp. 273–328.
- [13] ———, *Nonclassical stochastic flows and continuous products*, Probab. Surv. **1** (2004), 173–298.
- [14] William C. Waterhouse, *An empty inverse limit*, Proc. Amer. Math. Soc. **36** (1972), 618.

UNIVERSITY OF REGINA, DEPARTMENT OF MATHEMATICS, REGINA, SK, CANADA
Email address: Remus.Floricel@uregina.ca

UNIVERSITY OF REGINA, DEPARTMENT OF MATHEMATICS, REGINA, SK, CANADA
Email address: pdm456@uregina.ca