

Integral representations for the joint survival functions of the cumulated components of multinomial random vectors

Frédéric Ouimet^a

^a*Department of Mathematics and Statistics, McGill University, Montréal (Québec) Canada H3A 0B9*

Abstract

This paper presents a multivariate normal integral expression for the joint survival function of the cumulated components of any multinomial random vector. This result can be viewed as a multivariate analog of Equation (7) from Carter and Pollard [2], who improved Tusnády's inequality. Our findings are based on a crucial relationship between the joint survival function of the cumulated components of any multinomial random vector and the joint cumulative distribution function of a corresponding Dirichlet distribution. We offer two distinct proofs: the first expands the logarithm of the Dirichlet density, while the second employs Laplace's method applied to the Dirichlet integral.

Keywords: Dirichlet distribution, Gaussian approximation, Laplace's method, multinomial distribution, multivariate normal distribution, quantile coupling

2020 MSC: Primary: 62E17 Secondary: 62H10, 62H12, 62E20

1. Introduction

For any integer $d \in \mathbb{N}$, the d -dimensional simplex and its interior are defined by

$$\mathcal{S}_d = \{\mathbf{s} \in [0, 1]^d : \|\mathbf{s}\|_1 \leq 1\}, \quad \text{Int}(\mathcal{S}_d) = \{\mathbf{s} \in (0, 1)^d : \|\mathbf{s}\|_1 < 1\},$$

where $\|\mathbf{s}\|_1 = \sum_{i=1}^d |s_i|$ denotes the ℓ_1 -norm in \mathbb{R}^d . Given a set of probability weights $\mathbf{p} \in \text{Int}(\mathcal{S}_d)$, the Multinomial(n, \mathbf{p}) probability mass function is defined, for all $\mathbf{k} \in \mathbb{N}_0^d \cap n\mathcal{S}_d$, by

$$p_n(\mathbf{k}) = \frac{n!}{(n - \|\mathbf{k}\|_1)! \prod_{i=1}^d k_i!} p_{d+1}^{n - \|\mathbf{k}\|_1} \prod_{i=1}^d p_i^{k_i},$$

where $p_{d+1} = 1 - \|\mathbf{p}\|_1 \in (0, 1)$ and $n \in \mathbb{N}$. The covariance matrix of the multinomial distribution is well-known to be $n \Sigma_p$, where $\Sigma_p = \text{diag}(\mathbf{p}) - \mathbf{p}\mathbf{p}^\top$; see, e.g., [7, p. 377]. From Theorem 1 of Tanabe and Sagae [9], it is also well known that $\det(\Sigma_p) = p_1 \dots p_d p_{d+1}$. The centered multivariate normal density with the same covariance structure as the above multinomial distribution is defined, for all $\mathbf{x} \in \mathbb{R}^d$, by

$$\phi_{\Sigma_p}(\mathbf{x}) = \frac{\exp(-\mathbf{x}^\top \Sigma_p^{-1} \mathbf{x} / 2)}{\sqrt{(2\pi)^d \det(\Sigma_p)}}.$$

Email address: frederic.ouimet2@mcgill.ca (Frédéric Ouimet)

Let $[d] = \{1, \dots, d\}$. Let $I_1 = (0, p_1]$ and $I_j = (p_1 + \dots + p_{j-1}, p_1 + \dots + p_j]$ for all $j \in [d] \setminus \{1\}$. If $U_1, \dots, U_n \stackrel{\text{iid}}{\sim} \text{Uniform}(0, 1)$, then

$$\mathbf{X} = (X_1, \dots, X_d) \equiv \left(\sum_{i=1}^n \mathbb{1}_{\{U_i \in I_1\}}, \dots, \sum_{i=1}^n \mathbb{1}_{\{U_i \in I_d\}} \right) \sim \text{Multinomial}(n, \mathbf{p}).$$

If $U_{(1)} \leq \dots \leq U_{(n)}$ denote the order statistics, then

$$\begin{aligned} & \mathbb{P}(X_1 + \dots + X_i \geq k_1 + \dots + k_i, \forall i \in [d]) \\ &= \mathbb{P}(U_{(k_1 + \dots + k_i)} \leq p_1 + \dots + p_i, \forall i \in [d]) \\ &= \int_0^{p_1} \int_{u_1}^{p_1 + p_2} \dots \int_{u_{d-1}}^{p_1 + \dots + p_d} n! \prod_{i=1}^{d+1} \frac{(u_i - u_{i-1})^{k_i - k_{i-1} - 1}}{(k_i - k_{i-1} - 1)!} du_1 du_2 \dots du_d, \end{aligned}$$

where one defines $u_0 = 0$, $u_{d+1} = 1$, $k_0 = 0$, $k_{d+1} = n + 1$ in the multidimensional integral. After applying the change of variables $s_i = u_i - u_{i-1}$ and $j_i = k_i - k_{i-1}$, for all $i \in [d]$, together with $j_{d+1} = k_{d+1} - k_d = (n + 1) - \sum_{i=1}^d j_i$, the above can be rewritten as

$$\begin{aligned} & \mathbb{P}(X_1 + \dots + X_i \geq k_1 + \dots + k_i, \forall i \in [d]) \\ &= \int_0^{p_1} \int_0^{(p_1 - s_1) + p_2} \dots \int_0^{\sum_{k=1}^{d-1} (p_k - s_k) + p_d} n! \prod_{i=1}^{d+1} \frac{s_i^{j_i - 1}}{(j_i - 1)!} ds. \end{aligned}$$

This neat equation provides a direct relationship between the joint survival function of the cumulated components of any multinomial random vector and the joint cumulative distribution function of a corresponding Dirichlet distribution.

In turn, letting $N = n - d$, $J_i = j_i - 1$ for all $i \in [d + 1]$, and defining the region

$$\mathcal{R}_d = \left\{ \mathbf{s} \in \mathcal{S}_d : (s_1, s_2, \dots, s_i) \in \left(\sum_{k=1}^i p_k \right) \mathcal{S}_i \forall i \in [d] \right\}, \quad (1)$$

one can write

$$\mathbb{P}(X_1 + \dots + X_i \geq k_1 + \dots + k_i, \forall i \in [d]) = \int_{\mathcal{R}_d} \frac{(N + d)!}{N!} \times \frac{N!}{\prod_{i=1}^{d+1} J_i!} \prod_{i=1}^{d+1} s_i^{J_i} d\mathbf{s}. \quad (2)$$

2. Quantile coupling

For any integer $m \in \mathbb{N}$, let λ_m denote the error term in Stirling's approximation for $\ln(m!)$. It is well known that, for all $m \in \mathbb{N}$,

$$\ln(m!) = \frac{1}{2} \ln(2\pi m) + m \ln m - m + \lambda_m, \quad (3)$$

where $(12m + 1)^{-1} \leq \lambda_m \leq (12m)^{-1}$; see, e.g., Gronwall [3].

Together with the notation introduced in Section 1, define, for all $i \in [d + 1]$,

$$\varepsilon_i = \frac{J_i/N - p_i}{p_i}, \quad \tilde{\varepsilon}_i = p_i \varepsilon_i = J_i/N - p_i,$$

and also

$$\Lambda_N = \lambda_N - \sum_{i=1}^{d+1} \lambda_i, \quad \Delta_N = \ln \left\{ \frac{(N+d)!}{N!N^d} \right\} + \Lambda_N - \frac{1}{2} \sum_{i=1}^{d+1} \ln(1 + \varepsilon_i) - N\tilde{\gamma}(\boldsymbol{\varepsilon}), \quad (4)$$

$$\tilde{\gamma}(\boldsymbol{\varepsilon}) = \sum_{i=1}^{d+1} p_i(1 + \varepsilon_i) \ln(1 + \varepsilon_i) - \frac{1}{2} \tilde{\boldsymbol{\varepsilon}}^\top \Sigma_p^{-1} \tilde{\boldsymbol{\varepsilon}}, \quad (5)$$

$$\gamma^*(\mathbf{s}) = \sum_{i=1}^{d+1} p_i(1 + \varepsilon_i) \ln \left(\frac{s_i}{p_i} \right) - \left\{ \tilde{\boldsymbol{\varepsilon}}^\top \Sigma_p^{-1} (\mathbf{s} - \mathbf{p}) - \frac{1}{2} (\mathbf{s} - \mathbf{p})^\top \Sigma_p^{-1} (\mathbf{s} - \mathbf{p}) \right\}. \quad (6)$$

Theorem 1 below expresses the joint survival function of the cumulated components of any multinomial random vector in terms of a multivariate normal integral. The result can be seen as a multivariate analog of Eq. (7) of Carter and Pollard [2], who improved on Tusnády's inequality. For an overview of the literature on Tusnády's inequality and the most recent improvement in the bulk, see Massart [4] and Ouimet [6], respectively.

Theorem 1. *For all $\mathbf{k} \in \mathbb{N}_0^d \cap n\mathcal{S}_d$, one has*

$$\mathbb{P}(X_1 + \dots + X_i \geq k_1 + \dots + k_i, \forall i \in [d]) = e^{\Lambda_N} \int_{\mathcal{R}_d} \exp\{N\gamma^*(\mathbf{s})\} N^{d/2} \phi_{\Sigma_p}\{N^{1/2}(\mathbf{p} - \mathbf{s} + \tilde{\boldsymbol{\varepsilon}})\} d\mathbf{s}.$$

Useful alternative expressions for $\tilde{\gamma}(\boldsymbol{\varepsilon})$ and $\gamma^(\mathbf{s})$ can be found in (10) and (11), respectively.*

Remark 1. *Theorem 1 also complements the local limit theorem for the multinomial distribution developed independently by Siotani and Fujikoshi [8] and Ouimet [5], and the related non-asymptotic approximations for Pearson's chi square statistic by Bax and Ouimet [1].*

3. Proofs

First proof of Theorem 1. Note that $\sum_{i=1}^{d+1} J_i = N$. By taking the logarithm of the integrand in (2) and applying Stirling's formula (3), one obtains

$$\begin{aligned} & \ln \left\{ \frac{(N+d)!}{N!} \times \frac{N!}{\prod_{i=1}^{d+1} J_i!} \prod_{i=1}^{d+1} s_i^{J_i} \right\} \\ &= \ln \left\{ \frac{(N+d)!}{N!} \right\} + \ln(N!) - \sum_{i=1}^{d+1} \ln(J_i!) + \sum_{i=1}^{d+1} J_i \ln s_i \\ &= \ln \left\{ \frac{(N+d)!}{N!N^d} \right\} - \frac{1}{2} \ln \left\{ \left(\frac{2\pi}{N} \right)^d \prod_{i=1}^{d+1} (J_i/N) \right\} + \sum_{i=1}^{d+1} J_i \ln \left(\frac{s_i}{J_i/N} \right) + \lambda_N - \sum_{i=1}^{d+1} \lambda_i \quad (7) \\ &= \ln \left\{ \frac{(N+d)!}{N!N^d} \right\} - \frac{1}{2} \sum_{i=1}^{d+1} \ln(1 + \varepsilon_i) - \frac{1}{2} \ln \left\{ \left(\frac{2\pi}{N} \right)^d \prod_{i=1}^{d+1} p_i \right\} \\ & \quad + N \sum_{i=1}^{d+1} p_i(1 + \varepsilon_i) \ln \left(\frac{s_i}{p_i} \right) + N \sum_{i=1}^{d+1} (J_i/N) \ln \left(\frac{p_i}{J_i/N} \right) + \Lambda_N. \end{aligned}$$

Using the Taylor expansion

$$(1+x)\ln(1+x) = x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{12} + \mathcal{O}_\eta(x^5), \quad |x| \leq \eta < 1,$$

and the fact that $\tilde{\varepsilon}_{d+1} = -\sum_{i=1}^d \tilde{\varepsilon}_i$, one can write

$$\sum_{i=1}^{d+1} (J_i/N) \ln\left(\frac{p_i}{J_i/N}\right) = -\frac{1}{2} \tilde{\boldsymbol{\varepsilon}}^\top \Sigma_p^{-1} \tilde{\boldsymbol{\varepsilon}} - \tilde{\gamma}(\boldsymbol{\varepsilon}), \quad (8)$$

where the quantity $\tilde{\gamma}(\boldsymbol{\varepsilon})$, as defined in (5), can be expanded as follows:

$$\begin{aligned} \tilde{\gamma}(\boldsymbol{\varepsilon}) &\equiv \sum_{i=1}^{d+1} p_i(1+\varepsilon_i)\ln(1+\varepsilon_i) - \frac{1}{2} \tilde{\boldsymbol{\varepsilon}}^\top \Sigma_p^{-1} \tilde{\boldsymbol{\varepsilon}} \\ &= \frac{1}{2} \sum_{i,j=1}^d \tilde{\varepsilon}_i \tilde{\varepsilon}_j \left\{ \frac{1}{p_i} \mathbb{1}(i=j) + \frac{1}{p_{d+1}} \right\} - \frac{1}{2} \tilde{\boldsymbol{\varepsilon}}^\top \Sigma_p^{-1} \tilde{\boldsymbol{\varepsilon}} \\ &\quad - \frac{1}{6} \sum_{i,j,k=1}^d \tilde{\varepsilon}_i \tilde{\varepsilon}_j \tilde{\varepsilon}_k \left\{ \frac{1}{p_i^2} \mathbb{1}(i=j=k) - \frac{1}{p_{d+1}^2} \right\} \\ &\quad + \frac{1}{12} \sum_{i,j,k,\ell=1}^d \tilde{\varepsilon}_i \tilde{\varepsilon}_j \tilde{\varepsilon}_k \tilde{\varepsilon}_\ell \left\{ \frac{1}{p_i^3} \mathbb{1}(i=j=k=\ell) + \frac{1}{p_{d+1}^3} \right\} + \mathcal{O}_{d,p}(\|\boldsymbol{\varepsilon}\|_1^5). \end{aligned}$$

Moreover, by [9, Eq. 21], it is known that $(\Sigma_p^{-1})_{ij} = p_i^{-1} \mathbb{1}_{(i=j)} + p_{d+1}^{-1}$ for all $i, j \in [d]$, so

$$\frac{1}{2} \sum_{i,j=1}^d \tilde{\varepsilon}_i \tilde{\varepsilon}_j \left\{ \frac{1}{p_i} \mathbb{1}(i=j) + \frac{1}{p_{d+1}} \right\} - \frac{1}{2} \tilde{\boldsymbol{\varepsilon}}^\top \Sigma_p^{-1} \tilde{\boldsymbol{\varepsilon}} = 0. \quad (9)$$

Therefore, using the last two equations, one can rewrite $\tilde{\gamma}(\boldsymbol{\varepsilon})$ as follows:

$$\begin{aligned} \tilde{\gamma}(\boldsymbol{\varepsilon}) &= -\frac{1}{6} \sum_{i,j,k=1}^d \tilde{\varepsilon}_i \tilde{\varepsilon}_j \tilde{\varepsilon}_k \left\{ \frac{1}{p_i^2} \mathbb{1}(i=j=k) - \frac{1}{p_{d+1}^2} \right\} \\ &\quad + \frac{1}{12} \sum_{i,j,k,\ell=1}^d \tilde{\varepsilon}_i \tilde{\varepsilon}_j \tilde{\varepsilon}_k \tilde{\varepsilon}_\ell \left\{ \frac{1}{p_i^3} \mathbb{1}(i=j=k=\ell) + \frac{1}{p_{d+1}^3} \right\} + \mathcal{O}_{d,p}(\|\boldsymbol{\varepsilon}\|_1^5). \end{aligned} \quad (10)$$

Now, consider $\boldsymbol{s} = (s_1, \dots, s_d) \in \mathcal{R}_d$ as defined in (1), and let

$$\delta_i(\boldsymbol{s}) = \frac{s_i - p_i}{p_i}, \quad i \in [d], \quad \delta_{d+1}(\boldsymbol{s}) = \frac{s_{d+1} - p_{d+1}}{p_{d+1}} = -\frac{\sum_{i=1}^d (s_i - p_i)}{p_{d+1}}.$$

In particular, note that this definition yields $\sum_{i=1}^{d+1} p_i \delta_i(s_i) = 0$. Upon applying a second-order version of the Taylor expansion

$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} + \mathcal{O}_\eta(x^4), \quad |x| \leq \eta < 1,$$

with Lagrange remainder, there exists a vector $\mathbf{s}^* = (s_1^*, \dots, s_d^*)$ on the line segment joining \mathbf{s} and $\mathbf{p} = (p_1, \dots, p_d)$ such that

$$\begin{aligned}
\sum_{i=1}^{d+1} p_i(1 + \varepsilon_i) \ln\left(\frac{s_i}{p_i}\right) &= \sum_{i=1}^{d+1} p_i(1 + \varepsilon_i) \ln\{1 + \delta_i(s_i)\} \\
&= \sum_{i=1}^{d+1} p_i(1 + \varepsilon_i) \delta_i(s_i) - \frac{1}{2} \sum_{i=1}^{d+1} p_i(1 + \varepsilon_i) \{\delta_i(s_i)\}^2 + \frac{1}{3} \sum_{i=1}^{d+1} p_i(1 + \varepsilon_i) \frac{\{\delta_i(s_i)\}^3}{\{1 + \delta_i(s_i^*)\}^3} \\
&= \sum_{i=1}^{d+1} \tilde{\varepsilon}_i \delta_i(s_i) - \frac{1}{2} \sum_{i=1}^{d+1} p_i \{\delta_i(s_i)\}^2 - \frac{1}{2} \sum_{i=1}^{d+1} \tilde{\varepsilon}_i \{\delta_i(s_i)\}^2 + \frac{1}{3} \sum_{i=1}^{d+1} p_i(1 + \varepsilon_i) \frac{\{\delta_i(s_i)\}^3}{\{1 + \delta_i(s_i^*)\}^3} \\
&= \tilde{\boldsymbol{\varepsilon}}^\top \Sigma_p^{-1}(\mathbf{s} - \mathbf{p}) - \frac{1}{2}(\mathbf{s} - \mathbf{p})^\top \Sigma_p^{-1}(\mathbf{s} - \mathbf{p}) - \frac{1}{2} \sum_{i=1}^{d+1} \tilde{\varepsilon}_i \{\delta_i(s_i)\}^2 + \frac{1}{3} \sum_{i=1}^{d+1} p_i(1 + \varepsilon_i) \frac{\{\delta_i(s_i)\}^3}{\{1 + \delta_i(s_i^*)\}^3},
\end{aligned}$$

where $\tilde{\boldsymbol{\varepsilon}} = (\tilde{\varepsilon}_1, \dots, \tilde{\varepsilon}_d)$. Using the last equation, the quantity $\gamma^*(\mathbf{s})$, as defined in (6), can be rewritten as

$$\begin{aligned}
\gamma^*(\mathbf{s}) &\equiv \sum_{i=1}^{d+1} p_i(1 + \varepsilon_i) \ln\left(\frac{s_i}{p_i}\right) - \left\{ \tilde{\boldsymbol{\varepsilon}}^\top \Sigma_p^{-1}(\mathbf{s} - \mathbf{p}) - \frac{1}{2}(\mathbf{s} - \mathbf{p})^\top \Sigma_p^{-1}(\mathbf{s} - \mathbf{p}) \right\} \\
&= -\frac{1}{2} \sum_{i=1}^{d+1} \tilde{\varepsilon}_i \{\delta_i(s_i)\}^2 + \frac{1}{3} \sum_{i=1}^{d+1} p_i(1 + \varepsilon_i) \frac{\{\delta_i(s_i)\}^3}{\{1 + \delta_i(s_i^*)\}^3}.
\end{aligned} \tag{11}$$

It readily follows that

$$\begin{aligned}
\sum_{i=1}^{d+1} p_i(1 + \varepsilon_i) \ln\left(\frac{s_i}{p_i}\right) &= \tilde{\boldsymbol{\varepsilon}}^\top \Sigma_p^{-1}(\mathbf{s} - \mathbf{p}) - \frac{1}{2}(\mathbf{s} - \mathbf{p})^\top \Sigma_p^{-1}(\mathbf{s} - \mathbf{p}) + \gamma^*(\mathbf{s}) \\
&= \frac{1}{2} \tilde{\boldsymbol{\varepsilon}}^\top \Sigma_p^{-1} \tilde{\boldsymbol{\varepsilon}} - \frac{1}{2}(\mathbf{s} - \mathbf{J}/N)^\top \Sigma_p^{-1}(\mathbf{s} - \mathbf{J}/N) + \gamma^*(\mathbf{s}).
\end{aligned} \tag{12}$$

Therefore, by putting (8) and (12) back into (7), and exponentiating, one gets

$$\begin{aligned}
\frac{(N+d)!}{N!} \times \frac{N!}{\prod_{i=1}^{d+1} J_i!} \prod_{i=1}^{d+1} s_i^{J_i} &= \exp\{\Delta_N + N\gamma^*(\mathbf{s})\} \frac{\exp\{-N(\mathbf{s} - \mathbf{J}/N)^\top \Sigma_p^{-1}(\mathbf{s} - \mathbf{J}/N)/2\}}{\sqrt{(2\pi/N)^d \det(\Sigma_p)}} \\
&= \exp\{\Delta_N + N\gamma^*(\mathbf{s})\} N^{d/2} \phi_{\Sigma_p}\{N^{1/2}(\mathbf{J}/N - \mathbf{s})\} \\
&= \exp\{\Delta_N + N\gamma^*(\mathbf{s})\} N^{d/2} \phi_{\Sigma_p}\{N^{1/2}(\mathbf{p} - \mathbf{s} + \tilde{\boldsymbol{\varepsilon}})\},
\end{aligned}$$

where recall $\Delta_N = \ln\{(N+d)!/(N!N^d)\} + \Lambda_N - (1/2) \sum_{i=1}^{d+1} \ln(1 + \varepsilon_i) - N\tilde{\gamma}(\boldsymbol{\varepsilon})$ from (4). Putting this last equation in (2) yields

$$\begin{aligned}
&\mathbb{P}(X_1 + \dots + X_i \geq k_1 + \dots + k_i, \forall i \in [d]) \\
&= e^{\Delta_N} \int_{\mathcal{R}_d} \exp[N\gamma^*(\mathbf{s})] N^{d/2} \phi_{\Sigma_p}\{N^{1/2}(\mathbf{p} - \mathbf{s} + \tilde{\boldsymbol{\varepsilon}})\} d\mathbf{s}.
\end{aligned}$$

This concludes the proof. \square

Second proof of Theorem 1. For all $s \in \mathcal{S}_d$, define

$$H(s) = \sum_{i=1}^{d+1} p_i(1 + \varepsilon_i) \ln s_i = \sum_{i=1}^{d+1} \frac{J_i}{N} \ln s_i.$$

For all $\mathbf{k} \in \mathbb{N}_0^d \cap n\mathcal{S}_d$, the integral representation of the joint survival function of the cumulated components of any multinomial random vector derived in (2) can be rewritten as

$$\mathbf{P}(X_1 + \cdots + X_i \geq k_1 + \cdots + k_i, \forall i \in [d]) = \frac{(N+d)!}{N!} \times \frac{N!}{\prod_{i=1}^{d+1} J_i!} \int_{\mathcal{R}_d} \exp\{NH(s)\} ds. \quad (13)$$

By Stirling's formula (3), note that

$$m! = \sqrt{2\pi m} \exp(m \ln m - m + \lambda_m),$$

where $(12m+1)^{-1} \leq \lambda_m \leq (12m)^{-1}$. Since $\sum_{i=1}^{d+1} J_i = N$ and $\Lambda_N = \lambda_N - \sum_{i=1}^{d+1} \lambda_i$, one deduces that

$$\frac{N!}{\prod_{i=1}^{d+1} J_i!} = \frac{\exp(\Lambda_N)}{\sqrt{(2\pi N)^d \prod_{i=1}^{d+1} \{p_i(1 + \varepsilon_i)\}}} \exp\{-NH(\mathbf{J}/N)\},$$

Therefore, one can rewrite (13) as

$$\begin{aligned} & \mathbf{P}(X_1 + \cdots + X_i \geq k_1 + \cdots + k_i, \forall i \in [d]) \\ &= \frac{(N+d)!}{N!N^d} \times \frac{N^{d/2} \exp(\Lambda_N)}{\sqrt{(2\pi)^d \prod_{i=1}^{d+1} \{p_i(1 + \varepsilon_i)\}}} \int_{\mathcal{R}_d} \exp[N\{H(s) - H(\mathbf{J}/N)\}] ds. \end{aligned} \quad (14)$$

Decompose the integral above as

$$\int_{\mathcal{R}_d} \exp[N\{H(s) - H(\mathbf{p})\} + N\{H(\mathbf{p}) - H(\mathbf{J}/N)\}] ds. \quad (15)$$

Using the quantity $\tilde{\gamma}(\boldsymbol{\varepsilon})$ as defined in (5), one has

$$H(\mathbf{p}) - H(\mathbf{J}/N) = - \sum_{i=1}^{d+1} p_i(1 + \varepsilon_i) \ln(1 + \varepsilon_i) = -\frac{1}{2} \tilde{\boldsymbol{\varepsilon}}^\top \Sigma_p^{-1} \tilde{\boldsymbol{\varepsilon}} - \tilde{\gamma}(\boldsymbol{\varepsilon}), \quad (16)$$

where recall $\tilde{\boldsymbol{\varepsilon}} = (p_1 \varepsilon_1, \dots, p_d \varepsilon_d)^\top$, $\tilde{\boldsymbol{\varepsilon}}_{d+1} = p_{d+1} \varepsilon_{d+1}$, and $\Sigma_p^{-1} = \{p_i^{-1} \mathbf{1}(i = j) + p_{d+1}^{-1} \mathbf{1}_{1 \leq i, j \leq d}\}$. Recall from (4) that

$$\Delta_N = \ln \left\{ \frac{(N+d)!}{N!N^d} \right\} + \Lambda_N - \frac{1}{2} \sum_{i=1}^{d+1} \ln(1 + \varepsilon_i) - N\tilde{\gamma}(\boldsymbol{\varepsilon}),$$

so one can rewrite (14) as

$$\begin{aligned} & \mathbf{P}(X_1 + \cdots + X_i \geq k_1 + \cdots + k_i, \forall i \in [d]) \\ &= \frac{N^{d/2} \exp(\Delta_N)}{\sqrt{(2\pi)^d \det(\Sigma_p)}} \int_{\mathcal{R}_d} \exp \left[N\{H(s) - H(\mathbf{p})\} - \frac{N}{2} \tilde{\boldsymbol{\varepsilon}}^\top \Sigma_p^{-1} \tilde{\boldsymbol{\varepsilon}} \right] ds. \end{aligned}$$

Using (11) and (12), one has

$$\begin{aligned}
H(\mathbf{s}) - H(\mathbf{p}) &= \sum_{i=1}^{d+1} \frac{J_i}{N} \ln(s_i) - \sum_{i=1}^{d+1} \frac{J_i}{N} \ln(p_i) = \sum_{i=1}^{d+1} p_i(1 + \varepsilon_i) \ln\left(\frac{s_i}{p_i}\right) \\
&= \tilde{\boldsymbol{\varepsilon}}^\top \Sigma_p^{-1}(\mathbf{s} - \mathbf{p}) - \frac{1}{2}(\mathbf{s} - \mathbf{p})^\top \Sigma_p^{-1}(\mathbf{s} - \mathbf{p}) \\
&\quad - \frac{1}{2} \sum_{i=1}^{d+1} \tilde{\varepsilon}_i \{\delta_i(s_i)\}^2 + \frac{1}{3} \sum_{i=1}^{d+1} p_i(1 + \varepsilon_i) \frac{\{\delta_i(s_i)\}^3}{\{1 + \delta_i(s_i^*)\}^3} \\
&= \tilde{\boldsymbol{\varepsilon}}^\top \Sigma_p^{-1}(\mathbf{s} - \mathbf{p}) - \frac{1}{2}(\mathbf{s} - \mathbf{p})^\top \Sigma_p^{-1}(\mathbf{s} - \mathbf{p}) + \gamma^*(\mathbf{s}) \\
&= \frac{1}{2} \tilde{\boldsymbol{\varepsilon}}^\top \Sigma_p^{-1} \tilde{\boldsymbol{\varepsilon}} - \frac{1}{2}(\mathbf{s} - \mathbf{J}/N)^\top \Sigma_p^{-1}(\mathbf{s} - \mathbf{J}/N) + \gamma^*(\mathbf{s}).
\end{aligned}$$

Putting this in the previous equation yields

$$\begin{aligned}
&\mathbb{P}(X_1 + \dots + X_i \geq k_1 + \dots + k_i, \forall i \in [d]) \\
&= e^{\Delta_N} \int_{\mathcal{R}_d} \exp\{N\gamma^*(\mathbf{s})\} N^{d/2} \phi_{\Sigma_p}\{N^{1/2}(\mathbf{J}/N - \mathbf{s})\} d\mathbf{s} \\
&= e^{\Delta_N} \int_{\mathcal{R}_d} \exp\{N\gamma^*(\mathbf{s})\} N^{d/2} \phi_{\Sigma_p}\{N^{1/2}(\mathbf{p} - \mathbf{s} + \tilde{\boldsymbol{\varepsilon}})\} d\mathbf{s}.
\end{aligned}$$

This concludes the proof. \square

Remark 2. The choice of decomposition in (15) can be explained as follows. The Hessian matrix of H at \mathbf{s} is equal to

$$\left(-\frac{J_i}{N s_i^2} \mathbb{1}(i = j) - \frac{J_{d+1}}{N s_{d+1}^2} \right)_{1 \leq i, j \leq d},$$

which is (symmetric) negative definite, so that H is strictly concave on $\text{Int}(\mathcal{S}_d)$. Given the strict concavity and the fact that $\bar{\mathbf{s}} = \mathbf{J}/N \in \text{Int}(\mathcal{S}_d)$ is the unique point satisfying

$$\nabla H(\bar{\mathbf{s}}) = \left(\frac{J_i}{N \bar{s}_i} - \frac{J_{d+1}}{N \bar{s}_{d+1}} \right)_{1 \leq i \leq d} = \mathbf{0}$$

in the interior of the simplex, the global maximum of H on $\text{Int}(\mathcal{S}_d)$ is achieved at \mathbf{J}/N . When the point $\mathbf{k} \in \mathbb{N}_0 \cap n\text{Int}(\mathcal{S}_d)$ is such that \mathbf{J}/N falls outside the range of integration \mathcal{R}_d defined in (1), then H maximizes at $\mathbf{s} = \mathbf{p}$ inside the region of integration \mathcal{R}_d . The decomposition is therefore equivalent to an application of Laplace's method provided some restrictions on \mathbf{k} are met.

Funding

F. Ouimet's current postdoctoral fellowship is funded through the Canada Research Chairs Program (Grant 950-231937 to Christian Genest) and the Natural Sciences and Engineering Research Council of Canada (Grant RGPIN-2024-04088 to Christian Genest).

References

- [1] E. Bax and F. Ouimet. Non-asymptotic approximations for Pearson’s chi-square statistic and its application to confidence intervals for strictly convex functions of the probability weights of discrete distributions. *Preprint*, page 20 pp., 2023. [arXiv:2309.01882v1](#).
- [2] A. Carter and D. Pollard. Tusnády’s inequality revisited. *Ann. Statist.*, 32(6):2731–2741, 2004. [MR2154001](#).
- [3] T. H. Gronwall. The gamma function in the integral calculus. *Ann. of Math. (2)*, 20(2): 35–124, 1918. [MR1502544](#).
- [4] P. Massart. Tusnady’s lemma, 24 years later. *Ann. Inst. H. Poincaré Probab. Statist.*, 38(6): 991–1007, 2002. [MR1955348](#).
- [5] F. Ouimet. A precise local limit theorem for the multinomial distribution and some applications. *J. Statist. Plann. Inference*, 215:218–233, 2021. [MR4249129](#).
- [6] F. Ouimet. An improvement of Tusnády’s inequality in the bulk. *Adv. in Appl. Math.*, 133: Paper No. 102270, 24, 2022. ISSN 0196-8858. [MR4340237](#).
- [7] T. A. Severini. *Elements of Distribution Theory*, volume 17 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge, 2005. ISBN 978-0-521-84472-7; 0-521-84472-X. [MR2168237](#).
- [8] M. Siotani and Y. Fujikoshi. Asymptotic approximations for the distributions of multinomial goodness-of-fit statistics. *Hiroshima Math. J.*, 14(1):115–124, 1984. [MR750392](#).
- [9] K. Tanabe and M. Sagae. An exact Cholesky decomposition and the generalized inverse of the variance-covariance matrix of the multinomial distribution, with applications. *J. Roy. Statist. Soc. Ser. B*, 54(1):211–219, 1992. [MR1157720](#).