

This is a preprint of a paper whose final and definite form is with *Appl. Math. (Warsaw)*, ISSN: 1233-7234. Article accepted for publication 11-June-2024.

## UNIFORM STABILITY OF DYNAMIC SICA HIV TRANSMISSION MODELS ON TIME SCALES

ZAHRA BELARBI, BENAOUMEUR BAYOUR, AND DELFIM F. M. TORRES

**ABSTRACT.** We consider a SICA model for HIV transmission on time scales. We prove permanence of solutions and we derive sufficient conditions for the existence and uniform asymptotic stability of a unique positive almost periodic solution of the system in terms of a Lyapunov function.

### 1. INTRODUCTION

In 2015, the deterministic SICA model was first presented as a subsystem of a TB-HIV/AIDS co-infection model [22]. One of the primary objectives of SICA models is to demonstrate how some of the fundamental relationships between epidemiological variables and the general pattern of the AIDS epidemic can be clarified using a straightforward mathematical model [18]. The celebrated SICA mathematical model [5, 6, 16, 25, 27] divides the total human population into four compartments, namely

- $S(t)$ : susceptible individuals at time  $t$ ;
- $I(t)$ : HIV-infected individuals with no clinical symptoms of AIDS but able to transmit HIV to other individuals at time  $t$ ;
- $C(t)$ : HIV-infected individuals under antiretroviral therapy (ART), the so called chronic stage with a viral load remaining low at time  $t$ ;
- $A(t)$ : HIV-infected individuals with AIDS clinical symptoms at time  $t$ .

Under some realistic assumptions, the dynamics of the disease proliferation in a community is then translated into a mathematical model given by the

---

2020 *Mathematics Subject Classification.* Primary 34N05, 39A24; Secondary 92D25.

*Key words and phrases.* SICA model for HIV transmission, time scales, permanence, almost periodic solution, uniform asymptotic stability.

following system of four ordinary differential equations [22, 23, 24]:

$$(1) \quad \begin{cases} \dot{S}(t) = \Lambda - \beta\lambda(t)S(t) - \nu S(t), \\ \dot{I}(t) = \beta\lambda(t)S(t) - (\rho + \phi + \nu)I(t) + \gamma A(t) + \omega C(t), \\ \dot{C}(t) = \phi I(t) - (\omega + \nu)C(t), \\ \dot{A}(t) = \rho I(t) - (\gamma + \nu + d)A(t), \end{cases}$$

where  $\Lambda, \beta, \nu, \rho, \phi, \gamma, \omega$  and  $d$  are real positive rates,

- $\Lambda$  is the rate of new susceptible;
- $\beta$  is the HIV transmission rate;
- $\nu$  is the natural death rate;
- $\rho$  is the default treatment rate for  $I$  individuals;
- $\phi$  is the HIV treatment rate for  $I$  individuals;
- $\gamma$  is the AIDS treatment rate;
- $\omega$  is the default treatment rate for  $C$  individuals;
- $d$  is the AIDS induced death rate;

and where the effective contact rate with people infected with HIV is given by

$$\lambda(t) = \frac{\beta}{N(t)}(I(t) + \eta_C C(t) + \eta_A A(t))$$

with

- $N(t)$  the total population at time  $t$ , that is,

$$N(t) = S(t) + I(t) + C(t) + A(t);$$

- $0 \leq \eta_C \leq 1$  the modification parameter;
- $\eta_A \geq 1$  the partial restoration parameter of immune function of individuals with HIV infection that use correctly the ART treatment.

The study of dynamical systems on time scales is now a very active area of research. The books of Bohner and Peterson [2, 3] offer a good introduction with applications to the time scale calculus along with some advanced topics. Applications of the time scale calculus can be found in many areas, including economics [1, 7, 26], ecology [19, 20, 30] and epidemics [4, 21, 11]. Here we generalize the SICA model (1) by considering dynamic equations on time scales and study it using the time-scale theory. By doing it, we unify the continuous and discrete-time models [27], generalizing it also to other contexts like the quantum [12, 17] or mixed/hybrid settings [8, 28].

Recently, Prasad and Khuddush proved the existence and uniform asymptotic stability of positive and almost periodic solutions for a 3-species Lotka–Volterra competitive system on time scales [20]. Moreover, they also studied the permanence and positive almost periodic solutions of a  $n$ -species

Lotka–Volterra system on time scales [19]. Motivated by these works, here we investigate the permanence and uniform asymptotic stability of the unique positive almost periodic solution of the following SICA model on time scales:

$$(2) \quad \begin{cases} x_1^\Delta(t) = \Lambda - \beta\lambda(t)x_1^\sigma(t) - \nu x_1^\sigma(t), \\ x_2^\Delta(t) = \beta\lambda(t)x_1(t) - (\rho + \phi + \nu)x_2^\sigma(t) + \gamma x_4(t) + \omega x_3(t), \\ x_3^\Delta(t) = \phi x_2(t) - (\omega + \nu)x_3^\sigma(t), \\ x_4^\Delta(t) = \rho x_2(t) - (\gamma + \nu + d)x_4^\sigma(t), \end{cases}$$

where  $t \in \mathbb{T}^+$ , with  $\mathbb{T}^+$  a nonempty closed subset of  $\mathbb{R}^+ = ]0, +\infty[$ .

Note that, in our notation,  $(x_1(t), x_2(t), x_3(t), x_4(t))$  is interpreted as  $(S(t), I(t), C(t), A(t))$ .

The paper is organized as follows. Section 2 gives a brief introduction to the time-scale theory and we also recall there the definition of an almost periodic function with respect to time, uniformly for the state variables, and the lemma of Lyapunov that gives a sufficient condition for the existence of a unique almost periodic solution that is uniformly asymptotically stable. In Section 3, we derive sufficient conditions for system (2) to be permanent. Finally, in Section 4, we establish sufficient conditions for the existence and uniform asymptotic stability of the unique positive almost periodic solution to the SICA system (2). Our results seem to be new even for standard time scales. We end with an example (Example 4.2) and a brief conclusion (Section 5).

## 2. SOME PRELIMINARIES

For more background on the theory of time scales than the one given here we recommend the books [2, 3]. A time scale  $\mathbb{T}$  is a nonempty closed subset of the real numbers  $\mathbb{R}$ . The time scale  $\mathbb{T}$  has the topology that it inherits from the real numbers with the standard topology. Let  $f$  be a function defined on  $\mathbb{T}^+$ . We set

$$f^L = \inf \{f(t) : t \in \mathbb{T}^+\} \quad \text{and} \quad f^U = \sup \{f(t) : t \in \mathbb{T}^+\}.$$

**Definition 2.1** (See [2]). The jump operators  $\sigma, \rho : \mathbb{T} \rightarrow \mathbb{T}$  and the graininess function  $\mu : \mathbb{T} \rightarrow \mathbb{R}^+$  are defined, respectively, by

$\sigma(t) = \inf \{\tau \in \mathbb{T} : \tau > t\}$ ,  $\rho(t) = \sup \{\tau \in \mathbb{T} : \tau < t\}$ , and  $\mu(t) = \sigma(t) - t$  supplemented by  $\inf \emptyset := \sup \mathbb{T}$  and  $\sup \emptyset := \inf \mathbb{T}$ . A point  $t \in \mathbb{T}$  is left-dense, left-scattered, right-dense, or right-scattered if  $\rho(t) = t$ ,  $\rho(t) < t$ ,  $\sigma(t) = t$ , or  $\sigma(t) > t$ , respectively. If  $\mathbb{T}$  has a left-scattered maximum  $M$ , then we define  $\mathbb{T}^\kappa = \mathbb{T} \setminus \{M\}$ ; otherwise,  $\mathbb{T}^\kappa = \mathbb{T}$ . If  $\mathbb{T}$  has a right-scattered

minimum  $m$ , then we define  $\mathbb{T}_\kappa = \mathbb{T} \setminus \{m\}$ ; otherwise,  $\mathbb{T}_\kappa = \mathbb{T}$ . If  $f : \mathbb{T} \rightarrow \mathbb{R}$ , then  $f^\sigma : \mathbb{T} \rightarrow \mathbb{R}$  is given by  $f^\sigma(t) = f(\sigma(t))$  for all  $t \in \mathbb{T}$ .

**Definition 2.2.** Let  $f : \mathbb{T} \rightarrow \mathbb{R}$  and  $t \in \mathbb{T}^\kappa$ . We define  $f^\Delta(t)$  to be the number, provided it exists, with the property that given any  $\varepsilon > 0$  there is a neighborhood  $U$  of  $t$  (i.e.,  $U = (t - \delta, t + \delta) \cap \mathbb{T}$  for some  $\delta > 0$ ) such that

$$|[f^\sigma(t) - f(s)] - f^\Delta(t)[\sigma(t) - s]| \leq \varepsilon |\sigma(t) - s| \text{ for all } s \in U.$$

We call  $f^\Delta(t)$  the delta derivative of  $f$  at  $t$ . Moreover, we say that  $f$  is delta (or Hilger) differentiable on  $\mathbb{T}^\kappa$  provided  $f^\Delta(t)$  exists for all  $t \in \mathbb{T}^\kappa$ .

**Definition 2.3** (See [2]). A function  $f : \mathbb{T} \rightarrow \mathbb{R}$  is said to be regressive provided  $1 + \mu(t)f(t) \neq 0$  for all  $t \in \mathbb{T}^\kappa$ . The set of all regressive and rd-continuous functions (a function  $g : \mathbb{T} \rightarrow \mathbb{R}$  is called rd-continuous provided it is continuous at right-dense points in  $\mathbb{T}$  and its left-sided limits exist (finite) at left-dense points in  $\mathbb{T}$ )  $f : \mathbb{T} \rightarrow \mathbb{R}$  will be denoted by  $\mathcal{R} = \mathcal{R}(\mathbb{T}, \mathbb{R})$ . We define the set

$$\mathcal{R}^+ = \mathcal{R}^+(\mathbb{T}, \mathbb{R}) = \{f \in \mathcal{R} : 1 + \mu(t)f(t) > 0 \text{ for all } t \in \mathbb{T}\}.$$

Let now  $p$  and  $q \in \mathcal{R}$ . We define the circle plus addition  $\oplus$  on  $\mathcal{R}$  by

$$(p \oplus q)(t) := p(t) + q(t) + \mu(t)p(t)q(t) \quad \text{for all } t \in \mathbb{T}$$

and the circle minus subtraction  $\ominus$  on  $\mathcal{R}$  by

$$(p \ominus q)(t) := \frac{p(t) - q(t)}{1 + \mu(t)q(t)} \quad \text{for all } t \in \mathbb{T}.$$

The time scales exponential function  $e_p(\cdot, t_0)$  is defined for  $p \in \mathcal{R}$  and  $t_0 \in \mathbb{T}$  as the unique solution of the initial value problem

$$y^\Delta = p(t)y \text{ and } y(t_0) = 1 \text{ on } \mathbb{T}.$$

One has

$$e_p(\cdot, t_0)e_q(\cdot, t_0) = e_{p \oplus q}(\cdot, t_0), \quad \frac{e_p(\cdot, t_0)}{e_q(\cdot, t_0)} = e_{p \ominus q}(\cdot, t_0) \text{ and } e_{\ominus q}(\cdot, t_0) = \frac{1}{e_q(\cdot, t_0)}.$$

**Lemma 2.4** (See [10]). *Assume that  $a > 0$ ,  $b > 0$ , and that  $-a \in \mathcal{R}^+$ . Then,*

$$y^\Delta(t) \geq (\leq) b - ay^\sigma(t), \quad y(t) > 0, \quad t \in [t_0, \infty)_{\mathbb{T}}$$

*implies that*

$$y(t) \geq (\leq) \frac{b}{a} \left[ 1 + \left( \frac{ay(t_0)}{b} - 1 \right) e_{(\ominus a)}(t, t_0) \right], \quad t \in [t_0, \infty)_{\mathbb{T}}.$$

**Definition 2.5** (See [14]). A time scale  $\mathbb{T}$  is called an almost periodic time scale if

$$\Pi = \{\tau \in \mathbb{R} : t + \tau \in \mathbb{T} \text{ for all } t \in \mathbb{T}\} \neq \{0\}.$$

**Definition 2.6** (See [14]). Let  $\mathbb{T}$  be an almost periodic time scale. A function  $x \in C(\mathbb{T}, \mathbb{R}^n)$  is called an almost periodic function if the  $\varepsilon$ -translation set of  $x$ ,

$$E\{\varepsilon, x\} = \{\tau \in \Pi : |x(t + \tau) - x(t)| < \varepsilon \text{ for all } t \in \mathbb{T}\},$$

is a relatively dense set in  $\mathbb{T}$ , that is, for all  $\varepsilon > 0$  there exists a constant  $l(\varepsilon) > 0$  such that each interval of length  $l(\varepsilon)$  contains a  $\eta(\varepsilon) \in E\{\varepsilon, x\}$  such that  $|x(t + \tau) - x(t)| < \varepsilon$  for all  $t \in \mathbb{T}$ . Moreover,  $\tau$  is called the  $\varepsilon$ -translation number of  $x(t)$  and  $l(\varepsilon)$  is called the inclusion length of  $E\{\varepsilon, x\}$ .

**Definition 2.7** (See [14]). Let  $\mathbb{D}$  be an open set in  $\mathbb{R}^n$  and let  $\mathbb{T}$  be a positive almost periodic time scale. A function  $f \in C(\mathbb{T} \times \mathbb{D}, \mathbb{R}^n)$  is called an almost periodic function in  $t \in \mathbb{T}$ , uniformly for  $x \in \mathbb{D}$ , if the  $\varepsilon$ -translation set of  $f$ ,

$$E\{\varepsilon, f, \mathbb{S}\} = \{\tau \in \Pi : |f(t + \tau) - f(t)| < \varepsilon \text{ for all } (t, x) \in \mathbb{T} \times \mathbb{S}\},$$

is a relatively dense set in  $\mathbb{T}$  for all  $\varepsilon > 0$  and for each compact subset  $\mathbb{S}$  of  $\mathbb{D}$  there exists a constant  $l(\varepsilon, \mathbb{S}) > 0$  such that each interval of length  $l(\varepsilon, \mathbb{S})$  contains  $\tau(\varepsilon, \mathbb{S}) \in E\{\varepsilon, f, \mathbb{S}\}$  such that

$$|f(t + \tau, x) - f(t, x)| < \varepsilon \text{ for all } (t, x) \in \mathbb{T} \times \mathbb{S}.$$

Consider the system

$$(3) \quad x^\Delta(t) = h(t, x)$$

and its associate product system

$$x^\Delta(t) = h(t, x), \quad z^\Delta(t) = h(t, z),$$

where  $h : \mathbb{T}^+ \times \mathbb{S}_B \rightarrow \mathbb{R}^n$ ,  $\mathbb{S}_B = \{x \in \mathbb{R}^n : \|x\| < B\}$  and  $h(t, x)$  is almost periodic in  $t$  uniformly for  $x \in \mathbb{S}_B$  and continuous in  $x$ .

**Lemma 2.8** (See [29]). *Suppose that there exists a Lyapunov function  $V(t, x, z)$  defined on  $\mathbb{T}^+ \times \mathbb{S}_B \times \mathbb{S}_B$ , that is, there exists a function  $V(t, x, z)$  satisfying the following conditions:*

$$(1) \quad a(\|x - z\|) \leq V(t, x, z) \leq b(\|x - z\|), \quad \text{where } a, b \in \mathbb{K} \text{ with}$$

$$\mathbb{K} = \{\alpha \in C(\mathbb{R}^+, \mathbb{R}^+) : \alpha(0) = 0, \text{ and } \alpha \text{ increasing}\};$$

$$(2) \quad |V(t, x, z) - V(t, x_1, z_1)| \leq L(\|x - x_1\| + \|z - z_1\|), \text{ where } L > 0 \text{ is a constant};$$

$$(3) \quad D^+V^\Delta(t, x, z) \leq -cV(t, x, z), \text{ where } c > 0 \text{ and } -c \in \mathcal{R}^+.$$

Furthermore, if there exists a solution  $x(t) \in \mathbb{S}$  of system (3) for  $t \in \mathbb{T}^+$ , where  $\mathbb{S} \cup \mathbb{S}_B$  is a compact set, then there exist a unique almost periodic solution  $f(t) \in \mathbb{S}$  of system (3), which is uniformly asymptotically stable.

**Definition 2.9** (See [19]). System (2) is said to be permanent if there exist positive constants  $m$  and  $M$  such that

$$m \leq \liminf_{t \rightarrow \infty} x_i(t) \leq \limsup_{t \rightarrow \infty} x_i(t) \leq M, \quad i = 1, 2, 3, 4,$$

for any solution  $(x_1(t), x_2(t), x_3(t), x_4(t))$  of (2).

### 3. PERMANENCE OF POSITIVES SOLUTIONS

The principal objective of this section is to establish sufficient conditions for system (2) to be permanent. Let  $t_0 \in \mathbb{T}$  be a fixed positive initial time. We introduce the following assumption for (2):

( $H_1$ )  $\lambda(t)$  is a bounded and almost periodic function satisfying

$$0 < \lambda^L \leq \lambda(t) \leq \lambda^U.$$

**Lemma 3.1.** *Suppose hypothesis ( $H_1$ ) holds. Then, for any positive solution  $(x_1(t), x_2(t), x_3(t), x_4(t))$  of system (2), there exists positive constants  $M$  and  $T$  such that  $x_i(t) \leq M$ ,  $i = 1, 2, 3, 4$ , for  $t \geq T$ .*

*Proof.* Let  $Z(t) = (x_1(t), x_2(t), x_3(t), x_4(t))$  be any positive solution of system (2). From the  $i$ th equation of system (2), we have

$$(4) \quad \begin{cases} x_1^\Delta(t) \leq \Lambda - (\beta\lambda^L + \nu)x_1^\sigma(t), \\ x_2^\Delta(t) \leq \beta\lambda^U M_1 + (\gamma + \omega)\frac{\Lambda}{\nu} - (\rho + \phi + \nu)x_2^\sigma(t), \\ x_3^\Delta(t) \leq \phi M_2 - (\omega + \nu)x_3^\sigma(t), \\ x_4^\Delta(t) \leq \rho M_2 - (\gamma + \nu + d)x_4^\sigma(t). \end{cases}$$

Hence, by Lemma 2.4, there exist positive constants  $M_i$  and  $T_i$  such that for any positive solution  $(x_1(t), x_2(t), x_3(t), x_4(t))$  of system (2), we have

$$x_1(t) \leq \frac{\Lambda}{\beta\lambda^L + \nu} \left[ 1 + \left( \frac{(\beta\lambda^L + \nu)x_1(t_0)}{\Lambda} - 1 \right) e_{\ominus(\beta\lambda^L + \nu)}(t, t_0) \right].$$

If  $-(\beta\lambda^L + \nu) < 0$ , then  $e_{\ominus(\beta\lambda^L + \nu)}(t, t_0) \rightarrow 0$  as  $t \rightarrow \infty$  and

$$(5) \quad \begin{cases} x_1(t) \leq M_1 := \Lambda/(\beta\lambda^L + \nu), & \text{for } t \geq T_1, \\ x_2(t) \leq M_2 := \beta(\lambda^U M_1 + (\gamma + \omega)\frac{\Lambda}{\nu})/(\rho + \phi + \nu), & \text{for } t \geq T_2, \\ x_3(t) \leq M_3 := \phi M_2/(\omega + \nu), & \text{for } t \geq T_3, \\ x_4(t) \leq M_4 := \rho M_2/(\gamma + \nu + d), & \text{for } t \geq T_4. \end{cases}$$

Let  $M = \max_{1 \leq i \leq 4} \{M_i\}$  and  $T = \max_{1 \leq i \leq 4} \{T_i\}$ . Then,  $x_i(t) \leq M$ ,  $i = 1, 2, 3, 4$ , for all  $t \geq T$ .  $\square$

**Lemma 3.2.** *Suppose that ( $H_1$ ) holds. Then, system (2) is permanent.*

*Proof.* Let  $Z(t) = (x_1(t), x_2(t), x_3(t), x_4(t))$  be any positive solution of system (2). From the  $i$ th equation of system (2), we have

$$(6) \quad \begin{cases} x_1^\Delta(t) \geq \Lambda - (\beta\lambda^U + \nu)x_1^\sigma(t), \\ x_2^\Delta(t) \geq \beta\lambda^L m_1 - (\rho + \phi + \nu)x_2^\sigma(t), \\ x_3^\Delta(t) \geq \phi m_2 - (\omega + \nu)x_3^\sigma(t), \\ x_4^\Delta(t) \geq \rho m_2 - (\gamma + \nu + d)x_4^\sigma(t). \end{cases}$$

From hypothesis  $(H_1)$  and Lemma 2.4, there exists positive constants  $m_i > 0$  such that for any positive solution  $(x_1(t), x_2(t), x_3(t), x_4(t))$  of system (2) there exists  $\hat{T}_i$  such that

$$(7) \quad \begin{cases} x_1(t) \geq m_1 := \Lambda/(\beta\lambda^U + \nu), & \text{for } t \geq \hat{T}_1, \\ x_2(t) \geq m_2 := (\beta\lambda^L m_1)/(\rho + \phi + \nu), & \text{for } t \geq \hat{T}_2, \\ x_3(t) \geq m_3 := \phi m_2/(\omega + \nu), & \text{for } t \geq \hat{T}_3, \\ x_4(t) \geq m_4 := \rho m_2/(\gamma + \nu + d), & \text{for } t \geq \hat{T}_4. \end{cases}$$

Let  $m = \min_{1 \leq i \leq 4} \{m_i\}$  and  $\hat{T} = \max_{1 \leq i \leq 4} \{\hat{T}_i\}$ . We conclude that  $x_i(t) \geq m$ ,  $i = 1, 2, 3, 4$ , for all  $t \geq \hat{T}$ .  $\square$

#### 4. UNIFORM ASYMPTOTIC STABILITY

In this section, we prove sufficient conditions for the existence and uniform asymptotic stability of the unique positive almost periodic solution to system (2). Let us define

$$\Omega := \left\{ Z(t) = (x_1(t), x_2(t), x_3(t), x_4(t)) \in (\mathbb{R}^+)^4 : (x_1(t), x_2(t), x_3(t), x_4(t)) \right. \\ \left. \text{is a solution of (2) with } 0 < m \leq x_i \leq M, i = 1, \dots, 4, \text{ and } N(t) \leq \frac{\Lambda}{\nu} \right\}.$$

It is clear that  $\Omega$  is an invariant set of system (2) and, by Lemma 3.2, we have  $\Omega \neq \emptyset$ . We introduce some more notation. Let

$$\begin{aligned} a_1 &:= \beta\lambda^L + \nu, \\ a_2 &:= \rho + \phi + \nu, \\ a_3 &:= \omega + \nu, \\ a_4 &:= \gamma + \nu + d, \\ b_1 &:= \left( \beta\lambda^U + \frac{\beta^2 M}{2m} \right), \\ b_2 &:= (\rho + \phi), \\ b_3 &:= \left( \omega + \frac{\beta^2 M}{2m} \eta_C \right), \\ b_4 &:= \left( \gamma + \frac{\beta^2 M}{2m} \eta_A \right). \end{aligned}$$

Moreover, let  $\Gamma_1 := \min_{1 \leq i \leq 4} a_i$  and  $\Gamma_2 := \max_{1 \leq i \leq 4} b_i$ . In our next result (Theorem 4.1) we assume the following additional hypothesis:

$$(H_2) \quad \Gamma_2 < \Gamma_1 \text{ with } \Gamma_1, \Gamma_2 \in \mathcal{R}^+.$$

**Theorem 4.1.** *Suppose that  $(H_1)$  and  $(H_2)$  hold. Then the dynamic system (2) has a unique almost periodic solution  $Z(t) = (x_1(t), x_2(t), x_3(t), x_4(t)) \in \Omega$  that is uniformly asymptotically stable.*

*Proof.* According to Lemma 2.4, every solution

$$Z(t) = (x_1(t), x_2(t), x_3(t), x_4(t))$$

of system (2) satisfies  $x_i^L \leq x_i(t) \leq x_i^U, i = 1, \dots, 4$ , and  $|x_i| \leq K_i, i = 1, \dots, 4$ . Denote

$$\begin{aligned} \|Z\| &= \| (x_1(t), x_2(t), x_3(t), x_4(t)) \| \\ &= \sup_{t \in \mathbb{T}^+} (|x_1(t)| + |x_2(t)| + |x_3(t)| + |x_4(t)|). \end{aligned}$$

Let  $Z(t) = (x_1(t), x_2(t), x_3(t), x_4(t))$  and  $\hat{Z}(t) = (\hat{x}_1(t), \hat{x}_2(t), \hat{x}_3(t), \hat{x}_4(t))$  be two positive solutions of (2). Then  $\|Z\| \leq K$  and  $\|\hat{Z}\| \leq K$ , where

$$K = \sum_{i=1}^{i=4} K_i.$$

In view of (2), we have

$$\begin{cases} x_1^\Delta(t) = \Lambda - \beta\lambda(t)x_1^\sigma(t) - \nu x_1^\sigma(t), \\ x_2^\Delta(t) = \beta\lambda(t)x_1(t) - (\rho + \phi + \nu)x_2^\sigma(t) + \gamma x_4(t) + \omega x_3(t), \\ x_3^\Delta(t) = \phi x_2(t) - (\omega + \nu)x_3^\sigma(t), \\ x_4^\Delta(t) = \rho x_2(t) - (\gamma + \nu + d)x_4^\sigma(t), \end{cases}$$



and

$$\begin{cases} \hat{x}_1^\Delta(t) = \Lambda - \beta\hat{\lambda}(t)\hat{x}_1^\sigma(t) - \nu\hat{x}_1^\sigma(t), \\ \hat{x}_2^\Delta(t) = \beta\hat{\lambda}(t)\hat{x}_1(t) - (\rho + \phi + \nu)\hat{x}_2^\sigma(t) + \gamma\hat{x}_4(t) + \omega\hat{x}_3(t), \\ \hat{x}_3^\Delta(t) = \phi\hat{x}_2(t) - (\omega + \nu)\hat{x}_3^\sigma(t), \\ \hat{x}_4^\Delta(t) = \rho\hat{x}_2(t) - (\gamma + \nu + d)\hat{x}_4^\sigma(t). \end{cases}$$

Define the Lyapunov function  $V(t, Z, \hat{Z})$  on  $\mathbb{T}^+ \times \Omega \times \Omega$  as

$$V(t, Z, \hat{Z}) = \sum_{i=1}^{i=4} |x_i(t) - \hat{x}_i(t)| = \sum_{i=1}^{i=4} V_i(t),$$

where  $V_i(t) = |x_i(t) - \hat{x}_i(t)|$ . The two norms

$$\|Z(t) - \hat{Z}(t)\| = \sup_{t \in \mathbb{T}^+} \sum_{i=1}^{i=4} |x_i(t) - \hat{x}_i(t)|$$

and

$$\|Z(t) - \hat{Z}(t)\|_* = \sup_{t \in \mathbb{T}^+} \left( \sum_{i=1}^{i=4} (x_i(t) - \hat{x}_i(t))^2 \right)^{\frac{1}{2}}$$

are equivalent, that is, there exist two constants  $\eta_1$  and  $\eta_2 > 0$  such that

$$\eta_1 \|Z(t) - \hat{Z}(t)\|_* \leq \|Z(t) - \hat{Z}(t)\| \leq \eta_2 \|Z(t) - \hat{Z}(t)\|_*.$$

Hence,

$$\eta_1 \|Z(t) - \hat{Z}(t)\|_* \leq V(t, Z, \hat{Z}) \leq \eta_2 \|Z(t) - \hat{Z}(t)\|_*.$$

Let  $a, b \in C(\mathbb{R}^+, \mathbb{R}^+)$ ,  $a(x) = \eta_1 x$  and  $b(x) = \eta_2 x$ . Then the assumption (i) of Lemma 2.8 is satisfied. Moreover,

$$\begin{aligned} |V(t, Z, \hat{Z}) - V(t, Z^*, \hat{Z}^*)| &= \left| \sum_{i=1}^{i=4} |x_i(t) - \hat{x}_i(t)| - \sum_{i=1}^{i=4} |x_i^*(t) - \hat{x}_i^*(t)| \right| \\ &\leq \sum_{i=1}^{i=4} |(x_i(t) - \hat{x}_i(t)) - (x_i^*(t) - \hat{x}_i^*(t))| \\ &\leq \sum_{i=1}^{i=4} |(x_i(t) - x_i^*(t)) + (\hat{x}_i^*(t) - \hat{x}_i(t))| \\ &\leq \sum_{i=1}^{i=4} |x_i(t) - x_i^*(t)| + \sum_{i=1}^{i=4} |\hat{x}_i^*(t) - \hat{x}_i(t)| \\ &\leq L \left( \|Z(t) - Z^*(t)\| + \|\hat{Z}(t) - \hat{Z}^*(t)\| \right), \end{aligned}$$

where  $L = 1$ , so that condition (ii) of Lemma 2.8 is also satisfied. Now, let  $v_i(t) = x_i(t) - \hat{x}_i(t)$ ,  $i = 1, \dots, 4$ . We compute and estimate the Dini derivative  $D^+V^\Delta$  of  $V$  along the associated product system (2). Using Lemma 4.1

of [15], it follows that  $D^+V_1^\Delta(t) \leq \text{sign}(v_1^\sigma(t))(v_1(t))^\Delta$ . For more details on  $D^+V_1^\Delta(t)$  see [9, 13]. Now, let us begin computing

$$\begin{aligned}
D^+V_1^\Delta(t) &\leq \text{sign}(v_1^\sigma(t))(v_1(t))^\Delta \\
&= \text{sign}(v_1^\sigma(t))[-\beta\lambda(t)x_1^\sigma(t) - \nu x_1^\sigma(t) + \beta\hat{\lambda}(t)\hat{x}_1^\sigma(t) + \nu\hat{x}_1^\sigma(t)] \\
&= \text{sign}(v_1^\sigma(t))[-\beta\lambda(t)(x_1^\sigma(t) - \hat{x}_1^\sigma(t)) - \nu(x_1^\sigma(t) - \hat{x}_1^\sigma(t)) \\
&\quad - \beta\hat{x}_1^\sigma(t)(\lambda(t) - \hat{\lambda}(t))] \\
&= \text{sign}(v_1^\sigma(t))[-(\beta\lambda(t) + \nu)(x_1^\sigma(t) - \hat{x}_1^\sigma(t)) \\
&\quad - \beta\hat{x}_1^\sigma(t)(\lambda(t) - \hat{\lambda}(t))] \\
&\leq -(\beta\lambda(t) + \nu) | \hat{x}_1^\sigma(t) - x_1^\sigma(t) | + \beta\hat{x}_1^\sigma(t) | \lambda(t) - \hat{\lambda}(t) | \\
&\leq -(\beta\lambda^L(t) + \nu) | \hat{x}_1^\sigma(t) - x_1^\sigma(t) | + \beta\hat{x}_1^\sigma(t) | \lambda(t) - \hat{\lambda}(t) | \\
&\leq -(\beta\lambda^L(t) + \nu) | \hat{x}_1^\sigma(t) - x_1^\sigma(t) | + \beta M | \lambda(t) - \hat{\lambda}(t) |,
\end{aligned}$$

that is,

$$D^+V_1^\Delta(t) \leq -(\beta\lambda^L(t) + \nu) | \hat{x}_1^\sigma(t) - x_1^\sigma(t) | + (\beta M) | \lambda(t) - \hat{\lambda}(t) |.$$

Let  $v_2(t) = x_2(t) - \hat{x}_2(t)$ . Similarly, we have

$$\begin{aligned}
D^+V_2^\Delta(t) &\leq \text{sign}(v_2^\sigma(t))(v_2(t))^\Delta \\
&= \text{sign}(v_2^\sigma(t))[\beta\lambda(t)x_1(t) - (\rho + \phi + \nu)x_2^\sigma(t) + \gamma x_4(t) \\
&\quad + \omega x_3(t) - \beta\hat{\lambda}(t)\hat{x}_1(t) + (\rho + \phi + \nu)\hat{x}_2^\sigma(t) - \gamma\hat{x}_4(t) - \omega\hat{x}_3(t)] \\
&= \text{sign}(v_2^\sigma(t))[(\beta\lambda(t)x_1(t) - \beta\hat{\lambda}(t)\hat{x}_1(t)) - (\rho + \phi + \nu)(x_2^\sigma(t) - \hat{x}_2^\sigma(t)) \\
&\quad + \gamma(x_4(t) - \hat{x}_4(t)) + \omega(x_3(t) - \hat{x}_3(t))] \\
&= \text{sign}(v_2^\sigma(t))[\beta(\lambda(t)(x_1(t) - \hat{x}_1(t)) + \hat{x}_1(t)(\lambda(t) - \hat{\lambda}(t))) \\
&\quad - (\rho + \phi + \nu)(x_2^\sigma(t) - \hat{x}_2^\sigma(t)) + \gamma(x_4(t) - \hat{x}_4(t)) + \omega(x_3(t) - \hat{x}_3(t))] \\
&= \text{sign}(v_2^\sigma(t))[\beta\lambda(t)(x_1(t) - \hat{x}_1(t)) + \beta\hat{x}_1(t)(\lambda(t) - \hat{\lambda}(t)) \\
&\quad - (\rho + \phi + \nu)(x_2^\sigma(t) - \hat{x}_2^\sigma(t)) + \gamma(x_4(t) - \hat{x}_4(t)) + \omega(x_3(t) - \hat{x}_3(t))] \\
&\leq \beta\lambda^U | x_1(t) - \hat{x}_1(t) | + \beta M | \lambda(t) - \hat{\lambda}(t) | \\
&\quad - (\rho + \phi + \nu) | x_2^\sigma(t) - \hat{x}_2^\sigma(t) | + \gamma | x_4(t) - \hat{x}_4(t) | \\
&\quad + \omega | x_3(t) - \hat{x}_3(t) |;
\end{aligned}$$

for  $v_3(t) = x_3(t) - \hat{x}_3(t)$  one has

$$\begin{aligned}
D^+V_3^\Delta(t) &\leq \text{sign}(v_3^\sigma(t))(v_3(t))^\Delta \\
&= \text{sign}(v_3^\sigma(t))[\phi x_2(t) - (\omega + \nu)x_3^\sigma(t) - \phi\hat{x}_2(t) + (\omega + \nu)\hat{x}_3^\sigma(t)] \\
&\leq \phi | x_2(t) - \hat{x}_2(t) | - (\omega + \nu) | x_3^\sigma(t) - \hat{x}_3^\sigma(t) |;
\end{aligned}$$

and for  $v_4(t) = x_4(t) - \hat{x}_4(t)$  we have

$$D^+V_4^\Delta(t) \leq \rho | x_2(t) - \hat{x}_2(t) | - (\gamma + \nu + d) | x_4^\sigma(t) - \hat{x}_4^\sigma(t) |.$$

Since

$$\lambda(t) = \frac{\beta}{N(t)}(x_1(t) + \eta_C x_3(t) + \eta_A) x_4(t),$$

it follows that

$$(8) \quad \begin{aligned} D^+ V^\Delta(t) \leq & -(\beta \lambda^L(t) + \nu) | \hat{x}_1^\sigma(t) - \hat{x}_1^\sigma(t) | + (\beta M) | \lambda(t) - \hat{\lambda}(t) | \\ & + \beta \lambda^U | x_1(t) - \hat{x}_1(t) | + \beta M | \lambda(t) - \hat{\lambda}(t) | \\ & - (\rho + \phi + \nu) | x_2^\sigma(t) - \hat{x}_2^\sigma | + \gamma | x_4(t) - \hat{x}_4(t) | \\ & + \omega | x_3(t) - \hat{x}_3(t) | + \phi | x_2(t) - \hat{x}_2(t) | \\ & - (\omega + \nu) | x_3^\sigma(t) - \hat{x}_3^\sigma(t) | \\ & + \rho | x_2(t) - \hat{x}_2(t) | - (\gamma + \nu + d) | x_4^\sigma(t) - \hat{x}_4^\sigma(t) | . \end{aligned}$$

Therefore,

$$|\lambda(t) - \hat{\lambda}(t)| \leq \frac{\beta}{4m} [| x_1(t) - \hat{x}_1(t) | + \eta_C | x_3(t) - \hat{x}_3(t) | + \eta_A | x_4(t) - \hat{x}_4(t) |],$$

where  $0 \leq \eta_C \leq 1$ ,  $\eta_A \geq 1$ , and  $\beta > 0$ . The inequality (8) becomes

$$\begin{aligned} D^+ V^\Delta(t) \leq & -(\beta \lambda^L + \nu) | x_1^\sigma(t) - \hat{x}_1^\sigma(t) | - (\rho + \phi + \nu) | x_2^\sigma(t) - \hat{x}_2^\sigma(t) | \\ & - (\omega + \nu) | x_3^\sigma(t) - \hat{x}_3^\sigma(t) | - (\gamma + \nu + d) | x_4^\sigma(t) - \hat{x}_4^\sigma(t) | \\ & + 2\beta M | \lambda(t) - \hat{\lambda}(t) | + (\beta \lambda^U) | x_1(t) - \hat{x}_1(t) | + \gamma | x_4(t) - \hat{x}_4(t) | \\ & + \omega | x_3(t) - \hat{x}_3(t) | + (\rho + \phi) | x_2(t) - \hat{x}_2(t) | \\ = & -(\beta \lambda^L + \nu) | x_1^\sigma(t) - \hat{x}_1^\sigma(t) | - (\rho + \phi + \nu) | x_2^\sigma(t) - \hat{x}_2^\sigma(t) | \\ & - (\omega + \nu) | x_3^\sigma(t) - \hat{x}_3^\sigma(t) | - (\gamma + \nu + d) | x_4^\sigma(t) - \hat{x}_4^\sigma(t) | \\ & + \beta \left( \lambda^U + \frac{\beta M}{2m} \right) | x_1(t) - \hat{x}_1(t) | + (\rho + \phi) | x_2(t) - \hat{x}_2(t) | \\ & + \left( \omega + \frac{\beta^2 M}{2m} \eta_C \right) | x_3(t) - \hat{x}_3(t) | \\ & + \left( \gamma + \frac{\beta^2 M}{2m} \eta_A \right) | x_4(t) - \hat{x}_4(t) | . \end{aligned}$$

From (9) we can write that

$$\begin{aligned}
D^+V^\Delta(t) &\leq -(\beta\lambda^L + \nu) |x_1^\sigma(t) - \hat{x}_1^\sigma(t)| - (\rho + \phi + \nu) |x_2^\sigma(t) - \hat{x}_2^\sigma(t)| \\
&\quad - (\omega + \nu) |x_3^\sigma(t) - \hat{x}_3^\sigma(t)| - (\gamma + \nu + d) |x_4^\sigma(t) - \hat{x}_4^\sigma(t)| \\
&\quad + \left(\beta\lambda^U + \frac{\beta^2 M}{2m}\right) |\hat{x}_1(t) - \hat{x}_1(t)| \\
&\quad + (\phi + \rho) |x_2(t) - \hat{x}_2(t)| \\
&\quad + \left(\omega + \frac{\beta^2 M}{2m}\eta_C\right) |x_3(t) - \hat{x}_3(t)| \\
&\quad + \left(\gamma + \frac{\beta^2 M}{2m}\eta_A\right) |x_4(t) - \hat{x}_4(t)| \\
&= -a_1 |x_1^\sigma(t) - \hat{x}_1^\sigma(t)| - a_2 |x_2^\sigma(t) - \hat{x}_2^\sigma(t)| \\
&\quad - a_3 |x_3^\sigma(t) - \hat{x}_3^\sigma(t)| - a_4 |x_4^\sigma(t) - \hat{x}_4^\sigma(t)| \\
&\quad + b_1 |x_1(t) - \hat{x}_1(t)| + b_2 |x_2(t) - \hat{x}_2(t)| \\
&\quad + b_3 |x_3(t) - \hat{x}_3(t)| + b_4 |x_4(t) - \hat{x}_4(t)| \\
&= -\Gamma_1 V(\sigma(t)) + \Gamma_2 V(t) \\
&= (\Gamma_2 - \Gamma_1) V(t) - \Gamma_1 \mu(t) D^+V^\Delta(t)
\end{aligned}$$

and it follows that  $D^+V^\Delta(t) \leq \frac{\Gamma_2 - \Gamma_1}{(1 + \Gamma_1 \mu)} V(t) = -\Psi V(t)$ . By hypothesis

$(H_2)$ , one has  $-\Psi \in \mathcal{R}^+$  and  $\Psi = \frac{\Gamma_1 - \Gamma_2}{(1 + \Gamma_1 \mu)} > 0$ . Thus, the assumption (iii) of Lemma 2.8 is satisfied. Furthermore, the conditions (i) and (ii) of Lemma 2.8 also hold. For condition (i) we consider two functions  $a, b \in C(\mathbb{R}^+)$  with  $a(x) = 0.5x$  and  $b(x) = 2x$ . For condition (ii) we put  $L = 1$ . So there exists a unique uniformly asymptotically stable almost periodic solution  $Z(t) = (x_1(t), x_2(t), x_3(t), x_4(t))$  of the dynamic system (2) with  $Z(t) \in \Omega$ .  $\square$

**Example 4.2.** Motivated by [16] and the case of Morocco, we consider the following system for  $\mathbb{T} = \mathbb{Z}^+$ :

$$(9) \quad \begin{cases} x_1^\Delta(t) = \Lambda - \beta\lambda(t)x_1^\sigma(t) - \nu x_1^\sigma(t), \\ x_2^\Delta(t) = \beta\lambda(t)x_1(t) - (\rho + \phi + \nu)x_2^\sigma(t) + \gamma x_4(t) + \omega x_3(t), \\ x_3^\Delta(t) = \phi x_2(t) - (\omega + \nu)x_3^\sigma(t), \\ x_4^\Delta(t) = \rho x_2(t) - (\gamma + \nu + d)x_4^\sigma(t), \end{cases}$$

where

$$x_1(0) = 1 - \frac{11}{N_0}, \quad x_2(0) = \frac{2}{N_0}, \quad x_3(0) = 0, \quad x_4(0) = \frac{9}{N_0},$$

with  $N_0 = 325235$ ,  $\Lambda = 2190$ ,  $\beta = 2.710^{-7}$ ,  $\nu = 0.39$ ,  $\rho = 0.2$ ,  $\phi = 0.1$ ,  $\gamma = 0.33$ ,  $\omega = 0.09$ , and  $d = 1$ . The system (9) is permanent. Furthermore,  $m_1 = 5615.381462$ ,  $m_2 = 5.47870412010^{-9}$ ,  $m_3 = 1.14139669210^{-9}$ ,




FIGURE 1. Example 4.2: solution of (9) during 7 years.

$m_4 = 6.37058618610^{-10}$ ,  $M_1 = 5615.384615$ ,  $M_2 = 0.002773104793$ ,  $M_3 = 0.0005777301652$ , and  $M_4 = 0.0003224540457$ . The conditions of Theorem 4.1 are verified with  $0.37 = \Gamma_2 < \Gamma_1 = 0.39$  and  $\Psi = 0.01391941151$ . Therefore, system (9) has a unique positive almost periodic solution, which is uniformly asymptotic stable. In Figure 1 we plot the solution for the first 7 years with  $\eta_C = 0.5$  and  $\eta_A = 1.5$ . Computationally, it is difficult to plot the solution for longer periods of time.

## 5. CONCLUSION

We have investigated the uniform stability of the singular positive solution in an HIV/AIDS epidemic model, specifically the SICA model on an arbitrary time scale. The purpose of incorporating time scales is to integrate both continuous and discrete time models. We established the permanence of each solution and, using a suitable Lyapunov function, we derived a sufficient condition for uniform asymptotic stability of the solution. Additionally, we presented an illustrative example to substantiate our analytical results.

**Acknowledgments.** This research is part of first author's Ph.D. project, which is carried out at University Mustapha Stambouli of Mascara.

**Funding.** Torres was supported by CIDMA and is funded by the Fundação para a Ciência e a Tecnologia, I.P. (FCT, Funder ID = 50110000187) under Grants UIDB/04106/2020 and UIDP/04106/2020.

**Competing and Conflict of Interests.** The authors have no conflicts or competing of interests to declare.

**Data Availability.** No data is associated to the manuscript.

**Author contributions.** All authors contributed equally to this work and reviewed and approved the final manuscript.

#### REFERENCES

- [1] M. Bohner, J. Heim and A. Liu, *Solow models on time scales*, Cubo **15** (2013), no. 1, 13–31.
- [2] M. Bohner and A. Peterson, *Dynamic equations on time scales*, Birkhäuser Boston, Inc., Boston, MA, 2001.
- [3] M. Bohner and A. Peterson, *Advances in dynamic equations on time scales*, Birkhäuser Boston, Inc., Boston, MA, 2003.
- [4] M. Bohner, S. Streipert and D. F. M. Torres, *Exact solution to a dynamic SIR model*, Nonlinear Anal. Hybrid Syst. **32** (2019), 228–238. [arXiv:1812.09759](#)
- [5] A. Boukhouima, E. M. Lotfi, M. Mahrouf, S. Rosa, D. F. M. Torres and N. Yousfi, *Stability analysis and optimal control of a fractional HIV-AIDS epidemic model with memory and general incidence rate*, Eur. Phys. J. Plus **136** (2021), Art. 103, 20 pp. [arXiv:2012.04819](#)
- [6] J. Djordjevic, C. J. Silva and D. F. M. Torres, *A stochastic SICA epidemic model for HIV transmission*, Appl. Math. Lett. **84** (2018), 168–175. [arXiv:1805.01425](#)
- [7] M. Dryl and D. F. M. Torres, *A general delta-nabla calculus of variations on time scales with application to economics*, Int. J. Dyn. Syst. Differ. Equ. **5** (2014), no. 1, 42–71. [arXiv:1410.1190](#)
- [8] O. S. Fard, D. F. M. Torres and M. R. Zadeh, *A Hukuhara approach to the study of hybrid fuzzy systems on time scales*, Appl. Anal. Discrete Math. **10** (2016), no. 1, 152–167. [arXiv:1603.03737](#)
- [9] S. Hong, *Stability criteria for set dynamic equations in time scales*, Comput. Math. Appl. **59** (2010), no. 11, 3444–3457.
- [10] M. Hu and L. Wang, *Dynamic inequalities on time scales with applications in permanence of predator-prey system*, Discrete Dyn. Nat. Soc. **2012** (2012), Art. ID 281052, 15 pp.

- [11] R. Jankowski and A. Chmiel, *Role of time scales in the coupled epidemic-opinion dynamics on multiplex networks*, Entropy **24** (2022), no. 1, Paper No. 105, 14 pp.
- [12] V. Kac and P. Cheung, *Quantum calculus*, Universitext, Springer-Verlag, New York, 2002.
- [13] B. Kaymakçalan, *Lyapunov stability theory for dynamic systems on time scales*, J. Appl. Math. Stochastic Anal. **5** (1992), no. 3, 275–281.
- [14] Y. Li and C. Wang, *Uniformly almost periodic functions and almost periodic solutions to dynamic equations on time scales*, Abstr. Appl. Anal. **2011** (2011), Art. ID 341520, 22 pp.
- [15] Y. Li and C. Wang, *Almost periodic solutions to dynamic equations on time scales and applications*, J. Appl. Math. **2012**, Art. ID 463913, 19 pp.
- [16] E. M. Lotfi, M. Mahrouf, M. Maziane, C. J. Silva, D. F. M. Torres and N. Yousfi, *A minimal HIV-AIDS infection model with general incidence rate and application to Morocco data*, Stat. Optim. Inf. Comput. **7** (2019), no. 3, 588–603. [arXiv:1812.06965](https://arxiv.org/abs/1812.06965)
- [17] A. B. Malinowska and D. F. M. Torres, *Quantum variational calculus*, SpringerBriefs in Electrical and Computer Engineering, Springer, Cham, 2014.
- [18] R. M. May and R. M. Anderson, *Transmission dynamics of HIV infection*, Nature **326** (1987), 137–142.
- [19] K. R. Prasad and Md. Khuddush, *Stability of positive almost periodic solutions for a fishing model with multiple time varying variable delays on time scales*, Bull. Int. Math. Virtual Inst. **9** (2019), no. 3, 521–533.
- [20] K. R. Prasad and Md. Khuddush, *Existence and uniform asymptotic stability of positive almost periodic solutions for three-species Lotka-Volterra competitive system on time scales*, Asian-Eur. J. Math. **13** (2020), no. 3, 2050058, 24 pp.
- [21] K. R. Prasad, M. Khuddush and K. V. Vidyasagar, *Almost periodic positive solutions for a time-delayed SIR epidemic model with saturated treatment on time scales*, J. Math. Model. **9** (2021), no. 1, 45–60.
- [22] C. J. Silva and D. F. M. Torres, *A TB-HIV/AIDS coinfection model and optimal control treatment*, Discrete Contin. Dyn. Syst. **35** (2015), no. 9, 4639–4663. [arXiv:1501.03322](https://arxiv.org/abs/1501.03322)
- [23] C. J. Silva and D. F. M. Torres, *A SICA compartmental model in epidemiology with application to HIV/AIDS in Cape Verde*, Ecological Complexity **30** (2017), 70–75. [arXiv:1612.00732](https://arxiv.org/abs/1612.00732)

- [24] C. J. Silva and D. F. M. Torres, *Stability of a fractional HIV/AIDS model*, Math. Comput. Simulation **164** (2019), 180–190. [arXiv:1903.02534](#)
- [25] C. J. Silva and D. F. M. Torres, *On SICA models for HIV transmission*, in *Mathematical modelling and analysis of infectious diseases*, 155–179, Stud. Syst. Decis. Control, 302, Springer, Cham, 2020. [arXiv:2004.11903](#)
- [26] C. C. Tisdell and A. Zaidi, *Basic qualitative and quantitative results for solutions to nonlinear, dynamic equations on time scales with an application to economic modelling*, Nonlinear Anal. **68** (2008), no. 11, 3504–3524.
- [27] S. Vaz and D. F. M. Torres, *A dynamically-consistent nonstandard finite difference scheme for the SICA model*, Math. Biosci. Eng. **18** (2021), no. 4, 4552–4571. [arXiv:2105.10826](#)
- [28] C. Wang and R. P. Agarwal, *A survey of function analysis and applied dynamic equations on hybrid time scales*, Entropy **23** (2021), no. 4, Paper No. 450, 66 pp.
- [29] H. Zhang and Y. Li, *Almost periodic solutions to dynamic equations on time scales*, J. Egyptian Math. Soc. **21** (2013), no. 1, 3–10.
- [30] Y. Zhi, Z. Ding and Y. Li, *Permanence and almost periodic solution for an enterprise cluster model based on ecology theory with feedback controls on time scales*, Discrete Dyn. Nat. Soc. **2013** (2013), Art. ID 639138, 14 pp.

ZAHRA BELARBI

DEPARTMENT OF MATHEMATICS,  
UNIVERSITY MUSTAPHA STAMBOULI OF MASCARA, B.P. 305 MASCARA, ALGERIA;  
ECOLE NORMALE SUPÉRIEURE, MOSTAGANEM, ALGERIA  
*Email address:* [zahra.belarbi@univ-mascara.dz](mailto:zahra.belarbi@univ-mascara.dz)

BENAOUMEUR BAYOUR

DEPARTMENT OF MATHEMATICS,  
UNIVERSITY MUSTAPHA STAMBOULI OF MASCARA, B.P. 305 MASCARA, ALGERIA  
*Email address:* [b.bayour@univ-mascara.dz](mailto:b.bayour@univ-mascara.dz)

DELFIN F. M. TORRES

CENTER FOR RESEARCH AND DEVELOPMENT IN MATHEMATICS  
AND APPLICATIONS (CIDMA), DEPARTMENT OF MATHEMATICS,  
UNIVERSITY OF AVEIRO, 3810-193 AVEIRO, PORTUGAL;  
RESEARCH CENTER IN EXACT SCIENCES (CICE),  
FACULTY OF SCIENCES AND TECHNOLOGY,  
UNIVERSITY OF CAPE VERDE (UNI-CV), 7943-010 PRAIA, CABO VERDE  
*Email address:* [delfim@ua.pt](mailto:delfim@ua.pt); [delfim@unicv.cv](mailto:delfim@unicv.cv)  
ORCID: <http://orcid.org/0000-0001-8641-2505>