

Fast Convergence of Frank-Wolfe algorithms on polytopes

Elias Wirth* Javier Peña† Sebastian Pokutta‡

June 28, 2024

Abstract

We provide a template to derive convergence rates for the following popular versions of the Frank-Wolfe algorithm on polytopes: vanilla Frank-Wolfe, Frank-Wolfe with away steps, Frank-Wolfe with blended pairwise steps, and Frank-Wolfe with in-face directions. Our template shows how the convergence rates follow from two affine-invariant properties of the problem, namely, *error bound* and *extended curvature*. These properties depend solely on the polytope and objective function but not on any affine-dependent object like norms. For each one of the above algorithms, we derive rates of convergence ranging from sublinear to linear depending on the degree of the error bound.

1 Introduction

We consider several popular versions of the *Frank-Wolfe algorithm*, also known as conditional gradient method, for optimization problems of the form

$$\min_{\mathbf{x} \in \mathcal{C}} f(\mathbf{x}), \tag{1.1}$$

where $\mathcal{C} \subseteq \mathbb{R}^n$ is a polytope and $f: \mathcal{C} \rightarrow \mathbb{R}$ is a convex function differentiable in an open set containing \mathcal{C} . At iteration t , the *vanilla* Frank-Wolfe algorithm updates the current iterate $\mathbf{x}^{(t)} \in \mathcal{C}$ to

$$\mathbf{x}^{(t+1)} = \mathbf{x}^{(t)} + \eta_t(\mathbf{v}^{(t)} - \mathbf{x}^{(t)})$$

for $\mathbf{v}^{(t)} = \arg \min_{\mathbf{y} \in \mathcal{C}} \langle \nabla f(\mathbf{x}^{(t)}), \mathbf{y} \rangle$ and some stepsize $\eta_t \in [0, 1]$. This algorithmic scheme is attractive when projections onto \mathcal{C} are computationally expensive but a linear minimization oracle for the set \mathcal{C} is available.

When the pair (\mathcal{C}, f) satisfies a mild *curvature* property, the vanilla Frank-Wolfe algorithm has convergence rate $\mathcal{O}(t^{-1})$, that is,

$$f(\mathbf{x}^{(t)}) - \min_{\mathbf{x} \in \mathcal{C}} f(\mathbf{x}) = \mathcal{O}(t^{-1}),$$

*Technical University of Berlin

†Carnegie Mellon University

‡Technical University of Berlin

see, for example, [13]. There are various settings that yield faster convergence rates [4, 5, 8, 10, 15, 17, 19, 24, 25]. However, in the setting often referred to as *Wolfe’s lower bound* [26], the convergence rate of the Frank-Wolfe algorithm cannot be faster than $\Omega(t^{-1-\epsilon})$ for any $\epsilon > 0$. The Wolfe’s setting concerns the case when \mathcal{C} is a polytope, the optimal solution set lies in the relative interior of some proper face of \mathcal{C} , and the stepsizes are chosen via exact line-search or short-step.

Since optimization over polytopes is central to Frank-Wolfe research [3, 14, 21], this limitation has motivated the development of linearly convergent variants for settings similar to Wolfe’s [9, 12, 16, 22, 26]. In contrast to the update in the vanilla Frank-Wolfe algorithm, these variants perform an update of the form

$$\mathbf{x}^{(t+1)} = \mathbf{x}^{(t)} + \eta_t \mathbf{d}^{(t)}$$

for a more flexible choice of search direction $\mathbf{d}^{(t)}$ that relies on additional information about the current iterate $\mathbf{x}^{(t)}$. This flexibility facilitates linear convergence rates at the cost of some computational overhead.

Some popular variants are the Frank-Wolfe with away steps [1, 16], Frank-Wolfe with blended pairwise steps [22], and Frank-Wolfe with in-face directions [6, 7, 9]. These variants have stronger convergence properties than the vanilla Frank-Wolfe algorithm. In particular, under suitable smoothness and strong convexity assumptions, they have linear convergence, even in Wolfe’s setting. However, the linear convergence results derived in [1, 7, 9, 16, 22] depend on the dimension of \mathcal{C} , rely on additional difficult-to-verify assumptions, or are affine-dependent. Furthermore, all of these papers consider only two regimes of convergence: a linear rate when suitable favorable assumptions hold, or the iconic rate $\mathcal{O}(t^{-1})$ of first-order methods when only milder assumptions hold.

This paper develops a framework to derive convergence rates for the above variants of the Frank-Wolfe algorithm that circumvents the main limitations of previous approaches. Our framework is founded on two key properties of the pair (\mathcal{C}, f) , namely an *extended curvature property* and an *error bound property*. We derive rates of convergence ranging from sublinear to linear depending on the degree of the error bound property. Our approach yields the following main contributions. First, we provide a common affine-invariant template to derive convergence rates for vanilla Frank-Wolfe, as well as the Frank-Wolfe variants that include away, blended pairwise, and in-face directions. Second, we establish new convergence rates that interpolate between the $\mathcal{O}(t^{-1})$ and linear rates considered in [1, 9, 16, 22]. The specific rate of convergence is determined by the degree of the error bound property in the same spirit developed in the recent articles [19, 25] for vanilla Frank-Wolfe. Third, we sharpen the results in [16] by providing convergence results in terms of the local *facial distance*, in contrast to the dependence on the pyramidal width. The former depends only on local geometry of the optimal face of \mathcal{C} whereas the latter depends on the global geometry of the entire polytope \mathcal{C} . Fourth, we establish dimension-independent rates for the aforementioned variants of Frank-Wolfe without requiring any additional difficult-to-verify assumptions such as strict complementarity [7, 26]. Fifth, our developments enables us to subsume and extend the work of Garber and Meshi [9] for simplex-like polytopes.

1.1 Outline

The paper is structured as follows. In Section 2, we introduce the preliminaries and notation used throughout the paper. In particular, we introduce the two key building blocks underlying our approach, namely, the extended curvature and error bound properties. Section 3 presents our main developments, that is, the convergence results for vanilla Frank-Wolfe and Frank-Wolfe variants. The main convergence results are Theorem 3.2 for vanilla Frank-Wolfe, Theorem 3.6 for away-step and blended pairwise Frank-Wolfe, and Theorem 3.12 for in-face Frank-Wolfe. We deliberately use a similar format in each of the subsections of Section 3 to highlight how the convergence rates in all cases are a consequence of the extended curvature and error bound properties. When the polytope \mathcal{C} has a standard form description, Theorem 3.16 gives a stronger version of Theorem 3.12. When the polytope \mathcal{C} is a simplex-like polytope, we provide a specialized version of the in-face Frank-Wolfe and establish a much stronger convergence result, namely Theorem 3.18.

In Section 4, we develop some geometric properties of the facial structure of polytopes. These properties are the crux of sufficient conditions for the error bound property to hold.

2 Preliminaries

Throughout $\mathcal{C} \subseteq \mathbb{R}^n$ denotes a nonempty polytope equipped with a *linear oracle* that computes the mapping

$$\mathbf{g} \mapsto \arg \min_{\mathbf{y} \in \mathcal{C}} \langle \mathbf{g}, \mathbf{y} \rangle.$$

Furthermore, we shall assume that the minimizer returned by the above linear oracle is a vertex of \mathcal{C} .

We shall also assume that $f: \mathcal{C} \rightarrow \mathbb{R}$ is a convex function differentiable in an open set containing \mathcal{C} . For the optimization problem (1.1), we denote the *optimal value* by

$$f^* = \min_{\mathbf{x} \in \mathcal{C}} f(\mathbf{x})$$

and the *set of optimal solutions* by

$$X^* := \{\mathbf{x} \in \mathcal{C} \mid f(\mathbf{x}) = f^*\}.$$

Recall the following popular smoothness and Hölderian error bound properties. These concepts are defined in terms of a norm on \mathbb{R}^n and hence are typically affine-dependent.

Definition 2.1 (Smoothness). *Let \mathbb{R}^n be endowed with a norm $\|\cdot\|$, let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope, and let $f: \mathcal{C} \rightarrow \mathbb{R}$ be convex and differentiable in an open set containing \mathcal{C} . For $L > 0$ we say that f is L -smooth over \mathcal{C} with respect to $\|\cdot\|$ if for all $\mathbf{x}, \mathbf{y} \in \mathcal{C}$, it holds that*

$$f(\mathbf{y}) \leq f(\mathbf{x}) + \langle \nabla f(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle + \frac{L}{2} \|\mathbf{x} - \mathbf{y}\|^2.$$

Definition 2.2 (Hölderian error bound). *Let \mathbb{R}^n be endowed with a norm $\|\cdot\|$, let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope, and let $f: \mathcal{C} \rightarrow \mathbb{R}$ be convex and differentiable in an open set*

containing \mathcal{C} . For $\mu > 0$ and $\theta \in [0, 1/2]$ we say that (\mathcal{C}, f) satisfies the (μ, θ) -Hölderian error bound over \mathcal{C} with respect to $\|\cdot\|$ if for all $\mathbf{x} \in \mathcal{C}$ it holds that

$$\left(\frac{f(\mathbf{x}) - f^*}{\mu}\right)^\theta \geq \min_{\mathbf{x}^* \in X^*} \|\mathbf{x}^* - \mathbf{x}\|.$$

In order to derive affine-invariant accelerated convergence rates for FW variants, we rely on the following generalizations of smoothness and Hölderian error bound. As we detail in Section 3, the affine-invariant analysis of the methods presented in this paper relies on these concepts.

The following concepts of curvature and extended curvature were introduced by Jaggi [13] and by Lacoste-Julien and Jaggi [16] respectively.

Definition 2.3 (Curvature). *Let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope and $f: \mathcal{C} \rightarrow \mathbb{R}$ be convex and differentiable in an open set containing \mathcal{C} .*

- (a) *For $L > 0$ we say that (\mathcal{C}, f) has L -curvature if for all $\mathbf{x} \in \mathcal{C}$, $\mathbf{d} \in \mathcal{C} - \mathbf{x}$, and $\eta \in [0, 1]$ it holds that*

$$f(\mathbf{x} + \eta\mathbf{d}) \leq f(\mathbf{x}) + \eta\langle \nabla f(\mathbf{x}), \mathbf{d} \rangle + \frac{L\eta^2}{2}. \quad (2.1)$$

- (b) *For $L > 0$ we say that (\mathcal{C}, f) has extended L -curvature if for all $\mathbf{x} \in \mathcal{C}$, $\mathbf{d} \in \mathcal{C} - \mathcal{C}$, and $\eta \geq 0$ such that $\mathbf{x} + \eta\mathbf{d} \in \mathcal{C}$ it holds that*

$$f(\mathbf{x} + \eta\mathbf{d}) \leq f(\mathbf{x}) + \eta\langle \nabla f(\mathbf{x}), \mathbf{d} \rangle + \frac{L\eta^2}{2}.$$

It is easy to see that the above L -curvature and extended L -curvature properties are affine invariant. It is also evident that (f, \mathcal{C}) has L -curvature whenever (f, \mathcal{C}) has extended L -curvature. In addition, as noted by Jaggi [13] and by Lacoste-Julien and Jaggi [16], if f is L -smooth over \mathcal{C} with respect to a norm $\|\cdot\|$ then (f, \mathcal{C}) has extended $L \cdot \text{diam}(\mathcal{C})^2$ -curvature, where $\text{diam}(\mathcal{C})$ is the following *diameter* of the set \mathcal{C}

$$\text{diam}(\mathcal{C}) = \max_{\mathbf{x}, \mathbf{y} \in \mathcal{C}} \|\mathbf{x} - \mathbf{y}\|.$$

In addition to the above curvature concepts, we will rely on the following extended error bound concept.

Definition 2.4 (Error bound). *Let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope, $f: \mathcal{C} \rightarrow \mathbb{R}$ be convex and differentiable in an open set containing \mathcal{C} , and $\mathfrak{d}: \mathcal{C} \times \mathcal{C} \rightarrow [0, 1]$. For $\mu > 0$ and $\theta \in [0, 1/2]$, we say that (\mathcal{C}, f) satisfies the (μ, θ) -error bound relative to \mathfrak{d} if for all $\mathbf{x} \in \mathcal{C}$ it holds that*

$$\left(\frac{f(\mathbf{x}) - f^*}{\mu}\right)^\theta \geq \min_{\mathbf{x}^* \in X^*} \mathfrak{d}(\mathbf{x}^*, \mathbf{x}).$$

For ease of notation, we will write $\mathfrak{d}(X^, \mathbf{x})$ for $\min_{\mathbf{x}^* \in X^*} \mathfrak{d}(\mathbf{x}^*, \mathbf{x})$.*

It is easy to see that the above error bound property is affine invariant provided the function $\mathfrak{d} : \mathcal{C} \times \mathcal{C} \rightarrow [0, 1]$ is affine-invariant. Section 3 will feature the following instances of affine-invariant distance functions $\mathfrak{d} : \mathcal{C} \times \mathcal{C} \rightarrow [0, 1]$ for each of the algorithms we analyze: the *radial distance* for vanilla Frank-Wolfe, the *vertex distance* for away-step Frank-Wolfe and blended pairwise Frank-Wolfe, and the *face distance* for the in-face Frank-Wolfe.

We will also provide sufficient conditions for the error bound property to hold. To that end, we will rely on the following geometric features of a polytope that were introduced in [18].

Let \mathcal{C} be a polytope. Denote the sets of vertices and nonempty faces of \mathcal{C} by $\text{vert}(\mathcal{C})$ and $\text{faces}(\mathcal{C})$ respectively. Suppose $\|\cdot\|$ is a norm on \mathcal{C} . For $F, G \in \text{faces}(\mathcal{C})$ let

$$\text{dist}(F, G) = \min_{\mathbf{x} \in F, \mathbf{y} \in G} \|\mathbf{x} - \mathbf{y}\|.$$

Definition 2.5 (Inner and outer facial distances). *Let \mathbb{R}^n be endowed with a norm $\|\cdot\|$ and let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope. For $F \in \text{faces}(\mathcal{C})$ the inner facial distance from F is defined as*

$$\Phi(F, \mathcal{C}) := \min_{\substack{G \in \text{faces}(F) \\ G \subsetneq F}} \text{dist}(G, \text{conv}(\text{vert}(\mathcal{C}) \setminus G)),$$

and the outer facial distance from F is defined as

$$\bar{\Phi}(F, \mathcal{C}) := \min_{\substack{G \in \text{faces}(F), H \in \text{faces}(\mathcal{C}) \\ G \cap H = \emptyset}} \text{dist}(G, H).$$

It is easy to see that $\bar{\Phi}(F, \mathcal{C}) \geq \Phi(F, \mathcal{C}) \geq \Phi(\mathcal{C}, \mathcal{C}) > 0$ for all $F \in \text{faces}(\mathcal{C})$. Furthermore, in [18] it was shown that $\Phi(\mathcal{C}, \mathcal{C})$ coincides with the *pyramidal width* of \mathcal{C} introduced by Lacoste-Julien and Jaggi [16].

To derive sublinear convergence rates for FW variants, we will rely on the following lemma from [2, 20]:

Lemma 2.6. *Suppose that $p > 0$ and $(\beta_t)_{t \in \mathbb{N}}, (\sigma_t)_{t \in \mathbb{N}}$ are such that $\beta_t, \sigma_t \geq 0$ and $\beta_{t+1} \leq (1 - \sigma_t \beta_t^p) \beta_t$ for $t \in \mathbb{N}$. Then, for all $t \in \mathbb{N}$, it holds that*

$$\beta_t \leq \left(\beta_0^{-p} + p \sum_{i=0}^{t-1} \sigma_i \right)^{-\frac{1}{p}}.$$

3 Algorithms

In this section, we present affine-invariant convergence rates for the Frank-Wolfe algorithm (FW) in Section 3.1, the two active set variants away-step Frank-Wolfe algorithm (AFW) and blended-pairwise Frank-Wolfe algorithm (BPFW) in Section 3.2, and the in-face Frank-Wolfe algorithm (IFW) in Section 3.3. Although our discussion focuses on the case when $\mathcal{C} \subseteq \mathbb{R}^n$ is a polytope, we note that our developments for both FW and IFW apply in the more general context when $\mathcal{C} \subseteq \mathbb{R}^n$ is a compact convex set.

Algorithm 1 Frank-Wolfe algorithm (FW)

Input: $\mathbf{x}^{(0)} \in \mathcal{C}$.

Output: $\mathbf{x}^{(t)} \in \mathcal{C}$ for $t \in \mathbb{N}$.

for $t \in \mathbb{N}$ **do**

$\mathbf{v}^{(t)} \leftarrow \arg \min_{\mathbf{v} \in \mathcal{C}} \langle \nabla f(\mathbf{x}^{(t)}), \mathbf{v} \rangle$

$\mathbf{x}^{(t+1)} \leftarrow \mathbf{x}^{(t)} + \eta_t(\mathbf{v}^{(t)} - \mathbf{x}^{(t)})$ for some $\eta_t \in [0, 1]$

end for

3.1 Vanilla Frank-Wolfe algorithm

As a preamble to our main developments, we first discuss the conceptually more approachable case of the vanilla Frank-Wolfe algorithm as described in Algorithm 1. This case provides a blueprint for our subsequent discussion of Frank-Wolfe variants.

The *radial distance* introduced by Gutman and Peña [11] is the key ingredient to formalize the affine-invariant convergence rate of FW.

Definition 3.1 (Radial distance). *Let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope and $\mathbf{x}, \mathbf{y} \in \mathcal{C}$. The radial distance $\mathfrak{r}: \mathcal{C} \times \mathcal{C} \rightarrow [0, 1]$ between \mathbf{y} and \mathbf{x} is defined as*

$$\mathfrak{r}(\mathbf{y}, \mathbf{x}) = \min\{\rho \geq 0 \mid \mathbf{y} - \mathbf{x} = \rho(\mathbf{v} - \mathbf{x}) \text{ for some } \mathbf{v} \in \mathcal{C}\}.$$

Note that $\mathfrak{r}(\mathbf{y}, \mathbf{x}) = 0$ if and only if $\mathbf{y} = \mathbf{x}$. As detailed in Theorem 3.2 below, the convergence rate of Algorithm 1 follows from the curvature of (\mathcal{C}, f) and error bound of (\mathcal{C}, f) relative to \mathfrak{r} . Suppose that (\mathcal{C}, f) has L -curvature. We can minimize the left-hand-side of (2.1), a step-size rule referred to as *line-search*:

$$\eta_t = \arg \min_{\eta \in [0, 1]} f(\mathbf{x}^{(t)} + \eta(\mathbf{v}^{(t)} - \mathbf{x}^{(t)})).$$

Alternatively, note that the right-hand-side of (2.1) is a majorant for $f(\mathbf{x} + \eta \mathbf{d})$. Thus, another natural step-size rule minimizes the right-hand-side of (2.1) over $\eta \in [0, 1]$. The resulting majorant-minimization step is referred to as *short-step*:

$$\eta_t = \arg \min_{\eta \in [0, 1]} \left\{ f(\mathbf{x}^{(t)}) + \eta \langle \nabla f(\mathbf{x}^{(t)}), \mathbf{v}^{(t)} - \mathbf{x}^{(t)} \rangle + \frac{L\eta^2}{2} \right\}.$$

For either of the line-search or short-step rules, the following progress bound holds for every iteration of Algorithm 1:

$$f(\mathbf{x}^{(t+1)}) \leq f(\mathbf{x}^{(t)}) - \frac{\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{x}^{(t)} - \mathbf{v}^{(t)} \rangle^2}{2L}. \quad (3.1)$$

Our convergence analysis of FW relies on (3.1), and thus yields identical convergence rates for line-search and short-step.

Theorem 3.2 (FW). *Let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope, let $f: \mathcal{C} \rightarrow \mathbb{R}$ be convex and differentiable in an open set containing \mathcal{C} , and suppose that (\mathcal{C}, f) has L -curvature for some $L > 0$, and satisfies the (μ, θ) -error bound relative to \mathfrak{r} for some $\mu > 0$ and $\theta \in [0, 1/2]$. Then the iterates of Algorithm 1 with line-search or short-step satisfy the following convergence rates.*

- When $\theta = 1/2$, for all $t \in \mathbb{N}$ the following linear convergence rate holds:

$$f(\mathbf{x}^{(t)}) - f^* \leq (f(\mathbf{x}^{(0)}) - f^*) \left(1 - \frac{\mu}{2L}\right)^t. \quad (3.2)$$

- When $\theta \in [0, 1/2)$, for all $t \in \mathbb{N}$ the following sublinear convergence rate holds:

$$f(\mathbf{x}^{(t)}) - f^* \leq \left((f(\mathbf{x}^{(0)}) - f^*)^{(2\theta-1)} + \frac{(1-2\theta)\mu^{2\theta}t}{2L} \right)^{\frac{1}{2\theta-1}}. \quad (3.3)$$

The rates in Theorem 3.2 interpolate between the iconic rate $\mathcal{O}(t^{-1})$ [13] when $\theta = 0$ and the linear rate (3.2) when $\theta = 1/2$. The proof of Theorem 3.2 relies on the following *scaling lemma*.

Lemma 3.3. *Let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope and let $f: \mathcal{C} \rightarrow \mathbb{R}$ be convex and differentiable in an open set containing \mathcal{C} . Then, for all $\mathbf{x} \in \mathcal{C} \setminus X^*$, it holds that*

$$\max_{\mathbf{v} \in \mathcal{C}} \langle \nabla f(\mathbf{x}), \mathbf{x} - \mathbf{v} \rangle \geq \frac{f(\mathbf{x}) - f^*}{\mathfrak{r}(X^*, \mathbf{x})}. \quad (3.4)$$

In particular, if (\mathcal{C}, f) satisfies a (μ, θ) -error bound relative to \mathfrak{r} for some $\mu > 0$ and $\theta \in [0, 1/2]$, then for all $\mathbf{x} \in \mathcal{C}$ it holds that

$$\max_{\mathbf{v} \in \mathcal{C}} \langle \nabla f(\mathbf{x}), \mathbf{x} - \mathbf{v} \rangle \geq \mu^\theta (f(\mathbf{x}) - f^*)^{1-\theta}. \quad (3.5)$$

Proof. Suppose that $\mathbf{x} \in \mathcal{C} \setminus X^*$. Let $\mathbf{x}^* \in X^*$ be such that $\mathfrak{r}(\mathbf{x}^*, \mathbf{x}) = \mathfrak{r}(X^*, \mathbf{x})$. The definition of \mathfrak{r} implies that there exists $\mathbf{w} \in \mathcal{C}$ such that $\mathbf{x}^* - \mathbf{x} = \mathfrak{r}(X^*, \mathbf{x}) \cdot (\mathbf{w} - \mathbf{x})$. Thus,

$$\max_{\mathbf{v} \in \mathcal{C}} \langle \nabla f(\mathbf{x}), \mathbf{x} - \mathbf{v} \rangle \geq \langle \nabla f(\mathbf{x}), \mathbf{x} - \mathbf{w} \rangle = \frac{\langle \nabla f(\mathbf{x}), \mathbf{x} - \mathbf{x}^* \rangle}{\mathfrak{r}(X^*, \mathbf{x})} \geq \frac{f(\mathbf{x}) - f^*}{\mathfrak{r}(X^*, \mathbf{x})},$$

where the last inequality follows from convexity of f . Thus (3.4) follows. To show (3.5), suppose $\mathbf{x} \in \mathcal{C}$. If $\mathbf{x} \in X^*$ then (3.5) trivially holds. Otherwise, when $\mathbf{x} \in \mathcal{C} \setminus X^*$, observe that (3.4) and the (μ, θ) -radial error bound imply that

$$\max_{\mathbf{v} \in \mathcal{C}} \langle \nabla f(\mathbf{x}), \mathbf{x} - \mathbf{v} \rangle \geq \frac{f(\mathbf{x}) - f^*}{\mathfrak{r}(X^*, \mathbf{x})} = \frac{(f(\mathbf{x}) - f^*)^{1-\theta} (f(\mathbf{x}) - f^*)^\theta}{\mathfrak{r}(X^*, \mathbf{x})} \geq \mu^\theta (f(\mathbf{x}) - f^*)^{1-\theta}.$$

□

Proof of Theorem 3.2. The L -curvature and the choice of step-size imply that (3.1) holds. Thus, Lemma 3.3 implies that for each $t \in \mathbb{N}$

$$f(\mathbf{x}^{(t+1)}) \leq f(\mathbf{x}^{(t)}) - \frac{\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{x}^{(t)} - \mathbf{v}^{(t)} \rangle^2}{2L} \leq f(\mathbf{x}^{(t)}) - \frac{\mu^{2\theta} (f(\mathbf{x}^{(t)}) - f^*)^{2(1-\theta)}}{2L}.$$

Hence,

$$f(\mathbf{x}^{(t+1)}) - f^* \leq (f(\mathbf{x}^{(t)}) - f^*) \left(1 - \frac{\mu^{2\theta}}{2L} (f(\mathbf{x}^{(t)}) - f^*)^{1-2\theta} \right).$$

Thus (3.2) readily follows when $\theta = 1/2$. On the other hand, (3.3) follows by applying Lemma 2.6 with $p = 1 - 2\theta$, $\sigma_t = \frac{\mu^{2\theta}}{2L}$, and $\beta_t = f(\mathbf{x}^{(t)}) - f^*$.

□

We next discuss common sufficient conditions for the assumptions of Theorem 3.2 to hold. First, as noted in Section 2, the pair (\mathcal{C}, f) has L -curvature for some $L > 0$ whenever f is smooth with respect to a norm on \mathcal{C} . Let $\text{ri}(\mathcal{C})$ and $\text{rbd}(\mathcal{C})$ denote respectively the relative interior and relative boundary of \mathcal{C} . The following proposition shows that when $X^* \subseteq \text{ri}(\mathcal{C})$, an error bound relative to \mathfrak{r} holds provided f satisfies a Hölderian error bound over \mathcal{C} relative to some norm.

Proposition 3.4 (Radial error bound). *Let \mathbb{R}^n be endowed with a norm $\|\cdot\|$, let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope, and let $f: \mathcal{C} \rightarrow \mathbb{R}$ be convex. Suppose that $X^* \subseteq \text{ri}(\mathcal{C})$ and f satisfies a (μ, θ) -Hölderian error bound over \mathcal{C} with respect to the norm $\|\cdot\|$ for some $\mu > 0$ and $\theta \in [0, 1/2]$. Then (\mathcal{C}, f) satisfies a $(\tilde{\mu}, \theta)$ -error bound relative to \mathfrak{r} for*

$$\tilde{\mu} = \mu \cdot \text{dist}(X^*, \text{rbd}(\mathcal{C}))^{1/\theta}.$$

Proof. The construction of \mathfrak{r} implies that for all $\mathbf{x}^* \in X^*$ and $\mathbf{x} \in \mathcal{C}$

$$\|\mathbf{x}^* - \mathbf{x}\| = \mathfrak{r}(\mathbf{x}^*, \mathbf{x}) \cdot \|\mathbf{x} - \mathbf{u}\| \geq \mathfrak{r}(\mathbf{x}^*, \mathbf{x}) \cdot \|\mathbf{x}^* - \mathbf{u}\|$$

for some $\mathbf{u} \in \text{rbd}(\mathcal{C})$. Therefore for all $\mathbf{x} \in \mathcal{C}$

$$\min_{\mathbf{x}^* \in X^*} \|\mathbf{x}^* - \mathbf{x}\| \geq \mathfrak{r}(X^*, \mathbf{x}) \cdot \text{dist}(X^*, \text{rbd}(\mathcal{C})).$$

Since f satisfies a (μ, θ) -Hölderian error bound over \mathcal{C} with respect to the norm $\|\cdot\|$, it follows that for all $\mathbf{x} \in \mathcal{C}$

$$\left(\frac{f(\mathbf{x}) - f^*}{\mu} \right)^\theta \geq \min_{\mathbf{x}^* \in X^*} \|\mathbf{x}^* - \mathbf{x}\| \geq \mathfrak{r}(X^*, \mathbf{x}) \cdot \text{dist}(X^*, \text{rbd}(\mathcal{C})),$$

and thus

$$\left(\frac{f(\mathbf{x}) - f^*}{\tilde{\mu}} \right)^\theta \geq \mathfrak{r}(X^*, \mathbf{x}).$$

□

The lower bound in Proposition 3.4 is zero when $X^* \cap \text{rbd}(\mathcal{C}) \neq \emptyset$. Thus in this case Theorem 3.2 fails to show linear convergence even when f is smooth and (\mathcal{C}, f) satisfy the $(\mu, 1/2)$ Hölderian error bound relative to some norm in \mathbb{R}^n . This is consistent with the shortcoming of FW discovered and documented by Wolfe [26]. Indeed, Wolfe [26] characterizes instances of (1.1) where \mathcal{C} is a polytope and f is smooth and strongly convex for which FW with line-search or short-step cannot converge faster than $\Omega(t^{-1-\epsilon})$ for any $\epsilon > 0$. This phenomenon, referred to as *Wolfe's lower bound*, was the core motivation behind the study of FW variants to overcome the limitations of the vanilla version.

3.2 Away-step Frank-Wolfe algorithm and blended pairwise Frank-Wolfe algorithm

The main update in the vanilla Frank-Wolfe algorithm can be written as

$$\mathbf{x}^{(t+1)} \leftarrow \mathbf{x}^{(t)} + \eta_t \mathbf{d}^{(t)}$$

Algorithm 2 Away-step Frank-Wolfe algorithm (AFW) and blended pairwise Frank-Wolfe algorithm (BPFW)

Input: $\mathbf{x}^{(0)} \in \text{vert}(\mathcal{C})$, that is, $\mathbf{x}^{(0)} = \mathcal{V}\boldsymbol{\lambda}^{(0)} \in \mathcal{C}$ for some $\boldsymbol{\lambda}^{(0)} \in \Delta(\mathcal{V})$ with $|S(\boldsymbol{\lambda}^{(0)})| = 1$.

Output: $\mathbf{x}^{(t)} = \mathcal{V}\boldsymbol{\lambda}^{(t)} \in \mathcal{C}$ for some $\boldsymbol{\lambda}^{(t)} \in \Delta(\mathcal{V})$ for $t \in \mathbb{N}$.

for $t \in \mathbb{N}$ **do**

 compute $\mathbf{d}_{\text{FW}}^{(t)}, \mathbf{d}_{\text{AFW}}^{(t)}, \mathbf{d}_{\text{BPFW}}^{(t)}$ as in (3.6) for $\mathbf{x} = \mathbf{x}^{(t)}$ and $S = S(\boldsymbol{\lambda}^{(t)})$

 select either $\mathbf{d}^{(t)} \in \{\mathbf{d}_{\text{FW}}^{(t)}, \mathbf{d}_{\text{AFW}}^{(t)}\}$ in AFW or $\mathbf{d} \in \{\mathbf{d}_{\text{FW}}^{(t)}, \mathbf{d}_{\text{BPFW}}^{(t)}\}$ in BPFW that minimizes $\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}^{(t)} \rangle$

 compute $\eta_{t,\max}$ as in (3.7) for $\boldsymbol{\lambda} = \boldsymbol{\lambda}^{(t)}$

$\mathbf{x}^{(t+1)} \leftarrow \mathbf{x}^{(t)} + \eta_t \mathbf{d}^{(t)}$ for some $\eta_t \in [0, \eta_{t,\max}]$

 update $\boldsymbol{\lambda}^{(t+1)} \in \Delta(\mathcal{V})$ as in (3.8) for $\boldsymbol{\lambda} = \boldsymbol{\lambda}^{(t)}$

end for

where $\mathbf{d}^{(t)} = \mathbf{v}^{(t)} - \mathbf{x}^{(t)}$ is the *Frank-Wolfe direction* defined via $\mathbf{v}^{(t)} = \arg \min_{\mathbf{v} \in \mathcal{C}} \langle \nabla f(\mathbf{x}^{(t)}), \mathbf{v} \rangle$. As noted above, FW, limited to FW directions, does not admit fast convergence rates when there are optimal solutions on the relative boundary of \mathcal{C} . To overcome this limitation, FW variants have been studied that admit more flexible choices of $\mathbf{d}^{(t)}$. In this section, we focus on the away-step Frank-Wolfe algorithm (AFW) [1, 16, 26] and the blended pairwise Frank-Wolfe algorithm (BPFW) [22]. To facilitate a richer choice of search directions, both variants maintain and rely on a *vertex representation* of $\mathbf{x}^{(t)}$ as we next explain.

Let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope with vertex set $\mathcal{V} := \text{vert}(\mathcal{C})$ and let

$$\Delta(\mathcal{V}) := \left\{ \boldsymbol{\lambda} = (\lambda_{\mathbf{v}})_{\mathbf{v} \in \mathcal{V}} \mid \lambda_{\mathbf{v}} \geq 0 \text{ for } \mathbf{v} \in \mathcal{V} \text{ and } \sum_{\mathbf{v} \in \mathcal{V}} \lambda_{\mathbf{v}} = 1 \right\}.$$

Any $\mathbf{x} \in \mathcal{C}$ has a *vertex representation* of the form $\mathbf{x} = \sum_{\mathbf{v} \in \mathcal{V}} \lambda_{\mathbf{v}} \mathbf{v}$ for some $\boldsymbol{\lambda} \in \Delta(\mathcal{V})$. For ease of notation, we write $\mathcal{V}\boldsymbol{\lambda}$ for $\sum_{\mathbf{v} \in \mathcal{V}} \lambda_{\mathbf{v}} \mathbf{v}$. For $\boldsymbol{\lambda} \in \Delta(\mathcal{V})$, the *support set* of $\boldsymbol{\lambda}$ is defined as:

$$S(\boldsymbol{\lambda}) = \{\mathbf{v} \in \mathcal{V} \mid \lambda_{\mathbf{v}} > 0\}.$$

For $\mathbf{x} \in \mathcal{C}$, we further define

$$\mathbf{S}(\mathbf{x}) = \{S(\boldsymbol{\lambda}) \mid \mathbf{x} = \mathcal{V}\boldsymbol{\lambda} \text{ for some } \boldsymbol{\lambda} \in \Delta(\mathcal{V})\}.$$

Suppose that $\mathbf{x} \in \mathcal{C}$ and $\mathbf{x} = \mathcal{V}\boldsymbol{\lambda}$ for some $\Delta(\mathcal{V})$. Let $S = S(\boldsymbol{\lambda})$ and define the *away vertex* $\mathbf{a} \in S$, the *local FW vertex* $\mathbf{z} \in S$, and the *global FW vertex* $\mathbf{v} \in \mathcal{V}$ as follows:

$$\mathbf{a} = \arg \max_{\mathbf{y} \in S} \langle \nabla f(\mathbf{x}), \mathbf{y} \rangle, \quad \mathbf{z} = \arg \min_{\mathbf{y} \in S} \langle \nabla f(\mathbf{x}), \mathbf{y} \rangle, \quad \mathbf{v} = \arg \min_{\mathbf{y} \in \mathcal{C}} \langle \nabla f(\mathbf{x}), \mathbf{y} \rangle.$$

The above away, local FW, and global FW vertices in turn yield the following *FW*, *away*, and *blended pairwise* directions:

$$\mathbf{d}_{\text{FW}} = \mathbf{v} - \mathbf{x}, \quad \mathbf{d}_{\text{AFW}} = \mathbf{x} - \mathbf{a}, \quad \mathbf{d}_{\text{BPFW}} = \mathbf{z} - \mathbf{a}. \quad (3.6)$$

Notice that the away and blended pairwise directions reduce the weight of a vertex in the vertex representation of \mathbf{x} and increases the weight of other vertices. In contrast, the FW direction increases the weight of the vertex \mathbf{v} and reduces the weight of all other vertices equally. For $\mathbf{d} \in \{\mathbf{d}_{\text{FW}}, \mathbf{d}_{\text{AFW}}, \mathbf{d}_{\text{BPFW}}\}$, there is a natural maximum step-size η_{max} that ensures that $\mathbf{x} + \eta\mathbf{d} \in \mathcal{C}$ for $\eta \in [0, \eta_{\text{max}}]$. The precise calculation of η_{max} requires an explicit vertex representation of \mathbf{x} of the form $\mathbf{x} = \mathcal{V}\boldsymbol{\lambda}$ such that $S(\boldsymbol{\lambda}) = S$. For each of the directions in (3.6), the maximum step-size η_{max} is as follows, see [16, 22, 26]:

$$\eta_{\text{max}} = \begin{cases} 1, & \text{if } \mathbf{d} = \mathbf{d}_{\text{FW}} = \mathbf{v} - \mathbf{x} \\ \frac{\lambda_{\mathbf{a}}}{1 - \lambda_{\mathbf{a}}}, & \text{if } \mathbf{d} = \mathbf{d}_{\text{AFW}} = \mathbf{x} - \mathbf{a} \\ \lambda_{\mathbf{a}}, & \text{if } \mathbf{d} = \mathbf{d}_{\text{BPFW}} = \mathbf{z} - \mathbf{a}. \end{cases} \quad (3.7)$$

Furthermore, for $\eta \in [0, \eta_{\text{max}}]$ the point $\mathbf{x}^+ = \mathbf{x} + \eta\mathbf{d}$ has a vertex representation $\mathbf{x}^+ = \mathcal{V}\boldsymbol{\lambda}^+$ where

$$\lambda_{\mathbf{s}}^+ = \begin{cases} (1 - \eta)\lambda_{\mathbf{s}} + \eta, & \text{if } \mathbf{d} = \mathbf{d}_{\text{FW}} = \mathbf{v} - \mathbf{x} \text{ and } \mathbf{s} = \mathbf{v} \\ (1 - \eta)\lambda_{\mathbf{s}}, & \text{if } \mathbf{d} = \mathbf{d}_{\text{FW}} = \mathbf{v} - \mathbf{x} \text{ and } \mathbf{s} \neq \mathbf{v} \\ (1 + \eta)\lambda_{\mathbf{s}} - \eta, & \text{if } \mathbf{d} = \mathbf{d}_{\text{AFW}} = \mathbf{x} - \mathbf{a} \text{ and } \mathbf{s} = \mathbf{a} \\ (1 + \eta)\lambda_{\mathbf{s}}, & \text{if } \mathbf{d} = \mathbf{d}_{\text{AFW}} = \mathbf{x} - \mathbf{a} \text{ and } \mathbf{s} \neq \mathbf{a} \\ \lambda_{\mathbf{s}} - \eta, & \text{if } \mathbf{d} = \mathbf{d}_{\text{BPFW}} = \mathbf{z} - \mathbf{a} \text{ and } \mathbf{s} = \mathbf{a} \\ \lambda_{\mathbf{s}} + \eta, & \text{if } \mathbf{d} = \mathbf{d}_{\text{BPFW}} = \mathbf{z} - \mathbf{a} \text{ and } \mathbf{s} = \mathbf{z} \\ \lambda_{\mathbf{s}}, & \text{if } \mathbf{d} = \mathbf{d}_{\text{BPFW}} = \mathbf{z} - \mathbf{a} \text{ and } \mathbf{s} \notin \{\mathbf{a}, \mathbf{z}\}. \end{cases} \quad (3.8)$$

We next establish affine-invariant convergence results for Algorithm 2 analogous to Theorem 3.2. To do so, we rely on the following *vertex distance* in lieu of the radial distance.

Definition 3.5 (Vertex distance). *Let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope and $\mathbf{x}, \mathbf{y} \in \mathcal{C}$. The vertex distance $\mathbf{v}: \mathcal{C} \times \mathcal{C} \rightarrow [0, 1]$ between \mathbf{x} and \mathbf{y} is defined as*

$$\mathbf{v}(\mathbf{y}, \mathbf{x}) = \max_{S \in \mathcal{S}(\mathbf{x})} \min\{\gamma \geq 0 \mid \mathbf{y} - \mathbf{x} = \gamma(\mathbf{v} - \mathbf{u}) \text{ for some } \mathbf{v} \in C \text{ and } \mathbf{u} \in \text{conv}(S)\}.$$

Note that $\mathbf{v}(\mathbf{y}, \mathbf{x}) = 0$ if and only if $\mathbf{x} = \mathbf{y}$. Observe that if $\mathcal{C} \subseteq \mathbb{R}^n$ is a polytope, then for all $\mathbf{x}, \mathbf{y} \in \mathcal{C}$, it holds that $\mathbf{v}(\mathbf{y}, \mathbf{x}) \leq \mathbf{r}(\mathbf{y}, \mathbf{x})$. In particular, if (\mathcal{C}, f) satisfies the (μ, θ) -error bound relative to \mathbf{r} then (\mathcal{C}, f) also satisfies the (μ, θ) -error bound relative to \mathbf{v} and it would typically do so for a larger μ .

As we detail below in Theorem 3.6, the convergence of Algorithm 2 is a consequence of the extended curvature of (\mathcal{C}, f) and the error bound of (\mathcal{C}, f) relative to \mathbf{v} . Suppose that (\mathcal{C}, f) has extended L -curvature. We consider two possible choices of step-size η_t in Line 5 of Algorithm 2. The first one is *line-search*:

$$\eta_t = \arg \min_{\eta \in [0, \eta_{t, \text{max}}]} f(\mathbf{x}^{(t)} + \eta\mathbf{d}^{(t)}).$$

The second one is the *short-step*:

$$\eta_t = \arg \min_{\eta \in [0, \eta_{t, \text{max}}]} \left\{ f(\mathbf{x}^{(t)}) + \eta \langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}^{(t)} \rangle + \frac{L\eta^2}{2} \right\}.$$

It is easy to see that for either the line-search or short-step the following inequality holds when $\eta_t < \eta_{t,\max}$

$$f(\mathbf{x}^{(t+1)}) \leq f(\mathbf{x}^{(t)}) - \frac{\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}^{(t)} \rangle^2}{2L}, \quad (3.9)$$

and the following inequality holds when $\eta_t = \eta_{t,\max} \geq 1$

$$f(\mathbf{x}^{(t+1)}) \leq f(\mathbf{x}^{(t)}) + \frac{\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}^{(t)} \rangle}{2}. \quad (3.10)$$

Theorem 3.6 (AFW and BPFW). *Let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope, let $f: \mathcal{C} \rightarrow \mathbb{R}$ be convex and differentiable in an open set containing \mathcal{C} , and suppose that (\mathcal{C}, f) has extended L -curvature for some $L > 0$, and satisfies the (μ, θ) -error bound relative to \mathbf{v} for some $\mu > 0$ and $\theta \in [0, 1/2]$. Then, the iterates of Algorithm 2 with line-search or short-step satisfy the following convergence rates.*

- When $\theta = 1/2$ the following linear convergence rate holds for all $t \in \mathbb{N}$:

$$f(\mathbf{x}^{(t)}) - f^* \leq (f(\mathbf{x}^{(0)}) - f^*) \left(1 - \min \left\{ \frac{\mu}{8L}, \frac{1}{2} \right\} \right)^{\lceil t/2 \rceil}. \quad (3.11)$$

- When $\theta \in [0, 1/2)$, the following initial linear convergence rate holds for $t \in \{0, 1, \dots, t_0\}$

$$f(\mathbf{x}^{(t)}) - f^* \leq \frac{f(\mathbf{x}^{(0)}) - f^*}{2^{\lceil t/2 \rceil}}, \quad (3.12)$$

where $t_0 \in \mathbb{N}$ is the smallest t such that

$$\frac{\mu^{2\theta} (f(\mathbf{x}^{(t)}) - f^*)^{1-2\theta}}{8L} \leq \frac{1}{2}.$$

And for $t \in \mathbb{N}_{\geq t_0}$, the following sublinear convergence rate holds:

$$f(\mathbf{x}^{(t)}) - f^* \leq \left((f(\mathbf{x}^{(t_0)}) - f^*)^{(2\theta-1)} + \frac{(1-2\theta)\mu^{2\theta}}{8L} \cdot \left| \frac{t - |S(\boldsymbol{\lambda}^{(t_0)})| - t_0 + 1}{2} \right| \right)^{\frac{1}{2\theta-1}}. \quad (3.13)$$

Once again, the rates in Theorem 3.6 interpolate between the iconic rate $\mathcal{O}(t^{-1})$ when $\theta = 0$, to the linear rate (3.11) when $\theta = 1/2$. The proof of Theorem 3.6 relies on the following scaling lemma, resembling Lemma 3.3.

Lemma 3.7. *Let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope, let $f: \mathcal{C} \rightarrow \mathbb{R}$ be convex and differentiable in an open set containing \mathcal{C} . Then, for all $\mathbf{x} \in \mathcal{C} \setminus X^*$ and all $S \in \mathbf{S}(\mathbf{x})$, it holds that*

$$\max_{\mathbf{a} \in S, \mathbf{v} \in \mathcal{C}} \langle \nabla f(\mathbf{x}), \mathbf{a} - \mathbf{v} \rangle \geq \frac{f(\mathbf{x}) - f^*}{\mathbf{v}(X^*, \mathbf{x})}. \quad (3.14)$$

In particular, if (\mathcal{C}, f) satisfies a (μ, θ) -error bound relative to \mathbf{v} for some $\mu > 0$ and $\theta \in [0, 1/2]$, then for all $\mathbf{x} \in \mathcal{C}$

$$\max_{\mathbf{a} \in S, \mathbf{v} \in \mathcal{C}} \langle \nabla f(\mathbf{x}), \mathbf{a} - \mathbf{v} \rangle \geq \mu^\theta (f(\mathbf{x}) - f^*)^{1-\theta}. \quad (3.15)$$

Proof. Let $\gamma = \mathbf{v}(X^*, \mathbf{x}) > 0$ and $\mathbf{x}^* \in X^*$ be such that $\mathbf{v}(\mathbf{x}^*, \mathbf{x}) = \mathbf{v}(X^*, \mathbf{x}) = \gamma > 0$. The definition of \mathbf{v} implies that there exist $\mathbf{w} \in \mathcal{C}$, $\mathbf{u} \in \text{conv}(S)$ such that $\mathbf{x}^* - \mathbf{x} = \gamma(\mathbf{w} - \mathbf{u})$. Thus,

$$\max_{\mathbf{a} \in S, \mathbf{v} \in \mathcal{C}} \langle \nabla f(\mathbf{x}), \mathbf{a} - \mathbf{v} \rangle \geq \langle \nabla f(\mathbf{x}), \mathbf{u} - \mathbf{w} \rangle = \frac{\langle \nabla f(\mathbf{x}), \mathbf{x} - \mathbf{x}^* \rangle}{\gamma} \geq \frac{f(\mathbf{x}) - f^*}{\mathbf{v}(X^*, \mathbf{x})},$$

where the last inequality follows from convexity of f . Thus (3.14) follows. To show (3.15), suppose $\mathbf{x} \in \mathcal{C}$. If $\mathbf{x} \in X^*$ then (3.15) trivially holds. Otherwise, when $\mathbf{x} \in \mathcal{C} \setminus X^*$, observe that (3.14) and the (μ, θ) -error bound relative to \mathbf{v} imply that

$$\max_{\mathbf{a} \in S, \mathbf{v} \in \mathcal{C}} \langle \nabla f(\mathbf{x}), \mathbf{a} - \mathbf{v} \rangle \geq \frac{f(\mathbf{x}) - f^*}{\mathbf{v}(X^*, \mathbf{x})} = \frac{(f(\mathbf{x}) - f^*)^{1-\theta} (f(\mathbf{x}) - f^*)^\theta}{\mathbf{v}(X^*, \mathbf{x})} \geq \mu^\theta (f(\mathbf{x}) - f^*)^{1-\theta}.$$

□

Proof of Theorem 3.6. The choice of $\mathbf{d}^{(t)}$ implies that for $t \in \mathbb{N}_{\geq 0}$

$$\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}^{(t)} \rangle \leq \frac{\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{v}^{(t)} - \mathbf{a}^{(t)} \rangle}{2} < 0 \quad (3.16)$$

and

$$\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}^{(t)} \rangle \leq \langle \nabla f(\mathbf{x}^{(t)}), \mathbf{v}^{(t)} - \mathbf{x}^{(t)} \rangle \leq \langle \nabla f(\mathbf{x}^{(t)}), \mathbf{x}^* - \mathbf{x}^{(t)} \rangle \leq f^* - f(\mathbf{x}). \quad (3.17)$$

We next proceed by considering three possible cases:

1. $\eta_t < \eta_{t, \max}$. In this case, (3.9), (3.16), and (3.15) imply that

$$\begin{aligned} f(\mathbf{x}^{(t+1)}) &\leq f(\mathbf{x}^{(t)}) - \frac{\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}^{(t)} \rangle^2}{2L} \\ &\leq f(\mathbf{x}^{(t)}) - \frac{\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{v}^{(t)} - \mathbf{a}^{(t)} \rangle^2}{8L} \\ &\leq f(\mathbf{x}^{(t)}) - \frac{\mu^{2\theta} (f(\mathbf{x}^{(t)}) - f^*)^{2(1-\theta)}}{8L}. \end{aligned}$$

2. $\eta_t = \eta_{t, \max} \geq 1$. In this case, (3.10) and (3.17) imply that

$$f(\mathbf{x}^{(t+1)}) \leq f(\mathbf{x}^{(t)}) + \frac{\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}^{(t)} \rangle}{2} \leq f(\mathbf{x}^{(t)}) - \frac{f(\mathbf{x}^{(t)}) - f^*}{2}.$$

Thus, if either of Case 1 or Case 2 occurs, we get

$$f(\mathbf{x}^{(t+1)}) - f^* \leq (f(\mathbf{x}^{(t)}) - f^*) \left(1 - \min \left\{ \frac{1}{2}, \frac{\mu^{2\theta} (f(\mathbf{x}^{(t)}) - f^*)^{1-2\theta}}{8L} \right\} \right). \quad (3.18)$$

3. $\eta_t = \eta_{t, \max} < 1$. In this case, the choice of step-size η_t implies that

$$f(\mathbf{x}^{(t+1)}) - f^* \leq f(\mathbf{x}^{(t)}) - f^*.$$

Whenever Case 3 occurs we have $|S(\boldsymbol{\lambda}^{(t+1)})| \leq |S(\boldsymbol{\lambda}^{(t)})| - 1$ and whenever Case 1 or Case 2 occurs we have $|S(\boldsymbol{\lambda}^{(t+1)})| \leq |S(\boldsymbol{\lambda}^{(t)})| + 1$. Since $|S(\boldsymbol{\lambda}^{(t)})| \geq 1$ for all $t \in \mathbb{N}$, it follows that for any consecutive sequence of iterations $\{t_0, t_0 + 1, \dots, t\}$ Case 1 or Case 2 must occur at least $\lceil \frac{t - |S(\boldsymbol{\lambda}^{(t_0)})| - t_0 + 1}{2} \rceil$ times. Furthermore, the sequence $f(\mathbf{x}^{(t)}) - f^*$ is monotonically nonincreasing as this is guaranteed in any of the three cases. The proof is completed as follows. When $\theta = 1/2$, Inequality (3.18) and the monotonicity of $f(\mathbf{x}^{(t)}) - f^*$ imply that for $t \in \mathbb{N}$ whenever Case 1 or Case 2 occurs, it holds that

$$f(\mathbf{x}^{(t+1)}) - f^* \leq (f(\mathbf{x}^{(t)}) - f^*) \left(1 - \min \left\{ \frac{1}{2}, \frac{\mu}{8L} \right\} \right).$$

Thus (3.11) follows. When $\theta \in [0, 1/2)$, Inequality (3.18) and the monotonicity of $f(\mathbf{x}^{(t)}) - f^*$ imply that whenever Case 1 or Case 2 occurs and $t < t_0$, it holds that

$$f(\mathbf{x}^{(t+1)}) - f^* \leq \frac{1}{2}(f(\mathbf{x}^{(t)}) - f^*).$$

Therefore, (3.12) holds for $t = 0, 1, \dots, t_0$. Finally, when $\theta \in [0, 1/2)$, Inequality (3.18) and the monotonicity of $f(\mathbf{x}^{(t)}) - f^*$ imply that whenever Case 1 or Case 2 occurs and $t \geq t_0$, it holds that that

$$f(\mathbf{x}^{(t+1)}) - f^* \leq (f(\mathbf{x}^{(t)}) - f^*) \left(1 - \frac{\mu^{2\theta}}{8L} (f(\mathbf{x}^{(t)}) - f^*)^{1-2\theta} \right).$$

Thus Lemma 2.6 implies (3.13) for $t \geq t_0$. □

We next discuss common sufficient conditions for the the assumptions of Theorem 3.6 to hold. First, as noted in Section 2, the pair (\mathcal{C}, f) has extended L -curvature for some $L > 0$ whenever f is smooth with respect to a norm in \mathcal{C} . Proposition 3.10 below is an analogue of Proposition 3.4. It shows that (\mathcal{C}, f) satisfies an error bound relative to \mathbf{v} provided \mathcal{C} is a polytope and (\mathcal{C}, f) satisfies a Hölderian error bound with respect to some norm in \mathcal{C} .

The proof of Proposition 3.10 relies on the following definition and technical lemma.

Definition 3.8 (Minimal face). *Let $\mathcal{C} \subseteq \mathbb{R}^n$ be a nonempty polytope. For $\mathbf{x} \in \mathcal{C}$ let $F(\mathbf{x}) \in \text{faces}(\mathcal{C})$ denote the minimal face of \mathcal{C} that contains \mathbf{x} , that is, $F(\mathbf{x}) \in \text{faces}(\mathcal{C})$ satisfies $\mathbf{x} \in F(\mathbf{x})$ and $F(\mathbf{x}) \subseteq G$ for all $G \in \text{faces}(\mathcal{C})$ such that $\mathbf{x} \in G$. Similarly, for $\emptyset \neq X \subseteq \mathcal{C}$ let $F(X) \in \text{faces}(\mathcal{C})$ denote the minimal face of \mathcal{C} that contains X .*

Lemma 3.9. *Let \mathbb{R}^n be endowed with a norm $\|\cdot\|$ and let $\mathcal{C} \subseteq \mathbb{R}^n$ be a nonempty polytope. Then for all $\mathbf{x}, \mathbf{y} \in \mathcal{C}$*

$$\|\mathbf{y} - \mathbf{x}\| \geq \mathbf{v}(\mathbf{y}, \mathbf{x}) \cdot \Phi(F(\mathbf{y}), \mathcal{C}). \quad (3.19)$$

For ease of exposition, we defer the proof of Lemma 3.9 to Section 4.

Proposition 3.10 (Vertex distance error bound). *Let \mathbb{R}^n be endowed with a norm $\|\cdot\|$, let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope, and let $f: \mathcal{C} \rightarrow \mathbb{R}$ be convex. Suppose that (\mathcal{C}, f) satisfies a (μ, θ) -Hölderian error bound with respect to $\|\cdot\|$ for some $\mu > 0$ and $\theta \in [0, 1/2]$. Then (\mathcal{C}, f) satisfies a $(\tilde{\mu}, \theta)$ error bound relative to \mathbf{v} for*

$$\tilde{\mu} = \mu \cdot \Phi(F(X^*), \mathcal{C})^{1/\theta}.$$

Algorithm 3 In-face Frank-Wolfe algorithm (IFW)

Input: $\mathbf{x}^{(0)} \in \mathcal{C}$ **Output:** $\mathbf{x}^{(t)} \in \mathcal{C}$ for $t \in \mathbb{N}$.**for** $t \in \mathbb{N}$ **do** compute $\mathbf{d}_{\text{FW}}^{(t)}, \mathbf{d}_{\text{inAFW}}^{(t)}, \mathbf{d}_{\text{inBPFW}}^{(t)}$ as in (3.22) for $\mathbf{x} = \mathbf{x}^{(t)}$ select $\mathbf{d}^{(t)} \in \{\mathbf{d}_{\text{FW}}^{(t)}, \mathbf{d}_{\text{inAFW}}^{(t)}, \mathbf{d}_{\text{inBPFW}}^{(t)}\}$ that minimizes $\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}^{(t)} \rangle$ compute $\eta_{t,\max}$ as in (3.21) for $\mathbf{x} = \mathbf{x}^{(t)}$ and $\mathbf{d} = \mathbf{d}^{(t)}$ $\mathbf{x}^{(t+1)} \leftarrow \mathbf{x}^{(t)} + \eta_t \mathbf{d}^{(t)}$ for some $\eta_t \in [0, \eta_{t,\max}]$ **end for**

Proof. This proof is similar to that of Proposition 3.4. Lemma 3.9 implies that for all $\mathbf{x} \in \mathcal{C}$

$$\min_{\mathbf{x}^* \in X^*} \|\mathbf{x}^* - \mathbf{x}\| \geq \mathbf{v}(X^*, \mathbf{x}) \cdot \Phi(F(X^*), \mathcal{C}).$$

Since f satisfies a (μ, θ) -Hölderian error bound over \mathcal{C} with respect to the norm $\|\cdot\|$, it follows that for all $\mathbf{x} \in \mathcal{C}$

$$\left(\frac{f(\mathbf{x}) - f^*}{\mu} \right)^\theta \geq \min_{\mathbf{x}^* \in X^*} \|\mathbf{x}^* - \mathbf{x}\| \geq \mathbf{v}(X^*, \mathbf{x}) \cdot \Phi(F(X^*), \mathcal{C}),$$

and thus

$$\left(\frac{f(\mathbf{x}) - f^*}{\tilde{\mu}} \right)^\theta \geq \mathbf{v}(X^*, \mathbf{x}).$$

□

Note that in the seminal work of [16], the convergence rate depended on the global inner facial distance instead. The global facial distance generally depends on the dimension of the polytope \mathcal{C} . This has already been demonstrated, for example, for the ℓ_1 -ball [23] and the hypercube [18]. In Section 4.2, we demonstrate that the same does not hold true for the local facial distance. Instead, we will show that the local facial distance depends only on the dimension of the optimal face $F(X^*)$. Our theory is the first result that gives this without additional assumptions such as strict complementarity as in, for example, [7].

3.3 In-face Frank-Wolfe algorithm

We next describe another variant of the Frank-Wolfe algorithm based on *in-face away*, *in-face pairwise*, and *in-face blended pairwise* directions when a suitable *in-face* oracle is available. Variants of the Frank-Wolfe algorithm that incorporates in-face directions have been considered and analyzed by [6] for low-rank matrix completion and by [9] for minimization over simplex-like polytopes. An advantage of in-face variants is that they do not require an explicit vertex representation of the iterates.

Throughout this subsection we assume that $\mathcal{C} \subseteq \mathbb{R}^n$ is a polytope equipped with the following kind of *in-face linear minimization oracle*: for $\mathbf{x} \in \mathcal{C}$ and $\mathbf{g} \in \mathbb{R}^n$ there is

an oracle that computes

$$\arg \min_{\mathbf{y} \in F(\mathbf{x})} \langle \mathbf{g}, \mathbf{y} \rangle, \quad (3.20)$$

where $F(\mathbf{x}) \in \text{faces}(\mathcal{C})$ denotes the minimal face of \mathcal{C} that contains \mathbf{x} as detailed in Definition 3.8. We also assume that \mathcal{C} is equipped with the following kind of *maximum step-size oracle*: for $\mathbf{x} \in \mathcal{C} \subseteq \mathbb{R}^n$ and $\mathbf{d} \in \mathcal{C} - \mathcal{C}$ there is an oracle that computes

$$\max\{\eta \geq 0 \mid \mathbf{x} + \eta \mathbf{d} \in \mathcal{C}\}. \quad (3.21)$$

As it is discussed in [9], the above in-face linear minimization and maximum step-size oracles can be constructed when a linear minimization oracle for \mathcal{C} is available and \mathcal{C} has a polyhedral description of the form

$$\mathcal{C} = \{\mathbf{x} \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{b}, D\mathbf{x} \geq \mathbf{e}\},$$

provided there is an additional available oracle for the mapping $\mathbf{x} \mapsto D\mathbf{x}$. In particular, this is the case when \mathcal{C} has a standard form description

$$\mathcal{C} = \{\mathbf{x} \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}.$$

For $\mathbf{x} \in \mathcal{C}$ define the *in-face away* vertex $\mathbf{a} \in S$ and *in-face local minimizer* vertex $\mathbf{z} \in S$, and *global minimizer* vertex as follows

$$\mathbf{a} = \arg \max_{\mathbf{y} \in F(\mathbf{x})} \langle \nabla f(\mathbf{x}), \mathbf{y} \rangle, \quad \mathbf{z} = \arg \min_{\mathbf{y} \in F(\mathbf{x})} \langle \nabla f(\mathbf{x}), \mathbf{y} \rangle, \quad \mathbf{v} = \arg \min_{\mathbf{y} \in \mathcal{C}} \langle \nabla f(\mathbf{x}), \mathbf{y} \rangle.$$

The objects defined above in turn yield the following *Frank-Wolfe*, *in-face away*, and *in-face blended pairwise directions* for any $\mathbf{x} \in \mathcal{C}$:

$$\mathbf{d}_{\text{FW}} = \mathbf{v} - \mathbf{x}, \quad \mathbf{d}_{\text{inAFW}} = \mathbf{x} - \mathbf{a}, \quad \mathbf{d}_{\text{inBPFW}} = \mathbf{a} - \mathbf{z}. \quad (3.22)$$

We next establish affine-invariant convergence results for Algorithm 3 analogous to Theorem 3.2 and Theorem 3.6. To do so, we rely on the following *face distance*.

Definition 3.11 (Face distance). *Let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope and $\mathbf{x}, \mathbf{y} \in \mathcal{C}$. The face distance $\mathfrak{f}: \mathcal{C} \times \mathcal{C} \rightarrow [0, 1]$ between \mathbf{x} and \mathbf{y} is defined as*

$$\mathfrak{f}(\mathbf{y}, \mathbf{x}) = \min\{\gamma \geq 0 \mid \mathbf{y} - \mathbf{x} = \gamma(\mathbf{v} - \mathbf{u}) \text{ for some } \mathbf{v} \in \mathcal{C} \text{ and } \mathbf{u} \in F(\mathbf{x})\}.$$

Note that $\mathfrak{f}(\mathbf{x}, \mathbf{y}) = 0$ if and only if $\mathbf{x} = \mathbf{y}$. Observe that if $\mathcal{C} \subseteq \mathbb{R}^n$ is a polytope, then for all $\mathbf{x}, \mathbf{y} \in \mathcal{C}$, it holds that $\mathfrak{f}(\mathbf{y}, \mathbf{x}) \leq \mathfrak{v}(\mathbf{y}, \mathbf{x})$. In particular, if (\mathcal{C}, f) satisfies the (μ, θ) -error bound relative to \mathbf{v} then (\mathcal{C}, f) also satisfies the (μ, θ) -error bound relative to \mathfrak{f} and it would typically do so for a larger μ .

As we detail below, a suitable modification of Theorem 3.6 yields the following convergence rate of Algorithm 3.

Theorem 3.12 (IFW). *Let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope equipped with an in-face oracle (3.20) and a maximum step-size oracle (3.21), let $f: \mathcal{C} \rightarrow \mathbb{R}$ be convex and differentiable in an open set containing \mathcal{C} , and suppose that (\mathcal{C}, f) has extended L -curvature for some $L > 0$, and satisfies the (μ, θ) -error bound relative to \mathfrak{f} for some $\mu > 0$ and $\theta \in [0, 1/2]$. Then the iterates of Algorithm 3 with line-search or short-step satisfy the following convergence rates.*

- When $\theta = 1/2$ the following linear convergence rate holds for all $t \in \mathbb{N}$:

$$f(\mathbf{x}^{(t)}) - f^* \leq (f(\mathbf{x}^{(0)}) - f^*) \left(1 - \min \left\{ \frac{\mu}{8L}, \frac{1}{2} \right\}\right)^{\lceil t / \dim(\mathcal{C}) \rceil}. \quad (3.23)$$

- When $\theta \in [0, 1/2)$, the following initial linear convergence rate holds for $t \in \{0, 1, \dots, t_0\}$:

$$f(\mathbf{x}^{(t)}) - f^* \leq \frac{f(\mathbf{x}^{(0)}) - f^*}{2^{\lceil t / \dim(\mathcal{C}) \rceil}}, \quad (3.24)$$

where $t_0 \in \mathbb{N}$ is the smallest t such that

$$\frac{\mu^{2\theta} (f(\mathbf{x}^{(t)}) - f^*)^{1-2\theta}}{8L} \leq \frac{1}{2}.$$

And for $t \in \mathbb{N}_{\geq t_0}$, the following sublinear convergence rate holds:

$$f(\mathbf{x}^{(t)}) - f^* \leq \left((f(\mathbf{x}^{(t_0)}) - f^*)^{(2\theta-1)} + \frac{(1-2\theta)\mu^{2\theta}}{8L} \cdot \left\lceil \frac{t-t_0}{\dim(\mathcal{C})} \right\rceil \right)^{\frac{1}{2\theta-1}} \quad (3.25)$$

The proof of Theorem 3.12 relies on the following analogue of Lemma 3.7. We omit the proof of Lemma 3.13 as it is a straightforward modification of the proof of Lemma 3.7.

Lemma 3.13. *Let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope and let $f: \mathcal{C} \rightarrow \mathbb{R}$ be convex and differentiable in an open set containing \mathcal{C} . Let $\mathbf{x} \in \mathcal{C}$ such that $\mathfrak{f}(X^*, \mathbf{x}) > 0$. Then,*

$$\max_{\mathbf{a} \in F(\mathbf{x}), \mathbf{v} \in \mathcal{C}} \langle \nabla f(\mathbf{x}), \mathbf{a} - \mathbf{v} \rangle \geq \frac{f(\mathbf{x}) - f^*}{\mathfrak{f}(X^*, \mathbf{x})}. \quad (3.26)$$

In particular, if (\mathcal{C}, f) satisfies a (μ, θ) -error bound relative to \mathfrak{f} then for all $\mathbf{x} \in \mathcal{C}$

$$\max_{\mathbf{a} \in F(\mathbf{x}), \mathbf{v} \in \mathcal{C}} \langle \nabla f(\mathbf{x}), \mathbf{a} - \mathbf{v} \rangle \geq \mu^\theta (f(\mathbf{x}) - f^*)^{1-\theta}. \quad (3.27)$$

Proof of Theorem 3.12. This proof is similar that of Theorem 3.6 and thus we are deliberately succinct. The choice of $\mathbf{d}^{(t)}$ implies that for all $t \in \mathbb{N}_{\geq 0}$

$$\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}^{(t)} \rangle \leq \frac{\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{v}^{(t)} - \mathbf{a}^{(t)} \rangle}{2} < 0 \quad (3.28)$$

and

$$\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}^{(t)} \rangle \leq \langle \nabla f(\mathbf{x}^{(t)}), \mathbf{v}^{(t)} - \mathbf{x}^{(t)} \rangle \leq f^* - f(\mathbf{x}). \quad (3.29)$$

We proceed by considering three possible cases:

1. $\eta_t < \eta_{t, \max}$. In this case, (3.9), (3.28), and (3.27) imply that

$$f(\mathbf{x}^{(t+1)}) \leq f(\mathbf{x}^{(t)}) - \frac{\mu^{2\theta} (f(\mathbf{x}^{(t)}) - f^*)^{2(1-\theta)}}{8L}.$$

2. $\eta_t = \eta_{t,\max} \geq 1$. In this case, (3.10) and (3.29) imply that

$$f(\mathbf{x}^{(t+1)}) \leq f(\mathbf{x}^{(t)}) - \frac{f(\mathbf{x}^{(t)}) - f^*}{2}.$$

Thus if either of Case 1 or Case 2 occurs, then we get

$$f(\mathbf{x}^{(t+1)}) - f^* \leq \left(1 - \min \left\{ \frac{1}{2}, \frac{\mu^{2\theta} (f(\mathbf{x}^{(t)}) - f^*)^{1-2\theta}}{8L} \right\}\right) (f(\mathbf{x}^{(t)}) - f^*).$$

3. $\eta_t = \eta_{t,\max} < 1$. In this case the choice of step-size η_t implies that

$$f(\mathbf{x}^{(t+1)}) - f^* \leq f(\mathbf{x}^{(t)}) - f^*.$$

Whenever Case 3 occurs we have $\dim(F(\mathbf{x}^{(t+1)})) \leq \dim(F(\mathbf{x}^{(t)})) - 1$ and whenever Case 1 or Case 2 occurs we have $\dim(F(\mathbf{x}^{(t+1)})) \leq \dim(\mathcal{C})$. Since $\dim(F(\mathbf{x}^{(t)})) \geq 0$ for all $t \in \mathbb{N}$, Case 3 can occur at most $\dim(\mathcal{C})$ times per each occurrence of one of the first two cases. Thus for any consecutive sequence of iterations $\{t_0, t_0 + 1, \dots, t\}$ Case 1 or Case 2 must occur at least $\lceil \frac{t-t_0}{\dim(\mathcal{C})} \rceil$ times. The rest of the proof is a straightforward modification of the last part of the proof of Theorem 3.6. \square

Proposition 3.15 below is an analogue of Proposition 3.4 and 3.10. It shows that (\mathcal{C}, f) satisfies an error bound relative to \mathfrak{f} provided \mathcal{C} is a polytope and (\mathcal{C}, f) satisfies a Hölderian error bound with respect to some norm in \mathcal{C} . We omit the proof of Proposition 3.15 as it is a straightforward modification of the proof of Proposition 3.10 via the following analogue of Lemma 3.9.

Lemma 3.14. *Let \mathbb{R}^n be endowed with a norm $\|\cdot\|$ and let $\mathcal{C} \subseteq \mathbb{R}^n$ be a nonempty polytope. Then for all $\mathbf{x}, \mathbf{y} \in \mathcal{C}$*

$$\|\mathbf{y} - \mathbf{x}\| \geq \mathfrak{f}(\mathbf{y}, \mathbf{x}) \cdot \bar{\Phi}(F(\mathbf{y}), \mathcal{C}). \quad (3.30)$$

Again for ease of exposition, we defer the proof of Lemma 3.14 to Section 4.

Proposition 3.15 (Face distance error bound). *Let \mathbb{R}^n be endowed with a norm $\|\cdot\|$, let $\mathcal{C} \subseteq \mathbb{R}^n$ be a nonempty polytope, and let $f: \mathcal{C} \rightarrow \mathbb{R}$ be convex. Suppose that (\mathcal{C}, f) satisfies a (μ, θ) -Hölderian error bound with respect to $\|\cdot\|$ for some $\mu > 0$ and $\theta \in [0, 1/2]$. Then (\mathcal{C}, f) satisfies a $(\tilde{\mu}, \theta)$ error bound relative to \mathfrak{f} for*

$$\tilde{\mu} = \mu \cdot \bar{\Phi}(F(X^*), \mathcal{C})^{1/\theta}.$$

3.3.1 In-face FW for polytopes in standard form

So far, we discussed IFW for general polytopes. Notably, when a (μ, θ) -error bound holds with $\theta = 1/2$, we derived the convergence guarantee $\mathcal{O}(\exp(-\frac{t}{\dim(\mathcal{C})}))$. In the theorem below, when \mathcal{C} is a polytope in standard form, that is,

$$\mathcal{C} = \{\mathbf{x} \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}, \quad (3.31)$$

where $A \in \mathbb{R}^{m \times n}$, $\text{rank}(A) = m$, and $\mathbf{b} \in \mathbb{R}^m$, we derive an alternate convergence rate of order $\mathcal{O}(\exp(-\frac{t}{m}))$ which is sharper when $m < \dim(\mathcal{C})$.

Theorem 3.16 (IFW for polytopes in standard form). *Let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope of the form (3.31), let $f: \mathcal{C} \rightarrow \mathbb{R}$ be convex and differentiable in an open set containing \mathcal{C} , and suppose that (\mathcal{C}, f) has extended L -curvature for some $L > 0$, and satisfies the (μ, θ) -error bound with $\theta \in [0, 1/2]$ relative to \mathfrak{f} for some $\mu > 0$ and $\theta \in [0, 1/2]$. Suppose Algorithm 3 starts with $\mathbf{x}^{(0)} \in \text{vert}(\mathcal{C})$. Then the iterates of Algorithm 3 with line-search or short-step satisfy the following convergence rates.*

- When $\theta = 1/2$ the following linear convergence rate holds for all $t \in \mathbb{N}$:

$$f(\mathbf{x}^{(t)}) - f^* \leq (f(\mathbf{x}^{(0)}) - f^*) \left(1 - \min \left\{ \frac{\mu}{8L}, \frac{1}{2} \right\}\right)^{\lceil t/m \rceil}. \quad (3.32)$$

- When $\theta \in [0, 1/2)$ the following initial linear convergence rate holds for $t \in \{0, 1, \dots, t_0\}$:

$$f(\mathbf{x}^{(t)}) - f^* \leq \frac{f(\mathbf{x}^{(0)}) - f^*}{2^{\lceil t/m \rceil}}, \quad (3.33)$$

where $t_0 \in \mathbb{N}$ is the smallest t such that

$$\frac{\mu^{2\theta} (f(\mathbf{x}^{(t)}) - f^*)^{1-2\theta}}{8L} \leq \frac{1}{2}.$$

And for $t \in \mathbb{N}_{\geq t_0}$, the following sublinear convergence rate holds:

$$f(\mathbf{x}^{(t)}) - f^* \leq \left((f(\mathbf{x}^{(t_0)}) - f^*)^{(2\theta-1)} + \frac{(1-2\theta)\mu^{2\theta}}{8L} \cdot \left\lceil \frac{t - \dim(F(\mathbf{x}^{(t_0)})) - t_0}{m} \right\rceil \right)^{\frac{1}{2\theta-1}}. \quad (3.34)$$

Proof. The proof deviates from the proof of Theorem 3.12 only in the analysis of the number of drop steps. We focus on this aspect.

Whenever Case 3 occurs we have $\dim(F(\mathbf{x}^{(t+1)})) \leq \dim(F(\mathbf{x}^{(t)})) - 1$. On the other hand, whenever Case 1 or Case 2 occurs, the standard form description of \mathcal{C} implies that $\dim(F(\mathbf{x}^{(t+1)})) \leq \dim(F(\mathbf{x}^{(t)})) + m$ because the reduction in the number of binding inequalities from $F(\mathbf{x}^{(t)})$ to $F(\mathbf{x}^{(t+1)})$ is at most m . Since $\dim(F(\mathbf{x}^{(t)})) \geq 0$ for all $t \in \mathbb{N}$, it follows that for any consecutive sequence of iterations $\{t_0, t_0 + 1, \dots, t\}$ Case 1 or Case 2 must occur at least $\lceil \frac{t - \dim(F(\mathbf{x}^{(t_0)})) - t_0}{m} \rceil$ times. The rest of the proof is analogous to the proof of Theorem 3.12. \square

3.3.2 In-face pairwise Frank-Wolfe for simplex-like polytopes

This section features an interesting connection with [9] where Garber and Meshi introduced a decomposition-invariant pairwise Frank-Wolfe algorithm for *simplex-like polytopes* whose convergence properties depends on the sparsity of the optimal solution set.

Definition 3.17 (Simplex-like polytope). *A polytope $\mathcal{C} = \{\mathbf{x} \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{b}, \mathbf{x} \geq 0\}$ whose vertices are contained in $\{0, 1\}^n$, where $A \in \mathbb{R}^{m \times n}$ and $\mathbf{b} \in \mathbb{R}^m$, is called a simplex-like polytope.*

Algorithm 4 Frank-Wolfe algorithm with in-face pairwise directions for simplex-like polytopes (FWIPW)

Input: $\mathbf{x}^{(0)} \in \text{vert}(\mathcal{C})$

Output: $\mathbf{x}^{(t)} \in \mathcal{C}$ for $t \in \mathbb{N}$.

$\eta_{-1} \leftarrow 1$

for $t \in \mathbb{N}$ **do**

 compute $\mathbf{d}_{\text{PW}}^{(t)}$ as in (3.35) for $\mathbf{x} = \mathbf{x}^{(t)}$

$\gamma_t \leftarrow -\frac{\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}_{\text{PW}}^{(t)} \rangle}{L}$

$\eta_t \leftarrow \max\{2^{-k_t} \mid k_t \in \mathbb{N}, 2^{-k_t} \leq \min\{\gamma_t, \eta_{t-1}\}\}$

$\mathbf{x}^{(t+1)} \leftarrow \mathbf{x}^{(t)} + \eta_t \mathbf{d}_{\text{PW}}^{(t)}$

end for

We present a FW variant that is similar to the decomposition-invariant pairwise Frank-Wolfe algorithm of Garber and Meshi [9] in Algorithm 4. Like Algorithm 3, Algorithm 4 relies on the in-face linear minimization oracle (3.20) and the maximum step-size oracle (3.21). Unlike Algorithm 3, Algorithm 4 uses the *pairwise direction*:

$$\mathbf{d}_{\text{PW}} = \mathbf{v} - \mathbf{a}, \quad (3.35)$$

where $\mathbf{a} = \arg \max_{\mathbf{y} \in F(\mathbf{x})} \langle \nabla f(\mathbf{x}), \mathbf{y} \rangle$ and $\mathbf{v} = \arg \min_{\mathbf{y} \in \mathcal{C}} \langle \nabla f(\mathbf{x}), \mathbf{y} \rangle$. Algorithm 4 also uses the following target short-step rule

$$-\frac{\langle \nabla f(\mathbf{x}), \mathbf{d}_{\text{PW}} \rangle}{L} = \arg \min_{\gamma \geq 0} \left\{ f(\mathbf{x}) + \gamma \langle \nabla f(\mathbf{x}), \mathbf{d}_{\text{PW}} \rangle + \frac{L\gamma^2}{2} \right\}.$$

We next establish affine-invariant convergence results for Algorithm 4 analogous to but stronger than those in Theorems 3.6, 3.12, and 3.16.

Theorem 3.18 (FWIPW). *Let $\mathcal{C} \subseteq \mathbb{R}^n$ be a simplex-like polytope, let $f: \mathcal{C} \rightarrow \mathbb{R}$ be convex and differentiable in an open set containing \mathcal{C} , and suppose that (\mathcal{C}, f) has extended L -curvature for some $L > 0$, and satisfies the (μ, θ) -error bound relative to \mathfrak{f} for some $\mu > 0$ and $\theta \in [0, 1/2]$. Then the iterates of Algorithm 4 satisfy the following initial linear convergence rate for $t \in \{0, 1, \dots, t_0\}$:*

$$f(\mathbf{x}^{(t)}) - f^* \leq \frac{f(\mathbf{x}^{(0)}) - f^*}{2^t}, \quad (3.36)$$

where $t_0 \in \mathbb{N}$ is the smallest t such that

$$\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}_{\text{PW}}^{(t)} \rangle < L.$$

Then the following convergence rates hold.

- When $\theta = 1/2$ the following linear convergence rate holds for all $t \in \mathbb{N}_{\geq t_0}$:

$$f(\mathbf{x}^{(t)}) - f^* \leq (f(\mathbf{x}^{(t_0)}) - f^*) \left(1 - \frac{\mu}{4L}\right)^{t-t_0}.$$

- When $\theta \in [0, 1/2)$ the following sublinear convergence rate holds for all $t \in \mathbb{N}_{\geq t_0}$:

$$f(\mathbf{x}^{(t)}) - f^* \leq \left((f(\mathbf{x}^{(t_0)}) - f^*)^{(2\theta-1)} + \frac{(1-2\theta)\mu^{2\theta}(t-t_0)}{4L} \right)^{\frac{1}{2\theta-1}}.$$

Proof. We first prove that the iterates are feasible, that is, $\mathbf{x}^{(t)} \in \mathcal{C}$ for all $t \in \mathbb{N}$. We do so via the following two claims that are essentially identical to Observation 2 and Lemma 1 in [9].

Claim 1: Suppose that for some iteration $t \in \mathbb{N}$ of Algorithm 4, it holds that $\mathbf{x}^{(t)} \in \mathcal{C}$ and $x_i^{(t)} \geq \eta_t$ whenever $x_i^{(t)} > 0$. Then $\mathbf{x}^{(t+1)} \in \mathcal{C}$.

Proof of Claim 1: Since $\mathbf{a}^{(t)} = \arg \min_{\mathbf{y} \in F(\mathbf{x}^{(t)})} \langle \nabla f(\mathbf{x}^{(t)}), \mathbf{y} \rangle$, it follows that $a_i^{(t)} = 0$ whenever $x_i^{(t)} = 0$. Therefore since $\mathbf{a}^{(t)} \in \{0, 1\}^n$, our assumption on the positive entries of $\mathbf{x}^{(t)}$ implies that $\mathbf{x}^{(t)} - \eta_t \mathbf{a}^{(t)} \geq \mathbf{0}$. Since $\mathbf{v}^{(t)} \in \mathcal{C}$, in particular $\mathbf{v}^{(t)} \geq \mathbf{0}$ and thus $\mathbf{x}^{(t+1)} = \mathbf{x}^{(t)} - \eta_t \mathbf{a}^{(t)} + \eta_t \mathbf{v}^{(t)} \geq \mathbf{0}$. Furthermore, since $\mathbf{x}^{(t)}, \mathbf{a}^{(t)}, \mathbf{v}^{(t)} \in \mathcal{C}$, in particular $A\mathbf{x}^{(t)} = A\mathbf{a}^{(t)} = A\mathbf{v}^{(t)} = \mathbf{b}$ and hence $A\mathbf{x}^{(t+1)} = \mathbf{b}$ as well. Therefore, $\mathbf{x}^{(t+1)} \in \mathcal{C}$.

Claim 2: The iterates of Algorithm 4 are feasible, that is, $\mathbf{x}^{(t)} \in \mathcal{C}$ for all $t \in \mathbb{N}$.

Proof of Claim 2: We prove by induction that for all $t \in \mathbb{N}$, there exists $\boldsymbol{\alpha}^{(t)} \in \mathbb{N}^n$ such that $\mathbf{x}^{(t)} = \eta_t \boldsymbol{\alpha}^{(t)}$. The feasibility of the iterates then follows from Claim 1. For the base case $t = 0$ observe that since $\mathbf{x}^{(0)}$ is a vertex of \mathcal{C} , it holds that $x_i^{(0)} \in \{0, 1\}$ for each $i \in [n]$. Since $\eta_0 = 2^{-k_0}$ for some $k_0 \in \mathbb{N}$, it follows that $\mathbf{x}^{(0)} = \eta_0 \boldsymbol{\alpha}^{(0)}$ for $\boldsymbol{\alpha}^{(0)} = 2^{k_0} \mathbf{x}^{(0)} \in \mathbb{N}^n$. Suppose that the induction hypothesis holds for some $t \in \mathbb{N}$. Since subtracting $\eta_t \mathbf{a}^{(t)}$ from $\mathbf{x}^{(t)}$ can only decrease positive entries in $\mathbf{x}^{(t)}$, see the proof of Claim 1, and since both $\mathbf{a}^{(t)}$ and $\mathbf{v}^{(t)}$ are vertices of \mathcal{C} , the i th entry of $\mathbf{x}^{(t+1)}$ is given by

$$x_i^{(t+1)} = \eta_t \cdot \begin{cases} \alpha_i^{(t)}, & \text{if } \alpha_i^{(t)} \geq 1 \text{ and } a_i^{(t)} = v_i^{(t)} \in \{0, 1\} \\ \alpha_i^{(t)} - 1, & \text{if } \alpha_i^{(t)} \geq 1 \text{ and } a_i^{(t)} = 1 \text{ and } v_i^{(t)} = 0 \\ \alpha_i^{(t)} + 1, & \text{if } \alpha_i^{(t)} = 0 \text{ and } v_i^{(t)} = 1. \end{cases}$$

Since $\eta_{t+1} = 2^{-k_{t+1}} \leq 2^{-k_t} = \eta_t$ for some $k_{t+1} \in \mathbb{N}$, it follows that $\frac{\eta_t}{\eta_{t+1}} \in \mathbb{N}_{\geq 1}$. Thus, $\boldsymbol{\alpha}^{(t+1)} = \frac{\eta_t}{\eta_{t+1}} \boldsymbol{\alpha}^{(t)} \in \mathbb{N}^n$ is such that $\mathbf{x}^{(t+1)} = \eta_{t+1} \boldsymbol{\alpha}^{(t+1)}$, proving Claim 2.

It remains to prove the convergence rates. To do so we rely on the following result.

Claim 3: For all $t \in \mathbb{N}_{\geq t_0}$, it holds that

$$-\frac{\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}_{\text{PW}}^{(t)} \rangle}{L} \geq \eta_t \geq \frac{\mu^\theta (f(\mathbf{x}^{(t)}) - f^*)^{1-\theta}}{2L}, \quad (3.37)$$

which also implies that the suboptimality gap is nonincreasing, that is, $f(\mathbf{x}^{(t+1)}) \leq f(\mathbf{x}^{(t)})$ for all $t \in \mathbb{N}_{\geq t_0}$.

Proof of Claim 3: The upper bound in (3.37) readily follows from the choice of η_t :

$$\eta_t \leq \gamma_t = -\frac{\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}_{\text{PW}}^{(t)} \rangle}{L}.$$

Furthermore, since (\mathcal{C}, f) has extended L -curvature it also follows that

$$\begin{aligned}
f(\mathbf{x}^{(t+1)}) &= f(\mathbf{x}^{(t)} + \eta_t \mathbf{d}_{\text{PW}}^{(t)}) \\
&\leq f(\mathbf{x}^{(t)}) + \eta_t \langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}_{\text{PW}}^{(t)} \rangle + \frac{L\eta_t^2}{2} \\
&\leq f(\mathbf{x}^{(t)}) - \frac{L\eta_t^2}{2} \\
&\leq f(\mathbf{x}^{(t)}).
\end{aligned}$$

Thus the primal optimality gap is nonincreasing.

We prove the lower bound in (3.37) by induction. The choice of t_0 and (3.27) imply that

$$\eta_{t_0} \geq \frac{\gamma_{t_0}}{2} = -\frac{\langle \nabla f(\mathbf{x}^{(t_0)}), \mathbf{d}_{\text{PW}}^{(t_0)} \rangle}{2L} \geq \frac{\mu^\theta (f(\mathbf{x}^{(t_0)}) - f^*)^{1-\theta}}{2L}.$$

Suppose that the lower bound holds for $t \in \mathbb{N}$. In case $\gamma_{t+1} \leq \eta_t$, Claim 3 follows from (3.27):

$$\eta_{t+1} \geq \frac{\gamma_{t+1}}{2} = -\frac{\langle \nabla f(\mathbf{x}^{(t+1)}), \mathbf{d}_{\text{PW}}^{(t+1)} \rangle}{2L} \geq \frac{\mu^\theta (f(\mathbf{x}^{(t+1)}) - f^*)^{1-\theta}}{2L}.$$

In case $\gamma_{t+1} \geq \eta_t$, Claim 3 follows from the induction hypothesis and the the nonincreasing suboptimality gap:

$$\eta_{t+1} = \eta_t \geq \frac{\mu^\theta (f(\mathbf{x}^{(t)}) - f^*)^{1-\theta}}{2L} \geq \frac{\mu^\theta (f(\mathbf{x}^{(t+1)}) - f^*)^{1-\theta}}{2L}.$$

Finally, we derive the convergence guarantee. For $t \in \{0, \dots, t_0\}$ we have $\eta_t = 1 \leq -\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}_{\text{PW}}^{(t)} \rangle$. Thus the extended L -curvature of (\mathcal{C}, f) implies that

$$f(\mathbf{x}^{(t+1)}) \leq f(\mathbf{x}^{(t)}) + \frac{\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}_{\text{PW}}^{(t)} \rangle}{2} \leq f(\mathbf{x}^{(t)}) - \frac{f(\mathbf{x}^{(t)}) - f^*}{2}.$$

Thus (3.36) follows.

On the other hand, Claim 3, L -curvature, and (3.27) imply that for all $t \in \mathbb{N}_{\geq t_0}$ it holds that

$$\begin{aligned}
f(\mathbf{x}^{(t+1)}) &\leq f(\mathbf{x}^{(t)}) + \eta_t \langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}_{\text{PW}}^{(t)} \rangle + \frac{L\eta_t^2}{2} \\
&\leq f(\mathbf{x}^{(t)}) + \eta_t \left(\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}_{\text{PW}}^{(t)} \rangle + \frac{L\eta_t}{2} \right) \\
&\leq f(\mathbf{x}^{(t)}) + \eta_t \frac{\langle \nabla f(\mathbf{x}^{(t)}), \mathbf{d}_{\text{PW}}^{(t)} \rangle}{2} \\
&\leq f(\mathbf{x}^{(t)}) - \frac{\mu^{2\theta} (f(\mathbf{x}^{(t)}) - f^*)^{2(1-\theta)}}{4L}.
\end{aligned}$$

The theorem follows from repeated application of the above inequality when $\theta = 1/2$ and from Lemma 2.6 when $\theta \in [0, 1/2)$. \square

We conclude this section with the following result analogous to Proposition 3.4, Proposition 3.10, and Proposition 3.15 but specialized to the Euclidean norm. We rely on the following notation and terminology from [9]. Suppose $\mathcal{C} \subseteq \mathbb{R}^n$ be a simplex-like polytope and $\emptyset \neq X \subseteq \mathcal{C}$. Let $\text{card}(X)$ denote the cardinality of X , that is, the number of positive entries of the most dense point in X .

Proposition 3.19. *Let \mathbb{R}^n be endowed with the Euclidean norm $\|\cdot\|_2$, let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope, and let $f: \mathcal{C} \rightarrow \mathbb{R}$ be convex. Suppose that (\mathcal{C}, f) satisfies a (μ, θ) -Hölderian error bound with respect to $\|\cdot\|_2$ for some $\mu > 0$ and $\theta \in [0, 1/2]$. Then (\mathcal{C}, f) satisfies a $(\tilde{\mu}, \theta)$ error bound relative to \mathfrak{f} for*

$$\tilde{\mu} = \max \left\{ \frac{\mu}{\text{card}(X^*)^{1/(2\theta)}}, \frac{\mu}{(n - \text{card}(X^*))^{1/(2\theta)}} \right\}.$$

The bound in Proposition 3.19 is in the same spirit as the bound stated Lemma 2 in [9]. A difference is that the bound in Proposition 3.19 is sharper when the most dense point in X^* has more than $n/2$ non-zero components. We defer the proof of Proposition 3.19 to Section 4.

4 Facial geometry

In this section, we develop some geometric properties of the inner and outer facial distances. More precisely, we give proofs of Lemma 3.9 and Lemma 3.14 and give generic lower bounds on the inner and outer facial distances. In particular, we give a proof of Proposition 3.19 as an immediate consequence of more general bounds.

4.1 Proofs of Lemma 3.9 and Lemma 3.14

Proof of Lemma 3.9. Suppose that $\mathbf{x}, \mathbf{y} \in \mathcal{C}$ with $\mathbf{y} \neq \mathbf{x}$ as otherwise (3.19) trivially holds. Consider the following modified version of $\mathfrak{v}(\mathbf{y}, \mathbf{x})$:

$$\tilde{\mathfrak{v}}(\mathbf{y}, \mathbf{x}) := \max_{S \in \mathfrak{S}(\mathbf{x})} \min \{ \gamma \geq 0 \mid \mathbf{y} - \mathbf{x} = \gamma(\mathbf{v} - \mathbf{u}) \text{ for some } \mathbf{v} \in F(\mathbf{y}) \text{ and } \mathbf{u} \in \text{conv}(S) \}. \quad (4.1)$$

Suppose that $S, \mathbf{v}, \mathbf{u}$ attain the value $\gamma = \tilde{\mathfrak{v}}(\mathbf{y}, \mathbf{x})$ in (4.2) and S is a minimal set with respect to inclusion.

We claim that the construction of $\tilde{\mathfrak{v}}(\mathbf{y}, \mathbf{x})$ and the choice of \mathbf{u}, \mathbf{v} imply that $\mathbf{u} \in \text{conv}(\text{vert}(\mathcal{C}) \setminus F(\mathbf{v}))$. To show this claim it suffices to show that $S \cap \text{vert}(F(\mathbf{v})) = \emptyset$. To do so, we proceed by contradiction. Observe that the minimality of S implies that $\mathbf{u} = \sum_{\mathbf{s} \in S} \lambda_{\mathbf{s}} \mathbf{s}$ for some $\lambda_{\mathbf{s}} > 0, \mathbf{s} \in S$ with $\sum_{\mathbf{s} \in S} \lambda_{\mathbf{s}} = 1$. If $S \cap \text{vert}(F(\mathbf{v})) \neq \emptyset$ then for some $\mathbf{s} \in S \cap \text{vert}(F(\mathbf{v}))$ and $\epsilon > 0$ sufficiently small we have both $\mathbf{u}' := (1 + \epsilon)\mathbf{u} - \epsilon\mathbf{s} \in \text{conv}(S)$ because all $\lambda_{\mathbf{s}} > 0$ and $\mathbf{v}' := (1 + \epsilon)\mathbf{v} - \epsilon\mathbf{s} \in F(\mathbf{v})$ because $\mathbf{v} \in \text{ri}(F(\mathbf{v}))$. Hence we get

$$\mathbf{y} - \mathbf{x} = \frac{\gamma}{1 + \epsilon} (\mathbf{v}' - \mathbf{u}')$$

with $\gamma/(1 + \epsilon) < \gamma$. This contradicts the choice of $S, \mathbf{v}, \mathbf{u}$. Thus, $S \cap \text{vert}(F(\mathbf{v})) = \emptyset$.

To finish, observe that $\gamma = \tilde{\mathbf{v}}(\mathbf{y}, \mathbf{x}) \geq \mathbf{v}(\mathbf{y}, \mathbf{x})$ and thus

$$\begin{aligned} \|\mathbf{y} - \mathbf{x}\| &= \gamma \|\mathbf{v} - \mathbf{u}\| \\ &\geq \mathbf{v}(\mathbf{y}, \mathbf{x}) \cdot \text{dist}(F(\mathbf{v}), \text{conv}(\text{vert}(\mathcal{C}) \setminus F(\mathbf{v}))) \\ &\geq \mathbf{v}(\mathbf{y}, \mathbf{x}) \cdot \Phi(F(\mathbf{y}), \mathcal{C}). \end{aligned}$$

The second to last step holds because $\emptyset \neq F(\mathbf{v}) \in \text{faces}(F(\mathbf{y}))$ and $\mathbf{u} \in \text{conv}(\text{vert}(\mathcal{C}) \setminus F(\mathbf{v}))$. Since this holds for any $\mathbf{x}, \mathbf{y} \in \mathcal{C}$ with $\mathbf{y} \neq \mathbf{x}$, the bound (3.19) follows. \square

Proof of Lemma 3.14. This proof is similar, but simpler, to that of Lemma 3.9. Suppose that $\mathbf{x}, \mathbf{y} \in \mathcal{C}$ with $\mathbf{y} \neq \mathbf{x}$ as otherwise (3.30) trivially holds. Consider the following modified version of $\mathbf{f}(\mathbf{y}, \mathbf{x})$:

$$\bar{\mathbf{f}}(\mathbf{y}, \mathbf{x}) := \min\{\gamma \geq 0 \mid \mathbf{y} - \mathbf{x} = \gamma(\mathbf{v} - \mathbf{u}) \text{ for some } \mathbf{v} \in F(\mathbf{y}) \text{ and } \mathbf{u} \in F(\mathbf{x})\}, \quad (4.2)$$

and suppose \mathbf{v}, \mathbf{u} attain the value $\gamma = \bar{\mathbf{f}}(\mathbf{y}, \mathbf{x})$, that is, $\mathbf{v} \in F(\mathbf{y}), \mathbf{u} \in F(\mathbf{x})$ and $\mathbf{y} - \mathbf{x} = \gamma(\mathbf{v} - \mathbf{u})$. The construction of $\bar{\mathbf{f}}(\mathbf{y}, \mathbf{x})$ and the choice of $\gamma, \mathbf{u}, \mathbf{v}$ imply that $F(\mathbf{u}) \cap F(\mathbf{v}) = \emptyset$. To finish, observe that $\gamma = \bar{\mathbf{f}}(\mathbf{y}, \mathbf{x}) \geq \mathbf{f}(\mathbf{y}, \mathbf{x})$ and thus

$$\begin{aligned} \|\mathbf{y} - \mathbf{x}\| &= \gamma \|\mathbf{v} - \mathbf{u}\| \\ &\geq \mathbf{f}(\mathbf{y}, \mathbf{x}) \cdot \text{dist}(F(\mathbf{v}), F(\mathbf{u})) \\ &\geq \mathbf{f}(\mathbf{y}, \mathbf{x}) \cdot \bar{\Phi}(F(\mathbf{y}), \mathcal{C}). \end{aligned}$$

The last step holds because both $F(\mathbf{v}) \in \text{faces}(F(\mathbf{y}))$ and $F(\mathbf{u}) \in \text{faces}(F(\mathbf{x})) \subseteq \text{faces}(\mathcal{C})$ are evidently nonempty, and $F(\mathbf{u}) \cap F(\mathbf{v}) = \emptyset$. Since this holds for any $\mathbf{x}, \mathbf{y} \in \mathcal{C}$ with $\mathbf{y} \neq \mathbf{x}$, the bound (3.30) follows. \square

4.2 Lower bound on the inner and outer facial distances

We next provide lower bounds on the inner and outer facial distances of a polytope in terms of a polyhedral description of the polytope. The bounds are stated in Corollary 4.4 and Corollary 4.5 below. Throughout this section, we assume that $\mathcal{C} \subseteq \mathbb{R}^n$ is a polytope with the following polyhedral description

$$\mathcal{C} = \{\mathbf{x} \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{b}, D\mathbf{x} \geq \mathbf{e}\}, \quad (4.3)$$

where $A \in \mathbb{R}^{m \times n}$, $\mathbf{b} \in \mathbb{R}^m$, $D \in \mathbb{R}^{k \times n}$, and $\mathbf{e} \in \mathbb{R}^k$. We shall assume that the description (4.3) satisfies the following: for each $i \in [k]$, there exists an $\mathbf{x} \in \mathcal{C}$ such that $\langle \mathbf{d}_i, \mathbf{x} \rangle > e_i$. This can be assumed without loss of generality as otherwise the inequality $\langle \mathbf{d}_i, \mathbf{x} \rangle \geq e_i$ in (4.3) could be replaced with $\langle \mathbf{d}_i, \mathbf{x} \rangle = e_i$. We also assume without loss of generality that $\min_{\mathbf{x} \in \mathcal{C}} \langle \mathbf{d}_i, \mathbf{x} \rangle = e_i$, as otherwise the inequality $\langle \mathbf{d}_i, \mathbf{x} \rangle \geq e_i$ is redundant and can be dropped.

For $i \in [k]$ let $\mathbf{d}_i \in \mathbb{R}^n$ denote the i -th row of D so that for all $\mathbf{x} \in \mathbb{R}^n$

$$(D\mathbf{x})_i = \langle \mathbf{d}_i, \mathbf{x} \rangle.$$

Similarly, for $I \subseteq [k]$, let D_I denote the $|I| \times n$ submatrix of D obtained by selecting the rows of D indexed by I . Likewise, for $I \subseteq [k]$ let \mathbf{e}_I denote the $|I| \times n$ subvector of \mathbf{e} obtained by selecting the entries of \mathbf{e} indexed by I .

We also rely on the following notation.

Definition 4.1. Suppose that $\mathcal{C} \subseteq \mathbb{R}^n$ is a polytope with a polyhedral description (4.3). For each $i \in [k]$, let

$$\sigma_i := \min\{\langle \mathbf{d}_i, \mathbf{v} \rangle - e_i \mid \mathbf{v} \in \text{vert}(\mathcal{C}) \text{ and } \langle \mathbf{d}_i, \mathbf{v} \rangle > e_i\}$$

Suppose that $F \in \text{faces}(\mathcal{C})$ is such that $F \subsetneq \mathcal{C}$. Let

$$I_F := \{i \in [k] \mid \langle \mathbf{d}_i, \mathbf{x} \rangle = e_i \text{ for all } \mathbf{x} \in F\}.$$

Observe that $I_F \neq \emptyset$ because $F \in \text{faces}(\mathcal{C})$ and $F \neq \mathcal{C}$. Let D_{I_F} denote the $|I_F| \times n$ matrix whose rows are the rows from D indexed by I_F and let

$$\mathbb{I}(F) := \{I \subseteq I_F \mid D_I \text{ is full row rank and } I \text{ is maximal}\}.$$

In other words, $\mathbb{I}(F)$ is the collection of all maximal sets of rows from D_{I_F} that are linearly independent.

Lemma 4.2 (Minimal face for polytopes of the form (4.3)). Let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope of the form (4.3) and let $\emptyset \neq F \subsetneq \mathcal{C}$ be a face of \mathcal{C} . Then for all $I \in \mathbb{I}(F)$ it holds that

$$F = \mathcal{C} \cap \{\mathbf{x} \mid D_I \mathbf{x} = \mathbf{e}_I\} = \mathcal{C} \cap \{\mathbf{x} \mid D_{I_F} \mathbf{x} = \mathbf{e}_{I_F}\}.$$

Proof. The construction of I_F and $\mathbb{I}(F)$ immediately implies that $F \subseteq \mathcal{C} \cap \{\mathbf{x} \mid D_I \mathbf{x} = \mathbf{e}_I\} = \mathcal{C} \cap \{\mathbf{x} \mid D_{I_F} \mathbf{x} = \mathbf{e}_{I_F}\}$ for all $I \in \mathbb{I}(F)$. For the reverse inclusion, again the construction of I_F implies that if $\mathbf{x} \in \mathcal{C} \setminus F$ then $\langle \mathbf{d}_i, \mathbf{x} \rangle > e_i$ for some $i \in I_F$. Therefore $\mathcal{C} \setminus F \subseteq \mathcal{C} \setminus \{\mathbf{x} \mid D_{I_F} \mathbf{x} = \mathbf{e}_{I_F}\}$ and consequently $\mathcal{C} \cap \{\mathbf{x} \mid D_{I_F} \mathbf{x} = \mathbf{e}_{I_F}\} \subseteq \mathcal{C} \cap F$. \square

The following result provides the key components to bound the inner and outer facial distances that we subsequently state in Corollary 4.4 and Corollary 4.5.

Theorem 4.3 (Lower bound on facial distance). Let \mathbb{R}^n be endowed with a norm $\|\cdot\|$ and suppose that $\mathcal{C} \subseteq \mathbb{R}^n$ is a polytope of the form (4.3).

(a) If $F \in \text{faces}(\mathcal{C})$ and $F \subsetneq \mathcal{C}$ then

$$\text{dist}(F, \text{conv}(\text{vert}(\mathcal{C}) \setminus F)) \geq \max_{I \in \mathbb{I}(F)} \frac{1}{\|\sum_{i \in I} \mathbf{d}_i / \sigma_i\|^*}. \quad (4.4)$$

(b) If $F, G \in \text{faces}(\mathcal{C})$ and $F \cap G = \emptyset$ then

$$\text{dist}(F, G) \geq \max \left\{ \max_{I \in \mathbb{I}(F)} \frac{1}{\|\sum_{i \in I \cap I_G^c} \mathbf{d}_i / \sigma_i\|^*}, \max_{I \in \mathbb{I}(G)} \frac{1}{\|\sum_{i \in I \cap I_F^c} \mathbf{d}_i / \sigma_i\|^*} \right\}. \quad (4.5)$$

Proof. (a) Let $G := \text{conv}(\text{vert}(\mathcal{C}) \setminus F)$ and suppose $I \in \mathbb{I}(F)$. Lemma 4.2 implies for any vertex $\mathbf{u} \in \text{vert}(G)$ there exists $i \in I$ such that $\langle \mathbf{d}_i, \mathbf{u} \rangle - e_i \geq \sigma_i$ or equivalently $(\langle \mathbf{d}_i, \mathbf{u} \rangle - e_i) / \sigma_i \geq 1$.

Now suppose $\mathbf{x} \in F$ and $\mathbf{y} \in G$. Since \mathbf{y} is the convex combination of vertices of G and $\mathbf{x} \in F$, it follows that

$$\sum_{i \in I} \langle \mathbf{d}_i / \sigma_i, \mathbf{y} - \mathbf{x} \rangle = \sum_{i \in I} (\langle \mathbf{d}_i, \mathbf{y} \rangle - e_i) / \sigma_i \geq 1.$$

Thus, by Cauchy-Schwarz inequality,

$$\|\mathbf{y} - \mathbf{x}\| \geq \frac{1}{\|\sum_{i \in I} \mathbf{d}_i / \sigma_i\|^*}.$$

Since this holds for all $I \in \mathbb{I}(F)$ and all $\mathbf{x} \in F$ and $\mathbf{y} \in G$, (4.4) follows.

- (b) Once again, this follows via a similar but simpler argument to that used in part (a). Suppose that $F, G \in \text{faces}(\mathcal{C})$ and $F \cap G = \emptyset$. By symmetry, to prove (4.5) it suffices to show that for $I \in \mathbb{I}(F)$

$$\text{dist}(F, G) \geq \frac{1}{\|\sum_{i \in I \cap I_G^c} \mathbf{d}_i / \sigma_i\|^*}. \quad (4.6)$$

To that end, observe that Lemma 4.2 implies that for each $\mathbf{u} \in \text{vert}(G)$ there exists $i \in I \cap I_G^c$ such that

$$\langle \mathbf{d}_i, \mathbf{u} \rangle - e_i \geq \sigma_i.$$

It thus follows that for all $\mathbf{y} \in G$ and $\mathbf{x} \in F$

$$\sum_{i \in I \cap I_G^c} \langle \mathbf{d}_i / \sigma_i, \mathbf{y} - \mathbf{x} \rangle = \sum_{i \in I \cap I_G^c} (\langle \mathbf{d}_i, \mathbf{y} \rangle - e_i) / \sigma_i \geq 1.$$

Thus, by Cauchy-Schwarz inequality,

$$\|\mathbf{y} - \mathbf{x}\| \geq \frac{1}{\|\sum_{i \in I \cap I_G^c} \mathbf{d}_i / \sigma_i\|^*}.$$

Since this holds for all $I \in \mathbb{I}(F)$ and all $\mathbf{x} \in F$ and $\mathbf{y} \in G$, (4.6) follows. \square

Corollary 4.4. *Let \mathbb{R}^n be endowed with a norm $\|\cdot\|$, let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope of the form (4.3), and let $F \in \text{faces}(\mathcal{C})$. Then*

$$\bar{\Phi}(F, \mathcal{C}) \geq \Phi(F, \mathcal{C}) \geq \min_{\substack{G \in \text{faces}(F) \\ \emptyset \neq G \neq \mathcal{C}}} \max_{I \in \mathbb{I}(G)} \frac{1}{\|\sum_{i \in I} \mathbf{d}_i / \sigma_i\|^*}$$

For polytopes with a standard form description, that is, when $D = I$ and $\mathbf{e} = \mathbf{0}$, the bounds in Corollary 4.4 can be simplified and sharpened as in Corollary 4.5 below. The statement of Corollary 4.5 relies on the following terminology and notation.

Let $\boldsymbol{\sigma}^{-1} \in \mathbb{R}^k$ denote the vector with entries $1/\sigma_i$, $i \in [k]$. For $I \subseteq [k]$ let $\boldsymbol{\sigma}_I^{-1}$ denote the $|I| \times n$ subvector of $\boldsymbol{\sigma}^{-1}$ obtained by selecting the entries of $\boldsymbol{\sigma}^{-1}$ indexed by I . We shall write $\mathbf{1}$ in lieu of $\boldsymbol{\sigma}^{-1}$ in the special case when $\sigma_i = 1$ for all $i \in [k]$. We shall say that a norm $\|\cdot\|$ on \mathbb{R}^n satisfies the *compatibility assumption* if $\|\mathbf{x}\| \leq \|\mathbf{y}\|$ whenever $|\mathbf{x}| \leq |\mathbf{y}|$ componentwise. Observe that many popular norms on \mathbb{R}^n , including all ℓ_p -norms for $1 \leq p \leq \infty$, satisfy the compatibility assumption.

Corollary 4.5. *Let \mathbb{R}^n be endowed with a norm $\|\cdot\|$ such that $\|\cdot\|^*$ satisfies the compatibility assumption, let $\mathcal{C} \subseteq \mathbb{R}^n$ be a polytope of the form $\mathcal{C} = \{\mathbf{x} \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$. Then for all $F \in \text{faces}(\mathcal{C})$*

$$\Phi(F, \mathcal{C}) \geq \frac{1}{\|\boldsymbol{\sigma}_{I_F}^{-1}\|^*}$$

and

$$\bar{\Phi}(F, \mathcal{C}) \geq \max \left\{ \frac{1}{\|\boldsymbol{\sigma}_{I_F}^{-1}\|_*}, \frac{1}{\|\boldsymbol{\sigma}_{I_{F^c}}^{-1}\|_*} \right\}$$

where $I_F = \{i \in [n]: x_i = 0 \text{ for all } \mathbf{x} \in F\}$.

In particular, if \mathcal{C} is a simplex-like polytope then for all $F \in \text{faces}(\mathcal{C})$

$$\Phi(F, \mathcal{C}) \geq \frac{1}{\|\mathbf{1}_{I_F}\|_*} \quad \text{and} \quad \bar{\Phi}(F, \mathcal{C}) \geq \max \left\{ \frac{1}{\|\mathbf{1}_{I_F}\|_*}, \frac{1}{\|\mathbf{1}_{I_F^c}\|_*} \right\}.$$

Proof of Proposition 3.19. This readily follows from Proposition 3.15, Corollary 4.5, and the observation that

$$\|\mathbf{1}_{I_{F(X^*)^c}}\|_2 = \sqrt{\text{card}(X^*)} \quad \text{and} \quad \|\mathbf{1}_{I_{F(X^*)}}\|_2 = \sqrt{n - \text{card}(X^*)}.$$

□

Acknowledgements

Research reported in this paper was partially supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – The Berlin Mathematics Research Center MATH⁺ (EXC-2046/1, project ID 390685689, BMS Stipend). Research reported in this paper was also partially supported by the Bajaj Family Chair at the Tepper School of Business, Carnegie Mellon University.

References

- [1] Amir Beck and Shimrit Shtern. Linearly convergent away-step conditional gradient for non-strongly convex functions. *Mathematical Programming*, 164:1–27, 2017.
- [2] Jonathan M Borwein, Guoyin Li, and Liangjin Yao. Analysis of the convergence rate for the cyclic projection algorithm applied to basic semialgebraic convex sets. *SIAM Journal on Optimization*, 24(1):498–527, 2014.
- [3] Alejandro Carderera, Sebastian Pokutta, Christof Schütte, and Martin Weiser. CINDy: Conditional gradient-based identification of non-linear dynamics–noise-robust recovery. *arXiv preprint arXiv:2101.02630*, 2021.
- [4] Vladimir Fedorovich Demianov and Aleksandr Moiseevich Rubinov. *Approximate methods in optimization problems*. Number 32. Elsevier Publishing Company, 1970.
- [5] Joseph C Dunn. Rates of convergence for conditional gradient algorithms near singular and nonsingular extremals. *SIAM Journal on Control and Optimization*, 17(2):187–211, 1979.
- [6] Robert M Freund, Paul Grigas, and Rahul Mazumder. An extended Frank-Wolfe method with “in-face” directions, and its application to low-rank matrix completion. *SIAM Journal on optimization*, 27(1):319–346, 2017.

- [7] Dan Garber. Revisiting Frank-Wolfe for polytopes: Strict complementarity and sparsity. *Advances in Neural Information Processing Systems*, 33:18883–18893, 2020.
- [8] Dan Garber and Elad Hazan. Faster rates for the Frank-Wolfe method over strongly-convex sets. In *Proceedings of the International Conference on Machine Learning*. PMLR, 2015.
- [9] Dan Garber and Ofer Meshi. Linear-memory and decomposition-invariant linearly convergent conditional gradient algorithm for structured polytopes. In *Proceedings of the International Conference on Neural Information Processing Systems*, pages 1009–1017. PMLR, 2016.
- [10] Jacques Guélat and Patrice Marcotte. Some comments on Wolfe’s ‘away step’. *Mathematical Programming*, 35(1):110–119, 1986.
- [11] David H Gutman and Javier F Peña. The condition number of a function relative to a set. *Mathematical Programming*, 188:255–294, 2021.
- [12] Charles A Holloway. An extension of the Frank and Wolfe method of feasible directions. *Mathematical Programming*, 6(1):14–27, 1974.
- [13] Martin Jaggi. Revisiting Frank-Wolfe: Projection-free sparse convex optimization. In *Proceedings of the International Conference on Machine Learning*, pages 427–435. PMLR, 2013.
- [14] Armand Joulin, Kevin Tang, and Li Fei-Fei. Efficient image and video co-localization with Frank-Wolfe algorithm. In *Computer Vision – ECCV 2014*, pages 253–268, Cham, 2014. Springer International Publishing.
- [15] Thomas Kerdreux, Alexandre d’Aspremont, and Sebastian Pokutta. Projection-free optimization on uniformly convex sets. In *Proceedings of the International Conference on Artificial Intelligence and Statistics*, pages 19–27. PMLR, 2021.
- [16] Simon Lacoste-Julien and Martin Jaggi. On the global linear convergence of Frank-Wolfe optimization variants. In *Proceedings of Advances in Neural Information Processing Systems*, pages 496–504, 2015.
- [17] Evgeny S Levitin and Boris T Polyak. Constrained minimization methods. *USSR Computational Mathematics and Mathematical Physics*, 6(5):1–50, 1966.
- [18] Javier Peña and Daniel Rodriguez. Polytope conditioning and linear convergence of the Frank–Wolfe algorithm. *Mathematics of Operations Research*, 44(1):1–18, 2019.
- [19] Javier F Peña. Affine invariant convergence rates of the conditional gradient method. *SIAM Journal on Optimization*, 33(4):2654–2674, 2023.
- [20] Boris T Polyak. *Introduction to optimization*. New York, Optimization Software,, 1987.
- [21] Robert Tibshirani. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 58(1):267–288, 1996.
- [22] Kazuma K Tsuji, Ken’ichiro Tanaka, and Sebastian Pokutta. Pairwise conditional gradients without swap steps and sparser kernel herding. In *Proceedings of the International Conference on Machine Learning*, pages 21864–21883. PMLR, 2022.

- [23] Elias Wirth, Hiroshi Kera, and Sebastian Pokutta. Approximate vanishing ideal computations at scale. In *Proceedings of the International Conference on Learning Representations*, 2023.
- [24] Elias Wirth, Thomas Kerdreux, and Sebastian Pokutta. Acceleration of Frank-Wolfe algorithms with open-loop step-sizes. In *Proceedings of the International Conference on Artificial Intelligence and Statistics*, pages 77–100. PMLR, 2023.
- [25] Elias Wirth, Javier Peña, and Sebastian Pokutta. Accelerated affine-invariant convergence rates of the Frank-Wolfe algorithm with open-loop step-sizes. *arXiv preprint arXiv:2310.04096*, 2023.
- [26] Philip Wolfe. Convergence theory in nonlinear programming. *Integer and Non-linear Programming*, pages 1–36, 1970.