

MONOTONE SEQUENCES OF METRIC SPACES WITH COMPACT LIMITS

R. PERALES AND C. SORMANI

ABSTRACT. In this paper, we consider a fixed metric space (possibly an oriented Riemannian manifold with boundary) with an increasing sequence of distance functions and a uniform upper bound on diameter. When the metric space endowed with the pointwise limit of these distances is compact, then there is uniform and Gromov-Hausdorff (GH) convergence to this limit. When the metric space also has an integral current structure of uniformly bounded total mass (as is true for an oriented Riemannian manifold with boundary that has a uniform bound on total volume), we prove volume preserving intrinsic flat convergence to a subset of the GH limit whose closure is the whole GH limit. We provide a review of all notions and have a list of open questions at the end. *Dedicated to Xiaochun Rong.*

1. INTRODUCTION

Our goal is to teach the notions of Gromov-Hausdorff [GH] and Sormani-Wenger intrinsic flat [SWIF] convergence while proving a new theorem that has no assumptions on curvature. The notion of GH distance between metric spaces was first introduced by Edwards in [Edw75] and deeply explored by Gromov in [Gro07] and [Gro81]. See Rong’s 2010 survey [Ron10] for many applications of GH convergence to sequences of Riemannian manifolds with curvature bounds.

The notion of SWIF distance between integral current spaces was introduced by Sormani and Wenger in [SW11] applying the theory of Ambrosio-Kirchheim in [AK00] and work of Wenger in [Wen07]. Volume preserving intrinsic flat (\mathcal{VF}) convergence was introduced by Portegies in [Por15]. See Sormani’s survey [Sor17] for many applications of SWIF convergence to

Key words and phrases. Metric Spaces, Riemannian, Gromov-Hausdorff, Intrinsic Flat.

R. Perales’ research was partially funded by NSF DMS 1612049 on *Geometric Compactness Theorems*.

C. Sormani’s research was partially funded by NSF DMS 1612049 on *Geometric Compactness Theorems* and a PSC-CUNY grant.

sequences of Riemannian manifolds with lower bounds on their scalar curvature. See also papers by Allen, Bryden, Huang, Jauregui, Lakzian, Lee, Perales, Portegies, Sormani, and Wenger which prove SWIF and \mathcal{VF} convergence theorems and present counter examples [SW11][Wen11][LS13][AS19][AS20][HLS17][APS24][AP22][AB24][JL21][JPP22].

The work in this paper is inspired by a theorem in the appendix of [HLS17] by Huang-Lee-Sormani (which applies to sequences with biLipschitz bounds on their distances) and a theorem within [APS24] by Allen-Perales-Sormani (which assumes only volume converging from above and distance from below but requires the limit space to be a compact smooth oriented Riemannian manifold). Here we will only assume the limit space is a compact metric space but we add the hypothesis that the distances are monotone increasing. Our result will be applied by Sormani-Tian in [ST24] to prove SWIF and GH convergence to the extreme limits constructed by Sormani-Tian-Wang in [STW24].

Theorem 1.1. *Given a compact Riemannian manifold, (M^m, g_0) , possibly with boundary, with a monotone increasing sequence of Riemannian metric tensors g_j such that*

$$(1.1) \quad g_j(V, V) \geq g_{j-1}(V, V) \quad \forall V \in TM$$

with uniform bounded diameter,

$$(1.2) \quad \text{diam}_{g_j}(M) \leq D_0.$$

Then the induced length distance functions $d_j : M \times M \rightarrow [0, D_0]$ are monotone increasing and converge pointwise to a distance function, $d_\infty : M \times M \rightarrow [0, D_0]$ so that (M, d_∞) is a metric space.

If the metric space (M, d_∞) is a compact metric space, then $d_j \rightarrow d_\infty$ uniformly and

$$(1.3) \quad (M, d_j) \xrightarrow{GH} (M, d_\infty).$$

If, in addition, M is an oriented Riemannian manifold with uniformly bounded total volume,

$$(1.4) \quad \text{vol}_j(M) \leq V_0 \text{ and } \text{vol}_j(\partial M) \leq A_0$$

then we have volume preserving intrinsic flat convergence

$$(1.5) \quad (M, d_j, [[M]]) \xrightarrow{\mathcal{VF}} (M_\infty, d_\infty, T_\infty).$$

where T_∞ is an integral current on (M, d_∞) such that $T_\infty = [[M]]$ viewed as an integral current on (M, d_j) and

$$(1.6) \quad M_\infty = \text{set}_{d_\infty}(T_\infty) \subset M \text{ with closure } \overline{M}_\infty = M.$$

Note that Sormani-Wenger already proved that any GH converging sequence satisfying (1.4) has a subsequence converging to a SWIF limit that is a possibly empty subset of the GH limit in [SW11]. In the Riemannian theorem above we use the monotonicity to force the whole sequence to converge and to force the closure of the SWIF limit to agree with the GH limit. We will see in Example 4.14 within that an increasing sequence can converge to a manifold with a cusp singularity, and since a cusp singularity is not included in the definition of an integral current space, this example has $M_\infty = \text{set}(T_\infty) \neq M$.

We will prove this Riemannian theorem in stages as consequences of more general results concerning metric spaces and integral current spaces. We will review the various notions of convergence right before applying them, so that this article may be easily read by a novice. Note that GH, SWIF, and \mathcal{VF} limit spaces are defined using distance functions and do not necessarily agree with the metric completions of limits spaces found by taking smooth convergence of Riemannian metric tensors away from singular sets (see work of Allen-Sormani [AS19]).

In Section 2 we prove a few basic lemmas about metric spaces with monotone increasing sequences of distance functions including a review of the Riemannian distance functions. In Lemma 2.5 we prove pointwise convergence of the monotone increasing sequence of distance functions with a uniform upper bound on diameter. In Example 2.6 we see that we need not have uniform convergence when the limit is noncompact.

In Section 3 we review the definition of Gromov-Hausdorff convergence (Definition 3.1). We then prove the GH and uniform convergence part of Theorem 1.1. In fact we prove a more general theorem concerning a metric space with a monotone increasing sequence of distance functions that converge pointwise to a compact limit in Theorem 3.2. We show the sequence converges uniformly and in the GH sense. We prove this theorem using different techniques than would have been used by Gromov in order to prepare for the SWIF convergence part of the paper. In particular, we construct a common metric space for the converging sequence that has special properties in Proposition 3.9 and Lemma 3.10. In Remark 3.8 we explain that Example 2.6 does not have a Gromov-Hausdorff limit.

In Section 4 we prove the SWIF convergence part of Theorem 1.1. In fact we prove a more general SWIF convergence theorem concerning the convergence of an integral current space with increasing distance functions [Theorem 4.16]. We begin by reviewing De Giorgi's notion of a Lipschitz tuple [DG95], the notion of a Lipschitz chart, and Ambrosio-Kirchheim's notion of an integral current and weak convergence [AK00], proving lemmas about how these objects behave for monotone increasing sequences of distances. We review Sormani-Wenger's definition of an integral current space, including $(M, d, [[M]])$, and their definition of intrinsic flat distance. We then state Theorem 4.16 and apply it to complete the proof of Theorem 1.1 in Section 4.8. Finally we prove Theorem 4.16, first finding a converging subsequence and a limit current structure in Proposition 4.17 and completing the proof in Section 4.10.

In Section 5 we state open problems.

The authors would like to thank Wenchuan Tian and Changliang Wang for inspiring discussions of examples.

2. POINTWISE CONVERGENCE OF MONOTONE INCREASING DISTANCE FUNCTIONS

Definition 2.1. *A metric space, (X, d) , is a collection of points, X , paired with a distance function $d : X \times X \rightarrow [0, \infty)$ which is definite,*

$$(2.1) \quad d(x, y) = 0 \iff x = y,$$

symmetric,

$$(2.2) \quad d(x, y) = d(y, x) \quad \forall x, y \in X,$$

and satisfies the triangle inequality,

$$(2.3) \quad d(x, y) \leq d(x, z) + d(z, y) \quad \forall x, y, z \in X.$$

Here we consider metric spaces of finite diameter,

$$(2.4) \quad \text{diam}_d(X) = \sup\{d(x, y) : x, y \in X\} < \infty.$$

We also need the following definition:

Definition 2.2. *We say that $d_j : X \times X \rightarrow [0, \infty)$ is a monotone increasing sequence of distance functions on X , if for all $j \in \mathbb{N}$ one has that (X, d_j) is a metric space and*

$$(2.5) \quad d_{j+1}(x, y) \geq d_j(x, y) \quad \forall x, y \in X.$$

2.1. Riemannian Distances. Recall that a Riemannian manifold, (M, g) , has a natural metric space, (M, d_g) , defined using the induced Riemannian distance as follows. The lengths of piecewise smooth curves, $C : [0, 1] \rightarrow M$, are defined by

$$(2.6) \quad L_g(C) = \int_0^1 g(C'(t), C'(t))^{1/2} dt$$

and the Riemannian distance between points is defined by

$$(2.7) \quad d_g(x, y) = \inf\{L_g(C)\}$$

where the infimum is taken over all piecewise smooth curves, $C : [0, 1] \rightarrow M$, such that $C(0) = x$ and $C(1) = y$. The diameter of (M, d_g) is then defined as

$$(2.8) \quad \text{diam}_g(M) = \text{diam}_{d_g}(M) = \sup\{d_g(x, y) : x, y \in M\}.$$

Lemma 2.3. *If M is a Riemannian manifold with two Riemannian metric tensors g_a and g_b such that*

$$(2.9) \quad g_a(V, V) \leq g_b(V, V) \quad \forall V \in TM$$

then the corresponding induced Riemannian distances $d_a = d_{g_a}$ and $d_b = d_{g_b}$ satisfy

$$(2.10) \quad d_a(x, y) \leq d_b(x, y) \quad \forall x, y \in X.$$

The proof of Lemma 2.3 is an easy exercise applying 2.6 and 2.7.

Remark 2.4. *As a consequence of Lemma 2.3, a Riemannian manifold with a monotone increasing sequence of metric tensors as in (1.1), has a monotone increasing sequence of distance functions as in Definition 2.2. An easy way to construct such a sequence on the sphere, \mathbb{S}^m , is to take a sequence of graphs over spheres, $g_j = g_{\mathbb{S}^m} + dh_j^2$ where $g_{\mathbb{S}^m}$ is the standard round metric on the sphere and $h_j : \mathbb{S}^m \rightarrow [0, 1]$ has $|\nabla h_j| \leq |\nabla h_{j+1}|$ on \mathbb{S}^m , so that $L_{g_j}(C) \leq L_{g_{j+1}}(C)$ for all curves C .*

2.2. Pointwise convergence.

Lemma 2.5. *Suppose that $d_j : X \times X \rightarrow [0, \infty)$ is a monotone increasing sequence of distance functions on X as in Definition 2.2 and that there is a uniform upper bound on diameter,*

$$(2.11) \quad \text{diam}_j(X) = \sup_{x, y \in X} d_j(x, y) \leq D_0 \in (0, \infty)$$

then there is a limit distance function:

$$(2.12) \quad d_\infty(x, y) = \lim_{j \rightarrow \infty} d_j(x, y) = \sup_{j \in \mathbb{N}} d_j(x, y) \in [0, D_0]$$

and (X, d_∞) is a metric space with diameter $\text{diam}_\infty(X) \leq D_0$.

Proof. We know the pointwise limit function $d_\infty : X \times X \rightarrow [0, D_0]$ exists and satisfies (2.12) by the monotone convergence theorem for sequences of real numbers. We can confirm that d_∞ is positive definite as follows:

$$(2.13) \quad d_\infty(x, x) = \lim_{j \rightarrow \infty} d_j(x, x) = 0$$

and

$$(2.14) \quad d_\infty(x, y) = 0 \implies d_j(x, y) \leq 0 \implies x = y.$$

It is symmetric for all $x, y \in X$ by

$$(2.15) \quad d_\infty(x, y) = \lim_{j \rightarrow \infty} d_j(x, y) = \lim_{j \rightarrow \infty} d_j(y, x) = d_\infty(y, x) \quad \forall x, y \in X.$$

It satisfies the triangle inequality for any $x, y, z \in X$ by

$$(2.16) \quad d_\infty(x, y) = \lim_{j \rightarrow \infty} d_j(x, y)$$

$$(2.17) \quad \leq \lim_{j \rightarrow \infty} (d_j(x, z) + d_j(z, y))$$

$$(2.18) \quad = \lim_{j \rightarrow \infty} d_j(x, z) + \lim_{j \rightarrow \infty} d_j(z, y)$$

$$(2.19) \quad = d_\infty(x, z) + d_\infty(z, y) \quad \forall x, y, z \in X.$$

□

2.3. Example without Uniform Convergence. Even if (X, d_j) is a monotone increasing sequence of compact metric spaces with a uniform upper bound on diameter pointwise convergence does not imply uniform convergence. See Example 2.6 depicted in Figure 2.1.




FIGURE 2.1. This figure depicts Example 2.6.

Example 2.6. Here we adapt an example from Sormani-Wenger [SW11] as in Figure 2.1. Consider a sequence of distinct points $p_j \rightarrow p_\infty$ lying on the equator in the standard round sphere $(\mathbb{S}^2, g_{\mathbb{S}^2})$. Let $r_j > 0$ be decreasing radii such that the balls $B_{g_{\mathbb{S}^2}}(p_j, r_j)$ are disjoint and do not contain p_∞ . We define an increasing sequence of metric tensors

$$(2.20) \quad g_j = g_{\mathbb{S}^2} + dh_j^2$$

as in Remark 2.4 inductively starting with constant $h_0 = 0$ and taking

$$(2.21) \quad h_{j+1}(x) = h_j(x) \quad \text{for } x \in \mathbb{S}^2 \setminus B_{g_{\mathbb{S}^2}}(p_j, r_j)$$

$$(2.22) \quad = H(d_{\mathbb{S}^2}(p_j, x)/r_j) \quad \text{for } x \in B_{g_{\mathbb{S}^2}}(p_j, r_j)$$

where $H : (-1, 1) \rightarrow [0, H_0]$ is a fixed smooth even function of compact support such that $H(0) = H_0 > 0$ and $H'(r) \leq 0$ for $r \geq 0$.

So we have added increasingly thin wells in place of each of the balls. For all $i \leq j$, the g_j radial distance from p_i to any $q_i \in \partial B_{g_{\mathbb{S}^2}}(p_j, r_j)$,

$$(2.23) \quad d_{g_j}(p_i, q_i) = \int_0^{r_i} \sqrt{1 + (H'(r/r_i)(1/r_i))^2} dr.$$

Since $a \leq \sqrt{1 + a^2} \leq 1 + a$ we have

$$(2.24) \quad H_0 \leq d_{g_j}(p_i, q) \leq r_i + H_0 \quad \forall i \leq j.$$

Since any point in (\mathbb{S}^2, g_j) outside of the wells can be joined to a pole travelling a distance $\leq \pi/2$, and poles are a distance π apart, we can apply the triangle inequality to see that

$$(2.25) \quad \text{diam}_{g_j} \leq (H_0 + r_1 + \pi/2) + (H_0 + r_1 + \pi/2) + \pi.$$

This uniform upper bound and Lemma 2.5 implies there is a pointwise limit, $d_j \rightarrow d_\infty$. Taking the limit of (2.24) we have,

$$(2.26) \quad H_0 \leq d_\infty(p_i, p_\infty) \quad \forall i \in \mathbb{N}.$$

However, for $i > j$ sufficiently large

$$(2.27) \quad d_{g_j}(p_i, p_\infty) \leq d_{\mathbb{S}^2}(p_i, p_\infty) < H_0/2$$

so there is no uniform convergence. In fact, the limit distance function d_∞ has a sequence of disjoint g_j balls centered at the p_j of radius H_0 , so (\mathbb{S}^2, d_∞) is not compact.

3. GH CONVERGENCE TO COMPACT LIMITS

Recall the following definition of the Gromov-Hausdorff distance first introduced by Edwards in [Edw75] and then rediscovered and studied extensively by Gromov in [Gro07].

Definition 3.1. *Given a pair of metric spaces, (X_a, d_a) and (X_b, d_b) , the Gromov-Hausdorff distance between them is*

$$(3.1) \quad d_{GH}((X_a, d_a), (X_b, d_b)) = \inf\{d_H^Z(f_a(X_a), f_b(X_b))\}$$

where the infimum is taken over all common metric spaces, (Z, d_Z) , and over all distance preserving maps,

$$(3.2) \quad f_a : (X_a, d_a) \rightarrow (Z, d_Z) \text{ and } f_b : (X_b, d_b) \rightarrow (Z, d_Z)$$

Here d_H^Z denotes the Hausdorff distance,

$$(3.3) \quad d_H^Z(A, B) = \inf\{R : A \subset T_R(B) \text{ and } B \subset T_R(A)\}$$

between subsets $A, B \subset Z$ which is defined using tubular neighborhoods of a given radius R ,

$$(3.4) \quad T_R(A) = \{z \in Z : \exists p \in A \text{ s.t. } d_Z(p, z) < R\}.$$

In this section we prove the following simple theorem:

Theorem 3.2. *If (X, d_j) is a monotone increasing sequence as in Definition 2.2 with a uniform upper bound D_0 on diameter, and if d_j converges pointwise to d_∞ as in Lemma 2.5, and if (X, d_∞) is a compact metric space, then d_j converge uniformly to d_∞ and we have Gromov-Hausdorff convergence as well.*

Although this GH Convergence theorem can be proven quite easily, we will prove it through a sequence of lemmas and propositions that we will apply again to prove SWIF convergence later.

3.1. Distance on the Product Space. We introduce the following standard definition:

Definition 3.3. *Given a pair of metric spaces, (X_a, d_a) and (X_b, d_b) , we define the taxi product metric space $(X_a \times X_b, d_{sum,a,b})$ where*

$$(3.5) \quad d_{sum,a,b}((x_1, y_1), (x_2, y_2)) = d_a(x_1, x_2) + d_b(y_1, y_2)$$

for any $x_1, x_2 \in X_a$ and any $y_1, y_2 \in X_b$.

Lemma 3.4. *If (X, d_a) and (X, d_b) are compact then $(X \times X, d_{sum,a,b})$ is also compact.*

The proof is an exercise for the reader.

Lemma 3.5. *Given two metric spaces (X, d_a) and (X, d_b) , if*

$$(3.6) \quad d_a(x, y) \leq d_b(x, y) \quad \forall x, y \in X$$

then the function $d_a : X \times X \rightarrow [0, \infty)$ is 1-Lipschitz with respect to the distance $d_{sum,b,b}$ on $X \times X$ given in Definition 3.3. That is,

$$(3.7) \quad |d_a(x_1, y_1) - d_a(x_2, y_2)| \leq d_{sum,b,b}((x_1, y_1), (x_2, y_2)) \quad \forall (x_i, y_i) \in X \times X.$$

Proof. By the triangle inequality and then by (3.6) we have

$$(3.8) \quad |d_a(x_1, y_1) - d_a(x_2, y_2)| \leq d_a(x_1, x_2) + d_a(y_1, y_2)$$

$$(3.9) \quad \leq d_b(x_1, x_2) + d_b(y_1, y_2)$$

which gives our claim by (3.5) □

3.2. Uniform Convergence.

Proposition 3.6. *Under the hypotheses of Theorem 3.2, the distance functions, d_j , converge uniformly to d_∞ on $X \times X$.*

Proof. By Lemma 2.5, we have $d_j \leq d_\infty$ as functions on $X \times X$. By Lemma 3.5, the maps $d_j : X \times X \rightarrow [0, D_0]$ are bounded 1-Lipschitz functions on the metric space $(X \times X, d_{sum,\infty,\infty})$, which is compact by Lemma 3.4.

By the Arzela-Ascoli Theorem, there is a subsequence, $d_{j_k} : X \times X \rightarrow [0, D_0]$, which converges uniformly to a limit function. However, we already have a pointwise limit, so that limit function equals $d_\infty : X \times X \rightarrow [0, D_0]$. That is, $\forall \epsilon > 0 \exists N_\epsilon \in \mathbb{N}$ s.t. $\forall k \geq N_\epsilon$

$$(3.10) \quad d_\infty(x, y) - \epsilon < d_{j_k}(x, y) < d_\infty(x, y) + \epsilon \quad \forall x, y \in X \times X.$$

By the assumption of monotonicity, for all $j \geq j_k \geq j_{N_\epsilon}$,

$$(3.11) \quad d_\infty(x, y) - \epsilon < d_{j_k}(x, y) \leq d_j(x, y) \leq d_\infty(x, y) \quad \forall x, y \in X \times X.$$

□

3.3. Constructing a Common Metric Space. Recall that the definition of the Gromov-Hausdorff distance involves a common metric space Z and distance preserving maps [Definition 3.1]. The following lemma is not part of the original proof by Gromov that uniform convergence implies Gromov-Hausdorff convergence. Here we create a different common metric space Z that allows us to prove intrinsic flat convergence later as well. It is a simplification of the Z constructed by Allen-Perales-Sormani in [APS24].

Proposition 3.7. *Given two metric spaces (X, d_a) and (X, d_b) and an $\epsilon > 0$ such that*

$$(3.12) \quad d_b(x, y) - \epsilon \leq d_a(x, y) \leq d_b(x, y) \quad \forall x, y \in X$$

then there exists a metric space

$$(3.13) \quad (Z_{a,b} = X \times [0, h], d_{Z_{a,b}}) \text{ where } h = \epsilon/2,$$

with the distance between $z_1 = (x_1, t_1)$ and $z_2 = (x_2, t_2)$ defined by

$$(3.14) \quad d_{Z_{a,b}}(z_1, z_2) = \min\{d_{\text{taxi}_b}(z_1, z_2), (h - t_1) + (h - t_2) + d_a(x_1, x_2)\}.$$

$$(3.15) \quad \text{where } d_{\text{taxi}_b}(z_1, z_2) = d_b(x_1, x_2) + |t_1 - t_2|$$

so that the identity map,

$$(3.16) \quad \text{id}_{a,b} : (Z_{a,b}, d_{\text{taxi}_b}) \rightarrow (Z_{a,b}, d_{Z_{a,b}})$$

is 1-Lipschitz and there are distance preserving maps:

$$(3.17) \quad f_a = f_{a,(a,b)} : (X, d_a) \rightarrow Z_{a,b} \quad \text{s.t. } f_a(x) = (x, h),$$

$$(3.18) \quad f_b = f_{b,(a,b)} : (X, d_b) \rightarrow Z_{a,b} \quad \text{s.t. } f_b(x) = (x, 0),$$

so that

$$(3.19) \quad d_{GH}((X, d_a), (X, d_b)) \leq d_H^Z(f_a(X), f_b(X)) \leq h.$$

The intuition behind this construction is that we are taking the taxi product of (X, d_b) with the interval $([0, h], d_{\mathbb{R}})$ and then gluing (X, d_a) to this taxi product at $t = h$ which gives possibly shorter distances. See Figure 3.1.




FIGURE 3.1. On the left we see (X_a, d_a) above (X_b, d_b) mapping into $(Z_{a,b}, d_{Z_{a,b}})$ and on the right we see two paths between points in $Z_{a,b}$ where the first achieves d_{taxi_b} and the second takes a short cut through the image of X_a . The shorter of the two paths achieves the minimum in the definition of $d_{Z_{a,b}}$.

Proof. First we confirm that $d_{Z_{a,b}}$ is a metric. It is clearly definite and symmetric. To see that it satisfies the triangle inequality we have several cases. Let $z_i = (x_i, t_i) \in Z_{a,b}$, $i = 1, 2, 3$ and

$$(3.20) \quad D_{13} = d_{Z_{a,b}}(z_1, z_2) + d_{Z_{a,b}}(z_2, z_3).$$

Case I: Assume that $d_{Z_{a,b}}(z_1, z_2) = d_{taxib}(z_1, z_2)$ and $d_{Z_{a,b}}(z_2, z_3) = d_{taxib}(z_2, z_3)$. Then since d_{taxib} is a distance function, and thus satisfies the triangle inequality, and by definition of $d_{Z_{a,b}}$, we get

$$(3.21) \quad D_{13} = d_{taxib}(z_1, z_2) + d_{taxib}(z_2, z_3) \geq d_{taxib}(z_1, z_3) \geq d_{Z_{a,b}}(z_1, z_3).$$

Case II: Assume that we have $d_{Z_{a,b}}(z_1, z_2) = d_{taxib}(z_1, z_2)$ and

$$(3.22) \quad d_{Z_{a,b}}(z_2, z_3) = (h - t_2) + (h - t_3) + d_a(x_2, x_3).$$

Then by $d_b \geq d_a$, triangle inequality in \mathbb{R} and for d_a , we get

$$\begin{aligned} D_{13} &= d_{taxib}(z_1, z_2) + (h - t_2) + (h - t_3) + d_a(x_2, x_3) \\ &\geq d_a(x_1, x_2) + |t_1 - t_2| + (h - t_2) + (h - t_3) + d_a(x_2, x_3) \\ &\geq d_a(x_1, x_2) + (h - t_1) + (h - t_3) + d_a(x_2, x_3) \\ &\geq d_a(x_1, x_3) + (h - t_1) + (h - t_3) \geq d_{Z_{a,b}}(z_1, z_3). \end{aligned}$$

Case III: Assume that

$$(3.23) \quad d_{Z_{a,b}}(z_1, z_2) = (h - t_1) + (h - t_2) + d_a(x_1, x_2)$$

$$(3.24) \quad d_{Z_{a,b}}(z_2, z_3) = (h - t_2) + (h - t_3) + d_a(x_2, x_3).$$

Then, since $h - t_2 \geq 0$ and triangle inequality for d_a , we get

$$(3.25) \quad D_{13} = (h - t_1) + (h - t_2) + d_a(x_1, x_2)$$

$$(3.26) \quad \quad \quad + (h - t_2) + (h - t_3) + d_a(x_2, x_3)$$

$$(3.27) \quad \geq (h - t_1) + d_a(x_1, x_2) + (h - t_3) + d_a(x_2, x_3)$$

$$(3.28) \quad \geq d_a(x_1, x_3) + (h - t_1) + (h - t_3)$$

$$(3.29) \quad \geq d_{Z_{a,b}}(z_1, z_3),$$

which completes the proof of the triangle inequality on $(Z_{a,b}, d_{Z_{a,b}})$.

We have the claimed 1-Lipschitz identity map, because $d_{Z_{a,b}} \leq d_{taxib}$. We have the claims that f_a and f_b are distance preserving maps as follows. By the definition of $d_{Z_{a,b}}$, f_a and $d_b \geq d_a$,

$$d_{Z_{a,b}}(f_a(x), f_a(y)) = d_{Z_{a,b}}((x, h), (y, h)) = \min\{d_b(x, y), d_a(x, y)\} = d_a(x, y).$$

By definition of $d_{Z_{a,b}}$, f_b , (3.12) and $h = \epsilon/2$ we have

$$(3.30) \quad d_{Z_{a,b}}(f_b(x), f_b(y)) = \min\{d_b(x, y), 2h + d_a(x, y)\} = d_b(x, y).$$

Furthermore, by the definition of $d_{Z_{a,b}}$, for any $x \in X$, it holds

$$(3.31) \quad d_{Z_{a,b}}((x, 0), (x, h)) = h,$$

which implies that for any $R > h$,

$$(3.32) \quad f_b(X) \subset T_R(f_a(X)) \text{ and } f_a(X) \subset T_R(f_b(X)),$$

so the Hausdorff distance $d_H^Z(f_a(X), f_b(X)) \leq h$ which gives our final claim. \square

3.4. Proof of GH Convergence. Theorem 3.2 is now easy to prove:

Proof. First we apply Prop 3.6 to get uniform convergence from below. Then we apply Proposition 3.7 to (X, d_j) and (X, d_∞) to see that

$$(3.33) \quad d_{GH}((X, d_j), (X, d_\infty)) < \epsilon_j \rightarrow 0.$$

\square

Remark 3.8. *Note that in the proof of Theorem 3.2 it was essential that we assumed (X, d_∞) is compact, as seen in Example 2.6. This example has the monotonicity but no Gromov-Hausdorff limit. In fact Gromov proved that if a sequence of compact metric spaces has a GH limit then the entire sequence can be embedded via distance preserving maps into a common metric space, (Z, d_Z) [Gro81]. However, the Riemannian manifolds in Example 2.6 have an unbounded number of disjoint balls of radius 1, so this is impossible.*

3.5. A Common Compact Metric Space for the Whole Sequence. Gromov proved in [Gro81] that an entire GH-converging sequence of metric spaces can be embedded with distance preserving maps into a common compact metric space Z along with the limit so that the sequence of images converges in the Hausdorff sense inside Z to the image of the limit. Here we construct a specific Z which we will use later to prove SWIF convergence and Theorem 1.1. See Figure 3.2.

Proposition 3.9. *If (X, d_j) is uniformly converging to compact (X, d_∞)*

$$(3.34) \quad d_\infty(x, y) - \epsilon_j < d_j(x, y) \leq d_\infty(x, y) \quad \forall x, y \in X$$

with ϵ_j decreasing to 0 then taking Z_j of Proposition 3.7, we define

$$(3.35) \quad Z = \bigsqcup_{j=1}^{\infty} Z_j \sim \text{where } Z_j = Z_{j,\infty} = X_j \times [0, h_j]$$

where $(x, 0) \in Z_j$ is identified with $(x, 0) \in Z_k$ for all $x \in X$ and all $j, k \in \mathbb{N}$. Then $d_Z : Z \times Z \rightarrow [0, \infty)$ given by

$$(3.36) \quad d_Z(z_j, z_k) = \inf\{d_{Z_j}((x_j, t_j), (x, 0)) + d_{Z_k}((x, 0), (x_k, t_k)) \mid x \in X\}$$

for

$$(3.37) \quad z_j = (x_j, t_j) \in Z_j \text{ and } z_k = (x_k, t_k) \in Z_k \quad j \neq k,$$

and, otherwise,

$$(3.38) \quad d_Z(z_j, z_k) = d_{Z_j}(z_j, z_k),$$

so the inclusion maps

$$(3.39) \quad \zeta_j : (Z_j, d_{Z_j}) \rightarrow (Z, d_Z)$$

are distance preserving is a distance function. In addition, taking

$$(3.40) \quad f_j = f_{j,(j,\infty)} : (X, d_j) \rightarrow Z_j \text{ and } f_{\infty,(j,\infty)} : (X, d_{\infty}) \rightarrow Z_j$$

from Proposition 3.7, we have distance preserving maps:

$$(3.41) \quad \chi_j = \zeta_j \circ f_j : (X, d_j) \rightarrow (Z, d_Z)$$

and

$$(3.42) \quad \chi_{\infty} = \zeta_j \circ f_{\infty,(j,\infty)} : (X, d_{\infty}) \rightarrow (Z, d_Z),$$

where $\chi_{\infty}(x) = \zeta_j(x, 0)$ does not depend on j and is an isometry onto,

$$(3.43) \quad Z_0 = \chi_{\infty}(X) = \zeta_j(f_{\infty,(j,\infty)}(X)) \subset Z.$$

Furthermore, we have

$$(3.44) \quad d_Z(\chi_j(x), \chi_{\infty}(x)) = d_{Z_j}((x, h_j), (x, 0)) = h_j$$

and

$$(3.45) \quad d_H^Z(\chi_j(X), \chi_{\infty}(X)) \leq h_j.$$

Finally (Z, d_Z) is compact.

Intuitively we have glued the Z_j together like the pages of a book, $\zeta_j : Z_j \rightarrow Z$, along the spine, Z_0 , as in Figure 3.2.

Proof. It is clear that d_Z is symmetric and positive definite. To see that it satisfies the triangle inequality we have several cases. Let $z_{j_i} = (x_i, t_i) \in Z_{j_i}$, $j_i \in \mathbb{N}$ and $i = 1, 2, 3$,

$$(3.46) \quad D_{13} = d_Z(z_{j_1}, z_{j_2}) + d_Z(z_{j_2}, z_{j_3}).$$

Case I: Assume that $j_1 = j_2$ and $j_2 = j_3$. In this case, $d_Z = d_{Z_{j_i}}$ which satisfies the triangle inequality since $d_{Z_{j_i}}$ is a distance function.




FIGURE 3.2. On the left, we see the sequence (X, d_j) above with downward maps $f_{j,(j,\infty)} : (X, d_j) \rightarrow (Z_{j,\infty}, d_{Z_{j,\infty}})$ and upward maps $f_{\infty,(j,\infty)} : (X, d_\infty) \rightarrow (Z_{j,\infty}, d_{Z_{j,\infty}})$ as constructed in Proposition 3.7 and depicted in Figure 3.1. On the right, we glue together all the $(Z_{j,\infty}, d_{Z_{j,\infty}})$ to create (Z, d_Z) as in Proposition 3.9 with maps $\chi_j : X_j \rightarrow Z$ and $\chi_\infty : X_\infty \rightarrow Z$.

Case II: Assume that $j_1 = j_2$ and $j_2 \neq j_3$. By $j_1 = j_2$, the triangle inequality for $d_{Z_{j_2}}$ and definition of d_Z we get

$$(3.47) \quad D_{13} = d_{Z_{j_2}}(z_{j_1}, z_{j_2}) + \inf_{x \in X} \{d_{Z_{j_2}}(z_{j_2}, (x, 0)) + d_{Z_{j_3}}((x, 0), z_{j_3})\}$$

$$(3.48) \quad = \inf_{x \in X} \{d_{Z_{j_2}}(z_{j_1}, z_{j_2}) + d_{Z_{j_2}}(z_{j_2}, (x, 0)) + d_{Z_{j_3}}((x, 0), z_{j_3})\}$$

$$(3.49) \quad \geq \inf_{x \in X} \{d_{Z_{j_2}}(z_{j_1}, (x, 0)) + d_{Z_{j_3}}((x, 0), z_{j_3})\} = d_Z(z_{j_1}, z_{j_3}).$$

Case III: Assume that $j_1 \neq j_2$ and $j_2 \neq j_3$. Take $x, x' \in X$, using the triangle inequality for $d_{Z_{j_2}}$ and $d_{Z_{j_1}}$, and the definition of d_Z we get

$$\begin{aligned} & d_{Z_{j_1}}(z_{j_1}, (x, 0)) + d_{Z_{j_2}}((x, 0), z_{j_2}) + d_{Z_{j_2}}(z_{j_2}, (x', 0)) + d_{Z_{j_3}}((x', 0), z_{j_3}) \\ & \geq d_{Z_{j_1}}(z_{j_1}, (x, 0)) + d_{Z_{j_2}}((x, 0), (x', 0)) + d_{Z_{j_3}}((x', 0), z_{j_3}) \\ & \geq d_{Z_{j_1}}(z_{j_1}, (x, 0)) + d_{Z_{j_1}}((x, 0), (x', 0)) + d_{Z_{j_3}}((x', 0), z_{j_3}) \\ & \geq d_{Z_{j_1}}(z_{j_1}, (x', 0)) + d_{Z_{j_3}}((x', 0), z_{j_3}) \geq d_Z(z_{j_1}, z_{j_3}). \end{aligned}$$

Taking the infimum over $x, x' \in X$ in the previous expression we obtain $D_{13} \geq d_Z(z_{j_1}, z_{j_3})$. This concludes the proof that (Z, d_Z) is a metric space.

The inclusion maps ζ_j are distance preserving by their definition and the definition of d_Z . Since by Proposition 3.7 $f_j = f_{j,(j,\infty)}$ and $f_{\infty,(j,\infty)}$ are distance preserving maps, then the compositions $\chi_j = \zeta_j \circ f_j$ and $\chi_\infty = \zeta_j \circ f_{\infty,(j,\infty)}$ are distance preserving maps. Since we are identifying points of the form $(x, 0) \in Z_j$ with points $(x, 0) \in Z_k$ for all $x \in X$, then $\chi_\infty(x) = \zeta_j(x, 0)$ does not depend on j . Furthermore, χ_∞ is an isometry onto $Z_0 = \chi_\infty(X)$.

Since χ_j and χ_∞ are distance preserving maps and by (3.19) in Proposition 3.7 we have

$$(3.50) \quad d_Z(\chi_j(x), \chi_\infty(x)) \leq d_{Z_j}((x, h_j), (x, 0)) \leq h_j$$

so we have (3.44)-(3.44).

Finally we check that (Z, d_Z) is compact. Given any sequence $z_i \in Z$,

$$(3.51) \quad \exists j_i \in \mathbb{N} \exists (x_i, t_i) \in Z_{j_i} = X \times [0, h_{j_i}] \text{ such that } z_i = \zeta_{j_i}(x_i, t_i).$$

Since (X, d_∞) is compact, a subsequence $x_i \rightarrow x_\infty$. Since $[0, 1]$ is compact, a further subsequence can be taken such that $t_i \rightarrow t_\infty$. A further subsequence can be taken to guarantee that either $j_i \rightarrow \infty$ or j_i is constant $j_i = j_0$. In the diverging case we use $t_i \leq h_{j_i} \rightarrow 0$ to see that $t_\infty = 0$, thus our subsequence converges to

$$(3.52) \quad z_\infty = \chi_\infty(x_\infty) = \zeta_{j_i}(x_\infty, 0) \in Z_0$$

which we see as follows:

$$(3.53) \quad d_Z(z_i, z_\infty) = d_Z(\zeta_{j_i}(x_i, t_i), \zeta_{j_i}(x_\infty, 0))$$

$$(3.54) \quad = d_{Z_{j_i}}((x_i, t_i), (x_\infty, 0))$$

$$(3.55) \quad \leq d_\infty(x_i, x_\infty) + |t_i - 0| \rightarrow 0.$$

In the constant, $j_i = j_0$ case, our subsequence converges to

$$(3.56) \quad z_\infty = \zeta_{j_0}(x_\infty, t_\infty)$$

which we see as follows:

$$(3.57) \quad d_Z(z_i, z_\infty) = d_Z(\zeta_{j_0}(x_i, t_i), \zeta_{j_0}(x_\infty, t_\infty))$$

$$(3.58) \quad = d_{Z_{j_0}}((x_i, t_i), (x_\infty, t_\infty))$$

$$(3.59) \quad \leq d_\infty(x_i, x_\infty) + |t_i - t_\infty| \rightarrow 0.$$

Thus Z is compact. □

Lemma 3.10. *Assume the hypotheses of Proposition 3.9. For all $j \in \mathbb{N}$ there exists*

$$(3.60) \quad W_j = \bigcup_{k=j}^{\infty} \chi_k(Z_k) \subset Z$$

and there is a 1-Lipschitz map, $F_j : (W_j, d_Z) \rightarrow (X, d_j)$ where

$$(3.61) \quad F_j(\zeta_k(x, t)) = x \quad \forall k \geq j \forall x \in X, \forall t \in [0, h_k].$$

Proof. Let $z_1, z_2 \in W_j \subset Z$. Then

$$(3.62) \quad \exists k_i \geq j, \exists x_i \in X \exists t_i \in [0, h_{k_i}] \text{ s.t. } z_i = \chi_{k_i}(x_i, t_i).$$

If $k_1 = k_2$, since $d_\infty \geq d_{k_i} \geq d_j$, by the minimum in the definition of $d_{Z_{k_i}}$ and the monotonicity of the distance functions, we get

$$(3.63) \quad d_Z(z_1, z_2) \geq d_{k_i}(x_1, x_2) \geq d_j(x_1, x_2).$$

If $k_2 \neq k_1$, then observe that for all $x \in X$,

$$(3.64) \quad d_{Z_{k_i}}(z_i, (x, 0)) = d_{Z_{k_i}}((x_i, t_i), (x, 0)) \geq d_{k_i}(x_i, x) \geq d_j(x_i, x)$$

by the definition of $d_{Z_{k_i}}$ because $t_i \geq 0$ and $h_{k_i} - t_i + h_{k_i} \geq 0$, followed by the monotonicity of the distance functions. Thus

$$(3.65) \quad d_{Z_{k_1}}(z_1, (x, 0)) + d_{Z_{k_2}}((x, 0), z_2) \geq d_j(x_1, x) + d_j(x_2, x).$$

Taking the infimum over all $x \in X$ on the left and applying the triangle inequality for d_j on the right we have,

$$(3.66) \quad d_Z(z_1, z_2) \geq d_j(x_1, x_2).$$

□

4. INTRINSIC FLAT CONVERGENCE

In this section we review the definition of integral current spaces which include all oriented Riemannian manifolds of finite volume with boundary of finite volume. We then review the Sormani-Wenger Intrinsic Flat (SWIF) distance between these spaces. Finally we state and prove Theorem 4.16 concerning the convergence of monotone increasing sequences of integral current spaces which implies the SWIF convergence claimed in Theorem 1.1.

4.1. Lipschitz Functions and Tuples on Metric Spaces. We say that a function, $F : (X, d_X) \rightarrow (Y, d_Y)$ is K -Lipschitz if

$$(4.1) \quad d_Y(F(x_1, x_2)) \leq K d_X(x_1, x_2) \quad \forall x_1, x_2 \in X$$

We define the Lipschitz constant of F to be the smallest such K .

Lemma 4.1. *Suppose that (X, d_a) and (X, d_b) are two metric spaces and*

$$(4.2) \quad d_a(x_1, x_2) \leq d_b(x_1, x_2) \quad \forall x_1, x_2 \in X.$$

If a function $\pi : X \rightarrow Y$ is K -Lipschitz as a map $\pi : (X, d_a) \rightarrow (Y, d_Y)$ then it is also K -Lipschitz as a map $\pi : (X, d_b) \rightarrow (Y, d_Y)$.

Proof.

$$(4.3) \quad d_Y(\pi(x_1), \pi(x_2)) \leq K d_a(x_1, x_2) \leq K d_b(x_1, x_2) \quad \forall x_1, x_2 \in X.$$

□

By the definition in [AK00] an m -tuple, $(\pi_0, \pi_1, \dots, \pi_m)$, of Lipschitz functions with respect to d_a consists of a bounded Lipschitz function, $\pi_0 : (X, d_a) \rightarrow \mathbb{R}$, and m Lipschitz functions, $\pi_k : (X, d_a) \rightarrow \mathbb{R}$, for $k = 1, \dots, m$.

Corollary 4.2. *Suppose that (X, d_a) and (X, d_b) are two metric spaces and*

$$(4.4) \quad d_a(x, y) \leq d_b(x, y) \quad \forall x, y \in X.$$

Then tuples of Lipschitz functions with respect to d_a are tuples of Lipschitz functions with respect to d_b .

Lemma 4.3. *Given the hypotheses of Proposition 3.9 and Lemma 3.10. If $\pi : (X, d_j) \rightarrow Y$ is Lipschitz K_j , and $F_j : (W_j, d_Z) \rightarrow (X, d_j)$ of Lemma 3.10, then*

$$(4.5) \quad \tilde{\pi} : (W_j, d_Z) \rightarrow Y \text{ such that } \tilde{\pi}(z) = \pi \circ F_j(z)$$

is also Lipschitz K_j and satisfies

$$(4.6) \quad \tilde{\pi}(\chi_\infty(x)) = \pi(x) \quad \forall x \in X.$$

Remark 4.4. *Note that if we try to define $\tilde{\pi}$ as a function from all of (Z, d_Z) to \mathbb{R} as in (4.5) then it might not be Lipschitz. The proof depends strongly on the restriction to $W_j \subset Z$ and Lemma 3.10.*

Proof. Recall $F_j : (W_j, d_Z) \rightarrow (X, d_j)$ are Lipschitz one functions, so

$$(4.7) \quad d_Y(\tilde{\pi}(z_1), \tilde{\pi}(z_2)) \leq K_j d_j(F_j(z_1), F_j(z_2)) \leq K_j d_Z(z_1, z_2) \quad \forall z_1, z_2 \in Z.$$

□

4.2. Charts into our Metric Space. This is needed to understand integral structures on (X, d_j) vs (X, d_{j+1}) and (X, d_∞) .

Lemma 4.5. *Suppose (X, d_a) and (X, d_b) are two metric spaces and*

$$(4.8) \quad d_a(x, y) \leq d_b(x, y) \quad \forall x, y \in X.$$

If $\psi : U \subset \mathbb{R}^m \rightarrow X$ is a Lipschitz chart with respect to d_b on X , then it is Lipschitz with respect to d_a on X .

Proof. We have a Lipschitz constant $K_\psi \in (0, \infty)$ such that

$$(4.9) \quad d_b(\psi(a_1), \psi(a_2)) \leq K_\psi |a_1 - a_2| \quad \forall a_1, a_2 \in U.$$

By (4.8), we have

$$(4.10) \quad d_a(\psi(a_1), \psi(a_2)) \leq K_\psi |a_1 - a_2| \quad \forall a_1, a_2 \in U.$$

□

Remark 4.6. *Sometimes in the definition of a rectifiable space or current it is written that the charts $\psi_i : U_i \rightarrow X$ are bi-Lipschitz. This is true by Lemma 4 in Kirchheim's [Kir94] as soon as one has a countable collection of Lipschitz charts (cf. Lemma 4.1 of [AK00]). The bi-Lipschitz charts are not necessarily the same collection of charts. So even if the charts $\psi_i : (U_i \subset \mathbb{R}^m, d_{\mathbb{R}^m}) \rightarrow (X, d_b)$ are bi-Lipschitz onto their images, they are only Lipschitz as charts into (X, d_a) not bi-Lipschitz, and so one would need to apply Kirchheim's theorem to produce bi-Lipschitz charts into (X, d_b) .*

4.3. Ambrosio-Kirchheim's integral currents on complete metric spaces.

Here we review the work of Ambrosio-Kirchheim in [AK00]. They defined m dimensional currents on complete metric spaces, (Z, d_Z) as multilinear functionals, T , acting on Lipschitz tuples satisfying various hypotheses [AK00]. For example, given a compact m -dimensional oriented Riemannian manifold, (M, g) , we can define the current,

$$(4.11) \quad [[M]](\pi_0, \dots, \pi_m) = \int_M \pi_0 d\pi_1 \wedge \dots \wedge d\pi_m$$

which is well defined because Lipschitz maps are differentiable almost everywhere. The integration is actually defined by integrating over atlas of disjoint oriented charts and taking the sum.

An m dimensional *integer rectifiable current* on a complete metric space, (Z, d_Z) , is a multilinear functional, T , defined on Lipschitz tuples on (Z, d_Z) that has parametrization which is an atlas of Lipschitz charts

$$(4.12) \quad \psi_i : (U_i \subset \mathbb{R}^m, d_{\mathbb{R}^m}) \rightarrow (Z, d_Z)$$

defined on Borel sets, $U_i \subset \mathbb{R}^m$, with integer multiplicities a_i , such that total Hausdorff measures satisfies

$$(4.13) \quad \sum_{i=1}^{\infty} |a_i| \mathcal{H}_{d_Z}^m(\psi_i(U_i)) < \infty.$$

The integer rectifiable current T is defined by the weighted sum:

$$(4.14) \quad T = \sum_{i=1}^{\infty} a_i \psi_{\#}[[U_i]],$$

where

$$(4.15) \quad \psi_{\#}[[U]](\pi_0, \pi_1, \dots, \pi_m) = \int_U (\pi_0 \circ \psi) d(\pi_1 \circ \psi) \wedge \dots \wedge d(\pi_m \circ \psi).$$

The integral is well defined because charts are Lipschitz and thus differentiable almost everywhere. The weighted sum of the integrals defining $T(\pi_0, \dots, \pi_m)$ is finite because the weighted sum of the Hausdorff measures in (4.13), $\max\{\pi_0\}$, and the product $\text{Lip}(\pi_j)$ for $j = 1, \dots, m$ can be used to bound the integrals. Warning: the Lipschitz charts and the Hausdorff measures depend on the distance function on Z . See Lemma 4.7 below.

By Stoke's Theorem, an oriented Riemannian manifold of dimension m , M , has

$$(4.16) \quad \int_{\partial M} \pi_1 d\pi_2 \wedge \dots \wedge d\pi_m = \int_M 1 d\pi_1 \wedge \dots \wedge d\pi_m.$$

So it is natural to define the boundary of a current by

$$(4.17) \quad \partial T(\pi_1, \dots, \pi_m) = T(1, \pi_1, \dots, \pi_m).$$

An *integral current* is an integer rectifiable current whose boundary is also integer rectifiable. Be warned however that the parametrization of the boundary, ∂T , is not necessarily found by taking the boundaries of the atlas of charts parametrizing, T .

The *mass measure*, $\|T\|_{d_Z}$, of a current T on a complete metric space, (Z, d_Z) , is the smallest finite Borel measure such that

$$(4.18) \quad T(\pi_0, \pi_1, \dots, \pi_m) \leq \prod_{i=1}^m \text{Lip}(\pi_i) \int_Z |\pi_0| \|T\|_{d_Z}$$

for all tuples and the *mass* of T is

$$(4.19) \quad \mathbf{M}_{d_Z}(T) = \|T\|_{d_Z}(Z).$$

The support of a current is

$$(4.20) \quad \text{spt}_{d_Z}(T) = \{z \in Z : \|T\|_{d_Z}(B_{d_Z}(z, r)) > 0 \ \forall r > 0\}$$

and the set of an m dimensional current is

$$(4.21) \quad \text{set}_{d_Z}(T) = \left\{ z \in Z : \liminf_{r \rightarrow 0} \|T\|_{d_Z}(B_{d_Z}(z, r))/r^m > 0 \ \forall r > 0 \right\}.$$

Ambrosio-Kirchheim prove that $\text{set}_{d_Z}(T)$ is rectifiable and that the closure of this set is the support of T ,

$$(4.22) \quad \text{spt}(T) = \overline{\text{set}_{d_Z}(T)}.$$

Given a Lipschitz map, $F : (X, d_X) \rightarrow (Y, d_Y)$, the pushforward of an integral current, T , on (X, d_X) , to an integral current, $F_{\#}(T)$, on (Y, d_Y) is defined by

$$(4.23) \quad F_{\#}(T)(\pi_0, \dots, \pi_m) = T(\pi_0 \circ F, \dots, \pi_m \circ F)$$

for any Lipschitz tuple on (Y, d_Y) . To see that $F_{\#}(T)$ is an integral current, Ambrosio-Kirchheim prove that the mass is bounded as follows:

$$(4.24) \quad \mathbf{M}_{d_Y}(F_{\#}(T)) \leq (\text{Lip}(F))^m \mathbf{M}_{d_X}(T).$$

For our purposes we need the following lemma.

Lemma 4.7. *Suppose (X, d_a) and (X, d_b) are two metric spaces and*

$$(4.25) \quad d_a(x, y) \leq d_b(x, y) \quad \forall x, y \in X.$$

If S is an integral current on (X, d_b) then it is an integral current on (X, d_a) defined using the same collection of Lipschitz charts and weights with

$$(4.26) \quad \mathbf{M}_{d_a}(S) \leq \mathbf{M}_{d_b}(S) \text{ and } \mathbf{M}_{d_a}(\partial S) \leq \mathbf{M}_{d_b}(\partial S).$$

The mass measures also satisfy

$$(4.27) \quad \|S\|_{d_a} \leq \|S\|_{d_b} \text{ and } \|\partial S\|_{d_a} \leq \|\partial S\|_{d_b}.$$

Proof. By the definition of integral current structure on (X, d_b) , there is a countable collection of d_b -Lipschitz charts, $\psi_i : U_i \subset \mathbb{R}^m \rightarrow X$, and integers a_i such that

$$(4.28) \quad S(\omega) = \sum_{i=1}^{\infty} a_i \psi_{i\#}[[U_i]](\omega)$$

for any m tuple, ω , of d_b -Lipschitz functions. By Lemma 4.5 these charts are also d_a -Lipschitz charts, so the weighted sum in (4.28) is well defined acting on any m tuple, ω , of d_a -Lipschitz functions. Note that the same can be done for ∂S . Thus S is an integral current on (X, d_a) .

To estimate the inequalities of the mass measures, recall that in [AK00] the mass measure of a current, S , on a metric space (X, d) is defined as the smallest Borel measure, μ , on X such that

$$(4.29) \quad |S(\pi_0, \pi_1, \dots, \pi_m)| \leq \prod_{k=1}^m \text{Lip}_d(\pi_k) \int_X |\pi_0| d\mu$$

for all tuples, $(\pi_0, \pi_1, \dots, \pi_m)$, of d -Lipschitz functions. By Corollary 4.2, for any tuple, $(\pi_0, \pi_1, \dots, \pi_m)$, of d_a -Lipschitz functions, we also have (4.29) with $\mu = \|S\|_{d_b}$. Thus the smallest measure for the tuples of d_a -Lipschitz functions exists and is smaller,

$$(4.30) \quad \|S\|_{d_a} \leq \|S\|_{d_b}.$$

The same can be done for ∂S . This also gives us the total mass estimate in (4.26). \square

Note in particular the above lemma implies that distance preserving maps push forward currents conserving their masses, boundaries, and boundary masses.

Remark 4.8. *The converse of Lemma 4.7 does not hold. If T is an integral current on (X, d_a) , it does not necessarily define an integral current on (X, d_b) for $d_b \geq d_a$. It is a multilinear functional acting on tuples in (X, d_b) because these are also tuples on (X, d_a) by Lemma 4.1. However it does not necessarily have finite mass. In addition, a parametrization of T by Lipschitz charts into (X, d_a) might not give Lipschitz charts into (X, d_b) .*

Soon we will be considering a sequence of metric spaces (X, d_j) with an integral current T , and the limit, (X, d_∞) . Since $d_\infty \geq d_j$, we will need to overcome the lack of a converse to Lemma 4.7. We will see later in the proof of Lemma 4.11, that we can sometimes try to rewrite a parametrization of an integral current on (X, d_j) to obtain charts that are also Lipschitz into (X, d_∞) . We will see later in the proof of Theorem 4.16, which is needed to complete the proof of Theorem 1.1, that we can sometimes find an integral current, T_∞ , on (X, d_∞) and later show $T_\infty = T$ in the following sense:

Definition 4.9. *If we have an integral current T on (X, d_a) and there is an integral current, S , on (X, d_b) where $d_b \geq d_a$ such that*

$$(4.31) \quad S(\pi_0, \pi_1, \dots, \pi_m) = T(\pi_0, \pi_1, \dots, \pi_m).$$

for any tuple of d_a Lipschitz functions on X , then we say that $S = T$ viewed as an integral current on (M, d_a) . This means we can use any parametrizations of S and ∂S with charts into (X, d_b) as parametrizations for T and ∂T with the same charts and weights into (X, d_a) respectively. So we can view T as an integral current on (X, d_b) .

4.4. Weak Convergence of Currents. A sequence T_j converges weakly as currents in (Z, d_Z) to an integral current, T_∞ iff

$$(4.32) \quad T_j(\pi_0, \pi_1, \dots, \pi_m) \rightarrow T_\infty(\pi_0, \pi_1, \dots, \pi_m)$$

for all tuples, $(\pi_0, \pi_1, \dots, \pi_m)$ on (Z, d_Z) .

We will apply the following powerful theorem of Ambrosio-Kirchheim:

Theorem 4.10 (Ambrosio-Kirchheim). *If Z is compact and T_j have uniform upper bounds*

$$(4.33) \quad \mathbf{M}(T_j) \leq V_0 \text{ and } \mathbf{M}(\partial T_j) \leq A_0$$

then a subsequence T_{j_k} converges weakly to an integral current, T_∞ , and the mass measure is lower semicontinuous:

$$(4.34) \quad \|T_\infty\|(U) \leq \liminf_{k \rightarrow \infty} \|T_{j_k}\|(U)$$

for any open set $U \subset Z$.

4.5. Integral Current Spaces. In [SW11], Sormani and Wenger define an m -dimensional *integral current space*, (X, d, T) , as a metric space, (X, d) , with an integral current, T , defined on the closure of (X, d_X) such that the $\text{set}(T) = X$ where

$$(4.35) \quad \text{set}(T) = \left\{ x \in \bar{X} : \liminf_{r \rightarrow 0} \|T\|_d(B_d(x, r))/r^m > 0 \right\}.$$

Such metric spaces (X, d) are rectifiable and parametrized by the charts that parametrize their integral current structure, T .

Given a smooth compact oriented m -dimensional Riemannian manifold with boundary, (M, g) , it can be viewed as an integral current space, $(M, d_g, [[M]])$. Here the mass measure is the volume measure,

$$(4.36) \quad \mathbf{M}_{d_g}(M) = \text{vol}_g(M),$$

and

$$(4.37) \quad \text{set}_{d_g}([[M]]) = \left\{ x \in M : \liminf_{r \rightarrow 0} \text{vol}_g(B_{d_g}(x, r))/r^m > 0 \right\} = M.$$

The following lemma depicted in Figure 4.1 is useful for constructing examples:

Lemma 4.11. *Suppose the sphere \mathbb{S}^2 is endowed with the standard round metric, $g_{\mathbb{S}^2}$, and the singular metric, g_h ,*

$$(4.38) \quad g_h = dr^2 + \sin^2(r) d\theta^2 + (h'(r))^2 dr^2 \geq g_{\mathbb{S}^2} = dr^2 + \sin^2(r) d\theta^2$$

defined on a sphere parametrized by $r \in [0, \pi]$ and $\theta \in [0, 2\pi]$, where $h(r(x))$ is smooth on $r^{-1}(0, \pi)$, increasing on $[0, \pi/2]$ and decreasing on $[\pi/2, \pi]$ with $h(r) = h(\pi - r)$, but only continuous at the poles p_0, p_π where $r = 0, \pi$

respectively. We can define a unique compact metric space, (\mathbb{S}^2, d_h) , using g_h -lengths of curves, C , such that

$$(4.39) \quad C : (0, 1) \rightarrow \mathbb{S}^2 \setminus \{p_0, p_\pi\}$$

with bounded radial arclength functions from the poles $p = p_0, p_\pi$,

$$(4.40) \quad s_p(r(x)) = d_h(x, p) = \int_0^{r(x)} \sqrt{1 + h'(r)^2} dr \text{ with } \lim_{r \rightarrow 0} s_p(r) = 0.$$

So that (\mathbb{S}^2, d_h) is the metric completion of $(\mathbb{S}^2 \setminus \{p_0, p_\pi\}, d_{g_h})$.

We can find a parametrization of the standard integral current $[[\mathbb{S}^2]]$ on $(\mathbb{S}^2, d_{\mathbb{S}^2})$ by a countable collection of Lipschitz charts into (\mathbb{S}^2, d_h) which map onto $r^{-1}(0, \pi)$ so that we have an integral current space

$$(4.41) \quad (\text{set}_{d_h}([[\mathbb{S}^2]]), d_h, T_h) \text{ where } r^{-1}(0, \pi) \subset \text{set}_{d_h}(T_h) \subset \mathbb{S}^2,$$

and where $T_h = [[\mathbb{S}^2]]$ in the sense of Definition 4.9. So $\text{spt}_{d_h}(T_h) = \mathbb{S}^2$ and $\partial T_h = 0$. The poles are included in $\text{set}_{d_h}(T_h)$ iff

$$(4.42) \quad \lim_{r \rightarrow 0} |h'(r)| < \infty.$$




FIGURE 4.1. Lemma 4.11 applied with $h = h_{cone}$ of Example 4.12 on the left (which will include the singular poles) and with $h = h_{cusp}$ of Example 4.13 on the right (which will not include the singular poles due to the cusp).

Proof. We easily see that we can define d_h away from the poles using lengths of curves that avoid the poles. Near p_0 , we have radial arclengths,

$$(4.43) \quad s_{p_0}(x) = \int_0^{r(x)} \sqrt{1 + h'(r)^2} dr \leq \int_0^{r(x)} 1 + |h'(r)| dr < \infty$$

by the properties of h . So

$$(4.44) \quad \lim_{r \rightarrow 0} s_{p_0}(r) = \lim_{r \rightarrow 0} r + |h(r) - h(0)| = 0$$

and similarly for p_π . This allows us to define (\mathbb{S}^2, d_h) as in (4.39) so that the identity map to the standard round sphere is a homeomorphism and so $(\mathbb{S}^2, d_{\mathbb{S}^2})$ is compact with bounded diameter.

The usual integral current, $[[\mathbb{S}^2]]$, on the standard round sphere $(\mathbb{S}^2, d_{\mathbb{S}^2})$ can be defined using the standard single (r, θ) chart,

$$(4.45) \quad \psi : [0, \pi] \times [0, 2\pi] \rightarrow \mathbb{S}^2,$$

with multiplicity one, but this may not be a Lipschitz chart into (\mathbb{S}^2, d_h) due to the singularities at the poles.

Instead we define a new parametrization with a countable collection of multiplicity one charts avoiding the poles,

$$(4.46) \quad \psi_i : U_i \rightarrow \mathbb{S}^2 \text{ where } \psi_i(r, \theta) = \psi(r, \theta),$$

and where U_i are defined inductively by

$$(4.47) \quad U_1 = [\delta_1, \pi - \delta_1] \times [0, 2\pi) \text{ and } U_{i+1} = [\delta_{i+1}, \pi - \delta_{i+1}] \times [0, 2\pi) \setminus U_i$$

where δ_i decrease to 0 so that these are Lipschitz charts in (\mathbb{S}^2, d_h) . Then

$$(4.48) \quad T_h = \sum_{i=1}^{\infty} \psi_{i\#} [[U_i]]$$

is a rectifiable current on (\mathbb{S}^2, d_h) because

$$(4.49) \quad \sum_{i=1}^{\infty} \mathcal{H}^2(\psi_i(U_i)) \leq \text{vol}_{g_h}(\mathbb{S}^2) < \infty.$$

We see that $T_h = [[\mathbb{S}^2]]$ in the sense of Definition 4.9 because

$$(4.50) \quad T_h(\pi_0, \pi_1, \pi_2) = [[\mathbb{S}^2]](\pi_0, \pi_1, \pi_2)$$

for any tuple in $(\mathbb{S}^2, d_{\mathbb{S}^2})$, since missing two points does not affect the integration of tuples. So $(\text{set}_{d_h}(T_h), d_h, T_h)$ is an integral current space. Furthermore

$$(4.51) \quad \partial T_h = \partial [[\mathbb{S}^2]] = 0.$$

In order to determine which points lie in $\text{set}_{d_h}(T_h)$ we use (4.21) and the fact that the mass measure is the volume measure as stated above (4.36). It is easy to see that points where g_h is smooth lie in the set, so we conclude (4.41) holds. Taking either pole $p \in \{p_0, p_\pi\}$ and $r = r_p$, and applying

l'Hopital's rule twice, we have

$$(4.52) \quad \lim_{R \rightarrow 0} \frac{1}{R^2} \text{vol}_{g_h}(B_{d_h}(p, R)) = \lim_{R \rightarrow 0} \frac{1}{R^2} \int_0^R 2\pi r(s) ds = \lim_{s \rightarrow 0} \frac{2\pi r(s)}{2R}$$

$$(4.53) \quad = \lim_{s \rightarrow 0} \pi r'(s) = \lim_{s \rightarrow 0} \pi(1 + (h'(r))^2)^{-1/2}.$$

So $p \in \text{set}_{d_0}(T_h)$ iff this limit is > 0 which happens iff (4.42) holds. \square

Example 4.12. *If we consider g_h of Lemma 4.11 defined using*

$$(4.54) \quad h(r) = h_{\text{cone}}(r) = 1 - |\sin(r)|$$

then $(\mathbb{S}^2, d_{\text{cone}})$ is a compact metric space with conical singularities, and $(\text{set}_{d_{\text{cone}}}([\mathbb{S}^2]), d_h, [[\mathbb{S}^2]])$ is an integral current space with

$$(4.55) \quad \text{set}([\mathbb{S}^2]) = \mathbb{S}^2.$$

as depicted in Figure 4.1 because

$$(4.56) \quad \lim_{r \rightarrow 0} |h'(r)| = \lim_{r \rightarrow 0} \cos(r) = 1 < \infty.$$

Example 4.13. *If we consider $g_{\text{cusp}} = g_h$ of Lemma 4.11 defined using*

$$(4.57) \quad h(r) = h_{\text{cusp}}(r) = 1 - \sqrt{\sin(r)} = 1 - (\sin^2(r))^{1/4}$$

then $(\mathbb{S}^2, d_{\text{cusp}})$ is a compact metric space with cusp singularities at the poles, and the triple $(\text{set}_{d_{\text{cusp}}}([\mathbb{S}^2]), d_h, [[\mathbb{S}^2]])$ is an integral current space with

$$(4.58) \quad \text{set}_{d_{\text{cusp}}}([\mathbb{S}^2]) = \mathbb{S}^2 \setminus \{p_0, p_\pi\}.$$

as depicted in Figure 4.1 because

$$(4.59) \quad \lim_{r \rightarrow 0} |h'(r)| = \lim_{r \rightarrow 0} \frac{1}{2} \frac{\cos(r)}{\sqrt{\sin(r)}} = \infty.$$

Theorem 1.1 is careful with the description of the SWIF limit because the space, $(M, d_\infty, [[M]])$, might not be an integral current space, as it might fail to have $\text{set}([[M]]) = M$ as in Example 4.14 depicted in Figure 4.2.

Example 4.14. *Let us consider the sphere, \mathbb{S}^2 with a sequence of metric tensors, $g_j = g_{h_j}$ as in Lemma 4.11 defined using*

$$(4.60) \quad h_j(r) = 1 + \frac{1}{j} - \left(\frac{1}{j^4} + \sin^2(r)\right)^{1/4}.$$

Note that $h_j \circ r : \mathbb{S}^2 \rightarrow [0, 1]$ are smooth functions because $\sin^2(r)$ is a smooth nonnegative function on a sphere even at the poles where $r = 0, \pi$ and $u^{1/4}$ is smooth for $u > 0$. In fact

$$(4.61) \quad |h'_j(r)| = \frac{1}{4} \left(\frac{1}{r^4} + \sin^2(r) \right)^{-3/4} (2 \sin(r) \cos(r)).$$

so

$$(4.62) \quad |\nabla(h_j \circ r)| = |h'_j(r)| \leq |h'_{j+1}(r)| \leq |h'_{cusp}(r)|.$$

Thus we have a monotone increasing sequence of metric tensors as described in Remark 2.4. We have uniformly bounded radial arclength parameters. The radial arclengths,

$$(4.63) \quad s_j(x) = \int_0^{r(x)} \sqrt{1 + h'_j(r)^2} dr \leq \int_0^{r(x)} 1 + |h'_{cusp}(r)| dr,$$

are uniformly bounded, so we have a uniform upper bound on diameter.

Observe that $h_j \rightarrow h_{cusp}$ of Example 4.13 smoothly away from the poles. So $d_j \rightarrow d_{cusp}$ pointwise away from the poles. Since the arclength parameters also converge, we see that $d_j \rightarrow d_{cusp}$ on $\mathbb{S}^2 \times \mathbb{S}^2$. Since $(\mathbb{S}^2, d_{\mathbb{S}^2})$ is compact we have all the hypotheses of Theorem 1.1. This is an example where the SWIF limit is a proper subset of the GH limit:

$$(4.64) \quad \text{set}_{d_{cusp}}([\mathbb{S}^2]) \neq \mathbb{S}^2.$$




FIGURE 4.2. Example 4.2 has $d_j \leq d_{j+1}$ because $|h'_j| \leq |h'_{j+1}|$ and converges to Example 4.13 which has the poles with cusp singularities removed.

4.6. Intrinsic Flat Distance. In [SW11], Sormani and Wenger define the intrinsic flat distance between two integral current spaces as follows:

Definition 4.15.

$$(4.65) \quad d_{SWIF}((X_a, d_a, S_a), (X_b, d_b, S_b)) = \inf\{d_F^Z(f_{a\#}(S_a), f_{b\#}(S_b))\}$$

where the infimum is taken over all complete metric spaces, (Z, d_Z) , and over all distance preserving maps,

$$(4.66) \quad f_a : (X_a, d_a) \rightarrow (Z, d_Z) \text{ and } f_b : (X_b, d_b) \rightarrow (Z, d_Z).$$

Here d_F^Z denotes the Flat distance between integral currents T_a, T_b on (Z, d_Z) :

$$(4.67) \quad d_F^Z(T_a, T_b) = \inf\{\mathbf{M}(A) + \mathbf{M}(B) : A + \partial B = T_a - T_b\}$$

where the infimum is taken over all integral currents A, B on (Z, d_Z) .

We say that a sequence of integral current spaces converges in the intrinsic flat sense,

$$(4.68) \quad (X_j, d_j, S_j) \xrightarrow{\text{SWIF}} (X_\infty, d_\infty, S_\infty),$$

if

$$(4.69) \quad d_{\text{SWIF}}((X_j, d_j, S_j), (X_\infty, d_\infty, S_\infty)) \rightarrow 0.$$

In [SW11], Sormani-Wenger prove this implies there is a common complete metric space, (Z, d_Z) , and distance preserving maps, $f_j : X_j \rightarrow Z$ such that, $T_j = f_{j\#}S_j$ converge weakly to $T_\infty = f_{\infty\#}S_\infty$ as integral currents on Z . Thus by Ambrosio-Kirchheim semicontinuity of mass,

$$(4.70) \quad \liminf_{j \rightarrow \infty} \mathbf{M}_{d_j}(S_j) = \liminf_{j \rightarrow \infty} \mathbf{M}_{d_Z}(T_j) \geq \mathbf{M}_{d_Z}(T_\infty) = \mathbf{M}_{d_\infty}(S_\infty).$$

We say the sequence of integral current spaces converges in the *volume preserving intrinsic flat sense* if we have (4.68) and

$$(4.71) \quad \mathbf{M}_{d_j}(S_j) \rightarrow \mathbf{M}_{d_\infty}(S_\infty).$$

This notion was first studied by Portegies in [Por15]. Additional work was completed by Jauregui-Lee in [JL21]. The consequences of \mathcal{VF} convergence are most recently reviewed by Jauregui-Perales-Portegies in [JPP22].

4.7. Intrinsic Flat Convergence Theorem. We can now state our SWIF convergence theorem:

Theorem 4.16. *Suppose (X, d_j) is a monotone increasing sequence of metric spaces satisfying all the hypotheses of Theorem 3.2 so that $d_j \rightarrow d_\infty$ uniformly and*

$$(4.72) \quad (X, d_j) \xrightarrow{GH} (X, d_\infty) \text{ which is compact.}$$

If there is a current, T , which is an integral current structure for (X, d_j) ,

$$(4.73) \quad \text{set}_{d_j}(T) = X \quad \forall j \in \mathbb{N},$$

with uniform upper bounds on total mass,

$$(4.74) \quad \mathbf{M}_{d_j}(T) \leq V_0 \text{ and } \mathbf{M}_{d_j}(\partial T) \leq A_0 \quad \forall j \in \mathbb{N},$$

then

$$(4.75) \quad (X, d_j, T) \xrightarrow{\mathcal{F}} (M_\infty, d_\infty, T_\infty).$$

where $T_\infty = T$ in the sense defined in Definition 4.9 and

$$(4.76) \quad M_\infty = \text{set}_{d_\infty}(T_\infty) \subset X \text{ with } \overline{M_\infty} = X.$$

Furthermore,

$$(4.77) \quad \mathbf{M}_{d_\infty}(T_\infty) = \lim_{j \rightarrow \infty} \mathbf{M}_{d_j}(T).$$

so we have volume preserving SWIF convergence.

Recall that in Example 4.14 we saw that the SWIF limit, M_∞ , might be a proper subset of the Gromov-Hausdorff limit, X .

4.8. The Proof of Riemannian Theorem 1.1. Before we prove Theorem 4.16, we show in this subsection how it can be applied to prove Theorem 1.1:

Proof. Note that the hypotheses of Theorem 1.1 imply the hypotheses of Theorem 3.2 and Theorem 4.16 as follows:

- Monotonicity of metric tensors in (1.1) implies monotonicity of distances as in Definition 2.2 by Lemma 2.3 as in Remark 2.4.
- Adding the diameter bound in (1.2) implies the pointwise convergence of $d_j \rightarrow d_\infty$ in Lemma 2.5 and the diameter hypothesis of Theorem 3.2.
- The assumption that (M, d_∞) is compact in Theorem 1.1 implies the compactness assumption in Theorem 3.2, so that we have uniform convergence as in Proposition 3.6 and a common metric space Z_j for (M, d_j) and (M, d_∞) as in Proposition 3.9 and $(M, d_j) \xrightarrow{\text{GH}} (M, d_\infty)$ as in Theorem 3.2.
- The hypothesis that (M, g_j) are smooth oriented Riemannian manifolds in Theorem 1.1 implies $(M, d_{g_j}, [[M]])$ are integral current spaces with $T = [[M]]$ by (4.37).
- The volume bounds in (1.4) imply the mass bounds in (4.74) by (4.36).

Thus we may apply the Theorem 4.16 and (4.36) to conclude the volume preserving intrinsic flat convergence in (1.5) to a limit space satisfying (1.6).

□

4.9. Finding the Current Structure of the SWIF Limit for Theorem 4.16.

In [SW11], Sormani-Wenger proved that when one has a sequence of integral current spaces which converge in the Gromov-Hausdorff sense and have uniformly bounded total mass as in (4.74), then SWIF limit of the integral current spaces is a subset of the Gromov-Hausdorff limit. For Theorem 4.16 we wish to show the closure of the SWIF limit is the GH limit, so we need stronger control on the SWIF limit's current structure. To achieve this, we will repeat the steps of the argument in [SW11] using the special properties of the common metric space, (Z, d_Z) , that we constructed in Proposition 3.9. See Figure 4.3.




FIGURE 4.3. The distance preserving maps from Proposition 3.9.

Proposition 4.17. *Under the hypotheses Theorem 4.16, we have the metric space (Z, d_Z) as constructed in Proposition 3.9 depicted in Figure 4.3 with distance preserving maps,*

$$(4.78) \quad \chi_j = \zeta_j \circ f_{j,(j,\infty)} : (X, d_j) \rightarrow (Z_{j,\infty}, d_{Z_{j,\infty}}) \rightarrow (Z, d_Z),$$

$$(4.79) \quad \chi_\infty = \zeta_j \circ f_{\infty,(j,\infty)} : (X, d_\infty) \rightarrow (Z_{j,\infty}, d_{Z_{j,\infty}}) \rightarrow (Z, d_Z),$$

such that the pushforwards

$$(4.80) \quad T_j = \chi_{j\#}(T) = \zeta_{j\#}(f_{j\#}(T))$$

are integral currents in Z whose supports are

$$(4.81) \quad \text{spt}(T_j) = \zeta_j(f_j(X)) = \zeta_j(X \times \{h_j\}) \subset \zeta_j(Z_{j,\infty}) \subset Z$$

and have

$$(4.82) \quad \mathbf{M}_{d_Z}(T_j) = \mathbf{M}_{d_j}(T) \leq V_0 \text{ and } \mathbf{M}_{d_Z}(\partial T_j) = \mathbf{M}_{d_j}(\partial T) \leq A_0.$$

Furthermore, there is a subsequence T_{j_k} which converges weakly as currents in (Z, d_Z) to an integral current, T_∞ , whose support lies on the central spine in Z ,

$$(4.83) \quad \text{spt}(T_\infty) = Z_0 = \chi_\infty(X) = \zeta_j(X \times \{0\}) \subset \zeta_j(Z_{j,\infty}) \subset Z.$$

Finally $T = \chi_{\infty\#}^{-1}T_\infty$ is an integral current on (X, d_∞) as in Definition 4.9, and

$$(4.84) \quad \mathbf{M}_{d_\infty}(T) = \lim_{k \rightarrow \infty} \mathbf{M}_{d_{j_k}}(T).$$

Note that before we prove this proposition, we do not yet know that T is an integral current on (X, d_∞) . See Remark 4.8. After we prove this proposition we will apply it to prove the same statement holds without needing a subsequence in Corollary 4.18.

Proof. Since the maps in (4.78) are distance preserving, the pushforward integral currents T_j on (Z, d_Z) defined as in (4.80), satisfy (4.82) and (4.81) following the definitions of pushforward and mass (cf. [SW11]).

Since (Z, d_Z) is compact, we can apply the Ambrosio-Kirchheim Compactness Theorem, Theorem 4.10, to conclude that there is a subsequence T_{j_k} which converges weakly as currents in (Z, d_Z) to an integral current, T_∞ .




FIGURE 4.4. Here we see integral currents T_j in Z and $B_{d_Z}(z, t)$ of the proof that $\text{spt}(T_\infty) \subset Z_0 = \chi_\infty(X) \subset Z$ as claimed in (4.85).

First we claim

$$(4.85) \quad \text{spt}(T_\infty) \subset Z_0 = \chi_\infty(X) = \zeta_j(X \times \{0\}) \subset Z_{j_\infty} \subset Z.$$

Take any $z \in Z \setminus Z_0$ as in Figure 4.4. Then there exists $j_z \in \mathbb{N}$ such that $z = \zeta_{j_z}(f_{j_z}(x, t))$ where $t > 0$. Then by the definition of d_Z in Proposition 3.9, the ball:

$$(4.86) \quad B_{d_Z}(z, t) \subset \zeta_{j_z}(f_{j_z}(X \times (0, h_{j_z}])) \subset \zeta_{j_z}(Z_{j_z}) \setminus Z_0.$$

In particular

$$(4.87) \quad B_{d_Z}(z, t) \cap \zeta_j(f_j(X \times \{h_j\})) = \emptyset$$

for all $j > j_z$. By the lower semicontinuity of mass under weak convergence as in (4.34) we have

$$(4.88) \quad \|T_\infty\|(B_{d_Z}(z, t)) \leq \liminf_{j \rightarrow \infty} \|T_j\|(B_{d_Z}(z, t)) = 0$$

so $z \notin \text{spt}(T_\infty)$. Thus $\text{spt}(T) \subset Z_0$.

Since $\chi_\infty : (X, d_\infty) \rightarrow (Z_0, d_Z)$ is an isometry, we can define:

$$(4.89) \quad S = \chi_{\infty\#}^{-1} T_\infty$$

as an integral current on (X, d_∞) .

We claim that $S = T$ in the sense of Definition 4.9. Given any tuple, $(\pi_0, \pi_1, \dots, \pi_m)$ of Lipschitz functions on (X, d_j) , we need only show

$$(4.90) \quad T(\pi_0, \pi_1, \dots, \pi_m) = S(\pi_0, \pi_1, \dots, \pi_m).$$

By Lemma 4.1 these π_i are also tuples of Lipschitz functions on (X, d_∞) , so $S(\pi_0, \dots, \pi_m)$ is well defined.




FIGURE 4.5. Here we see $W_j \subset Z$ of Lemma 3.10 in dark gray, the rest of Z is lighter, and $Z_0 \subset W_j \subset Z$ is in black with the isometry, χ_∞^{-1} , to (X, d_∞) .

Recall the set $W_j \subset Z$ from Lemma 3.10, Lemma 4.3, and Remark 4.4 depicted in Figure 4.5. By Lemma 4.3, we define a tuple of Lipschitz functions, $(\tilde{\pi}_0, \tilde{\pi}_1, \dots, \tilde{\pi}_m)$, on the subset, (W_j, d_Z) , inside (Z, d_Z) , such that

$$(4.91) \quad \tilde{\pi}_i(\zeta_{j_k}(x, t)) = \pi_i(x) \quad \forall x \in X, t \in [0, h_{j_k}], j_k \geq j.$$

By (4.89) and the fact that

$$(4.92) \quad \tilde{\pi}_i(\zeta_{j_k}(x, t)) = \pi_i(x) = \tilde{\pi}_i(\chi_\infty(x)) \quad \forall x \in X, t \in [0, h_{j_k}], j_k \geq j,$$

we have by definition of pushforward that

$$(4.93) \quad S(\pi_0, \pi_1, \dots, \pi_m) = (\chi_{\infty\#}^{-1} T_\infty)(\tilde{\pi}_0 \circ \chi_\infty, \tilde{\pi}_1 \circ \chi_\infty, \dots, \tilde{\pi}_m \circ \chi_\infty)$$

$$(4.94) \quad = T_\infty(\tilde{\pi}_0, \tilde{\pi}_1, \dots, \tilde{\pi}_m).$$

Note that for any $j_k \geq j$ we have

$$(4.95) \quad \text{spt}(T_{j_k}) \subset \zeta_{j_k}(Z_{j_k}) \subset W_j \subset Z.$$

So T_{j_k} converge weakly as currents to T_∞ in (W_j, d_Z) : for any Lipschitz tuple, ω , on (W_j, d_Z) , we have

$$(4.96) \quad T_{j_k}(\omega) \rightarrow T_\infty(\omega).$$

Thus we have

$$(4.97) \quad S(\pi_0, \pi_1, \dots, \pi_m) = \lim_{k \rightarrow \infty} T_{j_k}(\tilde{\pi}_0, \tilde{\pi}_1, \dots, \tilde{\pi}_m).$$

On the other hand, by (4.91), we have,

$$(4.98) \quad \tilde{\pi}_i(\zeta_{j_k}(x, t)) = \pi_i(x) = \tilde{\pi}_i(\zeta_{j_k}(f_{j_k}(x))) \quad \forall x \in X, t \in [0, h_{j_k}], j_k \geq j.$$

Thus $T_{j_k} = \zeta_{j_k\#} f_{j_k\#} T$ and we have

$$(4.99) \quad T_{j_k}(\tilde{\pi}_0, \dots, \tilde{\pi}_m) = (\zeta_{j_k\#} f_{j_k\#} T)(\tilde{\pi}_0 \circ \zeta_{j_k} \circ f_{j_k}, \dots, \tilde{\pi}_m \circ \zeta_{j_k} \circ f_{j_k})$$

$$(4.100) \quad = T(\pi_0, \dots, \pi_m).$$

Substituting this into (4.97), and taking the limit $j_k \rightarrow \infty$, we reach our claim in (4.90). We conclude that $S = T$ in the sense of Definition 4.9.

Now we claim $\text{spt}_{d_Z}(T_\infty) = Z_0$. Since we know $T_\infty = \chi_{\infty\#} T$, this is the same as showing $\text{spt}_{d_\infty}(T) = X$. Given any $x \in X$ and any $r > 0$ we have

$$(4.101) \quad \|T\|_{d_\infty}(B_{d_\infty}(x, r)) \geq \|T\|_{d_j}(B_{d_j}(x, r))$$

by Lemma 4.7. By our hypotheses, we have uniform convergence of the distance functions, $|d_j - d_\infty| < \epsilon_j \rightarrow 0$, so for all j such that $\epsilon_j < r/2$

$$(4.102) \quad B_{d_j}(x, r) \supset B_{d_j}(x, r - \epsilon_j).$$

Combining this with the above,

$$(4.103) \quad \|T\|_{d_\infty}(B_{d_\infty}(x, r)) \geq \|T\|_{d_j}(B_{d_j}(x, r - \epsilon_j)) > 0.$$

Thus $\text{spt}_{d_\infty}(T) = X$ and $\text{spt}_{d_Z}(T_\infty) = Z_0$.

Finally, by Ambrosio-Kirchheim's semicontinuity of mass combined with the fact that ζ_∞ is distance preserving we have

$$(4.104) \quad \mathbf{M}_{d_\infty}(T) = \mathbf{M}_{d_Z}(T_\infty) \leq \liminf_{k \rightarrow \infty} \mathbf{M}_{d_Z}(T_{j_k}) = \liminf_{k \rightarrow \infty} \mathbf{M}_{d_{j_k}}(T).$$

Yet by Lemma 4.7, we know

$$(4.105) \quad \mathbf{M}_{d_\infty}(T) \geq \mathbf{M}_{d_{j_k}}(T)$$

so we have mass convergence. \square

Corollary 4.18. *Under the hypotheses of Proposition 4.17, we have the conclusions $T_j = \chi_{j\#} T$ converge weakly to $T_\infty = \chi_{\infty\#} T$ as integral currents in (Z, d_Z) and*

$$(4.106) \quad \mathbf{M}_{d_\infty}(T) = \lim_{j \rightarrow \infty} \mathbf{M}_{d_j}(T).$$

without taking a subsequence.

Proof. First note that (4.106) follows from (4.84) combined with the monotonicity and Lemma 4.7.

To prove weak convergence, assume on the contrary that there is a tuple, (π_0, \dots, π_m) , on (Z, d_Z) , and a $\delta > 0$ with a subsequence, we also denote T_j , such that

$$(4.107) \quad |T_j(\pi_0, \dots, \pi_m) - T_\infty(\pi_0, \dots, \pi_m)| < \delta \quad \forall j \in \mathbb{N}.$$

Note that this subsequence T_j satisfies the hypotheses of Proposition 4.17, so it has a subsequence which converges weakly to T , contradicting (4.107). \square

4.10. Proving SWIF Convergence. In this section we complete the proof of Theorem 4.16:

Proof. By Corollary 4.18 of Proposition 4.17, we have pushforwards, $T_j = \chi_{j\#}T$, converging weakly to $T_\infty = \chi_{\infty\#}T$ as integral currents in Z .

By the uniform upper bound on total mass, the compactness of (Z, d_Z) and Wenger's Flat=Weak Convergence theorem in [Wen07] (applied to metric spaces as explained in [SW11]) there exist integral currents, B_j, A_j , on (Z, d_Z) such that

$$(4.108) \quad \partial B_j + A_j = T_j - T_\infty \text{ and } \mathbf{M}_{d_Z}(B_j) + \mathbf{M}_{d_Z}(A_j) \rightarrow 0$$

so that

$$(4.109) \quad d_F^Z(\chi_{j\#}T, \chi_{\infty\#}T) \rightarrow 0.$$

Taking $M_\infty = \text{set}_{d_\infty}(T)$ and applying the fact that χ_j and χ_∞ are distance preserving maps, we have

$$(4.110) \quad d_{SWIF}((X, d_j, T), (M_\infty, d_\infty, T)) \rightarrow 0.$$

Since we have the convergence of masses in Corollary 4.18, we have \mathcal{VF} convergence as well. \square

Remark 4.19. *It is an exercise to circumvent applying Wenger's Flat=Weak Convergence Theorem in the above proof, by explicitly constructing a sequence of integral currents, A_j and B_j , on (Z, d_Z) using the special properties of (Z, d_Z) proven in Prop 3.9. Since we have shown T is an integral current on (X, d_∞) , there are integers, a_i , and charts*

$$(4.111) \quad \psi_i : (U_i \subset \mathbb{R}^m, d_{\mathbb{R}^m}) \rightarrow (X, d_\infty) \text{ with } \sum_{i=1}^{\infty} |a_i| \mathcal{H}_{d_\infty}^m(\psi_i(U_i)) < \infty$$

such that

$$(4.112) \quad T = \sum_{i=1}^{\infty} a_i \psi_{i\#}[[U_i]],$$

and the same holds for ∂T . Then it is natural to define the integral current, B_j , using the same integers a_i and charts,

$$(4.113) \quad \tilde{\psi}_{i,j} : (U_i \times [0, h_j] \subset \mathbb{R}^{m+1}, d_{\mathbb{R}^{m+1}}) \rightarrow (Z, d_Z)$$

such that

$$(4.114) \quad \tilde{\psi}_{i,j}(v, t) = \zeta_j(\psi_i(v), t).$$

We then similarly construct the integral current, A_j , using the charts for ∂T (which are not necessarily the boundaries of charts of T). Next we would verify that both atlases of charts are Lipschitz and that A_j and B_j have finite masses satisfying

$$(4.115) \quad \mathbf{M}_{d_Z}(A_j) \leq C h_j \mathbf{M}_{d_\infty}(T) \text{ and } \mathbf{M}_{d_Z}(B_j) \leq C h_j \mathbf{M}_{d_\infty}(T).$$

Most difficult is showing $A_j + \partial B_j = T_j - T_\infty$ by texting against arbitrary Lipschitz tuples on (Z, d_Z) . It may be necessary to restrict to W_j to get everything to work.

5. OPEN PROBLEMS

We know that without the compactness assumption in our Theorem 1.1 and Theorem 4.16, that we may not have Gromov-Hausdorff Convergence and a common compact metric space. However we know that by Wenger's Compactness Theorem the diameter and mass bounds imply that there is a subsequence converging to a SWIF limit,

$$(5.1) \quad (X, d_{j_k}, T) \xrightarrow{\text{SWIF}} (X'_\infty, d'_\infty, T'_\infty).$$

Is it possible that with monotonicity, $d_j \leq d_{j+1}$, one can show any of the following:

- the sequence itself converges,
- there is volume preserving intrinsic flat convergence,
- the limit, $X'_\infty \subset X$, with restricted metric, $d'_\infty = d_\infty$, possibly with closure, $\bar{X}_\infty = X$, and maybe $T_\infty = T$.

Note that assuming any of the following additional hypotheses might help:

- The integral current spaces, (X, d_j, T) , are Riemannian, $(M, d_{g_j}, [[M]])$, possibly without boundary.

- The integral current spaces, (X, d_j, T) , have empty boundary, $\partial T = 0$, so we don't have to use A_j in the proof of SWIF convergence.
- The integral current structure, T , is assumed to be an integral current on (X, d_∞) as an hypothesis, and that perhaps also that it has a bound on its multiplicity.

It would be interesting to find counter examples to any or all of these statements as well.

REFERENCES

- [AB24] Brian Allen and Edward Bryden. Volume above distance below with boundary II. *ArXiv:2406.11541*, 2024. [2](#)
- [AK00] Luigi Ambrosio and Bernd Kirchheim. Currents in metric spaces. *Acta Math.*, 185(1):1–80, 2000. [1](#), [4](#), [17](#), [18](#), [20](#)
- [AP22] Brian Allen and Raquel Perales. Intrinsic flat stability of manifolds with boundary where volume converges and distance is bounded below. *Arxiv: 2006.13030*, 2022. [2](#)
- [APS24] Brian Allen, Raquel Perales, and Christina Sormani. Volume above distance below. *Journal of Differential Geometry*, 126(3):837–874, 2024. [2](#), [9](#)
- [AS19] Brian Allen and Christina Sormani. Contrasting various notions of convergence in geometric analysis. *Pacific J. Math.*, 303(1):1–46, 2019. [2](#), [3](#)
- [AS20] Brian Allen and Christina Sormani. Relating notions of convergence in geometric analysis. *Nonlinear Anal.*, 200:111993, 33, 2020. [2](#)
- [DG95] Ennio De Giorgi. General Plateau problem and geodesic functionals. *Atti Sem. Mat. Fis. Univ. Modena*, 43(2):285–292, 1995. [4](#)
- [Edw75] David A. Edwards. The structure of superspace. In *Studies in Topology*, 1975. [1](#), [8](#)
- [Gro81] Mikhael Gromov. Groups of polynomial growth and expanding maps. *Inst. Hautes Études Sci. Publ. Math.*, 53:53–73, 1981. [1](#), [12](#)
- [Gro07] Misha Gromov. *Metric structures for Riemannian and non-Riemannian spaces*. Modern Birkhäuser Classics. Birkhäuser Boston, Inc., Boston, MA, english edition, 2007. Based on the 1981 French original, With appendices by M. Katz, P. Pansu and S. Semmes, Translated from the French by Sean Michael Bates. [1](#), [8](#)
- [HLS17] Lan-Hsuan Huang, Dan A. Lee, and Christina Sormani. Intrinsic flat stability of the positive mass theorem for graphical hypersurfaces of Euclidean space. *J. Reine Angew. Math.*, 727:269–299, 2017. [2](#)
- [JL21] Jeffrey L. Jauregui and Dan A. Lee. Lower semicontinuity of ADM mass under intrinsic flat convergence. *Calc. Var. Partial Differential Equations*, 60(5):Paper No. 193, 42, 2021. [2](#), [27](#)
- [JPP22] J. Jauregui, R. Perales, and J. Portegies. Semicontinuity of capacity under pointed intrinsic flat convergence. *arxiv*, 2022. [2](#), [27](#)
- [Kir94] Bernd Kirchheim. Rectifiable metric spaces: local structure and regularity of the Hausdorff measure. *Proc. Amer. Math. Soc.*, 121(1):113–123, 1994. [18](#)

- [LS13] Sajjad Lakzian and Christina Sormani. Smooth convergence away from singular sets. *Comm. Anal. Geom.*, 21(1):39–104, 2013. [2](#)
- [Por15] Jacobus W. Portegies. Semicontinuity of eigenvalues under intrinsic flat convergence. *Calc. Var. Partial Differential Equations*, 54(2):1725–1766, 2015. [1](#), [27](#)
- [Ron10] Xiaochun Rong. Convergence and collapsing theorems in Riemannian geometry. In *Handbook of geometric analysis, No. 2*, volume 13 of *Adv. Lect. Math. (ALM)*, pages 193–299. Int. Press, Somerville, MA, 2010. [1](#)
- [Sor17] Christina Sormani. *Scalar curvature and intrinsic flat convergence*, pages 288–338. Partial Differ. Equ. Meas. Theory. De Gruyter Open, Warsaw, 2017. [1](#)
- [ST24] Christina Sormani and Wenchuan Tian. Flat convergence for integral currents in metric spaces. *to appear soon*, 2024. [2](#)
- [STW24] Christina Sormani, Wenchuan Tian, and Changliang Wang. An extreme limit with nonnegative scalar. *Nonlinear Anal.*, 239:Paper No. 113427, 24, 2024. [2](#)
- [SW11] Christina Sormani and Stefan Wenger. The intrinsic flat distance between Riemannian manifolds and other integral current spaces. *J. Differential Geom.*, 87(1):117–199, 2011. [1](#), [2](#), [3](#), [7](#), [22](#), [26](#), [27](#), [29](#), [30](#), [33](#)
- [Wen07] Stefan Wenger. Flat convergence for integral currents in metric spaces. *Calc. Var. Partial Differential Equations*, 28(2):139–160, 2007. [1](#), [33](#)
- [Wen11] Stefan Wenger. Compactness for manifolds and integral currents with bounded diameter and volume. *Calc. Var. Partial Differential Equations*, 40(3-4):423–448, 2011. [2](#)

(R. Perales) CENTRO DE INVESTIGACIÓN EN MATEMÁTICAS, DE JALISCO S/N, VALENCIANA, GUANAJUATO, GTO. MEXICO. 36023

Email address: raquel.perales@cimat.mx

(C. Sormani) CUNY GRADUATE CENTER AND LEHMAN COLLEGE

Email address: sormanic@gmail.com