

# Chemical Continuous Time Random Walks under Anomalous Diffusion

Hong Zhang, Guohua Li

June 28, 2024

## Abstract

Chemical master equation plays an important role to describe the time evolution of homogeneous chemical system. In addition to the reaction process, it is also accompanied by physical diffusion of the reactants in complex system that is generally not homogeneous, which will result in non-exponential waiting times for particle reactions and diffusion. In this paper we shall introduce a chemical continuous time random walk under anomalous diffusion model based on renewal process to describe the general reaction-diffusion process in the heterogeneous system, where the waiting times are arbitrary distributed. According to this model, we will develop the systematic stochastic theory including generalizing the chemical diffusion master equation, deriving the corresponding mass action law, and extending the Gillespie algorithm. As an example, we analyze the monomolecular  $A \leftrightarrow B$  reaction-diffusion system for exponential and power-law waiting times respectively, and show the strong fractional memory effect of the concentration of the reactants on the history of the concentration in power-law case.

## 1 Introduction

Anomalous diffusion behaviors have attracted great interest in various fields of physics, biological, chemical, and environmental sciences. [1, 2] The main characteristics of anomalous diffusion is that the mean square displacement scales as a nonlinear power law in time, i.e.,  $\langle \Delta x^2 \rangle \sim t^\beta$ . Faster than linear scaling ( $\beta > 1$ ) is corresponding to as superdiffusion, slower than linear scaling  $0 < \beta < 1$  is referred to as subdiffusion, and when  $\beta = 1$  it is normal diffusion.

The continuous time random walk (CTRW) with long-tailed waiting time has been often evoked as a suitable description of anomalous diffusion. In this model, the particle begins its jump at time  $t = 0$  and then traps in a position for a random waiting time until it jumps away. In recent years there has been considerable effort to investigate the chemical reactions (e.g., reversible or irreversible conversion to a different species [3, 4, 5], spontaneous evanescence [6], fluorescence recovery after photobleaching [7], or linear reaction dynamics [8], etc.) under anomalous diffusion using different continuous time random walks models. In 2010, Fedotov considered three CTRW models: model  $A, B, C$  with nonlinear reactions under varying assumptions for the effect of reaction on random waiting time [9]. In 2013, Angstmann et al. generalized the model  $B$  where it is assumed that the created particles will draw a new waiting time to be in space- and time-dependent force fields, and applied this generalized model with space-dependent traps to obtain a fractional Fokker-Plank equation (FFPE) with space-dependent anomalous exponential for an ensemble of particles [10]. However, for the complex reaction-diffusion system, such as the system including more than three reactions each of which has many reactants and products performing anomalous diffusion, the CTRW approach is out

of work due to the technical difficulty that it is quite hard for the balance equation to remember all the effects of so many complex reactions on the probability evolution for every particle and of coupling relations between the reactions and anomalous diffusion.

Fortunately, there is a chemical master equation (CME) which can provide a stochastic approach to capture the time behavior of a spatially homogeneous chemical system in particle number space. The advantage of the CME is that it can deal with the system including many reactions, and has a firmer physical basis than the deterministic reaction-rate equations. Besides, the Gillespie algorithm based on the spatially homogeneous master equation is proved to be straightforward to implement on a digital computer. [11] Recently a generalized chemical diffusion master equation (CDME) is developed to describe the normal diffusion process for a given  $n$ -particle probability density coupled by the reaction dynamics similar in form to a chemical master equation by using creation and annihilation operators.[12] But up to now it is still a challenge to involve anomalous diffusion into CME.

On the other side, the system that both CME and CDME consider is the well-mixed chemical reaction system in which each reaction is generated by Poisson process, that corresponds to the exponential distributed waiting time. [14, 13] However, in various real-world systems the inter-reaction times typically obey non-Poissonian distribution. [15] In particular, inter-reaction times typically obey long-tailed distributions. Examples of long-tailed distributions of inter-reaction times include population dynamics, epidemic processes, finance, and so on. The non-Poissonian chemical process defines an inhomogeneous chemical continuous time random walk (CCTW) in particle number space. [16] Based on the CCTW Zhang et. al. proposed a stationary generalized chemical master equation which can model intracellular non-Markovian intracellular processes with molecular memory. [17-20]

In this paper we will consider the complex heterogeneous system involving both the arbitrary non-Poisson chemical process and the anomalous diffusion process of the particles. We shall overcome the technical difficulties to develop a systematic stochastic approach for the generalized CDME, the generalized mass law action, and the generalized Gillespie algorithm based on renewal processes used in CCTW, where inter-event times are independently generated from a given distribution [15, 21, 22], and on the continuous time random walk model, where it is assumed that the reactants can move from one space to the other, and at each space or site the reactants can react according to the chemical reaction law, but do not react in the process of jumping. Moreover, we will take the anomalous monomolecular  $A \leftrightarrow B$  reaction-diffusion system as an example, and obtain the corresponding mass action law for exponential and power-law distributed waiting times respectively, and show the strong fractional memory effect of the concentration of the reactants on the history of the concentration in the case of power-law distribution.

## 2 The renewal process

We assume that there are  $\gamma$  different species that participate in  $m$  different reactions and can diffuse in  $N$  positions. The chemical species are denoted by  $S_j$ , where  $j = 1, \dots, \gamma$ ; the  $v$  positions are denoted by  $x, y, z, \dots$ , the corresponding particle numbers at  $x$  at time  $t$  are denoted by  $k_{xj}$ . The state vector of particle numbers is a random  $v\gamma$ -dimensional vector  $k = (k_{x1}, k_{x2}, \dots, k_{x\gamma}, \dots, k_{z1}, k_{z2}, \dots, k_{z\gamma})^T$ , where the superscript  $T$  denotes the transpose. A single event of reaction and diffusion are characterized by the reaction and diffusion waiting times. We denote the probability density function (PDF) for the waiting time  $\tau_{ix}$  of reaction  $i$  at  $x$  by  $p_{ix}(t)$  and denote the PDF of the waiting time  $\tau_{dlx}$  for the particle  $l$  diffusing away from  $x$  by  $p_{dlx}(t)$ , respectively.

We now consider a special renewal process where  $N(t) = \sup\{n \in N : T_n \leq t\}$  is the number of renewal in  $[0, t]$  with  $N(0) = 0$  if  $X_1 < t$ , where  $T_n$  is the time after  $n$  renewal step and is the

sum  $T_n = \tau_1 + \tau_2 + \dots + \tau_n$ ,  $n \geq 1$  where  $\tau_1, \tau_2, \dots, \tau_n$  are a sequence of nonnegative independent distributions random variables. For each  $\tau_n, n \geq 1$  is the waiting time for the  $n$ -th update. It is assumed to be the minimum time of all waiting times of reactions and diffusions, that is,  $\tau_n = \min\{\tau_{ix}, \tau_{dlx}\} (i = 1, 2, \dots, m, l = 1, 2, \dots, \gamma, x = 1, 2, \dots, \nu)$ .

One can find  $N(t) = N_r(t) + N_d(t)$  where  $N_r(t)$  is the number of reaction renewals in  $[0, t]$ , and  $N_d(t)$  is the number of diffusion renewals in  $[0, t]$ . The state vector will change for each reaction renewal and each diffusion renewal. During a reaction  $i$  at  $x$  the loss (gain) in particle number  $k_{xj}$  is denoted by  $r_{ixj} \in N$  ( $p_{ixj} \in N$ ). These coefficients are typically, but need not be, given by the law of mass action. Thus, the impact of reaction  $i$  on the state space can be expressed as [16]

$$\sum_j r_{ixj} S_j \rightarrow \sum_j p_{ixj} S_j$$

The stoichiometric coefficients  $s_{ixj} = p_{ixj} - r_{ixj}$  denote the net change in each species  $j$  due to each reaction  $i$  at space  $x$ , and form the reaction transforming vector for the  $n$ -th step  $s_{ix} = (0, \dots, 0, s_{ix1}, s_{ix2}, \dots, s_{ix\gamma}, 0, \dots, 0)$ . For each particle  $j$ , its space at time  $t$  is  $Z_j(t) = Z_j(0) + \sum_{n=1}^{N_{d_j}} \xi_n$  where  $\xi_n$  is the jumping length for the  $n$ -th jumping step of particle  $j$ ,  $N_{d_j}(t)$  is the number of diffusion renewals of particle  $j$  in  $[0, t]$  and the relation with  $N_d(t)$  is  $\sum_j N_{d_j}(t) = N_d(t)$ . If in one step an  $i$  reactant moves from  $x$  to  $y$ , then after this step the number  $k_{xl}$  of the reactant  $l$  reduces 1 at  $x$  and  $k_{yl}$  adds 1 at  $y$ , so we can find the diffusion transforming vector  $s_{dlxy} = \binom{0, \dots, -1, 0, \dots, 1, 0, \dots, 0}{x_l, \dots, y_l}$  (all except  $(x-1)\gamma + l$ -th and  $(y-1)\gamma + l$ -th components are 0).

## 2.1 Distribution of the renewal waiting time

We now investigate the distribution of the renewal waiting time  $\tau_n$ . Recall that  $p_{ix}(t)$  is the waiting time PDF for one group of reaction  $i$  taking place at one site  $x$ , so the survival probability for such group does not react for time interval  $[0, t]$  is  $\int_t^\infty p_{ix}(t') dt'$ . Denoting the whole combinatorial number of groups of reactants for reaction  $i$  at space  $x$  by  $h_{ix}(k)$  [16], the PDF for only one group of reactants occurring at space  $x$  after waiting time  $t$  is as following

$$\psi_{ix}^r(t, k) = h_{ix}(k) p_{ix}(t) \left( \int_t^\infty p_{ix}(t') dt' \right)^{h_{ix}(k)-1}, \quad (1)$$

while the probability that the reaction  $i$  at space  $x$  does not occur after waiting  $t$  is

$$\Psi_{ix}^r(t, k) = \left( \int_t^\infty p_{ix}(t') dt' \right)^{h_{ix}(k)}. \quad (2)$$

Analogously,

$$\psi_{lx}^d(t, k) = k_{xl} p_{dlx}(t) \left( \int_t^\infty p_{dlx}(t') dt' \right)^{k_{xl}-1} \quad (3)$$

is the pdf that reactant  $l$  at space  $x$  occurs after waiting  $t$ , and

$$\Psi_{lx}^d(t, k) = \left( \int_t^\infty p_{dlx}(t') dt' \right)^{k_{xl}} \quad (4)$$

is the probability that all the reactants  $l$  at space  $x$  does not diffuse away after waiting time  $t$ . Note that in the above descriptions we assume that all random waiting times are independent, but once one reaction or diffusion occurs all waiting times in the system will be renewed.

If  $\tau_n$  is for a reaction  $i$  at  $x$ , then its PDF is

$$\begin{aligned} \phi_{ix}^r(t, k) &= \psi_{ix}^r(t) \prod_{j \neq i} \Psi_{jx}^r(t, k) \prod_{y \neq x; i=1,2,\dots,m} \Psi_{iy}^r(t, k) \\ &\cdot \prod_{x=1,2,\dots,v; l=1,2,\dots,\gamma} \Psi_{lx}^d(t, k). \end{aligned} \quad (5)$$

which denotes that when the system waits  $t$  the reaction  $i$  at space  $x$  first occurs but none of other reactions and diffusions have happened before this reaction, that is, the waiting time of this reaction in this renewal step is the minimum waiting time. Note that when the diffusion is not considered, and the chemical process focuses one space  $x$ ,  $\phi_{ix}^r(t, k)$  becomes  $\phi_{ix}^r(t, k) = \psi_{ix}^r(t) \prod_{j \neq i} \Psi_{jx}^r(t, k)$ , which is the non-Markovian case in Refs.[17, 16].

But if  $\tau_n$  is the waiting time of a diffusion  $i$  from  $x$ , then its PDF is

$$\begin{aligned} \phi_{lx}^d(t, k) &= \psi_{lx}^d(t, k) \prod_{l' \neq l} \Psi_{l'x}^d(t, k) \prod_{y \neq x; l=1,2,\dots,\gamma} \Psi_{ly}^d(t, k) \\ &\cdot \prod_{i=1,2,\dots,m; x=1,2,\dots,v} \Psi_{ix}^r(t, k) \end{aligned} \quad (6)$$

which denotes that when the system waits  $t$  the diffusion of  $i$  from  $x$  first occurs but none of other reactions and diffusion have happened before this diffusion, that is, the waiting time of this event in this renewal step is the minimum waiting time.

Besides, we also introduce

$$\begin{aligned} \Phi(t, k) &= \prod_{x=1,2,\dots,v; i=1,2,\dots,m} \Psi_{ix}^r(t, k) \\ &\cdot \prod_{x=1,2,\dots,v; l=1,2,\dots,\gamma} \Psi_{lx}^d(t, k), \end{aligned} \quad (7)$$

which is the probability that there is no diffusion or reaction in the system in the time interval  $[0, t]$ . One can find that

$$-\frac{\partial \Phi(t, k)}{\partial t} = \sum_{x=1}^v \sum_{i=1}^m \phi_{ix}^r(t, k) + \sum_{x=1}^v \sum_{l=1}^{\gamma} \phi_{lx}^d(t, k), \quad (8)$$

which means

$$\Phi(t, k) = \int_t^\infty \sum_{x=1}^v \sum_{i=1}^m \phi_{ix}^r(t', k) + \sum_{x=1}^v \sum_{l=1}^{\gamma} \phi_{lx}^d(t', k) dt', \quad (9)$$

and then

$$\begin{aligned} \Phi(u, k) &= \frac{1}{u} [1 - (\sum_{x=1}^v \sum_{i=1}^m \phi_{ix}^r(u, k) \\ &+ \sum_{x=1}^v \sum_{l=1}^{\gamma} \phi_{lx}^d(u, k))], \end{aligned} \quad (10)$$

where  $\Phi(u, k)$ ,  $\phi_{ix}^r(u, k)$  and  $\phi_{xl}^d(u, k)$  are the Laplace  $t \rightarrow u$  transform of  $\Phi(t, k)$ ,  $\phi_{ix}^r(t, k)$  and  $\phi_{xl}^d(t, k)$ , respectively.

### 3 Master equation

We then want to investigate the evolution of the state vector  $k$  at time  $t$  in this renewal process, that is, to obtain the master equation of the chemical continuous time random walks under anomalous diffusion model (see Figure.1).




Figure 1: The chemical continuous time random walks under anomalous diffusion model where the reactants wait some random time with arbitrary distribution at a space and then diffuse to other space with certain probability, and meanwhile, the reactants react according to chemical law at each space.

Let  $P(k, t)$  denote the probability distribution  $Prob[k(t) = k]$ . Firstly, we have the balance equation

$$P(k, t) = \int_0^t dt' \sum_{n=0}^{\infty} R_n(k, t') \Phi(t - t', k) \quad (11)$$

where  $R_n(k, t)$  is the joint density of arriving at  $k$  at time  $t$  after  $n$  steps, and satisfying

$$\begin{aligned} R_{n+1}(k, u) &= \sum_{i=1}^m \sum_{x=1}^v R_n(k - s_{ix}, u) \phi_{ix}^r(u, k - s_{ix}) \\ &+ \sum_{l=1}^{\gamma} \sum_{x=1}^v \sum_{y \neq x} [R_n(k - s_{dly}, u) \\ &\cdot \phi_{lx}^d(u, k - s_{dly}) \lambda(y - x)], \end{aligned} \quad (12)$$

for nonnegative integer  $n$  in Laplace space, and

$$R_0(k, t) = P(k, 0) \delta(t). \quad (13)$$

Here,  $R_n(k, u)$  is the Laplace  $t \rightarrow u$  transform of  $R_n(k, t)$ ,  $P(k, 0)$  is the initial distribution and  $\lambda(y - x)$  represents the transition probability for the reactant jumping from  $x$  to  $y$ .

Let  $R(k, u) = \sum_{n=0}^{\infty} R_n(k, u)$ . Then

$$\begin{aligned} R(k, u) &= \sum_{i=1}^m \sum_{x=1}^v R(k - s_{ix}, u) \phi_{ix}^r(u, k - s_{ix}) \\ &+ \sum_{l=1}^{\gamma} \sum_{x=1}^v \sum_{y \neq x} [R(k - s_{dly}, u) \\ &\cdot \phi_{lx}^d(u, k - s_{dly}) \lambda(y - x)] \\ &+ P(k, 0). \end{aligned} \quad (14)$$

Taking the Laplace transform of (14), and combining (10) we get

$$\begin{aligned}
P(k, u) &= \sum_{n=0}^{\infty} R_n(k, u) \Phi(u, k) = R(k, u) \Phi(u, k) \\
&= R(k, u) \frac{1}{u} \left[ 1 - \left( \sum_{x=1}^v \sum_{i=1}^m \phi_{ix}^r(u, k) \right. \right. \\
&\quad \left. \left. + \sum_{x=1}^v \sum_{l=1}^{\gamma} \phi_{xl}^d(u, k) \right) \right], \tag{15}
\end{aligned}$$

and so

$$R(k, u) = P(k, u) \frac{1}{\Phi(u, k)}, \tag{16}$$

where  $R(k, u)$  and  $P(k, u)$  are the Laplace transforms of  $R(k, t)$  and  $P(k, t)$ , respectively.

From Eq.(15) we finds

$$\begin{aligned}
uP(k, u) &= R(k, u) - R(k, u) \left[ \sum_{x=1}^v \sum_{i=1}^m \phi_{ix}^r(u, k) \right. \\
&\quad \left. + \sum_{x=1}^v \sum_{l=1}^{\gamma} \phi_{lx}^d(u, k) \right] \tag{17}
\end{aligned}$$

Substituting (14) and (16) into the above equation (17), and taking some algebraic operations, we finally obtain the master equation for the chemical continuous time random walks with anomalous diffusion in Laplace space

$$\begin{aligned}
uP(k, u) - P(k, 0) &= \sum_{i=1}^m \sum_{x=1}^v P(k - s_{ix}, u) \frac{\phi_{ix}^r(u, k - s_{ix})}{\Phi(u, k - s_{ix})} \\
&\quad + \sum_{l=1}^{\gamma} \sum_{x=1}^v \sum_{y \neq x} [P(k - s_{dly}, u) \\
&\quad \cdot \frac{\phi_{lx}^d(u, k - s_{dly})}{\Phi(u, k - s_{dly})} \lambda(y - x)] \\
&\quad - \sum_{x=1}^v \sum_{i=1}^m P(k, u) \frac{\phi_{ix}^r(u, k)}{\Phi(u, k)} \\
&\quad - \sum_{l=1}^{\gamma} \sum_{x=1}^v P(k, u) \frac{\phi_{xl}^d(u, k)}{\Phi(u, k)}. \tag{18}
\end{aligned}$$

Note that the first item on the right side is just the gain flux and the last two items is the loss

flux. Inverting it into time space we obtain the master equation in time space as following

$$\begin{aligned}
\frac{\partial P(k, t)}{\partial t} &= \sum_{i=1}^m \sum_{x=1}^v \int_0^t P(k - s_{ix}, t') \Theta_{ix}^r(t - t', k - s_{ix}) dt' \\
&+ \sum_{l=1}^{\gamma} \sum_{x=1}^v \sum_{y \neq x} \int_0^t [P(k - s_{dly}, t') \\
&\cdot \Theta_{lx}^d(t - t', k - s_{dly}) \lambda(y - x)] dt' \\
&- \sum_{x=1}^v \sum_{i=1}^m \int_0^t P(k, t') \Theta_{ix}^r(t - t', k) dt' \\
&- \sum_{l=1}^{\gamma} \sum_{x=1}^v \int_0^t P(k, t') \Theta_{xl}^d(t - t', k) dt', \tag{19}
\end{aligned}$$

where  $\Theta_{ix}^r(t, k)$  and  $\Theta_{lx}^d(t, k)$  are respectively the inverse Laplace transforms  $u \rightarrow t$  of

$$\Theta_{ix}^r(u, k) = \frac{\phi_{ix}^r(u, k)}{\Phi(u, k)},$$

and

$$\Theta_{lx}^d(u, k) = \frac{\phi_{xl}^d(u, k)}{\Phi(u, k)}.$$

Note that when the diffusion process disappears, the above master equation is just the CME for the original non-markovian reaction system in [17, 16], and  $\Theta_{ix}^r(t, k)$  is just the memory kernel [17]. Note also that in the steady state we get

$$\begin{aligned}
&\sum_{i=1}^m \sum_{x=1}^v \int_0^t P(k - s_{ix}, t') \Theta_{ix}^r(t - t', k - s_{ix}) dt' \\
&+ \sum_{l=1}^{\gamma} \sum_{x=1}^v \sum_{y \neq x} \int_0^t [P(k - s_{dly}, t') \\
&\cdot \Theta_{lx}^d(t - t', k - s_{dly}) \lambda(y - x)] dt' \\
&= \sum_{x=1}^v \sum_{i=1}^m \int_0^t P(k, t') \Theta_{ix}^r(t - t', k) dt' \\
&+ \sum_{l=1}^{\gamma} \sum_{x=1}^v \int_0^t P(k, t') \Theta_{xl}^d(t - t', k) dt'. \tag{20}
\end{aligned}$$

## 4 Mass action law

We will consider the rate law to describe the macroscopic behavior of reaction-diffusion system. We assume that the whole number  $\sum_{x=1}^v \sum_{l=1}^{\gamma} k_{xl}$  of the particles in this system is an invariable constant  $n_0$ . Let  $n_{xl}$  represent the number for the particle  $l$  at position  $x$ , and let  $P_l(c, x, t)$  represent the probability of the concentration  $c = \frac{n_{xl}}{n_0}$  for  $l$  at  $x$  at time  $t$ . Then we find  $P_l(c, x, t) = \sum_{k: k_{xl}=n_{xl}} P(k, t)$  which satisfies  $\int_0^1 P_l(c, x, t) dc = \sum_k P(k, t) = 1$ . Thus,

$$\frac{\partial P_l(c, x, t)}{\partial t} = \sum_{k: k_{xl}=n_{xl}} \frac{\partial P(k, t)}{\partial t}. \tag{21}$$

Combining with Eq.(19), we obtain

$$\begin{aligned}
\frac{\partial P_l(c, x, t)}{\partial t} &= \sum_{i:s_{ixl} \neq 0} \sum_{k:k_{xl}=n_{xl}} \left[ P(k - S_{ix}, t') \right. \\
&\quad \left. \cdot \Theta_{ix}^r(t - t', k - S_{ix}) - P(k, t') \Theta_{ix}^r(t - t', k) \right] \\
&+ \sum_{y \neq x} \sum_{k:k_{xl}=n_{xl}} \left[ P(k - S_{dlyx}, t') \Theta_{lx}^d(t - t', k - S_{dlyx}) \right. \\
&\quad \cdot \lambda(y - x) + P(k - S_{dlyx}, t') \Theta_{ly}^d(t - t', k - S_{dlyx}) \lambda(x - y) \\
&\quad - P(k, t') \Theta_{lx}^d(t - t', k) \lambda(y - x) \\
&\quad \left. - P(k, t') \Theta_{ly}^d(t - t', k) \lambda(x - y) \right]. \tag{22}
\end{aligned}$$

Let  $a_{ixl} = \frac{s_{ixl}}{n_0}$ . Then we get

$$\begin{aligned}
\frac{\partial P_l(c, x, t)}{\partial t} &= \int_0^t \left\{ \sum_{i:a_{ixl} \neq 0} \left[ P_l(c - a_{ixl}, x, t') \right. \right. \\
&\quad \left. \cdot \Theta_{ix}^r(t - t', c - a_{ixl}) - P_l(c, x, t') \Theta_{ix}^r(t - t', c) \right] \\
&\quad + \left[ P_l\left(c + \frac{1}{n_0}, x, t'\right) \Theta_{lx}^d(t - t', c + \frac{1}{n_0}) \right. \\
&\quad + \sum_{y \neq x} P\left(c - \frac{1}{n_0}, x, t'\right) \Theta_{ly}^d(t - t', c - \frac{1}{n_0}) \lambda(x - y) \\
&\quad - P_l(c, x, t') \Theta_{lx}^d(t - t', c) \\
&\quad \left. \left. - \sum_{y \neq x} P_l(c, x, t') \Theta_{ly}^d(t - t', c) \lambda(x - y) \right] \right\} dt'. \tag{23}
\end{aligned}$$

Since  $\langle C_l(x, t) \rangle = \sum_{n_{xl}=0}^{n_0} \frac{n_{xl}}{n_0} P_l(c, x, t)$ , then

$$\begin{aligned}
\left\langle \frac{\partial C_l(x, t)}{\partial t} \right\rangle &= \int_0^1 c \frac{\partial P_l(c, x, t)}{\partial t} dc \\
&= \int_0^1 c dc \int_0^t \left\{ \sum_{i:a_{ixl} \neq 0} \left[ P_l(c - a_{ixl}, x, t') \right. \right. \\
&\quad \left. \cdot \Theta_{ix}^r(t - t', c - a_{ixl}) - P_l(c, x, t') \Theta_{ix}^r(t - t', c) \right] \\
&\quad + \left[ P_l\left(c + \frac{1}{n_0}, x, t'\right) \Theta_{lx}^d(t - t', c + \frac{1}{n_0}) \right. \\
&\quad + \sum_{y \neq x} P\left(c - \frac{1}{n_0}, x, t'\right) \Theta_{ly}^d(t - t', c - \frac{1}{n_0}) \lambda(x - y) \\
&\quad - P_l(c, x, t') \Theta_{lx}^d(t - t', c) \\
&\quad \left. \left. - \sum_{y \neq x} P_l(c, x, t') \Theta_{ly}^d(t - t', c) \lambda(x - y) \right] \right\} dt', \tag{24}
\end{aligned}$$



which can be written as

$$\begin{aligned}
\left\langle \frac{\partial C_l(x, t)}{\partial t} \right\rangle &= \int_0^t dt' \int_0^1 \left\{ \left[ \sum_{i: a_{ixl} \neq 0} (c - a_{ixl} + a_{ixl}) P_l(c - a_{ixl}, x, t') \right. \right. \\
&\quad \cdot \Theta_{ix}^r(t - t', c - a_{ixl}) - c P_l(c, x, t') \Theta_{ix}^r(t - t', c) \left. \right] \\
&\quad + (c + \frac{1}{n_0} - \frac{1}{n_0}) P_l(c + \frac{1}{n_0}, x, t') \Theta_{lx}^d(t - t', c + \frac{1}{n_0}) \\
&\quad - c P_l(c, x, t') \Theta_{lx}^d(t - t', c) \\
&\quad + \sum_{y \neq x} \left[ (c - \frac{1}{n_0} + \frac{1}{n_0}) P_l(c - \frac{1}{n_0}, x, t') \Theta_{ly}^d(t - t', c - \frac{1}{n_0}) \lambda(x - y) \right. \\
&\quad \left. \left. - c P_l(c, x, t') \Theta_{ly}^d(t - t') \lambda(x - y) \right] \right\} dc. \tag{25}
\end{aligned}$$

We simplify the above equation, and get

$$\begin{aligned}
\left\langle \frac{\partial C_l(x, t)}{\partial t} \right\rangle &= \int_0^t \left[ \sum_{i: a_{ixl} \neq 0} a_{ixl} \langle \Theta_{ix}^r(t', t - t') \rangle \right. \\
&\quad \left. - \frac{1}{n_0} \langle \Theta_{lx}^d(t', t - t') \rangle + \sum_{y \neq x} \frac{1}{n_0} \langle \Theta_{ly}^d(t', t - t') \rangle \lambda(x - y) \right] dt'. \tag{26}
\end{aligned}$$

Here,  $\langle \Theta_{ix}^r(t', t - t') \rangle = \int_0^1 \Theta_{ix}^r(t - t', c) P_l(c, x, t') dc$ , and  $\langle \Theta_{lx}^d(t', t - t') \rangle = \int_0^1 \Theta_{lx}^d(t - t', c) P_l(c, x, t') dc$ . This equation is the mass action law for the one-dimensional reaction-diffusion system. If there is no diffusion in the system, the above equation recovers the mass action law in Ref.[16].

## 5 An example: monomolecular $A \leftrightarrow B$ reaction-diffusion system

As an example, we shall consider the monomolecular  $A \leftrightarrow B$  reaction-diffusion system in which the A and B particles can move in one-dimensional finite lattice. There are two reactions which make the change of the concentrations of A and B. The first reaction, we denote by 1, is the reaction from A to B, and the second we denote by 2 is the reaction from B to A. Then the one-dimensional mass action rate (26) becomes

$$\begin{aligned}
\frac{\partial \langle C_A(x, t) \rangle}{\partial t} &= \frac{1}{n_0} \int_0^t [-\langle \Theta_{1x}^r(t', t - t') \rangle \\
&\quad + \langle \Theta_{2x}^r(t', t - t') \rangle - \langle \Theta_{Ax}^d(t', t - t') \rangle \\
&\quad + \sum_{y \neq x} \int_0^t \langle \Theta_{Ay}^d(t', t - t') \rangle \lambda(x - y)] dt', \tag{27}
\end{aligned}$$

and

$$\begin{aligned}
\frac{\partial \langle C_B(x, t) \rangle}{\partial t} &= \frac{1}{n_0} \int_0^t [-\langle \Theta_{2x}^r(t', t - t') \rangle \\
&\quad + \langle \Theta_{1x}^r(t', t - t') \rangle - \langle \Theta_{Bx}^d(t', t - t') \rangle \\
&\quad + \sum_{y \neq x} \langle \Theta_{By}^d(t', t - t') \rangle \lambda(x - y)] dt', \tag{28}
\end{aligned}$$

where  $C_A(x, t) = \frac{n_{xA}}{n_0}$  and  $C_B(x, t) = \frac{n_{xB}}{n_0}$ .

## 5.1 Exponential case

For exponent waiting time, that is,  $p_{1x}(t) = \alpha_{rAx}e^{-\alpha_{rAx}t}$  and  $p_{2x}(t) = \alpha_{rBx}e^{-\alpha_{rBx}t}$ , and  $p_{dAx}(t) = \alpha_{dAx}e^{-\alpha_{dAx}t}$  and  $p_{dBx}(t) = \alpha_{dBx}e^{-\alpha_{dBx}t}$ , then we have

$$\psi_{1x}^r(t, k) = n_{xA}\alpha_{rAx} \exp(-\alpha_{rAx}n_{xA}t) \quad (29)$$

$$\psi_{2x}^r(t, k) = n_{xB}\alpha_{rBx} \exp(-\alpha_{rBx}n_{xB}t) \quad (30)$$

$$\Psi_{1x}^r(t, k) = \exp(-\alpha_{rAx}n_{xA}t) \quad (31)$$

$$\Psi_{2x}^r(t, k) = \exp(-\alpha_{rBx}n_{xB}t) \quad (32)$$

$$\psi_{1x}^d(t, k) = n_{xA}\alpha_{dAx} \exp(-\alpha_{dAx}n_{xA}t) \quad (33)$$

$$\psi_{2x}^d(t, k) = n_{xB}\alpha_{dBx} \exp(-\alpha_{dBx}n_{xB}t) \quad (34)$$

$$\Psi_{1x}^d(t, k) = \exp(-\alpha_{dAx}n_{xA}t) \quad (35)$$

$$\Psi_{2x}^d(t, k) = \exp(-\alpha_{dBx}n_{xB}t) \quad (36)$$

$$\begin{aligned} \phi_{1x}^r(t, k) &= n_{xA}\alpha_{rAx} \exp\left\{-\sum_{x=1,2,\dots,v} [\alpha_{rAx}n_{xA} + \alpha_{rBx}n_{xB} \right. \\ &\quad \left. + \alpha_{dAx}n_{xA} + \alpha_{dBx}n_{xB}]t\right\} \end{aligned} \quad (37)$$

$$\begin{aligned} \phi_{2x}^r(t, k) &= n_{xB}\alpha_{rBx} \exp\left\{-\sum_{x=1,2,\dots,v} [\alpha_{rAx}n_{xA} + \alpha_{rBx}n_{xB} \right. \\ &\quad \left. + \alpha_{dAx}n_{xA} + \alpha_{dBx}n_{xB}]t\right\} \end{aligned} \quad (38)$$

and

$$\begin{aligned} \phi_{1x}^d(t, k) &= n_{xA}\alpha_{dAx} \exp\left\{-\sum_{x=1,2,\dots,v} [\alpha_{rAx}n_{xA} + \alpha_{rBx}n_{xB} \right. \\ &\quad \left. + \alpha_{dAx}n_{xA} + \alpha_{dBx}n_{xB}]t\right\} \end{aligned} \quad (39)$$

$$\begin{aligned} \phi_{2x}^d(t, k) &= n_{xB}\alpha_{dBx} \exp\left\{-\sum_{x=1,2,\dots,v} [\alpha_{rAx}n_{xA} + \alpha_{rBx}n_{xB} \right. \\ &\quad \left. + \alpha_{dAx}n_{xA} + \alpha_{dBx}n_{xB}]t\right\} \end{aligned} \quad (40)$$

$$\begin{aligned} \Phi(t, k) &= \exp\left\{-\sum_{x=1,2,\dots,v} [\alpha_{rAx}n_{xA} + \alpha_{rBx}n_{xB} \right. \\ &\quad \left. + \alpha_{dAx}n_{xA} + \alpha_{dBx}n_{xB}]t\right\}, \end{aligned} \quad (41)$$

and then

$$\Theta_{1x}^r(u, k) = n_{xA}\alpha_{rAx},$$

$$\Theta_{2x}^r(u, k) = n_{xB}\alpha_{rBx},$$

and

$$\Theta_{Ax}^d(u, k) = n_{xA}\alpha_{dAx},$$

$$\Theta_{Bx}^d(u, k) = n_{xB}\alpha_{dBx}.$$

Inverting them to time space yields

$$\Theta_{1x}^r(t, k) = n_{xA}\alpha_{rAx}\delta(t),$$

$$\Theta_{2x}^r(t, k) = n_{xB}\alpha_{rBx}\delta(t),$$

and

$$\Theta_{Ax}^d(t, k) = n_{xA}\alpha_{dAx}\delta(t),$$

$$\Theta_{Bx}^d(t, k) = n_{xB}\alpha_{dBx}\delta(t). \quad (42)$$

Thus, we get

$$\frac{1}{n_0} \int_0^t \langle \Theta_{1x}^r(t', t-t') \rangle dt' = \alpha_{rAx} \langle C_A(x, t) \rangle,$$

$$\frac{1}{n_0} \int_0^t \langle \Theta_{2x}^r(t', t-t') \rangle dt' = \alpha_{rBx} \langle C_B(x, t) \rangle$$

$$\frac{1}{n_0} \int_0^t \langle \Theta_{Ax}^d(t', t-t') \rangle dt' = \alpha_{dAx} \langle C_A(x, t) \rangle,$$

$$\frac{1}{n_0} \int_0^t \langle \Theta_{Ay}^d(t', t-t') \rangle dt' = \alpha_{dAy} \langle C_A(y, t) \rangle,$$

$$\frac{1}{n_0} \int_0^t \langle \Theta_{Bx}^d(t', t-t') \rangle dt' = \alpha_{dBx} \langle C_B(x, t) \rangle,$$

and

$$\frac{1}{n_0} \int_0^t \langle \Theta_{By}^d(t', t-t') \rangle dt' = \alpha_{dBy} \langle C_B(y, t) \rangle.$$

We substitute them into the mass action law (27) and (28), and obtain

$$\begin{aligned} \frac{\partial \langle C_A(x, t) \rangle}{\partial t} &= -\alpha_{rAx} \langle C_A(x, t) \rangle \\ &+ \alpha_{rBx} \langle C_B(x, t) \rangle - \alpha_{dAx} \langle C_A(x, t) \rangle \\ &+ \sum_{y \neq x} \alpha_{dAy} \langle C_A(y, t) \rangle - \lambda(x-y), \end{aligned} \quad (43)$$

and

$$\begin{aligned} \frac{\partial \langle C_B(x, t) \rangle}{\partial t} &= -\alpha_{rBx} \langle C_B(x, t) \rangle \\ &+ \alpha_{rAx} \langle C_A(x, t) \rangle - \alpha_{dBx} \langle C_B(x, t) \rangle \\ &+ \sum_{y \neq x} \alpha_{dBy} \langle C_B(y, t) \rangle - \lambda(x-y), \end{aligned} \quad (44)$$

which are the classical mass action laws for the  $A \leftrightarrow B$  reaction-diffusion system.

## 5.2 Power-law case

In this case we consider the PDFs for waiting times obeying power-law distribution, that is,  $p_{1r}(t) \sim \frac{\tau_0^{\beta_{1r}} \beta_{1r}}{\Gamma(1-\beta_{1r})} \frac{1}{t^{1+\beta_{1r}}}$ ,  $p_{2r}(t) \sim \frac{\tau_0^{\beta_{2r}} \beta_{2r}}{\Gamma(1-\beta_{2r})} \frac{1}{t^{1+\beta_{2r}}}$ ,  $p_{dA}(t) = \frac{\tau_0^{\beta_{dA}} \beta_{dA}}{\Gamma(1-\beta_{dA})} \frac{1}{t^{1+\beta_{dA}}}$  and  $p_{dB}(t) = \frac{\tau_0^{\beta_{dB}} \beta_{dB}}{\Gamma(1-\beta_{dB})} \frac{1}{t^{1+\beta_{dB}}}$  where  $0 < \beta_{1r}, \beta_{2r}, \beta_{dA}, \beta_{dB} < 1$ .

Then we have

$$\phi_{1x}^r(t, k) = n_{xA} \beta_{1r} \frac{\tau_0^H}{F} \frac{1}{t^{1+H}} \quad (45)$$

Taking the Laplace transform  $t \rightarrow u$ , we obtain

$$\phi_{1x}^r(u, k) \sim \frac{n_{xA} \beta_{1r}}{H} \left[ 1 - \frac{\Gamma(1-H)}{F} (\tau_0 u)^H \right]. \quad (46)$$

Analogously, one can get

$$\Phi(u, k) \sim \frac{\Gamma(1-H)}{F} (\tau_0 u)^{H-1}, \quad (47)$$

Therefore,

$$\Theta_{1x}^r(u) \sim n_{xA} \beta_{1r} M u^{1-H}. \quad (48)$$

Taking the Laplace transform of  $\frac{1}{n_0} \int_0^t \langle \Theta_{1x}^r(t', t-t') \rangle dt'$ , we find

$$\begin{aligned} \mathbf{L} \left\{ \frac{1}{n_0} \int_0^t \langle \Theta_{1x}^r(t', t-t') \rangle dt' \right\} &= \frac{1}{n_0} \int_0^1 p(c, x, u) \Theta_{1x}^r(u, c) dc \\ &= \int_0^1 p(c, x, u) \frac{n_{xA}}{n_0} M u^{1-H} \beta_{1r} dc \\ &= \beta_{1r} M u^{1-H} \langle C_A(x, u) \rangle, \end{aligned} \quad (49)$$

whose inverse Laplace transform is

$$\frac{1}{n_0} \int_0^t \langle \Theta_{1x}^r(t', t-t') \rangle dt' = \beta_{1r} M \mathcal{D}_t^{1-H} \langle C_A(x, t) \rangle, \quad (50)$$

$$\frac{1}{n_0} \int_0^t \langle \Theta_{2x}^r(t', t-t') \rangle dt' = \beta_{2r} M \mathcal{D}_t^{1-H} \langle C_B(x, t) \rangle, \quad (51)$$

$$\frac{1}{n_0} \int_0^t \langle \Theta_{Ax}^d(t', t-t') \rangle dt' = \beta_{dA} M \mathcal{D}_t^{1-H} \langle C_A(x, t) \rangle, \quad (52)$$

$$\frac{1}{n_0} \int_0^t \langle \Theta_{Bx}^r(t', t-t') \rangle dt' = \beta_{dB} M \mathcal{D}_t^{1-H} \langle C_B(x, t) \rangle, \quad (53)$$

where  $M = \frac{1}{H} \frac{1}{\tau_0^H} \frac{F}{\Gamma(1-H)}$ , with  $F = \Gamma^{n_A}(1-\beta_{1r}) \Gamma^{n_B}(1-\beta_{2r}) \Gamma^{n_A}(1-\beta_{dA}) \Gamma^{n_B}(1-\beta_{dB})$ , and  $H = \beta_{1r} n_A + \beta_{2r} n_B + \beta_{dA} n_A + \beta_{dB} n_B$ ,  $0 < H < 1$ . We substitute them into the mass action laws (27) and (28), and obtain

$$\begin{aligned} \frac{\partial \langle C_A(x, t) \rangle}{\partial t} &= -\beta_{1r} M \mathcal{D}_t^{1-H} \langle C_A(x, t) \rangle \\ &\quad + \beta_{2r} M \mathcal{D}_t^{1-H} \langle C_B(x, t) \rangle - \beta_{dA} M \mathcal{D}_t^{1-H} \langle C_A(x, t) \rangle \\ &\quad + \sum_{y \neq x} \beta_{dA} M \mathcal{D}_t^{1-H} \langle C_A(y, t) \rangle \lambda(x-y), \end{aligned} \quad (54)$$

and

$$\begin{aligned}
\frac{\partial \langle C_B(x, t) \rangle}{\partial t} &= -\beta_{2r} M \mathcal{D}_t^{1-H} \langle C_B(x, t) \rangle \\
&+ \beta_{1r} M \mathcal{D}_t^{1-H} \langle C_A(x, t) \rangle - \beta_{dB} M \mathcal{D}_t^{1-H} \langle C_B(x, t) \rangle \\
&+ \sum_{y \neq x} \beta_{dB} M \mathcal{D}_t^{1-H} \langle C_B(y, t) \rangle \lambda(x - y).
\end{aligned} \tag{55}$$

Here,  $\mathcal{D}_t^{1-H}$  is the Riemann-Liouville fractional differential operator. This means that the concentration depends on the history evolution for A and B particles.

## 6 generalized Gillespie algorithm

Based on the chemical continuous time random walks under anomalous diffusion model, we can get the generalized Gillespie algorithm as follows.

(i) In the initial state  $n_0$  reactants are randomly placed at  $v$  places by uniform distribution.

(ii) The random waiting time  $\tau_{ix}$  as the internal clock to react for each reaction  $i$  at  $x$  is chosen from a series of values distributed according to  $p_{ix}(t)$  and the random waiting time  $\tau_{dlx}$  as the internal clock to make next jump for each reactant  $l$  at  $x$  is chosen from a series of values distributed according to  $p_{dlx}(t)$  [23-25]. If  $p_{ix}(t)$  is exponential distribution  $p_{ix}(t) = \alpha_{ix} e^{-\alpha_{ix} t}$ , then we get  $\tau_{ix} = -\frac{\ln u}{\alpha_{ix}}$ , where  $u$  is a random variate drawn from the uniform density on the interval  $[0, 1]$ . If the  $p_{ix}(t)$  is power-law distribution  $p_{ix}(t) = \alpha_{ix} \tau_0^{\alpha_{ix}} t^{-(1+\alpha_{ix})}$ , then we get  $\tau_{ix} = \tau_0 u^{-\frac{1}{\alpha_{ix}}}$ . The generation of  $\tau_{dlx}$  is as same as that of  $\tau_{ix}$ .

(iii) Find the minimum clock time  $\min\{\tau_{ix}, \tau_{dlx}\}$  ( $i = 1, 2, \dots, m; l = 1, 2, \dots, \gamma; x = 1, 2, \dots, v$ ). If the minimum waiting time is for one reaction  $i$  at  $x$  then we change the system according to this reaction [15], or the minimum waiting time is for one diffusion process for a reactant  $l$  at  $x$ , then we let  $l$  diffuse from  $x$  to one place  $y$  with the probability  $\lambda(y - x)$ , after which the internal clocks for reactions and diffusion are all reset by the distributions and repeat the procedure.

Note that this algorithm is different from the non-Markovian Gillespie algorithm for reactive system where the reaction  $i$  is chosen by the instantaneous probability after the time  $\Delta t$  for the next event drawn in [15].

## 7 Comparison with the mass action law derived from continuous time random walk (CTRW) models.

In this section we shall compare our mass action law with that derived from continuous time random walk (CTRW) for the state A to B and backward under anomalous transport by Campos e.t. in [26]. The corresponding master equations in [26] is as following

$$\begin{aligned}
\frac{dP_1(x, t)}{dt} &= \int_{-\infty}^{+\infty} \int_0^t P_1(x', t') \Theta_{1j}(t - t') \lambda_1(x - x') dt' dx' \\
&+ \int_0^t P_2(x, t') \Theta_{2r}(t - t') dt' - \int_0^t P_1(x, t') \Theta_{1j}(t - t') dt' \\
&- \int_0^t P_1(x, t') \Theta_{1r}(t - t') dt',
\end{aligned} \tag{56}$$

$$\begin{aligned}
\frac{dP_2(x, t)}{dt} &= \int_{-\infty}^{+\infty} \int_0^t P_2(x', t') \Theta_{2j}(t - t') \lambda_2(x - x') dt' dx' \\
&+ \int_0^t P_1(x, t') \Theta_{1r}(t - t') dt' - \int_0^t P_2(x, t') \Theta_{2j}(t - t') dt' \\
&- \int_0^t P_2(x, t') \Theta_{2r}(t - t') dt'.
\end{aligned} \tag{57}$$

Here,  $P_{1,2}(x, t) = \frac{N_{1,2}(x, t)}{N}$  respectively denote the PDFs for a particle being in state  $A$  and  $B$  at point  $x$  at time  $t$  where  $N_{1,2}(x, t)$  are the the densities of  $A$  and  $B$ , and  $N$  is the sum of  $N_1(x, t)$  and  $N_2(x, t)$  over the whole space. One can see that  $\int_{-\infty}^{+\infty} (P_1(x, t) + P_2(x, t)) dx = 1$ .  $\lambda_{1,2}(\Delta x)$  denote the jump length PDFs for  $A$  particle and  $B$  particle, respectively.  $\psi_{1j,2j}(t)$  denote the PDFs of random diffusion waiting times,  $\psi_{1r,2r}(t)$  denote the PDFs of random reaction waiting time of  $A \rightarrow B$  and  $B \rightarrow A$ ,  $\Psi_{1j,2j}(t) = 1 - \int_0^t \psi_{1j}(\tau) d\tau$  and  $\Psi_{1r,2r}(t) = 1 - \int_0^t \psi_{2j}(\tau) d\tau$  are the corresponding survival probabilities for  $\psi_{1j,2j}(t)$  and  $\psi_{1j,2j}(t)$ . Finally,  $\Theta_{1j}(u) = \frac{L(\Psi_{1r}(t)\psi_{1j}(t))}{L(\Psi_{1r}(t)\Psi_{1j}(t))}$  and  $\Theta_{1r}(u) = \frac{L(\Psi_{1j}(t)\psi_{1r}(t))}{L(\Psi_{1j}(t)\Psi_{1r}(t))}$  are the kernels.

When the random jump length is Gaussian satisfying

$$\lambda(k) \sim 1 - \frac{\sigma^2 k^2}{2} \tag{58}$$

and random waiting times for reaction and diffusion are both power-law distributed satisfying

$$\psi_{1r,2r}(t) = \beta_{1r,2r} \tau^{\beta_{1r,2r}} \frac{1}{t^{1+\beta_{1r,2r}}} \tag{59}$$

and

$$\psi_{1j,2j}(t) = \beta_{1j,2j} \tau^{\beta_{1j,2j}} \frac{1}{t^{1+\beta_{1j,2j}}}, \tag{60}$$

where  $\sigma^2$  is the jump length variance, one gets  $\Theta_{hj}(u) \sim \frac{\beta_{hj}}{\beta_{hr} + \beta_{hj}} \frac{u^{1-\beta_{hr}-\beta_{hj}}}{\tau^{\beta_{hr} + \beta_{hj}} \Gamma(1-\beta_{hr}-\beta_{hj})}$  and  $\Theta_{hr}(u) \sim \frac{\beta_{hr}}{\beta_{hr} + \beta_{hj}} \frac{u^{1-\beta_{hj}-\beta_{hr}}}{\tau^{\beta_{hj} + \beta_{hr}} \Gamma(1-\beta_{hj}-\beta_{hr})}$  for  $0 < \beta_{1r} + \beta_{1j}, \beta_{2r} + \beta_{2j} < 1$ .

Taking the Fourier-Laplace transform of the generalized master equations (56) and (57) yields

$$\begin{aligned}
uP_1(k, u) - P_1(k, 0) &= P_1(k, u) \Theta_{1j}(u) \left(-\frac{\sigma^2 k^2}{2}\right) \\
&+ P_2(k, u) \Theta_{2r}(u) - P_1(k, u) \Theta_{1r}(u),
\end{aligned} \tag{61}$$

and

$$\begin{aligned}
uP_2(k, u) - P_2(k, 0) &= P_2(k, u) \Theta_{2j}(u) \left(-\frac{\sigma^2 k^2}{2}\right) \\
&+ P_1(k, u) \Theta_{1r}(u) - P_2(k, u) \Theta_{2r}(u).
\end{aligned} \tag{62}$$

By substituting  $\Theta_{hj}(u)$  and  $\Theta_{hr}(u)$  into (61) and (62) and taking the inverse Fourier-Laplace transform, we finally obtain the generalized fractional reaction-diffusion equations

$$\begin{aligned}
\frac{dP_1(x, t)}{dt} &= C_{11} \frac{\partial^2 D_t^{1-\beta_{1r}-\beta_{1j}} P_1(x, t)}{\partial x^2} \\
&+ C_2 D_t^{1-\beta_{2j}-\beta_{2r}} P_2(x, t) \\
&- C_3 D_t^{1-\beta_{1j}-\beta_{1r}} P_1(x, t),
\end{aligned} \tag{63}$$

$$\begin{aligned} \frac{dP_2(x, t)}{dt} = & C_{21} \frac{\partial^2 D_t^{1-\beta_{2r}-\beta_{2j}} P_2(x, t)}{\partial x^2} \\ & - C_2 D_t^{1-\beta_{2j}-\beta_{2r}} P_2(x, t) \\ & + C_3 D_t^{1-\beta_{1j}-\beta_{1r}} P_1(x, t), \end{aligned} \quad (64)$$

with

$$\begin{aligned} C_{11} &= \frac{\sigma^2}{2} \frac{\beta_{1j}}{\beta_{1r} + \beta_{1j}} \frac{1}{\tau^{\beta_{1r}+\beta_{1j}} \Gamma(1 - \beta_{1r} - \beta_{1j})}, \\ C_{21} &= \frac{\sigma^2}{2} \frac{\beta_{2j}}{\beta_{2r} + \beta_{2j}} \frac{1}{\tau^{\beta_{2r}+\beta_{2j}} \Gamma(1 - \beta_{2r} - \beta_{2j})}, \\ C_2 &= \frac{\beta_{2r}}{\beta_{2r} + \beta_{2j}} \frac{1}{\tau^{\beta_{2r}+\beta_{2j}} \Gamma(1 - \beta_{2r} - \beta_{2j})}, \\ C_3 &= \frac{\beta_{1r}}{\beta_{1r} + \beta_{1j}} \frac{1}{\tau^{\beta_{1r}+\beta_{1j}} \Gamma(1 - \beta_{1r} - \beta_{1j})}. \end{aligned}$$

When we assume that there is only one particle in the reaction-diffusion system, the jump length is Gaussian whose PDF obeys (58) and the PDFs of reaction and diffusion waiting times is space-independent satisfying (59) and (60), we find that our mass action law (54) and (55) can recover (63) and (64).

## 8 Comparison with the chemical continuous time random walk (CTRW) models.

In Ref.[16] it defines an inhomogeneous continuous time random walk in particle number space, from which a generalized chemical master equation is derived as following

$$\begin{aligned} \frac{\partial P(n, t)}{\partial t} &= \sum_i \int_0^t dt' (\prod_j E_j^{-s_{ij}} - 1) \\ &\times P(n, t') M_i(t - t', n), \end{aligned} \quad (65)$$

where the step operator  $E_j^z$  acts on a function  $f(n)$  by incrementing the particle number  $n_j$  of species  $S_j$ .  $M_i$  is the memory function. This equation is the special case of the mass action law (26) without diffusion in the system.

Based on the derived master equation (65), the authors determine the modified chemical rate laws

$$\frac{\partial \langle C \rangle}{\partial t} = \sum_i s_i \int_0^t dt' M_i^C(t - t', C), \quad (66)$$

where  $M_i^C(t-t', C) = \frac{M_i^C(t-t', n_0 C)}{n_0}$ . Note that when the diffusion is ignored, namely,  $\lambda(y-x) = 0$  in our rate laws (26) we can recover (66).

Furthermore, a generalization of the Gillespie algorithm is proposed for the reaction system in [16]. Comparing with their algorithm, we find that their algorithm is specially for exponential reactive waiting times, while our algorithm is for arbitrary distributed and more general. Beside, we also take account of the diffusion for generalization while it is not considered in [16]. But they further consider the global delay that is added to inter-reaction waiting time.

## 9 Conclusion

To sum up, in order to describe the coupled non-Poissonian reaction and anomalous diffusion process in heterogeneous system, where the reactive and diffusive waiting times are both arbitrary distributions, we develop a chemical continuous time random walk under anomalous diffusion model. We firstly use the renewal process to describe the numbers of reaction and diffusion renewals, and also define the distribution of the minimum renewal waiting times. Based on the renewal process we obtain the generalized chemical diffusion master equation (19) for the probability of the numbers of reactants, and then derive the corresponding mass action law (26) for the average concentration of the reactants using the CDME. Moreover, we generalize the Gillespie algorithm to fit for the non-Markovian reaction-diffusion system. In the end, we take the  $A \leftrightarrow B$  reaction system in which the A and B particles for exponential and power-law waiting times respectively. We recover the mass law rate in the exponential waiting time case, and show the strong fractional memory effect of the concentration of the reactants on the history of the concentration in the power-law case.

## 10 Acknowledgements

This work was supported by the Natural Science Foundation of Sichuan Province (Grant No. 2022NSFSC1752).

## 11 Conflict of interest statement

The authors declare that there are no conflict of interests, we do not have any possible conflicts of interest.

## References

- [1] C. Tsallis and D. J. Bukman, *Phys. Rev. E* **54** R2197 (1996)
- [2] R. Metzler, J. Klafter, *Phys. Rep.* **339** 1 (2000)
- [3] I. M. Sokolov, M. G. W. Schmidt, F. Sagués, *Phys. Rev. E*, **73** 031102 (2006)
- [4] T. A. M. Langlands, B. I. Henry, S. L. Wearne, *Phys. Rev. E* **77** 021111 (2008)
- [5] M. G. W. Schmidt, F. Sagués, and I. M. Sokolov, *J. Phys. Condens. Matter* **19** 065118 (2007)
- [6] E. Abad, S. B. Yuste, K. Lindenberg, *Phys. Rev. E* **81** 031115 (2010)
- [7] S. B. Yuste, E. Abad, and K. Lindenberg, *J. Stat. Mech. -Theory and Exp.* **11** 014 (2014)
- [8] B. I. Henry, T. A. M. Langlands, S. L. Wearne, *Phys. Rev. E* **74** 031116 (2006)
- [9] S. Fedotov, *Phys. Rev. E* **81** 011117 (2010)
- [10] C. N. Angstmann, I. C. Donnelly, B. I. Henry, *Math. Model. Nat. Phenom.* **8** 17 (2013)
- [11] D. T. Gillespie *J. Phys. Chem.* **81**, 25 (1977)



- [12] M. J. del Razo, D. Frömberg, A. V. Straube, C. Schütte, F. Höfling, s. Winkelmann, *Lett. Math. Phys.* **112**, P1-59 (2022)
- [13] M. J. del Razo, S. Winkelmann, R. Klein, F. Höfling, *J. Math. Phys.* **64** , 013304 (2023)
- [14] H. Ge, H. Q., *Applications of mathematical dynamics models in biophysics and biochemistry*, Peking University Process (2017)
- [15] N. Masuda, L. E. C. Rocha, *SIAM Rev.* **60** , 95 (2018)
- [16] T. Aquino, M. Dentz, *Phys. Rev. Lett.* **119**, 230601 (2017)
- [17] J. J. Zhang, T. S. Zhou, *Proc. Natl. Acad. Sci. USA* **116**, 23542 (2019)
- [18] L. Y. Chen, Y. Wan, J. R. Liu, H. H. Wang *Biosystems* **214**, 104648 (2022)
- [19] M. L. Chen, S. H. Luo, M. F. Gao, C. J. Guo, T. S. Zhou, J. J. Zhang, *Phys. Rev. E* **105**, 014405 (2022)
- [20] H. W. Yin, S. Q. Liu, X. Q. Wen, *Phys. Rev. E* **103**, 022409 (2021)
- [21] D. R. Cox, *Renewal Theory*, Methuen Co. Ltd, (1962)
- [22] W. Feller, *An Introduction to Probability Theory and Its Applications*, John Wiley Sons, (1971)
- [23] S. J. Ni, and W. G. Weng, *Phys. Rev. E* **79** , 016111 (2009)
- [24] H. H. Schmidt-Martens, D. Froemberg, I. M. Sokolov, and F. Sagués, *Phys. Rev. E* **79** , 041135 (2009)
- [25] H. Zhang, G. H. Li, *Phys. Rev. E* **102** , 012315 (2020)
- [26] D. Campos, S. Fedotov, and V. Méndez. *Phys. Rev. E* **77** , 061130 (2008)