

A New Diophantine Approximation Inequality on Surfaces and Its Applications

KEPING HUANG, AARON LEVIN, AND ZHENG XIAO

Abstract

We prove a Diophantine approximation inequality for closed subschemes on surfaces which can be viewed as a joint generalization of recent inequalities of Ru-Vojta and Heier-Levin in this context. As applications, we study various Diophantine problems on affine surfaces given as the complement of three numerically parallel ample projective curves: inequalities involving greatest common divisors, degeneracy of integral points, and related Diophantine equations including families of S -unit equations. We state analogous results in the complex analytic setting, where our main result is an inequality of Second Main Theorem type for surfaces, with applications to the study and value distribution theory of holomorphic curves in surfaces.

1 Introduction

Schmidt's Subspace Theorem occupies a central place in the theory of Diophantine approximation, and it is known to have deep and far-reaching applications. Stated in the language of heights (and with improvements due to Schlickewei [Sch77] allowing for arbitrary finite sets of places), the Subspace Theorem may be stated as follows:

Theorem 1.1 (Schmidt Subspace Theorem). *Let S be a finite set of places of a number field K . For each $v \in S$, let $H_{0,v}, \dots, H_{n,v} \subset \mathbb{P}^n$ be hyperplanes over K in general position. Let $\varepsilon > 0$. Then there exists a finite union of hyperplanes $Z \subset \mathbb{P}^n$ such that for all points $P \in \mathbb{P}^n(K) \setminus Z$,*

$$\sum_{v \in S} \sum_{i=0}^n \lambda_{H_{i,v},v}(P) < (n+1+\varepsilon)h(P).$$

Here, $\lambda_{H_{i,v},v}$ is a local height function (also known as a local Weil function) associated to the hyperplane $H_{i,v}$ and place v in S , and h is the standard (logarithmic) height on \mathbb{P}^n .

The Subspace Theorem has been generalized to the setting of hypersurfaces in projective space by Corvaja and Zannier [CZ04a], and more generally, by Corvaja and Zannier [CZ04a] (for complete intersections) and Evertse and Ferretti [EF08] (for arbitrary projective varieties) to divisors which possess a common linearly equivalent

multiple. Building on work of Autissier [Aut11], Ru and Vojta [RV20] proved a general version of the Subspace Theorem in terms of beta constants (see Definition 3.1). Their inequality was subsequently extended to the context of closed subschemes by Ru and Wang [RW22] and by Vojta [Voj23] (under slightly different intersection conditions). We state the following general form of the inequality, due to Vojta [Voj23] (in fact, Vojta proves a stronger version of Theorem 1.2 with the condition “intersect properly” (Definition 2.10) replaced by “weakly intersect properly” [Voj23, Definition 4.1(c)]).

Theorem 1.2 (Ru-Vojta [RV20], Ru-Wang [RW22], Vojta [Voj23]). *Let X be a projective variety of dimension n defined over a number field K . Let S be a finite set of places of K . For each $v \in S$, let $Y_{0,v}, \dots, Y_{n,v}$ be closed subschemes of X , defined over K , that intersect properly. Let A be a big divisor on X , and let $\varepsilon > 0$. Then there exists a proper Zariski-closed subset $Z \subset X$ such that for all points $P \in X(K) \setminus Z$,*

$$\sum_{v \in S} \sum_{i=0}^n \beta(A, Y_{i,v}) \lambda_{Y_{i,v},v}(P) < (1 + \varepsilon) h_A(P).$$

Heier and the second author [HL21] proved an inequality using Seshadri constants (Definition 3.4) in place of beta constants, and with the proper intersection condition of Theorem 1.2 replaced by a flexible notion of general position for closed subschemes (Definition 2.12).

Theorem 1.3 (Heier-Levin [HL21]). *Let X be a projective variety of dimension n defined over a number field K . Let S be a finite set of places of K . For each $v \in S$, let $Y_{0,v}, \dots, Y_{n,v}$ be closed subschemes of X , defined over K , and in general position. Let A be an ample Cartier divisor on X , and $\varepsilon > 0$. Then there exists a proper Zariski-closed subset $Z \subset X$ such that for all points $P \in X(K) \setminus Z$,*

$$\sum_{v \in S} \sum_{i=0}^n \epsilon(A, Y_{i,v}) \lambda_{Y_{i,v},v}(P) < (n + 1 + \varepsilon) h_A(P).$$

In both Theorem 1.2 and Theorem 1.3, one may also apply the results with fewer than $n + 1$ chosen closed subschemes at v (e.g., by appropriately arbitrarily choosing the remaining closed subschemes and using positivity of the associated local heights outside a closed subset); we will use this fact without further remark.

Before stating our main result, we give some further remarks comparing Theorem 1.2 and Theorem 1.3. For simplicity and to avoid technical issues, we assume now for the discussion that X is nonsingular of dimension n and we fix an ample divisor A on X . If the closed subschemes $Y_{i,v} = D_{i,v}$ are all ample effective divisors, then by our nonsingularity assumption, $D_{0,v}, \dots, D_{n,v}$ intersect properly if and only if they are in general position (Remark 2.13). Furthermore, we note that if $D_{i,v} \sim d_{i,v}A$, then it follows easily from the asymptotic Riemann-Roch formula and the definitions that

$$\beta(A, D_{i,v}) = \frac{1}{(n + 1)d_{i,v}},$$

$$\epsilon(A, D_{i,v}) = \frac{1}{d_{i,v}}.$$

In this case, Theorem 1.2 and Theorem 1.3 coincide and yield the inequality (outside a Zariski-closed subset)

$$\sum_{v \in S} \sum_{i=0}^n \frac{\lambda_{D_{i,v},v}(P)}{d_{i,v}} < (n+1+\varepsilon)h_A(P),$$

which is precisely the inequality of the aforementioned theorem of Evertse and Ferretti.

Another interesting example comes from considering a single closed subscheme Y of X , which we assume additionally to be a local complete intersection (to satisfy the proper intersection hypothesis of Theorem 1.2). In this case, if $\text{codim } Y = r$, from [HL21] we have the inequality

$$\beta(A, Y) \geq \frac{r}{n+1} \epsilon(A, Y),$$

and so Theorem 1.2 yields (choosing $Y_{0,v} = Y$ for all v)

$$\sum_{v \in S} \epsilon(A, Y) \lambda_{Y,v}(P) < \left(\frac{n+1}{r} + \varepsilon \right) h_A(P). \quad (1.1)$$

This is a generalization of an inequality of McKinnon and Roth [MR15] (who proved the case when Y is a point). On the other hand, in Theorem 1.3 one may take $Y_{0,v} = \dots = Y_{r-1,v} = Y$ for all v (i.e., repeat Y as the choice of closed subscheme r times), as is allowed by the used notion of general position (Definition 2.12), and one again obtains the McKinnon-Roth type inequality (1.1). In short, there seems to be a tradeoff between the flexible notion of general position in Theorem 1.3 and the “better” beta constant of Theorem 1.2 (as compared to the Seshadri constant).

The goal of our main result is to provide, in the case of surfaces, an inequality which combines the separate advantages of Theorem 1.2 and Theorem 1.3; we prove an inequality in terms of beta constants of closed subschemes, under hypotheses which permit the use of nested closed subschemes (as allowed in Theorem 1.3, but not in Theorem 1.2).

Theorem 1.4 (Main Theorem). *Let X be a projective surface defined over a number field K . Let S be a finite set of places of K . For each $v \in S$, let $D_v \supset Y_v$ be a regular chain of nonempty closed subschemes of X (see Definition 2.14). Let A be a big Cartier divisor on X , and let $\varepsilon > 0$. Then there exists a proper Zariski-closed subset $Z \subset X$ such that for all $P \in X(K) \setminus Z$, we have*

$$\sum_{v \in S} \left(\beta(A, D_v) \lambda_{D_{v,v}}(P) + (\beta(A, Y_v) - \beta(A, D_v)) \lambda_{Y_{v,v}}(P) \right) < (1+\varepsilon)h_A(P). \quad (1.2)$$

In Section 4, we show that our main result implies the Ru-Vojta inequality for surfaces, and Theorem 1.3 for surfaces under regularity assumptions. Thus, we may view Theorem 1.4 as a kind of joint generalization, for surfaces, of the inequalities of Ru-Vojta and of Heier and the second author.

We remark that the regular chain assumption is quite natural from the point of view of classical Diophantine approximation, where one is interested in sums of local heights associated to divisors. Indeed, if D_1, \dots, D_q are effective divisors in general position on a nonsingular projective surface X , all defined over a number field K , S is a set of places of K , and we fix a point $P \in X(K)$, then from general position a point can be v -adically close to at most two divisors D_i . Thus,

$$\sum_{v \in S} \sum_{i=1}^q \lambda_{D_i, v}(P) = \sum_{v \in S} (\lambda_{D_{i_v}, v}(P) + \lambda_{D_{j_v}, v}(P)) + O(1),$$

for some choice of $i_v, j_v \in \{1, \dots, q\}$ depending on P and v (but with the constant in the $O(1)$ independent of P). Now assuming $\lambda_{D_{i_v}, v}(P) \geq \lambda_{D_{j_v}, v}(P)$ (as we may), from a basic property of heights associated to closed subschemes (see Subsection 2.1) we have (up to $O(1)$)

$$\lambda_{D_{i_v} \cap D_{j_v}, v}(P) = \min\{\lambda_{D_{i_v}, v}(P), \lambda_{D_{j_v}, v}(P)\} = \lambda_{D_{j_v}, v}(P).$$

Therefore, we may write

$$\sum_{v \in S} \sum_{i=1}^q \lambda_{D_i, v}(P) = \sum_{v \in S} (\lambda_{D_{i_v}, v}(P) + \lambda_{D_{i_v} \cap D_{j_v}, v}(P)) + O(1), \quad (1.3)$$

where now $D_{i_v} \supset D_{i_v} \cap D_{j_v}$ is a regular chain of closed subschemes of X for all $v \in S$. Although the inequality (1.3) is essentially elementary (relying at its base on Hilbert's Nullstellensatz), we will find the underlying argument useful in a number of proofs and applications.

Among the potential applications of our main inequality (1.2), in Section 5 we study inequalities related to greatest common divisors, the Diophantine equation $f(a^m, y) = b^n$, integral points on certain affine surfaces, and solutions to families of unit equations. For instance, we prove the following result for integral points on certain complements of three curves in \mathbb{P}^2 :

Theorem 1.5. *Let D_1, D_2, D_3 be distinct irreducible projective curves in \mathbb{P}^2 , defined over a number field K , of degrees d_1, d_2, d_3 , respectively, such that*

$$D_1 \cap D_2 \cap D_3 \neq \emptyset,$$

and for every point $Q \in (D_1 \cap D_2 \cap D_3)(\overline{K})$ and $1 \leq i < j \leq 3$,

$$(D_i \cdot D_j)_Q < \frac{4}{9} d_i d_j, \quad (1.4)$$

where $(D_i \cdot D_j)_Q$ denotes the local intersection multiplicity of D_i and D_j at Q . Let S be a finite set of places of K containing all the archimedean places. Then there exists a proper Zariski-closed subset $Z \subset \mathbb{P}^2$ such that for any set $R \subset X(K)$ of $(D_1 + D_2 + D_3, S)$ -integral points, the set $R \setminus Z$ is finite.

When $d_1 = d_2 = d_3 > 1$ and $D_1 \cap D_2 \cap D_3$ consists of a single point, at which the divisors intersect transversally, the result follows from work of Corvaja and Zannier [CZ06] (see Theorem 5.8). The factor $\frac{4}{9}$ in the intersection condition (1.4) of Theorem 1.5 cannot be replaced by anything larger than $\frac{1}{2}$ (Example 5.11), and in fact a more general version of the theorem stated in terms of beta constants (Theorem 5.9) can be shown to be sharp (Example 5.12). We note that in general it seems to be a difficult problem to prove the degeneracy of integral points on the complement of three curves in \mathbb{P}^2 . For instance, if $D = D_1 + D_2 + D_3$ is a normal crossings divisor of degree $\deg D \geq 4$ with D_1, D_2, D_3 geometrically integral curves in \mathbb{P}^2 (all defined over some number field K), then a suitable version of Vojta's conjecture predicts the same conclusion as in Theorem 1.5, but this does not appear to be known (for all K and S) for even a single such divisor D .

For a further application, by applying a general form of the above theorem (Theorem 5.9) to certain surfaces, we study families of unit equations of the form:

$$f_1(t)u + f_2(t)v = f_3(t), \quad t \in \mathcal{O}_{K,S}, u, v \in \mathcal{O}_{K,S}^*$$

where f_1, f_2, f_3 are polynomials in $K[t]$, and $\mathcal{O}_{K,S}^*$ is the group of S -units of a ring of S -integers $\mathcal{O}_{K,S}$ of a number field K . This equation was treated in the case $\deg f_1 = \deg f_2 = \deg f_3$ by Corvaja and Zannier [CZ06, CZ10], and in the case $\deg f_1 + \deg f_2 = \deg f_3$ by the second author [Lev06]. We prove a result whenever the degrees of the polynomials are roughly within a factor of 2 of each other.

Theorem 1.6. *Let $f_1, f_2, f_3 \in K[t]$ be nonconstant polynomials without a common zero of degrees d_1, d_2, d_3 , respectively, and suppose that*

$$\max_i (d_i + 1) < \frac{9}{5} \min_i d_i.$$

Then the set of solutions $(t, u, v) \in \mathbb{A}^3(K)$ of the equation

$$f_1(t)u + f_2(t)v = f_3(t), \quad t \in \mathcal{O}_{K,S}, u, v \in \mathcal{O}_{K,S}^*$$

is contained in a finite number of rational curves in \mathbb{A}^3 .

Excluding the linear case (proved in [CZ06]), this recovers Corvaja-Zannier's result [CZ10] (i.e., the case $d_1 = d_2 = d_3$).

Finally, we mention that via Vojta's dictionary [Voj87, Ch. 3] between Diophantine approximation and Nevanlinna theory, by substituting Vojta's version [Voj97] of Cartan's Second Main Theorem in place of Schmidt's Subspace Theorem, one can prove a result analogous to Theorem 1.4, giving the following inequality in the vein of the Second Main Theorem:

Theorem 1.7. *Let X be a complex projective surface. Let q be a positive integer and let $D_i \supset Y_i$ be a regular chain of nonempty closed subschemes of X for $i = 1, \dots, q$. Let A be a big Cartier divisor on X , and let $\varepsilon > 0$. Then there exists a proper Zariski-closed subset $Z \subset X$ such that for all holomorphic maps $f : \mathbb{C} \rightarrow X$ whose image is not contained in Z , the inequality*

$$\int_0^{2\pi} \max_i \left(\beta(A, D_i) \lambda_{D_i}(f(re^{i\theta})) + (\beta(A, Y_i) - \beta(A, D_i)) \lambda_{Y_i}(f(re^{i\theta})) \right) \frac{d\theta}{2\pi}$$

$$< (1 + \varepsilon)T_{f,A}(r).$$

holds for all $r \in (0, \infty)$ outside of a set of finite Lebesgue measure.

One may similarly apply Theorem 1.7 to obtain results on holomorphic curves. For instance, using Vojta’s dictionary, one may use Theorem 1.7 to prove the following analogue of Theorem 1.5 for holomorphic curves:

Theorem 1.8. *Let D_1, D_2, D_3 be distinct irreducible complex projective curves in \mathbb{P}^2 , of degrees d_1, d_2, d_3 , respectively, such that*

$$D_1 \cap D_2 \cap D_3 \neq \emptyset,$$

and for every point $Q \in (D_1 \cap D_2 \cap D_3)(\mathbb{C})$ and $1 \leq i < j \leq 3$,

$$(D_i \cdot D_j)_Q < \frac{4}{9}d_i d_j,$$

where $(D_i \cdot D_j)_Q$ denotes the local intersection multiplicity of D_i and D_j at Q . Then there exists a proper Zariski-closed subset $Z \subset \mathbb{P}^2$ such that every nonconstant holomorphic map $f : \mathbb{C} \rightarrow X \setminus (D_1 \cup D_2 \cup D_3)$ has image contained in Z .

As the proofs of these analogous results in the complex setting are similar to the proofs of their arithmetic counterparts (after making the appropriate “translations”), we omit the details.

The organization of this paper is as follows. Section 2 gives relevant background material, including a summary of Silverman’s theory of heights associated to closed subschemes, and some needed results in commutative algebra, algebraic geometry, and Diophantine approximation. In Section 3 we give some basic inequalities involving beta constants, and then in Section 4 we prove our main theorem and show how it implies, for surfaces, versions of the inequality of Ru-Vojta and the inequality of Heier and the second author. Finally, in Section 5 we give some illustrative Diophantine applications of our results.

2 Background

2.1 Heights Associated to Closed Subschemes

In [Sil87], Silverman generalized the Weil height machine for Cartier divisors to height functions on projective varieties with respect to closed subschemes. More precisely, let X be a projective variety over a number field K , and let $Z(X)$ denote the set of closed subschemes of X . Let M_K be the set of places of K . Note that the closed subschemes $Y \in Z(X)$ are in one-to-one correspondence with quasi-coherent ideal sheaves $\mathcal{I}_Y \subseteq \mathcal{O}_X$, and we identify a closed subscheme Y with its ideal sheaf \mathcal{I}_Y . Generalizing the Weil height machine for Cartier divisors, Silverman assigned to each $Y \in Z(X)$ and each place $v \in M_K$ a local height function $\lambda_{Y,v}$, and to each $Y \in Z(X)$ a global height function $h_Y = \sum_{v \in M_K} \lambda_{Y,v}$ (both uniquely determined up to a bounded function). We now summarize some of the basic properties of height functions associated to closed subschemes.

Theorem 2.1. ([Sil87]) Let X be a projective variety over a number field K . Let $Z(X)$ be the set of closed subschemes of X . There are maps

$$\begin{aligned} Z(X) \times M_K &\rightarrow \{\text{functions } X(K) \rightarrow [0, +\infty]\}, \\ (Y, v) &\mapsto \lambda_{Y,v}, \\ Z(X) &\rightarrow \{\text{functions } X(K) \rightarrow [0, +\infty]\}, \\ Y &\mapsto h_Y, \end{aligned}$$

satisfying the following properties (we also write $\lambda_{X,Y,v}$ and $h_{X,Y}$ for clarity in (6)):

1. If $D \in Z(X)$ is an effective Cartier divisor, then $\lambda_{D,v}$ and h_D agree with the classical height functions associated to D .
2. If $W, Y \in Z(X)$ satisfy $W \subseteq Y$, then $h_W \leq h_Y + O(1)$ and $\lambda_{W,v} \leq \lambda_{Y,v} + O(1)$ for all $v \in M_K$.
3. If $W, Y \in Z(X)$ satisfy $\text{Supp}(W) \subseteq \text{Supp}(Y)$, then there exists a constant C such that $h_W \leq C \cdot h_Y + O(1)$ and $\lambda_{W,v} \leq C \cdot \lambda_{Y,v} + O(1)$ for all $v \in M_K$.
4. For all $W, Y \in Z(X)$, $\lambda_{W \cap Y, v} = \min\{\lambda_{W,v}, \lambda_{Y,v}\} + O(1)$.
5. For all $W, Y \in Z(X)$, we have $h_{W+Y} = h_W + h_Y + O(1)$ and $\lambda_{W+Y, v} = \lambda_{W,v} + \lambda_{Y,v} + O(1)$ for all $v \in M_K$.
6. Let $\phi : X' \rightarrow X$ be a morphism of projective varieties over K , and let $Y \in Z(X)$. Then

$$\begin{aligned} h_{X', \phi^*Y} &= h_{X,Y} \circ \phi + O(1), \\ \lambda_{X', \phi^*Y, v} &= \lambda_{X,Y,v} \circ \phi + O(1), \end{aligned}$$

for all $v \in M_K$.

7. If D and E are numerically equivalent Cartier divisors on X and A is an ample divisor on X , then for any $\varepsilon > 0$, we have

$$|h_D(P) - h_E(P)| < \varepsilon h_A(P) + O(1)$$

for all $P \in X(K)$. [Voj87, Proposition 1.2.9(d)].

Here, $Y \subset Z$, $Y + Z$, and ϕ^*Y are all defined in terms of the associated ideal sheaves (see [Sil87]). For a closed subscheme Y and finite set of places S of K , we let $m_{Y,S}(P) = \sum_{v \in S} \lambda_{Y,v}(P)$. For Cartier divisors D and E on a variety X , we will also write $D \geq E$ (or $E \leq D$) if $D - E$ is an effective divisor.

Remark 2.2. If Y is a zero-dimensional closed subscheme of a projective variety X over a field K , with $\text{Supp } Y = \{Q_1, \dots, Q_r\}$, then we can write

$$Y = Y_{Q_1} + \dots + Y_{Q_r},$$

where Y_{Q_i} is a closed subscheme supported only at Q_i (see, e.g., [GW20, Prop. 5.11]). Suppose that K is a number field and fix $v \in M_K$. Since $Y_{Q_i} \cap Y_{Q_j} = \emptyset$, $i \neq j$, it follows from Theorem 2.1 (4) that

$$\min\{\lambda_{Y_{Q_i},v}, \lambda_{Y_{Q_j},v}\} = O(1).$$

Since we also have $\lambda_{Y,v} = \sum_{i=1}^r \lambda_{Y_{Q_i},v} + O(1)$, for any point $P \in X(K)$ there exists $j \in \{1, \dots, r\}$ (depending on P and v) such that

$$\lambda_{Y,v}(P) = \lambda_{Y_{Q_j},v}(P) + O(1),$$

where the $O(1)$ is independent of P .

Definition 2.3. If Y is a zero-dimensional closed subscheme of a projective variety X over K and $Q \in Y(\overline{K})$, we let Y_Q be the closed subscheme $(Y_{\overline{K}})_Q$ of Remark 2.2 (applied to the closed subscheme $Y_{\overline{K}}$ of $X_{\overline{K}}$).

When $Q \in Y(K)$, we also naturally identify Y_Q as a closed subscheme over K .

We also recall the notion of a set of (D, S) -integral points on a projective variety X (see [Voj87] for more details).

Definition 2.4. Let D be an effective Cartier divisor on a projective variety X , both defined over a number field K . Let R be a subset of $X(K) \setminus \text{Supp } D$, and let S be a finite set of places of K containing all the archimedean places. We say that R is a set of (D, S) -integral points on X if

$$\sum_{v \in S} \lambda_{D,v}(P) = h_D(P) + O(1)$$

for all $P \in R$.

More generally, this definition can be extended to an arbitrary closed subscheme Y of X by appropriately replacing D with Y everywhere.

2.2 Multiplicities, Regular Sequences, and the Filtration Lemma

In this section, we collect together some useful facts involving multiplicities and regular sequences, and state the Filtration Lemma which will play a basic role in many of the proofs.

Let R be a Noetherian local ring and let \mathfrak{q} be a parameter ideal for R . Let $L_{\mathfrak{q},R}(n) = \text{length}(R/\mathfrak{q}^n)$.

Theorem 2.5 ([Eis95, Section 12.4]). For $n \gg 0$, $L_{\mathfrak{q},R}(n)$ agrees with a polynomial of degree $\dim R$ in n .

The leading term of any polynomial of degree d in $\mathbb{Q}[n]$ which takes only integer values for $n \gg 0$ is of the form $cn^d/d!$ for some integer c . Then we have the following notions of multiplicity going back to Hilbert and Samuel.

Definition 2.6 ([Eis95]). We define the multiplicity of \mathfrak{q} in R to be the integer $e = e(\mathfrak{q}, R)$ such that $|L_{\mathfrak{q}, R}(n) - en^{\dim R}/(\dim R)!| = O(n^{\dim R-1})$.

Definition 2.7 ([Ful89, Ex. 4.3.4]). Let Z be a zero-dimensional closed subscheme of a projective variety X with ideal sheaf \mathcal{I} and let $P \in \text{Supp } Z$. We define the multiplicity of Z at P to be $e(\mathcal{I}_P, \mathcal{O}_{X, P})$, and denote it by $\mu_P(Z)$ or simply μ_P if the context is clear.

We also need some facts about certain ideal quotients.

Definition 2.8. Let I and J be ideals in R . The ideal quotient (or colon ideal) $(I : J)$ is defined by

$$(I : J) = \{r \in R \mid rJ \subset I\}.$$

Lemma 2.9. Let $I = (f_1, \dots, f_m)$ be an ideal of a ring R generated by a regular sequence f_1, \dots, f_m in R . Then for any positive integers $n \geq i > 0$ and $j \in \{1, \dots, m\}$, we have

$$(I^n : (f_j)^i) = I^{n-i}.$$

Proof. Since $f_j^i \in I^i$, $I^{n-i} \subset (I^n : (f_j)^i)$ is clear.

For the other inclusion, suppose that $af_j^i \in I^n$. Consider the associated graded ring $G_I(R) := \bigoplus_{r \geq 0} I^r/I^{r+1}$. By [Mat80, Theorem 27], $G_I(R)$ is isomorphic to the polynomial ring $(R/I)[x_1, \dots, x_m]$, where the indeterminates x_1, \dots, x_m correspond to the images of f_1, \dots, f_m in the first graded piece of $G_I(R)$.

Suppose $a \in I^\ell \setminus I^{\ell+1}$, so that $0 \neq \bar{a} \in I^\ell/I^{\ell+1}$. Since x_j is not a zero divisor in $(R/I)[x_1, \dots, x_m]$, we have $0 \neq \overline{af_j^i} \in I^{\ell+i}/I^{\ell+i+1}$. Therefore $af_j^i \notin I^{\ell+i+1}$ and it follows that $\ell + i + 1 \geq n + 1$. Hence $\ell \geq n - i$. \square

We now make definitions involving proper intersections, general position, and regular sequences, in the context of closed subschemes.

Definition 2.10. Let D_1, \dots, D_r be effective Cartier divisors on a projective variety X . We say that D_1, \dots, D_r intersect properly if for every nonempty subset $I \subset \{1, \dots, r\}$ and every point $P \in \text{Supp } \bigcap_{i \in I} D_i$, the sequence $(f_i)_{i \in I}$ is a regular sequence in the local ring $\mathcal{O}_{X, P}$, where f_i locally defines D_i at P , $i = 1, \dots, r$. More generally, the definition can be extended to closed subschemes Y_1, \dots, Y_r of X (see [RW22] or [Voj23]).

Remark 2.11. If D_1 and D_2 are Cartier divisors on X that intersect properly, then $\mathcal{O}_X(-D_1 - D_2) = \mathcal{O}_X(-D_1) \cap \mathcal{O}_X(-D_2)$ (see [Voj23, §3, §4] for a vast abstraction of this property).

Definition 2.12. Let X be a projective variety of dimension n . We say that closed subschemes Y_1, \dots, Y_q of X are in general position if for every subset $I \subset \{1, \dots, q\}$ with $|I| \leq n + 1$ we have $\text{codim } \bigcap_{i \in I} Y_i \geq |I|$, where we use the convention that $\dim \emptyset = -1$.

Remark 2.13. If the Cartier divisors D_1, \dots, D_r on X intersect properly, then they are in general position. If X is Cohen-Macaulay then the converse holds [Mat80, Theorem 17.4]. In particular, if $X \subset \mathbb{P}^n$ is a hypersurface, D_1, \dots, D_r are effective divisors on \mathbb{P}^n , and X, D_1, \dots, D_r are in general position (on \mathbb{P}^n), then X, D_1, \dots, D_r intersect properly, and from the definitions it follows that $D_1|_X, \dots, D_r|_X$ intersect properly, as divisors on X .

We make a related definition for a sequence of nested closed subschemes.

Definition 2.14. Let $Y_1 \supset Y_2 \supset \cdots \supset Y_r$ be closed subschemes of a projective variety X . We say that $Y_1 \supset \cdots \supset Y_r$ is a regular chain of closed subschemes of X if for every $i \in \{1, \dots, r\}$ and every point $P \in \text{Supp } Y_i$, there is a regular sequence $f_1, \dots, f_i \in \mathcal{O}_{X,P}$ such that for $1 \leq j \leq i$, the ideal sheaf of Y_j is locally given by the ideal (f_1, \dots, f_j) in $\mathcal{O}_{X,P}$.

Remark 2.15. If D_1, \dots, D_r intersect properly on X , then $D_1 \supset D_1 \cap D_2 \supset \cdots \supset D_1 \cap D_2 \cap \cdots \cap D_r$ is a regular chain of closed subschemes of X .

We will use the following consequence of Lemma 2.9.

Definition 2.16. For an effective Cartier divisor D and closed subscheme Y of X , let $n_Y(D)$ be the largest nonnegative integer such that $n_Y(D)Y \subset D$, as closed subschemes.

Lemma 2.17. Let D, D' be effective Cartier divisors on a surface X and let Y be a closed subscheme supported at a point P . Suppose that $mD \supset Y$ is a regular chain of closed subschemes for some positive integer m . Then for any positive integer i ,

$$|n_Y(iD) + n_Y(D') - n_Y(iD + D')| \leq 1, \quad (2.1)$$

and

$$n_Y(iD) = \lfloor i/m \rfloor.$$

Proof. Let \mathcal{I} be the ideal sheaf of Y and let $\mathcal{I}_P \subset \mathcal{O}_P$ be the localization. Let $n = n_Y(iD + D')$. Suppose that D and D' are represented by f and f' locally at P . Then since $nY \subset iD + D'$, we have $f^i f' \in \mathcal{I}_P^n$. It follows that $(f^m)^{\lfloor i/m \rfloor} f' \in \mathcal{I}_P^n$ and, since $mD \supset Y$ is a regular chain of closed subschemes, by Lemma 2.9, $f' \in \mathcal{I}_P^{n - \lfloor i/m \rfloor}$. On the other hand, since $f^i = (f^m)^{\lfloor i/m \rfloor} f^{i - m\lfloor i/m \rfloor} \in \mathcal{I}_P^{\lfloor i/m \rfloor}$, if $f' \in \mathcal{I}_P^{n - \lfloor i/m \rfloor + 1}$ then $f^i f' \in \mathcal{I}_P^{n+1}$, contradicting the definition of n . It follows that

$$n - \lfloor i/m \rfloor \leq n_Y(D') \leq n - \lfloor i/m \rfloor.$$

The inequality (2.1) (and the remainder of the lemma) now follows if we show that $n_Y(iD) = \lfloor i/m \rfloor$. If i divides m and $iD = \frac{i}{m}(mD)$ then it follows from Lemma 2.9 that $n_Y(iD) = i/m$. Since $\lfloor i/m \rfloor(mD) \subset iD \subset (\lfloor i/m \rfloor + 1)(mD)$, we have $\lfloor i/m \rfloor \leq n_Y(iD) \leq \lfloor i/m \rfloor + 1$. But if $n_Y(iD) = \lfloor i/m \rfloor + 1$, then

$$n_Y(miD) \geq mn_Y(iD) > i,$$

contradicting $n_Y(miD) = i$. Therefore we must have $n_Y(iD) = \lfloor i/m \rfloor$ as claimed. \square

Finally, we recall the elementary, but very useful, Filtration Lemma whose utility in Diophantine approximation was recognized and introduced by Corvaja and Zannier [CZ04b, Lemma 3.2].

Lemma 2.18 (Filtration Lemma). *Let V be a vector space of finite dimension d over a field K . Let $V = W_1 \supset W_2 \supset \cdots \supset W_h$ and $V = W_1^* \supset W_2^* \supset \cdots \supset W_h^*$ be two filtrations on V . There exists a basis v_1, \dots, v_d of V that contains a basis of each W_j and W_j^* .*

In the situation of Lemma 2.18, we say that the basis v_1, \dots, v_d of V is adapted to the two filtrations.

3 Beta constant estimates

Throughout this section we work over a field K of characteristic 0. For a line bundle \mathcal{L} on a projective variety X over K , we will write $h^0(X, \mathcal{L})$ (or simply $h^0(\mathcal{L})$) for $\dim_K H^0(X, \mathcal{L})$, and if D is a Cartier divisor on X , we write $h^0(D)$ for $h^0(\mathcal{O}(D))$. We first recall the definition of the beta constant.

Definition 3.1. *Let X be a projective variety over a field K . Let \mathcal{L} be a big line bundle on X and let Y be a closed subscheme of X with associated sheaf of ideals \mathcal{I} . Then*

$$\beta(\mathcal{L}, Y) = \liminf_{N \rightarrow \infty} \frac{\sum_{m=1}^{\infty} h^0(X, \mathcal{L}^N \otimes \mathcal{I}^m)}{N h^0(X, \mathcal{L}^N)},$$

where \mathcal{L}^N denotes the tensor product of N copies of \mathcal{L} .

By a result of Vojta [Voj20] (assuming as we do that K has characteristic 0) the \liminf in the definition can be replaced by a limit. By abuse of notation, if D is a big Cartier divisor on X , we also write $\beta(D, Y)$ for $\beta(\mathcal{O}(D), Y)$.

It follows easily that for any positive integer n ,

$$\begin{aligned} \beta(nD, Y) &= n\beta(D, Y), \\ \beta(D, nY) &= \frac{1}{n}\beta(D, Y). \end{aligned}$$

In particular, we can use these properties to canonically extend the definition of $\beta(D, E)$ to \mathbb{Q} -divisors D and E (with D big). We also show that $\beta(D, E)$ depends only on the numerical equivalence class of D and E .

Lemma 3.2. *Let D, D', E, E' be nonzero effective Cartier divisors on a projective variety X over K with D and D' big. Suppose that $D \equiv D'$ and $E \equiv E'$. Then*

$$\beta(D, E) = \beta(D', E').$$

Proof. We prove that $\beta(D, E) = \beta(D, E')$ (the proof that $\beta(D, E) = \beta(D', E)$ being similar). Let $n = \dim X$. Since $E \equiv E'$ and D is big, for any positive rational ε , the \mathbb{Q} -divisor $E - E' + \varepsilon D$ is big. It follows that for a sufficiently large positive integer N , $h^0(N(E - E' + \varepsilon D)) > 0$, and so $E + \varepsilon D \sim E' + F$ for some \mathbb{Q} -divisor F . Therefore,

$$\beta(D, E') \geq \beta(D, E' + F) = \beta(D, E + \varepsilon D).$$

Fix an ample divisor A on X and let $c > \frac{A^{n-1} \cdot D}{A^{n-1} \cdot E}$ be a positive integer. Then $(ND - NmE) \cdot A^{n-1} < 0$ for all $m \geq c$, and so $h^0(ND - NmE) = 0$ for all $m \geq c$. Let M be a positive integer such that $M(E + \varepsilon D)$ is an integral Cartier divisor and let N be a positive integer such that $M|N$. Then

$$\sum_{m=1}^{\infty} h^0(ND - mM(E + \varepsilon D)) = \sum_{m=1}^{cN/M} h^0(ND - mM(E + \varepsilon D))$$

$$\begin{aligned}
&\geq \sum_{m=1}^{cN/M} h^0((N - c\varepsilon N)D - mME) \\
&= \sum_{m=1}^{\infty} h^0(N(1 - c\varepsilon)D - mME),
\end{aligned}$$

and so

$$\frac{\sum_{m=1}^{\infty} h^0(ND - mM(E + \varepsilon D))}{Nh^0(ND)} \geq \frac{\sum_{m=1}^{\infty} h^0(N(1 - c\varepsilon)D - mME)}{Nh^0(N(1 - c\varepsilon)D)} \frac{h^0(N(1 - c\varepsilon)D)}{h^0(ND)}.$$

Taking limits (see Definition 3.6 and Remark 3.7) gives

$$\begin{aligned}
\beta(D, E + \varepsilon D) &= M\beta(D, M(E + \varepsilon D)) \geq M\beta((1 - c\varepsilon)D, ME) \frac{\text{vol}((1 - c\varepsilon)D)}{\text{vol}(D)} \\
&\geq (1 - c\varepsilon)^{n+1} \beta(D, E).
\end{aligned}$$

Thus,

$$\beta(D, E') \geq (1 - c\varepsilon)^{n+1} \beta(D, E).$$

Since $\varepsilon > 0$ was arbitrary, we find that $\beta(D, E') \geq \beta(D, E)$. Interchanging the roles of E and E' yields that $\beta(D, E) = \beta(D, E')$ \square

We now prove some useful inequalities involving the beta constant.

Lemma 3.3. *Let Y be a closed subscheme of a projective variety X . Let D, E be effective Cartier divisors on X intersecting properly and suppose that $mY \subset D$ and $nY \subset E$ (as closed subschemes). Let A be a big Cartier divisor on X . Then*

$$\beta(A, Y) \geq m\beta(A, D) + n\beta(A, E).$$

In particular,

$$\beta(A, D \cap E) \geq \beta(A, D) + \beta(A, E).$$

Proof. We first note that if (x_1, \dots, x_n) is a regular sequence, then for any positive integers i_1, \dots, i_n , $(x_1^{i_1}, \dots, x_n^{i_n})$ is again a regular sequence [Mat80, Theorem 26]. This implies that if i and j are positive integers, then iD and jE intersect properly and $\mathcal{O}_X(-iD) \cap \mathcal{O}_X(-jE) = \mathcal{O}_X(-iD - jE)$.

Let $N > 0$. We consider the two filtrations on $H^0(X, \mathcal{O}(NA))$ given by order of vanishing along D and E , respectively:

$$\begin{aligned}
&H^0(X, \mathcal{O}(NA)) \supset H^0(X, \mathcal{O}(NA - D)) \supset H^0(X, \mathcal{O}(NA - 2D)) \supset \dots \\
&H^0(X, \mathcal{O}(NA)) \supset H^0(X, \mathcal{O}(NA - E)) \supset H^0(X, \mathcal{O}(NA - 2E)) \supset \dots
\end{aligned}$$

Let $s_1, \dots, s_{h^0(NA)}$ be a basis of $H^0(X, \mathcal{O}(NA))$ that is adapted to both filtrations. Let \mathcal{I} denote the sheaf of ideals associated to Y .

For a nonzero section $s \in H^0(X, \mathcal{O}(NA))$, let $\mu_D(s)$, $\mu_E(s)$, and $\mu_Y(s)$, respectively, be the largest nonnegative integer μ such that $s \in H^0(X, \mathcal{O}(NA - \mu D))$, $s \in H^0(X, \mathcal{O}(NA - \mu E))$, and $s \in H^0(X, \mathcal{O}(NA) \otimes \mathcal{I}^\mu)$, respectively. Since $s_1, \dots, s_{h^0(NA)}$ is adapted to the filtrations above, one easily finds

$$\begin{aligned} \sum_{j=1}^{\infty} h^0(\mathcal{O}(NA - jD)) &= \sum_{l=1}^{h^0(NA)} \mu_D(s_l), \\ \sum_{j=1}^{\infty} h^0(\mathcal{O}(NA - jE)) &= \sum_{l=1}^{h^0(NA)} \mu_E(s_l). \end{aligned}$$

We also have the inequality

$$\sum_{j=1}^{\infty} h^0(\mathcal{O}(NA) \otimes \mathcal{I}^j) \geq \sum_{l=1}^{h^0(NA)} \mu_Y(s_l).$$

Since D and E intersect properly, if $s_l \in H^0(X, \mathcal{O}(NA - iD))$ and $s_l \in H^0(X, \mathcal{O}(NA - jE))$, then $s_l \in H^0(X, \mathcal{O}(NA - iD - jE)) \subset H^0(X, \mathcal{O}(NA) \otimes \mathcal{I}^{im+jn})$. Therefore, $\mu_Y(s_l) \geq m\mu_D(s_l) + n\mu_E(s_l)$. It follows that

$$\sum_{j=1}^{\infty} h^0(\mathcal{O}(NA) \otimes \mathcal{I}^j) \geq m \sum_{i=1}^{\infty} h^0(\mathcal{O}(NA - iD)) + n \sum_{j=1}^{\infty} h^0(\mathcal{O}(NA - jE)).$$

Dividing by $Nh^0(\mathcal{O}(NA))$ and taking a limit then gives the result. \square

Before stating and proving the next inequality, we recall the definition of the Seshadri constant.

Definition 3.4. *Let X be a projective variety over a field K . Let Y be a closed subscheme of X and $\pi : \tilde{X} \rightarrow X$ be the blowing-up along Y . Let A be a nef Cartier divisor on X . Define the Seshadri constant $\epsilon(A, Y)$ of Y with respect to A to be the real number*

$$\epsilon(A, Y) = \sup\{\gamma \in \mathbb{Q}^{\geq 0} \mid \pi^*A - \gamma E \text{ is } \mathbb{Q}\text{-nef}\},$$

where E is an effective Cartier divisor on \tilde{X} whose associated invertible sheaf is the dual of $\pi^{-1}\mathcal{I}_Y \cdot \mathcal{O}_{\tilde{X}}$.

Lemma 3.5. *Let X be a normal projective variety of dimension r . Let Y be a closed subscheme of X with $\text{codim } Y \geq 2$, let $D \supset Y$ be an effective Cartier divisor, and let A be an ample Cartier divisor on X . Then*

$$\beta(A, Y) \geq \beta(A, D) + \frac{1}{r+1}\epsilon(A, Y).$$

Proof. Let $\pi : \tilde{X} \rightarrow X$ be the blowup of X along Y , and let E be the associated exceptional divisor. Let $\epsilon'_Y \leq \epsilon(A, Y)$ be a positive rational number and let $\delta > 0$ be a rational number. By the same proof as in [HL21], for all sufficiently small $\varepsilon > 0$ (depending on δ) and sufficiently large and divisible N , we can find an effective divisor F such that

1. $N(1 + \delta)A \sim F$
2. $\pi^*F \geq N(\epsilon'_Y + \varepsilon)E$
3. D and F are in general position.

As X is normal, the last condition and Serre's criterion imply that D and F intersect properly. For all sufficiently large and divisible N , $\pi_*\mathcal{O}_{\tilde{X}}(-N(\epsilon'_Y + \varepsilon)E) = \mathcal{I}_Y^{N(\epsilon'_Y + \varepsilon)}$ [HL21] and since X is normal and π is birational, we have $\pi_*\pi^*F = F$ [Voj23, Lemma 6.2(a)]. Then $\beta(A, F) = \frac{1}{(r+1)N(1+\delta)}$, $N(\epsilon'_Y + \varepsilon)Y \subset F$ (for sufficiently large and divisible N), $Y \subset D$, and Lemma 3.3 gives

$$\beta(A, Y) \geq \beta(A, D) + N(\epsilon'_Y + \varepsilon) \frac{1}{(r+1)N(1+\delta)} = \beta(A, D) + (\epsilon'_Y + \varepsilon) \frac{1}{(r+1)(1+\delta)}.$$

Since we may choose ε, δ , and $\epsilon'_Y - \epsilon(A, Y)$ arbitrarily small, we find that

$$\beta(A, Y) \geq \beta(A, D) + \frac{1}{r+1}\epsilon(A, Y)$$

as desired. □

When Y is a closed subscheme supported at a point P , we will relate $\beta(A, Y)$, $\mu_P(Y)$, and the volume of A . We first recall the definition of the volume of a divisor.

Definition 3.6. *Let X be a projective variety of dimension n , and let D be a Cartier divisor on X . The volume of D is defined to be the non-negative real number*

$$\text{vol}(D) = \limsup_{m \rightarrow \infty} \frac{h^0(X, mD)}{m^n/n!}.$$

Remark 3.7. *The lim sup in Definition 3.6 can be replaced by a limit [Laz04, Remark 2.2.50], and the definition can be extended to \mathbb{Q} -divisors [Laz04, Remark 2.2.39].*

Remark 3.8. *We have $\text{vol}(D) > 0$ if and only if D is big.*

Remark 3.9. *If D is nef, then $\text{vol}(D) = D^n$ is the top self-intersection of D .*

We need the following lemma.

Lemma 3.10. *Let A be a big Cartier divisor on a projective variety X of dimension r . Let Y be a closed subscheme of X supported at a point P of X , let \mathcal{I} be the associated ideal sheaf, and let μ_P be the multiplicity of Y at P . Then*

$$\dim H^0(X, \mathcal{O}(nA) \otimes \mathcal{I}^m) \geq h^0(X, \mathcal{O}(nA)) - \mu_P \frac{m^r}{r!} + O(m^{r-1}). \quad (3.1)$$

In particular, let $c \in \mathbb{Q}$ be such that

$$0 < c < \sqrt[r]{\frac{\text{vol}(A)}{\mu_P}}.$$

Then for $n \gg 0$ there exists an effective divisor F such that

1. $F \sim nA$,
2. $[cn]Y \subset F$ (as closed subschemes).

Proof. From the definition of the multiplicity μ_P , we have

$$\dim \mathcal{O}_P/\mathcal{I}_P^m = \mu_P \frac{m^r}{r!} + O(m^{r-1}). \quad (3.2)$$

By definition of the ideal sheaf, we have an exact sequence

$$0 \rightarrow \mathcal{I}^m \rightarrow \mathcal{O}_X \rightarrow i_*\mathcal{O}_{mY} \rightarrow 0,$$

where $i : mY \rightarrow X$ is the inclusion map. Tensoring with $\mathcal{O}(nA)$, we have an exact sequence

$$0 \rightarrow \mathcal{O}(nA) \otimes \mathcal{I}^m \rightarrow \mathcal{O}(nA) \rightarrow \mathcal{O}(nA) \otimes i_*\mathcal{O}_{mY} = i_*\mathcal{O}_{mY} \rightarrow 0,$$

where the equality $\mathcal{O}(nA) \otimes i_*\mathcal{O}_{mY} = i_*\mathcal{O}_{mY}$ follows easily from the fact that $\mathcal{O}(nA)$ is locally free of rank 1 and $i_*\mathcal{O}_{mY}$ is supported at a point. Taking global sections, we find

$$\begin{aligned} 0 \rightarrow H^0(X, \mathcal{O}(nA) \otimes \mathcal{I}^m) \rightarrow H^0(X, \mathcal{O}(nA)) \rightarrow H^0(X, i_*\mathcal{O}_{mY}) \\ = H^0(mY, \mathcal{O}_{mY}) = \mathcal{O}_P/\mathcal{I}_P^m. \end{aligned}$$

It follows that

$$\dim H^0(X, \mathcal{O}(nA) \otimes \mathcal{I}^m) \geq \dim H^0(X, \mathcal{O}(nA)) - \dim \mathcal{O}_P/\mathcal{I}_P^m,$$

and then (3.1) follows from (3.2).

From (3.1), the definition of the volume, and Remark 3.7, it follows that if $c < \sqrt[r]{\frac{\text{vol}(A)}{\mu_P}}$, then there exists a nonzero global section s of $\mathcal{O}(nA) \otimes \mathcal{I}^{\lfloor cn \rfloor}$ for $n \gg 0$, and taking $F = \text{div}(s)$ gives the desired divisor. \square

We now prove a basic inequality between $\beta(A, Y)$, $\mu_P(Y)$, and $\text{vol}(A)$.

Lemma 3.11. *Let A be a big Cartier divisor on a projective variety X of dimension r . Let Y be a closed subscheme of X supported at a point P of X , and let μ_P be the multiplicity of Y at P . Then*

$$\beta(A, Y) \geq \frac{r}{r+1} \sqrt[r]{\frac{\text{vol}(A)}{\mu_P}}.$$

Proof. Let $c \in \mathbb{Q}$ be such that

$$0 < c < \sqrt[r]{\frac{\text{vol}(A)}{\mu_P}}.$$

Let N be a positive integer such that cN is an integer. Let \mathcal{I} be the sheaf of ideals associated to Y . Then we calculate

$$\begin{aligned} \sum_{m=1}^{cN} h^0(\mathcal{O}(NA) \otimes \mathcal{I}^m) &\geq \sum_{m=1}^{cN} \left(h^0(\mathcal{O}(NA)) - \mu_P \frac{m^r}{r!} + O(N^{r-1}) \right) \\ &\geq cN h^0(\mathcal{O}(NA)) - \mu_P \frac{(cN)^{r+1}}{(r+1)!} + O(N^r). \end{aligned}$$

Dividing by $Nh^0(\mathcal{O}(NA))$ we find that

$$\frac{\sum_{m=1}^{cN} h^0(\mathcal{O}(NA) \otimes \mathcal{I}^m)}{Nh^0(\mathcal{O}(NA))} \geq c - \mu_P \frac{c^{r+1} N^r}{(r+1)! h^0(\mathcal{O}(NA))} + O\left(\frac{N^{r-1}}{h^0(\mathcal{O}(NA))}\right).$$

Letting $N \rightarrow \infty$ gives

$$\beta(A, Y) \geq \frac{c}{(r+1) \text{vol}(A)} ((r+1) \text{vol}(A) - \mu_P c^r).$$

Since this is true for all appropriate values of c , letting $c \rightarrow \sqrt[r]{\frac{\text{vol}(A)}{\mu_P}}$ gives the inequality. \square

Under more hypotheses, we give another estimate involving $\beta(A, Y)$, $\mu_P(Y)$, and $\text{vol}(A)$. Towards this end, we first prove the following lemma.

Lemma 3.12. *Let A and D be big Cartier divisors on a normal projective surface X , with D a positive multiple of a prime Cartier divisor. Let Y be a closed subscheme of X supported at a point P , and let μ_P be the multiplicity of Y at P . Suppose that $D \supset Y$ is a regular chain of closed subschemes of X and*

$$\mu_P < \text{vol}(D).$$

Let $c \in \mathbb{Q}$ be such that

$$c < \sqrt[r]{\frac{\text{vol}(A)}{\mu_P}}.$$

Then for $n \gg 0$ there exists an effective divisor F such that

1. $F \sim nA$,
2. $\lfloor cn \rfloor Y \subset F$ (as closed subschemes),
3. D and F intersect properly.

Proof. For simplicity and ease of notation, we assume throughout that $cn \in \mathbb{Z}$. Let $D = mD_0$ where D_0 is a prime Cartier divisor on X . Since $\mu_P < \text{vol}(D)$, applying Lemma 3.10 with $A = D$ and an appropriate rational number $c' > 1$, it follows that for any sufficiently large and divisible integer n' , there exists an effective divisor F' such

that $F' \sim n'D$ and $n'c'Y \subset F'$. Since X is normal, we can write $F' = iD_0 + D'$, where i is a nonnegative integer, D' is effective, and $\text{Supp } D_0 \not\subset \text{Supp } D'$. By Lemma 2.17,

$$n_Y(D') \geq n_Y(F') - n_Y(iD_0) - 1 \geq n'c' - i/m - 1.$$

Let $n_0 = mn' - i$, and note also that $D' \sim n_0D_0$. Clearly $i \leq mn'$, and in fact $i < mn'$ for sufficiently large n' (or equivalently, $n_0 > 0$, as we now assume) since $c' > 1$ and therefore $n_Y(D') \geq n'c' - i/m - 1 \geq n'(c' - 1) - 1 > 0$ for large n' . Choosing n' sufficiently large, we also have $n_Y(D') > n' - i/m = n_0/m$.

By Lemma 3.10 again, there exists an effective divisor F'' such that $F'' \sim nA$ and $cnY \subset F''$. Write $F'' = jD_0 + D''$, where j is a nonnegative integer, D'' is effective, and $\text{Supp } D_0 \not\subset \text{Supp } D''$. Replacing F'' , n , j , and D'' by suitable multiples, we may assume that $n_0|j$. Then $F'' = jD_0 + D'' \sim \frac{j}{n_0}D' + D''$. By the same argument as before,

$$n_Y(D'') \geq n_Y(F'') - n_Y(jD_0) - 1 \geq cn - j/m - 1.$$

Let $F = \frac{j}{n_0}D' + D''$. Then

$$n_Y(F) \geq \frac{j}{n_0}n_Y(D') + n_Y(D'') \geq \frac{j}{n_0} \frac{n_0}{m} + cn - j/m - 1 = cn - 1.$$

Replacing c by a slightly large constant and taking n sufficiently large then gives

$$n_Y(F) \geq cn.$$

It follows that $cnY \subset F$, and we end by noting that $F \sim nA$ and $\text{Supp } D \not\subset \text{Supp } F = \text{Supp } D' \cup \text{Supp } D''$, which implies that D and F are in general position, and hence intersect properly (as a normal surface is Cohen-Macaulay). \square

We prove the following inequality, which will be used in Corollary 4.4. For simplicity, we state it under an ampleness assumption.

Lemma 3.13. *Let A and D be ample Cartier divisors on a normal projective surface X , with D a positive multiple of a prime Cartier divisor. Let Y be a closed subscheme of X supported at a point P , and let μ_P be the multiplicity of Y at P . Suppose that $D \supset Y$ is a regular chain of closed subschemes of X and*

$$\mu_P < D^2.$$

Then

$$\beta(A, Y) \geq \left(1 + \sqrt{\frac{D^2}{\mu_P}}\right) \beta(A, D) \tag{3.3}$$

and

$$\begin{aligned} \beta(A, Y) - \beta(A, D) &\geq \frac{2}{3} \sqrt{\frac{A^2}{\mu_P}} \frac{\sqrt{\frac{D^2}{\mu_P}}}{1 + \sqrt{\frac{D^2}{\mu_P}}} \\ &> \frac{1}{3} \sqrt{\frac{A^2}{\mu_P}}. \end{aligned}$$

Proof. Let $c \in \mathbb{Q}$ with $0 < c < \sqrt{\frac{D^2}{\mu_P}}$. Then by Lemma 3.12, for n sufficiently large with $nc \in \mathbb{Z}$, we can find an effective divisor F such that $F \sim nD$, $ncY \subset F$, and D and F intersect properly. Note that $\beta(A, F) = \beta(A, nD) = \frac{1}{n}\beta(A, D)$. Then by Lemma 3.3, we have

$$\beta(A, Y) \geq \beta(A, D) + cn\beta(A, F) = \beta(A, D) + c\beta(A, D).$$

Since this holds for every positive rational $c < \sqrt{\frac{D^2}{\mu_P}}$, the inequality (3.3) holds. This implies

$$\begin{aligned} \beta(A, Y) - \beta(A, D) &\geq \beta(A, Y) - \frac{1}{1 + \sqrt{\frac{D^2}{\mu_P}}}\beta(A, Y) \\ &\geq \frac{\sqrt{\frac{D^2}{\mu_P}}}{1 + \sqrt{\frac{D^2}{\mu_P}}}\beta(A, Y) \\ &\geq \frac{2}{3}\sqrt{\frac{A^2}{\mu_P}} \frac{\sqrt{\frac{D^2}{\mu_P}}}{1 + \sqrt{\frac{D^2}{\mu_P}}} \\ &> \frac{1}{3}\sqrt{\frac{A^2}{\mu_P}}, \end{aligned}$$

where the last line uses that $\mu_P < D^2$. \square

Finally, we prove a useful estimate for certain beta constants on the blowup of a projective variety X of dimension n at a point (the idea is to give a slight improvement to the equality $\beta(D, D) = \frac{1}{n+1}$ for an ample divisor D on X).

Lemma 3.14. *Let D be an ample effective Cartier divisor on a projective variety X over K of dimension n . Let $P \in X(K)$, and let $\pi : \tilde{X} \rightarrow X$ be the blowup at P , with associated exceptional divisor E . Then for all sufficiently small positive $\delta \in \mathbb{Q}$,*

$$\beta(\pi^*D - \delta E, \pi^*D - E) > \frac{1}{n+1}. \quad (3.4)$$

Proof. First, we recall that for some positive $\delta' \in \mathbb{Q}$, the \mathbb{Q} -divisor $\pi^*D - \delta E$ is ample for all $0 < \delta < \delta'$, $\delta \in \mathbb{Q}$ [Har77, II, Prop. 7.10(b)]. Let $\delta \in \mathbb{Q}$ satisfy $0 < \delta < \min\{\delta', 1\}$. Since $\pi^*D - \delta E$ is ample,

$$(\pi^*D - \delta E)^{n-1} \cdot E = (-\delta)^{n-1} E^n > 0.$$

Let

$$\gamma := (-1)^{n-1} E^n > 0.$$

Then

$$(a\pi^*D - bE)^n = a^n(\pi^*D)^n + (-1)^n b^n E^n$$

$$= a^n D^n - \gamma b^n.$$

Let N be such that $N\delta \in \mathbb{Z}$. We need to estimate

$$h^0\left(\tilde{X}, N(\pi^*D - \delta E) - m(\pi^*D - E)\right) = h^0\left(\tilde{X}, (N - m)\pi^*D - (N\delta - m)E\right)$$

for $m > 0$. We break the estimate into two cases. We first consider $0 < m \leq N\delta$. Since

$$\frac{N\delta - m}{N - m} < \delta < \delta',$$

$(N - m)\pi^*D - (N\delta - m)E$ is ample, and it follows from Riemann-Roch estimates (e.g., using that π^*D and $\pi^*D - \delta E$ are \mathbb{Q} -nef; see [Aut09, p. 233, Cas $k \leq n$]) that

$$\begin{aligned} h^0((N - m)\pi^*D - (N\delta - m)E) &= \frac{((N - m)\pi^*D - (N\delta - m)E)^n}{n!} + O(N^{n-1}) \\ &= \frac{(N - m)^n D^n - \gamma(N\delta - m)^n}{n!} + O(N^{n-1}) \end{aligned}$$

for $0 < m \leq N\delta$. Next, for $N\delta < m < N$, note that

$$(N - m)\pi^*D - (N\delta - m)E \geq (N - m)\pi^*D,$$

and so for $N\delta < m < N$,

$$h^0((N - m)\pi^*D - (N\delta - m)E) \geq h^0((N - m)\pi^*D) = \frac{(N - m)^n D^n}{n!} + O(N^{n-1}).$$

Therefore,

$$\begin{aligned} \sum_{m \geq 1} h^0(N(\pi^*D - \delta E) - m(\pi^*D - E)) \\ \geq \sum_{m=1}^{N\delta} \frac{(N - m)^n D^n - \gamma(N\delta - m)^n}{n!} + \sum_{m=N\delta+1}^N \frac{(N - m)^n D^n}{n!} + O(N^n). \end{aligned}$$

By standard formulas,

$$\sum_{m=1}^{N\delta} \frac{(N - m)^n D^n - \gamma(N\delta - m)^n}{n!} = \frac{N^{n+1}}{(n+1)!} (D^n - (1 - \delta)^{n+1} D^n - \gamma \delta^{n+1}) + O(N^n)$$

and

$$\sum_{m=N\delta+1}^N \frac{(N - m)^n D^n}{n!} = \frac{N^{n+1}}{(n+1)!} (1 - \delta)^{n+1} D^n + O(N^n).$$

Therefore,

$$\sum_{m \geq 1} h^0(N(\pi^*D - \delta E) - m(\pi^*D - E)) \geq \frac{N^{n+1}}{(n+1)!} (D^n - \gamma \delta^{n+1}) + O(N^n).$$

On the other hand,

$$\begin{aligned} Nh^0(N(\pi^*D - \delta E)) &= N \left(\frac{N^n}{n!} (D^n - \gamma\delta^n) + O(N^{n-1}) \right) \\ &= \frac{N^{n+1}}{n!} (D^n - \gamma\delta^n) + O(N^n). \end{aligned}$$

Note also that, in particular, $\gamma\delta^n < D^n$. Putting things together, we find

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{\sum_{m \geq 1} h^0(N(\pi^*D - \delta E) - m(\pi^*D - E))}{Nh^0(N(\pi^*D - \delta E))} &\geq \frac{1}{n+1} \frac{D^n - \gamma\delta^{n+1}}{D^n - \gamma\delta^n} \\ &> \frac{1}{n+1} \end{aligned}$$

as $\delta < 1$. □

4 Main Result and Some Consequences

We now prove (and restate) the Main Theorem.

Theorem 4.1. *Let X be a projective surface defined over a number field K . Let S be a finite set of places of K . For each $v \in S$, let $D_v \supset Y_v$ be a regular chain of nonempty closed subschemes of X . Let A be a big Cartier divisor on X , and let $\varepsilon > 0$. Then there exists a proper Zariski-closed subset $Z \subset X$ such that for all $P \in X(K) \setminus Z$, we have*

$$\sum_{v \in S} \left(\beta(A, D_v) \lambda_{D_v, v}(P) + (\beta(A, Y_v) - \beta(A, D_v)) \lambda_{Y_v, v}(P) \right) < (1 + \varepsilon) h_A(P).$$

Proof. From the regular chain assumption, for all $v \in S$, D_v is an effective Cartier divisor and $\dim Y_v = 0$. We may further reduce to the case that Y_v is supported at a single point: after replacing K by a finite extension L and replacing S by the set of places of L lying above S , we may assume that every point in the support of Y_v is K -rational. Then we may write $Y_v = Y_{1,v} + \cdots + Y_{r,v}$, where $Y_{1,v}, \dots, Y_{r,v}$ are closed subschemes supported at distinct K -rational points. Now the reduction follows from observing that $\beta(A, Y_{i,v}) \geq \beta(A, Y_v)$ for all i , and $\lambda_{Y_v, v}(P) = \max_i \lambda_{Y_{i,v}, v}(P) + O(1)$ for all $P \in X(K) \setminus Y_v$ (see Remark 2.2).

Let $N > 0$ and let $v \in S$. We consider the two filtrations on $H^0(X, \mathcal{O}(NA))$ given by order of vanishing along D_v and Y_v , respectively:

$$H^0(X, \mathcal{O}(NA)) \supset H^0(X, \mathcal{O}(NA - D_v)) \supset H^0(X, \mathcal{O}(NA - 2D_v)) \supset \cdots$$

and

$$H^0(X, \mathcal{O}(NA)) \supset H^0(X, \mathcal{O}(NA) \otimes \mathcal{I}_v) \supset H^0(X, \mathcal{O}(NA) \otimes \mathcal{I}_v^2) \supset \cdots.$$

By Lemma 2.18, there exists a basis $s_{1,v}, \dots, s_{h^0(NA),v}$ of $H^0(X, \mathcal{O}(NA))$ adapted to both filtrations. It follows that

$$\begin{aligned} \sum_{i=1}^{h^0(NA)} \operatorname{div}(s_{i,v}) &\geq \sum_{i=1}^{\infty} (h^0(NA - iD_v) - h^0(NA - (i+1)D_v)) iD_v \\ &\geq \left(\sum_{i=1}^{\infty} h^0(NA - iD_v) \right) D_v \end{aligned}$$

and similarly,

$$\sum_{i=1}^{h^0(NA)} \operatorname{div}(s_{i,v}) \supset \left(\sum_{i=1}^{\infty} h^0(\mathcal{O}(NA) \otimes \mathcal{I}_v^i) \right) Y_v.$$

as closed subschemes. Then we can write

$$\sum_{i=1}^{h^0(NA)} \operatorname{div}(s_{i,v}) = \left(\sum_{i=1}^{\infty} h^0(NA - iD_v) \right) D_v + F_v$$

for some effective divisor F_v . By Lemma 2.17,

$$\begin{aligned} n_{Y_v}(F_v) &\geq n_{Y_v} \left(\sum_{i=1}^{h^0(NA)} \operatorname{div}(s_{i,v}) \right) - \left(\sum_{i=1}^{\infty} h^0(NA - iD_v) \right) - 1 \\ &\geq \left(\sum_{i=1}^{\infty} h^0(\mathcal{O}(NA) \otimes \mathcal{I}_v^i) \right) - \left(\sum_{i=1}^{\infty} h^0(NA - iD_v) \right) - 1. \end{aligned}$$

It follows that

$$\begin{aligned} \lambda_{F_v,v}(P) &\geq n_{Y_v}(F_v) \lambda_{Y_v,v}(P) + O(1) \\ &\geq \left(\sum_{i=1}^{\infty} h^0(\mathcal{O}(NA) \otimes \mathcal{I}_v^i) - \sum_{i=1}^{\infty} h^0(NA - iD_v) - 1 \right) \lambda_{Y_v,v}(P) + O(1) \end{aligned}$$

for $P \in X(K) \setminus \operatorname{Supp} F_v$.

We now apply the Subspace Theorem in the form of [RV20, Theorem 2.10], with the union of the bases $s_{1,v}, \dots, s_{h^0(NA),v}, v \in S$, of $H^0(X, \mathcal{O}(NA))$ constructed above; we obtain that for $\varepsilon > 0$, there exists a proper Zariski-closed subset Z of X such that, up to $O(1)$,

$$\begin{aligned} \sum_{v \in S} \left(\sum_{i=1}^{\infty} h^0(NA - iD_v) \right) \lambda_{D_v,v}(P) &+ \left(\sum_{i=1}^{\infty} h^0(\mathcal{O}(NA) \otimes \mathcal{I}_v^i) - \sum_{i=1}^{\infty} h^0(NA - iD_v) - 1 \right) \lambda_{Y_v,v}(P) \\ &\leq \sum_{v \in S} \left(\sum_{i=1}^{\infty} h^0(NA - iD_v) \right) \lambda_{D_v,v}(P) + \lambda_{F_v,v}(P) \\ &\leq (h^0(NA) + \varepsilon) h_{NA}(P) \\ &\leq (Nh^0(NA) + N\varepsilon) h_A(P) \end{aligned}$$

for all $P \in X(K) \setminus Z$. Dividing by $Nh^0(NA)$ and taking N sufficiently large gives the desired inequality. \square

We show that Theorem 1.4 implies the Ru-Vojta inequality [RV20] for surfaces (a similar, but more complicated argument, could be used to derive Theorem 1.2 in the case of surfaces).

Corollary 4.2. *Let X be a projective surface over a number field K , and let D_1, \dots, D_q be nonzero effective Cartier divisors intersecting properly on X . Let \mathcal{L} be a big line bundle on X . Let S be a finite set of places of K and let $\varepsilon > 0$. Then there exists a proper Zariski-closed subset $Z \subset X$ such that for all $P \in X(K) \setminus Z$, we have*

$$\sum_{i=1}^q \beta(\mathcal{L}, D_i) m_{D_i, S}(P) \leq (1 + \varepsilon) h_{\mathcal{L}}(P).$$

Proof. For any point $P \in X(K)$, we have

$$\sum_{i=1}^q \beta(\mathcal{L}, D_i) m_{D_i, S}(P) \leq \sum_{v \in S} \left(\beta(\mathcal{L}, D_{P,1,v}) \lambda_{D_{P,1,v}, v}(P) + \beta(\mathcal{L}, D_{P,2,v}) \lambda_{D_{P,2,v}, v}(P) + O(1) \right),$$

for some divisors $D_{P,1,v}$ and $D_{P,2,v}$ depending on P and v (but with the $O(1)$ term independent of P). Then by considering the finitely many choices for $D_{P,1,v}$ and $D_{P,2,v}$, it suffices to study the sum

$$\sum_{v \in S} \left(\beta(\mathcal{L}, D_{1,v}) \lambda_{D_{1,v}, v}(P) + \beta(\mathcal{L}, D_{2,v}) \lambda_{D_{2,v}, v}(P) \right),$$

where for each $v \in S$ the divisors $D_{1,v}, D_{2,v}$ intersect properly. Again, by considering finitely many cases we may assume that $\lambda_{D_{1,v}, v}(P) \geq \lambda_{D_{2,v}, v}(P)$ for all $v \in S$ and thus

$$\lambda_{D_{2,v}, v}(P) = \min\{\lambda_{D_{1,v}, v}(P), \lambda_{D_{2,v}, v}(P)\} = \lambda_{D_{1,v} \cap D_{2,v}, v}(P) + O(1)$$

for all $v \in S$. By Lemma 3.3,

$$\beta(\mathcal{L}, D_{1,v} \cap D_{2,v}) \geq \beta(\mathcal{L}, D_{1,v}) + \beta(\mathcal{L}, D_{2,v}).$$

Therefore, by Theorem 1.4, up to $O(1)$,

$$\begin{aligned} & \sum_{v \in S} \left(\beta(\mathcal{L}, D_{1,v}) \lambda_{D_{1,v}, v}(P) + \beta(\mathcal{L}, D_{2,v}) \lambda_{D_{2,v}, v}(P) \right) \\ &= \sum_{v \in S} \left(\beta(\mathcal{L}, D_{1,v}) \lambda_{D_{1,v}, v}(P) + \beta(\mathcal{L}, D_{2,v}) \lambda_{D_{1,v} \cap D_{2,v}, v}(P) \right) \\ &\leq \sum_{v \in S} \left(\beta(\mathcal{L}, D_{1,v}) \lambda_{D_{1,v}, v}(P) + (\beta(\mathcal{L}, D_{1,v} \cap D_{2,v}) - \beta(\mathcal{L}, D_{1,v})) \lambda_{D_{1,v} \cap D_{2,v}, v}(P) \right) \\ &\leq (1 + \varepsilon) h_{\mathcal{L}}(P) \end{aligned}$$

for all K -rational points outside a proper Zariski-closed subset Z of X . Finally, we may omit any $O(1)$ term by enlarging Z . \square

From [HL21], if X is a nonsingular projective variety of dimension n and D is a divisor on X , then

$$\beta(A, D) \geq \frac{1}{n+1} \epsilon(A, D),$$

and in particular, on a surface X ,

$$\beta(A, D) \geq \frac{1}{3} \epsilon(A, D). \quad (4.1)$$

Using inequality (4.1) and Lemma 3.5, we immediately derive from Theorem 1.4 the inequality of Heier and the second author [HL21] for surfaces, under a smoothness and regular chain assumption:

Corollary 4.3. *Let X be a nonsingular projective surface defined over a number field K . Let S be a finite set of places of K . For each $v \in S$, let $D_v \supset Y_v$ be a regular chain of nonempty closed subschemes of X . Let A be an ample Cartier divisor on X , and let $\varepsilon > 0$. Then there exists a proper Zariski-closed subset $Z \subset X$ such that for all $P \in X(K) \setminus Z$, we have*

$$\sum_{v \in S} \left(\epsilon(A, D_v) \lambda_{D_v, v}(P) + \epsilon(A, Y_v) \lambda_{Y_v, v}(P) \right) < (3 + \varepsilon) h_A(P).$$

Under more hypotheses, as an immediate consequence of Lemma 3.13, we can replace $\epsilon(A, Y_v)$ by a quantity depending only on self-intersection numbers and the multiplicity of Y_v at a point.

Corollary 4.4. *Let X be a normal projective surface defined over a number field K . Let S be a finite set of places of K . For each $v \in S$, let $D_v \supset Y_v$ be a regular chain of closed subschemes of X such that D_v is a positive multiple of an ample prime Cartier divisor, Y_v is supported at a point $Q_v \in X(K)$, and*

$$\mu_v = \mu_{Q_v}(Y_v) < D_v^2.$$

Let A be an ample divisor on X , and let $\varepsilon > 0$. Then there exists a proper Zariski-closed subset $Z \subset X$ such that for all $P \in X(K) \setminus Z$, we have

$$\sum_{v \in S} \left(3\beta(A, D_v) \lambda_{D_v}(P) + 2 \sqrt{\frac{A^2}{\mu_v}} \frac{\sqrt{\frac{D_v^2}{\mu_v}}}{1 + \sqrt{\frac{D_v^2}{\mu_v}}} \lambda_{Y_v}(P) \right) < (3 + \varepsilon) h_A(P).$$

In particular, there exists a proper Zariski-closed subset $Z \subset X$ such that for all $P \in X(K) \setminus Z$, we have

$$\sum_{v \in S} \left(3\beta(A, D_v) \lambda_{D_v}(P) + \sqrt{\frac{A^2}{\mu_v}} \lambda_{Y_v}(P) \right) < (3 + \varepsilon) h_A(P).$$

Remark 4.5. For purposes of comparison, if X is a nonsingular projective surface over K and Y is supported at a point $Q \in X(K)$, then we have the inequality [CEL01, Remark 2.4]

$$\epsilon(A, Y) \leq \sqrt{\frac{A^2}{\mu_Q(Y)}}.$$

When $\mu_Q(Y) = 1$ and A^2 is not a perfect square, this inequality has been conjectured to always be strict (more precisely, it has been conjectured that the Seshadri constant at a point is always rational).

In the situation of Remark 4.5, it follows that we have the inequalities

$$\beta(A, Y) \geq \frac{2}{3} \sqrt{\frac{A^2}{\mu_Q(Y)}} \geq \frac{2}{3} \epsilon(A, Y).$$

We give a simple example where the inequalities are all strict.

Example 4.6. Let $X = \mathbb{P}^1 \times \mathbb{P}^1$, let A be a divisor of type $(1, 1)$ on X , and let $Y = Q \in X(K)$ be a point (viewed as a closed subscheme with the reduced induced structure). Then elementary computations give

$$\begin{aligned} \beta(A, Y) &= 1, \\ \sqrt{\frac{A^2}{\mu_Q(Y)}} &= \sqrt{2}, \\ \epsilon(A, Y) &= 1. \end{aligned}$$

Then we have strict inequalities

$$\beta(A, Y) > \frac{2}{3} \sqrt{\frac{A^2}{\mu_Q(Y)}} > \frac{2}{3} \epsilon(A, Y).$$

5 Applications

In the remaining sections, we investigate some Diophantine applications of our main result Theorem 1.4. In Section 5.1 we prove, under suitable conditions, an inequality for the “gcd height” $h_{D_i \cap D_j}(P)$, $i \neq j$, when P is an S -integral point with respect to three properly intersecting numerically parallel divisors D_1, D_2, D_3 on a surface. This inequality may be viewed as complementary to the gcd inequalities of Bugeaud-Corvaja-Zannier [BCZ03], Corvaja-Zannier [CZ05], and Wang-Yasufuku [WY21] (for surfaces) who study the height $h_Q(P)$ (under the same integrality assumption on P) when Q is *not* in the intersection of two of the divisors D_i . In Section 5.2, we apply our gcd inequalities to study the equation $f(a^m, y) = b^n$, which was previously studied by Corvaja and Zannier [CZ00]. In Section 5.3, we study integral points on surfaces on the complement of three numerically parallel divisors with nonempty intersection.

The results expand on earlier work of Corvaja and Zannier [CZ06] in a similar setting. Finally, using the results on integral points of Section 5.3, we study certain families of unit equations, proving a general result following work of Corvaja-Zannier [CZ06, CZ10] and the second author [Lev06].

5.1 Greatest Common Divisors on Surfaces

In 2003, Bugeaud, Corvaja, and Zannier [BCZ03] initiated a new line of results with the following inequality involving greatest common divisors:

Theorem 5.1 (Bugeaud-Corvaja-Zannier [BCZ03]). *Let $a, b \in \mathbb{Z}$ be multiplicatively independent integers. Then for every $\varepsilon > 0$,*

$$\log \gcd(a^n - 1, b^n - 1) \leq \varepsilon n \quad (5.1)$$

for all but finitely many positive integers n .

The inequality (5.1) was subsequently generalized by Corvaja and Zannier [CZ05], allowing a^n and b^n to be replaced by elements u and v , respectively, of a fixed finitely generated subgroup of $\overline{\mathbb{Q}}^*$, and replacing $u - 1$ and $v - 1$ by more general pairs of polynomials in u and v . Silverman [Sil05] interpreted these results in terms of heights and as a special case of Vojta's Conjecture. The second author [Lev19] further generalized these inequalities to multivariate polynomials, and Wang and Yasufuku proved the following general version of these results (see work of the first and second author [HL22] for an even more general version):

Theorem 5.2 (Wang-Yasufuku [WY21]). *Let X be a Cohen-Macaulay variety of dimension n defined over a number field K , and let S be a finite set of places of K . Let D_1, \dots, D_{n+1} be effective Cartier divisors defined over K and in general position. Suppose that there exists an ample Cartier divisor A on X and positive integers d_1, \dots, d_{n+1} such that $D_i \equiv d_i A$, $i = 1, \dots, n+1$. Let Y be a closed subscheme of X of codimension at least 2 that does not contain any point of the set*

$$\bigcup_{i=1}^{n+1} \bigcap_{j \neq i} \text{Supp} D_j. \quad (5.2)$$

Let $\varepsilon > 0$. Then there exists a proper Zariski-closed subset Z of X such that for any set R of $(\sum_{i=1}^n D_i, S)$ -integral points in $X(K)$, we have

$$h_Y(P) \leq \varepsilon h_A(P)$$

for all points $P \in R \setminus Z$.

When X is a surface, in Theorem 5.2 one may reduce to the case that $Y = Q$ is a point, and the condition (5.2) is simply the requirement that $Q \notin \cup_{i \neq j} D_i \cap D_j$. The main result of this section proves an inequality as in Theorem 5.2, but under the complementary condition $Q \in \cup_{i \neq j} D_i \cap D_j$, along with some mild additional hypotheses (necessary to exclude the case of lines in \mathbb{P}^2 where the analogous statement is false (Example 5.5)).

Theorem 5.3. *Let D_1, D_2, D_3 be effective divisors intersecting properly on a projective surface X , all defined over a number field K . Suppose that there exist positive integers a_1, a_2, a_3 such that a_1D_1, a_2D_2, a_3D_3 are all numerically equivalent to an ample divisor D . Suppose that for some $i_0, j_0 \in \{1, 2, 3\}$, $i_0 \neq j_0$, and for all $Q \in X(\overline{K})$,*

$$\beta(D, (a_{i_0}D_{i_0} \cap a_{j_0}D_{j_0})_Q) > \frac{2}{3}, \quad (5.3)$$

with $(a_{i_0}D_{i_0} \cap a_{j_0}D_{j_0})_Q$ as in Definition 2.3. Let S be a finite set of places of K containing all the archimedean ones and let $\varepsilon > 0$. Then there exists a proper Zariski-closed subset $Z \subset X$ such that for any set $R \subset X(K)$ of $(D_1 + D_2 + D_3, S)$ -integral points and all but finitely many points $P \in R \setminus Z$, we have

$$h_{D_{i_0} \cap D_{j_0}}(P) \leq \varepsilon h_D(P).$$

The condition (5.3) can be replaced by the simpler condition that $D_{i_0} \cap D_{j_0}$ contains more than one point:

Corollary 5.4. *Suppose that the same hypotheses as in Theorem 5.3 hold, except that (5.3) is replaced by the assumption that $D_{i_0} \cap D_{j_0}$ contains more than one point (over \overline{K}). Then there exists a proper Zariski-closed subset $Z \subset X$ such that for any set $R \subset X(K)$ of $(D_1 + D_2 + D_3, S)$ -integral points and all but finitely many points $P \in R \setminus Z$, we have*

$$h_{D_{i_0} \cap D_{j_0}}(P) \leq \varepsilon h_D(P).$$

Proof of Theorem 5.3. After replacing K by a finite extension, we may assume that every point in the support of $D_i \cap D_j$, $i \neq j$, is K -rational.

We note that since $a_i D_i \equiv D$, we have $\beta(D, a_i D_i) = \frac{1}{3}$ for all i . Moreover, by Lemma 3.3, for all $Q \in X(\overline{K})$ and $i \neq j$, we have

$$\beta(D, (a_i D_i \cap a_j D_j)_Q) \geq \beta(D, a_i D_i \cap a_j D_j) \geq \beta(D, a_i D_i) + \beta(D, a_j D_j) \geq \frac{2}{3}. \quad (5.4)$$

We let

$$\gamma = \min_{Q \in X(\overline{K})} 3\beta(D, (a_{i_0}D_{i_0} \cap a_{j_0}D_{j_0})_Q) - 2.$$

By assumption, $\gamma > 0$. By Remark 2.2, for any point $P \in X(K)$, $v \in S$, $i \neq j$, there is a point $Q \in \text{Supp}(a_i D_i \cap a_j D_j)$ (depending on P , v , and i and j) such that

$$\lambda_{a_i D_i \cap a_j D_j, v}(P) = \lambda_{(a_i D_i \cap a_j D_j)_Q, v}(P) + O(1) \quad (5.5)$$

where the constant in the $O(1)$ is independent of P .

For $v \in S$, let $i_v, j_v \in \{1, 2, 3\}$, $i_v \neq j_v$. Let $\varepsilon > 0$ and let

$$\gamma_v = \begin{cases} \gamma & \text{if } \{i_v, j_v\} = \{i_0, j_0\}, \\ 0 & \text{otherwise.} \end{cases}$$

By (5.4), (5.5), the definitions of γ and γ_v , and Theorem 1.4, there exists a proper Zariski-closed subset Z of X such that up to $O(1)$ (and after multiplying by 3)

$$\begin{aligned}
& \sum_{v \in S} \left(\lambda_{a_{i_v} D_{i_v}, v}(P) + (1 + \gamma_v) \lambda_{a_{i_v} D_{i_v} \cap a_{j_v} D_{j_v}, v}(P) \right) \\
&= \sum_{v \in S} \left(\lambda_{a_{i_v} D_{i_v}, v}(P) + (1 + \gamma_v) \max_Q \lambda_{(a_{i_v} D_{i_v} \cap a_{j_v} D_{j_v})_Q, v}(P) \right) \\
&\leq \sum_{v \in S} \left(\lambda_{a_{i_v} D_{i_v}, v}(P) + (3 \min_Q \beta(D, (a_{i_v} D_{i_v} \cap a_{j_v} D_{j_v})_Q) - 1) \max_Q \lambda_{(a_{i_v} D_{i_v} \cap a_{j_v} D_{j_v})_Q, v}(P) \right) \\
&\leq (3 + \varepsilon) h_D(P)
\end{aligned} \tag{5.6}$$

for all $P \in X(K) \setminus Z$. As there are only finitely many choices of i_v, j_v , we may find such a Z that works for all choices of $i_v, j_v, v \in S$ (with $i_v \neq j_v$).

Next we note that since D_1, D_2, D_3 intersect properly, we have $D_1 \cap D_2 \cap D_3 = \emptyset$, and by Theorem 2.1, for any $v \in S$,

$$\min\{\lambda_{a_1 D_1, v}(P), \lambda_{a_2 D_2, v}(P), \lambda_{a_3 D_3, v}(P)\} = O(1)$$

for all $P \in X(K)$. Let R be a set of $(D_1 + D_2 + D_3, S)$ -integral points. By definition and elementary properties of heights, for any $\varepsilon > 0$,

$$\begin{aligned}
\sum_{v \in S} (\lambda_{a_1 D_1, v}(P) + \lambda_{a_2 D_2, v}(P) + \lambda_{a_3 D_3, v}(P)) &= h_{a_1 D_1}(P) + h_{a_2 D_2}(P) + h_{a_3 D_3}(P) + O(1) \\
&\geq (3 - \varepsilon) h_D(P) + O(1),
\end{aligned}$$

where the $O(1)$ possibly depends on R (but not P).

Let $P \in R$. For $v \in S$, let $\{i_v, j_v, k_v\} = \{1, 2, 3\}$ be such that

$$\lambda_{a_{i_v} D_{i_v}, v}(P) \geq \lambda_{a_{j_v} D_{j_v}, v}(P) \geq \lambda_{a_{k_v} D_{k_v}, v}(P).$$

Then

$$\begin{aligned}
& \sum_{v \in S} (\lambda_{a_{i_v} D_{i_v}, v}(P) + \lambda_{a_{j_v} D_{j_v}, v}(P) + \lambda_{a_{k_v} D_{k_v}, v}(P)) \\
&= \sum_{v \in S} (\lambda_{a_{i_v} D_{i_v}, v}(P) + \lambda_{a_{j_v} D_{j_v}, v}(P)) + O(1) \\
&= \sum_{v \in S} (\lambda_{a_{i_v} D_{i_v}, v}(P) + \min\{\lambda_{a_{i_v} D_{i_v}, v}(P), \lambda_{a_{j_v} D_{j_v}, v}(P)\}) + O(1) \\
&= \sum_{v \in S} (\lambda_{a_{i_v} D_{i_v}, v}(P) + \lambda_{a_{i_v} D_{i_v} \cap a_{j_v} D_{j_v}, v}(P)) + O(1) \\
&\geq (3 - \varepsilon) h_D(P) + O(1).
\end{aligned}$$

Noting that if $\{i_0, j_0\} \neq \{i_v, j_v\}$ then

$$\lambda_{a_{i_0} D_{i_0} \cap a_{j_0} D_{j_0}, v}(P) = O(1)$$

and substituting into (5.6), we find that if $P \notin Z$, then

$$(3 - \varepsilon)h_D(P) + \gamma \sum_{v \in S} \lambda_{a_{i_0} D_{i_0} \cap a_{j_0} D_{j_0}, v}(P) \leq \sum_{v \in S} \lambda_{a_{i_v} D_{i_v}, v}(P) + (1 + \gamma_v) \lambda_{a_{i_v} D_{i_v} \cap a_{j_v} D_{j_v}, v}(P) + O(1) \\ < (3 + \varepsilon)h_D(P) + O(1).$$

Therefore, if $P \notin Z$,

$$\sum_{v \in S} \lambda_{a_{i_0} D_{i_0} \cap a_{j_0} D_{j_0}, v}(P) < \frac{2\varepsilon}{\gamma} h_D(P) + O(1).$$

Since $\min\{a_{i_0}, a_{j_0}\} \min\{\lambda_{D_{i_0}, v}, \lambda_{D_{j_0}, v}\} \leq \lambda_{a_{i_0} D_{i_0} \cap a_{j_0} D_{j_0}, v}$, we conclude that for any $\varepsilon > 0$, there exists a proper Zariski-closed subset Z of X such that for all $P \in R \setminus Z$,

$$\sum_{v \in S} \min(\lambda_{D_{i_0}, v}(P), \lambda_{D_{j_0}, v}(P)) \leq \varepsilon h_D(P) + O(1).$$

Since R is a set of $(D_1 + D_2 + D_3, S)$ -integral points, this is equivalent to

$$h_{D_{i_0} \cap D_{j_0}}(P) \leq \varepsilon h_D(P) + O(1)$$

for all $P \in R \setminus Z$. Finally, we note that we may remove the $O(1)$ in the inequality at the expense of excluding finitely many points of R , finishing the proof. \square

We now prove the corollary.

Proof of Corollary 5.4. The local intersection multiplicity and the local Hilbert-Samuel multiplicity coincide (see [Ful89, Ex. 2.4.8, Ex. 7.1.10]):

$$\mu_Q = \mu_Q((a_{i_0} D_{i_0} \cap a_{j_0} D_{j_0})_Q) = (a_{i_0} D_{i_0} \cdot a_{j_0} D_{j_0})_Q.$$

Since $(a_{i_0} D_{i_0} \cdot a_{j_0} D_{j_0}) = D^2$ is the sum of the local intersection multiplicities, and D_i and D_j intersect at more than one point, we must have

$$\mu_Q < D^2$$

for all $Q \in X(\overline{K})$.

Then by Lemma 3.11, for all $Q \in X(\overline{K})$,

$$\beta(D, (a_{i_0} D_{i_0} \cap a_{j_0} D_{j_0})_Q) \geq \frac{2}{3} \sqrt{\frac{D^2}{\mu_Q}} > \frac{2}{3},$$

and the desired result follows from Theorem 5.3. \square

We give an example to show that both the hypothesis (5.3) of Theorem 5.3 and the intersection condition of Corollary 5.4 are necessary.

Example 5.5. Let D_1, D_2, D_3 be the coordinate lines in \mathbb{P}^2 , let p, q be rational primes, and let $S = \{p, q, \infty\}$, a set of places of \mathbb{Q} . Then it follows from [Lev14, p. 707] that there exists a Zariski dense set of $(D_1 + D_2 + D_3, S)$ -integral points R in $\mathbb{P}^2(\mathbb{Q})$ such that

$$h_{D_i \cap D_j}(P) = \frac{1}{2} h(P) + O(1)$$

for all $P \in R$ and all $i, j \in \{1, 2, 3\}$, $i \neq j$. Note that in this case, if $i \neq j$, then $D_i \cap D_j$ consists of a single point and $\beta(\mathcal{O}(1), D_i \cap D_j) = \frac{2}{3}$.

5.2 On the Diophantine Equation $f(a^m, y) = b^n$

In this section, we provide an application of our result on greatest common divisors (Corollary 5.4) to study the exponential Diophantine equation $f(a^m, y) = b^n$, where $f(x, y)$ is a polynomial with rational coefficients and a and b are positive integers with a nontrivial common factor. Such an equation was studied by Corvaja and Zannier [CZ00], who noted that the equation did not seem to fall into prior treatments of Diophantine equations outside of very special situations (e.g., f is homogeneous (Thue-Mahler), or more generally f is homogeneous with respect to suitable weights). Corvaja and Zannier proved the following result:

Theorem 5.6 (Corvaja-Zannier [CZ00]). *Let $f(x, y) = a_0(x)y^d + a_1(x)y^{d-1} + \dots + a_d(x)$ be a polynomial with rational coefficients, of degree $d \geq 2$ in y ; let $a, b > 1$ be integers. Suppose that*

1. a_0 is constant,
2. the polynomial $f(0, y)$ has no repeated roots,
3. a and b are not relatively prime.

If the equation

$$f(a^m, y) = b^n$$

has an infinite sequence of solutions $(m, n, y) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{Z}$, such that $\min\{|m|, |n|\} \rightarrow \infty$, then there exist an integer $h \neq 0$ and a polynomial $p(x) \in \mathbb{Q}[x]$ such that $f(x^h, p(x))$ has only one term; furthermore, a and b are multiplicatively dependent.

Using Corollary 5.4, we derive a result along the lines of Theorem 5.6. For purposes of comparison, we note that our result is weaker in some respects (condition 1' below is much stronger than condition 1 in Theorem 5.6), but under our hypotheses we partially weaken condition 2 of Theorem 5.6. Possibly, condition 2 could also be weakened via the approach of Corvaja-Zannier in [CZ00]; our main purpose here is to illustrate how our general Diophantine approximation results may be used to shed new light on existing problems.

Theorem 5.7. *Let $F(X, Y, Z) \in \mathbb{Q}[X, Y, Z]$ be a homogeneous polynomial of degree $d \geq 2$ and let $a, b > 1$ be integers. Suppose that*

- 1'. $F(0, 1, 0) \neq 0$
- 2'. Neither $F(0, y, z)$ nor $F(x, y, 0)$ are powers of a linear form in $\overline{\mathbb{Q}}[x, y, z]$.
- 3'. a and b are not relatively prime.

Let $f(x, y) = F(x, y, 1)$ Then the set of points

$$\{(a^m, y) \in \mathbb{Z} \times \mathbb{Z} \mid f(a^m, y) = b^n, m, n \in \mathbb{Z}, m, n \geq 0\},$$

is not Zariski dense in \mathbb{A}^2 .

From the non-Zariski density statement one can derive a conclusion as in Theorem 5.6 (e.g., using the Lemma of [CZ00] and Siegel's theorem on integral points on curves); we leave the details to the interested reader.

Proof. Let V_i be the hypersurface of \mathbb{P}^3 defined by $b^i W^d = F(X, Y, Z)$, $i \in \{0, \dots, d-1\}$. If $f(a^m, y) = b^n$, for some $m, n, y \in \mathbb{Z}$, then writing $b^n = (b^{n'})^d b^i$ for some $n' \in \mathbb{Z}$ and $i \in \{0, \dots, d-1\}$, we have $[W : X : Y : Z] = [b^{n'} : a^m : y : 1] \in V_i(\mathbb{Q})$. Then after fixing i and letting $V = V_i$, it suffices to show that the set of points

$$R := \{[b^n : a^m : y : 1] \in V(\mathbb{Q}) \mid m, n, y \in \mathbb{Z}, m, n \geq 0\}.$$

is not Zariski dense in V (the statement in the theorem follows after taking an appropriate projection).

Let D_1, D_2, D_3 be the divisors on V defined by $W = 0$, $X = 0$, and $Z = 0$, respectively. Then clearly D_1, D_2 , and D_3 are linearly equivalent ample effective Cartier divisors; let D be any divisor in the same linear equivalence class. Since $F(0, 1, 0) \neq 0$, V and the hyperplanes defined by $W = 0$, $X = 0$, and $Z = 0$ are in general position on \mathbb{P}^3 . Then by Remark 2.13, D_1, D_2, D_3 intersect properly on V . The condition (2) implies that $\text{Supp}(D_1 \cap D_2)(\overline{\mathbb{Q}})$ and $\text{Supp}(D_1 \cap D_3)(\overline{\mathbb{Q}})$ both contain more than one point, while the conditions $F(0, 1, 0) \neq 0$ and $d \geq 2$ imply that $\text{Supp}(D_2 \cap D_3)(\overline{\mathbb{Q}})$ contains more than one point. Moreover, if S is the set of places of \mathbb{Q} given by

$$S = \{p \text{ prime} \mid p \mid ab\} \cup \{\infty\},$$

then R is a set of $(D_1 + D_2 + D_3, S)$ -integral points in $V(\mathbb{Q})$. By Corollary 5.4, there exists a proper Zariski-closed subset $Z \subset V$ such that if $i \neq j$ and $v \in S$, then

$$\min\{\lambda_{D_i, v}(P), \lambda_{D_j, v}(P)\} \leq \varepsilon h_D(P), \quad (5.7)$$

for all but finitely many $P \in R \setminus Z$.

Let $P = [b^n : a^m : y : 1] \in R$. Let $h = h(P) = \log \max\{|b^n|, |a^m|, |y|\} = h_D(P) + O(1)$ and let p be a prime dividing $\gcd(a, b)$. Then we compute

Divisor D_i	$\lambda_{D_i, \infty}(P)$	$\lambda_{D_i, p}(P)$
D_1	$h - n \log b$	$n(\log p)(\text{ord}_p b)$
D_2	$h - m \log a$	$m(\log p)(\text{ord}_p a)$
D_3	h	0

Then using (5.7) with $(i, j, v) = (1, 3, \infty), (2, 3, \infty), (1, 2, p)$, respectively, we find that for all $P \in R \setminus Z$,

$$\begin{aligned} n &\geq \frac{1}{\log b}(1 - \varepsilon)h_D(P) + O(1) \\ m &\geq \frac{1}{\log a}(1 - \varepsilon)h_D(P) + O(1) \\ \min\{m, n\} &\leq 2\varepsilon h_D(P) + O(1). \end{aligned}$$

Taking $0 < \varepsilon < \frac{1}{2 \max\{\log a, \log b\} + 1}$, the inequalities imply that $h_D(P)$ is bounded for $P \in R \setminus Z$. Since D is ample, we conclude that $R \setminus Z$ is a finite set and R is not Zariski dense in V . \square

5.3 Integral Points on the Complement of Three Numerically Parallel Curves Passing Through a Point

From the work of Corvaja-Zannier [CZ04b] and the second author [Lev09], it is known that the complement of any 4 ample divisors in general position on a projective surface does not contain a Zariski dense set of integral points; the number 4 here is sharp as $\mathbb{G}_m^2 \cong \mathbb{P}^2 \setminus \{xyz = 0\}$ and $\mathbb{G}_m^2(\mathcal{O}_{K,S}) \cong (\mathcal{O}_{K,S}^*)^2$ is Zariski dense in \mathbb{G}_m^2 as long as $|S| > 1$. As remarked after Theorem 1.5, it is already an open problem to prove the degeneracy of integral points on the complement of three plane curves forming a normal crossings divisor of degree at least 4. In contrast to this, it is sometimes possible to handle certain degenerate (i.e., non-normal crossings) configurations of three plane curves. For instance, the problem of integral points on the complement of a conic and two (distinct) tangent lines is easily reduced to Siegel's theorem for integral points on \mathbb{P}^1 . A deeper result, depending ultimately on the Subspace Theorem, is the following theorem of Corvaja and Zannier [CZ06].

Theorem 5.8 (Corvaja-Zannier). *Let D_1, D_2, D_3 be distinct, effective, irreducible, numerically equivalent divisors on a nonsingular projective surface X defined over a number field K , such that*

- (a) $D_1 \cap D_2 \cap D_3$ consists of a single point, at which the D_i intersect transversally.
- (b) $D_i \cdot D_j > 1$ for some i, j .

Let S be a finite set of places of K containing the archimedean places. Then no set of $(D_1 + D_2 + D_3, S)$ -integral points in $X(K)$ is Zariski-dense in X .

We prove a generalization of Theorem 5.8 where we greatly weaken the triple intersection condition, and we only require the three divisors to be numerically *parallel* rather than numerically equivalent (to be precise, we don't recover the case $D_i \cdot D_j = 2$ of Theorem 5.8; however, in our formulation, which allows more than one point in $D_1 \cap D_2 \cap D_3$, excluding this case is necessary by Example 5.11).

Theorem 5.9. *Let X be a projective surface over a number field K , and let D_1, D_2, D_3 be effective Cartier divisors on X pairwise intersecting properly. Suppose that there exist positive integers a_1, a_2, a_3 such that $a_1 D_1, a_2 D_2, a_3 D_3$ are all numerically equivalent to an ample divisor D , and that*

$$D_1 \cap D_2 \cap D_3 \neq \emptyset.$$

Let

$$\beta_0 = \min_{\substack{i \neq j \\ Q \in (D_1 \cap D_2 \cap D_3)(\overline{K})}} \beta(D, (a_i D_i \cap a_j D_j)_Q).$$

Furthermore, suppose that for every point $Q \in (D_1 \cap D_2 \cap D_3)(\overline{K})$ and every permutation i, j, k of the indices 1, 2, 3, we have

$$(\beta(D, (a_i D_i \cap a_j D_j)_Q) - 1) + (\beta(D, (a_i D_i \cap a_k D_k)_Q) - 1)(3\beta_0 - 2) > 0. \quad (5.8)$$

In particular, $\beta_0 \geq \frac{2}{3}$ and (5.8) holds if

$$\beta(D, (a_i D_i \cap a_j D_j)_Q) > 1 \quad (5.9)$$

or

$$(D_i \cdot D_j)_Q < \frac{4}{9}(D_i \cdot D_j) \quad (5.10)$$

for all $Q \in (D_1 \cap D_2 \cap D_3)(\overline{K})$ and all $i \neq j$ (where $(D_i \cdot D_j)_Q$ denotes the local intersection multiplicity of D_i and D_j at Q , and $(D_i \cdot D_j)$ the intersection multiplicity). Let S be a finite set of places of K containing all the archimedean places. Then there exists a proper Zariski-closed subset $Z \subset X$ such that for any set $R \subset X(K)$ of $(D_1 + D_2 + D_3, S)$ -integral points, the set $R \setminus Z$ is finite.

Theorem 1.5 from the introduction follows immediately (using (5.10)).

Proof. After replacing K by a finite extension, we can assume that every point in the support of $D_i \cap D_j$, $i \neq j$, is K -rational.

We first show that there exists a proper Zariski-closed subset $Z \subset X$ such that for any set $R \subset X(K)$ of $(D_1 + D_2 + D_3, S)$ -integral points, we have

$$\begin{aligned} h_{a_1 D_1 \cap a_2 D_2 \cap a_3 D_3}(P) &= \sum_{v \in S} \lambda_{a_1 D_1 \cap a_2 D_2 \cap a_3 D_3, v}(P) + O(1) \\ &= \sum_{v \in S} \min(\lambda_{a_1 D_1, v}(P), \lambda_{a_2 D_2, v}(P), \lambda_{a_3 D_3, v}(P)) + O(1) \\ &\leq \varepsilon h_D(P) + O(1) \end{aligned}$$

for all $P \in R \setminus Z$.

By definition and elementary properties of heights, for any $\varepsilon > 0$,

$$\begin{aligned} \sum_{v \in S} (\lambda_{a_1 D_1, v}(P) + \lambda_{a_2 D_2, v}(P) + \lambda_{a_3 D_3, v}(P)) &= h_{a_1 D_1}(P) + h_{a_2 D_2}(P) + h_{a_3 D_3}(P) + O(1) \\ &\geq (3 - \varepsilon)h_D(P) + O(1), \end{aligned}$$

for all $P \in R$, where the $O(1)$ possibly depends on R (but not P).

Let $P \in R$. For $v \in S$, let $\{i_v, j_v, k_v\} = \{1, 2, 3\}$ be such that

$$\lambda_{a_{i_v} D_{i_v}, v}(P) \geq \lambda_{a_{j_v} D_{j_v}, v}(P) \geq \lambda_{a_{k_v} D_{k_v}, v}(P).$$

For each $v \in S$, there exists a point $Q_v \in \text{Supp}(a_{i_v} D_{i_v} \cap a_{j_v} D_{j_v})$ (depending on P) such that

$$\lambda_{a_{i_v} D_{i_v} \cap a_{j_v} D_{j_v}, v}(P) = \lambda_{(a_{i_v} D_{i_v} \cap a_{j_v} D_{j_v})_{Q_v}, v}(P) + O(1),$$

where the constant in the $O(1)$ is independent of P .

If $Q_v \notin \text{Supp } D_1 \cap D_2 \cap D_3$, then

$$\lambda_{a_1 D_1 \cap a_2 D_2 \cap a_3 D_3, v}(P) = \min\{\lambda_{a_{k_v} D_{k_v}, v}(P), \lambda_{a_{i_v} D_{i_v} \cap a_{j_v} D_{j_v}, v}(P)\} + O(1)$$

$$\begin{aligned}
&= \min\{\lambda_{(a_{i_v}D_{i_v} \cap a_{j_v}D_{j_v})_{Q_v}, v}(P), \lambda_{a_{k_v}D_{k_v}, v}(P)\} + O(1) \\
&= \lambda_{(a_{i_v}D_{i_v} \cap a_{j_v}D_{j_v})_{Q_v} \cap (a_{k_v}D_{k_v}), v} \\
&= O(1)
\end{aligned}$$

since $(a_{i_v}D_{i_v} \cap a_{j_v}D_{j_v})_{Q_v} \cap (a_{k_v}D_{k_v})$ is empty. When $Q_v \in \text{Supp } D_1 \cap D_2 \cap D_3$, we use the estimate

$$\lambda_{a_{k_v}D_{k_v}, v}(P) = \lambda_{(a_1D_1 \cap a_2D_2 \cap a_3D_3)_{Q_v}, v}(P) + O(1) \leq \lambda_{(a_{i_v}D_{i_v} \cap a_{k_v}D_{k_v})_{Q_v}, v}(P) + O(1).$$

Let

$$\begin{aligned}
S' &= \{v \in S \mid Q_v \notin \text{Supp } D_1 \cap D_2 \cap D_3\}, \\
S'' &= S \setminus S'.
\end{aligned}$$

It follows that, up to $O(1)$,

$$\begin{aligned}
(3 - \varepsilon)h_D(P) &\leq \sum_{v \in S} (\lambda_{a_{i_v}D_{i_v}, v}(P) + \lambda_{a_{j_v}D_{j_v}, v}(P) + \lambda_{a_{k_v}D_{k_v}, v}(P)) \\
&\leq \sum_{v \in S'} (\lambda_{a_{i_v}D_{i_v}, v}(P) + \lambda_{(a_{i_v}D_{i_v} \cap a_{j_v}D_{j_v})_{Q_v}, v}(P)) \\
&\quad + \sum_{v \in S''} (\lambda_{a_{i_v}D_{i_v}, v}(P) + \lambda_{(a_{i_v}D_{i_v} \cap a_{j_v}D_{j_v})_{Q_v}, v}(P) + \lambda_{(a_{i_v}D_{i_v} \cap a_{k_v}D_{k_v})_{Q_v}, v}(P)).
\end{aligned}$$

Since $a_iD_i \equiv D$, we have $\beta(D, a_iD_i) = \frac{1}{3}$ for all i , and by Lemma 3.3, for all $Q \in X(\overline{K})$ and $i \neq j$, we have

$$\beta(D, (a_iD_i \cap a_jD_j)_Q) \geq \beta(D, a_iD_i \cap a_jD_j) \geq \beta(D, a_iD_i) + \beta(D, a_jD_j) \geq \frac{2}{3}.$$

Therefore $\beta_0 \geq \frac{2}{3}$.

By Theorem 1.4, there exists a proper Zariski-closed subset $Z \subset X$ such that $P \notin Z$,

$$\sum_{v \in S} \lambda_{a_{i_v}D_{i_v}, v}(P) + (3\beta(D, (a_{i_v}D_{i_v} \cap a_{j_v}D_{j_v})_{Q_v}) - 1)\lambda_{(a_{i_v}D_{i_v} \cap a_{j_v}D_{j_v})_{Q_v}, v}(P) \leq (3 + \varepsilon)h_D(P), \quad (5.11)$$

and

$$\begin{aligned}
&\sum_{v \in S'} \lambda_{a_{i_v}D_{i_v}, v}(P) + (3\beta(D, (a_{i_v}D_{i_v} \cap a_{j_v}D_{j_v})_{Q_v}) - 1)\lambda_{(a_{i_v}D_{i_v} \cap a_{j_v}D_{j_v})_{Q_v}, v}(P) \\
&+ \sum_{v \in S''} \lambda_{a_{i_v}D_{i_v}, v}(P) + (3\beta(D, (a_{i_v}D_{i_v} \cap a_{k_v}D_{k_v})_{Q_v}) - 1)\lambda_{(a_{i_v}D_{i_v} \cap a_{k_v}D_{k_v})_{Q_v}, v}(P) \quad (5.12) \\
&\leq (3 + \varepsilon)h_D(P).
\end{aligned}$$

Let $b = \frac{1}{3\beta_0 - 1}$. We have $0 < b \leq 1$ and by definition of β_0 , for any $i, j \in \{1, 2, 3\}, i \neq j$, and $Q \in (D_1 \cap D_2 \cap D_3)(\overline{K})$,

$$b \left(3\beta(D, (a_iD_i \cap a_jD_j)_Q) - 1 \right) \geq 1.$$

Note also that

$$\lambda_{(a_{i_v} D_{i_v} \cap a_{j_v} D_{j_v})_{Q_v}, v}(P) \geq \lambda_{(a_{i_v} D_{i_v} \cap a_{k_v} D_{k_v})_{Q_v}, v}(P) + O(1).$$

Let

$$\gamma = 3 \min_{\substack{\{i,j,k\}=\{1,2,3\} \\ Q \in (D_1 \cap D_2 \cap D_3)(\overline{K})}} \frac{(\beta(D, (a_i D_i \cap a_j D_j)_Q) - 1) + (\beta(D, (a_i D_i \cap a_k D_k)_Q) - 1)(3\beta_0 - 2)}{3\beta_0 - 1}.$$

By hypothesis, $\gamma > 0$. Fixing $Q \in (D_1 \cap D_2 \cap D_3)(\overline{K})$, P , and v , for $i, j \in \{1, 2, 3\}$ let

$$\begin{aligned} \beta_{ij} &= \beta(D, (a_i D_i \cap a_j D_j)_Q), \\ \lambda_{ij} &= \lambda_{(a_i D_i \cap a_j D_j)_Q, v}(P). \end{aligned}$$

Then assuming $\lambda_{ij} \geq \lambda_{ik}$, we compute

$$\begin{aligned} b(3\beta_{ij} - 1)\lambda_{ij} + (1 - b)(3\beta_{ik} - 1)\lambda_{ik} &\geq \lambda_{ij} + (b(3\beta_{ij} - 1) - 1 + (1 - b)(3\beta_{ik} - 1))\lambda_{ik} \\ &\geq \lambda_{ij} + \left(1 + \frac{3((\beta_{ij} - 1) + (\beta_{ik} - 1)(3\beta_0 - 2))}{3\beta_0 - 1}\right) \lambda_{ik} \\ &\geq \lambda_{ij} + (1 + \gamma)\lambda_{ik}. \end{aligned}$$

Multiplying (5.11) by b , (5.12) by $1 - b$, and summing and using the above calculation, we obtain

$$\begin{aligned} &\sum_{v \in S'} \lambda_{a_{i_v} D_{i_v}, v}(P) + (3\beta(D, (a_{i_v} D_{i_v} \cap a_{j_v} D_{j_v})_{Q_v}) - 1)\lambda_{(a_{i_v} D_{i_v} \cap a_{j_v} D_{j_v})_{Q_v}, v}(P) \\ &+ \sum_{v \in S''} \lambda_{a_{i_v} D_{i_v}, v}(P) + \lambda_{(a_{i_v} D_{i_v} \cap a_{j_v} D_{j_v})_{Q_v}, v}(P) + (1 + \gamma)\lambda_{(a_{i_v} D_{i_v} \cap a_{k_v} D_{k_v})_{Q_v}, v}(P) \\ &\leq (3 + \varepsilon)h_D(P) + O(1). \end{aligned}$$

Then there exists a proper Zariski-closed subset $Z \subset X$ such that if $P \notin Z$, up to $O(1)$,

$$\begin{aligned} &(3 - \varepsilon)h_D(P) + \gamma \sum_{v \in S} \lambda_{a_1 D_1 \cap a_2 D_2 \cap a_3 D_3, v}(P) \\ &\leq (3 - \varepsilon)h_D(P) + \gamma \sum_{v \in S''} \lambda_{(a_{i_v} D_{i_v} \cap a_{k_v} D_{k_v})_{Q_v}, v}(P) \\ &\leq \sum_{v \in S'} (\lambda_{a_{i_v} D_{i_v}, v}(P) + \lambda_{(a_{i_v} D_{i_v} \cap a_{j_v} D_{j_v})_{Q_v}, v}(P)) \\ &\quad + \sum_{v \in S''} (\lambda_{a_{i_v} D_{i_v}, v}(P) + \lambda_{(a_{i_v} D_{i_v} \cap a_{j_v} D_{j_v})_{Q_v}, v}(P) + (1 + \gamma)\lambda_{(a_{i_v} D_{i_v} \cap a_{k_v} D_{k_v})_{Q_v}, v}(P)) \\ &\leq (3 + \varepsilon)h_D(P). \end{aligned}$$

Therefore, if $P \in R \setminus Z$,

$$\sum_{v \in S} \lambda_{a_1 D_1 \cap a_2 D_2 \cap a_3 D_3, v}(P) < \frac{2\varepsilon}{\gamma} h_D(P) + O(1).$$

Equivalently, since R is a set of $(D_1 + D_2 + D_3, S)$ -integral points and $\text{Supp}(a_1 D_1 \cap a_2 D_2 \cap a_3 D_3) \subset \text{Supp}(D_1 + D_2 + D_3)$, for given $\varepsilon > 0$, there exists a proper Zariski-closed subset $Z \subset X$ such that for $P \in R \setminus Z$,

$$h_{a_1 D_1 \cap a_2 D_2 \cap a_3 D_3}(P) < \varepsilon h_D(P) + O(1), \quad (5.13)$$

finishing the proof of the claim.

Let Q be some point in $\text{Supp } D_1 \cap D_2 \cap D_3$ (which is nonempty by assumption). Let $\pi : \tilde{X} \rightarrow X$ be the blowup at Q , with exceptional divisor E . If R is a set of $(D_1 + D_2 + D_3, S)$ -integral points in $X(K)$, then $\pi^{-1}(R) \setminus E$ is a set of $(\pi^* D_1 + \pi^* D_2 + \pi^* D_3, S)$ -integral points in $\tilde{X}(K)$. So it suffices to show that there exists a proper Zariski-closed subset \tilde{Z} of \tilde{X} such that for any set \tilde{R} of $(\pi^* D_1 + \pi^* D_2 + \pi^* D_3, S)$ -integral points in $\tilde{X}(K)$, the set $\tilde{R} \setminus \tilde{Z}$ is finite. Define the effective Cartier divisors

$$D'_i = a_i \pi^* D_i - E, \quad i = 1, 2, 3.$$

Let \tilde{R} be a set of $(\pi^* D_1 + \pi^* D_2 + \pi^* D_3, S)$ -integral points in $\tilde{X}(K)$ (and hence a set of $(D'_1 + D'_2 + D'_3, S)$ -integral points). For $P \in \tilde{R}$ and $\varepsilon > 0$, we have

$$\begin{aligned} \sum_{v \in S} (\lambda_{D'_1, v}(P) + \lambda_{D'_2, v}(P) + \lambda_{D'_3, v}(P)) &= h_{D'_1}(P) + h_{D'_2}(P) + h_{D'_3}(P) + O(1) \quad (5.14) \\ &\geq (3 - \varepsilon) h_{\pi^* D}(P) - (3 - \varepsilon) h_E(P) + O(1), \end{aligned}$$

where the $O(1)$ possibly depends on \tilde{R} (but not P).

We now bound the left-hand side of the above equation. As in previous arguments, it suffices to bound a sum of the form

$$\sum_{v \in S} \left(\lambda_{D'_{i_v}, v}(P) + \lambda_{D'_{i_v} \cap D'_{j_v}, v}(P) + \lambda_{D'_{i_v} \cap D'_{j_v} \cap D'_{k_v}, v}(P) \right),$$

where $\{i_v, j_v, k_v\} = \{1, 2, 3\}$ for $v \in S$. We first note that it follows from (5.13) and functoriality that given $\varepsilon > 0$, there exists a proper Zariski-closed subset $\tilde{Z} \subset \tilde{X}$ such that

$$\sum_{v \in S} \lambda_{D'_{i_v} \cap D'_{j_v} \cap D'_{k_v}, v}(P) < \varepsilon h_{\pi^* D}(P) + O(1)$$

for all $P \in \tilde{R} \setminus \tilde{Z}$. For the same reasons, we may choose \tilde{Z} so that we also have

$$h_E(P) < \varepsilon h_{\pi^* D}(P) + O(1) \quad (5.15)$$

for all $P \in \tilde{R} \setminus \tilde{Z}$. We can write (as closed subschemes)

$$D'_{i_v} \cap D'_{j_v} = Y_{0, v} + Y_{1, v},$$

where $\text{Supp } \pi(Y_{1, v}) = Q$, $\dim Y_{0, v} = 0$, and $Y_{0, v} \cap E = \emptyset$. We have (for an appropriate \tilde{Z})

$$\sum_{v \in S} \lambda_{Y_{1, v}, v}(P) < \varepsilon h_{\pi^* D}(P) + O(1)$$

for all $P \in \tilde{R} \setminus \tilde{Z}$, and so

$$\lambda_{D'_{i_v} \cap D'_{j_v}, v}(P) \leq \lambda_{Y_{0,v}, v}(P) + \varepsilon h_{\pi^* D}(P) + O(1)$$

for all $P \in \tilde{R} \setminus \tilde{Z}$.

Let $\delta \in \mathbb{Q}, \delta > 0$, be chosen as in Lemma 3.14 so that (3.4) holds, and let

$$\gamma' = \beta(\pi^* D - \delta E, \pi^* D - E) - \frac{1}{3} > 0.$$

Note that

$$\beta(\pi^* D - \delta E, Y_{0,v}) \geq \beta(\pi^* D - \delta E, D'_{i_v}) + \beta(\pi^* D - \delta E, D'_{j_v}).$$

This does not follow directly from Lemma 3.3 (since D'_{i_v} and D'_{j_v} may not intersect properly above Q , along the component $Y_{1,v}$), but it follows from a slight modification to the proof of that lemma as D'_i and D'_j intersect properly in a neighborhood of the zero-dimensional closed subscheme $Y_{0,v}$.

Using Theorem 1.4, for any $\varepsilon > 0$ we find that for $P \in \tilde{R}$ outside a proper Zariski-closed subset of \tilde{X} (and up to $O(1)$)

$$\begin{aligned} & \left(\frac{1}{3} + \gamma' \right) \sum_{v \in S} \left(\lambda_{D'_{i_v}, v}(P) + \lambda_{D'_{i_v} \cap D'_{j_v}, v}(P) + \lambda_{D'_{i_v} \cap D'_{j_v} \cap D'_{k_v}, v}(P) \right) \\ & \leq \sum_{v \in S} \left(\beta(\pi^* D - \delta E, D'_{i_v}) \lambda_{D'_{i_v}, v}(P) + (\beta(\pi^* D - \delta E, Y_{0,v}) - \beta(\pi^* D - \delta E, D'_{i_v})) \lambda_{Y_{0,v}, v}(P) \right) \\ & \quad + \varepsilon h_{\pi^* D}(P) \\ & \leq (1 + \varepsilon) h_{\pi^* D - \delta E}(P) + \varepsilon h_{\pi^* D}(P) \\ & \leq (1 + 2\varepsilon) h_{\pi^* D}(P). \end{aligned}$$

Therefore, for some positive $\delta' > 0$, for $P \in \tilde{R}$ outside a proper Zariski-closed subset of \tilde{X} we have

$$\sum_{v \in S} (\lambda_{D'_1, v}(P) + \lambda_{D'_2, v}(P) + \lambda_{D'_3, v}(P)) \leq (3 - \delta') h_{\pi^* D}(P).$$

On the other hand, by (5.14) and (5.15) (taking ε sufficiently small), for $P \in \tilde{R}$ outside a proper Zariski-closed subset of \tilde{X} ,

$$\sum_{v \in S} (\lambda_{D'_1, v}(P) + \lambda_{D'_2, v}(P) + \lambda_{D'_3, v}(P)) \geq \left(3 - \frac{\delta'}{2} \right) h_{\pi^* D}(P).$$

Finally, since $\pi^* D$ is big, combining the above inequalities with an application of Northcott's theorem (for big divisors) gives that there exists a proper Zariski-closed subset \tilde{Z} of \tilde{X} such that $\tilde{R} \setminus \tilde{Z}$ is finite. \square

We now give two examples addressing the sharpness of conditions (5.9) and (5.10). Both examples are based on the following construction:

Example 5.10. Let K be a number field and let S be a finite set of places of K containing the archimedean places with $|S| \geq 2$. Let Λ be a linear pencil of curves in a projective surface X such that the general member of Λ is isomorphic to \mathbb{P}^1 , and the pencil Λ has exactly two base points Q_1, Q_2 , which, after possibly enlarging K , we may assume are K -rational (our construction would also work if there is a single base point). Let $D_1, D_2, D_3 \in \Lambda$ be distinct elements. Then we claim that the conclusion of Theorem 5.9 does not hold, i.e., there does not exist a proper Zariski-closed subset $Z \subset X$ such that for any set $R \subset X(K)$ of $(D_1 + D_2 + D_3, S)$ -integral points, the set $R \setminus Z$ is finite. Let $C \in \Lambda \setminus \{D_1, D_2, D_3\}$ be a general member. We claim that C contains an infinite set of $(D_1 + D_2 + D_3, S)$ -integral points. Indeed, any two elements of Λ intersect only at the points Q_1 and Q_2 , and therefore $C \cap (D_1 \cup D_2 \cup D_3) = \{Q_1, Q_2\}$. Since $C \cong \mathbb{P}^1$, we have $C \setminus (D_1 \cup D_2 \cup D_3) \cong \mathbb{G}_m$, and C will contain an infinite set R of $(D_1 + D_2 + D_3, S)$ -integral points (we use $|S| \geq 2$ here, so that $\mathcal{O}_{K,S}^*$ is infinite). Since the union of such elements $C \in \Lambda$ is Zariski dense in X , the conclusion of Theorem 5.9 does not hold.¹

We first give an example showing that the factor $\frac{4}{9}$ in the intersection condition (1.4) (or (5.10)) cannot be replaced by anything larger than $\frac{1}{2}$.

Example 5.11. Consider the pencil of plane conics $\Lambda = \{C_\lambda \mid \lambda \in K\}$, where $C_\lambda = \{y^2 - \lambda xz = 0\}$. Then any two distinct curves $C_\lambda, C'_\lambda \in \Lambda$ intersect precisely at the two points $Q_1 = [0 : 0 : 1], Q_2 = [1 : 0 : 0]$, each with multiplicity 2. By Example 5.10, if $D_1, D_2, D_3 \in \Lambda$ are distinct elements, then the conclusion of Theorem 5.9 does not hold (for appropriate K and S), and we note that

$$(D_i \cdot D_j)_{Q_1} = (D_i \cdot D_j)_{Q_2} = 2 = \frac{1}{2}(D_i \cdot D_j).$$

Somewhat surprisingly, the next example shows that the beta constant condition in Theorem 5.9 is sharp, in the sense that the condition

$$\beta(D, (a_i D_i \cap a_j D_j)_Q) > 1$$

cannot be replaced by the inequality

$$\beta(D, (a_i D_i \cap a_j D_j)_Q) \geq 1.$$

Example 5.12. Let D_1 and D_2 be two distinct irreducible curves of type $(1, 1)$ on $\mathbb{P}^1 \times \mathbb{P}^1$, defined over a number field K , intersecting in two distinct K -rational points Q_1 and Q_2 . Let Λ be the pencil of curves containing D_1 and D_2 , and let $D_3 \in \Lambda \setminus \{D_1, D_2\}$ be another irreducible element of the pencil.

For any point $Q \in (\mathbb{P}^1 \times \mathbb{P}^1)(K)$ (viewed as a closed subscheme with the reduced induced structure) an elementary computation gives

$$\beta(D_1, Q) = 1.$$

Since $D_i \cap D_j = Q_1 + Q_2$ (as closed subschemes) when $i \neq j$, we see from Example 5.10 (with this pencil) that the conclusion of Theorem 5.9 does not hold in general when (5.8) is replaced by $\beta(D, (a_i D_i \cap a_j D_j)_Q) \geq 1$.

¹We do not assert that there exists a Zariski dense set of $(D_1 + D_2 + D_3, S)$ -integral points; in fact, the pencil yields a morphism $X \setminus \{D_1 \cup D_2 \cup D_3\} \rightarrow \mathbb{P}^1 \setminus \{0, 1, \infty\}$ and Siegel's theorem shows that there is no Zariski dense set of $(D_1 + D_2 + D_3, S)$ -integral points in $X(K)$.

5.4 Families of Unit Equations

The unit equation theorem, proved by Siegel (when S is the set of archimedean places) and Mahler, is a fundamental and ubiquitous result in number theory:

Theorem 5.13 (Siegel-Mahler). *Let K be a number field and let S be a finite set of places of K containing the archimedean places. Let $\mathcal{O}_{K,S}$ be the ring of S -integers of K and let $\mathcal{O}_{K,S}^*$ be the group of S -units of K . Let $a, b, c \in K^*$. The S -unit equation*

$$au + bv = c, \quad u, v \in \mathcal{O}_{K,S}^*,$$

has only finitely many solutions.

From another viewpoint, the theorem is equivalent to Siegel's theorem on integral points on the affine curve $\mathbb{P}^1 \setminus \{0, 1, \infty\}$. In this section, we study the one-parameter family of unit equations

$$f_1(t)u + f_2(t)v = f_3(t), \quad t \in \mathcal{O}_{K,S}, u, v \in \mathcal{O}_{K,S}^*,$$

where f_1, f_2, f_3 are polynomials over a number field K . This equation was treated in the case $\deg f_1 = \deg f_2 = \deg f_3$ by Corvaja and Zannier [CZ06, CZ10], and in the case $\deg f_1 + \deg f_2 = \deg f_3$ by the second author [Lev06]. Applying the results of the previous section to certain surfaces, we handle a wide range of new values of the triple $(d_1, d_2, d_3) = (\deg f_1, \deg f_2, \deg f_3)$.

Theorem 5.14. *Let $f_1, f_2, f_3 \in K[t]$ be nonconstant polynomials without a common zero of degrees d_1, d_2, d_3 , respectively. Let $\{i_1, i_2, i_3\} = \{1, 2, 3\}$ be such that $d_{i_1} \geq d_{i_2} \geq d_{i_3}$ and suppose that*

$$\left(\sqrt{\frac{d_{i_1} + 1}{d_{i_1} + 1 - d_{i_2}}} - 1 \right) \left(\sqrt{\frac{d_{i_1} + 1}{d_{i_1} + 1 - d_{i_3}}} - 1 \right) > \frac{1}{4}. \quad (5.16)$$

In particular, (5.16) holds when

$$\max_i (d_i + 1) < \frac{9}{5} \min_i d_i.$$

Then the set of solutions $(t, u, v) \in \mathbb{A}^3(K)$ of the equation

$$f_1(t)u + f_2(t)v = f_3(t), \quad t \in \mathcal{O}_{K,S}, u, v \in \mathcal{O}_{K,S}^*, \quad (5.17)$$

is contained in a finite number of rational curves in \mathbb{A}^3 .

We note that if $d_{i_1} > d_{i_2}$ in Theorem 5.14, then by [Lev06, Lemma 5], one may drop the integrality condition on t in (5.17) (i.e., consider solutions with $t \in K$).

Proof. We first remark that the statement of the theorem is independent of the ordering of f_1, f_2, f_3 ; this follows from noting, for example, that if $(t_0, u_0, v_0) \in \mathcal{O}_{K,S} \times \mathcal{O}_{K,S}^* \times \mathcal{O}_{K,S}^*$ is a solution to $f_1(t)u + f_3(t)v = f_2(t)$, then $(t_0, -u_0/v_0, 1/v_0) \in \mathcal{O}_{K,S} \times \mathcal{O}_{K,S}^* \times \mathcal{O}_{K,S}^*$

is a solution to $f_1(t)u + f_2(t)v = f_3(t)$, and this relation corresponds to a birational automorphism of \mathbb{A}^3 . Thus, after permuting the f_i , we may assume that $d_1 \geq d_2 \geq d_3 > 0$, $d_{i_j} = d_j$, $j = 1, 2, 3$, and

$$\left(\sqrt{\frac{d_1 + 1}{d_1 + 1 - d_2}} - 1 \right) \left(\sqrt{\frac{d_1 + 1}{d_1 + 1 - d_3}} - 1 \right) > \frac{1}{4}.$$

In place of (5.17), we will actually study the slightly modified equation

$$f_1(t)u + f_2(t)v^{d_1+1-d_2} = f_3(t), \quad t \in \mathcal{O}_{K,S}, u, v \in \mathcal{O}_{K,S}^*. \quad (5.18)$$

Since $\mathcal{O}_{K,S}^*$ is finitely generated, we can find a number field L and a finite set of places S' of L such that every element of $\mathcal{O}_{K,S}^*$ has a $(d_1 + 1 - d_2)$ -th root in $\mathcal{O}_{L,S'}^*$. Then (5.17) reduces to studying the equation (5.18) (with (K, S) replaced by (L, S')).

We let $F_i \in K[x_0, x_1, x_2, x_3]$ be the homogeneous polynomial $F_i = f_i(x_0/x_3)x_3^{d_i}$, $i = 1, 2, 3$, and let X be the hypersurface in \mathbb{P}^3 defined by the equation

$$F := x_1 F_1 + x_2^{d_1+1-d_2} F_2 - x_3^{d_1+1-d_3} F_3 = 0.$$

Since the polynomials f_i do not have a common zero, it follows easily that F is irreducible in $\overline{K}[x_0, x_1, x_2, x_3]$ (and X is a projective surface).

Let H_i be the hyperplane of \mathbb{P}^3 defined by $x_i = 0$, $i = 1, 2, 3$, and let $D_i = H_i|_X$ be the divisor on X defined by $x_i = 0$, $i = 1, 2, 3$. Since X is a hypersurface in \mathbb{P}^3 and H_i, H_j, X , $i \neq j$, are in general position on \mathbb{P}^3 (as $F_i(1, 0, 0, 0) \neq 0$), by Remark 2.13, D_i and D_j intersect properly on X if $i \neq j$. Letting $P_0 = [1 : 0 : 0 : 0] \in X(K)$, we have $D_1 \cap D_2 \cap D_3 = \{P_0\} \neq \emptyset$, and by construction $D_1 \sim D_2 \sim D_3$. We also set $D = D_1$.

We note that P_0 is a nonsingular point of X , with maximal ideal (x_2, x_3) in the local ring at P_0 . Then if $\{i, j, k\} = \{1, 2, 3\}$, from the equation for F we find $(x_i, x_j)\mathcal{O}_{P_0} = (x_i, x_j, x_k^{d_1+1-d_k})\mathcal{O}_{P_0}$. It follows that

$$(D_i \cdot D_j)_{P_0} = d_1 + 1 - d_k.$$

Note also that $D^2 = \deg X = d_1 + 1$. Then by Lemma 3.11,

$$\begin{aligned} \beta(D, (D_i \cap D_j)_{P_0}) &\geq \frac{2}{3} \sqrt{\frac{D^2}{(D_i \cdot D_j)_{P_0}}} \\ &\geq \frac{2}{3} \sqrt{\frac{d_1 + 1}{d_1 + 1 - d_k}}. \end{aligned}$$

Then it is not hard to show that the condition (5.8) of Theorem 5.9 is satisfied if

$$\left(\frac{2}{3} \sqrt{\frac{d_1 + 1}{d_1 + 1 - d_3}} - 1 \right) + \left(\frac{2}{3} \sqrt{\frac{d_1 + 1}{d_1 + 1 - d_2}} - 1 \right) \left(2 \sqrt{\frac{d_1 + 1}{d_1 + 1 - d_3}} - 2 \right) > 0,$$

which is equivalent to (5.16). Now we let

$$R = \{[t : u : v : 1] \in X(K) \mid f_1(t)u + f_2(t)v^{d_1+1-d_2} = f_3(t), (t, u, v) \in \mathcal{O}_{K,S} \times \mathcal{O}_{K,S}^* \times \mathcal{O}_{K,S}^*\}.$$

Then R is easily seen to be a set of $(D_1 + D_2 + D_3, S)$ -integral points on X . Now under the assumption (5.16), we can apply Theorem 5.9 with the divisors D_1, D_2, D_3 on X and conclude that R lies in a finite union of curves on $X \subset \mathbb{P}^3$. Finally, we note that since the divisors D_i , $i = 1, 2, 3$ are ample on X , Siegel's theorem on integral points on curves implies that if C is a curve in X and $C \cap R$ is infinite, then C is a rational curve, and thus we may take a finite union of rational curves in the conclusion of the theorem. \square

Acknowledgment

We would like to thank Yizhen Zhao for helpful discussions.

References

- [Aut09] Pascal Autissier, *Géométries, points entiers et courbes entières*, Ann. Sci. Éc. Norm. Supér. (4) **42** (2009), no. 2, 221–239. MR 2518077
- [Aut11] ———, *Sur la non-densité des points entiers*, Duke Math. J. **158** (2011), no. 1, 13–27. MR 2794367
- [BCZ03] Yann Bugeaud, Pietro Corvaja, and Umberto Zannier, *An upper bound for the G.C.D. of $a^n - 1$ and $b^n - 1$* , Math. Z. **243** (2003), no. 1, 79–84. MR 1953049
- [CEL01] Steven Dale Cutkosky, Lawrence Ein, and Robert Lazarsfeld, *Positivity and complexity of ideal sheaves*, Math. Ann. **321** (2001), no. 2, 213–234. MR 1866486
- [CZ00] Pietro Corvaja and Umberto Zannier, *On the Diophantine equation $f(a^m, y) = b^n$* , Acta Arith. **94** (2000), no. 1, 25–40. MR 1762454
- [CZ04a] ———, *On a general Thue's equation*, Amer. J. Math. **126** (2004), no. 5, 1033–1055. MR 2089081
- [CZ04b] ———, *On integral points on surfaces*, Ann. of Math. (2) **160** (2004), no. 2, 705–726. MR 2123936
- [CZ05] ———, *A lower bound for the height of a rational function at S -unit points*, Monatsh. Math. **144** (2005), no. 3, 203–224. MR 2130274
- [CZ06] ———, *On the integral points on certain surfaces*, Int. Math. Res. Not. (2006), Art. ID 98623, 20. MR 2219222
- [CZ10] ———, *Integral points, divisibility between values of polynomials and entire curves on surfaces*, Adv. Math. **225** (2010), no. 2, 1095–1118. MR 2671189

- [EF08] Jan-Hendrik Evertse and Roberto G. Ferretti, *A generalization of the Subspace Theorem with polynomials of higher degree*, Diophantine approximation, Dev. Math., vol. 16, SpringerWienNewYork, Vienna, 2008, pp. 175–198. MR 2487693
- [Eis95] David Eisenbud, *Commutative algebra*, Graduate Texts in Mathematics, vol. 150, Springer-Verlag, New York, 1995, With a view toward algebraic geometry. MR 1322960
- [Ful89] William Fulton, *Algebraic curves*, Advanced Book Classics, Addison-Wesley Publishing Company, Advanced Book Program, Redwood City, CA, 1989, An introduction to algebraic geometry, Notes written with the collaboration of Richard Weiss, Reprint of 1969 original. MR 1042981
- [GW20] Ulrich Görtz and Torsten Wedhorn, *Algebraic geometry I. Schemes—with examples and exercises*, second ed., Springer Studium Mathematik—Master, Springer Spektrum, Wiesbaden, [2020] ©2020. MR 4225278
- [Har77] Robin Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, No. 52, Springer-Verlag, New York-Heidelberg, 1977. MR 0463157
- [HL21] Gordon Heier and Aaron Levin, *A generalized Schmidt subspace theorem for closed subschemes*, Amer. J. Math. **143** (2021), no. 1, 213–226. MR 4201783
- [HL22] Keping Huang and Aaron Levin, *Greatest common divisors on the complement of numerically parallel divisors*, 2022.
- [Laz04] Robert Lazarsfeld, *Positivity in algebraic geometry. I*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 48, Springer-Verlag, Berlin, 2004, Classical setting: line bundles and linear series. MR 2095471
- [Lev06] Aaron Levin, *One-parameter families of unit equations*, Math. Res. Lett. **13** (2006), no. 5-6, 935–945. MR 2280786
- [Lev09] ———, *Generalizations of Siegel’s and Picard’s theorems*, Ann. of Math. (2) **170** (2009), no. 2, 609–655. MR 2552103
- [Lev14] ———, *Wirsing-type inequalities*, Bull. Inst. Math. Acad. Sin. (N.S.) **9** (2014), no. 4, 685–710. MR 3309948
- [Lev19] ———, *Greatest common divisors and Vojta’s conjecture for blowups of algebraic tori*, Invent. Math. **215** (2019), no. 2, 493–533. MR 3910069
- [Mat80] Hideyuki Matsumura, *Commutative algebra*, second ed., Mathematics Lecture Note Series, vol. 56, Benjamin/Cummings Publishing Co., Inc., Reading, Mass., 1980. MR 575344

- [MR15] David McKinnon and Mike Roth, *Seshadri constants, diophantine approximation, and Roth's theorem for arbitrary varieties*, Invent. Math. **200** (2015), no. 2, 513–583. MR 3338009
- [RV20] Min Ru and Paul Vojta, *A birational Nevanlinna constant and its consequences*, Amer. J. Math. **142** (2020), no. 3, 957–991. MR 4101336
- [RW22] Min Ru and Julie Tzu-Yueh Wang, *The Ru-Vojta result for subvarieties*, Int. J. Number Theory **18** (2022), no. 1, 61–74. MR 4369792
- [Sch77] H. P. Schlickewei, *The p -adic Thue-Siegel-Roth-Schmidt theorem*, Arch. Math. (Basel) **29** (1977), no. 3, 267–270.
- [Sil87] Joseph H. Silverman, *Arithmetic distance functions and height functions in Diophantine geometry*, Math. Ann. **279** (1987), no. 2, 193–216. MR 919501
- [Sil05] ———, *Generalized greatest common divisors, divisibility sequences, and Vojta's conjecture for blowups*, Monatsh. Math. **145** (2005), no. 4, 333–350. MR 2162351
- [Voj87] Paul Vojta, *Diophantine approximations and value distribution theory*, Lecture Notes in Mathematics, vol. 1239, Springer-Verlag, Berlin, 1987.
- [Voj97] ———, *On Cartan's theorem and Cartan's conjecture*, Amer. J. Math. **119** (1997), no. 1, 1–17.
- [Voj20] ———, *Birational nevanlinna constants, beta constants, and diophantine approximation to closed subschemes*, arXiv preprint arXiv:2008.00405 (2020).
- [Voj23] ———, *Birational Nevanlinna constants, beta constants, and diophantine approximation to closed subschemes*, J. Théor. Nombres Bordeaux **35** (2023), no. 1, 17–61. MR 4596522
- [WY21] Julie Tzu-Yueh Wang and Yu Yasufuku, *Greatest common divisors of integral points of numerically equivalent divisors*, Algebra Number Theory **15** (2021), no. 1, 287–305. MR 4226990

Keping Huang, INSTITUTE FOR ADVANCED STUDIES IN MATHEMATICS, HARBIN INSTITUTE OF TECHNOLOGY, HARBIN, CHINA 150001
E-mail address: kphuang@hit.edu.cn

Aaron Levin, DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, USA 48824
E-mail address: levina@msu.edu

Zheng Xiao, BEIJING INTERNATIONAL CENTER FOR MATHEMATICAL RESEARCH, PEKING UNIVERSITY, BEIJING, CHINA 100871
E-mail address: xiaozheng@bicmr.pku.edu.cn