

Spectral dimension of p -adic integers

SURAJIT BISWAS, BIPUL SAURABH

Abstract

The notion of spectral dimension was introduced by Chakraborty and Pal in [4]. In this paper, we show that the spectral dimension of the ring of p -adic integers, \mathbb{Z}_p , is equal to its manifold dimension, which is 0. Finally, we determine the K -groups of \mathbb{Z}_p , and show that the generators of $K_0(\mathbb{Z}_p)$ can be expressed as finite span of the characters of \mathbb{Z}_p .

Keywords: p -adic Integers; Characters.

2020 MSC: 46L87

1. Introduction

Connes' formulation of noncommutative geometry centers around the concept of spectral triples [1]. These triples originate from fundamental properties of Dirac-type operators on manifolds. Importantly, a spectral triple for the algebra of continuous functions and Dirac operators on L^2 spinors has the remarkable ability to fully reconstruct a closed Riemannian manifold with a spin structure. This achievement is made possible through Connes' reconstruction theorem [2] and the contributions of Lord et al [8].

Noncommutative concepts, described by spectral triples, provide a distinct perspective on the group of p -adic integers, redefining them as differentiable spaces. Operators associated with p -adic numbers hold significant number-theoretic implications, potentially giving rise to captivating spectral functions. Of note, Connes' unpublished construction of a spectral triple for the algebra of continuous functions on the Cantor set is documented in [3]. Klimek et al [7] construct a spectral triple for the C^* -algebra of continuous functions on the space of p -adic integers. Their approach involves utilizing a rooted tree derived from a coarse-grained approximation of the space and applying the forward derivative on this tree. They not only validate the spectral triple's compliance with the traits of a compact spectral metric space but

also establish its equivalence to the traditional p -adic metric on the space of p -adic integers.

Motivated by Connes' definition of dimension for spectral triples, Chakraborty and Pal [4] introduced the spectral dimension as an invariant for ergodic C^* -dynamical systems. They conjectured that for a homogeneous space of a classical compact Lie group, the spectral dimension matches its dimension as a differentiable manifold. The spectral dimensions of $SU(2)$, quaternion sphere H^n , and sphere S^n computed in [4], [11] and [12] further bolster Chakraborty and Pal's conjecture. Furthermore, in [4], spectral dimensions were computed for the noncommutative torus, q -deformation of $SU(l + 1)$ ($l \geq 1$), and the Cuntz algebra $A_u(Q)$. The spectral dimension of the q -deformation of $U(2)$ was computed in [6]. In this article, Section 3 computes the spectral dimension of the group of p -adic integers. Section 4 describes the K -groups of \mathbb{Z}_p .

2. Preliminaries

In this section, we recall the notion of the spectral dimension of a homogeneous space from the reference [4].

Definition 2.1. Given an associative unital $*$ -algebra \mathcal{A} , a spectral triple for \mathcal{A} comprises a triple (\mathcal{H}, π, D) where

- (1) \mathcal{H} is a complex separable Hilbert space,
- (2) $\pi : \mathcal{A} \rightarrow \mathcal{L}(\mathcal{H})$ is a faithful $*$ -representation,
- (3) D is a self-adjoint operator with compact resolvent, satisfying $[D, \pi(a)] \in \mathcal{L}(\mathcal{H})$ for all $a \in \mathcal{A}$.

For clarity, we use (\mathcal{H}, A, D) when the representation π is clear from the context. A spectral triple is termed s -summable if $|I + D^2|^{-s/2}$ is in the ideal \mathcal{L}^1 of trace-class operators.

Since D has compact resolvent, this is equivalent to asserting that $|D|^{-s}$ is trace-class on the complement of its kernel.

Definition 2.2. [14] A compact quantum group $(C(G), \Delta)$ includes a unital C^* -algebra $C(G)$ and a unital $*$ -homomorphism $\Delta : C(G) \rightarrow C(G) \otimes C(G)$ satisfying

- (1) $(\Delta \otimes id)\Delta = (id \otimes \Delta)\Delta$,
- (2) Both $\{(a \otimes I)\Delta(b) : a, b \in C(G)\}$ and $\{(I \otimes a)\Delta(b) : a, b \in C(G)\}$ densely span $C(G) \otimes C(G)$.

Every compact quantum group possesses a unique invariant Haar state h . The Haar state's invariance is described as follows:

$$(h \otimes id)\Delta(a) = h(a)I = (id \otimes h)\Delta(a) \text{ for all } a \in A.$$

Definition 2.3. A compact quantum group $(C(G), \Delta)$ acts on a C^* -algebra A through a $*$ -homomorphism $\tau : A \rightarrow A \otimes C(G)$ satisfying

- (1) $(\tau \otimes id)\tau = (id \otimes \Delta)\tau$,
- (2) $\{(I \otimes b)\tau(a) : a \in A, b \in C(G)\}$ densely spans $A \otimes C(G)$.

A C^* -algebra A is called a homogeneous space of $C(G)$ when the fixed point subalgebra $a \in A : \tau(a) = a \otimes I$ is $\mathbb{C}I$. In such case, the action τ is called ergodic, and $(A, C(G), \tau)$ is called an ergodic C^* -dynamical system.

A covariant representation (π, u) of a C^* -dynamical system $(A, C(G), \tau)$ consists of a unital $*$ -representation $\pi : A \rightarrow \mathcal{L}(\mathcal{H})$, a unitary representation u of $(C(G), \Delta)$ on \mathcal{H} (i.e. a unitary element of the multiplier algebra $M(\mathcal{K}(\mathcal{H}) \otimes C(G))$ with $(id \otimes \Delta)(u) = u_{12}u_{13}$), fulfilling the condition $(\pi \otimes id)\tau(a) = u(\pi(a) \otimes I)u^*$ for all $a \in A$.

In the previous definition, for $j = 2, 3$, define u_{1j} as $\phi_{1j}(u)$, where $\phi_{1j} : \mathcal{K}(\mathcal{H}) \otimes C(G) \rightarrow \mathcal{K}(\mathcal{H}) \otimes C(G) \otimes C(G)$ is given by $\phi_{12}(T \otimes a) = T \otimes a \otimes I$ and $\phi_{13}(T \otimes a) = T \otimes I \otimes a$ for all $T \in \mathcal{K}(\mathcal{H})$ and $a \in C(G)$. For a compact Hausdorff space G , $M(\mathcal{K}(\mathcal{H}) \otimes C(G))$ is isomorphic to $C_b^{str}(G, \mathcal{L}(\mathcal{H}))$, where $C_b^{str}(G, \mathcal{L}(\mathcal{H}))$ represents the set of bounded continuous functions from G to $\mathcal{L}(\mathcal{H})$ equipped with the strict topology. The convergence of a net $\langle T_i \rangle_i$ to T in $\mathcal{L}(\mathcal{H})$ in the strict sense is characterized by $\|T_i S - T S\| \rightarrow 0$ and $\|S T_i - S T\| \rightarrow 0$ for all compact operators $S \in \mathcal{K}(\mathcal{H})$.

Definition 2.4. For a C^* -dynamical system $(A, C(G), \tau)$, an operator D acting on a Hilbert space \mathcal{H} is equivariant with respect to a covariant representation (π, u) of the system if $D \otimes I$ commutes with u . When (π, u) is a covariant representation of $(A, C(G), \tau)$ on a Hilbert space \mathcal{H} and (\mathcal{H}, π, D) is a spectral triple for a dense $*$ -subalgebra \mathcal{A} of A , (\mathcal{H}, π, D) is considered equivariant with respect to (π, u) if D is equivariant with respect to (π, u) .

A homogeneous space A for a compact quantum group $(C(G), \Delta)$ has an invariant state ρ satisfying

$$(\rho \otimes id)\tau(a) = \rho(a)I, \quad a \in A.$$

This invariant state ρ is unique and relates to the Haar state h on $(C(G), \Delta)$ through the equality

$$(id \otimes h)\tau(a) = \rho(a)I, \quad a \in A.$$

Given an ergodic C^* -dynamical system $(A, C(G), \tau)$ with unique invariant state ρ , let $(\mathcal{H}_\rho, \pi_\rho, \eta_\rho)$ denote the GNS representation associated with ρ , i.e. \mathcal{H}_ρ is a Hilbert space, $\eta_\rho : A \rightarrow \mathcal{H}_\rho$ is linear with $\eta_\rho(A)$ dense in \mathcal{H}_ρ , and $\langle \eta_\rho(a), \eta_\rho(b) \rangle = \rho(a^*b)$; and $\pi_\rho : A \rightarrow \mathcal{L}(\mathcal{H}_\rho)$ is the $*$ -representation of A on \mathcal{H}_ρ defined by $\pi_\rho(a)\eta_\rho(b) = \eta_\rho(ab)$. The action τ induces a unitary representation u_τ of $(C(G), \Delta)$ on \mathcal{H}_ρ , making (π_ρ, u_τ) a covariant representation of the system $(A, C(G), \tau)$. Let $\mathcal{O}(G)$ be the dense $*$ -subalgebra of $C(G)$ generated by the matrix entries of irreducible unitary representations of $(C(G), \Delta)$. We define

$$\mathcal{A} = \{a \in A : \tau(a) \in A \otimes_{alg} \mathcal{O}(G)\};$$

by [10, Theorem 1.5], \mathcal{A} is a dense $*$ -subalgebra of A . Let \mathcal{E} be the class of spectral triples for \mathcal{A} equivariant with respect to the covariant representation (π_ρ, u_τ) . We define the spectral dimension of the system $(A, C(G), \tau)$ as the quantity

$$\inf \{s > 0 : \exists D \text{ such that } (\mathcal{H}_\rho, \pi_\rho, D) \in \mathcal{E} \text{ and } D \text{ is } s\text{-summable}\}.$$

We will denote this number by $Sdim(A, C(G), \tau)$.

Proposition 2.5. *For a compact group G , in the ergodic C^* -dynamical system $(C(G), C(G), \Delta)$, we have $\mathcal{A} = \mathcal{O}(G)$.*

Proof. Let $a \in \mathcal{A}$ with $\Delta(a) = \sum_{i=1}^n a_i \otimes \chi_i$ for $a_i \in A$ and $\chi_i \in \mathcal{O}(G)$. Using the identity element 0 of G , we find that $a(y) = \Delta(a)(0, y) = \sum_{i=1}^n a_i(0) \cdot \chi_i(y)$ for all $y \in G$. Thus, $a = \sum_{i=1}^n a_i(0) \cdot \chi_i \in \mathcal{O}(G)$. \square

3. Computation of Spectral dimension

We begin by recalling the group of p -adic integers from [13], discussing its definition and key topological characteristics.

Definition 3.1. Consider the prime number p . For any nonzero rational number x , express it as $x = p^{v_p(x)}x_1$, where x_1 is a rational number coprime to p , meaning that when written in its simplest form, both the numerator and denominator are coprime to p . We define the p -adic absolute value of x , denoted as $|x|_p$, by the formula

$$|x|_p = p^{-v_p(x)}.$$

For the case of $x = 0$, we set $|0|_p = 0$. It can be readily verified that $|\cdot|_p$ constitutes a norm on \mathbb{Q} .

Let R be the set of all sequences $\langle x_n \rangle_{n=1}^\infty$ in \mathbb{Q} , which are Cauchy with respect to the norm $|\cdot|_p$. Addition and multiplication of sequences are defined pointwise:

$$\begin{aligned} \langle x_n \rangle_{n=1}^\infty + \langle y_n \rangle_{n=1}^\infty &= \langle x_n + y_n \rangle_{n=1}^\infty, \\ \langle x_n \rangle_{n=1}^\infty \cdot \langle y_n \rangle_{n=1}^\infty &= \langle x_n \cdot y_n \rangle_{n=1}^\infty. \end{aligned}$$

Hence, $(R, +, \cdot)$ forms a commutative ring. Moreover, the subset \mathfrak{m} of R that consists of null Cauchy sequences, i.e. sequences that converge to zero, is a maximal ideal. Consequently, the quotient ring R/\mathfrak{m} becomes a field. We can include \mathbb{Q} in R through the mapping $x \mapsto (x, x, \dots)$, which is clearly a Cauchy sequence. Thus, we regard \mathbb{Q} as a subfield of R/\mathfrak{m} . This completion of \mathbb{Q} with respect to $|\cdot|_p$ is denoted as \mathbb{Q}_p , and its elements are called p -adic numbers.

The set $\mathbb{Z}_p = \{x \in \mathbb{Q}_p : |x|_p \leq 1\}$ is a subring of \mathbb{Q}_p , referred to as the ring of p -adic integers.

Evidently, \mathbb{Z} (the set of integers) forms a dense subset within \mathbb{Z}_p . This implies that elements of \mathbb{Z}_p can be regarded as formal power series $\sum_{n=0}^\infty x_n p^n$, where $0 \leq x_n \leq p-1$. The topological group $(\mathbb{Z}_p, +)$ is well-established as compact, totally disconnected, and Hausdorff. As a compact abelian group, \mathbb{Z}_p possesses a unique probability Haar measure denoted as μ .

Definition 3.2. [5, Section 1.7] A character of a locally compact abelian group $(G, +)$ is a continuous group homomorphism $\chi : G \rightarrow \mathbb{T}$, where \mathbb{T}

is the circle group, i.e. the multiplicative group of all complex numbers of absolute value one.

The set of characters of the group G form a group under pointwise multiplication, called the dual group and denoted \widehat{G} .

Lemma 3.3. *Let $S = \{(1, 0)\} \cup \{(m, n) \in \mathbb{N} : m < p^n, p \nmid m\}$. The dual group $\widehat{\mathbb{Z}_p}$ is isomorphic to the group of p -power roots of unity, i.e., $\widehat{\mathbb{Z}_p} \cong \left\{ e^{\frac{2\pi im}{p^n}} : (m, n) \in S \right\}$. For any character χ of \mathbb{Z}_p , we have $\chi(1) = e^{\frac{2\pi im}{p^n}}$ for some $(m, n) \in S$.*

Proof. Let χ be a character of \mathbb{Z}_p . Since \mathbb{Z} is dense within \mathbb{Z}_p , the values of $\chi(\mathbb{Z})$ uniquely determine χ . Considering that \mathbb{Z} is a cyclic group generated by 1, this implies that χ is fully characterized by $\chi(1)$. As p^r tends to 0 within \mathbb{Z}_p as r approaches infinity, the sequence $\chi(p^r) = \chi(1)^{p^r}$ converges to $\chi(0) = 1$. Consequently, we find that $\chi(1) = e^{\frac{2\pi im}{p^n}}$ for some $(m, n) \in S$. \square

Henceforth, we adopt the notation $\chi_{m,n}$ for all $(m, n) \in S$ to represent the character of \mathbb{Z}_p , where $\chi_{m,n}(1) = e^{\frac{2\pi im}{p^n}}$. We consider $C(\mathbb{Z}_p)$ both as a compact quantum group and an unital C^* -algebra and compute the spectral dimension of the natural ergodic C^* -dynamical system $(C(\mathbb{Z}_p), C(\mathbb{Z}_p), \Delta)$ associated with \mathbb{Z}_p . The Haar state on $C(\mathbb{Z}_p)$ serves as an invariant Haar state for the homogeneous space $C(\mathbb{Z}_p)$. The relevant covariant representation of this system is given by the triple $(L^2(\mathbb{Z}_p), \pi, u)$, where $L^2(\mathbb{Z}_p)$ is the GNS Hilbert space corresponding to the Haar state on $C(\mathbb{Z}_p)$, π is the representation of $C(\mathbb{Z}_p)$ on $L^2(\mathbb{Z}_p)$ through left multiplication, and u is the right regular representation.

Theorem 3.4. *The spectral dimension of \mathbb{Z}_p is 0.*

Proof. Consider the equivariant self-adjoint operator D with compact resolvent specified by $D(\chi_{m,n}) = ((n+1)^2 p^{n+1})^{1/s} \chi_{m,n}$ for all $(m, n) \in S \setminus \{(1, 0)\}$. For $(m, n), (k, l) \in S$ with $n \leq l$, it follows that $[D, \pi(\chi_{m,n})](\chi_{k,l}) = 0$. This implies that $[D, \pi(\chi_{m,n})]$ is in $\mathcal{L}(L^2(\mathbb{Z}_p))$, being zero on the complement of the finite dimensional space spanned by $\chi_{k,l}$ where $l < n$.

Moreover, we have

$$\begin{aligned} \text{Tr}|D|^{-s} &= \sum_{(m,n) \in S} \frac{1}{(n+1)^2 p^{n+1}} = \frac{1}{p} + \sum_{n=2}^{\infty} \frac{1}{(n+1)^2 p^{n+1}} (p^n - p^{n-1}) \\ &= \frac{1}{p} + \left(1 - \frac{1}{p}\right) \frac{1}{p} \sum_{n=1}^{\infty} \frac{1}{n^2} < \infty, \end{aligned}$$

thus yielding $\mathcal{S}dim(C(\mathbb{Z}_p), C(\mathbb{Z}_p), \Delta) = 0$. \square

4. K -groups

\mathbb{Z}_p can alternatively be identified as the inverse (or projective) limit of the system $\{\langle \mathbb{Z}/p^n \mathbb{Z} \rangle_{n \in \mathbb{N}}, \langle \Phi_n \rangle_{n=2}^{\infty}\}$, where for each $n \in \mathbb{N}$ with $n \geq 2$, the transition map $\Phi_n : \mathbb{Z}/p^n \mathbb{Z} \rightarrow \mathbb{Z}/p^{n-1} \mathbb{Z}$ is defined as $\Phi_n(x \bmod p^n) := x \bmod p^{n-1}$ for all $x \in \mathbb{Z}$. This identification is precisely given by the isomorphism $\phi : \mathbb{Z}_p \rightarrow \varprojlim \mathbb{Z}/p^n \mathbb{Z}$ defined by $\phi(x) = \langle \phi_n(x) \rangle_{n \in \mathbb{N}}$ where for each $n \in \mathbb{N}$, $\phi_n : \mathbb{Z}_p \rightarrow \mathbb{Z}/p^n \mathbb{Z}$ is the projection map defined by $\phi(\sum_{k=0}^{\infty} x_k p^k) = \sum_{k=0}^{n-1} x_k p^k + p^n \mathbb{Z}$. Consequently, $C(\mathbb{Z}_p)$ can be seen as a C^* -algebraic inductive limit of the induced system $\{C(\mathbb{Z}/p^n \mathbb{Z}), \langle \Psi_n \rangle_{n=2}^{\infty}\}$, where $\Psi_n : C(\mathbb{Z}/p^{n-1} \mathbb{Z}) \rightarrow C(\mathbb{Z}/p^n \mathbb{Z})$ is the induced transition map defined by $(\Psi_n f)(x \bmod p^n) := f(x \bmod p^{n-1})$ for all $f \in C(\mathbb{Z}/p^{n-1} \mathbb{Z})$ and $x \in \mathbb{Z}$. The induced isomorphism $\psi : \varinjlim C(\mathbb{Z}/p^n \mathbb{Z}) \rightarrow C(\mathbb{Z}_p)$ is given by $\psi(\langle f_n \rangle_{n \in \mathbb{N}})(x) = \lim_{n \rightarrow \infty} f_n \phi_n(x)$ for all $\langle f_n \rangle_{n \in \mathbb{N}} \in \varinjlim C(\mathbb{Z}/p^n \mathbb{Z})$ and $x \in \mathbb{Z}_p$. As a result, this identification $C(\mathbb{Z}_p) = \varinjlim C(\mathbb{Z}/p^n \mathbb{Z})$ implies $K_i(C(\mathbb{Z}_p)) = \varinjlim K_i(C(\mathbb{Z}/p^n \mathbb{Z}))$ for $i = 1, 2$. Since $K_1(C(\mathbb{Z}/p^n \mathbb{Z})) = 0$ for all $n \in \mathbb{N}$, it follows that $K_1(C(\mathbb{Z}_p)) = 0$.

For every natural number r , \mathbb{Z}_p can be expressed as the disjoint union of p^r balls $x + p^r \mathbb{Z}_p$ for $x \in \{0, 1, \dots, p^r - 1\}$. Consequently, each ball $x + p^r \mathbb{Z}_p$ is both open and closed. For a subset B of a set X , we denote the characteristic function of B on X as $\mathbb{1}_B$.

Theorem 4.1. *The group $K_0(C(\mathbb{Z}_p))$ is generated by the equivalence classes of continuous functions $\mathbb{1}_{x+p^r \mathbb{Z}_p}$ for all $r \in \mathbb{N}$ and $x \in \{0, 1, \dots, p^r - 1\}$.*

Proof. Consider the generating subset \mathcal{K} of $\varinjlim K_0(C(\mathbb{Z}/p^n \mathbb{Z}))$ consisting of elements $\langle f_n \rangle_{n=1}^{\infty}$ such that there exists $r \in \mathbb{N}$ satisfying the recurrence relation $f_n = \Psi_n f_{n-1}$ for all $n > r$ with the initial condition $f_r = \mathbb{1}_{x+p^r \mathbb{Z}}$,

or equivalently, we can express f_n as $f_n = \sum_{l=1}^{n-r} \sum_{i_l=0}^{p-1} \mathbb{1}_{x + \sum_{l=1}^{n-r} i_l p^{r+l-1} + p^n \mathbb{Z}}$, where $x \in \{0, 1, \dots, p^r - 1\}$. Now, using the identification of $\varinjlim K_0(C(\mathbb{Z}/p^n\mathbb{Z}))$ with $K_0(C(\mathbb{Z}_p))$, we have

$$\mathcal{K} = \{f \in K_0(C(\mathbb{Z}_p)) : f = \mathbb{1}_{x+p^r\mathbb{Z}_p} \text{ for } r \in \mathbb{N} \text{ and } x \in \{0, 1, \dots, p^r - 1\}\}.$$

□

We recall a well-known continuous mapping from \mathbb{Z}_p to the interval $[0, 1]$, known as the Monna map [9]. This mapping, denoted as $T : \mathbb{Z}_p \rightarrow [0, 1]$, is defined by $T(\sum_{k=0}^{\infty} x_k p^k) = \sum_{k=0}^{\infty} \frac{x_k}{p^{k+1}}$, where $0 \leq x_k \leq p - 1$. The Monna map T is continuous with respect to the p -adic metric on \mathbb{Z}_p and the standard Euclidean metric on $[0, 1]$. Furthermore, it preserves measures, with T being measure-preserving concerning the probability Haar measure on \mathbb{Z}_p and the Lebesgue measure λ on $[0, 1]$. Additionally, we note that the set $E = \{x \in [0, 1] : x \text{ has multiple base } p \text{ representations}\}$ is countable and thus has measure zero. Consequently, when we restrict the map T to $\mathbb{Z}_p \setminus T^{-1}[E]$, it becomes a bijection. As a result, the induced map $\tilde{T} : L^2([0, 1]) \rightarrow L^2(\mathbb{Z}_p)$ defined as $\tilde{T}(f) := f \circ T$, where $f \in L^2([0, 1])$, is unitary.

Proposition 4.2. *For any $r \in \mathbb{N}$ and $x \in \{0, 1, \dots, p^r - 1\}$, we have*

$$\mathbb{1}_{x+p^r\mathbb{Z}_p} = \sum_{(m,n) \in S, n \leq r} \frac{e^{-\frac{2\pi i m x}{p^n}}}{p^r} \chi_{m,n}.$$

Proof. Let $r \in \mathbb{N}$ and $x \in \{0, 1, \dots, p^r - 1\}$, and note that $\tilde{T}^{-1}(\mathbb{1}_{x+p^r\mathbb{Z}_p}) = \mathbb{1}_{[T(x), T(x) + \frac{1}{p^r}]}$.

Let $(m, n) \in S$ and denote $\xi_{m,n} = e^{\frac{2\pi i m}{p^n}}$. We then have

$$\tilde{T}^{-1}(\chi_{m,n}) = \lim_{N \rightarrow \infty} \sum_{x_0, x_1, \dots, x_N=0}^{p-1} \xi_{m,n}^{x_0 + x_1 p + \dots + x_N p^N} \mathbb{1}_{\left[\frac{x_0}{p} + \frac{x_1}{p^2} + \dots + \frac{x_N}{p^{N+1}}, \frac{x_0}{p} + \frac{x_1}{p^2} + \dots + \frac{x_{N+1}}{p^{N+1}}\right]}.$$

The L^2 -inner product $\langle \mathbb{1}_{x+p^r\mathbb{Z}_p}, \chi_{m,n} \rangle$ simplifies to

$$\begin{aligned}
& \langle \mathbb{1}_{x+p^r\mathbb{Z}_p}, \chi_{m,n} \rangle \\
&= \left\langle \tilde{T}^{-1}(\mathbb{1}_{x+p^r\mathbb{Z}_p}), \tilde{T}^{-1}(\chi_{m,n}) \right\rangle \\
&= \left\langle \mathbb{1}_{[T(x), T(x) + \frac{1}{p^r}]}, \lim_{N \rightarrow \infty} \sum_{x_0, x_1, \dots, x_N=0}^{p-1} \xi_{m,n}^{x_0+x_1p+\dots+x_Np^N} \mathbb{1}_{\left[\frac{x_0}{p} + \dots + \frac{x_N}{p^{N+1}}, \frac{x_0}{p} + \dots + \frac{x_{N+1}}{p^{N+1}}\right]} \right\rangle \\
&= \lim_{N \rightarrow \infty} \int_{T(x)}^{T(x) + \frac{1}{p^r}} \sum_{x_0, x_1, \dots, x_N=0}^{p-1} \overline{\xi_{m,n}^{x_0+x_1p+\dots+x_Np^N}} \mathbb{1}_{\left[\frac{x_0}{p} + \dots + \frac{x_N}{p^{N+1}}, \frac{x_0}{p} + \dots + \frac{x_{N+1}}{p^{N+1}}\right]} d\lambda \\
&= \lim_{N \rightarrow \infty} \sum_{x_r, \dots, x_N=0}^{p-1} \overline{\xi_{m,n}^{x+\sum_{k=r}^N x_k p^k}} \int_{T(x)}^{T(x) + \frac{1}{p^r}} \mathbb{1}_{\left[T(x) + \sum_{k=r}^N \frac{x_k}{p^{k+1}}, T(x) + \sum_{k=r}^N \frac{x_k}{p^{k+1}} + \frac{1}{p^{N+1}}\right]} d\lambda \\
&= \overline{\xi_{m,n}^x} \lim_{N \rightarrow \infty} \frac{1}{p^{N+1}} \sum_{x_r, \dots, x_N=0}^{p-1} \overline{\xi_{m,n}^{\sum_{k=r}^N x_k p^k}} \\
&= \overline{\xi_{m,n}^x} \lim_{N \rightarrow \infty} \frac{1}{p^{N+1}} \prod_{k=r}^N \left(\sum_{l=0}^{p-1} \overline{\xi_{m,n}^{lp^k}} \right).
\end{aligned}$$

Now, we consider two cases.

Case 1: If $n \leq r$, then for each $k = r, r+1, \dots, N$ we have $\overline{\xi_{m,n}^{p^k}} = 1$, and therefore

$$\overline{\xi_{m,n}^x} \lim_{N \rightarrow \infty} \frac{1}{p^{N+1}} \prod_{k=r}^N \left(\sum_{l=0}^{p-1} \overline{\xi_{m,n}^{lp^k}} \right) = \overline{\xi_{m,n}^x} \lim_{N \rightarrow \infty} \frac{p^{N-r+1}}{p^{N+1}} = \frac{\overline{\xi_{m,n}^x}}{p^r},$$

i.e. $\langle \mathbb{1}_{x+p^r\mathbb{Z}_p}, \chi_{m,n} \rangle = \frac{\overline{\xi_{m,n}^x}}{p^r}$.

Case 2: If $n > r$, then $\sum_{l=0}^{p-1} \overline{\xi_{m,n}^{lp^{n-1}}} = \frac{1 - (\overline{\xi_{m,n}^{p^{n-1}}})^p}{1 - \overline{\xi_{m,n}^{p^{n-1}}}} = 0$, and therefore $\langle \mathbb{1}_{x+p^r\mathbb{Z}_p}, \chi_{m,n} \rangle = 0$.

This completes the proof. \square

Acknowledgement: The first named author would like to thank Professor Partha Sarathi Chakraborty for his valuable suggestions and discussions.

References

- [1] A. Connes, *Noncommutative geometry*, Academic Press, Inc., San Diego, CA, 1994. xiv+661 pp. ISBN:0-12-185860-X.
- [2] A. Connes, *On the spectral characterization of manifolds*, J. Noncommut. Geom. **7**(2013), no.1, 1–82.
- [3] E. Christensen, C. Ivan, *Spectral triples for AF C^* -algebras and metrics on the Cantor set*, J. Operator Theory **56**(2006), no.1, 17–46.
- [4] P.S. Chakraborty, A.K. Pal, *An invariant for homogeneous spaces of compact quantum groups*, Adv. Math. **301** (2016), 258–288.
- [5] A. Deitmar, S. Echterhoff, *Principles of harmonic analysis*, Second edition, Universitext, Springer, Cham, 2014. xiv+332 pp. ISBN:978-3-319-05791-0, ISBN:978-3-319-05792-7.
- [6] S. Guin, B. Saurabh, *Equivariant spectral triple for the quantum group $U_q(2)$ for complex deformation parameters*, J. Geom. Phys. **185**(2023), Paper No. 104748, 22 pp.
- [7] S. Klimek, M. McBride, S. Rathnayake, *A p -adic spectral triple*, J. Math. Phys. **55**(2014), no.11, 113502, 16 pp.
- [8] S. Lord, A. Rennie, J.C. Várilly, *Riemannian manifolds in noncommutative geometry*, J. Geom. Phys. **62**(2012), no.7, 1611–1638.
- [9] A.F. Monna, *Sur une transformation simple des nombres P -adiques en nombres réels* (French) Nederl. Akad. Wetensch. Proc. Ser. A 55 Indag. Math. **14** (1952), 1–9.
- [10] P. Podleś, *Symmetries of quantum spaces. Subgroups and quotient spaces of quantum $SU(2)$ and $SO(3)$ groups*, Comm. Math. Phys. 170 (1) (1995) 1–20.
- [11] B. Saurabh, *Spectral dimension of quaternion spheres*, Arch. Math. (Basel) **111**(2018), no.1, 47–55.
- [12] B. Saurabh, *Spectral dimension of spheres*, Comm. Algebra **48**(2020), no.6, 2539–2554.

- [13] V.S. Vladimirov, I.V. Volovich, E.I. Zelenov, *p-adic analysis and mathematical physics*, Ser. Soviet East European Math., 1, World Scientific Publishing Co., Inc., River Edge, NJ, 1994. xx+319 pp. ISBN:981-02-0880-4.
- [14] S.L. Woronowicz, *Compact quantum groups*, Symétries quantiques (Les Houches, 1995), 845–884. North-Holland Publishing Co., Amsterdam, 1998, ISBN:0-444-82867-2.

SURAJIT BISWAS (surajit.biswas@iitgn.ac.in, 241surajit@gmail.com)
Department of Mathematics,
Indian Institute of Technology, Gandhinagar,
Palaj, Gandhinagar 382055, India

BIPUL SAURABH (bipul.saurabh@iitgn.ac.in, saurabhbipul2@gmail.com)
Department of Mathematics,
Indian Institute of Technology, Gandhinagar,
Palaj, Gandhinagar 382055, India