

COUNTING $r \times s$ RECTANGLES IN NONDECREASING AND SMIRNOV WORDS

SELA FRIED*

ABSTRACT. The rectangle capacity, a word statistic that was recently introduced by the author and Mansour, counts, for two fixed positive integers r and s , the number of occurrences of a rectangle of size $r \times s$ in the bargraph representation of a word. In this work we find the bivariate generating function for the distribution on nondecreasing words of the number of $r \times s$ rectangles and the generating function for their total number over all nondecreasing words. We also obtain the analog results for Smirnov words, which are words that have no consecutive equal letters. This complements our recent results concerned with general words (i.e., not restricted) and Catalan words.

Keywords: bargraph, generating function, nondecreasing, Smirnov, word.

Math. Subj. Class.: 05A05, 05A15.

1. INTRODUCTION

Let n and k be two positive integers and set $[k] = \{1, 2, \dots, k\}$. A word over k of length n is any element of the set $[k]^n$. Words have a visual representation in terms of bargraphs (see Figure 1 below), giving rise to many natural statistics on words, such as their water capacity, number of lit cells, and perimeter (See [4] for a comprehensive review of the subject).

In this work we continue the study of a new statistic, that was introduced by the author and Mansour in [2], called *rectangle capacity*. This statistic, for two fixed positive integers r and s , counts the number of occurrences of a rectangle of size $r \times s$ in the bargraph representation of a word (see Figure 2 below). In [2] we studied general words (i.e., not restricted) and Catalan words. The purpose of this work is to explore additional families of words, namely nondecreasing words and Smirnov words. By the former we mean words $w_1 \cdots w_n \in [k]^n$ such that $w_{i+1} \geq w_i$, for each $1 \leq i \leq n-1$, and, by the latter, we mean words $w_1 \cdots w_n \in [k]^n$ such that $w_{i+1} \neq w_i$, for each $1 \leq i \leq n-1$ (cf. [1, Example 23 on p. 193]).

* Department of Computer Science, Israel Academic College, 52275 Ramat Gan, Israel.
friedsela@gmail.com.




FIGURE 1. The bargraph representation of the word 345134




FIGURE 2. There are four occurrences of a rectangle of size 3×2 in the bargraph representation of the word 345134.

Before we begin, let us fix three positive integers r, s , and k and let n be a nonnegative integer. If m is a positive integer, we denote by $[m]$ the set $\{1, 2, \dots, m\}$.

2. MAIN RESULTS - NONDECREASING WORDS

We denote by $\mathcal{W}_{n,k}^{\nearrow}$ the set of nondecreasing words over k of length n , i.e.

$$\mathcal{W}_{n,k}^{\nearrow} = \{w_1 \cdots w_n \in [k]^n : w_i \leq w_{i+1} \text{ for every } i \in [n-1]\}.$$

It is well known that

$$|\mathcal{W}_{n,k}^{\nearrow}| = \binom{n+k-1}{k-1}.$$

We distinguish between two cases, namely $r = 1$ and $r \geq 2$.

2.1. $1 \times s$ rectangles. Denote by $a_{n,k} = a_{n,k}(t)$ the distribution on $\mathcal{W}_{n,k}^{\nearrow}$ of the number of $1 \times s$ rectangles and let $A_k(x, t)$ denote the generating function of the numbers $a_{n,k}$. We shall need the following restrictions of $\mathcal{W}_{n,k}^{\nearrow}$: Let

$$\mathcal{W}_{n,k}^{\nearrow,(0)} = \left\{ w_1 \cdots w_n \in \mathcal{W}_{n,k}^{\nearrow} : w_j > 1 \text{ for every } j \in [n] \right\}$$

and, for $i \in [n]$, we define

$$\mathcal{W}_{n,k}^{\nearrow,(i)} = \left\{ w_1 \cdots w_n \in \mathcal{W}_{n,k}^{\nearrow} : w_i = 1 \text{ and } w_j > 1 \text{ for every } i < j \leq n \right\}.$$

For $0 \leq i \leq n$, let $a_{n,k}^{(i)}$ be the restriction of $a_{n,k}$ to $\mathcal{W}_{n,k}^{\nearrow,(i)}$. Notice that if $w_1 \cdots w_n \in \mathcal{W}_{n,k}^{\nearrow,(i)}$, then $w_1 = \cdots = w_i = 1$.

Theorem 1. *We have*

$$A_k(x, t) = \sum_{i=0}^{k-1} \frac{\alpha_{k-i}(t^i x, t)}{t^{i(s-1)} \prod_{j=1}^i (1 - t^j x)}, \quad (1)$$

where

$$\alpha_m(x, t) = \sum_{n=0}^{s-2} \binom{n+m-1}{m-1} x^n + \frac{x^{s-1}}{1-tx} \binom{s+m-3}{m-1} - \frac{1}{t^{s-1}(1-tx)} \sum_{n=0}^{s-2} \binom{n+m-2}{m-2} (tx)^n.$$

Proof. The set $\mathcal{W}_{n,1}^{\nearrow}$ consists solely of the word $1 \cdots 1$, containing $n - s + 1$ rectangles of size $1 \times s$. Thus, $a_{n,1} = t^{\max\{0, n-s+1\}}$ and it is easily verified that the corresponding generating function is given by

$$A_1(x, t) = \alpha_1(x, t) = \frac{1 - x^{s-1}}{1 - x} + \frac{x^{s-1}}{1 - tx}.$$

Assume now that $k \geq 2$. For $n \geq s$ we have

$$\begin{aligned} a_{n,k} &= a_{n,k}^{(0)} + \sum_{i=1}^n a_{n,k}^{(i)} \\ &= t^{n-s+1} a_{n,k-1} + t^{n-s+1} \sum_{i=1}^n a_{n-i,k-1}. \end{aligned}$$

Multiplying both sides of this equation by x^n , summing over $n \geq s$ and adding $\sum_{n=0}^{s-1} a_{n,k} x^n$ to both sides, with some algebra we obtain

$$\begin{aligned} A_k(x, t) &= \sum_{n=0}^{s-1} \binom{n+k-1}{k-1} x^n - \binom{s+k-3}{k-2} x^{s-1} \\ &\quad + \frac{1}{t^{s-1}(1-tx)} \left(A_{k-1}(tx, t) + \sum_{n=0}^{s-2} \binom{n+k-2}{k-2} ((tx)^s - (tx)^n) \right), \end{aligned}$$

from which (1) follows by induction. \square

Corollary 2. *The generating function for the total number $f_k(n)$ of $1 \times s$ rectangles over all words belonging to $\mathcal{W}_{n,k}^{\nearrow}$ is given by*

$$\frac{x^s \sum_{i=0}^{k-1} (-1)^i \binom{i+s-2}{i} \binom{k+s}{s+i+1} x^i}{(1-x)^{k+1}}.$$

Thus, for $n \geq s$,

$$f_k(n) = \sum_{i=0}^{\min\{k-1, n-s\}} (-1)^i \binom{i+s-2}{i} \binom{k+s}{s+i+1} \binom{n-s-i+k}{k}.$$

Proof. Let $0 \leq i \leq k-1$. We have

$$\begin{aligned} & \frac{\partial}{\partial t} \frac{\alpha_{k-i}(t^i x, t)}{t^{i(s-1)} \prod_{j=1}^i (1-t^j x)} \\ &= \frac{\left(\frac{\partial}{\partial t} \alpha_{k-i}(t^i x, t) \right) t^{i(s-1)} \prod_{j=1}^i (1-t^j x) - \alpha_{k-i}(t^i x, t) \left(\frac{\partial}{\partial t} t^{i(s-1)} \prod_{j=1}^i (1-t^j x) \right)}{\left(t^{i(s-1)} \prod_{j=1}^i (1-t^j x) \right)^2}. \end{aligned}$$

Now,

$$\begin{aligned} \left[t^{i(s-1)} \prod_{j=1}^i (1-t^j x) \right]_{|t=1} &= (1-x)^i, \\ \left[\frac{\partial}{\partial t} t^{i(s-1)} \prod_{j=1}^i (1-t^j x) \right]_{|t=1} &= i(1-x)^{i-1} \left((s-1)(1-x) - \frac{x(i+1)}{2} \right). \end{aligned}$$

It is not hard to see that

$$\begin{aligned} [\alpha_{k-i}(t^i x, t)]_{|t=1} &= \begin{cases} 0 & \text{if } i < k-1 \\ \frac{1}{1-x} & \text{if } i = k-1, \end{cases} \\ \left[\frac{\partial}{\partial t} \alpha_{k-i}(t^i x, t) \right]_{|t=1} &= \begin{cases} \sum_{n=0}^{s-2} (s-1-n) \binom{n+k-1-i}{k-1-i} x^n & \text{if } i < k-1 \\ \frac{x(x^{s-1}+k-1)}{(1-x)^2} & \text{if } i = k-1. \end{cases} \end{aligned}$$

It follows that

$$\begin{aligned} & \left[\frac{\partial}{\partial t} A_k(x, t) \right]_{|t=1} \\ &= \sum_{i=0}^{k-1} \left[\frac{\partial}{\partial t} \frac{\alpha_{k-i}(t^i x, t)}{t^{i(s-1)} \prod_{j=1}^i (1-t^j x)} \right]_{|t=1} \\ &= \sum_{i=0}^{k-2} \frac{\sum_{n=0}^{s-2} (s-1-n) \binom{n+k-1-i}{k-1-i} x^n}{(1-x)^i} + \frac{x \left(x^{s-1} + \frac{(k-1)(2s+k)}{2} \right) - (k-1)(s-1)}{(1-x)^{k+1}}, \end{aligned}$$

from which, with some algebra, the assertion follows. \square

Example 3. For $k=2$ we have

$$f_2(n) = \frac{3n^2 + (7-4s)n + (s-1)(s-4)}{2}$$

and Table 1 lists the corresponding sequences that we found in the On-Line Encyclopedia of Integer Sequences (OEIS) [5]. For $k = 3$ we have

$$f_3(n) = \frac{2n^3 - 3(s-3)n^2 + (s^2 - 10s + 13)n + 2(s-1)(s-3)}{2}$$

and it seems that no corresponding sequence is registered in the OEIS.

s	$f_2(n)$	OEIS
1	$(3n^2 - 3n)/2$	A045943
2	$(3n^2 - n - 2)/2$	A115067
3	$(3n^2 - 5n - 2)/2$	A140090
4	$(3n^2 - 9n)/2$	A140091
5	$(3n^2 - 13n + 4)/2$	A059845
6	$(3n^2 - 17n + 10)/2$	A140672
7	$(3n^2 - 21n + 18)/2$	A140673
8	$(3n^2 - 25n + 28)/2$	A140674
9	$(3n^2 - 29n + 40)/2$	A140675
10	$(3n^2 - 33n + 54)/2$	A151542
11	$(3n^2 - 37n + 70)/2$	A370238

TABLE 1. The total number $f_2(n)$ of $1 \times s$ rectangles over all words of length n , for several values of s .

2.2. $r \times s$ rectangles, where $r \geq 2$. Denote by $b_{n,k} = b_{n,k}(t)$ the distribution on $\mathcal{W}_{n,k}^\nearrow$ of the number of $r \times s$ rectangles and let $B_k(x, t)$ denote the generating function of the numbers $b_{n,k}$. We shall need the following restriction of $\mathcal{W}_{n,k}^\nearrow$: For $m \in [k]$, let

$$\mathcal{W}_{n,k}^{\nearrow, (\geq m)} = \left\{ w_1 \cdots w_n \in \mathcal{W}_{n,k}^\nearrow : w_j \geq m \text{ for each } j \in [n] \right\},$$

and let $b_{n,k}^{(\geq m)}$ be the restriction of $b_{n,k}$ to $\mathcal{W}_{n,k}^{\nearrow, (\geq m)}$. Let $B_k^{(\geq m)}(x, t)$ be the generating function of the numbers $b_{n,k}^{(\geq m)}$.

Lemma 4. *Assume that $k \geq r - 1$. Then*

$$B_k^{(\geq r-1)}(x, t) = \sum_{i=0}^{k-r+1} \frac{\alpha_{k-i}^{(\geq r-1)}(t^i x, t)}{t^{i(s-1)} \prod_{j=0}^{i-1} (1 - t^j x)}, \quad (2)$$

where

$$\alpha_m^{(\geq r-1)}(x, t)$$

$$= \sum_{n=0}^{s-1} \binom{n+m-r+1}{m-r+1} x^n + \frac{x^s}{1-x} \binom{s+m-r}{s-1} - \frac{1}{t^{s-1}(1-x)} \sum_{n=0}^{s-1} \binom{n+m-r}{m-r} (tx)^n.$$

Proof. The set $\mathcal{W}_{n,r-1}^{\nearrow,(\geq r-1)}$ consists solely of the word $(r-1) \cdots (r-1)$, containing no rectangles of size $r \times s$. Thus, $b_{n,r-1}^{(\geq r-1)} = 1$ and the corresponding generating function is given by $B_{r-1}^{(\geq r-1)}(x, t) = 1/(1-x)$. Assume now that $k \geq r$. For $n \geq s$ we have

$$\begin{aligned} b_{n,k}^{(\geq r-1)} &= b_{n,k}^{(\geq r)} + \sum_{i=1}^n b_{i,r-1}^{(\geq r-1)} b_{n-i,k}^{(\geq r)} \\ &= t^{n-s+1} b_{n,k-1}^{(\geq r-1)} + \sum_{i=1}^{n-s} t^{n-i-s+1} b_{n-i,k-1}^{(\geq r-1)} + \sum_{i=n-s+1}^n b_{n-i,k-1}^{(\geq r-1)}. \end{aligned}$$

Multiplying both sides of this equation by x^n , summing over $n \geq s$ and adding $\sum_{n=0}^{s-1} b_{n,k}^{(\geq r-1)} x^n$ to both sides, with some algebra we obtain the equation

$$\begin{aligned} B_k^{(\geq r-1)}(x, t) &= \frac{1}{t^{s-1}(1-x)} B_{k-1}^{(\geq r-1)}(tx, t) + \sum_{n=0}^{s-1} \binom{n+k-r+1}{k-r+1} x^n \\ &\quad - \frac{1}{t^{s-1}(1-x)} \sum_{n=0}^{s-1} \binom{n+k-r}{k-r} (tx)^n + \frac{x^s}{1-x} \sum_{n=0}^{s-1} \binom{n+k-r}{k-r}, \end{aligned}$$

from which the statement immediately follows by induction. \square

Theorem 5. *Assume that $k \geq r$. Then*

$$B_k(x, t) = \frac{1}{(1-x)^{r-1}} \sum_{i=1}^{k-r+1} \frac{\alpha_{k-i}^{(\geq r-1)}(t^i x, t)}{t^{i(s-1)} \prod_{j=1}^{i-1} (1-t^j x)} + \beta_k(x, t),$$

where

$$\begin{aligned} \beta_k(x, t) &= \sum_{i=0}^{s-1} \binom{i+k-1}{k-1} x^i + \frac{1}{(1-x)^{r-1}} \sum_{i=0}^{s-1} \binom{i+k-r}{k-r} x^i \\ &\quad - \sum_{i=0}^{s-1} \binom{i+k-r}{k-r} x^i \sum_{n=0}^{s-1-i} \binom{n+r-2}{r-2} x^n - \frac{1}{t^{s-1}(1-x)^{r-1}} \sum_{i=0}^{s-1} \binom{i+k-r}{k-r} (tx)^i. \end{aligned}$$

Proof. For $n \geq s$ we have

$$b_{n,k} = b_{n,k}^{(\geq r)} + \sum_{i=1}^n b_{i,r-1} b_{n-i,k}^{(\geq r)}$$

$$= t^{n-s+1} b_{n,k-1}^{(\geq r-1)} + \sum_{i=1}^{n-s} t^{n-i-s+1} b_{i,r-1} b_{n-i,k-1}^{(\geq r-1)} + \sum_{i=n-s+1}^n b_{i,r-1} b_{n-i,k-1}^{(\geq r-1)}.$$

Multiplying both sides of this equation by x^n , summing over $n \geq s$ and adding $\sum_{n=0}^{s-1} b_{n,k} x^n$ to both sides, with some algebra we obtain the equation

$$\begin{aligned} & B_k(x, t) \\ &= \frac{1}{t^{s-1}(1-x)^{r-1}} B_{k-1}^{(\geq r-1)}(tx, t) + \sum_{i=0}^{s-1} \binom{i+k-1}{k-1} x^i + \frac{1}{(1-x)^{r-1}} \sum_{i=0}^{s-1} \binom{i+k-r}{k-r} x^i \\ & - \sum_{i=0}^{s-1} \binom{i+k-r}{k-r} x^i \sum_{n=0}^{s-1-i} \binom{n+r-2}{r-2} x^n - \frac{1}{t^{s-1}(1-x)^{r-1}} \sum_{i=0}^{s-1} \binom{i+k-r}{k-r} (tx)^i, \end{aligned}$$

from which, together with (2), the statement immediately follows. \square

Corollary 6. *The generating function for the total number $g_k(n)$ of $r \times s$ rectangles over all words belonging to $\mathcal{W}_{n,k}^\nearrow$ is given by*

$$\frac{x^s \sum_{i=0}^{k-r+1} (-1)^i \binom{i+s-2}{i} \binom{k-r+1+s}{s+i+1} x^i}{(1-x)^{k+1}}.$$

Thus, for $n \geq s$,

$$g_k(n) = \sum_{i=0}^{\min\{k-r+1, n-s\}} (-1)^i \binom{i+s-2}{i} \binom{k-r+1+s}{s+i+1} \binom{n-s-i+k}{k}.$$

Proof. Let $i \in [k-r+1]$. We have

$$\begin{aligned} & \frac{\partial}{\partial t} \frac{\alpha_{k-i}^{(\geq r-1)}(tx, t)}{t^{i(s-1)} \prod_{j=1}^{i-1} (1-t^j x)} \\ &= \frac{\left(\frac{\partial}{\partial t} \alpha_{k-i}^{(\geq r-1)}(tx, t) \right) t^{i(s-1)} \prod_{j=1}^{i-1} (1-t^j x) - \alpha_{k-i}^{(\geq r-1)}(tx, t) \left(\frac{\partial}{\partial t} t^{i(s-1)} \prod_{j=1}^{i-1} (1-t^j x) \right)}{\left(t^{i(s-1)} \prod_{j=1}^{i-1} (1-t^j x) \right)^2}. \end{aligned}$$

Now,

$$\left[t^{i(s-1)} \prod_{j=1}^{i-1} (1-t^j x) \right]_{|t=1} = (1-x)^{i-1},$$

$$\left[\frac{\partial}{\partial t} t^{i(s-1)} \prod_{j=1}^{i-1} (1 - t^j x) \right]_{|t=1} = i(1-x)^{i-2} \left((s-1)(1-x) - \frac{x(i-1)}{2} \right).$$

It is not hard to see that

$$\begin{aligned} \left[\alpha_{k-i}^{(\geq r-1)}(t^i x, t) \right]_{|t=1} &= \begin{cases} 0 & \text{if } i < k - r + 1 \\ \frac{1}{1-x} & \text{if } i = k - r + 1, \end{cases} \\ \left[\frac{\partial}{\partial t} \alpha_{k-i}^{(\geq r-1)}(t^i x, t) \right]_{|t=1} &= \begin{cases} \frac{1}{1-x} \sum_{n=0}^{s-2} (s-1-n) \binom{n+k-r-i}{k-r-i} x^n & \text{if } i < k - r + 1 \\ \frac{x(k-r+1)}{(1-x)^2} & \text{if } i = k - r + 1, \end{cases} \\ \left[\frac{\partial}{\partial t} \beta_k(x, t) \right]_{|t=1} &= \frac{1}{(1-x)^{r-1}} \sum_{n=0}^{s-2} (s-1-n) \binom{n+k-r}{k-r} x^n. \end{aligned}$$

It follows that

$$\begin{aligned} & \left[\frac{\partial}{\partial t} A_k(x, t) \right]_{|t=1} \\ &= \frac{1}{(1-x)^{r-1}} \sum_{i=1}^{k-r+1} \left[\frac{\partial}{\partial t} \frac{\alpha_{k-i}^{(\geq r-1)}(t^i x, t)}{t^{i(s-1)} \prod_{j=1}^{i-1} (1 - t^j x)} \right]_{|t=1} + \left[\frac{\partial}{\partial t} \beta_k(x, t) \right]_{|t=1} \\ &= \frac{1}{(1-x)^{r-1}} \left(\sum_{i=1}^{k-r} \frac{\sum_{n=0}^{s-2} (s-1-n) \binom{n+k-r-i}{k-r-i} x^n}{(1-x)^i} + (k-r+1) \frac{x(k-r+2) - 2(s-1)(1-x)}{2(1-x)^{k-r+2}} \right) \\ &+ \frac{1}{(1-x)^{r-1}} \sum_{n=0}^{s-2} (s-1-n) \binom{n+k-r}{k-r} x^n \\ &= \frac{1}{(1-x)^{r-1}} \sum_{i=0}^{k-r} \frac{\sum_{n=0}^{s-2} (s-1-n) \binom{n+k-r-i}{k-r-i} x^n}{(1-x)^i} + (k-r+1) \frac{x(k-r+2) - 2(s-1)(1-x)}{2(1-x)^{k+1}} \end{aligned}$$

from which, with some algebra, the assertion follows. \square

Example 7. For $r = k = 2$ and arbitrary s we have

$$g_2(n) = \binom{n-s+2}{2}.$$

For $r = s = 2$ and arbitrary k we have

$$g_k(n) = \frac{n-1}{n+1} \binom{k}{2} \binom{n-1+k}{k},$$

which, for $k = 3, 4$ corresponds to [A077414](#) and [A105938](#) in the OEIS, respectively. For $s = 1$ and arbitrary k and r we have

$$g_k(n) = \binom{k-r+2}{2} \binom{n-1+k}{k}$$

and Table 2 lists the corresponding sequences that we found in OEIS.

k	r	OEIS
3	2	A027480
4	2	A033487
5	2	A266732
6	2	A240440
7	2	A266733
4	3	A050534
5	3	A253945
6	3	A271040

TABLE 2. The total number $g_k(n)$ of $r \times 1$ rectangles over all words of length n , for several values of k and r .

3. MAIN RESULTS - SMIRNOV WORDS

We denote by $\mathcal{S}_{n,k}$ the set of Smirnov words (e.g., [1, Example 23 on p. 193]), i.e.,

$$\mathcal{S}_{n,k} = \{w_1 \cdots w_n \in [k]^n : w_i \neq w_{i+1} \text{ for every } i \in [n-1]\}.$$

Since there are no Smirnov words for $k = 1$, we assume that $k \geq 2$. We distinguish between two cases, namely $r = 1$ and $r \geq 2$.

3.1. $1 \times s$ rectangles. Denote by $c_{n,k} = c_{n,k}(t)$ the distribution on $\mathcal{S}_{n,k}$ of the number of $1 \times s$ rectangles. Let $C_k(x, t)$ denote the generating function of the numbers $c_{n,k}$. We shall need the following restrictions of $\mathcal{S}_{n,k}$: Let

$$\mathcal{S}_{n,k}^{(0)} = \{w_1 \cdots w_n \in \mathcal{S}_{n,k} : w_j > 1 \text{ for every } j \in [n]\}$$

and, for $i \in [n]$, we define

$$\mathcal{S}_{n,k}^{(i)} = \{w_1 \cdots w_n \in \mathcal{S}_{n,k} : w_i = 1 \text{ and } w_j > 1 \text{ for every } j \in [i-1]\}.$$

For $0 \leq i \leq n$, let $c_{n,k}^{(i)}$ be the restriction of $c_{n,k}$ to $\mathcal{S}_{n,k}^{(i)}$. Clearly,

$$|\mathcal{S}_{n,k}| = \begin{cases} 1 & \text{if } n = 0 \\ k(k-1)^{n-1} & \text{otherwise,} \end{cases}$$

$$|\mathcal{S}_{n,k}^{(1)}| = \begin{cases} 1 & \text{if } n = 0 \\ (k-1)^{n-1} & \text{otherwise.} \end{cases}$$

Theorem 8. *We have*

$$C_2(x, t) = \frac{2(t-1)x^s}{(1-tx)(1-x)} + \frac{1+x}{1-x} \quad (3)$$

and, for $k \geq 3$,

$$C_k(x, t) = \frac{(1+tx)(\gamma_k(x, t) + \delta_k(x, t)C_{k-1}(tx, t))}{1+tx-txC_{k-1}(tx, t)}, \quad (4)$$

where

$$\begin{aligned} \gamma_k(x, t) &= \frac{1+x-k(k-1)^{s-1}x^s}{1-(k-1)x} - \frac{1+tx-k(k-1)^{s-1}(tx)^s}{t^{s-1}(1-(k-1)tx)}, \\ \delta_k(x, t) &= \frac{1-(k-2)tx-(k-1)^{s-1}(tx)^s}{t^{s-1}(1-(k-1)tx)} - \frac{tx}{1+tx} \frac{(1+x)(1-((k-1)x)^{s-1})}{1-(k-1)x}. \end{aligned}$$

Proof. For $n \geq 1$, the set $\mathcal{S}_{n,2}$ consists of two words, namely $1212 \cdots$ and $2121 \cdots$, containing exactly $n-s+1$ $1 \times s$ rectangles, each. It is easily verified that the corresponding generating function is given by (3). Assume now that $k \geq 3$. For $n \geq s$ we have

$$\begin{aligned} c_{n,k} &= c_{n,k}^{(0)} + \sum_{i=1}^n c_{n,k}^{(i)} \\ &= c_{n,k-1} + \sum_{i=1}^{n-s+1} t^{i-1} c_{i-1,k-1} c_{n-i+1,k}^{(1)} + \sum_{i=n-s+2}^n t^{n-s+1} c_{i-1,k-1} c_{n-i+1,k}^{(1)}, \quad (5) \end{aligned}$$

$$c_{n,k}^{(1)} = t \left(c_{n-1,k} - c_{n-1,k}^{(1)} \right). \quad (6)$$

Multiplying both sides of (6) by x^n , summing over $n \geq s$ and adding $\sum_{n=0}^{s-1} c_{n,k}^{(1)} x^n$ to both sides, with some algebra we obtain

$$C_k^{(1)}(x, t) = \frac{1}{1+tx} \left(\sum_{n=0}^{s-1} c_{n,k}^{(1)} x^n + tx \left(C_k(x, t) - \sum_{n=0}^{s-2} c_{n,k} x^n \right) + tx \sum_{n=0}^{s-2} c_{n,k}^{(1)} x^n \right). \quad (7)$$

Multiplying both sides of (5) by x^n , summing over $n \geq s$ and adding $\sum_{n=0}^{s-1} c_{n,k} x^n$ to both sides, with some algebra we obtain

$$C_k(x, t) = \sum_{n=0}^{s-1} c_{n,k} x^n + \frac{1}{t^{s-1}} \left(C_{k-1}(tx, t) - \sum_{n=0}^{s-1} c_{n,k-1} (tx)^n \right)$$

$$\begin{aligned}
& + C_{k-1}(tx, t) \left(C_k^{(1)}(x, t) - \sum_{n=0}^{s-1} c_{n,k}^{(1)} x^n \right) \\
& + \frac{1}{t^{s-1}} \sum_{i=2}^s c_{s-i+1,k}^{(1)} (tx)^{s-i+1} \left(C_{k-1}(tx, t) - \sum_{n=0}^{i-2} c_{n,k-1}(tx)^n \right). \tag{8}
\end{aligned}$$

Substituting (7) into (8), with some algebra we obtain (4). \square

Corollary 9. *The generating function for the total number $h_k(n)$ of $1 \times s$ rectangles over all words belonging to $\mathcal{S}_{n,k}$ is given by*

$$\frac{x^s \sum_{i=1}^{k-1} i^{s-1} (i+1)}{(1 - (k-1)x)^2}. \tag{9}$$

Thus, for $n \geq s$,

$$h_k(n) = (k-1)^{n-s} (n-s+1) \sum_{i=1}^{k-1} i^{s-1} (i+1).$$

Proof. We have

$$\left[\frac{\partial}{\partial t} C_2(x, t) \right]_{|t=1} = \frac{2x^s}{(1-tx)^2}.$$

Thus, (9) holds for $k=2$. Assume now that $k \geq 3$ and that (9) holds for $k-1$. It is not hard to see that

$$\begin{aligned}
& [\gamma_k(x, t)]_{|t=1} = 0, \\
& \left[\frac{\partial}{\partial t} \gamma_k(x, t) \right]_{|t=1} = \frac{x(s-1) + s}{1 - (k-1)x} - \frac{1+x - k(k-1)^{s-1}x^s}{(1 - (k-1)x)^2}, \\
& [\delta_k(x, t)]_{|t=1} = 1, \\
& \left[\frac{\partial}{\partial t} \delta_k(x, t) \right]_{|t=1} = 1 - s - \frac{x}{1+x} \left[\frac{\partial}{\partial t} \gamma_k(x, t) \right]_{|t=1}, \\
& [C_k(x, t)]_{|t=1} = \frac{1+x}{1 - (k-1)x}, \\
& \left[\frac{\partial}{\partial x} C_k(x, t) \right]_{|t=1} = \frac{k}{(1 - (k-1)x)^2}, \\
& \left[\frac{\partial}{\partial t} C_k(tx, t) \right]_{|t=1} = x \left[\frac{\partial}{\partial x} C_k(x, t) \right]_{|t=1} + \left[\frac{\partial}{\partial t} C_k(x, t) \right]_{|t=1}.
\end{aligned}$$

Differentiating (4) with respect to t and substituting $t = 1$, we obtain

$$\begin{aligned}
\left[\frac{\partial}{\partial t} C_k(x, t) \right]_{|t=1} &= \left[\frac{(1+tx)(\gamma_k(x, t) + \delta_k(x, t)C_{k-1}(tx, t))}{1+tx-txC_{k-1}(tx, t)} \right]_{|t=1} \\
&= \frac{(s-1)(k-1)x^2 + (2+(k-2)s)x - s + 1}{(1-(k-1)x)^2} \\
&\quad + \frac{(1-(k-2)x)^2}{(1-(k-1)x)^2} \left[\frac{\partial}{\partial t} C_{k-1}(x, t) \right]_{|t=1} + \left[\frac{\partial}{\partial t} \gamma_k(x, t) \right]_{|t=1} \\
&= \frac{k(k-1)^{s-1}x^s}{(1-(k-1)x)^2} + \frac{x^s \sum_{i=1}^{k-2} i^{s-1}(i+1)}{(1-(k-1)x)^2} \\
&= \frac{x^s \sum_{i=1}^{k-1} i^{s-1}(i+1)}{(1-(k-1)x)^2}. \quad \square
\end{aligned}$$

Example 10. For $k = 2$ and arbitrary s we have

$$h_2(n) = 2(n - s + 1).$$

For $k = 3$ and arbitrary s we have

$$h_3(n) = 2^{n-s}(n - s + 1)(2 + 3 \cdot 2^{s-1}),$$

which, for $s = 2, 3$ corresponds to [A241204](#) and [A281200](#) in the OEIS, respectively.

3.2. $r \times s$ rectangles, where $r \geq 2$. Denote by $d_{n,k} = d_{n,k}(t)$ the distribution on $\mathcal{S}_{n,k}$ of the number of $r \times s$ rectangles. Let $D_k(x, t)$ denote the generating function of the numbers $d_{n,k}$. We shall need the following restrictions of $\mathcal{S}_{n,k}$: For $m \in [k]$ let

$$\begin{aligned}
\mathcal{S}_{n,k}^{(\geq m)} &= \{w_1 \cdots w_n \in \mathcal{S}_{n,k} : w_i \geq m \text{ for every } i \in [n]\}, \\
\bar{\mathcal{S}}_{n,k}^{(\geq m)} &= \{w_1 \cdots w_n \in \mathcal{S}_{n,k}^{(\geq m)} : w_1 \neq m\}, \\
\bar{\mathcal{S}}_{n,k} &= \{w_1 \cdots w_n \in \mathcal{S}_{n,k} : w_1 \neq r - 1\}.
\end{aligned}$$

Denote by $d_{n,k}^{(\geq m)}$ (resp. $\bar{d}_{n,k}^{(\geq m)}$, $\bar{d}_{n,k}$) the restriction of $d_{n,k}$ to $\mathcal{S}_{n,k}^{(\geq m)}$ (resp. to $\bar{\mathcal{S}}_{n,k}^{(\geq m)}$, $\bar{\mathcal{S}}_{n,k}$). Let $D_k^{(\geq m)}(x, t)$ (resp. $\bar{D}_k^{(\geq m)}(x, t)$, $\bar{D}_k(x, t)$) denote the generating function of the numbers $d_{n,k}^{(\geq m)}$ (resp. $\bar{d}_{n,k}^{(\geq m)}$, $\bar{d}_{n,k}$). Clearly,

$$\begin{aligned}
\left| \mathcal{S}_{n,k}^{(\geq m)} \right| &= \begin{cases} 1 & \text{if } n = 0 \\ (k - m + 1)(k - m)^{n-1} & \text{otherwise,} \end{cases} \\
\left| \bar{\mathcal{S}}_{n,k}^{(\geq m)} \right| &= (k - m)^n.
\end{aligned}$$

Lemma 11. *We have*

$$D_r^{(\geq r-1)}(x, t) = \frac{1+x}{1-x} \quad (10)$$

and, for $k \geq r+1$,

$$D_k^{(\geq r-1)}(x, t) = (1+x) \frac{\frac{1}{t^{s-1}} D_{k-1}^{(\geq r-1)}(tx, t) + \sigma_k(x, t)}{-\frac{x}{t^{s-1}} D_{k-1}^{(\geq r-1)}(tx, t) + \rho_k(x, t)},$$

where

$$\begin{aligned} \sigma_k(x, t) &= \frac{1+x - (k-r+1)(k-r)^{s-1}x^s}{1-(k-r)x} - \frac{1+tx - (k-r+1)(k-r)^{s-1}(tx)^s}{t^{s-1}(1-(k-r)tx)}, \\ \rho_k(x, t) &= 1 - (k-r+1)x^2 \frac{1 - ((k-r)x)^{s-1}}{1-(k-r)x} + x \frac{1+tx - (k-r+1)(k-r)^{s-1}(tx)^s}{t^{s-1}(1-(k-r)tx)}. \end{aligned}$$

Proof. For $n \geq 1$, the set $\mathcal{S}_{n,r}^{(\geq r-1)}$ consists of two words, namely $r(r-1)r(r-1)\cdots$ and $(r-1)r(r-1)r(r-1)\cdots$, containing no $r \times s$ rectangles. Thus, (10) holds true in this case. Assume now that $k \geq r+1$. For $n \geq s$ we have

$$\begin{aligned} d_{n,k}^{(\geq r-1)} &= d_{n,k}^{(\geq r)} + \sum_{i=1}^n d_{i-1,k}^{(\geq r)} \bar{d}_{n-i,k}^{(\geq r-1)} \\ &= t^{n-s+1} d_{n,k-1}^{(\geq r-1)} + \sum_{i=1}^s d_{i-1,k-1}^{(\geq r-1)} \bar{d}_{n-i,k}^{(\geq r-1)} + \sum_{i=s+1}^n t^{i-s} d_{i-1,k-1}^{(\geq r-1)} \bar{d}_{n-i,k}^{(\geq r-1)}, \end{aligned} \quad (11)$$

$$\begin{aligned} \bar{d}_{n,k}^{(\geq r-1)} &= d_{n,k}^{(\geq r)} + \sum_{i=2}^n d_{i-1,k}^{(\geq r)} \bar{d}_{n-i,k}^{(\geq r-1)} \\ &= t^{n-s+1} d_{n,k-1}^{(\geq r-1)} + \sum_{i=2}^s d_{i-1,k-1}^{(\geq r-1)} \bar{d}_{n-i,k}^{(\geq r-1)} + \sum_{i=s+1}^n t^{i-s} d_{i-1,k-1}^{(\geq r-1)} \bar{d}_{n-i,k}^{(\geq r-1)}. \end{aligned} \quad (12)$$

Multiplying both sides of (11) by x^n , summing over $n \geq s$ and adding $\sum_{n=0}^{s-1} d_{n,k}^{(\geq r-1)} x^n$ to both sides, with some algebra we obtain the equation

$$\begin{aligned} &D_k^{(\geq r-1)}(x, t) \\ &= \sum_{n=0}^{s-1} d_{n,k}^{(\geq r-1)} x^n + \frac{1}{t^{s-1}} \left(D_{k-1}^{(\geq r-1)}(tx, t) - \sum_{n=0}^{s-1} d_{n,k-1}^{(\geq r-1)} (tx)^n \right) - \sum_{i=1}^s d_{i-1,k-1}^{(\geq r-1)} x^i \sum_{n=0}^{s-i-1} \bar{d}_{n,k}^{(\geq r-1)} x^n \\ &+ \bar{D}_k^{(\geq r-1)}(x, t) \left(\sum_{i=1}^s d_{i-1,k-1}^{(\geq r-1)} x^i + \frac{x}{t^{s-1}} \left(D_{k-1}^{(\geq r-1)}(tx, t) - \sum_{n=0}^{s-1} d_{n,k-1}^{(\geq r-1)} (tx)^n \right) \right). \end{aligned} \quad (13)$$

Multiplying both sides of (12) by x^n , summing over $n \geq s$ and adding $\sum_{n=0}^{s-1} \bar{d}_{n,k}^{(\geq r-1)} x^n$ to both sides, with some algebra we obtain the equation

$$\begin{aligned} & \bar{D}_k^{(\geq r-1)}(x, t) \left(1 - \frac{x}{t^{s-1}} D_{k-1}^{(\geq r-1)}(tx, t) - \sum_{i=2}^s d_{i-1, k-1}^{(\geq r-1)} x^i + \frac{x}{t^{s-1}} \sum_{n=0}^{s-1} d_{n, k-1}^{(\geq r-1)} (tx)^n \right) \\ &= \frac{1}{t^{s-1}} D_{k-1}^{(\geq r-1)}(tx, t) + \sum_{n=0}^{s-1} \bar{d}_{n, k}^{(\geq r-1)} x^n - \frac{1}{t^{s-1}} \sum_{n=0}^{s-1} d_{n, k-1}^{(\geq r-1)} (tx)^n \\ & - \sum_{i=2}^s d_{i-1, k-1}^{(\geq r-1)} x^i \sum_{n=0}^{s-i-1} \bar{d}_{n, k}^{(\geq r-1)} x^n. \end{aligned}$$

Substituting this into (13), with some algebra, the assertion follows. \square

Theorem 12. *We have*

$$D_r(x, t) = \frac{1+x}{1-x}$$

and, for $k \geq r+1$,

$$D_k(x, t) = (1+x) \frac{\frac{1}{t^{s-1}} D_{k-1}^{(\geq r-1)}(tx, t) + \sigma_k(x, t)}{-\frac{(r-1)x}{t^{s-1}} D_{k-1}^{(\geq r-1)}(tx, t) - (r-2)(1+x) + (r-1)\rho_k(x, t)}. \quad (14)$$

Proof. We have

$$\begin{aligned} d_{n, k} &= d_{n, k}^{(\geq r)} + (r-1) \sum_{i=1}^n d_{i-1, k}^{(\geq r)} \bar{d}_{n-i, k} \\ &= t^{n-s+1} d_{n, k-1}^{(\geq r-1)} + (r-1) \sum_{i=1}^s d_{i-1, k-1}^{(\geq r-1)} \bar{d}_{n-i, k} + (r-1) \sum_{i=s+1}^n t^{i-s} d_{i-1, k-1}^{(\geq r-1)} \bar{d}_{n-i, k}, \end{aligned} \quad (15)$$

$$\begin{aligned} \bar{d}_{n, k} &= d_{n, k}^{(\geq r)} + (r-1) \sum_{i=2}^n d_{i-1, k}^{(\geq r)} \bar{d}_{n-i, k} + (r-2) \bar{d}_{n-1, k} \\ &= t^{n-s+1} d_{n, k-1}^{(\geq r-1)} + (r-1) \sum_{i=2}^s d_{i-1, k-1}^{(\geq r-1)} \bar{d}_{n-i, k} \\ &+ (r-1) \sum_{i=s+1}^n t^{i-s} d_{i-1, k-1}^{(\geq r-1)} \bar{d}_{n-i, k} + (r-2) \bar{d}_{n-1, k}. \end{aligned} \quad (16)$$

Multiplying both sides of (15) by x^n , summing over $n \geq s$ and adding $\sum_{n=0}^{s-1} d_{n, k} x^n$ to both sides, with some algebra we obtain

$$D_k(x, t)$$

$$\begin{aligned}
&= \sum_{n=0}^{s-1} d_{n,k} x^n + \frac{1}{t^{s-1}} \left(D_{k-1}^{(\geq r-1)}(tx, t) - \sum_{n=0}^{s-1} d_{n,k-1}^{(\geq r-1)}(tx)^n \right) - (r-1) \sum_{i=1}^s d_{i-1,k-1}^{(\geq r-1)} x^i \sum_{n=0}^{s-i-1} \bar{d}_{n,k} x^n \\
&+ \bar{D}_k(x, t) \left((r-1) \sum_{i=1}^s d_{i-1,k-1}^{(\geq r-1)} x^i + \frac{(r-1)x}{t^{s-1}} \left(D_{k-1}^{(\geq r-1)}(tx, t) - \sum_{n=0}^{s-1} d_{n,k-1}^{(\geq r-1)}(tx)^n \right) \right). \tag{17}
\end{aligned}$$

Multiplying both sides of (16) by x^n , summing over $n \geq s$ and adding $\sum_{n=0}^{s-1} \bar{d}_{n,k} x^n$ to both sides, with some algebra we obtain

$$\begin{aligned}
&\bar{D}_k(x, t) \\
&= \sum_{n=0}^{s-1} \bar{d}_{n,k} x^n + \frac{1}{t^{s-1}} \left(D_{k-1}^{(\geq r-1)}(tx, t) - \sum_{n=0}^{s-1} d_{n,k-1}^{(\geq r-1)}(tx)^n \right) \\
&+ (r-1) \sum_{i=2}^s d_{i-1,k-1}^{(\geq r-1)} x^i \left(\bar{D}_k(x, t) - \sum_{n=0}^{s-i-1} \bar{d}_{n,k} x^n \right) \\
&+ \frac{(r-1)x}{t^{s-1}} \bar{D}_k(x, t) \left(D_{k-1}^{(\geq r-1)}(tx, t) - \sum_{n=0}^{s-1} d_{n,k-1}^{(\geq r-1)}(tx)^n \right) + (r-2)x \left(\bar{D}_k(x, t) - \sum_{n=0}^{s-2} \bar{d}_{n,k} x^n \right).
\end{aligned}$$

Solving for $\bar{D}_k(x, t)$ and substituting it into (17), with some algebra the assertion follows. \square

Corollary 13. *The generating function for the total number $i_k(n)$ of $r \times s$ rectangles over all words belonging to $\mathcal{S}_{n,k}$ is given by*

$$\frac{x^s \sum_{i=1}^{k-r} i^{s-1} (i+1)}{(1 - (k-1)x)^2}.$$

Thus, for $n \geq s$,

$$i_k(n) = (k-1)^{n-s} (n-s+1) \sum_{i=1}^{k-r} i^{s-1} (i+1).$$

Proof. It is not hard to see that

$$\begin{aligned}
\left[D_k^{(\geq r-1)}(x, t) \right]_{|t=1} &= \frac{1+x}{1 - (k-r+1)x}, \\
\left[\frac{\partial}{\partial x} D_k^{(\geq r-1)}(x, t) \right]_{|t=1} &= \frac{k-r+2}{(1 - (k-r+1)x)^2}, \\
\left[\frac{\partial}{\partial t} D_k^{(\geq r-1)}(x, t) \right]_{|t=1} &= \frac{x^s \sum_{n=0}^{k-r} n^{s-1} (n+1)}{(1 - (k-r+1)x)^2}.
\end{aligned}$$

We omit the rest of the details, that are very similar to those in the proof of Corollary 9.

□

REFERENCES

- [1] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press, 2009.
- [2] S. Fried and T. Mansour, Counting $r \times s$ rectangles in (Catalan) words, arxiv preprint arXiv:2405.06962 [math.CO], 2024. Available at <https://arxiv.org/abs/2405.06962>.
- [3] T. Mansour and A. S. Shabani, Bargraphs in bargraphs, *Turkish J. Math.* **42** (2018), 2763–2773.
- [4] T. Mansour and A. S. Shabani, Enumerations on bargraphs, *Discrete Math. Lett.* **2** (2019), 65–94.
- [5] N. J. A. Sloane, The On-Line Encyclopedia of Integer Sequences, OEIS Foundation Inc., <https://oeis.org>.