

INTEGRAL REPRESENTATION FOR RIEMANN-SIEGEL $Z(t)$ FUNCTION.

J. ARIAS DE REYNA

ABSTRACT. We apply Poisson formula for a strip to give a representation of $Z(t)$ by means of an integral.

$$F(t) = \int_{-\infty}^{\infty} \frac{h(x)\zeta(4+ix)}{7 \cosh \pi \frac{x-t}{7}} dx, \quad Z(t) = \frac{\operatorname{Re} F(t)}{\left(\frac{1}{4} + t^2\right)^{\frac{1}{2}} \left(\frac{25}{4} + t^2\right)^{\frac{1}{2}}}.$$

After that we get the estimate

$$Z(t) = \left(\frac{t}{2\pi}\right)^{\frac{7}{4}} \operatorname{Re}\{e^{i\vartheta(t)} H(t)\} + O(t^{-3/4}),$$

with

$$H(t) = \int_{-\infty}^{\infty} \left(\frac{t}{2\pi}\right)^{ix/2} \frac{\zeta(4+it+ix)}{7 \cosh(\pi x/7)} dx = \left(\frac{t}{2\pi}\right)^{-\frac{7}{4}} \sum_{n=1}^{\infty} \frac{1}{n^{\frac{1}{2}+it}} \frac{2}{1 + \left(\frac{t}{2\pi n^2}\right)^{-7/2}}.$$

We explain how the study of this function can lead to information about the zeros of the zeta function on the critical line.

1. INTRODUCTION

Riemann stated that the number $N_0(T)$ of zeros of $\zeta(s)$ on the critical line with ordinates between 0 and T is approximately equal to the total number of zeros $N(T)$ in the critical strip. In a letter to Weierstrass he said that this assertion follows from a new expansion of the function $\Xi(t)$ that he has not sufficiently simplified to communicate. Nobody knows what expansion Riemann is talking about. Siegel considered the possibility that it was his integral expansion studied in [10] but concluded that probably Riemann was wrong about the possibilities of this expansion. This work is an attempt to obtain a development following ideas available to Riemann.

It is somewhat surprising that the approximation obtained here is very similar to the one studied in our paper [2], obtained by an entirely different method.

2. INTEGRAL REPRESENTATION FOR $Z(t)$.

In this section, we define a real analytic function $F: \mathbf{R} \rightarrow \mathbf{C}$ such that its real part is equal to $Z(t)\sqrt{\frac{1}{4} + t^2}\sqrt{\frac{25}{4} + t^2}$. For this we will apply the solution of the Dirichlet problem for a strip.

The solution of Dirichlet problem in a strip was considered in [4] we quote its solution from Rademacher [7, § 31].

Theorem 1. *The solution $u(\sigma, t)$ of the Dirichlet problem on the strip $S(a, b) = \{s \in \mathbf{C}: a \leq \sigma \leq b\}$ with boundary values $u(a, t) = A(t)$ and $u(b, t) = B(t)$, where $A(t)$ and*

2020 *Mathematics Subject Classification.* Primary 11M06; Secondary 30D99.

Key words and phrases. función zeta, representation integral.

$B(t) = \mathcal{O}(e^{k|t|})$ with $0 < k < \pi/(b-a)$ is given by

$$(1) \quad U(\sigma, t) = \frac{1}{2(b-a)} \int_{-\infty}^{+\infty} A(x) \omega\left(\frac{\sigma-a}{b-a}, \frac{x-t}{b-a}\right) dx \\ + \frac{1}{2(b-a)} \int_{-\infty}^{+\infty} B(x) \omega\left(\frac{b-\sigma}{b-a}, \frac{x-t}{b-a}\right) dx.$$

where

$$(2) \quad \omega(\sigma, t) = \frac{\sin \pi \sigma}{\cosh \pi t - \cos \pi \sigma}.$$

Theorem 2. *There is a holomorphic function f in the strip $S(-3, 5)$ such that for $\frac{1}{2} < \sigma < 5$ and $t \in \mathbf{R}$*

$$(3) \quad Z(t) \sqrt{\frac{1}{4} + t^2} \sqrt{\frac{25}{4} + t^2} = \operatorname{Re} \left\{ \frac{1}{2\sigma - 1} \int_{-\infty}^{\infty} \frac{f(\sigma + ix)}{\cosh \pi \frac{x-t}{2\sigma-1}} dx \right\}.$$

Proof. Consider the function $\Phi(s) = (s+2)s(1-s)(3-s)\zeta(s)\zeta(1-s)$. It is clear that $\Phi(s)$ is an entire function that satisfies the functional equation $\Phi(s) = \Phi(1-s)$. The function $\Phi(s)$ have simple zeros at $-4, -6 \dots$ and at $5, 7 \dots$ due to the trivial zeros of $\zeta(s)$. The zeros on the strip $S(-4, 5)$ are all of even order, at $-2, 3$ and at each nontrivial zero ρ of $\zeta(s)$.

In the critical line, we have $\zeta(\frac{1}{2} + it) = Z(t)e^{-i\vartheta(t)}$ where $Z(t)$ and $\vartheta(t)$ real functions that are real analytic, and such that

$$(4) \quad Z(t) = Z(-t), \quad \vartheta(-t) = -\vartheta(t).$$

It follows that

$$(5) \quad \Phi(\frac{1}{2} + it) = (\frac{1}{4} + t^2)(\frac{25}{4} + t^2)Z(t)^2.$$

Since the strip $S(-3, 5)$ is simply connected and the zeros of $\Phi(s)$ are of even order, there is an analytic function $f: S(-4, 5) \rightarrow \mathbf{C}$ with $f(s)^2 = \Phi(s)$. We fix the sign of $f(s)$ so that

$$(6) \quad f(\frac{1}{2} + it) = Z(t) \sqrt{\frac{1}{4} + t^2} \sqrt{\frac{25}{4} + t^2}, \quad \text{for } t \in \mathbf{R},$$

where we take $\sqrt{\frac{1}{4} + t^2} > 0$ and $\sqrt{\frac{25}{4} + t^2} > 0$. Notice that $Z(0) = \zeta(1/2) = -1.46 < 0$, so that $f(\frac{1}{2}) < 0$.

We apply Theorem 1 to the harmonic function $f(s) = f(\sigma + it)$ on the strip $S(1-\sigma, \sigma)$. The boundary values are $A(t) = f(1-\sigma + it)$ and $B(t) = f(\sigma + it)$. Since $f(s)$ is real on the real axis and on the critical line, we have $f(1-\sigma + it) = \overline{f(\sigma + it)}$. Therefore, we obtain

$$f(\frac{1}{2} + it) = Z(t) \sqrt{\frac{1}{4} + t^2} \sqrt{\frac{25}{4} + t^2} \\ = \frac{1}{2(2\sigma - 1)} \int_{-\infty}^{\infty} A(x) \omega\left(\frac{\sigma-\frac{1}{2}}{2\sigma-1}, \frac{x-t}{2\sigma-1}\right) dx + \frac{1}{2(2\sigma - 1)} \int_{-\infty}^{+\infty} B(x) \omega\left(\frac{\sigma-\frac{1}{2}}{2\sigma-1}, \frac{x-t}{2\sigma-1}\right) dx.$$

The two integrals are complex conjugate numbers; therefore,

$$Z(t) \sqrt{\frac{1}{4} + t^2} \sqrt{\frac{25}{4} + t^2} = 2 \operatorname{Re} \frac{1}{2(2\sigma - 1)} \int_{-\infty}^{\infty} B(x) \omega\left(\frac{1}{2}, \frac{x-t}{2\sigma-1}\right) dx.$$

We have

$$\omega\left(\frac{1}{2}, \frac{x-t}{2\sigma-1}\right) = \frac{\sin \frac{\pi}{2}}{\cosh \pi \frac{x-t}{2\sigma-1} - \cos \frac{\pi}{2}} = \frac{1}{\cosh \pi \frac{x-t}{2\sigma-1}},$$

and the result follows. \square

There are several interesting cases of (3), but we start by trying to obtain the consequences of case $\sigma = 4$.

$$Z(t)\sqrt{\frac{1}{4} + t^2}\sqrt{\frac{25}{4} + t^2} = \operatorname{Re}\left\{\frac{1}{7} \int_{-\infty}^{\infty} \frac{f(4+ix)}{\cosh \pi \frac{x-t}{7}} dx\right\}.$$

Therefore, we define for $t \in \mathbf{R}$

$$(7) \quad F(t) = \int_{-\infty}^{\infty} \frac{f(4+ix)}{7 \cosh \pi \frac{x-t}{7}} dx.$$

It is easy to prove that $F(t)$ is analytic for $|\operatorname{Im}(t)| < \frac{7}{2}$.

2.1. The function $f(4+ix)$.

Proposition 3. *The function $f(4+ix) = h(x)\zeta(4+ix)$ where $h(x)$ is the real analytic function defined by*

$$(8) \quad h(x) = \frac{\sqrt{2}}{(2\pi)^2} \left\{ (6+ix)(4+ix)(3+ix)(1+ix)(2\pi)^{-ix} \cosh \frac{\pi x}{2} \Gamma(4+ix) \right\}^{1/2},$$

that satisfies $h(0) = \frac{3}{\pi^2} \sqrt{6} > 0$.

Proof. For $-2 < \sigma < 3$ the function $\Phi(\sigma)$ is continuous and does not vanish, so its sign coincides with the sign at $s = 1/2$. Therefore $\Phi(\sigma) > 0$ in this range. In the same range $f(\sigma) < 0$ because we took $f(\frac{1}{2}) < 0$ (take $t = 0$ in (6)).

$f(s)$ is defined on the strip $S(-4, 5)$ as the analytic square root of the function $\Phi(s)$ that is negative at $s = \frac{1}{2}$.

By the functional equation of $\zeta(s)$ we have

$$\begin{aligned} \Phi(s) &= (s+2)s(1-s)(3-s)\zeta(s)\zeta(1-s) \\ &= 2(s+2)s(1-s)(3-s)(2\pi)^{-s} \cos \frac{\pi s}{2} \Gamma(s)\zeta(s)^2. \end{aligned}$$

Notice that the function $2(s+2)s(1-s)(3-s)(2\pi)^{-s} \cos \frac{\pi s}{2} \Gamma(s)$ is analytic in $s = 4+ix$ with $x \in \mathbf{R}$ and does not vanish so that there is a real analytic function $h: \mathbf{R} \rightarrow \mathbf{C}$ defined by (8). By a simple computation we obtain

$$\frac{\Phi(4+ix)}{\zeta(4+ix)^2} = \frac{2}{(2\pi)^4} (6+ix)(4+ix)(3+ix)(1+ix)(2\pi)^{-ix} \cosh \frac{\pi x}{2} \Gamma(4+ix).$$

From this we see that $h(0) = \pm \frac{3}{\pi^2} \sqrt{6}$. We may define $h(x)$ as the only continuous square root with $h(0) = \frac{3}{\pi^2} \sqrt{6}$.

With these elections we have $f(4+ix) = \pm h(x)\zeta(4+ix)$. We know that $f(x) < 0$ for $-2 < x < 3$, and $f(4) > 0$, because f changes sign at $x = 3$. Therefore, taking $h(0) > 0$ we obtain the equality $f(4+ix) = h(x)\zeta(4+ix)$. \square

With the above definitions, we may write (7) as

$$(9) \quad F(t) = \int_{-\infty}^{\infty} \frac{h(x)\zeta(4+ix)}{7 \cosh \pi \frac{x-t}{7}} dx, \quad Z(t) = \frac{\operatorname{Re} F(t)}{\left(\frac{1}{4} + t^2\right)^{\frac{1}{2}} \left(\frac{25}{4} + t^2\right)^{\frac{1}{2}}}.$$

Proposition 4. *There are real analytic real functions $\rho: \mathbf{R} \rightarrow \mathbf{R}$ and $\alpha: \mathbf{R} \rightarrow \mathbf{R}$ such that for all $x \in \mathbf{R}$ we have $h(x) = \rho(x)e^{i\alpha(x)}$.*

Proof. Since $h(x) \neq 0$, we may use the result in [1, Cor.2.3] (see also Proposition 24 in Section 4) to show that there are a real analytic function $\alpha(x)$ such that $h(x) = |h(x)|e^{i\alpha(x)}$. The function $\rho(x) := |h(x)|$ is real analytic, because $h(x)$ does not vanish.

Since $\rho(0) > 0$ and $h(0) > 0$ we have $\alpha(0) = 2k\pi$. For definiteness, we take $\alpha(0) = 0$. \square

Proposition 5. *For $x \rightarrow +\infty$, we have the two asymptotic expansions*

$$(10) \quad \log \rho(x) \sim 2 \log x + \frac{7}{4} \log \frac{x}{2\pi} - \sum_{n=1}^{\infty} \frac{(-1)^n (6^{2n} + 4^{2n} + 2 \cdot 3^{2n} + 2^{2n} + 2)}{4nx^{2n}},$$

$$(11) \quad \alpha(x) \sim \frac{x}{2} \log \frac{x}{2\pi} - \frac{x}{2} + \frac{15\pi}{8} - \sum_{n=0}^{\infty} \frac{(-1)^n (3 \cdot 6^{2n} + 2 \cdot 4^{2n} + 3 \cdot 3^{2n} + 2^{2n} + 1)}{(2n+1)x^{2n+1}} + \sum_{n=1}^{\infty} \frac{(-1)^n B_{2n}}{4n(2n-1)x^{2n-1}}.$$

Proof. Consider the function between parenthesis in (8)

$$(6+ix)(4+ix)(3+ix)(1+ix)(2\pi)^{-ix} \cosh \frac{\pi x}{2} \Gamma(4+ix),$$

it is equal to

$$(6+ix)(4+ix)(3+ix)^2(2+ix)(1+ix)^2(2\pi)^{-ix} \cosh \frac{\pi x}{2} \Gamma(1+ix).$$

This function is equal to $2^4 3^3$ for $x = 0$ and its factors are all positive at $x = 0$, so we consider its logarithm as the sum of logarithms that are real at $x = 0$.

$$\begin{aligned} \log(6+ix) + \log(4+ix) + 2 \log(3+ix) + \log(2+ix) + 2 \log(1+ix) \\ + \log(\cosh \frac{\pi x}{2}) - ix \log(2\pi) + \log \Gamma(1+ix). \end{aligned}$$

Each term of the above sum has a well-known asymptotic expansion for $x \rightarrow +\infty$

$$\begin{aligned} \log(6+ix) &= \log x + \frac{\pi i}{2} - \sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{6i}{x}\right)^n, & \log(4+ix) &= \log x + \frac{\pi i}{2} - \sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{4i}{x}\right)^n, \\ 2 \log(3+ix) &= 2 \log x + \pi i - \sum_{n=1}^{\infty} \frac{2}{n} \left(\frac{3i}{x}\right)^n, & \log(2+ix) &= \log x + \frac{\pi i}{2} - \sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{2i}{x}\right)^n, \\ 2 \log(1+ix) &= 2 \log x + \pi i - \sum_{n=1}^{\infty} \frac{2}{n} \left(\frac{i}{x}\right)^n, \\ \log(\cosh \frac{\pi x}{2}) &= -\log 2 + \frac{\pi x}{2} + \log(1+e^{-\pi x}) \sim \frac{\pi x}{2} - \log 2, \end{aligned}$$

$$-ix \log(2\pi) + \log \Gamma(1+ix)$$

$$\sim -\frac{\pi x}{2} + \frac{1}{2} \log \frac{x}{2\pi} + \log(2\pi) + ix \log \frac{x}{2\pi} - ix + \frac{\pi i}{4} + i \sum_{n=1}^{\infty} \frac{(-1)^n B_{2n}}{2n(2n-1)x^{2n-1}}.$$

Then $\log h(x)$ is easily obtained from these values. From (8) we have to add $\frac{1}{2}$ of the above expansions and add the log of $\sqrt{2}/(2\pi)^2$. According to our election $\alpha(0) = 0$ we must take the logarithm of $\sqrt{2}/(2\pi)^2$ equal to $\log(\sqrt{2}/(2\pi)^2)$ the real logarithm. Therefore,

$$\begin{aligned} \log h(x) &= \frac{1}{2} \log 2 - 2 \log(2\pi) \\ &+ \frac{7}{2} \log x + \frac{7\pi i}{4} - \sum_{n=1}^{\infty} \frac{6^n + 4^n + 2 \cdot 3^n + 2^n + 2}{2n} \left(\frac{i}{x}\right)^n \\ &+ \frac{\pi x}{4} - \frac{1}{2} \log 2 - \frac{\pi x}{4} + \frac{1}{4} \log \frac{x}{2\pi} + \frac{1}{2} \log(2\pi) \\ &+ \frac{ix}{2} \log \frac{x}{2\pi} - \frac{ix}{2} + \frac{\pi i}{8} + i \sum_{n=1}^{\infty} \frac{(-1)^n B_{2n}}{4n(2n-1)x^{2n-1}}. \end{aligned}$$

Separating the real and imaginary parts, we obtain the asymptotic expansion of $\log \rho(x)$ and $\alpha(x)$

$$\begin{aligned} \alpha(x) &\sim \frac{x}{2} \log \frac{x}{2\pi} - \frac{x}{2} + \frac{15\pi}{8} \\ &- \sum_{n=0}^{\infty} \frac{(-1)^n (3 \cdot 6^{2n} + 2 \cdot 4^{2n} + 3 \cdot 3^{2n} + 2^{2n} + 1)}{(2n+1)x^{2n+1}} + \sum_{n=1}^{\infty} \frac{(-1)^n B_{2n}}{4n(2n-1)x^{2n-1}}. \end{aligned}$$

$$\log \rho(x) \sim -\frac{3}{2} \log(2\pi) + \frac{7}{2} \log x + \frac{1}{4} \log \frac{x}{2\pi} - \sum_{n=1}^{\infty} \frac{(-1)^n (6^{2n} + 4^{2n} + 2 \cdot 3^{2n} + 2^{2n} + 2)}{4nx^{2n}}.$$

or by summing the two convergent series we obtain

$$\log \rho(x) \sim 2 \log x + \frac{7}{4} \log \frac{x}{2\pi} + \frac{1}{4} \log \frac{(1+x^2)^2(4+x^2)(16+x^2)(9+x^2)^2(36+x^2)}{x^{14}}.$$

The difference between these two functions is exponentially small.

$$\begin{aligned} \alpha(x) &\sim \frac{x}{2} \log \frac{x}{2\pi} - \frac{x}{2} + \frac{15\pi}{8} - \arctan \frac{1}{x} - \frac{1}{2} \arctan \frac{2}{x} - \arctan \frac{3}{x} \\ &- \frac{1}{2} \arctan \frac{4}{x} - \frac{1}{2} \arctan \frac{6}{x} + \sum_{n=1}^{\infty} \frac{(-1)^n B_{2n}}{4n(2n-1)x^{2n-1}}. \end{aligned}$$

□

Lemma 6. *The first terms of the expansion of $\log \rho(x)$ are given by*

$$\log \rho(x) = -\frac{7}{4} \log(2\pi) + \frac{15}{4} \log x + \frac{19}{x^2} - \frac{433}{2x^4} + \frac{13069}{3x^6} - \frac{439633}{4x^8} + \dots$$

Therefore, we obtain the next Lemma.

Lemma 7. *We have*

$$\rho(x) = x^2 \left(\frac{x}{2\pi}\right)^{7/4} \left(1 + \frac{19}{x^2} + \mathcal{O}(x^{-4})\right), \quad x \rightarrow +\infty.$$

Analogously, the first terms of the asymptotic expansion of $\alpha(x)$ by the above reasoning are equal to

$$\alpha(x) = \frac{x}{2} \log \frac{x}{2\pi} - \frac{x}{2} + \frac{15\pi}{8} - \frac{241}{24x} + \frac{41279}{720x^3} - \frac{2348641}{2520x^5} + \dots$$

Therefore,

Lemma 8. *We have for $x \rightarrow +\infty$*

$$\alpha(x) = \frac{x}{2} \log \frac{x}{2\pi} - \frac{x}{2} + \frac{15\pi}{8} - \frac{241}{24x} + \mathcal{O}(x^{-3}).$$

3. FIRST APPROXIMATION TO $Z(t)$.

Let us define

$$(12) \quad \theta(x) = \frac{x}{2} \log \frac{x}{2\pi} - \frac{x}{2} + \frac{15\pi}{8} - \frac{241}{24x}, \quad x > 0,$$

$$(13) \quad \rho_0(x) = x^2 \left(\frac{x}{2\pi} \right)^{7/4} \left(1 + \frac{19}{x^2} \right) = (2\pi)^{-7/4} x^{7/4} (19 + x^2), \quad x > 0,$$

and

$$L_1(x, t) = 1 + \frac{15x}{4t} + \frac{ix^2}{4t} + \frac{165x^2}{32t^2} + \frac{241ix}{24t^2} + \frac{41ix^3}{48t^2} - \frac{x^4}{32t^2}.$$

The object of this section is to prove the following theorem

Theorem 9. *For $t \in \mathbf{R}$ let*

$$(14) \quad G(t) = e^{i\theta(t)} \rho_0(t) \int_{-\infty}^{\infty} L_1(x, t) \left(\frac{t}{2\pi} \right)^{ix/2} \frac{\zeta(4 + it + ix)}{7 \cosh \frac{\pi x}{7}} dx.$$

Then $G(t)$ is real analytic and for $t \rightarrow +\infty$ we have $F(t) = G(t) + \mathcal{O}(t^{3/4} \log^6 t)$, and consequently

$$(15) \quad Z(t) = \frac{\operatorname{Re}(G(t))}{\sqrt{\frac{1}{4} + t^2} \sqrt{\frac{25}{4} + t^2}} + \mathcal{O}(t^{-5/4} \log^6 t).$$

Proof. In the next Lemmas and Propositions we define the functions $F_1(t)$, $F_2(t)$, $F_3(t)$ and $F_4(t)$, such that $F_4(t) = G(t)$ and $F(t) = F_1(t) + \mathcal{O}(t^{-1/4})$, $F_1(t) = F_2(t) + \mathcal{O}(t^{3/4})$, $F_2(t) = F_3(t) + \mathcal{O}(t^{3/4} \log^6 t)$ and $F_3(t) - F_4(t) = \mathcal{O}(t^{-1/4})$. And the Theorem follows from this and (9). \square

Lemma 10 (Gabcke). *The incomplete Gamma functions satisfies the inequality*

$$\Gamma(a, x) := \int_x^{\infty} v^{a-1} e^{-v} dv \leq a e^{-x} x^{a-1}, \quad x > a \geq 1.$$

Proof. See [3, p.145, p.84 in the electronic version]. \square

Lemma 11. *Let $\alpha = 28/\pi$ and*

$$F_1(t) = \int_{t-\alpha \log t}^{t+\alpha \log t} \frac{h(x) \zeta(4 + ix)}{7 \cosh \frac{\pi}{7}(x-t)} dx.$$

Then for $t \rightarrow +\infty$ we have $F(t) = F_1(t) + \mathcal{O}(t^{-1/4})$.

Proof. The difference $F(t) - F_1(t)$ is the sum of two integrals, we bound each one. Let $b > 0$, we shall take at the end $b = \alpha \log t$. By Lemma 7 $\rho(x) \leq Cx^{15/4}$ for $x \geq x_0$. Also $|\zeta(4 + ix)|$ is bounded for x real. Therefore, we have for $t \rightarrow +\infty$

$$\begin{aligned} \left| \int_{t+b}^{\infty} \frac{h(x)\zeta(4 + ix)}{7 \cosh \frac{\pi}{7}(x-t)} dx \right| &\ll \int_{t+b}^{\infty} \frac{x^{15/4}}{7 \cosh \frac{\pi}{7}(x-t)} dx \\ &= \int_{b/t}^{\infty} \frac{t^{15/4}(1+y)^{15/4}}{7 \cosh \pi y t/7} t dy. \end{aligned}$$

We divide the integral into two other (assuming $t > b$)

$$\begin{aligned} &= t^{15/4} \int_{b/t}^1 \frac{(1+y)^{15/4} t}{7 \cosh \pi y t/7} dy + t^{15/4} \int_1^{\infty} \frac{(1+y)^{15/4} t}{7 \cosh \pi y t/7} dy \\ &\leq 2^{15/4} t^{15/4} \int_{b/t}^1 \frac{t}{\cosh \pi y t/7} dy + 2^{15/4} t^{15/4} \int_1^{\infty} \frac{y^{15/4} t}{\cosh \frac{\pi y t}{7}} dy \\ &\ll t^{15/4} \int_{b/t}^{\infty} t e^{-\frac{\pi y t}{7}} dy + t^{15/4} \int_1^{\infty} y^{15/4} t e^{-\frac{\pi y t}{7}} dy \\ &= \frac{7}{\pi} t^{15/4} e^{-\pi b/7} + t^{15/4} \int_{\pi t/7}^{\infty} \left(\frac{7x}{\pi t}\right)^{15/4} t e^{-x} \frac{7}{\pi t} dx \\ &\ll t^{15/4} e^{-\pi b/7} + \int_{\pi t/7}^{\infty} x^{15/4} e^{-x} dx \end{aligned}$$

The last integral can be bounded by Gabcke's lemma. In the first one we take as $b = \alpha \log t$ and obtain

$$\ll t^{15/4} t^{-4} + t^{15/4} e^{-\pi t/7} = \mathcal{O}(t^{-1/4}).$$

Now we turn to the second integral, for some positive $a < t/2$, (we will take at the end $a = \alpha \log t$)

$$\begin{aligned} \left| \int_{-\infty}^{t-a} \frac{h(x)\zeta(4 + ix)}{7 \cosh \frac{\pi}{7}(x-t)} dx \right| \\ \leq \left| \int_{t/2}^{t-a} \frac{h(x)\zeta(4 + ix)}{7 \cosh \frac{\pi}{7}(x-t)} dx \right| + \left| \int_{-\infty}^{t/2} \frac{h(x)\zeta(4 + ix)}{7 \cosh \frac{\pi}{7}(x-t)} dx \right| := I_1 + I_2. \end{aligned}$$

For the second integral here, we notice that $h(x)$ is bounded for $|x| < 1$ so that we have

$$\begin{aligned} I_2 = \left| \int_{-\infty}^{t/2} \dots \right| &\ll \int_{-1}^1 \frac{1}{\cosh \frac{\pi}{7}(x-t)} dx \\ &+ \int_1^{t/2} \frac{x^{15/4}}{\cosh \frac{\pi}{7}(x-t)} dx + \int_{-\infty}^{-1} \frac{|x|^{15/4}}{\cosh \frac{\pi}{7}(x-t)} dx := J_1 + J_2 + J_3. \end{aligned}$$

It is easy to see that $J_1 \ll e^{-\pi t/7}$. For J_2 we can obtain a better bound, but it is sufficient to bound the integral by the maximum of the function by the length of the interval

$$J_2 \ll t^{15/4} e^{-\pi t/14} \cdot t = \mathcal{O}(t^{-1/4}).$$

For J_3 we reduce it to an incomplete gamma integral

$$J_3 = \int_1^{\infty} \frac{x^{15/4}}{\cosh \frac{\pi}{7}(x+t)} dx \ll \int_1^{\infty} x^{15/4} e^{-\pi \frac{x+t}{7}} dx \ll e^{-\pi t/7}.$$

All is reduced to the bound of I_1

$$\begin{aligned} I_1 &= \left| \int_{t/2}^{t-a} \frac{h(x)\zeta(4+ix)}{7 \cosh \frac{\pi}{7}(x-t)} dx \right| \ll \int_{t/2}^{t-a} \frac{x^{15/4}}{\cosh \frac{\pi}{7}(x-t)} dx \\ &\ll t^{15/4} \int_{t/2}^{t-a} \frac{dx}{\cosh \frac{\pi}{7}(x-t)} \leq t^{15/4} \int_{-\infty}^{-a} \frac{dy}{\cosh \frac{\pi y}{7}} \\ &\leq 2t^{15/4} \int_{-\infty}^{-a} e^{\frac{\pi y}{7}} dy \ll t^{15/4} e^{-\pi a/7}. \end{aligned}$$

We see that this is $\mathcal{O}(t^{-1/4})$ if we choose $a = \alpha \log t$ with $\alpha = 28/\pi$, as we wanted. \square

Lemma 12. For $t - \alpha \log t > 0$ let us define

$$F_2(t) = \int_{t-\alpha \log t}^{t+\alpha \log t} \frac{\rho_0(x)e^{i\theta(x)}\zeta(4+ix)}{7 \cosh \frac{\pi}{7}(x-t)} dx,$$

where $\rho_0(x)$ and $\theta(x)$ are defined in (13) and (12). Then $F_1(t) = F_2(t) + \mathcal{O}(t^{3/4})$.

Proof. By Lemma 7 and the definition of $\rho_0(x)$ we have for $x \rightarrow +\infty$

$$\rho(x) = \rho_0(x) + \mathcal{O}(x^{-1/4}), \quad \rho_0(x) = \mathcal{O}(x^{15/4}).$$

By Lemma 8 and the definition of $\theta(x)$ we have

$$\alpha(x) = \theta(x) + \mathcal{O}(x^{-3}).$$

Then

$$\begin{aligned} h(x) &= \rho(x)e^{i\alpha(x)} \\ &= (\rho_0(x) + \mathcal{O}(x^{-1/4}))e^{i\theta(x) + \mathcal{O}(x^{-3})} \\ &= \rho_0(x)e^{i\theta(x)}(1 + \mathcal{O}(x^{-4}))(1 + \mathcal{O}(x^{-3})) \\ &= \rho_0(x)e^{i\theta(x)}(1 + \mathcal{O}(x^{-3})) \\ &= \rho_0(x)e^{i\theta(x)} + \mathcal{O}(x^{3/4}). \end{aligned}$$

It follows that the difference $F_1(t) - F_2(t)$ is bounded by

$$F_1(t) - F_2(t) \ll \int_{t-\alpha \log t}^{t+\alpha \log t} \frac{x^{3/4}}{7 \cosh \frac{\pi}{7}(x-t)} dx$$

In the integration interval $x^{3/4} \ll t^{3/4}$ and we have

$$F_1(t) - F_2(t) \ll t^{3/4} \int_{t-\alpha \log t}^{t+\alpha \log t} \frac{1}{7 \cosh \frac{\pi}{7}(x-t)} dx \ll t^{3/4},$$

since the integral on \mathbf{R} is bounded by a constant. \square

Lemma 13. For $t - \alpha \log t > 0$, define

$$F_3(t) = e^{i\theta(t)}\rho_0(t) \int_{-\alpha \log t}^{\alpha \log t} L_1(x, t) \left(\frac{t}{2\pi}\right)^{ix/2} \frac{\zeta(4+it+ix)}{7 \cosh(\pi x/7)} dx,$$

where

$$(16) \quad L_1(x, t) = 1 + \frac{15x}{4t} + \frac{ix^2}{4t} + \frac{165x^2}{32t^2} + \frac{241ix}{24t^2} + \frac{41ix^3}{48t^2} - \frac{x^4}{32t^2}.$$

Then we have $F_2(t) = F_3(t) + \mathcal{O}(t^{3/4} \log^6 t)$.

Proof. First, notice that

$$F_2(t) = \int_{t-\alpha \log t}^{t+\alpha \log t} \frac{\rho_0(x) e^{i\theta(x)} \zeta(4+ix)}{7 \cosh \frac{\pi}{7}(x-t)} dx = \int_{-\alpha \log t}^{\alpha \log t} \frac{\rho_0(t+x) e^{i\theta(t+x)} \zeta(4+it+ix)}{7 \cosh(\pi x/7)} dx.$$

Note also that $|x| \leq \alpha \log t$ when x is in the integration interval. We apply Taylor's theorem to the two functions (see Rudin [9, p. 110]).

For x in the integration interval there exists ξ between t and $x+t$, therefore $\xi \in (t-\alpha \log t, t+\alpha \log t)$ with

$$\begin{aligned} \theta(t+x) = & \left(\frac{t}{2} \log \frac{t}{2\pi} - \frac{t}{2} + \frac{15\pi}{8} - \frac{241}{24t} \right) + \left(\frac{1}{2} \log \frac{t}{2\pi} + \frac{241}{24t^2} \right) x + \left(\frac{1}{4t} - \frac{241}{24t^3} \right) x^2 \\ & - \left(\frac{1}{12t^2} - \frac{241}{24t^4} \right) x^3 + \left(\frac{1}{24\xi^3} - \frac{241}{24\xi^5} \right) x^4. \end{aligned}$$

So,

$$(17) \quad \theta(t+x) = \theta(t) + \left(\frac{1}{2} \log \frac{t}{2\pi} + \frac{241}{24t^2} \right) x + \frac{1}{4t} x^2 - \frac{1}{12t^2} x^3 + \mathcal{O}(t^{-3} \log^4 t).$$

Recall that

$$\rho_0(x) = x^2 \left(\frac{x}{2\pi} \right)^{7/4} \left(1 + \frac{19}{x^2} \right) = (2\pi)^{-7/4} x^{7/4} (19 + x^2).$$

For $t > 0$ large enough and $x \in (-\alpha \log t, \beta \log t)$, we have Taylor's expansion with $\xi \in (t-\alpha \log t, t+\alpha \log t)$

$$\begin{aligned} (2\pi)^{7/4} \rho_0(t+x) = & t^{7/4} (19 + t^2) + t^{7/4} \left(\frac{15t}{4} + \frac{133}{4t} \right) x \\ & + t^{7/4} \left(\frac{165}{32} + \frac{399}{32t^2} \right) x^2 + \xi^{7/4} \left(\frac{385}{128\xi} - \frac{133}{128\xi^3} \right) x^3. \end{aligned}$$

The last term is $\mathcal{O}(t^{3/4} (\log t)^3)$. Eliminating all terms of order less than or equal to this one, we obtain

$$(2\pi)^{7/4} \rho_0(t+x) = t^{7/4} (19 + t^2) + \frac{15t^{11/4}}{4} x + \frac{165t^{7/4}}{32} x^2 + \mathcal{O}(t^{3/4} (\log t)^3).$$

So that $(2\pi)^{7/4} \rho_0(t+x)$ is equal to

$$(18) \quad (2\pi)^{7/4} \rho_0(t) \left(1 + \frac{15t}{4(19+t^2)} x + \frac{165}{32(19+t^2)} x^2 + \mathcal{O}(t^{-3} (\log t)^3) \right).$$

By (17) and (18) we have

$$(19) \quad (2\pi)^{7/4} \rho_0(t+x) e^{i\theta(t+x)} = (2\pi)^{7/4} \rho_0(t) e^{i\theta(t)} \left(\frac{t}{2\pi} \right)^{ix/2} K(x, t),$$

where

$$\begin{aligned} K(x, t) = & \exp \left\{ i \left(\frac{241x}{24t^2} + \frac{x^2}{4t} - \frac{x^3}{12t^2} + \mathcal{O}(t^{-3} \log^4 t) \right) \right\} \\ & \cdot \left(1 + \frac{15t}{4(19+t^2)} x + \frac{165}{32(19+t^2)} x^2 + \mathcal{O}(t^{-3} (\log t)^3) \right). \end{aligned}$$

The exponential factor is equal to

$$\exp \left(\frac{241ix}{24t^2} + \frac{ix^2}{4t} - \frac{ix^3}{12t^2} \right) (1 + \mathcal{O}(t^{-3} \log^4 t)).$$

Expanding the exponential and retaining only terms greater than t^{-3} yields

$$\begin{aligned} & \left(1 + \frac{ix^2}{4t} + \frac{241ix}{24t^2} - \frac{ix^3}{12t^2} - \frac{x^4}{32t^2} + \mathcal{O}(t^{-3}(\log t)^6)\right)(1 + \mathcal{O}(t^{-3}\log^4 t)) \\ &= 1 + \frac{ix^2}{4t} + \frac{241ix}{24t^2} - \frac{ix^3}{12t^2} - \frac{x^4}{32t^2} + \mathcal{O}(t^{-3}(\log t)^6). \end{aligned}$$

Expanding also the second factor in the expression for $K(x, t)$ and retaining only the terms greater than t^{-3} , yields

$$1 + \frac{15x}{4t} + \frac{165x^2}{32t^2} + \mathcal{O}(t^{-3}(\log t)^3).$$

Therefore,

$$\begin{aligned} K(x, t) &= \left(1 + \frac{ix^2}{4t} + \frac{241ix}{24t^2} - \frac{ix^3}{12t^2} - \frac{x^4}{32t^2} + \mathcal{O}(t^{-3}(\log t)^6)\right) \\ &\quad \cdot \left(1 + \frac{15x}{4t} + \frac{165x^2}{32t^2} + \mathcal{O}(t^{-3}(\log t)^3)\right) \\ &= 1 + \frac{15x}{4t} + \frac{ix^2}{4t} + \frac{165x^2}{32t^2} + \frac{241ix}{24t^2} + \frac{41ix^3}{48t^2} - \frac{x^4}{32t^2} + \mathcal{O}(t^{-3}(\log t)^6). \end{aligned}$$

That is $K(x, t) = L_1(x, t) + \mathcal{O}(t^{-3}(\log t)^6)$ and therefore (19) can be written as

$$\rho_0(t+x)e^{i\theta(t+x)} = \rho_0(t)e^{i\theta(t)} \left(\frac{t}{2\pi}\right)^{ix/2} (L_1(x, t) + \mathcal{O}(t^{-3}(\log t)^6)),$$

It follows that the difference $F_2(t) - F_3(t)$ is bounded by

$$F_2(t) - F_3(t) \ll t^{15/4} \cdot t^{-3}(\log t)^6 \int_{-\alpha \log t}^{\alpha \log t} \frac{|\zeta(4+it+ix)|}{7 \cosh(\pi x/7)} dx = \mathcal{O}(t^{3/4}(\log t)^6),$$

since the integral is bounded by a constant. \square

To simplify $L_1(x, t)$ we need to consider again the complete integral.

Lemma 14. *Let*

$$G(t) = F_4(t) = e^{i\theta(t)} \rho_0(t) \int_{-\infty}^{+\infty} L_1(x, t) \left(\frac{t}{2\pi}\right)^{ix/2} \frac{\zeta(4+it+ix)}{7 \cosh(\pi x/7)} dx,$$

Then $F_3(t) - F_4(t) = \mathcal{O}(t^{-5/4} \log t)$.

Proof. We need only to bound the integrals extended to the intervals $(-\infty, -\alpha \log t)$ and $(\alpha \log t, +\infty)$. Since $|\zeta(4+ix)| \leq \zeta(4)$ and $\rho_0(t) \ll t^{15/4}$ we have for the first integral

$$I_1 := e^{i\theta(t)} \rho_0(t) \int_{-\infty}^{-\alpha \log t} L_1(x, t) \left(\frac{t}{2\pi}\right)^{ix/2} \frac{\zeta(4+it+ix)}{7 \cosh(\pi x/7)} dx \ll t^{15/4} \int_{-\infty}^{-\alpha \log t} \frac{|L_1(x, t)|}{\cosh(\pi x/7)} dx.$$

The definition (16) of $L_1(x, t)$ we have for x in the integration interval and $t > 0$ large

$$|L_1(x, t)| \ll 1 + \frac{x^2}{t} + \frac{x^4}{t^2}.$$

Therefore,

$$I_1 \ll t^{15/4} \int_{\alpha \log t}^{\infty} e^{-\pi x/7} dx + t^{11/4} \int_{\alpha \log t}^{\infty} x^2 e^{-\pi x/7} dx + t^{7/4} \int_{\alpha \log t}^{\infty} x^4 e^{-\pi x/7} dx.$$

Applying Gabcke's Lemma 10

$$I_1 \ll t^{-5/4} \log t + t^{-9/4} (\log t)^3 + t^{-13/4} (\log t)^5 \ll t^{-5/4} \log t.$$

The second integral is

$$I_2 := e^{i\theta(t)} \rho_0(t) \int_{\alpha \log t}^{\infty} L_1(x, t) \left(\frac{t}{2\pi} \right)^{ix/2} \frac{\zeta(4 + it + ix)}{7 \cosh(\pi x/7)} dx \ll t^{15/4} \int_{\alpha \log t}^{\infty} \frac{|L_1(x, t)|}{\cosh(\pi x/7)} dx,$$

which is bounded in the same way as I_1 . \square

4. SECOND APPROXIMATION TO $Z(t)$.

Proposition 15. *For all nonnegative integers n , we have*

$$(20) \quad \frac{d^n}{dy^n} \frac{1}{\cosh y} = (-1)^n \frac{2e^{-y} B_n(-e^{-2y})}{(e^{-2y} + 1)^{n+1}}.$$

where $B_n(x)$ are polynomials of degree n .

Proof. For $n = 0$ we have

$$\frac{1}{\cosh y} = \frac{2}{e^y + e^{-y}} = \frac{2e^{-y}}{1 + e^{-2y}},$$

so that $B_0(x) = 1$. By induction, we easily show that $B_n(x)$ satisfies the recurrence

$$B_n(x) = 2x(1-x)B'_{n-1}(x) + (1 + (2n-1)x)B_{n-1}(x).$$

From this equation it can be shown that they are polynomials of degree n . We will only need to know B_0 , B_1 and B_2 . The first few are

$$\begin{aligned} B_0(x) &= 1, & B_3(x) &= x^3 + 23x^2 + 23x + 1, \\ B_1(x) &= x + 1, & B_4(x) &= x^4 + 76x^3 + 230x^2 + 76x + 1, \\ B_2(x) &= x^2 + 6x + 1, & B_5(x) &= x^5 + 237x^4 + 1682x^3 + 1682x^2 + 237x + 1. \end{aligned} \quad \square$$

The coefficient of these polynomials appears in Oeis [A060187](#). These polynomials are known as *type B Eulerian polynomials* [6]. They have the following generating functions [5, eq. (13.3), (13.6)]

$$\sum_{j \geq 0} x^j (2j+1)^n = \frac{B_n(x)}{(1-x)^{n+1}}, \quad \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!} = \frac{(1-x)e^{(1-x)t}}{1 - xe^{2(1-x)t}}.$$

We will use the following integral

Proposition 16. *For n a nonnegative integer, α and t real numbers we have*

$$(21) \quad \int_{-\infty}^{\infty} \frac{e^{i\alpha x} x^n}{7 \cosh \frac{\pi x}{7}} dx = \frac{2e^{-y} B_n(-e^{-2y})}{(1 + e^{-2y})^{n+1}} \left(\frac{7i}{2} \right)^n, \quad \text{with } y = \frac{7\alpha}{2}.$$

Proof. It is known ([11, p. 81]) that for $y \in \mathbf{R}$

$$\int_{-\infty}^{\infty} \frac{e^{2\pi i y x}}{\cosh \pi x} dx = \frac{1}{\cosh \pi y}, \quad \text{so that} \quad \int_{-\infty}^{\infty} \frac{e^{2iyx}}{\cosh \pi x} dx = \frac{1}{\cosh y}.$$

Differentiating, we get

$$(2i)^n \int_{-\infty}^{\infty} \frac{x^n e^{2iyx}}{\cosh \pi x} dx = \frac{d^n}{dy^n} \frac{1}{\cosh y} = (-1)^n \frac{2e^{-y} B_n(-e^{-2y})}{(e^{-2y} + 1)^{n+1}}.$$

Therefore, our integral can be calculated, first changing variables $x \mapsto 7x$ and then taking $y = \frac{7\alpha}{2}$

$$\int_{-\infty}^{\infty} \frac{e^{i\alpha x} x^n}{7 \cosh \frac{\pi x}{7}} dx = 7^n \int_{-\infty}^{\infty} \frac{e^{i7\alpha x} x^n}{\cosh \pi x} dx = \left(-\frac{7}{2i}\right)^n \frac{2e^{-y} B_n(-e^{-2y})}{(e^{-2y} + 1)^{n+1}}. \quad \square$$

We now define the auxiliary functions $H_r(t)$.

Definition 17. For t real and r a nonnegative integer we define

$$H_r(t) = \int_{-\infty}^{\infty} e^{i\beta x} \frac{x^r \zeta(4 + it + ix)}{7 \cosh(\pi x/7)} dx, \quad \beta = \frac{1}{2} \log \frac{t}{2\pi},$$

Then the definition of $F_4(t) = G(t)$ implies that

$$(22) \quad G(t) = e^{i\theta(t)} \rho_0(t) \cdot \left(H_0(t) + \frac{15H_1(t)}{4t} + \frac{iH_2(t)}{4t} + \frac{165H_2(t)}{32t^2} + \frac{241iH_1(t)}{24t^2} + \frac{41iH_3(t)}{48t^2} - \frac{H_4(t)}{32t^2} \right).$$

Proposition 18. When $t \rightarrow +\infty$ and a fixed nonnegative integer r we have $H_r(t) = \mathcal{O}(t^{-3/2})$.

Proof. Since the Dirichlet series for $\zeta(4+it)$ converges uniformly in \mathbf{R} , we may interchange sum and integral, so that applying Proposition 16 we obtain

$$H_r(t) = \sum_{n=1}^{\infty} \frac{1}{n^{4+it}} \int_{-\infty}^{\infty} e^{i(\beta - \log n)x} \frac{x^r}{7 \cosh \frac{\pi x}{7}} dx = \left(\frac{7i}{2}\right)^r \sum_{n=1}^{\infty} \frac{1}{n^{4+it}} \frac{2e^{-y_n} B_r(-e^{-2y_n})}{(1 + e^{-2y_n})^{r+1}},$$

where

$$y_n = \frac{7}{2}(\beta - \log n) = \frac{7}{4} \log \frac{t}{2\pi n^2}, \quad e^{-y_n} = \left(\frac{2\pi n^2}{t}\right)^{7/4}.$$

Since $B_r(x)$ is a polynomial of degree r . There is a constant C_r such that

$$B_r(-x) \leq C_r(1+x)^r, \quad x \geq 0.$$

Then

$$\begin{aligned} |H_r(t)| &= \left| \left(\frac{7}{2i}\right)^r \sum_{n=1}^{\infty} \frac{1}{n^{4+it}} \frac{2e^{-y_n} B_r(-e^{-2y_n})}{(1 + e^{-2y_n})^{r+1}} \right| \leq C_r \left(\frac{7}{2}\right)^r \sum_{n=1}^{\infty} \frac{1}{n^4} \frac{2e^{-y_n}}{(1 + e^{-2y_n})^{r+1}} \\ &\leq C_r \left(\frac{7}{2}\right)^r \sum_{n=1}^{\infty} \frac{1}{n^4} \frac{2}{\left(\frac{2\pi n^2}{t}\right)^{7/4} + \left(\frac{2\pi n^2}{t}\right)^{-7/4}} \\ &= C_r \left(\frac{7}{2}\right)^r \sum_{n=1}^{\infty} \frac{2}{\left(\frac{2\pi}{t}\right)^{7/4} n^{15/2} + \left(\frac{2\pi}{t}\right)^{-7/4} n^{1/2}} \\ &\leq C_r \left(\frac{7}{2}\right)^r \int_0^{\infty} \frac{2 dx}{\left(\frac{2\pi}{t}\right)^{7/4} x^{15/2} + \left(\frac{2\pi}{t}\right)^{-7/4} x^{1/2}}. \end{aligned}$$

Change variables $x = \left(\frac{t}{2\pi}\right)^{1/2} y$ and we obtain

$$|H_r(t)| \leq C_r \left(\frac{7}{2}\right)^r \left(\frac{2\pi}{t}\right)^{3/2} \int_0^{\infty} \frac{2dx}{y^{15/2} + y^{1/2}} \ll t^{-3/2}. \quad \square$$

Theorem 19. For $t > 0$ let

$$(23) \quad H(t) = H_0(t) = \int_{-\infty}^{\infty} \left(\frac{t}{2\pi}\right)^{ix/2} \frac{\zeta(4+it+ix)}{7 \cosh(\pi x/7)} dx.$$

Then we have

$$(24) \quad Z(t) = \left(\frac{t}{2\pi}\right)^{\frac{7}{4}} \operatorname{Re}\{e^{i\theta(t)} H(t)\} + \mathcal{O}(t^{-3/4}).$$

Proof. By (22) we may write (15), as

$$Z(t) = \frac{\operatorname{Re}\{e^{i\theta(t)} \rho_0(t) H_0(t)\}}{\left(\left(\frac{1}{4} + t^2\right)\left(\frac{25}{4} + t^2\right)\right)^{1/2}} + \sum_{j=1}^6 T_j + \mathcal{O}(t^{-5/4} \log^6 t),$$

where each T_j is the real part of is of the form

$$T_j = C_j e^{i\theta(t)} \rho_0(t) \left(\left(\frac{1}{4} + t^2\right)\left(\frac{25}{4} + t^2\right)\right)^{-1/2} H_r(t) t^{-k}$$

where C_j is a constant for each j , $r \in \{1, 2, 3\}$ and $k \in \{1, 2\}$. Hence by Proposition 18

$$|T_j| \ll t^{2+\frac{7}{4}} t^{-2} t^{-3/2} t^{-k} = t^{\frac{1}{4}-k} \ll t^{-3/4}.$$

Since

$$\begin{aligned} \frac{\rho_0(t)}{\left(\left(\frac{1}{4} + t^2\right)\left(\frac{25}{4} + t^2\right)\right)^{1/2}} &= (2\pi)^{-7/4} t^{2+\frac{7}{4}} \left(1 + \frac{19}{t^2} + \mathcal{O}(t^{-4})\right) \left(\frac{1}{t^2} - \frac{13}{4t^4} + \mathcal{O}(t^{-6})\right) \\ &= \left(\frac{t}{2\pi}\right)^{\frac{7}{4}} + \mathcal{O}(t^{-\frac{1}{4}}), \end{aligned}$$

we have,

$$Z(t) = \left(\frac{t}{2\pi}\right)^{\frac{7}{4}} \operatorname{Re}\{e^{i\theta(t)} H(t)\} + \mathcal{O}(t^{-\frac{1}{4}} H_0(t)) + \mathcal{O}(t^{-3/4}).$$

Since $\mathcal{O}(t^{-\frac{1}{4}} H_0(t)) = \mathcal{O}(t^{-7/4})$ the result follows. \square

To give an idea of the value of this approximation, I give a table of values.

t	$Z(t)$	$\operatorname{Re}\{(t/2\pi)^{7/4} e^{i\theta(t)} H(t)\}$	t	$Z(t)$	$\operatorname{Re}\{(t/2\pi)^{7/4} e^{i\theta(t)} H(t)\}$
10	-1.5491945	-0.9983260	10^5	5.8795925	5.8790158
10^2	2.6926971	2.6269297	10^6	-2.8061339	-2.8061012
10^3	0.9977946	0.9849027	10^7	14.3525504	14.3525613
10^4	-0.3413947	-0.3452059	10^8	3.6454079	3.6454066

Proposition 20. For $t > 0$ we have

$$H(t) = \sum_{n=1}^{\infty} \frac{1}{n^{4+it}} \frac{2}{\left(\frac{t}{2\pi n^2}\right)^{7/4} + \left(\frac{t}{2\pi n^2}\right)^{-7/4}}.$$

Proof. In the integral (23) defining $H(t)$ we expand $\zeta(4 + it + ix)$ in series and integrate term by term. Then we apply (21).

$$\begin{aligned} H(t) &= \int_{-\infty}^{\infty} \left(\frac{t}{2\pi}\right)^{ix/2} \frac{\zeta(4 + it + ix)}{7 \cosh(\pi x/7)} dx = \sum_{n=1}^{\infty} \frac{1}{n^{4+it}} \int_{-\infty}^{\infty} \left(\frac{t}{2\pi n^2}\right)^{ix/2} \frac{1}{7 \cosh(\pi x/7)} dx \\ &= \sum_{n=1}^{\infty} \frac{1}{n^{4+it}} \frac{2}{\left(\frac{t}{2\pi n^2}\right)^{7/4} + \left(\frac{t}{2\pi n^2}\right)^{-7/4}}. \end{aligned} \quad \square$$

Remark 21. The above expression can be written as

$$H(t) = \left(\frac{t}{2\pi}\right)^{-7/4} \sum_{n=1}^{\infty} \frac{1}{n^{1/2+it}} \frac{2}{1 + \left(\frac{t}{2\pi n^2}\right)^{-7/2}}.$$

Remark 22. It appears that the phase of $H(t)$ decreases moderately with increasing t . For example, in the figure, we see that $\arg H(t)$ for $t \in (10000, 10020)$ start in the fourth quadrant and purely imaginary (or almost there), pass to the third quadrant, first quadrant and then




FIGURE 1. x-ray of $H(z)$ for $z \in (10000, 10020) \times (-2, 4)$

move between 1st and 4th quadrant (with two slight changes into 2nd and 3rd) ending in the fourth quadrant. Therefore, we pass from $-\pi/2$ to some point in the interval $(-2\pi, -2\pi - \pi/2)$. So the variation of the argument is at most -2π . In this rectangle there is only one zero of $H(z)$ with $\text{Im } z < 0$.

Remark 23. The computation of the continuous argument of $H(t)$ for $0 < t < 15000$ is consistent with

$$0 \leq -\arg H(t) \approx c \left(\frac{t}{2} \log \frac{t}{2\pi} - \frac{t}{2} \right), \quad c = \frac{17}{63} \approx 0.27.$$

If this continues to be true for larger values of t , it will imply 73% of zeros of zeta on the critical line.

APPENDIX. COUNTING ZEROS BY MEANS OF THE PHASE.

We show here how finding real analytic functions $F(t)$ such that $Z(t) = \text{Re } F(t)$ is relevant to counting zeros on the critical line.

Let $f: [a, b] \rightarrow \mathbf{C}$. We say that it is *real analytic* if f has an analytic extension to a neighborhood of $[a, b]$ in \mathbf{C} . Notice that f real analytic implies that $g(t) = \overline{f(t)} = \overline{f(\bar{t})}$ is real analytic also, so that

$$\operatorname{Re} f(t) = \frac{f(t) + g(t)}{2}, \quad \operatorname{Im} f(t) = \frac{f(t) - g(t)}{2i},$$

are also real analytic functions in $[a, b]$.

First we extend a Proposition from [1, p. 218]

Proposition 24. *If $f: [a, b] \rightarrow \mathbf{C}$ is real analytic not identically 0, then there are two real analytic functions $U: [a, b] \rightarrow \mathbf{R}$ and $\varphi: [a, b] \rightarrow \mathbf{R}$ such that*

$$f(t) = U(t)e^{i\varphi(t)}.$$

Given two such representations $f = U_1e^{i\varphi_1}$ and $f = U_2e^{i\varphi_2}$, we have either $U_1 = U_2$ and $\varphi_1 - \varphi_2 = 2k\pi$ or $U_1 = -U_2$ and $\varphi_1 - \varphi_2 = (2k + 1)\pi$ for some integer k .

Proof. If $f(t) \neq 0$ there is a simply connected neighborhood Ω of $[a, b]$ where f extends as an analytic function that is never 0 in Ω . Then we may define an analytic function $g: \Omega \rightarrow \mathbf{C}$ such that $e^{g(t)} = f(t)$. Then we may take $U(t) = e^{\operatorname{Re} g(t)}$ and $\alpha(t) = \operatorname{Im} g(t)$.

If f have zeros in $[a, b]$, they are a finite number (counting multiplicities). Let α_j , with $1 \leq j \leq n$ these zeros. Let $F(t) = f(t) \prod_{j=1}^n (t - \alpha_j)^{-1}$. Then F is real analytic and does not vanish, so it can be written as $F(t) = V(t)e^{i\alpha(t)}$. Putting $U(t) = V(t) \prod_j (t - \alpha_j)$ we obtain our desired representation.

Given two representations $f = U_1e^{i\varphi_1} = U_2e^{i\varphi_2}$, U_2 is not identically 0, so we will have $U_1/U_2 = e^{i(\varphi_2 - \varphi_1)}$. Therefore U_1/U_2 is real analytic, take only real values, and have modulus = 1. So $U_1/U_2 = 1$ or it is equal to -1 . Then the phases have to take the given values. \square

Given such a representation $f = Ue^{i\varphi}$ we say that φ is a phase of f . When $f(t)$ do not vanish in $[a, b]$ we can take $U(t) = |f(t)|$ and then for any $a \leq t \leq b$, the value of $\varphi(t)$ coincides with some argument of $f(t)$.

Given a real analytic function $f: [a, b] \rightarrow \mathbf{C}$, not identically 0, we define $N_f[a, b]$ as the number of zeros of f in the closed segment $[a, b]$ counting multiplicities.

Proposition 25. *Let $f: [a, b] \rightarrow \mathbf{R}$ and $g: [a, b] \rightarrow \mathbf{C}$ be real analytic functions with $f(t) = \operatorname{Re} g(t)$ not identically 0. Let $\varphi(t)$ a phase of $g(t)$. Then*

$$\frac{|\varphi(b) - \varphi(a)|}{\pi} < N_f(a, b) + 1.$$

Proof. Substituting g by \bar{g} that is also real analytic and has the same real part, we may assume that $\varphi(a) \leq \varphi(b)$.

We have something to prove only in the case $\varphi(b) - \varphi(a) > \pi$. Consider the odd integer multiples of $\pi/2$ contained between $\varphi(a)$ and $\varphi(b)$. Since $\varphi(b) - \varphi(a) > \pi$ there is at least one. Therefore, there exist $n \leq m$ integers such that

$$(2n - 1)\frac{\pi}{2} < \varphi(a) \leq (2n + 1)\frac{\pi}{2} < (2n + 3)\frac{\pi}{2} < \dots < (2m + 1)\frac{\pi}{2} \leq \varphi(b) < (2m + 3)\frac{\pi}{2}.$$

Since $\varphi(t)$ is continuous, it takes all values in the interval $[\varphi(a), \varphi(b)]$. Therefore, there are points $t_k \in [a, b]$ with $\varphi(t_k) = (2k + 1)\frac{\pi}{2}$ for $n \leq k \leq m$. At these points, we have

$f(t_k) = \operatorname{Re} g(t_k) = 0$. That is $N_f(a, b) \geq m - n + 1$. The interval $((2n - 1)\pi/2, (2m + 3)\pi/2) \supset [\varphi(a), \varphi(b)]$, comparing its lengths, yields

$$\pi(m - n) + 2\pi > \varphi(b) - \varphi(a).$$

The result follows. \square

Proposition 26. *Let f and $g: [a, b] \rightarrow \mathbf{C}$ real analytic functions such that*

$$(25) \quad |f(t) - g(t)| < |f(t)|, \quad \text{for all } a \leq t \leq b.$$

Then f and g do not vanish on $[a, b]$, so that we can write $f(t) = |f(t)|e^{i\alpha(t)}$ and $g(t) = |g(t)|e^{i\beta(t)}$ with α and β real analytic. Then we can choose $\beta(t)$ so that

$$|\alpha(t) - \beta(t)| < \pi, \quad a \leq t \leq b.$$

In particular

$$\left| \frac{\alpha(b) - \alpha(a)}{\pi} - \frac{\beta(b) - \beta(a)}{\pi} \right| < 2.$$

Proof. If $f(t) = 0$, equation (25) implies $|g(t)| < 0$ which is a contradiction. If $g(t) = 0$, then we arrive at the contradiction $|f(t)| < |f(t)|$. Since the functions do not vanish, we may apply Proposition 25 to get $f(t) = |f(t)|e^{i\alpha(t)}$ and $g(t) = |g(t)|e^{i\beta(t)}$. The function $\beta(t)$ is not unique, in fact $\beta(t) + 2k\pi$ for any integer k can be used instead of $\beta(t)$.

We have $|f(0) - g(0)| < |f(0)|$. Therefore $g(0)$ is in the open disc with center $f(0)$ and radius the distance from $f(0)$ to 0. $\alpha(0)$ is one of the determination of $\arg f(0)$, there is a value of $\arg g(0)$ such that $|\alpha(0) - \arg g(0)| < \pi$, since the two points are in some half plane. We may choose a k so that $|\beta(0) + 2k\pi - \alpha(0)| < \pi$, without loss of generality we may assume that $|\beta(0) - \alpha(0)| < \pi$. Since these functions are continuous, this will be true for some interval $|\beta(t) - \alpha(t)| < \pi$ if $a \leq t < a + \varepsilon$.

By contradiction we shall prove that $|\beta(t) - \alpha(t)| < \pi$ for all $t \in [a, b]$. Assume that there is some $|\beta(t) - \alpha(t)| \geq \pi$. The set of points that satisfies this inequality is a closed set of $[a, b]$ contained in $[a + \varepsilon, b]$. Let t_0 be the infimum of these closed sets. By continuity we will have $|\beta(t_0) - \alpha(t_0)| = \pi$. Therefore $f(t_0) = |f(t_0)|e^{i\alpha(t_0)}$ and

$$g(t_0) = |g(t_0)|e^{i\beta(t_0)} = |g(t_0)|e^{i\alpha(t_0) \pm \pi} = -|g(t_0)|e^{i\alpha(t_0)}.$$

And we have a contradiction with (25) because

$$|f(t_0)| > |f(t_0) - g(t_0)| = \left| |f(t_0)|e^{i\alpha(t_0)} + |g(t_0)|e^{i\alpha(t_0)} \right| = |f(t_0)| + |g(t_0)|.$$

The rest is an easy consequence of this. \square

REFERENCES

- [1] J. ARIAS DE REYNA & J. VAN DE LUNE, *On the exact location of the nontrivial zeros of Riemann's zeta function*, Acta Arith. **163** (2014) 215–245. DOI: [10.4064/aa163-3-3](https://doi.org/10.4064/aa163-3-3).
- [2] J. ARIAS DE REYNA, *Approximate formula for $Z(t)$* , arXiv:2406.18150.
- [3] W. GABCKE, *Neue Herleitung und explizite Restabschätzung der Riemann-Siegel Formel*, Ph.D. thesis, Georg-August-Universität zu Göttingen, 1979.
- [4] G. H. HARDY, A. E. INGHAM, G. PÓLYA, *Theorems concerning mean values of analytic functions*, Proc. Roy. Soc. London, **113** (1927) 542–569.
- [5] J. HORVÁTH, *Topological Vector Spaces I*, Addison-Wesley, Reading, Massachusetts, 1966.
- [6] T. K. PETERSEN, *Eulerian numbers*, Birkhäuser/Springer, New York, 2015.
- [7] H. RADEMACHER, *Topics in analytic number theory*, Edited by E. Grosswald, J. Lehner and M. Newman; Springer, New York-Heidelberg, 1973.
- [8] B. RIEMANN, *Gesammelte Mathematische Werke, Wissenschaftlicher Nachlass und Nachträge—Collected Papers*, Ed. Narasimham, R., Springer, Berlin-Heidelberg, 1990.

- [9] W. RUDIN, *Principles of Mathematical Analysis*, McGraw-Hill, 1976.
- [10] C. L. SIEGEL, *Über Riemann Nachlaß zur analytischen Zahlentheorie*, Quellen und Studien zur Geschichte der Mathematik Astronomie und Physik **2** (1932) 45–80. (Reprinted in [8] pp. 770–805.)
- [11] E. M. STEIN, R. SHAKARCHI, *Complex Analysis*, Princeton University Press, Princeton, NJ, 2003.

UNIVERSIDAD DE SEVILLA, FACULTAD DE MATEMÁTICAS, C/TARFIA, SN, 41012-SEVILLA, SPAIN.
Email address: arias@us.es, ariasdereyna1947@gmail.com