

# Multiple solutions for a class of nonhomogeneous elliptic systems with Dirichlet boundary or Neumann boundary

Xiaoli Yu<sup>1</sup>, Xingyong Zhang<sup>1,2\*</sup>

<sup>1</sup>Faculty of Science, Kunming University of Science and Technology, Kunming, Yunnan, 650500, P.R. China.

Kunming, Yunnan, 650500, P.R. China.

<sup>2</sup>Research Center for Mathematics and Interdisciplinary Sciences, Kunming University of Science and Technology,

Kunming, Yunnan, 650500, P.R. China.

**Abstract:** In this paper, we mainly establish the existence of at least three non-trivial solutions for a class of nonhomogeneous quasilinear elliptic systems with Dirichlet boundary value or Neumann boundary value in a bounded domain  $\Omega \subset \mathbb{R}^N$  and  $N \geq 1$ . We exploit the method which is based on [6]. This method let us obtain the concrete open interval about the parameter  $\lambda$ . Since the quasilinear term depends on  $u$  and  $\nabla u$ , it is necessary for our proofs to use the theory of monotone operators and the skill of adding one dimension to space.

**Keywords:** three non-trivial solutions; concrete open interval; Dirichlet boundary value; Neumann boundary value; quasilinear scalar field systems.

**2020 Mathematics Subject Classification.** 35J20; 35J50; 35J62.

## 1 Introduction and main results

Stuart [30] established the result of at least two non-negative weak solutions for the following nonhomogeneous problem:

$$\begin{cases} -\operatorname{div} \left( \phi \left( \frac{u^2 + |\nabla u|^2}{2} \right) \nabla u \right) + \phi \left( \frac{u^2 + |\nabla u|^2}{2} \right) u = \lambda u + h, & x \in \Omega, \\ u(x) = 0, & x \in \partial\Omega, \end{cases} \quad (1.1)$$

where  $h \in L^2(\Omega)$ ,  $h \geq 0$ ,  $\Omega$  is a bounded domain of  $\mathbb{R}^N$ ,  $N \geq 1$ ,  $\lambda \in \mathbb{R}$ ,  $\phi \in C([0, +\infty), \mathbb{R})$ ,  $\Phi(s) = \int_0^s \phi(\tau) d\tau$  and satisfies

( $\Phi_1$ )  $\Phi(0) = 0$ ,  $\phi = \Phi'$  is non-increasing on  $[0, +\infty)$  and  $\phi(\infty) = \lim_{s \rightarrow \infty} \phi(s) > 0$ ;

( $\Phi_2$ ) putting  $\Phi(s^2) = g(s)$ , there exists  $\rho > 0$  such that

$$g(t) \geq g(s) + g'(s)(t - s) + \rho(t - s)^2, \quad \text{for all } s, t \geq 0;$$

( $\Phi_3$ )  $\lim_{s \rightarrow \infty} (\Phi(s) - \Phi'(s)s) < +\infty$ .

( $\Phi_2$ ) is used to ensure the ellipticity of (1.1). More specifically, when  $\phi(\infty) + \phi(\infty)\lambda_1 < \lambda < \phi(0) + \phi(\infty)\lambda_1$ , he obtained results that the equation (1.1) has a trivial solution and another non-negative non-trivial weak solution for  $h \equiv 0$  and if there exists  $H_\lambda$  and  $0 < \|h\|_2 = [\int_\Omega h^2 dx]^{1/2} < H_\lambda$ , the equation (1.1) has at least two distinct non-negative weak solutions. He claimed that in both cases one is a strict local minimum solution and the other is a mountain pass type solution.

In the equation (1.1), the quasilinear term  $\phi$  depends on  $\frac{u^2 + |\nabla u|^2}{2}$  which drives from a nonlinear optical model. The optical model described the propagation about self-trapped transverse magnetic field modes in a cylindrical optical fiber made from a self-focusing dielectric material. For more details, see [26–29].

---

\*Corresponding author, E-mail address: zhangxingyong1@163.com

In recent years, there have been some studies on the particular quasilinear term  $\phi$ . For example, Jeanjean and Rădulescu [14] extended the results in [30]. They studied the following second order quasilinear elliptic equation:

$$\begin{cases} -\operatorname{div} \left( \phi \left( \frac{u^2 + |\nabla u|^2}{2} \right) \nabla u \right) + \phi \left( \frac{u^2 + |\nabla u|^2}{2} \right) u = f(u) + h, & x \in \Omega, \\ u = 0, & x \in \partial\Omega, \end{cases} \quad (1.2)$$

where  $h \in L^2(\Omega)$  is non-negative,  $\Omega$  is a bounded domain of  $\mathbb{R}^N$  with  $N \geq 1$ . They optimized the assumptions about the quasilinear part  $\phi$ . Specifically, they assumed that  $\phi$  satisfies the following conditions:

( $\phi_1$ )  $\phi \in C([0, +\infty), \mathbb{R})$  and there exist some constants  $0 < \phi_{\min} \leq \phi_{\max}$  such that

$$\phi_{\min} \leq \phi(s) \leq \phi_{\max}, \quad s \in [0, +\infty);$$

( $\phi_2$ ) the function  $s \mapsto \Phi(s^2)$  is convex on  $\mathbb{R}$ ,

where the condition ( $\phi_2$ ) ensures the ellipticity of (1.2) and is weaker than ( $\Phi_2$ ). They obtained that the problem (1.2) has a non-negative solution, when  $h \not\equiv 0$  and  $f$  satisfies sublinear growth. For  $h \equiv 0$ , they got the result that the equation (1.2) has a non-trivial solution if  $f$  satisfies sublinear growth. Meanwhile, they also claimed the nonexistence of solutions under some appropriate conditions. They deduced that the problem (1.2) has at least one non-negative solution for  $\|h\|_2$  sufficiently small and the solution is non-trivial if  $h \not\equiv 0$ , when  $f$  satisfies linear growth, and  $\Phi$  satisfies ( $\phi_2$ ) and the following condition,

( $\phi_3$ )  $\phi \in C([0, +\infty), \mathbb{R})$ , there exist some constants  $0 < \phi_{\min} \leq \phi_{\max}$  such that

$$\phi_{\min} \leq \phi(s) \leq \phi_{\max}, \quad s \in [0, +\infty),$$

and there exists  $\phi(\infty) > 0$  such that  $\phi(s) \rightarrow \phi(\infty)$  as  $s \rightarrow \infty$ .

They also claimed that the problem (1.2) has at least two non-trivial non-negative solutions for  $h \not\equiv 0$  and at least a non-trivial non-negative solution for  $h \equiv 0$  when  $\phi$  satisfies ( $\phi_3$ ) and the following condition:

( $\phi_4$ ) the function  $s \mapsto \Phi(s^2)$  is strictly convex on  $\mathbb{R}$ .

What's more, combining the mountain pass theorem and some analytical skills, Pomponio and Watanabe [18] got that the equation (1.2) has a radial ground state solution  $u$  and  $u \in C^{1,\sigma}$  for some  $\sigma \in (0, 1)$ , when  $h \equiv 0$  and the nonlinearity satisfies a variant of Berestycki-Lions' conditions.

In this paper, we first establish the existence of at least three non-trivial solutions for a class of quasilinear elliptic systems with Dirichlet boundary value:

$$\begin{cases} -\operatorname{div} \left( \phi_1 \left( \frac{u^2 + |\nabla u|^2}{2} \right) \nabla u \right) + \phi_1 \left( \frac{u^2 + |\nabla u|^2}{2} \right) u = \lambda G_u(x, u, v), & x \in \Omega, \\ -\operatorname{div} \left( \phi_2 \left( \frac{v^2 + |\nabla v|^2}{2} \right) \nabla v \right) + \phi_2 \left( \frac{v^2 + |\nabla v|^2}{2} \right) v = \lambda G_v(x, u, v), & x \in \Omega, \\ u = v = 0, & x \in \partial\Omega, \end{cases} \quad (1.3)$$

where  $\Omega$  is a non-empty bounded open subset in  $\mathbb{R}^N$ ,  $N \geq 1$ ,  $|\Omega| := \operatorname{vol}(\Omega) = \int_{\Omega} 1 dx < +\infty$ ,  $\lambda$  is a positive constant,  $(u, v) \in W := H_0^1(\Omega) \times H_0^1(\Omega)$  and  $\Phi_i(s) = \int_0^s \phi_i(\tau) d\tau$ ,  $i = 1, 2$ . We assume that  $\phi_i$  satisfies ( $\phi_4$ ) and the following conditions:

( $\phi_1$ )'  $\phi_i \in C([0, +\infty), \mathbb{R})$  and there exist constants  $0 < \phi_i^0 \leq \phi_i^m$ , such that

$$\phi_i^0 \leq \phi_i(s) \leq \phi_i^m, \quad \text{for all } s \in [0, +\infty).$$

And the nonlinearity  $G(x, \cdot, \cdot)$  is  $C^1$  in  $\mathbb{R} \times \mathbb{R}$  for a.e.  $x \in \Omega$  and satisfies  $G(\cdot, s, t)$  is measurable in  $\Omega$  for any  $(s, t) \in \mathbb{R} \times \mathbb{R}$  and the following assumptions:

( $G_1$ ) there exist non-negative constants  $a_1, a_2$  and  $p, q \in [1, 2^*)$  such that

$$|G_s(x, s, t)| \leq a_1 + a_2 |s|^{p-1}, \quad |G_t(x, s, t)| \leq a_1 + a_2 |t|^{q-1}, \quad \forall (x, s, t) \in \Omega \times \mathbb{R} \times \mathbb{R};$$

( $G_2$ )  $G(x, s, t) \geq 0$  for every  $(x, s, t) \in \Omega \times \mathbb{R}^+ \times \mathbb{R}^+$  and  $G(x, 0, 0) = 0$ ;

( $G_3$ ) there exist constants  $0 < \alpha, \beta < 2$  and  $b > 0$  such that

$$G(x, s, t) \leq b(1 + |s|^\alpha + |t|^\beta),$$

for almost every  $x \in \Omega$  and every  $s, t \in \mathbb{R}$ ;  
 $(G_4)$  there exist constants  $\gamma_1 > 0$  and  $\sigma_1 > 0$  with  $\sigma_1 > \gamma_1 \tau$ , such that

$$\frac{\inf_{x \in \Omega} G(x, \sigma_1, \sigma_1)}{\sigma_1^2} > 2a_1 \frac{A_1}{\gamma_1} + a_2 A_2 \gamma_1^{p-2} + a_2 A_3 \gamma_1^{q-2},$$

where

$$\begin{aligned} A_1 &= C_1 \phi_m \left( \frac{1}{\phi_0} \right)^{1/2}, \quad A_2 = \frac{(C_p)^p \phi_m}{p} \left( \frac{2}{\phi_0} \right)^{p/2}, \\ A_3 &= \frac{(C_q)^q \phi_m}{q} \left( \frac{2}{\phi_0} \right)^{q/2}, \quad \tau = \frac{\sigma_1}{\left( \left[ \Phi_1 \left( \frac{\sigma_1^2}{2} \right) + \Phi_2 \left( \frac{\sigma_1^2}{2} \right) \right] \frac{\pi^{N/2}}{\Gamma(1 + N/2)} D^N \right)^{1/2}}, \end{aligned} \quad (1.4)$$

$a_1, a_2$  are given in  $(G_1)$ ,  $\phi_0 = \min\{\phi_1^0, \phi_2^0\}$ ,  $\phi_m = \max\{\phi_1^m, \phi_2^m\}$ ,  $D := \sup_{x \in \Omega} \text{dist}(x, \partial\Omega)$ ,  $C_1, C_p$  and  $C_q$  denote the embedding constants. Specifically, fixing  $p \in [1, 2^*]$ , there exists the embedding  $H_0^1(\Omega) \hookrightarrow L^p(\Omega)$  which is continuous and there exists a positive constant  $C_p$  such that

$$\|u\|_p \leq C_p \|\nabla u\|_2. \quad (1.5)$$

In particular, the embedding  $H_0^1(\Omega) \hookrightarrow L^p(\Omega)$  is compact when  $p \in [1, 2^*)$ .  $H_0^1(\Omega)$  denotes the closure of  $C_0^\infty$  in the Sobolev space  $W^{1,2}(\Omega)$ , equipped with the inner product and norm

$$(u, v) = \int_{\Omega} \nabla u \cdot \nabla v dx, \quad \|u\|_0 = (u, u)^{1/2},$$

where  $W^{1,2}(\Omega)$  is defined as a space of all functions  $u : \Omega \rightarrow \mathbb{R}$  satisfying  $u \in L^2(\Omega)$  and the partial derivative  $D^\alpha u \in L^2(\Omega)$  for  $0 \leq |\alpha| \leq 1$ .  $L^p(\Omega)$  ( $1 \leq p < \infty$ ) denotes the Lebesgue space with the norm

$$\|u\|_p = \left( \int_{\Omega} |u|^p dx \right)^{1/p}.$$

By simple calculation, there exists  $x_0 \in \Omega$  such that the ball  $B(x_0, D) \subseteq \Omega$  and  $\text{meas}(B(x_0, D)) = \frac{\pi^{N/2}}{\Gamma(1 + N/2)} D^N$ .

Presently, the existence of three solutions for p-Laplacian equations and systems has been followed and many excellent results have been obtained. In [21], Ricceri studied a class of elliptic eigenvalue problems:

$$\begin{cases} -\Delta u = \lambda[f(u) + \mu g(u)], & x \in \Omega, \\ u(x) = 0, & x \in \partial\Omega, \end{cases} \quad (1.6)$$

where  $\Omega \subseteq \mathbb{R}^N$  is bounded,  $N \geq 3$ ,  $\lambda, \mu \in \mathbb{R}$ ,  $f, g : \mathbb{R} \rightarrow \mathbb{R}$  are two continuous functions. Combining [20, Theorem 1] and [19, Corollary 1], he obtained the existence of three non-trivial solutions for (1.6) when  $\lambda > 0$  and  $|\mu|$  is small enough. [21, Remarks 5.3 and 5.4] shows that there is a trivial solution of (1.6) for  $g \equiv 0$ . That is, there are only two non-trivial solutions for  $g \equiv 0$ . Then, Ricceri improved the result of [21, Theorem 3.1] to [22, Theorem 1] which shows that the conclusion of [21, Theorem 3.1] holds for each  $\lambda$  in an open sub-interval of an interval  $I \subseteq \mathbb{R}$  by replacing the convexity of the operator term with sequential weak lower semicontinuity. Under the additional assumption that the operator term is bounded on each bounded set of  $\Omega$ , Ricceri [23] reached the existence of three solutions for each  $\lambda$  in a non-empty open interval  $A$  and  $A \subseteq I \subseteq \mathbb{R}$ . However, neither [22] or [23] gave the further information on the size and location of the set of  $A$ . Hereafter, a new three critical points theorem was established in [24], which ensures that the conclusion of Theorem 1 in [23] actually holds for each compact interval  $A$  contained in  $(a, +\infty)$ , where  $a$  is given, please refer to [24] for details. Three critical points theorems for non-differentiable functionals were established in [4], when the functions  $f, g$  may be discontinuous on  $u$ , which extended the results previously in [22] from differentiable functionals to non-smooth functionals and determined a precise interval of parameter  $\lambda$ .

In [3, Theorem 3.5 and Theorem 3.6 ], Bonanno also got a well-determined interval of parameter  $\lambda$  in the smooth and the non-smooth framework for the functionals which have at least three critical points under weaker regularity and compactness conditions. In [6], the existence of three non-trivial solutions for non-autonomous elliptic Dirichlet problems without any small perturbation of the nonlinearity is presented:

$$\begin{cases} -\Delta u = \lambda f(x, u), & x \in \Omega, \\ u(x) = 0, & x \in \partial\Omega, \end{cases} \quad (1.7)$$

where  $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  is a function and satisfies the following assumptions:  
 $(F_1)$  there exist two non-negative constants  $a_1, a_2$  and  $q \in [1, 2^*)$  such that

$$|f(x, t)| \leq a_1 + a_2|t|^{q-1}, \quad \forall (x, t) \in \Omega \times \mathbb{R};$$

$(F_2)$   $F(x, \xi) \geq 0$  for every  $(x, \xi) \in \Omega \times \mathbb{R}^+$ ;  
 $(F_3)$  there exist constants  $0 < s < 2$  and  $b > 0$  such that

$$F(x, \xi) \leq b(1 + |\xi|^s),$$

for almost every  $x \in \Omega$  and every  $\xi \in \mathbb{R}$ ;

$(F_4)$  there exist constants  $\gamma > 0$  and  $\delta > 0$  with  $\delta > \gamma\kappa$ , such that

$$\frac{\inf_{x \in \Omega} F(x, \delta)}{\delta^2} > a_1 \frac{K_1}{\gamma} + a_2 K_2 \gamma^{q-2},$$

where  $a_1, a_2$  are given in  $(F_1)$  and  $\kappa, K_1$  and  $K_2$  are given in equation (10) of [6].

They reached that the equation (1.7) has at least three non-trivial solutions in  $H_0^1(\Omega)$  for each parameter  $\lambda$  belonging to

$$\Lambda_{\gamma, \delta} := \left( \frac{2(2^N - 1)}{D^2} \frac{\delta^2}{\inf_{x \in \Omega} F(x, \delta)}, \frac{2(2^N - 1)}{D^2} \frac{1}{a_1 K_1 / \gamma + a_2 K_2 \gamma^{q-2}} \right).$$

The method they used is based on a recent three critical points theorem [3, Theorem 3.6] for differentiable functional, which can be rewritten as [6, Theorem 2.1].

**Theorem I.** ([6, Theorem 2.1]) Let  $X$  be a reflexive real Banach space,  $\Phi : X \rightarrow \mathbb{R}$  be a coercive, continuously Gâteaux differentiable and sequentially weakly lower semicontinuous functional whose Gâteaux derivative admits a continuous inverse on  $X^*$ ,  $\Psi : X \rightarrow \mathbb{R}$  be a continuously Gâteaux differentiable functional whose Gâteaux derivative is compact such that

$$\Phi(0) = \Psi(0) = 0.$$

Assume that there exist  $r > 0$  and  $\bar{x} \in X$ , with  $r < \Phi(\bar{x})$ , such that:

$$(a_1) \frac{\sup_{\Phi(x) \leq r} \Psi(x)}{r} < \frac{\Psi(\bar{x})}{\Phi(\bar{x})};$$

$$(a_2) \text{ for each } \lambda \in \Lambda_\lambda := \left( \frac{\Phi(\bar{x})}{\Psi(\bar{x})}, \frac{r}{\sup_{\Phi(x) \leq r} \Psi(x)} \right), \text{ the functional } \Phi - \lambda\Psi \text{ is coercive.}$$

Then, for all  $\lambda \in \Lambda_\lambda$  the functional  $\Phi - \lambda\Psi$  has at least three distinct critical points in  $X$ .

Moreover, using variational methods and three non-trivial critical points theorem in [3, Theorem 3.6] or a more precise version of [6, Theorem 2.1], Bonanno-Heidarkhani-O'regan proved the existence of three non-zero solutions for a gradient nonlinear Dirichlet elliptic system driven by a  $(p, q)$ -Laplacian operator in [5], when  $\Omega \subseteq \mathbb{R}^N$  is bounded and  $p, q > N \geq 1$ . In [1], Averna and Bonanno investigated the existence of three non-trivial weak solutions for the Dirichlet elliptic system in [5], when  $N \geq 3, 1 < q \leq p < N$  and  $f$  satisfies some appropriate assumptions. The results in [1] determine an appropriate interval about the parameter  $\lambda$  which improve the conclusions in [2], [5] and [15].

Meanwhile, the existence of three solutions for the Neumann problem is also followed. In [9], applying Theorem 1 of [22], Bonanno and Candito proved the existence of three non-trivial weak solutions for the Neumann problem:

$$\begin{cases} -\Delta_p u + a(x)|u|^{p-2}u = \lambda f(x, u), & x \in \Omega, \\ \frac{\partial u}{\partial \nu} = 0, & x \in \partial\Omega, \end{cases} \quad (1.8)$$

where  $\Omega \subseteq \mathbb{R}^N$  is bounded,  $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ ,  $\lambda \in \mathbb{R}_+$ ,  $p > N \geq 1$ ,  $a \in L^\infty(\Omega)$ ,  $\text{ess inf}_{x \in \Omega} a(x) > 0$  and  $\nu$  is the outward unit normal to  $\partial\Omega$ . In [12], applying the result which is obtained in [3] and a more precise version of [4, Theorem 3.2], D'Agui and Bisci proved the existence of at least three non-zero solutions for the problem of (1.8) and determined a precise interval of values of the parameter  $\lambda$ , when  $\Omega \subseteq \mathbb{R}^N$  is bounded,  $N \geq p > 1$ . Using [4, Theorem 3.2], Bonanno-Bisci-Rădulescu [8] studied the following quasilinear elliptic Neumann problem:

$$\begin{cases} -\text{div}(\alpha(|\nabla u|)\nabla u) + \alpha(|u|)u = \lambda f(x, u), & x \in \Omega, \\ \frac{\partial u}{\partial \nu} = 0, & x \in \partial\Omega, \end{cases} \quad (1.9)$$

where  $\Omega \subseteq \mathbb{R}^N$  is bounded,  $N \geq 3$ , and

$$\varphi(t) := \begin{cases} \alpha(|t|)t, & \text{for } t \neq 0, \\ 0, & \text{for } t = 0 \end{cases}$$

They obtained that the Neumann problem of (1.9) has at least three nontrivial solutions, when  $p_0 > N \geq 3$ , where

$$p_0 := \inf_{t > 0} \frac{t\varphi(t)}{\Phi(t)}, \quad \Phi(t) = \int_0^t \varphi(s)ds.$$

For a class of quasilinear elliptic systems involving the  $p(x)$ -Laplace operator with Neumann boundary condition, the existence of at least three solutions is also established in [33] by using the three critical points theorem in [23].

Inspired by [5, 6, 8, 33], we investigate the following non-autonomous nonhomogeneous quasilinear elliptic Neumann problem:

$$\begin{cases} -\text{div} \left( \phi_1 \left( \frac{u^2 + |\nabla u|^2}{2} \right) \nabla u \right) + \phi_1 \left( \frac{u^2 + |\nabla u|^2}{2} \right) u = \lambda G_u(x, u, v), & x \in \Omega, \\ -\text{div} \left( \phi_2 \left( \frac{v^2 + |\nabla v|^2}{2} \right) \nabla v \right) + \phi_2 \left( \frac{v^2 + |\nabla v|^2}{2} \right) v = \lambda G_v(x, u, v), & x \in \Omega, \\ \frac{\partial u}{\partial \nu} = \frac{\partial v}{\partial \nu} = 0, & x \in \partial\Omega, \end{cases} \quad (1.10)$$

where  $\Omega$  is a non-empty bounded open subset of the Euclidean space  $\mathbb{R}^N$ ,  $N \geq 1$ ,  $\lambda$  is a positive constant,  $(u, v) \in X := H^1(\Omega) \times H^1(\Omega)$ . We mainly study the existence of at least three non-trivial solutions for (1.10), under the assumptions  $(\phi_1)'$ ,  $(\phi_4)$ ,  $(G_1)$ - $(G_3)$  and the following condition:

$(G_4)'$  there exist two positive constants  $\gamma_2$  and  $\sigma_2$ , with  $\sigma_2 > \gamma_2 \bar{\tau}$  such that

$$\begin{aligned} \frac{\inf_{x \in \Omega} G(x, \sigma_2, \sigma_2)}{\sigma_2^2} &> 2a_1 \frac{\bar{A}_1}{\gamma_2} + a_2 \bar{A}_2 \gamma_2^{p-2} + a_2 \bar{A}_3 \gamma_2^{q-2}, \\ \bar{A}_1 &= c_1 \phi_m \left( \frac{2}{\phi_0} \right)^{1/2}, \quad \bar{A}_2 = \frac{(c_p)^p \phi_m}{p} \left( \frac{2}{\phi_0} \right)^{p/2}, \\ \bar{A}_3 &= \frac{(c_q)^q \phi_m}{q} \left( \frac{2}{\phi_0} \right)^{q/2}, \quad \bar{\tau} = \tau = \frac{\sigma_2}{\left( \left[ \Phi_1 \left( \frac{\sigma_2^2}{2} \right) + \Phi_2 \left( \frac{\sigma_2^2}{2} \right) \right] \frac{\pi^{N/2}}{\Gamma(1+N/2)} D^N \right)^{1/2}}, \end{aligned} \quad (1.11)$$

where  $a_1, a_2$  are given in  $(G_1)$ ,  $\phi_0, \phi_m$  are given in  $(\phi_1)'$ ,  $c_1, c_p$  and  $c_q$  are the embedding constants which are related with the embedding  $H^1(\Omega) \hookrightarrow L^p(\Omega)$ . That is, when  $p \in [1, 2^*)$ ,  $H^1(\Omega) \hookrightarrow L^p(\Omega)$  and the following embedding inequality holds,

$$\|u\|_p \leq c_p \|u\|, \quad (1.12)$$

for every  $u \in H^1(\Omega)$ .  $H^1(\Omega)$  denotes the space  $W^{1,2}(\Omega)$  equipped with the norm

$$\|u\| = \left[ \int_{\Omega} (u^2 + |\nabla u|^2) dx \right]^{1/2}.$$

Next, we show our main results.

**Theorem 1.1.** *Assume that  $\phi_i$  ( $i = 1, 2$ ) and  $G$  satisfy  $(\phi_1)'$ ,  $(\phi_4)$  and  $(G_1)$ - $(G_4)$  respectively. Then for  $N \geq 1$  and each parameter  $\lambda$  belonging to*

$$\Lambda_{\gamma_1, \sigma_1} := \left( \frac{\phi_m \sigma_1^2}{\inf_{x \in \Omega} G(x, \sigma_1, \sigma_1)}, \frac{\phi_m}{\left( 2a_1 \frac{A_1}{\gamma_1} + a_2 A_2 \gamma_1^{p-2} + a_2 A_3 \gamma_1^{q-2} \right)} \right),$$

the problem (1.3) possesses at least three weak solutions in  $W$ , where  $A_1$ ,  $A_2$  and  $A_3$  are given in (1.4).

Particularly, for  $N = 1$ , since the embedding  $H_0^1(\Omega) \hookrightarrow C^0(\bar{\Omega})$  is compact, it holds

$$C := \sup_{u \in H_0^1(\Omega) \setminus \{0\}} \frac{\|u\|_\infty}{\|u\|_0} < \infty. \quad (1.13)$$

Without the assumption  $(G_1)$  and replacing  $(G_4)$  with the following condition  $(G_4)''$ ,  $(G_4)''$  there exist constants  $\gamma_1 > 0$  and  $\sigma_1 > 0$  with  $\sigma_1 > \gamma_1 \tau$ , such that

$$\frac{\inf_{x \in \Omega} G(x, \sigma_1, \sigma_1)}{\sigma_1^2} > b(A_{11} \gamma_1^{-2} + A_{21} \gamma_1^{\alpha-2} + A_{31} \gamma_1^{\beta-2}),$$

where

$$\begin{aligned} A_{11} &= \phi_m |\Omega|, & A_{21} &= \phi_m |\Omega| C^\alpha \left( \frac{2}{\phi_0} \right)^{\alpha/2}, \\ A_{31} &= b |\Omega| C^\beta \left( \frac{2}{\phi_0} \right)^{\beta/2}, & \tau &= \frac{\sigma_1}{\left( \left[ \Phi_1 \left( \frac{\sigma_1^2}{2} \right) + \Phi_2 \left( \frac{\sigma_1^2}{2} \right) \right] |\Omega| \right)^{1/2}}, \end{aligned} \quad (1.14)$$

we also get the existence of three non-zero solutions as the following.

**Theorem 1.2.** *Assume that  $\phi_i$  ( $i = 1, 2$ ) and  $G$  satisfy  $(\phi_1)'$ ,  $(\phi_4)$ ,  $(G_2)$ - $(G_3)$  and  $(G_4)''$  respectively. Then for  $N = 1$  and each parameter  $\lambda$  belonging to*

$$\bar{\Lambda}_{\gamma_1, \sigma_1} := \left( \frac{\phi_m \sigma_1^2}{\inf_{x \in \Omega} G(x, \sigma_1, \sigma_1)}, \frac{\phi_m}{b(A_{11} \gamma_1^{-2} + A_{21} \gamma_1^{\alpha-2} + A_{31} \gamma_1^{\beta-2})} \right),$$

the problem (1.3) possesses at least three weak solutions in  $W$ , where  $A_{11}$ ,  $A_{21}$  and  $A_{31}$  are given in (1.14).

For the problem (1.10), by replacing  $(G_4)$  with  $(G_4)'$ , we can reach the following conclusion.

**Theorem 1.3.** *Assume that  $\phi_i$  ( $i = 1, 2$ ) and  $G$  satisfy  $(\phi_1)'$ ,  $(\phi_4)$  and  $(G_1)$ - $(G_3)$  and  $(G_4)'$  respectively. Then for  $N \geq 1$  and each parameter  $\lambda$  belonging to*

$$\Lambda_{\gamma_2, \sigma_2} := \left( \frac{\phi_m \sigma_2^2}{\inf_{x \in \Omega} G(x, \sigma_2, \sigma_2)}, \frac{\phi_m}{\left( 2a_1 \frac{A_1}{\gamma_2} + a_2 \bar{A}_2 \gamma_2^{p-2} + a_2 \bar{A}_3 \gamma_2^{q-2} \right)} \right),$$

the problem (1.10) possesses at least three non-zero solutions in  $X$ , where  $\bar{A}_1$ ,  $\bar{A}_2$  and  $\bar{A}_3$  are given in (1.11).

Furthermore, for  $N = 1$ , the embedding  $H^1(\Omega) \hookrightarrow C^0(\bar{\Omega})$  is compact, and it holds

$$c := \sup_{u \in H^1(\Omega) \setminus \{0\}} \frac{\|u\|_\infty}{\|u\|} < \infty. \quad (1.15)$$

We deduce the existence of three non-zero solutions for the problem (1.10) when  $N = 1$ . The condition  $(G_1)$  is not needed and we replace  $(G_4)'$  with the following condition  $(G_4)'''$ ,  
 $(G_4)'''$  there exist two positive constants  $\gamma_3$  and  $\sigma_3$ , with  $\sigma_3 > \gamma_3 \hat{\tau}$  such that

$$\frac{\inf_{x \in \Omega} G(x, \sigma_3, \sigma_3)}{\sigma_3^2} > b(\hat{A}_1 \gamma_3^{-2} + \hat{A}_2 \gamma_3^{\alpha-2} + \hat{A}_3 \gamma_3^{\beta-2}),$$

where

$$\begin{aligned} \hat{A}_1 &= \phi_m |\Omega|, \quad \hat{A}_2 = \phi_m |\Omega| (c)^\alpha \left( \frac{2}{\phi_0} \right)^{\alpha/2}, \\ \hat{A}_3 &= \phi_m |\Omega| (c)^\beta \left( \frac{2}{\phi_0} \right)^{\beta/2}, \quad \hat{\tau} = \tau = \frac{\sigma_3}{\left( \left[ \Phi_1 \left( \frac{\sigma_3^2}{2} \right) + \Phi_2 \left( \frac{\sigma_3^2}{2} \right) \right] |\Omega| \right)^{1/2}}. \end{aligned} \quad (1.16)$$

and  $a_1, a_2$  are given in  $(G_1)$ .

**Theorem 1.4.** *Assume that  $\phi_i$  ( $i = 1, 2$ ) and  $G$  satisfy  $(\phi_1)'$ ,  $(\phi_4)$  and  $(G_1)$ - $(G_3)$  and  $(G_4)'''$  respectively. Then for  $N = 1$  and each parameter  $\lambda$  belonging to*

$$\Lambda_{\gamma_3, \sigma_3} := \left( \frac{\phi_m \sigma_3^2}{\inf_{x \in \Omega} G(x, \sigma_3, \sigma_3)}, \frac{\phi_m}{b(\hat{A}_1 \gamma_3^{-2} + \hat{A}_2 \gamma_3^{\alpha-2} + \hat{A}_3 \gamma_3^{\beta-2})} \right),$$

the problem (1.10) possesses at least three weak solutions in  $X$ , where  $\hat{A}_1, \hat{A}_2$  and  $\hat{A}_3$  are given in (1.16).

## 2 Preliminaries

Clearly, the corresponding functional of (1.3) is defined on  $W = H_0^1(\Omega) \times H_0^1(\Omega)$  by

$$J(u, v) = \int_{\Omega} \left[ \Phi_1 \left( \frac{u^2 + |\nabla u|^2}{2} \right) + \Phi_2 \left( \frac{v^2 + |\nabla v|^2}{2} \right) \right] dx - \lambda \int_{\Omega} G(x, u, v) dx, \quad (2.1)$$

where the space  $W$  equipped with the norm

$$\|(u, v)\|_W = \|u\|_0 + \|v\|_0 = \left( \int_{\Omega} |\nabla u|^2 dx \right)^{1/2} + \left( \int_{\Omega} |\nabla v|^2 dx \right)^{1/2}.$$

With the similar arguments in [35, Appendix A.1], from  $(\phi_1)'$  and  $(G_1)$ , one has  $J \in C^1(W, \mathbb{R})$ . For any  $(u, v)$  and  $(u_1, v_1) \in W$ , we have

$$\begin{aligned} \langle J'(u, v), (u_1, v_1) \rangle &= \int_{\Omega} \left[ \phi_1 \left( \frac{u^2 + |\nabla u|^2}{2} \right) (uu_1 + \nabla u \cdot \nabla u_1) + \phi_2 \left( \frac{v^2 + |\nabla v|^2}{2} \right) (vv_1 + \nabla v \cdot \nabla v_1) \right] dx \\ &\quad - \int_{\Omega} [G_u(x, u, v)u_1 + G_v(x, u, v)v_1] dx \\ &= \langle J_u(u, v), u_1 \rangle + \langle J_v(u, v), v_1 \rangle. \end{aligned} \quad (2.2)$$

Similarly, the corresponding functional of (1.10) is defined on  $X = H^1(\Omega) \times H^1(\Omega)$  by

$$L(u, v) = \int_{\Omega} \left[ \Phi_1 \left( \frac{u^2 + |\nabla u|^2}{2} \right) + \Phi_2 \left( \frac{v^2 + |\nabla v|^2}{2} \right) \right] dx - \lambda \int_{\Omega} G(x, u, v) dx.$$

The space  $X = H^1(\Omega) \times H^1(\Omega)$  equipped with the norm

$$\|(u, v)\|_X = \|u\| + \|v\| = \left[ \int_{\Omega} (u^2 + |\nabla u|^2) dx \right]^{1/2} + \left[ \int_{\Omega} (v^2 + |\nabla v|^2) dx \right]^{1/2}.$$

From  $(\phi_1)'$  and  $(G_1)$ , it is easy to know  $L \in C^1(X, \mathbb{R})$ . For any  $(\bar{u}, \bar{v})$  and  $(u_2, v_2) \in X$ , we have

$$\begin{aligned} \langle L'(\bar{u}, \bar{v}), (u_2, v_2) \rangle &= \int_{\Omega} \left[ \phi_1 \left( \frac{\bar{u}^2 + |\nabla \bar{u}|^2}{2} \right) (\bar{u}u_2 + \nabla \bar{u} \cdot \nabla u_2) + \phi_2 \left( \frac{\bar{v}^2 + |\nabla \bar{v}|^2}{2} \right) (\bar{v}v_2 + \nabla \bar{v} \cdot \nabla v_2) \right] dx \\ &\quad - \int_{\Omega} [G_u(x, \bar{u}, \bar{v})u_2 + G_v(x, \bar{u}, \bar{v})v_2] dx \\ &= \langle L_{\bar{u}}(\bar{u}, \bar{v}), u_2 \rangle + \langle L_{\bar{v}}(\bar{u}, \bar{v}), v_2 \rangle \end{aligned} \quad (2.3)$$

First, we recall a convergence result [16, Lemma 6] about the monotone operators.

**Proposition 2.1.** *Let  $X$  be a finite dimensional real Hilbert space with norm  $|\cdot|$  and inner product  $\langle \cdot, \cdot \rangle$ . Assume that  $\beta \in C(X, X)$  which is strictly monotone, that is*

$$\langle \beta(\eta) - \beta(\eta_1), \eta - \eta_1 \rangle > 0, \quad \text{for any } \eta, \eta_1 \in X \text{ with } \eta \neq \eta_1.$$

Setting  $\{\eta_n\} \subset X$ ,  $\eta \in X$  and

$$\lim_{n \rightarrow +\infty} \langle \beta(\eta_n) - \beta(\eta), \eta_n - \eta \rangle = 0.$$

Then  $\{\eta_n\}$  converges to  $\eta$  in  $X$ .

Similar to [14, Lemma 3.4], we have the following conclusion,

**Lemma 2.2.** *Assume that  $(\phi_1)'$  and  $(\phi_4)$  hold. Let  $\beta : \mathbb{R}^{N+1} \rightarrow \mathbb{R}^{N+1}$  given by*

$$\beta(\eta) = \phi \left( \frac{|\eta|^2}{2} \right) \eta.$$

Then  $\beta$  is strictly monotone.

**Proof.** Let  $\eta = (u, \nabla u)$  and  $\eta_1 = (\bar{u}, \nabla \bar{u})$ . Following from  $(\phi_4)$ , it is easy to see that  $\phi \left( \frac{t^2}{2} \right) t$  is strictly increasing on  $\mathbb{R}$ . In the view of [14, Lemma 3.4], it holds that

$$\left\langle \phi \left( \frac{|\eta|^2}{2} \right) \eta - \phi \left( \frac{|\eta_1|^2}{2} \right) \eta_1, \eta - \eta_1 \right\rangle > 0, \quad \text{for any } \eta, \eta_1 \in \mathbb{R}^{N+1}. \quad (2.4)$$

Then, we obtain that  $\beta$  strictly monotone. □

**Remark 2.3.** *In this paper, the quasilinear term depends on a function of  $\frac{u^2 + |\nabla u|^2}{2}$  but not  $|\nabla u|$ . This particular quasilinear term makes our problems are different from the problems about  $p$ -Laplacian[6, 15] and  $\Phi$ -Laplacian[8, 31]. To obtain the continuity of  $(I'_1)^{-1}$  and  $(I'_2)^{-1}$ , it is necessary for our arguments to use the theory of monotone operators and the skill of adding one dimension to space. The definition of  $(I'_1)^{-1}$  and  $(I'_2)^{-1}$  will be given in section 3 and section 4.*

### 3 Dirichlet boundary Problem

In this section, we will give the proofs of Theorem 1.1 and Theorem 1.2. Firstly, we drive some lemmas those are useful for the proofs.

In order to study problem (1.3), we put the functional  $I_1, \Upsilon_1 : H_0^1(\Omega) \times H_0^1(\Omega) \rightarrow \mathbb{R}$  and

$$I_1(u, v) = \int_{\Omega} \left[ \Phi_1 \left( \frac{u^2 + |\nabla u|^2}{2} \right) + \Phi_2 \left( \frac{v^2 + |\nabla v|^2}{2} \right) \right] dx \quad (3.1)$$

and

$$\Upsilon_1(u, v) = \int_{\Omega} G(x, u, v) dx \quad \forall (u, v) \in W. \quad (3.2)$$



### 3.1. Proofs of Theorem 1.1.

**Lemma 3.1.** *Assume that  $(\phi_1)'$  and  $(\phi_4)$  hold. Then functional  $I_1 \in C^1(W, \mathbb{R})$  is sequentially weakly lower semicontinuous, coercive and whose Gâteaux derivative  $I_1'$  admits a continuous inverse  $(I_1')^{-1}$  on the dual space  $W^*$  of  $W$ .*

**Proof.** Similar to [14, Lemma 3.2 (ii)], from  $(\phi_4)$ , it is easy to know that

$$I_1(u, v) - I_1(\bar{u}, \bar{v}) - \langle I_1'(\bar{u}, \bar{v}), (u, v) - (\bar{u}, \bar{v}) \rangle \geq 0, \quad \text{for any } (u, v), (\bar{u}, \bar{v}) \in W, \quad (3.3)$$

and  $I_1$  is sequentially weakly lower semicontinuous, that is

$$I_1(u, v) \leq \liminf_{n \rightarrow \infty} I_1(u_n, v_n), \quad \text{for } (u_n, v_n) \rightharpoonup (u, v) \text{ in } W.$$

Following from  $(\phi_1)'$ , (3.1) and the Mean value theorems for definite integrals, there exist  $\xi_1^x \in \left(0, \frac{u^2 + |\nabla u|^2}{2}\right)$  and  $\xi_2^x \in \left(0, \frac{v^2 + |\nabla v|^2}{2}\right)$  for every  $x \in \Omega$ , such that

$$\begin{aligned} I_1(u, v) &= \int_{\Omega} \left( \int_0^{(u^2 + |\nabla u|^2)/2} \phi_1(s) ds + \int_0^{(v^2 + |\nabla v|^2)/2} \phi_2(t) dt \right) dx \\ &= \int_{\Omega} \left[ \phi_1(\xi_1^x) \frac{u^2 + |\nabla u|^2}{2} + \phi_2(\xi_2^x) \frac{v^2 + |\nabla v|^2}{2} \right] dx \\ &\geq \frac{\phi_0}{2} (\|\nabla u\|_2^2 + \|\nabla v\|_2^2) \\ &\geq \frac{\phi_0}{4} \|(u, v)\|_W^2. \end{aligned}$$

Obviously,  $I_1(u, v) \rightarrow +\infty$  as  $\|(u, v)\|_W \rightarrow +\infty$ . Hence,  $I_1$  is coercive.

Next, we claim that  $(I_1')^{-1} \in C(W^*, W)$ . From  $(\phi_1)'$  and (2.2), we have

$$\begin{aligned} \|I_1'(u, v)\|_{W^*} &= \sup_{(u_1, v_1) \in W/\{0\}} \frac{\langle I_1'(u, v), (u_1, v_1) \rangle}{\|(u_1, v_1)\|_W} \\ &\geq \frac{\langle I_1'(u, v), (u, v) \rangle}{\|(u, v)\|_W} \\ &= \frac{\int_{\Omega} \left[ \phi_1 \left( \frac{u^2 + |\nabla u|^2}{2} \right) (u^2 + |\nabla u|^2) + \phi_2 \left( \frac{v^2 + |\nabla v|^2}{2} \right) (v^2 + |\nabla v|^2) \right] dx}{\|(u, v)\|_W} \\ &\geq \frac{\phi_0 \|u\|_0^2 + \phi_0 \|v\|_0^2}{\|(u, v)\|_W} \\ &= \frac{\phi_0}{2} \|(u, v)\|_W, \end{aligned}$$

for any  $(u, v) \in W$ . This shows that  $I_1'$  is coercive in  $W$ . Let  $\eta = (u, \nabla u)$ ,  $\eta_1 = (\bar{u}, \nabla \bar{u})$ ,  $\xi = (v, \nabla v)$  and  $\xi_1 = (\bar{v}, \nabla \bar{v})$ . Following from (2.2), (2.4) and (3.1), it holds that

$$\begin{aligned} &\langle I_1'(u, v) - I_1'(\bar{u}, \bar{v}), (u, v) - (\bar{u}, \bar{v}) \rangle \\ &= \int_{\Omega} \phi_1 \left( \frac{u^2 + |\nabla u|^2}{2} \right) [u(u - \bar{u}) + \nabla u \cdot (\nabla u - \nabla \bar{u})] dx \\ &\quad + \int_{\Omega} \phi_2 \left( \frac{v^2 + |\nabla v|^2}{2} \right) [v(v - \bar{v}) + \nabla v \cdot (\nabla v - \nabla \bar{v})] dx \\ &\quad - \int_{\Omega} \phi_1 \left( \frac{(\bar{u})^2 + |\nabla \bar{u}|^2}{2} \right) [\bar{u}(u - \bar{u}) + \nabla \bar{u} \cdot (\nabla u - \nabla \bar{u})] dx \end{aligned}$$

$$\begin{aligned}
& - \int_{\Omega} \phi_2 \left( \frac{(\bar{v})^2 + |\nabla \bar{v}|^2}{2} \right) [\bar{v}(v - \bar{v}) + \nabla \bar{v} \cdot (\nabla v - \nabla \bar{v})] dx \\
& = \int_{\Omega} \left[ \phi_1 \left( \frac{|\eta|^2}{2} \right) \eta - \phi_1 \left( \frac{|\eta_1|^2}{2} \right) \eta_1 \right] \cdot (\eta - \eta_1) dx + \int_{\Omega} \left[ \phi_2 \left( \frac{|\xi|^2}{2} \right) \xi - \phi_2 \left( \frac{|\xi_1|^2}{2} \right) \xi_1 \right] \cdot (\xi - \xi_1) dx \\
& > 0.
\end{aligned}$$

It holds that  $I'_1$  is strictly monotone in  $W$ . From  $(\phi_1)'$ , it is easy to see that  $I'_1$  is continuous. The continuity of  $I'_1$  shows  $I'_1$  is hemicontinuous. Together with [34, Theorem 26] and the fact that  $I'_1$  is strictly monotone in  $W$ , it holds that there exists the inverse  $(I'_1)^{-1}$  of  $I'_1$  and  $(I'_1)^{-1}$  is bounded in  $W^*$ . Next, we claim that  $(I'_1)^{-1}$  is continuous by proving the inverse  $(I'_1)^{-1}$  of  $I'_1$  is sequentially continuous. Put any sequence  $\{w_n\} \subset W^*$  such that  $w_n \rightarrow w \in W^*$ . Let  $(u_n, v_n) = (I'_1)^{-1}(w_n)$  and  $(u, v) = (I'_1)^{-1}(w)$ . We will prove that  $(u_n, v_n) \rightarrow (u, v)$  in  $W$ . Following from  $(I'_1)^{-1}$  is bounded and  $w_n \rightarrow w$  in  $W^*$ , it holds that  $\{(u_n, v_n)\}$  is bounded in  $W$ . From the fact that  $W$  is a reflexive Banach Space, we have  $(u_n, v_n) \rightharpoonup (u_0, v_0)$  in  $W$ , which shows that  $u_n \rightharpoonup u_0$  in  $H_0^1(\Omega)$  and  $v_n \rightharpoonup v_0$  in  $H_0^1(\Omega)$ , respectively. From the boundness of  $\{(u_n, v_n)\}$  and  $w_n \rightarrow w$  in  $W^*$ , it holds that

$$\langle w_n - w, (u_n, v_n) - (u_0, v_0) \rangle \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

that is,

$$\langle I'_1(u_n, v_n) - I'_1(u, v), (u_n, v_n) - (u_0, v_0) \rangle \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (3.4)$$

Since  $(u_n, v_n) \rightharpoonup (u_0, v_0)$  in  $W$ , one has

$$\langle I'_1(u, v), (u_n, v_n) - (u_0, v_0) \rangle \rightarrow 0 \quad \text{as } n \rightarrow \infty \quad (3.5)$$

and

$$\langle I'_1(u_0, v_0), (u_n, v_n) - (u_0, v_0) \rangle \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (3.6)$$

Combining (3.4), (3.5) and (3.6), we obtain that

$$\begin{aligned}
0 & = \lim_{n \rightarrow \infty} [\langle I'_1(u_n, v_n), (u_n, v_n) - (u_0, v_0) \rangle - \langle I'_1(u_0, v_0), (u_n, v_n) - (u_0, v_0) \rangle] \\
& = \lim_{n \rightarrow \infty} \langle I'_1(u_n, v_n) - I'_1(u_0, v_0), (u_n, v_n) - (u_0, v_0) \rangle \\
& = \lim_{n \rightarrow \infty} \left\{ \int_{\Omega} \left[ \phi_1 \left( \frac{u_n^2 + |\nabla u_n|^2}{2} \right) u_n - \phi_1 \left( \frac{u_0^2 + |\nabla u_0|^2}{2} \right) u_0 \right] (u_n - u_0) dx \right. \\
& \quad + \int_{\Omega} \left[ \phi_1 \left( \frac{u_n^2 + |\nabla u_n|^2}{2} \right) \nabla u_n - \phi_1 \left( \frac{u_0^2 + |\nabla u_0|^2}{2} \right) \nabla u_0 \right] \cdot (\nabla u_n - \nabla u_0) dx \\
& \quad + \int_{\Omega} \left[ \phi_2 \left( \frac{v_n^2 + |\nabla v_n|^2}{2} \right) v_n - \phi_2 \left( \frac{v_0^2 + |\nabla v_0|^2}{2} \right) v_0 \right] (v_n - v_0) dx \\
& \quad \left. + \int_{\Omega} \left[ \phi_2 \left( \frac{v_n^2 + |\nabla v_n|^2}{2} \right) \nabla v_n - \phi_2 \left( \frac{v_0^2 + |\nabla v_0|^2}{2} \right) \nabla v_0 \right] \cdot (\nabla v_n - \nabla v_0) dx \right\}. \quad (3.7)
\end{aligned}$$

Define operators  $\psi_i : H_0^1(\Omega) \rightarrow (H_0^1(\Omega))^*$ , where  $i = 1, 2$ , and

$$\langle \psi_1(u), u_1 \rangle := \int_{\Omega} \phi_1 \left( \frac{u^2 + |\nabla u|^2}{2} \right) (uu_1 + \nabla u \cdot \nabla u_1) dx \quad \forall u, u_1 \in H_0^1(\Omega)$$

and

$$\langle \psi_2(v), v_1 \rangle := \int_{\Omega} \phi_2 \left( \frac{v^2 + |\nabla v|^2}{2} \right) (vv_1 + \nabla v \cdot \nabla v_1) dx \quad \forall v, v_1 \in H_0^1(\Omega).$$

From Lemma 2.2, it has that both  $\psi_1$  and  $\psi_2$  are strictly monotone in  $H_0^1(\Omega)$ . Together with (3.7), we get

$$\begin{aligned} 0 &= \lim_{n \rightarrow \infty} \left\{ \int_{\Omega} \left[ \phi_1 \left( \frac{u_n^2 + |\nabla u_n|^2}{2} \right) u_n - \phi_1 \left( \frac{u_0^2 + |\nabla u_0|^2}{2} \right) u_0 \right] (u_n - u_0) dx \right. \\ &\quad \left. + \int_{\Omega} \left[ \phi_1 \left( \frac{u_n^2 + |\nabla u_n|^2}{2} \right) \nabla u_n - \phi_1 \left( \frac{u_0^2 + |\nabla u_0|^2}{2} \right) \nabla u_0 \right] \cdot (\nabla u_n - \nabla u_0) dx \right\} \end{aligned} \quad (3.8)$$

and

$$\begin{aligned} 0 &= \lim_{n \rightarrow \infty} \left\{ \int_{\Omega} \left[ \phi_2 \left( \frac{v_n^2 + |\nabla v_n|^2}{2} \right) v_n - \phi_2 \left( \frac{v_0^2 + |\nabla v_0|^2}{2} \right) v_0 \right] (v_n - v_0) dx \right. \\ &\quad \left. + \int_{\Omega} \left[ \phi_2 \left( \frac{v_n^2 + |\nabla v_n|^2}{2} \right) \nabla v_n - \phi_2 \left( \frac{v_0^2 + |\nabla v_0|^2}{2} \right) \nabla v_0 \right] \cdot (\nabla v_n - \nabla v_0) dx \right\}. \end{aligned} \quad (3.9)$$

From (3.6), (3.8) and (3.9), it is easy to obtain that

$$0 = \lim_{n \rightarrow \infty} \int_{\Omega} \left[ \phi_1 \left( \frac{u_n^2 + |\nabla u_n|^2}{2} \right) u_n (u_n - u_0) + \phi_1 \left( \frac{u_n^2 + |\nabla u_n|^2}{2} \right) \nabla u_n \cdot (\nabla u_n - \nabla u_0) \right] dx \quad (3.10)$$

and

$$0 = \lim_{n \rightarrow \infty} \int_{\Omega} \left[ \phi_2 \left( \frac{v_n^2 + |\nabla v_n|^2}{2} \right) v_n (v_n - v_0) + \phi_2 \left( \frac{v_n^2 + |\nabla v_n|^2}{2} \right) \nabla v_n \cdot (\nabla v_n - \nabla v_0) \right] dx. \quad (3.11)$$

From [14, Lemma 3.5], we get the conclusion that

$$\nabla u_n(x) \rightarrow \nabla u(x), \quad \text{a.e. in } \Omega. \quad (3.12)$$

In (3.3), let  $u = u_0$ ,  $v = v_0$ ,  $\bar{u} = u_n$  and  $\bar{v} = v_n$ . Combining (3.10) and (3.11), it holds that

$$\limsup_{n \rightarrow \infty} I_1(u_n, v_n) \leq I_1(u_0, v_0). \quad (3.13)$$

Similar to [14, Lemma 5.4] or [18, Lemma 2.3], let  $j(s, t) = \Phi_1(s^2) + \Phi_2(t^2)$ ,  $p_n = h_n - h$  and  $q_n = g_n - g$ , where

$$h_n = \left( \frac{u_n^2 + |\nabla u_n|^2}{2} \right)^{1/2}, \quad h = \left( \frac{u_0^2 + |\nabla u_0|^2}{2} \right)^{1/2}$$

and

$$g_n = \left( \frac{v_n^2 + |\nabla v_n|^2}{2} \right)^{1/2}, \quad g = \left( \frac{v_0^2 + |\nabla v_0|^2}{2} \right)^{1/2}.$$

Following from  $(\phi_1)'$  and  $(\phi_4)$ , it is easy to know that the function  $j$  is continuous, strictly convex on  $\mathbb{R}$  and  $j(0, 0) = 0$ . Using [14, Lemma 5.4] and (3.12), it holds that

$$\int_{\Omega} [j(h + p_n, g + q_n) - j(p_n, q_n) - j(h, g)] dx \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

That is,

$$I_1(u_n, v_n) - I_1(u_n - u_0, v_n - v_0) - I_1(u_0, v_0) \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Together with (3.13) and the fact that  $I_1 \geq 0$ , we obtain

$$\begin{aligned} 0 &\leq \liminf_{n \rightarrow \infty} I_1(u_n - u_0, v_n - v_0) \\ &\leq \limsup_{n \rightarrow \infty} [I_1(u_n, v_n) - I_1(u_0, v_0)] \end{aligned}$$

$$\begin{aligned}
&= \limsup_{n \rightarrow \infty} I_1(u_n, v_n) - I_1(u_0, v_0) \\
&\leq 0.
\end{aligned}$$

This implies

$$\lim_{n \rightarrow \infty} I_1(u_n - u_0, v_n - v_0) = 0. \quad (3.14)$$

From (3.1), it holds

$$I_1(u, v) \geq \frac{\phi_0}{2} \int_{\Omega} (u^2 + |\nabla u|^2 + v^2 + |\nabla v|^2) dx \geq 0, \quad \text{for any } (u, v) \in W.$$

Put  $u = u_n - u_0$  and  $v = v_n - v_0$ . Following (3.14), it has

$$\begin{aligned}
0 &= \lim_{n \rightarrow \infty} I_1(u_n - u_0, v_n - v_0) \\
&\geq \frac{\phi_0}{2} \lim_{n \rightarrow \infty} \int_{\Omega} [(u_n - u_0)^2 + |\nabla u_n - \nabla u_0|^2 + (v_n - v_0)^2 + |\nabla v_n - \nabla v_0|^2] dx \\
&\geq 0.
\end{aligned} \quad (3.15)$$

We get  $(u_n, v_n) \rightarrow (u_0, v_0)$  in  $W$ . Together with the continuity of  $I'_1$ , this shows  $I'_1(u_n, v_n) \rightarrow I'_1(u_0, v_0) = I'_1(u, v)$  in  $W^*$ , when  $n \rightarrow \infty$ . Since  $I'_1$  is continuous and strictly monotone in  $W^*$ , it has  $(u_0, v_0) = (u, v)$ . Therefore, we conclude that  $(I'_1)^{-1}$  is continuous.  $\square$

**Lemma 3.2.** *Assume that  $(G_1)$ - $(G_4)$  hold. Then the functional  $\Upsilon_1 \in C^1(W, \mathbb{R})$  with compact derivative. Moreover,*

$$\langle \Upsilon'_1(u, v), (u_1, v_1) \rangle = \int_{\Omega} \Upsilon_{1u}(u, v) u_1 dx + \int_{\Omega} \Upsilon_{1v}(u, v) v_1 dx \quad (3.16)$$

for all  $(u_1, v_1) \in W$ .

**Proof.** Following from  $(G_1)$ , (1.5) and (3.2), one has

$$\begin{aligned}
\Upsilon_1(u, v) &\leq \int_{\Omega} \left[ a_1(|u| + |v|) + a_2 \frac{|u|^p}{p} + a_2 \frac{|v|^q}{q} \right] dx \\
&\leq a_1 C_1 \|\nabla u\|_2 + a_1 C_1 \|\nabla v\|_2 + \frac{a_2 (C_p)^p}{p} \|\nabla u\|_2^p + \frac{a_2 (C_q)^q}{q} \|\nabla v\|_2^q.
\end{aligned}$$

Thus,  $\Upsilon_1(u, v)$  is well defined in  $W$ . From  $(G_1)$ , it is easy to know that  $\Upsilon_1 \in C^1(W, \mathbb{R})$  and the equation (3.16) holds.

To reach the compactness of  $\Upsilon'_1$ , we put any sequence  $\{(u_n, v_n)\} \subset W$  which is bounded. Following from the reflexivity of  $W$  and the Sobolev embedding Theorem, there exists a subsequence still denoted by  $\{(u_n, v_n)\}$ , such that  $(u_n, v_n) \rightharpoonup (u_0, v_0) \in W$ , and  $u_n \rightarrow u_0$  in  $L^p(\Omega)$  and  $v_n \rightarrow v_0$  in  $L^q(\Omega)$ , respectively. Similar to [32], following from (3.16), the Hölder's inequality and the Sobolev embedding inequality, we obtain

$$\begin{aligned}
&|\Upsilon'_1(u_n, v_n) - \Upsilon'_1(u_0, v_0), (u_1, v_1)| \\
&= \left| \int_{\Omega} [G_u(x, u_n, v_n) - G_u(x, u_0, v_0)] u_1 dx + \int_{\Omega} [G_v(x, u_n, v_n) - G_v(x, u_0, v_0)] v_1 dx \right| \\
&\leq \bar{C} \left\{ \left[ \int_{\Omega} |G_u(x, u_n, v_n) - G_u(x, u_0, v_0)|^{p/(p-1)} dx \right]^{(p-1)/p} \right. \\
&\quad \left. + \left[ \int_{\Omega} |G_v(x, u_n, v_n) - G_v(x, u_0, v_0)|^{q/(q-1)} dx \right]^{(q-1)/q} \right\} \|(u_1, v_1)\|_W,
\end{aligned} \quad (3.17)$$

for any  $(u_1, v_1) \in W$ , where  $\bar{C}$  is a positive constant. Meanwhile, by (3.17) and the continuity of  $G_u$  and  $G_v$ , one has

$$\int_{\Omega} |G_u(x, u_n, v_n) - G_u(x, u_0, v_0)|^{p/(p-1)} dx \rightarrow 0, \quad \text{as } n \rightarrow \infty \quad (3.18)$$

and

$$\int_{\Omega} |G_v(x, u_n, v_n) - G_v(x, u_0, v_0)|^{q/(q-1)} dx \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (3.19)$$

Combining (3.17), (3.18) and (3.19), we obtain

$$\begin{aligned} & \|\Upsilon'_1(u_n, v_n) - \Upsilon'_1(u_0, v_0)\|_* \\ = & \sup_{(\phi_1, \phi_2) \in W, \|(\phi_1, \phi_2)\|=1} |\langle \Upsilon'_1(u_n, v_n) - \Upsilon'_1(u_0, v_0), (\phi_1, \phi_2) \rangle| \\ = & \sup_{(\phi_1, \phi_2) \in W, \|(\phi_1, \phi_2)\|=1} \left| \int_{\Omega} [G_u(x, u_n, v_n) - G_u(x, u_0, v_0)] \phi_1 dx + \int_{\Omega} [G_v(x, u_n, v_n) - G_v(x, u_0, v_0)] \phi_2 dx \right| \\ \leq & \sup_{(\phi_1, \phi_2) \in W, \|(\phi_1, \phi_2)\|=1} \left\{ \int_{\Omega} |[G_u(x, u_n, v_n) - G_u(x, u_0, v_0)] \phi_1| dx + \int_{\Omega} |[G_v(x, u_n, v_n) - G_v(x, u_0, v_0)] \phi_2| dx \right\} \\ \leq & \sup_{(\phi_1, \phi_2) \in W, \|(\phi_1, \phi_2)\|=1} \left\{ \left[ \int_{\Omega} |G_u(x, u_n, v_n) - G_u(x, u_0, v_0)|^{p/(p-1)} dx \right]^{(p-1)/p} \|\phi_1\|_p \right. \\ & \left. + \left[ \int_{\Omega} |G_v(x, u_n, v_n) - G_v(x, u_0, v_0)|^{q/(q-1)} dx \right]^{(q-1)/q} \|\phi_2\|_q \right\} \\ \leq & \sup_{(\phi_1, \phi_2) \in W, \|(\phi_1, \phi_2)\|=1} \left\{ C_p \left[ \int_{\Omega} |G_u(x, u_n, v_n) - G_u(x, u_0, v_0)|^{p/(p-1)} dx \right]^{(p-1)/p} \|\nabla \phi_1\|_2 \right. \\ & \left. + C_q \left[ \int_{\Omega} |G_v(x, u_n, v_n) - G_v(x, u_0, v_0)|^{q/(q-1)} dx \right]^{(q-1)/q} \|\nabla \phi_2\|_2 \right\} \\ \leq & C \left\{ \left[ \int_{\Omega} |G_u(x, u_n, v_n) - G_u(x, u_0, v_0)|^{p/(p-1)} dx \right]^{(p-1)/p} \right. \\ & \left. + \left[ \int_{\Omega} |G_v(x, u_n, v_n) - G_v(x, u_0, v_0)|^{q/(q-1)} dx \right]^{(q-1)/q} \right\} \|(\phi_1, \phi_2)\|_W \\ = & 0. \end{aligned}$$

Then, we reach the conclusion that  $\Upsilon'_1$  is compact.  $\square$

Next, we will claim the assumptions  $(a_1)$  and  $(a_2)$  in Theorem I are satisfied. Following from  $(G_1)$ , one has

$$G(x, \xi, \zeta) \leq a_1(|\xi| + |\zeta|) + a_2 \frac{|\xi|^p}{p} + a_2 \frac{|\zeta|^q}{q}, \quad \text{for every } (x, \xi, \zeta) \in (\Omega \times \mathbb{R} \times \mathbb{R}). \quad (3.20)$$

Let  $r_1 \in (0, +\infty)$  and

$$\chi(r_1) := \frac{\sup_{(u,v) \in I_1^{-1}((-\infty, r_1))} \Upsilon_1(u, v)}{r_1}. \quad (3.21)$$

Following from (1.5) and (3.20), one has

$$\begin{aligned} \Upsilon_1(u, v) &= \int_{\Omega} G(x, u, v) dx \\ &\leq a_1 \|u\|_1 + a_1 \|v\|_1 + a_2 \frac{\|u\|_p^p}{p} + a_2 \frac{\|v\|_q^q}{q} \\ &\leq a_1 C_1 \|\nabla u\|_2 + a_1 C_1 \|\nabla v\|_2 + a_2 (C_p)^p \frac{\|\nabla u\|_2^p}{p} + a_2 (C_q)^q \frac{\|\nabla v\|_2^q}{q}. \end{aligned} \quad (3.22)$$

Following from (3.4), for every  $(u, v) \in W$  and  $I_1(u, v) \leq r_1$ , we have

$$I_1(u, v) \geq \frac{\phi_0}{2} (\|\nabla u\|_2^2 + \|\nabla v\|_2^2), \quad (3.23)$$

$$\frac{1}{2} (\|\nabla u\|_2 + \|\nabla v\|_2)^2 \leq \|\nabla u\|_2^2 + \|\nabla v\|_2^2 \leq \frac{2r_1}{\phi_0} \quad (3.24)$$

and

$$\|\nabla u\|_2 + \|\nabla v\|_2 \leq 2 \left( \frac{r_1}{\phi_0} \right)^{1/2}.$$

Together with (3.22), it holds

$$\sup_{(u,v) \in I_1^{-1}((-\infty, r_1])} \Upsilon_1(u, v) \leq 2a_1 C_1 \left( \frac{r_1}{\phi_0} \right)^{1/2} + \frac{a_2(C_p)^p}{p} \left( \frac{2r_1}{\phi_0} \right)^{p/2} + \frac{a_2(C_q)^q}{q} \left( \frac{2r_1}{\phi_0} \right)^{q/2}. \quad (3.25)$$

Following from (3.21) and (3.25), it holds that

$$\chi(r_1) \leq 2a_1 C_1 \left( \frac{1}{\phi_0} \right)^{1/2} r_1^{-1/2} + \frac{a_2(C_p)^p}{p} \left( \frac{2}{\phi_0} \right)^{p/2} r_1^{p/2-1} + \frac{a_2(C_q)^q}{q} \left( \frac{2}{\phi_0} \right)^{q/2} r_1^{q/2-1}, \quad r_1 > 0. \quad (3.26)$$

Put

$$u_{\sigma_1}(x) := \begin{cases} 0, & \text{if } x \in \Omega \setminus B(x_0, D), \\ \sigma_1, & \text{if } x \in B(x_0, D) \end{cases}$$

and

$$v_{\sigma_1}(x) := \begin{cases} 0, & \text{if } x \in \Omega \setminus B(x_0, D), \\ \sigma_1, & \text{if } x \in B(x_0, D). \end{cases}$$

Clearly,  $(u_{\sigma_1}, v_{\sigma_1}) \in W$  and

$$\begin{aligned} I_1(u_{\sigma_1}, v_{\sigma_1}) &= \int_{\Omega} \left[ \Phi_1 \left( \frac{u_{\sigma_1}^2 + |\nabla u_{\sigma_1}^2|}{2} \right) + \Phi_2 \left( \frac{v_{\sigma_1}^2 + |\nabla v_{\sigma_1}^2|}{2} \right) \right] dx \\ &= \int_{B(x_0, D)} \left[ \Phi_1 \left( \frac{\sigma_1^2}{2} \right) + \Phi_2 \left( \frac{\sigma_1^2}{2} \right) \right] dx \\ &= \left[ \Phi_1 \left( \frac{\sigma_1^2}{2} \right) + \Phi_2 \left( \frac{\sigma_1^2}{2} \right) \right] \text{meas}(B(x_0, D)) \\ &= \left[ \Phi_1 \left( \frac{\sigma_1^2}{2} \right) + \Phi_2 \left( \frac{\sigma_1^2}{2} \right) \right] \frac{\pi^{N/2}}{\Gamma(1 + N/2)} D^N. \end{aligned} \quad (3.27)$$

From  $(G_2)$ , we have

$$\begin{aligned} \Upsilon_1(u_{\sigma_1}, v_{\sigma_1}) &= \int_{\Omega} G(x, u_{\sigma_1}, v_{\sigma_1}) dx \\ &\geq \int_{B(x_0, D)} G(x, \sigma_1, \sigma_1) dx \\ &\geq \inf_{\Omega} G(x, \sigma_1, \sigma_1) \frac{\pi^{N/2}}{\Gamma(1 + N/2)} D^N. \end{aligned} \quad (3.28)$$

Combining (3.2), (3.27), (3.28) and  $(\phi_1)'$ , we get

$$\begin{aligned} \frac{\Upsilon_1(u_{\sigma_1}, v_{\sigma_1})}{I_1(u_{\sigma_1}, v_{\sigma_1})} &\geq \frac{\inf_{\Omega} G(x, \sigma_1, \sigma_1)}{\left[ \Phi_1\left(\frac{\sigma_1^2}{2}\right) + \Phi_2\left(\frac{\sigma_1^2}{2}\right) \right]} \\ &\geq \frac{1}{\phi_m} \frac{\inf_{\Omega} G(x, \sigma_1, \sigma_1)}{\sigma_1^2}. \end{aligned} \quad (3.29)$$

Note that  $\tau = \frac{\sigma_1}{[I_1(u_{\sigma_1}, v_{\sigma_1})]^{1/2}}$ . Then  $I_1(u_{\sigma_1}, v_{\sigma_1}) = \left(\frac{\sigma_1}{\tau}\right)^2$ . Taking into account that  $\sigma_1 > \gamma_1 \tau$ , it holds  $\gamma_1^2 < I_1(u_{\sigma_1}, v_{\sigma_1})$ . Together with (3.21), (3.26), (3.29) and  $(G_4)$ , we get

$$\begin{aligned} \chi(\gamma_1^2) &:= \frac{\sup_{(u,v) \in I_1^{-1}((-\infty, \gamma_1^2])} \Upsilon_1(u, v)}{\gamma_1^2} \\ &\leq \frac{2a_1 C_1}{\gamma_1} \left(\frac{1}{\phi_0}\right)^{1/2} + \frac{a_2 (C_p)^p}{p} \left(\frac{2}{\phi_0}\right)^{p/2} \gamma_1^{p-2} + \frac{a_2 (C_q)^q}{q} \left(\frac{2}{\phi_0}\right)^{q/2} \gamma_1^{q-2} \\ &= \frac{1}{\phi_m} \left(2a_1 \frac{A_1}{\gamma_1} + a_2 A_2 \gamma_1^{p-2} + a_2 A_3 \gamma_1^{q-2}\right) \\ &< \frac{1}{\phi_m} \frac{\inf_{\Omega} G(x, \sigma_1, \sigma_1)}{\sigma_1^2} \\ &\leq \frac{\Upsilon_1(u_{\sigma_1}, v_{\sigma_1})}{I_1(u_{\sigma_1}, v_{\sigma_1})}. \end{aligned}$$

Therefore, the condition  $(a_1)$  in Theorem I is satisfied.

Moreover, for every  $u, v \in H_0^1(\Omega)$ ,  $0 < \alpha < 2$  and  $0 < \beta < 2$ , it is easy to know that

$$|u|^\alpha \in L^{2/\alpha}(\Omega), \quad |v|^\beta \in L^{2/\beta}(\Omega).$$

Together with the Hölder's inequality, it has

$$\int_{\Omega} |u|^\alpha dx \leq \|u\|_2^\alpha |\Omega|^{(2-\alpha)/2}, \quad \forall u \in H_0^1(\Omega) \quad (3.30)$$

and

$$\int_{\Omega} |v|^\beta dx \leq \|v\|_2^\beta |\Omega|^{(2-\beta)/2}, \quad \forall v \in H_0^1(\Omega). \quad (3.31)$$

Following from (1.5), (2.1), (3.30), (3.31) and  $(G_3)$ , we obtain that

$$\begin{aligned} J(u, v) &\geq \frac{\phi_0}{2} (\|u\|_2^2 + \|\nabla u\|_2^2 + \|v\|_2^2 + \|\nabla v\|_2^2) - \lambda b \int_{\Omega} (1 + |u|^\alpha + |v|^\beta) dx \\ &\geq \frac{\phi_0}{2} (\|u\|_2^2 + \|\nabla u\|_2^2 + \|v\|_2^2 + \|\nabla v\|_2^2) - \lambda b |\Omega| - \lambda b \|u\|_2^\alpha |\Omega|^{(2-\alpha)/2} - \lambda b \|v\|_2^\beta |\Omega|^{(2-\beta)/2} \\ &\geq \frac{\phi_0}{2} (\|\nabla u\|_2^2 + \|\nabla v\|_2^2) - \lambda b |\Omega| - \lambda b (C_2)^\alpha |\Omega|^{(2-\alpha)/2} \|\nabla u\|_2^\alpha - \lambda b (C_2)^\beta |\Omega|^{(2-\beta)/2} \|\nabla v\|_2^\beta \\ &\geq \frac{\phi_0}{4} \|(u, v)\|_W^2 - \lambda b |\Omega| - \lambda b (C_2)^\alpha |\Omega|^{(2-\alpha)/2} \|\nabla u\|_2^\alpha - \lambda b (C_2)^\beta |\Omega|^{(2-\beta)/2} \|\nabla v\|_2^\beta, \end{aligned}$$

for any  $(u, v) \in W$ . Therefore,  $J(u, v)$  is coercive for every positive constant  $\lambda$ . Particularly,  $J(u, v)$  is coercive for every  $\lambda \in \Lambda_{\gamma_1, \sigma_1} \subseteq \left( \frac{I_1(u_{\sigma_1}, v_{\sigma_1})}{\Upsilon_1(u_{\sigma_1}, v_{\sigma_1})}, \frac{\gamma_1^2}{\sup_{(u,v) \in I_1^{-1}((-\infty, \gamma_1^2])} \Upsilon_1(u, v)} \right)$ . This shows that the condition  $(a_2)$  in Theorem

I is satisfied. Therefore, the conditions of Theorem I are satisfied. That is, we can reach the conclusion that the problem (1.3) has at least three non-trivial weak solutions which are the critical point of the functional  $J$  for any  $\lambda \in \Lambda_{\gamma_1, \sigma_1}$ .  $\square$

### 3.2. Proofs of Theorem 1.2.

Since  $H_0^1(\Omega) \hookrightarrow C^0(\overline{\Omega})$  is compact, when  $\Omega \in \mathbb{R}^N$  and is a bounded domain with  $N = 1$ . It holds the Sobolev embedding inequality

$$\|u\|_\infty \leq C\|u\|_0 < +\infty, \quad (3.32)$$

where  $u \in H_0^1(\Omega)$  and  $\|u\|_\infty := \sup_{x \in \overline{\Omega}} |u(x)|$ . It is easy to prove that both Lemma 3.1 and Lemma 3.2 hold. Following from  $(G_3)$  and (3.32), one has

$$\begin{aligned} \Upsilon_1(u, v) &= \int_{\Omega} G(x, u, v) dx \leq b \int_{\Omega} (1 + |u|^\alpha + |v|^\beta) dx \\ &\leq b|\Omega| + b|\Omega|\|u\|_\infty^\alpha + b|\Omega|\|v\|_\infty^\beta \\ &\leq b|\Omega| + b(C)^\alpha |\Omega| \|u\|_0^\alpha + b(C)^\beta |\Omega| \|v\|_0^\beta. \end{aligned} \quad (3.33)$$

Assume that  $I_1(u, v) \leq r_1$ . Following from (3.23) and (3.24), it holds

$$\|u\|_0 \leq \left(\frac{2}{\phi_0}\right)^{1/2} r_1^{1/2}, \quad \|v\|_0 \leq \left(\frac{2}{\phi_0}\right)^{1/2} r_1^{1/2}, \quad \|u\|_0 + \|v\|_0 \leq 2 \left(\frac{r_1}{\phi_0}\right)^{1/2}. \quad (3.34)$$

Combining (3.33), (3.34) and  $(G_3)$ , it follows that

$$\Upsilon_1(u, v) \leq b|\Omega| + b|\Omega|(C)^\alpha \left(\frac{2}{\phi_0}\right)^{\alpha/2} r_1^{\alpha/2} + b(C)^\beta |\Omega| \left(\frac{2}{\phi_0}\right)^{\beta/2} r_1^{\beta/2}$$

for  $I_1(u, v) \leq r_1$ . Thus, one has

$$\begin{aligned} \chi_1(r_1) &= \frac{\sup_{(u,v) \in I_1^{-1}((-\infty, r_1])} \Upsilon_1(u, v)}{r_1} \\ &\leq b|\Omega|r_1^{-1} + b|\Omega|(C)^\alpha \left(\frac{2}{\phi_0}\right)^{\alpha/2} r_1^{\alpha/2-1} + b|\Omega|(C)^\beta \left(\frac{2}{\phi_0}\right)^{\beta/2} r_1^{\beta/2-1}. \end{aligned}$$

Put

$$u_{\sigma_1}(x) := \sigma_1 \quad v_{\sigma_1}(x) := \sigma_1, \quad \text{for } x \in \Omega.$$

Clearly,  $(u_{\sigma_1}, v_{\sigma_1}) \in X$ . This implies that

$$\begin{aligned} I_1(u_{\sigma_1}, v_{\sigma_1}) &= \int_{\Omega} \left[ \Phi_1 \left( \frac{u_{\sigma_1}^2 + |\nabla u_{\sigma_1}|^2}{2} \right) + \Phi_2 \left( \frac{v_{\sigma_1}^2 + |\nabla v_{\sigma_1}|^2}{2} \right) \right] dx \\ &= \int_{\Omega} \left[ \Phi_1 \left( \frac{\sigma_1^2}{2} \right) + \Phi_2 \left( \frac{\sigma_1^2}{2} \right) \right] dx \\ &= \left[ \Phi_1 \left( \frac{\sigma_1^2}{2} \right) + \Phi_2 \left( \frac{\sigma_1^2}{2} \right) \right] |\Omega| \end{aligned}$$

and

$$\begin{aligned} \frac{\Upsilon_1(u_{\sigma_1}, v_{\sigma_1})}{I_1(u_{\sigma_1}, v_{\sigma_1})} &\geq \frac{\inf_{\Omega} G(x, \sigma_1, \sigma_1)}{\left[ \Phi_1 \left( \frac{\sigma_1^2}{2} \right) + \Phi_2 \left( \frac{\sigma_1^2}{2} \right) \right]} \\ &\geq \frac{1}{\phi_m} \frac{\inf_{\Omega} G(x, \sigma_1, \sigma_1)}{\sigma_1^2}. \end{aligned}$$



Following from  $(G_4)''$ , put  $r_1 = \gamma_1^2$ ,

$$\begin{aligned}
\chi(\gamma_1^2) &:= \frac{\sup_{(u,v) \in I_1^{-1}((-\infty, \gamma_1^2])} \Upsilon_1(u, v)}{\gamma_1^2} \\
&\leq b|\Omega|\gamma_1^{-2} + b|\Omega|(C)^\alpha \left(\frac{2}{\phi_0}\right)^{\alpha/2} \gamma_1^{\alpha-2} + b|\Omega|(C)^\beta \left(\frac{2}{\phi_0}\right)^{\beta/2} \gamma_1^{\beta-2} \\
&= \frac{1}{\phi_m} (bA_{11}\gamma_1^{-2} + bA_{21}\gamma_1^{\alpha-2} + bA_{31}\gamma_1^{\beta-2}) \\
&< \frac{1}{\phi_m} \frac{G(x, \sigma_1, \sigma_1)}{\sigma_1^2} \\
&\leq \frac{\Upsilon_1(u_{\sigma_1}, v_{\sigma_1})}{I_1(u_{\sigma_1}, v_{\sigma_1})}.
\end{aligned}$$

This claims that the assumption  $(a_1)$  in Theorem I holds.

Moreover, one has

$$\int_{\Omega} |u|^\alpha dx \leq \|u\|_\infty^\alpha |\Omega|, \quad \forall u \in H_0^1(\Omega)$$

and

$$\int_{\Omega} |v|^\beta dx \leq \|v\|_\infty^\beta |\Omega|, \quad \forall v \in H_0^1(\Omega).$$

Combining with (3.32), we have

$$\int_{\Omega} |u|^\alpha dx \leq \|u\|_\infty^\alpha |\Omega| \leq (C)^\alpha \|u\|_0^\alpha |\Omega|, \quad \forall u \in H_0^1(\Omega), \quad (3.35)$$

and

$$\int_{\Omega} |v|^\beta dx \leq \|v\|_\infty^\beta |\Omega| \leq (C)^\beta \|v\|_0^\beta |\Omega|, \quad \forall v \in H_0^1(\Omega). \quad (3.36)$$

Following from  $(\phi_1)'$ ,  $(G_3)$ , (2.1) (3.35) and (3.36), we obtain that

$$\begin{aligned}
J(u, v) &= \int_{\Omega} \left[ \Phi_1 \left( \frac{u^2 + |\nabla u|^2}{2} \right) + \Phi_2 \left( \frac{v^2 + |\nabla v|^2}{2} \right) \right] dx - \lambda \int_{\Omega} G(x, u, v) dx \\
&\geq \int_{\Omega} \left[ \Phi_1 \left( \frac{u^2 + |\nabla u|^2}{2} \right) + \Phi_2 \left( \frac{v^2 + |\nabla v|^2}{2} \right) \right] dx \\
&\quad - \int_{\Omega} \lambda b dx - \int_{\Omega} \lambda b |u|^\alpha dx - \int_{\Omega} \lambda b |v|^\beta dx \\
&\geq \frac{\phi_0}{2} \int_{\Omega} (u^2 + |\nabla u|^2 + v^2 + |\nabla v|^2) dx - \lambda b |\Omega| \\
&\quad - \lambda b \|u\|_\infty^\alpha |\Omega| - \lambda b \|v\|_\infty^\beta |\Omega| \\
&\geq \frac{\phi_0}{2} (\|u\|_0^2 + \|v\|_0^2) - \lambda b |\Omega| - \lambda b (C)^\alpha \|u\|_0^\alpha |\Omega| - \lambda b (C)^\beta \|v\|_0^\beta |\Omega|, \quad \forall (u, v) \in H_0^1(\Omega) \times H_0^1(\Omega).
\end{aligned}$$

Since  $\alpha, \beta < 2$ ,  $J$  is a coercive functional for every positive parameter  $\lambda$ , and so for every

$$\lambda \in \bar{\Lambda}_{\gamma_1, \sigma_1} = \left( \frac{\phi_m \sigma_1^2}{\inf_{x \in \Omega} G(x, \sigma_1, \sigma_1)}, \frac{\phi_m}{b(A_{11}\gamma_1^{-2} + A_{21}\gamma_1^{\alpha-2} + A_{31}\gamma_1^{\beta-2})} \right).$$

This shows that  $(a_2)$  in Theorem I holds. All these imply  $J$  has at least three non-trivial critical points for any  $\lambda \in \bar{\Lambda}_{\gamma_1, \sigma_1}$ . We complete the proofs of Theorem 1.2.  $\square$

## 4 Neuman boundary problem

In this section, we will give the proofs of Theorem 1.3 and Theorem 1.4 by replacing  $(G_4)$  with  $(G_4)'$  and  $(G_4)'''$  respectively. This proofs are similar to section 3, here we mainly state the proofs related to  $(X, \|\cdot\|)$  and the embedding inequality (1.12) or (1.15)

In order to study problem (1.10), we put the functional  $I_2, \Upsilon_2 : X \rightarrow \mathbb{R}$  and

$$I_2(u, v) = \int_{\Omega} \left[ \Phi_1 \left( \frac{u^2 + |\nabla u|^2}{2} \right) + \Phi_2 \left( \frac{v^2 + |\nabla v|^2}{2} \right) \right] dx \quad (4.1)$$

and

$$\Upsilon_2(u, v) = \int_{\Omega} G(x, u, v) dx, \quad \forall (u, v) \in X. \quad (4.2)$$

### 4.1. Proofs of Theorem 1.3.

**Lemma 4.1.** *Assume that  $(\phi_1)'$  and  $(\phi_4)$  hold. Then functional  $I_2 \in C^1(X, \mathbb{R})$  is sequentially weakly lower semicontinuous, coercive, bounded on each bounded subset of  $X$ , and whose Gâteaux derivative  $I_2'$  admits a continuous inverse  $(I_2')^{-1}$  on the dual space  $X^*$  of  $X$ .*

**Proof.** The arguments are similar to Lemma 3.1. Following from  $(\phi_4)$ , it holds that

$$I_2(u, v) \leq \liminf_{n \rightarrow \infty} I_2(u_n, v_n).$$

From (4.1) and  $(\phi_1)'$ , one has

$$\begin{aligned} I_2(u, v) &\geq \frac{\phi_0}{2} (\|u\|_2^2 + \|\nabla u\|_2^2 + \|v\|_2^2 + \|\nabla v\|_2^2) \\ &= \frac{\phi_0}{2} (\|u\|^2 + \|v\|^2) \\ &\geq \frac{\phi_0}{4} \|(u, v)\|_X^2. \end{aligned} \quad (4.3)$$

This implies  $I_2$  is coercive.

Next, we deduce that  $(I_2')^{-1} \in C(X^*, X)$ . From (2.3) and  $(\phi_1)'$ , we have

$$\begin{aligned} \frac{\langle I_2'(u, v), (u, v) \rangle}{\|(u, v)\|_X} &= \frac{\int_{\Omega} \left[ \phi_1 \left( \frac{u^2 + |\nabla u|^2}{2} \right) (u^2 + |\nabla u|^2) + \phi_2 \left( \frac{v^2 + |\nabla v|^2}{2} \right) (v^2 + |\nabla v|^2) \right] dx}{\|(u, v)\|_X} \\ &\geq \frac{\phi_0 \|u\|^2 + \phi_0 \|v\|^2}{\|(u, v)\|_X} \\ &\geq \frac{\phi_0}{2} \|(u, v)\|_X \end{aligned}$$

This shows that  $I_2'$  is coercive in  $X$ . Similar to Lemma 3.1, we also can obtain  $(I_2')^{-1}$  is continuous.  $\square$

By the similar arguments of Lemma 3.2, we can get the following result.

**Lemma 4.2.** *Assume that  $(G_1)$ - $(G_3)$  and  $(G_4)'$  hold. Then the functional  $\Upsilon_2 \in C^1(X, \mathbb{R})$  with compact derivative. Moreover,*

$$\langle \Upsilon_2'(u, v), (u_1, v_1) \rangle = \int_{\Omega} \Upsilon'_{2u}(u, v) u_1 dx + \int_{\Omega} \Upsilon'_{2v}(u, v) v_1 dx$$

for all  $(u_1, v_1) \in X$ .

Similar to (3.16), we can obtain

$$\langle \Upsilon'_2(u, v), (u_1, v_1) \rangle = \int_{\Omega} \Upsilon'_{2u}(u, v) u_1 dx + \int_{\Omega} \Upsilon'_{2v}(u, v) v_1 dx$$

for all  $(u_1, v_1) \in X$ . Next, we will show the conditions  $(a_1)$  and  $(a_2)$  in Theorem I are satisfied. Let  $r_2 \in (0, +\infty)$  and

$$\chi(r_2) := \frac{\sup_{(u,v) \in I_2^{-1}((-\infty, r_2])} \Upsilon_2(u, v)}{r_2} \quad (4.4)$$

Assume that  $I_2(u, v) \leq r_2$ , together with (4.3), it has

$$\frac{\phi_0}{2} (\|u\|^2 + \|v\|^2) \leq r_2. \quad (4.5)$$

Then, we have

$$\frac{1}{2} (\|u\| + \|v\|)^2 \leq \|u\|^2 + \|v\|^2 \leq \frac{2}{\phi_0} r_2 \quad (4.6)$$

and

$$\|u\| \leq \left(\frac{2}{\phi_0}\right)^{1/2} r_2^{1/2}, \quad \|v\| \leq \left(\frac{2}{\phi_0}\right)^{1/2} r_2^{1/2}, \quad \|u\| + \|v\| \leq 2 \left(\frac{r_2}{\phi_0}\right)^{1/2}. \quad (4.7)$$

Following from  $(G_1)$ , (1.12), (4.2) and (4.5)-(4.7), we have that

$$\begin{aligned} \Upsilon_2(u, v) &\leq a_1 \|u\|_1 + a_1 \|v\|_1 + \frac{a_2}{p} \|u\|_p^p + \frac{a_2}{q} \|v\|_q^q \\ &\leq a_1 c_1 (\|u\| + \|v\|) + \frac{a_2}{p} (c_p)^p \|u\|^p + \frac{a_2}{q} (c_q)^q \|v\|^q \\ &\leq 2a_1 c_1 \left(\frac{1}{\phi_0}\right)^{1/2} r_2^{1/2} + \frac{a_2}{p} (c_p)^p \left(\frac{2}{\phi_0}\right)^{p/2} r_2^{p/2} + \frac{a_2}{q} (c_q)^q \left(\frac{2}{\phi_0}\right)^{q/2} r_2^{q/2}, \end{aligned} \quad (4.8)$$

for  $I_2(u, v) \leq r_2$ . Combining (4.4) and (4.8), it holds that

$$\begin{aligned} \chi_2(r_2) &= \frac{\sup_{(u,v) \in I_2^{-1}((-\infty, r_2])} \Upsilon_2(u, v)}{r_2} \\ &\leq 2a_1 c_1 \left(\frac{1}{\phi_0}\right)^{1/2} r_2^{-1/2} + \frac{a_2}{p} (c_p)^p \left(\frac{2}{\phi_0}\right)^{p/2} r_2^{p/2-1} + \frac{a_2}{q} (c_q)^q \left(\frac{2}{\phi_0}\right)^{q/2} r_2^{q/2-1}. \end{aligned}$$

Put

$$u_{\sigma_2}(x) := \begin{cases} 0, & \text{if } x \in \Omega \setminus B(x_0, D), \\ \sigma_2, & \text{if } x \in B(x_0, D) \end{cases}$$

and

$$v_{\sigma_2}(x) := \begin{cases} 0, & \text{if } x \in \Omega \setminus B(x_0, D), \\ \sigma_2, & \text{if } x \in B(x_0, D). \end{cases}$$

Note that  $\bar{\tau} = \frac{\sigma_2}{[I_2(u_{\sigma_2}, v_{\sigma_2})]^{1/2}}$ . Then,  $I_2(u_{\sigma_2}, v_{\sigma_2}) = \left(\frac{\sigma_2}{\bar{\tau}}\right)^2$ . Taking into account that  $\sigma_2 > \gamma_2 \bar{\tau}$ , it holds  $\gamma_2^2 < I_2(u_{\sigma_2}, v_{\sigma_2})$ .

Similar to the proofs of Theorem 1.1, from  $(G_4)$ , we get

$$\chi_2(\gamma_2^2) := \frac{\sup_{(u,v) \in I_1^{-1}((-\infty, \gamma_2^2])} \Upsilon_1(u, v)}{\gamma_2^2} < \frac{\Upsilon_2(u_{\sigma_2}, v_{\sigma_2})}{I_2(u_{\sigma_2}, v_{\sigma_2})}.$$

Therefore, the condition  $(a_1)$  in Theorem I is satisfied.

Moreover, from the Hölder inequality and  $\alpha, \beta < 2$ , we obtain

$$\int_{\Omega} |u|^{\alpha} dx \leq \|u\|_2^{\alpha} |\Omega|^{(2-\alpha)/2}, \quad \forall u \in H^1(\Omega). \quad (4.9)$$

Combining with (1.12), we have

$$\int_{\Omega} |u|^{\alpha} dx \leq (c_2)^{\alpha} \|u\|^{\alpha} |\Omega|^{(2-\alpha)/2}, \quad \forall u \in H^1(\Omega). \quad (4.10)$$

Similar to (4.9) and (4.10), it also has

$$\int_{\Omega} |u|^{\beta} dx \leq (c_2)^{\beta} \|u\|^{\beta} |\Omega|^{(2-\beta)/2}, \quad \forall u \in H^1(\Omega). \quad (4.11)$$

Following from  $(\phi_1)'$ ,  $(G_3)$ , (2.1) (4.10) and (4.11), we obtain that

$$\begin{aligned} L(u, v) &= \int_{\Omega} \left[ \Phi_1 \left( \frac{u^2 + |\nabla u|^2}{2} \right) + \Phi_2 \left( \frac{v^2 + |\nabla v|^2}{2} \right) \right] dx - \lambda \int_{\Omega} G(x, u, v) dx \\ &\geq \int_{\Omega} \left[ \Phi_1 \left( \frac{u^2 + |\nabla u|^2}{2} \right) + \Phi_2 \left( \frac{v^2 + |\nabla v|^2}{2} \right) \right] dx \\ &\quad - \int_{\Omega} \lambda b dx - \int_{\Omega} \lambda b |u|^{\alpha} dx - \int_{\Omega} \lambda b |u|^{\beta} dx \\ &\geq \frac{\phi_0}{2} \int_{\Omega} [u^2 + |\nabla u|^2 + v^2 + |\nabla v|^2] dx - \lambda b |\Omega| \\ &\quad - \lambda b (c_2)^{\alpha} \|u\|^{\alpha} |\Omega|^{(2-\alpha)/2} - \lambda b (c_2)^{\beta} \|v\|^{\beta} |\Omega|^{(2-\beta)/2} \\ &\geq \frac{\phi_0}{4} \|(u, v)\|_X^2 - \lambda b |\Omega| - \lambda b (c_2)^{\alpha} \|u\|^{\alpha} |\Omega|^{(2-\alpha)/2} - \lambda b (c_2)^{\beta} \|v\|^{\beta} |\Omega|^{(2-\beta)/2}, \end{aligned}$$

for any  $(u, v) \in H^1(\Omega) \times H^1(\Omega)$ . Hence,  $L$  is a coercive functional for every positive parameter  $\lambda$ . So for every

$$\lambda \in \Lambda_{\gamma_2, \sigma_2} = \left( \frac{\phi_m \sigma_2^2}{\inf_{x \in \Omega} G(x, \sigma_2, \sigma_2)}, \frac{\phi_m}{\left( 2a_1 \frac{\bar{A}_1}{\gamma_2} + a_2 \bar{A}_2 \gamma_2^{p-2} + a_2 \bar{A}_3 \gamma_2^{q-2} \right)} \right)$$

the condition  $(a_2)$  in Theorem I holds. All these imply the problem (1.10) has at least three critical points for every  $\lambda \in \Lambda_{\gamma_2, \sigma_2}$ . We reach the result of Theorem 1.3.  $\square$

#### 4.2. Proofs of Theorem 1.4.

The proofs are similar to Theorem 1.2. We will omit them here.  $\square$

Similar to [6, Example 3.1], there are some examples about Theorem 1.1.

**Example.** Let  $\Omega$  be a non-empty bounded open subset of the Euclidean space  $\mathbb{R}^N$ . Put  $p, q \in (1, 2^*)$ ,  $\alpha, \beta < 2$  and

$$w := \max\{1, \tau, (A_1 + A_2 + A_3)^{1/(p-2)} p^{1/(p-2)}, (A_1 + A_2 + A_3)^{1/(q-2)} q^{1/(q-2)}\}.$$

Let  $r_1$  be a positive constant with  $r_1 > w$  and  $G : \Omega \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  and satisfies

$$G_u(x, y, z) := \begin{cases} 1 + |y|^{p-1}, & y \leq r_1, \\ 1 + r_1^{p-\alpha} |y|^{\alpha-1}, & y > r_1, \end{cases}$$

and

$$G_v(x, y, z) := \begin{cases} 1 + |z|^{q-1}, & z \leq r_1, \\ 1 + r_1^{q-\beta} |z|^{\beta-1}, & z > r_1, \end{cases}$$

Clearly,  $(G_1)$  and  $(G_2)$  hold. Moreover, for any  $(s, t) \in \mathbb{R} \times \mathbb{R}$ , it has

$$\begin{aligned} G(x, s, t) &= \int_0^s G_u(x, y, t) dy + \int_0^t G_v(x, 0, z) dz \\ &\leq \left( r_1 + \frac{r_1^\alpha}{r_1} \right) \left( 1 + |\xi|^{\max\{1, \alpha\}} \right) + \left( r_1 + \frac{r_1^\beta}{r_1} \right) \left( 1 + |\eta|^{\max\{1, \beta\}} \right) \\ &\leq 2 \max \left\{ \left( r_1 + \frac{r_1^\alpha}{r_1} \right), \left( r_1 + \frac{r_1^\beta}{r_1} \right) \right\} \left( 1 + |\xi|^{\max\{1, \alpha\}} + |\eta|^{\max\{1, \beta\}} \right). \end{aligned}$$

This shows that  $(G_3)$  holds. Since  $r_1 > w$ , it holds that

$$\frac{G(x, r_1, r_1)}{r_1^2} = \frac{r_1^{p-2}}{p} + \frac{1}{r_1} + \frac{r_1^{q-2}}{q} + \frac{1}{r_1} > A_1 + A_2 + A_3.$$

Particularly, let  $\Omega$  be an open ball with radius one in  $\mathbb{R}^N$ ,  $p = q = 3 \in [1, 2^*)$  and  $\alpha = \beta = \frac{3}{2} < 2$ . Consider the following systems,

$$\begin{cases} -\operatorname{div} \left( \left[ 1 + \left( 1 + \frac{u^2 + |\nabla u|^2}{2} \right)^{-1/2} \right] \nabla u \right) + \left[ 1 + \left( 1 + \frac{u^2 + |\nabla u|^2}{2} \right)^{-1/2} \right] u = \lambda G_u(x, u, v), & x \in \Omega, \\ -\operatorname{div} \left( \left[ 2 + \left( 1 + \frac{v^2 + |\nabla v|^2}{2} \right)^{-1/3} \right] \nabla v \right) + \left[ 2 + \left( 1 + \frac{v^2 + |\nabla v|^2}{2} \right)^{-1/3} \right] v = \lambda G_v(x, u, v), & x \in \Omega, \\ u = v = 0, & x \in \partial\Omega. \end{cases} \quad (4.12)$$

It is easy to see that  $\phi_0 = 1$  and  $\phi_m = 3$ . Following from the equations (6) and (7) in [6], we obtain that

$$\begin{aligned} & (A_1 + A_2 + A_3)^{1/(p-2)} p^{1/(p-2)} \\ &= (A_1 + A_2 + A_3)^{1/(q-2)} q^{1/(q-2)} \\ &= \left[ C_1 \phi_m \left( \frac{2}{\phi_0} \right)^{1/2} + \frac{\phi_m (C_p)^p}{p} \left( \frac{2}{\phi_0} \right)^{3/2} + \frac{\phi_m (C_q)^q}{q} \left( \frac{2}{\phi_0} \right)^{3/2} \right] \\ &= 3(A_1 + A_2 + A_3) \\ &= 3 \left[ 3\sqrt{2}C_1 + 2\sqrt[3]{2^3}(C_3)^3 \right] \\ &\leq 3 \left\{ 3\sqrt{2} \times \frac{\pi}{4} \sqrt[4]{3} + 4\sqrt{2} \times \left[ \left( \frac{\pi^2}{2} \right)^{1/4} \times (8\pi)^{-1/2} \times (6)^{1/4} \right]^3 \right\} \\ &\approx 14.9. \end{aligned}$$

Put  $r_1 = \sigma_1 = 20 > w$ , and  $G_u, G_v : \Omega \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  denoted by

$$G_u(x, y, z) := \begin{cases} 1 + y^2, & y \leq 20, \\ 1 + 20\sqrt{20y}, & y > 20, \end{cases}$$

and

$$G_v(x, y, z) := \begin{cases} 1 + z^2, & z \leq 20, \\ 1 + 20\sqrt{20z}, & z > 20. \end{cases}$$

Thus, for any  $\lambda \in \left( \frac{r_1^2}{G(x, r_1, r_1)}, \frac{1}{A_1 + A_2 + A_3} \right) = \left( \frac{3r_1}{2r_1^2 + 6}, \frac{1}{A_1 + A_2 + A_3} \right) \subseteq (0.07, 4.9)$ , the problem (4.12) has at least three non-trivial solution in  $H_0^1(\Omega) \times H_0^1(\Omega)$ .

The examples about Theorem 1.2, Theorem 1.3 and Theorem 1.4 can be obtained similarly.  $\square$

### Acknowledgement

This work is supported by Yunnan Fundamental Research Projects of China (grant No: 202301AT070465) and Xingdian Talent Support Program for Young Talents of Yunnan Province in China.

### Conflict of interest

On behalf of all authors, the authors states that there is no conflict of interest.

## References

- [1] Averna D, Bonanno G, Tornatore E. Gradient nonlinear elliptic systems driven by a  $(p, q)$ -Laplacian operator. *Dynamic Systems and Applications*, 2017, 26(1): 367-382.
- [2] Afrouzi G A, Heidarkhani S. Existence of three solutions for a class of Dirichlet quasilinear elliptic systems involving the  $p_1, \dots, p_n$ -Laplacian. *Nonlinear Analysis: Theory, Methods and Applications*, 2009, 70(1): 135-143.
- [3] Bonanno G, Marano S A. On the structure of the critical set of non-differentiable functions with a weak compactness condition. *Applicable Analysis*, 2010, 89(1): 1-10.
- [4] Bonanno G, Candito P. Non-differentiable functionals and applications to elliptic problems with discontinuous nonlinearities. *Journal of Differential Equations*, 2008, 244(12): 3031-3059.
- [5] Bonanno G, Heidarkhani S, O'regan D. Multiple solutions for a class of Dirichlet quasilinear elliptic systems driven by a  $(p, q)$ -Laplacian operator. *Dynamic Systems and Applications*, 2011, 20(1): 89.
- [6] Bonanno G, Bisci G M. Three weak solutions for elliptic Dirichlet problems. *Journal of Mathematical Analysis and Applications*, 2011, 382(1): 1-8.
- [7] Bonanno G. A critical point theorem via the Ekeland variational principle. *Nonlinear Analysis: Theory, Methods and Applications*, 2012, 75(5): 2992-3007.
- [8] Bonanno G, Bisci G M, Rădulescu V. Existence of three solutions for a non-homogeneous Neumann problem through Orlicz-Sobolev spaces. *Nonlinear Analysis: Theory, Methods and Applications*, 2011, 74(14): 4785-4795.
- [9] Bonanno G, Candito P. Three solutions to a Neumann problem for elliptic equations involving the  $p$ -Laplacian. *Archiv der Mathematik*, 2003, 80: 424-429.
- [10] Brézis H, Lieb E. A relation between pointwise convergence of functions and convergence of functionals. *Proceedings of the American Mathematical Society*, 1983, 88(3): 486-490.
- [11] Brézis H. *Functional analysis, Sobolev spaces and partial differential equations*. New York: Springer, 2011.
- [12] D'Agui G, Bisci G M. Three non-zero solutions for elliptic Neumann problems. *Analysis and Applications*, 2011, 9(4): 383-394.
- [13] Dal Maso G, Murat F. Almost everywhere convergence of gradients of solutions to nonlinear elliptic systems. *Nonlinear Analysis: Theory, Methods and Applications*, 1998, 31(3-4): 405-412.
- [14] Jeanjean L, Rădulescu V D. Nonhomogeneous quasilinear elliptic problems: linear and sublinear cases. *Journal d'analyse mathématique*, 2020: 1-24.
- [15] Li C, Tang C L. Three solutions for a class of quasilinear elliptic systems involving the  $(p, q)$ -Laplacian. *Nonlinear Analysis: Theory, Methods and Applications*, 2008, 69(10): 3322-3329.
- [16] Landes R. On Galerkin's method in the existence theory of quasilinear elliptic equations. *Journal of Functional Analysis*, 1980, 39(2): 123-148.

- [17] Mihăilescu M, Repovš D. Multiple solutions for a nonlinear and non-homogeneous problem in Orlicz-Sobolev spaces. *Applied Mathematics and Computation*, 2011, 217(14): 6624-6632.
- [18] Pomponio A, Watanabe T. Ground state solutions for quasilinear scalar field equations arising in nonlinear optics. *Nonlinear Differential Equations and Applications NoDEA*, 2021, 28: 1-33.
- [19] Pucci P, Serrin J. A mountain pass theorem. *Journal of Differential Equations*, 1985, 60(1): 142-149.
- [20] Ricceri B. A new method for the study of nonlinear eigenvalue problems. *Comptes Rendus de l'Académie des Sciences-Series I-Mathematics*, 1999, 328(3): 251-256.
- [21] Ricceri B. Existence of three solutions for a class of elliptic eigenvalue problems. *Mathematical and Computer Modelling*, 2000, 32(11-13): 1485-1494.
- [22] Ricceri B. On a three critical points theorem. *Archiv der Mathematik*, 2000, 75(3): 220-226.
- [23] Ricceri B. A three critical points theorem revisited. *Nonlinear Analysis: Theory, Methods and Applications*, 2009, 70(9): 3084-3089.
- [24] Ricceri B. A further three critical points theorem. *Nonlinear analysis: Theory, methods and applications*, 2009, 71(9): 4151-4157.
- [25] Ricceri B. A further refinement of a three critical points theorem. *Nonlinear Analysis: Theory, Methods and Applications*, 2011, 74(18): 7446-7454.
- [26] Stuart C A. Cylindrical TM-modes in a homogeneous self-focusing dielectric. *Mathematical Models and Methods in Applied Sciences*, 1996, 6(07): 977-1008.
- [27] Stuart C A. Magnetic field wave equations for TM-modes in nonlinear optical waveguides. *Reaction diffusion systems*. CRC Press, 2020: 377-400.
- [28] Stuart C A, Zhou H S. Existence of guided cylindrical TM-modes in a homogeneous self-focusing dielectric. *Annales de l'Institut Henri Poincaré C, Analyse non linéaire*. Elsevier Masson, 2001, 18(1): 69-96.
- [29] Stuart C A, Zhou H S. A constrained minimization problem and its application to guided cylindrical TM-modes in an anisotropic self-focusing dielectric. *Calculus of Variations and Partial Differential Equations*, 2003, 16(4): 335-373.
- [30] Stuart C A. Two positive solutions of a quasilinear elliptic Dirichlet problem. *Milan Journal of Mathematics*, 2011, 79(1): 327-341.
- [31] Shokooh S, Graef J R. Existence and multiplicity results for non-homogeneous Neumann problems in Orlicz-Sobolev spaces. *Rendiconti del Circolo Matematico di Palermo Series 2*, 2020, 69(2): 339-351.
- [32] Wang L, Zhang X, Fang H. Multiplicity of solutions for a class of quasilinear elliptic systems in Orlicz-Sobolev spaces. *Taiwanese Journal of Mathematics*, 2017, 21(4): 881-912.
- [33] Yin H, Yang Z. Three solutions for a class of quasilinear elliptic systems involving the  $p(x)$ -Laplace operator. *Boundary Value Problems*, 2012, 2012: 1-10.
- [34] Zeidler E. *Nonlinear functional analysis and its applications: II/B: nonlinear monotone operators*. Springer Science and Business Media, 2013.
- [35] Qi W, Zhang X. Multiplicity of solutions for a nonhomogeneous quasilinear elliptic equation with concave-convex nonlinearities. *arXiv: 2311.00283*.