HOLE PROBABILITIES OF RANDOM ZEROS ON COMPACT RIEMANN SURFACES

HAO WU AND SONG-YAN XIE

ABSTRACT. We establish a convergence speed estimate for hole probabilities of zeros of random holomorphic sections on compact Riemann surfaces.

Mathematics Subject Classification 2020: 31A15, 32C30, 32L10, 60B10, 60D05.

Keywords: line bundle, holomorphic section, hole probability, Abel-Jacobi map, quasipotential, Green function, Fekete points.

CONTENTS

1.	Introduction	1
2.	Preliminaries	5
3.	Regularity of the equilibrium measures	9
4.	Fekete points of the hole events	13
5.	Functional values with Fekete points	18
6.	Constructing holomorphic sections by Fekete points	24
7.	Difference between the two functionals	28
8.	Decay of the hole probabilities	31
9.	Asymptotic behavior of the hole probabilities	33
References		35

1. INTRODUCTION

Distribution of zeros of random polynomials is a classical subject with rich literature, initiated by the works of Waring [48], Bloch-Pólya [7], and henceforth developed intensively by the works of Littlewood-Offord, Hammersley, Kac, Erdös-Turán, see [5, 9, 38] for a review and references therein.

In this article, we study the distribution of zeros of random sections of holomorphic line bundles on compact Riemann surfaces. More precisely, based on the celebrated work [51] of Zelditch (see also [45, 50]) and a recent advance [12], we establish a convergence speed estimate of the hole probabilities. Some inspiration also takes from the approach of [31] on hole probabilities of Gaussian entire functions.

Here is our setting. Let X be a compact Riemann surface, equipped with a Kähler form ω_0 . Let \mathcal{L} be a positive holomorphic line bundle on X, *i.e.*, having degree $\deg(\mathcal{L}) > 0$. Fix a Hermitian metric \mathfrak{h} of \mathcal{L} with strictly positive curvature form $c_1(\mathcal{L}, \mathfrak{h})$. We will regard the normalized (1, 1)-form $\omega := c_1(\mathcal{L}, \mathfrak{h})/\deg(\mathcal{L})$ as a probability measure on X, since $\int_X \omega = 1$.

For every positive integer *n*, the *n*-th power $\mathcal{L}^n := \mathcal{L}^{\otimes n}$ of the line bundle \mathcal{L} inherits a natural metric \mathfrak{h}_n induced by \mathfrak{h} , *i.e.*, $\|s^{\otimes n}(x)\|_{\mathfrak{h}_n} := \|s(x)\|_{\mathfrak{h}}^n$ for local sections *s* of \mathcal{L} around a point $x \in X$. Let $(\cdot, \cdot)_n$ denote the pointwise Hermitian inner product corresponding to

the Hermitian metric \mathfrak{h}_n . On the space $H^0(X, \mathcal{L}^n)$ of global holomorphic sections of \mathcal{L}^n , we can define a global Hermitian inner product

$$\langle s_1, s_2 \rangle_n := \int_X (s_1, s_2)_n \,\omega_0 \quad \text{for} \quad s_1, s_2 \in H^0(X, \mathcal{L}^n).$$

By the Riemann-Roch theorem, the global sections of \mathcal{L}^n are abundant for $n \gg 1$, that is, $\dim_{\mathbb{C}} H^0(X, \mathcal{L}^n) = n \cdot \deg(\mathcal{L}) - g + 1 \gg 1$. On the projectivized space $\mathbb{P}H^0(X, \mathcal{L}^n)$, we consider the Fubini-Study volume form V_n^{FS} induced by $\langle \cdot, \cdot \rangle_n$.

The zero set Z_s of a section s in $H^0(X, \mathcal{L}^n) \setminus \{0\}$, or of an element [s] in $\mathbb{P}H^0(X, \mathcal{L}^n)$, contains $n \cdot \deg(\mathcal{L})$ points counting with multiplicities. Denote by $[Z_s]$ the sum of Dirac masses at the points in Z_s . The Poincaré-Lelong formula states that

$$[Z_s] = \mathrm{dd^c} \log \|s\|_{\mathfrak{h}_n} + n \cdot \mathrm{deg}(\mathcal{L}) \,\omega$$

Here, $d^c := \frac{i}{2\pi}(\overline{\partial} - \partial)$ and $dd^c = \frac{i}{\pi}\partial\overline{\partial}$. The *empirical probability measure* of Z_s is given by the unit mass normalization

$$\llbracket Z_s \rrbracket := (n \cdot \deg(\mathcal{L}))^{-1} [Z_s].$$

The distribution of $\llbracket Z_s \rrbracket$ for $s \in H^0(X, \mathcal{L}^n) \setminus \{0\}$ with respect to the complex Gaussian measure on $H^0(X, \mathcal{L}^n)$ associated with $\langle \cdot, \cdot \rangle_n$ is equal to the distribution of $\llbracket Z_s \rrbracket$ for $[s] \in \mathbb{P}H^0(X, \mathcal{L}^n)$ with respect to V_n^{FS} (cf. [42, Section 2]).

The study of the zero distribution of random holomorphic sections has rich literature, cf. [2, 3, 4, 6, 11, 13, 14, 15, 16, 26, 37, 39, 40, 42, 43, 44]. Notably, specialized in complex dimension one, a celebrated theorem of Shiffman and Zelditch [40] implies that, under the distribution V_n^{FS} , the zeros of sections in $\mathbb{P}H^0(X, \mathcal{L}^n)$ are equidistributed with respect to ω , *i.e.*, for any smooth test function ϕ on X, there holds

$$\lim_{n \to \infty} \int_{\mathbb{P}H^0(X,\mathcal{L}^n)} \langle \llbracket Z_s \rrbracket, \phi \rangle \, \mathrm{d}V_n^{\mathrm{FS}}(s) = \int_X \phi \, \omega.$$

Our goal is to study the hole probabilities of this distribution, in the vein of [12, 45, 50, 51]. Precisely, for an open subset D in X with $\overline{D} \neq X$, for every large n, we consider the hole event

$$H_{n,D} := \left\{ [s] \in \mathbb{P}H^0(X, \mathcal{L}^n), Z_s \cap D = \varnothing \right\}$$

about holomorphic sections of \mathcal{L}^n non-vanishing on D.

To state our main result, we need to employ a key functional $\mathcal{I}_{\omega,D}$ introduced in [12], whose initial model was proposed by Zeitouni-Zelditch [50] (see also [18, 19, 31, 51]). Let $\mathcal{M}(X \setminus D)$ be the set of probability measures supported on $X \setminus D$. Define

(1.1)
$$\mathcal{I}_{\omega,D}(\mu) := -\int_X U_\mu \,\omega - \int_X U_\mu \,\mathrm{d}\mu = -\int_X U'_\mu \,\mathrm{d}\mu + 2\max U'_\mu, \quad \forall \, \mu \in \mathcal{M}(X \setminus D).$$

Here, the function U_{μ} (resp. U'_{μ}) is the ω -potential of type M (resp. type I) of μ , see Section 2.2. When D is non-empty, $\mathcal{I}_{\omega,D}$ is strictly positive. We abbreviate $\mathcal{I}_D := \mathcal{I}_{\omega,D}$ for convenience, as ω is fixed throughout this article. One merit of \mathcal{I}_D is that it admits a unique minimizer $\nu_D \in \mathcal{M}(X \setminus D)$, which coincides with the limit of the conditional expectations $\mathbf{E}(\llbracket Z_s \rrbracket | H_{n,D})$ of the empirical probability measures $\llbracket Z_s \rrbracket$ of the hole events $[s] \in H_{n,D}$ as n tends to infinity. A surprising phenomenon is that the support of ν_D avoids an open "forbidden set" other than D (cf. [12]). Such phenomenon first appeared in the setting of Gaussian entire functions [19, 31].

An open set E in X is said to have smooth boundary ∂E , if for every $x \in \partial E$, there is some neighborhood B of x and a smooth diffeomorphism $\psi : E \to \mathbb{D}$, where \mathbb{D} is the unit disc in \mathbb{C} , such that

$$\psi(E \cap B) = \mathbb{D} \cap \{ \operatorname{Im}(z) > 0 \}, \text{ and } \psi(\partial E \cap B) = \mathbb{D} \cap \mathbb{R}.$$

For technical convenience, from now on, we assume that the hole D has **smooth boundary** and that ∂D **consists of only finitely many components**. Our main result is about the exponential decay of hole probabilities $\mathbf{P}_n(H_{n,D})$.

Theorem 1.1. There exist a $C_D > 0$ independent of n, and a C > 0 independent of n, D, such that

(1.2)
$$-\min \mathcal{I}_D - C_D \frac{\log n}{n} \le \frac{1}{n^2} \log \mathbf{P}_n(H_{n,D}) \le -\min \mathcal{I}_D + C \frac{\log n}{n}$$

for all sufficiently large n.

Here, we assume $n \gg 1$ to ensure that $\mathbf{P}_n(H_{n,D}) \neq 0$, since when $X \neq \mathbb{P}^1$ the hole event $H_{n,D}$ can possibly be empty for certain small $n \in \mathbb{Z}_+$. Note that (1.2) is a strengthen on the convergence speed of the following hole probability result

(1.3)
$$\lim_{n \to \infty} \frac{1}{n^2} \log \mathbf{P}_n(H_{n,D}) = -\min \mathcal{I}_D$$

established in [12, Lemma 7.2]. As a consequence of Theorem 1.1, in dimension one we can recover an estimate established by Shiffman-Zelditch-Zrebiec [45] on the exponential decay $\exp(-c_D n^{t+1})$ of the hole probabilities with respect to a fixed open domain D in a projective manifold of any dimension $t \ge 1$. For the case that $\mathbb{P}^t, t \ge 1$, equipped with the Fubini-Study metric, some good asymptotic formulas of the hole probabilities of polydiscs were established by Zrebiec [53] and Zhu [52]. Nevertheless, even in the simplest case that $(X, \mathcal{L}, \omega) = (\mathbb{P}^1, \mathcal{O}(1), \omega_{\text{FS}})$, the convergence speed estimate $O(\log n/n)$ in Theorem 1.1 is new (cf. [52, 53]).

Our second result concerns asymptotic behavior of the hole probabilities for discs with varying radii. Fix a radius $r_0 > 0$ such that $\overline{\mathbb{B}}(x, r_0) \neq X$ for any $x \in X$.

Theorem 1.2. There exist constants C > c > 0 independent of x, r, such that

(1.4)
$$-Cr^4 \le \lim_{n \to \infty} \frac{1}{n^2} \log \mathbf{P}_n(H_{n,\mathbb{B}(x,r)}) \le -cr^4, \qquad \forall x \in X, \quad \forall 0 < r \le r_0$$

Previously, such hole probabilities with a parameter r on the hole size were not known for zeros of random holomorphic sections, except in some very special cases for which *tour de force* calculation is possible, see [52, 53] for such exceptions in dimension one and higher. The estimate (1.4) is inspired by several classical works in recent decades. In the setting of Gaussian entire functions $F(z) := \sum_{n=1}^{\infty} a_n z^n / \sqrt{n!}$ with the i.i.d. standard complex Gaussian coefficients a_j , Sodin-Tsirelson [47] established that the hole probability with respect to a disc of radius r decays as $\exp(-cr^4)$ when r tends to infinity. An optimal constant c was obtained by Nishry [29]. Recently, Buckley-Nishry-Peled-Sodin [10] studied the hole probabilities for zeros of hyperbolic Gaussian Taylor series with finite radii of convergence. See also [8, 9, 18, 19, 20, 23, 24, 30, 31, 33, 41]. Now we discuss the insight and ideas in the proofs of Theorems 1.1, 1.2. First, let us explain more about the functional \mathcal{I}_D , since it is a crucial ingredient in our approach. Such a type of functional was first proposed by Zeitouni-Zelditch [50] and Zelditch [51] to study large deviations of random zeros of line bundle sections on compact Riemann surfaces. Later, Ghosh-Nishry [18, 19] modified it a bit to study the hole event problem for Gaussian entire functions on \mathbb{C} . Recently, Dinh, Ghosh and the first named author [12] finalized the functional \mathcal{I}_D to the current version, which is near but different to Zeitouni-Zelditch's, and used it to solve the equidistribution problem of hole events, and obtained the exponential decay (1.3) of the hole probabilities as a by-product. The proofs in this article heavily rely on the structure theorem of the equilibrium measure obtained in [12].

We shall emphasize that (1.2) is a strengthen of (1.3). For instance, the lower bound of (1.2) is equivalent to

(1.5)
$$\mathbf{P}_n(H_{n,D}) \ge e^{-C_D n \log n} \cdot e^{-\min \mathcal{I}_D \cdot n^2}, \qquad \forall n \gg 1$$

while (1.3) only means that

(1.6)
$$\mathbf{P}_n(H_{n,D}) \ge e^{-\epsilon_n n^2} \cdot e^{-\min \mathcal{I}_D \cdot n^2}, \qquad \forall n \gg 1$$

for some sequence $\{\epsilon_n\}_{n\gg 1}$ of positive numbers tending to zero.

The proofs of both (1.5) and (1.6) boil down to the following two considerations:

- (\heartsuit) to construct holomorphic sections $s \in H^0(X, \mathcal{L}^n)$ whose zeros avoid the hole D;
- (\diamondsuit) to show that such [s] constitute a subset of $\mathbb{P}H^0(X, \mathcal{L}^n)$ with "large" volume.

As a matter of fact, (1.5) is much more difficult to prove than (1.6). The method of [12] fails to generate enough $[s] \in H_{n,D}$ for (1.5) concerning (\diamondsuit) .

In general, constructing "desired" holomorphic sections is widely known to be a notoriously difficult task, and it lies in the heart of several prominent problems in algebraic and complex geometry (cf. [46]). Let us keep a broader scope in mind that, most of the time, one appeals to three kinds of classical tools:

- (1) methods motivated by electrostatic potentials;
- (2) differential geometric methods involving positive curvature;
- (3) L^2 -methods of solving $\overline{\partial}$ -equations.

Note that (2) and (3) work in any dimension, while (1) is useful only in dimension one.

Our approach is a reminiscence of (1). The starting point is Zelditch's density formula (2.12) (cf. [51, Theorem 2]), whose proof relies on certain bosonization formula [1] and ingenious calculation involving theta functions. In fact, the functional \mathcal{I}_D is obtained by exploring the key exponent in the formula (2.12). Therefore, \mathcal{I}_D serves as a rate function of a large deviation principle (LDP) for the empirical measures $[\![Z_s]\!]$ of zeros of random holomorphic sections $[s] \in H_{n,D}$, and such an idea goes back to Dinh, Ghosh and the first author [12].

The insight is that zeros of a random section in the hole event $H_{n,D}$ shall repel each other, like electrons. The zero sets Z_s of most $[s] \in H_{n,D}$ would produce empirical measures $[\![Z_s]\!]$ very close to the minimizer ν_D of the functional \mathcal{I}_D . This is due to the equidistribution result from [12, Theorem 2].

In practice, the upper bound in Theorem 1.1 is relatively easy to obtain, by means of Zelditch's density formula (2.12). The formidable difficulty lies in the the lower bound, *i.e.* (1.5). Our key strategies concerning (\heartsuit) and (\diamondsuit) consist of two steps.

- (\bigstar) First, to seek some $[s] \in H_{n,D}$ such that a certain necessary modification of \mathcal{I}_D takes value "extremely close" to $\mathcal{I}_D(\nu_D)$ on the empirical measure $[\![Z_s]\!]$. Here "necessary" is due to the trouble that $\mathcal{I}_D(\mu) = +\infty$ for any probability measure μ charging positive mass at some point.
- (\clubsuit) Next, to show that, by "arbitrary" mild perturbations $Z_{s'} \subset X \setminus D$ (but still subject to the Abel-Jacobi equation (2.5)) of the zero set Z_s of s, we can recover lots of $[s'] \in H_{n,D}$ which constitute "large" volume.

The reason for requiring the "extremely closeness" in (\blacklozenge) is to ensure that after mild perturbations $[\![Z_{s'}]\!]$ still take the modified-functional values "very close" to $\mathcal{I}_D(\nu_D)$ so that such $[s'] \in H_{n,D}$ could possibly constitute "large" volume according to Zelditch's density formula (2.12).

When g = 0, such a blueprint can be realized directly, see Remark 8.3 below. However, when $g \ge 1$, one basic difficulty appears as follows, as pointed out by Zelditch [51]. Say n > g and $\deg(\mathcal{L}) = 1$ for simplicity. Note that the zero number of every $[s] \in \mathbb{P}H^0(X, \mathcal{L}^n)$ is presicely n, which is strictly larger than the dimension n - g of $\mathbb{P}H^0(X, \mathcal{L}^n)$. Write the zeros of s in any order as $p_1, \ldots, p_{n-g}, q_1, \ldots, q_g$. For a generic s, by Abel's theorem, the last g zeros q_1, \ldots, q_g are uniquely determined by the first n - g zeros p_1, \ldots, p_{n-g} . However, even if all the p_j 's are outside the hole D, it is not certain whether q_j 's are in $X \setminus D$ or not, *i.e.*, there seems no direct way to parameterize $H_{n,D}$.

Let us highlight three major new ingredients along with the structure of this article.

- We adapt the Fekete points theory on compact Riemann surfaces (Section 4), inspired by Nishry-Wennman [31], so that (♠) can be achieved for every large n (Section 5).
- We show the Hölder regularity of ν_D (Section 3) to guarantee that the zeros of any section s obtained in (♠) are "sufficiently separated" with each other.
- We introduce a new perturbation method (Section 6) to accomplish (\$) (Section 7), in which the key engine is Lemma 6.1.

In the next Section 2, we will present some useful background. The proof of Theorem 1.1 will be reached in Section 8, and Theorem 1.2 in Section 9.

Lastly, we shall emphasize that our approach for Theorem 1.1 is not suitable in higher dimension, because it is based on the Abel-Jacobi theory. The hole probabilities in higher dimensional varieties remain a challenging research direction.

Notations: The symbols \leq and \geq stand for inequalities up to positive multiplicative constants, which may depend on X, ω_0, ω, D but not on any other parameter. The notation O(f), where f is some given positive function, means some term taking values less than $C \cdot f$ for some positive constant C which may depend on D. We denote by $\mathbb{D}(x,r) \subset \mathbb{C}$ the open disc centered at $x \in \mathbb{C}$ with radius r > 0. We abbreviate $\mathbb{D}(0,r)$ as \mathbb{D}_r , and \mathbb{D}_1 as \mathbb{D} . In a Riemann surface or a complex manifold, $\mathbb{B}(x,r)$ stands for the open ball centered at x having radius r with respect to some given metric.

2. PRELIMINARIES

2.1. Abel-Jacobi theory. On a compact Riemann surface X, a *divisor* is a finite formal linear combination $\mathcal{D} := \sum a_j x_j$ with $a_j \in \mathbb{Z}$ and $x_j \in X$. If $a_j \ge 0$ for all j, then we call

 \mathcal{D} an *effective divisor*. The degree deg(\mathcal{D}) of \mathcal{D} means the total sum $\sum a_j$. For a nonzero meromorphic function f on X (resp. a nontrivial holomorphic section s of \mathcal{L}), we write $(f) := (f)_0 - (f)_\infty$ (resp. (s)) the divisor defined by the zeros $(f)_0$ and the poles $(f)_\infty$ of f (resp. zeros of s) respectively. Such a divisor $\mathcal{D} = (f)$ is called a *principal divisor*, and it always has degree 0. A divisor \mathcal{D} on X can be associated with a line bundle $\mathscr{O}(\mathcal{D})$.

Take a canonical basis $\alpha_1, \ldots, \alpha_g, \beta_1, \ldots, \beta_g$ of $H_1(X, \mathbb{Z})$ with the intersection numbers $\alpha_i \cdot \beta_j = \delta_{ij}$ and $\alpha_{j_1} \cdot \alpha_{j_2} = \beta_{j_1} \cdot \beta_{j_2} = 0$ for $j_1 \neq j_2$. Let $\{\phi_1, \ldots, \phi_g\}$ be a basis for the complex vector space of holomorphic differential 1-forms on X. Denote by $\Phi = (\phi_1, \ldots, \phi_g)^T$ a column vector of length g. The $g \times 2g$ matrix

$$\Omega := \left(\int_{\alpha_1} \Phi, \ldots, \int_{\alpha_g} \Phi, \int_{\beta_1} \Phi, \ldots, \int_{\beta_g} \Phi\right).$$

is called a *period matrix*. The 2g columns of Ω are linearly independent over \mathbb{R} , hence generate a lattice Λ in \mathbb{C}^g (cf. [21, 22]). Define the *Jacobian variety* of X by

which is a *g*-dimensional complex torus. Fix a base point $p_{\star} \in X$. The *Abel-Jacobi map* $\mathcal{A}: X \to \text{Jac}(X)$ associated with p_{\star} is defined by

(2.2)
$$\mathcal{A}(x) := \left(\int_{p_{\star}}^{x} \phi_{1}, \dots, \int_{p_{\star}}^{x} \phi_{g}\right) \pmod{\Lambda}, \quad \forall x \in X.$$

The last vector is independent of the choices of path from p_{\star} to x after taking the modulo.

Let $X^{(m)}$ be the *m*-th symmetric product of *X*. Points in $X^{(m)}$ one-to-one correspond to effective divisors $p_1 + \cdots + p_m$ of degree *m*, where points $p_1, \ldots, p_m \in X$ are not necessarily distinct. It is well known that $X^{(m)}$ inherits a complex structure from X^n . Let

(2.3)
$$\pi_m: X^m \to X^{(m)}, \quad (p_1, \dots, p_m) \mapsto p_1 + \dots + p_m$$

be the canonical projection. Let $\mathscr{R}_m := \{(p_1, \ldots, p_m) : p_j = p_\ell \text{ for some } j \neq \ell\}$ be the ramification subvariety. Near a point $p_1 + \cdots + p_m \notin \pi_m(\mathscr{R}_m)$, the holomorphic coordinates of $X^{(m)}$ are given by the local coordinates of $p_1, \ldots, p_m \in X$.

For every $t \ge 1$, define a holomorphic map $\mathcal{A}_t : X^{(t)} \to \operatorname{Jac}(X)$ by

$$\mathcal{A}_t \left(p_1 + \dots + p_t \right) := \sum_{j=1}^t \mathcal{A}(p_j) = \sum_{j=1}^t \left(\int_{p_\star}^{p_j} \phi_1, \dots, \int_{p_\star}^{p_j} \phi_g \right) \pmod{\Lambda}.$$

Then $\mathcal{A}_1 = \mathcal{A}$ is injective, and \mathcal{A}_g is surjective by the Jacobi inversion theorem. Moreover, Riemann showed that \mathcal{A}_g is *birational*, *i.e.*, there exist two subvarieties $\mathscr{W} \subsetneq X^{(g)}$ and $W_q^1 = \mathcal{A}_t(\mathscr{W}) \subsetneq \operatorname{Jac}(X)$, both of codimension 2, such that the restriction

$$\mathcal{A}_g: X^{(g)} \setminus \mathscr{W} \longrightarrow \operatorname{Jac}(X) \setminus W_g^1$$

is biholomorphic. The subvariety \mathscr{W} consists of critical points of \mathcal{A}_g , and W_g^1 is called the *Wirtinger subvariety*. Both of them depend on p_* . For $q_1 + \cdots + q_g \in \mathscr{W}$, one has $\dim H^0(X, \mathscr{O}(q_1 + \cdots + q_g)) \geq 2$ (cf. [21, 22]). When g = 1, \mathscr{W} is empty.

Abel's theorem states that $\mathcal{A}_t(p_1 + \cdots + p_t) = \mathcal{A}_t(p'_1 + \cdots + p'_t)$ if and only if $p_1 + \cdots + p_t - p'_1 - \cdots - p'_t$ is a principal divisor. In particular, there is a holomorphic section $s \in H^0(X, \mathcal{O}(p_1 + \cdots + p_t))$ with $(s) = p'_1 + \cdots + p'_t$.

By the Riemann-Roch theorem, any line bundle on X of degree $\geq g$ has a nontrivial global section s, hence is linearly equivalent to $\mathscr{O}(\mathcal{D})$ for the effective divisor $\mathcal{D} = (s)$. For large n such that $m := n \cdot \deg(\mathcal{L}) - g > 0$, for any $p_1 + \cdots + p_m \in X^{(m)}$, we can apply this observation to the line bundle $\mathcal{L}^n \otimes \mathscr{O}(-(p_1 + \cdots + p_m))$ of degree g to see that there exists some $q_1 + \cdots + q_q \in X^{(g)}$ such that

(2.4)
$$\mathscr{O}(q_1 + \dots + q_g) \simeq \mathcal{L}^n \otimes \mathscr{O}(-(p_1 + \dots + p_m))$$

Equivalently, if $\mathcal{L}^n \simeq \mathscr{O}(w_1 + \cdots + w_{n \cdot \deg(\mathcal{L})})$, then

(2.5)
$$\mathcal{A}_m(p_1 + \dots + p_m) + \mathcal{A}_g(q_1 + \dots + q_g) = \mathcal{A}_n(w_1 + \dots + w_{n \cdot \deg(\mathcal{L})}) =: \mathcal{A}_n(\mathcal{L}^n)$$

Moreover, the choice of $q_1 + \cdots + q_g$ is unique if dim $H^0(X, \mathscr{O}(q_1 + \cdots + q_g)) = 1$, or equivalently, $\mathcal{A}_n(\mathcal{L}^n) - \mathcal{A}_m(p_1 + \cdots + p_m) \notin W_q^1$.

In this article, we will just assume $deg(\mathcal{L}) = 1$, because of the following well-known fact, which is a corollary of the Jacobi inversion theorem.

Lemma 2.1 (cf. [12]–Lemma 6.4). If \mathcal{L}' is positive line bundle on X of degree d > 1, then there exists a positive line bundle \mathcal{L} of degree 1 such that $\mathcal{L}^d \simeq \mathcal{L}'$.

2.2. Quasi-potentials. A function ϕ on X with values in $\mathbb{R} \cup \{-\infty\}$ is quasi-subharmonic if locally it is the difference of a subharmonic function and a smooth one. If ϕ is quasi-subharmonic, then there exists a constant $c \ge 0$ such that $\mathrm{dd}^c \phi \ge -c\omega$ in the sense of currents. When c = 1, ϕ is called an ω -subharmonic function, and $\mathrm{dd}^c \phi + \omega$ is a probability measure on X by Stoke's formula.

For any probability measure μ on X, we can write $\mu = \omega + dd^c U_{\mu}$, where U_{μ} is the unique quasi-subharmonic function such that $\max U_{\mu} = 0$. We call U_{μ} the ω -potential of type M of μ (M stands for "maximum"). There is an alternative way to normalize the potential U'_{μ} by requiring that $\int_X U'_{\nu} \omega = 0$. We call U'_{μ} the ω -potential of type I of μ (I stands for "integration"). To construct ω -potentials, we use the theory of Green functions.

Lemma 2.2 (cf. [25]–Theorem 2.2, [12]–Lemma 2.1). There exists a Green function G(x, y) of (X, ω) defined on $X \times X$, smooth outside the diagonal, such that for every $x \in X$, one has

$$\int_X G(x, \cdot) \, \omega = 0 \qquad \text{and} \qquad \mathrm{dd}^{\mathrm{c}} G(x, \cdot) = \delta_x - \omega$$

Moreover, the function $\rho(x, y) := G(x, y) - \log \operatorname{dist}(x, y)$ is Lipschitz in two variables x, y.

For a probability measure μ on X, the aforementioned ω -potentials are given by

(2.6)
$$U'_{\mu}(x) := \int_{X} G(x, \cdot) d\mu$$
 and $U_{\mu} := U'_{\mu} - \max U'_{\mu}$.

In fact, one can define the ω -potential U'_{ν} for any positive measure or sign measure μ on X by the above formula, which will be used in the study of Fekete points. For any two positive measures or sign measures μ_1, μ_2 on X, it is clear that $U'_{\mu_1+\mu_2} = U'_{\mu_1} + U'_{\mu_2}$. Moreover, one can show by Stoke's formula that (cf. [12, Lemma 2.3])

(2.7)
$$\int_X U'_{\mu_1} \, \mathrm{d}\mu_2 = \int_X U'_{\mu_2} \, \mathrm{d}\mu_1, \quad \text{and} \quad \int_X U'_{\mu_1} \, \mathrm{d}\mu_1 \le 0.$$

There is another geometric description for the ω -potentials. For every $p \in X$, the holomorphic line bundle $\mathscr{O}(p)$ associated with the one-point divisor p has a canonical

section $\mathbf{1}_{\mathscr{O}(p)}$. By the Poincaré-Lelong formula and by the $\partial \overline{\partial}$ -lemma, we can find a smooth Hermitian metric \mathfrak{h}_p for every p such that the curvature form is ω and such that the aforementioned potential reads as

$$U_{\delta_p}(z) = \log \left\| \mathbf{1}_{\mathscr{O}(p)}(z) \right\|_{\mathfrak{h}_p} + C_{\mathfrak{h}_p}$$

where $C_{\mathfrak{h}_p}$ is a normalized constant subject to the law $\max U_{\delta_p} = 0$.

For every point $\mathbf{p} := p_1 + \cdots + p_m \in X^{(m)}, m \ge 1$, we denote by

(2.8)
$$\delta_{\mathbf{p}} := (\delta_{p_1} + \dots + \delta_{p_m})/m$$

the empirical probability measure of p. Similarly, we have

(2.9)
$$U_{\delta_{\mathbf{p}}}(z) = \frac{1}{m} \log \left\| \mathbf{1}_{\mathscr{O}(\mathbf{p})}(z) \right\|_{\mathfrak{h}_{\mathbf{p}}} + C_{\mathfrak{h}_{\mathbf{p}}}$$

where $\mathbf{1}_{\mathscr{O}(\mathbf{p})} := \bigotimes_{j=1}^{m} \mathbf{1}_{\mathscr{O}(p_j)}, \, \mathfrak{h}_{\mathbf{p}} := \bigotimes_{j=1}^{m} \mathfrak{h}_{p_j}, \, \text{and} \, C_{\mathfrak{h}_{\mathbf{p}}} \text{ is a normalized constant subject to the law max } U_{\delta_{\mathbf{p}}} = 0.$

2.3. **Density of zeros.** From now on, we fix a positive line bundle \mathcal{L} on a compact Riemann surface X with degree 1. Let n > g be an integer. Set m := n - g. Define the *exceptional subvariety*

$$\mathbf{H}_m := \left\{ p_1 + \dots + p_m \in X^{(m)} : \mathcal{A}_n(\mathcal{L}^n) - \mathcal{A}_m(p_1, \dots, p_m) \in W_g^1 \right\}$$

of $X^{(m)}$ with codimension 2. Outside \mathbf{H}_m , we have a holomorphic map

$$\mathcal{B}_m: X^{(m)} \setminus \mathbf{H}_m \longrightarrow X^{(g)}$$

given by $\mathcal{B}_m(p_1 + \cdots + p_m) := q_1 + \cdots + q_g$ in (2.4) or (2.5). The image of \mathcal{B}_m is $X^{(g)} \setminus \mathcal{W}$. Consequently, we obtain a holomorphic map

$$\mathscr{A}_m: X^{(m)} \setminus \mathbf{H}_m \longrightarrow \mathbb{P}H^0(X, \mathcal{L}^n),$$

defined by

$$\mathscr{A}_m(p_1 + \dots + p_m) := [s]$$

where $s \in H^0(X, \mathcal{L}^n)$ satisfies $(s) = p_1 + \cdots + p_m + \mathcal{B}_m(p_1 + \cdots + p_m)$. The map \mathscr{A}_m is generically $\binom{n}{m}$ to one. For a measurable set $A \subset \mathcal{M}(X)$, if we set

$$A_{\mathbb{P}} := \left\{ [s] \in \mathbb{P}H^0(X, \mathcal{L}^n) : \llbracket Z_s \rrbracket \in A \right\},\$$

then

(2.10)
$$\mathbf{P}_n(A_{\mathbb{P}}) = \int_{A_{\mathbb{P}}} \mathrm{d}V_n^{\mathrm{FS}} = \binom{n}{m}^{-1} \int_{\mathscr{A}_m^{-1}(A_{\mathbb{P}})} \mathscr{A}_m^*(V_n^{\mathrm{FS}}).$$

Write $\mathbf{q} := \mathcal{B}_m(\mathbf{p}) = q_1 + \cdots + q_g \in X^{(g)}$. We define

(2.11)
$$\mathcal{E}_m(\mathbf{p}) := \frac{1}{m^2} \sum_{j_1 \neq j_2} G(p_{j_1}, p_{j_2}) \text{ and } \mathcal{F}_m(\mathbf{p}) := \log \left\| e^{U'_{\delta_{\mathbf{p}}} + m^{-1}gU'_{\delta_{\mathbf{q}}}} \right\|_{L^{2m}(\omega_0)}.$$

Here, G is the Green function of (X, ω) , and $U'_{\delta_{\mathbf{p}}}, U'_{\delta_{\mathbf{q}}}$ are the ω -potentials of type I.

Zelditch [51, Theorem 2] established the following explicit formula for $\mathscr{A}_m^*(V_n^{\text{FS}})$ (see also [12, Section 4]).

Proposition 2.3. For all $n \ge 2g$ and m = n-g, on $X^{(m)} \setminus \mathbf{H}_m$, the positive measures $\mathscr{A}_m^*(V_n^{FS})$ are of the shape

(2.12)
$$C_n \exp\left[m^2 \left(\mathcal{E}_m(\mathbf{p}) - \frac{2(m+1)}{m} \mathcal{F}_m(\mathbf{p})\right)\right] \kappa_n(\mathbf{p})$$

Here, $\{C_n\}$ are some normalized positive constants, and κ_n are positive continuous (m, m)-forms on $X^{(m)} \setminus \mathbf{H}_m$ given by

$$\kappa_n(\mathbf{p}) := \Upsilon(\mathbf{p})\xi(\mathbf{q}) \prod_{j=1}^m \prod_{l=1}^g \operatorname{dist}(p_j, q_l)^2 i \mathrm{d}z_1 \wedge \mathrm{d}\overline{z}_1 \wedge \cdots \wedge i \mathrm{d}z_m \wedge \mathrm{d}\overline{z}_m,$$

where $e^{-O(m)} \leq \Upsilon(\mathbf{p}) \leq e^{O(m)}$, and $\xi(\mathbf{q})$ is some strictly positive smooth function on $X^{(g)}$ independent of n.

Let us explain the above formula. If $\mathbf{p} \notin \pi_m(\mathscr{R}_m) \cup \mathbf{H}_m$ (see (2.3)), we can use the uniformizing coordinates z_j about p_j for j = 1, ..., m to construct coordinates $(z_1, ..., z_m)$ for $\mathbf{p} \in X^{(m)}$. For $\mathbf{p} \in \pi_m(\mathscr{R}_m) \setminus \mathbf{H}_m$, observing that $\mathcal{E}_m(\mathbf{p}) = -\infty$, we set $\mathscr{A}_m(V_n^{\text{FS}})$ to be 0 on $\pi_m(\mathscr{R}_m) \setminus \mathbf{H}_m$.

The following lemma says hole probability does not quite involve with the choice of ω_0 .

Lemma 2.4 ([12]–Lemma 4.2). One has

$$\sup_{\mathbf{p}\in X^{(m)}} \left| \mathcal{F}_m(\mathbf{p}) - \max U'_{\delta_{\mathbf{p}}} \right| = O\left(\frac{\log m}{m}\right) \quad as \quad m \to \infty,$$

and $U'_{\delta_{\mathbf{p}}} \leq C$ for some constant C independent of m and \mathbf{p} .

To apply Zelditch's density formula (2.12) to study hole probabilities, we need to compute its integration over the set $\mathscr{A}_m^{-1}(H_{n,D})$. Thus, we put $\mathcal{R}_{m,D} := \mathscr{A}_m^{-1}(H_{n,D})$, which can be rephrased as

$$\mathcal{R}_{m,D} = \big\{ \mathbf{p} \in (X \setminus D)^{(m)} \setminus \mathbf{H}_m : \, \mathbf{q} = \mathcal{B}_m(\mathbf{p}) \in (X \setminus D)^{(g)} \big\}.$$

We also define the following separating subset of $\mathcal{R}_{m,D}$:

(2.13)
$$\mathcal{S}_{m,D} := \left\{ \mathbf{p} \in \mathcal{R}_{m,D} : \operatorname{dist}(p_{j_1}, p_{j_2}) \ge m^{-4} \text{ for all } 1 \le j_1 \ne j_2 \le m, \\ \operatorname{dist}(p_j, q_l) \ge 1/m \text{ for all } 1 \le j \le m, 1 \le l \le g \right\},$$

where, $\mathbf{p} := p_1 + \cdots + p_m$ and $\mathbf{q} := q_1 + \cdots + q_g$. This $\mathcal{S}_{m,D}$ will play an important role in the proof of Theorem 1.1.

3. Regularity of the equilibrium measures

From [12, Theorem 1], we know that the minimizer of \mathcal{I}_D is $\nu_D := \omega|_{S_D} + \nu_{\partial D}$, where $S_D := \{U_{\nu_D} = 0\} \setminus \overline{D}$ and $\nu_{\partial D}$ is a non-vanishing positive measure on ∂D . Moreover, if ν_D has mass on some component of ∂D , then $\operatorname{supp}(\nu_D)$ contains this component and U_{ν_D} is strictly negative on this component of ∂D .

In this section, our goal is to prove the following regularity of $\nu_{\partial D}$, which is used for studying the Fekete points in the next section.

Proposition 3.1. For any arc A on ∂D , one has $\nu_{\partial D}(A) \leq c_D \cdot \text{length}(A)$ for some $c_D > 0$ independent of A.

Recall the following result from [12, Proposition 3.1].

Proposition 3.2. Let h be a continuous function on \overline{D} and harmonic on D. Assume that $h + U_{\nu_D} \leq 0$ on \overline{D} , then

$$\int_{\partial D} h \, \mathrm{d}\nu_D \le 0.$$

Suppose that Proposition 3.1 fails, by the following lemma we will construct a harmonic function h such that $h + U_{\nu_D} \leq 0$ on \overline{D} while $\int_{\partial D} h \, d\nu_D > 0$, which is absurd.

Lemma 3.3. Let μ be a probability measure on $\partial \mathbb{D}$ and z be a point in $\partial \mathbb{D}$. Suppose that there exists a sequence $\{r_j\}_{j\geq 1}$ of positive radii shrinking to 0 such that $\mu(\mathbb{D}(z,r_j))/r_j$ tends to infinity. Then there exist a positive sequence $\{\beta_j\}_{j\geq 1}$ tending to infinity and a sequence of functions $\{F_j\}_{j\geq 1}$ which are harmonic on \mathbb{D} and continuous on $\partial \mathbb{D}$, such that $\beta_j r_j$ tends to 0 and such that for j large enough, $F_j < 0$ on $\overline{\mathbb{D}} \setminus \mathbb{D}(z, \beta_j r_j)$ while $\int_{\partial \mathbb{D}} F_j \, \mathrm{d}\mu > 0$.

Proof. Denote $\alpha_j := \mu(\mathbb{D}(z, r_j))$. Take a sequence $\beta_j := 1/\sqrt{\alpha_j r_j}$ so that $\beta_j \alpha_j$ tends to infinity while $\beta_j r_j$ tends to zero. Let g_j be a smooth cut-off function on $\partial \mathbb{D}$ such that

$$g_j = 1$$
 on $\partial \mathbb{D} \cap \mathbb{D}(z, r_j)$ and $g_j = 0$ on $\partial \mathbb{D} \setminus \mathbb{D}(z, 2r_j)$

Set $f_j := g_j - \alpha_j/2$. Let F_j be the harmonic extensions of f_j to \mathbb{D} . It is clear that

$$\int_{\partial \mathbb{D}} F_j \,\mathrm{d}\mu = \int_{\partial \mathbb{D}} g_j \,\mathrm{d}\mu - \alpha_j/2 \ge \int_{\partial \mathbb{D} \cap \mathbb{D}(z,r_j)} \mathbf{1} \,\mathrm{d}\mu - \alpha_j/2 = \alpha_j - \alpha_j/2 > 0$$

for large *j* (in fact, for every *j*, after a moment of reflection).

Now we check that $F_j < 0$ on $\overline{\mathbb{D}} \setminus \mathbb{D}(z, \beta_j r_j)$ for large j. Since $\{\beta_j\}_{j\geq 1}$ tends to infinity, all but finite $\beta_j \geq 4$. In this case, $F_j = -\alpha_j/2 < 0$ on $\partial \mathbb{D} \setminus \mathbb{D}(z, \beta_j r_j)$. By the maximum principle, we only need to show that $F_j < 0$ on $\partial \mathbb{D}(z, \beta_j r_j) \cap \mathbb{D}$. The remaining argument is a standard application of the Poisson formula.

For every $w \in \partial \mathbb{D}(z, \beta_j r_j) \cap \mathbb{D}$, read

$$F_j(w) = \frac{1}{2\pi} \int_0^{2\pi} f_j(e^{i\theta}) \frac{1 - |w|^2}{|e^{i\theta} - w|^2} \,\mathrm{d}\theta = \frac{1}{2\pi} \int_0^{2\pi} g_j(e^{i\theta}) \frac{1 - |w|^2}{|e^{i\theta} - w|^2} \,\mathrm{d}\theta - \alpha_j/2.$$

The support of g_j is in $\partial \mathbb{D} \cap \mathbb{D}(z, 2r_j)$, whose arc length is $(4 + o(1))r_j \leq 5r_j$ for large j. Observe that $1 - |w|^2 \leq 2(1 - |w|) = 2(|z| - |w|) \leq 2|z - w| \leq 2\beta_j r_j$. Moreover, for every $e^{i\theta}$ in $\partial \mathbb{D} \cap \mathbb{D}(z, 2r_j)$, we have $|e^{i\theta} - w| \geq |w - z| - |e^{i\theta} - z| \geq \beta_j r_j - 2r_j \geq \beta_j r_j/2$. Thus we can conclude that

$$F_j(w) \le \frac{1}{2\pi} \cdot 5r_j \cdot \frac{2\beta_j r_j}{(\beta_j r_j/2)^2} - \alpha_j/2 = \frac{20}{\pi\beta_j} - \alpha_j/2 < 0$$

for large *j*.

We have a similar result for the hole *D*.

Lemma 3.4. Let μ be a probability measure on ∂D and z be a point in ∂D . Suppose that there exists a sequence $\{r_j\}_{j\geq 1}$ of positive radii shrinking to 0 such that $\mu(\mathbb{B}(z,r_j))/r_j$ tends to infinity. Then there exist a positive sequence $\{\beta_j\}_{j\geq 1}$ tending to infinity and a sequence of functions $\{F_j\}_{j\geq 1}$ which are harmonic on D and continuous on ∂D , such that $\beta_j r_j$ tends to 0 and such that for j large enough, $F_j < 0$ on $\overline{D} \setminus \mathbb{B}(z, \beta_j r_j)$ while $\int_{\partial D} F_j d\mu > 0$.

$$\Box$$

Proof. If D is simply connected, by the Riemann mapping theorem, we can find a biholomorphism $\phi : D \to \mathbb{D}$. By our assumption on the regularity of ∂D , the Kellogg theorem (cf. [34, p. 49]) guarantees that ϕ (resp. ϕ^{-1}) can be continuously extended to the boundary and is Lipschitz on \overline{D} (resp. $\overline{\mathbb{D}}$). Thus we can apply Lemma 3.3 to conclude.

In general, when D is not simply connected, take a small $\gamma > 0$ such that $\mathbb{B}(z,\gamma)$ intersects only one connected component of D. Reducing γ if necessary, we can draw a simply connected subset $D' \subset D$ with smooth boundary such that $D \cap \mathbb{B}(z,\gamma) \subset D'$.

Denote $\alpha_j := \mu(\mathbb{B}(z, r_j))$. Take $\beta_j := 1/\sqrt{\alpha_j r_j}$, so that $\beta_j \alpha_j$ tends to infinity, while $\beta_j r_j$ tends to zero. Let g_j be a smooth cut-off function on ∂D such that

$$g_j = 1$$
 on $\partial D \cap \mathbb{B}(z, r_j)$ and $g_j = 0$ on $\partial D \setminus \mathbb{B}(z, 2r_j)$

Set $f_j := g_j - \alpha_j/2$. Let F_j, G_j be the harmonic extensions of f_j, g_j respectively to D, whose existence is guaranteed by [49, Theorem I.12] since ∂D contains finitely many components. It is clear that

$$\int_{\partial D} F_j \,\mathrm{d}\mu = \int_{\partial D} g_j \,\mathrm{d}\mu - \alpha_j/2 \ge \alpha_j - \alpha_j/2 > 0$$

for every *j*.

Now we prove that $F_j < 0$ on $\overline{D} \setminus \mathbb{B}(z, \beta_j r_j)$ for large j by means of the maximum principle and the auxiliary simply connected subdomain $D' \subset D$. By throwing some finite j if necessary, we can assume that all $\beta_j r_j < \gamma$, hence $\partial \mathbb{B}(z_j, \beta_j r_j) \cap D$ coincides with $\partial \mathbb{B}(z_j, \beta_j r_j) \cap D'$ by our choice of D'.

Now we consider the restricted function of G_j on $\partial D'$, denoted by h_j . The key observation is that the support of h_j is contained in the union of $\partial D' \cap \mathbb{B}(z, 2r_j)$ and $\partial D' \setminus \partial D$, while these two segments keep relatively large distance compared with $\beta_j r_j$. This feature provides the opportunity for manipulating the Poisson formula on D' in the same spirit as using Katutani's formula on D.

By the Riemann mapping theorem and Kellogg's theorem, we can find a biholomorphism $\phi: D' \to \mathbb{D}$ such that both ϕ and ϕ^{-1} can be continuously extended to the boundaries and are K-Lipschitz with some positive constant K. Since G_j solves the Dirichlet problem on D' with the boundary data h_j , by the Poisson formula, we have

(3.1)
$$G_j(w) = \frac{1}{2\pi} \int_0^{2\pi} h_j \circ \phi^{-1}(e^{i\theta}) \frac{1 - |\phi(w)|^2}{|e^{i\theta} - \phi(w)|^2} \,\mathrm{d}\theta$$

for every w in $\partial \mathbb{B}(z_j, \beta_j r_j) \cap D'$. We only need to integral against two kinds of θ , either $\phi^{-1}(e^{i\theta}) \in \partial D' \cap \mathbb{B}(z, 2r_j)$ or $\phi^{-1}(e^{i\theta}) \in \partial D' \setminus \partial D$, since $h_j \circ \phi^{-1}(e^{i\theta})$ vanishes otherwise. The first kind of integration contributes $O(\beta_j^{-1})$ by the same argument as in the proof of Lemma 3.3, using the fact that ϕ is Lipschitz. For the second kind of θ with $\phi^{-1}(e^{i\theta}) \in \partial D' \setminus \partial D$, noting that the distance between $\partial D' \cap \mathbb{B}(z, \beta_j r_j)$ and $\partial D' \setminus \partial D$ is larger than a fixed positive number while $\beta_j r_j$ tends to zero, we can show that

$$\frac{1 - |\phi(w)|^2}{|e^{i\theta} - \phi(w)|^2} = O(\beta_j r_j).$$

Indeed, $1 - |\phi(w)|^2 \leq 2(|\phi(z)| - |\phi(w)|) \leq 2|\phi(z) - \phi(w)| \leq 2\mathsf{K}\beta_j r_j$, while $|e^{i\theta} - \phi(w)|^2 \geq \mathsf{K}^{-2} \operatorname{dist}(\phi^{-1}(e^{i\theta}), w)^2$, since both ϕ and ϕ^{-1} are K-Lipschitz. Thus the second kind of integration contributes $O(\beta_j r_j)$ because $0 \leq h_j \leq 1$ by the maximum

principle. Summarizing, for every w in $\partial \mathbb{B}(z_j, \beta_j r_j) \cap D'$, we have $G_j(w) = O(\beta_j^{-1}) + O(\beta_j r_j)$. Thus we can bound $F_j(w) = G_j(w) - \alpha_j/2$ from above by

$$O(\beta_j^{-1}) + O(\beta_j r_j) - \alpha_j/2 = \left(O(\beta_j^{-1}) - \alpha_j/4\right) + \left(O(\beta_j r_j) - \alpha_j/4\right).$$

Both brackets $(\cdot \cdot \cdot)$ are negative for large j by our choice of β_j . Hence we conclude the proof by the maximum principle.

Proof of Proposition 3.1. Assume on the contrary that such c_D does not exist. Then by Vitali's covering (e.g. [27, Chapter 2]), we can find some point $z \in \partial D$ with a sequence $\{r_j\}_{j\geq 1}$ of positive radii tending to zero such that $\nu_{\partial D}(\mathbb{B}(z,r_j))/r_j \to \infty$. By considering the probability measure $c\nu_{\partial D}$ for a suitable c and applying Lemma 3.3, we can find a positive sequence $\{\beta_j\}_{j\geq 1}$ tending to zero and a sequence of functions F_j , which are harmonic on D and continuous on ∂D , such that $\beta_j r_j$ tends to zero, $F_j < 0$ on $\overline{D} \setminus \mathbb{B}(z, \beta_j r_j)$, and that $\int_{\partial D} F_j \, d\nu_{\partial D} > 0$ for all j.

On the other hand, recall that U_{ν_D} is strictly negative on the component of ∂D having positive $\nu_{\partial D}$ measure. In particular, $U_{\nu_D}(z) < 0$. By the upper semi-continuity of U_{ν_D} , when $\beta_j r_j$ is small enough, for some small positive constant δ , the function δF_j satisfies that

$$\delta F_j + U_{\nu_D} \le 0 \text{ on } \overline{D} \text{ and } \int_{\partial D} \delta F_j \, \mathrm{d}\nu_D > 0.$$

This contradicts to Proposition 3.2.

Now we can state the regularity result for U_{ν_D} . The proof is standard, which is a consequence of Proposition 3.1. We provide it here for the convenience of the readers.

Proposition 3.5. The quasi-potential U_{ν_D} is 1/3-Hölder.

Proof. Recall the decomposition $\nu_D = \omega|_{S_D} + \nu_{\partial D}$ and $U'_{\nu_D} = U'_{\omega|_{S_D}} + U'_{\nu_{\partial D}}$. It is enough to show that $U'_{\nu_{\partial D}}$ is Hölder. Using (2.7), we have

$$U'_{\nu_{\partial D}}(z) - U'_{\nu_{\partial D}}(w) = \int U'_{\nu_{\partial D}} d(\delta_z - \delta_w) = \int (U'_{\delta_z} - U'_{\delta_w}) d\nu_{\partial D}.$$

By (2.6), we obtain that

$$U'_{\delta_z}(y) - U'_{\delta_w}(y) = \varrho(z, y) - \varrho(w, y) + \log \operatorname{dist}(z, y) - \log \operatorname{dist}(w, y),$$

where ρ is Lipschitz. Hence we only need to estimate the integration

$$\int \left(\log \operatorname{dist}(z, y) - \log \operatorname{dist}(w, y)\right) d\nu_{\partial D}(y)$$

Fix a small a > 0 to be determined. We only need to consider the case that $dist(z, w) \le a$. Let us separate the integration in two parts as

$$\Big(\int_{\operatorname{dist}(z,y)>2a} + \int_{\operatorname{dist}(z,y)\leq 2a}\Big)\Big(\log\operatorname{dist}(z,y) - \log\operatorname{dist}(w,y)\Big)\,\mathrm{d}\nu_{\partial D}(y)$$

If dist(z, y) > 2a, then dist(w, y) > a. Using Proposition 3.1 and mean value theorem, we can obtain that

$$\int_{\operatorname{dist}(z,y)>2a} \left(\log \operatorname{dist}(z,y) - \log \operatorname{dist}(w,y) \right) d\nu_{\partial D}(y) \Big|$$

$$\leq c'_D \int_{\operatorname{dist}(z,y)>2a} \operatorname{dist}(z,w) \cdot \operatorname{dist}(w,y)^{-1} \operatorname{dLeb}(\partial D) \leq c'_D a^{-1} \cdot \operatorname{dist}(z,w),$$

where $\text{Leb}(\partial D)$ is the probability measure on ∂D proportional to arc length. If $\text{dist}(z, y) \leq 2a$, then $\text{dist}(w, y) \leq 3a$. Using Proposition 3.1 again, we get

$$\begin{split} & \Big| \int_{\operatorname{dist}(z,y) \le 2a} \left(\log \operatorname{dist}(z,y) - \log \operatorname{dist}(w,y) \right) \mathrm{d}\nu_{\partial D}(y) \Big| \\ & \le c_D'' \Big| \int_{\operatorname{dist}(z,y) \le 2a} \log \operatorname{dist}(z,y) \operatorname{dLeb}(\partial D) \Big| + c_D'' \Big| \int_{\operatorname{dist}(w,y) \le 3a} \log \operatorname{dist}(w,y) \operatorname{dLeb}(\partial D) \Big|, \end{split}$$

which is $O(c''_D a | \log a |)$. Thus, by taking $a := \sqrt{\operatorname{dist}(z, w)}$, we finish the proof.

Remark 3.6. Actually we can show that U_{ν_D} is Lipschitz. Nevertheless, any γ -Hölder continuity of U_{ν_D} for some $\gamma > 0$ suffices for our purpose. Our boundary assumption on ∂D is also not optimal; as long as ∂D is $\mathscr{C}^{1+\epsilon}$ and does not contain cusps, U_{ν_D} is γ -Hölder continuous for some $\gamma > 0$. To keep this article in an economical length, we prefer not to go further in this direction.

4. FEKETE POINTS OF THE HOLE EVENTS

For proving Theorem 1.1, we need to investigate the minimum value of the functional $-\mathcal{E}_m(\mathbf{p}) + 2 \max U'_{\delta_{\mathbf{p}}}$ defined on $(X \setminus D)^{(m)}$. To this aim, we adapt the theory of logarithmic potentials with external fields (cf. [36, Chapter I]) to compact Riemann surfaces.

For a closed subset *E*, its ω -capacity is defined as

$$\operatorname{cap}_{\omega}(E) := \sup_{\mu \in \mathcal{M}(E)} \exp\Big[\int_X U'_{\mu} \,\mathrm{d}\mu\Big],$$

where U'_{ν} is the ω -potential of type I of ν , see (2.6). If a property holds outside a capacity zero set, then we say that it holds *quasi-everywhere*. It is a fact that if the ω -capacity of E is zero, then the ω' -capacity of E is again zero for any other smooth positive (1, 1)-form ω' on X giving unit area. Another fact is that a capacity zero set is a polar set.

Lemma 4.1. Let μ be a probability measure on X such that $\int_X U'_{\mu} d\mu$ is finite. If $\operatorname{cap}_{\omega}(E) = 0$, then $\mu(E) = 0$.

Proof. Assume that $\mu(E) > 0$ on the contrary. Then we can take a probability measure $\mu_E := c\mu|_E$ for some appropriate c > 0. Note that $\int_X U'_{\mu} d\mu > -\infty$ implies that $\int_X U'_{\mu_E} d\mu_E > -\infty$. Thus we have $\operatorname{cap}_{\omega}(E) \ge \exp\left[\int_X U'_{\mu_E} d\mu_E\right] > 0$, which is absurd. \Box

Let Q be a real continuous function on X. Define the functional

(4.1)
$$\mathcal{J}^Q(\mu) := -\int_X U'_\mu \,\mathrm{d}\mu + 2\int_X Q \,\mathrm{d}\mu$$

for every μ in $\mathcal{M}(X \setminus D)$. Here Q is called an *external field*. We have the following analog of [36, Theorem I.1.3].

Theorem 4.2. The functional \mathcal{J}^Q admits a unique minimizer $\nu_Q \in \mathcal{M}(X \setminus D)$ with continuous ω -potential U'_{ν_Q} of type I. Moreover, one has

(1) $-U'_{\nu_Q} + Q \ge \min \mathcal{J}^Q - \int_X Q \, \mathrm{d}\nu_Q$ on $X \setminus D$;

(2)
$$-U'_{\nu_Q} + Q = \min \mathcal{J}^Q - \int_X Q \,\mathrm{d}\nu_Q$$
 on $\mathrm{supp}(\nu_Q)$.

Proof. The existence of ν_Q follows from the continuity of Q. The uniqueness is based on a well-known fact that $\mu \mapsto -\int_X U'_{\mu} d\mu$ is strictly convex whenever it is finite, see e.g. [50, Proposition 12]. Now we verify the remaining two statements.

We first show that (1) holds on $X \setminus D$ quasi-everywhere. Suppose, on the contrary, that $E := \{ -U'_{\nu_Q} + Q < \min \mathcal{J}^Q - \int_X Q \, \mathrm{d}\nu_Q \} \setminus D$ has positive capacity. Then

(4.2)
$$E_a := \left\{ -U'_{\nu_Q} + Q \le \min \mathcal{J}^Q - \int_X Q \, \mathrm{d}\nu_Q - a \right\} \setminus D$$

also has positive capacity for some small a > 0. Hence we can find some probability measure σ supported on E_a such that $\int_X U'_{\sigma} d\sigma$ is finite. On the other hand, note that

$$\int_{X} (-U'_{\nu_{Q}} + Q) \, \mathrm{d}\nu_{Q} = -\int_{X} U'_{\nu_{Q}} \, \mathrm{d}\nu_{Q} + \int_{X} Q \, \mathrm{d}\nu_{Q} = \min \mathcal{J}^{Q} - \int_{X} Q \, \mathrm{d}\nu_{Q}.$$

Since $-U'_{\nu_O} + Q$ is lower semicontinuous, there exists a closed ball $B \subset X \setminus D$ such that

(4.3)
$$-U'_{\nu_Q} + Q > \min \mathcal{J}^Q - \int_X Q \, \mathrm{d}\nu_Q - a/2 \quad \text{on } B, \quad \text{and} \quad \nu_Q(B) > 0.$$

Set $c := \nu_Q(B) > 0$. Consider the probability measure $\sigma_{\varepsilon} := \nu_Q + \varepsilon(c\sigma - \nu_Q|_B)$ on X for small $\varepsilon > 0$ to be determined. Using (2.7), we can rewrite $\mathcal{J}^Q(\sigma_{\varepsilon})$ as

$$\mathcal{J}^{Q}(\nu_{Q}) - 2\int_{X} U'_{\nu_{Q}} d(\varepsilon c\sigma - \varepsilon \nu_{Q}|_{B}) + 2\int_{X} Q d(\varepsilon c\sigma - \varepsilon \nu_{Q}|_{B}) + O(\varepsilon^{2})$$

$$= \mathcal{J}^{Q}(\nu_{Q}) + 2\int_{X} (-U'_{\nu_{Q}} + Q) d(\varepsilon c\sigma) - 2\int_{X} (-U'_{\nu_{Q}} + Q) d(\varepsilon \nu_{Q}|_{B}) + O(\varepsilon^{2})$$

$$\leq \mathcal{J}^{Q}(\nu_{Q}) - 2 \cdot a \cdot \varepsilon c + 2 \cdot a/2 \cdot \varepsilon c + O(\varepsilon^{2}) \qquad [\text{use (4.2), (4.3)}]$$

$$= \mathcal{J}^{Q}(\nu_{Q}) - ca\varepsilon + O(\varepsilon^{2}).$$

Thus, for ε small enough, $\mathcal{J}^Q(\nu_{\varepsilon}) < \mathcal{J}^Q(\nu_Q)$. This is impossible, because ν_Q is the minimizer of \mathcal{J}^Q .

Next, we prove (2). Since $\int_X U'_{\nu_Q} d\nu_Q$ is finite, the set E of capacity zero must have ν_Q measure 0 by Lemma 4.1. Hence (1) holds ν_Q -almost everywhere. Moreover, $\int_X (-U'_{\nu_Q} + Q) d\nu_Q = \min \mathcal{J}^Q - \int_X Q d\nu_Q$ imples that (2) holds ν_Q -almost everywhere on $\operatorname{supp}(\nu_Q)$. The lower semi-continuity of $-U'_{\nu_Q}$ implies (2) holds everywhere on $\operatorname{supp}(\nu_Q)$.

Lastly, we come back to prove (1). From (2), we see that U'_{ν_Q} is continuous as a function on $\operatorname{supp}(\nu_Q)$. By a classical result [49, p. 54, Theorem III.2], U'_{ν_Q} is continuous on the whole X. Hence (1) holds everywhere on $X \setminus D$.

Theorem 4.3. One has

$$U'_{\nu_Q} + \min \mathcal{J}^Q - \int_X Q \, \mathrm{d}\nu_Q = \sup_{\phi} \left\{ \phi \text{ is } \omega \text{-subharmonic} : \phi \leq Q \text{ on } X \setminus D \right\}^*.$$

Here * means the upper semicontinuous regularization.

Proof. By Theorem 4.2, $U'_{\nu_Q} + \min \mathcal{J}^Q - \int_X Q \, d\nu_Q$ itself is an ω -subharmonic function which is bounded from above by Q on $X \setminus D$. Hence the direction " \leq " is done.

Denote the function on the right hand side by V, which is ω -subharmonic as well. For the probability measure $\nu := \mathrm{dd}^{\mathrm{c}}V + \omega$, we have $\mathrm{dd}^{\mathrm{c}}(V - U'_{\nu_{Q}}) = \nu - \nu_{Q}$. Thus $V - U'_{\nu_{Q}}$ is subharmonic on $X \setminus \text{supp}(\nu_Q)$. By Theorem 4.2, $Q = U'_{\nu_Q} + \min \mathcal{J}^Q - \int_X Q \, d\nu_Q$ on $\text{supp}(\nu_Q)$ while U'_{ν_Q} is continuous, hence $V = U'_{\nu_Q} + \min \mathcal{J}^Q - \int_X Q \, d\nu_Q$ on $\text{supp}(\nu_Q)$. By the maximum principle, $V - U'_{\nu_Q} \leq \min \mathcal{J}^Q - \int_X Q \, d\nu_Q$ on $X \setminus \text{supp}(\nu_Q)$. This shows " \geq ". \Box

For each m > 1 and $\mathbf{p} \in (X \setminus D)^{(m)}$, define (see (2.8), (2.11))

$$\mathcal{K}_m^Q(\mathbf{p}) := -\frac{m}{m-1} \mathcal{E}_m(\mathbf{p}) + 2 \int_X Q \,\mathrm{d}\delta_{\mathbf{p}}.$$

Let $\mathbf{f}_m = \sum_{j=1}^m f_{m,j}$ be a minimizer of \mathcal{K}_m^Q . It may not be unique and we just fix one. We call $\mathscr{F}_m := \{f_{m,1}, \ldots, f_{m,m}\}$ an *m*-th Fekete set and call $f_{m,j}$ Fekete points. Obviously, all the points in $\mathscr{F}_m \subset X \setminus D$ are pairwise distinct.

Theorem 4.4. One has $-U'_{\nu_Q}(x) + Q(x) = \min \mathcal{J}^Q - \int_X Q \, d\nu_Q$ for every x in \mathscr{F}_m .

Proof. By Theorem 4.2, we only need to show that $-U'_{\nu_Q}(x) + Q(x) \leq \min \mathcal{J}^Q - \int_X Q \, d\nu_Q$ every x in \mathscr{F}_m . Without loss of generality, we take $x = f_{m,1}$. Define the function

$$F(z) := \mathcal{K}_m^Q(z + f_{m,2} + \dots + f_{m,m})$$

for $z \in X \setminus D$, which attends a minimum value at $z = f_{m,1}$. Write $\mathbf{f}'_m := f_{m,2} + \cdots + f_{m,m}$. By the definition of \mathcal{E}_m , the new function (see (2.6))

$$\widetilde{F}(z) := -\frac{m}{m-1} \frac{2}{m^2} \sum_{j \ge 2} G(z, f_{m,j}) + \frac{2}{m} Q(z) = \frac{2}{m} \Big(-U'_{\delta_{\mathbf{f}'_m}}(z) + Q(z) \Big)$$

must attend its minimum on $X \setminus D$ at $z = f_{m,1}$. In other words,

(4.4)
$$-U'_{\delta_{\mathbf{f}'_m}}(f_{m,1}) + Q(f_{m,1}) \le -U'_{\delta_{\mathbf{f}'_m}}(z) + Q(z), \quad \forall z \in X \setminus D$$

Now we consider the auxiliary function $\phi(z) := U'_{\delta_{\mathbf{f}'_m}}(z) - U'_{\delta_{\mathbf{f}'}}(f_{m,1}) + Q(f_{m,1})$ on X. Clearly, it is ω -subharmonic and bounded from above by Q on $X \setminus D$. By Theorem 4.3,

$$\phi(z) \le U'_{\nu_Q}(z) + \min \mathcal{J}^Q - \int_X Q \,\mathrm{d}\nu_Q, \quad \forall \, z \in X.$$

Taking $z = f_{m,1}$, we obtain the desired inequality.

Using the theorems above, one can show that $\delta_{\mathbf{f}_m}$ converges to ν_Q weakly. Since we do not need this in this article, we left the proof to the readers. Now we show that any two Fekete points cannot be too close.

Proposition 4.5. If U'_{ν_Q} is γ -Hölder for some $0 < \gamma \leq 1$, then there exists a constant $c_Q > 0$ independent of m such that

$$dist(f_{m,j_1}, f_{m,j_2}) \ge c_Q m^{-1/\gamma}, \quad if \quad j_1 \neq j_2.$$

Some arguments in the proof are inspired by [31, Proposition A.2], though technically much more involved. One new difficulty is to find appropriate local holomorphic coordinate charts on X to apply the Schwarz lemma. Another new difficulty is to deal with the local weight functions of certain holomorphic line bundles. The next two lemmas will be useful for handling the mentioned two difficulties respectively.

Lemma 4.6. Let $g : \mathbb{D} \to \mathbb{C}$ be a holomorphic function. Let x, y be two points in \mathbb{D} . Assume that g(y) = 0, |g(x)| = a, and that $\sup_{U} |g| < b$ for some constant a, b > 0, where $U \subset \mathbb{D}$ is a simply connected region containing x. Then $|x - y| \ge \frac{a}{a+b} \cdot \operatorname{dist}(x, \partial U)$.

Proof. Assume on the contrary that $|x - y| < \frac{a}{a+b} \cdot \operatorname{dist}(x, \partial U)$. Then $y \in U$, and

$$\operatorname{dist}(y,\partial U) \ge \operatorname{dist}(x,\partial U) - |x-y| \ge \frac{b}{a+b} \cdot \operatorname{dist}(x,\partial U) > \frac{b}{a} \cdot |x-y|$$

Hence the disc $D' := \mathbb{D}(y, \frac{b}{a} \cdot |x - y|)$ centered at y with the radius $\frac{b}{a} \cdot |x - y|$ is contained in U. By the maximum principle, $\sup_{D'} |g| \leq \sup_{U} |g| < b$. Thus by the Schwarz lemma, noting that g(y) = 0, we obtain that

$$|g(x)| = |g(y + (x - y))| \le \frac{|x - y|}{\frac{b}{a} \cdot |x - y|} \cdot \sup_{D'} |g| < \frac{a}{b} \cdot b = a,$$

which is absurd.

Lemma 4.7. Let \mathcal{L} be a holomorphic line bundle on X, and let $U \subset X$ be a simply connected region. Let $\mathfrak{h}_1, \mathfrak{h}_2$ be two smooth metrics of \mathcal{L} over U, giving the same curvature. Then for any holomorphic section $s_1 \in \Gamma(U, \mathcal{L})$ of \mathcal{L} on U, one can find a holomorphic section $s_2 \in \Gamma(U, \mathcal{L})$ with $\|s_2\|_{\mathfrak{h}_2} = \|s_1\|_{\mathfrak{h}_1}$ on U.

Proof. We can define the ratio $\mathfrak{h}_2/\mathfrak{h}_1$ at every point of U, and check that it is a positive smooth function on U. Since the curvature of \mathfrak{h}_1 and \mathfrak{h}_2 coincide, $\log(\mathfrak{h}_2/\mathfrak{h}_1)$ is a harmonic function on U. As U is simply connected, we can write $\log(\mathfrak{h}_2/\mathfrak{h}_1) = \operatorname{Re}(u)$ as the real part of a holomorphic function $u \in \mathcal{O}(U)$. Hence $s_2 := e^u \cdot s_1$ suffices.

Proof of Proposition 4.5. Step 1. By symmetry, we only need to show the existence of $c_Q > 0$ and $M_Q \in \mathbb{Z}_+$ such that $\operatorname{dist}(f_{m,1}, f_{m,2}) \ge c_Q m^{-1/\gamma}$ for all $m \ge M_Q$.

Write $\mathbf{f}'_m := f_{m,2} + \cdots + f_{m,m}$. For $z \in \operatorname{supp}(\nu_Q)$, by Theorems 4.2, 4.4 and (4.4), we have

$$U'_{\delta_{\mathbf{f}'_m}}(z) - U'_{\delta_{\mathbf{f}'_m}}(f_{m,1}) \le Q(z) - Q(f_{m,1}) = U'_{\nu_Q}(z) - U'_{\nu_Q}(f_{m,1}).$$

Consequently, $U'_{\delta_{\mathbf{f}'_m}}(z) - U'_{\nu_Q}(z) \leq U'_{\delta_{\mathbf{f}'_m}}(f_{m,1}) - U'_{\nu_Q}(f_{m,1})$ for $z \in \operatorname{supp}(\nu_Q)$. Since U'_{ν_Q} is harmonic outside $\operatorname{supp}(\nu_Q)$, $U'_{\delta_{\mathbf{f}'_m}} - U'_{\nu_Q}$ is subharmonic on $X \setminus \operatorname{supp}(\nu_Q)$. By the maximum principle,

(4.5)
$$U'_{\delta_{\mathbf{f}'_m}}(z) - U'_{\delta_{\mathbf{f}'_m}}(f_{m,1}) \le U'_{\nu_Q}(z) - U'_{\nu_Q}(f_{m,1}) \quad \text{for all} \quad z \in X.$$

Since U'_{ν_Q} is γ -Hölder continuous with some constant $A := ||U'_{\nu_Q}||_{\mathscr{C}^{\gamma}}$, if $\operatorname{dist}(z, f_{m,1}) \leq (m-1)^{-1/\gamma}$, the right-hand-side of (4.5) is $\leq A \cdot (m-1)^{-1}$. Hence (2.9) implies that

$$\log \left\| \mathbf{1}_{\mathscr{O}(\mathbf{f}'_m)}(z) \right\|_{\mathfrak{h}_{\mathbf{f}'_m}} - \log \left\| \mathbf{1}_{\mathscr{O}(\mathbf{f}'_m)}(f_{m,1}) \right\|_{\mathfrak{h}_{\mathbf{f}'_m}} \le A \cdot (m-1)^{-1},$$

or equivalently (let \mathfrak{h}_j be the metric of $\mathscr{O}(f_{m,j})$)

(4.6)
$$\frac{\prod_{j=2}^{m} \|\mathbf{1}_{\mathscr{O}(f_{m,j})}\|_{\mathfrak{h}_{j}}(z)}{\prod_{j=2}^{m} \|\mathbf{1}_{\mathscr{O}(f_{m,j})}\|_{\mathfrak{h}_{j}}(f_{m,1})} \le e^{A}, \qquad \forall z \in \mathbf{B}_{m} := \mathbb{B}(f_{m,1}, (m-1)^{-1/\gamma}).$$

Step 2. The idea is to show that $f_{m,2}$ can not be too close to $f_{m,1}$ by exploring (4.6). First, we make some technical preparations which will be essential later.

For every point $y \in X$, we take a local holomorphic chart $g_y : \mathbb{D}_2 \to V_y$ with $g_y(0) = y$. Since $\{g_y(\mathbb{D})\}_{y \in X}$ covers X, by compactness, we can take finitely many points y_1, \ldots, y_M such that X is covered by the union of $W_j := g_{y_j}(\mathbb{D})$ for $j = 1, \ldots, M$.

Since each $g_{y_j} : \mathbb{D}_2 \to V_{y_j}$ is a biholomorphism, by compactness, both the restrictions $g_{y_j}|_{\overline{\mathbb{D}}}$ and $g_{y_j}^{-1}|_{\overline{W_j}}$ are K_j -Lipschitz for some large $\mathsf{K}_j > 0$ with respect to the Euclidean norm on \mathbb{D} and the given distance on X. Set $\mathsf{K} := \max\{\mathsf{K}_1, \ldots, \mathsf{K}_M\}$.

For each j = 1, 2, ..., m, we fix a smooth weight function ρ_j on V_{y_j} satisfying the equation $dd^c \rho_j = \omega$. By compactness, we can take some large constant K' > 0 such that all ρ_j are K'-Lipschitz on W_j for j = 1, 2, ..., M.

Step 3. By compactness argument, there exists some uniform radius r > 0 such that for any $z \in X$, the small disc $\mathbb{B}(z,r)$ is contained entirely in one of W_1, \ldots, W_M . Take a large integer M_Q so that $(m-1)^{-1/\gamma} < r$ for all $m \ge M_Q$. Without loss of generality, we can assume that $\mathbf{B}_m \subset W_1$.

Since W_1 is open, any holomorphic line bundle on W_1 is holomorphically trivial (cf. [17, p. 229]). In particular, for each line bundle $\mathcal{O}(f_{m,j})$, j = 2, ..., m, we can fix a nowhere vanishing section $e_j \in \Gamma(W_1, \mathcal{O}(f_{m,j}))$ and use it to trivialize $\mathcal{O}(f_{m,j})|_{W_1}$ by identifying every holomorphic section $s \in \Gamma(W_1, \mathcal{O}(f_{m,j}))$ with the holomorphic function $s/e_j \in \mathcal{O}(W_1)$. In accordance with this trivialization, the metric $\mathfrak{h}_j|_{W_1}$ of $\mathcal{O}(f_{m,j})|_{W_1}$ defines a smooth weight function φ_j on W_1 by the law

$$||s||_{\mathfrak{h}_j}(z) = |s/e_j|(z) \cdot e^{-\varphi_j(z)}, \qquad \forall z \in W_1.$$

Since the curvature of \mathfrak{h}_i is ω , we have $\mathrm{dd}^{\mathrm{c}}\varphi_i = \omega$.

In this way, each canonical section $\mathbf{1}_{\mathscr{O}(f_{m,j})} \in \Gamma(W_1, \mathscr{O}(f_{m,j}))$ for $j = 2, \ldots, m$ has norm

$$|\mathbf{1}_{\mathscr{O}(f_{m,j})}\|_{\mathfrak{h}_j}(z) = |h_j(z)| \cdot e^{-\varphi_j(z)}, \qquad \forall z \in W_1,$$

where $h_j := \mathbf{1}_{\mathscr{O}(f_{m,j})}/e_j$ is some holomorphic functions. Hence we can read (4.6) as

(4.7)
$$\frac{\prod_{j=2}^{m} |h_j(z)| \cdot \prod_{j=2}^{m} e^{-\varphi_j(z)}}{\prod_{j=2}^{m} |h_j(f_{m,1})| \cdot \prod_{j=2}^{m} e^{-\varphi_j(f_{m,1})}} \le e^A, \qquad \forall z \in \mathbf{B}_m$$

Step 4. For some technical reason to be seen later, we would like to have some uniform Lipschitz bound of each φ_j on W_1 . One way to achieve this is by using some singularity decomposition result (see Lemma (2.2)) of the Green kernel G(x, y) near the diagonal $\{x = y\} \subset X \times X$ and by choosing appropriate e_j, φ_j .

Nevertheless, here we present an alternative method. To apply Lemma (4.7), for every j = 2, ..., m, we define another metric \mathfrak{h}'_j of the line bundle $\mathscr{O}(f_{m,j})$ on W_1 , by evaluating any holomorphic section $s \in \Gamma(W_1, \mathscr{O}(f_{m,j}))$ with the norm $\|s\|_{\mathfrak{h}'_j}(z) = |s/e_j|(z) \cdot e^{-\rho_1(z)}$ at each point $z \in W_1$. Since $\mathrm{dd}^c \rho_1 = \omega$, the curvature of \mathfrak{h}'_j coincide with that of \mathfrak{h}_j on W_1 . By Lemma (4.7), we can find some holomorphic section $\mathbf{1}'_j \in \Gamma(W_1, \mathscr{O}(f_{m,j}))$ with the norm equality $\|\mathbf{1}_{\mathscr{O}(f_{m,j})}\|_{\mathfrak{h}_j} = \|\mathbf{1}'_j\|_{\mathfrak{h}'_j}$ on W_1 . Equivalent, we have

$$|h_j(z)| \cdot e^{-\varphi_j(z)} = |\widehat{h_j}(z)| \cdot e^{-\rho_1(z)}, \qquad \forall z \in W_1,$$

where $\hat{h_j} := \mathbf{1}'_j / e_j \in \mathscr{O}(W_1)$ is the representative holomorphic function of $\mathbf{1}'_j$ in the local trivialization. Thus we can rewrite (4.7) as

(4.8)
$$\frac{\prod_{j=2} \hat{h}_j(z)}{\prod_{j=2} \hat{h}_j(f_{m,1})} \cdot e^{\sum_{j=2}^m [\rho_1(f_{m,1}) - \rho_1(z)]} \le e^A, \qquad \forall z \in \mathbf{B}_m.$$

Note that by our preparation, the weight function ρ_1 is K'-Lipschitz on W_1 , which contains \mathbf{B}_m . Hence we have

$$\sum_{j=2}^{m} |\rho_1(f_{m,1}) - \rho_1(z)| \le (m-1) \cdot \mathsf{K}' \cdot (m-1)^{-1/\gamma} \le \mathsf{K}', \qquad \forall z \in \mathbf{B}_m.$$

By the above two estimates, the holomorphic function

$$H := \prod_{j=2}^{m} \frac{\widehat{h}_j}{\widehat{h}_j(f_{m,1})}$$

which is well-defined on W_1 , has absolute norm $\leq e^{A+\mathsf{K}'}$ on \mathbf{B}_m .

Step 5. Assume on the contrary that $dist(f_{m,1}, f_{m,2}) < \lambda \cdot (m-1)^{-1/\gamma}$ for some sufficiently small λ . It is clear that $f_{m,2} \in \mathbf{B}_m$, $H(f_{m,1}) = 1$, and that $H(f_{m,2}) = 0$ because the canonical section $\mathbf{1}_{\mathscr{O}(f_{m,2})}$ vanishes at $f_{m,2}$ by definition.

Now we are in a position to apply Lemma 4.6 upon

$$g := H \circ g_{y_1}, \ x := g_{y_1}^{-1}(f_{m,1}), \ y := g_{y_1}^{-1}(f_{m,2}), \ a = 1, \ b := e^{A + \mathsf{K}'}, \ U := g_{y_1}^{-1}(\mathbf{B}_m).$$

Since g_{y_1} is K-Lipschitz by our preparation,

$$(m-1)^{-1/\gamma} = \operatorname{dist}(f_{m,1}, \partial \mathbf{B}_m) = \operatorname{dist}(g_{y_1}(x), g_{y_1}(\partial U)) \le \mathsf{K} \cdot \operatorname{dist}(x, \partial U).$$

Thus Lemma 4.6 guarantees that

$$|x-y| \ge \frac{1}{1+e^{A+\mathsf{K}'}} \cdot \operatorname{dist}(x,\partial U) \ge \frac{1}{1+e^{A+\mathsf{K}'}} \cdot \mathsf{K}^{-1} \cdot (m-1)^{-1/\gamma}.$$

On the other hand, $g_{y_1}^{-1}$ is also K-Lipschitz. Hence

$$|x-y| = |g_{y_1}^{-1}(f_{m,1}) - g_{y_1}^{-1}(f_{m,2})| \le \mathsf{K} \cdot \operatorname{dist}(f_{m,1}, f_{m,2}).$$

Summarizing the above two inequalities, we obtain

dist
$$(f_{m,1}, f_{m,2}) \ge \frac{1}{1 + e^{A+B}} \cdot \mathsf{K}^{-2} \cdot (m-1)^{-1/\gamma}$$

Therefore, if $\lambda < \frac{K^{-2}}{1+e^{A+K'}}$, we get a contradiction. Hence $c_Q = \frac{K^{-2}}{1+e^{A+K'}}$ suffices.

5. FUNCTIONAL VALUES WITH FEKETE POINTS

This goal of this section is to find a $\mathbf{p} \in (X \setminus D)^{(m)}$ such that

(5.1)
$$-\mathcal{E}_m(\mathbf{p}) + 2\max U'_{\delta_{\mathbf{p}}} \le \min \mathcal{I}_D + O(\log m/m).$$

The minimizer of \mathcal{I}_D is ν_D . While we do not know much about $-\mathcal{E}_m(\mathbf{p}) + 2 \max U'_{\delta_{\mathbf{p}}}$. We will make use of the two functionals \mathcal{J}^Q and \mathcal{K}^Q_m defined in Section 4. These functionals are linked according through the following result.

Proposition 5.1. For $Q := U'_{\nu_D}$, one has $\nu_Q = \nu_D$.

Proof. Since $supp(\nu_D) \subset X \setminus D$, integrating the first inequality in Theorem 4.2 against ν , we get

$$-\int U'_{\nu_Q} \,\mathrm{d}\nu_D + \int U'_{\nu_D} \,\mathrm{d}\nu_D \ge \min \mathcal{J}^Q - \int_X U'_{\nu_D} \,\mathrm{d}\nu_Q.$$

Using (2.7), we obtain $\min \mathcal{J}^Q \leq \int U'_{\nu_D} d\nu_D$.

On the other hand, Q itself is an ω -subharmonic function and is bounded from above by Q on $X \setminus D$. By Theorem 4.3, we have

$$U'_{\nu_D} \le U'_{\nu_Q} + \min \mathcal{J}^Q - \int_X U'_{\nu_D} \,\mathrm{d}\nu_Q.$$

Integrating this inequality against ν_D , we get $\int U'_{\nu_D} d\nu_D \leq \min \mathcal{J}^Q$. By the uniqueness of the minimizer of the functional \mathcal{J}^Q (see Theorem 4.2), we conclude that $\nu_Q = \nu_D$.

The functional $\mathcal{E}_m(\mathbf{p})$ can be viewed as a discrete version of $\int U'_{\mu} d\mu$. Thus, the functionals $-\mathcal{E}_m(\mathbf{p}) + 2 \max U'_{\delta_{\mathbf{p}}}, \mathcal{K}^Q_m(\mathbf{p})$ are discrete versions of $\mathcal{I}_D(\mu), \mathcal{J}^Q(\mu)$ respectively. The advantage of introducing \mathcal{J}^Q is that under the setting of complex plane $\mathbb{C}, \mathcal{J}^Q$ was well-studied, see e.g. [36]. Thus, using Proposition 5.1, we will find a **p** satisfying (5.1), from the Fekete points of $\mathcal{K}^Q_m(\mathbf{p})$.

From now on, we fix the external field $Q := U'_{\nu_D}$, which is 1/3-Hölder continuous by Proposition 3.5. Hence Proposition 4.5 guarantees that distinct Fekete points $f_{m,1}, \ldots, f_{m,m}$ are separated for every large $m \gg 1$, with a distance estimate $\operatorname{dist}(f_{m,j_1}, f_{m,j_2}) \ge c_Q m^{-3}$ for $j_1 \neq j_2$ (we still use c_Q for convenience in this section, which is depending on D).

The following probability measure η will be used two times later.

Lemma 5.2. Let E be a subset of X (not necessary open) with with finitely many connected components, whose boundary $\overline{E} \setminus \operatorname{int}(E)$ consists of disjoint smooth curves. Then there exists a probability measure η supported on \overline{E} , such that U'_{η} is constant on \overline{E} and attends its minimum on \overline{E} .

Proof. Consider

$$\Psi := \sup_{\phi} \left\{ \phi \text{ is } \omega \text{-subharmonic} : \phi \leq 0 \text{ on } E \right\}^*.$$

Let $\eta := dd^c \Psi + \omega$. The remaining proof goes the same as [50, Theorem 21]. This η is also the minimizer of the functional

$$\mu \mapsto -\int_X U'_{\mu} d\mu \quad \text{for} \quad \mu \in \mathcal{M}(\overline{E}).$$

The assumptions on the boundary of E is to ensure the continuity of U'_n .

Let $\sigma_{m,j}$ be the probability measure on $\partial \mathbb{B}(f_{m,j}, c_Q m^{-5})$ induced by arc lengths. Define the probability measure

$$\sigma_m := \frac{1}{m} \sum_{j=1}^m \sigma_{m,j}.$$

We will use σ_m to approximate $\delta_{\mathbf{f}_m}$, because $\int U'_{\delta_{\mathbf{f}_m}} d\delta_{\mathbf{f}_m} = -\infty$, while $\int U'_{\sigma_m} d\sigma_m > -\infty$.

Lemma 5.3. There exists some constant $C_D > 0$, such that for all sufficiently large $m \gg 1$, for any $w \in \partial \mathbb{B}(f_{m,j}, c_Q m^{-5})$, one has

$$\left|U'_{\sigma_m}(w) - U'_{\mathbf{f}_m}(w)\right| \le C_D \log m/m.$$

Proof. We may assume $w \in \partial \mathbb{B}(f_{m,1}, c_Q m^{-5})$. The argument is by straightforward computation, using (2.6) and Lemma 2.2, $|U'_{\sigma_m}(w) - U'_{\mathbf{f}_m}(w)|$ is bounded by

$$\left|\int \log \operatorname{dist}(w, x) \operatorname{d}(\sigma_m - \mathbf{f}_m)(x)\right| + \left|\int \varrho(w, x) \operatorname{d}(\sigma_m - \mathbf{f}_m)(x)\right|.$$

Since ρ is Lipschitz, the second term is $O(m^{-5})$. By definition, the first term is bounded by

$$\frac{1}{m}\sum_{\ell=1}^{m}\int \left|\log\operatorname{dist}(w,x) - \log\operatorname{dist}(w,f_{m,\ell})\right| \mathrm{d}\sigma_{m,\ell}(x).$$

When $\ell \neq 1$, recall the support of $\sigma_{m,\ell}$ is $\partial \mathbb{B}(f_{m,\ell}, c_Q m^{-5})$ and $\operatorname{dist}(w, f_{m,\ell}) > c_Q m^{-3}/2$. By the monotonicity of the log function, for $x \in \operatorname{supp}(\sigma_{m,\ell})$,

$$\log \operatorname{dist}(w, x) - \log \operatorname{dist}(w, f_{m,\ell}) | \leq \\ \max \left(\log \frac{\operatorname{dist}(w, f_{m,\ell})}{\operatorname{dist}(w, f_{m,\ell}) - c_Q m^{-5}}, \log \frac{\operatorname{dist}(w, f_{m,\ell}) + c_Q m^{-5}}{\operatorname{dist}(w, f_{m,\ell})} \right) = \frac{O(1)}{m^2}.$$

It remains to estimate the first term with $\ell = 1$. Since $\sigma_{m,1}$ is the probability measure on $\mathbb{B}(f_{m,1}, c_Q m^{-5})$ induced by arc length, we have

$$\int \left|\log\operatorname{dist}(w,x) - \log\operatorname{dist}(w,f_{m,1})\right| \mathrm{d}\sigma_{m,1}(x) \leq \int \left|\log\operatorname{dist}(w,x)\right| \mathrm{d}\sigma_{m,1}(x) + \left|\log(c_Q m^{-5})\right|,$$

which is $O(\log m)$ by the singular integration formula

$$\int \log |w - z| \, \mathrm{dLeb}_{\partial \mathbb{D}(0, c_Q m^{-5})}(z) = \log(c_Q m^{-5}) \quad \text{for} \quad z \in \partial \mathbb{D}(0, c_Q m^{-5}),$$

where $\operatorname{Leb}_{\partial \mathbb{D}(0,c_Qm^{-5})}$ is the Lebesgue probability measure on $\partial \mathbb{D}(0,c_Qm^{-5})$. The proof of the lemma is finished.

Lemma 5.4. There exists a constant $C_D > 0$ independent of m such that

$$\int U_{\nu_D}' \,\mathrm{d}\nu_D - \int U_{\nu_D}' \,\mathrm{d}\delta_{\mathbf{f}_m} \ge -C_D \frac{\log m}{m}$$

for all sufficiently large $m \gg 1$.

Proof. By integrating (4.5) against $\nu_D = \nu_Q$, and by using (2.7), we obtain

$$\int U'_{\nu_D} \,\mathrm{d}\delta_{\mathbf{f}'_m} - U'_{\delta_{\mathbf{f}'_m}}(f_{m,1}) \le \int U'_{\nu_D} \,\mathrm{d}\nu_D - U'_{\nu_D}(f_{m,1})$$

Hence

(5.2)
$$\int U'_{\nu_D} \, \mathrm{d}\nu_D - \int U'_{\nu_D} \, \mathrm{d}\delta_{\mathbf{f}_m} \ge U'_{\nu_D}(f_{m,1}) - U'_{\delta_{\mathbf{f}'_m}}(f_{m,1}) + \varepsilon_{m,D},$$

where

$$\varepsilon_{m,D} := \int U'_{\nu_D} \,\mathrm{d}\delta_{\mathbf{f}'_m} - \int U'_{\nu_D} \,\mathrm{d}\delta_{\mathbf{f}_m} = \frac{1}{m(m-1)} \cdot \sum_{j=2}^m U'_{\nu_D}(f_{m,j}) - \frac{1}{m} \cdot U'_{\nu_D}(f_{m,1})$$

clearly satisfies that $|\varepsilon_{m,D}| \leq 2 \cdot m^{-1} \cdot \max |U'_{\nu_D}| = O(m^{-1})$. It remains to show that $U'_{\nu_D}(f_{m,1}) - U'_{\delta_{\mathbf{f}'_m}}(f_{m,1}) \geq -C_D \log m/m$.

Let $w \in X$ be a point near $f_{m,1}$ with $\operatorname{dist}(w, f_{m,1}) = c_Q m^{-5}$. Using (2.6) and Lemma 2.2, we can bound $\left|U'_{\delta_{\mathbf{f}'_m}}(f_{m,1}) - U'_{\delta_{\mathbf{f}_m}}(w)\right|$ from above by the sum of

(5.3)
$$\frac{1}{m} \left| \log \operatorname{dist}(w, f_{m,1}) \right| + \sum_{j=2}^{m} \left| \frac{1}{m} \log \operatorname{dist}(w, f_{m,j}) - \frac{1}{m-1} \log \operatorname{dist}(f_{m,1}, f_{m,j}) \right|$$

and

(5.4)
$$\frac{1}{m} |\varrho(w, f_{m,1})| + \sum_{j=2}^{m} \left| \frac{1}{m} \varrho(w, f_{m,j}) - \frac{1}{m-1} \varrho(f_{m,1}, f_{m,j}) \right|$$

Since ρ is Lipschitz, (5.4) is $O(m^{-1})$. By the monotonicity of the \log function,

$$\left|\log \operatorname{dist}(f_{m,1}, f_{m,j}) - \log \operatorname{dist}(w, f_{m,j})\right| \leq$$
(5.5)
$$\max\left(\log \frac{\operatorname{dist}(f_{m,1}, f_{m,j})}{\operatorname{dist}(f_{m,1}, f_{m,j}) - c_Q m^{-5}}, \log \frac{\operatorname{dist}(w, f_{m,j}) + c_Q m^{-5}}{\operatorname{dist}(w, f_{m,j})}\right) = \frac{O(1)}{m^2},$$

because ${\rm dist}(f_{m,1},f_{m,j}) \geq c_Q m^{-3}$ by Proposition 4.5. Hence (5.3) is $O(\log m/m).$ Summarizing, we have

(5.6)
$$\left| U'_{\delta_{\mathbf{f}'_m}}(f_{m,1}) - U'_{\delta_{\mathbf{f}_m}}(w) \right| = O(\log m/m)$$

On the other hand, the 1/3 -Hölder continuity of U_{ν_D}' yields

$$\left|U_{\nu_D}'(f_{m,1}) - U_{\nu_D}'(w)\right| = O(m^{-5/3}) < 1/m$$

for large $m \gg 1$. Thus we conclude that

$$U'_{\nu_D}(f_{m,1}) - U'_{\delta_{\mathbf{f}'_m}}(f_{m,1}) \ge U'_{\nu_D}(w) - U'_{\delta_{\mathbf{f}_m}}(w) - O(\log m/m)$$

by the above two estimates. Hence (5.2) and Lemma 5.3 implies that

(5.7)
$$\int U'_{\nu_D} \, \mathrm{d}\nu_D - \int U'_{\nu_D} \, \mathrm{d}\delta_{\mathbf{f}_m} \ge U'_{\nu_D}(w) - U'_{\sigma_m}(w) - O(\log m/m).$$

By repeating the same argument, or by symmetry, it is clear that (5.7) holds for any $w \in \partial \mathbb{B}(f_{m,j}, c_Q m^{-5}), j = 1, ..., m$.

Now take the probability measure η in Lemma 5.2 for $E := \bigcup_{j=1}^{m} \partial \mathbb{B}(f_{m,j}, c_Q m^{-5})$. Integrating (5.7) against η , we obtain

$$\int U'_{\nu_D} \,\mathrm{d}\nu_D - \int U'_{\nu_D} \,\mathrm{d}\delta_{\mathbf{f}_m} \ge \int U'_{\nu_D} \,\mathrm{d}\eta - \int U'_{\sigma_m} \,\mathrm{d}\eta - O(\log m/m)$$

[use (2.7)]
$$= \int U'_{\eta} \,\mathrm{d}\nu_D - \int U'_{\eta} \,\mathrm{d}\sigma_m - O(\log m/m)$$

By Lemma 5.2, U'_{η} equals constantly to its minimum on *E*. Hence $\int U'_{\eta} d\nu_D \geq \int U'_{\eta} d\sigma_m$, and the proof is done.

Lemma 5.5. There exists a constant $C_D > 0$ such that, for all sufficiently large m, one has

$$\left|\int U_{\nu_D}' \,\mathrm{d}\sigma_m - \int U_{\nu_D}' \,\mathrm{d}\delta_{\mathbf{f}_m}\right| \le \frac{C_D}{m}$$

and

$$\mathcal{E}_m(\mathbf{f}_m) - \int U'_{\sigma_m} \,\mathrm{d}\sigma_m \le C_D \frac{\log m}{m}.$$

Proof. Since U'_{ν_D} is 1/3-Hölder continuous with the constant $A := \|U'_{\nu_D}\|_{\mathscr{C}^{1/3}}$, we have

$$\left| \int U_{\nu_D}' \, \mathrm{d}\sigma_m - \int U_{\nu_D}' \, \mathrm{d}\delta_{\mathbf{f}_m} \right| \le \frac{1}{m} \sum_{j=1}^m \left| \int U_{\nu_D}' \, \mathrm{d}\sigma_{m,j} - U_{\nu_D}'(f_{m,j}) \right| \le A \left(\frac{c_Q}{2m^5}\right)^{1/3}$$

which implies the first inequality.

For the second one, we use Lemma 2.2 to rewrite (see (2.11))

$$\mathcal{E}_m(\mathbf{f}_m) = \frac{1}{m^2} \sum_{j \neq k} \Big(\log \operatorname{dist}(f_{m,j}, f_{m,k}) + \varrho(f_{m,j}, f_{m,k}) \Big),$$

and

$$\int U'_{\sigma_m} \,\mathrm{d}\sigma_m = \frac{1}{m^2} \sum_{j,k} \int \int \left(\log \operatorname{dist}(x,y) + \varrho(x,y) \right) \,\mathrm{d}\sigma_{m,j}(x) \,\mathrm{d}\sigma_{m,k}(y)$$

Denote by $\operatorname{Leb}_{\partial \mathbb{D}(0,m^{-5})}$ the probability Lebesgue measure on $\partial \mathbb{D}(0,m^{-5})$. For every j, we can estimate that

$$\begin{split} \left| \int \int \left(\log \operatorname{dist}(x, y) + \varrho(x, y) \right) \mathrm{d}\sigma_{m, j}(x) \mathrm{d}\sigma_{m, j}(y) \right| \\ &\leq O(1) \int \int \left| \log |x - y| \right| \mathrm{dLeb}_{\partial \mathbb{D}(0, m^{-5})}(x) \mathrm{dLeb}_{\partial \mathbb{D}(0, m^{-5})}(y) + \max \varrho \\ &= O(\log m) + \max \varrho + O(1), \end{split}$$

where the last inequality is based on the fact that the singular integration

$$\int \int \log |x-y| \, \mathrm{dLeb}_{\partial \mathbb{D}(0,m^{-5})}(x) \, \mathrm{dLeb}_{\partial \mathbb{D}(0,m^{-5})}(y) = \log(m^{-5}).$$

Therefore, to gain the desired estimate $\mathcal{E}_m(\mathbf{f}_m) - \int U'_{\sigma_m} d\sigma_m \leq C_D \log m/m$, we only need to show, for each pair $j \neq k$, that

$$\left(\log \operatorname{dist}(f_{m,j}, f_{m,k}) + \varrho(f_{m,j}, f_{m,k})\right) - \int \int \left(\log \operatorname{dist}(x, y) + \varrho(x, y)\right) \mathrm{d}\sigma_{m,j}(x) \mathrm{d}\sigma_{m,k}(y)$$

is $O(\log m/m)$. First, observe that the difference involving ρ is $O(m^{-5})$, because ρ is Lipschitz while $\sigma_{m,j}$, $\sigma_{m,k}$ are supported on the small circles around $f_{m,j}$ and $f_{m,k}$ respectively. For the remaining terms, by the monotonicity of the log function, we can show that

$$\left|\log \operatorname{dist}(f_{m,j}, f_{m,k}) - \int \int \log \operatorname{dist}(x, y) \, \mathrm{d}\sigma_{m,j}(x) \mathrm{d}\sigma_{m,k}(y)\right| = O(m^{-2}),$$

because for any $x \in \partial \mathbb{B}(f_{m,j}, c_Q m^{-5})$, $y \in \partial \mathbb{B}(f_{m,k}, c_Q m^{-5})$, we have

$$\left|\log \operatorname{dist}(f_{m,j}, f_{m,k}) - \log \operatorname{dist}(x, y)\right| = O(m^{-2})$$

by the same reasoning as (5.5).

Summarizing all the estimates above, we conclude the proof.

The following is the main result of this section, giving (5.1).

Proposition 5.6. There exists a constant $C_D > 0$, such that

$$-\mathcal{E}_m(\mathbf{f}_m) \le -\int_X U'_{\nu_D} \,\mathrm{d}\nu_D + C_D \frac{\log m}{m}$$

and

$$\max U_{\delta_{\mathbf{f}_m}}' \le \max U_{\nu_D}' + C_D \frac{\log m}{m}$$

for all sufficiently large m.

Proof. By symmetry, the inequality (4.4) remains true after replacing $(f_{m,1}, \mathbf{f}'_m)$ by $(f_{m,k}, \sum_{j \neq k} f_{m,j})$ for every $k = 1, 2, \dots, m$. Summing up all these *m* inequalities, we obtain that (see (2.6))

$$-\frac{1}{m-1}\sum_{j\neq k}G(f_{m,j}, f_{m,k}) + \sum_{j=1}^{m}Q(f_j) \le -\sum_{j=1}^{m}G(z, f_{m,j}) + mQ(z)$$

for all $z \in X \setminus D$, where $Q = U'_{\nu_D}$. Multiplying both sides by $(m-1)/m^2$, we receive

(5.8)
$$-\mathcal{E}_{m}(\mathbf{f}_{m}) + \frac{m-1}{m} \int U_{\nu_{D}}' \, \mathrm{d}\delta_{\mathbf{f}_{m}} \leq -\frac{m-1}{m} U_{\delta_{\mathbf{f}_{m}}}'(z) + \frac{m-1}{m} U_{\nu_{D}}'(z)$$

Integrating (5.8) against the probability measure η in Lemma 5.2 for $E := X \setminus D$, we get

$$-\mathcal{E}_{m}(\mathbf{f}_{m}) + \frac{m-1}{m} \int U_{\nu_{D}}^{\prime} d\delta_{\mathbf{f}_{m}} \leq -\frac{m-1}{m} \int U_{\delta_{\mathbf{f}_{m}}}^{\prime} d\eta + \int \frac{m-1}{m} U_{\nu_{D}}^{\prime} d\eta$$

[use (2.7)]
$$= -\frac{m-1}{m} \int U_{\eta}^{\prime} d\delta_{\mathbf{f}_{m}} + \frac{m-1}{m} \int U_{\eta}^{\prime} d\nu_{D}.$$

By Lemma 5.2, U'_{η} is constant on $X \setminus D$. Hence $-\int U'_{\eta} d\delta_{\mathbf{f}_m} + \int U'_{\eta} d\nu_D = 0$. Therefore

(5.9)
$$-\mathcal{E}_m(\mathbf{f}_m) \le -\frac{m-1}{m} \int U'_{\nu_D} \,\mathrm{d}\delta_{\mathbf{f}_m}.$$

Let σ_m be the probability measure in Lemma 5.5. By Stokes' formula, we have

$$\int (U'_{\sigma_m} - U'_{\nu_D}) \,\mathrm{d}(\sigma_m - \nu_D) \le 0 \qquad \text{(see (2.7))}.$$

Equivalently,

$$-\int U'_{\sigma_m} \,\mathrm{d}\sigma_m \ge \int U'_{\nu_D} \,\mathrm{d}\nu_D - 2\int U'_{\nu_D} \,\mathrm{d}\sigma_m$$

This combining with Lemma 5.5 gives

(5.10)
$$-\mathcal{E}_m(\mathbf{f}_m) + \int U'_{\nu_D} \,\mathrm{d}\delta_{\mathbf{f}_m} + O\left(\frac{\log m}{m}\right) \ge \int U'_{\nu_D} \,\mathrm{d}\nu_D - \int U'_{\nu_D} \,\mathrm{d}\delta_{\mathbf{f}_m} - O\left(\frac{1}{m}\right).$$

Taking (5.9) into account, we get

(5.11)
$$\int U'_{\nu_D} \,\mathrm{d}\nu_D - \int U'_{\nu_D} \,\mathrm{d}\delta_{\mathbf{f}_m} \leq -\frac{1}{m} \int U'_{\nu_D} \,\mathrm{d}\delta_{\mathbf{f}_m} + C'_D \frac{\log m}{m}.$$

Since U_{ν_D}' is uniformly bounded, we conclude the first assertion.

For the second one, it is enough to show that $U'_{\delta_{\mathbf{f}_m}}(z) \leq U'_{\nu_D}(z) + C_D \log m/m$ for all $z \in X$. Plugging (5.10) in the left-hand-side of (5.8), we obtain that for all $z \in X \setminus D$,

$$-\frac{1}{m}\int U'_{\nu_D} \,\mathrm{d}\delta_{\mathbf{f}_m} + \int U'_{\nu_D} \,\mathrm{d}\nu_D - \int U'_{\nu_D} \,\mathrm{d}\delta_{\mathbf{f}_m} - O\Big(\frac{\log m}{m}\Big) \le \frac{m-1}{m}\Big(U'_{\nu_D}(z) - U'_{\delta_{\mathbf{f}_m}}(z)\Big).$$

By Lemma 5.4 and the boundedness of U'_{ν_D} , we deduce that

$$U'_{\delta_{\mathbf{f}_m}}(z) \le U'_{\nu_D}(z) + C_D \log m/m, \quad \forall z \in X \setminus D.$$

Noting that $U'_{\nu_D} - U'_{\delta_{\mathbf{f}_m}}$ is harmonic on D, by the maximal principle, the above inequality holds for any $z \in D$ as well. Thus we conclude the proof.

6. CONSTRUCTING HOLOMORPHIC SECTIONS BY FEKETE POINTS

We have obtained a set of Fekete points $\mathscr{F}_m := \{f_{m,1}, \ldots, f_{m,m}\} \subset X \setminus D$. When $X = \mathbb{P}^1$, \mathscr{F}_m can be realized directly as the zero set of a degree *m* homogeneous polynomial in the hole event. However, in the higher genus case, there might not exist q_1, \ldots, q_g in $X \setminus D$ such that (2.5) holds for $p_j = f_{m,j}$, due to the restriction of the Abel-Jacobi equation (2.5). In other words, this *m*-th Fekete set may not be the zero set of one section in $H_{n,D}$. For this concern, we will construct a valid section s_n in the hole event $H_{n,D}$ whose zero set Z_{s_n} deviates from \mathscr{F}_m only mildly. The following lemma is instrumental in our approach.

Lemma 6.1. Let $\mathbb{B}(y, R) \subset X$ be the disc centered at $y \in X$ having radius R > 0 with respect to ω_0 . Then there exist small constants $0 < \zeta, \beta_1, \beta_2, \beta_3 \ll 1$ and a large integer $k \gg 1$, such that, for any $\mathbf{Z} \in \text{Jac}(X)$, for any $m \ge 1$ points $p_1, \ldots, p_m \in X$, one can find $\mathbf{P}^{[j]} = P_1^{[j]} + \cdots + P_g^{[j]} \in X^{(g)}$ $(1 \le j \le k)$ and $\mathbf{Q} = Q_1 + \cdots + Q_g \in X^{(g)}$ satisfying the following conditions

$$P_{\ell}^{[j]}, Q_{\ell} \in \mathbb{H}_{y, p_1, \dots, p_m}^{R, \zeta} := \mathbb{B}(y, R) \setminus \bigcup_{j=1}^m \mathbb{B}(p_j, \zeta/\sqrt{m}) \quad \text{for all } \ell, j,$$

(6.1)
$$\sum_{j=1}^{k} \mathcal{A}_{g}(\mathbf{P}^{[j]}) + \mathcal{A}_{g}(\mathbf{Q}) = \mathbf{Z},$$

(6.2)
$$\operatorname{dist}\left(\mathbf{Q}, \pi_g(\mathscr{R}_g) \cup \mathscr{W}\right) \geq \beta_1,$$

(6.3)
$$\operatorname{dist}(P_{\ell_1}^{[j_1]}, P_{\ell_2}^{[j_2]}) \ge \beta_2 \quad \text{for all } (j_1, \ell_1) \neq (j_2, \ell_2),$$

(6.4)
$$\operatorname{dist}(P_{\ell_1}^{[j]}, Q_{\ell_2}) \ge \beta_3 \quad \text{for all } j, \ell_1, \ell_2.$$

m

The proof is quite involved, and will be separated into 7 Steps with respect to the thinking process. For clarifying the independence of some uniform positive constants on \mathbb{Z} , m, p_1, \ldots, p_m , we will write $\mathsf{K}_1, \mathsf{K}_2, \mathsf{K}_3, \ldots$ instead of merely O(1).

Proof. Step 1. By demanding $\zeta \ll 1$ to be sufficiently small, we can ensure that the union of any m closed discs $\bigcup_{j=1}^{m} \overline{\mathbb{B}}(p_j, \zeta/\sqrt{m})$ does not cover the open annulus $\mathbb{B}(y, R) \setminus \overline{\mathbb{B}}(y, 2R/3)$ (we can assume that it is nonempty by shrinking R if necessary) by area comparison

(6.5)
$$\sum_{j=1} \operatorname{Area}_{\omega_0} \left(\overline{\mathbb{B}}(p_j, \zeta/\sqrt{m}) \right) \leq m \cdot \mathsf{K}_1 \cdot (\zeta/\sqrt{m})^2 = \mathsf{K}_1 \cdot \zeta^2 < \operatorname{Area}_{\omega_0} \left(\mathbb{B}(y, R) \setminus \overline{\mathbb{B}}(y, 2R/3) \right).$$

The existence of a uniform constant $K_1 > 0$ is guaranteed by the compactness of X. Thus we can choose g points Q_ℓ in the nonempty open set

$$\left(\mathbb{B}(y,R)\setminus\overline{\mathbb{B}}(y,2R/3)\right)\setminus \cup_{j=1}^{m}\overline{\mathbb{B}}(p_{j},\zeta/\sqrt{m})$$

for $\ell = 1, ..., g$, such that $\mathbf{Q} := Q_1 + \cdots + Q_g \notin \pi_g(\mathscr{R}_g) \cup \mathscr{W}$, since $\pi_g(\mathscr{R}_g) \cup \mathscr{W}$ is a proper subvariety of $X^{(g)}$. However, at the moment it is still not clear why we can choose $\beta_1 > 0$ independent of $\mathbf{Z}, m, p_1, ..., p_m$ such that the requirement (6.2) holds.

Step 2. To this aim, we improve the above reasoning as follows. First, we pick g points Q'_1, \ldots, Q'_q in $\mathbb{B}(y, R) \setminus \overline{\mathbb{B}}(y, 2R/3)$ such that $\mathbf{Q}' := Q'_1 + \cdots + Q'_q \notin \pi_g(\mathscr{R}_g) \cup \mathscr{W}$. Next,

by continuity argument, we can choose a sufficiently small positive constant $K_2 \ll 1$ such that $\overline{\mathbb{B}}(Q'_{\ell}, \mathsf{K}_2) \subset \mathbb{B}(y, R) \setminus \overline{\mathbb{B}}(y, 2R/3)$ for $\ell = 1, \ldots, g$ and such that $Q_1 + \cdots + Q_g \notin \pi_g(\mathscr{R}_g) \cup \mathscr{W}$ whenever Q_1, \ldots, Q_g are mild perturbations of Q'_1, \ldots, Q'_g respectively within distance K_2 , *i.e.*, dist $(Q_{\ell}, Q'_{\ell}) \leq \mathsf{K}_2$ for $\ell = 1, \ldots, g$. By compactness, for all such Q_1, \ldots, Q_g we have (6.2) for some uniform constant $\beta_1 > 0$. Lastly, we only need to make sure that $\cup_{j=1}^m \overline{\mathbb{B}}(p_j, \zeta/\sqrt{m})$ does not cover any of $\overline{\mathbb{B}}(Q_{\ell}, \mathsf{K}_2)$ for $\ell = 1, \ldots, g$. This can be achieved by modifying (6.5) as

(6.6)
$$\mathsf{K}_1 \cdot \zeta^2 < \operatorname{Area}_{\omega_0} \left(\overline{\mathbb{B}}(Q'_{\ell}, \mathsf{K}_2) \right) \text{ for every } \ell = 1, \dots, g.$$

Thus (6.2) is obtained.

Step 3. We will choose all the remaining points $P_{\bullet}^{[\bullet]}$ in $\mathbb{B}(y, R/2)$ so that (6.4) automatically holds for sufficiently small $\beta_3 \ll 1$. Now the only remaining obstacles are (6.1) and (6.3). Fix a local holomorphic coordinate chart $\varphi : \mathbb{D} \hookrightarrow \mathbb{B}(y, R/2)$ with

$$\|\mathrm{d}\varphi\| > \mathsf{K}_3 > 0$$

with respect to ω_0 and the Euclidean metric on \mathbb{D} . Recall the construction (2.1) of the Jacobian variety defined by the holomorphic 1-forms ϕ_1, \ldots, ϕ_q . Read

$$\varphi^* \phi_1(z) = \omega_1(z) dz, \quad \dots, \quad \varphi^* \phi_g(z) = \omega_g(z) dz$$

in the local chart, where z is the standard coordinate of the unit disc \mathbb{D} . By the common knowledge of Riemann surfaces (e.g. [28, p. 87]), the equation $\det(\omega_{\ell}(z_j))_{1 \leq \ell, j \leq g} = 0$ defines a proper subvariety of \mathbb{D}^g with coordinates (z_1, \ldots, z_g) . Fix a point $(z'_1, \ldots, z'_g) \in \mathbb{D}^g$ outside this subvariety, *i.e.*, $\det(\omega_{\ell}(z'_j))_{1 \leq \ell, j \leq g} \neq 0$. Therefore the fixed matrix $M := (\omega_{\ell}(z'_j))_{1 \leq \ell, j \leq g}$ is invertible, and the g fixed points z'_1, \ldots, z'_g are pairwise distinct.

Step 4. Fix a small positive constant

$$(6.8) \qquad \delta := \min\left\{ |z'_{\ell_1} - z'_{\ell_2}|/3, \, 1 - |z'_{\ell}| : \, 1 \le \ell_1 < \ell_2 \le g, \, \ell = 1, \dots, g \right\} > 0$$

which is obviously independent of **Z**, m, p_1, \ldots, p_m . Define a holomorphic map (see (2.2))

$$F: \mathbb{D}^g_{\delta} \to \operatorname{Jac}(X), \quad (z_1, \dots, z_g) \mapsto \mathcal{A}\big(\varphi(z_1' + z_1)\big) + \dots + \mathcal{A}\big(\varphi(z_g' + z_g)\big)$$

Then the differential dF of F at the origin $(0, \ldots, 0) \in \mathbb{D}^g$ has the Jacobian matrix M. Select another system $(\widetilde{\phi}_1, \ldots, \widetilde{\phi}_g) := (\phi_1, \ldots, \phi_g) \cdot M^{-1}$ of \mathbb{C} -linearly independent holomorphic 1-forms on X. The Jacobian variety $\widetilde{Jac}(X)$ associated with $(\widetilde{\phi}_1, \ldots, \widetilde{\phi}_g)$ is clearly $\widetilde{Jac}(X) := \mathbb{C}^g/\Lambda \cdot M^{-1}$ by (2.1), and the corresponding Abel-Jacobi map $\widetilde{\mathcal{A}} : X \to \widetilde{Jac}(X)$ defined as

$$\widetilde{\mathcal{A}}(x) := \left(\int_{p_{\star}}^{x} \widetilde{\phi}_{1}, \dots, \int_{p_{\star}}^{x} \widetilde{\phi}_{g}\right) \pmod{\Lambda \cdot M^{-1}}, \quad \forall x \in X$$

satisfies that $\widetilde{\mathcal{A}} = \mathcal{A} \cdot M^{-1}$, where in abuse of notation we also denote M^{-1} as the isomorphism from $\operatorname{Jac}(X) = \mathbb{C}^g / \Lambda$ to $\widetilde{\operatorname{Jac}}(X) = \mathbb{C}^g / \Lambda \cdot M^{-1}$. Define another holomorphic map

(6.9)
$$\widetilde{F}: \mathbb{D}^g_{\delta} \to \widetilde{\operatorname{Jac}}(X), \quad (z_1, \dots, z_g) \mapsto \widetilde{\mathcal{A}}\big(\varphi(z_1' + z_1)\big) + \dots + \widetilde{\mathcal{A}}\big(\varphi(z_g' + z_g)\big).$$

It is clear that $\widetilde{F} = F \cdot M^{-1}$. Once we identify both the tangent spaces of \mathbb{D}^g_{δ} and $\widetilde{\text{Jac}}(X)$ at the origins as \mathbb{C}^g , the differential $d\widetilde{F} : \mathbb{C}^g \to \mathbb{C}^g$ of \widetilde{F} at $(0, \ldots, 0) \in \mathbb{D}^g_{\delta}$ is nothing but the identity map.

Step 5. We plan to choose

(6.10)
$$P_{\ell}^{[j]} := \varphi(z_{\ell}' + z_{\ell}^{[j]}) \text{ for some } z_{\ell}^{[j]} \in \mathbb{D}_{\delta} \text{ for all } j, \ell$$

Let $\widetilde{G} := \widetilde{F} - \widetilde{F}(0, ..., 0)$ be the translation of \widetilde{F} so that \widetilde{G} maps the origin of \mathbb{D}^g_{δ} to that of $\widetilde{\operatorname{Jac}}(X)$. First of all, we transform the equation (6.1) equivalently to

(6.11)
$$\sum_{j=1}^{\kappa} G(z_1^{[j]}, \dots, z_g^{[j]}) = \left(\mathbf{Z} - \mathcal{A}_g(\mathbf{Q})\right) \cdot M^{-1} - k\widetilde{F}(0, \dots, 0) =: [\widetilde{\mathbf{Z}}] \in \widetilde{\operatorname{Jac}}(X)$$

for some $\widetilde{\mathbf{Z}} \in \mathbb{C}^g$ in a fundamental domain of $\widetilde{\operatorname{Jac}}(X) = \mathbb{C}^g / \Lambda \cdot M^{-1}$. Note that the Euclidean norm of $\widetilde{\mathbf{Z}}$ is uniformly bounded from above by some finite constant K₄.

Step 6. The upshot is that the differential $d\tilde{G}: \mathbb{C}^g \to \mathbb{C}^g$ at the origin of \mathbb{D}^g_{δ} is the identity map. By the inverse function theorem, we can define the inverse \tilde{G}^{-1} of \tilde{G} on a small neighborhood U of the origin of $\tilde{\text{Jac}}(X)$. The key idea is to show, by means of volume comparison, that for some large integer k and for sufficiently small $0 < \zeta \ll 1$, we can choose certain k points $\tilde{\mathbf{P}}^{[j]} \in \mathbb{C}^g$ near the origin for $j = 1, \ldots, k$, such that $\sum_{j=1}^k \tilde{\mathbf{P}}^{[j]} = 0$, and such that

(6.12)
$$(z_1^{[j]}, \dots, z_g^{[j]}) := \widetilde{G}^{-1} ([k^{-1} \cdot \widetilde{\mathbf{Z}} + \widetilde{\mathbf{P}}^{[j]}])$$

meets the requirement (6.3).

Now we fulfill some technical details. For convenience of notation, we regard U as an open neighborhood of the origin of \mathbb{C}^g . It is clear that the differential $\mathrm{d}\widetilde{G}^{-1}: \mathbb{C}^g \to \mathbb{C}^g$ at the origin is the identity map. Hence we can fix a sufficiently small constant $\epsilon > 0$, such that for every w in the polydisc $\mathbb{D}^g_{\epsilon} \subset U$, we have

(6.13)
$$\widetilde{G}^{-1}(\mathbf{w}) = \mathbf{w} + O(\|\mathbf{w}\|^2),$$

where the error term $O(||\mathbf{w}||^2)$ has Euclidean norm $\leq K_5 \cdot ||\mathbf{w}||^2$.

Denote the finite square lattice

$$\mathcal{W}_{t,\rho} := \left\{ a\rho + b\rho \cdot i \in \mathbb{C} : a, b \in \mathbb{Z} \cap [1, t] \right\}$$

for some $t \in \mathbb{Z}_+, \rho > 0$ to be determined. Let $\mathfrak{I} : \mathbb{C} \to \mathbb{C}^g$ be the diagonal map, sending z to (z, z, \ldots, z) . We will take $k = 2t^2$, and take the aforementioned points $\widetilde{\mathbf{P}}^{[j]} \in \mathbb{C}^g$ for $j = 1, \ldots, t^2$ as mild perturbations of the t^2 points in $\mathfrak{I}(\mathcal{W}_{t,\rho})$, and for $j = 1 + t^2, \ldots, 2t^2$ as mild perturbations of the t^2 points in $-\mathfrak{I}(\mathcal{W}_{t,\rho})$. Here, perturbations mean that the differences are within $\mathbb{D}^g_{\rho/3}$. In practice, we will take

(6.14)
$$\widetilde{\mathbf{P}}^{[j+t^2]} = -\widetilde{\mathbf{P}}^{[j]} \quad \text{for} \quad j = 1, \dots, t^2.$$

First of all, we need to ensure that all possible $\{k^{-1} \cdot \widetilde{\mathbf{Z}} + \widetilde{\mathbf{P}}^{[j]}\}_{j=1}^k$ chosen in this way are contained in \mathbb{D}^g_{ϵ} . For this goal, we require that

(6.15)
$$||k^{-1} \cdot \widetilde{\mathbf{Z}}|| = ||2^{-1} \cdot t^{-2} \cdot \widetilde{\mathbf{Z}}|| \le \epsilon/100 \text{ and } t \cdot \rho \le \epsilon/2.$$

Next, we verify (6.14) for each j. The only risk is that one of g coordinates of $\widetilde{G}^{-1}(k^{-1} \cdot \widetilde{\mathbf{Z}} + \widetilde{\mathbf{P}}^{[j]})$ or of $\widetilde{G}^{-1}(k^{-1} \cdot \widetilde{\mathbf{Z}} + \widetilde{\mathbf{P}}^{[j+t^2]}) = \widetilde{G}^{-1}(k^{-1} \cdot \widetilde{\mathbf{Z}} - \widetilde{\mathbf{P}}^{[j]})$ lies in $\varphi^{-1}(\bigcup_{j=1}^m \mathbb{B}(p_j, \zeta/\sqrt{m})) =: \mathscr{B}$. Note that such "bad" choices of $\widetilde{\mathbf{P}}^{[j]}$ are contained either in

$$\bigcup_{j=0}^{g} \left(\widetilde{G}(\mathbb{D}^{j}_{\delta} \times \mathscr{B} \times \mathbb{D}^{g-1-j}_{\delta}) - k^{-1} \cdot \widetilde{\mathbf{Z}} \right) \quad \text{or in} \quad \bigcup_{j=0}^{g} \left(k^{-1} \cdot \widetilde{\mathbf{Z}} - \widetilde{G}(\mathbb{D}^{j}_{\delta} \times \mathscr{B} \times \mathbb{D}^{g-1-j}_{\delta}) \right),$$

whose total Euclidean volume is $\leq K_6 \cdot \zeta^2$ (see (6.7)). Since all possible choices of $\widetilde{\mathbf{P}}^{[j]}$ has the Euclidean volume equal to that of $\mathbb{D}^g_{\rho/3}$, as long as

(6.16)
$$\mathsf{K}_6 \cdot \zeta^2 < \operatorname{Vol}(\mathbb{D}^g_{\rho/3}) = \mathsf{K}_7 \cdot \rho^{2g},$$

we can take $\widetilde{\mathbf{P}}^{[j]}$ not "bad". Thus (6.1) is satisfied because of (6.11), (6.12) and (6.14).

Step 7. In the end, we check the requirement (6.3). By the definition (6.12), for any $\ell = 1, \ldots, g$, for any two distinct $j_1, j_2 \in \{1, \ldots, k\}$, using (6.13) we can obtain that

(6.17)
$$|z_{\ell}^{[j_1]} - z_{\ell}^{[j_2]}| \ge (\rho - \rho/3 - \rho/3) - 2 \cdot \mathsf{K}_5 \cdot g\epsilon^2 = \rho/3 - 2g\mathsf{K}_5 \cdot \epsilon^2.$$

Here $g\epsilon^2$ stands for an upper bound for both $||k^{-1} \cdot \widetilde{\mathbf{Z}} + \widetilde{\mathbf{P}}^{[j_1]}||^2$ and $||k^{-1} \cdot \widetilde{\mathbf{Z}} + \widetilde{\mathbf{P}}^{[j_2]}||^2$. Note that under the restriction (6.15) we can still make

$$\rho/3 - 2g\mathsf{K}_5 \cdot \epsilon^2 \ge \rho/6.$$

Indeed, (6.15) is a consequence of

(6.19)
$$2^{-1} \cdot t^{-2} \cdot \mathsf{K}_4 \le \epsilon/100, \quad t \cdot \rho \le \epsilon/2$$

By shrinking $\epsilon > 0$ if necessary, we can check that by selecting

$$t := \left[\sqrt{50\mathsf{K}_4/\epsilon}\right] + 1, \qquad \rho := 12g\mathsf{K}_5\epsilon^2 > 0,$$

both the requirements (6.18) and (6.19) are satisfied. Thus (6.17) implies that

 $|z_{\ell}^{[j_1]} - z_{\ell}^{[j_2]}| \geq \rho/6 > 0, \quad \text{hence} \quad \mathrm{dist}(P_{\ell}^{[j_1]}, P_{\ell}^{[j_2]}) \geq \mathsf{K}_8 \cdot \rho/6 > 0.$

On the other hand, for any $1 \le j_1, j_2 \le k$, for any two distinct $\ell_1, \ell_2 \in \{1, \ldots, g\}$, by the construction (6.8) and (6.10), we have

$$\left| (z'_{\ell_1} + z^{[j_1]}_{\ell_1}) - (z'_{\ell_2} + z^{[j_2]}_{\ell_2}) \right| \ge \delta$$
, hence $\operatorname{dist}(P^{[j_1]}_{\ell_1}, P^{[j_2]}_{\ell_2}) \ge \mathsf{K}_8 \cdot \delta > 0$.

Summarizing, we can take $\beta_2 := \min\{\mathsf{K}_8 \cdot \rho/6, \mathsf{K}_8 \cdot \delta\}$, and demand $\zeta > 0$ to obey (6.6) and (6.16), so that all the requirements are satisfied.

Using Lemma 6.1, we can construct a valid section in $H_{n,D}$, whose zeros are mostly coming from the *m*-th Fekete set.

Proposition 6.2. For every *n* large enough, one can find a holomorphic section $s_n \in H_{n,D}$ with zero set $Z_{s_n} = \{p_1, \ldots, p_m, q_1, \ldots, q_g\}$, such that

$$(6.20) p_j = f_{m,j} for all j \le m - k_D g_j$$

(6.21)
$$\operatorname{dist}(p_{j_1}, p_{j_2}) \ge c_D m^{-3} \text{ for all } j_1 \neq j_2,$$

(6.22)
$$\operatorname{dist}(p_j, q_l) \ge c_D / \sqrt{m} \quad \text{for all } j, l,$$

(6.23)
$$\operatorname{dist}(q_l, D) \ge c_D \quad \text{for all } l,$$

(6.24)
$$\operatorname{dist}\left(q_1 + \dots + q_g, \pi_g(\mathscr{R}_g) \cup \mathscr{W}\right) \ge c_D,$$

where $k_D \in \mathbb{Z}_+, c_D > 0$ are independent of n.

Proof. We apply Lemma 6.1 upon an open disc $\mathbb{B}(y, R) \Subset X \setminus \overline{D}$ to get a $k_D := k \in \mathbb{N}^*$. Now we start with the data

$$p_j := f_{m,j}$$
 for $j \le m - kg$, $\mathbf{Z} := \mathcal{A}_n(\mathcal{L}^n) - \sum_{j=1}^{m-\kappa g} \mathcal{A}(f_{m,j})$,

to obtain the remaining kg + g points

$$p_{m-kg+1} + \dots + p_m := \mathbf{P}^{[1]} + \dots + \mathbf{P}^{[k]}$$
 and $q_1 + \dots + q_g := \mathbf{Q}$.

These *n* points are the zeros of some holomorphic section s_n because of equation (6.1). The desired distance inequalities come from the definition of $\mathbb{H}_{y,p_1,\ldots,p_m}^{R,\zeta}$ and (6.2),(6.3), (6.4).

The zeros of s_n also gives an element in $S_{m,D}$ (see (2.13)) for all m large enough. The last inequality of the corollary implies that $p_1 + \cdots + p_m \notin \mathbf{H}_m$, *i.e.*, $q_1 + \cdots + q_q$ is unique.

7. DIFFERENCE BETWEEN THE TWO FUNCTIONALS

In this section, our goal is to approximate the minimum of the functional \mathcal{I}_D closely by $-\mathcal{E}_m(\mathbf{p}) + 2 \max U'_{\delta_{\mathbf{p}}}$, and show that such $\mathbf{p} \in X^{(m)}$ can constitute "large enough" volume.

Proposition 7.1. There exist constants $m_0 \in \mathbb{Z}_+, C_D > 0$, such that for every large integer $m \geq m_0$, one can find an open subset $\Omega_{m,D} \subset S_{m,D} \subset X^{(m)}$ (see (2.13)) with volume $\operatorname{Vol}(\Omega_{m,D}) \geq m^{-11m}$ and that

(7.1)
$$-\mathcal{E}_m(\mathbf{p}) + \frac{2(m+1)}{m} \mathcal{F}_m(\mathbf{p}) \le \min \mathcal{I}_D + C_D \frac{\log m}{m}, \qquad \forall \mathbf{p} \in \Omega_{m,D}.$$

First, we need the following two lemmas, which guarantees the stability of the functional values $-\mathcal{E}_m(\mathbf{p}) + 2 \max U'_{\delta_{\mathbf{p}}}$ under mild perturbations of $\mathbf{p} := p_1 + \cdots + p_m \in \mathcal{S}_{m,D}$.

Lemma 7.2. For every sufficiently large m, for any $\mathbf{p}' := p'_1 + \cdots + p'_m$ with $p'_j \in \mathbb{B}(p_j, m^{-5})$, one has

$$\left|\mathcal{E}_{m}(\mathbf{p})-\mathcal{E}_{m}(\mathbf{p}')\right|=O(m^{-1}) \quad and \quad \left|\max U'_{\delta_{\mathbf{p}}}-\max U'_{\delta_{\mathbf{p}'}}\right|=O(m^{-1}).$$

Proof. By (2.11) and Lemma 2.2, $|\mathcal{E}_m(\mathbf{p}) - \mathcal{E}_m(\mathbf{p}')|$ is bounded from above by

$$\frac{1}{m^2} \sum_{j \neq \ell} \left| \log \operatorname{dist}(p_j, p_\ell) - \log \operatorname{dist}(p'_j, p'_\ell) \right| + \frac{1}{m^2} \sum_{j \neq \ell} \left| \varrho(p_j, p_\ell) - \varrho(p'_j, p'_\ell) \right|.$$

Since ρ is Lipschitz, the second term is $O(m^{-5})$. For the first one, using the monotonicity of the log function, every term $|\cdots|$ is less than

(7.2)
$$\max\left(\log\frac{\operatorname{dist}(p_j, p_\ell)}{\operatorname{dist}(p_j, p_\ell) - 2m^{-5}}, \log\frac{\operatorname{dist}(p_j, p_\ell) + 2m^{-5}}{\operatorname{dist}(p_j, p_\ell)}\right) = O(m^{-1}),$$

because $dist(p_j, p_\ell) \ge m^{-4}$ for $j \ne \ell$ by (2.13). Hence their summation (m(m-1) terms), divided by m^2 , is also $O(m^{-1})$. Thus the first assertion is proved.

For the second assertion, we only need to show the following three estimates:

(i)
$$|U'_{\delta_{\mathbf{D}}}(w) - U'_{\delta_{\mathbf{D}'}}(w)| = O(m^{-1})$$
, if dist $(w, p_j) \ge m^{-6}$ and dist $(w, p'_j) \ge m^{-6}$;

(ii) $\max_{X \setminus \mathbb{B}(p_j, m^{-6})} U'_{\delta_{\mathbf{p}}} \ge \sup_{\mathbb{B}(p_j, m^{-6})} U'_{\delta_{\mathbf{p}}} - O(m^{-1});$

- (iii) $\max_{X \setminus \mathbb{B}(p'_j, m^{-6})} U'_{\delta_{\mathbf{p}'}} \ge \sup_{\mathbb{B}(p'_j, m^{-6})} U'_{\delta_{\mathbf{p}'}} O(m^{-1}).$
- By (2.6) and Lemma 2.2, the left-hand-side of (i) is bounded from above by

$$\frac{1}{m}\sum_{j=1}^{m} \left|\log\operatorname{dist}(w, p_j) - \log\operatorname{dist}(w, p'_j)\right| + \frac{1}{m}\sum_{j=1}^{m} \left|\varrho(w, p_j) - \varrho(w, p'_j)\right|.$$

The remaining argument goes the same as the first assertion above.

Now we prove (ii) by showing that

$$\max_{\partial \mathbb{B}(p_j,m^{-6})} U'_{\delta_{\mathbf{p}}} \ge \sup_{\mathbb{B}(p_j,m^{-6})} U'_{\delta_{\mathbf{p}}} - O(\log m/m).$$

Take any $w \in \partial \mathbb{B}(p_j, m^{-6}), y \in \mathbb{B}(p_j, m^{-6})$ for e.g. j = 1. By (2.6), we have

$$U'_{\delta_{\mathbf{p}}}(w) - U'_{\delta_{\mathbf{p}}}(y) = \frac{1}{m} \big(G(w, p_1) - G(y, p_1) \big) + \frac{1}{m} \sum_{\ell=2}^{m} \big(G(w, p_\ell) - G(y, p_\ell) \big)$$

Using Lemma 2.2 again, we can estimate

$$\frac{1}{m} (G(w, p_1) - G(y, p_1)) = \frac{1}{m} (\log \operatorname{dist}(w, p_1) - \log \operatorname{dist}(y, p_1)) + \frac{1}{m} (\varrho(w, p_1) - \varrho(y, p_1)) \\ \ge 0 - O(1) \operatorname{dist}(w, y)/m \ge -O(m^{-7}).$$

By similar computations, and using the fact that

$$\left|\log \operatorname{dist}(w, p_{\ell}) - \log \operatorname{dist}(y, p_{\ell})\right| \le \log \frac{\operatorname{dist}(p_1, p_{\ell}) + m^{-6}}{\operatorname{dist}(p_1, p_{\ell}) - m^{-6}} = O(m^{-2}), \quad \forall \ell = 2, \dots, m,$$

we can show that

$$\sum_{\ell=2}^{m} \left| G(w, p_{\ell}) - G(y, p_{\ell}) \right| = O(m^{-1})$$

Hence we conclude (ii). The proof of (iii) is likewise.

Lemma 7.3. For every sufficiently large integer $m \gg 1$, for any $\mathbf{p}' := p'_1 + \cdots + p'_m \in X^{(m)}$ with $\operatorname{dist}(p'_{j_1}, p'_{j_2}) \ge c_D m^{-3}$ for $j_1 \neq j_2$, and with $p'_j = p_j$ for $1 \le j \le m - k_D g$, one has

$$\left|\mathcal{E}_{m}(\mathbf{p}) - \mathcal{E}_{m}(\mathbf{p}')\right| = O(\log m/m) \quad and \quad \left|\max U'_{\delta_{\mathbf{p}}} - \max U'_{\delta_{\mathbf{p}'}}\right| = O(\log m/m)$$

Proof. Since p and p' only differ in the choice of $k_D g$ points, by (2.11) and Lemma 2.2, we can bound $|\mathcal{E}_m(\mathbf{p}) - \mathcal{E}_m(\mathbf{p}')|$ from above by

(7.3)
$$\frac{1}{m^2} \sum_{j=1}^m \sum_{\substack{\ell=m-k_Dg+1,\\\ell\neq j}}^m \left(\left| \log \operatorname{dist}(p_j, p_\ell) - \log \operatorname{dist}(p'_j, p'_\ell) \right| + \left| \varrho(p_j, p_\ell) - \varrho(p'_j, p'_\ell) \right| \right).$$

By the assumptions, each $\log \operatorname{dist}(\cdot, \cdot)$ is $O(\log m)$. Using the boundedness of ρ , it is clear that (7.3) is $O(k_D \log m/m)$. This proves the first assertion.

For the second one, using (ii) and (iii) in the preceding proof, we only need to show that, if $w \in X$ satisfies $dist(w, p_j), dist(w, p'_j) \ge m^{-6}$ for all j = 1, ..., m, then

$$\left|U_{\delta_{\mathbf{p}}}'(w) - U_{\delta_{\mathbf{p}'}}'(w)\right| = O(\log m/m).$$

Applying (2.6) and Lemma 2.2 again, we can bound $|U'_{\delta_{\mathbf{D}}}(w) - U'_{\delta_{\mathbf{L}'}}(w)|$ from above by

$$\frac{1}{m} \sum_{j=m-k_Dg+1}^{m} \left| \log \operatorname{dist}(w, p_j) - \log \operatorname{dist}(w, p'_j) \right| + \frac{1}{m} \sum_{j=m-k_Dg+1}^{m} \left| \varrho(w, p_j) - \varrho(w, p'_j) \right|.$$

The remaining estimate goes the same as (7.3).

Now we are ready to prove Proposition 7.1. The idea is to fix a set $\{f_{m,1}, \ldots, f_{m,m}\}$ of Fekete points for every large m, and then to apply Proposition 6.2 to show that certain mild perturbations $\mathbf{p} \in X^{(m)}$ of $f_{m,1} + \cdots + f_{m,m}$, which correspond to zero sets of some sections in $H_{n,D}$, constitute "large" volume, and that their functional values $-\mathcal{E}_m(\mathbf{p}) + 2 \max U'_{\delta_{\mathbf{p}}}$ are close to $\min \mathcal{I}_D$.

Proof of Proposition 7.1. Take an *m*-th Fekete set $\{f_{m,1}, \ldots, f_{m,m}\}$ for every $m \ge 1$. Applying Proposition 6.2 with the same notation, we obtain a holomorphic section $s_n \in H_{n,D}$ whose zero set reads in two separated parts as $\mathbf{p} := p_1 + \cdots + p_m$ and $\mathbf{q} := q_1 + \cdots + q_g$. Define

$$\Omega_{m,D} := \bigotimes_{j=1}^{m} \left(\mathbb{B}(p_j, m^{-5}) \setminus D \right) \subset X^{(m)}.$$

Since *D* has smooth boundary and one half of the unit disc has Euclidean area $\pi/2 > 1$, the volume of $\mathbb{B}(p_j, m^{-5}) \setminus D$ is $> m^{-10}$ for sufficiently large *m*, no matter where is p_j in the compact complement $X \setminus D$. Thus the volume of $\Omega_{m,D}$ is $> m^{-11m}$ for $m \gg 1$.

Now we check that $\Omega_{m,D} \subset S_{m,D}$. Take any point $\mathbf{p}' := p'_1 + \cdots + p'_m \in \Omega_{m,D}$, *i.e.*, $p'_j \in \mathbb{B}(p_j, m^{-5})$. Let $\mathbf{q}' = q'_1 \cdots + q'_g$ be a solution to the Abel-Jacobi equation $\mathcal{A}_m(\mathbf{p}') + \mathcal{A}_g(\mathbf{q}') = \mathcal{A}_n(\mathcal{L}^n)$. We need to show that $\operatorname{dist}(p'_{j_1}, p'_{j_2}) \ge m^{-4}$ for all $j_1 \neq j_2$, $\operatorname{dist}(p'_j, q'_l) \ge 1/m$ for all j, l, and $q'_l \in X \setminus D$ for all l. The first inequality follows from (6.21) directly when $m \gg 1$. For the second and third assertions, since \mathcal{A}_1 is holomorphic, for a fixed Euclidean distance in $\operatorname{Jac}(X)$,

dist
$$(\mathcal{A}_m(\mathbf{p}), \mathcal{A}_m(\mathbf{p}')) = \text{dist}\left(\sum_{j=1}^m \mathcal{A}_1(p_j), \sum_{j=1}^m \mathcal{A}_1(p_j')\right)$$

 $\leq \sum_{j=1}^m \text{dist}\left(\mathcal{A}_1(p_j), \mathcal{A}_1(p_j')\right) \leq \sum_{j=1}^m O(1) \text{dist}(p_j, p_j') = O(m^{-4}).$

Consequently, dist $(\mathcal{A}_g(\mathbf{q}), \mathcal{A}_g(\mathbf{q}')) \leq O(m^{-4})$. Recalling that dist $(\mathbf{q}, \pi_g(\mathscr{R}_g) \cup \mathscr{W})$ is bounded below by a constant from (6.24), by the inverse function theorem and the compactness argument, \mathcal{A}_g is an isomorphism in a small ball centered at \mathbf{q} with some uniform radius. For sufficiently large m, $\mathcal{A}_g(\mathbf{q}')$ is in the image of such a ball under \mathcal{A}_g . Thus \mathbf{q}' is unique, and we can read \mathbf{q}' as $q'_1 + \cdots + q'_g$ in a suitable order so that dist $(q_l, q'_l) \leq O(m^{-4})$. Whence dist $(p'_i, q'_l) \geq 1/m$ by (6.22) and $q'_l \in X \setminus D$ by (6.23).

It remains to verify (7.1). Now we apply Lemma 7.3 upon the two points \mathbf{p} and $\mathbf{f}_m := f_{m,1} + \cdots + f_{m,m}$, which clearly meet the criterion by (6.20). Taking also Proposition 5.6 into consideration, we see that

$$-\mathcal{E}_m(\mathbf{p}) + 2 \max U'_{\delta_{\mathbf{p}}} \le \min \mathcal{I}_D + O(\log m/m).$$

This combining with Lemma 7.2 yields

$$-\mathcal{E}_m(\mathbf{p}') + 2\max U'_{\delta_{\mathbf{p}'}} \le \min \mathcal{I}_D + O(\log m/m), \qquad \forall \, \mathbf{p}' \in \Omega_{m,D}$$

Thus we can conclude the proof by Lemma 2.4.

8. DECAY OF THE HOLE PROBABILITIES

In this section, we will prove Theorem 1.1 in the difficult case $g \ge 1$ (see Remark 8.3 below for the easy case g = 0).

Let μ be a probability measure on X. Define

$$\mathcal{E}_{\Delta}(\mu) := \int_{(X \times X) \setminus \Delta} G(x, y) \, \mathrm{d}\mu(x) \mathrm{d}\mu(y).$$

Observe that $\mathcal{E}_m(\mathbf{p}) \leq \mathcal{E}_{\Delta}(\delta_{\mathbf{p}})$ for every $\mathbf{p} = p_1 + \cdots + p_m \in X^{(m)}$. The equality holds if and only if p_1, \ldots, p_m are pairwise distinct.

Lemma 8.1. For any open set D (can be empty) in X, one has

$$\inf_{\mu \in \mathcal{M}(X \setminus D)} \left(-\mathcal{E}_{\Delta}(\mu) + 2 \max U'_{\mu} \right) = \min \mathcal{I}_D = \mathcal{I}_D(\nu_D).$$

Proof. Recall (2.6) that

$$\mathcal{I}_D(\mu) = -\int_{X \times X} G(x, y) \,\mathrm{d}\mu(x) \mathrm{d}\mu(y) + 2 \max U'_{\mu}$$

Noting that $G(x, y) = -\infty$ on the diagonal $\Delta := \{x = y\}$ of $X \times X$, it is clearly that $\nu_D \times \nu_D$ charges no mass on Δ , for otherwise $\mathcal{I}_D(\nu_D) = +\infty!$ Hence we conclude the proof.

Proof of Theorem 1.1. By (2.10) and Proposition 2.3, we can compute

$$\mathbf{P}_{n}(H_{n,D}) = {\binom{n}{m}}^{-1} \int_{\mathcal{R}_{m,D}} \mathscr{A}_{m}^{*}(V_{n}^{\mathrm{FS}})$$
$$= {\binom{n}{m}}^{-1} \int_{\mathbf{p}\in\mathcal{R}_{m,D}} C_{n} \exp\left[m^{2} \left(\mathcal{E}_{m}(\mathbf{p}) - \frac{2(m+1)}{m} \mathcal{F}_{m}(\mathbf{p})\right)\right] \kappa_{n}(\mathbf{p}).$$

Taking logarithm, we receive that

(8.1)
$$\log \mathbf{P}_n(H_{n,D}) \leq \log C_n + m^2 \sup_{\mathbf{p} \in (X \setminus D)^{(m)}} \left(\mathcal{E}_m(\mathbf{p}) - \frac{2(m+1)}{m} \mathcal{F}_m(\mathbf{p}) \right) + \log \int_{X^{(m)}} \kappa_n.$$

By Lemma 2.4, as $m \to \infty$, we have $\mathcal{F}_m(\mathbf{p}) \ge \max U'_{\delta_{\mathbf{p}}} - O(\log m/m)$. Together with the uniformly boundedness of $\max U'_{\delta_{\mathbf{p}}}$ for all \mathbf{p} , we conclude that

$$\sup_{\mathbf{p}\in(X\setminus D)^{(m)}}\left(\mathcal{E}_m(\mathbf{p})-\frac{2(m+1)}{m}\mathcal{F}_m(\mathbf{p})\right)\leq O(\log m/m)+\sup_{\mathbf{p}\in(X\setminus D)^{(m)}}\left(\mathcal{E}_m(\mathbf{p})-2\max U'_{\delta_{\mathbf{p}}}\right).$$

Since $\mathcal{E}_m(\mathbf{p}) \leq \mathcal{E}_{\Delta}(\delta_{\mathbf{p}})$, the last term is bounded from above by

$$\sup_{\mu \in \mathcal{M}(X \setminus D)} \left(\mathcal{E}_{\Delta}(\mu) - 2 \max U'_{\mu} \right) = -\min \mathcal{I}_D$$

due to Lemma 8.1.

Now we estimate the integration of κ_n . Using Proposition 2.3, we get

(8.2)
$$\log \int_{X^{(m)}} \kappa_n \lesssim \log \int_{X^{(m)}} e^{O(m)} \operatorname{diam}(X)^{2mg} i \mathrm{d}z_1 \wedge \mathrm{d}\overline{z}_1 \wedge \cdots \wedge i \mathrm{d}z_m \wedge \mathrm{d}\overline{z}_m \lesssim m \log m.$$

Therefore, we conclude by Lemma 8.2 below that

$$\log \mathbf{P}_n(H_{n,D}) \le -m^2 \min \mathcal{I}_D + Cm \log m$$

Notice that C can be chosen independent of D.

For the lower bound, we apply Proposition 7.1 to get $\Omega_{m,D}$, so that

$$\mathbf{P}_{n}(H_{n,D}) = {\binom{n}{m}}^{-1} \int_{\mathcal{R}_{m,D}} \mathscr{A}_{m}^{*}(V_{n}^{\mathrm{FS}}) \ge {\binom{n}{m}}^{-1} \int_{\Omega_{m,D}} \mathscr{A}_{m}^{*}(V_{n}^{\mathrm{FS}})$$
$$= {\binom{n}{m}}^{-1} \int_{\mathbf{p}\in\Omega_{m,D}} C_{n} \exp\left[m^{2} \left(\mathcal{E}_{m}(\mathbf{p}) - \frac{2(m+1)}{m} \mathcal{F}_{m}(\mathbf{p})\right)\right] \kappa_{n}(\mathbf{p}).$$

It follows from (7.1) that

(8.3)
$$\log \mathbf{P}_n(H_{n,D}) \ge -\log \binom{n}{m} + \log C_n - m^2 \min \mathcal{I}_D - C_D m \log m + \log \int_{\mathbf{p} \in \Omega_{m,D}} \kappa_n.$$

It remains to estimate the integration of κ_n . Since $\Omega_{m,D} \subset S_{m,D}$ (see (2.13)), for any point $\mathbf{p} = p_1 + \cdots + p_m \in S_{m,D}$ and $\mathbf{q} := \mathcal{B}_m(\mathbf{p}) = q_1 + \cdots + q_g$, we have

$$\prod_{j=1}^{m} \prod_{l=1}^{g} \operatorname{dist}(p_j, q_l)^2 \ge \prod_{j=1}^{m} \prod_{l=1}^{g} \frac{1}{m^2} = m^{-2mg}$$

Thus, applying Proposition 2.3 and reminding that $\operatorname{Vol}(\Omega_{m,D}) \geq m^{-11m}$ from Proposition 7.1, the logarithm of the integration of κ_n is no less than

$$O(m) + \log \int_{\mathbf{p} \in \Omega_{m,D}} m^{-2mg} \, i \mathrm{d} z_1 \wedge \mathrm{d} \overline{z}_1 \wedge \dots \wedge i \mathrm{d} z_m \wedge \mathrm{d} \overline{z}_m \ge -O(m \log m).$$

Therefore, we deduce from Lemma 8.2 below that

$$\log \mathbf{P}_n(H_{n,D}) \ge -m^2 \min \mathcal{I}_D - C'_D m \log m.$$

This completes the proof of Theorem 1.1.

Some useful estimate of C_n might be known in literature. For the sake of completeness, we provide a proof here.

Lemma 8.2. There exists a constant c > 0 independent of n and D such that

$$-c n \log n \le \log C_n \le c n \log n.$$

Proof. Note that $\mathbf{P}_n(H_{n,\varnothing}) = 1$ and $\min \mathcal{I}_{\varnothing} = 0$. Repeating the argument in the proof of Theorem 1.1 above for $D = \emptyset$, we can deduce from (8.1) that

$$\log C_n \ge m^2 \min \mathcal{I}_{\varnothing} - c_1 m \log m = -c_1 m \log m$$

for some c_1 independent of n.

On the other hand, using the method of Fekete points upon the case $D = \emptyset$, we receive $\Omega_{m,\emptyset} \subset S_{m,\emptyset}$. Repeating the same reasoning as (8.3), we can obtain that

$$\log \mathbf{P}_n(H_{n,\emptyset}) \ge -\log \binom{n}{m} + \log C_n - m^2 \min \mathcal{I}_{\emptyset} - c_2 m \log m + \log \int_{\mathbf{p} \in \Omega_{m,\emptyset}} \kappa_n$$

for some c_2 independent of n. This combining with (8.2) yields

$$\log C_n \le m^2 \min \mathcal{I}_{\varnothing} + c_3 m \log m = c_3 m \log m$$

Hence we finish the proof.

Remark 8.3. For the proof of the case g = 0, *i.e.*, $X = \mathbb{P}^1$, we can repeat the same strategy, by using a density formula of the zeros, cf. [50, Proposition 3]. The proof is much simpler because there is no q any more and we only need to deal with the functional $\mathcal{E}_m(\mathbf{p}) - \frac{2(m+1)}{m} \mathcal{F}_m(\mathbf{p})$ with m = n, where \mathcal{F}_m is given by $\mathcal{F}_m(\mathbf{p}) := \log \|e^{U'_{\delta_{\mathbf{p}}}}\|_{L^{2m}(\omega_0)}$.

9. Asymptotic behavior of the hole probabilities

To prove Theorem 1.2, it is enough to show the inequality for a fixed x_0 and all small r. The theorem will follow by a standard compactness argument.

Fix a point $x_0 \in X$. For a radius r > 0, we denote $\mathcal{I}_r := \mathcal{I}_{\mathbb{B}(x_0,r)}$ and $\nu_r := \nu_{\mathbb{B}(x_0,r)}$ for abbreviation. This section is devoted to prove the following proposition, which implies Theorem 1.2 by Theorem 1.1.

Proposition 9.1. There exist constants C > c > 0 independent of r such that

 $cr^4 \leq \min \mathcal{I}_r \leq Cr^4$ for every $0 < r \ll 1$.

We decompose the proof into several lemmas below.

Lemma 9.2. For any probability measure μ supported on $X \setminus \mathbb{B}(x_0, r)$, one has $\mathcal{I}_r(\mu) \ge cr^4$ for some c > 0 independent of r and μ .

Proof 1. Take a local coordinate function $\phi : V \to \mathbb{C}$ with $\phi(x_0) = 0$, where V contains $\overline{\mathbb{B}}(x_0, r_0)$ for some small $r_0 > 0$. By compactness, we can find a sufficiently small $\epsilon_0 > 0$ such that both ω_0 and ω are greater than $\epsilon_0^2 \cdot \phi^* \mathrm{dd}^c |z|^2$ on $\overline{\mathbb{B}}(x_0, r_0)$, where z is the coordinate of \mathbb{C} . For any probability measure μ supported on $X \setminus \mathbb{B}(x_0, r)$ with $r < r_0$, the ω -potential U_{μ} of μ satisfies the equation $\mathrm{dd}^c U_{\mu} = -\omega$ on $\mathbb{B}(x_0, r)$. Note that for any $r \in [0, r_0]$, the range $\phi(\overline{\mathbb{B}}(x, r))$ contains $\overline{\mathbb{D}}(0, \epsilon_1 \cdot r) \subset \mathbb{C}$ for some $\epsilon_1 = \mathrm{const} \cdot \epsilon_0$. We can check that $\widehat{U}_{\mu} := U_{\mu} \circ \phi^{-1} : \overline{\mathbb{D}}(0, \epsilon_1 \cdot r_0) \to \mathbb{R}_{\leq 0}$ is strictly supharmonic

$$-\mathrm{dd}^{c}\widehat{U}_{\mu} = -(\phi^{-1})^{*}\mathrm{dd}^{c}U_{\mu} = (\phi^{-1})^{*}\omega \ge (\phi^{-1})^{*}(\epsilon_{0}^{2} \cdot \phi^{*}\mathrm{dd}^{c}|z|^{2}) = \epsilon_{0}^{2} \cdot \mathrm{dd}^{c}|z|^{2}.$$

Using Jensen's formula (cf. e.g. [32, 35]), for any $0 < t \le \epsilon_1 \cdot r_0$, we have

$$\widehat{U}_{\mu}(0) - \frac{1}{2\pi} \int_{|z|=t} \widehat{U}_{\mu} \mathrm{d}\theta = -2 \int_{x=0}^{t} \frac{\mathrm{d}x}{x} \int_{\mathbb{D}_{x}} \mathrm{d}\mathrm{d}^{\mathrm{c}} \widehat{U}_{\mu} \ge 2 \int_{x=0}^{t} \frac{\mathrm{d}x}{x} \int_{\mathbb{D}_{x}} \epsilon_{0}^{2} \cdot \mathrm{d}\mathrm{d}^{\mathrm{c}} |z|^{2}.$$

Discarding the first term $\widehat{U}_{\mu}(0) \leq 0$, we obtain that

$$-\frac{1}{2\pi} \int_{|z|=t} \widehat{U}_{\mu} \mathrm{d}\theta \ge 2 \int_{x=0}^{t} \frac{\mathrm{d}x}{x} \int_{\mathbb{D}_{x}} \epsilon_{0}^{2} \cdot \mathrm{d}\mathrm{d}^{\mathrm{c}} |z|^{2} = \mathrm{const} \cdot \epsilon_{0}^{2} \cdot t^{2}$$

for any $0 < t \le \epsilon_1 \cdot r_0$. Thus we conclude that

$$-\int_{\mathbb{D}_r} \widehat{U}_{\mu} \, \mathrm{dd}^{\mathrm{c}} |z|^2 = -\mathrm{const} \cdot \int_{t=0}^r t \, \mathrm{d}t \int_{|z|=t} \widehat{U}_{\mu} \mathrm{d}\theta \ge -\mathrm{const} \cdot \epsilon_0^2 \cdot r^4$$

for any $0 < r \le \epsilon_1 \cdot r_0$. Since $U_{\mu} \le 0$, we can bound $\mathcal{I}_r(\mu) \ge -\int_{\mathbb{B}(x,r)} U_{\mu} \cdot \omega$ from below by

$$-\int_{\phi(\mathbb{B}(x,r))} U_{\mu} \circ \phi^{-1} \cdot (\phi^{-1})^{*} \omega \ge -\int_{\mathbb{D}(0,\epsilon_{1}\cdot r)} \widehat{U}_{\mu} \cdot \epsilon_{0}^{2} \,\mathrm{dd}^{c} |z|^{2} \ge \operatorname{const} \cdot \epsilon_{0}^{4} \cdot r^{4}$$

for any $r \in (0, r_0)$.

Proof 2. In the definition of \mathcal{I}_r , U_{μ} is non-positive. It suffices to show that $\int_X U_{\mu} \omega \ge cr^4$. The problem is local, we can work on $\mathbb{D} \subset \mathbb{C}$. More precisely, we will show that for any $\omega_{\mathbb{D}}$ -subharmonic function U_r with $U_r \le 0$ on \mathbb{D} and $\mathrm{dd}^c U_r = -\omega_{\mathbb{D}}$ on \mathbb{D}_r , one has $-\int_{\mathbb{D}_r} U_r \omega_{\mathbb{D}} \ge cr^4$, where $\omega_{\mathbb{D}} := f(z) i \mathrm{d} z \wedge \mathrm{d} \overline{z}$ is some fixed Kähler form on \mathbb{D} .

Suppose for contradiction, there exists a sequence (U_j, r_j) with $dd^c U_j = -\omega_{\mathbb{D}}, U_j \leq 0$ and $\lim_{j\to\infty} r_j = 0$, such that

$$\lim_{j \to \infty} r^{-4} \int_{\mathbb{D}(0,r_j)} U_j \,\omega_{\mathbb{D}} = 0.$$

Consider the dilation map $\tau_r : \mathbb{D} \to \mathbb{D}(0, r), z \mapsto rz$. Define

$$\Phi_j := -r_j^{-2} \tau_{r_j}^* U_j.$$

Then on \mathbb{D} , for *j* large, we have $\Phi_j \ge 0$ and

$$\mathrm{dd}^{\mathrm{c}}\Phi_{j} = r_{j}^{-2}\tau_{r_{j}}^{*}(-\mathrm{dd}^{\mathrm{c}}U_{j}) = r^{-2}\tau_{r_{j}}^{*}\omega_{\mathbb{D}} = f(r_{j}z)\,\mathrm{id}z\wedge\mathrm{d}\overline{z}\gtrsim\mathrm{id}z\wedge\mathrm{d}\overline{z}.$$

On the other hand,

$$\int_{\mathbb{D}} \Phi_j \, i \mathrm{d}z \wedge \mathrm{d}\overline{z} \lesssim \int_{\mathbb{D}} \Phi_j \, r^{-2} \tau_{r_j}^* \omega_{\mathbb{D}} = \int_{\mathbb{D}} r^{-2} (\tau_{r_j})_* (\Phi_j) \, \omega_{\mathbb{D}} = -r^{-4} \int_{\mathbb{D}(0,r_j)} U_j \, \omega_{\mathbb{D}}$$

This says the L^1 -norm of Φ_j converges to 0. It is not possible because $dd^c \Phi_j$ does not converge to 0. So we get a contradiction.

To prove the upper bound, we construct a local quasi-subharmonic function first.

Lemma 9.3. Let $\omega_{\mathbb{D}}$ be a Kähler form on \mathbb{D} . There exist an integer $\ell > 2$ and a constant $\alpha > 0$, such that for every small $0 < r \ll 1$, there exists a continuous $\omega_{\mathbb{D}}$ -subharmonic function Φ_r on \mathbb{D} , such that $\Phi_r = 0$ on $\mathbb{D} \setminus \mathbb{D}_{\ell r}$, $-\alpha r^2 \leq \Phi_r \leq 0$ on $\mathbb{D}_{\ell r}$ and $\mathrm{dd}^c \Phi_r = -\omega_{\mathbb{D}}$ on \mathbb{D}_r .

Proof. Take two positive constants $\beta_1, \beta_2 > 0$ such that $\beta_1 \operatorname{dd^c} |z|^2 \le \omega_{\mathbb{D}} \le \beta_2 \operatorname{dd^c} |z|^2$ on $\mathbb{D}_{1/2}$. Let ℓ be a large positive integer to be determined. Define an auxiliary function

$$\Phi_r(z) := -\beta_1 (|z| - \ell r)^2$$

on $\mathbb{D}_{\ell r} \setminus \mathbb{D}_r$, where $r \in (0, 1/(2\ell)]$. Direct computation shows that

$$\mathrm{dd}^{\mathrm{c}}\Phi_r = -\beta_1 \,\mathrm{dd}^{\mathrm{c}}|z|^2 + 2\beta_1 \ell r \,\mathrm{dd}^{\mathrm{c}}|z| \ge -\beta_1 \,\mathrm{dd}^{\mathrm{c}}|z|^2 \ge -\omega_{\mathbb{D}}$$

on $\mathbb{D}_{\ell r} \setminus \mathbb{D}_r \subset \mathbb{D}_{1/2}$. Next, we trivially extend $\Phi_r = 0$ outside $\mathbb{D}_{\ell r}$, and we extend Φ_r inside \mathbb{D}_r as the unique solution to the elliptic equation $\mathrm{dd}^c \Phi_r = -\omega_{\mathbb{D}}$ with the boundary condition $\Phi_r|_{\partial \mathbb{D}_r} = -\beta_1 (\ell - 1)^2 r^2$.

Now we show that $\Phi_r \leq 0$ on \mathbb{D}_r . To do this, we introduce a comparison function Ψ_r which solves the equation $\mathrm{dd}^c \Psi_r = -\beta_2 \mathrm{dd}^c |z|^2$ with the same boundary condition that $\Psi_r|_{\partial \mathbb{D}_r} = -\beta_1 (\ell-1)^2 r^2$. Noting that $\mathrm{dd}^c (\Phi_r - \Psi_r) = -\omega_{\mathbb{D}} + \beta_2 \mathrm{dd}^c |z|^2 \geq 0$ with the vanishing boundary values $(\Phi_r - \Psi_r)|_{\partial \mathbb{D}_r} = 0$, by the maximal principle, $\Phi_r - \Psi_r \leq 0$ on \mathbb{D}_r . Thus we only need to check that $\Psi_r \leq 0$. Indeed, it is easy to see that

$$\Psi(z) = -\beta_1(\ell - 1)^2 r^2 + \beta_2 \cdot (r^2 - |z|^2) \le (-\beta_1(\ell - 1)^2 + \beta_2) \cdot r^2 \le 0$$

for $z \in \mathbb{D}_r$ as long as $\ell \ge \sqrt{\beta_2/\beta_1} + 1$. It is clear that Φ_r is a continuous $\omega_{\mathbb{D}}$ -subharmonic function and satisfies all the desired properties with $\alpha = -\beta_1(\ell - 1)^2$.

Lemma 9.4. For every small $0 < r \ll 1$, there exists a probability measure μ supported outside $\mathbb{B}(x_0, r)$ such that $\mathcal{I}_r(\mu) \leq Cr^4$ for some C > 0 independent of r.

Proof. Using Lemma 9.3, we can find an ω -subharmonic function Φ_r on X such that $\Phi_r = 0$ on $X \setminus \mathbb{B}(x_0, \ell r)$, $-\alpha r^2 \leq \Phi_r \leq 0$ on $\mathbb{B}(x_0, \ell r)$ and $\mathrm{dd}^c \Phi_r = -\omega$ on $\mathbb{B}(x_0, r)$, where ℓ, α are independent of r. Let $\mu := \mathrm{dd}^c \Phi_r + \omega$. Obviously, $\mathrm{supp}(\mu) \subset X \setminus \mathbb{B}(x_0, r)$. Observe that Φ_r is actually the ω -potential of type M of μ . Furthermore, since $\mu = \omega$ outside $\overline{\mathbb{B}}(x_0, \ell r)$, the mass of μ on $\overline{\mathbb{B}}(x_0, \ell r)$ is equal to $\omega(\overline{\mathbb{B}}(x_0, \ell r))$. Hence

$$\mathcal{I}_r(\mu) = -\int_{\overline{\mathbb{B}}(x_0,\ell r)} \Phi_r \,\mathrm{d}\mu - \int_{\overline{\mathbb{B}}(x_0,\ell r)} \Phi_r \,\omega \le \alpha r^2 \cdot \int_{\overline{\mathbb{B}}(x_0,\ell r)} (\mathrm{d}\mu + \omega) \lesssim r^4.$$

This ends the proof.

Acknowledgements. Hao Wu wants to thank Professor Tien-Cuong Dinh for helpful discussions around the functional \mathcal{I}_D . Song-Yan Xie would like to thank Dr. Bingxiao Liu for giving an online short course on higher dimensional hole probabilities in AMSS SCV seminar.

Funding. Song-Yan Xie is partially supported by National Key R&D Program of China Grants No. 2021YFA1003100, No. 2023YFA1010500 and National Natural Science Foundation of China Grant No. 12288201. Hao Wu is supported by the Singapore Ministry of Education Grant MOE-T2EP20121-0013.

REFERENCES

- [1] Luis Alvarez-Gaumé, Jean-Benoît Bost, Gregory Moore, Philip Nelson, and Cumrun Vafa. Bosonization on higher genus Riemann surfaces. *Comm. Math. Phys.*, 112(3):503–552, 1987.
- [2] Turgay Bayraktar. Equidistribution of zeros of random holomorphic sections. *Indiana Univ. Math. J.*, 65(5):1759–1793, 2016.
- [3] Turgay Bayraktar, Dan Coman, Hendrik Herrmann, and George Marinescu. A survey on zeros of random holomorphic sections. *Dolomites Res. Notes Approx.*, 11:1–19, 2018.
- [4] Turgay Bayraktar, Dan Coman, and George Marinescu. Universality results for zeros of random holomorphic sections. *Trans. Amer. Math. Soc.*, 373(6):3765–3791, 2020.
- [5] Pavel Bleher and Xiaojun Di. Correlations between zeros of a random polynomial. J. Statist. Phys., 88(1-2):269–305, 1997.
- [6] Pavel Bleher, Bernard Shiffman, and Steve Zelditch. Universality and scaling of correlations between zeros on complex manifolds. *Invent. Math.*, 142(2):351–395, 2000.
- [7] André Bloch and George Polyá. On the roots of certain algebraic equations. *Proc. London Math. Soc.*, 33:102–114 (1932), 1932.
- [8] Thomas Bloom and Duncan Dauvergne. Asymptotic zero distribution of random orthogonal polynomials. *Ann. Probab.*, 47(5):3202–3230, 2019.
- [9] Thomas Bloom and Bernard Shiffman. Zeros of random polynomials on \mathbb{C}^m . Math. Res. Lett., 14(3):469–479, 2007.
- [10] Jeremiah Buckley, Alon Nishry, Ron Peled, and Mikhail Sodin. Hole probability for zeroes of Gaussian Taylor series with finite radii of convergence. *Probab. Theory Related Fields*, 171(1-2):377–430, 2018.
- [11] Dan Coman, George Marinescu, and Viêt-Anh Nguyên. Holomorphic sections of line bundles vanishing along subvarieties. *Indiana Univ. Math. J.*, 73(2):493–536, 2024.
- [12] Tien-Cuong Dinh, Subhroshekhar Ghosh, and Hao Wu. Hole event for random holomorphic sections on compact riemann surfaces. arXiv:2402.11672, 2024.
- [13] Tien-Cuong Dinh, Xiaonan Ma, and George Marinescu. Equidistribution and convergence speed for zeros of holomorphic sections of singular Hermitian line bundles. J. Funct. Anal., 271(11):3082–3110, 2016.
- [14] Tien-Cuong Dinh, Xiaonan Ma, and Viêt-Anh Nguyên. Equidistribution speed for Fekete points associated with an ample line bundle. *Ann. Sci. Éc. Norm. Supér.* (4), 50(3):545–578, 2017.

- [15] Tien-Cuong Dinh and Nessim Sibony. Distribution des valeurs de transformations méromorphes et applications. *Comment. Math. Helv.*, 81(1):221–258, 2006.
- [16] Alexander Drewitz, Bingxiao Liu, and George Marinescu. Large deviations for zeros of holomorphic sections on punctured riemann surfaces. arXiv:2109.09156, to appear in Michigan Math. J., 2021.
- [17] Otto Forster. *Lectures on Riemann surfaces*, volume 81 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1981. Translated from the German by Bruce Gilligan.
- [18] Subhroshekhar Ghosh and Alon Nishry. Point processes, hole events, and large deviations: random complex zeros and Coulomb gases. *Constr. Approx.*, 48(1):101–136, 2018.
- [19] Subhroshekhar Ghosh and Alon Nishry. Gaussian complex zeros on the hole event: the emergence of a forbidden region. *Comm. Pure Appl. Math.*, 72(1):3–62, 2019.
- [20] Subhroshekhar Ghosh and Ofer Zeitouni. Large deviations for zeros of random polynomials with i.i.d. exponential coefficients. *Int. Math. Res. Not. IMRN*, 2016(5):1308–1347, 2016.
- [21] Phillip Griffiths and Joseph Harris. *Principles of algebraic geometry*. Wiley Classics Library. John Wiley & Sons, Inc., New York, 1994. Reprint of the 1978 original.
- [22] Robert C. Gunning. *Lectures on Riemann surfaces, Jacobi varieties*. Mathematical Notes, No. 12. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1972.
- [23] Manjunath Krishnapur. Overcrowding estimates for zeroes of planar and hyperbolic Gaussian analytic functions. *J. Stat. Phys.*, 124(6):1399–1423, 2006.
- [24] Andriy O. Kuryliak and Oleh B. Skaskiv. Analytic Gaussian functions in the unit disc: probability of zeros absence. *Mat. Stud.*, 59(1):29–45, 2023.
- [25] Serge Lang. Introduction to Arakelov theory. Springer-Verlag, New York, 1988.
- [26] George Marinescu and Duc-Viet Vu. Bergman kernel functions associated to measures supported on totally real submanifolds. *J. Reine Angew. Math.*, 810:217–251, 2024.
- [27] Pertti Mattila. *Geometry of sets and measures in Euclidean spaces*, volume 44 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1995. Fractals and rectifiability.
- [28] Raghavan Narasimhan. *Compact Riemann surfaces*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 1992.
- [29] Alon Nishry. Asymptotics of the hole probability for zeros of random entire functions. *Int. Math. Res. Not. IMRN*, 2010(15):2925–2946, 2010.
- [30] Alon Nishry. Hole probability for entire functions represented by Gaussian Taylor series. J. Anal. Math., 118(2):493–507, 2012.
- [31] Alon Nishry and Aron Wennman. The forbidden region for random zeros: appearance of quadrature domains. *Comm. Pure Appl. Math.*, 77(3):1766–1849, 2024.
- [32] Junjiro Noguchi and Jörg Winkelmann. Nevanlinna theory in several complex variables and Diophantine approximation, volume 350 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer, Tokyo, 2014.
- [33] Yuval Peres and Bálint Virág. Zeros of the i.i.d. Gaussian power series: a conformally invariant determinantal process. *Acta Math.*, 194(1):1–35, 2005.
- [34] Christian Pommerenke. Boundary behaviour of conformal maps, volume 299 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 1992.
- [35] Min Ru. *Nevanlinna theory and its relation to Diophantine approximation*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, second edition, [2021] ©2021.
- [36] Edward B. Saff and Vilmos Totik. *Logarithmic potentials with external fields*, volume 316 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1997. Appendix B by Thomas Bloom.
- [37] Guokuan Shao. Equidistribution of zeros of random holomorphic sections for moderate measures. *Math. Z.*, 283(3-4):791–806, 2016.
- [38] Larry A. Shepp and Robert J. Vanderbei. The complex zeros of random polynomials. *Trans. Amer. Math. Soc.*, 347(11):4365–4384, 1995.
- [39] Bernard Shiffman. Asymptotic expansion of the variance of random zeros on complex manifolds. J. *Geom. Anal.*, 31(8):8607–8631, 2021.

- [40] Bernard Shiffman and Steve Zelditch. Distribution of zeros of random and quantum chaotic sections of positive line bundles. *Comm. Math. Phys.*, 200(3):661–683, 1999.
- [41] Bernard Shiffman and Steve Zelditch. Equilibrium distribution of zeros of random polynomials. Int. Math. Res. Not., 2003(1):25–49, 2003.
- [42] Bernard Shiffman and Steve Zelditch. Number variance of random zeros on complex manifolds. *Geom. Funct. Anal.*, 18(4):1422–1475, 2008.
- [43] Bernard Shiffman and Steve Zelditch. Number variance of random zeros on complex manifolds, II: smooth statistics. *Pure Appl. Math. Q.*, 6(4):1145–1167, 2010.
- [44] Bernard Shiffman, Steve Zelditch, and Qi Zhong. Random zeros of complex manifolds: conditional expectations. J. Inst. Math. Jussieu, 10(3):753–783, 2011.
- [45] Bernard Shiffman, Steve Zelditch, and Scott Zrebiec. Overcrowding and hole probabilities for random zeros on complex manifolds. *Indiana Univ. Math. J.*, 57(5):1977–1997, 2008.
- [46] Yum-Tong Siu. Some recent transcendental techniques in algebraic and complex geometry. In Proceedings of the International Congress of Mathematicians, Vol. I (Beijing, 2002), pages 439–448. Higher Ed. Press, Beijing, 2002.
- [47] Mikhail Sodin and Boris Tsirelson. Random complex zeroes. III. Decay of the hole probability. *Israel J. Math.*, 147:371–379, 2005.
- [48] Isaac Todhunter. A history of the mathematical theory of probability. Stechert, New York, 1931.
- [49] Masatsugu Tsuji. *Potential theory in modern function theory*. Chelsea Publishing Co., New York, 1975. Reprinting of the 1959 original.
- [50] Ofer Zeitouni and Steve Zelditch. Large deviations of empirical measures of zeros of random polynomials. *Int. Math. Res. Not. IMRN*, 2010(20):3935–3992, 2010.
- [51] Steve Zelditch. Large deviations of empirical measures of zeros on Riemann surfaces. Int. Math. Res. Not. IMRN, 2013(3):592–664, 2013.
- [52] Junyan Zhu. Hole probabilities of SU(m + 1) Gaussian random polynomials. Anal. PDE, 7(8):1923–1968, 2014.
- [53] Scott Zrebiec. The zeros of flat Gaussian random holomorphic functions on \mathbb{C}^n , and hole probability. *Michigan Math. J.*, 55(2):269–284, 2007.

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE - 10, LOWER KENT RIDGE ROAD - SINGAPORE 119076

Email address: e0011551@u.nus.edu

ACADEMY OF MATHEMATICS AND SYSTEM SCIENCE & HUA LOO-KENG KEY LABORATORY OF MATHEMATICS, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA; SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, CHINA

Email address: xiesongyan@amss.ac.cn