Aritra Banik 🖂

National Institute of Science, Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar 752050, Odisha, India

Fedor V. Fomin ⊠ **●** University of Bergen, Norway

Petr A. Golovach \square

University of Bergen, Norway

Tanmay Inamdar ⊠ Indian Institute of Technology Jodhpur, Jodhpur, India

Satyabrata Jana ⊠ Duriversity of Warwick, UK

Saket Saurabh 🖂 🛽

The Institute of Mathematical Sciences, HBNI, Chennai, India University of Bergen, Norway

— Abstract

VERTEX (s, t)-CUT and VERTEX MULTIWAY CUT are two fundamental graph separation problems in algorithmic graph theory. We study matroidal generalizations of these problems, where in addition to the usual input, we are given a representation $R \in \mathbb{F}^{r \times n}$ of a linear matroid $\mathcal{M} = (V(G), \mathcal{I})$ of rank r in the input, and the goal is to determine whether there exists a vertex subset $S \subseteq V(G)$ that has the required cut properties, as well as is independent in the matroid \mathcal{M} . We refer to these problems as INDEPENDENT VERTEX (s, t)-CUT, and INDEPENDENT MULTIWAY CUT, respectively. We show that these problems are fixed-parameter tractable (FPT) when parameterized by the solution size (which can be assumed to be equal to the rank of the matroid \mathcal{M}). These results are obtained by exploiting the recent technique of flow augmentation [Kim et al. STOC '22], combined with a dynamic programming algorithm on flow-paths á la [Feige and Mahdian, STOC '06] that maintains a representative family of solutions w.r.t. the given matroid [Marx, TCS '06; Fomin et al., JACM]. As a corollary, we also obtain FPT algorithms for the independent version of ODD CYCLE TRANSVERSAL. Further, our results can be generalized to other variants of the problems, e.g., weighted versions, or edge-deletion versions.

2012 ACM Subject Classification Theory of computation \rightarrow Design and analysis of algorithms

Keywords and phrases *s-t*-cut, multiway Cut, matroid, odd cycle transversal, feedback vertex set, fixed-parameter tractability

Funding The research leading to these results has received funding from the Research Council of Norway via the project BWCA (grant no. 314528).

1 Introduction

Studying problems in graphs and sets with additional constraints given by algebraic structures such as the matroid or submodularity is one of the well-established research directions. These versions not only generalize classical problems, but also show combinatorial interplay between graph structure and linear algebra. Some of the well-known problems in this direction include subsmodular versions of VERTEX COVER, SHORTEST PATH, MATCHING, or MIN-(s, t)-CUT [12, 15] as well as coverage problems such as MAX COVERAGE with matroid constraints [2, 34]. In this paper we study some of the classical cut problems in the realm of parameterized complexity in the presence of constraints given by matroids.

It is customary to begin the discussion on our framework with VERTEX COVER (given a graph G and a positive integer k, find a subset of size at most k that covers all edges), which is a poster-child problem in parameterized complexity. Among its myriad of generalizations studied in the literature, one of particular relevance to the present discussion is ranked version of VERTEX COVER arising in the works of Lovász [22, 23]. He defined RANK VERTEX COVER in his study of critical graphs having k-sized vertex cover, by defining the notion of a framework, also known as pregeometric graphs, which refers to a pair (G, \mathcal{M}) , where G is a graph and $\mathcal{M} = (V(G), \mathcal{I})$ is a matroid on the vertex set of G. In RANK VERTEX COVER, we are given a framework (G, \mathcal{M}) and a parameter k, and the goal is to determine whether G has a vertex cover S whose rank—which is defined as the maximum-size independent subset of S—is upper bounded by k. An FPT algorithm for RANK VERTEX COVER (or indeed RANK d-HITTING SET) parameterized by the rank is folklore. Meesum et al. [30] used RANK VERTEX COVER as a natural target problem for giving a compression from a different parameterization for VERTEX COVER. However, even for a slightly more involved problem, such as FEEDBACK VERTEX SET (FVS), the ranked version immediately becomes W[1]-hard parameterized by the rank. This is similar to the phenomena seen when we consider weighted problems such as WEIGHTED VERTEX COVER parameterized by weight [31, 33].

Having faced with such a obstacle, we study an alternate generalization of FVS in frameworks. In INDEPENDENT FVS (IFVS), where we are given a framework (G, \mathcal{M}) , we want to determine whether there exists some $S \subseteq V(G)$ such that (i) S is an independent set in the matroid, and (ii) G - S is acyclic. This happens to be a "sweet spot" between the vanilla version and the ranked version – we can adopt the textbook algorithms, and obtain a polynomial kernel and $\mathcal{O}^*(c^k)$ time FPT algorithm when parameterized by k ($\mathcal{O}^*(\cdot)$) suppresses polynomial factors in the input size.), which is equal to the solution-size (and can also be assumed to be equal to the rank of the matroid, by appropriately truncating it). This was the starting point of our result. Note that IFVS reduces to vanilla FVS when the matroid \mathcal{M} is a uniform matroid of rank k, and hence is NP-hard. Indeed, this gives us a recipe for defining the "independent" versions of classical vertex-subset problems in graphs. We give a formal definition of a prototypical problem below.

- INDEPENDENT Π (where Π is a vertex-subset problem)

A graph G = (V, E), a matroid $\mathcal{M} = (V, \mathcal{I})$ of rank r. Input: **Question:** Does there exist a set $S \subseteq V(G)$ such that

- **S** is an independent set in \mathcal{M} , i.e., $S \in \mathcal{I}$, and
- **—** S is a feasible solution for Π

On the other hand, even for a classically polynomial-time solvable problem such as VERTEX (s, t)-CUT, the corresponding independent version can be shown to be NP-hard even when \mathcal{M} is a partition matroid [14, 24].

1.1 Our Results and Methods.

Let G be a graph and s, t be a distinct pair of vertices in G. A vertex (s,t)-cut is a set S of vertices such that G - S has no path from s to t. Given a graph G, a distinct pair of vertices $s, t \in V(G)$, and an integer k, the (s,t)-cut problem asks if a vertex (s,t)-cut of size at most k exists. Note that (VERTEX) (s,t)-CUT forms the bedrock of classical graph theory and algorithms via Menger's theorem and cut-flow duality. Thus, numerous constrained versions of (s,t)-CUT have been studied in the literature, and matroid independence is a fundamental constraint that remains unexplored to the best of our knowledge. Another motivation for studying INDEPENDENT VERTEX (s,t)-CUT (IVstC) comes from the fact that it appears as a natural subproblem while trying to solve the independent version of ODD CYCLE TRANSVERSAL (aka OCT – given G, k, check whether we can delete at most k vertices from G such that the remaining graph is bipartite) via the so-called iterative compression technique [32]. Thus, we would like to design an FPT algorithm for IVstC, and in turn IOCT, with running times that are as close to the running respective vanilla versions (note that (s, t)-cut is in P, whereas OCT admits an $\mathcal{O}^*(3^k)$ time algorithm, see [32]).

There are two possible approaches that one could take for designing an FPT algorithm for a constrained (s, t)-cut problem, such as IVstC. First approach is to reduce the treewidth of the input graph by using the methodology of Marx, O'Sullivan, and Razgon [28, 29], and then perform a dynamic programming on a bounded treewidth graph. While designing a dynamic programming algorithm for IVstC on bounded-treewidth graphs appears to be straightforward using the technique of representative sets ([27, 11]), it is not obvious that we could use this result as a black-box-we have to be careful while performing any modifications to the graph, since we also have a matroid on the vertex-set. Further, since the treewidth bound obtained via this result is exponential in the size of the solution we are trying to find, it seems that such an approach would result in a double-exponential running time. Thus, we do not pursue this approach further, and the following question remains.

Question 1. Does INDEPENDENT VERTEX (s, t)-CUT (and in turn, INDEPENDENT ODD CYCLE TRANSVERSAL) admit a single-exponential, preferably $2^{\mathcal{O}(k)} \cdot n^{\mathcal{O}(1)}$ FPT algorithm?

Another promising approach to tackle this question is given by Feige and Mahdian [8], who give a very general recipe for solving constrained minimum (s, t)-cut problems. This is done via performing dynamic programming on the flow-paths obtained via Menger's theorem. Naturally, in order to use this approach, it is crucial that we are looking for a *minimum* (s,t)-cut with certain additional constraints. However, this appears to be a roadblock since a minimum independent (s, t)-cut can be much larger than the minimum (not-necessarilyindependent) (s,t)-cut, and hence Menger's theorem is not applicable. This is where the recent technique of flow augmentation [16, 18] comes to the rescue. At a high level, this a powerful tool that can "augment" the graph by adding additional edges, thus lifting the size of minimum (s,t)-cut to a higher value (hence, flow augmentation), such that an unknown minimal (s,t)-cut becomes a minimum (s,t)-cut. Note that minimality of a solution can be assumed w.l.o.g. since it is compatible with independence as well as cut property. Indeed, all of the aforementioned problems, namely, FVS, OCT, (s, t)-CUT satisfy the minimality condition. Hence, using flow augmentation, at the overhead of $2^{\mathcal{O}(k \log k)} \cdot n^{\mathcal{O}(1)}$, we can assume that we are looking for an independent set that is a minimum (s, t)-cut in the augmented graph. However, due to certain technical obstacles while applying flow augmentation, we

Π in Independent Π	Matroid access	Result
Feedback Vertex Set	Oracle	$\mathcal{O}^*(10^k)$ and poly kernel
Vertex/Edge (s, t) -cut	Representation	$\mathcal{O}^*(2^{\mathcal{O}(k^4\log k)})$
Vertex (s, t) -Cut	Oracle	No $f(k) \cdot n^{o(k)}$ algorithm
Odd Cycle Transversal	Representation	$\mathcal{O}^*(2^{\mathcal{O}(k^4\log k)})$
Multiway Cut	Representation	$\mathcal{O}^*(2^{\mathcal{O}(k^4 \log k)})$

Table 1 Our Results.

need to solve the *directed version* of (s, t)-cut using the approach of [8]. Now, we perform a dynamic programming on the flow-paths given by a decomposition theorem from [8] (which needs to be extended to directed graphs), and maintain a representative family of all partial solutions in the DP table. Note that for this, we assume that the matroid \mathcal{M} is given as a representation matrix $R \in \mathbb{F}^{r \times n}$, and the field operations can be performed in polynomial-time¹. Thus, in fact, our algorithm solves a more general problem where it outputs the representative family of all minimum independent (s, t)-cuts. This property is crucial when we want to use this problem as a subroutine to solve a more general problem, to which we return in the next paragraph. Note that for the sake of conceptual simplicity and to avoid references to success probabilities, we only give the deterministic running times arising from $\mathcal{O}^*(2^{\mathcal{O}(k^4 \log k)})$ -time deterministic flow augmentation in the table and in the paper. However, note that by instead using the randomized version, we can improve the running time to $2^{\mathcal{O}(k \log k)}$, which takes us closer to our initial goal. Furthermore, due to versatility of the representative sets framework, our algorithms also extend to the weighted versions of these problems, where we want to find a minimum-weight independent set that is a feasible solution to Π . Note that one of the original motivations for the flow augmentation technique was solving one such problem, i.e., BI-OBJECTIVE DIRECTED (or weighted) (s, t)-CUT [18]. Next, having (almost) answered **Question 1**, we look toward an even more general problem.

Independent Multiway Cut. Let G be a graph and $T \subseteq V(G)$ be a set of terminals. A vertex multiway cut is a set S of vertices such that every component of G-S contains at most one vertex of T. Given a graph G, terminals T, and an integer k, the (VERTEX) MULTIWAY CUT (MWC) problem asks if a vertex multiway cut of size at most k exists. MWC is a natural generalization of (s, t)-cut. This been studied extensively in the domain of parameterized and approximation algorithms. Note that the vanilla version of the problem was classically shown to be FPT using the technique of important separators [25]. On the other hand, it is not clear whether this technique is any useful in presence of matroid constraints, i.e., for INDEPENDENT MULTIWAY CUT (IMWC). Even when armed with an FPT algorithm for INDEPENDENT (s, t)-CUT as a subroutine, it is not trivial to design an FPT algorithm for IMWC.

¹ A matroid \mathcal{M} is representable over a field \mathbb{F} if there exists a matrix $R \in \mathbb{F}^{r \times n}$, and a bijection between the ground set of \mathcal{M} and the columns of R, such that a set is independent in \mathcal{M} iff the corresponding set of columns is independent in \mathbb{F} .

The algorithm initially begins by trying to find a multiway cut (separator) S of size at most k for the terminal set T in G – here we completely ignore independence constraints, and such a separator can be found in $\mathcal{O}^*(2^k)$ using the result of Cygan et al. [6]. Note that if the graph lacks such a separator, it also cannot have an independent separator. W.l.o.g. suppose that we find a separator S that is also minimal. Hence, for every vertex v in the set S, there exists a path that connects a pair of terminal vertices and intersecting the set S solely at v. This fact brings us to the conclusion that every independent separator either includes v or some vertices of terminal components (we say that a connected component in G-Sis a *terminal component* if it contains some terminal). Moreover, each vertex of S has a neighbor in at least two terminal components. With this information, we attempt to design a recursive algorithm by assuming that the "true solution" O to IMC of size k either intersects S, or some terminal component (that can be identified). However, it is also possible that a terminal component contains the entire solution O. In this case, we cannot make a recursive call to the same algorithm, since the parameter does not decrease. To address this scenario, we introduce the concept of a strong separator (Definition 36), which is a minimal separator S with the following properties: any minimal solution O satisfies (i) $O \cap S \neq \emptyset$, or (ii) there exists a pair of terminal components—that can be identified in polynomial time—such that O intersects at least one of them, with the size of intersection being between 1 and k-1. Further, such a strong separator can be found in time $\mathcal{O}^*(2^k)$ time (Lemma 39) by applying "pushing arguments". This lets us design a recursive divide-and-conquer algorithm, and due to property (ii), we know that in each recursive call, the parameter always strictly decreases. For handling the matroid independence constraints, in fact we need to strengthen the algorithm to return a representative family of all possible multiway cuts. This is where it is useful that our algorithm for IVstC also returns such a family, since this algorithm is used in the base case of the algorithm (when |T| = 2). This strategy results $\mathcal{O}^*(2^{\mathcal{O}(k^4 \log k)})$ time deterministic (or $\mathcal{O}^*(2^{\mathcal{O}(k \log k)})$ time randomized) algorithm for IMWC, where the running time is dominated by the flow augmentation step required in the base case of IVstC. The Table 1 depicts all the results in this paper.

Motivation for independence constraints. One motivation for studying such independent versions of classical graph problems comes from the fact that matroid-independence capture several interesting constraints on the solution vertices. One prominent example of this is the colorful versions of the problem that arise naturally from the color-coding technique introduced by Alon, Yuster and Zwick [1], where a problem such as k-PATH, i.e., determining whether the given graph contains a path of length at least k. While k-PATH itself is difficult to approach using combinatorial techniques, there is a simple dynamic programming algorithm, if we want to find a colorful k-path, i.e., where we are given a graph whose vertices are colored from $\{1, 2, \ldots, k\}$, and we want to determine whether the graph contains a path of length k whose vertices are colored with distinct colors [1]. Then, a standard argument shows that a hypothetical unknown solution will get colored with distinct colors if each vertex is assigned a color uniformly at random. On the flip side, colorful versions of the problems such as INDEPENDENT SET, CLIQUE, are also convenient starting points in various W[1]-hardness reductions. Note that colorful constraints is one of the simplest examples of matroid constraints – it is simply a partition matroid, where the coloring of vertices corresponds to a partition of the vertex set, and the solution may contain (at most) one vertex from each partition. Slightly more general partition matroid constriants can capture "local budgets" on the solution from each class of vertices, which can be used to model different fairness constraints, i.e., we cannot delete too many vertices of the same type to

achieve certain graph property, which can be "unfair" for that type. Such constraints on solution are rather popular in clustering [13, 19], which are indeed motivated from such fairness considerations.

2 **Preliminaries**

Notations. Let [n] be the set of integers $\{1, \ldots, n\}$. For a (directed/undirected) graph G, we denote the set of vertices of G by V(G) and the set of edges by E(G). For an undirected graph, we denote the set of non-edges by $\overline{E}(G)$, that is, $\overline{E}(G) = \binom{V(G)}{2} \setminus E(G)$, where $\binom{V(G)}{2}$ is the set of all unordered pairs of distinct vertices in V(G); whereas for a directed graph $G.\overline{E}(G) := (V(G) \times V(G)) \setminus E(G)$. For notational convenience, even in the undirected setting, we use the notation (a, b) instead of $\{a, b\}$ – note that due to this convention, (a, b) = (b, a) in undirected graphs. For a un/directed graph G = (V, E) and a subset S of un/directed edges in $\overline{E}(G)$, we use the notation G + S to mean the graph $G' = (V, E \cup S)$. In an un/directed graph G, and $A, B \in V(G)$, we say that B is reachable from A if there is an un/directed path from some $u \in A$ to $v \in B$ in G. Note that reachability is symmetric in undirected graphs. For a vertex-subset $X \subseteq V(G) \setminus (A \cup B)$ (resp. edge-subset $X \subseteq E(G)$), we say that X is an (A, B)-cut if B is not reachable from A in G - X. Further, we say that X is an inclusion-minimal (A, B)-cut (or simply, a minimal (A, B)-cut), if no strict subset of X is also an (A, B)-cut. More generally, let $\mathcal{A} = \{A_1, A_2, \ldots, A_t\}$ be a collection of vertex-subsets. We say that X is a multiway cut for \mathcal{A} if X is an (A_i, A_j) -cut for each pair $A_i, A_j \in \mathcal{A}$, and a minimal multiway cut is defined analogously. For vertex multiway cut $X \subseteq V(G)$ for \mathcal{A} (resp. (A, B)-cut), we also require that $X \cap A_i = \emptyset$ for all $A_i \in \mathcal{A}$ (resp. $X \cap A = X \cap B = \emptyset$). Finally, when $A = \{s\}, B = \{t\}$, we may slightly abuse the notation and use (s, t)-cut or s-t cut instead of $(\{s\}, \{t\})$ -cut. When we have a graph G and a subset of vertices $X \subseteq V(G)$, the operation of *identifying* the vertices of X is defined as follows: we first eliminate the vertex set X, then introduce a new vertex v_X , make v_X adjacent to all vertices in the open neighborhood of X, and finally remove any multi-edges.

Matroids.

Let $\mathcal{M} = (U, \mathcal{I})$ be a matroid. Each set $S \in \mathcal{I}$ is called an independent set and an inclusionwise maximal independent set is known as a basis of the matroid \mathcal{M} . From the third property of matroids, it is easy to observe that every inclusion-wise maximal independent set has the same size; this size is referred to as the rank of \mathcal{M} , and denoted by $\mathsf{rank}(\mathcal{M})$. In the following, we will use r to denote $rank(\mathcal{M})$.

The rank of a set $P \subseteq U$ is defined as the size of the largest subset of P that is independent, and is denoted by $\mathsf{rank}(P)$. Note that $\mathsf{rank}(P) \leq \mathsf{rank}(Q)$ if $P \subseteq Q$. For any $P \subseteq U$, define the closure of P, $\mathsf{cl}(P) \coloneqq \{x \in U : \mathsf{rank}(P \cup \{x\}) = \mathsf{rank}(P)\}$. Note that $\mathsf{cl}(P) \subseteq \mathsf{cl}(Q)$ if $P \subseteq Q$.

Finally, we will use $U(\mathcal{M})$ and $\mathcal{I}(\mathcal{M})$ to refer to the ground set and the independent-set collection of \mathcal{M} , respectively.

A matroid $\mathcal{M} = (U, \mathcal{I})$ is said to be *representable* over a field \mathbb{F} if there exists a matrix $R \in \mathbb{F}^{r \times n}$ where $r = \operatorname{rank}(\mathcal{M}), |U| = n$, and a bijection from U to the columns of R, such that for any $S \subseteq U, S \in \mathcal{I}$ iff the corresponding set of columns is linearly independent over F.

For a matroid $\mathcal{M} = (U, \mathcal{I})$ and an element $u \in U$, the matroid obtained by contracting u is represented by $\mathcal{M}' = \mathcal{M}/u$, where $\mathcal{M}' = (U \setminus \{u\}, \mathcal{I}')$, where $\mathcal{I}' =$ $\{S \subseteq U \setminus \{u\} : S \cup \{u\} \in \mathcal{I}\}$. Note that $\mathsf{rank}(\mathcal{M}') = \mathsf{rank}(\mathcal{M}) - 1$. For any $0 \le k \le r$,

7

one can define a truncated version of the matroid as follows: $\mathcal{M}_k = (U, \mathcal{I}_{k'})$, where $\mathcal{I}_k = \{S \in \mathcal{I} : |S| \leq k\}$. Note that $\operatorname{rank}(\mathcal{M}_k) = k$, and one can also think of \mathcal{M}_k as an intersection ² of \mathcal{M} with a uniform matroid of rank k. It is easy to verify that the contraction as well as truncation operation results in a matroid. Furthermore, given a linear representation of a matroid \mathcal{M} , a linear representation of the matroid resulting from contraction/truncation can be computed in (randomized) polynomial time [26, 20].

We call a family of subsets (of U) of size $p \ge 0$ as a *p*-family. Next, we state the following crucial definition of representative families.

▶ **Definition 1** ([9, 4]). Let $\mathcal{M} = (U, \mathcal{I})$ be a matroid. For two subsets $A, B \subseteq U$, we say that A fits B (or equivalently B fits A), if $A \cap B = \emptyset$ and $A \cup B \in \mathcal{I}$.

Let $\mathcal{A} \subseteq \mathcal{I}$ be a p-family for some $0 \leq p \leq \operatorname{rank}(\mathcal{M})$. A subfamily $\mathcal{A}' \subseteq \mathcal{A}$ is said to q-represent \mathcal{A} if for every set B of size q such that there is an $A \in \mathcal{A}$ that fits B, then there is an $A' \in \mathcal{A}'$ that also fits B. If \mathcal{A}' q-represents \mathcal{A} , we write $\mathcal{A}' \subseteq_{rep}^q \mathcal{A}$.

The following proposition lets us compute representative families efficiently.

▶ **Proposition 2** ([9, 4]). There is an algorithm that, given a matrix R over a field \mathbb{F} , representing a matroid $\mathcal{M} = (U, \mathcal{I})$ of rank k, a p-family \mathcal{A} of independent sets in \mathcal{M} , and an integer q such that p + q = k, computes a q-representative family $\mathcal{A}' \subseteq_{rep}^q \mathcal{A}$ of size at most $\binom{p+q}{p}$ using at most $\mathcal{O}(|\mathcal{A}|(\binom{p+q}{p})p^{\omega} + (\binom{p+q}{p})^{\omega-1})$ operations over \mathbb{F} . Furthermore, an analogous result holds even if $p + q \leq k$, albeit depending on the field \mathbb{F} , the algorithm may be randomized. ³

For a *p*-family \mathcal{P} and a *q*-family \mathcal{Q} , we define $\mathcal{P} \bullet \mathcal{Q} \coloneqq \{P \cup Q : P \in \mathcal{P}, Q \in \mathcal{Q}, P \text{ fits } Q\}$ (this operation is also known as subset convolution). The following properties of representative families follow straightfordly from the definitions, and formal proofs can be found in [5].

▶ Proposition 3 ([5]). Let $\mathcal{M} = (U, \mathcal{I})$ be a matroid and \mathcal{A} be a p-family.

- $= If \mathcal{A}_1 \subseteq_{rep}^{r-p} \mathcal{A}, and \mathcal{A}_2 \subseteq_{rep}^{r-p} \mathcal{A}_1, then, \mathcal{A}_2 \subseteq_{rep}^{r-p} \mathcal{A}.$
- $= If \mathcal{A}_1 \subseteq_{rep}^{r-p} \mathcal{A}, then \mathcal{A} = \emptyset iff \mathcal{A}_1 = \emptyset.$
- = Let \mathcal{P} be a *p*-family and \mathcal{Q} be a *q*-family. Let $\mathcal{P}' \subseteq_{rep}^{r-p} \mathcal{P}$, and $\mathcal{Q}' \subseteq_{rep}^{r-q} \mathcal{Q}$. Then, $\mathcal{P}' \bullet \mathcal{Q}' \subseteq_{rep}^{r-p-q} \mathcal{P} \bullet \mathcal{Q}$.

3 FPT Algorithm for INDEPENDENT VERTEX (s, t)-CUT

In this section, we design an FPT algorithm for INDEPENDENT VERTEX (s, t)-CUT. To this end, we define some notation. Consider a graph G = (V, E), a matroid $\mathcal{M} = (U, \mathcal{I})$ of rank r where $V(G) \setminus \{s, t\} \subseteq U$, and two vertices $s, t \in V(G)$. By iteratively checking values $k = 1, 2, \ldots$, and running the algorithm described below, we may inductively assume that we are working with a value $0 \leq k \leq r$, such that there is no independent (s, t)-cut in Gfor any k' < k. This follows from the fact that, our generalized problem, as defined below, requires the algorithm to return a set corresponding to k' whose emptiness is equivalent to

² Intersection of two matroids $\mathcal{M}_1 = (U, \mathcal{I}_1)$ and $\mathcal{M}_2 = (U, \mathcal{I}_2)$ defined over common ground set U is the set family (U, \mathcal{I}) , where $\mathcal{I} = \{X \subseteq U : X \in \mathcal{I}_1 \cap \mathcal{I}_2\}$. Note that (U, \mathcal{I}) is not necessarily a matroid in general.

³ Here, we need to compute the representation of an appropriately truncated matroid. For this, only randomized polynomial algorithm is known in general [27]; however for certain kinds of fields, it can be done in polynomial time [21].

concluding that the there is no independent (s, t)-cut of size k'. It is important to note that, depending on whether we are addressing INDEPENDENT VERTEX (s, t)-CUT as a standalone problem or as a sub-procedure, we may either obtain the smallest value of k for free, or we may need to perform iterations. Regardless, we can safely assume that we are operating with a value of this nature. For any $0 \le k \le r$, let

$$\mathcal{F}(s,t,k) \coloneqq \{S \subseteq V(G) \setminus \{s,t\} : S \in \mathcal{I}, |S| = k, S \text{ is a minimal } (s,t)\text{-cut.}\}$$

In fact, we design an FPT algorithm for a following generalization of the problem, stated below.

– Generalized Independent Vertex (s, t)-Cut

Input: A graph G, a vertex set $Q \subseteq V(G)$ containing s, t, a representation $R \in \mathbb{F}^{r \times n}$ of a matroid $\mathcal{M} = (U, \mathcal{I})$ of rank r where $V \setminus Q \subseteq U$, an integer $0 \leq k \leq r$. **Output:** $\mathcal{F}'(s, t, k) \subseteq_{\text{rep}}^{r-k} \mathcal{F}(s, t, k)$.

Note that this generalizes INDEPENDENT VERTEX (s, t)-CUT, since the latter problem is equivalent to determining whether $\mathcal{F}(s, t, k)$ (or, equivalently, $\mathcal{F}'(s, t, k) \subseteq_{\mathrm{rep}}^{r-k} \mathcal{F}(s, t, k)$, via Proposition 3) is non-empty for some $0 \leq k \leq r$.



Figure 1 Schematic depiction of sequence of reductions from undirected vertex (s, t)-cut to directed vertex (s, t)-cut via flow augmentation. These details are omitted from the short version.

▶ **Theorem 4.** Consider an instance $(G, \mathcal{M}, s, t, Q, k)$ of GENERALIZED INDEPENDENT VERTEX (s,t)-CUT, where there exists no independent (s,t)-cut in G of size less than k. Then, there exists a deterministic algorithm for GENERALIZED INDEPENDENT VERTEX (s,t)-CUT that runs in time $2^{\mathcal{O}(k^4 \log k)} \cdot 2^{\mathcal{O}(r)} \cdot n^{\mathcal{O}(1)}$, and outputs $\mathcal{F}'(s,t,k) \subseteq_{rep}^{r-k} \mathcal{F}(s,t,k)$ such that $|\mathcal{F}'(s,t,k)| \leq 2^r$. Here, $\mathcal{F}(s,t,k)$ is the collection of all independent (s,t)-cuts of size k.

Flow augmentation.

Below we present the recent results regarding the so-called *flow-augmentation* technique [17, 18].

▶ Proposition 5 (Flow-augmentation (randomized) for Directed Graphs, [16, 18]). There exists a randomized polynomial-time algorithm that, given a directed graph G, two vertices $s, t \in V(G)$, and an integer k, outputs a set $A \subseteq V(G) \times V(G)$ such that the following holds: for every minimal (s,t)-cut $Z \subseteq E(G)$ of size at most k, with probability $2^{-\mathcal{O}(k \log k)}$ the edge set Z remains an st-cut in G + A and, furthermore, Z is a minimum (s,t)-cut in G + A.

▶ **Proposition 6** (Flow-augmentation (deterministic) for Directed Graphs [16, 18]). There exists an algorithm that, given a directed graph G, two vertices $s, t \in V(G)$, and an integer k, in time $2^{\mathcal{O}(k^4 \log k)} \cdot |V(G)|^{\mathcal{O}(1)}$ outputs a set $\mathcal{A} \subseteq 2^{V(G) \times V(G)}$ of size $2^{\mathcal{O}(k^4 \log k)} \cdot (\log n)^{\mathcal{O}(k^3)}$ such that the following holds: for every minimal (s, t)-cut $Z \subseteq E(G)$ of size at most k, there exists $A \in \mathcal{A}$ such that the edge set Z remains an (s, t)-cut in G + A and, furthermore, Z is a minimum (s, t)-cut in G + A.

3.1 Transforming the solution into a (directed) minimum cut via Flow Augmentation

In this section we give a reduction from undirected vertex (s,t)-cut to directed vertex (s,t)-cut. This process consists of four phases. We adopt the following convention. Let k denote the size of the minimum (s,t)-cut and let r denote the rank of the matroid \mathcal{M} . If k > r, then we immediately terminate and output No. Therefore, assume that $0 \le k \le r$.

Phase I. Undirected Vertex (s,t)-**cut** \Rightarrow **Directed Vertex** (s,t)-**cut**. We are given an instance $(G, s, t, \mathcal{M}, Q, k)$ of INDEPENDENT UNDIRECTED VERTEX (s,t)-CUT to an instance of $(G_1, s, t, \mathcal{M}_1, Q_1, k)$ of INDEPENDENT DIRECTED VERTEX (s,t)-CUT. First, given G, we define a directed graph G_1 as follows.

The vertex set remains the same, that is, $V(G_1) = V(G), Q_1 = Q$.

For each edge e = uv in G, add two arcs (u, v) and (v, u) in $A(G_1)$.

It is easy to see that S is a vertex (s, t)-cut in G if and only if S is a vertex (s, t)-cut in G_1 .

Matroid Transformation. Here we construct a matroid \mathcal{M}_1 for instance (G_1, k) . Initially, the matroid \mathcal{M} is given as an input with G. We keep the same matroid for the new instance (G_1, k) , that is, $\mathcal{M}_1 := \mathcal{M}$. Now, toward the correctness of the transformation we have the following claim. Let $\mathcal{F}_1(s, t, k)$ denote the set of all minimal independent (w.r.t. \mathcal{M}_1) (s, t)-cuts of size k in G_1 . The following claim is straightforward.

 \triangleright Claim 7. There is a natural bijection $f_1 : \mathcal{F}(s,t,k) \to \mathcal{F}_1(s,t,k)$ that can be computed in polynomial-time. In particular, $\mathcal{F}(s,t,k) \neq \emptyset \iff \mathcal{F}_1(s,t,k) \neq \emptyset$.

Proof. The proof is derived from the observation that the vertex set remains unchanged in both graphs G and G_1 . Therefore, any independent set in \mathcal{M} will also be an independent set in \mathcal{M}_1 , and vice versa (since $\mathcal{M} = \mathcal{M}_1$).

Let $\mathcal{F}_1(s,t,k) := \{X \subseteq V(G_1) \setminus \{s,t\} : X \in \mathcal{I}(\mathcal{M}_1), |X| = k, X \text{ is a directed } (s,t)\text{-cut}\}.$ It is straightforward to see that the task of finding $\mathcal{F}'(s,t,k) \subseteq_{\operatorname{rep}}^q \mathcal{F}(s,t,k)$ is equivalent to finding $\mathcal{F}'_1(s,t,k) \subseteq_{\operatorname{rep}}^q \mathcal{F}_1(s,t,k)$, using the bijection f_1 in Claim 7. Then, by using Lemma 15, we obtain the proof of Theorem 4.



Figure 2 Illustration of Phase I.

Phase II. Directed Vertex (s,t)-**cut** \Rightarrow **Directed Edge** (s,t)-**cut**. We are given an instance $(G_1, s, t, \mathcal{M}_1, Q_1, k)$ of GENERALIZED INDEPENDENT DIRECTED VERTEX (s,t)-CUT. We show how to reduce it to an instance of $(G_2, s, t, \mathcal{M}_2, Q_2, k)$ of GENERALIZED INDEPENDENT DIRECTED EDGE (s,t)-CUT. Note that here $Q_2 \subseteq A(G_2)$. Given a directed graph G_1 , we construct a directed graph G_2 as follows.

- For each vertex v in G_1 , we introduce two vertices v_{in} and v_{out} in G_2 . And add the arcs (v_{in}, v_{out}) to $A(G_2)$. We call them edges in G_2 corresponding to $V(G_1)$, or *v*-edges.
- For each arc (u, v) in $A(G_1)$, we add the arc (u_{out}, v_{in}) in $A(G_2)$. We call them edges in G_2 corresponding to $A(G_1)$, or *e-edges*
- The set Q_2 is the v-edges corresponding to the set $Q_1 \subseteq V(G)$.



Figure 3 Illustration of Phase II.

Matroid Transformation. Here we construct a matroid \mathcal{M}_2 for the instance $(G_2, s, t, \mathcal{M}_2, k)$. Assume that the in the input we are given a graph G_1 along with a matroid \mathcal{M}_1 . The ground set of \mathcal{M}_2 is equal to $A(G_2)$. Any edge subset that contains a e-edge is deemed to be dependent. Further, using the natural bijection between a vetex v in $V(G_1)$ and the corresponding v-edge in $V(G_2)$, we obtain a bijection between independent vertex-subsets in \mathcal{M}_1 , and make the corresponding v-edge-subsets independent in \mathcal{M}_2 . It is straightforward to obtain a representation of \mathcal{M}_2 , given a representation of \mathcal{M}_1 – we use the same vectors for v-edges, whereas other edges is assigned a zero vector.

Let $\mathcal{F}_1(s,t,k) \coloneqq \{S \subseteq A(G_1) : S \text{ is a minimal edge } (s,t)\text{-cut in } G_1, |S| = k, S \in \mathcal{I}(\mathcal{M}_1)\},\$ and $\mathcal{F}_2(s,t,k) \coloneqq \{S \subseteq V(G_2) : S \text{ is a minimal vertex } (s,t)\text{-cut in } G_2, |S| = k, S \in \mathcal{I}(\mathcal{M}_2)\}.\$ We prove the following claim.

 \triangleright Claim 8. There is a natural bijection $f_{12} : \mathcal{F}_1(s,t,k) \to \mathcal{F}_2(s,t,k)$ that can be computed in polynomial time. In particular, $\mathcal{F}_1(s,t,k) \neq \emptyset \iff \mathcal{F}_2(s,t,k) \neq \emptyset$.

Proof. In the forward direction, assume that $S \subseteq V(G_1)$ is a solution of size at most k. We show that the arcs in G_2 corresponding to the vertices in $S \subseteq G_1$, more specifically, the set of arcs $\{(v_{in}, v_{out}) : v \in S\}$ (denoted by A_S) of size at most k is a solution of size at most k. On the contrary, assume that there is a path P from s to t in $G_2 - A_S$. Now, every arc on the path P, of the form (v_{in}, v_{out}) (Type I) for some $v \in V(G_1)$ or of the form (u_{out}, v_{in}) (Type II) for some $u, v \in V(G_1)$. Now, the existence of arcs of Type I implies that there is an arc $(u, v) \in A(G_1)$. Note that any pair of consecutive arcs on the path P is of different type. With this we can find a path from s to t in $G_1 - S$. In the backward direction, assume that $Z \subseteq A(G_2)$ is a solution for G_2 . Now we find a set of vertex V_Z in G_1 in the following way: If there is an arc $(u_{out}, v_{in}) \in Z$ (Type I) for some $v \in V(G_1)$, we add v to V_Z , if there is an arc of the form $(u_{out}, v_{in}) \in Z$ (Type II) for some $u, v \in V(G_1)$, we add u in V_Z . We

show that $G_1 - V_Z$ does not have a path from s to t. On the contrary, assume that there is a path P from s to t in $G_1 - V_Z$. Now, corresponding to the path P in $G_1 - V_Z$, we can find a path from s to t in $G_2 - Z$ in the following way. For every arc (u, v) in P we have an arc (u_{out}, v_{in}) . Note that by our process, as the arc (u, v) is present in $G_1 - V_Z$, so both the arcs (u_{in}, u_{out}) and (u_{out}, v_{in}) are present in $G_2 - Z$. So for every arc (u, v) in P we have a path from u_{in} to v_{out} in $G_2 - Z$. With this we can find the path from s to t in $G_2 - Z$. A contradiction.

We now demonstrate that the solution maintains the independence property when transitioning from G_1 to G_2 , and similarly in the reverse direction. In the forward direction, assume that S is an independent (s, t)-cut of size at most k in G_1 . Notice that the arcs of G_2 corresponding to the vertices in S (denoted by A_S) are indeed a (s,t)-cut in G_2 . Now, according to our construction, the vector set corresponding to A_S in \mathcal{M}_2 is exactly the same as the vector set corresponding to S in \mathcal{M}_1 . Therefore, A_S eventually becomes independent in M_2 . In the reverse direction, assume that Z is an independent (s, t)-cut of size at most k in G_2 . Now Z is independent in \mathcal{M}_2 . This implies that Z does not contain any edge of the form (u_{out}, v_{in}) for some $u, v \in V(G_1)$ (since no independent set of vectors contains a zero vector and the vector corresponding to the edge (u_{out}, v_{in}) is a zero vector in \mathcal{M}_2). So, every edge Z is of the form (v_{in}, v_{out}) for some $v \in V(G_1)$. Notice that the vertices of G_1 corresponding to the arcs of Z (denoted by V_Z) are indeed a (s,t)-cut in G_2 . Now, according to our construction, the vector set corresponding to V_Z in \mathcal{M}_1 is exactly the same as the vector set corresponding to Z in \mathcal{M}_2 . Therefore, V_Z eventually becomes independent in \mathcal{M}_1 . Hence, the claim. \triangleleft

Algorithm. Given an instance $(G_1, s, t, \mathcal{M}_1, Q_1, k)$, we compute the instance $(G_2, s, t, \mathcal{M}_2, Q_2, k)$, as mentioned above. From Claim 8, the task of finding $\mathcal{F}'_1(s, t, k) \subseteq_{\mathrm{rep}}^{r-k} \mathcal{F}_1(s, t, k)$ is reduced to $\mathcal{F}'_2(s, t, k) \subseteq_{\mathrm{rep}}^{r-k} \mathcal{F}_2(s, t, k)$, for which we use the algorithm from Lemma 13. Thus, we obtain the following lemma.

▶ Lemma 9. There exists a deterministic algorithm that takes an input $(G_1, s, t, \mathcal{M}_1, Q_1, k)$ of GENERALIZED INDEPENDENT DIRECTED VERTEX (s, t)-CUT, runs in time $2^{\mathcal{O}(r)} \cdot 2^{\mathcal{O}(k^4)} \cdot n^{\mathcal{O}(1)}$, where $r = \operatorname{rank}(\mathcal{M}_1)$, and returns $\mathcal{F}'_1(s, t, k) \subseteq_{rep}^{r-k} \mathcal{F}_1(s, t, k)$ of size at most 2^r .

Phase III. Flow augmentation

▶ Definition 10 (Compatible augmenting set). Let $Z \in E(G)$ be a minimal (s,t)-cut in G. We say that $A \in V(G) \times V(G)$ is a compatible augmenting edge set for Z in G if (i) Zremains an edge (s,t)-cut in G + A, and (ii) Z is a minimum edge (s,t)-cut in G + A. Let $\mathcal{A} = \{A_1, A_2, \ldots\}$ is a collection of subsets of $V(G) \times V(G)$. If for $Z \in E(G)$, there exists some $A_i \in \mathcal{A}$ that is a compatible augmentating edge set for Z in G, then we use $A_{\mathcal{A}}(Z)$ to denote such an A_i – if there are multiple such A_i 's, then we pick arbitrarily; whereas if there is no such $A_i \in \mathcal{A}$, then we define $A_{\mathcal{A}}(Z) = \bot$. When \mathcal{A} is clear from the context, we will omit it from the subscript.

Assume that we have an instance G_2 for directed edge (s, t)-cut. Using Proposition 6, in time $2^{\mathcal{O}(k^4 \log k)} \cdot |V(G)|^{\mathcal{O}(1)}$ we will find a set $\mathcal{A} \subseteq 2^{V(G) \times V(G)}$ of compatible augmenting edge-sets with $\alpha := |\mathcal{A}| = 2^{\mathcal{O}(k^4 \log k)} \cdot (\log n)^{\mathcal{O}(k^3)}$. Now, from G_2 , we create α instances in the following way: for each $\mathcal{A} \in \mathcal{A}$, we create an instance $G_2^{\mathcal{A}} = G_2 + \mathcal{A}$.

 \triangleright Claim 11. For any $Z \subseteq E(G_2)$ of size $k' \leq k$, Z is an edge (s,t)-cut in G_2 if and only if there exists an $A_{\mathcal{A}} \in \mathcal{A}$ such that Z is a minimum edge (s,t)-cut of size k' in G_2^A .

Proof. The forward direction can be derived from the assertion made in Proposition 6. In the reverse direction, since G_2^A is created by adding some edges to G_2 , it is evident that any edge (s, t)-cut in G_2^A also remains an edge (s, t)-cut in G_2 .

Matroid Transformation. Now, fix some $A \in \mathcal{A}$ and we describe the matroid associated with the instance corresponding G_2^A , called \mathcal{M}^A . The ground set $U(\mathcal{M}_3^A)$ is equal to $U(\mathcal{M}_2) \cup A = E(G_2^A)$. The collection of independent sets in \mathcal{M}^A is same as that in \mathcal{M}_2 – any set $S \subseteq E(G_2^A)$ with $S \cap A \neq \emptyset$ is deemed dependent in \mathcal{M}_2^A . It is straightforward to verify that \mathcal{M}_2^A is a matroid. Given a representation for \mathcal{M}_2 , it is easy to come up with a representation for \mathcal{M}_2^A – we simply associate each edge in A with a zero vector. Let $\mathcal{F}_2^A(s,t,k) = \{X \subseteq E(G_2^A) : X \text{ is an independent minimum edge } (s,t)$ -cut in $G_2^A, |X| = k\}$. Now, from Claim 11 and the above discussion, we obtain the following corollary.

▶ **Observation 12.** For each $Z \in \mathcal{F}_2(s, t, k)$, $Z \in \mathcal{F}_2^A(s, t, k)$, where $A = A_A(Z)$.

Algorithm. For each $A \in \mathcal{A}$, we create an instance $(G_2^A, s, t, \mathcal{M}^A, Q^A, k)$, and compute $\mathcal{F}_2^{'A}(s, t, k) \subseteq_{\operatorname{rep}}^{r-k} \mathcal{F}_2^A(s, t, k)$ using the algorithm of Lemma 15. Then, we compute $\tilde{\mathcal{F}}_2(s, t, k) := \bigcup_{A \in \mathcal{A}} \mathcal{F}_2^{'A}(s, t, k)$, and then we compute $\mathcal{F}_2'(s, t, k) \subseteq_{\operatorname{rep}}^{r-k} \tilde{\mathcal{F}}_2(s, t, k)$. We show the following.

▶ Lemma 13. There exists a deterministic algorithm that takes input an instance $(G_2, s, t, \mathcal{M}_2, Q_2, k)$ of GENERALIZED INDEPENDENT DIRECTED EDGE (s, t)-CUT, runs in time $2^{\mathcal{O}(r)} \cdot 2^{\mathcal{O}(k^4)} \cdot n^{\mathcal{O}(1)}$, where $r = \operatorname{rank}(\mathcal{M}_2)$, and returns $\mathcal{F}'_2(s, t, k) \subseteq_{rep}^{r-k} \mathcal{F}_2(s, t, k)$ of size at most 2^r .

Proof. Consider any $Z \in \mathcal{F}_2(s,t,k)$, i.e., $Z \subseteq E(G_2)$ is a minimal independent edge (s,t)-cut of size k. From Observation 12, it follows that $Z \in \mathcal{F}_2^A(s,t,k)$, where $A = A_A(Z)$. Now consider any $Y \subseteq U(\mathcal{M}_2) \subseteq U(\mathcal{M}_3)$ such that Y fits Z. Then, there exists some $Z' \in \mathcal{F}_2'^A(s,t,k)$ such that Y fits Z'. Clearly, $Z' \subseteq E(G_2)$, since Z' is independent in \mathcal{M}_2^A , and Z' is of size k. Further, Z' is a minimum edge (s,t)-cut in G_2^A of size k. Then, by Claim 11, it follows that $Z' \in \mathcal{F}_2(s,t,k)$, which shows the lemma.

Phase IV. Directed Edge (s,t)-cut \Rightarrow **Directed Vertex** (s,t)-cut. We are given an instance $(G_3, s, t, \mathcal{M}_3, Q_3, k)$ of GENERALIZED INDEPENDENT DIRECTED EDGE (s,t)-CUT. We show how to reduce it to an instance of GENERALIZED INDEPENDENT DIRECTED VERTEX (s,t)-CUT. Given a directed graph G_3 , define a directed graph G_4 , and set $Q_4 \subseteq V(G_4)$ as follows.

- for each vertex u in $V(G_3)$, we add k + 1 vertices $u_1, u_2, \ldots, u_{k+1}$. We call them vertices in G_4 corresponding to $V(G_3)$, or *v*-vertices.
- For each arc $(u, v) \in A(G_3)$, we add a vertex e_{uv} in $V(G_4)$, and arcs (u_i, e_{uv}) and (e_{uv}, v_i) to $A(G_4)$ for every $i \in [k+1]$. We call them vertices in G_4 corresponding to $A(G_3)$, or *e-vertices*.
- The set Q_4 is the e-vertices corresponding to the set $Q_3 \subseteq A(G_3)$.

Matroid Transformation. Here we construct a matroid called \mathcal{M}_4 for the instance $(G_4, s, t, \mathcal{M}_4, k)$ using a similar approach to what we did in phase II. The ground set of \mathcal{M}_4 is equal to $V(G_4)$. Since $s, t \in V(G_4)$ cannot be part of any solution, we prevent them from being part of any independent set. Any vertex subset that contains a v-vertex is deemed to be dependent. Further, using the natural bijection between an edge in $A(G_3)$ and the corresponding



Figure 4 Illustration of Phase IV.

e-vertex in $V(G_4)$, we obtain a bijection between independent edge-subsets in \mathcal{M}_3 , and make the corresponding e-vertex-subsets independent in \mathcal{M}_4 . It is straightforward to obtain a representation of \mathcal{M}_4 , given a representation of \mathcal{M}_3 – we use the same vectors for e-vertices, whereas each v-vertex is assigned a zero vector.

Let $\mathcal{F}_3(s,t,k) \coloneqq \{S \subseteq E(G_3) : S \text{ is a minimal edge } (s,t)\text{-cut in } G_3, |S| = k, S \in \mathcal{I}(\mathcal{M}_3)\},\$ and $\mathcal{F}_4(s,t,k) \coloneqq \{S \subseteq V(G_4) : S \text{ is a minimal edge } (s,t)\text{-cut in } G_4, |S| = k, S \in \mathcal{I}(\mathcal{M}_4)\}.\$ We prove the following claim.

 \triangleright Claim 14. There is a natural bijection $f_{34} : \mathcal{F}_3(s,t,k) \to \mathcal{F}_4(s,t,k)$ that can be computed in polynomial time. In particular, $\mathcal{F}_3(s,t,k) \neq \emptyset \iff \mathcal{F}_4(s,t,k) \neq \emptyset$.

Proof. In the forward direction, assume that $Z \subseteq A(G_3)$ is a solution of size at most k. We show that the vertices in G_4 corresponding to the arcs in Z in G_3 , more specifically, the set of vertices $\{e_{uv}: (u, v) \in Z\}$ (denoted by V_Z) is a solution of size at most k in G_4 . On the contrary, assume that there is a path P from s to t in $G_4 - V_Z$. Now, every arc on the path P, either of the form (u_i, e_{uv}) or (e_{uv}, v_j) for some $u, v \in V(G_3)$ and $i, j \in [k+1]$. Now if the vertex e_{uv} is present in the path P, that means $(u, v) \notin Z$. Now we get a path from s to t in $G_3 - Z$ using the arcs $\{e_{uv} : e_{uv} \in V(P)\}$, which is a contradiction. In the backward direction, the proof is similar to the forward direction. Aassume that $S \subseteq V(G_4)$ is a solution for G_4 . Let there be a vertex u_i for some $i \in [k+1]$ such that $u_i \in S$. We claim that $S \setminus u_i$ remains a solution for G_4 . Note that since there are k+1 many vertices (copy) for each vertex $u \in G_3$, we must have $\{u_j : j \in [k+1]\} \setminus Z \neq \emptyset$. Let $u_j \in \{u_j : j \in [k+1]\} \setminus Z$. On the contrary, if $S \setminus u_i$ is not a solution, then there is a path from s to t in $G_4 - (S \setminus u_i)$. And that path must pass through the vertex u_i . Now, from this we can find a path from s to t in $G_4 - S$ just by passing the path replacing u_i by u_j (since both vertices have the same neighbor sets). A contradiction. So $S \setminus u_i$ remains a solution for G_4 . Repeating this argument, we can obtain that the set of all vertices in S that correspond to some arc in G_3 (denoted by S') remains a solution for G_4 . Now we show that the arcs in G_3 corresponding to the vertices in S' in G_4 , more specifically, the set of arcs $\{(u, v): e_{uv} \in S'\}$ (denoted by $A_{S'}$) is a solution of size at most k in G_3 .

We now demonstrate that the solution maintains the independence property when transitioning from G_1 to G_2 , and similarly in the reverse direction. Since the vectors for all the edges in G_3 remain the same for all the vertices in G_4 that correspond to those edges, any independent set in \mathcal{M}_3 will still be an independent set in \mathcal{M}_4 . On the contrary, since no independent set in a matroid contains a zero vector, any independent set in G_4 cannot include the vertex in G_4 that corresponds to a vertex in $V(G_3)$. Therefore, any independent set in \mathcal{M}_4 is also an independent set in \mathcal{M}_3 . This completes the proof.

Algorithm. We take the instance $(G_3, s, t, \mathcal{M}_3, Q_3, k)$ of GENERALIZED INDEPENDENT DIRECTED EDGE (s, t)-CUT and reduce it to an instance $(G_4, s, t, \mathcal{M}_4, Q_4, k)$ of GENERALIZED

INDEPENDENT DIRECTED VERTEX (s,t)-CUT. Then, we use Theorem 26 and compute $\mathcal{F}'_4(s,t,k) \subseteq_{\operatorname{rep}}^{r-k} \mathcal{F}_4(s,t,k)$ in time $2^{\mathcal{O}(r)} \cdot n^{\mathcal{O}(1)}$, such that $|\mathcal{F}'_4(s,t,k)|$. Then, by using the bijection-inverse f_{34}^{-1} from Claim 14 we obtain $\mathcal{F}'_3(s,t,k)$. We observe, due to the properties of the bijection f_{34} , it holds that $\mathcal{F}'_3(s,t,k) \subseteq_{\operatorname{rep}}^{r-k} \mathcal{F}_3(s,t,k)$. We summarize this in the following lemma.

▶ Lemma 15. There exists a deterministic algorithm that takes input an instance $(G_3, s, t, \mathcal{M}_3, Q_3, k)$, where $E(G_3) \subseteq U(\mathcal{M}_3)$ and $r = \operatorname{rank}(\mathcal{M}_3)$, and computes $\mathcal{F}'_3(s, t, k) \subseteq_{rep}^{r-k} \mathcal{F}_3(s, t, k)$ of size at most 2^r in time $2^{\mathcal{O}(r)} \cdot n^{\mathcal{O}(1)}$.

3.2 Algorithm For Directed Vertex (s, t)-cut

Assume that we are given a directed graph G = (V, E), two specified vertices s and t, a vertex set $Q \subseteq V(G)$ containing s, t, a matroid $\mathcal{M} = (U, \mathcal{I})$ such that $V \setminus Q \subseteq U$. In this section, path always refers to a directed path unless mentioned otherwise. We say that a vertex set $S \subseteq V(G) \setminus \{s, t\}$ is a (directed) (s, t)-cut in G if G - S does not contain any s to t path. Further, we say that an (s, t)-cut $S \subseteq V(G) \setminus \{s, t\}$ is an independent (s, t)-cut if $S \in \mathcal{I}$.

Similar to the previous sections, for $0 \le k \le r$, we define

 $\mathcal{F}(s,t,k) \coloneqq \{ X \subseteq V(G) \setminus \{s,t\} : X \text{ is an independent } (s,t) \text{-cut of size } k \}.$

Here is a formal description of the problem we consider.

- Generalized Independent Directed Vertex (s,t)-cut (GIDVC) -

Input: A directed graph G, a linear matroid $\mathcal{M} = (U, \mathcal{I})$, a set $Q \subseteq V(G)$ containing s, t, such that $V(G) \setminus Q \subseteq U$, and two non-negative integers k, q such that $k + q \leq r = \mathsf{rank}(\mathcal{M})$, and k is the size of minimum vertex (s, t)-cut. **Output:** Return $\mathcal{F}'(s, t, k) \subseteq_{\mathrm{rep}}^q \mathcal{F}(s, t, k)$ of bounded size.

In this section, we adopt the approach of Feige and Mahdian [8] to our end. Many of the definitions are directly borrowed from the same (albeit for directed graphs).

▶ Definition 16 (Critical and non-critical vertex). A vertex v of G is called *critical* if every collection of k vertex-disjoint paths from s to t contains v. A vertex is *non-critical* if it is not critical.

We now have the following fact, which directly follows from the above definition.

▶ **Proposition 17.** A vertex v is critical if and only if there is a directed vertex (s,t)-cut of size k containing v.

Definition 18 (Connecting pairs). For a pair of vertices u and v in G, we say that u is connected to v in G if there is a directed path from u to v such that every internal vertex of the path is non-critical.

Fix a collection of k vertex-disjoint paths P_1, P_2, \ldots, P_k from s to t. By definition, each critical vertex must be on one of these paths. In addition, Proposition 17 implies the following.

▶ **Observation 19.** For every $S \in \mathcal{F}(s,t,k)$, all $v \in S$ are critical vertices. Furthermore, distinct vertices of S belong to distinct paths from $\{P_1, P_2, \ldots, P_k\}$.

A. Banik, F. V. Fomin, P. A. Golovach, T. Inamdar, S. Jana, S. Saurabh

For each P_i , let $v_{i,1}, \ldots, v_{i,z_i}$ be the sequence of critical vertices of P_i in the order in which they appear on this path from s to t. Here, z_i denotes the number of critical vertices in P_i . To simplify notation, we define $v_{i,0} = s$ and $v_{i,z_i+1} = t$ for every i, and treat the vertices s and t as critical. Note that in the problem definition, we are not allowed to be chosen as solution vertices (i.e., cut vertices). Let Ω be the set of all k-tuples $\boldsymbol{a} = (a_1, \ldots, a_k)$ where $1 \leq a_i \leq z_i$ for every i. In other words, each $\boldsymbol{a} \in \Omega$ corresponds to one way of selecting a critical vertex from each P_i . Note that this does not have to correspond to an (s, t)-cut in G, since there are edges and vertices in G that are not in the paths P_i 's.

▶ Definition 20 (Prefix subgraph). The prefix subgraph G[a] defined by $a \in \Omega$ is an induced subgraph of G with the vertex set defined as follows:

- a critical vertex $v_{i,j}$ is in $G[\mathbf{a}]$ if and only if $j \leq a_i$;
- a non-critical vertex u is in G[a] if and only if all critical vertices that are connected to u are in G[a].

The last two layers of $G[\mathbf{a}]$ are the set of critical vertices $v_{i,j}$ such that $a_{i-1} \leq j \leq a_i$.

▶ Lemma 21 (Decomposition lemma, [8]). There is a sequence $a^1, \ldots, a^p \in \Omega$ such that

- (a) $a^1 = (1, ..., 1)$ and $a^p = (z_1, ..., z_k);$
- (b) for every h = 2, ..., p, $a^h a^{h-1}$ is a vector with exactly one entry equal to one, and zero elsewhere; and
- (c) for every h = 1, ..., p, every critical vertex of $G[\mathbf{a}^h]$, except possibly the vertices in the last two layers of $G[\mathbf{a}^h]$, is not connected to any critical vertex not in $G[\mathbf{a}^h]$.

Proof. We construct the sequence inductively. It is clear that $G[a^1]$ satisfies condition (c) above. Assume that we have constructed the sequence up to a^{h-1} . We show that there is some a^h that satisfies the conditions of the lemma. To this end, we show that there is an $i \in [k]$ such that $v_{i,a_i^{h-1}-1}$ is not connected to any critical vertex outside $G[a^{h-1}]$. Assume, for contradiction, that such an i does not exist. This means that for each i, there is a critical vertex $v_{j,\ell}$ outside $G[a^{h-1}]$ (that is, with $\ell > a_j^{h-1}$) such that $v_{i,a_i^{h-1}-1}$ is connected to $v_{j,\ell}$.

Now, we construct an auxiliary directed graph H with vertex set [k]. For each $i, j \in [k]$, there is a directed edge from i to j in H if $v_{i,a_i^{h-1}-1}$ is connected to a critical vertex $v_{j,\ell}$ on P_j with $\ell > a_j^{h-1}$. By our assumption, every vertex in H has outdegree at least one, and therefore H has a cycle. Consider the shortest cycle $C = i_0, \ldots, i_{f-1}$ in H. Each edge $i_b i_{(b+1) \mod f}$ of this cycle corresponds to a path Q_b from $v_{i_b, a_{i_i}^{h-1}-1}$ to $v_{i_{b+1}, \ell_{b+1}}$ for some $\ell_{b+1} > a_{i_{b+1}}^{h-1}$, such that all internal vertices of this path are non-critical. We show that the existence of this cycle contradicts the fact that $v_{i_0, a_{i_0}^{h-1}}$ is a critical vertex. As $v_{i_0, a_{i_0}^{h-1}}$ is a critical vertex, there exists a vertex st-cut S of size k that contains the vertex $v_{i_0, a_i^{h-1}}$. The removal of S divides the graph into several connected components. We call the vertices that can be reached from s on the graph G - S silver vertices, and the vertices from which we can reach t in G - S tan vertices. Since S contains exactly one vertex from each P_i and $\ell_0 > a_{i_0}^{h-1}$ the vertex v_{i_0,ℓ_0} must be a tan vertex. This vertex is connected by the path Q_{f-1} from the vertex $v_{i_{f-1}, a_{i_{f-1}}^{h-1}-1}$. Since all the vertices of Q_{f-1} are non-critical and hence not in S, the vertex $v_{i_{f-1}, a_{i_{f-1}}^{h-1}-1}$ is either tan or in S. Therefore, the vertex $v_{i_{f-1}, \ell_{f-1}}$ must be tan. Similarly, we can argue that the vertices $v_{i_{f-2},\ell_{f-2}},\ldots,v_{i_1,\ell_1}$ are all tan. However, v_{i_1,ℓ_1} is connected by the path Q_0 from $v_{i_0, \boldsymbol{a}_{i_0}^{h-1}-1}$, and the latter vertex must be silver, since the only vertex on P_{i_0} that is in S is $v_{i_0, \boldsymbol{a}_{i_0}^{h-1}}$. This gives us the desired contradiction.

The above argument shows that there is a vertex $v_{i,\boldsymbol{a}_{i}^{h-1}-1}$ that is not connected to any critical vertex outside $G[\boldsymbol{a}^{h-1}]$. Such a vertex can be found efficiently by trying all possibilities. Now, we simply let $\boldsymbol{a}_{i}^{h} = a_{i}^{h-1} + 1$ and $\boldsymbol{a}_{j}^{h} = \boldsymbol{a}_{j}^{h-1}$ for every $j \neq i$. It is easy to see that this choice of \boldsymbol{a}^{h} satisfies the conditions of the lemma.

▶ Remark 22. The generating sequence in the decomposition lemma maintains the invariant that at any step, every critical vertex outside the current subgraph is not connected from any critical vertex other than the ones in the last two layers of the current subgraph. In other words, the last two layers of the current prefix subgraph act as the *interface* of this subgraph to the rest of G.

Next, we define the notion of a valid coloring that essentially captures partial solutions that will be used to perform dynamic programming.

▶ Definition 23 (Valid coloring). A valid coloring of a prefix subgraph G[a] is a partial coloring of the vertices of G[a] with colors silver (s), tan (t), and black (b) such that

- (black vertices) for each i, there is at most one vertex of $G[\mathbf{a}]$ on P_i that is colored black; furthermore, s and t cannot be colored black;
- (silver and tan critical vertices) for each i and $j \leq a_i$, if there is no $j' \leq a_i$ such that $v_{i,j'}$ is colored black, then $v_{i,j}$ must be colored silver; if there is such a j', then $v_{i,j}$ must be colored silver if j < j' and tan if j > j';
- No silver-colored critical vertex is connected to a tan-colored critical vertex; and
- (coloring for non-critical vertices) every non-critical vertex that is connected from at least one silver critical vertex is colored silver; every non-critical vertex that is connected to at least one tan critical vertex is colored tan;
- (some vertex remains uncolored) every non-critical vertex that is neither connected from any silver critical vertex nor connected to any tan critical vertex remains uncolored.

We use col to denote the set of all possible valid colorings of G that color vertex t tan. We use S to denote all minimum vertex (s, t)-cuts in G.

▶ Remark 24. At a high level, the first condition in Definition 23 says that, each path contains one black vertex (must be critical by Observation 19). The second condition says, in each path the colors of critical vertices follow the pattern either "silver $< \cdots <$ silver < black $< \tan < \cdots < \tan$ ". The third condition in the above definition guarantees that no non-critical vertex is connected from a silver critical vertex as well as a tan critical vertex. The fourth and fifth conditions specifies a well-defined coloring for non-critical vertices. Notice that due to third condition, among two exactly at most one situation can occur for any non-critical vertex.



Figure 5 An illustration of a valid coloring on critical vertices.

We start with following lemma.

▶ Lemma 25. There is a bijection $f : col \to S$, i.e. between the set of all valid colorings of G that color vertex t tan, and the set of all minimum vertex (s,t)-cuts in G.

Proof. Let S denote the set of all minimum vertex (s, t)-cuts in G and col denote the set of all possible valid colorings of G that color vertex t tan. We show that there is a bijection $f: \operatorname{col} \to S$.

Consider a set $S \in S$. As |S| = k, this means that S intersects exactly one vertex on every path in $\{P_i : i \in [k]\}$. Now we provide a coloring in the following way. We color **s** for each vertex in S. For the remaining vertices, for each $i \in [k]$ if there is a vertex v in the path P_i such that v is critical and v lies on the subpath of P_i from s and the black vertex (which is the solution vertex), then we color **s** to the critical vertex. Similarly, for each $i \in [k]$ if there is a vertex v on the path P_i such that v is critical and v lies on the subpath of P_i from s and the black vertex (which is the solution vertex) and t then we color **t** to the critical vertex. We color the vertex t to **t**. Now there is no non-critical vertex, as in this case we get a (s, t)-path in G - S, a contradiction. Now we can give a coloring to some non-critical vertices with the following way. For every non-critical vertex that is connected from at least one silver critical vertex we color that vertex **s**; every non-critical vertex that is connected to at least one tan critical vertex we color that vertex **t**. Now as per the Definition 23, we have obtained a valid coloring.

Now consider a valid coloring c with t color to the vertex t. Let S' be the set defined by all the black vertices in the coloring. Now we use the property of valid coloring. In the first property, we know that each path contains at most one black vertex, that means $|S'| \leq k$. As the vertex t obtained the color s (as our consideration), and no silver-colored critical vertex is connected to a tan-colored critical vertex (third property), maintaining the the second property we have that each path contains exactly one black vertices, so |S'| = k. In a contrary assume that S' is not a vertex (s, t)-cut then there is a path P from s to t such that no vertex in V(P) is colored black. So, every critical vertex on the path is colored either sor t. As the vertex s and t gets the color s and t, respectively, there must be a path from the s colored critical vertex to the t colored critical vertex, which is a contradiction, as it violates the third property of valid coloring. Hence, the proof.

▶ **Theorem 26.** There exists a deterministic algorithm for GENERALIZED DIRECTED IN-DEPENDENT VERTEX (s,t)-CUT that runs in time $2^{\mathcal{O}(r)}$ time, and returns $\mathcal{F}'(s,t,k) \subseteq_{rep}^{r-k} \mathcal{F}(s,t,k)$ of size at most 2^r . Here, k denotes the size of the minimum (s,t)-cut and r denotes the rank of the matroid \mathcal{M} .

Proof. Note that by Lemma 25 any valid coloring of the graph G that colors the vertex t tan (t) corresponds to a (s,t)-cut in G. Conversely, any rank (s,t)-cut in G gives rise to a valid coloring of G that colors t tan. Recall that our goal is to efficiently compute $\mathcal{F}'(s,t,k) \subseteq_{\mathrm{rep}}^{r-k} \mathcal{F}(s,t,k)$ of bounded size. By Lemma 25, there is a bijection between set of all minimum vertex (s,t)-cuts in G and valid colorings with t is colored tan. Therefore, we design a dynamic programming-based algorithm along the decomposition lemma (Lemma 21) to compute "partial representative family" that represents (partial) valid coloring. We describe this DP below.

We use Lemma 21 to construct the sequence a^1, \ldots, a^p in polynomial time. Based on this sequence, we define a table A of dimension $p \times 3^{2k} \times k A$ as follows.

• The entry A[h, y, i] is indexed an integer $h \in [p]$, a string $y \in \{s, t, b\}^{2k}$, and an integer $i \in [k]$.

• Fix some $h \in [p], y \in \{\mathbf{s}, \mathbf{t}, \mathbf{b}\}^{2k}, i \in [k]$. We say that a valid coloring $\chi : V(G[\mathbf{a}^h]) \to \{\mathbf{s}, \mathbf{t}, \mathbf{b}\}$ respects the tuple (h, y, i) if it colors the vertices in the last two layers of $G[\mathbf{a}^h]$ according to y, and $|B_{\chi}| = i$, where $B_{\chi} \coloneqq \chi^{-1}(\mathbf{b})$. We define $\mathcal{S}(h, y, i)$ to be the set of all such subsets B_{χ} over all valid colorings χ of $G[\mathbf{a}^h]$ that respect (h, y, i). In the entry A(h, y, i), we will store $\widehat{\mathcal{S}} \subseteq_{\mathrm{rep}}^{r-i} \mathcal{S}(h, y, i)$.

The base entries $A[1, \cdot, \cdot]$ can be computed by inspection. We now give a procedure to compute A[h, y, i], based on the entries $A[h - 1, \cdot, \cdot]$. The last two layers of $G[a^h]$ differ from the last two layers of $G[a^{h-1}]$ in that one vertex (say v) was added and one vertex (say u) was removed. Let w denote the vertex which is in the last layer of $G[a^{h-1}]$ but in the second last layer of $G[a^h]$. Note that the vertex w always exists. Technically, if u = s, then s may still belong to the last two layers of $G[a^h]$, but the treatment of this case is only simpler than in the case $u \neq s$ and is omitted. We try all three possible colors for u. For each color, we first check if in combination with y it violates the condition for a valid coloring. If 'not, then we do the next step. We compute the color of all the vertices that are in $G[a^h]$ but not in $G[a^{h-1}]$ using the fourth condition in Definition 23. Note that by condition (c) of Lemma 21, the color of any such vertex can be uniquely specified. We perform a case analysis based on y(v) and define a family F that we will eventually store in A[h, y, i]. Let y' be a coloring of the vertices in the last two layers of $G[a^{h-1}]$. We say that y and y' are compatible with each other, if they agree on coloring of the vertices that are in common in their respective last two layers. We denote this by $y \propto y'$.

- (i) y(v) = b As y is a valid coloring we have y(u) = y(w) = s. In this case, if y' ∝ y then we must have y'(u) = y'(w) = s. Therefore, there is a unique coloring in the last two layers in the vertices of G[a^{h-1}] that is compatible with y. Let F = {Q ∪ {v}: Q ∈ A[h-1, y', i-1], y' ∝ y}. So here |F| = |A[h-1, y', i-1]| ≤ 2^r.
- (ii) y(v) = s As y is a valid coloring, we have y(u) = y(w) = s. In this case, if y' ∝ y then we must have y'(u) = y'(w) = s. Therefore, there is a unique coloring in the last two layers in the vertices of G[a^{h-1}] that is compatible with y. Let F = A[h − 1, y', i]. So, here |F| = |A[h − 1, y', i]| ≤ 2^r.
- (iii) y(v) = t As y is a valid coloring, we have either y(u) = y(w) = t, or y(u) = b, y(w) = t, or y(u) = s, y(w) = b. In this case, if $y' \propto y$ then we must have either y'(u) = y'(w) = t, or y'(u) = b, y'(w) = t, or y'(u) = s, y'(w) = b. Here we distinguish these by two subcases based on y'(w). Recall that y(w) = y'(w).
 - If y'(w) = b then we must have y'(u) = s. So in this case, there is a unique coloring in the last two layers in the vertices of $G[\mathbf{a}^{h-1}]$ which is compatible with y. Let F = A[h-1, y', i]. So here $|F| = |A[h-1, y', i]| \le 2^r$.
 - If y'(w) = t then u can take color either black or tan. So there are at most two different colorings in the last two layers in the vertices of $G[\mathbf{a}^{h-1}]$ which is compatible with y. Let $F' = \bigcup_{y' \propto y} A[h-1, y', i]$. So here $|F'| = |\bigcup_{y' \propto y} A[h-1, y', i]| \leq 2 \cdot 2^r$. Now we use the tool of *representative sets* (Proposition 2) to find a set $F \subseteq_{\text{rep}}^{r-i} F'$ of size at most 2^r .

Finally, we will let $A[h, y, i] \leftarrow F$. Let \mathcal{Y} denote the set of all strings that color the vertex t tan in $G[\mathbf{a}^p]$. Note that $|\mathcal{Y}| \leq 2^k$. Given the table A, one can easily check the existence of a desired (s, t)-cut by checking the entries A[p, y, k] for all strings $y \in \mathcal{Y}$. Formally if $A[p, y, k] \neq \emptyset$ for some string y where y(t) = t then we return YES to our problem on the

instance G. For the more general problem, when we seek the representative family, we compute $\mathcal{F}' \subseteq_{\mathrm{rep}}^{r-k} \bigcup_{u \in \mathcal{V}} A[p, y, k]$ and return \mathcal{F}' .

▶ **Theorem 4.** Consider an instance $(G, \mathcal{M}, s, t, Q, k)$ of GENERALIZED INDEPENDENT VERTEX (s,t)-CUT, where there exists no independent (s,t)-cut in G of size less than k. Then, there exists a deterministic algorithm for GENERALIZED INDEPENDENT VERTEX (s,t)-CUT that runs in time $2^{\mathcal{O}(k^4 \log k)} \cdot 2^{\mathcal{O}(r)} \cdot n^{\mathcal{O}(1)}$, and outputs $\mathcal{F}'(s,t,k) \subseteq_{rep}^{r-k} \mathcal{F}(s,t,k)$ such that $|\mathcal{F}'(s,t,k)| \leq 2^r$. Here, $\mathcal{F}(s,t,k)$ is the collection of all independent (s,t)-cuts of size k.

4 (Odd) Cycle Hitting

In this section, we consider matroidal generalizations of two quintessential cycle hitting problems in parameterized complexity, namely FEEDBACK VERTEX SET (FVS), and ODD CYCLE TRANSVERSAL (OCT), respectively.

4.1 INDEPENDENT FVS

Here, we sketch an argument that gives a polynomial kernel for INDEPENDENT FVS parameterized by the solution size k, which we can assume to be equal to the rank r of the matroid, even in the oracle access model. However, we assume that the memory used to store oracles does not contribute to the input size, and this is important for our kernelization result. The standard degree 0 and 1 rules are also applicable here. Note that in standard FVS, we can replace an internal vertex v from an induced degree-2 path P by a vertex u of degree at least 3 that is at an end of P, since any cycle that passes through v, also passes through u. However, in the presence of matroid constraint, such an argument no longer works – since replacing v by u in a solution may not result in an independent set. Nevertheless, we can bound the relevant vertices from a long induced path of degree 2 by r + 1, using the following proposition.

▶ Lemma 27. Let \mathcal{M} be a matroid of rank r given via oracle access, and let \mathcal{A} be a 1-family of subsets of $U(\mathcal{M})$. Then, $\mathcal{A}' \subseteq_{rep}^{r-1} \mathcal{A}$ can be computed in polynomial time, where $|\mathcal{A}'| \leq r$.

Proof. Let $A = \{x : \{x\} \in \mathcal{A}\}$ be the set of underlying elements corresponding to the sets of \mathcal{A} . We compute an inclusion-wise maximal subset $R \subseteq A$ that is independent in \mathcal{M} . Note that this subset can be computed using O(|U|r) queries, where $r = \operatorname{rank}(\mathcal{M})$. We claim that $\mathcal{A}' = \{\{y\} : y \in R\}$ is a 1-representative set of \mathcal{A} .

Consider any $u \in A$ and $X \subseteq U$ of size r such that $u \in X$ and $X \in \mathcal{I}$. We will show that there exists some $u' \notin R$ such that $X \setminus \{u\} \cup \{u'\} \in \mathcal{I}$. Since R is an inclusion-wise maximal independent subset of A, it follows that $u \in \mathsf{cl}(R) \subseteq \mathsf{cl}(R \cup (X \setminus \{u\}))$. This means that $\mathsf{rank}(R \cup (X \setminus \{u\})) = \mathsf{rank}(R \cup (X \setminus \{u\})) = \mathsf{rank}(R \cup X)$. However, $\mathsf{rank}(R \cup X) \ge \mathsf{rank}(X)$. Therefore, we obtain that $\mathsf{rank}(R \cup (X \setminus \{u\})) \ge \mathsf{rank}(X)$. This implies that there exists some $u' \in R$ such that $X \setminus \{u\} \cup \{u'\} \in \mathcal{I}$.

Polynomial kernel. Consider any set P that induces a maximal induced degree 2 path in G. We replace P with a degree-2 path induced by P', where P' is obtained via Lemma 27. After performing each such replacement, we obtain a graph G', and let $V' \subseteq V(G')$ be the set of vertices of degree at least 3. Now consider a graph H' that is obtained by G' by replacing each maximal induced degree-2 path in G' by a single edge, and we call such an edge a *special edge*. Note that H' is exactly the graph we would obtain by using standard

polynomial kernelization algorithm on G. By standard arguments, we know that the number of vertices and edges in H' of degree at least 3 is bounded by $\mathcal{O}(r^2)$. Since each edge in H' may correspond to a path of length k induced by degree-2 vertices, it follows that the number of vertices and edges in G' is bounded by $\mathcal{O}(r^3)$.

FPT algorithm. Brute-forcing on the $\mathcal{O}(r^3)$ kernel immediately shows that INDEPENDENT FVS is FPT parameterized w.r.t. r with running time $r^{\mathcal{O}(r)} + n^{\mathcal{O}(1)}$ – note that this also requires only an oracle access to the matroid. However, we can design a faster, single exponential FPT algorithm for the problem via adapting a couple of approaches for standard FVS.

We first use the kernelization algorithm and obtain an equivalent instance of size $\mathcal{O}(r^3)$. In fact, instead of working with the graph G', we work with the graph H' where the number of vertices and edges is $\mathcal{O}(r^2)$, where we denote the subset of special edges in E(H) by E_{sp} . For simplicity, we use G and H instead of G' and H'. Now we use the iterative compression technique, where we are working with an induced subgraph H[X], for some $X \subseteq V(H)$. Note here that X is a subset of vertices of degree at least 3 in G. Further, let $Z := V(G) \setminus V(H)$ denote the set of vertices of degree 2 in G, and for $X \subseteq V(H)$, let $Z_X \subseteq Z$ denote the subset of vertices that appear on a path P connecting two vertices $u, v \in X$. Note that such a path is represented by a special edge $uv \in E(H)$, and indeed the special edge uv is present in H[X].

Inductively we assume that we have an FVS $S \subseteq X$ of size r + 1 in the graph H[X], and we want to determine whether there exists a vertex-subset $O \subseteq X$, and $F \subseteq E(H[X]) \cap E_{sp}$ of special edges, such that: (i) $|O \cup F| \leq r$, and (ii) for each edge $e \in F$, there exists a degree-2 vertex v_e in the corresponding degree-2 path, such that $O \cup \bigcup_{e \in F} v_e \in \mathcal{I}$.

We try to guess $Y := O \cap S$, assuming such an O indeed exists. In this step, in addition to deleting Y from the graph, we additionally contract the matroid \mathcal{M} on the set Y of size $p \leq r$, and obtain another matroid \mathcal{M}/Y . Since \mathcal{M} is provided to us using its representation matrix, we can obtain that of \mathcal{M}_Y in polynomial-time (see, e.g., [27]). Thus, we can reduce the problem to DISJOINT INDEPENDENT FVS, where we have an FVS $N := S \setminus Y$ for the graph H[W] = G - Y, and we want to find an independent FVS of size t := r - p in G' that is disjoint from N. For this problem, we can emulate a standard branching algorithm, albeit with some modifications. In the course of the algorithm, we also maintain a set $\mathcal{E} \subseteq E_S$ of special edges, which is initialized to an empty set. Let $W = X \setminus N$, and note that H[W] is a forest.

First, if there exists a vertex $v \in W$ such that v has two special neighbors $u_1, u_2 \in N$ (by this, we mean that u_1v, u_2v are special edges) such that u_1, u_2 belong to the same connected component in H[N], then we make three recursive calls, say 1, 2 and 3: In the recursive call $i \in \{1, 2\}$, the graph is obtained by deleting the special edge u_iv and adding it to \mathcal{E} . In the third recursive call, we add v to the solution, contract the matroid \mathcal{M} on v, and remove vfrom the graph. In each of the three recursive calls, the parameter t drops by 1. If a recursive call 1 or 2 returns a solution, then we return any of those. Otherwise, if call 3 returns a solution S, then we return $S \cup \{v\}$. If none of the calls returns a solution, then we return \perp .

Now assume that there is no such vertex $v \in W$ with two special neighbors $u_1, u_2 \in N$ from the same connected component of H[N]. In this case, we proceed with the same kind of branching strategy as for the standard FVS. We sketch it for completeness. Since H[W] is a forest, it contains some leaf x. However, since each vertex has degree at least 3 in H, it follows that x must have at least two neighbors in N. In addition, these neighbors must be in different components. We make two recursive calls. In the first recursive call, we add x **Algorithm 1** DisjointFVS $(H = (X, E), \mathcal{M}, W, N, \mathcal{E}, t)$ 1: **if** t = 0 **then** 2: if H contains a cycle then return \perp 3: else Find common base (if exists) for \mathcal{M} and \mathcal{M}_1 using [3], where 4: $\mathcal{M}_1 = (V(G), \mathcal{I}_1)$ where $\mathcal{I}_1 = \{S \subseteq V(G) | S \cap P_e | \le 1 : e \in \mathcal{E}\}$ 5:**return** common base S if it exists, else \perp 6: 7: if $\exists v \in W$ with neighbors $u_1, u_2 \in N$ in the same component of H[N] then if $u_1v \in E_s$ then $S_1 \leftarrow \texttt{DisjointFVS}(H, \mathcal{M}, W, N, \mathcal{E} \cup \{u_1v\}, t-1)$ else $S_1 \leftarrow \bot$ 8: if $u_2v \in E_s$ then $S_2 \leftarrow \texttt{DisjointFVS}(H, \mathcal{M}, W, N, \mathcal{E} \cup \{u_2v\}, t-1)$ else $S_2 \leftarrow \bot$ 9: $S_3 \leftarrow \texttt{DisjointFVS}(H-v, \mathcal{M}/v, W-v, N, t-1)$ 10: if $S_1 \neq \bot$ then return S_1 11: else if $S_2 \neq \bot$ then return S_2 12:else if $S_3 \neq \bot$ then return $S_3 \cup \{v\}$ 13:else return \perp 14:15: else Let $x \in W$ be a leaf in H[W]16: $S_4 \leftarrow \texttt{DisjointFVS}(H-x, \mathcal{M}/x, W-x, N, t-1)$ 17: $S_5 \leftarrow \texttt{DisjointFVS}(H, \mathcal{M}, W - x, N \cup \{x\}, t - 1)$ 18:19:if $S_4 \neq \bot$ then return $S_4 \cup \{x\}$ else if $S_5 \neq \bot$ then return S_5 20:21:else return \perp

to the solution, contract \mathcal{M} on x, and delete it from the graph. In this call, the parameter t is reduced by 1. If this call returns a solution S, then we return $S \cup \{x\}$. In the second recursive call, we add x to W, and the parameter t remains unchanged. If this call returns a solution, then we return the same solution. Otherwise, we return \perp .

At the bottom of the recursion, when t = 0, if the graph G' is not acyclic, then we return \bot . Otherwise, we look at the special edges that have been added to \mathcal{E} over the course of the recursion. Recall that these edges correspond to the induced degree-2 paths in the original graph. Thus, we have a YES-instance if and only if for each $e \in \mathcal{E}$, we can select some $v_e \in P_e$, where P_e is the set of vertices along the induced degree-2 path corresponding to e, such that $\bigcup_{e \in \mathcal{E}} v_e \in \mathcal{I}$. At this point, we observe that this can be reduced to a matroid intersection problem between the current matroid $\mathcal{M} = (V(G'), \mathcal{I}')$ (where $V(G') \subseteq V(G)$ – here G refers to the original kernelized graph) and a partition matroid $\mathcal{M}_1 = (V(G), \mathcal{I}_1)$, where \mathcal{M}_1 is a matroid of rank $|\mathcal{E}|$, such that a subset $S \subseteq V(G)$ is independent if and only if $|S \cap P_e| \leq 1$. This problem can be solved in polynomial time even using oracle access to \mathcal{M} using, e.g., the algorithm of Cunningham [3] (note that \mathcal{M}_1 is a partition matroid, and thus oracle queries can be easily simulated in polynomial time). It is also straightforward to show that this leads to an algorithm of running time $9^t \cdot n^{\mathcal{O}(1)}$. Due to the iterative compression step, we obtain an overall running time of $10^r \cdot n^{\mathcal{O}(1)}$. We conclude with the following theorem, by recalling that the solution size $k \leq r$.

▶ **Theorem 28.** INDEPENDENT FEEDBACK VERTEX SET admits a kernel of $\mathcal{O}(k^3)$ vertices and edges that can be computed in polynomial time in the oracle access model, where k denotes the solution size, with $k \leq r = \operatorname{rank}(\mathcal{M})$. Further, the problem admits a deterministic algorithm running in time $10^k \cdot n^{\mathcal{O}(1)}$ in the oracle access model.

4.2 INDEPENDENT OCT

This is a standard application of the iterative compression technique. Suppose we have an OCT S of size k + 1 of a graph G, and we want to determine whether G has an OCT O that is independent in \mathcal{M} . We try to guess $Y \coloneqq O \cap S$, assuming such an O indeed exists. In this step, in addition to deleting Y from the graph, we additionally contract the matroid \mathcal{M} on the set Y of size $p \leq k$, and obtain another matroid \mathcal{M}/Y . Since \mathcal{M} is provided to us using its representation matrix, we can obtain that of \mathcal{M}_Y in polynomial-time (see, e.g., [27]). Thus, we can reduce the problem to DISJOINT INDEPENDENT ODD CYCLE TRANSVERSAL, where we have an OCT $N \coloneqq S \setminus Y$ for the graph G' = G - Y, and we want to find an independent OCT of size k - p in G' that is disjoint from N. Then, by standard techniques (see, e.g., [32] or Lemma 4.14 of [5]), this problem can be reduced to solving 2^{k-p} instances of INDEPENDENT (s, t)-CUT in G', where the underlying matroid is now \mathcal{M}_Y . Then, by using Theorem 4, we obtain the following theorem.

▶ **Theorem 29.** INDEPENDENT ODD CYCLE TRANSVERSAL is FPT parameterized by k, which denotes the rank of the given linear matroid. Specifically, it admits a $2^{\mathcal{O}(k^4 \log k)} \cdot n^{\mathcal{O}(1)}$ time algorithm.

5 Vertex Multiway Cut

Like INDEPENDENT VERTEX (s, t)-CUT, our aim here is also to design an algorithm that solves a more general version of the problem, called GENERALIZED INDEPENDENT MULTIWAY CUT, or GIMC for short. By iteratively checking values k = 1, 2, ..., and running the algorithm described below, we may inductively assume that we are working with a value $0 \le k \le r$, such that there is no independent multiway cut for T in G for any k' < k. This follows from the fact that, our generalized problem, as defined below, requires the algorithm to return a set corresponding to k' whose emptiness is equivalent (via Proposition 3, item 2) to concluding that the there is no independent multiway cut of size k'.

Then, let $\mathcal{F}(T, k, G) \coloneqq \{S \subseteq V(G) \setminus T : |S| = k, S \in \mathcal{I}, \text{ and } S \text{ is a multiway cut for } T\}$. Note that, if there is no independent multiway cut of size k, then $\mathcal{F}(T, k, G) = \emptyset$. On the other hand, if $\mathcal{F}(T, k, G) \neq \emptyset$, then each $S \in \mathcal{F}(T, k, G)$ must also be a *minimal* multiway cut for G – otherwise, there would exist some $S' \subset S$ of size k' < k, and due to matroid properties $S' \in \mathcal{I}$. Therefore, k would not be the smallest such value. Next, we give a formal definition of the problem.

- Generalized Independent Multiway Cut(GIMC) -

Input: An instance $(G, \mathcal{M}, T, Q, k)$, where: = G = (V, E) is a graph, $= T \subseteq V(G) \text{ is a set of terminals}, Q \subseteq V(G) \text{ is a set of special vertices, with}$ $T \subseteq Q,$ $= \text{Matroid } \mathcal{M} = (U, \mathcal{I}) \text{ of rank } r, \text{ where } V(G) \setminus Q \subseteq U,$ $= 0 \leq k \leq r$ **Output:** $\mathcal{F}'(T, k, G) \subseteq_{\text{rep}}^{r-k} \mathcal{F}(T, k, G).$

First, in Section 5.1, we analyze certain structural properties of solutions of GIMC, and use it to design an FPT algorithm to find a so-called *strong separator* for the instance, or to conclude that we have a No-instance. Then, in Section 5.2, we use such a strong separator,

and use to design a recursive algorithm for GIMC, and then show its correctness and that it is FPT in k.

5.1 Finding a "Strong Separator" for GIMC

In this subsection, we define the notion of a "strong separator" and how it interacts with any minimal multiway cut. Let G be a graph and $T \subseteq V(G)$ be a set of terminals. Throughout the section, we will assume that G is connected. Let $S \subseteq V(G)$ be a minimal multiway cut for T, and let C_S denote the set of connected components of G - S. If a connected component $C \in C_S$ contains a terminal vertex then we call it a *terminal component*; otherwise, we call it a *non-terminal component*. Let \mathcal{T}_S denote the terminal components and \mathcal{N}_S denote the non-terminal components in G - S. Note that $\mathcal{C}_S = \mathcal{T}_S \uplus \mathcal{N}_S$. We say that a $v \in S$ is adjacent to a component $C \in \mathcal{C}_S$ if there exists some $u \in C$ such that uv is an edge. For each $v \in S$, let $\mathcal{T}(v, S) \subseteq \mathcal{T}$ denote the set of terminal components that are adjacent to v. If the set S is obvious from the context, then we denote $\mathcal{T}(v, S)$ by \mathcal{T}_v . Given two components C_1 and C_2 we say C_1 is adjacent to C_2 (and, vice versa) if there is a vertex in C_1 which has a neighbour in C_2 . Because such a multiway cut S (with certain additional properties, as stated below) will play a central role while designing the algorithm, we will often refer to it as a *separator*, which is essentially a synonym for *multiway cut for* T. We now define the notion of good and bad configuration corresponding to a minimal separator S.

▶ Definition 30 (Good and bad configurations). Let G be a graph, $T \subseteq V(G)$ be a terminal set with $|T| \geq 3$, and S be a minimal multiway cut S for T. We refer to $\langle G, T, S \rangle$ as a configuration. If there is a component C in G - S that is adjacent to each connected component in G[S], then we say that C is a large component. We say $\langle G, T, S \rangle$ is a bad configuration, if (i) each component in G[S] is adjacent to exactly two terminal components in G - S, and (ii) there is a large component ⁴. Otherwise, we say that $\langle G, T, S \rangle$ is a good configuration.

The definition of good and bad components easily lends itself to design a straightforward polynomial-time algorithm to check whether, corresponding to a given separator S, whether $\langle G, T, S \rangle$ is a good or a bad configuration. We state this in the following observation.

▶ **Observation 31.** There exists a polynomial-time algorithm that takes input a graph G, and a set of terminals $T \subseteq V(G)$, and a minimal multiway cut $S \subseteq V(G) \setminus T$, and outputs whether $\langle G, T, S \rangle$ is a good or a bad configuration.

In the following claims, we prove an important property of a good and a bad configuration, respectively.

 \triangleright Claim 32. If $\langle G, T, S \rangle$ is a good configuration, then there is no multiway cut that is fully contained in a terminal component in G - S.

Proof. For a contradiction, assume that there is a multiway cut S' and a terminal component C in G - S such that $S' \subseteq V(C)$. We know that each component in G[S] is adjacent to at least two terminal components. As $\langle G, T, S \rangle$ is a good configuration, that means that either (i) there is a component C_S in G[S] such that C_S is adjacent to at least three terminal components, or (ii) each component in G[S] is adjacent to exactly two terminal components,

⁴ Observe that, since $|T| \ge 3$, in a bad configuration, there is exactly one large component.



Figure 6 Examples of good (left) and bad (right) configurations. The blue and red vertices denote the separator and terminal vertices, respectively.

but there is no large component. We analyze each case separately, and show that we arrive at a contradiction.

In case (i), let C_1, C_2, C_3 be such three components in G - S that are adjacent to the vertices of C_S . Among C_1, C_2, C_3 , at most one can be C – hence at least two of them, say C_1 , and C_2 must have empty intersection with the set S'. However, since $S' \subseteq C$, it follows that $S' \cap S = \emptyset$. This implies that, in G - S', there is a path between the two terminals contained inside the components C_1 and C_2 , respectively. This is contradiction to the fact that S' is a multiway cut for T. In case (ii), we know that each component in G[S] is adjacent to exactly two terminal components, but there is no large component. Recall that $S' \subseteq V(C)$ for some terminal component C. By the case assumption C is not a large component, which implies that there is a component C'_S in G[S] such that C_1, C_2 , and none of them is C. As $S' \subseteq V(C)$ and $S \cap S' \neq \emptyset$, it follows that, in G - S', there is path between two terminals contained in C_1, C_2 , respectively, which again contradicts that S' is a multiway cut for T. Hence, the claim follows.

 \triangleright Claim 33. If $\langle G, T, S \rangle$ is a bad configuration then, except possibly the large component, no other terminal component contain a multiway cut.

Proof. Assume that $\langle G, T, S \rangle$ is a bad configuration with C as large component. Suppose for the contradiction, assume that some C_1 , distinct from C is a terminal component that contains a multiway cut, say S'. Let C_2 be a terminal component (arbitrary choosen) other than C and C_1 (note that such a component must exist since $|T| \ge 3$). As C is adjacent to each component in G[S], hence in G - S', there is path between the two terminals that are contained in C and C_2 , respectively, which contradicts that S' is a multiway cut. Hence, there is no multiway cut in G - S that is fully contained in a terminal component that is not large.

Next, we define the notion of type 1 and type 2 scenarios corresponding to a minimal separator S.

▶ Definition 34 (type 1 and type 2 scenarios). Given a graph G, a terminal set $T \subseteq V(G)$, and a minimal multiway cut S, we say the separator S creates a type 1 scenario when no terminal component in G - S can contain a multiway cut of size at most |S|. Otherwise, we say it creates a type 2 scenario.

 \triangleright Claim 35. Given a graph G, a terminal set $T \subseteq V(G)$, and a minimal multiway cut S of size k, in $2^k \cdot n^{\mathcal{O}(1)}$ time we can check whether S creates an type 1 scenario or not.



Figure 7 Example of a type 1 (left) and type 2 (right) scenario. Blue and red vertices denote separator, and terminal vertices, respectively.

Proof. If $\langle G, T, S \rangle$ is a good configuration, then by Claim 32 we know that there is no multiway cut that is fully contained in a terminal component in G - S. Hence, in this case S creates type 1 scenario. Now, assume that $\langle G, T, S \rangle$ is a bad configuration. Now by Claim 33 we see that except possibly the large component no other terminal component contains a multiway cut. Now to check whether large component contain a multiway cut we do the following. Let C denote the large component. We take the subgraph $G_1 := G[S \cup V(C)]$. Let S_1, S_2, \ldots, S_q denote the partition of S such that each S_i corresponds to a connected component of G[S]. In the subgraph G_1 , we contract each S_i , $1 \le i \le q$ into a single vertex s_i . Let G_2 be the new graph after contraction. Let $T' = \{s_1, \ldots, s_q, t\}$. Note that $q := |S'| \le |S| \le k$. Now we check whether there exists a multiway cut of size at most k in the graph G_2 with terminal set T'. If NO then we conclude that S creates a type 1 scenario in G for the terminal set T. Otherwise, we have type 2 scenario. The correctness of the algorithm follows from the observation that G has a multiway cut of size k that is fully contained in C, if and only if G_2 has multiway cut of size at most k with terminal set T'.

Now we define a notion of a special type of multiway cut which we refer as strong separator which we use later in our algorithmic purpose.

▶ Definition 36 (Strong separator). Given a configuration $\langle G, T, S \rangle$, we say S is strong separator for T in G if (i) $\langle G, T, S \rangle$ is a good configuration, or (ii) $\langle G, T, S \rangle$ is a bad configuration, but S creates a type 1 scenario.

Now we prove an important result that leads to the design of a branching algorithm.

▶ Lemma 37. Let $(G, \mathcal{M}, T, Q, k)$ be an instance of GIMC with $k \ge 2$, $|T| \ge 3$, and let $S \subseteq V$ be a strong separator of size at most k. There exists a polynomial-time algorithm to find a pair of terminal components in G - S such that for any solution O, we either have $S \cap O \neq \emptyset$ or one of the components must contain at least one and at most k - 1 vertices of O.

Proof. Assume that $(G, \mathcal{M}, T, Q, k, q)$ be an instance of INDEPENDENT MULTIWAY CUT and $S \subseteq V$ be a strong separator of size at most k. Due to the minimality of S, for each vertex $v \in S$ the set $S \setminus \{v\}$ is not a separator. Fix an arbitrary $v \in S$. It follows that, there exists a pair of terminals t_1, t_2 , contained in terminal components T_1, T_2 respectively, such that we have a path P connecting t_1 and t_2 in G such that the path intersects exactly once which is exactly at the vertex v. Furthermore, $V(P) \subseteq V(T_1) \cup V(T_2) \cup \{v\}$. Since O is also a multiway cut for $T, O \cap V(P) \neq \emptyset$, which implies that $O \cap (V(T_1) \cup V(T_2) \cup \{v\}) \neq \emptyset$. If $v \in O$ then we are done, as in this case $S \cap O \neq \emptyset$. Otherwise, $O \subseteq V(T_1 \cup T_2)$. However, since S is strong

separator, we know that $O \not\subseteq V(T_1)$, and $O \not\subseteq V(T_2)$. Hence, either $1 \leq |V(T_1) \cap O| \leq k-1$ or $1 \leq |V(T_2) \cap O| \leq k-1$. This completes the proof.

5.1.0.1 Finding a Strong Separator

In this section, we show that given a graph G a terminal set T, an integer k how to find a strong separator of size at most t in $2^k \cdot n^{\mathcal{O}(1)}$ time. First, we use a known FPT algorithm of Cygan et al. [7]—that runs in time $2^k \cdot n^{\mathcal{O}(1)}$ —to check whether G has a minimal multiway cut of size at most k. Either this algorithm returns that there is no multiway cut of size k, or it returns a (w.l.o.g. minimal) multiway cut $S \subseteq V(G)$ of size at most k. In the former case, since there is no multiway cut of size k, there is no strong separator of size k either – hence we return that G does not have a strong separator. Otherwise, in the latter case, we proceed to find a strong separator using the minimal separator S returned by the algorithm, as described below. First, using Observation 31, we can check in polynomial-time whether $\langle G, T, S \rangle$ is a good or a bad configuration, and proceed to either of the following two cases.

- Case A. ($\langle G, T, S \rangle$ is a good configuration) In this case, we are done as per, that is, S is a strong separator, as per our Definition 36.
- Case B. ($\langle G, T, S \rangle$ is a bad configuration) In this case, we can use Claim 35 to check in time $2^k \cdot n^{\mathcal{O}(1)}$ whether S creates a type 1 or a type 2 scenario. In the former case, S is a strong separator, and we are done. Thus, we are left with the case when S creates a type 2 scenario, i.e., each component in G[S] is adjacent to exactly two terminal components in G-S, and there is a large terminal component in G-S, say C_L . Assume that the component C_L contains the terminal t_L . Note that via If type 1 then we say that S is strong separator. Else We try to find another separator of size k which is fully contained in the large component using following procedure. Let $G_1 \coloneqq G - C_L$ and $G_2 \coloneqq G[S \cup C_L]$. Now consider the graph G_2 . For a terminal $t \in T \setminus \{t_L\}$, let $S_t \subseteq S$ denote the set of vertices in S that are reachable from t in the graph G_1 . Now for each terminal vertex $t \in T \setminus \{t_L\}$ we identify all the vertices in S_t to obtain a single merged vertex v_t . The merging operation preserves adjacencies towards the terminal components, i.e., there was a vertex $u \in V(G_2)$ such that u has a neighbour in S_t if and only if u and v_t are neighbors in the resulting graph. Consider a modified graph, say H, with vertex set $\{v_t: t \in T \setminus \{t_L\}\} \cup V(C(t_L))$. We then find another minimal multiway cut of size |S|, with the new terminal set $T' := \{t_L\} \cup \{v_t : t \in T \setminus \{t_L\}\}$. The outcome is one of the following two possibilities: (i) either we find that there is no separator for T' in H, or (ii) we find another separator S^1 for T' in H. Again, we return to the beginning of this algorithm, by treating $S \leftarrow S^1$, $T \leftarrow T'$, and $H \leftarrow G$, and check whether the new $\langle G, T, S \rangle$, i.e., $\langle H, T', S^1 \rangle$ is a good and bad configuration, and so on. Suppose that in a particular iteration, let n_T denote the number of terminal vertices in current graph. Then in next iteration, the size of the vertex set strictly decreases, more specifically it decreases by at least $(n_T - 1)$. Hence, this algorithm runs for at most n iterations, and eventually we find a vertex set S'. Below we show that the set S' is indeed a strong separator for the original set of terminals T in the original graph G.

 \triangleright Claim 38. Let S' be the vertex set returned by the aforementioned procedure. Then S' must be a strong separator for T in G.

Proof. First, we show that S' is the separator for T in G. Assume that our procedure runs for ℓ iterations, and let $S^0 = S, S^1, \ldots, S^{\ell} = S'$ be the separators found in each iteration. We show by induction that S' is a separator for T. In the base case, trivially $S^0 = S$ is a

separator for T in G. Assume that for some $0 \le i \le \ell - 1$, S^i is the separator in G. Clearly S^i is not a strong separator, otherwise we have returned S^i in our procedure. As S^i is not a strong separator, then by definition $\langle G, T, S^i \rangle$ is a bad configuration and S^i creates a type 2 scenario. Now in our process, while working with the separator S^i , we consider a modified graph, say H, with a set of vertices $\{v_t : t \in T \setminus \{t_L\}\} \cup V(C(t_L))$ and find another minimal multiway cut S^{i+1} with the set of terminals $T' := \{t_L\} \cup \{v_t : t \in T \setminus \{t_L\}\}$. For the sake of contradiction, assume that S^{i+1} is not a separator for T in G, which means that there is a pair of terminal vertices s and t such that there is a path P_{st} between s and t in $G - S^{i+1}$. First, we consider the case where none of the s and t vertices is t_L . As $\langle G, T, S^i \rangle$ is a bad configuration, the path must consist of a pair of vertices of S_s^{i+1} and S_t^{i+1} . Hence we got a subpath connecting v_t and v_s of the path P_{st} in $H - S^{i+1}$, which is a contradiction to the fact that S^{i+1} is a separator for the terminal set $\{v_t : t \in T \setminus \{t_L\}\}$ in H. The same argument holds when $t_L \in \{s, t\}$. Therefore, by induction, we have shown that S^{ℓ} is a separator for T in G. Now, it remains to show that S^{ℓ} is a strong separator for T in G. Suppose not, then $\langle G, T, S^{\ell} \rangle$ must be a bad configuration, and S^{ℓ} must create a type 2 scenario. And our process must be able to find another separator S'' that is a contradiction to our termination condition. \triangleleft

▶ Lemma 39. There exists an algorithm that takes an input an instance $(G, \mathcal{M}, T, Q, k)$ of GIMC, where G is connected, $k \geq 2$, and $|T| \geq 3$, and in time $2^k \cdot n^{\mathcal{O}(1)}$, either returns that $\mathcal{F}(T, k, G) = \emptyset$, or returns a strong separator S corresponding to the instance, and two components C_s, C_t satisfying the property stated in Lemma 37. Here, n = |V(G)|.

From the above discussion, we obtain the following corollary.

▶ **Corollary 40.** An instance $(G, \mathcal{M}, T, Q, k)$ of GIMC, where G is connected, $k \ge 2$, and $|T| \ge 3$, if $\mathcal{F}(T, k, G) \neq \emptyset$ then G has a strong separator of size at most k.

5.2 A Recursive Algorithm for GIMC

We design a recursive algorithm called IMWCut for the problem GENERALIZED INDEPENDENT MULTIWAY CUT (GIMC) that takes the following as input.

- \blacksquare A graph G = (V, E),
- \blacksquare a set $T \subseteq V(G)$ of terminals,
- \blacksquare a set $Q \subseteq V(G)$ of special vertices, with $T \subseteq Q$,
- Matroid $\mathcal{M} = (U, \mathcal{I})$ of rank r, where $V(G) \setminus Q \subseteq U$, and
- two non-negative integers k, q where $k + q \leq r$.

We use $\mathcal{J} = (G, \mathcal{M}, T, Q, k, q)$ to denote the input instance to the algorithm. Given this instance \mathcal{J} , our goal of algorithm IMWCut is to find $\mathcal{F}'(T, k, G) \subseteq_{\mathrm{rep}}^{q} \mathcal{F}(T, k, G)$ of bounded size. For the values $k \leftarrow r$ and $q \leftarrow 0$, the solution of the instance $(G, \mathcal{M}, T, Q, r, 0)$ corresponding to the representative set of the entire solution of size r for INDEPENDENT MULTIWAY CUT in G for the terminal set T.

Step 0: Preprocessing We perform the following preprocessing steps exhaustively in the order in which they are mentioned. The soundness of these rules is immediate.

Rule 0.a. If there exists a non-terminal vertex $u \in V(G)$ that has no path to any terminal vertex in T, then make a recursive call IMWCut $(G - u, \mathcal{M} - u, T, Q, k, q)$. If the recursive call returns NO, then we return NO to original instance. Otherwise, return the solution returned by the recursive call.

Rule 0.b. If there exists a terminal vertex $t \in T$ that is disconnected from all other terminals in $T \setminus \{t\}$ in G, then make a recursive call IMWCut $(G - t, \mathcal{M} - t, T - t, Q, k, q)$. If the recursive call returns NO, then we return NO to original instance. Otherwise, return the solution returned by the recursive call.

Rule 0.c. If there exists a pair of adjacent terminal vertices, then return No. The following observation is immediate.

▶ **Observation 1.** After an exhaustive application of Rule 0.a and 0.b, each connected component of G contains at least two terminals of T.

Rule 0.c. At this point we have exhaustively applied the previous two rules. Now, if the number of connected components (each of which contains at least two terminals by Observation 1) of G is more than k, then we return NO.

Step 1: Base Case. Here we have following two cases.

Case a: If k = 1, then $\mathcal{F}(G, T, 1)$ consists of singleton vertex-sets $\{v\}$, such that $\{v\} \in \mathcal{I}$, and $\{v\}$ is a multiway cut for T (of size 1). For each $v \in V(G) \setminus T$, we can easily check whether $v \in \mathcal{F}(G, T, 1)$ in polynomial time; this implies that the set $\mathcal{F}(G, T, 1)$ can be found in polynomial time. Then, we compute $\mathcal{F}'(G, T, 1) \subseteq_{\mathrm{rep}}^q \mathcal{F}(G, T, 1)$ in $n^{\mathcal{O}(1)}$ time, and return $\mathcal{F}'(G, T, 1)$. The correctness of this step is immediate.

Case b: If |T| = 2, then we solve the instance $(G, \mathcal{M}, Q, s, t, k, q)$ of INDEPENDENT VERTEX (s, t)-CUT using Theorem 4, and return the family returned by the algorithm. The correctness of this step follows from that of Theorem 4, and the theorem implies that this step runs in time $f(r, k) \cdot n^{\mathcal{O}(1)}$ and returns a family $\mathcal{F}'(G, \{s, t\}, k)$ of size 2^r . The correctness of this step is immediate.

 \triangleright Claim 41. Let $\mathcal{F}'(G,T,k)$ denote the set that we return in the base case. Then $\mathcal{F}'(G,T,k) \subseteq_{\mathrm{rep}}^{q} \mathcal{F}(T,k,G).$

Step 2: Solving the problem in a disconnected graph. In this step, we describe how to deal with the problem when the underlying graph is not connected. Let $(G, \mathcal{M}, T, Q, k, q)$ be the instance and G be the union of ℓ connected components G_1, \ldots, G_ℓ . Observe that after an exhaustive application of the preprocessing (Rule 0.b), each component in G has at least two terminal components, so each minimal solution must have a non-empty intersection with each terminal component. Hence, we can assume that $\ell \leq k$. Let $T_i := T \cap V(G_i)$ and $Q_i \coloneqq Q \cap V(G_i)$. Now for each component G_i we try to find the smallest $k_i \leq k$ such that G_i has an independent multiway cut of size k_i – which can be found by making a recursive call to the algorithm for each component separately, starting from $k'_i = 1, 2, \ldots, k-1$, in order to find the smallest value k_i such that the recursive call for all $k'_i < k_i$ returned NO, and k_i returns a non-empty family. Note that, assuming the number of connected components is more than 1, if for some $i \in [\ell]$, we find that for $k'_i = k - 1$, the instance $(G_i, \mathcal{M}, T_i, Q_i, k_i, q+k-\beta_i)$ is a NO-instance, then we conclude that we have a NO-instance. The correctness of this follows from the fact that any $S \in \mathcal{F}(T, k, G)$ must have a non-empty intersection with each connected component. Hence, in each recursive call, the parameter strictly decreases. Note that we make a recursive call to most k^2 many sub-problems. Let k_i be the smallest value for each component G_i such that we find non-empty solution for $(G_i, \mathcal{M}, T_i, Q_i, k_i, q+k-k_i)$. Note that the k_i 's form a natural partition \mathcal{P} of k such that $k = k_1 + \ldots + k_\ell$. Let \mathcal{F}'_i be the output of the *i*th call with parameter k_i . We compute $\mathcal{F}'_1 \bullet \mathcal{F}'_2 \bullet \cdots \bullet \mathcal{F}'_{\ell}$, and we filter out any sets from the resultant that do not constitute a minimal independent multiway cut of size k for T. Let the remaining collection of sets be $\mathcal{F}^1(T,k,G)$. Finally, we compute and return $\mathcal{F}'(T,k,G) \subseteq_{\mathrm{rep}}^q \mathcal{F}^1(T,k,G)$.

 \triangleright Claim 42. $\mathcal{F}'(T,k,G) \subseteq_{\mathrm{rep}}^{q} \mathcal{F}(T,k,G).$

Proof. We show the claim in two parts. First, we show that $\mathcal{F}'(T, k, G) \subseteq \mathcal{F}(T, k, G)$, and then we show that the former also q-represents the latter. For the first part, consider an $R \in \mathcal{F}'(T, k, G)$. Hence, for each $i \in [\ell]$, there exists some $R_i \in \mathcal{F}'(T_i, k_i, G_i) \subseteq_{\mathrm{rep}}^{q+k-k_i}$ $\mathcal{F}(T_i, k_i, G_i)$, such that, (i) each R_i is a minimal independent multiway cut of size k_i for T_i in G_i , (ii) $R = R_1 \cup R_2 \cup \ldots \cup R_\ell$. It follows that R is a minimal multiway cut of size k for T in G, and thus $R \in \mathcal{F}(T, k, G)$.

Now consider $Y \subseteq U(\mathcal{M})$ be a set of size q such that there is a set $O \in \mathcal{F}(T, k, G)$ (i.e., O is a minimal multiway cut of size k for T), such that O fits Y. We have to show that there is a $\hat{X} \in \mathcal{F}'(T, k, G)$, such that \hat{X} fits Y, that is also a minimal multiway cut of size k for T.

We know that \mathcal{F}'_i is the output of the recursive call on $(G_i, \mathcal{M}, T_i, Q_i, k_i, q + k - k_i)$. By induction (recall that $k_i < k$), we know that $\mathcal{F}'_i \subseteq_{\operatorname{rep}}^{q+k-k_i} \mathcal{F}(G_i, k_i, T_i)$. Let $O_i := O \cap V(G_i)$, $Y_i := Y \cap V(G_i)$, and $Y_{out} = Y \setminus V(G)$. Note that $Y = Y_1 \cup Y_2 \cup \ldots \cup Y_\ell \cup Y_{out}$. Now we provide a recursive definition of \hat{X}_i for each $i \in [\ell]$. The set \hat{X}_1 is a representative for the set O_1 in the graph G_1 for the $k + q - k_1$ sized set $(O \cup Y) \setminus O_1$. Now for i > 1

$$Z_i \coloneqq \left(\bigcup_{j < i} (\hat{X_j} \cup Y_j)\right) \cup Y_i \cup \left(\bigcup_{j > i} (O_j \cup Y_j)\right) \cup Y_{out}.$$

Clearly, $|Z_i| = k + q - k_i$. By the property of \mathcal{F}'_i , there exists some $\hat{X}_i \in \mathcal{F}'_i$, such that \hat{X}_i is a minimum multiway cut of size k_i for T_i in G_i , and \hat{X}_i fits Z_i . Now, let $\hat{X} \coloneqq \hat{X}_1 \cup \cdots \cup \hat{X}_\ell$. Clearly $|\hat{X}| = \sum_{i=1}^{\ell} k_i = k$. As \hat{X}_i is a solution for G_i and G consists of ℓ components $G_i, i \in [\ell]$, so $\hat{X} \coloneqq \hat{X}_1 \cup \cdots \cup \hat{X}_\ell$ is a solution for G. By our recursive definition, we can conclude that $\hat{X}_\ell \cup Z_\ell$ is an independent set. Since $\hat{X}_\ell \cup Z_\ell = \hat{X} \cup Y$, we are done.

- **Step 3:** Finding a Strong Separator. Since Step 2 is not applicable, we know that the graph G is connected. Then we use our FPT algorithm of Lemma 39 to check whether G has a strong separator corresponding to the instance $(G, \mathcal{M}, T, Q, k, q)$ that runs in time $2^k \cdot n^{\mathcal{O}(1)}$. Either this algorithm returns a (w.l.o.g. minimal) separator $S \subseteq V(G)$ of size at most k, or it returns NO. In the former case, we proceed to the next step, i.e., Step 4. In the later case, we return NO to our original instance $(G, \mathcal{M}, T, Q, k, q)$. Because if G does not admit a multiway cut for T of size k, then it also does not admit an independent vertex multiway cut of size k. This shows the soundness of returning NO.
- Step 4: Branching procedure. Let $\mathcal{F} \coloneqq \mathcal{F}(G, T, k)$ denote the family of feasible solutions for the given instance $\mathcal{J} = (G, \mathcal{M}, T, Q, k, q)$. If $\mathcal{F} \neq \emptyset$, then Lemma 37 implies that, there exists a pair of components C_s and C_t , such that for any $O \in \mathcal{F}$, for each strong separator S of size k corresponding to the terminal set T, we have that, either (i) $S \cap O \neq \emptyset$, or (ii) either $1 \leq |C_s \cap O| \leq k - 1$, or $1 \leq |C_t \cap O| \leq k - 1$. Furthermore, as stated in Lemma 37, given a strong separator S, such a pair of components can be found in polynomial-time. However, since a-priori we do not know whether case (A) or (B) is applicable, we design a branching strategy that considers both possibilities. We describe this branching in the following.
 - Handling case (A): Branching on the vertices in S: We branch on each vertex v on S assuming that $v \in S \cap O$. More specifically, we do the following: For each vertex $v \in S$ we make a recursive call to IMWCut $(G v, \mathcal{M}, T, Q \setminus \{v\}, k 1, q)$. Let $\mathcal{F}'(v)$ be the set returned by the recursive call. Then $\mathcal{F}^1(v) \coloneqq \mathcal{F}'_v \bullet \{v\}$.

 \triangleright Claim 43. $\mathcal{F}^1(v) \subseteq_{\operatorname{rep}}^q \mathcal{F}(T, k, v, G)$, where $\mathcal{F}(T, k, v, G) \subseteq \mathcal{F}(T, k, G)$ is the set of all independent minimal multiway cuts for T that contain v.

Proof. First we argue that $\mathcal{F}^1(v) \subset \mathcal{F}(T, k, v, G)$, and then we show that $\mathcal{F}^1(v)$ also q-represents $\mathcal{F}(T, k, v, G)$. Consider some $R \in \mathcal{F}^1(v)$, and R must be equal to $Q \cup \{v\}$ for some $Q \in \mathcal{F}'_v$. By induction hypothesis, $Q \in \mathcal{F}(G-v, T, k-1)$, i.e., Q is a minimal multiway cut for T of size k-1 in G-v. It follows that $R=Q\cup\{v\}$ is also a minimal multiway cut of size k for T in G. Since $v \in R$, this shows that $\mathcal{F}^1(v) \subseteq \mathcal{F}(T, k, v, G)$. Now, consider some $Y \subseteq U(\mathcal{M})$ of size q for which there is a set $O \in \mathcal{F}(T, k, v, G)$ such that Y fits O. First, by the induction hypothesis, we have $\mathcal{F}'(v) \subseteq_{\mathrm{rep}}^{q+1} \mathcal{F}(T, k-1, G-v)$. This implies, in particular, that each $X \in \mathcal{F}'(v)$, is a minimal multiway cut of size k-1for T in G - v. We have to show that there is a $\hat{X} \in \mathcal{F}^1(v)$, such that \hat{X} is a minimal multiway cut of size k in such that \hat{X} fits Y. As $\mathcal{F}(T, k, v, G) \subseteq \mathcal{F}(T, k, G)$ is the set of all independent minimal multiway cuts for T that contain v, the vertex v must belong to O. Consider the set $O_v := O \setminus \{v\}$ and $Y_v := Y \cup \{v\}$. Clearly $O_v \cup Y_v = O \cup Y$. and for any $X \in \mathcal{F}'(v), v \notin X$. So for the set $Y_v \subseteq U(\mathcal{M})$, of size q+1 there is a set $O_v \in \mathcal{F}(T, k-1, G-v)$, such that O_v fits Y_v . Hence, there exists $X_v \in \mathcal{F}'(v)$ such that X_v also fits O_v . Further, $X_v \in \mathcal{F}(T, k - 1, G - v)$ as well. Now, consider the set $\dot{X} := X_v \cup \{v\}$. Obviously, \dot{X} is disjoint from Y. As X_v is an independent multiway cut for T in G-v, so $\hat{X} = X_v \cup \{v\}$ is an independent multiway cut in G. Finally, since $\hat{X} \cup Y = X_v \cup Y_v \in \mathcal{I}$, the claim follows. \leq

Handling case (B): Branching on terminal components in G - S: For a terminal vertex $x \in T$, we use the notation C_x to denote the terminal component in G - S containing the vertex x. Recall the components C_s and C_t returned by Lemma 37 with the property that, if \mathcal{J} is a YES-instance, and for some $O \in \mathcal{F}(T, k, G)$, it holds that $O \cap S = \emptyset$, then either $1 \leq |O \cap V(C_s)| \leq k - 1$, or $1 \leq |O \cap V(C_t)| \leq k - 1$. First we guess (i.e., branch) which components among C_t and C_s have the above mentioned property. WLOG, assume that the component C_t has such property, and the other case is symmetric.

Let $\operatorname{\mathsf{Reach}}(t)$ denote the set of vertices in S which is reachable (connected) from the vertex t in the induced subgraph $G[C_t \cup S]$. Next, we guess the following two things (i) the value $|V(C_t) \cap O|$ and (ii) the connectivity among the vertices in $S \cup \{t\}$ in the subgraph $G[V(C_t) \cup \operatorname{\mathsf{Reach}}(t)] - O$. More specifically, we guess a partition of the vertices in $\operatorname{\mathsf{Reach}}(t) \cup \{t\}$ such that it holds the following property: a pair of vertices $x, y \in \operatorname{\mathsf{Reach}}(t) \cup \{t\}$ belongs to the same set of the partition if and only if there is a path between x and y in $G[V(C_t) \cup \operatorname{\mathsf{Reach}}(t)] - O$. We introduce the following notation.

▶ Definition 44. Given a graph H and partition Π of vertex set $U \subseteq V(H)$, the graph obtained in the following procedure is denoted by H^*/Π : for each part $\pi \in \Pi$, we identify all the vertices of π into a single vertex, say v_{π} (we refer preliminary section to detail about the identify operation).

Now in our algorithm, for the component C_t , for each partition \mathcal{P} of the vertex set $\operatorname{Reach}(t) \cup \{t\}$, for each integer $1 \leq k_1 \leq k-1$ (let $k_2 = k-k_1$), we create two instances of INDEPENDENT MULTIWAY CUT and recursively call IMWCut on each of them. We highlight that the parameters for both the instances satisfy $1 \leq k_1, k_2 \leq k-1$, hence the parameter strictly decreases in both the recursive calls. These two instances are defined as follows.

■ Instance 1 is $\mathcal{J}_1 := (G_1, \mathcal{M}, T_1, Q_1, k_1, q + k_2)$. Here G_1 is H^*/\mathcal{P} , where $H := G[V(C_t) \cup \mathsf{Reach}(t)]$. That is $V(G_1) = V(C_t) \cup V(\mathcal{P})$ where $V(\mathcal{P})$ is the set of all new vertices added after processing identify operation on each $P \in \mathcal{P}$. And the terminal set $T_1 := V(\mathcal{P})$,



Figure 8 An illustration of recursive algorithm when branching on terminal components.

 $Q_1 := T_1$. Let $\mathcal{F}'(T_1, k_1, G_1) \subseteq_{\operatorname{rep}}^{q+k_2} \mathcal{F}(T_1, k_1, G_1)$ is the output returned by recursive call. By induction this is true.

Instance 2 is $\mathcal{J}_2 := (G_2, \mathcal{M}, T_2, Q_2, k_2, q+k_1)$. Here, the description of the graph G_2 and terminal set T_2 is based on the partition \mathcal{P} . We have the following two cases depending on whether the vertex t is identified with some other vertices in $\mathsf{Reach}(t)$ in \mathcal{P} or not. Let P_t denotes the set in the partition in \mathcal{P} which contains t.

Case 4.a. $|P_t| = 1$ Here t is not in same part in \mathcal{P} with any $v \in \mathsf{Reach}(t)$. Note that this implies that we require that t is disconnected from all $v \in \mathsf{Reach}(t)$ in each solution of \mathcal{J}_1 . Since any path between t and t' in G must use a vertex of $\mathsf{Reach}(t)$, it follows that G - X already disconnects t from all $t' \in T \setminus \{t\}$, for any solution X of \mathcal{J}_1 . Hence, in \mathcal{J}_2 , we may remove t from T.

More formally, we define $G_2 \coloneqq G^* / \mathcal{P} - V(C_t), T_2 \coloneqq T \setminus \{t\}$, and $Q_2 \coloneqq T_2 \cup (V(\mathcal{P}) \setminus \{t\})$. That is, $V(G_2) = (V(G) \setminus (V(C_t) \cup \mathsf{Reach}(t))) \cup (V(\mathcal{P}) \setminus \{t\})$.

Case 4.b. $|P_t| \ge 2$ In this case, the terminal t is identified with at least one vertex from Reach(t). Let v_t denote the result of the identify operation on the set P_t .

In this case $G_2 \coloneqq G^* / \mathcal{P} - (V(C_t) \setminus \{t\}), T_2 \coloneqq T \setminus \{t\} \cup \{v_t\}$, and $Q_2 \coloneqq T_2 \cup V(\mathcal{P})$. That is $V(G_2) = (V(G) \setminus (V(C_t) \cup \mathsf{Reach}(t))) \cup V(\mathcal{P})$.

Let $\mathcal{F}'(T_2, k_2, G_2) \subseteq_{\mathrm{rep}}^{q+k_1} \mathcal{F}(T_2, k_2, G_2)$ is the output returned by recursive call. By induction this is true.

More formally, let \mathcal{G}_t as the collection of all tuples (C_t, \mathcal{P}, k_1) of guesses corresponding to component C_t – analogously, we also define the set of guesses \mathcal{G}_s corresponding to C_s , and also recursively call the algorithm for each guess in \mathcal{G}_s , the pair sub-instances obtained in a similar way (we omit this description).

Now to return the solution for branching on terminal components, we do the following. First we compute $\mathcal{F}'(T_1, k_1, G_1) \bullet \mathcal{F}'(T_2, k_2, G_2)$, and from the resultant, we filter out any sets that do not form an independent minimal multiway cut for T in G. Let us refer to the remaining k-family as $\mathcal{F}^1(C_t, \mathcal{P}, k_1)$. We say a minimal independent multiway cut O is *compatible* with the guess (C_t, \mathcal{P}, k_1) if following two conditions hold: (i) $|O \cap V(C_t)| = k_1$, (ii) for any pair of vertices $u, v \in \mathsf{Reach}(t) \cup \{t\}$, if u and v belong to two different sets in the partition \mathcal{P} , then there is no path from u to v in the subgraph $G[V(C_t) \cup \mathsf{Reach}(t) \setminus O]$.

 \triangleright Claim 45. $\mathcal{F}^1(C_t, \mathcal{P}, k_1) \subseteq_{\operatorname{rep}}^q \mathcal{F}(T, k, G, C_t, \mathcal{P}, k_1)$, where $\mathcal{F}(T, k, G, C_t, \mathcal{P}, k_1) \subseteq \mathcal{F}(T, k, G)$ is the set of minimal independent multiway cuts that are compatible with the guess (C_t, \mathcal{P}, k_1) .

Proof. First, we show that $\mathcal{F}^1(C_t, \mathcal{P}, k_1) \subseteq \mathcal{F}(T, k, G, C_t, \mathcal{P}, k_1)$. To this end, consider any $R \in \mathcal{F}^1(C_t, \mathcal{P}, k_1)$, which implies that there exists some $P \in \mathcal{F}'(T_1, k_1, G_1), Q \in \mathcal{F}'(T_2, k_2, G_2)$, such that $R = P \cup Q, P \cap Q = \emptyset$, and R is indeed a minimal multiway cut of size k for T in G. Firstly, note $R \cap V(C_t) = P$, which implies that $|R \cap V(C_t)| =$ $|P| = k_1$. Secondly, since $P \in \mathcal{F}'(T_1, k_1, G_1) \subseteq_{\operatorname{rep}}^{q+k_2} \mathcal{F}(T_1, k_1, G_1)$, it follows that, for any $u, v \in \operatorname{Reach}(t) \cup \{t\}$, such that u, v belong to different parts of \mathcal{P} , there is no path from u to v in $G_1 - P$. Therefore, there is no path from u to v in $G[(V(C_t) \cup \operatorname{Reach}(t))] - P =$ $G[(V(C_t) \cup \operatorname{Reach}(t)) \setminus R]$. This shows that R is indeed compatible with the guess (C_t, \mathcal{P}, k_1) , i.e., $R \in \mathcal{F}(T, k, G, C_t, \mathcal{P}, k_1)$.

Next, we show that $\mathcal{F}^1(C_t, \mathcal{P}, k_1)$ also q-represents $\mathcal{F}(T, k, G, C_t, \mathcal{P}, k_1)$. To this end, consider $Y \subseteq U(\mathcal{M})$ of size q, for which there exists some $O \in \mathcal{F}(T, k, G, C_1, \mathcal{P}, k_1)$ such that O fits Y. We show that there exists some $\hat{X} \in \mathcal{F}^1(C_t, \mathcal{P}, k_1)$ such that \hat{X} also fits Y. Let $O_i \coloneqq O \cap V(G_i), Y_i \coloneqq Y \cap V(G_i)$, and $Y_{out} \coloneqq Y \setminus V(G)$. Note that $Y = Y_1 \cup Y_2 \cup Y_{out}$. Now we define the set \hat{X}_i for each $i \in [2]$. The set \hat{X}_1 is a representative for the set O_1 in the graph G_1 for the $q + k_2$ sized set $(O \cup Y) \setminus O_1$. Let $X_2 = \hat{X}_1 \cup Y$. As $|X_2| = q + k_1$ by induction, there exists some $\hat{X}_2 \in \mathcal{F}'(T_2, k_2, G_2)$, such that \hat{X}_2 is a minimum multiway cut of size k_2 for T_2 in G_2 , and \hat{X}_2 fits O_2 . Now, let $\hat{X} \coloneqq \hat{X}_1 \cup \hat{X}_2$. Clearly $|\hat{X}| = |\hat{X}_1| + |\hat{X}_2| = k_1 + k_2$. By our argument, we can conclude that $\hat{X}_2 \cup X_2$ is an independent set. Since $\hat{X}_2 \cup X_2 = \hat{X}_2 \cup \hat{X}_1 \cup Y = \hat{X} \cup Y$, we have that $\hat{X} \cup Y$ is an independent set.

It remains to show that \hat{X} is a minimal multiway cut for T in G. Consider a shortest path P_{12} between two terminal vertices, say $t_1, t_2 \in T$ in $G - \hat{X}$.

Case 1 First, we analyze the case where $t \notin \{t_1, t_2\}$. Naturally, P_{12} is also a shortest path between t_1 and t_2 . In this case, the path P_{12} must use a vertex from $V(C_t)$, otherwise \hat{X}_2 is not a solution for \mathcal{J}_1 . We first claim that, $|V(P_{12}) \cap \mathsf{Reach}(t)| \ge 2$. Suppose to the contrary that $|V(P_{12} \cap \mathsf{Reach}(t)| \le 1$, then there exists a path P'_{12} in $G_2 - \hat{X}_2$ between t_1 and t_2 , which contradicts the fact that \hat{X}_2 is a solution for \mathcal{J}_2 . Thus, $|V(P_{12}) \cap \mathsf{Reach}(t)| \ge 2$, Now consider distinct $x_1, x_2 \in V(P_{12}) \cap \mathsf{Reach}(t)$, such that the internal vertices of the subpath P'_{12} , say I, between x_1 and x_2 satisfies that $\emptyset \neq I \subseteq V(C_t)$. Now, suppose that x_1 and x_2 are adjacent in G, then we can obtain a path between t_1 and t_2 by replacing $x_1, V(I), x_2$ by the edge x_1x_2 in P_{12} , which yields a shorter path between t_1 and t_2 , which is a contradiction to the assumption that P_{12} is a shortest path. In summary, we have a subpath P'_{12} of P_{12} such that the endpoints of P'_{12} , that is, $x_1, x_2 \in \mathsf{Reach}(t)$, and $\emptyset \neq I \subseteq V(C_t)$. Now we consider different cases and arrive at a contradiction.

Case (1a): x_1 and x_2 are in the same part $P \in \mathcal{P}$. Then P is contracted into a single vertex v_P in G_2 . This implies that there is also a path between t_1 and t_2 in $G_2 - \hat{X}_2$, which is a contradiction.

Case (1b): x_1 and x_2 are in different parts P_1 and P_2 respectively, which are identified into the vertices v_1, v_2 in G_1 , respectively. However, this implies that there is a path P'_{12} between v_1 and v_2 in $G_1 - \hat{X}_1$, which contradicts that X_1 is a solution for \mathcal{J}_1 .

A. Banik, F. V. Fomin, P. A. Golovach, T. Inamdar, S. Jana, S. Saurabh

Case 2 Now, we analyze the case where $t \in \{t_1, t_2\}$. WLOG, assume that $t = t_1$; therefore, P_{12} is the shortest path between t and t_2 in $G - \hat{X}$. In this case, as $t \in V(C_t)$ the path P_{12} must use a vertex from $(V(C_t) \setminus \{t\}) \cup \text{Reach}(t)$. In path P_{12} , let x_2 denote the vertex in Reach(t) such that all the internal vertices of the subpath connecting x_2 to t along P_2 are from $V(C_t)$. Now we consider different cases and arrive at a contradiction.

Case (2a): x_2 and t are in the same part $P \in \mathcal{P}$. Then it follows that there is a path from t_2 to v_P in $G - \hat{X}_2$, which contradicts the assumption that \hat{X}_2 is a solution for \mathcal{J}_2 .

Case (2b): t and x_2 are in different parts. Now, we note that there is a path between t and x_2 in $G[V(C_t) \cup \mathsf{Reach}(t)] - \hat{X}_1$, it contradicts the assumption that \hat{X}_1 is a solution for \mathcal{J}_1 .

This completes the claim.

Here is a simple observation.

 \triangleright Claim 46. Consider any $O \in \mathcal{F}(T, k, G)$ such that $O \cap S = \emptyset$. Then, there is some guess (C_t, \mathcal{P}, k_1) with which O is compatible.

Proof. Assuming that $O \cap S = \emptyset$, we know that at least one of the components C_s or C_t has a nonempty intersection with the vertices of O. Now we obtain a guess that is compatible with O. Assume that $|V(C_t) \cap O| = k_1 > 0$. Let \mathcal{P} be the partition of the vertices in $\operatorname{Reach}(t) \cup \{t\}$ defined as follows. A pair of vertices u and v from $\operatorname{Reach}(t) \cup \{t\}$ belongs to a different part if they are not connected in the graph $G[V[C_t] \cup \operatorname{Reach}(t)] = O$ and the graph $G - (V(C_t) \cup O)$. It is easy to see that guess (C_t, \mathcal{P}, k_1) is compatible with O.

Step 5: Return as output We return NO if each of the branching steps returns NO to their corresponding instance. Otherwise, we do the following. Notice that in each branching step, we have at most $k + 2(k - 1)d_{k+1} = k^{\mathcal{O}(k)}$ many recursive calls, where d_{k+1} denotes the (k + 1) th Bell's number: number of possible partitions of a set of (k + 1) elements. First we take union of all the solution sets corresponding to every recursive calls, specifically, we compute

$$\mathcal{F}^{1}(T,k,G) \coloneqq \bigcup_{v \in S} \mathcal{F}^{1}_{v} \cup \bigcup_{(C_{t},\mathcal{P},k_{1}) \in \mathcal{G}_{t}} \mathcal{F}^{1}(C_{t},\mathcal{P},k_{1}) \cup \bigcup_{(C_{s},\mathcal{P},k_{1}) \in \mathcal{G}_{s}} \mathcal{F}^{1}(C_{t},\mathcal{P},k_{1}).$$
(1)

Then we compute

$$\mathcal{F}'(T,k,G) \subseteq_{\mathrm{rep}}^{q} \mathcal{F}^1(T,k,G) \tag{2}$$

using Proposition 2 and return the set $\mathcal{F}'(T, k, G)$ as the output for the instance $\mathcal{J} = (G, \mathcal{M}, T, Q, k, q)$.

This finishes the description of IMWCut.

Correctness.

▶ Lemma 47 (Safeness of IMWCut). Our branching algorithm returns the set $\mathcal{F}'(T, k, G) \subseteq_{rep}^{q} \mathcal{F}(T, k, G)$ of size 2^{k+q} .

Proof. Recall that $\mathcal{F}(T, k, G)$ is the set of all minimal independent multiway cuts for T of size k in G. First, we consider base case, then the correctness follows from Claim 41. Now, suppose that neither of the base cases is applicable. If the graph G is disconnected, then again the correctness of the algorithm follows directly from Claim 42. Hence, finally let us consider the case when G is connected, $k \geq 2$, and $|T| \geq 3$. First, by Lemma 39, we either correctly conclude that $\mathcal{J} = (G, \mathcal{M}, T, Q, k, q)$ is a NO-instance, or we find a strong separator S and a pair of components C_s, C_t in time $2^k \cdot n^{\mathcal{O}(1)}$, such that for any minimal solution $O \in \mathcal{F}(G, T, k)$ of size k, either (i) $S \cap O = \emptyset$, (ii) $1 \leq |C_s \cap O| \leq k - 1$, or (iii) $1 \leq |C_t \cap O| \leq k - 1$. Further, any O satisfying (ii) (resp. (iii)) must be compatible with some guess $(C_t, \mathcal{P}, k_1) \in \mathcal{G}_t$ (resp. $(C_s, \mathcal{P}, k_1) \in \mathcal{G}_s$) (by claim 46). This implies that,

$$\mathcal{F}(T,k,G) = \bigcup_{v \in S} \mathcal{F}(T,k,v,G) \cup \bigcup_{(C_t,\mathcal{P},k_1) \in \mathcal{G}_t} \mathcal{F}(T,k,G,C_t,\mathcal{P},k_1)$$
$$\cup \bigcup_{(C_s,\mathcal{P},k_1) \in \mathcal{G}_s} \mathcal{F}(T,k,G,C_s,\mathcal{P},k_1).$$
(3)

Then, by Claims 43 and 45, it follows that $\mathcal{F}^1(T, k, G) \subseteq_{\operatorname{rep}}^q \mathcal{F}(T, k, G)$, where the former is defined in (1). Then, since $\mathcal{F}'(T, k, G) \subseteq_{\operatorname{rep}}^q \mathcal{F}^1(T, k, G)$ as computed in (2), by Proposition 3, it follows that $\mathcal{F}'(T, k, G) \subseteq_{\operatorname{rep}}^q \mathcal{F}(T, k, G)$. Finally, by Proposition 2, it follows that $|\mathcal{F}'(T, k, G)| \leq 2^{k+q}$.

Time analysis. Note that in each of the recursive calls to our branching steps, the parameter k decreases by at least 1. The number of partition of the vertices in S is bounded by k^k . So the degree at each node of the branching tree is bounded by $k^{\mathcal{O}(k)}$. In the base case where |T| = 2 or k = 1 we found the corresponding solution in time $f(r, k) \cdot n^{\mathcal{O}(1)}$. The number of nodes in the tree is bounded by $k^{\mathcal{O}(k)}$. Now we calculate the running to find a solution to each node assuming that we have all the solution to its child node. For this, we have to take union of at most $(k+2(k-1)(k+1)^{k+1})^k$ many sets where the size of each set is at most $\binom{k+q}{k}$. This union takes $\binom{k+q}{k} \cdot (k+2(k-1)(k+1)^{k+1})^k$ time. Then finding the q-representative set will take time $\binom{k+q}{k}$ using at most $\mathcal{O}(|\mathcal{A}|(\binom{k+q}{k})k^{\omega} + (\binom{k+q}{k})^{\omega-1})$ (by Proposition 2, where \mathcal{A} is the size of the set after taking the union.

Hence, we have the following theorem.

▶ **Theorem 48.** There exists a deterministic algorithm for GENERALIZED INDEPENDENT MULTIWAY CUT that runs in time $f(r,k) \cdot n^{\mathcal{O}(1)}$, and outputs $\mathcal{F}'(T,k,G) \subseteq_{rep}^{q} \mathcal{F}(T,k,G)$ such that $|\mathcal{F}'(T,k,G)| = 2^{k+q}$. Here $\mathcal{F}(T,k,G)$ denotes the set of all k size multiway cuts for T in G, assuming that there is no independent multiway cut of size k' < k.

6 Lower bound

In this section, we show an unconditional lower bound for the INDEPENDENT (s, t)-CUT problem when the input matroid is given by the independence oracle. The proof is inspired by the reductions used for the similar statements in [10, Theorem 1].

▶ **Theorem 49.** There is no algorithm solving INDEPENDENT (s, t)-CUT with matroids of rank k represented by the independence oracles using $f(k) \cdot n^{o(k)}$ oracle calls for any computable function f.

Proof. Let p and q be positive integers. We construct the graph $G_{p,q}$ as follows (see Figure 9).



Figure 9 Construction of $G_{p,q}$.

- \blacksquare Construct two vertices s and t.
- For each $i \in [p]$, construct a set $A_i = \{a_1^i, \ldots, a_q^i\}$ of q vertices and a set $B_i = \{b_1^i, \ldots, b_q^i\}$ of q vertices.
- For each $i \in [p]$, make s adjacent to a_1^i and b_1^i , and make t adjacent to a_q^i and b_q^i .
- For each $i \in [p]$ and $j \in [q-1]$, make a_j^i adjacent to a_{j+1}^i and b_{j+1}^i , and make b_j^i adjacent to a_{j+1}^i and b_{j+1}^i .

We have that G_{pq} has 2pq + 2 vertices. Notice that each minimum (s, t)-cut in $G_{p,q}$ has size 2p and is of form $\bigcup_{i=1}^{p} \{a_{j_i}^i, b_{j_i}^i\}$ for some $j_1, \ldots, j_p \in [q]$.

Consider a family of indices $j_1, \ldots, j_p \in [q]$ and set $W = \bigcup_{i=1}^p \{a_{j_i}^i, b_{j_i}^i\}$. We define the matroid \mathcal{M}_W with the ground set $V(G_{p,q})$ where each independent set is a subset of $V(G_{p,q}) \setminus \{s,t\}$ as follows for k = 2p:

- Each set $X \subseteq V(G_{p,q}) \setminus \{s,t\}$ of size at most k-1 is independent and any set of size at least k+1 is not independent.
- A set $X \subseteq V(G_{p,q}) \setminus \{s,t\}$ of size k is independent if and only if either X = W or there is $i \in \{1, \ldots, p\}$ such that $|A_i \cap X| \ge 2$ or $|B_i \cap X| \ge 2$ or there are distinct $h, j \in [q]$ such that $a_h^i, b_i^i \in X$.

We denote by \mathcal{I}_W the constructed family of independent sets. In fact, \mathcal{M}_W is the same as the matroid constructed in the proof of [10, Theorem 1]. We provide the proof of the following claim showing that \mathcal{M}_W is a matroid for completeness even if it identical to the proof of [10, Claim 1.1] in the proof of Theorem 1 therein.

 \triangleright Claim 50 ([10, Claim 1.1]). $\mathcal{M}_W = (V(G_{p,q}), \mathcal{I}_W)$ is a matroid of rank k = 2p.

Proof of Claim 50. We have to verify that \mathcal{I}_W satisfies the independence axioms (1)–(3). The axioms (1) and (2) for \mathcal{I}_W follow directly from the definition of \mathcal{I}_W . To establish (3), consider arbitrary $X, Y \in \mathcal{I}_W$ such that |X| < |Y|. If |X| < k - 1 then for any $v \in Y \setminus X$, $Z = X \cup \{v\} \in \mathcal{I}_W$ because $|Z| \le k - 1$.

Suppose |X| = k - 1 and |Y| = k. If there is $i \in [p]$ such that $|A_i \cap X| \ge 2$ or $|B_i \cap X| \ge 2$ or there are distinct $h, j \in [q]$ such that $a_h^i, b_j^i \in X$ then for any $v \in Y \setminus X$, the set $Z = X \cup \{v\}$ has the same property and, therefore, $Z \in \mathcal{I}_W$. Assume that this is not the case. By the construction of $G_{p,q}$, we have that for each $i \in [p], |X \cap A_i| \le 1$ and $|X \cap B_i| \le 1$, and, furthermore, there is $j \in [q]$ such that $X \cap (A_i \cup B_i) \subseteq \{a_j^i, b_j^i\}$. Because |X| = k - 1, we can assume without loss of generality that there are indices $h_1, \ldots, h_p \in [q]$

such that $X \cap (A_i \cup B_i) = \{a_{h_i}^i, b_{h_i}^i\}$ for $i \in [p]$ and $X \cap (A_p \cup B_p) = \{a_{h_p}^p\}$. Recall that $W = \bigcup_{i=1}^p \{a_{j_i}^i, b_{j_i}^i\}$ for $j_1, \ldots, j_p \in [q]$. If there is $v \in Y \setminus X$ such that $v \neq b_{j_p}^p$ then consider $Z = X \cup \{v\}$. We have that there is $i \in [p]$ such that $|A_i \cap Z| \ge 2$ or $|B_i \cap Z| \ge 2$ or there are distinct $h, j \in [q]$ such that $a_h^i, b_j^i \in Z$, that is, $Z \in \mathcal{I}_W$. Now we assume that $Y \setminus X = \{b_{j_p}^p\}$. Then Y = W and we can take $v = b_{j_p}^p$. We obtain that $X \cup \{v\} = Y \in \mathcal{I}_W$.

By the construction of \mathcal{I} , the rank of \mathcal{M} is k. This concludes the proof.

We show the following lower bound for the number of oracle queries for frameworks $(G_{p,q}, \mathcal{M}_W)$; the proof is similar to the proof of [10, Claim 1.2] and we provide it for completeness.

 \triangleright Claim 51. Solving INDEPENDENT (s,t)-CUT for instances $(G_{p,q}, \mathcal{M}_W, s, t)$ with the matroids \mathcal{M}_W defined by the independence oracle for an (unknown) stable set W of $G_{p,q}$ of size k = 2p demands at least $q^p - 1$ oracle queries.

Proof of Claim 51. Recall that any (s,t)-cut in $G_{p,q}$ has size at least k and all sets $X \subseteq$ $V(G_{p,q}) \setminus \{s,t\}$ of size at least k+1 are not independent with respect to \mathcal{M} . Thus, we are looking for an (s,t)-cut of size k. For every (s,t)-cut X in $G_{p,q}$ of size $k, X = \bigcup_{i=1}^{p} \{a_{h_i}^i, b_{h_i}^i\}$ for some $h_1, \ldots, h_p \in [q]$. By the definition of \mathcal{M} , any X of this form is independent if and only if X = W. This means that the problem boils down to identifying unknown W using oracle queries. Querying the oracle for sets X of size at most k-1 or at least k+1 does not provide any information about W. Also, querying the oracle for X of size k with the property that there is $i \in [p]$ such that $|A_i \cap X| \geq 2$ or $|B_i \cap X| \geq 2$ or there are distinct $h, j \in [q]$ such that $a_h^i, b_i^i \in X$ also does not give any information because all these are independent. Hence, we can assume that the oracle is queried only for sets X of size k with the property that for each $i \in [p]$, there is $j \in [q]$ such that $X \cap V(G_i) = \{a_i^i, b_i^i\}$, that, is the oracle is queried for (s,k)-cuts of size k. The graph $G_{p,q}$ has q^p such sets. Suppose that the oracle is queried for at most $q^p - 2$ stable sets of size k with the answer no. Then there are two distinct stable sets W and W' of size k such that the oracle was queried neither for W nor W'. The previous queries do not help to distinguish between W and W'. Hence, at least one more query is needed. This proves the claim.

To show the theorem, suppose that there is an algorithm \mathcal{A} solving INDEPENDENT (s, t)-CUT with at most $f(k) \cdot n^{g(k)}$ oracle calls on matroids of rank k for computable functions f and g such that g(k) = o(k). Without loss of generality, we assume that f and g are monotone non-decreasing functions. Because g(k) = o(k), there is a positive integer K such that g(k) < k/2 for all $k \ge K$. Then for each $k \ge K$, there is a positive integer N_k such that for every $n \ge N_k$, $(f(k) \cdot n^{g(k)} + 1)k^{k/2} < (n-2)^{k/2}$.

Consider instances $(G_{p,q}, \mathcal{M}_W)$ for even $k \ge K$ where p = k/2 and $q \ge (N_k - 2)/k$. We have that k = 2p and n = 2pq + 2. Then \mathcal{A} applied to such instances would use at most $f(k) \cdot n^{g(k)} < \left(\frac{n-2}{k}\right)^{k/2} - 1 = q^p - 1$ oracle queries contradicting Claim 51. This completes the proof.

7 Conclusion

In this paper we studied matroidal generalizations of several fundamental graph separation problems, such as VERTEX (s, t)-CUT and VERTEX MULTIWAY CUT. We showed that these problems are FPT when parameterized by the size of the solution. These results were obtained by combining several recent and old techniques in the world of parameterized complexity. Our paper leaves several interesting questions open. These include obtaining $2^{\mathcal{O}(k)} \cdot n^{\mathcal{O}(1)}$ time algorithm for all the problems studied in this paper. In addition, designing FPT algorithms for matroid versions of other problems such as VERTEX MULTI-CUT, DIRECTED FEEDBACK VERTEX SET remains an interesting research direction.

— References -

- 1 Noga Alon, Raphael Yuster, and Uri Zwick. Color-coding. J. ACM, 42(4):844-856, 1995. URL: http://dx.doi.org/10.1145/210332.210337, doi:10.1145/210332.210337.
- 2 Gruia Călinescu, Chandra Chekuri, Martin Pál, and Jan Vondrák. Maximizing a monotone submodular function subject to a matroid constraint. SIAM J. Comput., 40(6):1740–1766, 2011. doi:10.1137/080733991.
- 3 William H. Cunningham. Improved bounds for matroid partition and intersection algorithms. SIAM J. Comput., 15(4):948–957, 1986. doi:10.1137/0215066.
- 4 M. Cygan, F. Fomin, L. Kowalik, D. Lokshtanov, D. Marx, M. Pilipczuk, M. Pilipczuk, and S. Saurabh. *Parameterized Algorithms*. Springer, 2015.
- 5 Marek Cygan, Fedor V. Fomin, Lukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michal Pilipczuk, and Saket Saurabh. *Parameterized Algorithms*. Springer, 2015. doi:10.1007/978-3-319-21275-3.
- 6 Marek Cygan, Fabrizio Grandoni, and Danny Hermelin. Tight kernel bounds for problems on graphs with small degeneracy. ACM Trans. Algorithms, 13(3):43:1–43:22, 2017. doi: 10.1145/3108239.
- 7 Marek Cygan, Marcin Pilipczuk, Michal Pilipczuk, and Jakub Onufry Wojtaszczyk. On multiway cut parameterized above lower bounds. ACM Trans. Comput. Theory, 5(1):3:1–3:11, 2013. doi:10.1145/2462896.2462899.
- 8 Uriel Feige and Mohammad Mahdian. Finding small balanced separators. In Jon M. Kleinberg, editor, Proceedings of the 38th Annual ACM Symposium on Theory of Computing, Seattle, WA, USA, May 21-23, 2006, pages 375–384. ACM, 2006. doi:10.1145/1132516.1132573.
- 9 F. V. Fomin, D. Lokshtanov, and S. Saurabh. Efficient computation of representative sets with applications in parameterized and exact algorithms. In *proceedings of SODA*, pages 142–151, 2014.
- 10 Fedor V. Fomin, Petr A. Golovach, Tuukka Korhonen, and Saket Saurabh. Stability in graphs with matroid constraints, 2024. arXiv:2404.03979.
- 11 Fedor V. Fomin, Daniel Lokshtanov, Fahad Panolan, and Saket Saurabh. Efficient computation of representative families with applications in parameterized and exact algorithms. J. ACM, 63(4):29:1–29:60, 2016. doi:10.1145/2886094.
- 12 Gagan Goel, Chinmay Karande, Pushkar Tripathi, and Lei Wang. Approximability of combinatorial problems with multi-agent submodular cost functions. In 50th Annual IEEE Symposium on Foundations of Computer Science, FOCS 2009, October 25-27, 2009, Atlanta, Georgia, USA, pages 755–764. IEEE Computer Society, 2009. doi:10.1109/F0CS.2009.81.
- 13 MohammadTaghi Hajiaghayi, Rohit Khandekar, and Guy Kortsarz. Local search algorithms for the red-blue median problem. *Algorithmica*, 63(4):795-814, 2012. URL: https://doi.org/ 10.1007/s00453-011-9547-9, doi:10.1007/S00453-011-9547-9.
- 14 Zhong Huang and Xueliang Li. Hardness results for rainbow disconnection of graphs. CoRR, abs/1811.11939, 2018. URL: http://arxiv.org/abs/1811.11939, arXiv:1811.11939.
- 15 Stefanie Jegelka and Jeff Bilmes. Notes on graph cuts with submodular edge weights. In NIPS 2009 Workshop on Discrete Optimization in Machine Learning: Submodularity, Sparsity Polyhedra (DISCML), pages 1–6, 2009.
- Eun Jung Kim, Stefan Kratsch, Marcin Pilipczuk, and Magnus Wahlström. Flow-augmentation
 i: Directed graphs. arXiv preprint arXiv:2111.03450, 2021.
- 17 Eun Jung Kim, Stefan Kratsch, Marcin Pilipczuk, and Magnus Wahlström. Solving hard cut problems via flow-augmentation. In Dániel Marx, editor, *Proceedings of the 2021 ACM-SIAM*

Symposium on Discrete Algorithms, SODA 2021, Virtual Conference, January 10 - 13, 2021, pages 149–168. SIAM, 2021. doi:10.1137/1.9781611976465.11.

- 18 Eun Jung Kim, Stefan Kratsch, Marcin Pilipczuk, and Magnus Wahlström. Directed flowaugmentation. In Stefano Leonardi and Anupam Gupta, editors, STOC '22: 54th Annual ACM SIGACT Symposium on Theory of Computing, Rome, Italy, June 20 - 24, 2022, pages 938–947. ACM, 2022. doi:10.1145/3519935.3520018.
- 19 Ravishankar Krishnaswamy, Shi Li, and Sai Sandeep. Constant approximation for k-median and k-means with outliers via iterative rounding. In Ilias Diakonikolas, David Kempe, and Monika Henzinger, editors, Proceedings of the 50th Annual ACM SIGACT Symposium on Theory of Computing, STOC 2018, Los Angeles, CA, USA, June 25-29, 2018, pages 646–659. ACM, 2018. doi:10.1145/3188745.3188882.
- 20 Daniel Lokshtanov, Pranabendu Misra, Fahad Panolan, and Saket Saurabh. Deterministic truncation of linear matroids. In Magnús M. Halldórsson, Kazuo Iwama, Naoki Kobayashi, and Bettina Speckmann, editors, *ICALP 2015*, pages 922–934, 2015. doi: 10.1007/978-3-662-47672-7_75.
- 21 Daniel Lokshtanov, Pranabendu Misra, Fahad Panolan, and Saket Saurabh. Deterministic truncation of linear matroids. ACM Trans. Algorithms, 14(2):14:1–14:20, 2018. doi:10.1145/ 3170444.
- 22 László Lovász. Flats in matroids and geometric graphs. In Combinatorial Surveys (Proc. 6th British Combinatorial Conference, pages 45–86, 1977.
- 23 László Lovász. Graphs and geometry, volume 65 of American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, RI, 2019. doi:10.1090/coll/065.
- 24 Pasin Manurangsi, Erel Segal-Halevi, and Warut Suksompong. On maximum bipartite matching with separation. Inf. Process. Lett., 182:106388, 2023. URL: https://doi.org/10. 1016/j.ipl.2023.106388, doi:10.1016/J.IPL.2023.106388.
- Dániel Marx. Parameterized graph separation problems. Theor. Comput. Sci., 351(3):394–406, 2006. URL: https://doi.org/10.1016/j.tcs.2005.10.007, doi:10.1016/J.TCS.2005.10.007.
- 26 Dániel Marx. A parameterized view on matroid optimization problems. In Hajo Broersma, Stefan S. Dantchev, Matthew Johnson, and Stefan Szeider, editors, ACiD 2006, volume 7 of Texts in Algorithmics, page 158. King's College, London, 2006.
- 27 Dániel Marx. A parameterized view on matroid optimization problems. Theor. Comput. Sci., 410(44):4471-4479, 2009. doi:10.1016/j.tcs.2009.07.027.
- 28 Dániel Marx, Barry O'Sullivan, and Igor Razgon. Treewidth reduction for constrained separation and bipartization problems. In Jean-Yves Marion and Thomas Schwentick, editors, 27th International Symposium on Theoretical Aspects of Computer Science, STACS 2010, March 4-6, 2010, Nancy, France, volume 5 of LIPIcs, pages 561–572. Schloss Dagstuhl Leibniz-Zentrum für Informatik, 2010. URL: https://doi.org/10.4230/LIPIcs.STACS.2010.2485, doi:10.4230/LIPICS.STACS.2010.2485.
- 29 Dániel Marx, Barry O'Sullivan, and Igor Razgon. Finding small separators in linear time via treewidth reduction. ACM Trans. Algorithms, 9(4):30:1–30:35, 2013. doi:10.1145/2500119.
- 30 Syed Mohammad Meesum, Fahad Panolan, Saket Saurabh, and Meirav Zehavi. Rank vertex cover as a natural problem for algebraic compression. SIAM J. Discret. Math., 33(3):1277–1296, 2019. doi:10.1137/17M1154370.
- 31 Rolf Niedermeier and Peter Rossmanith. On efficient fixed-parameter algorithms for weighted vertex cover. J. Algorithms, 47(2):63–77, 2003. doi:10.1016/S0196-6774(03)00005-1.
- 32 Bruce A. Reed, Kaleigh Smith, and Adrian Vetta. Finding odd cycle transversals. Oper. Res. Lett., 32(4):299-301, 2004. URL: https://doi.org/10.1016/j.orl.2003.10.009, doi: 10.1016/J.ORL.2003.10.009.
- 33 Hadas Shachnai and Meirav Zehavi. A multivariate framework for weighted FPT algorithms. J. Comput. Syst. Sci., 89:157–189, 2017. URL: https://doi.org/10.1016/j.jcss.2017.05.003, doi:10.1016/J.JCSS.2017.05.003.

34 Laurence A Wolsey. An analysis of the greedy algorithm for the submodular set covering problem. *Combinatorica*, 2(4):385–393, 1982.