# LIMITS OF DEFINABLE FAMILIES AND DILATIONS IN NILMANIFOLDS

#### YA'ACOV PETERZIL AND SERGEI STARCHENKO

ABSTRACT. Let G be a unipotent group and  $\mathcal{F} = \{F_t : t \in (0, \infty)\}$  a family of subsets of G, with  $\mathcal{F}$  definable in an o-minimal expansion of the real field. Given a lattice  $\Gamma \subseteq G$ , we study the possible Hausdorff limits of  $\pi(\mathcal{F})$  in  $G/\Gamma$  as t tends to  $\infty$  (here  $\pi: G \to G/\Gamma$  is the canonical projection). Towards a solution, we associate to  $\mathcal{F}$  finitely many real algebraic subgroups  $L \subseteq G$ , and, uniformly in  $\Gamma$ , determine if the only Hausdorff limit at  $\infty$  is  $G/\Gamma$ , depending on whether  $L^{\Gamma} = G$  or not. The special case of polynomial dilations of a definable set is treated in details.

# 1. Introduction Contents

1	. Inti	roduction	1
2	. Lor	ng types and $\mu$ -stabilizers	4
	2.1.	Model theoretic preliminaries	4
	2.2.	Long and short sets	7
	2.3.	The $\mu$ -stabilizer of a type	9
3	. Nea	arest cosets	11
	3.1.	Nearest co-commutative cosets	11
	3.2.	The set $\mathscr{L}_{\max}(\mathcal{X})$	14
4	. Γ-d	ense types in unipotent groups	16
	4.1.	Preliminaries on unipotent groups	16
	4.2.	$\Gamma$ -dense sets in unipotent groups	17
	4.3.	$\Gamma$ -dense types in abelian groups	18
	4.4.	Abelianization and density	20
5	. Γ-d	ense definable subsets of unipotent groups	22
6	. Inte	erpreting the results as Hausdorff limits	27
	6.1.	Hausdorff limits	27
	6.2.	Haudorff Limits via the standard part map	28
	6.3.	Hausdorff limits in $G/H$	29
	6.4.	Hausdorff limits in nilmanifolds	29

The first author was partially supported by ISF grant 290/19.

The second author was partially supported by NSF research grant DMS-1800806.

6.5. T	'he abelian case	32		
7. Polynomial dilations				
7.1. T	The setting	35		
7.2. P	olynomial dilations in vector spaces	35		
7.3. P	olynomial dilations in unipotent groups	42		
References				

Let G be  $\langle \mathbb{R}^n, + \rangle$  or more generally a real unipotent group, and let  $X \subseteq G$  be a definable set in some o-minimal structure over  $\mathbb{R}$ . In [8] and [9] we examined the following problem: For a lattice  $\Gamma \subseteq G$ , and  $\pi: G \to G/\Gamma$ , what is the topological closure of  $\pi(X)$  in  $G/\Gamma$ ?

Using model theoretic machinery, we described the frontier of  $\operatorname{cl}(\pi(X))$  as the projection of finitely many definable families of cosets of positive dimensional subgroups associated to X. The answer can be seen, in a certain sense, as uniform in  $\Gamma$ .

Here we consider an extension of the problem:

For G as above, let  $\{F_s : s \in S\}$  be a family of subsets of G that is definable in an o-minimal structure over the reals, let  $\Gamma \subseteq G$  be a lattice and  $\pi : G \to G/\Gamma$  the projection. What are the possible Hausdorff limits of the family  $\{\pi(F_s) : s \in S\}$  in  $G/\Gamma$ ? How does the answer vary with  $\Gamma$ ?

Some results of this paper can be seen as an extension of work [3] and [1] on polynomial dilations in nilmanifolds. But instead of considering equidistribution of certain measures linked to these dilations, we focus here on topological properties (see Section 7 below).

Our precise setting is as follows: Let  $\mathbb{R}_{om}$  be an o-minimal expansion of the field of reals. Let  $\mathcal{F} = \{F_t : t \in (0, \infty)\}$  be an  $\mathbb{R}_{om}$ -definable family of subsets of G, and let  $\Gamma$  be a lattice in G. We study the possible Hausdorff limits of the family  $\{\pi(F_t) : t \in (0, \infty)\}$ , as t tends to  $\infty$ . Using model theory, we replace the Hausdroff limits question by a question on non-standard members of the family in an elementary extension. More precisely, we consider an elementary extension  $\mathfrak{R}$  of  $\langle \mathbb{R}_{om}, \Gamma \rangle$  where for every definable set Z in  $\langle \mathbb{R}, \Gamma \rangle$  we denote by  $Z^{\sharp}$  its realization in  $\mathfrak{R}$  (see Section 2.1 for details). Now every Hausdorff limit at  $\infty$  of  $\pi(\mathcal{F})$  is the standard part of  $\pi(F_{\tau}^{\sharp} \cdot \Gamma^{\sharp})$  for  $\tau > \mathbb{R}$  a non-standard parameter in  $\mathfrak{R}$  (see Section 6.1.1). Thus, the problem reduces to the study of sets of the form st $(F_{\tau}^{\sharp} \cdot \Gamma^{\sharp})$ .

Similarly to the answers to the closure problem, we associate to the family  $\mathcal{F}$  finitely many normal co-commutative subgroups  $L_i \subseteq G$ , and

then for every  $\Gamma$ , the answers depend on whether one of the  $L_i$  is a  $\Gamma$ dense subgroup or not. More precisely, (see Section 4.1 for details), let  $L^{\Gamma}$  be the smallest  $\Gamma$ -rational real algebraic subgroup of G containing L. We prove: (Theorem 6.12):

**Theorem** (see Theorem 6.12). Let G be a unipotent group,  $\mathcal{F} = \{F_t :$  $t \in (0,\infty)$  an  $\mathbb{R}_{om}$ -definable family of subsets of G.

Then, there exists a finite collection  $\mathcal{L}(\mathcal{F})$  of normal co-commutative subgroups of G, such that for every lattice  $\Gamma \subseteq G$  and  $\pi: G \to G/\Gamma$ , we have:

- (1)  $L^{\Gamma} = G$  for some  $L \in \mathcal{L}(\mathcal{F})$  if and only if  $\pi(\mathcal{F})$  converges strongly to  $G/\Gamma$  at  $\infty$  (i.e.  $G/\Gamma$  is the only Hausdroff limit at  $\infty$  of  $\pi(\mathcal{F})$ and this remains true for every lattice in G commensurable with  $\Gamma$ ).
- (2)  $L^{\Gamma} \neq G$  for all  $L \in \mathcal{L}(\mathcal{F})$  if and only if there exists a subgroup  $\Gamma_0 \subseteq \Gamma$  of finite index such that all Hausdorff limits at  $\infty$  of  $\pi_0(\mathcal{F})$ are proper subsets of  $G/\Gamma_0$  (here  $\pi_0: G \to G/\Gamma_0$  is the quotient map).

Note that the theorem above does not identify all the possible Hausdorff limits of families  $\pi(\mathcal{F})$  in  $G/\Gamma$ . However, we can do it when G is a abelian and  $\mathcal{F}$  is a family of polynomial dilations with no constant term (see 7). We prove:

**Theorem** (see Corollary 7.6). Let  $\{\rho_t \colon \mathbb{R}^k \to \mathbb{R}^m \colon t \in (0,\infty)\}$  be a family of polynomial dilations with no constant term, and  $X \subseteq \mathbb{R}^k$  an  $\mathbb{R}_{om}$ -definable set.

Then there are linear subspaces  $L_1, \ldots, L_n \subseteq \mathbb{R}^n$ , and there is a coset of a linear space  $\bar{c} + V \subseteq (\mathbb{R}^m)^n$  such that set of Hausdorff limits at  $\infty$ of the family  $\{\pi_{\Gamma} \circ \rho_t(X) : t \in (0, \infty)\}$  is exactly the family

$$\left\{ \pi_{\Gamma} \left( \bigcup_{i=1}^{n} (d_i + L_i^{\Gamma}) \right) : (d_1, \dots, d_n) \in \bar{c} + V^{\Gamma^n} \right\}.$$

In particular, it is the projection under  $\pi_{\Gamma}$  of a definable family of subsets of  $\mathbb{R}^n$ .

Our work on dilations was motivated by [3], of Kra, Shah and Sun.

The structure of the paper. From a model theoretic point of view, the main complexity of this work over the closure theorems in [9] is the fact that we study sets defined over  $\mathbb{R}\langle \tau \rangle$ , where  $\tau$  is a non-standard parameter as above. This requires several adjustments to our previous work in [8] and [9]. In Section 2 we develop the notion of short and long types (which replace bounded and unbounded types over  $\mathbb{R}$ ). In addition, we modify the theory of  $\mu$ -stabilizers developed in [7], so it fits our setting. In Section 3 we study types and their nearest co-commutative subgroups (again, the results in [9] need adjustments since the types are over  $\mathbb{R}\langle\tau\rangle$ ). In Section 4, lattices come in and we prove the main theorems about the  $\Gamma$ -closure of types. In Section 5, we study definable sets over  $\mathbb{R}\langle\tau\rangle$  and formulate conditions under which such sets are  $\Gamma$ -dense. In Section 6 we translate the results obtained thus far back to the original problem of Hausdorff limits, and in Section 7 we study in more details families given by polynomial dilations.

**Acknowledgements.** We thank Amos Nevo for suggesting this problem and explaining its ergodic theoretic origins.

## 2. Long types and $\mu$ -stabilizers

2.1. **Model theoretic preliminaries.** As background on model theory and o-minimality we refer to [13] and [4]. We follow the set-up from [8, Section 2] and [7, Section 2.3].

We fix an o-minimal structure  $\mathbb{R}_{om} = \langle \mathbb{R}, <, +, \cdot, \cdots \rangle$  expanding the real field and denote by  $\mathcal{L}_{om}$  its language. For convenience we add to  $\mathcal{L}_{om}$  a constant symbol for every real number.

We use  $\mathcal{L}_{\text{full}}$  for a language in which every subset of  $\mathbb{R}^n$ ,  $n \in \mathbb{N}$ , has a predicate symbol, and denote the corresponding structure on  $\mathbb{R}$  by  $\mathbb{R}_{\text{full}}$ . This will allow us to talk about lattices as definable sets.

We let  $\mathfrak{R}_{\text{full}} = \langle \mathfrak{R}, < \ldots \rangle$  be an elementary extension of  $\mathbb{R}_{\text{full}}$  which is  $|\mathbb{R}|^+$ -saturated and strongly- $|\mathbb{R}|^+$ -homogeneous, and let  $\mathfrak{R}_{\text{om}}$  be the reduct to  $\mathcal{L}_{\text{om}}$ . Clearly,  $\mathfrak{R}_{\text{om}}$  is an elementary extensions of  $\mathbb{R}_{\text{om}}$ .

We use Roman letters X, Y, Z to denote subsets of  $\mathbb{R}^n$  and let  $X^{\sharp}, Y^{\sharp}, Z^{\sharp}$  denote their realizations in  $\mathfrak{R}_{\text{full}}$ . We use script  $\mathcal{X}$  to denote subsets of  $\mathfrak{R}^n$  which are not necessarily of the form  $X^{\sharp}$ . When we write  $A \subseteq \mathfrak{R}$ , for a parameter set over which definable sets and types are considered, we mean that  $|A| \leq |\mathbb{R}|$ .

For  $\mathcal{L} = \mathcal{L}_{om}$  or  $\mathcal{L} = \mathcal{L}_{full}$ , as usual, a complete  $\mathcal{L}$ -type over A is an ultrafilter on sets which are  $\mathcal{L}$ -definable using parameters in A. For  $A \subseteq \mathfrak{R}$  and  $\mathcal{X} \subseteq \mathfrak{R}^n$  an  $\mathbb{R}_{om}$ -definable set over A, we let  $S_{\mathcal{X}}(A)$  be the collection of all complete  $\mathcal{L}_{om}$ -types over A, containing the set  $\mathcal{X}$ . If  $\mathcal{X} = X^{\sharp}$  for some  $\mathcal{L}_{om}$  definable  $X \subseteq \mathbb{R}^n$  then instead of  $S_{X^{\sharp}}(A)$  we write  $S_X(A)$ . For  $p \in S_{\mathcal{X}}(A)$  we let  $p(\mathfrak{R})$  denote the set of its realizations in  $\mathfrak{R}_{om}$ .

Unless otherwise stated, by "definable" we mean " $\mathcal{L}_{om}$ -definable". In particular dcl denotes the definable closure in the structure  $\mathfrak{R}_{om}$ . Note that by our assumptions,  $\mathcal{L}_{om}$  contains constant symbols for real numbers, hence, by definability of Skolem functions, for any set  $A \subseteq \mathfrak{R}$ ,

the definable closure dcl(A) is an elementary substructure of  $\mathfrak{R}_{om}$  which contains  $\mathbb{R}_{om}$  as an elementary substructure.

A type-definable subset of  $\mathfrak{R}^n$ , over A, is the intersection of (possibly infinitely many) definable sets over A, which by our convention means  $\mathcal{L}_{om}$ -definable sets. The notion of a  $\mathcal{L}_{full}$  type-definable set is similarly defined. Since  $|A| \leq |\mathbb{R}|$ , every collection of such definable sets is bounded in size. A subset of  $\mathfrak{R}^n$  is said to be type-definable if it is type-definable over some  $A \subseteq \Re$ .

We let  $\mathcal{O}$  be the convex hull of  $\mathbb{R}$  in  $\mathfrak{R}$ , namely,

$$\mathcal{O} = \{ \alpha \in \mathfrak{R} : \exists r \in \mathbb{R}^{>0} \, |\alpha| < r \},\$$

It is a valuation ring of  $\Re$ , whose associated maximal ideal is

$$\mathbf{m} = \{ \alpha \in \mathfrak{R} : \forall r \in \mathbb{R}^{>0} \, |\alpha| < r \}.$$

The ring homomorphism  $\mathcal{O} \to \mathcal{O}/\mathbf{m}$  restricts to an isomorphism between  $\mathbb{R}$  and  $\mathcal{O}/\mathbf{m}$ . The corresponding ring homomorphism st:  $\mathcal{O} \to \mathcal{O}$  $\mathbb{R} \simeq \mathcal{O}/\mathbf{m}$  is called the standard part map, and we extend it coordinatewise to st :  $\mathcal{O}^n \to \mathbb{R}^n$ . For  $\mathcal{X} \subseteq \mathfrak{R}^n$ , we write st( $\mathcal{X}$ ) instead of st( $\mathcal{X} \cap$  $\mathcal{O}^n$ ).

For  $a \in \mathbb{R}^n$  and r > 0, we let  $B_r(a) = \{x \in \mathbb{R}^n : |x - a| < r\}$ . We need the following lemma.

- **Lemma 2.1.** (1) If  $\mathcal{X} \subseteq \mathfrak{R}^n$  is an  $\mathcal{L}_{\text{full}}$  type-definable set then  $\operatorname{st}(\mathcal{X})$ is a closed subset of  $\mathbb{R}^n$ .
- (2) For a set  $X \subseteq \mathbb{R}^n$ , we have  $\operatorname{cl}(X) = \operatorname{st}(X^{\sharp})$  (where  $\operatorname{cl}(X)$  is the topological closure of X).
- (3) Let  $\Sigma$  be a collection of  $\mathcal{L}_{full}$ -definable subsets of  $\mathfrak{R}^n$  with  $|\Sigma| \leq |\mathbb{R}|$ . If  $\Sigma$  is closed under finite intersections, then

$$\operatorname{st}\left(\bigcap \Sigma\right) = \bigcap_{\mathcal{X} \in \Sigma} \operatorname{st}(\mathcal{X}).$$

In particular  $\operatorname{st}(\bigcap \Sigma)$  is closed.

*Proof.* (1) If  $a \in \operatorname{cl}(\operatorname{st}(\mathcal{X}))$  then for every  $r \in \mathbb{R}^{>0}$ ,  $B_r(a)^{\sharp} \cap \operatorname{st}(\mathcal{X}) \neq \emptyset$ and therefore also for every  $r \in \mathbb{R}^{>0}$ ,  $B_r(a)^{\sharp} \cap \mathcal{X} \neq \emptyset$ . By saturation, there is  $b \in \mathcal{X}$  such that  $b \in a + \mathbf{m}$ , so  $a \in \operatorname{st}(\mathcal{X})$ .

(2) is easy and (3) is just 
$$[8, \text{Claim } 3.1]$$

The following standard fact is easy to prove.

Fact 2.2. Let  $X \subseteq \mathbb{R}^m$  and  $Y \subseteq \mathbb{R}^n$  be closed subsets and  $f: X \to Y$ a continuous function. Let  $\operatorname{st}_m \colon \mathcal{O}^m \to \mathbb{R}^m$  and  $\operatorname{st}_n \colon \mathcal{O}^n \to \mathbb{R}^n$  be the corresponding standard part maps. For every  $\gamma \in \mathcal{O}^m \cap X^{\sharp}$  we have  $f(\operatorname{st}_m(\gamma)) = \operatorname{st}_n(f(\gamma))$ . In particular, for every  $\mathcal{X} \subseteq X^{\sharp}$  we have  $f(\operatorname{st}_m(\mathcal{X})) \subseteq \operatorname{st}_n(f(\mathcal{X})).$ 

We will need the following important result:

Fact 2.3 ([14, Proposition 8.1]). If  $\mathcal{X}$  is definable in  $\mathfrak{R}_{om}$  then  $st(\mathcal{X})$  is definable in  $\mathbb{R}_{om}$ .

Let  $G \subseteq \mathbb{R}^m$  be an  $\mathbb{R}_{om}$ -definable group, namely the universe of G and the group operation are definable in  $\mathbb{R}_{om}$ . For example, any real algebraic, or more generally semi-algebraic group is definable in  $\mathbb{R}_{om}$  (for more on definable groups in o-minimal structures see [6]). The group G can be endowed with a group topology with a definable basis (see [10]). This topology might disagree with the natural o-minimal topology, coming from the fact that G is a subset of  $\mathbb{R}^m$ . However, as we observed in [7, Claim 3.1], we may embed G definably as a closed subset of  $\mathbb{R}^n$  for some n, such that the above group topology agrees with the induce  $\mathbb{R}^n$ -topology. Thus, whenever  $G \subseteq \mathbb{R}^n$  is an  $\mathbb{R}_{om}$ -definable group we assume it to be closed in  $\mathbb{R}^n$ , and in addition assume that the Euclidean topology makes G a topological group.

We will use very often the following.

Fact 2.4 ([10]). Let G be an  $\mathbb{R}_{om}$ -definable group.

- (1) If  $H \subseteq G$  is an  $\mathbb{R}_{om}$ -definable subgroup then H is closed in G.
- (2) If H is an  $\mathbb{R}_{om}$ -definable group and  $f: G \to H$  an  $\mathbb{R}_{om}$ -definable homomorphism then f is continuous.

For  $G \subseteq \mathbb{R}^n$  as above, we consider two distinguished subgroups of  $G^{\sharp}$ . The first is the infinitesimal group  $\mu_G$ , defined as follows:

$$\mu_G = \bigcap \{X^{\sharp} : X \subseteq G \text{ an } \mathbb{R}_{om}\text{-definable open neighborhood of } e\}.$$

It is a type-definable subgroup and under our assumption on G it equals  $e + \mathbf{m}^n \cap G^{\sharp}$ , with  $\mathbf{m} \subseteq \mathcal{O} \subseteq \mathfrak{R}$  the infinitesimal ideal defined above. The second subgroup is  $\mathcal{O}_G$ , defined by:

$$\mathcal{O}_G = \bigcup \{X^{\sharp} : X \subseteq G \text{ an } \mathbb{R}_{om}\text{-definable compact neighborhood of } e\}.$$

 $\mathcal{O}_G$  is a V-definable (or Ind-definable) subgroup of  $G^{\sharp}$ , which equals, under our assumptions on G, to  $\mathcal{O}^n \cap G^{\sharp}$ . In particular,  $G \subseteq \mathcal{O}_G$ . The group  $\mu_G$  is a normal subgroup of  $\mathcal{O}_G$  and the latter can be written as the semi-direct product  $\mathcal{O}_G = \mu_G \rtimes G$ . We identify the quotient  $\mathcal{O}_G/\mu_G$  with G and call the quotient map  $\operatorname{st}_G : \mathcal{O}_G \to G$  the standard part map. As before, we extend it coordinate-wise to  $\operatorname{st}_G : \mathcal{O}_G^n \to G^n$ . By our assumptions on G, for every  $a \in \mathcal{O}_G$ , we have  $\operatorname{st}(a) = \operatorname{st}_G(a) \in G$ . In particular, Lemma 2.1 holds if one restricts to subsets of G and to  $\operatorname{st}_G$ .

When the underlying group G is fixed we omit the subscript G and just use  $\mathcal{O}$ ,  $\mu$  and st.

As before, given an arbitrary set  $\mathcal{X} \subseteq G^{\sharp}$ , we let  $\operatorname{st}(\mathcal{X})$  denote the set st $(\mathcal{X} \cap \mathcal{O}_G) \subseteq G$ .

**Fact 2.5.** Assume that  $G_1, G_2$  are definable in  $\mathbb{R}_{om}$ . Then,

- (1)  $\mu_{G_1 \times G_2} = \mu_{G_1} \times \mu_{G_2}$  and  $\mathcal{O}_{G_1 \times G_2} = \mathcal{O}_{G_1} \times \mathcal{O}_{G_2}$ .
- (2) If  $f: G_1 \to G_2$  is an  $\mathbb{R}_{om}$ -definable surjective homomorphism then  $f(\mu_{G_1}) = \mu_{G_2}$  and  $f(\mathcal{O}_{G_1}) = \mathcal{O}_{G_2}$

*Proof.* (1) follows from the fact that the topology on  $G_1 \times G_2$  is the product topology. For (2), see [9, Lemma 3.10].

2.2. Long and short sets. Let  $G \subseteq \mathbb{R}^n$  be an  $\mathbb{R}_{om}$ -definable group.

**Definition 2.6.** A subset  $\mathcal{X} \subseteq G^{\sharp}$  is called *left-short in G* if there exists some compact set  $K \subseteq G$  and  $g \in G^{\sharp}$  such that  $\mathcal{X} \subseteq K^{\sharp} \cdot g$  (since G can be written as an increasing union of relatively compact  $\mathbb{R}_{om}$ -definable open sets, we may always take K to be  $\mathbb{R}_{om}$ -definable).

Otherwise,  $\mathcal{X}$  is called *left-long in G*. For  $A \subseteq \mathfrak{R}$ , we say that a type  $p \in S_G(A)$  is left-short (left-long) in G if  $p(\mathfrak{R})$  is left-short (left-long) in G.

The following are easy to verify:

Lemma 2.7. Given  $\mathcal{X} \subseteq G^{\sharp}$ ,

- (1)  $\mathcal{X}$  is left-short in G if and only if  $\mathcal{X} \cdot \mathcal{X}^{-1} \subseteq K^{\sharp}$  for some compact set  $K \subseteq G$ .
- (2) For every  $g \in G^{\sharp}$ ,  $\mathcal{X}$  is left-short in G if and only if  $\mathcal{X}g$  is left-short
- (3) For every  $g \in G$ ,  $\mathcal{X}$  is left-short in G if and only if  $g\mathcal{X}$  is left-short
- (4) If  $\mathcal{X} = X^{\sharp}$ , for  $X \subseteq G$  an  $\mathbb{R}_{\text{full}}$ -definable set, then  $\mathcal{X}$  is left-short in G if and only if X is bounded in  $\mathbb{R}^n$ .

We may similarly define right-short and right-long in G and in general these notions are different. However, for the rest of the paper we use *short* and *long* to refer only to left-short and left-long.

If  $H \subseteq G$  is an  $\mathbb{R}_{om}$ -definable subgroup and  $\mathcal{X}$  a subset of  $H^{\sharp}$  then, by Lemma 2.7(1),  $\mathcal{X}$  is short in H if and only if it is short in G, hence we omit the reference to the group when the context is clear.

Note that by saturation, an  $\mathcal{L}_{\text{full}}$  type-definable set  $\mathcal{X} \subseteq G^{\sharp}$  is short if and only if  $\mathcal{X} \subseteq \mathcal{O} \cdot q$  for some  $q \in G^{\sharp}$ .

**Lemma 2.8.** Let  $G_1, G_2, G$  be  $\mathbb{R}_{om}$ -definable groups.

- (1) If  $\mathcal{X} \subseteq G_1^{\sharp}$  is short and  $f: G_1 \to G_2$  is an  $\mathbb{R}_{om}$ -definable homomorphism then  $f(\mathcal{X})$  is short in  $G_2$ .
- (2) If  $\mathcal{X}_1 \subseteq G_1^{\sharp}$  is short and  $\mathcal{X}_2 \subseteq G_2^{\sharp}$  is short then  $\mathcal{X}_1 \times \mathcal{X}_2$  is short in  $G_1 \times G_2$ .
- (3) If  $H_1$ ,  $H_2$  are two normal  $\mathbb{R}_{om}$ -definable subgroups of G and  $\mathcal{X} \subseteq G^{\sharp}$  an arbitrary set then the image of  $\mathcal{X}$  in  $G^{\sharp}/(H_1^{\sharp} \cap H_2^{\sharp})$  is short if and only if its images in  $G^{\sharp}/H_1^{\sharp}$  and in  $G^{\sharp}/H_2^{\sharp}$  are short.

*Proof.* (1) and (2) are immediate since the image of a compact set under a  $\mathbb{R}_{om}$ -definable homomorphism is compact, and similarly the direct product of such sets is compact.

For (3), let  $\pi_i: G \to G/H_i$ , i=1,2, be the natural projections, and let  $\pi: G \to G/H_1 \times G/H_2$  be the map  $\pi(g) = (\pi_1(g), \pi_2(g))$ . The kernel of  $\pi$  is  $H_1 \cap H_2$  hence the image is isomorphic to  $G/(H_1 \cap H_2)$ . If  $\pi_1(\mathcal{X})$  and  $\pi_2(\mathcal{X})$  are both short then by (2), so is  $\pi_1(\mathcal{X}) \times \pi_2(\mathcal{X}) \subseteq G^{\sharp}/H_1^{\sharp} \times G^{\sharp}/H_2^{\sharp}$ . But  $\pi(\mathcal{X})$  is contained in  $\pi_1(\mathcal{X}) \times \pi_2(\mathcal{X})$  so also short. The converse follows from (1) using the natural homomorphisms from  $G/(H_1 \cap H_2)$  onto  $G/H_i$ , i=1,2.

# **Lemma 2.9.** For $A \subseteq \mathfrak{R}$ , and $p \in S_G(A)$ , we have:

- (1) p is short in G if and only if there exists  $a \in dcl(A) \cap G^{\sharp}$  such that  $p \vdash \mu \cdot a$ , namely  $p(\mathfrak{R}) \subseteq \mu \cdot a$ .
- (2) Let  $H \subseteq G$  be a  $\mathbb{R}_{om}$ -definable normal subgroup and  $\pi : G \to G/H$  the quotient map. Then  $\pi(p) \in S_{G/H}(A)$  is short in G/H if and only if there exists  $a \in dcl(A) \cap G^{\sharp}$  such that  $p(\mathfrak{R}) \subseteq \mu \cdot aH^{\sharp}$ .
- Proof. (1) Assume that p is short. Hence  $p(\mathfrak{R}) \cdot p(\mathfrak{R})^{-1} \subseteq K^{\sharp}$  for some  $\mathbb{R}_{om}$ -definable compact set  $K \subseteq G$ . By logical compactness, there exists an A-definable set  $\mathcal{X}$  in p such that  $\mathcal{X} \cdot \mathcal{X}^{-1} \subseteq K$ . By definability of Skolem functions in o-minimal structures, the set  $\mathcal{X}$  contains a point  $a \in dcl(A)$ . Consider the complete  $\mathbb{R}_{om}$ -type over A,  $p \cdot a^{-1}$ . We have  $p(\mathfrak{R}) \cdot a^{-1} \subseteq \mathcal{X} \cdot a^{-1} \subseteq K^{\sharp}$ . Let  $\beta \models p \cdot a^{-1}$  and  $g = st(\beta)$  (this is defined since  $\beta \in \mathcal{O}$ ). Because  $p \cdot a^{-1}$  is a complete A-type and  $\beta \in \mu \cdot g$ , we have  $p \cdot a^{-1} \vdash \mu \cdot g$ . Hence  $p(\mathfrak{R}) \subseteq \mu \cdot g \cdot a$ . Clearly,  $g \cdot a \in dcl(A)$ .

The converse is clear.

For (2), notice that if p is short then by Lemma 2.8(1),  $\pi(p)$  is short and hence by (1) there exists  $b \in \operatorname{dcl}(A) \cap (G/H)^{\sharp}$  such that  $\pi(p(\mathfrak{R})) \in \mu_{G/H} \cdot b$ .

We now take any  $a \in dcl(A)$  in the A-definable set  $\pi^{-1}(b)$ , and we have  $p(\mathfrak{R}) \subseteq \pi^{-1}(\mu_{G/H} \cdot b) \subseteq \mu_G \cdot aH^{\sharp}$ .

For the converse, notice that by Fact 2.5,  $\pi(\mu_G) = \mu_{G/H}$ . Hence,  $\pi(\mu_G \cdot a) = \mu_{G/H} \cdot \pi(a)$ , so if  $p(\mathfrak{R}) \subseteq \mu_G \cdot aH^{\sharp}$  then  $\pi(p)(\mathfrak{R}) \subseteq \mu_{G/H} \cdot \pi(a)$  is short.

2.3. The  $\mu$ -stabilizer of a type. We fix an  $\mathbb{R}_{om}$ -definable group G. In [7] we developed a theory for  $\mu$ -stabilizers of types over  $\mathbb{R}$ . Here we take a more general viewpoint which we now explain.

Consider the set  $S_G(A)$ . Given  $p \in S_G(A)$  we let  $\mu \cdot p$  (below written as  $\mu p$ ) denote the partial type over A whose realization is the set  $\mu \cdot p(\mathfrak{R})$ .

Remark 2.10. We note that when we consider here types over arbitrary  $A \subseteq \mathfrak{R}$ , then, unlike [7], we still keep  $\mu = \mu_G$  fixed and not change it to a smaller infinitesimal group (namely, the intersection of all A-definable open neighborhoods of e).

Since our point of view here is slightly different from [7], we go briefly through the results we need and explain how their proofs differ from the analogous results in [7].

Given  $p, q \in S_G(A)$ , we say that p and q are  $\mu$ -equivalent,  $p \sim_{\mu} q$ , if  $\mu p = \mu q$ , i.e.  $\mu p(\mathfrak{R}) = \mu q(\mathfrak{R})$ .

**Fact 2.11.** For  $p, q \in S_G(A)$ , the following are equivalent:

- (1)  $p \sim_{\mu} q$
- (2)  $\mu p(\mathfrak{R}) \cap \mu q(\mathfrak{R}) \neq \emptyset$ .

(see [7, Claim 2.7] for an identical argument).

It is easy to verify that if  $p \sim_{\mu} q$  then p is a long if and only if q is

Let  $S_G^{\mu}(A) = \{\mu p : p \in S_G(A)\}$ . The group G acts from the left on  $S_G^{\mu}(A)$  by  $g \cdot \mu p = \mu(gp)$ . The following subgroup plays a crucial role in our analysis:

**Definition 2.12.** Given  $p \in S_G(A)$ , the left stabilizer of  $\mu p$  is defined as:

$$\operatorname{Stab}^{\mu}(p) = \{ g \in G : g \cdot \mu p = \mu p \}.$$

Since the definition of  $\operatorname{Stab}^{\mu}(p)$  depends only on  $\mu p$ , if  $p \sim_{\mu} q$  then  $\operatorname{Stab}^{\mu}(p) = \operatorname{Stab}^{\mu}(q).$ 

Our main focus in [7] was on unbounded definable types. For  $\mathbb{R}_{om}$ definable groups, and  $\mathcal{L}_{om}$ -types over  $\mathbb{R}$  these are types which do not contain any formula over  $\mathbb{R}$  defining a compact subset of G. Since we are considering here types which are not only over  $\mathbb{R}$  our focus is shifted to long types.

Recall that for p an  $\mathcal{L}_{om}$ -type we let  $\dim(p)$  be the smallest o-minimal dimension of the formulas in p.

**Definition 2.13.** We say that a type  $p \in S_G(A)$  is  $\mu$ -reduced if for all  $q \in S_G(A)$ , if  $p \sim_{\mu} q$  then  $\dim(p) \leq \dim(q)$ .

Clearly, every  $p \in S_G(A)$  is  $\mu$ -equivalent to a  $\mu$ -reduced type in  $S_G(A)$ : just take a  $\mu$ -equivalent type of minimal dimension (but there might be more than one such). Notice that by Lemma 2.9, if p is short and  $\mu$ -reduced then dim p = 0 and  $p = \operatorname{tp}(a/A)$  for some  $a \in \operatorname{dcl}(A)$ .

Our main goal in this section is to prove:

## **Proposition 2.14.** Let $p \in S_G(A)$ . Then

- (1)  $\operatorname{Stab}^{\mu}(p)$  is  $\mathbb{R}_{om}$ -definable and can be written as  $\operatorname{st}(S \cdot \alpha^{-1})$ , for some definable S in p and  $\alpha \models p$ .
- (2) If p is a long type then  $\dim(\operatorname{Stab}^{\mu}(p)) > 0$ . Moreover, in this case  $\operatorname{Stab}^{\mu}(p)$  is a torsion-free solvable group.

The proof is very similar to the proof of [7, Theorem 3.10] so we only point out the differences. As we noted above, we may assume that p is  $\mu$ -reduced and if p is short that  $\mu p = \mu \cdot a$  for some  $a \in dcl(A)$  so  $\operatorname{Stab}^{\mu}(p)$  is trivial. Thus, we fix a long  $\mu$ -reduced type  $p \in S_G(A)$  and  $\alpha \in p(\mathfrak{R})$ .

We start with an analogue of [7, Claim 3.12]:

Claim 2.15. If  $Y \subseteq G^{\sharp}$  is A-definable and dim  $Y < \dim p$  then  $\mathcal{O} \cdot \alpha \cap Y = \emptyset$ .

*Proof.* Assume towards contradiction that  $\beta \in Y \cap \mathcal{O} \cdot \alpha$ . Then  $\beta \in \mu \cdot r\alpha$ , for some  $r \in G$ , hence  $r^{-1}\beta \in \mu p$ . But  $\dim(r^{-1}\beta/A) \leq \dim(Y) < \dim p$ , contradicting the fact that p is  $\mu$ -reduced.

Next, we note, just like [7, Claim 3.8], that for every A-definable set S in p,  $\operatorname{Stab}^{\mu}(p) \subseteq \operatorname{st}(S \cdot \alpha^{-1})$ . Indeed, if  $g \in \operatorname{Stab}^{\mu}(p)$  then there exists  $\beta \models p$  and  $\epsilon \in \mu$  such that  $g\alpha = \epsilon\beta$ . It follows that  $\beta \in S$  and  $\beta\alpha^{-1} \in \mathcal{O}$ , thus  $g = \operatorname{st}(\beta\alpha^{-1}) \in \operatorname{st}(S \cdot \alpha^{-1})$ .

The next claim is similar to [7, Claim 3.13].

Claim 2.16. There exists an A-definable set S in p such that every element in  $S \cap \mathcal{O} \cdot \alpha$  realizes p.

Let us explain the proof: As in [7], for every A-definable set S in p, the set  $S \cdot \alpha^{-1} \cap \mathcal{O}$  is a relatively definable subset of  $\mathcal{O}$ . Hence, by [7, Theorem B.2] it has finitely many connected components (see precise definition of connectedness there). We choose an A-definable such cell S in p with dim  $S = \dim(p)$ , for which the number of components of  $S \cdot \alpha^{-1} \cap \mathcal{O}$  is minimal. Using Claim 2.15, we can prove, just as in [7, Claim 3.13], that any  $\beta \in S \cap \mathcal{O} \cdot \alpha$  must realize p.

Finally, we prove an analogue of [7, Claim 3.14]:

Claim 2.17. For S as in Claim 2.16, we have  $\operatorname{Stab}^{\mu}(p) = \operatorname{st}(S \cdot \alpha^{-1})$ .

*Proof.* It is sufficient to show that  $\operatorname{st}(\mathcal{S} \cdot \alpha^{-1}) \subseteq \operatorname{Stab}^{\mu}(p)$ , so we take  $g \in \operatorname{st}(\mathcal{S} \cdot \alpha^{-1})$  and note that for some  $\epsilon \in \mu$ , we have  $\epsilon g \alpha \in \mathcal{S} \cap \mathcal{O} \cdot \alpha$ , so by our choice of  $\mathcal{S}$ ,  $\epsilon g \alpha \models p$ . It follows that  $\mu g p(\mathfrak{R}) \cap p(\mathfrak{R}) \neq \emptyset$ , so by Fact 2.11,  $g \cdot \mu p = \mu p$ .

Thus, by Fact 2.3,  $\operatorname{Stab}^{\mu}(p) = \operatorname{st}(\mathcal{S} \cdot \alpha^{-1})$  is definable.

Since p is long the set  $S \cdot \alpha^{-1}$  is not contained in  $\mathcal{O}$ , thus  $\operatorname{Stab}^{\mu}(p)$  is unbounded in G.

To see that  $\operatorname{Stab}^{\mu}(p)$  is solvable, torsion-free we repeat the argument from [7, Theorem 3.6]: By [7, Fact 3.25], G can be written as a product of two sets  $G = C \cdot H$ , with  $C \subseteq G$  a  $\mathcal{L}_{om}$ -definable compact set and H a  $\mathcal{L}_{om}$ -definable torsion-free solvable group. Thus,  $\alpha \in G^{\sharp}$  as above can be written as  $\alpha = \epsilon \cdot g \cdot h^*$  for  $\epsilon \in \mu_G, g \in C, h^* \in H^{\sharp}$ . It follows that  $\alpha \in \mu \cdot (H^g)^{\sharp} \cdot g$ , so  $\operatorname{tp}(\alpha/A)$  is  $\mu$ -equivalent to a type  $q \vdash (H^g)^{\sharp} \cdot g$ . But then  $\operatorname{Stab}^{\mu}(p) = \operatorname{Stab}^{\mu}(pg) \subseteq H^g$  so  $\operatorname{Stab}^{\mu}(p)$  is a torsion-free solvable group.

This ends the proof of Proposition 2.14.

**Remark 2.18.** In fact, the remainder of the proof of [7, Theorem 3.12] goes through identically and thus we could have proved the stronger result, saying that for p a  $\mu$ -reduced type over A, the dimension of  $\operatorname{Stab}^{\mu}(p)$  equals to  $\dim(p)$ . However, this will not be needed here.

## 3. Nearest cosets

We now assume again that G is a definable group in  $\mathbb{R}_{om}$ .

#### 3.1. Nearest co-commutative cosets.

**Definition 3.1.** Given a type  $p \in S_G(A)$ , an  $\mathbb{R}_{om}$ -definable subgroup  $H \subseteq G$  and  $a \in dcl(A) \cap G^{\sharp}$ , we say that the coset  $aH^{\sharp}$  is near p if  $p(\mathfrak{R}) \subseteq \mu \cdot aH^{\sharp}$ .

Sometimes we omit  $\sharp$ , write  $p \vdash \mu aH$ , and say that aH is near p.

Notice that in the above definition the subgroup H is defined over  $\mathbb{R}$ , but the element a is taken from  $dcl(A) \subseteq \mathfrak{R}$ .

**Remark 3.2.** By Lemma 2.9, a type  $p \in S_G(A)$  is short if and only if, for the trivial subgroup  $\{e\}$ , a coset  $a \cdot e$ , is near p.

Also, for a normal  $\mathbb{R}_{om}$ -definable subgroup  $H \subseteq G$ , some coset aH is near p if and only if the image of p in G/H is short.

**Lemma 3.3.** Let  $p \in S_G(A)$ ,  $H_1, H_2 \subseteq G$  be two  $\mathbb{R}_{om}$ -definable normal subgroups,  $a_1, a_2 \in dcl(A)$ , and assume both  $aH_1$  and  $aH_2$  are near p. Then there exists  $d \in dcl(A)$  such that the coset  $d(H_1 \cap H_2)$  is near p.

*Proof.* Let  $G_i = G/H_i$ , i = 1, 2, and  $\pi_i : G \to G_i$  the natural projection. Let  $f : G \to G_1 \times G_2$  be the definable homomorphism  $f(g) = (\pi_1(g), \pi_2(g))$ . We have  $\ker(f) = H_1 \cap H_2$ .

By Lemma 2.9(2), the images of  $p(\mathfrak{R})$  in both  $G/H_1$  and in  $G/H_2$  are short. Hence, by Lemma 2.8(3), its image in  $G/(H_1 \cap H_2)$  is also short. By Lemma 2.9 (2), there exists  $d \in \operatorname{dcl}(A)$  such that  $d(H_1 \cap H_2)$  is near p.

In the case of a unipotent group G and  $A = \mathbb{R}$ , the above lemma holds without assuming normality of  $H_1$  and  $H_2$  (see [9, Theorem 3.7]). Unfortunately, in general, this fails for an arbitrary A:

**Example 3.4.** We consider the Heisenberg group, identified with  $\mathbb{R}^3$ , as

$$[a, b, c] \cdot [d, e, f] = [a + d, b + e, ae + c + f].$$

We let  $H_1 = Z(G) = \{[0,0,x] : x \in \mathbb{R}\}$ , and  $H_2 = \{[0,t,t] : t \in \mathbb{R}\}$ . We now consider  $H_1^{\sharp}$  and  $H_2^{\sharp}$  in  $G^{\sharp}$ . Fix  $\tau \in \mathfrak{R}$  with  $\tau > \mathbb{R}$  and let  $A = \operatorname{dcl}(\tau)$ .

Consider the 1-type over dcl(A):

$$q(t) = \{ r < t < c : r \in \mathbb{R}, c \in \operatorname{dcl}(A) \text{ with } c > \mathbb{R} \}.$$

Let p(t) be the type over dcl(A) given by  $\{[\tau, 0, t] : t \models q\}$ . For  $\alpha = [\tau, 0, 0]$ , the realizations of p are contained in the coset  $\alpha H_1$ . It is easy to see that p is a long type and we claim that  $\alpha H_2$  is near p.

Indeed, consider the type  $q_0 = (1/\tau)q$ . It is also a 1-type over dcl(A), whose realizations are contained in  $\mu \subseteq \mathfrak{R}$ , and, for every  $\beta \models q_0$ , the element  $g_{\beta} = [0, \beta, \beta]$  is in  $H_2^{\sharp}$ . Now, for  $\varepsilon_{\beta} = [0, -\beta, -\beta] \in \mu_G$ , we have

$$\varepsilon_{\beta} \cdot \alpha \cdot g_{\beta} = [0, -\beta, -\beta] \cdot [\tau, \beta, \tau\beta + \beta] = [\tau, 0, \tau\beta] \models p,$$

and also  $\varepsilon_{\beta} \cdot \alpha \cdot g_{\beta} \in \mu_{G} \cdot \alpha \cdot H_{2}^{\sharp}$ . Hence the coset  $\alpha H_{2}$  is near p.

Thus both  $\alpha H_1$  and  $\alpha H_2$  are near p. However, since p is long, a coset of  $H_1 \cap H_2 = \{e\}$  can not be near p.

The main part of this paper deals with unipotent groups, and, in the unipotent case, instead of nearest cosets, as in [9], it is more convenient to work with nearest co-commutative cosets.

**Definition 3.5.** We say that a subgroup  $H \subseteq G$  is *co-commutative* if it is normal and the quotient G/H is abelian (equivalently H contains [G,G]).

Since the intersection of two co-commutative subgroups is co-commutative, using Lemma 3.3, we may conclude:

Corollary 3.6. Given  $p \in S_G(A)$  there exists a smallest (by inclusion)  $\mathbb{R}_{om}$ -definable co-commutative subgroup  $L \subseteq G$  such that for some  $a \in dcl(A)$  the coset aL is near p.

We can now define:

**Definition 3.7.** Given  $p \in S_G(A)$ , a nearest co-commutative coset to p is a coset of the form aL, where  $a \in dcl(A)$  and  $L \subseteq G$  is an  $\mathbb{R}_{om}$ -definable co-commutative subgroup as in Corollary 3.6. It is unique up to  $\mu_G$ , namely if  $a_1L_1$  and  $a_1L_1$ , are both nearest co-commutative cosets to p then  $L_1 = L_2$  and  $\mu_G \cdot a_1L_1^{\sharp} = \mu_G \cdot a_2L_2^{\sharp}$ .

We will denote this subgroup L as  $L_p$ .

We now prove some basic properties of nearest co-commutative cosets.

**Lemma 3.8.** Let  $p \in S_G(A)$ . If  $K \subseteq G$  is a compact definable set,  $a \in \operatorname{dcl}(A)$  and  $p(\mathfrak{R}) \subseteq K^{\sharp} \cdot a \cdot L^{\sharp}$  for some  $\mathbb{R}_{om}$ -definable co-commutative  $L \subseteq G$  then  $L_p \subseteq L$ .

*Proof.* Clearly  $\pi(p)$  is short in G/L, where  $\pi: G \to G/L$  is the quotient map. By Lemma 2.9(2), there exists  $a' \in \operatorname{dcl}(A)$  such that  $a' \cdot L$  is near p, hence  $L_p \subseteq L$ .

**Lemma 3.9.** Let  $f: G \to H$  be an  $\mathbb{R}_{om}$ -definable surjective homomorphism of definable groups and  $A \subseteq \mathfrak{R}$ . For a type  $p \in S_G(A)$  and q = f(p), if  $D_p = aL_p$  is a nearest co-commutative coset to p then  $f(D_p)$  is a nearest co-commutative coset to q, and in particular,  $L_q = f(L_p)$ .

*Proof.* Since f is surjective, it maps a co-commutative subgroup onto a co-commutative subgroup.

Let  $D_q$  be a nearest co-commutative coset to q. It is sufficient to see that  $\mu_H D_q = \mu_H f(D_p)$ . We have  $p \vdash \mu_G D_p$ , so by Fact 2.5,  $q \vdash \mu_H f(D_p)$ , hence  $D_q \subseteq \mu_H f(D_p)$ . Conversely, since  $q \vdash \mu_H D_q$  then  $p \vdash \mu_G f^{-1}(D_q)$ , so  $D_p \subseteq \mu_G f^{-1}(D_q)$ , hence  $f(D_p) \subseteq \mu_H D_q$ .

**Lemma 3.10.** For  $p \in S_G(A)$ , let  $H \subseteq G$  be the  $\mu$ -stabilizer of p, and let  $aL_p$  be a nearest co-commutative coset to p. Then  $H \subseteq L_p$ .

*Proof.* Fix  $\beta \models p$ . Then by assumption, there exists  $\varepsilon \in \mu$  and  $\ell \in L_p^{\sharp}$  such that  $\beta = \varepsilon a \ell$ . Given  $h \in H$ , we have  $h\beta \in \mu p(\mathfrak{R})$ , hence  $h\beta = \varepsilon' a \ell'$ , with  $\varepsilon' \in \mu$  and  $\ell' \in L_p^{\sharp}$ . Thus

$$h = h\beta\beta^{-1} = \varepsilon' a\ell' \ell^{-1} a^{-1} \varepsilon^{-1}.$$

Since  $L_p$  is normal in G (so  $L_p^{\sharp}$  normal in  $G^{\sharp}$ ), it follows that  $h \in \mu \cdot L_p^{\sharp}$ . However, h is in G and  $L_p$  is closed in G, therefore  $h \in L_p$ . 3.2. The set  $\mathcal{L}_{\max}(\mathcal{X})$ . Again we fix a group G definable in  $\mathbb{R}_{\text{om}}$ .

Recall that by our assumption,  $G \subseteq \mathbb{R}^n$  is a closed subset. For r > 0, we will denote by  $\overline{B}_r \subseteq G$  the set  $\overline{B}_r(e) \cap G$ , where  $\overline{B}_r(e)$  is the closed ball of radius r centered at e. Clearly, each  $\overline{B}_r$  is a compact subset of G, definable in  $\mathbb{R}_{om}$ , with  $\mu_G = \bigcap_{r \in \mathbb{R}} \overline{B}_r^{\sharp}$ .

**Lemma 3.11.** For  $A \subseteq B \subseteq \mathfrak{R}$ , let  $\mathcal{X} \subseteq G$  be a set  $\mathcal{L}_{om}$ -definable over A, and  $p \in S_{\mathcal{X}}(A)$ .

- (1) If  $q \in S_{\mathcal{X}}(B)$  is an extension of p then  $L_q \subseteq L_p$ .
- (2) There is  $q \in S_{\mathcal{X}}(B)$  extending p such that  $L_q = L_p$ .

*Proof.* (1). Choose  $a_p \in \operatorname{dcl}(A)$  such that  $p(\mathfrak{R}) \subseteq \mu \cdot a_p \cdot L_p^{\sharp}$ . We have  $q(\mathfrak{R}) \subseteq p(\mathfrak{R}) \subseteq \mu \cdot a_p \cdot L_p^{\sharp}$ . Hence the coset  $a_p L_p$  is near q and  $L_q \subseteq L_p$ .

(2). Let  $\mathcal{Q}$  be the set of all  $q \in S_{\mathcal{X}}(B)$  extending p. For each  $q \in \mathcal{Q}$  we choose  $b_q \in \operatorname{dcl}(B)$  such that  $q(\mathfrak{R}) \subseteq \mu \cdot b_q \cdot L_q^{\sharp}$ . We have

$$p(\mathfrak{R})\subseteq\bigcup_{q\in\mathcal{Q}}q(\mathfrak{R})\subseteq\bigcup_{q\in\mathcal{Q}}\mu\cdot b_q\cdot L_q^\sharp\subseteq\bigcup_{q\in\mathcal{Q}}\overline{B}_1^\sharp\cdot b_q\cdot L_q^\sharp.$$

Thus the type definable set  $p(\mathfrak{R})$  is covered by a bounded family of definable sets of the form  $\overline{B}_1^{\sharp} \cdot b_q \cdot L_q^{\sharp}$ . Hence, by logical compactness, we can find a set  $\mathcal{X}_0 \in p$ , definable over A, and a finite subset  $\mathcal{Q}_0 \subseteq \mathcal{Q}$  such that

$$\mathcal{X}_0 \subseteq \bigcup_{q \in \mathcal{Q}_0} \overline{B}_1^{\sharp} \cdot b_q \cdot L_q^{\sharp}.$$

Since dcl(A) is an elementary substructure of  $\mathfrak{R}_{om}$ , we can find  $a_q \in dcl(A)$ , for each  $q \in \mathcal{Q}_0$ , such that  $\mathcal{X}_0 \subseteq \bigcup_{q \in \mathcal{Q}_0} \overline{B}_1^{\sharp} \cdot a_q \cdot L_q^{\sharp}$ . Since p is a complete over A, there is  $q \in \mathcal{Q}_0$  with  $p(\mathfrak{R}) \subseteq \overline{B}_1^{\sharp} \cdot a_q \cdot L_q^{\sharp}$ . By Lemma 3.8,  $L_p \subseteq L_q$ , hence by (1), we have  $L_p = L_q$ .

For  $A \subseteq \mathfrak{R}$  and a set  $\mathcal{X} \subseteq \mathfrak{R}^n$  definable over A, we denote by  $\mathscr{L}_A(\mathcal{X})$  the set

$$\mathscr{L}_A(\mathcal{X}) = \{L_p \colon p \in S_{\mathcal{X}}(A)\}.$$

Corollary 3.12. For  $A \subseteq B \subseteq \mathfrak{R}$ , let  $\mathcal{X} \subseteq G^{\sharp}$  be definable over A. Then,

- (1)  $\mathscr{L}_A(\mathcal{X}) \subseteq \mathscr{L}_B(\mathcal{X})$ .
- (2) An  $\mathbb{R}_{om}$ -definable co-commutative subgroup L of G is maximal (by inclusion) in  $\mathcal{L}_A(\mathcal{X})$  if an only if it is maximal in  $\mathcal{L}_B(\mathcal{X})$

*Proof.* Follows from Lemma 3.11.

Remark 3.13. In general, for  $A \subseteq B$  we do not have equality of sets,  $\mathscr{L}_A(\mathcal{X}) = \mathscr{L}_B(\mathcal{X})$ . As an example, consider the group  $G = (\mathbb{R}^2, +)$  with  $\mathcal{X} = \{(x, y) \in \mathfrak{R}^2 \colon x \geq 0, y = x^2\}$ . For  $A = \mathbb{R}$  there is only one unbounded type in  $S_{\mathcal{X}}(A)$ , whose a nearest co-commutative coset is the whole  $\mathbb{R}^2$ . Thus  $\mathcal{L}_A(\mathcal{X}) = \{\{0\}, \mathbb{R}^2\}$ . However it is not hard to see that in any proper elementary extension B of  $\mathbb{R}$  there are types in  $S_{\mathcal{X}}(B)$  whose nearest co-commutative cosets are translates of  $L = \{0\} \times \mathbb{R}$ , and  $\mathscr{L}_B(\mathcal{X}) = \{\{0\}, L, \mathbb{R}^2\}$ .

**Definition 3.14.** For  $\mathcal{X} \subseteq G^{\sharp}$  an  $\mathcal{L}_{om}$ -definable set over A, we denote by  $\mathscr{L}_{max}(\mathcal{X})$  the set of maximal subgroups, by inclusion, in  $\mathscr{L}_{A}(\mathcal{X})$ . By Corollary 3.12, it does not depend on A.

We now have:

**Theorem 3.15.** Let G be an  $\mathbb{R}_{om}$ -definable group,  $A \subseteq \mathfrak{R}$ , and let  $\mathcal{X} \subseteq G^{\sharp}$  be  $\mathcal{L}_{om}$ -definable over A.

For every  $r \in \mathbb{R}^{>0}$ , there are definable co-commutative subgroups  $L_1, \ldots, L_k \subseteq G$ , possibly with repetitions, and  $a_1, \ldots, a_k \in \operatorname{dcl}(A)$  such that each  $a_iL_i$  is a nearest co-commutative coset to some  $p_i \in S_{\mathcal{X}}(A)$ , and

$$\mathcal{X} \subseteq \overline{B}_r^{\sharp} \cdot \bigcup_{i=1}^k a_i \cdot L_i^{\sharp}.$$

In addition, every  $\mathcal{L}_{\max}(\mathcal{X})$  appears at least once among  $L_1, \ldots, L_k$ . Proof. For each  $p \in S_{\mathcal{X}}(A)$ , we choose  $a_p \in \operatorname{dcl}(A)$  such that  $p(\mathfrak{R}) \subseteq \mu \cdot a_p \cdot L_p^{\sharp}$ .

We have

$$\mathcal{X} \subseteq \bigcup_{p \in S_{\mathcal{X}}(A)} p(\mathfrak{R}) \subseteq \bigcup_{p \in S_{\mathcal{X}}(A)} \mu \cdot a_p \cdot L_p^{\sharp} \subseteq \bigcup_{p \in S_{\mathcal{X}}(A)} \overline{B}_r^{\sharp} \cdot a_p \cdot L_p^{\sharp}.$$

Using logical compactness, we obtain finitely many  $p_1, \ldots, p_k \in S_{\mathcal{X}}(A)$  such that

$$\mathcal{X} \subseteq \bigcup_{i=1}^k \overline{B}_r^{\sharp} \cdot a_{p_i} \cdot L_{p_i}^{\sharp} = \overline{B}_r^{\sharp} \cdot \bigcup_{i=1}^k a_{p_i} \cdot L_{p_i}^{\sharp}.$$

This proves the main part.

In addition, let  $L \in \mathcal{L}_{\text{max}}$ . Choose  $p \in S_{\mathcal{X}}(A)$  such that  $L = L_p$  and also choose  $a \in \text{dcl}(A)$  such that  $p(\mathfrak{R}) \subseteq \mu \cdot a \cdot L^{\sharp}$ . We have

$$p(\mathfrak{R}) \subseteq \mathcal{X} \subseteq \bigcup_{i=1}^k \overline{B}_r^{\sharp} \cdot a_{p_i} \cdot L_{p_i}^{\sharp}.$$

Since p is a complete type over A, there is  $1 \leq j \leq k$  such that  $p(\mathfrak{R}) \subseteq \overline{B}_r^{\sharp} \cdot a_{p_j} \cdot L_{p_j}^{\sharp}$ . Since aL is a nearest co-commutative coset to p,

by Lemma 3.8, we conclude  $L \subseteq L_{p_j}$ . By maximality of L we get  $L = L_{p_i}$ .

## 4. Γ-DENSE TYPES IN UNIPOTENT GROUPS

4.1. Preliminaries on unipotent groups. As in [9], by a unipotent group we mean a real algebraic subgroup of the group of real  $n \times n$  upper triangular matrices with 1 on the diagonal.

We list below some properties of unipotent groups that we need and refer to [9] and [2] for more details.

We fix a unipotent group G.

**Fact 4.1.** For a subgroup H of G, the following are equivalent.

- (1) H is a closed connected subgroup of G.
- (2) H is a real algebraic subgroup of G.
- (3) H is definable in  $\mathbb{R}_{om}$ .

A lattice in G is a discrete subgroup  $\Gamma$  such that  $G/\Gamma$  is compact. Let  $\Gamma \subseteq G$  be a lattice. A real algebraic subgroup H of G is called  $\Gamma$ -rational if  $\Gamma \cap H$  is a lattice in H.

#### **Fact 4.2.** Let $\Gamma$ be a lattice in G.

- (1) The center Z(G) is  $\Gamma$ -rational.
- (2) The commutator subgroup [G, G] is closed and  $\Gamma$ -rational.
- (3) If H is a  $\Gamma$ -rational normal subgroup of G and  $\pi: G \to G/H$  is the quotient map then  $\pi(\Gamma)$  is a lattice in G/H. In addition, for every  $\pi(\Gamma)$ -rational subgroup  $K \subseteq G/H$ , the preimage  $\pi^{-1}(K)$  is  $\Gamma$ -rational.
- (4) If  $H_1$  and  $H_2$  are  $\Gamma$ -rational subgroups of G then  $H_1 \cap H_2$  is  $\Gamma$ -rational as well.

It follows from the above fact that for any real algebraic subgroup H of G there is the smallest  $\Gamma$ -rational subgroup containing H. We call it the  $\Gamma$ -rational closure of H and denote by  $H^{\Gamma}$ .

The next fact easily follows from Fact 4.2(3).

**Fact 4.3.** Assume that H is a  $\Gamma$ -rational normal subgroup of G,  $\pi \colon G \to G/H$  the quotient map and  $\Gamma_0 = \pi(\Gamma)$ . Then for every real algebraic subgroup  $L \subseteq G$ ,  $\pi(L^{\Gamma}) = \pi(L)^{\Gamma_0}$ .

We will need the following fact.

**Fact 4.4.** Let  $\Gamma$  be a lattice in G and H be a real algebraic subgroup of G. If H is a normal subgroup then  $H^{\Gamma}$  is normal as well.

The following is a restatement of Ratner's Orbit Closure Theorem in the case of unipotent groups.

**Fact 4.5.** [11] Let  $\Gamma$  be a lattice in G and H be a real algebraic subgroup of G. The topological closure of  $H \cdot \Gamma$  in G is  $H^{\Gamma} \cdot \Gamma$ .

We will be using the following well-known fact.

- **Fact 4.6.** Let H be a real algebraic subgroup of G and  $\Gamma_1$ ,  $\Gamma_2$  be lattices in G. If  $\Gamma_1$  and  $\Gamma_2$  are commensurable, i.e.  $\Gamma_1 \cap \Gamma_2$  is of finite index in both  $\Gamma_1$  and  $\Gamma_2$ , then  $H^{\Gamma_1} = H^{\Gamma_2}$ .
- 4.2.  $\Gamma$ -dense sets in unipotent groups. Let G be a unipotent group and  $\Gamma \subseteq G$  be a lattice. We say that a subset  $X \subseteq G$  is  $\Gamma$ -dense in G if the set  $X \cdot \Gamma$  is dense in G, i.e.  $\operatorname{cl}(X \cdot \Gamma) = G$ . Using Lemma 2.1(2), we conclude that a subset  $X \subseteq G$  is  $\Gamma$ -dense in G if and only if  $\operatorname{st}(X^{\sharp} \cdot \Gamma^{\sharp}) = G$ . We use this fact to extend the notion of  $\Gamma$ -density to arbitrary subsets of  $G^{\sharp}$ .

**Definition 4.7.** Let G be a unipotent group,  $\Gamma \subseteq G$  a lattice and  $\mathcal{X} \subseteq G^{\sharp}$  be an arbitrary set.

- (1) We say that  $\mathcal{X}$  is  $\Gamma$ -dense in G if  $\operatorname{st}(\mathcal{X} \cdot \Gamma^{\sharp}) = G$ .
- (2) We say that  $\mathcal{X}$  is strongly  $\Gamma$ -dense in G if  $\operatorname{st}(\mathcal{X} \cdot \Gamma_1^{\sharp}) = G$  for every lattice  $\Gamma_1$  commensurable with  $\Gamma$ .
- (3) We say that a type  $p \in S_G(A)$  is (strongly)  $\Gamma$ -dense in G if the set  $p(\mathfrak{R})$  is (strongly)  $\Gamma$ -dense in G.

Remark 4.8. Let G is be a unipotent group,  $\Gamma \subseteq G$  a lattice and  $\mathcal{X} \subseteq G^{\sharp}$ . It is easy to see that  $\mathcal{X}$  is strongly  $\Gamma$ -dense in G if and only if it is  $\Gamma_0$ -dense for every subgroup  $\Gamma_0 \subseteq \Gamma$  of finite index.

**Example 4.9.** Let  $G = (\mathbb{R}, +)$ ,  $\Gamma = \mathbb{Z}$  and let X be the closed interval [0, 1]. The set  $X^{\sharp}$  is  $\Gamma$ -dense in G, but not strongly  $\Gamma$ -dense.

The following fact follows from Facts 4.5 and 4.6.

**Fact 4.10.** Let G be a unipotent group and  $L \subseteq G$  a real algebraic subgroup. For a lattice  $\Gamma \subseteq G$  the following are equivalent.

- (1) L is  $\Gamma$ -dense in G.
- (2)  $L^{\Gamma} = G$
- (3) L is strongly  $\Gamma$ -dense in G.

We observe:

**Lemma 4.11.** Let  $\Gamma$  be a lattice in a unipotent group G. A subset  $\mathcal{X} \subseteq G^{\sharp}$  is  $\Gamma$ -dense in G if and only if  $\mu \cdot \mathcal{X} \cdot \Gamma^{\sharp} = G^{\sharp}$ .

*Proof.* The "if" part is clear.

For "the only if" part, since  $G/\Gamma$  is compact, given  $g \in G^{\sharp}$  there is  $\gamma \in \Gamma^{\sharp}$  such that  $g\gamma \in \mathcal{O}$ . Thus, since  $\mathcal{X}$  is  $\Gamma$ -dense in G, there is  $a \in \mathcal{X}$  such that  $\operatorname{st}(g\gamma) = \operatorname{st}(a)$ . It follows that  $g \in \mu \cdot a \cdot \Gamma^{\sharp}$ .

We will need the following fact.

Fact 4.12 ([9, Lemma 5.1]). Let  $\pi: G \to H$  be a real algebraic surjective homomorphism of unipotent groups, and  $\mathcal{X}$  a subset of  $G^{\sharp}$ . Then, for every lattice  $\Gamma \subseteq G$ , if  $\pi(\Gamma)$  is closed in H then

$$\pi(\operatorname{st}(\mathcal{X}\cdot\Gamma^{\sharp})) = \operatorname{st}(\pi(\mathcal{X})\cdot\pi(\Gamma^{\sharp})).$$

Our main goal is to describe  $\Gamma$ -dense types. We will consider the abelian case first.

4.3.  $\Gamma$ -dense types in abelian groups. Since every abelian unipotent group is algebraically isomorphic to  $(\mathbb{R}^m, +)$  for some m, we often identify an abelian unipotent group with an  $\mathbb{R}$ -vector space  $(\mathbb{R}^m, +)$ .

In the abelian case, every subgroup is co-commutative, hence for a set  $A \subseteq \mathfrak{R}$  and a type  $p(x) \in S(A)$  on  $\mathbb{R}^m$ , instead of a nearest co-commutative coset to p we say a nearest coset to p.

Our first goal of this section is to prove the following:

**Proposition 4.13.** Let G be an abelian unipotent group,  $A \subseteq \mathfrak{R}$ ,  $p \in S_G(A)$ , and  $a_p \in \operatorname{dcl}(A)$  be such that  $a_p + L_p$  is a nearest coset to p. For a lattice  $\Gamma \subseteq G$ , the following are equivalent.

- (1) The type p is  $\Gamma$ -dense in G.
- $(2) L_p^{\Gamma} = G.$
- (3) The type p is strongly  $\Gamma$ -dense in G.

*Proof.* Since, by Fact 4.6,  $L^{\Gamma} = L^{\Gamma_0}$  for any subgroup  $\Gamma_0 \subseteq \Gamma$  of finite index, it is sufficient to show  $(1) \Leftrightarrow (2)$ .

For simplicity we denote  $L_p^{\Gamma}$  by L, and assume  $A = \operatorname{dcl}(A)$ .

 $(1)\Rightarrow (2)$ . Assume that  $L\neq G$ , hence, by Fact 4.5, the set  $L+\Gamma$  is a closed proper subgroup of G. By Lemma 2.1(2),  $\mu+L^{\sharp}+\Gamma^{\sharp}$  is a proper subgroup of  $G^{\sharp}$ , hence the coset  $a_p+\mu+L^{\sharp}+\Gamma^{\sharp}$  is a proper subset of  $G^{\sharp}$ .

Since  $\mu + p(\mathfrak{R}) + \Gamma^{\sharp} \subseteq a_p + \mu + L^{\sharp} + \Gamma^{\sharp}$ , the set  $\mu + p(\mathfrak{R}) + \Gamma^{\sharp}$  is a proper subset of  $G^{\sharp}$  and, by Fact 4.11,  $p(\mathfrak{R})$  is not  $\Gamma$ -dense, so (1) fails.

 $(2) \Rightarrow (1)$ . Assume L = G and we prove that  $\operatorname{st}(p(\mathfrak{R}) + \Gamma^{\sharp}) = G$ , The proof is similar to [9, Proposition 5.3].

We use induction on  $\dim G$ .

If  $\dim(G) = 0$  then there is nothing to prove.

Assume  $\dim(G) = n > 0$  and the result holds for all abelian unipotent groups of dimension less than n.

We have  $\dim(L) > 0$ , hence, by Remark 3.2, p is a long type. Let P be the  $\mu$ -stabilizer of p. By Proposition 2.14, P is a real algebraic

subgroup of G of positive dimension, and by Lemma 3.10,  $P \subseteq L_p$ , hence  $P^{\Gamma} \subseteq L_p^{\Gamma} = L$ .

Let  $\pi: G \to G_0 := G/P^{\Gamma}$  be the quotient map,  $\Gamma_0 = \pi(\Gamma)$ , and  $q = \pi(p)$ . By Fact 4.2,  $\Gamma_0$  is a lattice in  $G_0$ . Notice that  $\dim(G_0) < \dim(G)$ .

Let  $a_q + L_q$  be a nearest coset to q. It follows from Lemma 3.9 that  $L_q = \pi(L_p)$ . Since  $G = L_p^{\Gamma}$ , by Fact 4.2(3),  $G_0 = L_q^{\Gamma_0}$ , hence, by induction hypothesis, the type q is  $\Gamma_0$ -dense in  $G_0^{\sharp}$ , and

$$\operatorname{st}(q(\mathfrak{R}) + \Gamma_0^{\sharp}) = G_0.$$

Applying Fact 4.12, we obtain

$$\pi(\operatorname{st}(p(\mathfrak{R}) + \Gamma^{\sharp})) = \operatorname{st}(q(\mathfrak{R}) + \Gamma_0^{\sharp}) = G_0.$$

Let  $D = \operatorname{st}(q(\mathfrak{R}) + \Gamma^{\sharp})$ . By Lemma 2.1(3), it is a closed subset of G. It is not hard to see that D is invariant ander the action of both P and  $\Gamma$ , hence it is invariant under  $P + \Gamma$ . Since D is closed, it is invariant under the action of the topological closure of  $P + \Gamma$ . By Fact 4.5,  $P^{\Gamma}$  is contained in  $\operatorname{cl}(P+\Gamma)$ , hence D is invariant under  $P^{\Gamma}$ . Since  $\pi(D) = G_0$  and  $\operatorname{ker}(\pi) = P^{\Gamma}$ , it follows then D = G, hence p is  $\Gamma$ -dense in G.

This finishes the proof of Proposition 4.13.

As a corollary we obtain the following theorem.

**Theorem 4.14.** Let  $A \subseteq \mathfrak{R}$ ,  $G = (\mathbb{R}^n, +)$ ,  $p \in S_G(A)$ , and let  $a_p + L_p$  be a nearest coset to p (so  $a_p \in \operatorname{dcl}(A)$ ). Then for every lattice  $\Gamma \subseteq \mathbb{R}^n$ , we have

$$\mu + p(\mathfrak{R}) + \Gamma^{\sharp} = \mu + a_p + L_p^{\sharp} + \Gamma^{\sharp}.$$

Proof. Consider the type  $p_1 = -a_p + p$ . It is a complete  $\mathcal{L}_{om}$ -type over A. Clearly  $L_p$  is a nearest coset to  $p_1$ , hence there exists a type  $p_2 \in S_{L_p}(A)$  which is  $\mu$ -equivalent to  $p_1$ , and therefore  $L_p$  is also a nearest coset to  $p_2$ . Let  $G_0 = L_p^{\Gamma}$  and  $\Gamma_0 = \Gamma \cap G_0$ , a lattice in  $G_0$ .

Working in  $G_0$  we have that  $L_p^{\Gamma_0} = G_0$ , hence by Proposition 4.13, the type  $p_2$  is  $\Gamma_0$ -dense in  $G_0$ , so, by Lemma 4.11,

$$(\mu \cap G_0^{\sharp}) + p_2(\mathcal{R}) + \Gamma_0^{\sharp} = G_0^{\sharp}.$$

Obviously,  $L_p$  is also  $\Gamma_0$ -dense in  $G_0$ , hence  $G_0^{\sharp} = (\mu \cap G_0^{\sharp}) + L_p^{\sharp} + \Gamma_0^{\sharp}$ . We conclude

$$\mu + p(\mathcal{R}) + \Gamma^{\sharp} = \mu + a_p + \mu + p_2(\mathcal{R}) + \Gamma^{\sharp} = \mu + a_p + L_p^{\sharp} + \Gamma^{\sharp}.$$

4.4. Abelianization and density. For a unipotent group G we will denote by  $G_{ab}$  the abelianization of G, i.e. the group  $G_{ab} = G/[G, G]$ , and by  $\pi_{ab}$  the quotient map  $\pi_{ab} : G \to G_{ab}$ . The group  $G_{ab}$  is also unipotent and dim G > 0 if and only if dim  $G_{ab} > 0$ .

If  $\Gamma \subseteq G$  is a lattice then we denote by  $\Gamma_{ab}$  the group  $\Gamma_{ab} = \pi_{ab}(\Gamma)$ . By Fact 4.2,  $\Gamma_{ab}$  is a lattice in  $G_{ab}$ . Our main goal in this section to show that a type  $p \in S_G(A)$  is  $\Gamma$ -dense in G if and only if its abelianization  $\pi_{ab}(p)$  is  $\Gamma_{ab}$ -dense in  $G_{ab}$ .

The next proposition is a key.

**Proposition 4.15.** Let G be a unipotent group,  $A \subseteq \mathfrak{R}$ ,  $\Gamma$  a lattice in G and  $p \in S_G(A)$ . Assume p is not  $\Gamma$ -dense in G. Then there is a co-commutative  $\Gamma$ -rational subgroup  $H \subseteq G$  such that for the projection  $\pi \colon G \to G/H$  the type  $\pi(p)$  is not  $\pi(\Gamma)$ -dense in G/H.

*Proof.* By induction on  $\dim(G)$ .

If dim(G) = 0 then there is nothing to prove.

Assume  $\dim(G) = n > 0$  and the proposition holds for all unipotent groups of dimension less than n.

If the type p is short then, by Lemma 2.8(1), the type  $\pi_{ab}(p)$  is short as well, and it is easy too see, e.g. using Lemma 2.9(1), that a short type is not  $\Gamma_{ab}$ -dense in  $G_{ab}$ . We can take H = [G, G] that is  $\Gamma$ -rational by Fact 4.2(2).

Thus we may assume that p is a long type. Let P be the  $\mu$ -stabilizer of p. By Proposition 2.14, P is an  $\mathbb{R}_{om}$ -definable subgroup of G of positive dimension.

As in [9, Propositioni 5.3], we consider the smallest  $\mathbb{R}_{om}$ -definable, normal  $\Gamma$ -rational subgroup of G containing P and denote it by  $N(P)^{\Gamma}$ . Let  $N_0$  be the intersection of  $N(P)^{\Gamma}$  with the center of G. Since G is unipotent and  $N(P)^{\Gamma}$  has positive dimension, the group  $N_0$  is also of positive dimension (see, for example, [12, Proposition 7.13]), and, by Fact 4.2, it is  $\Gamma$ -rational.

Let  $\pi: G \to G_0 := G/N_0$  be the quotient map,  $\Gamma_0 = \pi_0(\Gamma)$ , and  $q = \pi(p)$ . By Fact 4.2,  $\Gamma_0$  is a lattice in  $G_0$ .

We claim that the type q is not  $\Gamma_0$ -dense in  $G_0$ .

Indeed, assume towards contradiction that q is  $\Gamma_0$ -dense in  $G_0$ . Then, by Fact 4.12,

$$\pi_0(\operatorname{st}(p(\mathfrak{R})\cdot\Gamma^{\sharp}))=G_0.$$

Let  $D_{p,\Gamma} = \operatorname{st}(p(\mathfrak{R}) \cdot \Gamma^{\sharp})$  It follows from the above equation that

$$(4.1) D_{p,\Gamma} \cdot N_0 = G.$$

Our aim is to show that  $D_{p,\Gamma} = G$ , contradicting the fact that p is not  $\Gamma$ -dense in G.

Since, by Lemma 2.1(2), the set  $D_{p,\Gamma}$  is a closed subset of G, it is sufficient to show that it is dense in G.

Claim A. The set  $D_{p,\Gamma}$  is left invariant under the  $\mu$ -stabilizer P of p.

*Proof.* Note that  $D_{p,\Gamma} = \operatorname{st}(\mu \cdot p(\mathfrak{R}) \cdot \Gamma^{\sharp})$ , and  $\mu p$  is left-invariant under P. Thus, for  $g \in P$ ,

$$g \cdot D_{p,\Gamma} = g \cdot \operatorname{st}(\mu \cdot p(\mathfrak{R}) \cdot \Gamma^{\sharp}) = \operatorname{st}(g \cdot \mu \cdot p(\mathfrak{R}) \cdot \Gamma^{\sharp}) = \operatorname{st}(\mu \cdot p(\mathfrak{R}) \cdot \Gamma^{\sharp}) = D_{p,\Gamma}.$$

Clearly  $D_{p,\Gamma}$  is right-invariant under action of  $\Gamma$ . Thus,  $P \cdot D_{p,\Gamma} \cdot \Gamma = D_{p,\Gamma}$ , and, in addition, by equation (4.1),  $D_{p,\Gamma} N_0 = G$ .

Because  $N_0 = N(P)^{\Gamma} \cap Z(G)$ , our goal,  $D_{p,\Gamma} = G$ , follows from the following general result:

Claim B. For a unipotent group G, assume that  $D \subseteq G$  is a closed set, left invariant under a real algebraic subgroup  $P \subseteq G$  and right invariant under a lattice  $\Gamma \subseteq G$ . Let  $N_0 \subseteq N(P)^{\Gamma} \cap N_G(P)$  be a subgroup of G. If  $DN_0 = G$  then D = G.

*Proof.* Let  $Y = \{g \in G : (P^g)^{\Gamma} = N(P)^{\Gamma}\}$ . This is not, in general, a definable set, but, by [9, Proposition 4.3], it is dense in G. Thus, it is sufficient to prove that  $Y \subseteq D$ .

First note that Y is left invariant under  $N_0$ . Indeed, assume that  $a \in N_0 b$ . Since  $ab^{-1} \in N_0 \subseteq N_G(P)$ , then  $P^a = P^b$ , implying that  $b \in Y$  if and only if  $a \in Y$ .

Let  $b \in Y$ . Since  $DN_0 = G$ , there is  $a \in D$  such that  $b \in aN_0$ , and therefore  $a \in Y$ . Using the definition of Y and the fact that  $aP^a = Pa$ , we obtain

$$b \in a \cdot N_0 \subseteq a \cdot N(P)^{\Gamma} = a \cdot (P^a)^{\Gamma} = a \cdot \operatorname{cl}(P^a \cdot \Gamma) \subseteq \operatorname{cl}(a \cdot P^a \cdot \Gamma) = \operatorname{cl}(P \cdot a \cdot \Gamma).$$

Since  $a \in D$ , by the invariance properties of D, also  $b \in D$ . Hence  $Y \subseteq D$ , so D = G, a contradiction.

This ends the proof of Claim B, and thus we conclude that q is not  $\Gamma_0$ -dense in  $G_0$ .

Applying the induction hypothesis to  $G_0$ ,  $\Gamma_0$  and q, we obtain a co-commutative  $\Gamma_0$ -rational subgroup  $H_0 \subseteq G_0$  such that the image of q in  $G_0/H_0$  is not  $\Gamma_0/H_0$ -dense. It is not hard to see that  $H = \pi_0^{-1}(H_0)$  is a co-commutative  $\Gamma$ -rational subgroup of G satisfying the conclusion of the proposition.

This finishes the proof of Proposition 4.15

We can now prove the main theorem of this section.

**Theorem 4.16.** Let G be a unipotent group,  $A \subseteq \Re$  and  $\Gamma$  a lattice in G. A type  $p \in S_G(A)$  is  $\Gamma$ -dense in G if and only if the type  $\pi_{ab}(p)$  is  $\Gamma_{ab}$ -dense in  $G_{ab}$ .

*Proof.* Let  $q = \pi_{ab}(p)$ . We write additively the group operation in  $G_{ab}$ . By Fact 4.12,

$$\pi_{\rm ab}(\operatorname{st}(p(\mathfrak{R})\cdot\Gamma^{\sharp})) = \operatorname{st}(q(\mathfrak{R}) + \Gamma_{\rm ab}^{\sharp}).$$

This implies the "only if" part.

We prove the "if part" part by contraposition, using Proposition 4.15. Indeed, assume the type p is not  $\Gamma$ -dense in G, and we derive that  $\pi_{ab}(p)$  is not  $G_{ab}$ -dense in  $G_{ab}$ .

Let H and  $\pi: G \to G/H$  be as in Proposition 4.15. Since H contains [G, G], the map  $\pi$  factors through  $G_{ab}$ , i.e. there is  $\pi': G_{ab} \to G/H$  with  $\pi = \pi' \circ \pi_{ab}$ . By Fact 4.12

$$\pi' \Big( \mathrm{st} \big( \pi_{\mathrm{ab}}(p)(\mathfrak{R}) + \Gamma_{\mathrm{ab}}^{\sharp} \big) \Big) = \mathrm{st} \Big( \pi(p)(\mathfrak{R}) + \pi(\Gamma)^{\sharp} \Big).$$

Since  $\pi(p)$  is not  $\pi(\Gamma)$ -dense in G/H, the type  $\pi_{ab}(p)$  is not  $\Gamma_{ab}$ -dense in  $G_{ab}$ .

The following summarizes main results of this section.

**Theorem 4.17.** Let G be a unipotent group,  $A \subseteq \mathfrak{R}$  and  $p \in S_G(A)$ . For a lattice  $\Gamma \subseteq G$ , the following are equivalent.

- (1) The type p is  $\Gamma$ -dense in G.
- (2) The type p is strongly  $\Gamma$ -dense in G.
- (3)  $L_p^{\Gamma} = G$ .
- (4) The type  $\pi_{ab}(p)$  is  $\Gamma_{ab}$ -dense in  $G_{ab}$ .
- (5) The type  $\pi_{ab}(p)$  is strongly  $\Gamma_{ab}$ -dense in  $G_{ab}$ .
- (6)  $\pi_{ab}(L_p)^{\Gamma_{ab}} = G_{ab}$ .

#### 5. Γ-dense definable subsets of unipotent groups

We fix a unipotent group G.

In this section we obtain a description of  $\Gamma$ -dense definable subsets of G similar to that of Theorem 4.17.

First an elementary lemma.

**Lemma 5.1.** For  $A \subseteq \mathfrak{R}$ , let  $\mathcal{X} \subseteq G^{\sharp}$  be  $\mathcal{L}_{om}$ -definable over A. For a lattice  $\Gamma \subseteq G$ , if some type  $p \in S_{\mathcal{X}}(A)$  is  $\Gamma$ -dense in G then  $\mathcal{X}$  is strongly  $\Gamma$ -dense G.

Proof. Assume a type  $p \in S_{\mathcal{X}}(A)$  is Γ-dense, hence, by Theorem 4.17, it is strongly Γ-dense. Since  $p(\mathfrak{R}) \subseteq \mathcal{X}$ , obviously  $\mathcal{X}$  is strongly Γ-dense.

The main goal of this section (Theorem 5.5) is to show that an appropriate converse of the above lemma holds. Namely, an  $\mathcal{L}_{om}$ -definable subset  $\mathcal{X} \subseteq G^{\sharp}$  is strongly  $\Gamma$ -dense in G if and only if some type on  $\mathcal{X}$  is  $\Gamma$ -dense.

We need two lemmas and a proposition.

**Lemma 5.2.** Let  $L \subseteq G$  be a normal real algebraic subgroup and  $\alpha \in G^{\sharp}$ .

(1) For any lattice  $\Gamma \subseteq G$  the set  $\mathcal{O} \cap \alpha \Gamma^{\sharp}$  is nonempty, and for every  $g \in \operatorname{st}(\alpha \Gamma^{\sharp})$  we have

$$\operatorname{st}(\alpha \cdot L^{\sharp} \cdot \Gamma^{\sharp}) = g \cdot L^{\Gamma} \cdot \Gamma.$$

(2) If L is co-commutative then  $\alpha L$  is a nearest co-commutative coset to some type  $p \in S(\alpha)$  on  $\alpha L^{\sharp}$ , hence  $\mathscr{L}_{\max}(\alpha L^{\sharp}) = \{L\}$ .

*Proof.* (1) Because  $\Gamma$  is co-compact, the set  $\mathcal{O} \cap \alpha \Gamma^{\sharp}$  is not empty. Let  $g \in \operatorname{st}(\alpha \Gamma^{\sharp})$ , and we choose  $\gamma \in \Gamma^{\sharp}$  with  $\alpha \cdot \gamma \in \mu g$ . Since L is normal, we have

$$\operatorname{st}(\alpha \cdot L^{\sharp} \cdot \Gamma^{\sharp}) = \operatorname{st}(\alpha \cdot \Gamma^{\sharp} \cdot L^{\sharp}) = \operatorname{st}(g \cdot \Gamma^{\sharp} \cdot L^{\sharp})$$
$$= \operatorname{st}(g \cdot L^{\sharp} \cdot \Gamma^{\sharp}) = \operatorname{cl}(g \cdot L \cdot \Gamma) = g \cdot L^{\Gamma} \cdot \Gamma.$$

- (2) We first consider the abelian case, namely we assume  $G = (\mathbb{R}^n, +)$  and  $L \subseteq \mathbb{R}^n$  a linear subspace. We need to show that there is  $\beta \in \alpha + L^{\sharp}$  such that  $\beta \notin \mu(\gamma + L_0^{\sharp})$ , for any  $\gamma \in \operatorname{dcl}(\alpha)$  and a proper subspace  $L_0 \subseteq L$ . It is thus sufficient to show
- $(5.1) \quad \alpha + L^{\sharp} \not\subseteq \bigcup \{ \overline{B}_{1}^{\sharp} + \gamma + L_{0}^{\sharp} \colon \gamma \in \operatorname{dcl}(\alpha), L_{0} \subsetneq L \text{ a subspace} \}.$

By logical compactnes, (5.1) follows from the following claim.

Claim. Let  $r \in \mathbb{R}^{\geq 0}$ , and  $L_1, \ldots, L_k \subseteq L$  be proper subspaces. Then for any  $\alpha, \gamma_1, \ldots, \gamma_k \in \mathfrak{R}^n$  we have

$$\alpha + L^{\sharp} \not\subseteq \overline{B}_r^{\sharp} + \bigcup_{i=1}^k (\gamma_i + L_i^{\sharp}).$$

*Proof of Claim.* Since, for fixed r and  $L_1, \ldots, L_k$ , the conclusion of the claim can be expressed by a first-order formula, we can work in  $\mathbb{R}$  instead of  $\mathfrak{R}$ ; and also, subtracting  $\alpha$  from both sides, we only need to consider the case  $\alpha = 0$ .

We fix proper subspaces  $L_1, \ldots, L_k \subseteq L$ , and show that for all  $\geq 0$  and  $b_1, \ldots, b_k \in \mathbb{R}^n$ , we have  $L \not\subseteq \overline{B}_r + \bigcup_{i=1}^k (b_i + L_i)$ .

Clearly, by the dimension assumptions,  $L \neq L_1 \cup \ldots \cup L_k$ .

Next, let us see that  $L \not\subseteq \overline{B}_r + \bigcup_{i=1}^k L_i$  for any  $r \in \mathbb{R}^{\geq 0}$ . Indeed, choose  $c \in L \setminus (\bigcup_{i=1}^k L_i)$ . Then, for every  $i = 1, \ldots, k$ , we have  $d(c, L_i) > 0$ , where  $d(\cdot, \cdot)$  denotes the Euclidean distance in  $\mathbb{R}^n$ . Since  $d(tc, L_i) = td(c, L_i)$  for  $t \in \mathbb{R}^{>0}$ , we obtain that for given  $r \in \mathbb{R}^{\geq}$  and  $i = 1, \ldots, k$ , for t large enough,  $d(tc, L_i) > r$ , and hence  $tc \in L \setminus (\overline{B}_r + \bigcup_{i=1}^k L_i)$ .

Finally, assume that for some r > 0 and  $b_1, \ldots, b_k \in \mathbb{R}^n$  we would have  $L \subseteq \overline{B}_r + \bigcup_{i=1}^k (b_i + L_i)$ . Then, choosing r' > 0 big enough so that  $b_i + \overline{B}_r \subseteq \overline{B}_{r'}$  for  $i = 1, \ldots, k$ , we would have  $L \subseteq \overline{B}_{r'} + \bigcup_{i=1}^k L_i$ , a contradiction.

This finishes the proof of Claim, and hence the lemma, in the case that G is abelian.

When G is nilpotent and  $L \subseteq G$  is co-commutative we first apply the above to  $\pi_{ab}(\alpha L^{\sharp}) \subseteq G_{ab}^{\sharp}$  and find a type over  $\alpha$  with  $q \vdash \pi_{ab}(\alpha L^{\sharp})$ , such that  $\pi_{ab}(\alpha L)$  is a nearest coset to q. Now, choose a type p over  $\alpha$  such that  $p \vdash \alpha L$  and  $\pi_{ab}(p) = q$ . A nearest co-commutative coset to p is contained in  $\alpha L$ , and projects via  $\pi_{ab}$  onto  $\pi_{ab}(\alpha L)$  (see Lemma 3.9). Since  $L \supseteq [G, G]$ , it follows that  $\alpha L$  is a nearest co-commutative coset to p. This finishes the proof of the claim.

End of the proof of the lemma.

We are going to need the following result.

**Lemma 5.3.** If H is a proper real algebraic subgroup of G then  $\pi_{ab}(H)$  is a proper subgroup of  $G_{ab}$ 

*Proof.* By [2, Theorem 1.1.13] there is a chain of real algebraic subgroups

$$\{e\} = H_0 \subsetneq \cdots \subsetneq H = H_m \subsetneq H_{m+1} \subsetneq \cdots \subsetneq H_n = G,$$

with  $n = \dim(G)$  and  $\dim H_{i+1} = \dim H_i + 1$ . By [2, Lemma 1.1.8],  $[G, G] \subseteq H_{n-1}$ . Hence  $H_{n-1}/[G, G]$  is a proper subgroup of  $G_{ab}$  and so is H/[G, G].

**Proposition 5.4.** Let  $\Gamma \subseteq G$  be a lattice, and  $L_1, \ldots, L_k$  proper  $\Gamma$ -rational subgroups of G. If  $K \subset G$  is a compact set then there is a subgroup  $\Gamma_0 \subseteq \Gamma$  of finite index such that for any  $g_1, \ldots, g_k \in G$ , we have

$$K \cdot \bigcup_{i=1}^{k} g_i \cdot L_i \cdot \Gamma_0 \neq G.$$

*Proof.* We first consider the case when G is abelian. So we assume  $G = (\mathbb{R}^n, +)$ .

**Claim.** Let  $\Gamma \subseteq \mathbb{R}^n$  be a lattice,  $K \subseteq \mathbb{R}^n$  a compact set, and  $L \subseteq \mathbb{R}^n$  be a proper  $\Gamma$ -rational subspace. Then for any  $m \in \mathbb{N}$ , there is a subgroup  $\Gamma' \subseteq \Gamma$  of finite index such that for some  $b_1, \ldots, b_m \in \mathbb{R}^n$ , the translates  $b_i + K + L + \Gamma'$ ,  $i = 1, \ldots, m$ , are pairwise disjoint.

*Proof.* Replacing  $\mathbb{R}^n$  by  $\mathbb{R}^n/L$  if needed, we may assume that L is the trivial subspace  $\{0\}$ .

Since K is compact, it is bounded, hence there are  $b_1, \ldots b_m \in \mathbb{R}^n$  such that the translates  $b_1 + K, \ldots, b_m + K$  are pair-wise disjoint. Let  $B = \bigcup_{i=1}^m (b_i + K)$ . Obviously B is compact and hence the set  $B' = B - B = \{b - b' : b, b' \in B\}$  is compact as well.

Since  $\Gamma$  is discrete, the intersection  $\Gamma \cap B'$  is finite. Every finitely generated abelian group is residually finite, i.e. the intersection of all subgroups of finite index is trivial, hence there is a subgroup  $\Gamma' \subseteq \Gamma$  of finite index with  $\Gamma' \cap B' = \{0\}$ . It is not hard to see that the sets  $b_i + K + \Gamma'$ ,  $i = 1, \ldots m$ , are pairwise disjoint.

This finishes the proof of the claim.

We return to the proof of the proposition for  $G = (\mathbb{R}^n, +)$ . We apply the above claim to each  $L_i$  with m = k + 1, and for each  $i = 1, \ldots, k$ , obtain a subgroup  $\Gamma_i \subseteq G$  of finite index such that  $K + L_i + \Gamma_i$  has k + 1 disjoint translates.

Since every abelian group is amenable, there is a G-invariant finitely additive probability measure  $\lambda \colon \mathscr{P}(G) \to [0,1]$ . By our choice of  $\Gamma_i$ , we have  $\lambda(K + L_i + \Gamma_i) \leq 1/(k+1)$ .

We take  $\Gamma_0 = \bigcap_{i=1}^k \Gamma_i$ . For any  $g_1, \ldots, g_k \in G$  we have

$$\lambda \Big( \bigcup_{i=1}^{k} g_i + K + L_i + \Gamma_0 \Big) \le \sum_{i=1}^{k} \lambda (g_i + K + L_i + \Gamma_0) \le \sum_{i=1}^{k} \lambda (g_i + K + L_i + \Gamma_i) \le k/(k+1) < 1.$$

Hence  $\bigcup_{i=1}^{k} g_i + K + L_i + \Gamma_0 \neq G$ . This finishes the proof of the abelian case.

Assume now that G is an arbitrary unipotent group. Let  $K_{ab} = \pi_{ab}(K)$ ,  $\Gamma_{ab} = \pi_{ab}(\Gamma)$ , and, for i = 1, ..., k, let  $L_i^{ab} = \pi_{ab}(L_i)$ . Obviously  $K_{ab}$  is a compact subset of  $G_{ab}$ ,  $\Gamma_{ab}$  is a lattice in  $G_{ab}$  by Fact 4.2, and it is not hard to see that each  $L_i^{ab}$  is  $\widetilde{\Gamma}$ -rational subgroup of  $G_{ab}$ .

It also follows from Lemma 5.3 that each  $L_i^{\text{ab}}$  is a proper subgroup of  $G_{\text{ab}}$ .

We now use the abelian case and find a subgroup  $\Gamma'_0 \subseteq \Gamma_{ab}$  such that for any  $b_1, \ldots, b_m \in G_{ab}$  we have  $K_{ab} \cdot \bigcup_{i=1}^k b_i \cdot L_i^{ab} \cdot \Gamma'_0 \neq G_{ab}$ .

We take 
$$\Gamma_0 = \pi_{ab}^{-1}(\Gamma_0') \cap \Gamma$$
.

We are now ready to prove one of the main theorems of this paper.

**Theorem 5.5.** Let G be a unipotent group,  $A \subseteq \mathfrak{R}$ , and let  $\mathcal{X} \subseteq G^{\sharp}$  be a set  $\mathcal{L}_{om}$ -definable over A.

For a lattice  $\Gamma \subseteq G$ , the following are equivalent:

- (a) The set  $\mathcal{X}$  is strongly  $\Gamma$ -dense in G.
- (b)  $L^{\Gamma} = G$  for some  $L \in \mathcal{L}_{\max}(\mathcal{X})$ .
- (c) Some type  $p \in S_{\mathcal{X}}(A)$  is  $\Gamma$ -dense.

*Proof.* By Theorem 4.17,  $(b) \Leftrightarrow (c)$ , and, by Lemma 5.1,  $(c) \Rightarrow (a)$ .

Let us show that  $(a) \Rightarrow (b)$ .

Let  $\Gamma \subseteq G$  be a lattice. We choose  $L_i$  and  $a_i$ , i = 1, ..., k, as in Theorem 3.15 with r = 1.

Assume (b) fails, namely,  $L^{\Gamma} \neq G$  for all  $L \in \mathcal{L}_{\max}(\mathcal{X})$ . Then clearly,  $L_i^{\Gamma} \neq G$ , for all i = 1, ..., k. For any subgroup  $\Gamma_0 \subseteq G$  of finite index we have

$$\mathcal{X} \cdot \Gamma_0^{\sharp} \subseteq \overline{B}_1^{\sharp} \cdot \bigcup_{i=1}^k a_i \cdot L_i^{\sharp} \cdot \Gamma_0^{\sharp},$$

hence

$$\operatorname{st}(\mathcal{X} \cdot \Gamma_0^{\sharp}) \subseteq \operatorname{st}\left(\overline{B}_1^{\sharp} \cdot \bigcup_{i=1}^k a_i \cdot L_i^{\sharp} \cdot \Gamma_0^{\sharp}\right) = \bigcup_{i=1}^k \operatorname{st}(\overline{B}_1^{\sharp} \cdot a_i \cdot L_i^{\sharp} \cdot \Gamma_0^{\sharp}).$$

Using Lemma 5.2(1) we choose  $g_1, \ldots, g_k \in G$  such that  $\operatorname{st}(\overline{B}_r^{\sharp} \cdot a_i \cdot L_i^{\sharp} \cdot \Gamma_0^{\sharp}) = \overline{B}_r \cdot g_i \cdot L_i^{\Gamma_0} \cdot \Gamma_0$ . By fact 4.6,  $L_i^{\Gamma_0} = L_i^{\Gamma}$ , hence

$$\operatorname{st}(\mathcal{X}\cdot\Gamma_0^{\sharp})\subseteq \overline{B}_r\cdot\bigcup_{i=1}^k g\cdot L_i^{\Gamma}\cdot\Gamma_0.$$

By Proposition 5.4, there exists  $\Gamma_0 \subseteq \Gamma$  of finite index for which the set on the right is a proper subset of G, hence (a) fails. Thus,  $(a) \Rightarrow (b)$ .

Recall that for  $\pi_{ab}: G \to G_{ab}$ , and  $\Gamma \subseteq G$  a lattice, we let  $\Gamma_{ab} = \pi_{ab}(\Gamma)$ .

Corollary 5.6. Let G be a unipotent group, and let  $\mathcal{X} \subseteq G^{\sharp}$  be an  $\mathcal{L}_{om}$ -definable set. For a lattice  $\Gamma \subseteq G$ , the set  $\mathcal{X}$  is strongly  $\Gamma$ -dense in G if and only if  $\pi_{ab}(\mathcal{X})$  is strongly  $\Gamma_{ab}$ -dense in  $G_{ab}$ .

*Proof.* The "only if" part follows from Fact 4.12.

For the "if" part, assume that  $\pi_{ab}(\mathcal{X})$  is strongly  $\Gamma_{ab}$ -dense in  $G_{ab}$ . Choose a set  $A \subseteq \mathfrak{R}$  such that  $\mathcal{X}$  is  $\mathcal{L}_{om}$ -definable over A. Applying Theorem 5.5(2) to  $\pi_{ab}(\mathcal{X})$ , we obtain a type  $q(x) \in S_{\pi_{ab}(\mathcal{X})}(A)$  that is  $\Gamma_{ab}$ -dense in  $G_{ab}$ . Let  $p \in S_{\mathcal{X}}(A)$  be a type on  $\mathcal{X}$  with  $\pi_{ab}(p) = q$ . By Theorem 4.17, the type p is  $\Gamma$ -dense in G, hence  $\mathcal{X}$  is  $\Gamma$ -dense in G as well.

## 6. Interpreting the results as Hausdorff limits

### 6.1. **Hausdorff limits.** We first recall some definitions.

Let (M, d) be a compact metric space, and  $X_1, X_2 \subseteq X$ . The Hausdorff distance  $d_H(X_1, X_2)$  between  $X_1$  and  $X_2$  is defined as follows: First, for  $x \in M$ , we let  $d(x, X_i) = \inf_{y \in X_i} d(x, y)$ . Next,

$$d_H(X_1, X_2) = \max\{ \sup_{x \in X_1} d(x, X_2), \sup_{x \in X_2} d(x, X_1) \}.$$

An equivalent definition is given by:

$$d_H(X_1, X_2) = \inf\{r \ge 0 : \forall x_i \in X_i, i = 1, 2, d(x_1, X_2), d(x_2, X_1) \le r\}.$$

**Remark 6.1.** For  $X_1, X_2 \subseteq M$  we have  $d_H(X_1, X_2) = 0$  if and only if  $cl(X_1) = cl(X_2)$ .

Denoting by  $\mathcal{K}(M)$  the set of all compact subsets of M, it is known that the restriction of  $d_H$  to  $\mathcal{K}(M)$  makes it into a compact metric space.

The topology induced by  $d_H$  on  $\mathcal{K}(M)$  does not depend on the metric d but only on the topology of M. It coincides with the Vietoris topology.

Given a family  $\mathcal{F} \subseteq \mathcal{K}(M)$ , a set  $Y \in \mathcal{K}(M)$  is a Hausdorff limit of  $\mathcal{F}$  if for every  $\varepsilon > 0$  there is  $F \in \mathcal{F}$  with  $d_H(Y, F) < \varepsilon$ . Using Remark 6.1, we extend this definition to a family  $\mathcal{F} \subseteq \mathcal{P}(M)$  of arbitrary subsets of M by saying that  $Y \in \mathcal{K}(M)$  is a Hausdorff limit of  $\mathcal{F}$  if it is a Hausdorff limit of the family  $\{\operatorname{cl}(F) \colon F \in \mathcal{F}\}$ .

6.1.1. Limits at infinity. We denote by  $I_{\infty}$  the interval  $(0, +\infty) \subseteq \mathbb{R}$ . We abreviate "for all sufficiently large t" by " $t \gg 0$ ". We define:

**Definition 6.2.** Let  $\mathcal{F} = \{F_t : t \in I_{\infty}\}$  be a family of subsets of M.

- (1) A set  $Y \in \mathcal{K}(M)$  is a Hausdorff limit at  $\infty$  of the family  $\mathcal{F}$  if for all  $\epsilon > 0$  and r > 0 there is t > r with  $d_H(Y, F_t) < \varepsilon$ .
- (2) We say that the family  $\mathcal{F}$  converges to a set  $Y \in \mathcal{K}(M)$  at  $\infty$  if Y is the unique Hausdorff limit of  $\mathcal{F}$  at  $\infty$ . In this case, since  $\mathcal{K}(M)$

is compact, Y is **the** limit of  $\mathcal{F}$  as t goes to  $\infty$ , namely for any  $\epsilon > 0$  there is  $R \in \mathbb{R}$  such that for all t > R,  $d_H(Y, F_t) < \varepsilon$ .

6.2. Haudorff Limits via the standard part map. We fix a compact set  $M \subseteq \mathbb{R}^n$  with the metric d induced by the Euclidean metric of  $\mathbb{R}^n$ , and we view both M and d as definable in  $\mathbb{R}_{\text{full}}$ .

Since M is compact,  $M^{\sharp} \subseteq \mathcal{O}^n$ , and we denote by  $\operatorname{st}_M$  the restriction of the standrad part map  $\operatorname{st} \colon \mathcal{O}^n \to \mathbb{R}^n$  to  $M^{\sharp}$ . It is not hard to see that  $\operatorname{st}_M \colon M^{\sharp} \to M$  maps  $\alpha \in M^{\sharp}$  to the unique  $a \in M$  such that  $\alpha \in U^{\sharp}$  for every neighborhood U of a.

Let  $\mathcal{F} = \{F_t : t \in T\}$  be a family of subsets of M indexed by a set  $T \subseteq \mathbb{R}^m$ . We can view this family also as the family of fibers of the set  $F = \{(x,t) \in X \times T : x \in F_t\}$ , with respect to the second projection, and hence as a family definable in  $\mathbb{R}_{\text{full}}$ . Thus for  $\tau \in T^{\sharp}$  we also have a "non-standard" fiber  $F_{\tau}^{\sharp} = \{x \in M^{\sharp} : (x,\tau) \in F^{\sharp}\}$ . Using [5, Theorem 4.4] we obtain:

**Fact 6.3.** In the above setting, a set  $Y \in \mathcal{K}(M)$  is a Hausdorff limit of the family  $\mathcal{F} = \{F_t : t \in T\}$  if and only if there is  $\tau \in T^{\sharp}$  such that  $Y = \operatorname{st}_M(F_{\tau}^{\sharp})$ .

Using the above, we conclude:

**Lemma 6.4.** For a family  $\mathcal{F} = \{F_t : t \in I_\infty\}$  of subsets of M indexed by  $I_\infty$ , a set  $Y \in \mathcal{K}(M)$  is a Hausdorff limit at  $\infty$  of  $\mathcal{F}$  if and only if there is  $\tau \in \mathfrak{R}$  with  $\tau > \mathbb{R}$ , such that  $Y = \operatorname{st}_M(F_\tau^\sharp)$ .

*Proof.* For  $r \in \mathbb{R}^{>0}$  let  $I_{>r}$  be the interval  $(r, +\infty) \subseteq \mathbb{R}$  and  $\mathcal{F}_r$  be the family  $\mathcal{F}_r = \{F_t : t \in I_{>r}\}.$ 

It is easy to see that a set  $Y \in \mathcal{K}(M)$  is a Hausdorf limit at  $\infty$  of the family  $\mathcal{F}$  if and only if for every  $r \in \mathbb{R}^{>0}$  the set Y is a Hausdorf limit of the family  $\mathcal{F}_r$ .

By Fact 6.3, the latter condition is equivalent to the following: for every  $r \in \mathbb{R}^{>0}$ , there is  $\tau_r \in I_{>r}^{\sharp}$  with  $Y = \operatorname{st}_M(F_{\tau_n}^{\sharp})$ .

Thus the conclusion of the lemma can be restated as follows:

For a set  $Y \in \mathcal{K}(M)$  the following are equivalent:

- (a) For every  $r \in \mathbb{R}^{>0}$ , there is  $\tau_r \in I_{>r}^{\sharp}$  with  $Y = \operatorname{st}_M(F_{\tau_r}^{\sharp})$ .
- (b) There is  $\tau \in \mathfrak{R}$  with  $\tau > \mathbb{R}$  such that  $Y = \operatorname{st}_M(F_{\tau}^{\sharp})$ .

The direction  $(b) \Rightarrow (a)$  is obvious, and the opposite direction follows from the  $|\mathbb{R}|^+$ -saturation of  $\mathfrak{R}_{\text{full}}$ .

6.3. Hausdorff limits in G/H. Let G be a connected Lie group and  $H \subseteq G$  a closed subgroup such that the space of the left cosets N = G/H is compact, with respect to the quotient topology. We denote by  $\pi \colon G \to N$  the quotient map. Using Whitney embedding theorem we embed G into some  $\mathbb{R}^m$  and N into some  $\mathbb{R}^n$  as closed subsets, and view G, N and  $\pi$  as definable in  $\mathbb{R}_{\text{full}}$ .

Given a family  $\mathcal{F} = \{F_t : t \in T\}$  of subsets of G, we let

$$\pi(\mathcal{F}) = \{ \pi(F_t) \colon t \in T \}$$

be the corresponding family of subsets of N.

**Proposition 6.5.** Let  $\mathcal{F} = \{F_t : t \in I_\infty\}$  be a family of subsets of G. A set  $Y \in \mathcal{K}(N)$  is a Hausdorf limit at  $\infty$  of the family  $\pi(\mathcal{F})$  if and only if there is  $\tau \in \mathfrak{R}$  with  $\tau > \mathbb{R}$  such that  $Y = \pi(\operatorname{st}_G(F_\tau^{\sharp} \cdot H^{\sharp}))$ .

*Proof.* For  $\tau \in \mathfrak{R}_{\text{full}}$ , by Claim 6.4, it is sufficient to show that

$$\operatorname{st}_N(\pi^{\sharp}(F_{\tau}^{\sharp})) = \pi(\operatorname{st}_G(F_{\tau}^{\sharp} \cdot H^{\sharp})).$$

For  $\alpha \in G^{\sharp}$ , we will show that  $\operatorname{st}_{N}(\pi^{\sharp}(\alpha)) = \pi(\operatorname{st}_{G}(\alpha \cdot H^{\sharp}))$ . That is clearly enough.

Since G/H is compact, there is  $\beta \in \mathcal{O}^m \cap G^{\sharp}$  with  $\beta \in \alpha \cdot H^{\sharp}$ . Let  $b = \operatorname{st}_G(\beta)$ . Since H is a closed subgroup, the set  $b \cdot H$  is closed and, by Lemma 2.1(2), we have

$$b \cdot H = \operatorname{st}_G(b \cdot H^{\sharp}) = \operatorname{st}_G(\beta \cdot H^{\sharp}).$$

Since  $\pi^{\sharp}$  is invariant under the action of  $H^{\sharp}$  on the right we also have  $\pi^{\sharp}(\alpha) = \pi^{\sharp}(\beta)$  and we are left to show

$$\operatorname{st}_N(\pi^{\sharp}(\beta)) = \pi(b).$$

Since  $\pi$  is continuous, the latter follows from Fact 2.2.

6.4. Hausdorff limits in nilmanifolds. We go back to our o-minimal structure  $\mathbb{R}_{om}$  and fix a unipotent group G.

For a lattice  $\Gamma \subseteq G$ , we use  $\pi_{\Gamma}$  to denote the projection  $\pi_{\Gamma} \colon G \to G/\Gamma$ . When no confusion arises, we omit the subscript  $\Gamma$ . Also, whenever  $\Gamma_0 \subseteq \Gamma$  is a subgroup of finite index, we let  $\pi_0 \colon G \to G/\Gamma_0$  denote the natural projection.

Given an  $\mathbb{R}_{om}$ -definable family  $\mathcal{F} = \{F_t : t \in I_{\infty}\}$ , for a lattice  $\Gamma \subseteq G$  we consider the possible Hausdorff limits at  $\infty$  of the family  $\pi(\mathcal{F}) \subseteq G/\Gamma$ . Notice that if  $\mathcal{F}$  is a constant family  $F_t = F$  then the only Hausdorff limit at  $\infty$  is the closure of  $\pi(F)$  and this case was handled in [9].

**Example 6.6.** (1) Consider first  $G = (\mathbb{R}^2, +)$  and  $\Gamma = \mathbb{Z}^2$ .

Let  $L_0$  be the line  $L_0 = \{(x,0) \in \mathbb{R}^2 : x \in \mathbb{R}\}$ , and  $\mathcal{F}_1$  be the family of  $L_0$ -translates:  $\mathcal{F}_1 = \{L_0 + (0,t) : t \in I_\infty\}$ . It is not hard to see that the Hausdorff limits at  $\infty$  of  $\pi(\mathcal{F}_1)$  are exactly the sets  $\pi(L_0 + q)$  for  $q \in G$ .

Let  $\mathcal{F}_2 = \{L_t : t \in I_\infty\}$  be the family of lines in G where  $L_t$  is the line  $L_t = \{(x, y) \in G : y = tx\}$ . It is not hard to see that the only Hausdorff limit at  $\infty$  is the whole  $G/\Gamma$ .

(2) Assume now that  $G = \mathbb{R}$ ,  $\Gamma = \mathbb{Z}$ , and let  $\mathcal{F} = \{t + [0, 2] : t \in I_{\infty}\}$ . The family  $\pi(\mathcal{F})$  is the constant family  $\pi(F_t) = \mathbb{R}/\mathbb{Z}$ , and hence this is the only Hausdorff limit at  $\infty$ . However, for any lattice  $\Gamma_0 \subseteq \mathbb{Z}$  with  $|\mathbb{Z} : \Gamma_0| \geq 3$ , the Hausdorff limits at  $\infty$  of  $\pi_0(\mathcal{F})$  are the sets of the form  $g + \pi_0([0, 2])$ , for  $g \in \mathbb{R}/\Gamma_0$ , and none of these equals  $\mathbb{R}/\Gamma_0$ .

**Definition 6.7.** Let  $\mathcal{F} = \{F_t : t \in I_{\infty}\}$  be an  $\mathbb{R}_{om}$ -definable family of subsets of a unipotent group G, and  $\Gamma \subseteq G$  be a lattice.

We say that the family  $\pi(\mathcal{F})$  converges strongly to  $G/\Gamma$  at  $\infty$  if  $\pi_0(\mathcal{F})$  converges to  $G/\Gamma_0$  at  $\infty$ , for any subgroup  $\Gamma_0 \subseteq \Gamma$  of finite index, as in Definition 6.2.

The next observation immediately follows from Proposition 6.5:

**Proposition 6.8.** Let  $\mathcal{F} = \{F_t : t \in I_\infty\}$  be an  $\mathbb{R}_{om}$ -definable family of subsets of a unipotent group G, and  $\Gamma \subseteq G$  be a lattice. Then:

- (1)  $G/\Gamma$  is a Hausdorff limit of  $\pi(\mathcal{F}) \Leftrightarrow \text{there exists } \tau \in \mathfrak{R}, \ \tau > \mathbb{R},$  such that  $F_{\tau}^{\sharp}$  is  $\Gamma$ -dense in G.
- (2)  $\pi(\mathcal{F})$  converges to  $G/\Gamma \Leftrightarrow \text{for all } \tau \in \mathfrak{R} \text{ with } \tau > \mathbb{R}, F_{\tau}^{\sharp} \text{ is } \Gamma\text{-dense} \text{ in } G.$
- (3)  $\pi(\mathcal{F})$  converges strongly to  $G/\Gamma \Leftrightarrow \text{for all } \tau \in \mathfrak{R}, \text{ with } \tau > \mathbb{R}, F_{\tau}^{\sharp}$  is strongly  $\Gamma$ -dense in G.

For the next result, recall the notation at the beginning of Section 4.4, regarding the abelianization of G.

Corollary 6.9. Let G be a unipotent group and let  $\mathcal{F} = \{F_t : t \in I_\infty\}$  be an  $\mathcal{L}_{om}$ -definable family of subsets of G. We denote by  $\mathcal{F}_{ab}$  the family  $\pi_{ab}(\mathcal{F})$  of subsets of  $G_{ab}$ . For a lattice  $\Gamma \subseteq G$ , we let  $\pi : G \to G/\Gamma$ , and  $\pi^* : G_{ab} \to G_{ab}/\Gamma_{ab}$  be the quotient maps.

Then, the family  $\pi(\mathcal{F})$  converges strongly to  $G/\Gamma$  at  $\infty$  if and only if  $\pi^*(\mathcal{F})$  converges strongly to  $G_{ab}/\Gamma_{ab}$  at  $\infty$ .

Proof. By Proposition 6.8,  $\pi(\mathcal{F})$  converges strongly to  $G/\Gamma$  at  $\infty$  if and only if for all  $\tau > \mathbb{R}$  in  $\mathfrak{R}$ ,  $F_{\tau}^{\sharp}$  is strongly Γ-dense in G. By Corollary 5.6, this is equivalent to  $\pi_{ab}(F_{\tau}^{\sharp})$  being strongly  $\Gamma_{ab}$ -dense in  $G_{ab}$ , for

all  $\tau > \mathbb{R}$ , which again, by Proposition 6.8, is equivalent to  $\pi_{ab}(\mathcal{F})$  strongly converging to  $G_{ab}/\Gamma_{ab}$  at  $\infty$ .

Before the next theorem we observe:

**Lemma 6.10.** Let G be a unipotent group and let  $\mathcal{F} = \{F_t : t \in I_\infty\}$  be an  $\mathbb{R}_{om}$ -definable family of subsets of G. Then, for all  $\tau, \tau' \in \mathfrak{R}$ , with  $\tau, \tau' > \mathbb{R}$ ,  $\mathscr{L}_{max}(F_{\tau}^{\sharp}) = \mathscr{L}_{max}(F_{\tau'}^{\sharp})$ .

*Proof.* We use the fact that  $\tau$  and  $\tau'$  have the same  $\mathcal{L}_{om}$ -type over  $\mathbb{R}$ . Clearly, it is enough to show  $\mathscr{L}(F_{\tau}^{\sharp}) = \mathscr{L}(F_{\tau'}^{\sharp})$ , and, by symmetry, it is sufficient to show  $\mathscr{L}(F_{\tau}^{\sharp}) \subseteq \mathscr{L}(F_{\tau'}^{\sharp})$ .

Let  $L \in \mathcal{L}(F_{\tau}^{\sharp})$ . We choose  $\alpha \in \operatorname{dcl}(\tau)$  such that the coset  $\alpha L$  is a nearest co-commutative coset to some type on  $F_{\tau}^{\sharp}$ . Let a(t) be an  $\mathbb{R}_{\text{om}}$ -definable function with  $a(\tau) = \alpha$ ,

By saturation of  $\mathfrak{R}_{om}$ , the coset  $a(\tau')L$  is a nearest co-commutative coset to some type on  $F_{\tau'}^{\sharp}$ .

The above lemma justifies the following definition

**Definition 6.11.** For an  $\mathbb{R}_{om}$ -definable family  $\mathcal{F} = \{F_t : t \in I_{\infty}\}$  of a unipotent group G, we denote by  $\mathscr{L}_{max}(\mathcal{F})$  the finite set of co-commutative subgroups  $\mathscr{L}_{max}(F_{\tau}^{\sharp})$ , for some (any)  $\tau > \mathbb{R}$ .

The next theorem is one of our main results.

**Theorem 6.12.** Let G be a unipotent group,  $\mathcal{F} = \{F_t : t \in I_\infty\}$  an  $\mathbb{R}_{om}$ -definable family of subsets of G.

For every lattice  $\Gamma \subseteq G$  we have:

- (1)  $L^{\Gamma} = G$  for some  $L \in \mathcal{L}_{max}(\mathcal{F})$  if and only if  $\pi(\mathcal{F})$  converges strongly to  $G/\Gamma$  at  $\infty$ .
- (2)  $L^{\Gamma} \neq G$  for all  $L \in \mathcal{L}_{\max}(\mathcal{F})$  if and only if there exists a subgroup  $\Gamma_0 \subseteq \Gamma$  of finite index such that all Hausdorff limits at  $\infty$  of  $\pi_0(\mathcal{F})$  are proper subsets of  $G/\Gamma_0$ .

Note that the condition given in (2) is formally stronger than the negation of the condition in (1), thus both need to be proved separately.

- Proof. (1) By Proposition 6.8,  $\pi(\mathcal{F})$  converges strongly to  $G/\Gamma$  if and only if for all  $\tau > \mathbb{R}$ ,  $F_{\tau}^{\sharp}$  is strongly Γ-dense in G, which by Theorem 5.5, is equivalent to  $L^{\Gamma} = G$  for some  $L \in \mathcal{L}_{\max}(\mathcal{F})$ .
- (2) Assume that for all  $L \in \mathcal{L}_{\text{max}}(\mathcal{F})$  we have  $L^{\Gamma} \neq G$ , and for contradiction assume that for every subgroup  $\Gamma_0 \subseteq \Gamma$  of finite index,  $G/\Gamma_0$  is one of the Hausdorff limits at  $\infty$ , of the family  $\pi_0(\mathcal{F})$ . Equivalently,

it follows from Proposition 6.5 and Lemma 4.11, that for every subgroup  $\Gamma_0 \subseteq \Gamma$  of finite index, there is a  $\tau' \in \mathfrak{R}$  with  $\tau' > \mathbb{R}$ , such that  $G = \operatorname{st}(F_{\tau'}^{\sharp} \cdot \Gamma_0^{\sharp})$ .

Claim 6.13. There exists  $\tau^* \in \mathfrak{R}$  with  $\tau^* > \mathbb{R}$  such that for every  $\Gamma_0 \subseteq \Gamma$  of finite index,  $\operatorname{st}(F_{\tau^*}^{\sharp} \cdot \Gamma_0^{\sharp}) = G$ .

Proof of the claim. For every  $g \in G = G(\mathbb{R})$ ,  $r \in \mathbb{R}^{>0}$  and  $\Gamma_0 \subseteq \Gamma$  of finite index, we consider the following formula  $\phi_{q,r,\Gamma_0}(t)$ :

$$t > r \& g \in B_{1/r}(e) \cdot F_t \cdot \Gamma_0$$
.

We let p(t) be the type consisting of all  $\phi_{g,r,\Gamma_0}$ , as  $g,r,\Gamma_0$  vary over all  $g \in G$ ,  $r \in \mathbb{R}^{>0}$  and  $\Gamma_0 \subseteq \Gamma$  of finite index, respectively. We claim that p is finitely consistent. Indeed, given finitely many subgroups of  $\Gamma$  of finite index, let  $\Gamma_1$  be their intersection. Clearly  $\Gamma_1$  has finite index in  $\Gamma$ . By our assumption, there is  $\tau > \mathbb{R}$  such that  $G(\mathbb{R}) \subseteq \mu \cdot F_{\tau}^{\sharp} \cdot \Gamma_1^{\sharp}$ , which implies that for any  $g_1, \ldots, g_k \in G$  and  $r_1, \ldots, r_k \in \mathbb{R}$ ,  $\phi_{g_i, r_i, \Gamma_1}(\tau)$  holds. It follows that p(t) is finitely consistent, so by the saturation of  $\mathfrak{R}_{\text{full}}$ , there exists  $\tau^* \in \mathfrak{R}$  realizing p(t).

Now, given  $\Gamma_0 \subseteq \Gamma$  of finite index, and  $g \in G$ , we have  $g \in B_{\epsilon}(e)^{\sharp} \cdot F_{\tau *}^{\sharp} \cdot \Gamma_0^{\sharp}$ , for all  $\epsilon \in \mathbb{R}^{>0}$ . Using saturation again, it follows that  $g \in \mu \cdot F_{\tau *}^{\sharp} \cdot \Gamma_0^{\sharp}$ , and hence  $G = \operatorname{st}(F_{\tau *}^{\sharp} \cdot \Gamma_0^{\sharp})$ , proving the claim.

For  $\tau^*$  as in the above claim,  $F_{\tau^*}^{\sharp}$  is strongly  $\Gamma$ -dense in G and therefore, by Theorem 5.5, there is  $L \in \mathscr{L}_{\max}(\mathcal{F})$  with  $L^{\Gamma} = G$ , contradiction.

The opposite implication of (2) follows from (1).

6.5. The abelian case. In the unipotent case, Theorem 6.12 tells us when the family  $\pi(\mathcal{F})$  converges (strongly) to  $G/\Gamma$ . In the abelian case we can say more about the possible Hausdorff limits of  $\pi(\mathcal{F})$ , due to the following theorem.

**Theorem 6.14.** Let  $G = (\mathbb{R}^m, +)$  and let  $\mathcal{F} = \{F_t : t \in I_\infty\}$  be an  $\mathbb{R}_{om}$ -definable family of subsets of G.

For every  $r \in \mathbb{R}^{>0}$ , there are subspaces  $L_1, \ldots, L_k \subseteq G$  (possibly with repetitions and with k depending on r), with  $\mathcal{L}_{\max}(\mathcal{F}) \subseteq \{L_1, \ldots, L_k\}$ , and there are  $\mathbb{R}_{\text{om}}$ -definable functions  $a_1(t), \ldots, a_k(t) \colon I_{\infty} \to G$ , such that

(1) For some  $\tau \in \mathfrak{R}, \tau > \mathbb{R}$ , each  $a_i(\tau) + L_i$  is a nearest coset to some type  $p \in S_{F^{\sharp}}(\tau)$ .

(2) For  $t \gg 0$ ,

$$F_t \subseteq \overline{B}_r + \bigcup_{i=1}^k a_i(t) + L_i.$$

(3) Let  $\Gamma \subseteq G$  be a lattice and  $\pi : G \to G/\Gamma$  the projection. For every  $s \in \mathbb{R}^{>0}$ , there exists  $t_s > 0$  such that for all  $t > t_s$ , and all  $i = 1, \ldots, k$ .

$$\pi(a_i(t) + L_i) \subseteq \pi(\overline{B}_s) + \pi(F_t).$$

*Proof.* Let  $\tau \in I_{\infty}^{\sharp}$  with  $\tau > \mathbb{R}$ , and let  $A = \operatorname{dcl}(\tau)$ .

Fix  $r \in \mathbb{R}^{>0}$ . Let  $L_1, \ldots, L_k \in \mathcal{L}_A(F_\tau^{\sharp})$  be subgroups and  $\alpha_1, \ldots, \alpha_k \in dcl(A)$  be as in Theorem 3.15. Thus we have

$$F_{\tau}^{\sharp} \subseteq \overline{B}_{r}^{\sharp} + \bigcup_{i=1}^{k} \alpha_{i} + L_{i}^{\sharp}, \quad \text{with } \mathscr{L}_{\max}(F_{\tau}^{\sharp}) \subseteq \{L_{1}, \dots, L_{k}\},$$

and, by Theorem 4.14, for every i = 1, ..., k, and s > 0 we also have

$$\alpha_i + L_i^{\sharp} \subseteq \overline{B}_s^{\sharp} + F_{\tau}^{\sharp} + \Gamma^{\sharp}.$$

Since each  $\alpha_i \in \operatorname{dcl}(\tau)$ , for  $i = 1, \ldots, k$ , we choose  $\mathbb{R}_{om}$ -definable functions  $a_i(t) \colon \mathbb{R} \to G$ , such that  $\alpha_i = a_i(\tau)$ . We have

$$F_{\tau}^{\sharp} \subseteq \bigcup_{i=1}^{k} \overline{B}_{r}^{\sharp} + a_{i}(\tau) + L_{i}^{\sharp}.$$

Since the  $\mathcal{L}_{om}$ -type of  $\tau$  over  $\mathbb{R}$  is implied by  $\{x > r \colon r \in \mathbb{R}\}$  and the above inclusion can be expressed by an  $\mathcal{L}_{om}$ -formula over  $\tau$ , we obtain that for  $t \gg 0$ ,

$$F_t \subseteq \bigcup_{i=1}^k \overline{B}_r + a_i(t) + L_i.$$

This proves (1) and (2).

Assume (3) fails. Then, there is a lattice  $\Gamma \subseteq G$  and  $s \in \mathbb{R}^{>0}$ , such that for some  $i_0 = 1, \ldots, k$ , the set

$$\{t \in I_{\infty} : \pi(a_i(t) + L_i) \nsubseteq \pi(\overline{B}_s) + \pi(F_t)\}$$

is unbounded in  $\mathbb{R}$ . Without loss of generality, we assume  $i_0 = 1$ .

Using saturation of  $\mathfrak{R}_{\text{full}}$ , we can find  $\tau' \in \mathfrak{R}$  with  $\tau' > \mathbb{R}$ , such that  $\pi(a_1(\tau') + L_1^{\sharp}) \nsubseteq \pi(\overline{B}_s^{\sharp}) + \pi(F_{\tau'}^{\sharp})$ , so in particular,

$$a_1(\tau') + L_1^{\sharp} \nsubseteq \overline{B}_s^{\sharp} + F_{\tau'}^{\sharp} + \Gamma^{\sharp}.$$

Since  $\tau$  and  $\tau'$  realize the same  $\mathcal{L}_{om}$ -type over  $\mathbb{R}$ , and  $a_1(\tau) + L_1^{\sharp}$  is the nearest coset to some type on  $F_{\tau}^{\sharp}$ , the coset  $a_1(\tau') + L_1^{\sharp}$  is the nearest coset to a type, call it p, on  $F_{\tau'}^{\sharp}$ . However, using Theorem 4.14, we have

$$a_1(\tau') + L_1^{\sharp} \subseteq \mu + a_1(\tau') + L_1^{\sharp} = \mu + p(\mathfrak{R}) + \Gamma^{\sharp} \subseteq \overline{B}_s^{\sharp} + F_{\tau'}^{\sharp} + \Gamma^{\sharp},$$

contradiction.

We can now show that, in the abelian case, every Hausdorff limit of  $\pi(\mathcal{F})$  at  $\infty$  is trapped between a finite union of cosets and a "thickening" of it.

Corollary 6.15. Let  $G = (\mathbb{R}^m, +)$  and let  $\mathcal{F} = \{F_t : t \in I_\infty\}$  be an  $\mathbb{R}_{om}$ -definable family of subsets of G.

For every  $r \in \mathbb{R}^{>0}$ , there are subspaces  $L_1, \ldots, L_k \subseteq G$  (possibly with repetitions and with k depending on r), with  $\mathcal{L}_{\max}(\mathcal{F}) \subseteq \{L_1, \ldots, L_k\}$ , such that for any lattice  $\Gamma \subseteq G$ , and any Hausdorff limit X of the family  $\pi(\mathcal{F})$  at  $\infty$ , there are  $g_1, \ldots, g_k \in G$  with

$$\pi\left(\bigcup_{i=1}^{k}(g_i+L_i)\right)\subseteq X\subseteq \pi(\overline{B}_r)+\pi\left(\bigcup_{i=1}^{k}(g_i+L_i)\right).$$

*Proof.* By Proposition 6.5,  $X = \pi(\operatorname{st}(F_{\tau}^{\sharp} + \Gamma^{\sharp}))$ , for some  $\tau > \mathbb{R}$ . We now apply Theorem 6.14 and obtain linear subspaces  $L_1, \ldots, L_k \subseteq \mathbb{R}^m$  and definable functions  $a_1(t), \ldots, a_k(t) \colon I_{\infty} \to G$  as in the theorem. Given a lattice  $\Gamma \subseteq G$ , clause (2) of that theorem implies that

$$F_{\tau}^{\sharp} + \Gamma^{\sharp} \subseteq \overline{B}_{r}^{\sharp} + \bigcup_{i=1}^{k} a_{i}(\tau) + L_{i}^{\sharp} + \Gamma^{\sharp},$$

while (3) (applied for all  $r \in \mathbb{R}^{>0}$ ) implies

$$\bigcup_{i=1}^{k} a_i(\tau) + L_i^{\sharp} + \Gamma^{\sharp} \subseteq \mu + F_{\tau}^{\sharp} + \Gamma^{\sharp}.$$

Using Lemma 5.2, for  $g_i \in \operatorname{st}(a_i(\tau) + \Gamma^{\sharp})$ , we obtain

$$\bigcup_{i=1}^k g_i + L_i^{\Gamma} + \Gamma = \bigcup_{i=1}^k \operatorname{st}(a_i(\tau) + L_i^{\sharp} + \Gamma^{\sharp}) \subseteq \operatorname{st}(F_{\tau}^{\sharp} + \Gamma^{\sharp}) \subseteq \overline{B}_r + \bigcup_{i=1}^k g_i + L_i^{\Gamma} + \Gamma.$$

Applying  $\pi$  the result follows.

#### 7. Polynomial dilations

Let  $G \subseteq GL(n, \mathbb{R})$  be a unipotent group. In several articles ([3], [1]) a particular type of families of subsets of G, given by dilations of an initial curve, was considered in the unipotent setting. We first make some definitions.

## 7.1. The setting.

**Definition 7.1.** A polynomial  $m \times k$  matrix  $M_t$  (over  $\mathbb{R}$ ) is a matrix  $M_t \in M_{m \times k}(\mathbb{R}[t])$ , namely, a matrix all of whose entries are polynomials in  $\mathbb{R}[t]$ . It can be written as  $\sum_{j=0}^{d} t^j A_j$ , with each  $A_j$  is an  $m \times k$  matrix over  $\mathbb{R}$ .

Let G be a unipotent group of dimension m. We identify the underlying vector space of its Lie algebra  $\mathfrak{g}$  with  $\mathbb{R}^m$ , and we let  $\exp: \mathfrak{g} \to G$  be the exponential map (which is a polynomial bijection with a polynomial inverse, see [2]). For a polynomial  $m \times k$  matrix  $M_t$  we consider the family of "dilations"  $\rho_t \colon \mathbb{R}^k \to G$  given by  $x \mapsto \exp(M_t x)$  and for a set  $X \subseteq \mathbb{R}^k$ , the family  $\{\rho_t(X) : t \in I_\infty\}$  of subsets of G.

In [3], the authors start with a measure  $\nu$  on  $\mathfrak{g}$  given as the pushforward of the Lebesgue measure on (0,1) via a real analytic map  $\phi:(0,1)\to\mathfrak{g}$ , then "dilate" the measure  $\nu$  using multiplication by a polynomial  $m\times m$  matrix  $M_t$ , and consider the limit of the measures as  $t\to\infty$ . In [3, Theorem 1.1] the authors prove that under some assumptions on  $\Phi=\mathrm{Image}(\phi)$ , given in terms of kernels of particular characters of G, the associated family of measures  $\mu_t$ , on  $G/\Gamma$  is "equidistributed", roughly saying that for any Borel  $D\subseteq G/\Gamma$ , the family  $\mu_t(D)$  converges to the canonical Haar measure of D. Translated to the topological language this would imply (but a-priori might not be equivalent) that the family  $\{\pi_{\Gamma}(\rho_t(\Phi)): t \in I_{\infty}\}$  converges to  $G/\Gamma$  at  $\infty$ .

In the current section we extend the topological corollary by replacing the one-dimensional set  $\Phi$  with an  $\mathbb{R}_{om}$ -definable set  $X \subseteq \mathbb{R}^k$  of arbitrary dimension. Using Theorem 6.12, one can formulate conditions, similar to those in [3], as to when the family  $\{\pi(\exp(M_t \cdot X)): t \in I_\infty\}$  converges strongly to  $G/\Gamma$  at  $\infty$ . Instead, we consider the special case, when the polynomial matrix  $M_t$  does not have a constant term, and describe first in full, in the abelian case, all possible Hausdroff limits at  $\infty$  of the family.

## 7.2. Polynomial dilations in vector spaces. We fix some notations.

- **Notation 7.2.** (1) We call a map  $a(t) : \mathbb{R} \to \mathbb{R}^m$  a polynomial map if  $a(t) = \sum_{i=0}^d t^i a_i$  for some  $a_0, \ldots, a_d \in \mathbb{R}^m$ . We say that the polynomial map is *proper* if  $a_0 = 0$ .
- (2) By a polynomial family of dilations we mean a family of maps  $\{\rho_t \colon \mathbb{R}^k \to \mathbb{R}^m \colon t \in I\}$  with  $I \subseteq \mathbb{R}$ , such that for some polynomial  $m \times k$ -matrix  $M_t$ , for all  $t \in I$  and  $x \in \mathbb{R}^k$ , we have  $\rho_t(x) = M_t x$ . We say that the family is proper if the constant term of  $M_t$  is the zero  $m \times k$ -matrix.
- (3) By a a polynomial family of cosets we mean a family  $\{a(t) + L : t \in I\}$ , where  $I \subseteq \mathbb{R}$ ,  $L \subseteq \mathbb{R}^m$  is a subspace, and  $a(t) : \mathbb{R} \to \mathbb{R}^m$  a polynomial map, We say that the family is proper if a(t) is a proper polynomial map.
- (4) By a polynomial family of multi-cosets we mean a family of the form  $\{\bigcup_{j=1}^{n} (a_j(t) + L_j) : t \in I\}$ , where  $I \subseteq \mathbb{R}$ , each  $a_j(t) : \mathbb{R} \to \mathbb{R}^m$  is a polynomial map and each  $L_j \subseteq \mathbb{R}^m$  is a subspace. We say that the polynomial family of multi-cosets is *proper* if each  $a_j(t)$  is proper.

We refer the readers to [3, Remark 2.5] for an example explaining the reason to work with proper polynomial dilations instead of general ones.

**Remark 7.3.** Let  $\mathcal{M} = \{\bigcup_{j=1}^{n} (a_j(t) + L_j) : t \in I\}$  be a polynomial family of multi-cosets. It is not hard to see, using Lemma 5.2(2), that  $\mathcal{L}_{\max}(\mathcal{M})$  is exactly the set of maximal (by inclusion) subspaces in  $\{L_1, \ldots, L_n\}$ .

Our main result is:

**Theorem 7.4.** Let  $\{\rho_t \colon \mathbb{R}^k \to \mathbb{R}^m \colon t \in I_\infty\}$  be a proper family of polynomial dilations,  $X \subseteq \mathbb{R}^k$  an  $\mathbb{R}_{om}$ -definable set, and  $\mathcal{F} = \{\rho_t(X) \colon t \in I_\infty\}$ .

Then, there is a proper polynomial family of multi-cosets

$$\mathcal{M} = \left\{ \bigcup_{j=1}^{n} (p_j(t) + L_j) \colon t \in I_{\infty} \right\}$$

such that

- (1)  $\rho_t(X) \subseteq \bigcup_{j=1}^n (p_j(t) + L_j)$  for all  $t \in \mathbb{R}$ ;
- (2)  $\mathscr{L}_{\max}(\mathcal{F}) = \mathscr{L}_{\max}(\mathcal{M});$
- (3) for any lattice  $\Gamma \subseteq \mathbb{R}^m$ , the families  $\pi(\mathcal{F})$  and  $\pi(\mathcal{M})$  have the same Hausdorff limits at  $\infty$ .

*Proof.* Let  $A_1, \ldots, A_d$  be  $m \times k$ -matrices such that  $\rho_t(x) = \sum_{i=1}^d t^i A_i x$ . We fix  $\tau \in \mathfrak{R}$  with  $\tau > \mathbb{R}$ , and let  $\mathcal{Y} = \rho_{\tau}(X^{\sharp})$ .

Applying Theorem 6.14 with r = 1, we obtain subspaces  $L_1, \ldots L_n \subseteq \mathbb{R}^m$  and  $\mathbb{R}_{om}$ -definable maps  $a_1(t), \ldots, a_n(t) \colon I_\infty \to \mathbb{R}^m$  such that

each  $a_i(\tau) + L_i$  is a nearest coset to some type in  $S_{\mathcal{Y}}(\tau)$ , and

(7.1) 
$$\rho_t(X) \subseteq \overline{B}_1 + \bigcup_{j=1}^n (a_j(t) + L_j) \text{ for } t \gg 0.$$

We pick  $a_i(t)$  and  $L_i$ ,  $j=1,\ldots,n$ , as above, with the minimal possible n.

We claim that  $\mathscr{L}_{\max}(\mathcal{F})$  is exactly the set of maximal elements of  $\{L_1, \ldots, L_n\}$ . Indeed, assume  $L \in \mathscr{L}_{\max}(\mathcal{F})$  and  $p \in S_{\mathcal{Y}}(\underline{\tau})$  is such that  $L_p = L$ . Then, for some  $j = 1, \ldots, n, p \vdash a_j(\tau) + L_j + \overline{B}_1$ . It follows from Lemma 3.8 that  $L \subseteq L_j$ , so by maximality  $L = L_j$ . Since every  $L_i$  is contained in some  $L \in \mathscr{L}_{\max}(\mathcal{F})$ , the claim follows.

For each j = 1, ..., n, let  $L_j^{\perp} \subseteq \mathbb{R}^m$  be the orthogonal complement to  $L_j$  with respect to the standard inner product on  $\mathbb{R}^m$ , and  $\pi_j^{\perp} : \mathbb{R}^m \to L_j^{\perp}$  be the corresponding projection, whose kernel is  $L_j$ .

Replacing each  $a_j(t)$  with  $\pi_j^{\perp}(a_j(t))$ , if needed, we assume  $a_j(t) \in L_j^{\perp}$  for all  $t \in I_{\infty}$ .

For  $j = 1, \ldots, n$ , let

$$X_j = \{x \in X : \text{ for } t \gg 0, \ \rho_t(x) \in \overline{B}_1 + a_j(t) + L_j\}.$$

Notice that each  $X_j$  is  $\mathbb{R}_{om}$ -definable,

$$X = \bigcup_{j=1}^{n} X_j,$$

and, by the minimality of n, each  $X_j$  is non-empty.

Claim A. For each j = 1, ..., n, and every i = 1, ..., d, the set  $A_iX_j$  is contained in a single coset of  $L_i$ .

Proof of Claim A. We fix  $j \in \{1, ..., n\}$ .

Clearly it is sufficient to show that for any i = 1, ..., d, every  $\mathbb{R}$ -linear function  $F: \mathbb{R}^m \to \mathbb{R}$  that vanishes on  $L_j$  is constant on  $A_i X_j$ .

Assume not, and for some  $\mathbb{R}$ -linear function  $F \colon \mathbb{R}^m \to \mathbb{R}$  vanishing on  $L_j$  there are  $r \in \{1, \ldots, d\}$  and  $b_1, b_2 \in X_j$  with  $F(A_r b_1) \neq F(A_r b_2)$ . Consider the map  $q(t) = F(\rho_t(b_1) - \rho_t(b_2))$ .

Since  $\rho_t(b_1) - \rho_t(b_2) = \sum_{i=1}^d t^i A_i(b_1 - b_2)$ , by linearity of F, we have

$$q(t) = \sum_{i=1}^{d} t^{i} F(A_{i}(b_{1} - b_{2})),$$

so q(t) is a polynomial map from  $\mathbb{R}^k$  to  $\mathbb{R}$ .

By our assumptions, the coefficient of  $t^r$  in q(t) is not zero, hence q(t) is a non-zero polynomial and  $\lim_{t\to\infty} |q(t)| = \infty$ .

On the other hand, by the definition of  $X_j$ , we have  $\rho_t(b_1) - \rho_t(b_2) \in L_j + \overline{B}_1 - \overline{B}_1$  for  $t \gg 0$ . Since F vanishes on  $L_j$ , we obtain  $q(t) \in F(\overline{B}_1 - \overline{B}_1)$ , which is compact set. A contradiction with  $\lim_{t\to\infty} |q(t)| = \infty$ . This finishes the proof of the claim.

Let  $j=1,\ldots,n$ . By Claim A, for  $i=1,\ldots,d$ , we may choose  $a_{ij} \in L_j^{\perp}$  with  $A_i X_j \subseteq a_{ij} + L_j$ . For  $x \in X_j$  and  $t \in I_{\infty}$  we have

$$\rho_t(x) = \sum_{i=1}^d t^i A_i x \in \sum_{i=1}^d t^i (a_{ij} + L_j) = \sum_{i=1}^d t^i a_{ij} + \sum_{i=1}^d t^i L_j.$$

Since  $L_j$  is closed under multiplication by scalars, setting  $p_j(t) = \sum_{i=1}^d t^i a_{ij}$  we obtain

(7.2) 
$$\rho_t(X_j) \subseteq p_j(t) + L_j$$
 and  $\rho_t(X) \subseteq \bigcup_{j=1}^n (p_j(t) + L_j)$  for all  $t \in \mathbb{R}$ .

Notice that each  $p_j(t)$  is a proper polynomial map, and takes values in  $L_j^{\perp}$ .

Let  $\mathcal{M}$  be the proper polynomial family of multi-cosets

$$\mathcal{M} = \left\{ \bigcup_{j=1}^{n} (p_j(t) + L_j) : t \in I_{\infty} \right\}.$$

By (7.2),  $\mathcal{M}$  satisfies clause (1) of the theorem, and by the explanation right after (7.1),  $\mathscr{L}_{\text{max}}(\mathcal{F}) = \mathscr{L}_{\text{max}}(\mathcal{M})$ , implying clause (2). Let us see that clause (3) also holds.

Claim B. For every j = 1, ..., n, there exists  $c_j \in L_j^{\perp}$ , such that  $c_j = \lim_{t \to \infty} (a_j(t) - p_j(t))$ .

*Proof of Claim B.* We fix  $j=1,\ldots,n$ , and choose  $b\in X_j$ . By the definition of  $X_j$ , we have

$$\rho_t(b) \in \overline{B}_1 + a_i(t) + L_i \text{ for } t \gg 0,$$

and by (7.2),

$$\rho_t(b) \in p_j(t) + L_j.$$

Since both  $a_i(t)$  and  $p_i(t)$  take values in  $L_i^{\perp}$ , we have

$$p_j(t) \in a_j(t) + \pi_j^{\perp}(\overline{B}_1).$$

Thus  $a_j(t) - p_j(t) \in \pi_j^{\perp}(\overline{B}_1)$ . The set  $\pi_j^{\perp}(\overline{B}_1)$  is compact, hence, by o-minimality,  $\lim_{t\to+\infty}(a_j(t)-p_j(t))$  exists and it belongs to  $L_j^{\perp}$ , call it  $c_j$ . This finishes the proof of Claim B.

Recall that each  $a_i(\tau) + L_i$  is a nearest coset to some  $q_i \in S_{\mathcal{Y}}(\tau)$ . It follows that  $p_i(\tau) + c_i + L_i$  is also a nearest coset to  $q_i$ .

Claim C. For every j = 1..., n, we have  $c_j = 0$ , namely the coset  $p_i(\tau) + L_i$  is a nearest coset to  $q_i$ .

*Proof of Claim C.* We proceed by reverse induction on  $\dim(L_i)$ , so we consider  $j_0 \in \{1, ..., n\}$  and assume that for all j with  $\dim(L_i) >$  $\dim(L_{j_0})$ , we already know that  $c_j = 0$ , so  $p_j(\tau) + L_j$  is a nearest coset

Since  $X = \bigcup_{j=1}^n X_j$ , there exists  $j_1 \in \{1, \ldots, n\}$  such that  $q_{j_0}$  lies on  $\rho_{\tau}(X_{i_1}^{\sharp})$ . From (7.2) we conclude

$$(7.3) q_{j_0} \vdash p_{j_1}(\tau) + L_{j_1}^{\sharp},$$

hence, by the definition of a nearest coset  $L_{i_0} \subseteq L_{i_1}$ .

If  $L_{j_0} \neq L_{j_1}$ , then  $\dim(L_{j_1}) > \dim(L_{j_0})$ , so by our induction assumption, the coset  $p_{j_1}(\tau) + L_{j_1}^{\sharp}$  is a nearest coset to  $q_{j_1}$ . Thus  $\mu + p_{j_1}(\tau) +$  $L_{j_1}^{\sharp} = \mu + a_{j_1}(\tau) + L_{j_1}$ , hence

$$a_{j_0}(t) + L_{j_0} \subseteq \overline{B}_1 + a_{j_1}(t) + L_{j_1}$$
, for  $t \gg 0$ .

It follows that

$$\rho_t(X) \subseteq \bigcup_{j=1}^n (a_j(t) + L_j + \overline{B}_1) \subseteq \bigcup_{\substack{1 \le j \le n \\ i \ne j_0}} (a_j(t) + L_j + \overline{B}_1),$$

contradicting minimality of n.

Thus, we must have  $L_{j_0} = L_{j_1}$ . From the definition of a nearest coset and (7.3), we get

$$\mu + p_{j_0}(\tau) + c_{j_0} + L_{j_0}^{\sharp} = \mu + p_{j_1}(\tau) + L_{j_1}^{\sharp}.$$

Since  $c_{j_0} \in L_{j_1}^{\perp}$ , and both  $p_{j_1}(t)$ ,  $p_{j_0}(t)$  take values in  $L_{j_1}^{\perp}$ , it follows that  $\lim_{t\to\infty}(p_{j_1}(t)-p_{j_0}(t))=c_{j_0}$ . Recall that both polynomial  $p_{j_1}(t)$  and  $p_{j_0}(t)$  have zero constant terms, so  $c_{j_0} = 0$ . 

This finishes the proof of Claim C.

In order to prove that  $\mathcal{M}$  satisfies clause (3), we need to show that for every lattice  $\Gamma \subseteq \mathbb{R}^n$ , and  $\pi : \mathbb{R}^n \to \mathbb{R}^n/\Gamma$ , the families  $\pi(\mathcal{F})$  and  $\pi(\mathcal{M})$ have the same Hausdorff limits at  $\infty$ . Equivalently, by Proposition 6.5, it is sufficient to show, for any  $\tau > \mathbb{R}$ ,

$$\operatorname{st}(\rho_{\tau}(X^{\sharp}) + \Gamma^{\sharp}) = \operatorname{st}\left(\bigcup_{j=1}^{n} p_{j}(\tau) + L_{j}^{\sharp} + \Gamma^{\sharp}\right).$$

We fix  $\tau > \mathbb{R}$ .

By clause (1), we clearly have the left-to-right inclusion. For the opposite inclusion, for each  $j=1,\ldots,n$ , by Claim C, there exists a type  $q_j \in S_{\mathcal{Y}}(\tau)$ , whose nearest coset is  $p_j(\tau) + L_j$ . By Theorem 4.14,  $\operatorname{st}(q_j(\mathfrak{R}) + \Gamma^{\sharp}) = \operatorname{st}(p_j(\tau) + L_j + \Gamma^{\sharp})$ , thus  $\operatorname{st}(\mathcal{Y} + \Gamma^{\sharp}) \supseteq \operatorname{st}(\bigcup_{j=1}^n p_j(\tau) + L_j^{\sharp} + \Gamma^{\sharp})$ .

This ends the proof of Theorem 7.4.

7.2.1. Hausdorff limits of families of multi-cosets. In this section we describe Hausdorff limits of families of multi-cosets in real tori. Together with Theorem 7.4 it provides a complete description of the Hausdorff limits at  $\infty$  of proper families of polynomial dilations.

Let  $a_1(t), \ldots, a_n(t) \colon I_{\infty} \to \mathbb{R}^m$  be  $\mathbb{R}_{om}$ -definable functions, and let  $L_1, \ldots, L_n \subseteq \mathbb{R}^m$  be linear subspaces.

For  $t \in I_{\infty}$ , let  $\mathcal{M}_t$  be the multi-coset  $\mathcal{M}_t = \bigcup_{i=1}^n (a_i(t) + L_i)$ , and let  $\mathcal{M} = \{\mathcal{M}_t : t \in I_{\infty}\}$  be the corresponding family of multi-cosets. Notice, we do not assume that  $a_i(t)$  are polynomials.

Let  $\Gamma \subseteq \mathbb{R}^m$  be a lattice. We denote by  $\Gamma^n \subseteq (\mathbb{R}^m)^n$  the lattice obtained by the *n*-fold cartesian power of  $\Gamma$ , and as before, for a subspace  $V \subseteq (\mathbb{R}^m)^n$ , we denote by  $V^{\Gamma^n}$  the smallest linear  $\Gamma^n$ -rational subspace of  $(\mathbb{R}^m)^n$  containing V. For the quotient map  $\pi \colon \mathbb{R}^m \to \mathbb{R}^m/\Gamma$ , we denote by  $\pi^{(n)}$  the quotient map  $\pi^{(n)} \colon (\mathbb{R}^m)^n \to (\mathbb{R}^m)^n/\Gamma^n$ .

**Theorem 7.5.** Let  $a_1(t), \ldots, a_n(t), L_1, \ldots, L_n$  and  $\mathcal{M}$  be as above. Then, there exists a coset  $\bar{c} + V$  of a linear subspace  $V \subseteq (\mathbb{R}^m)^n$ , such that for every lattice  $\Gamma \subseteq \mathbb{R}^m$ , the family of Hausdorff limits of  $\pi_{\Gamma}(\mathcal{M})$  at  $\infty$  is exactly the family

$$\left\{ \pi_{\Gamma} \left( \bigcup_{i=1}^{n} (d_i + L_i^{\Gamma}) \right) : (d_1, \dots, d_n) \in \bar{c} + V^{\Gamma^n} \right\}.$$

*Proof.* Consider the  $\mathbb{R}_{om}$ -definable curve  $\sigma: I_{\infty} \to (\mathbb{R}^m)^n$ , given by  $\sigma(t) = (a_1(t), \ldots, a_n(t))$ . Let  $p(x) \in S(\mathbb{R})$  be the unique o-minimal type on  $\sigma(t)$  at  $\infty$ , whose realization is the set

$$p(\mathfrak{R}) = \{ \sigma(\tau) \colon \tau \in \mathfrak{R}, \tau > \mathbb{R} \} \subseteq (\mathfrak{R}^m)^n.$$

Let  $\bar{c} + V$  be a nearest coset to p(x). Since p is a type over  $\mathbb{R}$ , we have  $\bar{c} = (c_1, \ldots, c_n)$ , with each  $c_i$  in  $\mathbb{R}^m$ , and  $V \subseteq (\mathbb{R}^m)^n$  is a subspace. We claim that this coset satisfies the conclusion of the theorem.

Let  $\Gamma \subseteq \mathbb{R}^m$  be a lattice, and  $X \subseteq \mathbb{R}^m/\Gamma$ . We let  $\pi = \pi_{\Gamma}$ . We denote by  $\mathcal{H}_{\Gamma}$  the set of Hausdorff limits at  $\infty$  of  $\pi(\mathcal{M})$ 

Since  $\pi$  is  $\Gamma$ -invariant, it is sufficient to show that  $X \in \mathcal{H}_{\Gamma}$  if and only if

$$X = \pi \left( \bigcup_{i=1}^{n} (d_i + L_i^{\Gamma}) \right) \text{ for some } (d_1, \dots, d_n) \in \bar{c} + V^{\Gamma^n} + \Gamma^n.$$

Using Proposition 6.5, we have that  $X \in \mathcal{H}_{\Gamma}$  if and only if

$$X = \pi \left( \operatorname{st} \bigcup_{i=1}^{n} (a_i(\tau) + L_i^{\sharp} + \Gamma^{\sharp}) \right) \text{ for some } \tau \in \mathfrak{R} \text{ with } \tau > \mathbb{R}.$$

Thus  $X \in \mathcal{H}_{\Gamma}$  if and only if

$$X = \pi \left( \bigcup_{i=1}^{n} \operatorname{st}(\alpha_{i} + L_{i}^{\sharp} + \Gamma^{\sharp}) \right) \text{ for some } (\alpha_{1}, \dots, \alpha_{n}) \in p(\mathfrak{R}).$$

Let  $\alpha_1, \ldots, \alpha_n \in \Re^n$ . By Lemma 5.2, for every  $i = 1, \ldots, n$ , the set  $\operatorname{st}(\alpha_i + \Gamma^{\sharp})$  is non-empty, and for any  $d_i \in \operatorname{st}(\alpha_i + \Gamma^{\sharp})$  we have  $\operatorname{st}(\alpha_i + L_i^{\sharp} + \Gamma^{\sharp}) = d_i + L_i^{\Gamma} + \Gamma.$ Also, clearly, for any  $d_1, \ldots, d_n \in \mathbb{R}^m$  and  $\alpha_1, \ldots, \alpha_n \in \mathfrak{R}^m$  we have

$$\bigwedge_{i=1}^{n} d_i \in \operatorname{st}(\alpha_i + \Gamma^{\sharp}) \iff (d_1, \dots, d_n) \in \operatorname{st}((\alpha_1, \dots, \alpha_n) + (\Gamma^n)^{\sharp}).$$

It follows that  $X \in \mathcal{H}_{\Gamma}$  if and only if

$$X = \pi \left( \bigcup_{i=1}^{n} (d_i + L_i^{\Gamma}) \right) \text{ for some } (d_1, \dots, d_n) \in \operatorname{st} \left( p(\mathfrak{R}) + (\Gamma^n)^{\sharp} \right).$$

By Theorem 4.14 (over the parameter set  $\mathbb{R}$ ),

$$\operatorname{st}(p(\mathfrak{R}) + (\Gamma^n)^{\sharp}) = \operatorname{st}(\bar{c} + V^{\sharp} + (\Gamma^n)^{\sharp}),$$

and by Lemma 2.1(2), the set on the right equals  $cl(\bar{c} + V + \Gamma)$ . By Fact 4.5, we have

$$\operatorname{st}(p(\mathfrak{R}) + (\Gamma^n)^{\sharp}) = \bar{c} + V^{\Gamma^n} + \Gamma^n.$$

This finishes the proof of the theorem.

Theorem 7.4 and Theorem 7.5 immediately yield the definability of the family of Hausodrff limits at  $\infty$  of proper families of polynomial dilations, in the following sense.

Corollary 7.6. Let  $\{\rho_t \colon \mathbb{R}^k \to \mathbb{R}^m \colon t \in I_\infty\}$  be a proper family of polynomial dilations, and  $X \subseteq \mathbb{R}^k$  an  $\mathbb{R}_{om}$ -definable set.

Then there are linear subspaces  $L_1, \ldots, L_n \subseteq \mathbb{R}^n$ , and there is a coset of a linear space  $\bar{c} + V \subseteq (\mathbb{R}^m)^n$  such that for any lattice  $\Gamma \subseteq \mathbb{R}^m$ , the set of Hausdorff limits at  $\infty$  of the family  $\{\pi \circ \rho_t(X) : t \in I_\infty\}$  is exactly the family

$$\left\{\pi\left(\bigcup_{i=1}^{n}(d_i+L_i^{\Gamma})\right):(d_1,\ldots,d_n)\in\bar{c}+V^{\Gamma^n}\right\}.$$

In particular, it is the projection under  $\pi$  of a definable family of subsets of  $\mathbb{R}^m$ .

- 7.3. Polynomial dilations in unipotent groups. We now consider the case of general unipotent groups. For the setting we refer to Section 7.1
- 7.3.1. Abelinization of a family of dilations. Following [3, Section 1.5] we introduce the abelinization of a family of dilations.

Let G be a unipotent group of dimension d,  $G_{ab} = G/[G, G]$  its abelianization and  $\pi_{ab} \colon G \to G_{ab}$  the projection map. Since  $G_{ab}$  is an abelian unipotent group, we identify it with  $(\mathbb{R}^m, +)$  for  $m = \dim(G_{ab})$ . We also identify the Lie algebra  $\mathfrak{g}_{ab}$  with  $(\mathbb{R}^m, +)$ , and assume that the exponential map  $\exp: \mathfrak{g}_{ab} \to G_{ab}$  is the identity map.

Let  $d\pi_{ab}$  be the differential of  $\pi_{ab}$  at the identity  $e \in G$ . We have the following commutative diagram with polynomial maps.

$$G \xrightarrow{\pi_{ab}} G_{ab}$$

$$exp \qquad id$$

$$g \xrightarrow{d\pi_{ab}} g = \mathbb{R}^m$$

Let  $M_t$  be a polynomial  $d \times k$  matrix  $M_t$ , and  $\{\rho_t \colon \mathbb{R}_k \to G \colon t \in I_\infty\}$  the corresponding polynomial family of dilations. For  $t \in I_\infty$  we denote by  $L_t \colon \mathbb{R}^k \to \mathbb{R}^d$  the linear map  $x \mapsto M_t x$ . Thus  $\rho_t = \exp \circ L_t$ , and the following diagram is commutative

with  $\rho_t^{ab} = \pi_{ab} \circ \rho_t = d\pi_{ab} \circ L_t$ .

Since  $d\pi_{ab}$  is a linear map, there is a  $d \times m$  matrix D such that  $d\pi_{ab} : x \mapsto Dx$ . Thus

$$\rho_t^{\rm ab} \colon x \to D(M_t x) = (DM_t) x.$$

It is not hard to see that  $DM_t$  is a polynomial matrix, hence we obtain that the family  $\{\rho_t^{ab} \colon \mathbb{R}^k \to G_{ab} \colon t \in I_{\infty}\}$  is a polynomial family of dilations, and it is proper if the original family  $\{\rho_t \colon t \in I_{\infty}\}$  was.

We call the family  $\{\rho_t^{ab}: t \in I_{\infty}\}$  the abelinization of the family  $\{\rho_t: t \in I_{\infty}\}.$ 

We are now ready to prove a strong version of Theorem 6.12 for proper polynomial dilations.

**Theorem 7.7.** Let G be a unipotent group,  $\{\rho_t \colon \mathbb{R}^k \to G \colon t \in I_\infty\}$  a proper family of polynomial dilations,  $X \subseteq \mathbb{R}^k$  an  $\mathbb{R}_{om}$ -definable set, and let  $\mathcal{F}$  be the family

$$\mathcal{F} = \{ \rho_t(X) \subseteq G \colon t \in I_{\infty} \}.$$

Then, for every lattice  $\Gamma \subseteq G$  the following conditions are equivalent:

- (a)  $L^G = G$  for some  $L \in \mathscr{L}_{\max}(\mathcal{F})$ .
- (b)  $\pi(\mathcal{F})$  converges strongly to  $G/\Gamma$  at  $\infty$ .
- (c)  $\pi(\mathcal{F})$  converges to  $G/\Gamma$  at  $\infty$ .
- (d)  $G/\Gamma$  is a Hausdorff limit at  $\infty$  of the family  $\pi(\mathcal{F})$ .

*Proof.* By Theorem 6.12, (a) snd (b) are equivalent.

 $(b) \Rightarrow (c)$  and  $(c) \Rightarrow (d)$  are obvious.

We are left to show that  $(d) \Rightarrow (a)$ .

Assume (d) holds. i.e.  $G/\Gamma$  is one of the Hausdorff limits at  $\infty$  of the family  $\pi(\mathcal{F})$ .

Let  $\pi^*$  be the quotient map  $\pi^*: G_{ab} \to \Gamma_{ab}$ , and  $\mathcal{F}^{ab}$  be the family

$$\mathcal{F}^{ab} = \{ \rho_t^{ab}(X) \subseteq G_{ab} \colon t \in I_{\infty} \}.$$

Applying abelianization, we obtain that  $G_{ab}/\Gamma_{ab}$  is a Hausdorff limit at  $\infty$  of the family  $\pi^*(\mathcal{F}^{ab})$ .

Let  $\mathcal{M} = \{\bigcup_{j=1}^{n} (p_j(t) + L_j) : t \in I_{\infty}\}$  be a proper polynomial family of multi-cosets of  $G_{ab}$ , as in Theorem 7.5 applied to the family  $\mathcal{F}^{ab}$ . Then  $G_{ab}/\Gamma_{ab}$  is a Hausdorff limit at  $\infty$  of  $\mathcal{M}$ , and hence (e,g. by Lemma 5.2(1) and Lemma 6.4)

$$G_{\mathrm{ab}} = \Gamma_{\mathrm{ab}} + \bigcup_{j=1}^{n} (a_j + L_j^{\Gamma_{\mathrm{ab}}}),$$

for some  $a_1, \ldots, a_n \in G_{ab}$ .

Since  $\Gamma_{ab}$  is a discrete subgroup, it follows that  $G_{ab} = L_k^{\Gamma_{ab}}$  for some  $k \in \{1, ..., n\}$ . By Fact 4.4, it follows then that  $G_{ab} = L_k^{\Gamma_0}$  for any subgroup  $\Gamma_0 \subseteq \Gamma_{ab}$  of finite index. Thus  $\pi^*(\mathcal{M})$  converges strongly to  $G_{ab}/\Gamma_{ab}$  at  $\infty$ , and hence, by the choice of  $\mathcal{M}$ , the family  $\pi^*(\mathcal{F}^{ab})$  converges strongly at  $\infty$  to  $G_{ab}/\Gamma_{ab}$  as well.

By Corollary 6.9,  $\pi(\mathcal{F})$  converges strongly at  $\infty$  to  $G/\Gamma$ . Thus (b) holds.

This finishes the proof of the theorem.

Remark 7.8. Theorem 7.7 can be compared to [3, Theorem 1.3]. The latter is an equidistribution result on measures which are associated to polynomial dilations of real analytic curves in nilmanifolds. The set  $\mathcal{L}_{\text{max}}$  in our analysis is replaced there by (a-priori infinitely many) kernels of characters. The equidistribution of the measures implies the convergence of the family to  $G/\Gamma$ , under the appropriate assumptions. Our additional input, under the assumption of  $\mathbb{R}_{\text{om}}$ -definability, is the treatment of higher dimensional sets, as well as the fact that the sets in  $\mathcal{L}_{\text{max}}$  work for all lattices.

#### References

- [1] Michael Björklund and Alexander Fish, Equidistribution of dilations of polynomial curves in nilmanifolds, Proc. Amer. Math. Soc. 137 (2009), no. 6, 2111–2123. MR2480293
- [2] Lawrence J. Corwin and Frederick P. Greenleaf, Representations of nilpotent Lie groups and their applications. Part I, Cambridge Studies in Advanced Mathematics, vol. 18, Cambridge University Press, Cambridge, 1990. Basic theory and examples. MR1070979
- [3] Bryna Kra, Nimish A. Shah, and Wenbo Sun, Equidistribution of dilated curves on nilmanifolds, J. Lond. Math. Soc. (2) 98 (2018), no. 3, 708–732. MR3893198
- [4] David Marker, Model theory, Graduate Texts in Mathematics, vol. 217, Springer-Verlag, New York, 2002. An introduction. MR1924282
- [5] Louis Narens, Topologies of closed subsets, Trans. Amer. Math. Soc. 174 (1972), 55-76. MR312450
- [6] Margarita Otero, A survey on groups definable in o-minimal structures, Model theory with applications to algebra and analysis. Vol. 2, 2008, pp. 177–206. MR2436142
- [7] Ya'acov Peterzil and Sergei Starchenko, Topological groups,  $\mu$ -types and their stabilizers, J. Eur. Math. Soc. (JEMS) **19** (2017), no. 10, 2965–2995. MR3712999
- [8] \_\_\_\_\_\_, Algebraic and o-minimal flows on complex and real tori, Adv. Math. 333 (2018), 539–569. MR3818086
- [9] \_\_\_\_\_, o-minimal flows on nilmanifolds, Duke Math. J. 170 (2021), no. 18, 3935–3976. MR4348231
- [10] Anand Pillay, On groups and fields definable in o-minimal structures, J. Pure Appl. Algebra 53 (1988), no. 3, 239–255. MR961362

- [11] Marina Ratner, Strict measure rigidity for unipotent subgroups of solvable groups, Invent. Math. 101 (1990), no. 2, 449–482. MR1062971
- [12] Markus Stroppel, *Locally compact groups*, EMS Textbooks in Mathematics, European Mathematical Society (EMS), Zürich, 2006. MR2226087
- [13] Lou van den Dries, *Tame topology and o-minimal structures*, London Mathematical Society Lecture Note Series, vol. 248, Cambridge University Press, Cambridge, 1998. MR1633348
- [14] \_\_\_\_\_\_, Limit sets in o-minimal structues, O-minimal structures: Lisbon 2003; proceedings of a summer school by the european research and training network raag, 2005.

University of Haifa

Email address: kobi@math.haifa.ac.il

University of Notre Dame *Email address*: sstarche@nd.edu