

ON EXACT PRODUCTS OF TWO DIHEDRAL GROUPS

KAN HU AND HAO YU*

ABSTRACT. An exact product of two finite groups H and K is a finite group X which contains H and K as subgroups, satisfying $X = HK$ and $H \cap K = \{1_X\}$. In this paper, we provide a classification of the exact products of two dihedral groups of orders $2m$ and $2n$ for all odd numbers $m, n \geq 3$.

1. INTRODUCTION

Throughout the paper, groups considered are all finite. A group X is *factorizable* if X contains two subgroups H and K such that $X = HK$. In this case, we say that H and K furnish a *factorization*. The factorization $X = HK$ is *proper* if both H and K are non-trivial subgroups of X , and *maximal* if both H and K are maximal subgroups of X . Moreover, the factorization is *exact* if $H \cap K = \{1_X\}$, and in this case we also say that X is an *exact product* of H and K . The following factorization problem arises naturally: For a given group X , determine whether or not X admits a proper factorization, and if so, find all proper factorizations of X ; in particular, identify all maximal or exact factorizations of X .

In this paper we consider the factorization problem formulated in a slightly different way:

Problem 1. For two specified finite groups H and K , describe and classify the products, or more specifically, the exact products $X = HK$ of H and K .

Several famous results on this problem were obtained by some pioneering group theorists in the 20th century. Itô established that products of two abelian groups are metabelian [19], Douglas proved that products of two cyclic groups are supersolvable [8]. Meanwhile Wielandt and Kegel [24, 34] showed that products of two nilpotent groups are necessarily solvable.

Recent interest in exact products of two finite groups has focused on primitive permutation groups containing a regular subgroup and factorizations of finite almost simple groups [30, 28, 29, 35]. Hering, Liebeck and Saxl classified factorizations of exceptional groups of Lie type [12], and the landmark work by Liebeck, Praeger, and Saxl classified maximal factorizations of almost simple groups [29]. Recently, all exact factorizations of the almost simple groups have been classified by Li, Wang and Xia [28].

A group is *bicyclic* if it is a product of two cyclic subgroups. One of the longstanding unsolved problems in this field is the classification of finite bicyclic groups. Besides the aforementioned classical result by Douglas on supersolvability of bicyclic groups, Huppert established that every bicyclic p -group is metacyclic if $p > 2$ is an odd prime [17]. However, it is well known that non-metacyclic bicyclic 2-groups exist. The metacyclic

2010 *Mathematics Subject Classification.* 05C10, 05C25, 57M15.

Key words and phrases. group factorization, exact product, dihedral group, regular subgroup.

This research was supported by National Natural Science Foundation of China (11801507, 12071312).

* Corresponding author e-mail: 3485676673@qq.com .

p -groups were classified by Xu and Newman for odd prime p [36], and by Xu and Zhang for $p = 2$. The bicyclic 2-groups were eventually classified by Janko [21]. Some results directly related to bicyclic groups can be found in the context of regular embeddings of complete bipartite graphs into orientable surfaces [23], and in the context of Hopf algebras arising from exact factorizations of bicyclic groups [1, 2]. To the best of our knowledge it remains a challenging open problem in the general case.

The interest in exact factorization with a cyclic factor extends to combinatorics. A group X is called a *cyclic complementary extension* of H if X is an exact product of H and a cyclic complement $K = \langle c \rangle$; X is also said to admit a *cyclic complementary factorization*. Cyclic complementary extensions of a given group H have been intensively investigated in the context of regular Cayley maps, and more generally, within the theory of skew morphisms. Without going into the details, a *Cayley map* $M := \text{CM}(H, S, \rho)$ is an embedding of an (undirected, simple, and connected) Cayley graph $\Gamma := \text{Cay}(H, S)$ on a finite group H into an oriented surface, where the local rotation R at each vertex is defined uniformly by setting

$$R(h, hs) = R(h, h\rho(s)), h \in H, s \in S,$$

where S is an inverse-closed generating set of H with $1_H \notin S$, and ρ is a full cycle on S . It is well known that every Cayley map contains a vertex-regular subgroup of automorphisms. In the case where it is also arc-regular, it is termed a *regular Cayley map*. It turns out that the automorphism group $X := \text{Aut}(M)$ of a regular Cayley map M has a cyclic complementary factorization $X = H_L X_{1_H}$, where H_L is the left regular representation of H , and X_{1_H} is the (cyclic) stabilizer of the vertex 1_H of the map. Therefore, if we identify H with H_L , the group X can be regarded as a cyclic complementary extension of H .

Every cyclic complementary extension $X = HK$ of a finite group H by a cyclic complement $K = \langle c \rangle \cong C_n$ gives rise to a permutation φ on H , called a *skew morphism* of H , and an integer-valued function $\Pi : H \rightarrow \mathbb{Z}_n$ on H , called the *extended power function* of φ , which satisfy the following defining conditions:

- (a) $\varphi(1_H) = 1_H$, and $\varphi(gh) = \varphi(g)\varphi^{\Pi(g)}(h)$ for all $g, h \in H$;
- (b) $\Pi(1_H) = 1_H$, and $\Pi(gh) = \sum_{i=1}^{\Pi(g)} \Pi(\varphi^{i-1}(h)) \pmod{n}$ for all $g, h \in H$.

Remarkably, the cyclic complementary extension of H by K can be recovered from φ and Π in a canonical way [13]. In particular, a Cayley map $\text{CM}(H, S, \rho)$ is regular if and only if ρ extends to a skew morphism of H [20, Theorem 1]. In this context the cyclic complementary extensions of various families of groups have been investigated, including cyclic groups [6, 7, 16], dihedral groups [14, 26, 39], simple groups [3] and characteristically simple groups [5, 9, 10, 11], among others.

The main concern of the current paper is exact products of two dihedral groups, which have important applications in unoriented regular Cayley maps [25, 16, 38]. Our main result is the following theorem, which determines the exact products of two non-abelian dihedral groups $H \cong D_{2n}$ and $K \cong D_{2m}$ for all odd numbers $m, n \geq 3$

Theorem 1. *If $n, m \geq 3$ are odd numbers, then every exact product $X = HK$ of two dihedral groups $H = \langle x \rangle \rtimes \langle y \rangle \cong D_{2n}$ and $K = \langle z \rangle \rtimes \langle w \rangle \cong D_{2m}$ has a presentation*

$$\begin{aligned} X = \langle x, y, z, w \mid x^n = y^2 = z^m = w^2 = [x, z] = 1, x^y = x^{-1}, z^w = z^{-1}, \\ [x, w] = x^{sn_1} z^{bm_1}, [z, y] = x^{rn_1} z^{am_1}, [y, w] = x^{tn_1} z^{cm_1} \rangle, \end{aligned} \quad (1)$$

where m_1 and n_1 are positive divisors of m and n such that $\langle z \rangle_X = \langle z^{m_1} \rangle$ and $\langle x \rangle_X = \langle x^{n_1} \rangle$, the integers $a, b, c \in \mathbb{Z}_{m/m_1}$ and $r, s, t \in \mathbb{Z}_{n/n_1}$ satisfy the following conditions:

- (a) $a(2 + am_1) \equiv b(2 + am_1) \equiv c(2 + am_1) \equiv 0 \pmod{m/m_1}$,
- (b) $r(2 + sn_1) \equiv s(2 + sn_1) \equiv t(2 + sn_1) \equiv 0 \pmod{n/n_1}$,
- (c) $bk \equiv 0 \pmod{m/m_1} \iff k \equiv 0 \pmod{n_1}$, for any integer k ,
- (d) $rk \equiv 0 \pmod{n/n_1} \iff k \equiv 0 \pmod{m_1}$, for any integer k .

The paper is organized as follows. In Section 2 we gather all necessary notations, terminology, and known results from group theory for future reference. Section 3 presents important characterizations on exact products of two dihedral groups, while the last Section 4 is dedicated to proving Theorem 1.

2. PRELIMINARIES

Throughout the paper, we denote the ring of residue classes of integers modulo a positive integer n by \mathbb{Z}_n . We also use C_n, D_{2n}, A_n and S_n to represent the cyclic group of order n , the dihedral group of order $2n$, the alternating group, and the symmetric group of degree n , respectively. Additionally, we employ $GL(n, q)$, $PGL(n, q)$, $PSL(n, q)$, and $AGL(n, q)$ to denote the general linear group, the projective linear group, the projective special linear group, and the affine group, respectively, of dimension n over a finite field \mathbb{F}_q with $q := p^e$ elements.

If G is a group and H is a subset of G , we denote the normalizer and centralizer of H in G by $N_G(H)$ and $C_G(H)$, respectively. The following N/C theorem is well known.

Proposition 2 ([17]). *If H is a subgroup of a group G , then $N_G(H)/C_G(H)$ is isomorphic to a subgroup of $\text{Aut}(H)$.*

A finite group G is *solvable* if it contains a finite sequence of (subnormal) subgroups

$$1 = N_0 \leq N_1 \leq \dots \leq N_s = G$$

such that $N_{i-1} \triangleleft N_i$ and N_i/N_{i-1} is cyclic of prime order for all $1 \leq i \leq s$. In general, if $N \triangleleft G$ is a normal subgroup of a group G with $F := G/N$, then we say that G is an *extension* of N by F . In particular, if the quotient group F is cyclic, then the extension is *cyclic*. The following theorem on cyclic extensions is important for solvable groups.

Proposition 3 ([18, Theorem 3.36]). *Let m be a positive integer, and N a group with $a \in N$ and $\sigma \in \text{Aut}(N)$. If $a^\sigma = a$ and $x^{\sigma^m} = x^a$ for all $x \in N$, then there exists a group G , unique up to isomorphism, and having N as a normal subgroup with the following properties:*

- (a) $G/N = \langle gN \rangle$ is cyclic of order m ;
- (b) $g^m = a$;
- (c) $x^\sigma = x^g$.

Let G be a finite permutation group acting on a finite set Ω . If for any $\alpha, \beta \in \Omega$ there exists an element $g \in G$ such that $\beta = \alpha^g$, then G is *transitive* on Ω . Additionally, if such an element g exists uniquely, then G is *regular* on Ω . A subset $\Delta \subseteq \Omega$ is a *block* for G if $\{\Delta^g | g \in G\}$ is a partition of Ω . A transitive permutation group G on Ω is *primitive* if the only blocks for G are the singletons and the whole set Ω , and *quasiprimitive* if every nontrivial normal subgroup of G is transitive. It is well known that every primitive group is quasiprimitive.

Proposition 4. [23, 27] *Let X be a quasiprimitive permutation group of degree n . If X contains a cyclic regular subgroup H . Then exactly one of the following holds:*

- (a) $C_p \cong H \leq X \leq \text{AGL}(1, p)$, where $n = p$ is a prime.
- (b) $X = A_n$ where n is odd, or $X = S_n$, where $n \geq 4$;
- (c) $\text{PGL}(d, q) \leq X \leq \text{P}\Gamma\text{L}(d, q)$, H is a Singer subgroup of X , and $n = (q^d - 1)/(q - 1)$.
- (d) $X = \text{P}\Gamma\text{L}(2, 8)$, $H = \langle s\sigma \rangle \cong C_9$, where $\langle s \rangle$ is a Singer subgroup of X and $\sigma \in X \setminus \text{PSL}(2, 8)$ with $|\sigma| = 3$.
- (e) $(X, n) = (\text{PSL}(2, 11), 11)$, $(M_{11}, 11)$, or $(M_{23}, 23)$.

Proposition 5. [27, Theorem 1.5] *Let X be a quasiprimitive permutation group on a finite set Ω . If X contains a dihedral regular subgroup H , then X is 2-transitive, and exactly one of the following holds for some $\omega \in \Omega$:*

- (a) $(X, H) = (A_4, D_4), (S_4, D_4), (\text{AGL}(3, 2), D_8), (\text{AGL}(4, 2), D_{16}), (C_2^4 \rtimes A_6, D_{16}),$ or $(C_2^4 \rtimes A_7, D_{16})$.
- (b) $(X, H, X_\omega) = (M_{12}, D_{12}, M_{11}), (M_{22}.2, D_{22}, \text{PSL}(3, 4).2),$ or (M_{24}, D_{24}, M_{23}) .
- (c) $(X, H, X_\omega) = (S_{2m}, D_{2m}, S_{2m-1}),$ or $(A_{4m}, D_{4m}, A_{4m-1})$.
- (d) $X = \text{PSL}(2, p^e).O, H = D_{p^e+1},$ and $X_\omega \supseteq C_p^e \rtimes C_{\frac{p^e-1}{2}}.O,$ where $p^e \equiv 3 \pmod{4}$, and $O \leq \text{Out}(\text{PSL}(2, p^e)) \cong C_2 \times C_e$.
- (e) $X = \text{PGL}(2, p^e)C_f, H = D_{p^e+1},$ and $X_\omega = C_p^e \rtimes C_{p^e-1},$ where $p^e \equiv 1 \pmod{4}$, and $f \mid e$.

Corollary 6. *Let X be a quasiprimitive permutation group with a stabilizer H and a regular subgroup K . If X is solvable, then one of the the following holds true:*

- (a) *If K is cyclic, then either $(X, H, K) = (S_4, D_6, C_4),$ or $C_p \cong K \leq X \leq \text{AGL}(1, p)$ for some prime p .*
- (b) *If K is dihedral and H is either cyclic or dihedral, then $(X, H, K) = (A_4, C_3, D_4)$ or (S_4, D_6, D_4) .*

The following three results on exact products related to dihedral groups will be useful.

Proposition 7. [31, Theorem 1] *If a finite group X admits a factorization $X = HK$ of two subgroups H and K , each having a cyclic subgroup of index at most 2, then X is solvable.*

Proposition 8. [11] *Let X be an exact product of a dihedral subgroup $H = \langle x \rangle \rtimes \langle y \rangle \cong D_{2n}$ and a cyclic subgroup $C = \langle z \rangle \cong C_m$, where $m, n \geq 2$. Suppose that M is the subgroup of maximal order in X such that $\langle z \rangle \leq M \subseteq \langle x \rangle \langle z \rangle$. Then M, M_X and X/M_X fall into one of the following classes:*

TABLE 1. Structure of M

Case	M	M_X	X/M_X
1	$\langle x \rangle \langle z \rangle$	$\langle x \rangle \langle z \rangle$	C_2
2	$\langle x^2 \rangle \langle z \rangle$	$\langle x^2 \rangle \langle z^2 \rangle$	D_8
3	$\langle x^2 \rangle \langle z \rangle$	$\langle x^2 \rangle \langle z^3 \rangle$	A_4
4	$\langle x^4 \rangle \langle z \rangle$	$\langle x^4 \rangle \langle z^3 \rangle$	S_4
5	$\langle x^3 \rangle \langle z \rangle$	$\langle x^3 \rangle \langle z^4 \rangle$	S_4

Proposition 9. [11, 39] *Let X be an exact product of a dihedral subgroup $H = \langle x \rangle \rtimes \langle y \rangle \cong D_{2n}$ and a cyclic subgroup $\langle z \rangle \cong C_m$, where $m, n \geq 2$. If n is odd, $\langle z \rangle_X = 1$ and $\langle x \rangle \langle z \rangle$ is a subgroup of X , then $H \triangleleft X$.*

3. CHARACTERIZATIONS

In this section, we give some structural characterizations of the exact products of two non-abelian dihedral groups. To simplify the statements, in the remainder of this paper, we use notation X to denote an exact product of two dihedral subgroups $H = \langle x \rangle \rtimes \langle y \rangle \cong D_{2n}$ and $K = \langle z \rangle \rtimes \langle w \rangle \cong D_{2m}$, where $m, n \geq 3$.

Lemma 10. *If X has a maximal subgroup G such that $H \leq G$ and $G \cap K \leq \langle z \rangle$, then*

- (a) *either $G = G_X = H\langle z \rangle$, $X = (H\langle z \rangle) \rtimes \langle w \rangle$, $X/G_X \cong C_2$, or*
- (b) *$G = H\langle z^2 \rangle$, $G_X = \langle x^3, z^2 \rangle$, $X = (K\langle x \rangle) \rtimes \langle y \rangle$ and $X/G_X \cong S_4$. In particular, $3|n$ and $2|m$.*

Proof. By hypothesis, we may assume $G \cap K = \langle z_1 \rangle$ for some $z_1 \in \langle z \rangle$. Using Dedekind's modular law, we have $G = G \cap X = G \cap HK = H(G \cap K) = H\langle z_1 \rangle$. Since

$$\langle z_1 \rangle \leq \bigcap_{k,l \in \mathbb{Z}} G^{z^k w^l} = \bigcap_{i,j,k,l \in \mathbb{Z}} G^{x^i y^j z^k w^l} = G_X,$$

we have $G_X = G_X \cap G = G_X \cap H\langle z_1 \rangle = (G_X \cap H)\langle z_1 \rangle$ and $G = HG_X$.

Consider now the quotient group $\overline{X} := X/G_X$. We have $\overline{G} = G/G_X = HG_X/G_X = \overline{H}$ and $\overline{X} = \overline{HK}$, where $\overline{K} = KG_X/G_X \cong K/\langle z_1 \rangle$. Since G is a maximal subgroup of X and $\overline{G_X} = \overline{G_X} = 1$, the (faithful) transitive permutation representation of \overline{X} on the cosets $[\overline{X} : \overline{G}]$ gives rise to a permutation group with a stabilizer $\overline{G} = \overline{H}$ and a regular subgroup \overline{K} . Being homomorphic images of the dihedral subgroups H and K , both \overline{H} and \overline{K} are either dihedral or cyclic. Note that in the latter case, we have $|\overline{H}| = 1$ or 2 , and $|\overline{K}| = 2$. By checking Corollary 6, only two possible cases may occur: either $C_p \cong \overline{K} \leq \overline{X} \leq \text{AGL}(1, p)$ with $p = 2$, or $(\overline{X}, \overline{H}, \overline{K}) = (S_4, D_6, D_4)$. In the former case, we have $z_1 = z$, and hence (a) holds true. In the latter case, we have $z_1 = z^2$, which gives (b), as required. \square

By reversing the roles of H and K in the above lemma, we obtain the following result.

Lemma 11. *If X has a maximal subgroup G such that $K \leq G$ and $G \cap H \leq \langle x \rangle$, then*

- (a) *either $G = G_X = K\langle x \rangle$, $X = (K\langle x \rangle) \rtimes \langle y \rangle$, $X/G_X \cong C_2$, or*
- (b) *$G = K\langle x^2 \rangle$, $G_X = \langle z^3, x^2 \rangle$, $X = (H\langle z \rangle) \rtimes \langle w \rangle$ and $X/G_X \cong S_4$. In particular, $3|m$ and $2|n$.*

Lemma 12. *If both m and n are odd, then $X = (H\langle z \rangle) \rtimes \langle w \rangle = (\langle x \rangle K) \rtimes \langle y \rangle$.*

Proof. The hypothesis implies that the roles of H and K in X are symmetric, so it suffices to show $X = (H\langle z \rangle) \rtimes \langle w \rangle$.

Let G be a maximal subgroup G containing H . Then $G = G \cap X = G \cap HK = H(G \cap K)$. If $G \cap K \leq \langle z \rangle$, then the result follows from Lemma 10 (note that the case (b) cannot appear since both m and n are odd).

Suppose now $G \cap K \not\leq \langle z \rangle$. Then $G \cap K = \langle z_1, w_1 \rangle$ for some $z_1 \in \langle z \rangle$ and $w_1 \in K \setminus \langle z \rangle$. We may assume $w_1 = w$, and so $G = H\langle z_1, w \rangle$. As before, we have

$$\langle z_1 \rangle \leq \bigcap_{k,l \in \mathbb{Z}} G^{z^k w^l} = \bigcap_{i,j,k,l \in \mathbb{Z}} G^{x^i y^j z^k w^l} = G_X,$$

and the quotient group $\overline{X} := X/G_X$ is a primitive permutation group on $[\overline{X}, \overline{G}]$ with a stabilizer \overline{G} . Since $\langle \overline{z} \rangle$ is cyclic and transitive, it is a regular subgroup of \overline{X} . Now

Corollary 6(a) is applicable, we have

$$C_p \cong \langle \bar{z} \rangle \leq \bar{X} \leq \text{AGL}(1, p) \quad \text{for some odd prime } p.$$

Thus, $z_1 = z^p$ and the stabilizer \bar{G} is cyclic. Since \bar{H} is a cyclic homomorphic image of the dihedral group H and $\bar{H} \leq \bar{G}$, we have $\bar{H} = \bar{1}$ or $\bar{H} = \langle \bar{y} \rangle \cong C_2$. In either case, $x \in G_X$.

If $\bar{H} = 1$, then $G_X = H\langle z^p \rangle$, and so $\bar{X} = \langle \bar{z}^p \rangle \rtimes \langle \bar{w} \rangle$. Therefore, $X = (H\langle z \rangle) \rtimes \langle w \rangle$. If $\bar{H} = \langle \bar{y} \rangle$, then $G_X = \langle x \rangle \langle z^p \rangle$ and $\bar{X} = \langle \bar{z}, \bar{w} \rangle \rtimes \langle \bar{y} \rangle$. Note that $\langle \bar{z} \rangle \text{ char } \langle \bar{z}, \bar{w} \rangle \triangleleft \bar{X}$, we have $\langle \bar{z} \rangle \triangleleft \bar{X}$. Therefore, $X = (H\langle z \rangle) \rtimes \langle w \rangle$, as desired. \square

Corollary 13. *If both m and n are odd, then $X = (\langle z \rangle_X \rtimes H).K = (\langle x \rangle_X \rtimes K).H$*

Proof. By Lemma 12, $X = (H\langle z \rangle) \rtimes \langle w \rangle$ and the maximal subgroup $G := H\langle z \rangle$ of X is an exact product of the dihedral subgroup $H = \langle x \rangle \rtimes \langle y \rangle \cong D_{2n}$ and the cyclic subgroup $\langle z \rangle \cong C_m$. Since both m and n are odd, by Lemma 8 we get $H\langle z \rangle = (\langle x \rangle \langle z \rangle) \rtimes \langle y \rangle$, and so $X = (\langle x \rangle \langle z \rangle) \rtimes \langle y \rangle \rtimes \langle w \rangle$. It is clear that $\langle z \rangle_X = \langle z \rangle_G$. Thus, the quotient group $\bar{G} := G/\langle z \rangle_X$ contains a dihedral subgroup $\bar{H} = \langle \bar{x}, \bar{y} \rangle$ with a core-free cyclic complement $\langle \bar{z} \rangle$. Since $\langle \bar{x} \rangle \langle \bar{z} \rangle \leq \bar{G}$, by Proposition 9, we get $\bar{H} \triangleleft \bar{G}$, and so $\bar{G} = \bar{H} \rtimes \langle \bar{z} \rangle$. Since \bar{z} has odd order, $\bar{g} \in H$ for all $\bar{g} \in \bar{G}$ with $\bar{g}^2 = 1$, so $\bar{H} = \langle \bar{g} \in \bar{G} \mid \bar{g}^2 = 1 \rangle$. Thus $\bar{H} \text{ char } \bar{G} \triangleleft \bar{X}$, and so $\bar{H} \triangleleft \bar{X}$. Therefore, $X = (\langle z \rangle_X \rtimes H).K$, as claimed. \square

4. CLASSIFICATION

In this section, we utilize the results from the previous section to provide a proof of Theorem 1.

Proof of Theorem 1. By Corollary 13, we have $X = (\langle z \rangle_X \rtimes H).K = (\langle x \rangle_X \rtimes K).H$. Therefore, X contains the following two chains of subnormal subgroups with cyclic factors:

$$1 \leq \langle z \rangle_X < \langle z \rangle_X \rtimes \langle x \rangle \leq \langle z \rangle_X \rtimes H \leq (\langle z \rangle_X \rtimes H).\langle z \rangle < ((\langle z \rangle_X \rtimes H).\langle z \rangle) \rtimes \langle w \rangle = X. \quad (2)$$

and

$$1 \leq \langle x \rangle_X < \langle x \rangle_X \rtimes \langle z \rangle \leq \langle x \rangle_X \rtimes K \leq (\langle x \rangle_X \rtimes K).\langle x \rangle < ((\langle x \rangle_X \rtimes K).\langle z \rangle) \rtimes \langle y \rangle = X. \quad (3)$$

Let $z_1 := z^{m_1}$ be a generator of $\langle z \rangle_X$, where m_1 is a positive divisor of m . By N/C theorem, $X/C_X(z_1)$ is abelian, so its subgroup $HC_X(z_1)/C_X(z_1)$ is also abelian. Note that $HC_X(z_1)/C_X(z_1)$ is a homomorphic image of the dihedral group $H = \langle x \rangle \rtimes \langle y \rangle \cong D_{2n}$ with n odd, we must have $x \in C_X(z_1)$. Thus,

$$z_1^x = z_1, \quad z_1^y = z_1^u \quad (4)$$

for some $u \in \mathbb{Z}_{m/m_1}$.

Consider $\bar{X} := X/\langle z_1 \rangle = \bar{H} \rtimes \bar{K}$. Since $\langle \bar{x} \rangle \text{ char } \bar{H} \triangleleft \bar{X}$, we have $\langle \bar{x} \rangle \triangleleft \bar{X}$. Thus, by the N/C theorem again, $\bar{z} \leq C_{\bar{X}}(\bar{x})$. It follows that the following relations hold in \bar{X} :

$$\bar{x}^{\bar{z}} = \bar{x}, \quad \bar{y}^{\bar{z}} = \bar{x}^R \bar{y}, \quad \bar{x}^{\bar{w}} = \bar{x}^S, \quad \bar{y}^{\bar{w}} = \bar{x}^T \bar{y},$$

where $R, S, T \in \mathbb{Z}_n$. Consequently, we obtain the following relations in X :

$$x^z = xz_1^e, \quad y^z = z_1^a x^R y, \quad x^w = x^S z_1^b, \quad y^w = yx^T z_1^c \quad (5)$$

or written in an equivalent form:

$$[x, z] = z_1^e, \quad [z, y] = x^R z_1^a, \quad [x, w] = x^{S-1} z_1^b, \quad [y, w] = x^T z_1^c, \quad (6)$$

where $e, a, b, c \in \mathbb{Z}_{m/m_1}$.

On the other hand, due to the formal symmetry between H and K in X , we can also consider the core $\langle x \rangle_X$ of $\langle x \rangle$ in X and the corresponding quotient group $\overline{X} := X/\langle x \rangle_X$. Let $x_1 := x^{n_1}$ be a generator of $\langle x \rangle_X$, where n_1 is a positive divisor of n . Using a similar argument as before, it is straightforward to show that $z \in C_X(x_1)$ and $\langle \bar{z} \rangle \triangleleft \overline{X}$. Now we can deduce from (3) and (6) that $R, T \equiv 0 \pmod{n_1}$ and $S \equiv 1 \pmod{n_1}$. Therefore, we may set

$$R = n_1 r, \quad S = 1 + n_1 s \quad \text{and} \quad T = n_1 t, \quad \text{where } r, s, t \in \mathbb{Z}_{n/n_1}.$$

Then the relations in (6) can be expressed in the following symmetric form:

$$[x, z] = z_1^e, \quad [z, y] = x_1^r z_1^a, \quad [x, w] = x_1^s z_1^b, \quad [y, w] = x_1^t z_1^c,$$

or written in an equivalent form as follows:

$$x^z = x z_1^e, \quad y^z = x_1^r z_1^a y, \quad x^w = x x_1^s z_1^b, \quad y^w = y x_1^t z_1^c.$$

Therefore, $X = \langle x, y, z, w \rangle$ has the following defining relations

$$\begin{aligned} x^n = y^2 = z^m = w^2 = 1, \quad x_1 = x^{n_1}, \quad z_1 = z^{m_1}, \\ z_1^x = z_1, \quad z_1^y = z_1^u, \quad z_1^z = z_1, \quad z_1^w = z_1^{-1}, \\ x^y = x^{-1}, \quad x^z = x z_1^e, \quad x^w = x x_1^s z_1^b, \\ y^z = x_1^r z_1^a y, \quad y^w = y x_1^t z_1^c, \quad z^w = z^{-1}. \end{aligned} \tag{7}$$

In the subsequent analysis, we utilize Proposition 3 to exam the relations in (7) and establish the numerical conditions on the parameters.

Since $y^2 = 1$ and $z_1^y = z_1^u$, we have $z_1 = z_1^{y^2} = (z_1^y)^y = (z_1^u)^u = z_1^{u^2}$, and so

$$u^2 \equiv 1 \pmod{m/m_1}. \tag{8}$$

Moreover, from $x^y = x^{-1}$, $z_1 = z_1^{m_1}$, $x^z = x z_1^e$ and $y^z = x_1^r z_1^a y$ we derive that

$$\begin{aligned} 1 = x^n = (x^n)^z = (x^z)^n = x^n z_1^{ne}; \\ 1 = y^2 = (y^2)^z = (y^z)^2 = (x^r z_1^a y)^2 = z_1^{a(u+1)}; \\ x = x^{z_1} = x^{z^{m_1}} = (x z_1^e)^{z^{m_1-1}} = x^{z^{m_1-1}} z_1^e = x z_1^{m_1 e}; \\ z_1^{u-1} y = y^{z_1} = y^{z^{m_1}} = (x_1^r z_1^a y)^{z^{m_1-1}} = x_1^r z_1^a y^{z^{m_1-1}} = x_1^{m_1 r} z_1^{m_1 a} y; \\ x^{-1} z_1^{-e} = (x^{-1})^z = (x^y)^z = (x^z)^{y^z} = (x z_1^e)^{x_1^r z_1^a y} = (x z_1^e)^y = x^{-1} z_1^{ue}, \end{aligned}$$

which imply the following congruences

$$ne \equiv 0 \pmod{m/m_1}, \tag{9}$$

$$a(u+1) \equiv 0 \pmod{m/m_1}, \tag{10}$$

$$m_1 e \equiv 0 \pmod{m/m_1}, \tag{11}$$

$$m_1 r \equiv 0 \pmod{n/n_1}, \tag{12}$$

$$u \equiv 1 + m_1 a \pmod{m/m_1}, \tag{13}$$

$$(u+1)e \equiv 0 \pmod{m/m_1}. \tag{14}$$

Combining equations (11), (13) with (14), we obtain

$$2e \equiv (u+1)e + (u-1)e \equiv (1-u)e \stackrel{(13)}{\equiv} (u+1)e + m_1 a e \stackrel{(11)}{\equiv} 0 \pmod{m/m_1}.$$

Since m is odd, we deduce that $e \equiv 0 \pmod{m/m_1}$, which implies $[x, z] = 1$. Consequently, the congruences (8)–(14) are reduced to

$$a(u+1) \equiv 0, \quad u \equiv 1 + m_1 a \pmod{m/m_1} \quad \text{and} \quad m_1 r \equiv 0 \pmod{n/n_1}. \quad (15)$$

Furthermore, from $x^w = xx_1^s z_1^b$ and $y^w = yx^t z_1^c$ we infer that

$$\begin{aligned} 1 &= x^n = (x^n)^w = (x^w)^n = (xx_1^s z_1^b)^n = z_1^{nb}; \\ 1 &= y^2 = (y^2)^w = (y^w)^2 = (yx^t z_1^c)^2 = z_1^{c(u+1)}; \\ x &= x^{w^2} = (x^{1+n_1 s} z_1^b)^w = (x^w)^{1+n_1 s} z_1^{-b} \\ &= (xx_1^s z_1^b)^{1+n_1 s} z_1^{-b} = x^{1+n_1 s(2+n_1 s)} z_1^{n_1 s b}; \\ x^{-1} x_1^{-s} z_1^{-b} &= (x^{-1})^w = (x^y)^w = (x^w)^{y^w} = (xx_1^s z_1^b)^{yx_1^t z_1^c} \\ &= (x^{-1} x_1^{-s} z_1^{bu}) x_1^t z_1^c = x^{-1} x_1^{-s} z_1^{bu}; \\ yx_1^t z_1^c &= y^w = (y^w)^{z^w} = (y^z)^{wz} = (x_1^r z_1^a y)^{wz} \\ &= ((x_1^{1+sn_1} z_1^{bn_1})^r z_1^{-a} yx_1^t z_1^c)^z = x_1^{r(1+sn_1)} z_1^{bn_1 r - a} y^z x_1^t z_1^c \\ &= x_1^{r(1+sn_1)} z_1^{bn_1 r - a} (x_1^r z_1^a y) x_1^t z_1^c = yx_1^{t-r(2+sn_1)} z_1^{ubn_1 r + c}. \end{aligned}$$

Thus, we obtain

$$\begin{aligned} nb &\equiv 0 \pmod{m/m_1}, \\ c(u+1) &\equiv 0 \pmod{m/m_1}, \\ s(2+n_1 s) &\equiv 0 \pmod{n/n_1}, \\ n_1 s b &\equiv 0 \pmod{m/m_1}, \\ b(u+1) &\equiv 0 \pmod{m/m_1}, \\ r(2+sn_1) &\equiv 0 \pmod{n/n_1}, \\ n_1 r b &\equiv 0 \pmod{m/m_1}. \end{aligned} \quad (16)$$

In addition, for any integer k , we have

$$(x^k)^w = (xx_1^s z_1^b)^k = x^k x_1^{sk} z_1^{bk} \quad \text{and} \quad (z^k)^y = (x_1^r z_1^a)^k = x_1^{rk} z_1^{ak}.$$

Observe that $[x, z] = 1$, $x^y = x^{-1}$ and $z^w = z^{-1}$. Using the assumption $\langle x \rangle_X = \langle x^{n_1} \rangle$ and $\langle z \rangle_X = \langle z^{m_1} \rangle$ we infer from the above equations that

$$\begin{aligned} rk &\equiv 0 \pmod{n/n_1} \iff k \equiv 0 \pmod{m_1}, \\ bk &\equiv 0 \pmod{m/m_1} \iff k \equiv 0 \pmod{n_1}. \end{aligned} \quad (17)$$

Finally, we see from (15) that $u = 1 + m_1 a \pmod{m/m_1}$. Therefore, upon substitution, the congruences in (15), (16) and (17) are reduced to the stated form given in the theorem. By Proposition 3, the group X defined by the stated presentation and numerical conditions is well-defined, it contains two dihedral subgroups $H = \langle x, y \rangle \cong D_{2n}$ and $K = \langle z, w \rangle \cong D_{2m}$ with $H \cap K = \{1_X\}$. In particular, it is straightforward to verify that $\langle x \rangle_X = \langle x^{n_1} \rangle$ and $\langle z \rangle_X = \langle z^{m_1} \rangle$, as required. \square

REFERENCES

- [1] A.L. Agore, C.G. Bontea and G. Militaru, Classifying bicrossed products of Hopf Algebras, *Algebr Represent Theory* 17 (2014) 227–264.
- [2] A.L. Agore, A. Chirvăsitu, B. Ion and G. Militaru, Bicrossed products for finite groups, *Algebr. and Represent. Theory*, 12(2-5) (2009) 481–488.

- [3] M. Bachratý, M. Conder, and G. Verret. Skew product groups for monolithic groups, *Algebraic Combinatorics* **5**(5) (2022), 785–802.
- [4] T.C. Burness, and C.H. Li, On solvable factors of almost simple groups, *Advances in Mathematics* **377** (2021) 107499.
- [5] J.Y. Chen, S.F. Du and C.H. Li. Skew-morphisms of nonabelian characteristically simple groups. *J. Combin. Theory Ser. A* **185** (2022), 105539.
- [6] M. Conder, R. Jajcay and T. Tucker, Cyclic complements and skew morphisms of groups, *J. Algebra* **453** (2016), 68–100.
- [7] M. Conder and T. Tucker, Regular Cayley maps for cyclic groups, *Trans. Amer. Math. Soc.* **366** (2014), 3585–3609.
- [8] J. Douglas, On the supersolvability of bicyclic groups, *Proceedings of the National Academy of Sciences* **47**(9) (1961), 1493–1495.
- [9] S.F. Du, W.J. Luo, H. Yu and J.Y. Zhang, Skew-morphisms of elementary abelian p -groups, *J. Group Theory*, 2024, <https://doi.org/10.1515/jgth-2022-0092>
- [10] S.F. Du, W.J. Luo and H. Yu, Regular Cayley maps of elementary abelian p -groups: Classification and enumeration, *J. Combin. Theory, Ser. A*, 2023
- [11] S.F. Du, H. Yu and W.J. Luo, The Product of a generalized quaternion group and a cyclic group, arXiv preprint, 2023.
- [12] C. Hering, M.W. Liebeck and J. Saxl, The factorizations of the finite exceptional groups of Lie type, *J. Algebra* **106** (2) (1987) 517–527.
- [13] K. Hu and R. Jajcay, Cyclic complementary extensions and skew morphisms, preprint, 2023.
- [14] K. Hu, I. Kovács and Y.S. Kwon, A classification of skew morphisms of dihedral groups, *J. Group Theory* **26** (3)(2023) 547–569.
- [15] K. Hu, I. Kovács and Y.S. Kwon, Classification of cyclic groups underlying only smooth skew morphisms, *J. Algebraic Combin.* 2024, <https://doi.org/10.1007/s10801-024-01311-4>.
- [16] K. Hu, I. Kovács and Y.S. Kwon, On exact products of a dihedral group and a cyclic group, preprint, 2024.
- [17] B. Huppert, Endliche Gruppen. I, Grundlehren Math. Wiss. 134, Springer, Berlin, 1967.
- [18] I.M. Isaacs, Finite Group Theory, Grad. Stud. Math. 92, American Mathematical Society, Providence, 2008.
- [19] N. Itô, Über das Produkt von zwei abelschen Gruppen, *Math. Z.* **62** (1955) 400–401.
- [20] R. Jajcay and J. Širáň, Skew morphisms of regular Cayley maps, *Discrete Math.* **224** (2002), 167–179.
- [21] Z. Janko, Finite 2-groups with exactly one nonmetacyclic maximal subgroup, *Israel J. Math.*, **166** (2008) 313–347.
- [22] G.A. Jones, Cyclic regular subgroups of primitive permutation groups. *J. Group Theory* **5** (4) (2002) 403–407.
- [23] G.A. Jones, Regular embeddings of complete bipartite graphs: classification and enumeration, *Proc. Lond. Math. Soc.*, **101** (3) (2010) 427–453.
- [24] O.H. Kegel, Produkte nilpotenter Gruppen, *Arch. Math. (Basel)* **12** (1961), 90–93.
- [25] J. H. Kwak and Y. S. Kwon. Unoriented Cayley maps. *Studia Scientiarum Mathematicarum Hungarica* **43**(2) (2006), 137–157.
- [26] I. Kovács and Y. S. Kwon. Regular Cayley maps for dihedral groups. *J. Combin. Theory Ser. B* **148** (2021), 84–124.
- [27] C.H. Li, Finite edge-transitive Cayley graphs and rotary Cayley maps, *Transactions of the American Mathematical Society* **358**, no. 10 (2006) 4605–4635.
- [28] C.H. Li, L. Wang and B. Xia, The exact factorizations of almost simple groups, *J. London Math. Soc.*, 2023, <https://doi.org/10.1112/jlms.12784>.
- [29] M.W. Liebeck, C.E. Praeger, and J. Saxl, On factorizations of almost simple groups, *J. Algebra* **185** (2) (1996) 409–419.
- [30] M.W. Liebeck, C.E. Praeger, and J. Saxl, Regular subgroups of primitive permutation groups, *Mem. Amer. Math. Soc.* **203** (952) (2010) vi+74.
- [31] V. S. Monakhov, The product of two groups, one of which contains a cyclic subgroup of index ≤ 2 , *Mathematical Notes of the Academy of Sciences of the USSR*, **16**(2) (1974) 757–762.
- [32] O Ore, Structures and group theory. I, *Duke Math. J.*, **3** (2)(1937) 149–174.

- [33] M. Takeuchi, Matched pairs of groups and bismash products of Hopf algebras, *Comm. Algebra*, 9(8) (1981) 841–882.
- [34] H. Wielandt, Über das Produkt von paarweise vertauschbaren nilpotenten Gruppen, *Math. Z.* **55** (1951), 1–7.
- [35] B. Xia, Quasiprimitive groups containing a transitive alternating group, *Journal of Algebra* 490 (2017) 555–567.
- [36] M.F. Newman and M.Y. Xu, Metacyclic groups of prime-power order (Research announcement), *Adv. in Math. (China)* 17 (1988) 106–107.
- [37] M.Y. Xu and Q.H. Zhang, A classification of metacyclic 2-groups, *Alg. Colloq.* 13 (2006) 25–34.
- [38] H. Yu, Unoriented regular Cayley maps of elementary abelian groups p -groups, preprint, 2024
- [39] J.Y. Zhang and S.-F. Du, On the skew morphisms of dihedral groups, *J. Group Theory* **19** (2016), 993–1016.

K. HU,
DEPARTMENT OF MATHEMATICS, ZHEJIANG OCEAN UNIVERSITY, ZHOUSHAN, ZHEJIANG 316022,
P.R. CHINA
Email address: hukan@zjou.edu.cn

H. YU
SCHOOL OF MATHEMATICAL SCIENCES, CAPITAL NORMAL UNIVERSITY, BEIJING 100048, P.R.
CHINA
Email address: 3485676673@qq.com