# Many-body Fu-Kane-Mele index

Sven Bachmann  $^{*1}$ , Alex Bols  $^{\dagger 2}$ , and Mahsa Rahnama  $^{\ddagger 3}$ 

<sup>1</sup>Institute for Theoretical Physics, ETH Zürich <sup>2, 3</sup>Department of Mathematics, University of British Columbia, Vancouver

### July 1, 2024

#### Abstract

We define a  $\mathbb{Z}_2$ -valued index for stably short-range entangled states of two-dimensional fermionic lattice systems with charge conservation and time reversal symmetry. The index takes its non-trivial value precisely if the 'fluxon', the state obtained by inserting a  $\pi$ -flux through the system, transforms under time reversal as part of a Kramers pair. This index extends the Fu-Kane-Mele index of free fermions to interacting systems.

### Contents

1	Introduction			
	1.1	Idea behind the construction	2	
	1.2	Organisation of the paper	í	
2	Set	up and results	5	
	2.1	Observable algebras	5	
	2.2	Locally generated automorphisms	5	
	2.3	Symmetries	;	
	2.4	Symmetric short-range entangled states	7	
	2.5	Examples: Non-interacting electrons with spin	7	
	2.6	Symmetry protected phases	)	
	2.7	The many-body $\mathbb{Z}_2$ -index	)	
3	Def	inition of the index	)	
	3.1	Flux insertion	)	
	3.2	Defect states at $\pm \pi$ flux		
	3.3	Definition of the index		
4	Pro	operties of the index 18	3	
	4.1	Independence of choices	3	
	4.2	Stacking		
	4.3	Invariance under symmetry preserving LGAs		
	4.4	Extension of the index to symmetric stably SRE states		

\*email: sbach@math.ubc.ca  $^{\dagger}$ email: abols01@phys.ethz.ch  $^{\ddagger}$ email: mahsa.r@math.ubc.ca

5	Free fermion examples	24
	5.1 Proof that spin Hall states are stably SRE	24
	5.2 Index of spin Hall states	26
6	Proof of the main theorem	
$\mathbf{A}$	Almost local transitivity	31
В	Flux insertion for free fermions	36

### 1 Introduction

The discovery of the integer quantum Hall effect [39] and its understanding in terms of the topology of band insulators [51, 3] set the stage for the study of topological phases of matter. The discovery of the quantum spin Hall effect for time reversal symmetric insulators [32, 31] quickly led to the recognition that the landscape of topological phases of free fermion insulators and superconductors is extremely rich, eventually leading to a complete classification in terms of K-theory, summarised in the periodic table of topological invariants [38, 49].

A natural question is whether the phases identified and classified by the periodic table of topological invariants survive if interactions are allowed. The answer is no in general [21, 22], but yes at least for those phases for which the topological invariant has a clear physical interpretation. This is the case for the integer quantum Hall effect where the invariant is a linear response coefficient. The fact that the Hall conductance is a well-defined quantised invariant of topological phases of interacting matter was explained early [40] and it is by now very well understood, in particular in the framework of quantum lattice systems [29, 5, 6, 43, 7, 33]. Of course, the stability of the non-interacting phases comes in general with the appearance of new, purely interacting ones such as the fractional Hall phases that can also be classified [24].

The invariant that characterises the quantum spin Hall effect is the  $\mathbb{Z}_2$ -valued Fu-Kane-Mele index [32, 31, 25, 26]. It was argued in [48] [47] that this invariant has a physical interpretation in terms of time reversal transformation properties of defects bound to  $\pm \pi$  magnetic fluxes. If the invariant is non-trivial then these defects form a Kramers pair for the time reversal symmetry. We review the argument below. While these early works placed some importance on the role of spin, they all recognized that the invariant is in fact only relying on fermionic time reversal symmetry: no component of spin must be conserved for the invariant to be meaningful.

Following these ideas we construct a  $\mathbb{Z}_2$ -valued index for short-range entangled states of two-dimensional interacting lattice fermion systems that are invariant under a fermionic time reversal symmetry that commutes with a U(1)-symmetry. We show that this index agrees with the Fu-Kane-Mele index of non-interacting systems, namely for symmetric quasi-free states, and is stable under locally generated automorphisms that respect the symmetries. In particular, we show that the quantum spin Hall phase indeed persists in the interacting regime.

### 1.1 Idea behind the construction

Let us now summarise the Kramers pair binding idea of [48, 47] and describe how our many-body index is constructed based on this idea.

The systems we consider are interacting generalisations of gapped free fermion Hamiltonians on the two-dimensional square lattice with a time reversal symmetry  $\mathcal{T}$  that squares to -1 (Altland-Zirnbauer class AII [1]). Such Hamiltonians come in two topological phases distinguished from each other by the  $\mathbb{Z}_2$ -valued Fu-Kane-Mele index. A nice characterisation of the index is obtained by piercing magnetic flux through the system [17] as follows. Let  $\mathscr{H}$  be a gapped free fermion Hamiltonian such that  $\mathscr{TH}\mathscr{T}^* = \mathscr{H}$ , and let  $\mathscr{H}(\phi)$  be the free fermion Hamiltonian obtained from  $\mathscr{H}$  by inserting flux  $\phi$ . Then the spectrum of  $\mathscr{H}(\phi)$  inherits the

bulk spectrum of  $\mathscr{H}$ , but may in addition have eigenvalues in the bulk gap. As a function of  $\phi$ , these eigenvalues form bands within the gap of  $\mathscr{H}$  corresponding to modes bound to the magnetic flux, see Figure 1. The time reversal symmetry forces the spectra of  $\mathscr{H}(\phi)$  and  $\mathscr{H}(-\phi)$  to be equal and for the time reversal invariant fluxes  $\phi_* = 0, \pm \pi$  the time reversal symmetry forces all eigenvalues of  $\mathscr{H}(\phi_*)$  to be doubly degenerate due to Kramers pairing. Energy bands satisfying these constraints come in two classes distinguished from each other as follows. Let  $E_F$  be a generic Fermi energy in the bulk gap of the Hamiltonian  $\mathscr{H}$  (for simplicity we assume that  $E_F$  is not an eigenvalue of  $\mathscr{H}(\phi_*)$ ) and let  $N \in \mathbb{Z}_2$  be the parity of the number of times a band crosses the Fermi energy in the interval  $\phi \in [0,\pi)$ . Then N is invariant under any homotopy of the bands that respects the time reversal symmetry constraints. The Fu-Kane-Mele index of  $\mathscr{H}$  (w.r.t. the chosen gap) is equal to N. The spectrum of Figure 1 corresponds to an  $\mathscr{H}$  with a non-trivial Fu-Kane-Mele index. This characterisation of the Fu-Kane-Mele index parallels the characterisation for translation invariant free fermion systems through the bulk-boundary correspondence [27, 23, 12] where the bands associated to modes bound to the magnetic flux are replaced by bands formed by edge modes.





Figure 1: The spectrum of a gapped free fermion Hamiltonian  $\mathcal{H}(\phi)$  as a function of pierced flux. The filled grey regions are bulk spectrum. The bands represent spectrum associated to modes bound to the magnetic flux. The bands cross at time reversal invariant values of the flux  $\phi = 0, \pm \pi$  forming Kramers pairs represented by dots. The spectrum depicted here corresponds to a non-trivial insulator because for any Fermi energy  $E_F$  in the bulk gap, the eigenvalue bands cross  $E_F$  an odd number of times (in fact,  $\pm 1$ ) in the interval  $\phi \in [0, \pi)$ .

We now imagine preparing the system at  $\phi = 0$  in its ground state where all single particle states with energy below  $E_F$  in the bulk gap are occupied, and increasing the flux  $\phi$  adiabatically from 0 to  $\pi$ . The state is adiabatically transported along this path and at each value of  $\phi \in [0, \pi]$  the occupied single particle states are indicated in the left panel of Figure 2. We see that one mode of the Kramers pair at  $\phi = \pi$  is occupied while the other mode of the pair remains unoccupied. I instead the flux is decreased from  $\phi = 0$  to  $\phi = -\pi$  we get a similar picture, except that now the other mode of the Kramers pair becomes occupied, see the right panel of Figure 2. The two states obtained from the ground state by either increasing to flux from 0 to  $\pi$  or decreasing the flux from 0 to  $-\pi$  are mapped into each other by time reversal.

Since these two states differ from each other in the occupation of a single Kramers pair it seems reasonable to suspect that the corresponding many-body states form a Kramers pair. The results of this paper provide a rigorous justification of this intuition by showing that the two states obtained by  $\pm \pi$  flux insertion are unitary equivalent, and are therefore given by unit vectors in the same irreducible representation of the observable algebra. We prove that the time reversal symmetry is implemented in this representation by an antiunitary operator T, and that the vector representatives of the two states are indeed a Kramers pair for T.

To define a many-body  $\mathbb{Z}_2$  index for a time reversal and U(1) symmetric state  $\omega$  we therefore want to construct states  $\omega^{\pm}$  corresponding to adiabatically inserting  $\pi$  or  $-\pi$  magnetic flux in the system. Under the assumption that  $\omega$  is a symmetric short-range entangled state one can construct a Hamiltonian  $H = \sum_x H_x$  with symmetric terms  $H_x$  localized near the site x and






Figure 2: The occupied single particle spectrum (dark) as the ground state at  $\phi=0$  is adiabatically transported from  $\phi=0$  to  $\phi=\pi$  (left panel) or to  $\phi=-\pi$  (right panel). The states at  $\pm\pi$  differ only in the occupation of the Kramers pair of modes bound to the  $\pm\pi$  flux. The mode of this Kramers pair that is occupied in the state obtained by adiabatic transport from 0 to  $\pi$  flux is unoccupied in the state obtained by adiabatic transport from 0 to  $-\pi$  flux and vice versa.

such that  $\omega$  is the unique gapped ground state of H. Let  $H(\phi) = \sum_x H_x(\phi)$  with  $H_x(\phi)$  obtained from  $H_x$  by conjugation with the U(1) symmetry restricted to the upper half plane. Since the  $H_x$  are symmetric the  $H_x(\phi)$  substantially differ from  $H_x$  only for those x near the horizontal axis. The Hamiltonians  $H(\phi)$  have unique gapped ground states  $\omega_{\phi}$  related to  $\omega$  by conjugation by the U(1) symmetry restricted to the upper half plane. The states  $\omega_{\phi}$  differ from  $\omega$  only near the horizontal axis. Moreover,  $\omega_{\pi} = \omega_{-\pi}$  since the symmetry is U(1). We then consider the quasi-adiabatic flow [30] for the family  $H(\phi)$  (Section 3.1.3). Since  $H(\phi)$  only changes as a function of  $\phi$  near the horizontal axis, the quasi-adiabatic flow acts non-trivially only near this axis. An important technical merit of the quasi-adiabatic flow is that it is generated by a local interaction localised near the horizontal axis and so it can be restricted to act non-trivially only near the left horizontal axis (Section 3.1.4). We define the states  $\omega^{\pm}$  by acting with this restricted quasi-adiabatic flow for flux  $\pm \pi$  on the ground state. Because we work in the half-line gauge, both states  $\omega^+$  and  $\omega^-$  look like  $\omega_{\pi} = \omega_{-\pi}$  far to the left of the origin near the left horizontal axis, and look like  $\omega$  far from the origin and away from the left axis, see Figure 3. Under the assumption that  $\omega$  is short-range entangled it follows that the states  $\omega^{\pm}$  are




Figure 3: The geometry of flux insertion. In the half-line gauge, the electric field E generated by the time-dependent flux  $\phi(t)$  exists only along the left half-line, affecting the state only in the vicinity of that line. The states  $\omega^{\pm}$  are the result of the process  $\phi = 0 \rightarrow \phi = \pm \pi$ .

unitarily equivalent (Proposition 3.8). The states  $\omega^{\pm}$  therefore have unit vector representatives  $\Omega^{\pm} \in \mathcal{H}$  in the GNS representation  $(\pi, \mathcal{H})$  of either of them. Since they are time reversal images of each other, time reversal is implemented in the representation by an anti-unitary operator T (Lemma 3.10), and  $T^2$  implements fermion parity. It turns out that either  $T^2\Omega^+ = \Omega^+$  or  $T^2\Omega^+ = -\Omega^+$  (3.14). The first possibility correspond to a trivial index while the second possibility corresponds to a non-trivial  $\mathbb{Z}_2$  index. In the non-trivial case we have that  $\Omega^- = T\Omega^+$  and  $\Omega^+$  form an orthogonal Kramers pair for T.

### 1.2 Organisation of the paper

The paper is organised as follows. In Section 2 we give precise definitions of the fermionic lattice systems and the class of pure states on these systems that we will study. We then state our main theorem: That there is a  $\mathbb{Z}_2$ -valued index associated to these states, that this index is stable under locally generated automorphisms which respect the symmetries, and that this index extends the Fu-Kane-Mele index for non-interacting quantum spin Hall effect. The rest of the paper is devoted to proving the main theorem. In Section 3 we define the index of a symmetric short-range entangled (SRE) state by carrying out the construction sketched above. In Section 4 we first prove that the index is well-defined in the sense that it is independent of various choices made in the definition. We then turn to its properties, namely multiplicativity under stacking and stability under locally generated automorphisms that respect the symmetries. We also show that symmetric product states have trivial index which allows us to extend the index to stably SRE states. Section 5 is devoted to the non-interacting situation. We first show that all translation-invariant, time reversal invariant quasi-free states are stably SRE and therefore have a well defined index. We then prove that our index agrees with the Fu-Kane-Mele index for translation invariant non-interacting spin Hall states. The next Section 6 collects these results to prove the main theorem. The appendices are devoted to some technical results. In Appendix A we prove a transitivity result for fermionic SRE states that is used in Section 3 to show that the  $\pm \pi$  flux states are unitarily equivalent. In Appendix B we describe the free fermion analogue of the adiabatic flux insertion used in Section 5.

### 2 Setup and results

### 2.1 Observable algebras

Let  $\mathcal{A}$  be the CAR algebra over  $l^2(\mathbb{Z}^2; \mathbb{C}^n)$ , i.e.  $\mathcal{A}$  is the  $C^*$ -algebra generated by an identity element and annihilation operators  $\{a_{x,i}\}_{x\in\mathbb{Z}^2, i=1,\cdots,n}$  that satisfy the canonical anticommutation relations

$$\{a_{x,i}, a_{y,j}\} = \{a_{x,i}^*, a_{y,j}^*\} = 0, \quad \{a_{x,i}, a_{y,j}^*\} = \delta_{x,y}\delta_{i,j}$$
(2.1)

for all sites  $x,y\in\mathbb{Z}^2$  and all  $i,j=1,\cdots,n$ . For any  $\Gamma\subset\mathbb{Z}^2$  we denote by  $\mathcal{A}_{\Gamma}$  the unital  $C^*$ -subalgebra of  $\mathcal{A}$  generated by the  $\{a_{x,i}\}_{x\in\Gamma,i=1,\cdots,n}$ . We also write  $\mathcal{A}_x=\mathcal{A}_{\{x\}}$  for any site  $x\in\mathbb{Z}^2$ . The fermion parity is the \*-automorphism  $\theta$  of  $\mathcal{A}$  uniquely determined by  $\theta(a_{x,i})=-a_{x,i}$  for all  $x\in\mathbb{Z}^2$  and all  $i=1,\cdots,n$ . We have  $\theta(\mathcal{A}_{\Gamma})=\mathcal{A}_{\Gamma}$  for all  $\Gamma\subset\mathbb{Z}^2$ . An operator  $O\in\mathcal{A}$  is called even if  $\theta(O)=O$ , odd if  $\theta(O)=-O$  and homogeneous if it is either even or odd. For any  $\Gamma\subset\mathbb{Z}^2$  the subset of even operators of  $\mathcal{A}_{\Gamma}$  is a  $C^*$ -subalgebra of  $\mathcal{A}_{\Gamma}$  which we denote by  $\mathcal{A}_{\Gamma}^+$ . We will write  $\Gamma\subset\mathbb{Z}^2$  to mean that  $\Gamma$  is a finite subset of  $\mathbb{Z}^2$ . If  $A\in\mathcal{A}_{\Gamma}$  for some  $\Gamma\subset\mathbb{Z}^2$  then A is said to be a local operator. The union of all local operators is a \*-subalgebra of  $\mathcal{A}$  which we denote by  $\mathcal{A}^{\text{loc}}$ . It is dense in the topology of the  $C^*$ -norm, see [14].

If  $\mathcal{A}$  is the CAR algebra over  $l^2(\mathbb{Z}^2; \mathbb{C}^n)$  and  $\mathcal{A}'$  is the CAR algebra over  $l^2(\mathbb{Z}^2; \mathbb{C}^{n'})$  then we denote by  $\mathcal{A} \hat{\otimes} \mathcal{A}'$  the CAR algebra over  $l^2(\mathbb{Z}^2; \mathbb{C}^{n+n'})$  which contains  $\mathcal{A}$  and  $\mathcal{A}'$  as unital subalgebras. This algebra describes the system obtained by stacking the systems described by  $\mathcal{A}$  and  $\mathcal{A}'$  on top of each other.

### 2.2 Locally generated automorphisms

For any  $x \in \mathbb{Z}^2$  and any  $r \in \mathbb{N}$  we let  $B_x(r) := \{y \in \mathbb{Z}^2 : \operatorname{dist}(x,y) \leq r\}$  be the ball of radius r centered at x. Let  $\mathcal{F}$  be the collection of non-increasing, strictly positive functions  $f : \mathbb{R}^+ \to \mathbb{R}^+$  such that  $\lim_{r \uparrow \infty} r^p f(r) = 0$  for all p > 0. An operator  $A \in \mathcal{A}$  is said to be f-localised near a site  $x \in \mathbb{Z}^2$  if there is a sequence of operators  $A_r \in \mathcal{A}_{\mathcal{B}_x(r)}$  such that  $||A - A_r|| \leq f(r)||A||$  for all  $r \in \mathbb{N}$ . Any operator that is f-localised near some site for some  $f \in \mathcal{F}$  is called an almost local

operator. The almost local operators form a \*-subalgebra of  $\mathcal{A}$  which we denote by  $\mathcal{A}^{al}$ . Since  $\mathcal{A}^{\text{loc}} \subset \mathcal{A}^{\text{al}}$ , the algebra of almost local operators is norm-dense in  $\mathcal{A}$ .

A 0-chain F is a map  $\mathbb{Z}^2 \ni x \to F_x \in \mathcal{A}^{al}$  such that  $F_x$  is even and self-adjoint,  $\sup_x \|F_x\| < \infty$ and such that each  $F_x$  is f-localised near x for the same  $f \in \mathcal{F}$ . We say that F is an f-local 0-chain. For any  $A \in \mathcal{A}^{al}$  the sum  $[F, A] := \sum_{x \in \mathbb{Z}^2} [F_x, A]$  converges in norm to an element of  $\mathcal{A}^{al}$ . Hence, to every 0-chain corresponds a derivation  $\delta^F$  of the dense subalgebra  $\mathcal{A}^{al}$  which generates a one-parameter family of automorphisms  $\alpha_s^F$  with  $s \in \mathbb{R}$  called the time evolution generated by F and defined by  $\alpha_0^F = \operatorname{id}$  and  $-\operatorname{i}(\operatorname{d}\alpha_s^F(A)/\operatorname{d}s) = \alpha_s^F([F,A])$  for all  $A \in \mathcal{A}^{\operatorname{al}}$  and

Let us generalise this to a time-dependent setting. A family of 0-chains F(s) for  $s \in \mathbb{R}$  is called a time-dependent interaction (TDI) if  $\sup_{s,x} ||F_x(s)|| < \infty$  and for any  $s_0 \in \mathbb{R}$ , there is an  $f \in \mathcal{F}$  such that F(s) is an f-local 0-chain for all  $|s| \leq s_0$ . We assume moreover that the map  $s \mapsto F_x(s)$  is norm continuous for all  $x \in \mathbb{Z}^2$ . As above, a TDI F generates a strongly continuous family of automorphisms  $\alpha_s^F$  of  $\mathcal{A}$  for  $s \in \mathbb{R}$  which is given by

$$\alpha_s^F(A) = A + i \int_0^s du \, \alpha_u^F([F(u), A])$$
 (2.2)

for all  $A \in \mathcal{A}^{\mathrm{al}}$ . Equivalently,  $\alpha_s^F(A)$  is the unique solution of the initial value problem  $-\mathrm{i}\frac{\mathrm{d}}{\mathrm{d}s}\alpha_s^F(A) = \alpha_s^F([F(s),A])$  with  $\alpha_0^F(A) = A$ . An important consequence of the Lieb-Robinson bound [42] is that

$$\alpha_s^F \left( \mathcal{A}^{\text{al}} \right) \subset \mathcal{A}^{\text{al}},$$
 (2.3)

see e.g. [9, 46].

**Definition 2.1.** An automorphism  $\alpha$  of  $\mathcal{A}$  is a locally generated automorphism (LGA) if it is of the form  $\alpha = \alpha_s^F$  for a finite  $s \ge 0$  and a TDI F.

For any  $\Gamma \subset \mathbb{Z}^2$  the map  $F^{\Gamma}$  defined by  $F_x^{\Gamma} = \chi_{\Gamma}(x)F_x$  is again a TDI. Since TDIs are formal sums of even operators we have that  $\alpha_s^F \circ \theta = \theta \circ \alpha_s^F$  for any TDI F and any  $s \in \mathbb{R}$ .

The set of LGAs is a group since  $(\alpha_t^F)^{-1}$  and  $\alpha_t^F \circ \alpha_t^G$  are LGAs, generated by the TDIs

 $-\alpha_t^F(F(t))$  and  $G(t) + (\alpha_t^G)^{-1}(F(t))$  respectively. The group is non-abelian with

$$\alpha_t^F \circ \alpha_t^G = \alpha_t^{\alpha^F(G)} \circ \alpha_t^F, \tag{2.4}$$

see [8, 34].

#### 2.3 Symmetries

The charge at site  $x \in \mathbb{Z}^2$  is the self-adjoint operator  $Q_x := \sum_{i=1}^n a_{x,i}^* a_{x,i}$ . Clearly, each  $Q_x$ is an even self-adjoint operator of norm n supported on  $\{x\}$  so Q is a 0-chain. We denote by  $\rho_{\phi} = \alpha_{\phi}^{Q}$  the corresponding LGAs. Similarly, we let  $\rho_{\phi}^{\Gamma} = \alpha_{\phi}^{Q^{\Gamma}}$  for any  $\Gamma \subset \mathbb{Z}^{2}$  and note that  $\rho_{\phi}^{\Gamma}(\mathcal{A}_{\Gamma'}) = \mathcal{A}_{\Gamma'}$  for any  $\Gamma, \Gamma' \subset \mathbb{Z}^2$ . Moreover, since  $Q_x$  has integer spectrum,

$$\rho_{2\pi}^{\Gamma} = id \tag{2.5}$$

and hence  $\rho_{\phi+2\pi}^{\Gamma} = \rho_{\phi}^{\Gamma}$  for any  $\phi \in \mathbb{R}$ . We shall refer to these automorphisms as U(1) transformations. We note that the commutation relation  $[Q_x, a_{x,i}] = -a_{x,i}$  implies that

$$\rho_{\pi} = \theta. \tag{2.6}$$

We also equip A with an antilinear automorphism  $\tau$  such that

$$\tau^2 = \theta, \tag{2.7}$$

which we call *(fermionic) time reversal.* We further assume that its action is local in the sense that  $\tau(\mathcal{A}_{\Gamma}) = \mathcal{A}_{\Gamma}$  for all  $\Gamma \subset \mathbb{Z}^2$ . Finally, we assume that  $\tau(Q_x) = Q_x$  for all  $x \in \mathbb{Z}^2$ . It follows from this and the antilinearity of  $\tau$  that

$$\tau \circ \rho_{\phi} = \rho_{-\phi} \circ \tau \tag{2.8}$$

for all  $\phi \in \mathbb{R}$ .

We call  $(\mathcal{A}, \tau)$  a fermion system with time reversal. If  $(\mathcal{A}, \tau)$  and  $(\mathcal{A}', \tau')$  are fermion systems with time reversal, then  $(\mathcal{A} \hat{\otimes} \mathcal{A}', \tau \hat{\otimes} \tau')$  is again a fermion system with time reversal.

We say that a TDI F is time reversal symmetry preserving if  $\tau(F_x(s)) = -F_x(s)$  for all  $s \in \mathbb{R}$  and all  $x \in \mathbb{Z}^2$ . Similarly, we say the TDI F is U(1) symmetry preserving if  $\rho_{\phi}(F_x(s)) = F_x(s)$  for all  $\phi \in \mathbb{R}$ , all  $s \in \mathbb{R}$  and  $x \in \mathbb{Z}^2$ . A TDI that is simultaneously time reversal and U(1) symmetry preserving is called a symmetry preserving TDI. If F is a symmetry preserving TDI then the associated family of automorphisms  $\alpha_s^F$  are called symmetry preserving LGAs, and satisfy  $\rho_{\phi} \circ \alpha_s^F \circ \rho_{\phi}^{-1} = \tau \circ \alpha_s^F \circ \tau^{-1} = \alpha_s^F$ .

### 2.4 Symmetric short-range entangled states

Let  $(\mathcal{A}, \tau)$  be a fermion system with time reversal. A state  $\psi : \mathcal{A} \to \mathbb{C}$  is called homogeneous if

$$\psi \circ \theta = \psi$$
,

time reversal invariant if

$$\psi\circ\tau=\bar{\psi}$$

where  $\bar{\psi}$  is the antilinear map  $A \ni A \mapsto \bar{\psi}(A) = \psi(A^*) \in \mathbb{C}$ , and U(1)-invariant if

$$\psi \circ \rho_{\phi} = \psi$$

for all  $\phi \in \mathbb{R}$ . If a state is simultaneously homogeneous, time reversal invariant, and U(1)-invariant, then we call the state *symmetric*. A pure state  $\psi_0$  is said to be a pure product state if the restriction of  $\psi_0$  to each  $\mathcal{A}_x$  is pure.

**Definition 2.2.** (i) A state  $\psi : \mathcal{A} \to \mathbb{C}$  is short-range entangled (SRE) if there is a homogeneous pure product state  $\psi_0$  and an LGA  $\alpha$  such that  $\psi = \psi_0 \circ \alpha$ .

(ii) A state  $\psi : \mathcal{A} \to \mathbb{C}$  is stably SRE if there is a homogeneous product state  $\psi_0$  on an auxiliary fermion system  $\mathcal{A}'$  such that  $\psi \hat{\otimes} \psi_0$  is SRE.

Note that a (stably) SRE state is necessarily pure and homogeneous.

If  $\mathcal{A}$  is equipped with a time reversal and the SRE state is symmetric, we call it a symmetric SRE state on  $(\mathcal{A}, \tau)$ . A stably SRE state is called symmetric if, additionally, the stabilising product state  $\psi_0$  can be chosen symmetric w.r.t. a time reversal  $\tau'$  on  $\mathcal{A}'$ . If  $\psi$  is a symmetric SRE state on  $(\mathcal{A}, \tau)$  and  $\psi'$  is a symmetric SRE state on  $(\mathcal{A}', \tau')$  then  $\psi \hat{\otimes} \psi'$  is a symmetric SRE state on  $(\mathcal{A} \hat{\otimes} \mathcal{A}', \tau \hat{\otimes} \tau')$ .

### 2.5 Examples: Non-interacting electrons with spin

We consider a single-particle Hilbert space  $\mathcal{K}_m = l^2(\mathbb{Z}^2; \mathbb{C}^{2m}) \simeq l^2(\mathbb{Z}^2) \otimes \mathbb{C}^{2m}$  for some positive integer m. We write the internal space as  $\mathbb{C}^{2m} = \mathbb{C}^2 \otimes \mathbb{C}^m$  and fix an orthonormal tensor product basis  $\{e_{\sigma,i}\}_{\sigma \in \{\uparrow,\downarrow\}, i \in \{1,\cdots,m\}}$ . We think of the label  $\sigma$  as a spin-1/2 degree of freedom. Denote by  $\{\delta_x\}_{x \in \mathbb{Z}^2}$  the orthonormal position basis of  $l^2(\mathbb{Z}^2)$ . Then the vectors  $\{\delta_x \otimes e_{\sigma,i}\}$  form an orthonormal basis of  $\mathcal{K}_m$ . Denote by  $\mathscr{K}$  the complex conjugation on  $\mathcal{K}_m$  with respect to this basis, and let  $\mathscr{U}_T$  be the unitary given by

$$\mathscr{U}_T = \mathbb{1} \otimes \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \otimes \mathbb{1} \tag{2.9}$$

with respect to the tensor product decomposition  $\mathcal{K}_m = l^2(\mathbb{Z}^2) \otimes \mathbb{C}^2 \otimes \mathbb{C}^m$ . We define an antiunitary time-reversal by

$$\mathcal{T} = \mathcal{K} \mathcal{U}_T \tag{2.10}$$

which satisfies  $\mathcal{T}^2 = -1$ .

Let  $\mathcal{A}$  be the CAR algebra over  $\mathcal{K}_m$ . Then  $\tau(a(f)) = a(Tf)$  defines a fermionic time reversal on  $\mathcal{A}$ , i.e.  $\tau^2 = \theta$  and for any  $\Gamma \subset \mathbb{Z}^2$  we have  $\tau(\mathcal{A}_{\Gamma}) = \mathcal{A}_{\Gamma}$ , so  $(\mathcal{A}, \tau)$  is a fermion system with time reversal.

Let  $\mathscr{P}$  be an orthogonal projection on  $\mathcal{K}_m$ . The functional defined by

$$\omega_P(a^*(f_{n'})\cdots a^*(f_1)a(g_1)\cdots a(g_n)) = \delta_{n',n} \det\left(\langle g_i, \mathscr{P}f_j \rangle_{i,j=1}^n\right)$$
(2.11)

and extended to  $\mathcal{A}$  by linearity is a pure U(1)-invariant quasi-free state which we call the state corresponding to  $\mathscr{P}$ .

### 2.5.1 The empty product state

Consider the trivial projection  $\mathscr{P} = 0$  on the single particle space  $\mathcal{K}_m$ . Then the corresponding quasi-free state  $\omega_{\text{empty}} = \omega_{\mathscr{P}}$  is characterized by  $\omega_{\text{empty}}(Q_x) = 0$  for all  $x \in \mathbb{Z}^2$ . It follows that  $\omega_{\text{empty}}$  is a product state because it restricts to each  $\mathcal{A}_x$  as the unique state on  $\mathcal{A}_x$  with zero expectation value for  $Q_x$ . Since  $\mathscr{TPT}^* = \mathscr{P}$  we see that  $\omega_{\text{empty}}$  is time reversal invariant, so  $\omega_{\text{empty}}$  is a symmetric product state.

### 2.5.2 Quasi-free translation invariant spin Hall states

Let us consider the single particle Hilbert space  $\mathcal{K}_2 = l^2(\mathbb{Z}^2; \mathbb{C}^2 \otimes \mathbb{C}^2)$  where the first  $\mathbb{C}^2$  is the spin-1/2 degree of freedom, and the second  $\mathbb{C}^2$  is some additional internal degree of freedom.

Using the Fourier transform we identify  $\mathcal{K}_2 = l^2(\mathbb{Z}^2; \mathbb{C}^2 \otimes \mathbb{C}^2)$  with  $L^2(\mathbb{T}; \mathbb{C}^2 \otimes \mathbb{C}^2)$  where  $\mathbb{T}$  is the Brillouin torus. Let us further identify  $L^2(\mathbb{T}; \mathbb{C}^2 \otimes \mathbb{C}^2) \simeq L^2(\mathbb{T}; \mathbb{C}^2) \oplus L^2(\mathbb{T}; \mathbb{C}^2)$  where the first summand corresponds to  $\sigma = \uparrow$  and the second summand corresponds to  $\sigma = \downarrow$ .

Let  $\hat{\mathscr{P}}^{\uparrow}: \mathbb{T} \to \operatorname{Proj}(\mathbb{C}^2)$  be an analytic projection valued map of rank 1, which can be interpreted as a projection on  $L^2(\mathbb{T}; \mathbb{C}^2)$ . Note that  $E = \{(k, v) \in \mathbb{T} \times \mathbb{C}^2 : v \in \operatorname{Ran} \hat{\mathscr{P}}^{\uparrow}(k)\}$  describes a line bundle over the torus. We define the Chern number  $\operatorname{Ch}(\hat{\mathscr{P}}^{\uparrow})$  of  $\hat{\mathscr{P}}^{\uparrow}$  as the Chern number of this line bundle. Denote by  $\hat{\mathscr{P}}^{\downarrow} = \overline{\hat{\mathscr{P}}}^{\uparrow}$  the complex conjugate of  $\hat{\mathscr{P}}^{\uparrow}$  using the conjugation  $\mathscr{K}$ . Then

$$\hat{\mathscr{P}}_{SH}(k) = \hat{\mathscr{P}}^{\uparrow}(k) \oplus \hat{\mathscr{P}}^{\downarrow}(k) \tag{2.12}$$

defines an analytic projection valued map of rank 2 which satisfies  $\mathscr{T}\hat{\mathscr{P}}_{SH}(k)\mathscr{T}^* = \hat{\mathscr{P}}_{SH}(-k)$  by construction. By Fourier transform we obtain the corresponding projection  $\mathscr{P}_{SH} = \mathscr{P}^{\uparrow} \oplus \mathscr{P}^{\downarrow}$  on  $\mathcal{K}_2$  which satisfies  $\mathscr{T}\mathscr{P}_{SH}\mathscr{T}^* = \mathscr{P}_{SH}$ . The corresponding quasi-free state  $\omega_{SH} = \omega_{\mathscr{P}_{SH}}$  on  $\mathcal{A}$  is time reversal invariant. We call any state of this form a quasi-free translation invariant spin Hall state. To avoid possible confusion here, we note that with this definition, a spin Hall state may be topologically trivial in the sense that it has both vanishing Chern number and trivial Fu-Kane-Mele  $\mathbb{Z}_2$  index, see Section 5.

For later use we note that the analyticity of  $\hat{\mathscr{P}}_{SH}$  implies that  $\mathscr{P}_{SH}$  is exponentially local, i.e. there are constants  $C, \eta > 0$  such that

$$\left| \left\langle \delta_x \otimes e_{\sigma,i}, \, \mathscr{P}_{SH} \, \delta_y \otimes e_{\sigma',j} \right\rangle \right| \leqslant C e^{-\eta \|x - y\|_1}$$
 (2.13)

for all  $x, y \in \mathbb{Z}^2$ , all  $\sigma, \sigma' \in \{\uparrow\downarrow\}$ , and all  $i, j \in \{1, 2\}$ . We use the norm  $||x||_1 = |x_1| + |x_2|$  to measure spatial distances.

A typical example of the above setting is the Kane-Mele model without Rashba term and staggered chemical potential, see [27, Section 3.2]. In Section 5.1 we will show that

**Lemma 2.3.** Quasi-free translation invariant spin Hall states are stably SRE.

### 2.6 Symmetry protected phases

**Definition 2.4.** Two symmetric states  $\omega_1$  and  $\omega_2$  defined on the same system  $(\mathcal{A}, \tau)$  are called equivalent if there is a symmetry preserving LGA  $\alpha$  of  $\mathcal{A}$  such that  $\omega_2 = \omega_1 \circ \alpha$ .

We say that a symmetric SRE state  $(\omega_0, \mathcal{A}, \tau)$  is a symmetric product state if  $\omega_0$  is a product state. We call two symmetric states  $(\omega_1, \mathcal{A}_1, \tau_1)$  and  $(\omega_2, \mathcal{A}_2, \tau_2)$  stably equivalent if there are symmetric product states  $(\omega_1', \mathcal{A}_1', \tau_1')$  and  $(\omega_2', \mathcal{A}_2', \tau_2')$  such that  $(\omega_1 \hat{\otimes} \omega_1', \mathcal{A}_1 \hat{\otimes} \mathcal{A}_1', \tau_1 \hat{\otimes} \tau_1')$  and  $(\omega_2 \hat{\otimes} \omega_2', \mathcal{A}_2 \hat{\otimes} \mathcal{A}_2', \tau_2, \hat{\otimes} \tau_2')$  are equivalent.

Denote by  $\mathcal{P}$  the class of all symmetric stably SRE states. Equivalence and stable equivalence are equivalence relations on  $\mathcal{P}$ . If  $\omega_1, \omega_2 \in \mathcal{P}$  are stably equivalent then we write  $\omega_1 \sim \omega_2$ . The stable equivalence classes of  $\mathcal{P}$  are called symmetry protected topological (SPT) phases. Note that any two symmetric product states are stably equivalent. The equivalence class containing any symmetric product state is called the *trivial phase*.

### 2.7 The many-body $\mathbb{Z}_2$ -index

With these definitions, we can now state our main theorem which posits the existence of a  $\mathbb{Z}_2$ -valued index associated with any symmetric stably SRE state. The empty and spin Hall states described in Section 2.5 give examples of symmetric stably SRE states for which the index takes its two possible values. The index is multiplicative under stacking, and it is an invariant under the action of symmetry preserving LGAs and therefore constant on SPT phases. The theorem thus shows that there are two distinct such phases.

**Theorem 2.5.** There is a map  $\operatorname{Ind}_2 : \mathcal{P} \to \mathbb{Z}_2 = \{+1, -1\}$  such that:

- 1. If  $\omega_0$  is a product state then  $\operatorname{Ind}_2(\omega_0) = +1$ . If  $\omega_{SH}$  is a spin Hall state, then  $\operatorname{Ind}_2(\omega_{SH}) = (-1)^{\operatorname{Ch}(\hat{\mathscr{P}}^{\uparrow})}$ .
- 2. For any  $\omega_1, \omega_2 \in \mathcal{P}$ ,  $\operatorname{Ind}_2(\omega_1 \otimes \omega_2) = \operatorname{Ind}_2(\omega_1) \times \operatorname{Ind}_2(\omega_2)$ .
- 3. Let  $\omega_1, \omega_2 \in \mathcal{P}$ . If  $\omega_1 \sim \omega_2$  then  $\operatorname{Ind}_2(\omega_1) = \operatorname{Ind}_2(\omega_2)$ .

While we prove that our index generalizes the Fu-Kane-Mele index of the translation-invariant quantum spin Hall states, the same methods of Section 5 could be used to extend the equivalence to translation-invariant cases without spin conservation since the K-theory classification is known to be complete [38, 16].

### 3 Definition of the index

Throughout this section we fix a symmetric SRE state  $(\omega, \mathcal{A}, \tau)$  on a fermion system with time reversal and let  $\alpha = \alpha_1^F$  be an LGA and  $\omega_0$  a homogeneous product state such that  $\omega = \omega_0 \circ \alpha$ .

#### 3.1 Flux insertion

We first describe a construction which models the adiabatic insertion of a magnetic flux  $\phi$  piercing the plane.

### 3.1.1 Unique gapped ground states

**Definition 3.1.** A pure state  $\omega$  is a ground state of a 0-chain F if

$$\omega(A^*[F,A]) \geqslant 0, \quad \forall A \in \mathcal{A}^{loc}.$$
 (3.1)

The pure state  $\omega$  is a locally unique gapped ground state of F with gap  $\Delta$  if moreover

$$\omega(A^*[F,A]) \geqslant \Delta \,\omega(A^*A) \tag{3.2}$$

for all  $A \in \mathcal{A}^{loc}$  for which  $\omega(A) = 0$ .

It follows by density that if  $\omega$  is a locally unique gapped ground state with gap  $\Delta$ , then (3.1,3.2) hold for all A in the domain of the derivation  $\delta^F$ , and in particular for all  $A \in \mathcal{A}^{al}$ .

### 3.1.2 Symmetric parent Hamiltonian for $\omega$

Here we show the existence of a symmetric parent Hamiltonian for the symmetric SRE state  $\omega$  which we can use to describe the flux insertion process.

**Definition 3.2.** A 0-chain H is a symmetric parent Hamiltonian for a symmetric pure state  $\omega$  if  $\omega$  is a locally unique gapped ground state of H and  $H_x = \rho_{\phi}(H_x) = \tau(H_x)$  for all x and all  $\phi \in \mathbb{R}$ .

**Lemma 3.3.** The symmetric SRE state  $\omega$  has a symmetric parent Hamiltonian.

Proof. Since  $\omega_0$  is a homogeneous product state there is a 0-chain  $H^{(0)}$  such that  $H_x^{(0)} \in \mathcal{A}_{\{x\}}$  is even for each  $x \in \mathbb{Z}^2$  and  $\omega_0$  is the unique gapped ground state of  $H^{(0)}$  with gap one. Since  $\omega = \omega_0 \circ \alpha$  we claim that  $\omega$  is the unique gapped ground state of the 0-chain H' with gap one, where  $H_x' := \alpha^{-1}(H_x^{(0)})$ . Indeed, since  $\alpha(A) \in \mathcal{A}^{\mathrm{al}}$ , the series  $\sum_{x \in \mathbb{Z}^2} [H_x^{(0)}, \alpha(A)]$  converges in norm, and therefore  $[H', A] = \sum_{x \in \mathbb{Z}^2} [\alpha^{-1}(H_x^{(0)}), A] = \sum_{x \in \mathbb{Z}^2} \alpha([H_x^{(0)}, \alpha(A)])$  is also convergent since  $\alpha$  is an automorphism. But then  $\omega(A^*[H', A]) = \omega_0(\alpha(A)^*[H^{(0)}, \alpha(A)]) \geqslant 0$  for all  $A \in \mathcal{A}^{\mathrm{loc}}$  by the remark after Definition 3.1. Hence  $\omega$  is a ground state of H'. If H' has another ground state  $\omega'$ , then  $\omega' \circ \alpha^{-1}$  would be a ground state of  $H_0$ , and so  $\omega$  is the unique ground state of H'. That it is gapped follows by a similar argument. Note that the  $H_x'$  are still even, but they may not be symmetric since  $\alpha$  is in general not symmetric. Therefore, we average  $H_x'$  over the time reversal and U(1) symmetries to obtain a symmetric 0-chain H:

$$H_x := \frac{1}{4\pi} \int_0^{2\pi} d\phi \, \rho_\phi \left( H_x' + \tau (H_x') \right). \tag{3.3}$$

Since  $\omega$  is a symmetric state,

$$\omega(A^*[H,A]) = \frac{1}{4\pi} \int_0^{2\pi} d\phi \,\omega\Big(\rho_{-\phi}(A)^* \left[ \left( H_x' + \tau(H_x') \right), \rho_{-\phi}(A) \right] \Big)$$
(3.4)

and the integrand is greater than  $2\omega(A^*A)$  for all A such that  $\omega(A) = 0$  since  $\omega$  is a gapped ground state of H'. Hence  $\omega$  is a locally unique gapped ground state of H with gap one.  $\square$ 

### 3.1.3 U(1) transformations on half-planes and quasi-adiabatic generators

A half-line in  $\mathbb{R}^2$  is determined by a base point  $a \in \mathbb{R}^2$  and a non-zero direction vector  $v \in \mathbb{R}^2$ . To any half-line l = (a, v) we can associate a half-plane (with marked boundary)

$$h_l := \{ x \in \mathbb{R}^2 : (x - a) \cdot (Rv) > 0 \}$$
(3.5)

where R is the clockwise rotation by  $\pi/2$ . In the following, we will use the same notation for a subset  $S \subset \mathbb{R}^2$  and  $S \cap \mathbb{Z}^2 \subset \mathbb{Z}^2$ .

Let  $h=h_l$  be the half-plane associated to the half-line l=(a,v) and consider for any symmetric 0-chain H the 0-chains  $H^{h,\phi}$  with

$$H_x^{h,\phi} = \rho_{-\phi}^h(H_x). \tag{3.6}$$

Since each  $H_x$  is time reversal and U(1) invariant, and using  $\tau \circ \rho_{\phi}^h = \rho_{-\phi}^h \circ \tau$ , we find that the  $H_x^{h,\phi}$  are U(1) invariant and  $\tau(H_x^{h,\phi}) = H_x^{h,-\phi}$ .

If H is a symmetric parent Hamiltonian of a symmetric pure state  $\omega$  then  $\omega^{h,\phi} := \omega \circ \rho_{\phi}^{h}$  is the locally unique gapped ground state of  $H^{h,\phi}$  with gap one for all  $\phi \in \mathbb{R}$ . It follows that the states  $\omega^{h,\phi}$  are related to  $\omega = \omega^{h,0}$  by the quasi-adiabatic flow [46], see also [30, 28, 10, 45], which we briefly recall. Let

$$K_x^{h,H}(\phi) := \int_{\mathbb{R}} dt \, W(t) \, \alpha_t^{H^{h,\phi}} \left( \frac{dH_x^{h,\phi}}{d\phi} \right) \tag{3.7}$$

where  $W : \mathbb{R} \to \mathbb{R}$  is an odd bounded function such that  $\lim_{t \uparrow \infty} |t|^p W(t) = \lim_{t \uparrow \infty} |t|^p W(-t) = 0$  for all  $p \in \mathbb{N}$  and such that its Fourier transform  $\hat{W}(E)$  equals -i/E whenever |E| > 1. Then:

- 1. The family  $K_x^{h,H}(\phi)$  defines a TDI  $K^{h,H}$ .
- 2. There is  $g \in \mathcal{F}$  such that

$$||K_x^{h,H}(\phi)|| \le g(\operatorname{dist}(x,\partial h)) \tag{3.8}$$

for all  $x \in \mathbb{Z}^2$  and all  $\phi \in \mathbb{R}$ .

3. We have

$$K_x^{h,H}(\phi) = \rho_{\phi'}(K_x^{h,H}(\phi)) = \theta(K_x^{h,H}(\phi)) = \tau(K_x^{h,H}(-\phi))$$
(3.9)

for all  $x \in \mathbb{Z}^2$  and all  $\phi, \phi' \in \mathbb{R}$ .

Points 1 and 2 are standard consequences of the Lieb-Robinson bound, the fast decay of W and the U(1) invariance of all  $H_x$ . The first two equalities of 3 follow from U(1) invariance while the last equality follows from the antilinearity of  $\tau$  through  $\tau \left(W(t)\alpha_t^{H^{h,\phi}}(\partial_\phi H_x^{h,\phi})\right) = W(t)\alpha_{-t}^{H^{h,-\phi}}(\partial_\phi H_x^{h,-\phi})$  and the fact that W is odd.

 $W(t)\alpha_{-t}^{H^{h,-\phi}}(\partial_{\phi}H_{x}^{h,-\phi})$  and the fact that W is odd. The TDI  $K^{h,H}$  is called the quasi adiabatic generator for the family  $H^{h,\phi}$ . It generates a quasi-adiabatic flow  $\alpha_{\phi}^{h,H}:=\alpha_{\phi}^{K^{h,H}}$  which has the following crucial property:

$$\omega^{h,\phi} = \omega \circ \alpha_{\phi}^{h,H} \tag{3.10}$$

for all  $\phi$ , where again  $\omega = \omega^{h,0}$ . In other words, the quasi-adiabatic flow  $\alpha_{\phi}^{h,H}$  implements the U(1) transformations  $\rho_{\phi}^{h}$  on the state  $\omega$ . Unlike the U(1) transformations  $\rho_{\phi}^{h}$  whose generator is supported on the full half-plane h, the generator of the quasi-adiabatic flow  $K^{h,H}$  is supported only near the line  $\partial h$ , see (3.8).

### 3.1.4 Defect states

The above construction allows us to define an automorphism that corresponds to inserting a magnetic flux at the point a in the plane. Let l = (a, v) be any half-line with base point a. We define a TDI  $K^{l,H}$  as a restriction of the TDI  $K^{h,H}$  as follows:

$$K_x^{l,H}(\phi) = \begin{cases} K_x^{h,H}(\phi) & \text{if } (x-a) \cdot v > 0\\ 0 & \text{otherwise} \end{cases}$$
 (3.11)

for all  $\phi \in \mathbb{R}$ . We denote by

$$\gamma_\phi^{l,H} := \alpha_\phi^{K^{l,H}}$$

the LGA it generates, and we shall refer to such LGAs as flux insertion automorphisms. We define defect states by

$$\omega^{l,H,\phi} := \omega \circ \gamma_{\phi}^{l,H}. \tag{3.12}$$

As emphasized by the notation, the states  $\omega^{l,H,\phi}$  depend on the symmetric parent Hamiltonian H, on the choice of half-line l, and on the flux  $\phi$ .

**Lemma 3.4.** Let H be a symmetric parent Hamiltonian of  $\omega$  and let l be a half-line. For any  $\phi, \phi' \in \mathbb{R}$  we have

$$\gamma_{\phi}^{l,H} \circ \rho_{\phi'} = \rho_{\phi'} \circ \gamma_{\phi}^{l,H}, \quad \gamma_{\phi}^{l,H} \circ \theta = \theta \circ \gamma_{\phi}^{l,H} \quad and \quad \gamma_{\phi}^{l,H} \circ \tau = \tau \circ \gamma_{-\phi}^{l,H}. \tag{3.13}$$

It follows in particular that

$$\omega^{l,H,\phi} = \omega^{l,H,\phi} \circ \rho_{\phi'} = \overline{\omega^{l,H,-\phi}} \circ \tau. \tag{3.14}$$

*Proof.* The first two identities are immediate consequences of (3.9). For the last one, we let  $\tilde{\gamma}_{\phi}^{l,H} := \tau^{-1} \circ \gamma_{\phi}^{l,H} \circ \tau$ . Then for any  $A \in \mathcal{A}$ 

$$\tilde{\gamma}_{\phi}^{l,H}(A) = A - i \int_{0}^{\phi} d\phi' \, \tilde{\gamma}_{\phi'}^{l,H} \left( [\tau^{-1}(K^{l,H}(\phi')), A] \right)$$
(3.15)

where we used the antilinearity of  $\tau^{-1}$ . By (3.9),  $\tau^{-1}(K^{l,H}(\phi')) = \tau(K^{l,H}(\phi')) = K^{l,H}(-\phi')$ . Hence,  $\tilde{\gamma}_{\phi}^{l,H}(A)$  solves the same equation (2.2) as  $\gamma_{-\phi}^{l,H}(A)$ , and since A is arbitrary we conclude that  $\tau^{-1} \circ \gamma_{\phi}^{l,H} \circ \tau = \gamma_{-\phi}^{l,H}$  indeed. The identity  $\omega^{l,H,\phi} = \omega^{l,H,\phi} \circ \rho_{\phi'}$  is now immediate. As for the last one, we use the invariance of  $\omega$  under parity  $\theta = \tau^2$  to conclude that

$$\omega^{l,H,\phi} = \omega \circ \tau^2 \circ \gamma_{\phi}^{l,H} = \omega \circ \tau \circ \gamma_{-\phi}^{l,H} \circ \tau = \bar{\omega} \circ \gamma_{-\phi}^{l,H} \circ \tau \tag{3.16}$$

since  $\omega$  is time reversal invariant.

#### 3.1.5 Locality

We conclude this section with a locality result for LGAs generated by TDIs satisfying (3.8). Let l = (a, v) be a half-line. For any  $\varsigma \in (0, 2\pi)$ , the open subset of  $\mathbb{R}^2$  given by

$$\Lambda_{l,\varsigma} := \{ x \in \mathbb{R}^2 : (x - a) \cdot v > ||x - a|| \cos(\varsigma/2) \}. \tag{3.17}$$

will be called the *cone* with apex at  $a \in \mathbb{R}^2$ , axis  $v \in \mathbb{R}^2$  of unit length, and opening angle  $\varsigma$ .

**Lemma 3.5.** Let l be a half-line and let  $\Lambda_{l,\varsigma}$  be a cone. Let F be an f-local TDI. Assume moreover that there is  $g \in \mathcal{F}$  such that  $||F_x|| \leq g(\operatorname{dist}(x,l))$  for all  $x \in \mathbb{Z}^2$ . Then for any  $s_0 \in \mathbb{R}$  and any  $\varsigma \in (0, 2\pi)$ , there is  $h \in \mathcal{F}$  such that

$$\|\alpha_s^F(A) - A\| \le \|A\|h(r)$$
 (3.18)

for all  $|s| \leq s_0$  and all  $A \in \mathcal{A}_{\Lambda_{l,\varsigma}^c \cap B_a(r)^c}$ . Moreover, the same holds if A is replaced by  $\alpha_{s'}^G(A)$  for any TDI G.

*Proof.* For the duration of the proof, we write  $\Lambda = \Lambda_{l,\varsigma}$  and  $\tilde{\Lambda} = \Lambda_{l,\varsigma/2}$ , see Figure 4. If




Figure 4: The sets forming the keyhole.

 $x \in \tilde{\Lambda} \cup B_a(\frac{r}{2})$  belongs to the 'keyhole' then

$$||[F_x, A]|| = ||[F_x - F_{x, \operatorname{dist}(x, \Lambda^c \cap B_a(r)^c)}, A]|| \le 2||A|| f(\operatorname{dist}(x, \Lambda^c \cap B_a(r)^c))$$
(3.19)

since  $F_x$  is f-localized near x, see Section 2.2. On the other hand, if  $x \in \tilde{\Lambda}^c \cap B_a(\frac{r}{2})^c$ , then

$$||[F_x, A]|| \le 2||F_x|| ||A|| \le 2||A|| g(\operatorname{dist}(x, l))$$
(3.20)

We now sum these estimates over x. If  $x \in B_a(\frac{r}{2})$ , then  $(\operatorname{dist}(x, \Lambda^c \cap B_a(r)^c) \ge \frac{r}{2}$ . If  $x \in \tilde{\Lambda} \setminus B_a(\frac{r}{2})$ , then  $(\operatorname{dist}(x, \Lambda^c \cap B_a(r)^c) \ge \frac{\theta}{4} ||x||$ . Hence,

$$\sum_{x \in \tilde{\Lambda} \cup B_a(\frac{r}{2})} \| [F_x, A] \| \le 2 \|A\| \left( cr^2 f(\frac{r}{2}) + \sum_{x \in \tilde{\Lambda}, \|x\| > r/2} f(\frac{\theta}{4} \|x\|) \right)$$
(3.21)

which is bounded above by  $2\|A\|\tilde{f}(r)$  for some  $\tilde{f} \in \mathcal{F}$ . For the complement of the keyhole, we have that  $\operatorname{dist}(x,l) \geq \frac{\theta}{4}\|x\|$  and so by assumption

$$\sum_{x \in \tilde{\Lambda}^c \cap B_a(\frac{r}{2})^c} \| [F_x, A] \| \leqslant 2 \| A \| \sum_{x \in \tilde{\Lambda}^c, \|x\| > r/2} g(\frac{\theta}{4} \| x \|)$$
 (3.22)

which is again bounded by  $2||A||\tilde{g}(r)$  for some  $\tilde{g} \in \mathcal{F}$ . We conclude that there is  $\tilde{h} \in \mathcal{F}$  such that

$$||[F, A]|| \le ||A||\tilde{h}(r).$$
 (3.23)

The first claim of the lemma now follows from (2.2). For the second claim, it suffices to use (2.4) to conclude that

$$\|\alpha_s^F(\alpha_r^G(A)) - \alpha_r^G(A)\| = \|(\alpha_r^G)^{-1} \circ \alpha_s^F \circ \alpha_r^G(A) - A\| = \|\alpha_s^{(\alpha_r^G)^{-1}(F)}(A) - A\|$$
(3.24)

and note that  $(\alpha_r^G)^{-1}(F)$  is a TDI satisfying the assumptions of the lemma.

#### 3.2 Defect states at $\pm \pi$ flux

For the remainder of this section we fix a half-line l and a symmetric parent Hamiltonian H of  $\omega$ , and write  $\omega^{\pm} := \omega^{l,H,\pm\pi}$ . These two states correspond to inserting a  $\pm\pi$  magnetic flux defect into  $\omega$ . Our main goal is to show that these states are unitary equivalent and related to each other by time reversal. We can then study the transformation properties of vector representatives of these states under time reversal, finally leading to the definition of the index.

### 3.2.1 Almost local equivalence

We show that the states  $\omega^{\pm}$  are almost local perturbations of each other, which will imply by the results of Appendix A that they are unitarily equivalent through an operator that is almost local.

**Definition 3.6.** Let  $f \in \mathcal{F}$  and let  $\psi$  and  $\psi'$  be states on  $\mathcal{A}$ . We say  $\psi$  and  $\psi'$  are f-close if for all r > 0 we have

$$\left|\psi(A) - \psi'(A)\right| \leqslant f(r) \|A\| \quad \forall A \in \mathcal{A}_{B(r)^c}^{\mathrm{loc}}. \tag{3.25}$$

The state  $\psi'$  is an almost local perturbation of the state  $\psi$  if  $\psi, \psi'$  are f-close for some  $f \in \mathcal{F}$ .

Recall that  $\omega = \omega_0 \circ \alpha$  for some  $\alpha = \alpha_1^F$  and define homogeneous states

$$\tilde{\omega}^{\pm} := \omega^{\pm} \circ (\alpha \circ \gamma_{\pi}^{l,H})^{-1}. \tag{3.26}$$

It follows immediately from these definitions that

$$\tilde{\omega}^+ = \omega_0, \qquad \tilde{\omega}^- = \omega_0 \circ \alpha \circ \gamma_{-\pi}^{l,H} \circ \left(\gamma_{\pi}^{l,H}\right)^{-1} \circ \alpha^{-1}.$$
 (3.27)

**Lemma 3.7.** Let  $\Lambda = \Lambda_{l,\pi}$ . There is  $f \in \mathcal{F}$  such that for any  $A \in \mathcal{A}^{loc}_{\Lambda \cap B_a(r)^c} \cup \mathcal{A}^{loc}_{\Lambda^c \cap B_a(r)^c}$  we have

$$\left|\tilde{\omega}^{-}(A) - \tilde{\omega}^{+}(A)\right| \leqslant f(r)\|A\|. \tag{3.28}$$

for all r > 0.

*Proof.* In this proof, f stands for a function in  $\mathcal{F}$  and it changes from equation to equation. We write  $\tilde{\omega}^- = \omega_0 \circ \tilde{\beta}$ , where  $\tilde{\beta} := \alpha \circ \gamma_{-\pi}^{l,H} \circ \left(\gamma_{\pi}^{l,H}\right)^{-1} \circ \alpha^{-1}$ . Let first  $A \in \Lambda \cap B_a(r)^c$ . Let  $h = h_l$  be the half-plane associated to the half-line l = (a, v) and let

$$\beta = \alpha \circ \alpha_{-\pi}^{h,H} \circ (\alpha_{\pi}^{h,H})^{-1} \circ \alpha^{-1}. \tag{3.29}$$

We claim that

$$\|\tilde{\beta}(A) - \beta(A)\| \le f(r)\|A\| \tag{3.30}$$

where f is independent of A. We first prove the claim with the replacements  $\tilde{\beta} \to \gamma_{-\pi}^{l,H} \circ \left(\gamma_{\pi}^{l,H}\right)^{-1}$  and  $\beta \to \alpha_{-\pi}^{h,H} \circ \left(\alpha_{\pi}^{h,H}\right)^{-1}$  since this yields the claim by the second part of Lemma 3.5. The family of automorphisms  $\left(\gamma_{\phi}^{l,H}\right)^{-1} \circ \alpha_{\phi}^{h,H}$  is an LGA generated by

$$(K^{h,H}(\phi) - K^{l,H}(\phi)) + i(\alpha_{\phi}^{h,H})^{-1} \left[ \int_{0}^{\phi} du \, \alpha_{u}^{h,H} \left( \left[ K^{h,H}(u) - K^{l,H}(u), \gamma_{\phi,u}^{l,H}(K^{l,H}(u)) \right] \right) \right], (3.31)$$

see e.g. [8, Lemma 5.4]. If we denote by -l the half-line -l = (a, -v), we have that  $K^{l,H} - K^{h,H} = K^{-l,H}$  which satisfies the assumptions of Lemma 3.5, hence so does the TDI (3.31). The Lemma (with  $\Lambda$  replacing  $\Lambda^c$  since it is applied to the cone  $\Lambda^c$ ) shows that

$$\left\| \left( \gamma_{-\pi}^{l,H} \right)^{-1} \circ \alpha_{-\pi}^{h,H} \left( (\alpha_{\pi}^{h,H})^{-1}(A) \right) - \left( (\alpha_{\pi}^{h,H})^{-1}(A) \right) \right\| \leqslant f(r) \|A\| \tag{3.32}$$

for all  $A \in \Lambda \cap B_a(r)^c$ . Similarly,  $\gamma_{\phi}^{l,H} \circ (\alpha_{\phi}^{h,H})^{-1}$  is generated by  $-\alpha_{\phi}^{h,H}(K^{h,H}(\phi) - K^{l,H}(\phi))$  which implies that

$$||A - \gamma_{\pi}^{l,H} \circ (\alpha_{\pi}^{h,H})^{-1}(A)|| \le f(r)||A||.$$
 (3.33)

These estimates yield the claim (3.30) since

$$\left\| \gamma_{-\pi}^{l,H} \circ \left( \gamma_{\pi}^{l,H} \right)^{-1}(A) - \alpha_{-\pi}^{h,H} \circ \left( \alpha_{\pi}^{h,H} \right)^{-1}(A) \right\| = \left\| A - \gamma_{\pi}^{l,H} \circ \left( \gamma_{-\pi}^{l,H} \right)^{-1} \circ \alpha_{-\pi}^{h,H} \circ \left( \alpha_{\pi}^{h,H} \right)^{-1}(A) \right\|$$

$$\leq \left\| A - \gamma_{\pi}^{l,H} \circ \left( \alpha_{\pi}^{h,H} \right)^{-1}(A) \right\| + \left\| \gamma_{\pi}^{l,H} \circ \left( \alpha_{\pi}^{h,H} \right)^{-1}(A) - \gamma_{\pi}^{l,H} \circ \left( \gamma_{-\pi}^{l,H} \right)^{-1} \circ \alpha_{-\pi}^{h,H} \circ \left( \alpha_{\pi}^{h,H} \right)^{-1}(A) \right\| .$$

Now note that

$$\omega \circ \alpha_{-\pi}^{h,H} \circ \left(\alpha_{\pi}^{h,H}\right)^{-1} = \omega^{h,-\pi} \circ \left(\alpha_{\pi}^{h,H}\right)^{-1} = \omega^{h,\pi} \circ \left(\alpha_{\pi}^{h,H}\right)^{-1} = \omega, \tag{3.34}$$

where the first and last equalities are by the definition of the states (3.10) and the second one follows from  $\omega^{h,-\pi} = \omega \circ \rho_{\pi} = \omega \circ \rho_{-\pi} = \omega^{h,\pi}$ . It follows that  $\omega_0 \circ \beta = \omega_0 = \tilde{\omega}^+$  and so for all  $A \in \Lambda \cap B_a(r)^c$  we find

$$\left|\tilde{\omega}^{-}(A) - \tilde{\omega}^{+}(A)\right| = \left|\omega_{0} \circ \left((\tilde{\beta} - \beta)(A)\right)\right| \leqslant f(r) \|A\|. \tag{3.35}$$

If  $A \in \Lambda^c \cap B_a(r)^c$ , then Lemma 3.5 implies that  $\|(\gamma_\phi^{l,H})^{-1}(A) - A\| \leq f(r)\|A\|$  by similar arguments and therefore that

$$\|\tilde{\beta}(A) - A\| \leqslant f(r)\|A\| \tag{3.36}$$

instead of (3.30), with the same conclusion as above since  $\omega_0 = \tilde{\omega}^+$ .

**Proposition 3.8.** There exists a homogeneous unitary  $U \in \mathcal{A}^{al}$  such that  $\omega^+ = \omega^- \circ Ad[U]$ .

*Proof.* Since  $\tilde{\omega}^+ = \omega_0$  is a product state, it follows from Lemma 3.7 and Lemma A.1 that the homogeneous state  $\tilde{\omega}^+$  is an almost local perturbation of  $\tilde{\omega}^-$ , see Definition 3.6. By Lemma A.2 this implies that there is a homogeneous unitary  $\tilde{U} \in \mathcal{A}^{\text{al}}$  such that  $\tilde{\omega}^+ = \tilde{\omega}^- \circ \text{Ad}[\tilde{U}]$ . Recalling that  $\omega^{\pm} = \tilde{\omega}^{\pm} \circ (\alpha \circ \gamma_{\pi}^{l,H})$ , we have that

$$\omega^+ = \tilde{\omega}^+ \circ (\alpha \circ \gamma_\pi^{l,H}) = \tilde{\omega}^- \circ \operatorname{Ad}[\widetilde{U}] \circ (\alpha \circ \gamma_\pi^{l,H}) = \tilde{\omega}^- \circ (\alpha \circ \gamma_\pi^{l,H}) \circ \operatorname{Ad}[U] = \omega^- \circ \operatorname{Ad}[U] \ (3.37)$$

where  $U = (\alpha \circ \gamma_{\pi}^{l,H})^{-1}(\widetilde{U})$  is again almost local by (2.3), and is homogeneous since  $\alpha$  and  $\gamma_{\pi}^{l,H}$  commute with parity.

### 3.2.2 GNS representation and time reversal

Let  $(\Pi, \mathcal{H}, \Omega^+)$  be the GNS triple of  $\omega^+$  and fix a homogeneous unitary  $U \in \mathcal{A}^{al}$  such that  $\omega^+ = \omega^- \circ' Ad[U]$ , which exists be Proposition 3.8. Then the unit vector  $\Omega^- := \Pi(U) \Omega^+$  represents the state  $\omega^-$ :

$$\omega^{-}(A) = \langle \Omega^{-}, \Pi(A) \Omega^{-} \rangle \tag{3.38}$$

for all  $A \in \mathcal{A}$ . We now investigate the transformation properties of the vectors  $\Omega^{\pm}$  under time reversal. To this end we first note the following general fact.

**Lemma 3.9.** Let  $\nu : \mathcal{A} \to \mathbb{C}$  be a pure state that is invariant under an antilinear automorphism  $\tau : \mathcal{A} \to \mathcal{A}$ , namely  $\nu \circ \tau = \bar{\nu}$ . Let  $(\Pi_{\nu}, \mathcal{H}_{\nu}, \Omega_{\nu})$  be the GNS triple of  $\nu$ . Then there exists a unique antiunitary operator T acting on  $\mathcal{H}_{\nu}$  such that  $T\Omega_{\nu} = \Omega_{\nu}$  and

$$T \Pi_{\nu}(A) T^* = \Pi_{\nu}(\tau(A))$$
 (3.39)

for all  $A \in \mathcal{A}$ .

*Proof.* Let K be a conjugation on  $\mathcal{H}_{\nu}$  such that  $K\Omega_{\nu} = \Omega_{\nu}$ . Then the map  $\tilde{\Pi}_{\nu} = \operatorname{Ad}[K] \circ \Pi_{\nu} \circ \tau$ :  $\mathcal{A} \to \mathcal{B}(\mathcal{H}_{\nu})$  is a (linear) \*-representation of  $\mathcal{A}$  on  $\mathcal{H}_{\nu}$ . Moreover,  $\Omega_{\nu}$  is cyclic for  $\tilde{\Pi}_{\nu}$  and

$$\langle \Omega_{\nu}, \tilde{\Pi}_{\nu}(A) \Omega_{\nu} \rangle = \langle \Pi_{\nu}(\tau(A)) K \Omega_{\nu}, K \Omega_{\nu} \rangle = \overline{\nu(\tau(A))} = \nu(A)$$
 (3.40)

for all  $A \in \mathcal{A}$ . It follows that  $(\tilde{\Pi}_{\nu}, \mathcal{H}_{\nu}, \Omega_{\nu})$  is a GNS triple for  $\nu$ . By uniqueness of the GNS representation, there is a unique unitary V such that  $V\Omega_{\nu} = \Omega_{\nu}$  and

$$\tilde{\Pi}_{\nu}(A) = V \Pi_{\nu}(A) V^*.$$
 (3.41)

This is equivalent to

$$\Pi_{\nu}(\tau(A)) = (KV) \Pi_{\nu}(A) (KV)^*,$$
(3.42)

and  $(KV)\Omega_{\nu} = \Omega_{\nu}$ . Hence (3.39) holds for the choice T = KV.

If T' were another antiunitary that satisfies the conditions of the lemma, then T' = KV' for a unitary V' = KT', and  $V'\Omega_{\nu} = \Omega_{\nu}$ . Moreover,

$$V' \Pi_{\nu}(A) (V')^* = KT' \Pi_{\nu}(A) (T')^* K = (\text{Ad}[K] \circ \Pi_{\nu} \circ \tau)(A) = \tilde{\Pi}_{\nu}(A). \tag{3.43}$$

Since V was the unique unitary with these properties, we conclude that V' = V and therefore T' = KV' = KV = T, showing uniqueness.

With this in hand, we come back to our specific setting with  $(\Pi, \mathcal{H}, \Omega^+)$  the GNS triple of  $\omega^+$  and  $\Omega^- \in \mathcal{H}$  given by  $\Omega^- = \Pi(U)\Omega^+$ .

**Lemma 3.10.** There is a unique antiunitary T acting on the GNS Hilbert space  $\mathcal{H}$  such that  $\Omega^- = T\Omega^+$  and such that T implements time reversal, i.e.

$$\Pi(\tau(A)) = T\Pi(A)T^* \tag{3.44}$$

for all  $A \in \mathcal{A}$ .

*Proof.* The unitary  $U \in \mathcal{A}^{al}$  is such that

$$\omega^{-} = \omega^{+} \circ \operatorname{Ad}[U^{*}]. \tag{3.45}$$

Since  $\omega^+ = \overline{\omega^-} \circ \tau$  by Lemma 3.4, it follows that

$$\omega^{-} = \overline{\omega^{-}} \circ \tau \circ \operatorname{Ad}[U^{*}]. \tag{3.46}$$

In other words, the antilinear automorphism  $\tilde{\tau} = \tau \circ \operatorname{Ad}[U^*]$  leaves the state  $\omega^-$  invariant. By Lemma 3.9 it follows that there is a unique antiunitary  $\tilde{T}$  such that

$$\Pi(\tilde{\tau}(A)) = \tilde{T}\Pi(A)\tilde{T}^* \tag{3.47}$$

for all  $A \in \mathcal{A}$ , and  $\tilde{T}\Omega^- = \Omega^-$ . It follows that  $\tilde{T}\Pi(U)\Omega^+ = \tilde{T}\Omega^- = \Omega^-$  and

$$\Pi(\tau(A)) = \Pi(\tilde{\tau} \circ \operatorname{Ad}[U](A)) = \tilde{T}\Pi(U)\Pi(A)\Pi(U^*)\tilde{T}^*$$
(3.48)

for all  $A \in \mathcal{A}$ . We see that  $T := \tilde{T}\Pi(U)$  satisfies the requirements of the lemma. Suppose now that T' is another such operator. Then  $\tilde{T}' = T'\Pi(U^*)$  satisfies

$$\tilde{T}'\Pi(A)(\tilde{T}')^* = \Pi((\tau \circ \operatorname{Ad}[U^*])(A)) = \Pi(\tilde{\tau}(A))$$
(3.49)

and  $\tilde{T}'\Omega^- = T'\Pi(U^*)\Omega^- = T'\Omega^+ = \Omega^-$ . The uniqueness of  $\tilde{T}$  implies that  $\tilde{T}' = \tilde{T}$  and hence T' = T.

### 3.2.3 The states $\omega^+$ and $\omega^-$ have the same parity

With these preliminary constructions, we can now prove a property that is essential to the definition of the index: that the defect states  $\omega^{\pm}$  have the same parity.

Recall that  $(\Pi, \mathcal{H}, \Omega^+)$  is the GNS triple of  $\omega^+$  and we fixed a homogeneous unitary  $U \in \mathcal{A}^{\mathrm{al}}$  such that  $\omega^+ = \omega^- \circ \mathrm{Ad}[U]$ , whose existence is guaranteed by Proposition 3.8. Let  $\Omega^- = \Pi(U)\Omega^+$  be the corresponding vector representative of  $\omega^-$ . Since  $\omega^+$  is homogeneous there is a unique unitary  $\Theta \in \mathcal{B}(\mathcal{H})$  such that

$$\Theta\Omega^+ = \Omega^+, \qquad \Pi(\theta(A)) = \Theta\Pi(A)\Theta^*$$
 (3.50)

for all  $A \in \mathcal{A}$ . The fact that  $\omega^-$  is also homogeneous implies that  $\Omega^-$  is an eigenvector of the operator  $\Theta^* = \Theta$  (since  $\Theta^2 = \mathbb{1}$ ). We conclude that  $\Theta\Omega^- = \pm \Omega^-$ . We note that the property  $\Theta\Omega^+ = \Omega^+$  is really a choice of the overall sign of  $\Theta$ , so the eigenvalue  $\pm 1$  above should be understood as relative to that choice.

**Lemma 3.11.** Let  $\Theta$  be such that (3.50) hold. Then  $\Theta\Omega^- = \Omega^-$  and there is an even unitary  $U \in \mathcal{A}^{al}$  such that  $\omega^+ = \omega^- \circ Ad[U]$ .

In the following, we shall say that two states have the same parity whenever they are, as above, unitarily equivalent through an even unitary element of the algebra.

Proof. Let T be the antiunitary operator on  $\mathcal{H}$  provided by Lemma 3.10. Since  $\omega^+$  is invariant under the  $2\pi$ -periodic family of automorphisms  $\rho_{\phi}$  for  $\phi \in \mathbb{R}$  there is a unique strongly continuous 1-parameter group of unitaries  $V_{\phi} \in \mathcal{B}(\mathcal{H})$  that implement the U(1) transformations, namely  $V_{\phi}\Omega^+ = \Omega^+$  and

$$V_{\phi}\Pi(A)V_{\phi}^* = \Pi(\rho_{\phi}(A)) \tag{3.51}$$

for all  $A \in \mathcal{A}$ . The map  $\phi \mapsto V_{\phi}$  is  $2\pi$ -periodic and such that  $V_0 = \mathbb{I}$ . Since  $\omega^-$  is also invariant under  $\rho_{\phi}$ , and  $\Pi$  is irreducible since  $\omega^+$  is pure, we have  $V_{\phi}\Omega^- = e^{in\phi}\Omega^-$  for all  $\phi \in \mathbb{R}$  and some  $n \in \mathbb{Z}$ .

The  $2\pi$ -periodic family of unitaries  $\tilde{V}_{\phi} := e^{-in\phi}T^*V_{\phi}T$  implements  $\tau^{-1} \circ \rho_{\phi} \circ \tau = \rho_{-\phi}$  in the GNS representation:

$$\tilde{V}_{\phi}\Pi(A)\tilde{V}_{\phi}^{*} = T^{*}V_{\phi}T\Pi(A)T^{*}V_{\phi}^{*}T = \Pi(\tau^{-1} \circ \rho_{\phi} \circ \tau(A)) = \Pi(\rho_{-\phi}(A)). \tag{3.52}$$

Moreover,

$$\tilde{V}_{-\phi}\Omega^{+} = e^{in\phi} T^{*} V_{-\phi} T \Omega^{+} = e^{in\phi} T^{*} V_{-\phi} \Omega^{-} = e^{in\phi} T^{*} e^{in\phi} \Omega^{-} = \Omega^{+}, \tag{3.53}$$

where we used antilinearity of  $T^*$ . It follows from uniqueness of the  $V_{\phi}$  that  $\tilde{V}_{-\phi} = V_{\phi}$ , in other words,

$$V_{\phi} = e^{in\phi} T^* V_{-\phi} T. \tag{3.54}$$

Now note that  $V_{\pi} = \Theta$  by uniqueness of  $\Theta$ , and since  $\tau^2 = \theta$  we have that the unitary  $T^2$  also implements parity in the GNS representation. It follows that  $T^2 = \lambda \Theta$  for some  $\lambda \in U(1)$ . In particular,  $T^2 = \lambda V_{\pi}$  commutes with  $V_{\phi}$  for all  $\phi \in \mathbb{R}$ . Using (3.54) twice yields

$$V_{\phi} = e^{in\phi} T^* \left( e^{-in\phi} T^* V_{\phi} T \right) T = e^{2in\phi} (T^*)^2 V_{\phi} T^2 = e^{2in\phi} V_{\phi}$$
 (3.55)

for all  $\phi \in \mathbb{R}$ . We conclude that n = 0, so  $\Theta\Omega^- = V_\pi\Omega^- = \Omega^-$ , as required.

The existence of a homogeneous unitary  $U \in \mathcal{A}^{al}$  such that  $\omega^+ = \omega^- \circ \operatorname{Ad}[U]$  is guaranteed by Proposition 3.8, so we only need to show that this unitary is even. Since  $\Omega^-$  was chosen so that  $\Omega^- = \Pi(U)\Omega^+$ . By the above we conclude that  $\Theta^*\Omega^- = \Pi(U)\Theta^*\Omega^+$  and hence  $\Omega^- = \Pi(\theta(U))\Omega^+$ . Since U is homogeneous, this is compatible with  $\Omega^- = \Pi(U)\Omega^+$  only if U is even.

#### 3.3 Definition of the index

**Definition 3.12.** Let  $(\Pi, \mathcal{H})$  be an irreducible representation of a fermion system with time reversal  $(\mathcal{A}, \tau)$  such that the time reversal is implemented by an antiunitary operator T acting on  $\mathcal{H}$ , namely  $\Pi(\tau(A)) = T\Pi(A)T^*$  for all  $A \in \mathcal{A}$ .

A vector  $\Psi \in \mathcal{H}$  is a Kramers singlet if  $T^2\Psi = \Psi$ , and  $\Psi$  belongs to a Kramers pair if  $T^2\Psi = -\Psi$ . If  $(\Pi, \mathcal{H}, \Psi)$  is a GNS representation of a pure state  $\psi$ , then we say that  $\psi$  is a Kramers singlet if  $\Psi$  is, and we say that  $\psi$  belongs to a Kramers pair if  $\Psi$  does.

**Remark 3.13.** If  $\Psi$  belongs to a Kramers pair then if  $\Psi' = T\Psi$  we have

$$\langle \Psi, \Psi' \rangle = \langle \Psi, T\Psi \rangle = \langle T^2 \Psi, T\Psi \rangle = -\langle \Psi, \Psi' \rangle \tag{3.56}$$

which shows that  $\Psi$  and  $\Psi' = T\Psi$  are orthogonal. The 'Kramers pair' to which  $\Psi$  belongs is the pair of vectors  $\Psi$  and  $\Psi'$ .

We now show that the defect state  $\omega^+$  is either a Kramers singlet, or it belongs to a Kramers pair with the defect state  $\omega^-$ . Note first that this statement makes sense since Lemma 3.10 shows that time reversal is implemented by an antiunitary operator T in the GNS representation  $(\Pi, \mathcal{H}, \Omega^+)$  of  $\omega^+$ .

**Lemma 3.14.** The defect state  $\omega^+$  is either a Kramers singlet or it belongs to a Kramers pair.

*Proof.* Let  $\Theta$  be as in (3.50). Since  $\tau^2 = \theta$  we must have  $T^2 = \lambda \Theta$  for some phase  $\lambda \in U(1)$ . It follows that

$$T^2\Omega^+ = \lambda\Theta\Omega^+ = \lambda\Omega^+. \tag{3.57}$$

By Lemma 3.11 the vector  $\Omega^- = T\Omega^+$  satisfies  $\Theta\Omega^- = \Omega^-$ . By acting on both sides of (3.57) with T we obtain

$$T^3\Omega^+ = \bar{\lambda}T\Omega^+ = \bar{\lambda}\Omega^-. \tag{3.58}$$

On the other hand,

$$T^{3}\Omega^{+} = \lambda \Theta T \Omega^{+} = \lambda \Theta \Omega^{-} = \lambda \Omega^{-}. \tag{3.59}$$

We conclude that  $\lambda$  is real, hence  $\lambda \in \{+1, -1\}$  as claimed.

**Definition 3.15.** Let  $\omega$  be a symmetric SRE state with symmetric parent Hamiltonian H. For any half-line l we define

$$\operatorname{Ind}_{2}(\omega, H, l) := \begin{cases} +1 & \text{if } \omega^{+} \text{ is a Kramers singlet,} \\ -1 & \text{if } \omega^{+} \text{ belongs to a Kramers pair.} \end{cases}$$
(3.60)

We will show in Section 4.1 that  $\operatorname{Ind}_2(\omega, H, l)$  is actually independent of the choice of symmetric parent Hamiltonian H and the choice of half-line l. We will anticipate this result and denote the index from now on by  $\operatorname{Ind}_2(\omega)$ .

Note that the index is for now defined only for SRE states and not yet for the larger class of stably SRE states. This extension will be carried out in Section 4.4 as a consequence of the multiplicativity under stacking and the triviality of the index for product states.

Before we study the properties of the index we have just defined, we first show that it has an expression that does not rely on the GNS representation at all. While this expression loses it transparency to physical interpretation, this more 'algebraic' formulation hints at the announced invariance under symmetry preserving LGAs. Both formulations can be used in the proofs of the next section but we have chosen to consistency to rely only on Definition 3.15.

**Proposition 3.16.** Let  $\omega^-$  be the  $-\pi$  flux state and let  $U \in \mathcal{A}$  be a homogeneous unitary such that  $\omega^+ = \omega^- \circ \operatorname{Ad}[U]$ . Then

$$\operatorname{Ind}_{2}(\omega) = \omega^{-}(U\tau(U)). \tag{3.61}$$

*Proof.* We recall first that the index is the value of  $T^2$  in the subspace spanned by the GNS vectors  $\Omega^{\pm}$  for  $\omega^{\pm}$  and secondly that we can choose the phases of  $\Omega^{\pm}$  and of U so that  $\Omega^{-} = T\Omega^{+}$  and  $\Omega^{-} = \Pi(U) \Omega^{+}$ . It follows that

$$\operatorname{Ind}_{2}(\omega) = \langle \Omega^{+}, T^{2}\Omega^{+} \rangle = \langle T\Omega^{+}, T^{*}\Omega^{+} \rangle = \langle \Omega^{-}, T^{*}\Pi(U)^{*}\Omega^{-} \rangle = \langle \Pi(U)\Omega^{+}, T^{*}\Pi(U)^{*}\Omega^{-} \rangle$$

$$(3.62)$$

$$= \langle T^* \Omega^-, \Pi(U)^* T^* \Pi(U)^* \Omega^- \rangle = \langle \Omega^-, \Pi(U) T \Pi(U) T^* \Omega^- \rangle = \omega^- (U \tau(U))$$
 (3.63)

where we used repeatedly the antiunitarity of T.

## 4 Properties of the index

### 4.1 Independence of choices

As pointed out earlier, the construction leading to the definition of  $\operatorname{Ind}_2(\omega)$  requires many choices that are not unique. We shall now show that these choices do not affect the value of the index. More importantly, we prove in this section that the index is a bonafide topological invariant, namely that it is constant under deformations by symmetry preserving LGAs.

### 4.1.1 Kramers pairing is invariant under even unitaries

We first note that 'being a Kramers singlet' or 'belonging to a Kramers pair' is invariant under even unitaries, a property that we will use repeatedly in this section.

**Lemma 4.1.** Let  $\psi$  be a homogeneous pure state on a fermion system with time reversal  $(\mathcal{A}, \tau)$  that is a Kramers singlet or belongs to a Kramers pair. Let  $V \in \mathcal{A}$  be an even unitary. Then  $\psi' = \psi \circ \operatorname{Ad}[V^*]$  is a Kramers singlet if  $\psi$  is, and  $\psi'$  belongs to a Kramers pair if  $\psi$  does.

*Proof.* Let  $(\Pi, \mathcal{H}, \Psi)$  be the GNS triple of  $\psi$ . The state  $\psi'$  is represented by the vector  $\Psi' = \Pi(V)\Psi$ . Since  $\tau^2 = \theta$  we have that the unitary  $T^2$  implements fermion parity. Since V is even we therefore find

$$T^{2}\Psi' = T^{2}\Pi(V)\Psi = \Pi(\theta(V))T^{2}\Psi = \Pi(V)T^{2}\Psi = \pm \Psi'$$
(4.1)

by Definiton 
$$3.12$$

### 4.1.2 Independence from symmetric parent Hamiltonian

The role played by the parent Hamiltonian H constructed in Section 3.1.2 is to allow for a definition of the flux insertion automorphism and therefore of the defect states  $\omega^{\pm}$ . Different parent Hamiltonians yield different defect states, but they all result in the same index.

**Proposition 4.2.** Let  $\omega$  be a symmetric SRE state and let  $H_1, H_2$  be two parent Hamiltonians for  $\omega$ . Then

$$\operatorname{Ind}_{2}(\omega, H_{1}, l) = \operatorname{Ind}_{2}(\omega, H_{2}, l). \tag{4.2}$$

*Proof.* We claim that the two defect states  $\omega_j^- = \omega \circ \gamma_{-\pi}^{l,H_j}, j = 1,2$  are unitarily equivalent through an even unitary. The argument follows closely the proof of Lemma 3.7 and Proposition 3.8. Instead of considering  $\gamma_{-\pi}^{l,H} \circ \left(\gamma_{\pi}^{l,H}\right)^{-1}$ , we have here  $\gamma_{-\pi}^{l,H_1} \circ \left(\gamma_{-\pi}^{l,H_2}\right)^{-1}$ . The same argument allows one to replace  $\gamma_{-\phi}^{l,H_1} \circ \left(\gamma_{-\phi}^{l,H_2}\right)^{-1}$  by  $\alpha_{-\phi}^{h,H_1} \circ \left(\alpha_{-\phi}^{h,H_2}\right)^{-1}$  for all  $\phi \in [-\pi,\pi]$  and all  $A \in \Lambda \cap B_a(r)^c$ , see (3.30). We also have that

$$\omega \circ \alpha_{-\phi}^{h,H_1} \circ \left(\alpha_{-\phi}^{h,H_2}\right)^{-1} = \omega^{h,-\phi} \circ \left(\alpha_{-\phi}^{h,H_2}\right)^{-1} = \omega \tag{4.3}$$

and hence again

$$|\omega_0(A) - \omega_0 \circ \alpha \circ \gamma_{-\phi}^{l, H_1} \circ \left(\gamma_{-\phi}^{l, H_2}\right)^{-1} \circ \alpha^{-1}(A)| \leqslant f(r) ||A|| \tag{4.4}$$

for all  $\phi \in [-\pi, \pi]$  and all  $A \in \Lambda \cap B_a(r)^c$ . The case  $A \in \Lambda^c \cap B_a(r)^c$  follows similarly with  $\gamma_{-\phi}^{l,H_1} \circ \left(\gamma_{-\phi}^{l,H_2}\right)^{-1}$  being close to the identity, which yields the same estimate. By Lemma A.1, we conclude that the two states are local perturbations of each other, see Definition 3.6, and hence by Proposition A.2 that there is a unitary  $W_{\phi}$  such that  $\omega_0 \circ \alpha \circ \gamma_{-\phi}^{l,H_1} \circ \left(\gamma_{-\phi}^{l,H_2}\right)^{-1} \circ (\alpha)^{-1} = \omega_0 \circ \operatorname{Ad}[W_{-\phi}]$ . The family of homogeneous states  $\omega_0 \circ \alpha \circ \gamma_{-\phi}^{l,H_1} \circ \left(\gamma_{-\phi}^{l,H_2}\right)^{-1} \circ \alpha^{-1}$  is weak-\* continuous so the  $W_{-\phi}$  are all even by Lemma A.7.

Altogether, we conclude that

$$\omega_{1}^{-} = \omega_{0} \circ \operatorname{Ad}[W_{-\pi}] \circ \alpha \circ \gamma_{-\pi}^{l,H_{2}} = \omega_{2}^{-} \circ \operatorname{Ad}[(\alpha \circ \gamma_{-\pi}^{l,H_{2}})^{-1}(W_{-\pi})]. \tag{4.5}$$

where the unitary  $V = (\alpha \circ \gamma_{-\pi}^{l,H_2})^{-1}(W_{-\pi})$  is even. We conclude by Lemma 4.1.

### 4.1.3 Independence of the half-line

We finally turn to the irrelevance of the choice of half-line l.

An argument similar to the above yields invariance under moving the endpoint of the half-line along the half-line. If  $l_1=(a,v)$  and  $l_2=(a+bv,v)$  then  $\gamma_{\phi}^{l_1,H}\circ (\gamma_{\phi}^{l_2,H})^{-1}$  is generated by  $-\gamma_{\phi}^{l_2,H}(K^{l_1,H}(\phi)-K^{l_2,H}(\phi))$  which is an almost local even observable and hence  $\gamma_{\phi}^{l_1,H}\circ (\gamma_{\phi}^{l_2,H})^{-1}=\mathrm{Ad}[V_{\phi}]$  for a family of even unitaries  $V_{\phi}$ . Invariance of the index then follows from Lemma 4.1.

Let us now consider half-lines  $l_1 = (a, v_1)$  and  $l_2 = (a, v_2)$  with common endpoint a. Now the corresponding defect states are truly distinct at infinity. However, they differ at infinity by a U(1) transformation with phase  $\pi$  that does not impact the flux, which explains why the index is invariant:

**Lemma 4.3.** Let  $\Gamma \subset \mathbb{Z}^2$ . If  $\psi$  is a Kramers singlet or belongs to a Kramers pair then the state  $\psi' = \psi \circ \rho_{\pi}^{\Gamma}$  is a Kramers singlet if  $\psi$  is, and  $\psi'$  belongs to a Kramers pair of  $\psi$  did.

*Proof.* Let  $(\Pi, \mathcal{H}, \Psi)$  be the GNS triple of  $\psi$  and let T implement time reversal. Then  $T^2\Psi = \pm \Psi$  depending on whether  $\psi$  is a Kramers singlet, or belongs to a Kramers pair. Since  $\psi' = \psi \circ \rho^{\Gamma_{\pi}}$  we have that  $(\Pi', \mathcal{H}, \Psi)$  with  $\Pi' = \Pi \circ \rho_{\pi}^{\Gamma}$  is the GNS triple of  $\psi$ . Using (2.8) and  $\rho_{-\pi}^{\Gamma} = \rho_{\pi}^{\Gamma}$  we see that time reversal is still implemented in  $\Pi'$  by T. The claim follows immediately.

**Proposition 4.4.** Let  $\omega$  be a symmetric SRE state. Then

$$\operatorname{Ind}_2(\omega, H, l_1) = \operatorname{Ind}_2(\omega, H, l_2) \tag{4.6}$$

for any two half-lines  $l_1, l_2$ .

*Proof.* By the remarks at the beginning of this section we can restrict our attention to  $l_1 = (a, v_1)$  and  $l_2 = (a, v_2)$ . Let  $\varsigma = \arccos v_1 \cdot v_2$  be the angle between  $l_1$  and  $l_2$ . It is sufficient to consider the case  $\varsigma < \pi$ , then  $v = \frac{1}{2}(v_1 + v_2)$  is non-zero and we put l = (a, v). We denote

$$\Lambda = \Lambda_{l,\varsigma}, \qquad \Lambda_1 = \Lambda_{l_1,\varsigma}. \tag{4.7}$$

see Figure 5




Figure 5: The U(1) transformation in  $\Lambda$  slides the half-line  $l_1$  onto  $l_2$ .

First of all, if  $A \in \mathcal{A}_{B_a(r)^c \cap \Lambda_1^c}$ , then  $\|\gamma_{-\phi}^{l_1,H} \circ \rho_{\phi}^{\Lambda}(A) - \rho_{\phi}^{\Lambda}(A)\| \leqslant f(r)\|A\|$  by Lemma 3.5. Hence

$$|\omega \circ \gamma_{-\phi}^{l_1, H} \circ \rho_{\phi}^{\Lambda}(A) - \omega \circ \rho_{\phi}^{\Lambda}(A)| \leqslant f(r) ||A||. \tag{4.8}$$

We now use that  $\rho_{\phi}^{\Lambda} = \rho_{\phi} \circ \rho_{-\phi}^{\Lambda^c}$  and the U(1) invariance of  $\omega$  to conclude that

$$\omega \circ \rho_{\phi}^{\Lambda} = \omega \circ \rho_{-\phi}^{\Lambda^{c}} = \omega \circ \alpha_{-\phi}^{K^{\Lambda^{c}, H}} \tag{4.9}$$

where  $K^{\Lambda^c,H}(\phi)$  is defined as in (3.7) but with the cone  $\Lambda^c$  replacing the half-plane h. Lemma 3.5 yields that  $\|\alpha_{-\phi}^{K^{\Lambda^c,H}}(A) - \gamma_{-\phi}^{l_2,H}(A)\| \leq f(r)\|A\|$  by the choice of  $\Lambda$ . Altogether, we have that

$$|\omega \circ \gamma_{-\phi}^{l_1,H} \circ \rho_{\phi}^{\Lambda}(A) - \omega \circ \gamma_{-\phi}^{l_2,H}(A)| \leqslant f(r) \|A\| \tag{4.10}$$

for all  $A \in \mathcal{A}_{B_a(r)^c \cap \Lambda_1^c}$ . Secondly, if  $A \in \mathcal{A}_{B_a(r)^c \cap \Lambda_1}$  we use that  $\rho_{\phi}^{\Lambda}(A)$  is almost localized in the same set to conclude that  $\|\gamma_{-\phi}^{l_1,H} \circ \rho_{\phi}^{\Lambda}(A) - \rho_{-\phi}^{\Lambda} \circ \rho_{\phi}^{\Lambda}(A)\| \leq f(r)\|A\|$  and hence

$$|\omega \circ \gamma_{-\phi}^{l_1, H} \circ \rho_{\phi}^{\Lambda}(A) - \omega \circ \rho_{-\phi}^{\Lambda} \circ \rho_{\phi}^{\Lambda}(A)| \leqslant f(r) ||A||, \tag{4.11}$$

namely  $|\omega \circ \gamma_{-\phi}^{l_1,H} \circ \rho_{\phi}^{\Lambda}(A) - \omega(A)| \leq f(r) \|A\|$ . It remains to note that  $\|\gamma_{-\phi}^{l_2,H}(A) - A\| \leq f(r) \|A\|$  to conclude that

$$|\omega \circ \gamma_{-\phi}^{l_1,H} \circ \rho_{\phi}^{\Lambda}(A) - \omega \circ \gamma_{-\phi}^{l_2,H}(A)| \leqslant f(r)||A|| \tag{4.12}$$

for all  $A \in \mathcal{A}_{B_a(r)^c \cap \Lambda_1}$ . The two estimates (4.10,4.12) and Lemma A.1 imply that  $\omega_0 \circ \alpha \circ \gamma_{-\phi}^{l_1,H} \circ \rho_{\phi}^{\Lambda} \circ \left(\gamma_{-\phi}^{l_2,H}\right)^{-1} \circ \alpha^{-1}$  is an almost local perturbation of  $\omega_0$  and so Lemma A.2 yields a unitary  $W_{-\phi}$  such that

$$\omega \circ \gamma_{-\phi}^{l_1,H} \circ \rho_{\phi}^{\Lambda} = \omega \circ \gamma_{-\phi}^{l_2,H} \circ \operatorname{Ad}[W_{-\phi}]. \tag{4.13}$$

By continuity, the unitaries are even, see Lemma A.7. The proof is concluded by putting  $\phi = -\pi$  and applying Lemmas 4.3 and 4.1.

Propositions 4.2 and 4.4 allow us to make the following definition announced earlier.

**Definition 4.5.** Let  $\omega$  be a symmetric SRE state. We define

$$\operatorname{Ind}_{2}(\omega) := \operatorname{Ind}_{2}(\omega, H, l) \tag{4.14}$$

where H is any symmetric parent Hamiltonian for  $\omega$  and l is any half-line.

Recall that the existence of a symmetric parent Hamiltonian is guaranteed by Lemma 3.3.

With this, we have concluded the proof that the index of a symmetric SRE state is a well-defined property of the state. We conclude this section with two fundamental properties of the index: its multiplicativity under stacking and its 'topological invariance', namely the fact that it is invariant under the action of symmetry preserving LGAs. This makes it an index of the symmetry protected phases of symmetric stably SRE states.

### 4.2 Stacking

**Proposition 4.6.** Let  $(\omega_1, \mathcal{A}_1, \tau_1)$  and  $(\omega_2, \mathcal{A}_2, \tau_2)$  be symmetric SRE states. Then  $\omega_1 \otimes \omega_2$  is a symmetric SRE state on the fermion system with time-reversal  $(\mathcal{A}_1 \otimes \mathcal{A}_2, \tau_1 \otimes \tau_2)$  and

$$\operatorname{Ind}_{2}(\omega_{1} \otimes \omega_{2}) = \operatorname{Ind}_{2}(\omega_{1}) \times \operatorname{Ind}_{2}(\omega_{2}). \tag{4.15}$$

*Proof.* Let  $(\Pi_i, \mathcal{H}_i, \Omega_i^+)$  be the GNS representation of  $\omega_i^+$  and let  $T_i$  be the antiunitary that implements time-reversal in the representation  $(\Pi_i, \mathcal{H}_i)$ . The index  $\operatorname{Ind}_2(\omega_i)$  is defined by  $T_i^2 \Omega_i^+ = \operatorname{Ind}_2(\omega_i) \Omega_i^+$ , see Lemma 3.14.

Let  $\omega = \omega_1 \hat{\otimes} \omega_2$ . Then the 0-chain H with  $H_x = H_x^{(1)} \hat{\otimes} \mathbb{1} + \mathbb{1} \hat{\otimes} H_x^{(2)}$  is a symmetric parent Hamiltonian for  $\omega$  which we can use to construct a defect state  $\omega^+ = \omega \circ \gamma_\pi^{l,H}$ . Let  $\Theta_1 \in \mathcal{B}(\mathcal{H}_1)$  be the unique unitary that implements parity in the representation  $(\Pi_1, \mathcal{H}_1)$  such that  $\Theta_1 \Omega_1^+ = \Omega_1^+$ . One easily checks that  $\omega^+ = \omega_1^+ \hat{\otimes} \omega_2^+$ , and the GNS representation of  $\omega^+$  can be identified with  $(\Pi, \mathcal{H}, \Omega^+)$  where  $\mathcal{H} = \mathcal{H}_1 \otimes \mathcal{H}_2$ ,  $\Omega^+ = \Omega_1^+ \otimes \Omega_2^+$ , and  $\Pi$  is determined by  $\Pi(A_1 \hat{\otimes} A_2) = \Pi(A_1)\Theta_1^{\sigma(A_2)} \otimes \Pi_2(A_2)$  for homogeneous  $A_1 \in \mathcal{A}_1$  and  $A_2 \in \mathcal{A}_2$  and where  $\sigma(A) = 0$  if A is even, and  $\sigma(A) = 1$  if A is odd. Time reversal  $\tau = \tau_1 \hat{\otimes} \tau_2$  is implemented in this representation by  $T = T_1 \otimes T_2$ . The index  $\operatorname{Ind}_2(\omega)$  is defined by  $T^2 \Omega^+ = \operatorname{Ind}_2(\omega) \Omega^+$ , but

$$T^{2}\Omega^{+} = (T_{1}^{2}\Omega_{1}^{+}) \otimes (T_{2}^{2}\Omega_{2}^{+}) = (\operatorname{Ind}_{2}(\omega_{1})\Omega_{1}^{+}) \otimes (\operatorname{Ind}_{2}(\omega_{2})\Omega_{2}^{+}) = \operatorname{Ind}_{2}(\omega_{1}) \times \operatorname{Ind}_{2}(\omega_{2})\Omega^{+}.$$
(4.16)

This proves the claim.

### Invariance under symmetry preserving LGAs

We now turn to showing that  $\operatorname{Ind}_2(\omega)$  is a bonafide topological invariant.

In order to show invariance of the index under symmetry preserving LGAs we recall Definition 3.12 and first prove the following lemma:

**Lemma 4.7.** Let  $\psi$  be a pure state on a fermionic system with time reversal  $(\mathcal{A}, \tau)$  and let  $\beta$ be an automorphism of A that commutes with  $\tau$ . If  $\psi$  is a Kramers singlet then so is  $\psi \circ \beta$ , and if  $\psi$  belongs to a Kramers pair then so does  $\psi \circ \beta$ .

*Proof.* Let  $(\Pi, \mathcal{H}, \Psi)$  be the GNS triple of  $\psi$  and suppose time reversal is implemented by an antiunitary operator T acting on  $\mathcal{H}$ . The GNS triple of  $\psi' = \psi \circ \beta$  is  $(\Pi', \mathcal{H}, \Psi)$  with  $\Pi' = \Pi \circ \beta$ . Since  $\tau^{-1} \circ \beta \circ \tau = \beta$  we have  $\Pi'(\tau(A)) = T\Pi'(A)T^*$ , so time reversal is also implemented by Tin the representation  $\Pi'$ . The claim follows immediately.

**Proposition 4.8.** Let  $\omega$  a symmetric SRE state and let G be a symmetry preserving TDI. Then

$$\operatorname{Ind}_{2}(\omega) = \operatorname{Ind}_{2}(\omega \circ \alpha_{1}^{G}). \tag{4.17}$$

*Proof.* Let  $G^R$  be the restriction of the TDI G to the right half plane  $R = \{x \in \mathbb{Z}^2 : x_1 > 0\}$  and put  $G^L = G - G^R$ . Then  $\alpha_s^G \circ (\alpha_s^{G^R})^{-1}$  is an LGA generated by  $\alpha_s^{G^R}(G^L(s))$ . The TDIs  $G_s^R$ and  $\tilde{G}^L = \alpha_s^{G^R}(G^L(s))$  satisfy the assumptions of Lemma 3.5 for the left half plane and right half plane respectively, yielding a decomposition

$$\alpha_s^G = \alpha_s^{\tilde{G}^L} \circ \alpha_s^{G^R}. \tag{4.18}$$

We will write  $\alpha^G = \alpha_1^G$ ,  $\alpha^L = \alpha_1^{\tilde{G}^L}$  and  $\alpha^R = \alpha_1^{G^R}$  so  $\alpha^G = \alpha^L \circ \alpha^R$ . Let furthermore  $\tilde{\omega} = \omega \circ \alpha^G$  and consider the state  $\tilde{\omega}_L = \omega \circ \alpha^L$  which looks like  $\omega$  far to the right and like  $\tilde{\omega}$  for to the left of the vertical axis. We will prove the proposition by showing that the index of  $\omega$  is equal to the index of  $\tilde{\omega}_L$ , and that the index of  $\tilde{\omega}_L$  is equal to the index of  $\tilde{\omega}$ .

If H is a symmetric parent Hamiltonian for  $\omega$  then the collection of almost local operators  $(\alpha^L)^{-1}(H_x)$  for  $x \in \mathbb{Z}^2$  define a symmetric parent Hamiltonian  $\tilde{H}^L$  for  $\tilde{\omega}_L$ . The  $\pi$ -flux defect states for  $\omega$  and  $\tilde{\omega}_L$  may be taken to be

$$\omega^{+} = \omega \circ \gamma^{+}, \qquad \tilde{\omega}_{L}^{+} = \tilde{\omega}_{L} \circ \tilde{\gamma}^{+} \tag{4.19}$$

where  $\gamma^+ = \gamma_\pi^{l,H}$ ,  $\tilde{\gamma}^+ = \gamma_\pi^{l,\tilde{H}_L}$  and l = (0,(1,0)) is the right horizontal axis. Lemma 3.5 implies that  $\tilde{H}^L - H$  is a TDI that is almost localized on the left half plane. Denoting by h the upper half-plane, it follows from this and the definition (3.6) that  $K^{h,\tilde{H}^L}$  –  $K^{h,H}$  is localized along the left horizontal axis (0, (-1, 0)). The LGA  $\gamma_{\phi}^{l,\tilde{H}^L} \circ (\gamma_{\phi}^{l,H})^{-1}$  is generated by  $\gamma_{\phi}^{l,H}(K^{l,\tilde{H}^L} - K^{l,H})$ . Since  $K^{l,\cdot}$  is the restriction of  $K^{h,\cdot}$  to the right half-plane, it follows by the above that  $K^{l,\tilde{H}^L} - K^{l,H}$  is in fact an almost local even operator so there is a family of even unitaries  $\tilde{V}_{\phi}$  such that  $\gamma_{\phi}^{l,\tilde{H}^{L}} = \operatorname{Ad}[\tilde{V}_{\phi}] \circ \gamma_{\phi}^{l,H}$ , hence

$$\gamma_{\phi}^{l,\tilde{H}^L} = \gamma_{\phi}^{l,H} \circ \operatorname{Ad}[V_{\phi}], \tag{4.20}$$

where the  $V_{\phi} = (\gamma_{\phi}^{l,H})^{-1}(\tilde{V}_{\phi})$  are again even. We therefore find

$$\tilde{\omega}_L^+ = \omega^+ \circ \beta_L^+ \circ \operatorname{Ad}[V_\pi] \tag{4.21}$$

where  $\beta_L^+ = (\gamma^+)^{-1} \circ \alpha_L \circ \gamma^+$ .

In order to unburden the notation, let us write  $K(s) = K^{l,H}(\pi s)$  so  $\gamma^+ = \alpha_1^K$  and put  $\widetilde{K} = \alpha_1^{\widetilde{G}^L}(K)$ . Using (2.4) we find that

$$\beta_L^+ = (\alpha_1^K)^{-1} \circ \alpha_1^{\widetilde{K}} \circ \alpha_L. \tag{4.22}$$

We will show that the automorphism  $(\alpha_1^K)^{-1} \circ \alpha_1^{\widetilde{K}}$  is given by conjugation with an almost local even unitary. Indeed we have  $\alpha_1^{\widetilde{K}} = \alpha_1^{\widetilde{K}-K+K} = \alpha_1^J \circ \alpha_1^K$  with  $J(s) = \alpha_s^K(\widetilde{K}(s) - K(s))$ . Now,

$$\widetilde{K}(s) - K(s) = i \int_{0}^{1} ds' \, \alpha_{s'}^{\widetilde{G}^{L}} ([\widetilde{G}^{L}(s'), K(s)])$$
 (4.23)

is an even almost local self-adjoint operator because  $\tilde{G}^L$  is supported near the left half plane while K is supported near the right horizontal axis. It follows that J(s) is an almost local even self-adjoint operator for all s and so  $\alpha_1^J = \operatorname{Ad}[\widetilde{W}]$  for an even almost local unitary  $\widetilde{W}$ . We therefore obtain that

$$\beta_L^+ = (\alpha_1^K)^{-1} \circ \operatorname{Ad}[\widetilde{W}] \circ \alpha_1^K \circ \alpha_L = \alpha_L \circ \operatorname{Ad}[W]$$
(4.24)

where  $W = (\alpha_1^K \circ \alpha_L)^{-1}(\widetilde{W})$  is also an almost local even unitary. Recalling (4.21) this yields  $\widetilde{\omega}_L^+ = \omega^+ \circ \alpha^L \circ \operatorname{Ad}[WV_{\pi}]$ . Since  $WV_{\pi}$  is an even unitary and  $\alpha^L$  commutes with  $\tau$ , we conclude by Lemma 4.1 and Lemma 4.7 that the defect state  $\widetilde{\omega}_L$  is a Kramers singlet/belongs to a Kramers pair if and only if  $\omega^+$  does.

Repeating the same argument with the replacements  $\omega \to \tilde{\omega}_L$  and  $\tilde{\omega}_L \to \tilde{\omega} = \omega \circ \alpha^G$  concludes the proof.

### 4.4 Extension of the index to symmetric stably SRE states

In order to extend the index to symmetric stably SRE states we must first show that the index of symmetric product states is trivial.

**Lemma 4.9.** Let  $\omega_0$  be a symmetric product state, then  $\operatorname{Ind}_2(\omega) = 1$ .

Proof. Since  $\omega_0$  is a symmetric product state we can choose a symmetric parent Hamiltonian  $H^{(0)}$  that consists of purely on-site terms, see the proof of Lemma 3.3. It follows that  $H_x^{h,\phi} = H_x^{h,0}$  for all  $\phi$ , see (3.6), and in turn that the generator  $K_x^{l,H}(\phi)$  of the flux insertion automorphism vanishes for any half-line l, see (3.7) and (3.11). It follows that  $\omega^+ = \omega^-$  so the vectors  $\Omega^+, \Omega^-$  in the GNS representation must be proportional to each other. They can therefore not form a Kramers pair and  $T^2\Omega^+ = \Omega^+$ . This means by definition that  $\operatorname{Ind}_2(\omega_0) = 1$ .  $\square$ 

**Lemma 4.10.** Let  $\omega$  be a symmetric stably SRE state and suppose  $\omega'$  and  $\omega''$  are symmetric product states such that  $\omega \hat{\otimes} \omega'$  and  $\omega \hat{\otimes} \omega''$  are symmetric SRE states. Then

$$\operatorname{Ind}_{2}(\omega \hat{\otimes} \omega') = \operatorname{Ind}_{2}(\omega \hat{\otimes} \omega''). \tag{4.25}$$

*Proof.* Since the index of a symmetric product state is trivial (Lemma 4.9) and the index is multiplicative under stacking (Proposition 4.6) we have

$$\operatorname{Ind}_{2}(\omega \hat{\otimes} \omega') = \operatorname{Ind}_{2}(\omega \hat{\otimes} \omega' \hat{\otimes} \omega'') = \operatorname{Ind}_{2}(\omega \hat{\otimes} \omega'' \hat{\otimes} \omega') = \operatorname{Ind}_{2}(\omega \hat{\otimes} \omega'')$$
(4.26)

where we used Proposition 4.8 and the fact that  $\omega' \, \hat{\otimes} \, \omega''$  and  $\omega'' \, \hat{\otimes} \, \omega'$ , being symmetric product states on isomorphic fermion systems with time reversal, are in the same SPT phase and therefore related by a symmetry preserving LGA.

We can now make the following extension of the index to symmetric stably SRE states:

**Definition 4.11.** Let  $\omega$  be a symmetric stably SRE state, and let  $\omega'$  be any symmetric product state such that  $\omega \hat{\otimes} \omega'$  is symmetric SRE. We define

$$\operatorname{Ind}_{2}(\omega) := \operatorname{Ind}_{2}(\omega \,\hat{\otimes} \, \omega'). \tag{4.27}$$

By Lemma 4.10, this definition is independent of the choice of  $\omega'$ .

### 5 Free fermion examples

In this section, we establish that the quasi-free spin Hall states  $\omega_{SH}$  described in Example 2.5.2 are stably SRE and have a non-trivial value of the index, as claimed in Theorem 2.5.

### 5.1 Proof that spin Hall states are stably SRE

We will first prove a more general fact and then specialise to the setting of Lemma 2.3.

For any Hilbert space V we denote by  $\operatorname{Proj}_r(V)$  the space of rank r orthogonal projections in  $\mathcal{B}(V)$ . Recall that any continuous map  $\hat{\mathscr{P}}: \mathbb{T} \to \operatorname{Proj}_r(V)$  yields a rank r complex vector bundle  $E_{\hat{\mathscr{P}}} = \{(k,v) \in \mathbb{T} \times V : v \in \operatorname{Ran} \hat{\mathscr{P}}(k)\}$  over  $\mathbb{T}$  which is a subbundle of the trivial bundle  $\mathbb{T} \times V$ . We define the Chern number  $\operatorname{Ch}(\hat{\mathscr{P}})$  of the map  $\hat{\mathscr{P}}$  to be the Chern number of this vector bundle.

**Lemma 5.1.** Let  $0 \le r \le n$  be integers and let  $\hat{\mathscr{P}}: \mathbb{T} \to \operatorname{Proj}_r(\mathbb{C}^n)$  be a smooth map. If  $\operatorname{Ch}(\hat{\mathscr{P}}) = 0$  and  $n \ge 7r$  then there is a smooth homotopy  $\hat{\mathscr{P}}: [0,1] \times \mathbb{T} \to \operatorname{Proj}_r(\mathbb{C}^n): (t,k) \mapsto \hat{\mathscr{P}}_t(k)$  such that  $\hat{\mathscr{P}}_0 = \hat{\mathscr{P}}$  and  $\hat{\mathscr{P}}_1$  is constant.

*Proof.* There is a one-to-one correspondence between elements of  $\operatorname{Proj}_r(\mathbb{C}^n)$  and elements of the Grassmannian  $\operatorname{Gr}_r(\mathbb{C}^n)$ , which consists of the r-dimensional subspaces of  $\mathbb{C}^n$ . So the map  $\widehat{\mathscr{P}}$  can equivalently be regarded as a smooth map from  $\mathbb{T}$  to  $\operatorname{Gr}_r(\mathbb{C}^n)$ , and we shall use this identification throughout the proof.

Since isomorphism classes of complex vector bundles over the torus are completely classified by the Chern number, our assumption  $\operatorname{Ch}(\hat{\mathscr{P}})=0$  implies that the bundle  $E_{\hat{\mathscr{P}}}$  is isomorphic to a trivial bundle. It now follows from [13, Theorem 23.10], and noting that the torus has a good open cover of 7 sets ([36, Theorem 5.3]), that there is a homotopy  $\hat{\mathscr{P}}:[0,1]\times\mathbb{T}\to\operatorname{Gr}_r(\mathbb{C}^n)$  such that  $\hat{\mathscr{P}}_0=\hat{P}$  and  $\hat{\mathscr{P}}_1$  is constant. By [41, Theorem 10.22] we can moreover take this homotopy to be smooth.

Any smooth matrix-valued map  $\hat{\mathscr{A}}: \mathbb{T} \to \mathcal{B}(\mathbb{C}^n)$  determines a translation invariant operator  $\mathscr{A}$  on  $l^2(\mathbb{Z}^2; \mathbb{C}^n)$  by inverse Fourier transform through

$$(\mathscr{A}f)(x) = \frac{1}{4\pi^2} \int_{\mathbb{T}} dk \, e^{ik \cdot x} \hat{\mathscr{A}}(k) \hat{f}(k)$$
 (5.1)

where  $\hat{f}(k) = \sum_{x \in \mathbb{Z}^2} e^{-ik \cdot x} f(x) \in \mathbb{C}^n$  for any  $f \in l^2(\mathbb{Z}^2; \mathbb{C}^n)$ . The smoothness of  $\hat{\mathscr{A}}$  implies moreover that  $\mathscr{A}$  has super-polynomial off-diagonal decay in the sense that there is a  $g_A \in \mathcal{F}$  such that

$$|\langle \delta_x \otimes u, \mathscr{A} \delta_y \otimes v \rangle| \leqslant g_A(\operatorname{dist}(x, y)) \tag{5.2}$$

where  $\{\delta_x\}_{x\in\mathbb{Z}^2}$  is the position basis of  $l^2(\mathbb{Z}^2)$  and  $u,v\in\mathbb{C}^n$  are arbitrary. A fortiori, a smooth projection valued map  $\hat{\mathscr{P}}:\mathbb{T}\to\operatorname{Proj}_r(\mathbb{C}^n)$  determines a translation invariant projection  $\mathscr{P}$  on  $l^2(\mathbb{Z}^2;\mathbb{C}^n)$ .

### Lemma 5.2. Let

$$\hat{\mathscr{P}}: [0,1] \times \mathbb{T} \to \operatorname{Proj}_r(\mathbb{C}^n): (t,k) \mapsto \hat{\mathscr{P}}_t(k)$$
(5.3)

be a smooth homotopy and denote by  $\mathscr{P}_t$  the corresponding projections on  $l^2(\mathbb{Z}^2;\mathbb{C}^n)$ . Then there is a one-parameter norm-continuous family of unitaries  $[0,1] \mapsto \mathscr{V}_t$  on  $l^2(\mathbb{Z}^2;\mathbb{C}^n)$  and a family of bounded self-adjoint operators  $[0,1] \mapsto \mathscr{G}_t$  on  $l^2(\mathbb{Z}^2;\mathbb{C}^n)$  such that

$$i\frac{d\mathcal{V}_t}{dt} = \mathcal{G}_t\mathcal{V}_t, \qquad \mathcal{V}_0 = 1, \qquad and \quad \mathcal{P}_t = \mathcal{V}_t\mathcal{P}_0\mathcal{V}_t^*.$$
 (5.4)

The generators  $\mathcal{G}_t$  have super-polynomial off-diagonal decay in the sense of (5.2), where  $g \in \mathcal{F}$  is independent of t.

*Proof.* Let

$$\hat{\mathscr{G}}_t(k) = i \left[ \frac{\partial}{\partial t} \hat{\mathscr{P}}_t(k), \hat{\mathscr{P}}_t(k) \right]$$
(5.5)

be the well-known self-adjoint Kato generator [37] and let  $\hat{\mathscr{V}}_t(k)$  be the unique family of unitaries that solve

$$i\frac{\partial}{\partial t}\hat{\mathscr{V}}_t(k) = \hat{\mathscr{G}}_t(k)\hat{\mathscr{V}}_t(k), \qquad \hat{\mathscr{V}}_0(k) = 1.$$
(5.6)

Then

$$\hat{\mathscr{P}}_t(k) = \hat{\mathscr{V}}_t(k)\hat{\mathscr{P}}_0(k)\hat{\mathscr{V}}_t(k)^*. \tag{5.7}$$

for all  $t \in [0,1]$  and all  $k \in \mathbb{T}$ . Since the family of projections is smooth, so is the map  $(t,k) \mapsto \hat{\mathcal{G}}_t(k)$ . All the claims of the lemma now follow by inverse Fourier transform.

Finally we show that the data provided by the previous lemma yields an equivalence of symmetric quasi-free states.

**Lemma 5.3.** Let  $[0,1] \ni t \mapsto \mathscr{G}_t$  be a norm-continuous family of self-adjoint operators on  $l^2(\mathbb{Z}^2; \mathbb{C}^n)$  such that (5.2) holds uniformly in t. Let  $\mathscr{V}_t$ ,  $\mathscr{P}_t$  be the corresponding one-parameter family of unitaries given by (5.4). Let  $\mathcal{A}$  be the CAR algebra over  $l^2(\mathbb{Z}^2; \mathbb{C}^n)$  and let  $\omega_t = \omega_{\mathscr{P}_t}$  be the gauge-invariant quasi-free state on  $\mathcal{A}$  corresponding to the projection  $\mathscr{P}_t$ , see (2.11). Then there is a U(1) symmetry preserving TDI F such that  $\omega_t = \omega_0 \circ \alpha_t^F$  for all  $t \in [0,1]$ .

*Proof.* Let  $\{\delta_x\}_{x\in\mathbb{Z}^2}$  be the position basis of  $l^2(\mathbb{Z}^2)$  and let  $\{e_i\}_{i=1}^n$  be an orthonormal basis of  $\mathbb{C}^n$ . Let us write

$$\mathscr{G}_t(x,i;y,j) := \langle \delta_x \otimes e_i, \mathscr{G}_t \, \delta_y \otimes e_j \rangle \tag{5.8}$$

for the matrix elements of  $\mathscr{G}_t$  with respect to the orthonormal basis  $\{\delta_x \otimes e_i\}$ . We take

$$F_x(t) = \frac{1}{2} \sum_{y \in \mathbb{Z}^2} \sum_{i,j=1}^n \left( \mathcal{G}_t(x,i;y,j) a_{x,i}^* a_{y,j} + \overline{\mathcal{G}_t(x,i;y,j)} a_{y,j}^* a_{x,i} \right)$$
(5.9)

for all  $x \in \mathbb{Z}^2$ . These are homogeneous self-adjoint elements of  $\mathcal{A}$ , and they define a TDI by the super-polynomial decay of the  $\mathcal{G}_t(x, i; y, j)$ . A short calculation yields

$$\sum_{x \in \mathbb{Z}^2} i \left[ F_x(t), a(f) \right] = a(i\mathcal{G}_t f)$$
(5.10)

which implies that

$$\alpha_t^F(a(f)) = a(\mathcal{V}_t^* f). \tag{5.11}$$

for any  $f \in l^2(\mathbb{Z}^2; \mathbb{C}^n)$ .

With this, we check that

$$(\omega_0 \circ \alpha_t^F) (a^*(f)a(g)) = \omega_0 (a^*(\mathcal{V}_t^* f)a(\mathcal{V}_t^* g)) = \langle \mathcal{V}_t^* g, \mathcal{P}_0 \mathcal{V}_t^* f \rangle = \langle g, \mathcal{P}_t f \rangle = \omega_t (a^*(g)a(f)).$$

$$(5.12)$$

Since the  $\omega_t$  are quasi-free we conclude that  $\omega_t = \omega_0 \circ \alpha_t^F$ , as required.

We are now equipped to prove Lemma 2.3. We briefly recall the setting described in Section 2.5.2. The one particle Hilbert space is  $\mathcal{K}_2 = L^2(\mathbb{T}; \mathbb{C}^2) \oplus L^2(\mathbb{T}; \mathbb{C}^2)$  and the state  $\omega_{SH}$  corresponds to the projection  $\mathscr{P}_{SH} = \mathscr{P}^{\uparrow} \oplus \mathscr{P}^{\downarrow}$  whose Fourier transform is a smooth projection valued map of the form

$$\hat{\mathscr{P}}_{\mathrm{SH}}(k) = \hat{\mathscr{P}}^{\uparrow}(k) \oplus \hat{\mathscr{P}}^{\downarrow}(k) \tag{5.13}$$

Here  $\hat{\mathscr{P}}^{\downarrow}(k)$  is the complex conjugate of  $\hat{\mathscr{P}}^{\uparrow}(k)$ . It follows that the projection valued maps  $\hat{\mathscr{P}}^{\uparrow}$  and  $\hat{\mathscr{P}}^{\downarrow}$  have opposite Chern numbers, and since the Chern number is additive under direct sums of bundles, this implies  $\operatorname{Ch}(\hat{\mathscr{P}}_{\operatorname{SH}}) = \operatorname{Ch}(\hat{\mathscr{P}}^{\uparrow}) + \operatorname{Ch}(\hat{\mathscr{P}}^{\downarrow}) = 0$ .

Proof of Lemma 2.3. In order to apply Lemma 5.1, we expand the 'ambient dimension' from 4 to 14 by considering the algebra  $\mathcal{A}'$  over  $l^2(\mathbb{Z}^2; \mathbb{C}^4) \oplus l^2(\mathbb{Z}^2; \mathbb{C}^{10})$  and the symmetric state  $\omega' = \omega_{\text{SH}} \, \hat{\otimes} \, \omega_{\text{empty}}$  corresponding to the projection  $\mathscr{P} = \mathscr{P}_{\text{SH}} \oplus 0$  and consequently  $\hat{\mathscr{P}} = \hat{\mathscr{P}}_{\text{SH}} \oplus 0$ . The projection-valued map  $\hat{\mathscr{P}} = \hat{\mathscr{P}}_{\text{SH}} \oplus 0$  takes values in  $\text{Proj}_2(\mathbb{C}^{14})$  and has  $\text{Ch}(\hat{\mathscr{P}}) = 0$ . Now Lemma 5.1 provides a smooth homotopy  $(t, k) \mapsto \hat{\mathscr{P}}_t(k) \in \text{Proj}_2(\mathbb{C}^{12})$  interpolating between  $\hat{\mathscr{P}}$  and a constant projection.

Let  $\mathscr{P}_t$  be the inverse Fourier transform of  $\hat{\mathscr{P}}_t$ . We then apply Lemma 5.2 to obtain unitaries  $\mathscr{V}_t$  and generators  $\mathscr{G}_t$  satisfying the assumption of Lemma 5.3. The latter lemma now provides a U(1)-invariant TDI F such that  $\omega'_t = \omega_0 \circ \alpha_t^F$  where  $\omega'_t = \omega'_{\mathscr{P}_t}$  is the quasi-free state on  $\mathscr{A}'$  determined by the projection  $\mathscr{P}_t$ . This family of states interpolates between  $\omega_{\mathrm{SH}} \hat{\otimes} \omega_{\mathrm{empty}}$  and  $\omega_{\mathscr{P}_1 \oplus 0}$ . Since  $\hat{\mathscr{P}}_1$  is constant, its Fourier transform  $\mathscr{P}_1$  is strictly local and  $\omega_1 = \omega_{\mathscr{P}_1 \oplus 0}$  is a product state. We conclude that  $\omega_{\mathrm{SH}} \hat{\otimes} \omega_{\mathrm{empty}}$  is LGA equivalent to a product state. Since  $\omega_{\mathrm{empty}}$  is also a product state, we conclude that  $\omega_{\mathrm{SH}}$  is stably SRE, see Section 2.4.

### 5.2 Index of spin Hall states

Let  $\omega_{\mathrm{SH}}$  be a spin Hall state on the CAR algebra  $\mathcal{A}$  over  $l^2(\mathbb{Z}^2; \mathbb{C}^4)$ . We have just proved that  $\omega_{\mathrm{SH}}$  is stably SRE; Concretely,  $\omega = \omega_{\mathrm{SH}} \, \hat{\otimes} \, \omega_{\mathrm{empty}}$  is a symmetric SRE state and it corresponds to the Fermi projection  $\mathscr{P} = \mathscr{P}_{\mathrm{SH}} \oplus 0$  on  $l^2(\mathbb{Z}^2; \mathbb{C}^4) \oplus l^2(\mathbb{Z}^2; \mathbb{C}^{10})$  where  $\mathscr{P}_{\mathrm{SH}} = \mathscr{P}^{\uparrow} \oplus \mathscr{P}^{\downarrow}$  is the Fermi projection of the spin Hall state.

### 5.2.1 Flux insertion for spin Hall states in the free fermion formalism

The charge operator in a region  $\Lambda$  is given by  $\Pi_{\Lambda} = (\chi_{\Lambda} \otimes \mathbb{1}) \oplus (\chi_{\Lambda} \otimes \mathbb{1})$ , where  $\chi_{\Lambda}$  is the multiplication operator by the characteristic function of the set  $\Lambda$ . Let l = (0, (-1, 0)) be the left horizontal axis so  $h = h_l$  is the upper half plane. The analog of the many-body U(1)-transformation of the upper half-plane h is now

$$\mathscr{P}_{\phi}^{h} := e^{i\phi\Pi_{h}} \mathscr{P} e^{-i\phi\Pi_{h}}. \tag{5.14}$$

Since  $\mathscr{P}^h_\phi(\partial_\phi\mathscr{P}^h_\phi)\mathscr{P}^h_\phi=(\mathscr{P}^h_\phi)^\perp(\partial_\phi\mathscr{P}^h_\phi)(\mathscr{P}^h_\phi)^\perp=0$  we have

$$\left[ \left[ \partial_{\phi} \mathcal{P}_{\phi}^{h}, \mathcal{P}_{\phi}^{h} \right], \mathcal{P}_{\phi}^{h} \right] = \partial_{\phi} \mathcal{P}_{\phi}^{h}, \tag{5.15}$$

and therefore the flow  $\phi \mapsto \mathscr{P}^h_{\phi}$  is also generated by

$$\mathcal{K}_{\phi}^{h} := -\mathrm{i}[\partial_{\phi}\mathcal{P}_{\phi}^{h}, \mathcal{P}_{\phi}^{h}]. \tag{5.16}$$

If we let  $\mathscr{U}_{\phi}^{h}$  be the unitary solution to

$$\mathscr{U}_0^h = 1, \qquad \partial_\phi \mathscr{U}_\phi^h = i\mathscr{K}_\phi^h \mathscr{U}_\phi^h, \tag{5.17}$$

then  $\mathscr{P}_{\phi}^{h} = \mathscr{U}_{\phi}^{h} \mathscr{P}(\mathscr{U}_{\phi}^{h})^{*}$ . Let now  $\Pi_{L}$  be the projection on the left half-plane. Adiabatic flux insertion in the left horizontal half-line gauge is generated by the restriction

$$\mathscr{K}_{\phi} := \Pi_L \mathscr{K}_{\phi}^h \Pi_L, \tag{5.18}$$

which we call the Kato generator for flux insertion. Namely, if  $\mathscr{U}_{\phi}$  is the solution to

$$\mathcal{U}_0 = 1, \qquad i \frac{\mathrm{d}\mathcal{U}_\phi}{\mathrm{d}\phi} = \mathcal{K}_\phi \mathcal{U}_\phi.$$
 (5.19)

then the projection

$$\mathscr{P}_{\phi} := \mathscr{U}_{\phi} \mathscr{P} \mathscr{U}_{\phi}^{*} \tag{5.20}$$

corresponds to a state obtained from  $\mathscr{P}$  by inserting a  $\phi$  flux at the origin.

In particular, the Fermi projections obtained from  $\mathscr P$  by inserting  $\pm \pi$  flux in the half-line gauge are given by

$$\mathscr{P}_{\pm} = \mathscr{P}_{+}^{\uparrow} \oplus \mathscr{P}_{+}^{\downarrow} \oplus 0. \tag{5.21}$$

Since

$$\mathcal{T}\mathcal{P}_{-}\mathcal{T}^* = \mathcal{P}_{+},\tag{5.22}$$

we have that

$$\mathcal{K}\mathcal{P}_{-}^{\downarrow} = \mathcal{P}_{+}^{\downarrow}\mathcal{K}, \qquad \mathcal{K}\mathcal{P}_{-}^{\uparrow} = \mathcal{P}_{+}^{\downarrow}\mathcal{K}, \tag{5.23}$$

where  $\mathcal{K}$  is the complex conjugation appearing in the definition of  $\mathcal{T}$ .

It is shown in Appendix B, Proposition B.6 that  $\mathscr{P}_+^{\uparrow} - \mathscr{P}_-^{\uparrow} \in \mathcal{J}_1$  is trace class and that the Chern number of  $\mathscr{P}_+^{\uparrow}$  is equal to the index of the pair of projections  $\mathscr{P}_+^{\uparrow}$  and  $\mathscr{P}_-^{\uparrow}$ :

$$Ch(\mathscr{P}^{\uparrow}) = ind(\mathscr{P}^{\uparrow}_{+}, \mathscr{P}^{\uparrow}_{-}). \tag{5.24}$$

We recall that the index of a pair of projection  $\mathscr{P}_1, \mathscr{P}_2$  acting on the same Hilbert space and such that  $\mathscr{P}_1 - \mathscr{P}_2$  is compact is given by

$$\operatorname{ind}(\mathscr{P}_1, \mathscr{P}_2) = \operatorname{dim} \operatorname{Ker}(\mathscr{P}_1 - \mathscr{P}_2 - 1) - \operatorname{dim} \operatorname{Ker}(\mathscr{P}_1 - \mathscr{P}_2 + 1), \tag{5.25}$$

see [4].

Let  $\mathscr{N}_{\pm}$  be the orthogonal projections onto  $\operatorname{Ker}(\mathscr{P}_{+}^{\uparrow} - \mathscr{P}_{-}^{\uparrow} \mp \mathbb{1})$  and let  $n_{\pm} = \dim \mathscr{N}_{\pm}$  be the dimensions of these kernels. With this, (5.24) reads  $n_{+} - n_{-} = \operatorname{Ch}(\mathscr{P}^{\uparrow})$ . We define

$$\mathcal{Q}_{+}^{\uparrow} := \mathcal{P}_{+}^{\uparrow} - \mathcal{N}_{+} \tag{5.26}$$

and note that

**Lemma 5.4.** The operators  $\mathcal{Q}_{\pm}^{\uparrow}$  are projections and  $\mathcal{Q}_{+}^{\uparrow} - \mathcal{Q}_{-}^{\uparrow} \in \mathcal{J}_{1}$ . Moreover,  $\operatorname{ind}(\mathcal{Q}_{+}^{\uparrow}, \mathcal{Q}_{-}^{\uparrow}) = 0$ .

Proof. We first show that the projection  $\mathcal{N}_{+}$  is a subprojection of  $\mathscr{P}_{+}^{\uparrow}$ , and similarly for  $\mathscr{N}_{-}$  and  $\mathscr{P}_{-}^{\uparrow}$ . i.e. we want to show that  $\mathscr{N}_{\pm}\mathscr{P}_{\pm}^{\uparrow}\mathscr{N}_{\pm} = \mathscr{N}_{\pm}$ . First,  $0 \leq \mathscr{N}_{\pm}\mathscr{P}_{\pm}^{\uparrow}\mathscr{N}_{\pm} \leq \mathscr{N}_{\pm}$  follows from  $0 \leq \mathscr{P}_{\pm}^{\uparrow} \leq \mathbb{1}$  and the fact that  $\mathscr{N}_{\pm}$  is self-adjoint. Second, the definition of  $\mathscr{N}_{\pm}$  is equivalent to  $\mathscr{N}_{+}(\mathscr{P}_{+}^{\uparrow} - \mathscr{P}_{-}^{\uparrow})\mathscr{N}_{+} = \pm \mathscr{N}_{\pm}$ , which implies that

$$\mathcal{N}_{\pm} \mathcal{P}_{+}^{\uparrow} \mathcal{N}_{\pm} = \mathcal{N}_{\pm} (\mathbb{1} + \mathcal{P}_{\mp}^{\uparrow}) \mathcal{N}_{\pm} \geqslant \mathcal{N}_{\pm}. \tag{5.27}$$

We conclude that  $\mathscr{N}_{\pm}\mathscr{P}_{\pm} = \mathscr{P}_{\pm}\mathscr{N}_{\pm} = \mathscr{N}_{\pm}$ . It follow immediately that  $\mathscr{Q}_{\pm}^{\uparrow} = \mathscr{P}_{\pm}^{\uparrow} - \mathscr{N}_{\pm}$  is a projection. Since  $\mathscr{N}_{\pm}$  are finite range projections the claim that  $\mathscr{Q}_{+}^{\uparrow} - \mathscr{Q}_{-}^{\uparrow}$  is trace class follows immediately from the fact that  $\mathscr{P}_{+}^{\uparrow} - \mathscr{P}_{-}^{\uparrow}$  is trace class (Proposition B.6). It follows that  $\operatorname{ind}(\mathscr{Q}_{+}^{\uparrow}, \mathscr{Q}_{-}^{\uparrow})$  is well-defined and we compute it as follows:

$$\operatorname{ind}(\mathcal{Q}_{\pm}^{\uparrow}, \mathcal{P}_{\pm}^{\uparrow}) = \operatorname{dim}\operatorname{Ker}(-\mathcal{N}_{\pm} - \mathbb{1}) - \operatorname{dim}\operatorname{Ker}(-\mathcal{N}_{\pm} + \mathbb{1}) = 0 - \operatorname{dim}\mathcal{N}_{\pm}. \tag{5.28}$$

Hence, by the additivity of the index of a pair of projections [4, Theorem 3.4],

$$\operatorname{ind}(\mathcal{Q}_{+}^{\uparrow}, \mathcal{Q}_{-}^{\uparrow}) = \operatorname{ind}(\mathcal{Q}_{+}^{\uparrow}, \mathcal{P}_{+}^{\uparrow}) + \operatorname{ind}(\mathcal{P}_{+}^{\uparrow}, \mathcal{P}_{-}^{\uparrow}) - \operatorname{ind}(\mathcal{Q}_{-}^{\uparrow}, \mathcal{P}_{-}^{\uparrow})$$

$$(5.29)$$

$$= -n_{+} + \operatorname{Ch}(\mathscr{P}^{\uparrow}) + n_{-} = 0 \tag{5.30}$$

where we used  $\operatorname{ind}(\mathscr{P}_{+}^{\uparrow}, \mathscr{P}_{-}^{\uparrow}) = \operatorname{Ch}(\mathscr{P}^{\uparrow}) = n_{+} - n_{-}.$ 

note that  $\mathscr{K}$  leaves  $\mathcal{H}^{\uparrow}$  and  $\mathcal{H}^{\downarrow}$  invariant, and we identify these two spaces in this equality by  $|x,\uparrow\rangle \leftrightarrow |x,\downarrow\rangle$ 

Since  $\mathcal{Q}_{+}^{\uparrow} - \mathcal{Q}_{-}^{\uparrow} \in \mathcal{J}_{1}$  we can apply to the following Proposition:

**Proposition 5.5.** Let  $\mathscr{P}, \mathscr{Q}$  be projections acting on a Hilbert space  $\mathcal{H}$  with  $\mathscr{P} - \mathscr{Q} \in \mathcal{J}_1$  and  $\operatorname{ind}(\mathscr{P}, \mathscr{Q}) = 0$ . Then there exists a unitary  $\mathscr{V}$  such that  $\mathscr{P} = \mathscr{V} \mathscr{Q} \mathscr{V}^*$  and  $\mathscr{V} - \mathbb{1} \in \mathcal{J}_1$ .

The proof can be found in Section VII B of [15] (without the trace class statement) or in Section 4.2.1 of [11] (including the trace class statement).

From Lemma 5.4 and Proposition 5.5 we get a unitary  $\mathscr V$  such that

$$\mathcal{Q}_{+}^{\uparrow} = \mathcal{V}\mathcal{Q}_{-}^{\uparrow}\mathcal{V}^{*}, \qquad \mathcal{V} - \mathbb{1} \in \mathcal{J}_{1}. \tag{5.31}$$

Using this and (5.23) we get for  $\mathscr{P}_+ = \mathscr{P}_+^{\uparrow} \oplus \mathscr{P}_+^{\downarrow} \oplus 0$  that

$$\mathscr{P}_{+} = (\mathscr{Q}_{+}^{\uparrow} + \mathscr{N}_{+}) \oplus (\mathscr{K} \mathscr{P}_{-}^{\uparrow} \mathscr{K}) \oplus 0 \tag{5.32}$$

$$= (\mathcal{V}\mathcal{Q}_{-}^{\uparrow}\mathcal{V}^{*} + \mathcal{N}_{+}) \oplus \mathcal{K}(\mathcal{Q}_{-}^{\uparrow} + \mathcal{N}_{-})\mathcal{K} \oplus 0$$

$$(5.33)$$

$$= (\mathcal{V} \oplus \mathbb{1} \oplus \mathbb{1})(\mathcal{Q}_{-}^{\uparrow} \oplus \mathcal{K} \mathcal{Q}_{-}^{\uparrow} \mathcal{K} \oplus 0)(\mathcal{V} \oplus \mathbb{1} \oplus \mathbb{1})^{*} + (\mathcal{N}_{+} \oplus (\mathcal{K} \mathcal{N}_{-} \mathcal{K}) \oplus 0). \tag{5.34}$$

The projections  $\mathscr{P}_+$  and  $(\mathscr{V} \oplus \mathbb{1} \oplus \mathbb{1})(\mathscr{Q}_-^{\uparrow} \oplus \mathscr{K} \mathscr{Q}_-^{\uparrow} \mathscr{K} \oplus 0)(\mathscr{V} \oplus \mathbb{1} \oplus \mathbb{1})^*$  differ by a finite rank operator and hence the quasi-free states associated with  $\mathscr{P}_+$  and  $\mathscr{Q}_-^{\uparrow} \oplus \mathscr{K} \mathscr{Q}_-^{\uparrow} \mathscr{K} \oplus 0$  are unitarily equivalent. Moreover,

$$[(\mathscr{V} \oplus \mathbb{1} \oplus \mathbb{1}), (\mathscr{Q}_{-}^{\uparrow} \oplus \mathscr{K} \mathscr{Q}_{-}^{\uparrow} \mathscr{K} \oplus 0)] = [\mathscr{V}, \mathscr{Q}_{-}^{\uparrow}] \in \mathcal{J}_{1}$$

$$(5.35)$$

because  $\mathscr{V} - \mathbb{1} \in \mathcal{J}_1$ . The Shale-Stinespring criterion [50], see also [20, Theorem 6.16], implies that the Bogoliubov transformation  $a(f) \mapsto a((\mathscr{V} \oplus \mathbb{1} \oplus \mathbb{1})f)$  is unitarily implementable in the GNS representation of the quasi-free state associated with the projection  $\mathscr{Q}_{-}^{\uparrow} \oplus \mathscr{K} \mathscr{Q}_{-}^{\uparrow} \mathscr{K} \oplus 0$ . We denote the corresponding unitary by  $\Gamma(\mathscr{V} \oplus \mathbb{1} \oplus \mathbb{1})$ .

To the projection  $\mathscr{R} = \mathscr{Q}_{-}^{\uparrow} \oplus \mathscr{K} \mathscr{Q}_{-}^{\uparrow} \mathscr{K} \oplus 0$  corresponds a quasi-free state  $\omega_{\mathscr{R}}$  whose GNS triple we denote by  $(\Pi, \mathcal{H}, |\Phi\rangle)$ . One then checks using the definition (2.10) of  $\mathscr{T}$  that  $\mathscr{T}(\mathscr{Q}_{-}^{\uparrow} \oplus \mathscr{K} \mathscr{Q}_{-}^{\uparrow} \mathscr{K} \oplus 0) \mathscr{T}^{*} = \mathscr{Q}_{-}^{\uparrow} \oplus \mathscr{K} \mathscr{Q}_{-}^{\uparrow} \mathscr{K} \oplus 0$ , so that  $\omega_{\mathscr{R}}$  is time reversal invariant. It follows that the time reversal  $\tau$  induced by the single particle time reversal  $\mathscr{T}$  is implemented in the GNS representation by an anti-unitary operator T which we may choose such that

$$T|\Phi\rangle = |\Phi\rangle \tag{5.36}$$

and therefore

$$T^{2}|\Phi\rangle = |\Phi\rangle. \tag{5.37}$$

We see that  $|\Phi\rangle$  is a Kramers singlet. By the remarks above, the quasi-free state corresponding to  $\mathscr{P}_+$  is represented by a vector  $|\Psi_+\rangle \in \mathcal{H}$  given by

$$|\Psi_{+}\rangle := A^{*}\Gamma(\mathscr{V} \oplus \mathbb{1} \oplus \mathbb{1})|\Phi\rangle \tag{5.38}$$

where

$$A^* = \Pi \Big( \prod_{i=1}^{n_+} a_{\uparrow}^*(v_i) \prod_{j=1}^{n_-} a_{\downarrow}^*(\mathcal{K}w_i) \Big).$$
 (5.39)

Here  $\{v_i\}_{i=1}^{n_+}$  and  $\{w_i\}_{i=1}^{n_-}$  are orthonormal bases of Ran $\mathcal{N}_+$  and Ran $\mathcal{N}_-$  respectively. Being a product of creation operators,  $A^*$  is homogeneous with

$$T^2 A^* T^2 = (-1)^{n_+ + n_-} A^* (5.40)$$

since  $T^2$  implements fermion parity. Since  $(-1)^{n_++n_-}=(-1)^{n_+-n_-}$  and  $n_+-n_-$  equals the Chern number of  $\mathscr{P}^{\uparrow}$  we conclude that

$$T^{2}|\Psi_{+}\rangle = (-1)^{\operatorname{Ch}(\mathscr{P}^{\uparrow})}A^{*}T^{2}\Gamma(\mathscr{V} \oplus \mathbb{1} \oplus \mathbb{1})|\Phi\rangle = (-1)^{\operatorname{Ch}(\mathscr{P}^{\uparrow})}|\Psi_{+}\rangle$$
 (5.41)

where we used that  $\Gamma(\mathcal{V} \oplus \mathbb{1} \oplus \mathbb{1})|\Phi\rangle$  is even under  $T^2$  because  $\Gamma(\mathcal{V} \oplus \mathbb{1} \oplus \mathbb{1})$  is an even unitary and  $|\Phi\rangle$  is a Kramers singlet, see [44] and (5.37). This shows that the quasi-free state  $\omega_{\mathscr{P}_+}$  corresponding to  $\mathscr{P}_+$  is a Kramers singlet if  $Ch(\mathscr{P}^{\uparrow})$  is even, while it belongs to a Kramers pair if  $Ch(\mathscr{P}^{\uparrow})$  is odd.

### 5.2.2 Non-triviality of the index for spin Hall states

The state  $\psi^+ := \omega_{\mathscr{P}_+}$  was obtained from  $\omega = \omega_{SH} \hat{\otimes} \omega_{empty}$  by inserting a  $\pi$  flux defect using a Kato generator (5.18) in the single particle Hilbert space. Since we defined our index using defect states obtained through a quasi-adiabatic evolution rather than Kato's parallel transport, it remains to show that both flux insertion protocols lead to the same index. The argument follows the strategy used in the proof of Proposition 3.8 to show that  $\omega^+$  and  $\psi^+$  have the same parity.

In order to compare the defect state  $\psi^+$  obtained using the Kato generator with a defect state  $\omega^+$  obtained by quasi-adiabatic evolution we first give a many-body description of the Kato generator (5.18). It is shown in the Appendix, Lemma B.1, that  $\mathscr{K}_{\phi}^h$  is exponentially local, and exponentially localised near the horizontal axis. i.e. writing

$$\mathscr{K}_{\phi}^{h}(x,i;y,j) := \langle \delta_{x} \otimes e_{i}, \mathscr{K}_{\phi}^{h} \delta_{y} \otimes e_{j} \rangle$$

$$(5.42)$$

we have

$$\left| \mathcal{K}_{\phi}^{h}(x,i;y,j) \right| \le c e^{-\xi(|x_1 - y_1| + |x_2| + |y_2|)}$$
 (5.43)

for some positive constants c and  $\xi$ . It follows immediately that the Kato generator  $\mathcal{K}_{\phi}$ , see (5.18), is exponentially localised near the left horizontal axis l = (0, -(1, 0)).

In particular,  $\mathcal{K}_{\phi}$  satisfies the assumptions of Lemma 5.3 so that the projections  $\mathscr{P}_{\phi}$  defined in (5.20) correspond to quasi-free pure states  $\psi^{\phi}$  such that  $\psi^{+} = \psi^{\pi}$  and there is a TDI  $K^{l}$  such that

$$\psi^{\phi} = \omega \circ \alpha_{\phi}^{K^l} \tag{5.44}$$

for all  $\phi \in [-\pi, \pi]$ . The TDI may be taken to be (see the proof of Lemma 5.3)

$$K_x^l(\phi) = \frac{1}{2} \sum_{y \in \mathbb{Z}^2} \sum_{i,j=1}^n \left( \mathscr{K}_{\phi}(x,i;y,j) a_{x,i}^* a_{y,j} + \overline{\mathscr{K}_{\phi}(x,i;y,j)} a_{y,j}^* a_{x,i} \right).$$
 (5.45)

Let H be any symmetric parent Hamiltonian for  $\omega$  (which exists by Lemma 3.3) and denote by  $\omega^{\phi} = \omega \circ \gamma_{\phi}^{l,H}$  the corresponding defect states obtained by quasi-adiabatic evolution described in Section 3.1.4.

**Lemma 5.6.** There is an  $f \in \mathcal{F}$  for each  $\phi \in [0, \pi]$  an f-localized even unitary  $V_{\phi} \in \mathcal{A}^{al}$  such that  $\omega^{\phi} = \psi^{\phi} \circ \operatorname{Ad}[V_{\phi}]$ .

*Proof.* Since the argument is essentially the same as the ones proving Lemma 3.7 and Proposition 3.8, we allow ourselves to gloss over some of the technical details.

Let l be the left horizontal axis and  $h=h_l$  be the upper half plane. The projections  $\mathscr{P}^h_{\phi}$  correspond to the states  $\omega^{h,\phi}:=\omega\circ\rho^h_{\phi}$ . By quasi-adiabatic evolution, we have that  $\omega^{h,\phi}:=\omega\circ\alpha^{h,H}_{\phi}$ , see (3.10).

Similarly to (5.45) we can introduce a TDI  $K^h$  given by

$$K_x^h(\phi) = \frac{1}{2} \sum_{y \in \mathbb{Z}^2} \sum_{i,j=1}^n \left( \mathcal{K}_{\phi}^h(x,i;y,j) a_{x,i}^* a_{y,j} + \overline{\mathcal{K}_{\phi}^h(x,i;y,j)} a_{y,j}^* a_{x,i} \right), \tag{5.46}$$

which is so that

$$\omega^{h,\phi} = \omega \circ \alpha_{\phi}^{K^h} \tag{5.47}$$

for all  $\phi \in \mathbb{R}$ .

Since  $\omega$  is SRE there is a homogeneous product state  $\omega_0$  and an LGA  $\alpha$  such that  $\omega = \omega_0 \circ \alpha$ . For all  $\phi \in \mathbb{R}$  we now put

$$\tilde{\psi}^{\phi} = \psi^{\phi} \circ (\alpha \circ \alpha_{\phi}^{K})^{-1} \tag{5.48}$$

$$\tilde{\omega}^{\phi} = \omega^{\phi} \circ (\alpha \circ \alpha_{\phi}^{K})^{-1}, \tag{5.49}$$

then

$$\tilde{\psi}^{\phi} = \omega_0, \quad \text{and} \quad \tilde{\omega}^{\phi} = \omega_0 \circ \tilde{\beta}^{\phi}$$
 (5.50)

with

$$\tilde{\beta}^{\phi} = \alpha \circ \gamma_{\phi}^{l,H} \circ (\alpha_{\phi}^{K})^{-1} \circ \alpha^{-1}. \tag{5.51}$$

Let

$$\beta^{\phi} = \alpha \circ \alpha_{\phi}^{h,H} \circ (\alpha_{\phi}^{K^h})^{-1} \circ \alpha^{-1}. \tag{5.52}$$

Arguing as in the first part of the proof of Lemma 3.7, we can find  $g \in \mathcal{F}$  uniform in  $\phi \in [0, \pi]$  such that

$$\|\tilde{\beta}^{\phi}(A) - \beta^{\phi}(A)\| \leqslant g(r)\|A\| \tag{5.53}$$

for all  $\phi \in [0, \pi]$  and all  $A \in \Lambda_{\pi,l} \cap B(r)^c$ . Since  $\tilde{\psi}^{\phi} = \omega_0$ , we have  $\tilde{\psi}^{\phi} \circ \beta^{\phi} = \tilde{\psi}^{\phi}$ . It follows that

$$\left| \tilde{\psi}^{\phi}(A) - \tilde{\omega}^{\phi}(A) \right| = \left| \omega_0 \left( \beta^{\phi}(A) - \tilde{\beta}^{\phi}(A) \right) \right| \leqslant g(r) |A| \tag{5.54}$$

for all  $A \in \Lambda_{\pi,l} \cap B(r)^c$ . Similarly we have  $\|\tilde{\beta}(A) - A\| \leq g(r)\|A\|$  for  $A \in \Lambda_{\pi,-l} \cap B(r)^c$  which implies  $\left|\tilde{\psi}^{\phi}(A) - \tilde{\omega}^{\phi}(A)\right| \leq g(r)\|A\|$  for  $A \in \Lambda_{\pi,l} \cap B(r)^c$ .

Since  $\tilde{\psi}^{\phi} = \omega_0$  is a homogeneous product state it now follows from Lemma A.1 that the states  $\tilde{\omega}^{\phi}$  are all h-close to  $\omega_0$  for  $\phi \in [0, \pi]$  for an  $h \in \mathcal{F}$  that depends only on g, and by Proposition A.2 there is an  $\tilde{f} \in \mathcal{F}$  depending only on h, and  $\tilde{f}$ -local homogeneous unitaries  $W_{\phi} \in \mathcal{A}^{\text{al}}$  such that  $\tilde{\omega}^{\phi} = \omega_0 \circ \text{Ad}[W_{\phi}]$ . It follows that  $\omega^{\phi} = \psi^{\phi} \circ \text{Ad}[V_{\phi}]$  with  $V_{\phi} = (\alpha \circ \alpha_{\phi}^K)^{-1}(W_{\phi})$ , which are uniformly f-local for some  $f \in \mathcal{F}$ .

The family of states  $\tilde{\omega}^{\phi}$  are related to the pure homogeneous product state  $\omega_0$  by conjugation with the homogeneous almost local unitaries unitaries  $W_{\phi}$  which satisfy the assumptions of Lemma A.7. Since  $\tilde{\omega}^0 = \omega_0$  we must have that  $W_0$  is even and so all the  $W_{\phi}$  are even. Since  $\alpha$  and  $\alpha_{\phi}^K$  are parity preserving, we conclude that also all the  $V_{\phi}$  are even.

**Proposition 5.7.**  $\operatorname{Ind}_2(\omega_{SH}) = (-1)^{\operatorname{Ch}(\mathscr{P}^{\uparrow})}$ .

Proof. We must show that the symmetric SRE state  $\omega = \omega_{SH} \hat{\otimes} \omega_{empty}$  has non-trivial index. We have shown above that the defect state  $\psi^+$  obtained by the Kato generator is either a Kramers singlet or belongs to a Kramers pair. Now, Lemma 5.6 yields in particular an even unitary  $V \in \mathcal{A}^{al}$  such that  $\psi^+ = \omega^+ \circ \operatorname{Ad}[V]$ . By Lemma 4.1 we conclude that  $\omega^+$  is itself either a Kramers singlet or belongs to a Kramers pair. The claim follows by Definitions 3.15 and 4.11.

### 6 Proof of the main theorem

We collect the results obtained in the previous sections to give a proof of Theorem 2.5.

*Proof.* **Proof of (1).** Triviality of the index for symmetric product states is Lemma 4.9. The claim about the spin Hall states is Proposition 5.7.

**Proof of (2).** Multiplicativity of the index under stacking is the statement of Proposition 4.6.

**Proof of (3).** By items (1) and (2), the index is invariant under stacking with a product state. With this, the claim (3) follows from the invariance under symmetry preserving LGAs, Proposition 4.8.

Acknowledgements. The work of S.B. and M.R. is supported by NSERC of Canada.

### A Almost local transitivity

The classical Kadison transitivity theorem implies that two GNS equivalent representations associated with two pure states are necessarily unitarily equivalent through a unitary element of the algebra, see e.g. [18]. Our goal in this section is two-fold. First to prove that if  $\psi_0$  is a product state and  $\psi_0$ ,  $\psi$  are almost local perturbations of each other when restricted to a cone and its complement, then  $\psi_0$ ,  $\psi$  are almost local perturbations of each other, see Definition 3.6. Second of all to show that if they are moreover pure, then they are unitarily equivalent through a unitary in the almost local algebra.

### A.1 Lifting closeness at infinity on cones to closeness at infinity

We start with the first goal, a version of which can also be found in Appendix C of [33]. It is a rather classical argument using relative entropy, except for the fact that we need Araki's extension [2] for the subadditivity of the quantum entropy. Recall the definition of a cone given in Section 3.1.5, and Definition 3.6 for two states to be almost local perturbations of each other.

**Lemma A.1.** Let  $\Lambda$  be a cone. Let  $f \in \mathcal{F}$  and suppose  $\psi_0$  is a homogeneous product state and  $\psi$  a homogeneous state on  $\mathcal{A}$  such that

$$|\psi_0(A) - \psi(A)| \le f(r) ||A|| \quad \forall A \in \mathcal{A}_{\Lambda \cap B_a(r)^c}^{\text{loc}}$$
(A.1)

and

$$|\psi_0(A) - \psi(A)| \leqslant f(r)||A|| \quad \forall A \in \mathcal{A}_{\Lambda^c \cap B_a(r)^c}^{\mathrm{loc}}.$$
 (A.2)

Then  $\psi$  is an almost local perturbation of  $\psi$ .

Proof. For any  $x \in \mathbb{Z}^2$ , we denote by  $\rho^{(x)}$  and  $\rho^{(x)}_0$  the density matrices of the restrictions to  $\mathcal{A}_x$  of  $\psi$  and  $\psi_0$  respectively. By assumption we have for any  $A \in \mathcal{A}_x$  that  $|\psi(A) - \psi_0(A)| \leq f(|x|) ||A||$ , which implies  $||\rho^{(x)} - \rho^{(x)}_0||_1 \leq f(|x|)$ . Let  $S^{(x)} = -\operatorname{Tr}(\rho^{(x)}\log\rho^{(x)})$  be the von Neumann entropy of  $\rho^{(x)}$ . Since  $\psi_0$  is pure, its restriction is pure as well by definition and so  $S_0^{(x)} = 0$ . If |x| is large enough so that f(|x|) < 1/e then Fannes' inequality yields

$$S^{(x)} \le f(|x|)\log(2^n/f(|x|)),$$
 (A.3)

where  $2^n$  is the dimension of the local Fock space for  $A_x$ . Since f decays faster than any polynomial we have that

$$S := \sum_{x \in \Lambda} S^{(x)} < \infty. \tag{A.4}$$

By subadditivity of the von Neumann entropy (see Theorem 3.7 of [2] for subadditivity for fermions), the quantity S is an upper bound for the entanglement entropy of any finite  $\Gamma \subset \mathbb{Z}^2$  in the state  $\psi$ .

We now show that  $\psi_0$  and  $\psi$  are g-close for some  $g \in \mathcal{F}$ . Let  $\Gamma \subseteq \overline{B(r)^c}$  be a finite region outside the ball B(r) and denote by  $\rho^{(\Gamma)}$  and  $\rho_0^{(\Gamma)}$  the density matrices of the restrictions to  $\mathcal{A}_{\Gamma}$  of the states  $\psi$  and  $\psi_0$  respectively. Similarly, let  $\rho^{(\Gamma \cap \Lambda)}$  and  $\rho^{(\Gamma \cap \Lambda^c)}$  be the density matrices of the restrictions of  $\psi$  to  $\mathcal{A}_{\Gamma \cap \Lambda}$  and to  $\mathcal{A}_{\Gamma \cap \Lambda^c}$  respectively. Equivalently,  $\rho^{(\Gamma \cap \Lambda)}$  is the restriction of  $\rho^{(\Gamma)}$  to  $\mathcal{A}_{\Gamma \cap \Lambda}$ , and similarly for  $\rho^{(\Gamma \cap \Lambda^c)}$ . Let

$$S(\rho^{(\Gamma)} \| \rho^{(\Gamma \cap \Lambda)} \hat{\otimes} \rho^{(\Gamma \cap \Lambda^c)}) = S(\rho^{(\Gamma)}) - S(\rho^{(\Gamma \cap \Lambda)}) - S(\rho^{(\Gamma \cap \Lambda^c)})$$
(A.5)

be the relative entropy of  $\rho^{(\Gamma)}$  and  $\rho^{(\Gamma \cap \Lambda)} \hat{\otimes} \rho^{(\Gamma \cap \Lambda^c)}$ . By non-negativity and subadditivity of the von Neumann entropy, and the bound (A.3) on  $S^{(x)}$  obtained above,

$$S(\rho^{(\Gamma)} \| \rho^{(\Gamma \cap \Lambda)} \, \hat{\otimes} \, \rho^{(\Gamma \cap \Lambda^c)}) \leqslant S(\rho^{(\Gamma)}) \leqslant \sum_{x: |x| > r} S^{(x)} \leqslant \frac{h(r)}{2} \tag{A.6}$$

for some  $h \in \mathcal{F}$  that only depends on f and n. The quantum Pinsker inequality now yields

$$\|\rho^{(\Gamma)} - \rho^{(\Gamma \cap \Lambda)} \, \hat{\otimes} \, \rho^{(\Gamma \cap \Lambda^c)}\|_1^2 \leqslant 2S(\rho^{(\Gamma)} \|\rho^{(\Gamma \cap \Lambda)} \otimes \rho^{(\Gamma \cap \Lambda^c)}) \leqslant h(r). \tag{A.7}$$

It follows from the assumptions (A.1,A.2) that

$$\|\rho^{(\Gamma \cap \Lambda)} \, \hat{\otimes} \, \rho^{(\Gamma \cap \Lambda^c)} - \rho_0^{\Gamma}\|_1 \leqslant f(r). \tag{A.8}$$

Combining the two last inequalities finally yields

$$\|\rho^{(\Gamma)} - \rho_0^{(\Gamma)}\| \le f(r) + h(r),$$
 (A.9)

thus proving the claim since  $f + h \in \mathcal{F}$ .

### A.2 Closeness at infinity implies almost local unitary equivalence

We now move to the transitivity itself. A similar result in the context of one-dimensional spin chains can be found in [35], and extended to the two-dimensional setting in [33]

**Proposition A.2.** Let v be a homogeneous pure product state. Assume that  $\psi$  is a homogeneous pure state that is f-close to v. Then there is  $\tilde{f} \in \mathcal{F}$  which depends only on f, and a homogeneous  $\tilde{f}$ -local unitary  $U \in \mathcal{A}^{al}$  such that  $\psi = v \circ Ad_U$ . The unitary U may be taken even if  $\psi$  has the same parity as v, and odd of  $\psi$  has opposite parity to v.

We first establish several lemmas which will be combined into a proof of the proposition at the end of Section A.2.2 below.

### A.2.1 GNS representation for homogeneous pure product states

Let  $v = \hat{\otimes}_{x \in \mathbb{Z}^2} v_x$  be a homogeneous pure product state on  $\mathcal{A}$ . For each  $x \in \mathbb{Z}^2$  let  $\pi_x : \mathcal{A}_x \to \mathcal{B}(\mathcal{H}_x)$  be the defining representation of  $\mathcal{A}_x$ . Since we assumed that each  $\mathcal{A}_x$  is a CAR algebra on n fermion modes we have  $\mathcal{H}_x \simeq \mathbb{C}^{2^n}$ . Let  $v_x \in \mathcal{H}_x$  be a unit vector representing the state  $v_x$  in  $\pi_x$  and let  $\Theta_x \in \mathcal{A}_x$  be the unique unitary such that  $\theta(A) = \mathrm{Ad}_{\Theta_x}(A)$  for all  $A \in \mathcal{A}_x$  and  $\pi_x(\Theta_x)\Psi_x = \Psi_x$ . Note that  $\Theta_x$  is even and  $\Theta_x^2 = \mathbb{1}$ . We call a vector  $\Psi \in \mathcal{H}_x$  homogeneous if it is an eigenvector of  $\pi_x(\Theta_x)$ , equivalently if  $\Psi$  represents a homogeneous state on  $\mathcal{A}_x$ .

Let  $\{\Upsilon_x^{(i)}\}_{i=1}^{2^n}$  be an orthonormal basis of  $\mathcal{H}_x$  consisting of homogeneous vectors, and such that  $\Upsilon_x^{(1)} = \Upsilon_x$ .

Let  $\mathcal{K}$  be the linear space of sequences  $\{\Psi_x\}_{x\in\mathbb{Z}^2}$  such that  $\Psi_x\in\mathcal{H}_x$  and such that  $\Psi_x\neq\Upsilon_x$  for only finitely many  $x\in\mathbb{Z}^2$ . Then the inner product

$$\langle \{\Psi_x\}, \{\Psi_x'\} \rangle = \prod_{x \in \mathbb{Z}^2} \langle \Psi_x, \Psi_x' \rangle \tag{A.10}$$

is well-defined for all  $\{\Psi_x\}$ ,  $\{\Psi_x'\} \in \mathcal{K}$  and extends to the whole of  $\mathcal{K}$  by linearity. We let  $\mathcal{H}$  be the Hilbert space obtained by closing  $\mathcal{K}$  w.r.t. the norm induced by this inner product. The set of sequences  $\{\Upsilon_x^{(i_x)}\}$  with only finitely many  $i_x \neq 1$  form an orthonormal basis for  $\mathcal{H}$ .

We write  $\bigotimes_{x \in \mathbb{Z}^2} \Psi_x := \{\Psi_x\}$  for any sequence in  $\mathcal{H}_v$ . It is clear that for any  $X \in \mathbb{Z}^2$ , the Hilbert space  $\mathcal{H}$  decomposes as a tensor product  $\mathcal{H} = \mathcal{H}_X \otimes \mathcal{H}_{X^c}$  and if X is finite then  $\mathcal{H}_X$  is identified with  $\bigotimes_{x \in X} \mathcal{H}_x$ .

We now represent  $\mathcal{A}$  on  $\mathcal{H}$  using a Jordan-Wigner construction. Fix an enumeration  $x:\mathbb{N}\to\mathbb{Z}^2$  of the sites And let  $X_N=\{x(n):1\leqslant n\leqslant N\}$  for all  $N\in\mathbb{N}$ . Assume moreover that the enumeration is such that  $B_0(r)=X_{N_r}$  for all  $r\in\mathbb{N}$  and some  $N_r\in\mathbb{N}$ . For any finite  $X\subseteq\mathbb{Z}^2$  and any homogeneous operator  $A\in\mathcal{A}_{x(N)}$  we let

$$\pi(A) = \begin{cases} \mathbb{1}_{\mathcal{H}_{X_{N-1}}} \otimes \pi_{x(N)}(A) \otimes \mathbb{1}_{\mathcal{H}_{X_N}^c} & \text{if } A \text{ is even} \\ \left( \bigotimes_{x \in X_{N-1}} \pi_x(\Theta_x) \right) \otimes \pi_{x(N)}(A) \otimes \mathbb{1}_{\mathcal{H}_{X_N^c}} & \text{if } A \text{ is odd.} \end{cases}$$
(A.11)

The 'tail' of grading operators  $\Theta_x$  ensures that odd operators supported on disjoint regions anticommute in the representation. Since the on-site homogeneous operators generate the whole algebra, this defines the representation uniquely.

To see that this is a GNS representation for v, let  $A_{x_1} \cdots A_{x_l}$  be a monomial of homogeneous operators  $A_{x_i} \in \mathcal{A}_{x_i}$ , writing  $\Upsilon := \bigotimes_x \Upsilon_x$ , and use the fact that the  $\Upsilon_x$  are even to find

$$\langle \Upsilon, \pi(A_{x_1} \cdots A_{x_l}) \Upsilon \rangle = \upsilon(A_{x_1}) \cdots \upsilon(A_{x_l}) = \upsilon(A_{x_1} \cdots A_{x_l}). \tag{A.12}$$

By linearity and continuity it follows that  $\Upsilon(A) = \langle \Upsilon, \pi(A) \Upsilon \rangle$  for any  $A \in \mathcal{A}$ . Finally, the vector  $\Upsilon \in \mathcal{H}$  is cyclic for the representation  $\pi$  by construction.

### A.2.2 Construction of a unitary

Let  $\psi$  be a homogeneous pure state on  $\mathcal{A}$  such that the hypotheses of Proposition A.2 hold w.r.t. the product state v. i.e. the state  $\psi$  is f-close to v for some  $f \in \mathcal{F}$ .

For any  $X \subseteq \mathbb{Z}^2$  let  $P_X$  be the projection on the state  $\Upsilon_X = \bigotimes_{x \in X} \Upsilon_x \in \mathcal{H}_X$ . Note that each  $P_X$  is even. Let r' > r and let  $P_{rr'} = P_{B_0(r') \setminus B_0(r)}$ . Denote by  $\rho_r$  the density matrix acting on  $\mathcal{H}_r := \mathcal{H}_{B_0(r)}$  corresponding to the restriction of  $\psi$  to  $\mathcal{A}_{B_0(r)}$ . Note that for all  $A \in \mathcal{A}_{B_0(r)}$ ,

$$\operatorname{Tr}(\rho_r \pi(A)) = \psi(A) = \psi \circ \theta(A) = \operatorname{Tr}(\Theta_r^* \rho_r \Theta_r \pi(A)), \tag{A.13}$$

by cyclicity of the trace, where  $\Theta_r$  is the restriction of  $\Theta$  to the  $\mathcal{H}_r$ . Since the density matrix is unique, we conclude that  $\rho_r$  is even.

**Lemma A.3.** For any  $r, r' \in \mathbb{N}$  such that r' > r we have

$$\|\rho_{r'}(\mathbb{1} \hat{\otimes} P_{rr'}^{\perp})\|_{1}^{2} \leqslant f(r)$$
 (A.14)

where we interpret all operators as operators on  $\mathcal{H}_{r'}$  by tensoring with the identity if needed.

*Proof.* Since  $P_{rr'}^{\perp}$  is even we have  $\mathbb{1} \otimes P_{rr'}^{\perp} = \mathbb{1} \otimes P_{rr'}^{\perp}$ . Then using Hölder's inequality we get

$$\|\rho_{r'}P_{rr'}^{\perp}\|_{1}^{2} \leq \|\sqrt{\rho_{r'}}\|_{2}^{2} \|\sqrt{\rho_{r'}P_{rr'}^{\perp}}\|_{2}^{2} = \operatorname{Tr}\{\rho_{r'}\} \times \operatorname{Tr}\{\rho_{r'}P_{rr'}^{\perp}\}$$
(A.15)

$$= \psi(P_{rr'}^{\perp}) = (\psi - v)(P_{rr'}^{\perp}) \leqslant f(r) \tag{A.16}$$

by the assumption that  $\psi$  and v are f-close.

Now let  $Q_r = P_{B_0(r)^c}$ , which is an even rank 1 projection on  $\mathcal{H}_{B_r^c}$ . We have

**Lemma A.4.** The sequence  $\{\rho_r \otimes Q_r\}_{r \in \mathbb{N}}$  is a Cauchy sequence in trace norm.

*Proof.* Note first that  $Q_r = P_{rr'} \otimes Q_{r'}$  so

$$\|\rho_{r'} \otimes Q_{r'} - \rho_r \otimes Q_r\|_1 \le \|\rho_{r'} - \rho_r \otimes P_{rr'}\|_1.$$
 (A.17)

Let us denote by  $\operatorname{Tr}_{rr'}$  the partial trace over  $\mathcal{H}_{rr'} = \mathcal{H}_{B_0(r')\setminus B_0(r)}$ . We bound the right-hand side of (A.17) by noting that

$$\rho_r \otimes P_{rr'} = \text{Tr}_{rr'} \{ P_{rr'} \rho_{r'} P_{rr'} \} \otimes P_{rr'} + \text{Tr}_{rr'} \{ P_{rr'}^{\perp} \rho_{r'} P_{rr'}^{\perp} \} \otimes P_{rr'}$$
(A.18)

$$= P_{rr'}\rho_{r'}P_{rr'} + \text{Tr}_{rr'}\{P_{rr'}^{\perp}\rho_{r'}P_{rr'}^{\perp}\} \otimes P_{rr'}$$
(A.19)

$$= \rho_{r'} - P_{rr'}\rho_{r'}P_{rr'}^{\perp} - P_{rr'}^{\perp}\rho_{r'}P_{rr'} - P_{rr'}^{\perp}\rho_{r'}P_{rr'}^{\perp}$$
(A.20)

$$+\operatorname{Tr}_{rr'}\{P_{rr'}^{\perp}\rho_{r'}P_{rr'}^{\perp}\}\otimes P_{rr'}.\tag{A.21}$$

Here the last four terms are bounded in trace norm by  $\sqrt{f(r)}$  using Hölder's inequality and Lemma A.3 yielding

$$\|\rho_{r'} \hat{\otimes} Q_{r'} - \rho_r \hat{\otimes} Q_r\|_1 \leqslant 4\sqrt{f(r)}. \tag{A.22}$$

We conclude that the sequence  $\{\rho_r \otimes Q_r\}_r$  is indeed Cauchy.

Let us denote by  $\rho \in \mathcal{B}(\mathcal{H})$  the limit of the Cauchy sequence  $\{\rho_r \hat{\otimes} Q_r\}$ . By construction  $\rho$  represents the state  $\psi$  which is homogeneous and pure, so  $\rho$  is actually an even rank one projection  $\rho = |\Psi\rangle\langle\Psi|$  for some homogeneous unit vector  $|\Psi\rangle \in \mathcal{H}$ . The next step towards proving Proposition A.2 is to construct a sequence of pure density matrices  $\tilde{\rho}_r$  that converge to  $\rho$ , and such that  $\tilde{\rho}_r$  agrees with product state  $\Upsilon$  outside the ball  $B_0(r)$ .

**Lemma A.5.** Let  $r_0 \in \mathbb{N}$  be the smallest natural number such that  $36\sqrt{f(r)} < 1$  for all  $r \ge r_0$ . Then for all  $r \ge r_0$  we have that the pure density matrices

$$\tilde{\rho}_r := \frac{(\mathbb{1} \otimes Q_r) \rho(\mathbb{1} \otimes Q_r)}{\text{Tr}\{(\mathbb{1} \otimes Q_r) \rho(\mathbb{1} \otimes Q_r)\}}$$
(A.23)

are well defined and

$$\|\rho - \tilde{\rho}_r\|_1 \leqslant 9\sqrt{f(r)}.\tag{A.24}$$

Proof. First of all,

$$\rho_{r'} P_{rr'}^{\perp} Q_{r'} - \rho Q_r^{\perp} = (\rho_{r'} Q_{r'} - \rho) + (\rho - \rho_{r'} Q_{r'}) Q_r \tag{A.25}$$

converges to zero as  $r' \to \infty$  in trace norm by Lemma A.4. (A.14) then implies that

$$\|\rho(\mathbb{1} \otimes Q_r^{\perp})\|_1 \leqslant \sqrt{f(r)}. \tag{A.26}$$

From

$$1 = \|\rho\|_1 \le \|\rho(\mathbb{1} \otimes Q_r)\|_1 + \|\rho(\mathbb{1} \otimes Q_r^{\perp})\|_1 \tag{A.27}$$

we then obtain

$$\|\rho(\mathbb{1} \otimes Q_r)\|_1 \geqslant 1 - \sqrt{f(r)}. \tag{A.28}$$

Using that  $\rho$  is a rank one projection it follows that

$$\operatorname{Tr}\{(\mathbb{1} \otimes Q_r)\rho(\mathbb{1} \otimes Q_r)\} = \|\rho(\mathbb{1} \otimes Q_r)\|_2^2 \leq \|\rho(\mathbb{1} \otimes Q_r)\|_1^2 \leq (1 - \sqrt{f(r)})^2. \tag{A.29}$$

For  $r \ge r_0$  we have  $36\sqrt{f(r)} < 1$  so certainly  $\text{Tr}\{(\mathbb{1} \otimes Q_r)\rho(\mathbb{1} \otimes Q_r)\} > 1/2$  and therefore  $\tilde{\rho}_r$  is well-defined. (Note that  $\tilde{\rho}_r$  is the rank one projection onto the span of  $(\mathbb{1} \otimes Q_r)|\Psi\rangle$ ).

To show the bound (A.24) we note that (A.26) implies  $\|\rho - Q_r \rho Q_r\|_1 \leq \|Q_r^{\perp} \rho Q_r\|_1 + \|Q_r \rho Q_r^{\perp}\|_1 + \|Q_r^{\perp} \rho Q_r^{\perp}\|_1 \leq 3\sqrt{f(r)}$ . Combining this with  $|1 - \text{Tr}\{Q_r \rho Q_r\}| \leq \|\rho - Q_r \rho Q_r\|_1$  and  $\text{Tr}\{Q_r \rho Q_r\} > 1/2$  for all  $r \geq r_0$  we find

$$\|\rho - \tilde{\rho}_r\|_1 = \|\rho - Q_r \rho Q_r\|_1 + \frac{1}{\text{Tr}\{Q_r \rho Q_r\}} \|\left(1 - \text{Tr}\{Q_r \rho Q_r\}\right) Q_r \rho Q_r\|_1 \leqslant 9\sqrt{f(r)}$$
(A.30)

as required. 
$$\Box$$

Let us denote by  $\tilde{\psi}_r$  the pure states corresponding to the density matrices  $\tilde{\rho}_r$ . i.e.  $\tilde{\psi}_r(A) = \text{Tr}\{\tilde{\rho}_r\pi(A)\}$  for all  $A\in\mathcal{A}$ . Furthermore, (A.24) implies that the density matrices  $\tilde{\rho}_r$  are all even, for r large enough. Indeed, vectors with different parity are orthogonal to each other, so the trace norm of the difference of their density matrices equals 2, but the bound is  $9\sqrt{f(r)} < 2$  for all r large enough.

**Lemma A.6.** For all  $r \ge r_0$  there is an even unitary  $U_r \in \mathcal{A}_{B_0(r+1)}$  such that  $\tilde{\psi}_{r+1} = \tilde{\psi}_r \circ \operatorname{Ad}_{U_r}$  and such that

$$\| \mathbb{1} - U_r \| \le 6f(r)^{1/4}.$$
 (A.31)

*Proof.* From (A.24) we have

$$\|\tilde{\rho}_{r+1} - \tilde{\rho}_r\|_1 \le \|\tilde{\rho}_{r+1} - \rho\|_1 + \|\rho - \tilde{\rho}_r\|_1 \le 18\sqrt{f(r)}.$$
 (A.32)

for all  $r \ge r_0$ . We can choose normalised vector representatives  $|\Psi_r\rangle$  of  $\tilde{\rho}_r$ , all with the same parity and such that  $a_r := \langle \Psi_r, \Psi_{r+1} \rangle > 0$  are real and positive. Note that once  $\Psi_{r_0}$  is chosen, this uniquely fixes all the other  $\Psi_r$ . Then

$$\cos(\theta_r)^2 := \left| \langle \Psi_r, \Psi_{r+1} \rangle \right|^2 = \text{Tr}\{\tilde{\rho}_{r+1}\tilde{\rho}_r\} = 1 + \text{Tr}\{(\tilde{\rho}_{r+1} - \tilde{\rho}_r)\tilde{\rho}_r\}$$
(A.33)

hence

$$\cos(\theta_r)^2 \geqslant 1 - 18\sqrt{f(r)}.\tag{A.34}$$

We can decompose  $\Psi_{r+1}$  as

$$\Psi_{r+1} = \cos(\theta_r)\Psi_r + \sin(\theta_r)\Psi_r^{\perp} \tag{A.35}$$

where  $\Psi_r^{\perp}$  is the normalized component of  $\Psi_r$  that is perpendicular to  $\Psi_{r+1}$ .

By construction the states corresponding to the  $\tilde{\rho}_r$  agree with the product state v outside of  $B_0(r)$ , so we have  $\Psi_r = \Psi_r' \otimes \Upsilon_{B_0(r)^c}$  for all  $r \geq r_0$ . Here  $\Psi_r' \in \mathcal{H}_r$  is a homogeneous vector and  $\Upsilon_{B_0(r)^c} = \bigotimes_{x \in B_0(r)^c} \Upsilon_x \in \mathcal{H}_{B_0(r)^c}$  is the restriction of  $\Upsilon$  to the complement of  $B_0(r)$ . Let  $V_r$  be the unitary which acts on the subspace spanned by  $\Psi_r$  and  $\Psi_{r+1}$  as the  $\theta_r$ -rotation which rotates  $\Psi_{r+1}$  into  $\Psi_r$ , and acts as the identity on the complement of this subspace. Then  $V_r$  is even and  $V_r = \pi(U_r)$  for a unique even unitary  $U_r \in \mathcal{A}_{B_0(r+1)}$ . Since  $\pi(U_r)\Psi_{r+1} = \Psi_r$  we have  $\tilde{\psi}_{r+1} = \tilde{\psi}_r \circ \mathrm{Ad}_{U_r}$  as required. With (A.34) we find that

$$\| \mathbb{1} - U_r \|^2 = \| \mathbb{1} - V_r \|^2 \leqslant \theta_r^2 \leqslant \frac{18\sqrt{f(r)}}{1 - 18\sqrt{f}} \leqslant 36\sqrt{f(r)}$$
(A.36)

which establishes the bound (A.31).

Let us finally combine the unitary steps provided by Lemma A.6 into a single unitary, providing a proof of the main proposition.

Proof of Proposition A.2. Let  $g_r := 6f(r)^{1/4}$  so the unitaries  $U_r \in \mathcal{A}_{B_0(r+1)}$  provided by Lemma A.6 are even and satisfy  $||U_r - 1|| \leq g_r$ . Since  $f \in \mathcal{F}$  the sequence  $\{g_r\}_{r \geq r_0}$  is summable. For  $R \geq r_0$ , define  $U^{(R)} := \prod_{r=r_0}^R U_r$ , which is even. Then for  $R' > R \geq r_0$  we have

$$||U^{(R')} - U^{(R)}|| \le \sum_{r=R+1}^{R'} ||U^{(r)} - U^{(r-1)}|| = \sum_{r=R+1}^{R'} g_r.$$
(A.37)

Since  $\{g_r\}_{r\geqslant r_0}$  is summable, this shows that  $\{U^{(R)}\}_{R\geqslant r_0}$  is Cauchy and converges to a unitary  $U^{(\infty)}$  which satisfies

$$\tilde{\psi}_{r_0} \circ \operatorname{Ad}_{U^{(\infty)}} = \lim_{R \uparrow \infty} \ \tilde{\psi}_{r_0} \circ \operatorname{Ad}_{U^{(R)}} = \lim_{R \uparrow \infty} \ \tilde{\psi}_R = \psi. \tag{A.38}$$

Note moreover that  $U^{(\infty)}$  is almost local because  $||U^{(\infty)} - U^{(R)}|| \le G(R)$  with  $G(R) = \sum_{r=R}^{\infty} g_r$ , which defines a rapidly decaying function in  $\mathcal{F}$ . It is also even as the norm limit of even operators.

Finally, since  $\tilde{\psi}_{r_0}$  and v are both homogeneous and differ only in the region  $B_0(r_0)$ , there is a homogeneous unitary  $V \in \mathcal{A}_{B_0(r_0)}$  such that  $\tilde{\psi}_{r_0} = v \circ \mathrm{Ad}_V$ . It can be chosen even provided  $\tilde{\psi}_{r_0}, v$ , and therefore  $\psi, v$ , have the same parity. It follows that the claim of the proposition holds true with  $U = VU^{(\infty)}$ .

### A.3 A continuity result for parity of states

**Lemma A.7.** Let a < b and let  $[a,b] \ni \phi \mapsto \omega_{\phi}$  be a weakly-\* continuous family of homogeneous pure states that are all unitarily equivalent to a homogeneous pure product state  $\omega_0$ . Suppose the unitaries  $V_{\phi}$  are homogeneous and f-localised near the origin, where f is independent of  $\phi$ . Then all  $\omega_{\phi}$  have the same parity and so do all the  $V_{\phi}$ .

*Proof.* Let  $(\Pi, \mathcal{H}, \Omega)$  be the GNS representation of  $\omega_0$ . Since  $\omega_0$  is homogeneous, there is a unique unitary  $\Theta \in \mathcal{B}(\mathcal{H})$  such that

$$\Theta\Omega = \Omega, \quad \Pi(\theta(A)) = \Theta^* \Pi(A) \Theta, \quad \forall A \in \mathcal{A}.$$
 (A.39)

Since  $\Omega$  is cyclic and  $\theta^2 = \mathrm{id}$ , the first equality above implies that  $\Theta^2 = \mathbb{1}$ . Moreover, the purity of  $\omega_0$  implies that  $\mathcal{B}(\mathcal{H})$  is the weak closure of  $\Pi(\mathcal{A})$  and so the adjoint action of  $\Theta$  extends the action of  $\theta$  to all of  $\mathcal{B}(\mathcal{H})$ .

Let  $\theta_L$  be the restriction of  $\theta$  to the algebra  $\mathcal{A}_{B(L)}$ . Since  $\omega_0$  is a product state, it is invariant under  $\theta_L$  and so there is a unitary  $\Theta_L$  implementing that partial parity such that  $\Theta_L\Omega = \Omega$ . For any  $A, B \in \mathcal{A}_{loc}$ , there is L large enough such that  $\theta_L(A) = \theta(A)$  and so

$$\langle \Pi(A)\Omega, \Theta_L \Pi(B)\Omega \rangle = \langle \Theta_L^* \Pi(A)\Theta_L \Omega, \Pi(B)\Omega \rangle = \langle \Pi(\theta(A))\Omega, \Pi(B)\Omega \rangle = \langle \Pi(A)\Omega, \Theta\Pi(B)\Omega \rangle. \tag{A.40}$$

It now follows by density that  $\Theta_L$  converges to  $\Theta$  in the weak operator topology.

By assumption, the vectors  $\Omega_{\phi} = \Pi(V_{\phi})\Omega$  are unit vector representatives of the states  $\omega_{\phi}$  in the GNS space  $\mathcal{H}$ . Since all  $\omega_{\phi}$  are homogeneous,

$$\Theta\Omega_{\phi} = \lambda_{\phi}\Omega_{\phi}, \qquad \lambda_{\phi} \in \{+1, -1\}. \tag{A.41}$$

We claim that  $\phi \mapsto \lambda_{\phi}$  is constant. For any  $\phi, \phi'$ ,

$$\lambda_{\phi} - \lambda_{\phi'} = \langle \Omega_{\phi}, \Theta \Omega_{\phi} \rangle - \langle \Omega_{\phi'}, \Theta \Omega_{\phi'} \rangle = \langle \Omega, \Pi(V_{\phi})^* \Theta \Pi(V_{\phi}) \Omega \rangle - \langle \Omega, \Pi(V_{\phi'})^* \Theta \Pi(V_{\phi'}) \Omega \rangle$$
(A.42)

$$= \langle \Omega, \Pi(\theta(V_{\phi}))^* \Pi(V_{\phi}) \Omega \rangle - \langle \Omega, \Pi(\theta(V_{\phi'}))^* \Pi(U_{\phi'}) \Omega \rangle$$
(A.43)

since  $\Theta\Omega = \Omega$ . By the uniform almost locality of the family  $V_{\phi}$ , we can replace  $\theta$  by its restriction  $\theta_L$ , run the sequence of equalities in reverse order with  $\Theta \to \Theta_L$  and conclude that

$$\left|\lambda_{\phi} - \lambda_{\phi'}\right| \le \left|\omega_{\phi}(\Theta_L) - \omega_{\phi'}(\Theta_L)\right| + cf(L). \tag{A.44}$$

Taking L large enough such that f(L) < 1 and choosing then  $\phi$  and  $\phi'$  close enough such that  $|\omega_{\phi}(\Theta_L) - \omega_{\phi'}(\Theta_L)| < 1$  (which is possible by weak-\* continuity) yields  $|\lambda_{\phi} - \lambda_{\phi'}| < 1$ . Since  $\lambda_{\phi}$  and  $\lambda_{\phi'}$  take values in  $\{+1, -1\}$ , it follows that  $\lambda_{\phi} = \lambda_{\phi'}$  if  $\phi$  and  $\phi'$  are close enough, which yields the claim.

### B Flux insertion for free fermions

In this appendix we will study the spin up component  $\mathscr{P}^{\uparrow}$  of a spin Hall state  $\mathscr{P}_{SH} = \mathscr{P}^{\uparrow} \oplus \mathscr{P}^{\downarrow}$ . Recall that  $\mathscr{P}^{\uparrow}$  is a projection on the Hilbert space  $\mathcal{K}_1 = l^2(\mathbb{Z}^2; \mathbb{C}^2)$ . We fix an orthonormal basis  $\{e_i\}_{i=1,2}$  of  $\mathbb{C}^2$  so that  $\{\delta_x \otimes e_i\}_{x,i}$  is an orthonormal basis of  $\mathcal{K}_1$ . For any operator  $\mathscr{A}$  on  $\mathcal{K}_1$  we denote by

$$\mathscr{A}(x,i;y,j) := \langle \delta_x \otimes e_i, \mathscr{A} \delta_y \otimes e_j \rangle \tag{B.1}$$

its matrix elements with respect to this basis.

Since in this appendix we will be exclusively concerned with the spin up component we write  $\mathscr{P} = \mathscr{P}^{\uparrow}$  here and only here, dropping the spin label from the notation.

Throughout this appendix we will use the following notations:  $L = \{x \in \mathbb{Z}^2 : x_1 \leq 0\}$  is the left half plane and  $R = \mathbb{Z}^2 \setminus L$  is the right half plane. For any subset  $S \subset \mathbb{Z}^2$  we write  $\chi_S$  for the indicator function on S, and  $\Pi_S$  for the projection on S, i.e.  $\Pi_S | \delta_x \otimes i \rangle = \delta_{x \in S} | \delta_x \otimes i \rangle$ . We write  $l_L$  and  $l_R$  for the left and right horizontal axes respectively, regarded as subsets of  $\mathbb{R}^2$ . For any  $S \subset \mathbb{R}^2$  and any  $x \in \mathbb{Z}^2$  we write  $\mathrm{dist}(x,S) = \inf_{y \in S} \|x - y\|_1$  for the distance from x to the set S.

### B.1 Locality estimates for Fermi projections

Recall that the Fermi projection  $\mathscr{P}_{SH}$  describing the spin Hall state is exponentially local and so

$$|\mathscr{P}(x,i;y,j)| \leqslant Ce^{-\eta \|x-y\|_1} \tag{B.2}$$

where  $C, \eta > 0$  are the constants introduced in (2.13).

Lemma B.1. We have

$$\left| \mathscr{P}_{\phi}^{h}(x,i;y,j) \right| \leq C e^{-\eta \|x-y\|_{1}} \quad and \quad \left| \mathscr{K}_{\phi}^{h}(x,i;y,j) \right| \leq c e^{-\xi(|x_{1}-y_{1}|+|x_{2}|+|y_{2}|)}$$
 (B.3)

for some constants  $c, \xi > 0$  and for all  $\phi \in \mathbb{R}$ , all  $x = (x_1, x_2), y = (y_1, y_2) \in \mathbb{Z}^2$  and all  $i, j \in \{1, 2\}$ .

Proof. The first claim follows immediately from Eq (B.2) and the definition  $\mathscr{P}^h_{\phi} = \mathrm{e}^{\mathrm{i}\phi\Pi_h}\mathscr{P}\mathrm{e}^{-\mathrm{i}\phi\Pi_h}$ . In order to bound the matrix elements of  $\mathscr{K}^h_{\phi} = \mathrm{i}[\partial_{\phi}\mathscr{P}^h_{\phi},\mathscr{P}^h_{\phi}]$  we first bound the matrix elements of  $\partial_{\phi}\mathscr{P}^h_{\phi} = \mathrm{i}[\Pi_h,\mathscr{P}^h_{\phi}]$ . If  $x,y\in h$  of  $x,y\notin h$  then  $(\partial_{\phi}\mathscr{P}^h_{\phi})(x,i;y,j)=0$ . If  $x\in h$  and  $y\notin h$  then using the first claim of this lemma we obtain

$$\left| (\partial_{\phi} \mathscr{P}_{\phi}^{h})(x, i; y, j) \right| = |\mathscr{P}_{\phi}(x, i; y, j)| \le C e^{-\eta ||x - y||_{1}} \le C e^{-\eta (|x_{1} - y_{1}| + |x_{2}| + |y_{2}|)}.$$
(B.4)

where for the last inequality we used  $x \in h$  and  $y \notin h$  so  $||x - y||_1 = |x_1 - y_1| + |x_2 - y_2| = |x_1 - y_1| + |x_2| + |y_2|$ . The same result is obtained if  $x \notin h$  and  $y \in h$ . The final claim of the lemma now follows straightforwardly from the definition  $\mathcal{K}_{\phi}^h = \mathrm{i} [\partial_{\phi} \mathcal{P}_{\phi}^h, \mathcal{P}_{\phi}^h]$  and the bounds on the matrix elements of  $\partial_{\phi} \mathcal{P}_{\phi}^h$  and  $\mathcal{P}_{\phi}^h$ .

We now turn to locality properties of the defect states  $\mathscr{P}_{\phi}$  defined in (5.20). It will be useful to consider the following quantities:

$$a_n(x,y) := \sup_{\substack{\vec{\phi} \in [-\pi,\pi]^n \\ i,j \in \{1,2\}}} \left| (\mathscr{K}_{\phi_1}^h \cdots \mathscr{K}_{\phi_n}^h)(x,i;y,j) \right|$$
(B.5)

$$b_n(x,y) := \sup_{\substack{\vec{\phi} \in [-\pi,\pi]^n \\ i,j \in \{1,2\}}} |(\mathscr{K}_{\phi_1} \cdots \mathscr{K}_{\phi_n})(x,i;y,j)|.$$
(B.6)

**Lemma B.2.** There are constants  $c, \kappa, \xi > 0$  such that

$$a_n(x,y) \le c^n \kappa^{n-1} e^{-\xi(|x_1-y_1|+|x_2|+|y_2|)}$$
 (B.7)

$$b_n(x,y) \le c^n \kappa^{n-1} e^{-\xi(|x_1-y_1|+|x_2|+|y_2|)} \chi_L(x) \chi_L(y)$$
 (B.8)

for all  $x, y \in \mathbb{Z}^2$ .

 $\textit{Proof.} \text{ We take } c, \xi > 0 \text{ such that } \left| \mathscr{K}^h_\phi(x,i;y,j) \right| \leqslant c \mathrm{e}^{-2\xi(|x_1-y_1|+|x_2|+|y_2|)} \text{ (cf. Lemma B.1) and }$ 

$$\kappa = 2 \sup_{x_1 \in \mathbb{Z}} \sum_{z \in \mathbb{Z}^2} e^{-\xi |z_1 - x_1| - 3\xi |z_2|}.$$
 (B.9)

Then the claimed bound on  $a_n(x,y)$  holds for n=1 by that same Lemma. We proceed by induction using the again second claim of Lemma B.1 to obtain

$$a_n(x,y) \le \sup_{\phi \in [-\pi,\pi]} \sup_{i \in \{1,2\}} \sum_{z,i'} |\mathscr{K}_{\phi}(x,i;z,i')| \times a_{n-1}(z,y)$$
 (B.10)

$$\leq c^n \kappa^{n-2} \sum_{z,j'} e^{-2\xi(|x_1 - z_1| + |x_2| + |z_2|)} \times e^{-\xi(|z_1 - y_1| + |z_2| + |y_2|)}$$
(B.11)

noting that  $e^{-\xi|z_1-y_1|}e^{-2\xi|x_1-z_1|} \le e^{-\xi|x_1-y_1|}e^{-\xi|x_1-z_1|}$  this is easily bounded by

$$\leq c^n \kappa^{n-1} e^{-\xi(|x_1 - y_1| + |x_2| + |y_2|)}$$
 (B.12)

which proves the bound on  $a_n$  for all n. The bound on  $b_n$  is obtained in exactly the same way and recalling that  $\mathscr{K}_{\phi} = \Pi_L \mathscr{K}_{\phi}^h \Pi_L$ .

Recall that  $\mathscr{U}_{\phi}$  is the unique solution to  $i(d\mathscr{U}_{\phi}/d\phi) = \mathscr{K}_{\phi}\mathscr{U}_{\phi}$  with initial condition  $\mathscr{U}_{0} = \mathbb{1}$ . Similarly we let  $\mathscr{U}_{\phi}^{h}$  be the unique solution to  $\mathrm{i}(\mathrm{d}\mathscr{U}_{\phi}^{h}/\mathrm{d}\phi) = \mathscr{K}_{\phi}^{h}\mathscr{U}_{\phi}^{h}$  with initial condition  $\mathcal{U}_0^h = 1.$ 

**Lemma B.3.** There are constants  $c, \xi > 0$  such that

$$\left| (\mathcal{U}_{\phi}^{h} - 1)(x, i; y, j) \right| \le c e^{-\xi(|x_1 - y_1| + |x_2| + |y_2|)}$$
 (B.13)

$$|(\mathcal{U}_{\phi} - 1)(x, i; y, j)| \le c e^{-\xi(|x_1 - y_1| + |x_2| + |y_2|)} \chi_{\le 0}(x_1) \chi_{\le 0}(x_2)$$
(B.14)

for all  $x = (x_1, x_2), y = (y_1, y_2) \in \mathbb{Z}^2$ , all  $i, j \in \{1, 2\}$ , and all  $\phi \in [-\pi, \pi]$ .

*Proof.* We have

$$\left| (\mathcal{U}_{\phi}^{h} - \mathbb{1})(x, i; y, j) \right| \leqslant \sum_{n=1}^{\infty} \frac{|\phi|^{n}}{n!} a_{n}(x, y)$$
(B.15)

where  $a_n(x,y)$  was defined in (B.5). The first claim of the Lemma now follows straightforwardly from the bound on  $a_n(x,y)$  provided by Lemma B.2. The second claim follows in the same way using  $b_n(x,y)$  instead. 

 $\mathscr{U}_{\phi}$  is a restriction of  $\mathscr{U}_{\phi}^{h}$  to the left horizontal axis. Since  $\mathscr{U}_{\phi}^{h}$  acts non-trivially only near the horizontal axis so we expect  $\mathscr{U}_{\phi}^{h} - \mathscr{U}_{\phi}$  to be localised near the right horizontal axis.

Let us introduce a new family of projections

$$\mathcal{Q}_{\phi} := e^{i(\phi + \pi)\Pi_h} \mathscr{P}_{-} e^{-i(\phi + \pi)\Pi_h}$$
(B.16)

which satisfies  $\partial_{\phi} \mathcal{Q}_{\phi} = -i[\mathcal{L}_{\phi}, \mathcal{Q}_{\phi}]$  with  $\mathcal{L}_{\phi} = i[\partial_{\phi} \mathcal{Q}_{\phi}, \mathcal{Q}_{\phi}]$ . We let  $\mathcal{V}_{(\phi_2, \phi_1)}$  be the unique unitaries such that  $i(d\mathcal{V}_{(\phi_2,\phi_1)}/d\phi_2) = \mathcal{L}_{\phi_2}\mathcal{V}_{(\phi_2,\phi_1)}$  and  $\mathcal{V}_{(\phi_1,\phi_1)} = \mathbb{1}$ , then  $\mathcal{Q}_{\phi_2} = \mathcal{V}_{(\phi_2,\phi_1)}\mathcal{Q}_{\phi_1}\mathcal{V}_{(\phi_1,\phi_2)}$ .

We want to compare the unitaries  $\mathscr{V}_{(\phi_1,\phi_2)}$  generated by the  $\mathscr{L}_{\phi}$  with the unitaries  $\mathscr{U}_{\phi}^h$ generated by the  $\mathscr{K}_{\phi}^{h}$ . It is therefore convenient to extend the notation by letting  $\mathscr{U}_{(\phi_{2},\phi_{1})}^{h}$  be the solution to  $i(d\mathscr{U}_{(\phi_2,\phi_1)}^h/d\phi_2) = \mathscr{K}_{\phi_2}^h\mathscr{U}_{(\phi_2,\phi_1)}^h$  and  $\mathscr{U}_{(\phi_1,\phi_1)} = \mathbb{1}$ . Then  $\mathscr{U}_{\phi}^h = \mathscr{U}_{(\phi,0)}^h$ . By construction, the projections  $\mathscr{Q}_{\phi}$  differ from  $\mathscr{P}_{\phi}^h$  only near the right horizontal axis, and

so the same holds for the generators  $\mathscr{L}_{\phi}$  and  $\mathscr{K}_{\phi}^{h}$  and for the unitaries  $\mathscr{V}_{\phi_{2},\phi_{1}}$  and  $\mathscr{U}^{h}(\phi_{2},\phi_{1})$ .

**Lemma B.4.** There are constants  $c, \xi > 0$  such that

$$\left| (\mathcal{U}_{\phi}^{h} - \mathcal{U}_{\phi})(x, i; y, j) \right| \leqslant c e^{-\xi(\|x - y\|_{1} + \operatorname{dist}(x, l_{R}))}$$
(B.17)

$$\left| (\mathcal{Q}_{\phi} - \mathscr{P}_{\phi}^{h})(x, i; y, j) \right| \leqslant c e^{-\xi(\|x - y\|_{1} + \operatorname{dist}(x, l_{R}))}$$
(B.18)

$$\left| (\mathcal{L}_{\phi} - \mathcal{K}_{\phi}^{h})(x, i; y, j) \right| \leqslant c e^{-\xi(\|x - y\|_1 + \operatorname{dist}(x, l_R))}$$
(B.19)

$$\left| (\mathscr{V}_{(\phi_2,\phi_1)} - \mathscr{U}_{(\phi_2,\phi_1)}^h)(x,i;y,j) \right| \le c e^{-\xi(\|x-y\|_1 + \operatorname{dist}(x,l_R))}$$
(B.20)

for all  $x = (x_1, x_2), y = (y_1, y_2) \in \mathbb{Z}^2$ , all  $i, j \in \{1, 2\}$ , and all  $\phi, \phi_1, \phi_2 \in [-\pi, \pi]$ .

*Proof.* The first bound: If  $x, y \in R$  this follows immediately from Lemma B.3. To deal with the other cases, note first that

$$\left| (\mathcal{U}_{\phi}^{h} - \mathcal{U}_{\phi})(x, i; y, j) \right| \leq \sum_{n \geq 1} \frac{|\phi|^{n}}{n!} c_{n}(x, y)$$
(B.21)

with

$$c_n(x,y) = \sup_{\substack{\vec{\phi} \in [-\pi,\pi]^n \\ i,j \in \{1,2\}}} \left| (\mathcal{K}_{\phi_1}^h \cdots \mathcal{K}_{\phi_n}^h)(x,i;y,j) - (\mathcal{K}_{\phi_1} \cdots \mathcal{K}_{\phi_n})(x,i;y,j) \right|.$$
(B.22)

For arbitrary  $x \in L$  and  $y \in \mathbb{Z}^2$  we find

$$c_{n}(x,y) = \sup_{\substack{\vec{\phi} \in [-\pi,\pi]^{n} \\ i,j \in \{1,2\}}} \left| \sum_{\substack{\vec{A} \in \{L,R\}^{n-1} \\ \text{not all } L}} (\mathcal{K}_{\phi_{1}}^{h} \Pi_{A_{1}} \cdots \Pi_{A_{n-1}} \mathcal{K}_{\phi_{n}}^{h})(x,i;y,j) \right|$$
(B.23)

noting that there are  $2^n-1$  terms in the sum, that each summand contains at least one projection  $\Pi_R$ , and using Lemma B.2,

$$\leq 2^{n} \sup_{k+l=n} \sum_{z \in R} a_{k}(x, z) a_{l}(z, y) \leq (2c)^{n} \kappa^{n-2} \sum_{z \in R} e^{-\xi(|x_{1}-z_{1}|+|z_{1}-y_{1}|+|x_{2}|+|y_{2}|+2|z_{2}|)}$$
(B.24)

since  $|x_1 - z_1| = |x_1| + |z_1|$ , using the triangle inequality and performing the sum over z there is a constant D > 0 such that

$$\leq D^n e^{-\frac{\xi}{2}(\|x-y\|_1 + |x_1| + |x_2|)}$$
 (B.25)

It follows that

$$\left| (\mathcal{U}_{\phi}^{h} - \mathcal{U}_{\phi})(x, i; y, j) \right| \leq (e^{2\pi D} - 1) e^{-\frac{\xi}{2}(\|x - y\|_{1} + |x_{1}| + |x_{2}|)}$$
(B.26)

which proves the claim for  $x \in L$  and  $y \in \mathbb{Z}$  arbitrary. The case where x is arbitrary and  $y \in L$  follows in the same way.

The second bound: We have

$$\mathcal{Q}_{\phi} - \mathscr{P}_{\phi}^{h} = e^{i(\phi + \pi)\Pi_{h}} \mathscr{U}_{-\pi} \mathscr{P} \mathscr{U}_{-\pi}^{*} e^{-i(\phi + \pi)\Pi_{h}} - e^{i\phi\Pi_{h}} \mathscr{P} e^{-i\phi\Pi_{h}}$$
(B.27)

so it is sufficient to establish the bound for  $\mathscr{U}_{-\pi} \mathscr{P} \mathscr{U}_{-\pi}^* - e^{-i\pi\Pi_h} \mathscr{P} e^{i\pi\Pi_h} = \mathscr{U}_{-\pi} \mathscr{P} \mathscr{U}_{-\pi}^* - \mathscr{U}_{-\pi}^h \mathscr{P} (\mathscr{U}_{-\pi}^h)^*$  which follows easily from the first bound.

The third bound: follows from

$$\mathcal{L}_{\phi} - \mathcal{K}_{\phi}^{h} = -[[\Pi_{h}, \mathcal{Q}_{\phi}], (\mathcal{Q}_{\phi} - \mathcal{P}_{\phi}^{h})] - [[\Pi_{h}, (\mathcal{P}_{\phi}^{h} - \mathcal{Q}_{\phi})], \mathcal{P}_{\phi}^{h}]$$
(B.28)

and the second bound.

The fourth bound : follows straightforwardly from the analogous third bound on the generators.  $\hfill\Box$ 

**Proposition B.5.** The difference of projections  $\mathscr{P}_{+} - \mathscr{P}_{-}$  is trace class.

*Proof.* Fix  $x, y \in \mathbb{Z}^2$  and  $i, j \in \{1, 2\}$ . Since  $\mathscr{P}_+ - \mathscr{P}_- = \mathscr{U}_\pi \mathscr{P} \mathscr{U}_\pi^* - \mathscr{U}_{-\pi} \mathscr{P} \mathscr{U}_{-\pi}^*$  we get from Lemma B.3 that

$$|(\mathscr{P}_{+} - \mathscr{P}_{-})(x, i; y, j)| \le c e^{-\xi(||x - y||_1 + \operatorname{dist}(x, l_L))}$$
 (B.29)

for some constants  $c, \xi > 0$ . This shows that  $\mathscr{P}_+ - \mathscr{P}_-$  is localized near the left horizontal axis, but it is also localized near the right horizontal axis. Indeed since  $\mathscr{P}_- = \mathscr{Q}_\pi =$ 

 $\mathcal{V}_{(\pi,-\pi)}\mathcal{Q}_{-\pi}\mathcal{V}_{(\pi,-\pi)}^* = \mathcal{V}_{(\pi,-\pi)}\mathcal{P}_{-}\mathcal{V}_{(\pi,-\pi)}^* \text{ we have } \mathcal{P}_{+} - \mathcal{P}_{-} = \mathcal{U}_{(\pi,-\pi)}\mathcal{P}_{-}\mathcal{U}_{(\pi,-\pi)}^* - \mathcal{V}_{(\pi,-\pi)}\mathcal{P}_{-}\mathcal{V}_{(\pi,-\pi)}^*$  so using the fourth bound of Lemma B.4 we obtain

$$|(\mathscr{P}_{+} - \mathscr{P}_{-})(x, i; y, j)| \le ce^{-\xi(\|x - y\|_{1} + \operatorname{dist}(x, l_{R}))}$$
 (B.30)

for some constants  $c, \xi > 0$ .

Combining the bounds (B.29) and (B.30) we get

$$|(\mathscr{P}_{+} - \mathscr{P}_{-})(x, i; y, j)| \leq c e^{-\frac{\xi}{2}(2\|x - y\|_{1} + \operatorname{dist}(x, l_{L}) + \operatorname{dist}(x, l_{R}))} \leq c e^{-\xi(\|x - y\|_{1} + \|x\|_{1}/2)}.$$
 (B.31)

The trace class claim follows immediately.

### B.2 The spin Hall Chern number as an index of a pair of projections

By Proposition B.5 the difference  $\mathscr{P}_+ - \mathscr{P}_-$  is trace class so the projections  $\mathscr{P}_+$  and  $\mathscr{P}_-$  have a well-defined index of a pair of projections which is given by

$$\operatorname{ind}(\mathscr{P}_{+},\mathscr{P}_{-}) = \operatorname{Tr}\{\mathscr{P}_{+} - \mathscr{P}_{-}\}. \tag{B.32}$$

We will show that this index is equal to the Chern number of the Fermi projection  $\mathscr{P}$ . This Chern number may be expressed as (see for example Eq. 1.4 of [19])

$$Ch(\mathscr{P}) = 2\pi i \operatorname{Tr} \{ \mathscr{P}[[\Pi_h, \mathscr{P}], [\Pi_R, \mathscr{P}]] \}$$
(B.33)

where  $\Pi_h$  and  $\Pi_R$  are the projections onto the upper and right half-planes respectively.

Let  $\Pi(r)$  be the projection onto the box  $\{(x_1, x_2) \in \mathbb{Z}^2 : |x_1|, |x_2| \leq r\}$ , then

$$\operatorname{ind}(\mathscr{P}_{+},\mathscr{P}_{-}) = \lim_{r \to \infty} \operatorname{Tr}\{\Pi(r)(\mathscr{P}_{+} - \mathscr{P}_{-})\Pi(r)\}$$
(B.34)

Since both terms are now separately trace class,

$$= -i \lim_{r \to \infty} \int_{-\pi}^{\pi} d\phi \operatorname{Tr} \{ \mathscr{P}_{\phi} [\mathscr{K}_{\phi}, \Pi(r)] \}$$
 (B.35)

using repeatedly the cyclicity of the trace. Since  $\mathscr{K}_{\phi}$  is localised near the left horizontal axis, where it is equal to  $\mathscr{K}_{\phi}^{h}$  (Lemma B.1 and  $\mathscr{K}_{\phi} = \Pi_{L} \mathscr{K}_{\phi}^{h} \Pi_{L}$ ) this becomes

$$= -i \lim_{r \to \infty} \int_{-\pi}^{\pi} d\phi \operatorname{Tr} \{ \mathscr{P}_{\phi} [\mathscr{K}_{\phi}^{h}, \Pi_{\geqslant -r}] \}$$
 (B.36)

where  $\Pi_{\geqslant -r}$  is the projection onto the half-plane  $\{(x_1, x_2) \in \mathbb{Z}^2 : x_1 \geqslant -r\}$ . Similarly since  $\mathscr{P}_{\phi} - \mathscr{P}_{\phi}^h$  is localised on the right half axis (immediate consequence of the first bound of Lemma B.4) and  $\mathscr{K}_{\phi}^h = -\mathrm{i}[\partial_{\phi}\mathscr{P}_{\phi}^h, \mathscr{P}_{\phi}^h]$ ,

$$= -\lim_{r \to \infty} \int_{-\pi}^{\pi} d\phi \operatorname{Tr} \{ \mathscr{P}_{\phi}^{h} [[\hat{\sigma}_{\phi} \mathscr{P}_{\phi}^{h}, \mathscr{P}_{\phi}^{h}], \Pi_{\geq -r}] \}.$$
 (B.37)

Jacobi's identity reads

$$-[[\partial_{\phi} \mathscr{P}_{\phi}^{h}, \mathscr{P}_{\phi}^{h}], \Pi_{\geqslant -r}] = [[\mathscr{P}_{\phi}^{h}, \Pi_{\geqslant -r}], \partial_{\phi} \mathscr{P}_{\phi}^{h}] + [[\Pi_{\geqslant -r}, \partial_{\phi} \mathscr{P}_{\phi}^{h}], \mathscr{P}_{\phi}^{h}]. \tag{B.38}$$

Since  $\mathscr{P}^h_{\phi}$  is translation invariant in the horizontal direction, the first term is a translate of  $[[\mathscr{P}^h_{\phi},\Pi_R],\partial_{\phi}\mathscr{P}^h_{\phi}]$  and can therefore be replaced by the latter in the trace. The second term is off-diagonal with respect to  $\mathscr{P}^h_{\phi}$  so its contribution vanishes in the trace. Noting that the derivative is given by  $\partial_{\phi}\mathscr{P}^h_{\phi}=\mathrm{i}[\Pi_h,\mathscr{P}^h_{\phi}]$  we therefore obtain

$$\operatorname{ind}(\mathscr{P}_{+},\mathscr{P}_{-}) = \operatorname{i} \int_{-\pi}^{\pi} d\phi \operatorname{Tr}\{\mathscr{P}_{\phi}^{h}[[\mathscr{P}_{\phi}^{h},\Pi_{R}],[\Pi_{h},\mathscr{P}_{\phi}^{h}]]\}, \tag{B.39}$$

and since the integrand does not depend on  $\phi$  (by unitary invariance of the trace and  $[\Pi_h, \Pi_R] = 0$ ), we conclude that

$$\operatorname{ind}(\mathscr{P}_{+},\mathscr{P}_{-}) = -2\pi i \operatorname{Tr} \left\{ \mathscr{P} \left[ \left[ \Pi_{R}, \mathscr{P} \right], \left[ \Pi_{h}, \mathscr{P} \right] \right] \right\}. \tag{B.40}$$

Recalling that we wrote  $\mathscr{P} = \mathscr{P}^{\uparrow}$  and  $\mathscr{P}_{\pm} = \mathscr{P}_{\pm}^{\uparrow}$ , and comparing with (B.33) we have proven

**Proposition B.6.** The difference  $\mathscr{P}_+^{\uparrow} - \mathscr{P}_-^{\uparrow}$  is trace class and the Chern number  $\operatorname{Ch}(\mathscr{P}^{\uparrow})$  of  $\mathscr{P}_+^{\uparrow}$  is equal to the index of a pair of projections  $\operatorname{ind}(\mathscr{P}_+^{\uparrow}, \mathscr{P}_-^{\uparrow})$ .

### References

- [1] A. Altland and M. R. Zirnbauer. Nonstandard symmetry classes in mesoscopic normal-superconducting hybrid structures. *Physical Review B*, 55(2):1142, 1997. arXiv:cond-mat/9602137.
- [2] H. Araki and H. Moriya. Equilibrium statistical mechanics of fermion lattice systems. Reviews in Mathematical Physics, 15(02):93–198, 2003. arXiv:math-ph/0211016.
- [3] J. E. Avron, R. Seiler, and B. Simon. Homotopy and quantization in condensed matter physics. *Physical Review Letters*, 51(1):51, 1983.
- [4] J. E. Avron, R. Seiler, and B. Simon. The index of a pair of projections. *Journal of Functional Analysis*, 120(1):220–237, 1994.
- [5] S. Bachmann, A. Bols, W. De Roeck, and M. Fraas. Quantization of conductance in gapped interacting systems. In *Annales Henri Poincaré*, volume 19, pages 695–708. Springer, 2018. arXiv:1707.06491.
- [6] S. Bachmann, A. Bols, W. De Roeck, and M. Fraas. A many-body index for quantum charge transport. Communications in Mathematical Physics, 375:1249–1272, 2020. arXiv:1810.07351.
- [7] S. Bachmann, A. Bols, W. De Roeck, and M. Fraas. Note on linear response for interacting hall insulators. *Analytic Trends in Mathematical Physics*, 741:23–41, 2020. arXiv:1811.08699.
- [8] S. Bachmann, W. De Roeck, M. Fraas, and T. Jappens. A classification of G-charge Thouless pumps in 1D invertible states. *Communications in Mathematical Physics*, to appear, 2024. arXiv:2204.03763.
- [9] S. Bachmann, W. Dybalski, and P. Naaijkens. Lieb-Robinson bounds, Arveson spectrum and Haag-Ruelle scattering theory for gapped quantum spin systems. *Annales Henri Poincaré*, 17:1737–1791, 2016. arXiv:1412.2970.
- [10] S. Bachmann, S. Michalakis, B. Nachtergaele, and R. Sims. Automorphic equivalence within gapped phases of quantum lattice systems. *Communications in Mathematical Physics*, 309(3):835–871, 2012. arXiv:1102.0842.
- [11] A. Bols and C. Cedzich. Absolutely continuous edge spectrum of topological insulators with an odd time-reversal symmetry. 2022. arXiv:2203.05474.
- [12] A. Bols, J. Schenker, and J. Shapiro. Fredholm homotopies for strongly-disordered 2d insulators. *Communications in Mathematical Physics*, 397(3):1163–1190, 2023. arXiv:2110.07068.

- [13] R. Bott, L. W. Tu, et al. *Differential forms in algebraic topology*, volume 82. Springer, 1982.
- [14] O. Bratteli and D. W. Robinson. Operator algebras and quantum statistical mechanics: Volume 2: Equilibrium States, Models in Quantum Statistical Mechanics. Springer Berlin, Heidelberg, 2013.
- [15] C. Cedzich, T. Geib, F. A. Grünbaum, C. Stahl, L. Velázquez, A. H. Werner, and R. F. Werner. The topological classification of one-dimensional symmetric quantum walks. *Annales Henri Poincaré*, 19:325–383, 2018. arXiv:1611.04439.
- [16] G. De Nittis and K. Gomi. Classification of "quaternionic" Bloch-bundles: Topological quantum systems of type AII. *Communications in Mathematical Physics*, 339:1–55, 2015. arXiv:1404.5804.
- [17] G. De Nittis and H. Schulz-Baldes. Spectral flows associated to flux tubes. *Annales Henri Poincaré*, 17(1):1–35, 2016. arXiv:1405.2054.
- [18] J. Dixmier. Les C\*-algèbres et leurs représentations. Gauthier-Villars, Paris, 1964.
- [19] A. Elgart, G. Graf, and J. Schenker. Equality of the bulk and edge Hall conductances in a mobility gap. *Communications in Mathematical Physics*, 261(2):545–546, 2006. arXiv:math-ph/0409017.
- [20] D. E. Evans and Y. Kawahigashi. Quantum symmetries on operator algebras. Oxford University Press, 1998.
- [21] L. Fidkowski and A. Kitaev. Effects of interactions on the topological classification of free fermion systems. *Physical Review B*, 81(13):134509, 2010. arXiv:0904.2197.
- [22] L. Fidkowski and A. Kitaev. Topological phases of fermions in one dimension. *Physical Review B*, 83(7):075103, 2011. arXiv:1008.4138.
- [23] E. Fonseca, J. Shapiro, A. Sheta, A. Wang, and K. Yamakawa. Two-dimensional time-reversal-invariant topological insulators via fredholm theory. *Mathematical Physics, Analysis and Geometry*, 23:1–22, 2020. arXiv:1908.00910.
- [24] J. Fröhlich, U. M. Studer, and E. Thiran. A classification of quantum Hall fluids. *Journal of statistical physics*, 86:821–897, 1997.
- [25] L. Fu and C. L. Kane. Time reversal polarization and a  $\mathbb{Z}_2$  adiabatic spin pump. *Physical Review B*, 74(19):195312, 2006. arXiv:cond-mat/0606336.
- [26] L. Fu, C. L. Kane, and E. J. Mele. Topological insulators in three dimensions. *Physical Review Letters*, 98(10):106803, 2007. arXiv:cond-mat/0607699.
- [27] G. M. Graf and M. Porta. Bulk-edge correspondence for two-dimensional topological insulators. *Communications in Mathematical Physics*, 324:851–895, 2013. arXiv:1207.5989.
- [28] M. B. Hastings. Lieb-Schultz-Mattis in higher dimensions. *Physical Review B*, 69(10):104431, 2004. arXiv:cond-mat/0305505.
- [29] M. B. Hastings and S. Michalakis. Quantization of Hall conductance for interacting electrons on a torus. *Communications in Mathematical Physics*, 334:433–471, 2015. arXiv:0911.4706.

- [30] M. B. Hastings and X.-G. Wen. Quasiadiabatic continuation of quantum states: The stability of topological ground-state degeneracy and emergent gauge invariance. *Physical Review B*, 72(4):045141, 2005. arXiv:cond-mat/0503554.
- [31] C. L. Kane and E. J. Mele.  $\mathbb{Z}_2$  topological order and the quantum spin hall effect. *Physical Review Letters*, 95(14):146802, 2005. arXiv:cond-mat/0506581.
- [32] C. L. Kane and E. J. Mele. Quantum spin Hall effect in graphene. *Physical Review Letters*, 95(22):226801, 2005. arXiv:cond-mat/0411737.
- [33] A. Kapustin and N. Sopenko. Hall conductance and the statistics of flux insertions in gapped interacting lattice systems. *Journal of Mathematical Physics*, 61(10), 2020. arXiv:2006.14151.
- [34] A. Kapustin and N. Sopenko. Local Noether theorem for quantum lattice systems and topological invariants of gapped states. *Journal of Mathematical Physics*, 63(9), 2022. arXiv:2201.01327.
- [35] A. Kapustin, N. Sopenko, and B. Yang. A classification of invertible phases of bosonic quantum lattice systems in one dimension. *Journal of Mathematical Physics*, 62(8), 2021. arXiv:2012.15491.
- [36] M. Karoubi and C. A. Weibel. On the covering type of a space. L'Enseignement Mathématique, 62(3):457-474, 2017. arXiv:1612.00532.
- [37] T. Kato. On the adiabatic theorem of quantum mechanics. *Journal of the Physical Society of Japan*, 5(6):435–439, 1950.
- [38] A. Kitaev. Periodic table for topological insulators and superconductors. AIP conference proceedings, 1134(1):22–30, 2009. arXiv:0901.2686.
- [39] K. v. Klitzing, G. Dorda, and M. Pepper. New method for high-accuracy determination of the fine-structure constant based on quantized hall resistance. *Physical Review Letters*, 45(6):494, 1980.
- [40] R. B. Laughlin. Quantized Hall conductivity in two dimensions. *Physical Review B*, 23(10):5632, 1981.
- [41] J. Lee. Introduction to Smooth Manifolds, volume 218 of Graduate Texts in Mathematics. Springer, 2003.
- [42] E. H. Lieb and D. W. Robinson. The finite group velocity of quantum spin systems. Communications in mathematical physics, 28(3):251–257, 1972.
- [43] Y.-M. Lu, Y. Ran, and M. Oshikawa. Filling-enforced constraint on the quantized hall conductivity on a periodic lattice. *Annals of Physics*, 413:168060, 2020. arXiv:1705.09298.
- [44] L.-E. Lundberg. Quasi-free "second quantization". Communications in Mathematical Physics, 50:103–112, 1976.
- [45] A. Moon and Y. Ogata. Automorphic equivalence within gapped phases in the bulk. Journal of Functional Analysis, 278(8):108422, 2020. arXiv:1906.05479.
- [46] B. Nachtergaele, R. Sims, and A. Young. Lieb-Robinson bounds, the spectral flow, and stability of the spectral gap for lattice fermion systems. In *Mathematical Problems in Quantum Physics, Proceedings of QMATH13*, volume 717 of *Contemporary Mathematics*, pages 93–115. American Mathematical Society, Providence, RI, 2018. arXiv:1705.08553.

- [47] X.-L. Qi and S.-C. Zhang. Spin-charge separation in the quantum spin Hall state. *Physical Review Letters*, 101(8):086802, 2008. arXiv:0801.0252.
- [48] Y. Ran, A. Vishwanath, and D.-H. Lee. Spin-charge separated solitons in a topological band insulator. *Physical Review Letters*, 101(8):086801, 2008. arXiv:0801.0627.
- [49] S. Ryu, A. P. Schnyder, A. Furusaki, and A. W. Ludwig. Topological insulators and superconductors: tenfold way and dimensional hierarchy. New Journal of Physics, 12(6):065010, 2010. arXiv:0912.2157.
- [50] D. Shale and W. F. Stinespring. Spinor representations of infinite orthogonal groups. Journal of Mathematics and Mechanics, 14(2):315–322, 1965.
- [51] D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. den Nijs. Quantized Hall conductance in a two-dimensional periodic potential. *Physical Review Letters*, 49(6):405, 1982.