

The Galois-equivariant K -theory of finite fields

David Chan and Chase Vogeli

Abstract

We compute the $RO(G)$ -graded equivariant algebraic K -groups of a finite field with an action by its Galois group G . Specifically, we show these K -groups split as the sum of an explicitly computable term and the well-studied $RO(G)$ -graded coefficient groups of the equivariant Eilenberg–MacLane spectrum $H\mathbb{Z}$. Our comparison between the equivariant K -theory spectrum and $H\mathbb{Z}$ further shows they share the same Tate spectra and geometric fixed point spectra. In the case where G has prime order, we provide an explicit presentation of the equivariant K -groups.

Contents

| | | |
|----------|--|-----------|
| 1 | Introduction | 2 |
| 1.1 | Main result | 2 |
| 1.2 | Overview of key ideas | 3 |
| 1.3 | Future directions | 5 |
| 2 | The Tate diagram | 6 |
| 3 | Equivariant algebra | 8 |
| 3.1 | Mackey functors | 8 |
| 3.2 | Extensions of Mackey functors | 9 |
| 4 | The Galois action on K-theory | 10 |
| 4.1 | Galois cohomology Mackey functors | 10 |
| 4.2 | The norm cofiber sequence | 14 |
| 5 | Proofs of the main theorems | 18 |
| 5.1 | Comparison of Tate diagrams | 19 |
| 5.2 | The $RO(G)$ -graded K -groups of finite fields | 20 |
| 5.3 | Geometric fixed points | 23 |
| 6 | Extensions of prime degree | 25 |
| 6.1 | Quadratic extensions | 25 |
| 6.2 | Extensions of odd prime degree | 29 |

1 Introduction

Algebraic K -theory is an important invariant of rings which provides a natural home for constructions in a range of subjects, from number theory to geometric topology. Classically, low dimensional K -groups of rings were defined first in terms of concrete algebraic constructions in work of Grothendieck, Bass, Milnor, and others. Over time it was realized that these groups fit into a larger picture: there should be a sequence of such groups, indexed on the natural numbers. This idea was first implemented in the work of Quillen, who constructed a space whose homotopy groups agreed with the already-known K -groups and whose higher homotopy groups give the correct generalization of “higher algebraic K -theory.” Using this machinery, Quillen gave a complete computation of the higher algebraic K -theory of finite fields [Qui72].

Despite Quillen’s success in computing the algebraic K -theory of finite fields, the task of computing higher algebraic K -theory for other rings remains difficult. Indeed, the algebraic K -theory of the integers is still not entirely known, and its computation would resolve the longstanding Kummer–Vandiver conjecture in number theory [Kur92].

The work of this paper exists in the context of *equivariant algebraic K -theory*, a variant of algebraic K -theory defined for rings with an action by a group G . The most familiar examples of such rings come to us from Galois theory, where the group G is the Galois group of a field extension. Work of Merling [Mer17] provides a construction of algebraic K -theory for rings with G -action which produces a *genuine G -spectrum* – an enhancement of a spectrum with G -action – with many desirable properties. In particular, Merling shows that this construction is naturally related to the (now proven) Quillen–Lichtenbaum conjecture and provides a natural home for studying the K -theory of Galois extensions.

In addition to Merling’s work, Barwick and Barwick–Glasman–Shah provide a different of construction of equivariant algebraic K -theory using the language of ∞ -categories and spectral Mackey functors [Bar17, BGS20]. Malkiewich and Merling, in on-going work with Goodwillie and Igusa, are using techniques of equivariant algebraic K -theory to prove an equivariant refinement of Waldhausen’s A -theory and the stable parametrized h -cobordism theorem [MM19, MM22, GIMM23]. A connection between equivariant algebraic K -groups, equivariant A -theory, and Wall’s finiteness obstruction is provided in work of the first-named author, Calle, and Mejia [CCM23].

1.1 Main result

One feature of genuine G -spectra is that their homotopy groups are graded on $RO(G)$, the real orthogonal representation ring of the group G . The ordinary \mathbb{Z} -grading is recovered by considering the trivial G -representations of various dimensions. The computation of $RO(G)$ -graded equivariant homotopy groups, even in tractable cases like Eilenberg–MacLane spectra, can be challenging. This work can be quite valuable, however. For example, in Hill, Hopkins, and Ravenel’s solution to the Kervaire invariant one problem they make critical use of these $RO(G)$ -graded homotopy groups for the group $G = C_8$ [HHR16].

Let us fix a field k with G -action and denote the equivariant algebraic K -theory of k by $K_G(k)$. Because $K_G(k)$ is a genuine G -spectrum, the algebraic K -groups are naturally

graded on the group $RO(G)$. Our main result extends Quillen’s computation of the K -groups of finite fields to a computation of the equivariant algebraic K -theory of finite fields with action by their Galois groups.

Theorem A (Theorem 5.2.3 and Proposition 5.2.1). *Let k/\mathbb{F}_q be any finite extension of finite fields with Galois group G , and let V be any finite-dimensional virtual real orthogonal G -representation. Then the equivariant algebraic K -groups of k split as*

$$\pi_V^G(K_G(k)) \cong \begin{cases} H^0(G; \pi_{|V|}(K(k))) \oplus \pi_V^G(H\mathbb{Z}), & |V| > 0, \\ \pi_V^G(H\mathbb{Z}), & |V| \leq 0, \end{cases}$$

where $H\mathbb{Z}$ is the equivariant Eilenberg–MacLane spectrum on the constant Mackey functor \mathbb{Z} .

This is one of the first systematic computations of equivariant algebraic K -groups. Indeed, to our knowledge, the only prior computation of $RO(G)$ -graded equivariant K -groups to appear in the literature is in the work of Elmanto and Zhang [EZ24]. There, the authors give a partial computation of the $RO(C_2)$ -graded K -groups of finite quadratic field extensions, and show how these groups are related to the study of Artin L -functions.

1.2 Overview of key ideas

Our computation of the $RO(G)$ -graded K -groups of finite fields hinges on a comparison of $K_G(k)$ with $H\mathbb{Z}$, the equivariant Eilenberg–MacLane spectrum on the constant \mathbb{Z} -Mackey functor. The choice of \mathbb{Z} comes from the fact that there is an isomorphism of Mackey functors $\pi_0(K_G(k)) \cong \mathbb{Z}$. Accordingly, there is always a truncation map of genuine G -spectra $K_G(k) \rightarrow H\mathbb{Z}$ which realizes this isomorphism.

Our analysis of this map begins with the observation that it is fully computable after applying a construction called geometric completion. If X is a G -spectrum, its *geometric completion* is the G -spectrum

$$X^h = F(EG_+, X).$$

The G -fixed points of X^h are the familiar *homotopy fixed points* X^{hG} of X . As such, this is computable via a homotopy fixed-point spectral sequence which begins with group cohomology. We directly compute the following.

Theorem B (Proposition 5.2.1). *For any extension of finite fields k/\mathbb{F}_q with Galois group G , the fiber of the truncation map on geometric completions has $RO(G)$ -graded homotopy groups*

$$\pi_V^G \text{hofib}(K_G(k)^h \rightarrow H\mathbb{Z}^h) \cong \begin{cases} H^0(G; K_{|V|}(k)), & |V| > 0, \\ 0 & \text{otherwise.} \end{cases}$$

Denote by

$$F = \text{hofib}(K_G(k) \rightarrow H\mathbb{Z})$$

the fiber of the original truncation map. The proof of Theorem A proceeds by showing that F is in fact equivalent to the previously computed fiber.

Theorem C (Theorem 5.0.1). *For any extension of finite fields k/\mathbb{F}_q with Galois group G there is a homotopy pullback of genuine G -spectra*

$$\begin{array}{ccc} K_G(k) & \longrightarrow & H\underline{\mathbb{Z}} \\ \downarrow & & \downarrow \\ K_G(k)^h & \longrightarrow & H\underline{\mathbb{Z}}^h. \end{array}$$

Consequently, there is an equivalence of fibers

$$F = \text{hofib}(K_G(k) \rightarrow H\underline{\mathbb{Z}}) \simeq \text{hofib}(K_G(k)^h \rightarrow H\underline{\mathbb{Z}}^h).$$

In conclusion, we have a fiber sequence of G -spectra

$$F \rightarrow K_G(k) \rightarrow H\underline{\mathbb{Z}} \tag{1.2.1}$$

where we completely understand the homotopy groups of the fiber. This fiber sequence gives the decomposition in Theorem A.

Our work reduces the $RO(G)$ -graded K -groups of finite fields to the computation of the $RO(G)$ -graded coefficient ring of $H\underline{\mathbb{Z}}$. While these groups are complicated, they are known in many cases where G is a finite cyclic group – the only possibility for a Galois extension of finite fields. The first such computation is due to Stong for $G = C_p$, and is recorded in Ferland–Lewis [FL04]. The computation for $G = C_{2^n}$ plays an important role in the work of Hill–Hopkins–Ravenel [HHR16, HHR17]. Work of Zeng handles the case of $G = C_{p^2}$ [Zen18], and work of Georgakopoulos revisits this case as well as providing a computer program for $G = C_{p^n}$ [Geo21]. A recent preprint of Basu–Dey addresses the general case of $G = C_n$ [BD24]. In Section 6 we specialize to the case $G = C_\ell$, for ℓ a prime, and write out the K -groups explicitly.

1.2.1 Proof of Theorem C

Let us briefly outline our approach to proving Theorem C. The main method we employ is a comparison of the *Tate squares* associated to the G -spectra $K_G(k)$ and $H\underline{\mathbb{Z}}$. To any genuine G -spectrum X there is a functorially associated homotopy pullback diagram of the form

$$\begin{array}{ccc} X & \longrightarrow & \tilde{X} \\ \downarrow & & \downarrow \\ X^h & \longrightarrow & X^t \end{array}$$

where X^t is called the Tate G -spectrum of X . As the name suggests, the homotopy groups of X^t can be computed using a spectral sequence whose E_2 -page is computed using Tate cohomology. The map $K_G(k) \rightarrow H\underline{\mathbb{Z}}$ induces a map on Tate diagrams and the following theorem is proved using direct computation.

Theorem D (Proposition 4.2.4 and Proposition 5.1.5). *For any extension of finite fields k/\mathbb{F}_q with Galois group G the map $K_G(k) \rightarrow H\underline{\mathbb{Z}}$ induces equivalences of genuine G -spectra*

$$\widetilde{K_G(k)} \rightarrow \widetilde{H\underline{\mathbb{Z}}} \quad \text{and} \quad K_G(k)^t \rightarrow H\underline{\mathbb{Z}}^t.$$

This result, together with the naturality of the Tate square, is enough to prove Theorem C.

1.2.2 Geometric fixed points

As a further demonstration of our methods, we show $K_G(k)$ and $H\underline{\mathbb{Z}}$ have the same geometric fixed points. Using Theorem A, and the computation of the homotopy groups of the fiber, we prove the following.

Theorem E (Corollary 5.3.4 and Propositions 5.3.5 and 5.3.6). *There is an equivalence of spectra $\Phi^G(K_G(k)) \rightarrow \Phi^G(H\underline{\mathbb{Z}})$. The homotopy groups of these spectra are given as graded rings by*

$$\pi_*(\Phi^G(H\underline{\mathbb{Z}})) \cong \begin{cases} \mathbb{Z}/p[x], & G \cong C_{p^n}, n > 0, \\ 0, & \text{else,} \end{cases}$$

where $|x| = 2$.

We note that the computation of the groups $\pi_*(\Phi^G(H\underline{\mathbb{Z}}))$ is well-known to experts, but we did not know a reference in the literature so the computation is given in Propositions 5.3.5 and 5.3.6 below.

1.3 Future directions

We end the introduction with a short discussion of possible future computations for equivariant algebraic K -theory of other Galois extensions. To any Galois extension E/F of fields there is a natural map

$$K(F) \rightarrow K(E)^{hG}.$$

Underlying the analysis in this paper is the fact that for finite fields, this map is a connective cover [Qui72, final remark]. This behavior, often called Galois descent, is very atypical of Galois extensions of rings; indeed we are aware of no other examples where Galois descent holds for field extensions. In particular, a direct application of the techniques in this paper is unlikely to apply in other examples.

On the other hand, the work of Clausen–Mathew–Naumann–Noel [CMNN20] (see also [Tho85, TT90]) shows that algebraic K -theory satisfies Galois descent after chromatic localization at the spectra $T(n)$. Given this, we expect that after $T(n)$ -localization it should be possible to use the ideas of this paper for further computations of equivariant algebraic K -theory for other Galois ring extensions. We intend to explore this idea in future work.

Outline

The paper is organized as follows. In Section 2 we recall the Tate diagram associated to a G -spectrum, which serves as a guiding framework to our work. In Section 3 we recall the definition of a Mackey functor and prove results concerning extensions of Mackey functors. In Section 4 we make a number of group (co)homology computations which provide the input to homotopy fixed point and Tate spectral sequences. These computations set the stage for the proofs of the main theorems which are in Section 5. We conclude the paper in Section 6 with explicit computations in the case G is a cyclic group of prime order.

Acknowledgments

The authors would like to thank Liam Keenan, Tomas Mejia Gomez, J.D. Quigley, and Ben Spitz for helpful conversations. We would additionally like to thank Maxine Calle, Teena Gerhardt, Maxiemiilien Péroux, and Inna Zakharevich for reading parts of earlier drafts of this paper. The first-named author was partially supported by NSF grant DMS-2135960.

2 The Tate diagram

In this short section we recall the Tate diagram from [GM95], which forms the basis for our work in later sections. For any finite group G let EG be a contractible CW-complex with free G -action. An explicit model is given by taking the unit sphere in the G -representation obtained by taking the infinite direct sum of copies of the reduced regular representation of G . Define a space \widetilde{EG} by the cofiber sequence

$$EG_+ \xrightarrow{\varepsilon} S^0 \rightarrow \widetilde{EG}$$

and note that for any G -spectrum X we have an induced cofiber sequence

$$EG_+ \wedge X \rightarrow X \rightarrow \widetilde{EG} \wedge X$$

which is natural in X .

We write $X^h := F(EG_+, X)$, where F denotes the internal mapping spectrum, and call this spectrum the *geometric completion* of X . The map ε induces a map $X \cong F(S^0, X) \rightarrow X^h$ which gives a map on cofiber sequences

$$\begin{array}{ccccc} EG_+ \wedge X & \longrightarrow & X & \longrightarrow & \widetilde{EG} \wedge X \\ \downarrow & & \downarrow & & \downarrow \\ EG_+ \wedge X^h & \longrightarrow & X^h & \longrightarrow & \widetilde{EG} \wedge X^h \end{array}$$

called the *Tate diagram*.

Theorem 2.0.1 ([GM95, Proposition 1.2]). *The left vertical map in the Tate diagram is an equivalence. Equivalently, the right square is a homotopy pullback of G -spectra.*

For the remainder of this paper we use the following abbreviations:

$$\begin{aligned} X_h &:= EG_+ \wedge X \simeq EG_+ \wedge X^h \\ X^t &:= \widetilde{EG} \wedge X^h \\ \widetilde{X} &:= \widetilde{EG} \wedge X \end{aligned}$$

so that the Tate diagram takes the abbreviated form

$$\begin{array}{ccccc} X_h & \longrightarrow & X & \longrightarrow & \widetilde{X} \\ \simeq \downarrow & & \downarrow & \lrcorner & \downarrow \\ X_h & \longrightarrow & X^h & \longrightarrow & X^t \end{array} .$$

The bottom row of the Tate diagram is called the *norm cofiber sequence*. The three G -spectra in the norm cofiber sequence come equipped with spectral sequences which compute their $RO(G)$ -graded homotopy groups. These spectral sequences are recalled in Section 4.2 and used to compute the homotopy groups appearing in the norm cofiber sequence for equivariant algebraic K -theory spectra. From there, the computation of $RO(G)$ -graded K -groups is achieved by comparing the top and bottom rows of the Tate diagram.

We end this section with a lemma we will need in Section 5.

Lemma 2.0.2. *Suppose that $f: X \rightarrow Y$ is a map of genuine G -spectra which induces equivalences $\widetilde{X} \rightarrow \widetilde{Y}$ and $X^t \rightarrow Y^t$. Then the square*

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow & & \downarrow \\ X^h & \xrightarrow{f^h} & Y^h \end{array}$$

is a homotopy pullback square.

Proof. Consider the commutative diagram

$$\begin{array}{ccccc} X_h & \longrightarrow & X & & \\ \simeq \downarrow & \searrow^{f_h} & \downarrow & \searrow^f & \\ X_h & \longrightarrow & X^h & \longrightarrow & Y \\ \downarrow & \searrow^{f_h} & \downarrow & \searrow^{f^h} & \\ X_h & \longrightarrow & X^h & \longrightarrow & Y^h \end{array}$$

obtained by naturality of the Tate square. It suffices to show that the induced map $\text{hofib}(f) \rightarrow \text{hofib}(f^h)$ is a weak equivalence. Taking fibers of the labeled maps in the

cube above produces a commutative square

$$\begin{array}{ccc} \mathrm{hofib}(f_h) & \xrightarrow{\simeq} & \mathrm{hofib}(f) \\ \downarrow \simeq & & \downarrow \\ \mathrm{hofib}(f_h) & \xrightarrow{\simeq} & \mathrm{hofib}(f^h) \end{array}$$

where the horizontal maps are equivalences by the assumptions on the map f . Thus the map $\mathrm{hofib}(f) \rightarrow \mathrm{hofib}(f^h)$ is a weak equivalence which completes the proof. \square

3 Equivariant algebra

In this section we lay the algebraic groundwork for the computation of $RO(G)$ -graded K -groups. The computation is achieved using tools from equivariant algebra. We first recall the definition of Mackey functors and some relevant constructions in Section 3.1. In Section 3.2, we prove a useful result for showing extensions of Mackey functors are trivial in Proposition 3.2.1.

3.1 Mackey functors

For a finite group G , the *Burnside category* \mathcal{A}^G is the additive category whose objects are finite G -sets. The abelian group $\mathcal{A}^G(X, Y)$ is the group completion of the commutative monoid of spans X to Y under disjoint union. The category Mack_G of *Mackey functors* for the group G is the category of additive functors $\mathcal{A}^G \rightarrow \mathrm{Ab}$.

It can often be helpful to give an explicit description of what data is entailed by a Mackey functor. In this paper, we will only be interested in the case where G is abelian (in fact cyclic), where we can give the following ‘‘piecemeal’’ definition.

Lemma 3.1.1 ([Maz13, Definition 1.1.2]). *A Mackey functor \underline{M} for an abelian finite group G is equivalent to the data of*

- an abelian group $\underline{M}(G/H)$ for each $H \leq G$,
- a transfer homomorphism $T_H^K: \underline{M}(G/H) \rightarrow \underline{M}(G/K)$ for $H < K \leq G$,
- a restriction homomorphism $R_H^K: \underline{M}(G/K) \rightarrow \underline{M}(G/H)$ for $H < K \leq G$,
- an action of the Weyl group $W_K H = K/H$ on $\underline{M}(G/H)$ for $H < K \leq G$,

which are subject to the relations

- (a) $T_H^K T_L^H = T_L^K$ and $R_L^H R_H^K = R_L^K$ for $L < H < K \leq G$,
- (b) $T_H^K(\gamma \cdot x) = T_H^K(x)$ for $x \in \underline{M}(G/H), \gamma \in W_K H$,
- (c) $\gamma \cdot R_H^K(x) = R_H^K(x)$ for $x \in \underline{M}(G/K), \gamma \in W_K H$,
- (d) $R_{H'}^K T_H^K(x) = \sum_{\gamma \in W_K H'} \gamma \cdot T_{H \cap H'}^H(x)$ for $x \in \underline{M}(G/(H \cap H'))$

Remark 3.1.2. It is also possible to define Mackey functors for nonabelian G , but it necessarily involves the additional data of maps between conjugate pairs of subgroups, and a more complicated summation formula over double cosets [TW95, Section 2].

The Weyl group action on $\underline{M}(G/e)$ turns $\underline{M}(G/e)$ into a G -module for any G -Mackey functor \underline{M} . The functor taking a Mackey functor \underline{M} to the G -module $\underline{M}(G/e)$ admits left and right adjoints

$$\begin{array}{ccc} & L & \\ \swarrow & & \searrow \\ \text{Mack}_G & \longrightarrow & \text{Mod}_{\mathbb{Z}G} \\ \nwarrow & & \nearrow \\ & R & \end{array}$$

which we denote by L and R , respectively. For a G -module P , $L(P)$ is its *orbit Mackey functor* and $R(P)$ is its *fixed point Mackey functor*. Explicitly, we have

$$L(P)(G/H) = P_H, \quad \text{and} \quad R(P)(G/H) = P^H.$$

There is a natural transformation $N_P: L(P) \rightarrow R(P)$ which at each level G/H is given by the norm map $P_H \rightarrow P^H$,

$$x \mapsto \sum_{h \in H} h \cdot x$$

for any $x \in P_H$.

3.2 Extensions of Mackey functors

We prove an Ext vanishing result which will be used to resolve extension problems later.

Proposition 3.2.1. *For any G -Mackey functor N with $M(G/e) = 0$ and any a G -module N ,*

$$\text{Ext}^1(M, R(N)) = 0.$$

Proof. We will show that any short exact sequence of Mackey functors

$$0 \rightarrow R(N) \xrightarrow{f} P \rightarrow M \rightarrow 0$$

must be split. Note that since $M(C_n/e) = 0$, we must have $f_{C_n/e}: R(N)(C_n/e) \rightarrow P(C_n/e)$ is an isomorphism. Let $\varphi: P(C_n/e) \rightarrow R(N)(C_n/e)$ denote the inverse.

Since R is the right adjoint to evaluation at C_n/e , the map φ uniquely determines a map of Mackey functors $\Phi: P \rightarrow R(N)$ which is given by φ at level C_n/e . Thus the composite $\Phi \circ f$ is the identity on level C_n/e , and therefore must be the identity on the entire Mackey functor. Thus every such short exact sequence is split and $\text{Ext}^1(M, R(N)) = 0$. \square

A dual argument can be applied to see that $\text{Ext}^1(L(N), M) = 0$ for any N .

4 The Galois action on K -theory

In [Mer17], Merling defines the algebraic K -theory of rings with G -action. This invariant assigns to every ring R with action of a finite group G a genuine G -spectrum $K_G(R)$. A theorem of Merling tells us how K_G behaves when our G -rings arise from Galois extensions; we refer the reader to Merling's paper [Mer17] for proofs and further discussion.

Theorem 4.0.1 ([Mer17, Theorem 1.2, (4) and (6)]). *For a finite Galois extension of fields E/F with $G = \text{Gal}(E/F)$ there are equivalences of spectra*

$$K(F) \simeq K_G(E)^G$$

where $K(F)$ is the non-equivariant algebraic K theory of F .

We are interested in Merling's construction when the ring $R = k$ is a finite field which is a finite Galois extension of a finite field with G the Galois group. Let p be a prime, let $q = p^r$ for some $r \geq 1$, and let \mathbb{F}_q be the field with q -elements. Quillen showed \mathbb{F}_q has its higher K -groups concentrated in odd degrees, where we have

$$K_{2i-1}(\mathbb{F}_q) \cong \mathbb{Z}/(q^i - 1)$$

for $i > 0$ [Qui72, Theorem 8(i)]. We will be interested in the case where $k = \mathbb{F}_{q^n}$ is a degree n extension of \mathbb{F}_q . The Galois group $G = \text{Gal}(k/\mathbb{F}_q)$ is cyclic of order n , generated by the Frobenius automorphism φ specified by $\varphi(x) = x^q$ of k . Quillen furthermore calculated the action of G on the higher K -groups.

Theorem 4.0.2 ([Qui72, Theorem 8(iii)]). *Let $k = \mathbb{F}_{q^n}$ and let $G = \text{Gal}(\mathbb{F}_{q^n}, \mathbb{F}_q)$. For $i > 0$, the action of the Frobenius automorphism φ on the group*

$$K_{2i-1}(k) = \mathbb{Z}/(q^{ni} - 1)$$

is multiplication by q^i .

In this section, we use these two results to perform the calculations underlying our main results. Our goal is to explicitly compute the $RO(G)$ -graded homotopy Mackey functors of the terms appearing in the norm cofiber sequence for $K_G(k)$. In Section 4.1, we enumerate all the possible Mackey functors appearing on the E^2 -pages of spectral sequences converging to these terms, and in Section 4.2, we analyze these spectral sequences.

4.1 Galois cohomology Mackey functors

In this section, we determine the Mackey functors appearing on the E^2 -pages of the homotopy orbit and homotopy fixed point spectral sequences for $K_G(k)$. The E_2 -pages of these spectral sequences are recalled in Section 4.2 below. We first extend Theorem 4.0.2 to a computation of the G -modules $\pi_V^e K_G(k)$ for $V \in RO(G)$, and then we obtain Mackey functors by applying the functors

$$L, R: \text{Mod}_{\mathbb{Z}G} \rightarrow \text{Mack}_G$$

defined in the last subsection.

Since these functors are given by taking orbits and fixed points respectively, they form the input for the filtration-zero line of the homotopy orbit and fixed point spectral sequences. The remainder of the input for these spectral sequences amounts to computing the higher group (co)homology of these G -modules. Since G is cyclic, its (co)homology with coefficients in any G -module P is 2-periodic, and is fully determined by the norm map $N: P_G \rightarrow P^G$. Specifically, we have

$$H_s(G; P) \cong \begin{cases} P_G & s = 0, \\ \text{coker } N_P & s > 0, s \text{ odd}, \\ \ker N_P & s > 0, s \text{ even}, \end{cases} \quad H^s(G; P) \cong \begin{cases} P^G & s = 0, \\ \ker N_P & s > 0, s \text{ odd}, \\ \text{coker } N_P & s > 0, s \text{ even}. \end{cases}$$

Thus, we need to compute the kernel and cokernel of the norm transformation $N: L \Rightarrow R$.

Recall that $K_G(k)^e \simeq K(k)$ [Mer17, Theorem 6.4], so $\pi_V^e K_G(k) \cong K_{|V|}(k)$ with G -action determined by Quillen's computation, Theorem 4.0.2. The action of G on $K_0(k) \cong \mathbb{Z}$ is necessarily trivial, as there are no nontrivial ring automorphisms of \mathbb{Z} . Note, however, that the action of G on $\pi_V^e(K_G(k))$ might be non-trivial even when $|V| = 0$, since the identification of abelian groups $\pi_V^e K_G(k) \cong K_{|V|}(k)$ need not respect the G -action.

If n is even, it is possible for $G = C_n$ to act by a "twist" of -1 on these G -modules. We use the notation \mathbb{Z}^σ to denote the sign representation, i.e., the group \mathbb{Z} with G acting by -1 . For a G -module M , we write

$$M^\sigma = M \otimes \mathbb{Z}^\sigma$$

for M twisted by the sign action. We write $K_{2i-1}(k)$ for the G -module $\mathbb{Z}/(q^{ni} - 1)$ with the implicit G -action by q^i as in Theorem 4.0.2.

Notation 4.1.1. We call a G -representation V orientation-preserving (resp. reversing) if the induced map on S^V is orientation-preserving (resp. reversing). By extending linearly to virtual representations, this gives a well defined group homomorphism $RO(G) \rightarrow \mathbb{Z}/2$ where the kernel is all orientation preserving virtual representations.

Lemma 4.1.2. *Let k be any finite field. For $V \in RO(G)$, G -modules $\pi_V^e K_G(k)$ are given by*

$$\pi_V^e K_G(k) \cong \begin{cases} \mathbb{Z} & |V| = 0, V \text{ orientation-preserving}, \\ \mathbb{Z}^\sigma & |V| = 0, V \text{ orientation-reversing}, \\ K_{2i-1}(k) & |V| = 2i - 1 > 0, V \text{ orientation-preserving}, \\ K_{2i-1}(k)^\sigma & |V| = 2i - 1 > 0, V \text{ orientation-reversing}, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. The action on the homotopy groups is given by the conjugation action on maps. Since the action of G on S^V is the sign representation when V is orientation-reversing and is the trivial representation when V is orientation preserving the claimed computation follows from Theorem 4.0.2. \square

Writing M for any of the G -modules in Lemma 4.1.2, the rest of this section is devoted to computing the Mackey functors in the exact sequence:

$$0 \rightarrow \ker(N) \rightarrow L(M) \xrightarrow{N} R(M) \rightarrow \operatorname{coker}(N) \rightarrow 0.$$

We treat the four cases of Lemma 4.1.2 in order. A summary of these calculations can be found in Table 1.

4.1.1 The trivial G -module \mathbb{Z} .

We denote by

$$\begin{aligned} R(\mathbb{Z}) &= \underline{\mathbb{Z}} = \square, \\ L(\mathbb{Z}) &= \dot{\square}, \end{aligned}$$

the fixed points and orbit Mackey functors, respectively, of trivial C_n -module \mathbb{Z} . We define the Mackey functor \circ as the cokernel of the norm map. This is depicted by the short exact sequence of Mackey functors

$$\begin{array}{ccccccc} 0 & \longrightarrow & \dot{\square} & \longrightarrow & \square & \longrightarrow & \circ & \longrightarrow & 0 \\ C_n/C_a : & & 0 & \longrightarrow & \mathbb{Z} & \xrightarrow{a} & \mathbb{Z} & \twoheadrightarrow & \mathbb{Z}/a & \longrightarrow & 0 \\ & & \left(\begin{array}{c} \uparrow \\ \downarrow \end{array}\right) & & a/b \left(\begin{array}{c} \uparrow \\ \downarrow \end{array}\right) & & 1 \left(\begin{array}{c} \uparrow \\ \downarrow \end{array}\right) & & a/b & & 1 \left(\begin{array}{c} \uparrow \\ \downarrow \end{array}\right) & & a/b & & \left(\begin{array}{c} \uparrow \\ \downarrow \end{array}\right) \\ C_n/C_b : & & 0 & \longrightarrow & \mathbb{Z} & \xrightarrow{b} & \mathbb{Z} & \twoheadrightarrow & \mathbb{Z}/b & \longrightarrow & 0. \end{array}$$

4.1.2 The sign representation \mathbb{Z}^σ .

In the case where n is even, there is the C_n -module \mathbb{Z}^σ . We denote by

$$\begin{aligned} R(\mathbb{Z}^\sigma) &= \overline{\square}, \\ L(\mathbb{Z}^\sigma) &= \dot{\square}, \end{aligned}$$

the fixed points and orbit Mackey functors, respectively. A subgroup $C_m \leq C_n$ acts nontrivially on \mathbb{Z}^σ if and only if its index n/m is odd. The norm map $L_- \rightarrow R_-$ has both a nontrivial kernel \bullet and cokernel $\overline{\bullet}$, which have components given by:

$$\begin{aligned} \dot{\square}(C_n/C_m) &\cong \begin{cases} \mathbb{Z}/2 & n/m \text{ odd} \\ \mathbb{Z}^\sigma & n/m \text{ even,} \end{cases} & \overline{\square}(C_n/C_m) &\cong \begin{cases} 0 & n/m \text{ odd} \\ \mathbb{Z}^\sigma & n/m \text{ even,} \end{cases} \\ \bullet(C_n/C_m) &\cong \begin{cases} \mathbb{Z}/2 & n/m \text{ odd} \\ 0 & n/m \text{ even,} \end{cases} & \overline{\bullet}(C_n/C_m) &\cong \begin{cases} 0 & n/m \text{ odd} \\ (\mathbb{Z}/m)^\sigma & n/m \text{ even.} \end{cases} \end{aligned}$$

4.1.3 Higher K -groups with the Galois action.

Recall that the G -module $K_{2i-1}(k)$ is given by $\mathbb{Z}/(q^{ni} - 1)$ with G acting by q^i .

Lemma 4.1.3. *Let $k = \mathbb{F}_{q^n}$ and $G = C_n$. The norm map*

$$N: L(K_{2i-1}(k)) \rightarrow R(K_{2i-1}(k))$$

is an isomorphism of C_n -Mackey functors, and thus has trivial kernel and cokernel. The Mackey functor

$$\ominus^i = L(K_{2i-1}(k)) \cong R(K_{2i-1}(k))$$

is specified by $\ominus^i(C_n/C_m) = \mathbb{Z}/(q^{ni/m} - 1)$. For $C_a < C_b \leq C_n$, the transfers $T_{C_a}^{C_b}$ map $1 \mapsto 1$, and the restrictions $R_{C_a}^{C_b}: \mathbb{Z}/(q^{ni/b} - 1) \rightarrow \mathbb{Z}/(q^{ni/a} - 1)$ map

$$1 \mapsto 1 + q^{ni/b} + q^{2ni/b} + \dots + q^{(ni/a)-(ni/b)}.$$

The Weyl group action of C_n/C_m on $\mathbb{Z}/(q^{ni/m} - 1)$ is given by multiplication by $q^{ni/m}$.

Proof. The coinvariants and invariants of K_{2i-1} relative to the subgroup

$$C_m = \langle q^{n/m} \rangle \leq C_n,$$

are computed as follows.

The coinvariants $K_{2i-1}(k)_{C_m}$ are obtained by adding the relations $q^{ni/m}x = x$ for all x to the group $\mathbb{Z}/(q^{ni} - 1)$. In other words, this is the quotient by the subgroup generated by $q^{ni/m} - 1 \in \mathbb{Z}/(q^{ni} - 1)$. This quotient is cyclic of order $q^{ni/m} - 1$, generated by the coset of $1 \in \mathbb{Z}/(q^{ni} - 1)$.

The invariants $K_{2i-1}(k)^{C_m}$ are the subgroup of $\mathbb{Z}/(q^{ni} - 1)$ of elements x such that $q^{ni/m}x = x$. In other words, it is the subgroup annihilated by $q^{ni/m} - 1$, which is cyclic of order $q^{ni/m} - 1$ and is generated by the element

$$\frac{q^{ni} - 1}{q^{ni/m} - 1} \in \mathbb{Z}/(q^{ni} - 1).$$

Thus, we see that the invariants and coinvariants are abstractly isomorphic. Moreover, the norm map is multiplication by

$$N = 1 + q^{ni/m} + q^{2ni/m} + \dots + q^{(m-1)ni/m} = \frac{q^{ni} - 1}{q^{ni/m} - 1}.$$

Under the above identifications, this takes the generator of $K_{2i-1}(k)_{C_m}$ to a generator of $K_{2i_1}(k)^{C_m}$, from which the claim follows. \square

4.1.4 Higher K -groups with the twisted Galois action.

In the last section we computed the group (co)homology of the G -modules $K_{2i-1}(k)$ where k is a finite field. In this section we state the group (co)homology of the twisted G -modules $K_{2i-1}(k)^\sigma = K_{2i-1}(k) \otimes \mathbb{Z}^\sigma$. This case is almost identical to the last, but with $(-1)^{n/m} q^{ni/m}$ in place of $q^{ni/m}$. We leave the details to the reader.

Lemma 4.1.4. *Recall that C_n acts on $K_{2i-1}(k)^\sigma \cong \mathbb{Z}/(q^{ni} - 1)$ by $-q^i$. For $C_m \subset C_n$ we have*

- (a) *the coinvariants $(K_{2i-1}(k)^\sigma)_{C_m}$ are cyclic of order $(-1)^{n/m} q^{ni/m} - 1$,*
- (b) *the invariants $(K_{2i-1}(k)^\sigma)^{C_m}$ are cyclic of order $(-1)^{n/m} q^{ni/m} - 1$,*
- (c) *the norm map*

$$N : (K_{2i-1}(k)^\sigma)_{C_m} \rightarrow (K_{2i-1}(k)^\sigma)^{C_m}$$

is an isomorphism.

Thus we have an isomorphism of Mackey functors $L(K_{2i-1}(k)^\sigma) \cong R(K_{2i-1}(k)^\sigma)$ and we write \oplus^i for this Mackey functor.

4.1.5 Summary

The results of the computations above are summarized in Table 1. For space reasons, we abbreviate ‘‘orientation preserving’’ (resp. reversing) to o.p. (resp. o.r.).

| | $\underline{K}_V(k)$ | $L(\underline{K}_V(k))$ | $R(\underline{K}_V(k))$ | $\ker(N)$ | $\operatorname{coker}(N)$ |
|----------------------------------|------------------------------------|-------------------------|-------------------------|-----------|---------------------------|
| $ V = 0, \text{ o.p.}$ | \mathbb{Z} | \boxtimes | \square | 0 | \circ |
| $ V = 0, \text{ o.r.}$ | \mathbb{Z}^σ | $\dot{\square}$ | $\bar{\square}$ | \bullet | $\bar{\bullet}$ |
| $ V = 2i - 1 > 0, \text{ o.p.}$ | $\mathbb{Z}/(q^{ni} - 1)$ | \ominus^i | \ominus^i | 0 | 0 |
| $ V = 2i - 1 > 0, \text{ o.r.}$ | $\mathbb{Z}^\sigma / (q^{ni} - 1)$ | \oplus^i | \oplus^i | 0 | 0 |

Table 1: The G -modules $\underline{K}_V(k)$ and associated Mackey functors

4.2 The norm cofiber sequence

In this section, we determine the $RO(G)$ -graded homotopy Mackey functors of the G -spectra in the norm cofiber sequence

$$K_G(k)_h \xrightarrow{N} K_G(k)^h \longrightarrow K_G(k)^t,$$

which forms the bottom row of the Tate diagram. The main result is Proposition 4.2.4, which says that the map $K_G(k)^t \rightarrow H\underline{\mathbb{Z}}^t$ is a weak equivalence of G -spectra.

Our analysis is centered around the $RO(G)$ -graded, Mackey functor-valued homotopy orbit spectral sequence (HOSS)

$$E_{s,V}^2 = \underline{H}_s(G; \pi_V^e K_G(k)) \Rightarrow \underline{\pi}_{s+V} K_G(k)_h,$$

and homotopy fixed point spectral sequence (HFPS) [BBHS20, §2.2]

$$\underline{E}_{s,V}^2 = \underline{H}^{-s}(G; \pi_V^e K_G(k)) \Rightarrow \underline{\pi}_{s+V} K_G(k)^h.$$

The group homology Mackey functors $\underline{H}_*(G; M)$ appearing on the \underline{E}^2 -page are given by

$$\underline{H}_*(G; M)(G/K) = H_*(K; \text{res}_K^G M)$$

for a G -module M . The group cohomology Mackey $\underline{H}^*(G; M)$ functors are defined analogously. The differentials have grading

$$d^r : \underline{E}_{s,V}^r \rightarrow \underline{E}_{s-r, V+r-1}^r.$$

Proposition 4.2.1. *The $RO(G)$ -graded homotopy Mackey functors of $K_G(k)_h$ are given on orientation preserving representations V by*

$$\underline{\pi}_V K_G(k)_h \cong \begin{cases} \boxplus & |V| = 0, \\ \oplus^i \oplus \circ & |V| = 2i - 1 > 0, \\ 0 & \text{otherwise,} \end{cases}$$

and on orientation reversing V by

$$\underline{\pi}_V K_G(k)_h \cong \begin{cases} \boxminus & |V| = 0 \\ \oplus^i \oplus \bar{\circ} & |V| = 2i - 1 > 0, \\ \bullet & |V| = 2i > 0, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Fix a $V \in RO(G)$ and let $W = V - |V|$. The homotopy orbit spectral sequence, based at W , has the form

$$\underline{E}_{s,W+t}^w = \underline{H}_s(G; \pi_{W+t}^e(K_G(k))) \Rightarrow \underline{\pi}_{W+s+t}(K_G(k))$$

and we compute $\underline{\pi}_V(K_G(k))$ by looking at the line in this spectral sequence with $s + t = |V|$. Note that by Lemma 4.1.2, the Mackey functors in this spectral sequence depend only on the number $|W| + t$ and whether or not W is orientation preserving or reversing. Since W and V differ by a trivial representation they are either both orientation preserving or orientation reversing.

The spectral sequences vanish for negative s because negative group homology is zero. When t is negative, the spectral sequence vanishes because $K(k)$ is a connective spectrum. Thus we have a first quadrant spectral sequence. In fact this spectral sequence vanishes except when s or t is equal to zero.

To see this, suppose first that W is orientation preserving. Note that when t is not zero we have

$$E_{s,t+W}^2 \cong H_s(G; K_t(k))$$

and this group is zero since the groups $K_t(k)$ are either zero or the groups $\mathbb{Z}/(q^{in} - 1)$ with C_n -action by q^i . These homology groups vanish for $s > 0$ since, by Lemma 4.1.3,

the norm map from orbits to fixed points is an isomorphism. The case of W orientation reversing is essentially the same.

The spectral sequence is displayed in Figure 1, with two cases depending on whether W is orientation preserving or reversing. There are no possible differentials when W is orientation preserving. There are possible differentials when W is orientation reversing. These differentials have the form $\bullet \rightarrow \oplus^i$ for various i . Note that these maps must be zero since $\bullet(C_n/e) = 0$ and thus there are no non-zero maps of Mackey functors $\bullet \rightarrow \oplus^i$ since \oplus^i is in the image of the functor R . All possible extensions are trivial by Proposition 3.2.1. \square




Figure 1: The \underline{E}^2 -pages of the homotopy orbit spectral sequence (HOSS) for $K_G(k)$. Dashed lines indicate potential extensions, and arrows indicate potential differentials.

Remark 4.2.2. The Ext computations afforded by Proposition 3.2.1 inform our choice to work with Mackey functors throughout this paper. Even if one were interested only in the C_n/C_n -level of the computations, it is important to make use of the entire Mackey functor structure because it helps us to easily resolve extension problems which come from spectral sequences.

For a concrete example, consider the Galois extension $\mathbb{F}_9/\mathbb{F}_3$, with Galois group C_2 , and $i = 1$. If we tried to solve extension problems only at the level C_2/C_2 , we would arrive at an extensions problem which, a priori, has two possible solutions. Indeed,

$$\text{Ext}_{\mathbb{Z}}^1(\circ(C_2/C_2), \ominus^1(C_2/C_2)) = \text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}/2, \mathbb{Z}/2) = \mathbb{Z}/2.$$

Of course, one can often resolve such extension problems by other means, but using the additional structure afforded by Mackey functors provides a systematic way to do so.

The next proposition follows from essentially the same arguments as the last, using the homotopy fixed points spectral sequence in place of the homotopy orbits spectral sequence. This spectral sequence is displayed in Figure 2. Note that in this case there is no room for either non-trivial differentials or extensions.

Proposition 4.2.3. *The $RO(G)$ -graded homotopy Mackey functors of $K_G(k)^h$ are given on orientation preserving representations V by*

$$\pi_V K_G(k)^h \cong \begin{cases} \square & |V| = 0, \\ \ominus^i & |V| = 2i - 1 > 0, \\ \circ & |V| = 2i < 0, \\ 0 & \text{otherwise,} \end{cases}$$

and on orientation reversing V by

$$\pi_V K_G(k)^h \cong \begin{cases} \square & |V| = 0, \\ \oplus^i & |V| = 2i - 1 > 0, \\ \bullet & |V| = 2i - 1 < 0 \\ \bar{\circ} & |V| = 2i < 0, \\ 0 & \text{otherwise.} \end{cases}$$




Figure 2: The E^2 -pages of the homotopy fixed point spectral sequence (HFPSS) for $K_G(k)$.

With this computation in hand, we prove the first part of Theorem D

Proposition 4.2.4. *There is an equivalence of G -spectra $K_G(k)^t \xrightarrow{\simeq} H\underline{\mathbb{Z}}^t$.*

Proof. We consider the map of Tate spectral sequences

$$\begin{array}{ccc} \underline{E}_{s,V}^2 = \widehat{H}^s(G; \pi_V^e K_G(k)) & \Longrightarrow & \underline{\pi}_{s+V} K_G(k)^t \\ \downarrow & & \downarrow \\ \underline{E}_{s,V}^2 = \widehat{H}^s(G; \pi_V^e H\underline{\mathbb{Z}}) & \Longrightarrow & \underline{\pi}_{s+V} H\underline{\mathbb{Z}}^t \end{array}$$

induced by the zeroth Postnikov section map $K_G(k) \rightarrow H\underline{\mathbb{Z}}$, and show that there is an equivalence of \underline{E}^2 -pages.

The effect of the computations in Lemmas 4.1.3 and 4.1.4 is that

$$\widehat{H}^*(G; \pi_V^e K_G(k)) = 0$$

for $|V| \neq 0$. Indeed, a G -module for which the norm map is an isomorphism has vanishing Tate cohomology.

As a result, the \underline{E}^2 -page for $K_G(k)$ is supported in the region where $|V| = 0$. The \underline{E}^2 -page for $H\underline{\mathbb{Z}}$ is also supported in this range, since $\pi_*^e H\underline{\mathbb{Z}}$ is concentrated in total degree 0. Within the $|V| = 0$ region of the \underline{E}^2 -page, the zero Postnikov section map induces an isomorphism

$$\widehat{H}^*(G; \pi_V^e K_G(k)) \cong \widehat{H}^*(G; \pi_V^e H\underline{\mathbb{Z}}),$$

which finishes the proof of the claim. \square

5 Proofs of the main theorems

As in the last section, we fix a field $k = \mathbb{F}_{q^n}$ for q a positive power of a prime p and let $G \cong C_n$ be the Galois group of the extension k/\mathbb{F}_q . In this section we prove the main results of the paper, starting with Theorem C which we now recall.

Theorem 5.0.1. *There is a homotopy pullback of genuine G -spectra*

$$\begin{array}{ccc} K_G(k) & \longrightarrow & H\underline{\mathbb{Z}} \\ \downarrow & & \downarrow \\ K_G(k)^h & \longrightarrow & H\underline{\mathbb{Z}}^h \end{array}$$

where the horizontal maps are the zeroth Postnikov truncations and the vertical maps are the geometric completion maps from the Tate diagram.

The homotopy groups of the bottom horizontal arrow can be computed effectively using the homotopy orbit spectral sequences. We use this to compute the homotopy groups of the fiber of both horizontal maps in Proposition 5.2.1 below.

To prove Theorem 5.0.1, note that Lemma 2.0.2 reduces the theorem to proving that the map $K_G(k) \rightarrow H\underline{\mathbb{Z}}$ induces equivalences

$$\widetilde{K_G(k)} \rightarrow \widetilde{H\underline{\mathbb{Z}}} \quad \text{and} \quad K_G(k)^t \rightarrow H\underline{\mathbb{Z}}^t.$$

The equivalence on Tate spectra was proven in the previous section as Proposition 4.2.4. The fact that the map $\widetilde{K}_G(k) \rightarrow \widetilde{H}\mathbb{Z}$ is an equivalence then follows from the observation, Corollary 5.1.4 below, that the maps $\widetilde{K}_G(k) \rightarrow K_G(k)^t$ and $\widetilde{H}\mathbb{Z} \rightarrow H\mathbb{Z}^t$ are both connective covers.

5.1 Comparison of Tate diagrams

Proposition 5.1.1. *The map $K_G(k) \rightarrow K_G(k)^h$ is a connective covering.*

Proof. By Merling's result, Theorem 4.0.1, we know that for all $C_m \leq C_n$ we have

$$K_G(\mathbb{F}_{p^n})^{C_n} \cong K((\mathbb{F}_{p^n})^{C_m}) \cong K(\mathbb{F}_{p^{n/m}})$$

and so by Quillen's computation, Theorem 4.0.2, we have

$$\pi_i^{C_m}(K_G(k)) \cong \pi_i(K(\mathbb{F}_{p^{n/m}})) \cong \mathbb{Z}/(q^{ni/m} - 1) \cong \ominus^i(C_n/C_m)$$

and so there are isomorphisms $\pi_i(K_G(k)) \cong \ominus^i$ for all $i \geq 0$. Comparing with Proposition 4.2.3 for $V = i$, we see that the positive integral homotopy Mackey functors are abstractly isomorphic and it remains to check that the map $K_G(k) \rightarrow K_G(k)^h$ from the Tate diagram actually induces an isomorphism.

Since the map of underlying spectra $K_G(k) \rightarrow K_G(k)_h$ is an equivalence it gives an isomorphism at C_n/e level of \ominus^i . Since

$$\ominus^i \cong R(\mathbb{Z}/q^{ni} - 1),$$

an endomorphism which is an isomorphism at the bottom level is an isomorphism. Thus for all $i \geq 0$ the map $\pi_i(K_G(k)) \rightarrow \pi_i(K_G(k)^h)$ is an isomorphism of Mackey functors. \square

Proposition 5.1.2. *The map $H\mathbb{Z} \rightarrow H\mathbb{Z}^h$ is a connective covering.*

Proof. On integer-graded homotopy groups, we have

$$\pi_s H\mathbb{Z}^h \cong \underline{H}^{-s}(G; \mathbb{Z}),$$

which is concentrated in nonpositive degrees. In degree zero,

$$\pi_0 H\mathbb{Z}^h \cong \underline{H}^0(G; \mathbb{Z}) \cong \mathbb{Z},$$

and $\pi_0 H\mathbb{Z} \rightarrow \pi_0 H\mathbb{Z}_h$ is an isomorphism. \square

Lemma 5.1.3 (c.f. [GM17, Lemma 11.2]). *For any connective G -spectrum X , the map $X \rightarrow X^h$ is a connective cover if and only if $\widetilde{X} \rightarrow X^t$ is a connective cover.*

Proof. Because X is connective the homotopy orbit spectral sequence shows that X_h is also connective. Thus the Tate diagram induces a map of long exact sequences of homotopy Mackey functors which, near degree zero, looks like:

$$\begin{array}{ccccccccc} \cdots & \longrightarrow & \pi_0(X_h) & \longrightarrow & \pi_0(X) & \longrightarrow & \pi_0(\widetilde{X}) & \longrightarrow & 0 & \longrightarrow & \cdots \\ & & \downarrow \cong & & \downarrow & & \downarrow & & \downarrow & & \\ \cdots & \longrightarrow & \pi_0(X_h) & \longrightarrow & \pi_0(X^h) & \longrightarrow & \pi_0(X^t) & \longrightarrow & 0 & \longrightarrow & \cdots \end{array}$$

For degree zero the claim follows from this portion of the diagram and the fact that there is a column of zeros. For degrees above zero the claim follows inductively from the five lemma. \square

The following is an immediate corollary of Propositions 5.1.1 and 5.1.2 and Lemma 5.1.3.

Corollary 5.1.4. *The maps $\widetilde{K_G(k)} \rightarrow K_G(k)^t$ and $\widetilde{HZ} \rightarrow HZ^t$ are connective coverings.*

We now prove the second part of Theorem D.

Proposition 5.1.5. *The Postnikov truncation $K_G(k) \rightarrow HZ$ induces an equivalence $\widetilde{K_G(k)} \rightarrow \widetilde{HZ}$.*

Proof. There is a commutative square of G -spectra

$$\begin{array}{ccc} \widetilde{K_G(k)} & \longrightarrow & \widetilde{HZ} \\ \downarrow & & \downarrow \\ K_G(k)^t & \longrightarrow & HZ^t \end{array}$$

where both vertical maps are connective covers by Corollary 5.1.4. The result now follows from that fact that the bottom arrow is an equivalence by Proposition 4.2.4. \square

This completes the proof of Theorem 5.0.1, which follows immediately from Propositions 4.2.4 and 5.1.5 and Lemma 2.0.2.

5.2 The $RO(G)$ -graded K -groups of finite fields

Theorem 5.0.1 tells us there is a homotopy pullback diagram

$$\begin{array}{ccc} K_G(k) & \longrightarrow & HZ \\ \downarrow & & \downarrow \\ K_G(k)^h & \longrightarrow & HZ^h, \end{array}$$

which we use in this section to reduce the $RO(G)$ -graded K -groups of finite fields to those of $H\mathbb{Z}$. This reduction proceeds by noticing that the pullback square give us an equivalence of fibers

$$\text{hofib}(K_G(k) \rightarrow H\mathbb{Z}) \simeq \text{hofib}(K_G(k)^h \rightarrow H\mathbb{Z}^h).$$

The homotopy Mackey functors of the fiber are straightforward to compute, giving us Theorem B from the introduction.

Proposition 5.2.1. *Let F denote the fiber of the map $K_G(k)^h \rightarrow H\mathbb{Z}^h$. The $RO(G)$ -graded homotopy groups of F are given on orientation preserving representations V by*

$$\underline{\pi}_V(F) \cong \begin{cases} \ominus^i & |V| = 2i - 1 > 0, \\ 0 & \text{otherwise,} \end{cases}$$

and on orientation reversing V by

$$\underline{\pi}_V(F) \cong \begin{cases} \oplus^i & |V| = 2i - 1 > 0, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. The homotopy fixed points spectral sequences shows that the map $K_G(k)^h \rightarrow H\mathbb{Z}^h$ is an equivalence for all $V \in RO(G)$ with $|V| \leq 0$. Thus for $|V| < 0$ the long exact sequence associated to the fibration

$$F \rightarrow K_G(k)^h \rightarrow H\mathbb{Z}^h$$

takes the form

$$\cdots \rightarrow \underline{\pi}_{V+1}K_G(K)^h \xrightarrow{\cong} \underline{\pi}_{V+1}H\mathbb{Z}^h \rightarrow \underline{\pi}_V F \rightarrow \underline{\pi}_V K_G(k)^h \xrightarrow{\cong} \underline{\pi}_V H\mathbb{Z}^h \rightarrow \cdots$$

which shows that $\underline{\pi}_V F = 0$ when $|V| < 0$. When $|V| = 0$ the claim holds because the Mackey functor $\underline{\pi}_{V+1}H\mathbb{Z}^h = 0$.

For $|V| > 0$, the long exact sequence takes the form

$$\cdots \rightarrow 0 \rightarrow \underline{\pi}_V F \rightarrow \underline{\pi}_V K_G(k)^h \rightarrow 0 \rightarrow \cdots$$

so we have $\underline{\pi}_V F \cong \underline{\pi}_V K_G(k)^h$ for $|V| > 0$. Putting all this together with Proposition 4.2.3 we obtain the result. □

Note that the groups appearing in the Mackey functors \oplus^i and \ominus^i consist of torsion abelian groups with q invertible. It follows that the q -completion of all these Mackey functors are zero, hence $F_q^\wedge \simeq 0$.

Corollary 5.2.2. *For all primes q there is an equivalence $K_{C_n}(\mathbb{F}_{q^n})_q^\wedge \simeq H\mathbb{Z}_q^\wedge$.*

Theorem 5.2.3. *The $RO(C_n)$ -graded homotopy groups of $K_{C_n}(k)$ are given by*

$$\underline{\pi}_V(K_{C_n}(k)) \cong \underline{\pi}_V(F) \oplus \underline{\pi}_V(H\underline{\mathbb{Z}}).$$

for all $V \in RO(C_n)$.

Proof. We consider the long exact sequence

$$\cdots \rightarrow \underline{\pi}_{V+1}H\underline{\mathbb{Z}} \rightarrow \underline{\pi}_V F \rightarrow \underline{\pi}_V K_G(k) \rightarrow \underline{\pi}_V H\underline{\mathbb{Z}} \rightarrow \underline{\pi}_{V-1} F \rightarrow \cdots$$

associated to the fibration

$$F \rightarrow K_G(k) \rightarrow H\underline{\mathbb{Z}}$$

for various choices of V . Note that because $\underline{\pi}_V(F) = 0$ for all V with $|V| \leq 0$ we immediately obtain the result for V with non-positive total degree.

When $|V| = 2i > 0$ we have an exact sequence

$$\underline{\pi}_{V+1}H\underline{\mathbb{Z}} \rightarrow 0 \rightarrow \underline{\pi}_V K_G(k) \rightarrow \underline{\pi}_V H\underline{\mathbb{Z}} \rightarrow M^i$$

where M^i is either \oplus^i or \ominus^i . Note that we have

$$\underline{\pi}_V(H\underline{\mathbb{Z}})(G/e) = \pi_V^e(H\underline{\mathbb{Z}}) = \pi_{|V|}H\underline{\mathbb{Z}} = 0.$$

Since both \oplus^i and \ominus^i are in the image of the right adjoint the the evaluation functor $\text{Mack}_G \rightarrow \text{Mod}_{\mathbb{Z}G}$, we see that the map $\underline{\pi}_V H\underline{\mathbb{Z}} \rightarrow M^i$ must be the zero map and we have an isomorphism

$$\underline{\pi}_V K_G(k) \rightarrow \underline{\pi}_V H\underline{\mathbb{Z}} \cong \underline{\pi}_V(F) \oplus \underline{\pi}_V H\underline{\mathbb{Z}}$$

for $|V| = 2i > 0$.

For $|V| = 2i - 1 > 0$ the long exact sequence has the form

$$\underline{\pi}_{V+1}H\underline{\mathbb{Z}} \rightarrow \underline{\pi}_V(F) \rightarrow \underline{\pi}_V K_G(k) \rightarrow \underline{\pi}_V H\underline{\mathbb{Z}} \rightarrow 0.$$

The same reasoning as the last case tells us that the leftmost map is the zero map so this is really an extension problem of the form

$$0 \rightarrow \underline{\pi}_V(F) \rightarrow \underline{\pi}_V K_G(k) \rightarrow \underline{\pi}_V H\underline{\mathbb{Z}} \rightarrow 0.$$

and by Proposition 3.2.1 this extension problem is trivial, giving us the desired splitting. \square

Remark 5.2.4. While we obtain a splitting of K -groups at every degree, we stress that this is not coming from a splitting of G -spectra. Indeed, looking at underlying spectra this would be equivalent to $K(k)$ splitting at $H\underline{\mathbb{Z}} \vee F^e$, which is not true.

5.3 Geometric fixed points

In this section, we show that calculating the geometric fixed points of $K_G(k)$ can also be reduced to computing the geometric fixed points of $H\mathbb{Z}$. For any finite group G let EP denote a G -space with

$$(EP)^H \cong \begin{cases} * & H \neq G \\ \emptyset & H = G. \end{cases}$$

This space is unique up to G -homotopy equivalence and we write \widetilde{EP} for the unreduced suspension of EP . For any G -spectrum X we write $X^\Phi = \widetilde{EP} \wedge X$. Note that when G is a cyclic group of order p there is a G -homotopy equivalence $EP \simeq EG$ and thus $X^\Phi \cong \widetilde{X}$, as defined in Section 2.

Definition 5.3.1. The *geometric fixed points* of X are the spectrum $\Phi^G(X) = (X^\Phi)^G$.

The equivariant homotopy groups of X^Φ can be understood as a localization of the homotopy groups of X . Let $\tilde{\rho}$ denote the reduced regular representation of G and let $\alpha: S^0 \rightarrow S^{\tilde{\rho}}$ be the inclusion of the fixed points $\{0, \infty\}$. Since the reduced regular representation has trivial fixed points the map α is not null homotopic and represents a non-trivial class in $\pi_{-\tilde{\rho}}^G(S_G)$.

Lemma 5.3.2. For any G -spectrum X the homotopy groups of $\Phi^G(X)$ are given by the localization

$$\pi_*(\Phi(X)) \cong \alpha^{-1} \pi_*^G(X).$$

Proof. The statement when $G = C_{2^n}$ is [HHR16, Proposition 3.18]. The proof in the general case is identical, except that we replace the role of the sign representation σ from [HHR16] with $\tilde{\rho}$. \square

Proposition 5.3.3. Let F denote the fiber of the map $K_G(k) \rightarrow H\mathbb{Z}$. The geometric fixed points $\Phi^G(F)$ are contractible.

Proof. The homotopy groups of F^Φ are $\alpha^{-1} \pi_*(F)$. Note that multiplication by α annihilates all elements in the image of transfer because

$$\alpha \cdot T_H^G(x) = T_H^G(R_H^G(\alpha) \cdot x) = T_H^G(0 \cdot x) = 0$$

for any proper subgroup $H < G$. By inspection of the homotopy groups, we see that the G/G -level of the Mackey functors \oplus^i and \ominus^i consist entirely of elements in the image of proper transfer, and thus α acts by zero on these levels. Accordingly, we have $\alpha^{-1} \pi_*(F)$ is a graded Mackey functor which vanishes at the G/G -level and the result follows. \square

Corollary 5.3.4. There is an equivalence of spectra $\Phi^G(K_G(k)) \rightarrow \Phi^G(H\mathbb{Z})$.

Proof. Geometric fixed points, being the composite of smashing with a space and categorical fixed points, preserves (co)fiber sequences. \square

Therefore, the computation of $\Phi^G K_G(k)$ reduces to the following well-known (e.g., for $p = 2$ [HHR16, Proposition 3.18]) computation of $\Phi^G H\mathbb{Z}$.

Proposition 5.3.5. *If G is a cyclic p -group, then as a graded ring,*

$$\pi_* \Phi^G H\underline{\mathbb{Z}} \cong \mathbb{Z}/p[x], \quad |x| = 2.$$

We learned of the following proof technique from a MathOverflow answer by Justin Noel.

Proof. First suppose that $G = C_{p^n}$ and for all n write \widetilde{EP}_{C_n} for unreduced suspension of the associated universal space. For all $n > 1$ let $f: C_{p^n} \rightarrow C_p$ denote the quotient by the subgroup $C_{p^{n-1}}$. Checking the fixed points for all subgroups, we see that $f^* \widetilde{EP}_{C_p}$ is a model for $\widetilde{EP}_{C_{p^n}}$. We have an isomorphism

$$\Phi^{C_{p^n}}(H\underline{\mathbb{Z}})_* \cong H\underline{\mathbb{Z}}_*^{C_{p^n}}(\widetilde{EP}_{C_{p^n}}) = H\underline{\mathbb{Z}}_*^{C_{p^n}}(f^* \widetilde{EP}_{C_p}) \cong H\underline{\mathbb{Z}}_*^{C_p}(\widetilde{EP}_{C_p})$$

where the last isomorphism comes from an isomorphism of Bredon chains

$$C_*^{C_{p^n}}(f^* \widetilde{EP}_{C_p}; \underline{\mathbb{Z}}) \cong C_*^{C_p}(\widetilde{EP}_{C_p}; \underline{\mathbb{Z}}).$$

Thus the computation for $G = C_{p^n}$ is reduced to the case $n = 1$.

When $n = 1$ we have $\widetilde{EC}_p = \widetilde{EP}_{C_p}$ and so there is an equality

$$\Phi^G(H\underline{\mathbb{Z}}) = \widetilde{H\underline{\mathbb{Z}}}^G.$$

By Corollary 5.1.4, the right hand side is the connective cover of $H\underline{\mathbb{Z}}^{tG}$. Using the Tate spectral sequence ([GM95, Theorem 10.3])

$$E_{r,s}^2 = \widehat{H}^r(C_p; \pi_s^{C_p} H\underline{\mathbb{Z}}) \Rightarrow \pi_{r+s}^{C_p} H\underline{\mathbb{Z}}^{tC_p}$$

we see that $\pi_*^{C_p} H\underline{\mathbb{Z}}^{tC_p}$ is isomorphic, as a graded ring, to the Tate cohomology of C_p . Taking the non-negative part, we see that $\pi_* \Phi^{C_p}(H\underline{\mathbb{Z}})$ is isomorphic to the group cohomology ring of C_p , as claimed. \square

Proposition 5.3.6. *If G is a cyclic group whose order has at least two prime factors, then*

$$\Phi^G H\underline{\mathbb{Z}} \simeq 0.$$

Proof. Now suppose that G is any finite group with at least two distinct prime factors, say p and q . So there exists some subgroups $C_p, C_q \subset G$. Let $\tilde{\rho}$ denote the reduced regular representation of G and let λ_p and λ_q denote the 2-dimensional representations of C_p and C_q , respectively, which rotate the plane by $\frac{2\pi}{p}$ and $\frac{2\pi}{q}$ radians. Note that λ_p and λ_q have trivial fixed points and thus the induced representations $\Lambda_p = \text{Ind}_{C_p}^G(\lambda_p)$ and $\Lambda_q = \text{Ind}_{C_q}^G(\lambda_q)$ have trivial G -fixed points. It follows that for a sufficiently large integer n there is an equivariant embedding of Λ_p and Λ_q into $n\tilde{\rho}$.

For any two G -representations V and W we have $\alpha_{V \oplus W} = \alpha_V \alpha_W$ so we must have that α_{Λ_p} and α_{Λ_q} divide $\alpha_{n\tilde{\rho}}$. Thus if we invert $\alpha_{\tilde{\rho}}$ we also invert α_{Λ_p} and α_{Λ_q} . We claim that, as

elements in the ring $H\underline{\mathbb{Z}}_*$, α_{Λ_p} is $p^{|G/C_p|}$ -torsion and α_{Λ_q} is $q^{|G/C_q|}$ -torsion. It follows that inverting both kills everything and thus $\Phi^G(H\underline{\mathbb{Z}}) \simeq *$.

To prove the claim, note that α_{λ_p} and α_{λ_q} are p and q -torsion, in $H\underline{\mathbb{Z}}_*$. This follows, for example, from Theorem 6.2.1 and Theorem 6.1.1 below which imply that $H\underline{\mathbb{Z}}_{-\lambda_p}^{C_p} \cong \mathbb{Z}/p$. Since $H\underline{\mathbb{Z}}$ is a G - E_∞ ring spectrum we can take norms and we have $\alpha_{\Lambda_p} = N_{C_p}^G(\alpha_{\lambda_p})$. The claim now follows from the multiplicative property of the norms. \square

6 Extensions of prime degree

In this section we specialize to the case of $K_{C_\ell}(\mathbb{F}_{q^\ell})$ for ℓ a prime. We recall the computation of $H\underline{\mathbb{Z}}_*$ and use this to give an explicit identification of the $RO(C_\ell)$ -graded K -groups. The computation is slightly different when $\ell = 2$ so we treat this case separately.

6.1 Quadratic extensions

Our first example is the case of quadratic extensions $\mathbb{F}_{q^2}/\mathbb{F}_q$ where the Galois group is C_2 . Table 2 below gives Lewis diagrams for all the relevant C_2 -Mackey functors.

| | | | |
|-------------------------------|---------------------------|-------------------------------|---------------------------|
| \square | $\overline{\square}$ | \circ | \boxtimes |
| \mathbb{Z} | 0 | $\mathbb{Z}/2$ | \mathbb{Z} |
| $1 \downarrow \uparrow 2$ | $0 \downarrow \uparrow 0$ | $0 \downarrow \uparrow 0$ | $2 \downarrow \uparrow 1$ |
| \mathbb{Z} | \mathbb{Z}^σ | 0 | \mathbb{Z} |
| \ominus^i | | \oplus^i | |
| $\mathbb{Z}/(q^i - 1)$ | | $\mathbb{Z}/(-q^i - 1)$ | |
| $1 \downarrow \uparrow 1+q^i$ | | $1 \downarrow \uparrow 1-q^i$ | |
| $\mathbb{Z}/(q^{2i} - 1)$ | | $\mathbb{Z}/(q^{2i} - 1)$ | |

Table 2: C_2 -Mackey functors which appear in the $RO(C_2)$ -graded homotopy of $K_{C_2}(\mathbb{F}_{q^2})$.

We begin by recalling the $RO(C_2)$ -graded homotopy groups of $H\underline{\mathbb{Z}}$. This computation is originally due to unpublished work of Stong; we use [FL04] as a reference.

Theorem 6.1.1 ([FL04, Theorem 8.1]). *Let $V \in RO(C_2)$. The $RO(C_2)$ -graded homotopy groups of $H\underline{\mathbb{Z}}$ are given by the following rules:*

(a) *If $|V^{C_2}| = 0$ then*

$$\pi_V H\underline{\mathbb{Z}} \cong \begin{cases} 0 & |V| > 0, \\ \square & |V| = 0, \\ \circ & |V| < 0. \end{cases}$$

(b) If $|V^{C_2}| > 0$ and $|V| = 0$ then

$$\pi_V H\underline{\mathbb{Z}} \cong \begin{cases} \square & |V^{C_2}| \text{ even,} \\ \overline{\square} & |V^{C_2}| \text{ odd,} \end{cases}$$

(c) If $|V^{C_2}| < 0$ and $|V| = 0$ then

$$\pi_V H\underline{\mathbb{Z}} \cong \begin{cases} \boxtimes & |V^{C_2}| \text{ even,} \\ \circ & |V^{C_2}| \leq -3 \text{ odd,} \\ \overline{\square} & |V^{C_2}| = -1 \end{cases}$$

(d) If $|V^{C_2}| > 0$ and $|V| < 0$ then

$$\pi_V H\underline{\mathbb{Z}} \cong \begin{cases} \circ & |V^{C_2}| \text{ even,} \\ 0 & |V^{C_2}| \text{ odd,} \end{cases}$$

(e) If $|V^{C_2}| < 0$ and $|V| > 0$ then

$$\pi_V H\underline{\mathbb{Z}} \cong \begin{cases} \circ & |V^{C_2}| \leq -3 \text{ odd,} \\ 0 & \text{else} \end{cases}$$

(f) If $|V^{C_2}|$ and $|V|$ are both positive or both negative then $\pi_V H\underline{\mathbb{Z}} = 0$.

We depict this result graphically in Figure 3, following the *motivic* grading convention. There are two irreducible real orthogonal C_2 -representations: the 1-dimensional trivial representation 1 and the 1-dimensional sign representation σ . As a result, $\pi_* H\underline{\mathbb{Z}}$ is a bigraded Mackey functor. In bidegree (x, y) is the representation $(x - y) + y\sigma$, whereby x is the *total degree* and y is the *twisted degree*. The *trivial degree* $x - y$ appeared in Theorem 6.1.1.

With the homotopy groups of $H\underline{\mathbb{Z}}$ and F in hand we can read off the $RO(C_2)$ -graded homotopy groups of $K_{C_2}(k)$ using Theorem 5.2.3.

Theorem 6.1.2. *The $RO(C_2)$ -graded homotopy Mackey functors of $K_{C_2}(\mathbb{F}_{q^2})$ admit a direct sum decomposition*

$$\pi_V K_{C_2}(\mathbb{F}_{q^2}) \cong \pi_V F \oplus \pi_V H\underline{\mathbb{Z}}.$$

Explicitly, they are given by

$$\pi_V K_{C_2}(\mathbb{F}_{q^2}) = \begin{cases} 0 & |V| = 2i > 0 \text{ and } |V^{C_2}| > -3 \text{ or even} \\ \circ & |V| = 2i > 0 \text{ and } |V^{C_2}| < -3 \text{ and odd} \\ \circ \oplus \ominus^i & |V| = 2i - 1 > 0 \text{ and } |V^{C_2}| \leq -3 \text{ and odd} \\ \ominus^i & |V| = 2i - 1 > 0 \text{ and } |V^{C_2}| > -3 \text{ and odd} \\ \oplus^i & |V| = 2i - 1 > 0 \text{ and } |V^{C_2}| \text{ even} \\ \pi_V H\underline{\mathbb{Z}} & \text{else} \end{cases}$$

We depict this result graphically in Figure 4. Bidegrees (e.g., $(1, 4)$) in which two Mackey functors appear represent the sum of the two Mackey functors, as was proven to be the case in Theorem 5.2.3.

$$\pi_{(x-y)+y\sigma} H\underline{\mathbb{Z}}$$




Figure 3: The $RO(C_2)$ -graded homotopy Mackey functors of $H\underline{\mathbb{Z}}$.

$$\underline{\pi}_{(x-y)+y\sigma} K_{C_2}(\mathbb{F}_{q^2})$$

| | | | | | | | | | | | | | |
|----|----|----|----|----|----|----|---|-----------------|---|-----------------|---|----------------|---|
| 6 | | | | | | | ◻ | ○⊖ ¹ | | ○⊖ ² | | ⊖ ³ | |
| 5 | | | | | | | ○ | ⊕ ¹ | ○ | ⊕ ² | | ⊕ ³ | |
| 4 | | | | | | | ◻ | ○⊖ ¹ | | ⊖ ² | | ⊖ ³ | |
| 3 | | | | | | | ○ | ⊕ ¹ | | ⊕ ² | | ⊕ ³ | |
| 2 | | | | | | | ◻ | ⊖ ¹ | | ⊖ ² | | ⊖ ³ | |
| 1 | | | | | | | ◻ | ⊕ ¹ | | ⊕ ² | | ⊕ ³ | |
| 0 | | | | | | | ◻ | ⊖ ¹ | | ⊖ ² | | ⊖ ³ | |
| -1 | | | | | ○ | | ◻ | ⊕ ¹ | | ⊕ ² | | ⊕ ³ | |
| -2 | | | | ○ | | | ◻ | ⊖ ¹ | | ⊖ ² | | ⊖ ³ | |
| -3 | | | ○ | | ○ | | ◻ | ⊕ ¹ | | ⊕ ² | | ⊕ ³ | |
| -4 | | ○ | | ○ | | | ◻ | ⊖ ¹ | | ⊖ ² | | ⊖ ³ | |
| -5 | | ○ | | ○ | | ○ | ◻ | ⊕ ¹ | | ⊕ ² | | ⊕ ³ | |
| -6 | ○ | | ○ | | ○ | | ◻ | ⊖ ¹ | | ⊖ ² | | ⊖ ³ | |
| | -6 | -5 | -4 | -3 | -2 | -1 | 0 | 1 | 2 | 3 | 4 | 5 | 6 |

Figure 4: The $RO(C_2)$ -graded homotopy Mackey functors of $K_{C_2}(\mathbb{F}_{q^2})$. The contribution from the summand $\underline{\pi}_* F$ is blue.

6.2 Extensions of odd prime degree

Here we compute the $RO(C_\ell)$ -graded homotopy of $K_{C_\ell}(\mathbb{F}_{q^\ell})$ for an odd prime ℓ . The $RO(C_\ell)$ -graded homotopy of $H\mathbb{Z}$ is similar to the case $\ell = 2$, except it is a bit simpler because all non-trivial irreducible representations have dimension 2. Lewis diagrams for the relevant Mackey functors can be found in Table 3

| \square | \circ | \boxplus | \ominus^i |
|------------------------------|---------------------------|------------------------------|--|
| \mathbb{Z} | \mathbb{Z}/ℓ | \mathbb{Z} | $\mathbb{Z}/(q^i - 1)$ |
| $1 \downarrow \uparrow \ell$ | $0 \downarrow \uparrow 0$ | $\ell \downarrow \uparrow 1$ | $1 \downarrow \uparrow \sum_{a=0}^{\ell-1} q^{ai}$ |
| \mathbb{Z} | 0 | \mathbb{Z} | $\mathbb{Z}/(q^{\ell i} - 1)$ |

Table 3: C_ℓ -Mackey functors which appear in the $RO(C_\ell)$ -graded homotopy of $K_{C_\ell}(\mathbb{F}_{q^\ell})$ for ℓ an odd prime.

Theorem 6.2.1 ([FL04, Theorem 8.1]). *Let ℓ be an odd prime and let $V \in RO(C_\ell)$. The $RO(C_\ell)$ -graded homotopy groups of $H\mathbb{Z}$ are given by the following rules:*

(a) *If $|V^{C_\ell}| = 0$ then*

$$\pi_V H\mathbb{Z} \cong \begin{cases} 0 & |V| > 0, \\ \square & |V| = 0, \\ \circ & |V| < 0. \end{cases}$$

(b) *If $|V^{C_\ell}| > 0$ and $|V| = 0$ then $\pi_V H\mathbb{Z} \cong \square$.*

(c) *If $|V^{C_\ell}| < 0$ and $|V| = 0$ then $\pi_V H\mathbb{Z} \cong \boxplus$.*

(d) *If $|V^{C_\ell}| > 0$ and $|V| < 0$ then*

$$\pi_V H\mathbb{Z} \cong \begin{cases} \circ & |V^{C_\ell}| \text{ even,} \\ 0 & |V^{C_\ell}| \text{ odd,} \end{cases}$$

(e) *If $|V^{C_\ell}| < 0$ and $|V| > 0$ then*

$$\pi_V H\mathbb{Z} \cong \begin{cases} \circ & |V^{C_\ell}| \leq -3 \text{ odd,} \\ 0 & \text{else} \end{cases}$$

(f) *If $|V^{C_\ell}|$ and $|V|$ are both positive or both negative then $\pi_V H\mathbb{Z} = 0$.*

Once again we can immediately obtain the $RO(C_\ell)$ -graded K -groups using Theorem 5.2.3.

Theorem 6.2.2. *The $RO(C_\ell)$ -graded homotopy Mackey functors of $K_{C_\ell}(\mathbb{F}_{q^\ell})$ admit a direct sum decomposition*

$$\pi_V K_{C_\ell}(\mathbb{F}_{q^\ell}) \cong \pi_V F \oplus \pi_V H\mathbb{Z}.$$

Explicitly, they are given by

$$\underline{\pi}_V K_{C_\ell}(\mathbb{F}_{q^\ell}) = \begin{cases} 0 & |V| = 2i > 0 \text{ and } |V^{C_\ell}| > -3 \text{ or even} \\ \circ & |V| = 2i > 0 \text{ and } |V^{C_\ell}| < -3 \text{ and odd} \\ \circ \oplus \ominus^i & |V| = 2i - 1 > 0 \text{ and } |V^{C_\ell}| \leq -3 \text{ and odd} \\ \ominus^i & |V| = 2i - 1 > 0 \text{ and } |V^{C_\ell}| > -3 \text{ and odd} \\ \underline{\pi}_V H\mathbb{Z} & \text{else.} \end{cases}$$

References

- [Bar17] Clark Barwick. Spectral Mackey functors and equivariant algebraic K -theory (I). *Adv. Math.*, 304:646–727, 2017.
- [BBHS20] Agnès Beaudry, Irina Bobkova, Michael Hill, and Vesna Stojanoska. Invertible $K(2)$ -local E -modules in C_4 -spectra. *Algebr. Geom. Topol.*, 20(7):3423–3503, 2020.
- [BD24] Samik Basu and Pinka Dey. Equivariant cohomology for cyclic groups, 2024.
- [BGS20] Clark Barwick, Saul Glasman, and Jay Shah. Spectral Mackey functors and equivariant algebraic K -theory, II. *Tunis. J. Math.*, 2(1):97–146, 2020.
- [CCM23] Maxine Calle, David Chan, and Andres Mejia. A linearization map for genuine equivariant algebraic k -theory, 2023.
- [CMNN20] Dustin Clausen, Akhil Mathew, Niko Naumann, and Justin Noel. Descent in algebraic K -theory and a conjecture of Ausoni-Rognes. *J. Eur. Math. Soc. (JEMS)*, 22(4):1149–1200, 2020.
- [EZ24] Elden Elmanto and Ningchuan Zhang. Equivariant algebraic K -theory and artin L -functions, 2024.
- [FL04] Kevin K. Ferland and L. Gaunce Lewis, Jr. The $RO(G)$ -graded equivariant ordinary homology of G -cell complexes with even-dimensional cells for $G = \mathbb{Z}/p$. *Mem. Amer. Math. Soc.*, 167(794):viii+129, 2004.
- [Geo21] Nick Georgakopoulos. The $RO(C_4)$ integral homology of a point, 2021.
- [GIMM23] Thomas Goodwillie, Kiyoshi Igusa, Cary Malkiewich, and Mona Merling. On the functoriality of the space of equivariant smooth h -cobordisms, 2023.
- [GM95] J. P. C. Greenlees and J. P. May. Generalized Tate cohomology. *Mem. Amer. Math. Soc.*, 113(543), 1995.
- [GM17] J. P. C. Greenlees and Lennart Meier. Gorenstein duality for real spectra. *Algebr. Geom. Topol.*, 17(6):3547–3619, 2017.

- [HHR16] M. A. Hill, M. J. Hopkins, and D. C. Ravenel. On the nonexistence of elements of Kervaire invariant one. *Ann. of Math. (2)*, 184(1):1–262, 2016.
- [HHR17] Michael A. Hill, Michael J. Hopkins, and Douglas C. Ravenel. The slice spectral sequence for the C_4 analog of real K -theory. *Forum Math.*, 29(2):383–447, 2017.
- [Kur92] Masato Kurihara. Some remarks on conjectures about cyclotomic fields and K -groups of \mathbf{Z} . *Compositio Math.*, 81(2):223–236, 1992.
- [Maz13] Kristen Luise Mazur. *On the Structure of Mackey Functors and Tambara Functors*. PhD thesis, University of Virginia, 2013.
- [Mer17] Mona Merling. Equivariant algebraic K -theory of G -rings. *Math. Z.*, 285(3-4):1205–1248, 2017.
- [MM19] Cary Malkiewich and Mona Merling. Equivariant A -theory. *Doc. Math.*, 24:815–855, 2019.
- [MM22] Cary Malkiewich and Mona Merling. The equivariant parametrized h -cobordism theorem, the non-manifold part. *Adv. Math.*, 399, 2022.
- [Qui72] Daniel Quillen. On the cohomology and K -theory of the general linear groups over a finite field. *Ann. of Math. (2)*, 96:552–586, 1972.
- [Tho85] R. W. Thomason. Algebraic K -theory and étale cohomology. *Ann. Sci. École Norm. Sup. (4)*, 18(3):437–552, 1985.
- [TT90] R. W. Thomason and Thomas Trobaugh. Higher algebraic K -theory of schemes and of derived categories. In *The Grothendieck Festschrift, Vol. III*, volume 88 of *Progr. Math.*, pages 247–435. Birkhäuser Boston, Boston, MA, 1990.
- [TW95] Jacques Thévenaz and Peter Webb. The structure of Mackey functors. *Trans. Amer. Math. Soc.*, 347(6):1865–1961, 1995.
- [Zen18] Mingcong Zeng. Equivariant Eilenberg-Mac Lane spectra in cyclic p -groups, 2018.