

Deparametrization of General Relativity by Space-Time Filling Unstable D9-Brane with Arbitrary Value of Tachyon

J. Klusoň ¹

*Department of Theoretical Physics and Astrophysics, Faculty of Science,
Masaryk University, Kotlářská 2, 611 37, Brno, Czech Republic*

Abstract

We calculate algebra of constraints of deparametrized General Relativity with space-time filling unstable D-brane for arbitrary value of tachyon field T . We also propose observables that have vanishing Poisson brackets with all first class constraints.

¹Email addresses: klu@physics.muni.cz

1 Introduction and Summary

In our previous paper [1] we analyzed space-time filling unstable D9-brane in Type IIA theory [2, 3, 4, 5] in general space-time. We performed deparametrization of theory and calculated algebra of constraints at the asymptotic region of tachyon field space where we showed that such a system is natural for deparametrization of gravity [11]. Deparametrization is general procedure when the original Hamiltonian constraint is replaced with following one in the form [14] $\mathcal{C} = \pi + H \approx 0$ where π is momentum conjugate to scalar field ϕ and where H is positive function of remaining phase space degrees of freedom which does not depend on π . This procedure was used by T. Thiemann in [14] when he suggested possible solution of problem of time in gravity.

In more details, the problem of time has following origin. According to Dirac [6] all observables have to be constant along gauge orbits and therefore have vanishing Poisson brackets with all first class constraints that are generators of gauge transformations. In case of General Relativity it can be shown that the Hamiltonian is given as sum of the first class constraints and hence Hamiltonian vanishes on the constraint surface [7, 8]. Then it is clear that all proper observables do not evolve dynamically which is known as problem of time. The fact that there is no evolution with respect to the time is in conflict with our everyday experience. On the other hand the canonical Hamiltonian describes evolution with respect to coordinate time which really does not have physical meaning thanks to the manifest diffeomorphism invariance of any theory coupled to gravity. What really makes sense is evolution with respect to other fields.

In our previous paper [1] we applied procedure suggested in [14] to the case of space-time filling non-BPS D9-brane in Type IIA theory where the tachyon field is used for deparametrization of General Relativity in the regime of large tachyon field so that tachyon is directly related to the time evolution. We should stress that the idea that open string tachyon could be related to the physical time was suggested by A. Sen in seminal paper [15] and recently in [16] where it was again emphasized that tachyon could be related to physical time when tachyon approaches its vacuum value. It is well known that open string tachyon is special since the vacuum corresponds to the vanishing tachyon potential, for review, see [27].

The crucial presumption of the paper [1] was that the tachyon is large and hence we can neglect all terms proportional to V^2 where V is tachyon

potential that appears in the Hamiltonian constraint. In the present work we relax this condition and study the situation when tachyon takes any value. We would like to stress that even in this case we can perform deparametrization of gravity however it is not clear whether Poisson bracket between new Hamiltonian constraints is strongly zero or whether it is proportional to linear combinations of primary constraints. We explicitly perform calculations of these Poisson brackets and we show that they are really zero ².

Having derived this important result it is natural to proceed to the construction of Dirac variables in the same way as in case [1], following [14]. However now we immediately find that such a naive construction cannot work due to the fact that $H(\mathbf{x})$ does not have vanishing Poisson bracket with Hamiltonian constraint thanks to the explicit dependence of H on T . In other words the case of finite T needs more general treatment which is based on relational observables, for review see [9, 10]. We try to implement this procedure in our case following seminal papers [25, 26] but it turns out that this general mechanism cannot be applied directly to the case of the system of unstable D9-brane coupled to gravity from following reason. For construction of Dirac variables [25, 26] it is necessary to have the same number of clock fields as the number of first class constraints which is ten in case of space-time filling non-BPS D9-brane. On the other hand there is only one scalar field on the world-volume of space-time filling unstable D9-brane, which is tachyon, that can serve as "clock" for Hamiltonian constraint while there are no clock fields for nine spatial diffeomorphism constraints. For that reason we propose Dirac observable whose construction will not be as general as the one developed in [25, 26]. Explicitly, we construct Dirac observable using the Hamiltonian constraint $\mathcal{G} = \frac{1}{\sqrt{\lambda}}p_T + H$. We will also presume that the parameter that appears in Dirac observable is constant over the whole spatial section in the same way as in [14]. Finally we define this observable with partial observable that is also invariant under spatial diffeomorphism which is again crucial presumption. Then we will show that this observable strongly Poisson commutes with smeared form of Hamiltonian constraint. We also show that it Poisson commutes with spatial diffeomorphism constraint and we determine evolution equation for her. Now due to the fact that Hamiltonian constraint explicitly depends on T for finite T this evolution equation cannot have the form of Poisson bracket between this Dirac observable and some Hamiltonian function. We also show that the right side of evolution

²For previous work, see [12].

equation has the form of Poisson bracket between this Dirac observable and Hamiltonian in the asymptotic region $T \rightarrow \infty$ in agreement with [14].

Let us outline our results and suggest possible extension of this work. We study deparametrized theory for space-time filling unstable D9-brane coupled to gravity for arbitrary value of tachyon. As the main result we explicitly check that the Poisson bracket between deparametrized Hamiltonian constraints is zero even in this general case. We mean that this is really remarkable result which is especially important when we construct Dirac observable. We argue that this observable has to be defined using Hamiltonian constraint instead of the Hamiltonian function that was used in case of asymptotic large value of tachyon field. Then we show that this observable has vanishing Poisson brackets with all first class constraints and hence it is true Dirac observable.

The present analysis suggests that open string tachyon has natural interpretation as time variable even in case of its finite value. Then it would be interesting to study cosmological consequences of this model. We hope to return to this problem in future.

This paper is organized as follows. In the next section (2) we review basic facts about space-time filling unstable D9-brane and determine its Hamiltonian formulation in deparametrized form. In section (3) we calculate Poisson brackets between smeared form of Hamiltonian constraints and we show that they strongly vanish. In section (3) we define observables as in case of the vanishing tachyon field and we show that this procedure does not lead to Dirac observables. Finally in section (4) we introduce alternative form of the extended variable and we show that it is true Dirac observable.

2 Review of the Basic Facts About Space-Time Filling Non-BPS D9-Brane

In this section we review basic facts about space-time filling non-BPS D9-brane coupled to gravity. Recall that the action for this system has the form

$$S = S_{GR} + S_{nonD} , \tag{1}$$

where

$$S_{GR} = \frac{1}{\kappa} \int d^{10}x \sqrt{-g} R(g) \tag{2}$$

and where

$$S_{nonD} = - \int d^{10}x V(T) \sqrt{-\det \mathbf{A}_{MN}} , \quad (3)$$

where

$$\mathbf{A}_{MN} = g_{MN} + \lambda F_{MN} + \lambda \partial_M T \partial_N T , \quad (4)$$

where $\lambda = 2\pi\alpha' = l_s^2$, where l_s is string length, and $F_{MN} = \partial_M A_N - \partial_N A_M$, where A_M is a gauge field living on the world-volume of D9-brane, T is the tachyon field with the potential $V(T)$ with the property that T has two stable minima for $T_{min\pm} = \pm\infty$ where $V(T_{min\pm}) = 0$ while it has an unstable maximum at $T_{max} = 0$ where $V(T_{max}) = \tau_{nonBPS}$ where $\tau_{nonBPS} = \frac{\sqrt{2}2\pi}{(2\pi l_s)^{10}}$ is a tension of unstable Dp-brane.

In [1] we derived canonical form of the action for this system that has the form

$$\begin{aligned} S &= S_{GR} + S_{nonD} = \\ &= \int d^{10}x (\pi^{ij} \partial_0 h_{ij} + p_T \partial_0 T + \pi^i \partial_0 A_i - N\mathcal{C} - N^i \mathcal{C}_i + A^0 \mathcal{E}) , \end{aligned} \quad (5)$$

where

$$\begin{aligned} \mathcal{C} &= \frac{\kappa}{\sqrt{h}} (\pi^{ij} h_{ik} h_{jl} \pi^{kl} - (\pi^{ij} h_{ij})^2) - \frac{1}{\kappa} \sqrt{hr} + \sqrt{\mathcal{D} + \mathcal{H}_i h^{ij} \mathcal{H}_j} \equiv \\ &\equiv \mathcal{H}_\perp^G + \mathcal{C}^{matt} , \\ \mathcal{D} &= \frac{1}{\lambda} p_T^2 + \frac{1}{\lambda^2} \pi^i \mathbf{A}_{ij}^S \pi^j + V^2 \det \mathbf{A}_{ij} , \\ \mathbf{A}_{ij} &= h_{ij} + \lambda \partial_i T \partial_j T + \lambda F_{ij} , \mathbf{A}_{ij}^S = \frac{1}{2} (\mathbf{A}_{ij} + \mathbf{A}_{ji}) = h_{ij} + \lambda T \partial_j T , \\ \mathcal{C}_i &= -2\nabla_l \pi^{kl} h_{ki} + \mathcal{H}_i \equiv \mathcal{H}_i^G + \mathcal{H}_i \equiv \tilde{\mathcal{H}}_i + p_T \partial_i T \approx 0 , \\ \mathcal{E} &= \partial_i \pi^i \approx 0 , \quad \mathcal{H}_i = p_T \partial_i T + F_{ij} \pi^j , \quad \tilde{\mathcal{H}}_i = \mathcal{H}_i^G + F_{ij} \pi^j , \end{aligned} \quad (6)$$

and where we introduced 9 + 1 formalism for the background gravity³. We considered 10-dimensional manifold \mathcal{M} with the coordinates x^M , $M = 0, \dots, 9$ and where $x^M = (t, \mathbf{x})$, $\mathbf{x} = (x^1, x^2, \dots, x^9)$. This space-time is endowed with the metric $g_{MN}(x^\rho)$ with signature $(-, +, \dots, +)$ and it is foliated by a family of space-like surfaces Σ defined by $t = x^0 = \text{const}$. h_{ij} , $i, j =$

³For review, see [17].

$1, 2, \dots, 9$ denotes the metric on Σ with inverse h^{ij} so that $h_{ij}h^{jk} = \delta_i^k$. We further introduced the operator ∇_i that is covariant derivative defined with the metric h_{ij} . We also defined the lapse function $N = 1/\sqrt{-g^{00}}$ and the shift functions $N^i = -g^{0i}/g^{00}$. In terms of these variables the components of the metric g_{MN} were written as

$$\begin{aligned} g_{00} &= -N^2 + N_i h^{ij} N_j, & g_{0i} &= N_i, & g_{ij} &= h_{ij}, \\ g^{00} &= -\frac{1}{N^2}, & g^{0i} &= \frac{N^i}{N^2}, & g^{ij} &= h^{ij} - \frac{N^i N^j}{N^2}. \end{aligned} \quad (7)$$

We further also used the fact that requirement of the preservation of the constraint $\pi^0 \approx 0$ implies the constraint $\mathcal{E} \approx \partial_i \pi^i \approx 0$ ⁴. Note that π^{ij} are momenta conjugate to h_{ij} and r is scalar curvature calculated with the metric h_{ij} . Finally the requirement of preservation of constraints $\pi_N \approx 0, \pi_{N^i} \approx 0$ where π_N and π_{N^i} are momenta conjugate to N and N^i respectively led to an existence of the constraints

$$\mathcal{C} \approx 0, \quad \mathcal{C}_i \approx 0. \quad (8)$$

We proved in [1] that it is possible to replace $\mathcal{C} \approx 0$ with new constraint $\mathcal{G} \approx 0$ that has the form

$$\begin{aligned} \mathcal{G} &\equiv \frac{1}{\sqrt{\lambda}} p_T + \sqrt{\frac{1}{2}((\mathcal{H}_\perp^G)^2 - \mathcal{S})} + \sqrt{\frac{1}{4}(\mathcal{S} - (\mathcal{H}_\perp^G)^2) - \mathcal{R}} \equiv \\ &\equiv \frac{1}{\sqrt{\lambda}} p_T + H \approx 0, \end{aligned} \quad (9)$$

and where \mathcal{S} and \mathcal{R} are defined as

$$\begin{aligned} \mathcal{S} &\equiv \frac{1}{\lambda^2} \pi^i h_{ij} \pi^j + V^2 \det \mathbf{a} + \mathcal{H}_i^G h^{ij} \mathcal{H}_j^G, \\ \mathcal{R} &= \frac{1}{\lambda^2} \pi^i \tilde{\mathcal{H}}_i \tilde{\mathcal{H}}_j \pi^j + V^2 \det \mathbf{a} a^{ij} \tilde{\mathcal{H}}_i \tilde{\mathcal{H}}_j \approx 0, \quad \mathbf{a}_{ij} = h_{ij} + \lambda F_{ij}. \end{aligned} \quad (10)$$

⁴In can be easily shown that $\mathcal{E} \approx 0$ is first class constraint and it is generator of gauge transformations for the gauge field A_i . In what follows we will presume that all observables are invariant under this gauge transformation.

Note that $\pi^i \tilde{\mathcal{H}}_i = \pi^i \mathcal{H}_i^G$ due to the fact that $\pi^i F_{ij} \pi^j = 0$. Further, \mathbf{a}^{ij} is matrix inverse to \mathbf{a}_{ij}

$$\mathbf{a}_{ij} \mathbf{a}^{jk} = \delta_i^k . \quad (11)$$

One of the goals of this paper is to calculate Poisson brackets between constraints $\mathcal{G}(\mathbf{x})$ and $\mathcal{G}(\mathbf{y})$. In our previous paper [1] we shown that in the case of asymptotic large T , when we can neglect terms proportional to V^2 , the Poisson brackets between constraints $\mathcal{G}(\mathbf{x}), \mathcal{G}(\mathbf{y})$ vanish strongly. In this paper we would like to extend this analysis to the case of arbitrary value of tachyon. We will explicitly calculate algebra of constraints $\mathcal{G}(\mathbf{x}), \mathcal{G}(\mathbf{y})$ with the help of canonical Poisson brackets

$$\begin{aligned} \{h_{ij}(\mathbf{x}), \pi^{kl}(\mathbf{y})\} &= \frac{1}{2}(\delta_i^k \delta_j^l + \delta_i^l \delta_j^k) \delta(\mathbf{x} - \mathbf{y}) , \quad \{T(\mathbf{x}), p_T(\mathbf{y})\} = \delta(\mathbf{x} - \mathbf{y}) , \\ \{A_\alpha(\mathbf{x}), \pi^\beta(\mathbf{y})\} &= \delta_\alpha^\beta \delta(\mathbf{x} - \mathbf{y}) , \quad \alpha, \beta = 0, 1, \dots, 9 . \end{aligned} \quad (12)$$

First of all we define following object

$$G(\mathbf{x}) = (H_\perp^G)^2 - H_i^G h^{ij} H_j^G , \quad \mathbf{G}(X) = \int d^9 \mathbf{x} X(\mathbf{x}) G(\mathbf{x}) , \quad (13)$$

whose importance was firstly stressed in [11] and further studied in [12, 13, 14]. Note that it is function of gravitational variables only. In our previous paper we reproduced an important result [11]

$$\{\mathbf{G}(X), \mathbf{G}(Y)\} = 0 . \quad (14)$$

Further, due to the complex form of the expression H we will calculate very carefully Poisson brackets between individual terms that appear in H . We start with \mathcal{M} where \mathcal{M} is defined as

$$\mathcal{M} = (\mathcal{H}_\perp^G)^2 - \mathcal{S} . \quad (15)$$

It is convenient to introduce smeared form of this expression which again allows us to avoid to work with partial derivatives of delta functions. Explicitly,

let us introduce $\mathbf{M}(X) = \int d^9 \mathbf{x} X \mathcal{M}$ and calculate

$$\begin{aligned}
\{\mathbf{M}(X), \mathbf{M}(Y)\} &= \{\mathbf{G}(X), \mathbf{G}(Y)\} - \\
&\left\{ \mathbf{G}(X), \int d^9 \mathbf{y} Y \left(\frac{1}{\lambda^2} \pi^i h_{ij} \pi^j + V^2 \det \mathbf{a} \right) \right\} - \\
&- \left\{ \int d^9 \mathbf{x} X \left(\frac{1}{\lambda^2} \pi^i h_{ij} \pi^j + V^2 \det \mathbf{a} \right), \mathbf{G}(Y) \right\} + \\
&\left\{ \int d^9 \mathbf{x} X \left(\frac{1}{\lambda^2} \pi^i h_{ij} \pi^j + V^2 \det \mathbf{a} \right), \int d^9 \mathbf{y} Y \left(\frac{1}{\lambda^2} \pi^i h_{ij} \pi^j + V^2 \det \mathbf{a} \right) \right\} \\
&= 4 \int d^9 \mathbf{x} (X \partial_i Y - Y \partial_i X) \frac{1}{\lambda^2} \pi^i \tilde{\mathcal{H}}_j \pi^j + \\
&2 \int d^9 \mathbf{x} (X \partial_k Y - Y \partial_k X) (\mathcal{H}_l^G \mathbf{a}^{lk} + \mathbf{a}^{kl} \mathcal{H}_l^G) V^2 \det \mathbf{a} + \\
&+ 2 \int d^9 \mathbf{x} (X \partial_i Y - Y \partial_i X) (\mathbf{a}^{ik} + \mathbf{a}^{ki}) F_{km} \pi^m V^2 \det \mathbf{a} ,
\end{aligned} \tag{16}$$

where we used the fact that

$$\pi^i \tilde{\mathcal{H}}_i = \pi^i (\mathcal{H}_i^G + F_{ij} \pi^j) = \pi^i \mathcal{H}_i^G . \tag{17}$$

and also

$$\begin{aligned}
&\left\{ \int d^9 \mathbf{x} X \mathcal{H}_i^G h^{ij} \mathcal{H}_j^G, \int d^9 \mathbf{y} Y V^2 \det \mathbf{a} \right\} + \left\{ \int d^9 \mathbf{y} X V^2 \det \mathbf{a}, \int d^9 \mathbf{x} Y \mathcal{H}_i^G h^{ij} \mathcal{H}_j^G \right\} = \\
&= -2 \int d^9 \mathbf{x} (\partial_k X Y - X \partial_k Y) (\mathcal{H}_l^G \mathbf{a}^{lk} + \mathbf{a}^{kl} \mathcal{H}_l^G) V^2 \det \mathbf{a} ,
\end{aligned} \tag{18}$$

where we have also used

$$\left\{ \int d^9 \mathbf{x} X^i \mathcal{H}_i^G, h_{kl} \right\} = -X^m \partial_m h_{kl} - \partial_k X^m h_{ml} - h_{km} \partial_l X^m . \tag{19}$$

Finally we calculate the last expression in (16)

$$\begin{aligned}
& \left\{ \int d^9 \mathbf{x} X \left(\frac{1}{\lambda^2} \pi^i h_{ij} \pi^j + V^2 \det \mathbf{a} \right), \int d^9 \mathbf{y} Y \left(\frac{1}{\lambda^2} \pi^i h_{ij} \pi^j + V^2 \det \mathbf{a} \right) \right\} = \\
& = \frac{2}{\lambda} \int d^9 \mathbf{x} (X \partial_k Y - Y \partial_k X) \pi^j h_{ji} (\mathbf{a}^{ik} - \mathbf{a}^{ki}) V^2 \det \mathbf{a} \\
& = -2 \int d^9 \mathbf{x} (X \partial_k Y - Y \partial_k X) \pi^j F_{ji} (\mathbf{a}^{ik} + \mathbf{a}^{ki}) V^2 \det \mathbf{a} ,
\end{aligned} \tag{20}$$

where in the final step we used the fact that $\mathbf{a}_{ji} \mathbf{a}^{ik} = \delta_j^k$ so that

$$(h_{ji} + \lambda F_{ji}) \mathbf{a}^{ik} = \delta_j^k , \quad \mathbf{a}^{ki} (h_{ij} + \lambda F_{ij}) = \delta_j^k \tag{21}$$

that when we combine together we obtain an important result

$$h_{ji} (\mathbf{a}^{ik} - \mathbf{a}^{ki}) = -\lambda F_{ji} (\mathbf{a}^{ik} + \mathbf{a}^{ki}) .$$

Before we proceed further we return to the definition of \mathcal{R} that can be written as

$$\mathcal{R} = \tilde{\mathcal{H}}_i \mathbf{A}^{ij} \tilde{\mathcal{H}}_j , \tag{22}$$

where the symmetric matrix $\mathbf{A}^{ij} = \mathbf{A}^{ji}$ is defined as

$$\mathbf{A}^{ij} = \frac{1}{\lambda^2} \pi^i \pi^j + \frac{1}{2} (\mathbf{a}^{ij} + \mathbf{a}^{ji}) V^2 \det \mathbf{a} . \tag{23}$$

Collecting all these results together we can write (16) in compact form

$$\{\mathbf{M}(X), \mathbf{M}(Y)\} = 4 \int d^9 \mathbf{x} (X \partial_i Y - Y \partial_i X) \mathbf{A}^{ij} \tilde{\mathcal{H}}_j . \tag{24}$$

As the second step we calculate following expression

$$\begin{aligned}
& \{\mathbf{R}(X), \mathbf{M}(Y)\} + \{\mathbf{M}(X), \mathbf{R}(Y)\} = \\
& = 2 \int d^9 \mathbf{x} d^9 \mathbf{y} X \tilde{\mathcal{H}}_i \mathbf{A}^{ij}(\mathbf{x}) \left\{ \tilde{\mathcal{H}}_j(\mathbf{x}), \mathcal{M}(\mathbf{y}) \right\} Y(\mathbf{y}) - \\
& - 2 \int d^9 \mathbf{x} d^9 \mathbf{y} Y \tilde{\mathcal{H}}_i \mathbf{A}^{ij}(\mathbf{x}) \left\{ \tilde{\mathcal{H}}_j(\mathbf{x}), \mathcal{M}(\mathbf{y}) \right\} X(\mathbf{y}) + \\
& + \int d^9 \mathbf{x} d^9 \mathbf{y} X \tilde{\mathcal{H}}_i \tilde{\mathcal{H}}_j(\mathbf{x}) \left\{ \mathbf{A}^{ij}(\mathbf{x}), \mathcal{M}(\mathbf{y}) \right\} Y(\mathbf{y}) - \\
& - \int d^9 \mathbf{x} d^9 \mathbf{y} Y \tilde{\mathcal{H}}_i \tilde{\mathcal{H}}_j(\mathbf{x}) \left\{ \mathbf{A}^{ij}(\mathbf{x}), \mathcal{M}(\mathbf{y}) \right\} X(\mathbf{y}) ,
\end{aligned} \tag{25}$$

where we again introduced smeared form of \mathcal{R} defined as

$$\mathbf{R}(X) \equiv \int d^9 \mathbf{x} X(\mathbf{x}) \mathcal{R}(\mathbf{x}) .$$

First of all we consider formulas on the first two lines in (25) where explicit calculation gives

$$\begin{aligned}
& 2 \int d^9 \mathbf{x} d^9 \mathbf{y} X \tilde{\mathcal{H}}_i \mathbf{A}^{ij}(\mathbf{x}) \left\{ \tilde{\mathcal{H}}_j(\mathbf{x}), \mathcal{M}(\mathbf{y}) \right\} Y(\mathbf{y}) - \\
& - 2 \int d^9 \mathbf{x} d^9 \mathbf{y} Y \tilde{\mathcal{H}}_i \mathbf{A}^{ij}(\mathbf{x}) \left\{ \tilde{\mathcal{H}}_j(\mathbf{x}), \mathcal{M}(\mathbf{y}) \right\} X(\mathbf{y}) = \\
& = 4 \int d^9 \mathbf{x} (X \partial_m Y - Y \partial_m X) \mathbf{A}^{mi} \tilde{\mathcal{H}}_i \mathcal{M} .
\end{aligned} \tag{26}$$

On the other hand calculations on the last two lines in (25) is much more

involved

$$\begin{aligned}
& \int d^9 \mathbf{x} d^9 \mathbf{y} X \tilde{\mathcal{H}}_i \tilde{\mathcal{H}}_j(\mathbf{x}) \{ \mathbf{A}^{ij}(\mathbf{x}), \mathcal{M}(\mathbf{y}) \} Y(\mathbf{y}) - \\
& - \int d^9 \mathbf{x} d^9 \mathbf{y} Y \tilde{\mathcal{H}}_i \tilde{\mathcal{H}}_j(\mathbf{x}) \{ \mathbf{A}^{ij}(\mathbf{x}), \mathcal{M}(\mathbf{y}) \} X(\mathbf{y}) = \\
& = 2 \int d^9 \mathbf{x} d^9 \mathbf{y} Y \mathcal{H}_i^G h^{ij}(\mathbf{x}) \left\{ \mathcal{H}_j^G(\mathbf{x}), \det \mathbf{a} \frac{1}{2} (\mathbf{a}^{kl} + \mathbf{a}^{lk})(\mathbf{y}) \right\} V^2 X \tilde{\mathcal{H}}_k \tilde{\mathcal{H}}_l(\mathbf{y}) - \\
& - 2 \int d^9 \mathbf{x} d^9 \mathbf{y} X \mathcal{H}_i^G h^{ij}(\mathbf{x}) \left\{ \mathcal{H}_j^G(\mathbf{x}), \det \mathbf{a} \frac{1}{2} (\mathbf{a}^{kl} + \mathbf{a}^{lk})(\mathbf{y}) \right\} V^2 Y \tilde{\mathcal{H}}_k \tilde{\mathcal{H}}_l(\mathbf{y}) - \\
& - \int d^9 \mathbf{x} d^9 \mathbf{y} X \tilde{\mathcal{H}}_i \tilde{\mathcal{H}}_j(\mathbf{x}) \left\{ \left(\frac{1}{\lambda^2} \pi^i \pi^j + V^2 \det \mathbf{a} \mathbf{a}_S^{ij}(\mathbf{x}) \right), \left(\frac{1}{\lambda^2} \pi^k h_{kl} \pi^l + V^2 \det \mathbf{a}(\mathbf{y}) \right) \right\} Y(\mathbf{y}) + \\
& + \int d^9 \mathbf{x} d^9 \mathbf{y} Y \tilde{\mathcal{H}}_i \tilde{\mathcal{H}}_j(\mathbf{x}) \left\{ \left(\frac{1}{\lambda^2} \pi^i \pi^j + V^2 \det \mathbf{a} \mathbf{a}_S^{ij}(\mathbf{x}) \right), \left(\frac{1}{\lambda^2} \pi^k h_{kl} \pi^l + V^2 \det \mathbf{a}(\mathbf{y}) \right) \right\} X(\mathbf{y}) = \\
& = -4 \int d^9 \mathbf{x} (X \partial_m Y - Y \partial_m X) \mathbf{a}_S^{mn} \tilde{\mathcal{H}}_n V^2 \det \mathbf{a} \mathbf{a}^{kl} \tilde{\mathcal{H}}_k \tilde{\mathcal{H}}_l \\
& + 2 \int d^9 \mathbf{x} (X \partial_m Y - Y \partial_m X) (\mathbf{a}^{jm} + \mathbf{a}^{mj}) \tilde{\mathcal{H}}_j \tilde{\mathcal{H}}_i \mathbf{a}^{ik} \tilde{\mathcal{H}}_k V^2 \det \mathbf{a} = 0 ,
\end{aligned} \tag{27}$$

where we again used

$$h_{lk} \mathbf{a}^{km} - \mathbf{a}^{mk} h_{kl} = -\lambda F_{lk} (\mathbf{a}^{km} + \mathbf{a}^{mk}) , \tag{28}$$

and where $\mathbf{a}_S^{ij} = \frac{1}{2} (\mathbf{a}^{ij} + \mathbf{a}^{ji})$. Collecting (26) and (27) together we obtain final result

$$\{ \mathbf{R}(X), \mathbf{M}(Y) \} + \{ \mathbf{M}(X), \mathbf{R}(Y) \} = 4 \int d^9 \mathbf{x} (X \partial_m Y - Y \partial_m X) \mathbf{A}^{mn} \tilde{\mathcal{H}}_n \mathcal{M} . \tag{29}$$

Finally we proceed to the calculation of Poisson bracket between smeared

forms of \mathcal{R}

$$\begin{aligned}
\{\mathbf{R}(X), \mathbf{R}(Y)\} &= 4 \int d^9 \mathbf{x} d^9 \mathbf{y} \tilde{\mathcal{H}}_i \mathbf{A}^{ij}(\mathbf{x}) \left\{ \tilde{\mathcal{H}}_j(\mathbf{x}), \tilde{\mathcal{H}}_k(\mathbf{y}) \right\} \mathbf{A}^{kl} \tilde{\mathcal{H}}_l(\mathbf{y}) + \\
&+ 2 \int d^9 \mathbf{x} d^9 \mathbf{y} X \tilde{\mathcal{H}}_i \mathbf{A}^{ij}(\mathbf{x}) \left\{ \tilde{\mathcal{H}}_j(\mathbf{x}), \mathbf{A}^{kl}(\mathbf{y}) \right\} \tilde{\mathcal{H}}_k \tilde{\mathcal{H}}_l(\mathbf{y}) + \\
&+ 2 \int d^9 \mathbf{x} d^9 \mathbf{y} X \tilde{\mathcal{H}}_i \tilde{\mathcal{H}}_j(\mathbf{x}) \left\{ \mathbf{A}^{ij}(\mathbf{x}), \tilde{\mathcal{H}}_k(\mathbf{y}) \right\} \mathbf{A}^{kl} \tilde{\mathcal{H}}_l(\mathbf{y}) + \\
&+ \int d^9 \mathbf{x} d^9 \mathbf{y} \tilde{\mathcal{H}}_i \tilde{\mathcal{H}}_j(\mathbf{x}) \left\{ \mathbf{A}^{ij}(\mathbf{x}), \mathbf{A}^{kl}(\mathbf{y}) \right\} \tilde{\mathcal{H}}_k \tilde{\mathcal{H}}_l(\mathbf{y}) .
\end{aligned} \tag{30}$$

First three terms in (30) can be easily calculated using

$$\begin{aligned}
\left\{ \int d^9 \mathbf{x} X^i \tilde{\mathcal{H}}_i, \int d^9 \mathbf{y} Y^j \tilde{\mathcal{H}}_j \right\} &= \int d^9 \mathbf{x} (X^m \partial_m Y^j - Y^m \partial_m X^j) \tilde{\mathcal{H}}_j , \\
\left\{ \int d^9 \mathbf{x} X^i \tilde{\mathcal{H}}_i, \mathbf{A}^{ij} \right\} &= -2 \partial_m (X^m \mathbf{A}^{ij}) + \partial_m X^i \mathbf{A}^{mj} + \mathbf{A}^{im} \partial_m X^j .
\end{aligned} \tag{31}$$

In case of the last term in (30) the situation is more involved. For that reason we define $\mathbf{A}(Y_{ij})$ as

$$\mathbf{A}(Y_{ij}) = \int d^9 \mathbf{x} Y_{ij} \mathbf{A}^{ij} , \tag{32}$$

where Y_{ij} is arbitrary tensor function. Then we obtain

$$\begin{aligned}
\{\mathbf{A}(X_{ij}), \mathbf{A}(Y_{kl})\} &= 2 \int d^9 \mathbf{x} [(X_{ij} \pi^i \partial_m Y_{kl} - Y_{ij} \pi^i \partial_m X_{kl}) (\mathbf{a}^{jm} - \mathbf{a}^{mj}) V^2 \det \mathbf{a} \mathbf{a}^{kl} + \\
&+ 2 \int d^9 \mathbf{x} [(X_{ij} \partial_m Y_{kl} - Y_{ij} \partial_m X_{kl}) \pi^i (\mathbf{a}^{mk} \mathbf{a}^{lj} - \mathbf{a}^{km} \mathbf{a}^{jl})] V^2 \det \mathbf{a} .
\end{aligned} \tag{33}$$

Collecting (31) together with (33) we obtain that (30) is equal to

$$\{\mathbf{R}(X), \mathbf{R}(Y)\} = 4 \int d^9 \mathbf{x} (X \partial_m Y - Y \partial_m X) \mathbf{A}^{mn} \tilde{\mathcal{H}}_n \mathcal{R} . \tag{34}$$

Now we have all ingredient for calculations of Poisson brackets between $\mathbf{H}(X), \mathbf{H}(Y)$ where $\mathbf{H}(X)$ is defined as

$$\mathbf{H}(X) = \int d^9 \mathbf{x} X H(\mathbf{x}), \quad H = \sqrt{\frac{\mathcal{M}}{2} + \sqrt{\frac{1}{4} \mathcal{M}^2 - \mathcal{R}}} \quad (35)$$

and hence

$$\begin{aligned} & \{\mathbf{H}(X), \mathbf{H}(Y)\} = \\ &= \frac{1}{16} \int d^9 \mathbf{x} d^9 \mathbf{y} \frac{X(\mathbf{x})}{\sqrt{\frac{1}{2} \mathcal{M} + \sqrt{\frac{1}{4} \mathcal{M}^2 - \mathcal{R}}}} \{\mathcal{M}(\mathbf{x}), \mathcal{M}(\mathbf{y})\} \frac{Y(\mathbf{y})}{\sqrt{\frac{1}{2} \mathcal{M} + \sqrt{\frac{1}{4} \mathcal{M}^2 - \mathcal{R}}}} + \\ &+ \frac{1}{16} \int d^9 \mathbf{x} d^9 \mathbf{y} \frac{X(\mathbf{x})}{\sqrt{\frac{1}{2} \mathcal{M} + \sqrt{\frac{1}{4} \mathcal{M}^2 - \mathcal{R}}}} \times \\ & \left\{ \mathcal{M}(\mathbf{x}), \left(\frac{1}{4} \mathcal{M}^2 - \mathcal{R} \right)(\mathbf{y}) \right\} \frac{Y(\mathbf{y})}{\sqrt{\frac{1}{2} \mathcal{M} + \sqrt{\frac{1}{4} \mathcal{M}^2 - \mathcal{R}}} \sqrt{\frac{1}{4} \mathcal{M}^2 - \mathcal{R}}} + \\ &+ \frac{1}{16} \int d^9 \mathbf{x} d^9 \mathbf{y} \frac{X(\mathbf{x})}{\sqrt{\frac{1}{2} \mathcal{M} + \sqrt{\frac{1}{4} \mathcal{M}^2 - \mathcal{R}}} \sqrt{\frac{1}{4} \mathcal{M}^2 - \mathcal{R}}} \times \\ & \left\{ \left(\frac{1}{4} \mathcal{M}^2 - \mathcal{R} \right)(\mathbf{x}), \mathcal{M}(\mathbf{y}) \right\} \frac{Y(\mathbf{y})}{\sqrt{\frac{1}{2} \mathcal{M} + \sqrt{\frac{1}{4} \mathcal{M}^2 - \mathcal{R}}}} + \\ &+ \frac{1}{16} \int d^9 \mathbf{x} d^9 \mathbf{y} \frac{X(\mathbf{x})}{\sqrt{\frac{1}{2} \mathcal{M} + \sqrt{\frac{1}{4} \mathcal{M}^2 - \mathcal{R}}} \sqrt{\frac{1}{4} \mathcal{M}^2 - \mathcal{R}}} \times \\ & \left\{ \left(\frac{1}{4} \mathcal{M}^2 - \mathcal{R} \right)(\mathbf{x}), \left(\frac{1}{4} \mathcal{M}^2 - \mathcal{R} \right)(\mathbf{y}) \right\} \frac{Y(\mathbf{y})}{\sqrt{\frac{1}{2} \mathcal{M} + \sqrt{\frac{1}{4} \mathcal{M}^2 - \mathcal{R}}} \sqrt{\frac{1}{4} \mathcal{M}^2 - \mathcal{R}}} = 0 \end{aligned} \quad (36)$$

using Poisson brackets (24),(29) and (34). In other words we got an important result

$$\{H(\mathbf{x}), H(\mathbf{y})\} = 0 \quad (37)$$

which holds even in case of finite tachyon field. Finally using this result we

obtain

$$\{\mathcal{G}(\mathbf{x}), \mathcal{G}(\mathbf{y})\} = \frac{1}{\sqrt{\lambda}} \{p_T(\mathbf{x}), H(\mathbf{y})\} + \frac{1}{\sqrt{\lambda}} \{H(\mathbf{x}), p_T(\mathbf{y})\} + \{H(\mathbf{x}), H(\mathbf{y})\} = 0 \quad (38)$$

due to the fact that Poisson bracket $\{p_T(\mathbf{x}), H(\mathbf{y})\}$ is ultralocal.

Formulas (36) and (38) are one of the most important results presented in this paper. Explicitly they say that deparametrized theory of space-time filling non-BPS D9-brane with any value of tachyon has vanishing Poisson brackets between new Hamiltonian constraints. This fact has an important consequence for the construction of Dirac variables as we will see in the next section.

Finally we will calculate Poisson bracket between $\mathbf{C}_S(X^i)$ and \mathcal{G} . Using

$$\{\mathbf{C}_S(X^i), H\} = -X^m \partial_m H - \partial_m X^m H \quad (39)$$

we obtain

$$\{\mathbf{C}_S(X^i), \mathcal{G}\} = -X^m \partial_m \mathcal{G} - \partial_m X^m \mathcal{G} \quad (40)$$

that shows that \mathcal{G} transforms as tensor density. Finally note that it is easy to see that Poisson bracket between smeared form of spatial diffeomorphism constraints is equal to

$$\{\mathbf{C}_S(X^i), \mathbf{C}_S(Y^j)\} = \mathbf{C}_S(X^j \partial_j Y^i - Y^j \partial_j X^i) . \quad (41)$$

In other words we derived that $\mathcal{G} \approx 0, \mathcal{C}_i \approx 0$ are first class constraints and that Poisson bracket between Hamiltonian constraints $\mathcal{G}(\mathbf{x}) \approx 0$ vanishes and hence we have all ingredients for construction of Dirac observables which will be performed in next sections.

3 Deparametrization of General Relativity Coupled to Tachyon-Naive Treatment

We showed in [1], following seminal work [14], that in the case of asymptotic large tachyon field when we neglect terms proportional to V^2 in \mathcal{G} we can define Dirac observable in very natural way. Explicitly, let us define \mathbf{H}_τ as [14]

$$\mathbf{H}_\tau = \int d^9 \mathbf{x} [\tau - \sqrt{\lambda} T(\mathbf{x})] H(\mathbf{x}) , \quad (42)$$

where τ has physical dimension of length which is appropriate for time variable. Further, let f is spatial diffeomorphism invariant quantity that does not depend on T and p_T . Then let us define $O_f(\tau)$ by following prescription

$$O_f(\tau) = \sum_{n=0}^{\infty} \frac{1}{n!} \{f, \mathbf{H}_\tau\}_{(n)} , \quad (43)$$

where multiple Poisson brackets are defined as

$$\begin{aligned} \{f, \mathbf{H}_\tau\}_{(0)} &= f , \quad \{f, \mathbf{H}_\tau\}_{(1)} = \{\{f, \mathbf{H}_\tau\}, \mathbf{H}_\tau\}_{(0)} = \{f, \mathbf{H}_\tau\} , \\ \{f, \mathbf{H}_\tau\}_{(n+1)} &= \left\{ \{f, \mathbf{H}_\tau\}_{(n)}, \mathbf{H}_\tau \right\} . \end{aligned} \quad (44)$$

Then it can be shown that evolution with respect to the parameter τ has the form

$$\frac{d}{d\tau} O_f(\tau) = \{O_f, \mathbf{H}\} , \quad \mathbf{H} \equiv \int d^9 \mathbf{x} H \quad (45)$$

which has the form of the Hamiltonian equation that expresses true evolution with respect to the time parameter τ generated by Hamiltonian \mathbf{H} . However the crucial question is whether $O_f(\tau)$ defined above is true Dirac observable which means that it Poisson commutes with all first class constraints $\mathcal{C}_i \approx 0, \mathcal{G} \approx 0$. In fact it is easy to demonstrate that such defined observable cannot be Dirac one simply from the fact that there is non-zero contribution from Poisson bracket between $\mathcal{G}(\mathbf{x})$ and \mathbf{H}_τ that follows from explicit dependence of H on T . In more details, let us calculate Poisson bracket between p_T and H

$$\begin{aligned} \left\{ \frac{1}{\sqrt{\lambda}} p_T(\mathbf{x}), H(\mathbf{y}) \right\} &= \frac{1}{\sqrt{\lambda}} \frac{V \frac{dV}{dT}(\mathbf{x})}{H(\mathbf{x})} \det \mathbf{a}(\mathbf{x}) \delta(\mathbf{x} - \mathbf{y}) + \\ &+ \frac{1}{2\sqrt{\lambda}} \frac{1}{H(\mathbf{x})} \frac{1}{2\sqrt{\frac{\mathcal{M}^2}{4} - \mathcal{R}}} V \frac{dV}{dT} \mathcal{M} \det \mathbf{a} \delta(\mathbf{x} - \mathbf{y}) + \\ &+ \frac{1}{2\sqrt{\lambda}} \frac{1}{H(\mathbf{x})} \frac{1}{\sqrt{\frac{\mathcal{M}^2}{4} - \mathcal{R}}} \frac{dV}{dT} V \det \mathbf{a} \mathbf{a}^{ij} \tilde{\mathcal{H}}_i \tilde{\mathcal{H}}_j \delta(\mathbf{x} - \mathbf{y}) \end{aligned} \quad (46)$$

which is non-zero for finite T . Then it is easy to see that

$$\{O_f(\tau), \mathbf{G}(M)\} \neq 0 \quad (47)$$

and hence $O_f(\tau)$ as was defined above is not Dirac observable. This is a consequence of the explicit dependence of H on the tachyon field. On the other hand we can still define Dirac observables when we proceed in the similar way as in the case of relational observables.

4 Gauge Invariant Observables for General T

Let us outline general procedure of construction of relational observables when we mainly follow excellent papers by Dittrich [25, 26]. Recall that we have ten first class constraints $\mathcal{C}_i \approx 0, \mathcal{G} \approx 0$ so that, following [25, 26], we should consider their smeared form

$$\mathbf{C}(\Lambda) \equiv \int d^9 \mathbf{x} (\Lambda^G \mathcal{G} + \Lambda^i \mathcal{C}_i) \equiv \int d^9 \mathbf{x} \Lambda^K \tilde{\mathcal{C}}_K, \quad K = 0, 1, \dots, 9, \quad (48)$$

where Λ^K are ten smeared functions. The gauge transformations generated by the function $\mathbf{C}(\Lambda)$ has the form

$$\alpha_{\mathbf{C}(\Lambda)}(f(\mathbf{x})) = \sum_{r=0}^{\infty} \frac{1}{r!} \{f(\mathbf{x}), \mathbf{C}(\Lambda)\}_r, \quad (49)$$

where

$$\begin{aligned} \{f, \mathbf{C}(\Lambda)\}_0 &= f, & \{f, \mathbf{C}(\Lambda)\}_1 &= \{f, \mathbf{C}(\Lambda)\}, \\ \{f, \mathbf{C}(\Lambda)\}_{r+1} &= \{\{f, \mathbf{C}(\Lambda)\}_r, \mathbf{C}(\Lambda)\}. \end{aligned} \quad (50)$$

Observables, which are invariant under these gauge transformations, are called Dirac observables. Partial observables are phase space functions which are not invariant under gauge transformations. It is convenient to split these partial variables into clocks variables $T_K(\mathbf{x})$ where K label first class constraints and remaining ones that we denote as f . Note that generally we need as many partial variables T_K as there are first class constraints.

The complete observable $F_{[f;T]}(\tau)(P)$ associated to the partial observable f and the clock variables $T^K(\mathbf{x})$ will generally depend on infinite many parameters $\tau^K(\mathbf{x})$. It gives value of the phase space function at the phase space

point Q in the gauge orbit through the phase space point P when the clock variables give the values $T^K(\mathbf{x})(Q) = \tau^K(\mathbf{x})$ for all $\mathbf{x} \in \Sigma$ and for all K .

Generally the calculation of complete observable is very difficult. First of all we find point Q on the gauge orbit through the point P at which $[T^K(\mathbf{x})](Q) = \tau^K(\mathbf{x})$. In other words we calculate the flow

$$[\alpha_{\mathbf{C}(\Lambda)}(T(\mathbf{x}))](P) = \sum_{r=0}^{\infty} \frac{1}{r!} \{T(\mathbf{x}), \mathbf{C}(\Lambda)\}_r \quad (51)$$

and find function $\beta(\mathbf{x})$ such that

$$[\alpha_{\mathbf{C}(\Lambda)}(T^K(\mathbf{x}))]_{\Lambda \rightarrow \beta(P)}(P) \approx \tau^K(\mathbf{x}) \quad (52)$$

for all $\mathbf{x} \in \Sigma$ where \approx means that this equation holds on the constraint surface only. It is important to stress the difference between smeared function Λ which depends on \mathbf{x} only, and $\beta(P)$ that generally depends on the phase space point P . For that reason in the equations above we should firstly calculate all Poisson brackets and only then to replace Λ with β . The value of complete observable is then given by expression [25, 26]

$$F_{[f;T]}(\tau, P) = [\alpha_{\mathbf{C}(\Lambda)}(f)]_{\Lambda \rightarrow \beta(P)}(P) . \quad (53)$$

As was shown in [25, 26] it is very difficult to find explicit form of complete observable due to the fact that (52) leads to functional differential equations.

In case of the constraint $\mathcal{G} \approx 0$ the situation is simpler since

$$\{T(\mathbf{x}), \mathcal{G}(\mathbf{y})\}_0 = T(\mathbf{x}) , \quad \{T(\mathbf{y}), \mathcal{G}(\mathbf{x})\}_1 = \frac{1}{\sqrt{\lambda}} \delta(\mathbf{x} - \mathbf{y}) \quad (54)$$

while we also have

$$\{T(\mathbf{x}), \mathbf{C}(\Lambda^i)\} = \Lambda^i \partial_i T \quad (55)$$

and hence

$$[\alpha_{\mathbf{C}(\Lambda)}(T^K(\mathbf{x}))](P) = \sum_{r=0}^{\infty} \frac{1}{r!} \{T^K(\mathbf{x}), \mathbf{C}(\Lambda)\}_r = T(\mathbf{x}) + \frac{1}{\sqrt{\lambda}} \Lambda^G(\mathbf{x}) + \Lambda^i \partial_i T . \quad (56)$$

It is important to stress that in case of world-volume of space-time filling non-BPS D9-brane there are no additional modes that could serve as spatial "clock" variables $T^i(\mathbf{x})$. In other words the general procedure of construction

of Dirac variables is not directly applicable in case of space-time filling non-BPS D9-brane. For that reason we slightly generalize procedure suggested in [14]. Explicitly, let us define following observable $F_{[f;T]}$

$$F_{[f;T]} = \sum_{r=0}^{\infty} \frac{1}{r!} \int_{\Sigma} d\mathbf{x}_1 \dots d\mathbf{x}_r (\tau - \sqrt{\lambda}T(\mathbf{x}_1)) \times \dots (\tau - \sqrt{\lambda}T(\mathbf{x}_r)) \times \{f, \mathcal{G}(\mathbf{x})\}_r . \quad (57)$$

where f is phase space function that is invariant under spatial diffeomorphism so that it has the form of integral of space density over spatial section. Further we should also stress that parameter τ that appears in (57) does not depend on \mathbf{x} .

Let us now calculate derivative of $F_{[f;T]}$ with respect to τ

$$\begin{aligned} \frac{dF_{[f;T]}}{d\tau} &= \sum_{r=0}^{\infty} \int_{\Sigma} d^9\mathbf{x}_1 \dots d\mathbf{x}_n \frac{1}{(r-1)!} (\tau - \sqrt{\lambda}T(\mathbf{x}_2)) \times \dots \times (\tau - \sqrt{\lambda}T(\mathbf{x}_{(r-1)})) \times \\ &\times \{ \dots \{ \{f, \mathcal{G}(\mathbf{x}_1)\}, \mathcal{G}(\mathbf{x}_2)\}, \dots, \mathcal{G}(\mathbf{x}_r) \} = F_{[\{f, \mathbf{G}(1)\}; T]} , \end{aligned} \quad (58)$$

where

$$\mathbf{G}[1] = \int d^9\mathbf{x} \mathcal{G}(\mathbf{x}) , \quad (59)$$

and where we used Jacobi identity

$$\{\mathcal{G}(\mathbf{x}_1), \{\mathcal{G}(\mathbf{x}_2), f\}\} + \{\mathcal{G}(\mathbf{x}_2), \{f, \mathcal{G}(\mathbf{x}_1)\}\} + \{f, \{\mathcal{G}(\mathbf{x}_1), \mathcal{G}(\mathbf{x}_2)\}\} = 0$$

that using the fact that $\{\mathcal{G}(\mathbf{x}_1), \mathcal{G}(\mathbf{x}_2)\} = \{\mathcal{H}(\mathbf{x}_1), \mathcal{H}(\mathbf{x}_2)\} = 0$ implies

$$\{\mathcal{G}(\mathbf{x}_1), \{\mathcal{G}(\mathbf{x}_2), f\}\} = \{\mathcal{G}(\mathbf{x}_2), \{\mathcal{G}(\mathbf{x}_1), f\}\} . \quad (60)$$

We see that (61) does not have the form of the Poisson bracket between $F_{[f;T]}$ and Hamiltonian function which was the case when Hamiltonian constraint does not depend on T explicitly. On the other hand it is clear that in the limit $T \rightarrow \infty$ when we can neglect V^2 in H we find that p_T Poisson commutes with H so that

$$\begin{aligned} \lim_{T \rightarrow \infty} \frac{dF_{[f;T]}}{d\tau} &= \sum_{r=0}^{\infty} \int_{\Sigma} d^9\mathbf{x}_1 \dots d\mathbf{x}_n \frac{1}{(r-1)!} (\tau - \sqrt{\lambda}T(\mathbf{x}_2)) \times \dots \times (\tau - \sqrt{\lambda}T(\mathbf{x}_{(r-1)})) \times \\ &\times \{ \dots \{ \{f, H(\mathbf{x}_1)\}, H(\mathbf{x}_2)\}, \dots, H(\mathbf{x}_r) \} = \{F_{[f;T]}, \mathbf{H}\} \end{aligned} \quad (61)$$

so that we reproduce result derived in [1] which is nice consistency check. Let us also show that $F_{[f;T]}$ has vanishing Poisson bracket with $\mathbf{G}(M)$. In order to simplify notation let us define function $F_r(\mathbf{x}_1, \dots, \mathbf{x}_r)$ as

$$\begin{aligned} F_r(\mathbf{x}_1, \dots, \mathbf{x}_r) &= \{\dots \{f, \mathcal{G}(\mathbf{x}_1)\}, \mathcal{G}(\mathbf{x}_2)\}, \dots, \mathcal{G}(\mathbf{x}_r)\} \equiv \\ &\equiv \{F_{(r-1)}(\mathbf{x}_1, \dots, \mathbf{x}_{r-1}), \mathcal{G}(\mathbf{x}_r)\} . \end{aligned} \tag{62}$$

Then we have

$$\begin{aligned} &\int d^9 \mathbf{x}_1 \dots d^9 \mathbf{x}_r \left\{ (\tau - \sqrt{\lambda} T(\mathbf{x}_1)) \times \dots \times (\tau - \sqrt{\lambda} T(\mathbf{x}_r)) F_r(\mathbf{x}_1, \dots, \mathbf{x}_r), \mathbf{G}(M) \right\} = \\ &-r \int d^9 \mathbf{x}_1 \dots d^9 \mathbf{x}_{(r-1)} (\tau - \sqrt{\lambda} T(\mathbf{x}_1)) \times \\ &\dots \times (\tau - \sqrt{\lambda} T(\mathbf{x}_{(r-1)})) \{F_{r-1}(\mathbf{x}_1, \dots, \mathbf{x}_{r-1}), \mathbf{G}(M)\} + \\ &+ \int d^9 \mathbf{x}_1 \dots d^9 \mathbf{x}_r (\tau - \sqrt{\lambda} T(\mathbf{x}_1)) \times \dots \times (\tau - \sqrt{\lambda} T(\mathbf{x}_r)) \{F_r(\mathbf{x}_1, \dots, \mathbf{x}_r), \mathbf{G}(M)\} \end{aligned} \tag{63}$$

where we again used Jacobi identity

$$\{\{X, \mathcal{G}(\mathbf{x})\}, \mathbf{G}(M)\} + \{\{\mathcal{G}(\mathbf{x}), \mathbf{G}(M)\}, X\} + \{\{\mathbf{G}(M), X\}, \mathcal{G}(\mathbf{x})\} = 0 \tag{64}$$

that holds for any phase space function X . Since $\{\mathcal{G}(\mathbf{x}), \mathbf{G}(M)\} = 0$ we see that (64) implies

$$\{\{X, \mathcal{G}(\mathbf{x})\}, \mathbf{G}(M)\} = \{\{X, \mathbf{G}(M)\}, \mathcal{G}(\mathbf{x})\} . \tag{65}$$

Using these results we obtain

$$\begin{aligned}
& \{F_{[f;T]}, \mathbf{G}(M)\} = \{f, \mathbf{G}(M)\} + \\
& + \sum_{r=1}^{\infty} \frac{1}{r!} \int_{\Sigma} d\mathbf{x}_1 \dots \mathbf{x}_r \left\{ (\tau - \sqrt{\lambda}T(\mathbf{x}_1))(\tau - \sqrt{\lambda}T(\mathbf{x}_r)) F_r(\mathbf{x}_1, \dots, \mathbf{x}_r), \mathbf{G}(M) \right\} = \\
& \{f, \mathbf{G}(M)\} - \{f, \mathbf{G}(M)\} - \sum_{r=2}^{\infty} \frac{1}{(r-1)!} \int d^9 \mathbf{x}_1 \dots d^9 \mathbf{x}_{(r-1)} (\tau - \sqrt{\lambda}T(\mathbf{x}_1)) \times \\
& \dots \times (\tau - \sqrt{\lambda}T(\mathbf{x}_{(r-1)})) \{F_{r-1}(\mathbf{x}_1, \dots, \mathbf{x}_{(r-1)}), \mathbf{G}(M)\} + \\
& + \sum_{r=1}^{\infty} \int d^9 \mathbf{x}_1 \dots d^9 \mathbf{x}_r (\tau - \sqrt{\lambda}T(\mathbf{x}_1)) \times \\
& \dots \times (\tau - \sqrt{\lambda}T(\mathbf{x}_r)) \{F_r(\mathbf{x}_1, \dots, \mathbf{x}_r), \mathbf{G}(M)\} = 0
\end{aligned} \tag{66}$$

which is desired result.

4.1 Spatial Diffeomorphism Constraint

Finally we will check that $F_{[f;T]}$ Poisson commutes with $\mathbf{C}_S(M^i)$. Since f is spatially invariant function by definition we have

$$\{\mathbf{C}_S(M^i), f\} = 0. \tag{67}$$

Further, since $\mathcal{G}(\mathbf{x})$ is tensor density we immediately obtain

$$\{\mathbf{C}(M^i), \mathcal{G}(\mathbf{x})\} = -\partial_i M^i \mathcal{G}(\mathbf{x}) - M^i \partial_i \mathcal{G}(\mathbf{x}). \tag{68}$$

Let us further calculate Poisson bracket between the function $F_r(\mathbf{x}_1, \dots, \mathbf{x}_r)$ and $\mathbf{C}(M^i)$

$$\begin{aligned}
& \{F_r(\mathbf{x}_1, \dots, \mathbf{x}_r), \mathbf{C}(M^i)\} = \{\{F_{r-1}(\mathbf{x}_1, \dots, \mathbf{x}_{(r-1)})\}, \mathcal{G}(\mathbf{x}_r), \mathbf{C}(M^i)\} = \\
& = -\{\{\mathcal{G}(\mathbf{x}_r), \mathbf{G}(M)\}, F_{r-1}(\mathbf{x}_1, \dots, \mathbf{x}_{(r-1)})\} - \{\{\mathbf{C}(M^i), F_{r-1}(\mathbf{x}_1, \dots, \mathbf{x}_{(r-1)})\}, \mathcal{G}(\mathbf{x}_r)\} = \\
& = M^i(\mathbf{x}_r) \frac{\partial}{\partial x_r^i} F_r(\mathbf{x}_1, \dots, \mathbf{x}_r) + \frac{\partial M^i(\mathbf{x}_r)}{\partial x_r^i} F_r(\mathbf{x}_1, \dots, \mathbf{x}_r) + \\
& + M^i(\mathbf{x}_{r-1}) \frac{\partial}{\partial x_{r-1}^i} F_r(\mathbf{x}_1, \dots, \mathbf{x}_r) + \frac{\partial M^i(\mathbf{x}_{r-1})}{\partial x_{r-1}^i} F_r(\mathbf{x}_1, \dots, \mathbf{x}_r) + \dots \\
& + M^i(\mathbf{x}_1) \frac{\partial}{\partial x_1^i} F_r(\mathbf{x}_1, \dots, \mathbf{x}_r) + \frac{\partial M^i(\mathbf{x}_1)}{\partial x_1^i} F_r(\mathbf{x}_1, \dots, \mathbf{x}_r),
\end{aligned} \tag{69}$$

where we used the fact that

$$\{\partial_{x^i}\mathcal{G}(\mathbf{x}), X(\mathbf{y})\} = \partial_{x^i}\{\mathcal{G}(\mathbf{x}), X(\mathbf{y})\} \quad (70)$$

that holds for any phase space function X that does not depend on \mathbf{x} . Then it is easy to see that

$$\begin{aligned} & \left\{ \int d^9\mathbf{x}_1 \dots d^9\mathbf{x}_r (\tau - T(\mathbf{x}_1)) \times (\tau - T(\mathbf{x}_r)) F_r(\mathbf{x}_1, \dots, \mathbf{x}_r), \mathbf{C}(M^i) \right\} = \\ & = \int d^9\mathbf{x}_1 \dots d^9\mathbf{x}_r (\sqrt{\lambda} M^i(\mathbf{x}_1) \frac{T}{\partial x_1^i} (\tau - \sqrt{T}(\mathbf{x}_2) \times \dots \times (\tau - \sqrt{\lambda} T(\mathbf{x}_r))) + \dots \\ & M^i(\mathbf{x}_r) \frac{\partial T}{\partial x_r^i} (\tau - T(\mathbf{x}_1)) \dots (\tau - T(\mathbf{x}_{(r-1)}))) F_r(\mathbf{x}_1, \dots, \mathbf{x}_r) + \\ & + \int d^9\mathbf{x}_1 \dots d^9\mathbf{x}_r (\tau - T(\mathbf{x}_1)) \times (\tau - T(\mathbf{x}_r)) \times (\\ & | \times M^i(\mathbf{x}_r) \frac{\partial}{\partial x_r^i} F_r(\mathbf{x}_1, \dots, \mathbf{x}_r) + \frac{\partial M^i(\mathbf{x}_r)}{\partial x_r^i} F_r(\mathbf{x}_1, \dots, \mathbf{x}_r) + \\ & + M^i(\mathbf{x}_{r-1}) \frac{\partial}{\partial x_{r-1}^i} F_r(\mathbf{x}_1, \dots, \mathbf{x}_r) + \frac{\partial M^i(\mathbf{x}_{r-1})}{\partial x_{r-1}^i} F_r(\mathbf{x}_1, \dots, \mathbf{x}_r) + \dots \\ & + M^i(\mathbf{x}_1) \frac{\partial}{\partial x_1^i} F_r(\mathbf{x}_1, \dots, \mathbf{x}_r) + \frac{\partial M^i(\mathbf{x}_1)}{\partial x_1^i} F_r(\mathbf{x}_1, \dots, \mathbf{x}_r)) = 0 \end{aligned} \quad (71)$$

and consequently

$$\{F_{[f;T]}, \mathbf{C}_S(M^i)\} = 0. \quad (72)$$

We see that the observable $F_{[f;T]}$ has vanishing Poisson brackets with all first class constraints and it is true Dirac observable. This result shows that we can still define these variables even in case of finite tachyon however now the time evolution of these variables is more complicated than in case of asymptotic large tachyon. In fact, it is easy to see that the observable $F_{[f;T]}$ reduces to observable introduced in [1] for large tachyon T . However the main message of this analysis is the fact that open string tachyon is natural time variable even in case when we cannot neglect term proportional to V^2 in Hamiltonian constraint. We mean that this result is further support for A. Sen's proposal [16, 15].

Acknowledgment:

This work is supported by the grant ‘‘Dualities and higher order derivatives’’ (GA23-06498S) from the Czech Science Foundation (GACR).

References

- [1] J. Kluson, “*Brown-Kuchar Mechanism for Unstable D-Brane at Large Tachyon Regime*,” [arXiv:2404.19080 [hep-th]].
- [2] A. Sen, “*Supersymmetric world volume action for nonBPS D-branes*,” JHEP **10** (1999), 008 doi:10.1088/1126-6708/1999/10/008 [arXiv:hep-th/9909062 [hep-th]].
- [3] M. R. Garousi, “*Tachyon couplings on nonBPS D-branes and Dirac-Born-Infeld action*,” Nucl. Phys. B **584** (2000), 284-299 doi:10.1016/S0550-3213(00)00361-8 [arXiv:hep-th/0003122 [hep-th]].
- [4] E. A. Bergshoeff, M. de Roo, T. C. de Wit, E. Eyras and S. Panda, “*T duality and actions for nonBPS D-branes*,” JHEP **05** (2000), 009 doi:10.1088/1126-6708/2000/05/009 [arXiv:hep-th/0003221 [hep-th]].
- [5] J. Kluson, “*Proposal for nonBPS D-brane action*,” Phys. Rev. D **62** (2000), 126003 doi:10.1103/PhysRevD.62.126003 [arXiv:hep-th/0004106 [hep-th]].
- [6] Dirac P. A. M., “*Lectures on quantum mechanics*”, Belfer Graduate School of Science Monographs Series, Vol. 2, Belfer Graduate School of Science, New York, 1967.
- [7] P. A. M. Dirac, “*The Theory of gravitation in Hamiltonian form*,” Proc. Roy. Soc. Lond. A **246** (1958), 333-343 doi:10.1098/rspa.1958.0142
- [8] R. L. Arnowitt, S. Deser and C. W. Misner, “*The Dynamics of general relativity*,” Gen. Rel. Grav. **40** (2008), 1997-2027 doi:10.1007/s10714-008-0661-1 [arXiv:gr-qc/0405109 [gr-qc]].
- [9] J. Tambornino, “*Relational Observables in Gravity: a Review*,” SIGMA **8** (2012), 017 doi:10.3842/SIGMA.2012.017 [arXiv:1109.0740 [gr-qc]].
- [10] K. Giesel and A. Herzog, “*Gauge invariant canonical cosmological perturbation theory with geometrical clocks in extended phase-space — A review and applications*,” Int. J. Mod. Phys. D **27** (2018) no.08, 1830005 doi:10.1142/S0218271818300057 [arXiv:1712.09878 [gr-qc]].

- [11] J. D. Brown and K. V. Kuchar, “*Dust as a standard of space and time in canonical quantum gravity*,” Phys. Rev. D **51** (1995), 5600-5629 doi:10.1103/PhysRevD.51.5600 [arXiv:gr-qc/9409001 [gr-qc]].
- [12] K. V. Kuchar and J. D. Romano, “*Gravitational constraints which generate a lie algebra*,” Phys. Rev. D **51** (1995), 5579-5582 doi:10.1103/PhysRevD.51.5579 [arXiv:gr-qc/9501005 [gr-qc]].
- [13] F. G. Markopoulou, “*Gravitational constraint combinations generate a Lie algebra*,” Class. Quant. Grav. **13** (1996), 2577-2584 doi:10.1088/0264-9381/13/9/021 [arXiv:gr-qc/9601038 [gr-qc]].
- [14] T. Thiemann, “*Solving the Problem of Time in General Relativity and Cosmology with Phantoms and k-Essence*,” [arXiv:astro-ph/0607380 [astro-ph]].
- [15] A. Sen, “*Time and tachyon*,” Int. J. Mod. Phys. A **18** (2003), 4869-4888 doi:10.1142/S0217751X03015313 [arXiv:hep-th/0209122 [hep-th]].
- [16] A. Sen, “*String Theory in Rolling Tachyon Vacuum*,” [arXiv:2312.11634 [hep-th]].
- [17] E.ourgoulhon, “*3+1 formalism and bases of numerical relativity*,” [arXiv:gr-qc/0703035 [gr-qc]].
- [18] A. Sen, “*Open and closed strings from unstable D-branes*,” Phys. Rev. D **68** (2003), 106003 doi:10.1103/PhysRevD.68.106003 [arXiv:hep-th/0305011 [hep-th]].
- [19] A. Sen, “*Fundamental strings in open string theory at the tachyonic vacuum*,” J. Math. Phys. **42** (2001), 2844-2853 doi:10.1063/1.1377037 [arXiv:hep-th/0010240 [hep-th]].
- [20] H. U. Yee and P. Yi, “*Open / closed duality, unstable D-branes, and coarse grained closed strings*,” Nucl. Phys. B **686** (2004), 31-52 doi:10.1016/j.nuclphysb.2004.03.010 [arXiv:hep-th/0402027 [hep-th]].
- [21] G. W. Gibbons, K. Hori and P. Yi, “*String fluid from unstable D-branes*,” Nucl. Phys. B **596** (2001), 136-150 doi:10.1016/S0550-3213(00)00716-1 [arXiv:hep-th/0009061 [hep-th]].

- [22] A. Sen, “*Dirac-Born-Infeld action on the tachyon kink and vortex,*” Phys. Rev. D **68** (2003), 066008 doi:10.1103/PhysRevD.68.066008 [arXiv:hep-th/0303057 [hep-th]].
- [23] M. R. Garousi, “*D-brane anti-D-brane effective action and brane interaction in open string channel,*” JHEP **01** (2005), 029 doi:10.1088/1126-6708/2005/01/029 [arXiv:hep-th/0411222 [hep-th]].
- [24] J. Kluson, “*Proposal for D0-brane anti-D0-brane action,*” J. Phys. A **56** (2023) no.1, 015401 doi:10.1088/1751-8121/acb025 [arXiv:2207.02548 [hep-th]].
- [25] B. Dittrich, “*Partial and complete observables for Hamiltonian constrained systems,*” Gen. Rel. Grav. **39** (2007), 1891-1927 doi:10.1007/s10714-007-0495-2 [arXiv:gr-qc/0411013 [gr-qc]].
- [26] B. Dittrich, “*Partial and complete observables for canonical general relativity,*” Class. Quant. Grav. **23** (2006), 6155-6184 doi:10.1088/0264-9381/23/22/006 [arXiv:gr-qc/0507106 [gr-qc]].
- [27] A. Sen, “*Tachyon dynamics in open string theory,*” Int. J. Mod. Phys. A **20** (2005), 5513-5656 doi:10.1142/S0217751X0502519X [arXiv:hep-th/0410103 [hep-th]].