

# STRICT LYAPUNOV FUNCTIONS AND ENERGY DECAY IN HAMILTONIAN CHAINS WITH DEGENERATE DAMPING

ANDREY DYMOV, LEV LOKUTSIEVSKIY, AND ANDREY SARYCHEV

ABSTRACT. We consider a Hamiltonian chain of rotators (in general nonlinear) in which the first rotator is damped. Being motivated by problems of nonequilibrium statistical mechanics of crystals, we construct a strict Lyapunov function that allows us to find a lower bound for the total energy dissipation rate when the energy and time are large. Our construction is explicit and its analysis is rather straightforward. We rely on a method going back to Matrosov, Malisoff and Mazenc, which we review in our paper. The method is rather universal and we show that it is applicable to a chain of oscillators as well.

## 1. INTRODUCTION

**1.1. Problem setting.** In this paper we consider a Hamiltonian chain of particles — rotators or oscillators, in general nonlinear, and add dissipation to the first particle. More specifically, the chain of rotators is given by the Hamiltonian

$$(1.1) \quad H(p, q) = \sum_{j=1}^N \frac{p_j^2}{2} + \sum_{j=1}^{N-1} V_j(q_j - q_{j+1}),$$

where  $N \geq 2$ ,  $p = (p_1, \dots, p_N) \in \mathbb{R}^N$  denote the moments of rotators and  $q = (q_1, \dots, q_N) \in \mathbb{T}^N$  stand for their coordinates, each one living on the one-torus  $\mathbb{T} \ni q_j$ . The periodic potentials  $V_j : \mathbb{T} \mapsto \mathbb{R}$  are assumed to be sufficiently smooth and satisfy non-degeneracy assumptions to be specified later. The Hamiltonian of the chain of oscillators is given by (4.1). The equations of motion are

$$(1.2) \quad \dot{q}_j = p_j, \quad \dot{p}_j = -\partial_{q_j} H - \delta_{1j} p_1,$$

where  $\delta_{1j}$  is the Kronecker symbol and the dot stands for the time derivative.

Being motivated by problems of nonequilibrium statistical mechanics (see Section 1.2), we are studying the total energy decay in this system when its energy is high. In the case of rotators, we construct a suitable Lyapunov function for equation (1.2), strict outside a compact set. Using it, we find a lower bound  $C_N H^{-(2N-3)}$  for the energy dissipation rate when the energy  $H$  and time are sufficiently large.

More specifically, our main result is Theorem 3.1, in which we explicitly construct a Lyapunov-type function  $W = W(p, q)$ , that behaves as  $C_N H^{2N-1}$  when  $H$  is large, satisfying

$$\dot{W}_t \leq -C_{1,N} W_t^{1/(2N-1)} + C_{2,N}.$$

Here  $W_t := W(p, q)(t)$  and  $(p, q)(t)$  is a solution to (1.1)-(1.2), while  $C_N$ ,  $C_{1,N}$  and  $C_{2,N}$  are positive constants, depending on  $N$ . Then, in Corollary 3.2 we deduce that once the initial energy  $H_0 := H(p, q)(0)$  is sufficiently large, the energy  $H_t := H(p, q)(t)$  satisfies

$$(1.3) \quad H_t \leq H_0 - C_N t H_0^{-(2N-3)}$$

on time interval  $H_0^{2N-\frac{7}{2}} \ll t \ll H_0^{2N-2}$ . Proofs of Theorem 3.1 and Corollary 3.2 are rather short and given in Section 3 which can be read independently from the rest of the paper.

In [7] the authors gave numerical evidence that the lower bound (1.3) is optimal in the sense that it is achieved at least for some potentials  $V_j$  and initial conditions, but they studied it on shorter times  $t \sim H_0^{(2N-3)/2}$ ; see below. This indicates that the energy decay in the chain of rotators can become extremely slow when the energy  $H$  is large. The reason for this phenomenon is that in the case when a  $k$ -th rotator has large energy but the energy of the  $k-1$ -st rotator is not so large, the argument  $q_{k-1} - q_k$  of the interaction potential  $V_{k-1}$  oscillates very fast. Then effective interaction of these rotators almost vanishes due to an averaging-type effect, which leads to very slow energy transport through the chain.

For the chain of oscillators we prove the existence of a strict Lyapunov function for the high energy regime, but we do not find explicitly a lower bound for the energy decay rate. These computations can be performed by analogy with those for the chain of rotators, and we hope to provide them elsewhere.

To get our result, we follow an approach going back to Matrosov, Malisoff and Mazenc, see [17], allowing one under mild conditions to construct a strict Lyapunov function once a non-strict one is known (in our case the latter is the Hamiltonian  $H$ ). Roughly speaking, it suggests a way to construct functions  $f_j(p, q)$  with  $0 \leq j \leq r$ ,  $r \geq 1$ , satisfying the following properties. The function  $f_0$  has the form  $f_0 = g(H)$  for a sufficiently large function  $g$ . The remaining functions  $f_j$  are such that their Lie derivatives  $Lf_j$  along the vector field from the r.h.s. of (1.2) satisfy  $Lf_j|_{\{Lf_{j-1}=0\}} \leq 0$ , outside the surface  $\{Lf_{j-1} = 0\}$  the derivatives  $Lf_j$  are dominated by negative parts of the derivatives  $Lf_k$ ,  $0 \leq k \leq r$ , and the surface  $\{Lf_j = 0\}$  has positive codimension in the surface  $\{Lf_{j-1} = 0\}$ . Then the desired strict Lyapunov function is given by  $\sum_{j=0}^r f_j$ .

Under additional restrictions the lower bound (1.3) was previously obtained in [7], but for times  $t \sim H_0^{(2N-3)/2}$ . Namely, the authors considered

only a specific regime when the most part of the total energy is concentrated in a single rotator <sup>1</sup>, under the additional assumption that the potentials  $V_j$  are trigonometric polynomials. As we have mentioned above, they presented numerical evidence of optimality of their lower bound.

In [7] they used a completely different, technically more complicated approach, viewing the inverse momentum of the "fast" rotator as a small parameter and using a KAM-like machinery. Namely, they performed a finite number of canonical transformations making explicit the discussed above decoupling of the "fast" rotator from its "slow" neighbours and propagating the dissipation along the chain. Adaptation of this approach to the general case when the energy is distributed arbitrarily among the rotators looks problematic, in particular because of the resonances arising when both neighbouring rotators are "fast". On the other hand, in contrast to our method, approach used in [7] allowed in [6, 4] to establish the mixing property for chains of 3 and 4 rotators, coupled via their ends to thermal bath. See a brief discussion in the next subsection and a more detailed one in Section 5.

**1.2. Motivation: mixing in chains.** Stability of Hamiltonian systems, sometimes with added dissipation, is a problem widely discussed in the literature, see e.g. excellent introduction to [7] and references therein. However, our primary motivation comes from nonequilibrium statistical mechanics in which the chains of oscillators and rotators, coupled via the ends with heat baths, are classical models to study energy transport in crystals. The heat baths are usually modelled by adding a dissipation (as in equation (1.2)) and stochastic perturbation to the equations of motion of the first and the last particles in the chain. In particular, one is interested in proving that the system approaches a stationary state when time goes to infinity, i.e. exhibits the *mixing* property. Once the mixing property is established, the question of validity of the Fourier law arises — a famous completely open problem [2, 15]. Despite many efforts by the community of mathematical physicists, some progress was achieved only in the presence of additional stochastic perturbation (see e.g. [1, 16, 10] and references therein) or assuming hyperbolicity of uncoupled dynamics [9, 20].

For the moment of writing, the mixing property is proven only for the chains of rotators of length  $N = 3, 4$  [6, 4]. A source of difficulty here is in the already discussed extremely slow energy transport through the chain. In the case of oscillators the situation is much better but the problem is not yet closed at all [12, 11, 19, 3, 13, 5].

One of the most effective approaches to this problem consist in constructing a strict Lyapunov function for high energy regime, that controls the energy decay rate in the chain provided by dissipation at its ends; our paper

---

<sup>1</sup>The authors obtained the lower bound  $C_k H^{-(2k-3)}$  for the energy dissipation rate, where  $k$  is a number of the "fast" rotator. In the case when  $k = N$  this bound coincides with ours.

is devoted to studying of this question.<sup>2</sup> Unfortunately, the construction we present gives a Lyapunov function only for the deterministic system but not for its white noise stochastic perturbation, used in the discussed above questions of nonequilibrium statistical mechanics. It does not allow to prove mixing even for the chain of 3 rotators coupled to thermal baths, in contrast to the approach suggested in [6, 4, 7]. However, our construction is simple, universal, requires only minimal assumptions and allows to get the rate of energy dissipation which seems to be optimal. We believe that it is interesting by itself and hope that its suitable modification could allow to make a progress in understanding of the mixing problem.

**1.3. Organization of the text.** In Section 2 we recall some basic facts from the Lyapunov function theory. In Proposition 2.6 we present an adapted for our purpose construction from [17], allowing to get a strict Lyapunov function (outside a compact set) once a non-strict one is given, under mild assumptions. This general construction applied to the system of rotators does not lead to the desired energy dissipation rate, but clarifies the result from Section 3, obtained by its significant modification. There, in Theorem 3.1 which is our main result, we find a strict Lyapunov function for the system of rotators leading to the claimed estimate of the energy dissipation rate. Section 3 can be read independently, but Section 2 could help to understand the presented there construction. In Section 4 we introduce the system of oscillators and check that it satisfies Assumption 2 from Section 2, crucial for our construction and allowing to apply Proposition 2.6 to get a strict Lyapunov function. In Section 5 we discuss in more details the problem of mixing in chains we concerned with in Section 1.2.

**Acknowledgments.** We are grateful to Noé Cuneo and Sergei B. Kuksin for discussions.

## 2. LASALLE'S SYSTEMS AND STRICT LYAPUNOV FUNCTIONS

### 2.1. Barbashin-Krasovskii Theorem, Lasalle's Invariance Principle.

We start by recalling a number of results on Lyapunov's direct method of establishing asymptotic stability of equilibrium for a nonlinear system via existence of Lyapunov function.

Consider a smooth  $d$ -dimensional manifold<sup>3</sup>  $\mathcal{M}$  and a  $C^m$ -smooth function  $F : \mathcal{M} \mapsto \mathbb{R}$ ,  $m \geq 1$ . Let  $\hat{x} \in \mathcal{M}$  be an equilibrium point for the system

$$(2.1) \quad \dot{x} = F(x), \quad x \in \mathcal{M}.$$

<sup>2</sup>Although we put dissipation only at one end of the chain, our results can be straightforwardly extended to the case when the dissipation acts at the both ends.

<sup>3</sup>In applications below  $\mathcal{M}$  will be chosen as a phase space of considered system, i.e.  $\mathcal{M} = \mathbb{R}^{2N}$  for the chain of  $N$  oscillators and  $\mathcal{M} = \mathbb{R}^N \times \mathbb{T}^N$  in the case of rotators.

Below we denote by  $L_F W$  the Lie derivative of a function  $W$  along the vector field  $F$ ,

$$L_F W := \text{grad}(W) \cdot F.$$

We recall that a function  $W : \mathcal{M} \mapsto \mathbb{R}$  is called *proper* if sets  $\{x : W(x) \leq c\}$  are compact for any  $c \in \mathbb{R}$ .

**Definition 2.1.** A  $C^1$ -smooth function  $W : \mathcal{M} \rightarrow \mathbb{R}$  is called *strict Lyapunov function for the equilibrium  $\hat{x}$*  of (2.1), if i)  $W$  is proper; ii)  $W(x)$  is positive in  $\mathcal{M} \setminus \hat{x}$ ; iii)  $W(\hat{x}) = 0$ , and

$$(2.2) \quad \text{iv) } L_F W(x) < 0, \quad \forall x \in \mathcal{M} \setminus \hat{x}.$$

Classical A.M. Lyapunov's theorem claims that if a strict Lyapunov function exists, then  $\hat{x}$  is asymptotically stable.

Assume now that  $W(x)$  is a *non-strict* Lyapunov function, i.e. it satisfies the non-strict version of assumption (2.2)

$$(2.3) \quad L_F W(x) \leq 0, \quad \forall x \in \mathcal{M}.$$

Asymptotic stability does not necessarily hold in this case, unless the set in which  $L_F W(x) = 0$  satisfies additional assumptions. Most well-known criteria of asymptotic stability in this case are the two related results: Barbashin-Krasovskii theorem and Lasalle's Invariance Principle.

**Proposition 2.2** (Barbashin-Krasovskii-LaSalle theorem). *Let  $\hat{x}$  be an equilibrium of system (2.1) and  $W : \mathcal{M} \rightarrow \mathbb{R}$  be a non-strict Lyapunov function, which satisfies (2.3). Let  $\mathcal{P} \subset \mathcal{M}$  be a closed bounded neighbourhood of  $\hat{x}$  which is positively invariant for (2.1). If there is no entire trajectory of the system, which is contained in  $\mathcal{P} \setminus \hat{x}$  and on which  $W$  is constant (the same, inequality (2.3) turns equality), then  $\hat{x}$  is asymptotically stable and  $\mathcal{P}$  is contained in the basin of attraction of  $\hat{x}$  for the system.*

See e.g. [14, Section 4.2]. Note that if an entire trajectory  $x(t)$  of system (2.1) is contained in  $\mathcal{P} \setminus \hat{x}$  and  $W(x(t))$  is constant, then there holds

$$(2.4) \quad L_F^k W(x(t)) = 0, \quad k = 1, 2, \dots$$

at each point of the trajectory  $x(t)$ . The following assumption would preclude the fulfilment of (2.4). Recall that the vector field  $F$  is assumed to be  $C^m$ -smooth.

**Assumption 1.** There is  $1 \leq r \leq m + 1$  such that the non-strict Lyapunov function  $W$  is  $C^r$ -smooth and

$$(2.5) \quad \left( L_F W(x), L_F^2 W(x), \dots, L_F^r W(x) \right) \neq 0, \quad \forall x \in \mathcal{P} \setminus \hat{x}.$$

One gets an obvious corollary of Proposition 2.2.

**Corollary 2.3.** *Let  $W : \mathcal{M} \rightarrow \mathbb{R}$  be a non-strict Lyapunov function for the equilibrium  $\hat{x}$  of system (2.1). Let  $\mathcal{P} \in \mathcal{M}$  be a closed bounded neighbourhood of  $\hat{x}$  which is positively invariant for the system. Then, under Assumption 1, the equilibrium  $\hat{x}$  is asymptotically stable and  $\mathcal{P}$  is contained in the basin of attraction of  $\hat{x}$  for the system.*

**2.2. Strict Lyapunov Function for Lasalle's System.** Converse Lyapunov theorem claims that for each asymptotically stable equilibrium one can find a strict Lyapunov function [18]. In this section we discuss a "constructive version" of this result appropriate for our needs, which holds under a version of Assumption 1. To this end we first extend Definition 2.1 as follows.

**Definition 2.4.** *i)* A  $C^1$ -smooth function  $W(x)$  is called *strict Lyapunov function for (2.1) outside a compact set*, if it is proper and there is  $Q \in \mathbb{R}$  such that

$$(2.6) \quad L_F W(x) < 0, \quad \forall x \in \mathcal{M} \text{ satisfying } W(x) > Q.$$

*ii)* It is called a *non-strict Lyapunov function outside a compact set*, if it is proper and for some  $Q \in \mathbb{R}$  satisfies the non-strict version inequality (2.6)

$$L_F W(x) \leq 0, \quad \forall x \in \mathcal{M} \text{ satisfying } W(x) > Q.$$

Given a proper function  $f$ , we define compact sets

$$\mathcal{K}_Q^f := \{x \in \mathcal{M} : f(x) \leq Q\}, \quad Q \in \mathbb{R}.$$

If in Definition 2.4 we wish to specify  $Q$ , we will write that  $W$  is a strict or non-strict Lyapunov function outside the compact set  $\mathcal{K}_Q^W$ .

For example,

- a strict Lyapunov function for the equilibrium  $\hat{x}$  of system (2.1) is a strict Lyapunov function outside any compact set  $\mathcal{K}_Q^W$  with  $Q \geq 0$ ;
- any proper function  $W$  satisfying  $L_F W(x) \leq 0$  for all  $x \in \mathcal{M}$  is a non-strict Lyapunov function outside any compact set  $\mathcal{K}_Q^W$ ,  $Q \in \mathbb{R}$ .

We have the following immediate result.

**Lemma 2.5.** *If  $W$  is a strict Lyapunov function outside a compact set  $\mathcal{K}_Q^W$  then the distance*

$$\text{dist}(x(t), \mathcal{K}_Q^W) \rightarrow 0 \quad \text{as } t \rightarrow \infty$$

*for any solution  $x(t)$  to (2.1).*

*Proof.* Since  $L_F W < 0$  outside  $\mathcal{K}_Q^W$ , the latter set is positively invariant. Thus, if  $x(t) \in \mathcal{K}_Q^W$  for some  $t$  then we are done. Otherwise,  $W(x(t))$  monotonically decays to some  $W_\infty \geq Q$ , so that

$$(2.7) \quad Q \leq W_\infty \leq W(x(t)) \leq W(x(0)) \quad \text{for all } t \geq 0.$$

Suppose that  $W_\infty \neq Q$ . Then  $x(t)$  for all  $t \geq 0$  belongs to a compact set which does not intersect with  $\mathcal{K}_Q^W$ . Then,  $\inf_{t \geq 0} L_F W(x(t)) < 0$  which contradicts (2.7).  $\square$

Now let  $Q \in \mathbb{R}$  and  $W : \mathcal{M} \rightarrow \mathbb{R}$  be a non-strict Lyapunov function outside a compact set  $\mathcal{K}_Q^W$  for system (2.1). We will need the following version of Assumption 1.

**Assumption 2.** There exists  $1 \leq r \leq m$  such that the Lyapunov function  $W$  is  $C^{r+1}$ -smooth and satisfies relation (2.5) of Assumption 1 outside the compact set  $\mathcal{K}_Q^W$ ,

$$\left( L_F W(x), L_F^2 W(x), \dots, L_F^r W(x) \right) \neq 0, \quad \forall x \notin \mathcal{K}_Q^W.$$

Under this assumption there exist smooth positive functions  $\varphi, \Phi : (Q, \infty) \mapsto \mathbb{R}_{>0}$ , for any  $x \notin \mathcal{K}_Q^W$  satisfying

$$(2.8) \quad |L_F W(x)| + \sum_{k=2}^r |L_F^k W(x)|^2 \geq \varphi(W(x)),$$

$$(2.9) \quad |L_F^k W(x)| \leq \Phi(W(x)) \quad \text{for all } 1 \leq k \leq r+1,$$

and

$$(2.10) \quad \varphi \leq \Phi^2.$$

Indeed, one can take

$$\tilde{\varphi}(w) = \inf \left\{ |L_F W(x)| + \sum_{k=2}^r |L_F^k W(x)|^2 \quad \middle| \quad x \in \mathcal{M} : W(x) = w \right\}$$

and

$$\tilde{\Phi}(w) = \sup \left\{ |L_F^k W(x)| \quad \middle| \quad k = 1, \dots, r+1; x \in \mathcal{M} : W(x) = w \right\},$$

and then minorize and majorize functions  $\tilde{\varphi}$  and  $\tilde{\Phi}$  by smooth positive functions  $\varphi$  and  $\Phi$  correspondingly, satisfying (2.10).

Below we fix a pair of functions  $\varphi$  and  $\Phi$  satisfying (2.8), (2.9) and (2.10).

**Proposition 2.6.** *Let  $Q \in \mathbb{R}$  and  $W : \mathcal{M} \rightarrow \mathbb{R}$  be a non-strict Lyapunov function outside a compact set  $\mathcal{K}_Q^W$  for system (2.1). Then, under Assumption 2, there exist smooth functions  $A, B_k : (Q, \infty) \mapsto \mathbb{R}$ ,  $k = 2, \dots, r$ , such that  $B_k \geq 1$  and for any  $\varepsilon > 0$  the function*

$$(2.11) \quad W^\#(x) := A(W(x)) - \sum_{k=2}^r B_k(W(x)) L_F^{k-1} W(x) L_F^k W(x), \quad x \notin \mathcal{K}_{Q+\varepsilon}^W,$$

*continued anyhow to a  $C^1$ -smooth function on  $\mathcal{M}$ , is a strict Lyapunov function for (2.1) outside  $\mathcal{K}_{Q+\varepsilon}^W$  (the  $C^1$ -smooth continuation always exists).*

*The functions  $B_k$  can be chosen in the form*

$$(2.12) \quad B_k = 2^{(r-k)(r-k+1)} (\Phi^2 / \varphi)^{r-k}, \quad k = 2, \dots, r,$$

*while the function  $A$  can be taken arbitrary, growing at infinity so fast that  $W^\#$  is proper, with derivative satisfying*

$$(2.13) \quad A' > \Phi^2 \sum_{k=2}^r |B_k'| + \Phi B_2 + 1.$$

With this choice,

$$L_F W^\#(x) \leq -\varphi(W(x))/4, \quad \forall x \notin \mathcal{K}_{Q+\varepsilon}^W.$$

This is a version of result established in [17, Th. 5.1]. We provide a proof which is more succinct than the original one.

**2.3. Proof of Proposition 2.6.** Differentiating function (2.11) along the direction of the vector field  $F$  we get

$$(2.14) \quad L_F W^\#(x) = \left[ A'(W(x)) - \sum_{k=2}^r B'_k(W(x)) L_F^{k-1} W(x) L_F^k W(x) \right] L_F W(x) - \sum_{k=2}^r B_k(W(x)) (L_F^k W(x))^2 -$$

$$(2.15) \quad B_2(W(x)) L_F W(x) L_F^3 W(x) - \sum_{k=3}^r B_k(W(x)) L_F^{k-1} W(x) L_F^{k+1} W(x) = A_1(x) L_F W(x) - \sum_{k=2}^r B_k(W(x)) (L_F^k W(x))^2 -$$

$$(2.16) \quad \sum_{k=3}^r B_k(W(x)) L_F^{k-1} W(x) L_F^{k+1} W(x),$$

where

$$A_1(x) = \left[ A'(W(x)) - \sum_{k=2}^r B'_k(W(x)) L_F^{k-1} W(x) L_F^k W(x) - B_2(W(x)) L_F^3 W(x) \right]$$

Since we suppose  $B_k \geq 1$  all  $k$ , the sum in (2.15) is non-negative. By (2.9),

$$A_1 \geq A'(W) - \sum_{k=2}^r |B'_k(W)| \Phi(W)^2 - B_2(W) \Phi(W).$$

Choosing the function  $A$  satisfying (2.13), we get  $A_1 \geq 1$ . In particular, the term (2.15) is non-positive, since  $L_F W \leq 0$ .

Next, we choose  $B_k$  in such a way that (2.16) is bounded by (2.15). More specifically, we assume that

$$(2.17) \quad B_k \leq \frac{(B_{k-1})^{1/2} (B_{k+1})^{1/2}}{\sqrt{2}}, \quad 3 \leq k \leq r-1.$$

Then

$$\left| B_k(W) L_F^{k-1} W L_F^{k+1} W \right| \leq \frac{B_{k-1}(W)}{4} (L_F^{k-1} W)^2 + \frac{B_{k+1}(W)}{4} (L_F^{k+1} W)^2,$$

so that

$$\begin{aligned} \sum_{k=3}^{r-1} |B_k(W) L_F^{k-1} W L_F^{k+1} W| &\leq \sum_{k=4}^{r-2} \frac{B_k(W)}{2} (L_F^k W)^2 + \sum_{k=2,3,r-1,r} \frac{B_k(W)}{4} (L_F^k W)^2 \\ &\leq \sum_{k=2}^r \frac{B_k(W)}{2} (L_F^k W)^2 - \frac{B_{r-1}(W)}{4} (L_F^{r-1} W)^2. \end{aligned}$$

Similarly,

$$|B_r(W) L_F^{r-1} W L_F^{r+1} W| \leq \frac{B_{r-1}(W)}{4} (L_F^{r-1} W)^2 + \frac{(B_r(W))^2}{B_{r-1}(W)} (L_F^{r+1} W)^2.$$

Inserting these estimate into (2.15)-(2.16), we find

$$L_F W^\# \leq A_1 L_F W - \frac{1}{2} \sum_{k=2}^r B_k(W) (L_F^k W)^2 + \frac{(B_r(W))^2}{B_{r-1}(W)} (L_F^{r+1} W)^2.$$

Now we set  $B_r := 1$ . Then, according to (2.8)

$$A_1 L_F W - \frac{1}{2} \sum_{k=2}^r B_k(W) (L_F^k W)^2 \leq -\frac{\varphi(W)}{2},$$

since  $A_1, B_k \geq 1$ . On the other hand,

$$\frac{(B_r(W))^2}{B_{r-1}(W)} (L_F^{r+1} W)^2 \leq \frac{(\Phi(W))^2}{B_{r-1}(W)}.$$

We choose  $B_{r-1}$  in such a way that  $\frac{(\Phi(W))^2}{B_{r-1}(W)} = \frac{\varphi(W)}{4}$ , i.e.

$$(2.18) \quad B_{r-1} = \frac{4\Phi^2}{\varphi}.$$

Note that  $B_{r-1} \geq 1$ , due to (2.10). Then,

$$L_F W^\# \leq -\frac{\varphi(W)}{4}.$$

It remains to choose the functions  $B_k \geq 1$ ,  $k < r-1$ , satisfying (2.17), which is equivalent to

$$B_{k-1} \geq \frac{4B_k^2}{B_{k+1}}.$$

We choose them to be minimal, that is replace the inequality above by the equality. Since  $B_r = 1$  and  $B_{r-1}$  is given by (2.18), we find

$$(2.19) \quad B_k = 2^{(r-k)(r-k+1)} \left( \frac{\Phi^2}{\varphi} \right)^{r-k}.$$

Finally, we note that the constructed in such a way function  $W^\#$  is defined outside the compact set  $\mathcal{K}_Q^W$  and is  $C^1$ -smooth. Indeed,  $W(x) > Q$  for  $x \notin \mathcal{K}_Q^W$ , so the functions  $\varphi(W(x))$  and  $\Phi(W(x))$  are well-defined and the function  $\varphi(W(x))$  appearing in the denominator of (2.19) does not vanish.

To get from  $W^\sharp$  the desired strict Lyapunov function, which we denote  $W_\varepsilon^\sharp(x)$ , we restrict  $W^\sharp$  to the set  $\mathcal{K}_{Q+\varepsilon}^W$  and  $C^1$ -smoothly continue it to the whole manifold  $\mathcal{M}$ . Such continuation exists since we can take

$$W_\varepsilon^\sharp(x) := \begin{cases} W^\sharp(x)\chi_\varepsilon(W(x)) & \text{if } x \notin \mathcal{K}_Q^W, \\ 0 & \text{if } x \in \mathcal{K}_Q^W, \end{cases}$$

where  $\chi$  is a smooth function satisfying  $\chi(w) = 1$  for  $w \geq Q + \varepsilon$  and  $\chi(w) = 0$  for  $w \leq Q + \varepsilon/2$ .  $\square$

### 3. CHAIN OF ROTATORS

**3.1. Lyapunov function and energy decay rate.** In this section we study the chain of rotators given by (1.1)-(1.2). We construct a strict Lyapunov function outside a compact set that provides lower bound (1.3) for the energy dissipation rate.

Writing the equations of motion (1.2) in more details, we get

$$(3.1) \quad \dot{q}_j = p_j, \quad \dot{p}_j = -\delta_{j1}p_1 + V_{j-1}'(q_{j-1} - q_j) - V_j'(q_j - q_{j+1}), \quad 1 \leq j \leq N,$$

where  $V_0 = V_N := 0$ . Clearly, the Hamiltonian  $H$  is a non-strict Lyapunov function for system (3.1) outside a compact set  $\mathcal{K}_Q^H$  with arbitrary  $Q \in \mathbb{R}$ , since it is proper and

$$(3.2) \quad L_F H = -p_1^2 \leq 0.$$

Here  $F$  denotes the vector field from the r.h.s. of (3.1) and  $L_F$  is the Lie derivative along its direction. We assume that the interaction potentials  $V_j$  satisfy

**Assumption 3.** Potentials  $V_i : \mathbb{T} \mapsto \mathbb{R}$  are  $C^3$ -smooth and  $(V_i'(x))^2 + (V_i''(x))^2 \neq 0$  for every  $x \in \mathbb{T}$  and  $1 \leq i \leq N - 1$ . Moreover, without loss of generality we assume that  $V_i \geq 1$  for any  $i$ , so  $H \geq 1$  as well.

By simplifying proof of Proposition 4.1 below, one can easily check that Assumption 3 implies Assumption 2 with appropriate  $Q$  and  $r = 4N - 3$ , once the potentials  $V_j$  are  $C^{r+1}$ -smooth. So, Proposition 2.6 with  $W = H$  provides a strict Lyapunov function  $W^\sharp$  outside a compact set for system (3.1). However, it is difficult to identify the function  $\varphi$  from (2.8), and consequently  $W^\sharp$ . Moreover, the function  $\Phi$  in (2.9) grows at least as  $Cp_1^r$  when  $|p_1| \rightarrow \infty$  since  $L_F^{r+1}H$  does, so the function  $B_2$  from (2.12) grows at least as  $C_1p_1^{2r(r-2)}/\varphi^{r-2}$ . Our estimations for the function  $\varphi$  show that this gives too large upper bound for the function  $W^\sharp$ , leading to a bad lower bound for the energy decay rate.

To sharpen the bound, below we suggest another strict Lyapunov function outside a compact set, construction of which is motivated by (2.11), but instead of the derivatives  $L_F^k H$  we employ their appropriate parts. This construction is also related to the one presented in [17, Theorem 3.1].

Let us denote

$$(3.3) \quad \xi_j(q) := -V_j'(q_j - q_{j+1}), \quad \text{so that} \quad L_F p_j = \xi_j - \xi_{j-1} - \delta_{1j} p_1,$$

for  $1 \leq j \leq N-1$ , where  $\xi_0 := 0$ . Consider the function

$$(3.4) \quad W(p, q) = a_0 H^{\gamma_0} - \sum_{j=1}^{N-1} (a_{2j-1} H^{\alpha_{2j-1}} p_j \xi_j + a_{2j} H^{\alpha_{2j}} \xi_j L_F \xi_j),$$

where the constants  $a_k \geq 1$  and

$$(3.5) \quad \gamma_0 := 2N-1, \quad \alpha_k := 2(N-1) - k \quad \text{for} \quad 1 \leq k \leq 2(N-1).$$

Due to the bounds

$$(3.6) \quad |p_j \xi_j| \leq C\sqrt{H}, \quad |\xi_j L_F \xi_j| \leq C\sqrt{H},$$

the function  $W$  satisfies

$$(3.7) \quad \frac{a_0}{2} H^{\gamma_0} \leq a_0 H^{\gamma_0} (1 - CH^{-3/2}) \leq W \leq a_0 H^{\gamma_0} (1 + CH^{-3/2}) \leq 2a_0 H^{\gamma_0},$$

since we assume  $a_0$  to be sufficiently large. Here and below by  $C, C_1, \dots$  we denote various positive constants which may depend on  $N$  and change from line to line.

**Theorem 3.1.** *Let  $N \geq 2$ , the constants  $a_{2N-2}$  and  $a_k - a_{k+1}$  are sufficiently large for all  $k$ , and Assumption 3 is satisfied. Then  $W$  is a strict Lyapunov function outside a compact set and*

$$(3.8) \quad L_F W \leq -H + C_1 \leq -C_2 W^{1/\gamma_0} + C_1,$$

for appropriate constants  $C_{1,2} = C_{1,2}(N; a_0, \dots, a_{2N-2})$ .

In the following corollary we analyse the energy decay rate in the system of rotators. Let  $(p, q)(t)$  be a solution to (3.1). Below we denote  $W_t := W(p, q)(t)$  and  $H_t := H(p, q)(t)$ . Recall that  $\gamma_0 = 2N-1$ .

**Corollary 3.2.** *Under Assumption 3, there are  $N$ -dependent constants  $h_0 \geq 1$ ,  $0 < \varepsilon < 1$  and  $C > 0$  such that for  $H_0 \geq h_0$ ,*

$$(3.9) \quad W_t \leq W_0 - Ct W_0^{1/\gamma_0} \quad \text{for any} \quad 0 \leq t \leq \varepsilon H_0^{\gamma_0-1}$$

and

$$(3.10) \quad H_t \leq H_0 - \frac{Ct}{H_0^{\gamma_0-2}} \quad \text{for any} \quad \varepsilon^{-1} H_0^{\gamma_0-\frac{5}{2}} \leq t \leq \varepsilon H_0^{\gamma_0-1}.$$

We call  $CH^{-(\gamma_0-2)}$  energy decay rate of system (3.1) on the time interval from (3.10).

*Proof.* Let

$$t_{stop} = \max\{0 \leq t \leq \varepsilon H_0^{\gamma_0-1} : W_t \geq C_{stop}^{-1} W_0\}$$

for sufficiently large constant  $C_{stop}$ . Since by (3.7)  $W_0 \geq Ch_0^{\gamma_0}$ , for sufficiently large  $h_0$  by (3.8) we have

$$\dot{W}_t \leq -C W_t^{1/\gamma_0} \quad \text{for} \quad 0 \leq t \leq t_{stop}.$$

Integrating this inequality, we get

$$W_t^{(\gamma_0-1)/\gamma_0} \leq W_0^{(\gamma_0-1)/\gamma_0} - Ct, \quad t \leq t_{stop}.$$

Accordingly,

$$(3.11) \quad W_t \leq W_0 \left(1 - \frac{Ct}{W_0^{(\gamma_0-1)/\gamma_0}}\right)^{\gamma_0/(\gamma_0-1)} \leq W_0 \left(1 - \frac{C_1 t}{W_0^{(\gamma_0-1)/\gamma_0}}\right),$$

since the ratio  $\frac{C_1 t}{W_0^{(\gamma_0-1)/\gamma_0}}$  is small for  $t \leq t_{stop} \leq \varepsilon H_0^{\gamma_0-1}$  once  $\varepsilon$  is. Thus, we

get (3.9) for  $0 \leq t \leq t_{stop}$ . For  $t_{stop} \leq t \leq \varepsilon H_0^{\gamma_0-1}$  we have  $t \leq \varepsilon C W_0^{(\gamma_0-1)/\gamma_0}$ , so (3.9) also holds once  $\varepsilon$  is sufficiently small, since  $W_t \leq C_{stop}^{-1} W_0$  for  $t \geq t_{stop}$ .

Due to (3.7),  $H_t \leq H_0/2$  for  $t \geq t_{stop}$  once the constant  $C_{stop}$  is sufficiently large. Then, (3.10) holds true for  $t_{stop} \leq t \leq \varepsilon H_0^{\gamma_0-1}$ . By (3.7) and (3.11), for  $0 \leq t \leq t_{stop}$ ,

$$(3.12) \quad H_t \leq H_0 \left(\frac{1 + C H_0^{-3/2}}{1 - C H_t^{-3/2}}\right)^{1/\gamma_0} \left(1 - \frac{Ct}{H_0^{\gamma_0-1}}\right)^{1/\gamma_0} \\ \leq \left(1 + C_1 H_t^{-3/2}\right) \left(H_0 - \frac{C_2 t}{H_0^{\gamma_0-2}}\right) \leq H_0 (1 + C_3 H_0^{-3/2}) - \frac{C_2 t}{H_0^{\gamma_0-2}},$$

where we have used that  $C^{-1} H_0 \leq H_t \leq H_0$  in the considered time interval. Then, for  $t \geq \varepsilon^{-1} H_0^{\gamma_0 - \frac{5}{2}}$  we conclude (3.10).  $\square$

**3.2. Proof of the theorem.** We abbreviate  $V_j := V_j(q_j - q_{j+1})$ . Unless otherwise is explicitly stated, in this proof constants  $C, C_1, C_2, \dots$  never depend on the parameters  $a_i$ . Without loss of generality we assume that the energy  $H$  satisfies  $H \geq h_0$  for sufficiently large  $h_0$ . Indeed, for  $(p, q)$  such that  $H(p, q) < h_0$  the r.h.s. of equation (3.13) below is bounded by a constant  $C = C(h_0, a_0, \dots, a_{2N-2})$ .

*Step 1.* By (3.2),

$$L_F W = \\ - \left( a_0 \gamma_0 H^{\gamma_0-1} - \sum_{j=1}^{N-1} \left( a_{2j-1} \alpha_{2j-1} H^{\alpha_{2j-1}-1} p_j \xi_j + a_{2j} \alpha_{2j} H^{\alpha_{2j}-1} \xi_j L_f \xi_j \right) \right) p_1^2 \\ (3.13) \quad - \sum_{j=1}^{N-1} \left( a_{2j-1} H^{\alpha_{2j-1}} L_F(p_j \xi_j) + a_{2j} H^{\alpha_{2j}} L_F(\xi_j L_F \xi_j) \right).$$

We start with estimating the terms of the last sum in (3.13). By (3.3),

$$L_F(p_j \xi_j) = \xi_j L_F p_j + p_j L_F \xi_j = \xi_j (\xi_j - \xi_{j-1} - \delta_{j1} p_1) + p_j L_F \xi_j.$$

Let us bound  $|p_j L_F \xi_j| \leq \Gamma_j H p_j^2 + \Gamma_j^{-1} H^{-1} (L_F \xi_j)^2$ , where constants  $\Gamma_j$  will be chosen later to be sufficiently large. Then, applying twice the inequality  $|xy| \leq x^2/4 + y^2$  to  $xy = \xi_j \xi_{j-1}$  and  $xy = \xi_j (\delta_{j1} p_1)$ , we find

$$(3.14) \quad L_F(p_j \xi_j) \geq \frac{\xi_j^2}{2} - \xi_{j-1}^2 - \delta_{j1} p_1^2 - \Gamma_j H p_j^2 - \Gamma_j^{-1} H^{-1} (L_F \xi_j)^2.$$

Next, we note that

$$(3.15) \quad L_F \xi_j = -(p_j - p_{j+1}) V_j''.$$

Differentiation  $L_F$  applied to (3.15) gives us

$$L_F^2 \xi_j = -(p_j - p_{j+1})^2 V_j''' - (L_F p_j - L_F p_{j+1}) V_j''$$

and by (3.3),

$$(L_F^2 \xi_j)^2 \leq C(p_j^4 + p_{j+1}^4 + \xi_{j-1}^2 + \xi_j^2 + \xi_{j+1}^2 + \delta_{j1} p_1^2).$$

Then,

$$(3.16) \quad \begin{aligned} L_F(\xi_j L_F \xi_j) &= (L_F \xi_j)^2 + \xi_j L_F^2 \xi_j \geq (L_F \xi_j)^2 - a_{2j} H \xi_j^2 - (a_{2j} H)^{-1} (L_F^2 \xi_j)^2 \\ &\geq (L_F \xi_j)^2 - a_{2j} H \xi_j^2 \\ &\quad - C(a_{2j} H)^{-1} (p_j^4 + p_{j+1}^4 + \xi_{j-1}^2 + \xi_j^2 + \xi_{j+1}^2 + \delta_{j1} p_1^2). \end{aligned}$$

*Step 2.* Substituting the r.h.s. of estimates (3.14) and (3.16) into the last sum in (3.13), we bring together positive terms of the result into

$$(3.17) \quad I_{pos} := \sum_{j=1}^{N-1} \left( a_{2j-1} H^{\alpha_{2j-1}} \frac{\xi_j^2}{2} + a_{2j} H^{\alpha_{2j}} (L_F \xi_j)^2 \right).$$

Applying the inequality  $(x - y)^2 \geq x^2/2 - y^2$  to the squared (3.15), we find

$$(3.18) \quad (L_F \xi_j)^2 \geq \frac{(V_j'')^2}{2} p_{j+1}^2 - C p_j^2.$$

Then, using that  $\xi_j = -V_j'$  and  $H \geq p_{j+1}^2/2$ , we obtain by (3.18)

$$H \xi_j^2 + (L_F \xi_j)^2 \geq \frac{(V_j')^2 + (V_j'')^2}{2} p_{j+1}^2 - C p_j^2 \geq C_1^{-1} p_{j+1}^2 - C p_j^2,$$

according to Assumption 3. Recalling that  $\alpha_{2j-1} = \alpha_{2j} + 1$  and assuming  $a_{2j-1}/2 \geq a_{2j}$ , we find

$$I_{pos} \geq \sum_{j=1}^{N-1} a_{2j} H^{\alpha_{2j}} (C_1^{-1} p_{j+1}^2 - C p_j^2) \geq \frac{C_1^{-1}}{2} \sum_{j=2}^N a_{2j-2} H^{\alpha_{2j-2}} p_j^2 - C a_2 H^{\alpha_2} p_1^2.$$

Finally, we bound a half of the term  $I_{pos}$  by the estimate above and keep another half as it is,

$$(3.19) \quad I_{pos} \geq \frac{1}{2} \sum_{j=1}^{N-1} \left( a_{2j-1} H^{\alpha_{2j-1}} \frac{\xi_j^2}{2} + a_{2j} H^{\alpha_{2j}} (L_F \xi_j)^2 \right) \\ + C^{-1} \sum_{j=2}^N a_{2j-2} H^{\alpha_{2j-2}} p_j^2 - C a_2 H^{\alpha_2} p_1^2.$$

*Step 3.* Now we estimate the r.h.s. of (3.13) using bounds (3.6), (3.14), (3.16) and (3.19):

$$L_F W \leq -(I_{p_1} + I_\xi + I_{\tilde{\xi}} + I_p),$$

where

$$I_{p_1} = p_1^2 \left( a_0 \gamma_0 H^{\gamma_0-1} - C \sum_{j=1}^{N-1} (a_{2j-1} \alpha_{2j-1} H^{\alpha_{2j-1}-1/2} + a_{2j} \alpha_{2j} H^{\alpha_{2j}-1/2}) \right. \\ \left. - a_1 H^{\alpha_1} - \Gamma_1 a_1 H^{\alpha_1+1} - C H^{\alpha_2-1} (p_1^2 + 1) - C a_2 H^{\alpha_2} \right),$$

the term  $I_\xi$  is given by

$$I_\xi = \sum_{j=1}^{N-1} \xi_j^2 \left( \frac{a_{2j-1}}{4} H^{\alpha_{2j-1}} - a_{2j+1} H^{\alpha_{2j+1}} - a_{2j}^2 H^{\alpha_{2j}+1} - C (H^{\alpha_{2j-2}-1} + H^{\alpha_{2j}-1} + H^{\alpha_{2j+2}-1}) \right);$$

here and below we denote  $a_{2N-1} = a_{2N} = \Gamma_N := 0$  and  $\alpha_{2N-1} = \alpha_{2N} = \alpha_0 := -\infty$ . The term  $I_{\tilde{\xi}}$  has the form

$$I_{\tilde{\xi}} = \sum_{j=1}^{N-1} (L_F \xi_j)^2 \left( \frac{a_{2j}}{2} H^{\alpha_{2j}} - \Gamma_j^{-1} a_{2j-1} H^{\alpha_{2j-1}-1} \right)$$

and finally

$$I_p = \sum_{j=2}^N p_j^2 \left( C^{-1} a_{2j-2} H^{\alpha_{2j-2}} - \Gamma_j a_{2j-1} H^{\alpha_{2j-1}+1} - C p_j^2 (H^{\alpha_{2j-2}-1} + H^{\alpha_{2j}-1}) \right).$$

Let  $\Gamma_j = 2a_{2j-1}/a_{2j}$ . Then  $I_{\tilde{\xi}} = 0$ , since  $\alpha_{2j-1} - 1 = \alpha_{2j}$ . Next, using that  $p_j^2 \leq 2H$ , we find

$$I_p \geq \sum_{j=2}^N p_j^2 H^{\alpha_{2j-2}} \left( C^{-1} a_{2j-2} - \Gamma_j a_{2j-1} - C(1 + H^{-2}) \right) \\ \geq \sum_{j=2}^N p_j^2 H^{\alpha_{2j-2}} \geq H - C - \frac{p_1^2}{2},$$

once  $a_{2j-1}$  is sufficiently large in terms of  $\Gamma_j a_{2j-1} = 2a_{2j-1}^2/a_{2j}$ . Similarly,

$$I_\xi \geq \sum_{j=1}^{N-1} \xi_j^2 H^{\alpha_{2j-1}} \left( \frac{a_{2j-1}}{4} - a_{2j+1} H^{-2} - a_{2j}^2 - C(1 + H^{-2} + H^{-4}) \right) \geq 0,$$

once  $a_{2j-1}$  is sufficiently large in terms of  $a_{2j+1}$  and  $a_{2j}$ . Finally,

$$I_{p_1} \geq H^{\gamma_0-1} \frac{p_1^2}{2}$$

once  $a_0$  is sufficiently large.

The coefficients  $a_k$  satisfying the restrictions above can be chosen iteratively. Then, collecting together the estimates above, we get  $I_{p_1} + I_\xi + I_\xi + I_p \geq H - C$ , so

$$L_F W \leq -H + C.$$

The last inequality in (3.8) holds due to (3.7).  $\square$

#### 4. CHAIN OF OSCILLATORS

**4.1. The setting.** In this section we establish the existence of a strict Lyapunov function outside a compact set for the chain of oscillators in presence of friction in the first oscillator. To this end we show that Proposition 2.6 applies to the system under natural assumptions. We do not construct a Lyapunov function satisfying good bounds as we did for the chain of rotators in Section 3, since in the present case it can be done similarly.

The Hamiltonian of the chain of oscillators is

$$(4.1) \quad H(p, q) = \sum_{j=1}^N \frac{p_j^2}{2} + \sum_{j=1}^N U_j(q_j) + \sum_{j=1}^{N-1} V_j(q_j - q_{j+1}),$$

where  $N \geq 2$ ,  $p = (p_1, \dots, p_N) \in \mathbb{R}^N$  denote moments of the oscillators while  $q = (q_1, \dots, q_N) \in \mathbb{R}^N$  stand for their coordinates. For simplicity, we assume the interaction potentials  $V_k$  as well as the so-called pinning potentials  $U_k$  to be smooth. The equations of motion are given by (1.2), or, in more details, (4.2)

$$\dot{q}_j = p_j, \quad \dot{p}_j = -\delta_{j1} p_1 - U_j'(q_j) + V_{j-1}'(q_{j-1} - q_j) - V_j'(q_j - q_{j+1}), \quad 1 \leq j \leq N,$$

where  $V_0 = V_N := 0$ . If the Hamiltonian  $H$  is proper, it gives a non-strict Lyapunov function outside a compact set  $\mathcal{K}_Q^H$  with any  $Q \in \mathbb{R}$  since

$$L_F H = -p_1^2,$$

where  $F$  stands for the vector field from the r.h.s. of equation (4.2).

**4.2. Strictly convex case.** We start our analysis with a simple case of strictly convex potentials.

**Proposition 4.1.** *Suppose that for every  $k$  the potentials  $U_k(x)$  and  $V_k(x)$  are strictly convex and have minimum at  $x = 0$ . Then for any  $(p, q) \in \mathbb{R}^{2N} \setminus \{0\}$  there exists  $1 \leq p \leq 4N - 1$  such that  $L_F^p H \neq 0$ .*

**Corollary 4.2.** *Under assumptions of Proposition 4.1, system (4.2) satisfies Assumption 2 with  $W = H$ ,  $r = 4N - 1$  and  $Q = H(0, 0)$ . Hence it admits a strict Lyapunov function  $W^\sharp$  outside a compact set, constructed in Proposition 2.6.*

*Proof of Proposition 4.1.* Suppose that for some  $(p, q)$  we have  $L_F^k H = 0$  for any  $k \leq 4N - 1$ . Since

$$L_F^{k+1} H = - \sum_{j=0}^k \binom{k}{j} L_F^j p_1 L_F^{k-j} p_1,$$

the following vanishings are equivalent:

$$\forall 1 \leq k \leq 4N - 1 : L_F^k H = 0 \iff \forall 0 \leq k \leq 2N - 1 : L_F^k p_1 = 0.$$

Denote  $U_k = U_k(q_k)$  and  $V_k = V_k(q_k - q_{k+1})$  for short. Then

$$L_F p_1 = -U_1' - V_1' = 0 \quad \text{and} \quad L_F^2 p_1 = -(U_1'' + V_1'') p_1 + V_1'' p_2.$$

Since  $V_1'' \neq 0$ , if  $p_1 = L_F p_1 = 0$  then

$$\forall 2 \leq j \leq 2N - 1 : L_F^j p_1 = 0 \implies \forall 0 \leq j \leq 2N - 3 : L_F^j p_2 = 0.$$

Similarly,

$$L_F p_2 = -U_2' + V_1' - V_2' \quad \text{and} \quad L_F^2 p_2 = V_1'' p_1 - (U_2'' + V_2'' + V_1'') p_2 + V_2'' p_3.$$

Again, since  $V_2'' \neq 0$ , if  $p_1 = L_F p_1 = p_2 = L_F p_2 = 0$ , then

$$\forall 2 \leq j \leq 2N - 3 : L_F^j p_2 = 0 \implies \forall 0 \leq j \leq 2N - 5 : L_F^j p_3 = 0.$$

Proceeding by induction, we obtain

$$\forall 0 \leq j \leq 2N - 1 : L_F^j p_1 = 0 \implies \forall 1 \leq k \leq N : p_k = L_F p_k = 0.$$

So  $p = 0$  and equations  $L_F p_k = 0$  form the following system:

$$(4.3) \quad \begin{cases} U_1' + V_1' = 0 \\ U_2' - V_1' + V_2' = 0 \\ \dots \\ U_{N-1}' - V_{N-2}' + V_{N-1}' = 0 \\ U_N' - V_{N-1}' = 0. \end{cases}$$

Since  $x = 0$  is the unique minimum for the potentials  $U_k$ ,  $U_k'(x) > 0$  for  $x > 0$  and vice versa. Suppose that  $q_1 \geq 0$  (the case  $q_1 \leq 0$  is similar). Then  $U_1' \geq 0$  and the first equation implies

$$V_1' \leq 0 \implies q_2 \geq q_1 \geq 0 \implies U_2' \geq 0 \implies U_2' - V_1' \geq 0.$$

So the second equation implies

$$V_2' \leq 0 \implies q_3 \geq q_2 \geq 0 \implies U_3' \geq 0 \implies U_3' - V_2' \geq 0,$$

and so on. The penultimate equation implies

$$V_{N-1}' \leq 0 \implies q_N \geq q_{N-1} \geq 0 \implies U_N' \geq 0 \implies U_N' - V_{N-1}' \geq 0,$$

And the last equation is  $U_N' - V_{N-1}' = 0$ . Hence  $U_N' = 0$  and  $q_N = 0$ . Therefore  $0 = q_N \geq q_{N-1} \geq \dots \geq q_1 \geq 0$ . So  $q = 0$ .  $\square$

**4.3. General case.** Now we extend the argument to a more general case when potentials  $U_k$  and  $V_k$  are strictly convex outside a compact set.

**Theorem 4.3.** *Suppose that  $V_k''(x) > 0$  for  $|x| \geq R$  with some  $R \geq 0$ ,  $U_k'(x) \rightarrow \pm\infty$  as  $x \rightarrow \pm\infty$ , and  $V_k''(x) \neq 0$  or  $V_k'''(x) \neq 0$  at every  $x \in \mathbb{R}$ . Then there is a bounded set  $K \subset \mathbb{R}^{2N}$  such that for any  $(p, q) \in \mathbb{R}^{2N} \setminus K$  there exists  $1 \leq j \leq 3 \cdot 2^{N+1} - 5$  for which  $L_F^j H \neq 0$ .*

Note that under assumptions of the theorem, the Hamiltonian  $H$  is proper.

From the proof of the theorem it can be seen that

$$K \subset \{p = 0, -b < q_k < a \forall k\}$$

with  $a, b$  defined in (4.6), (4.7).

**Corollary 4.4.** *Under conditions of Theorem 4.3, system (4.2) satisfies Assumption 2 with  $W = H$ ,  $r = 3 \cdot 2^{N+1} - 5$  and  $Q$  such that  $\mathcal{K}_Q^H \supset K$ . Hence it admits a strict Lyapunov function  $W^\sharp$  outside a compact set, constructed in Proposition 2.6.*

Proof of Theorem 4.3 is based on the following three lemmata.

**Definition 4.5.** Let  $g : \mathbb{R}^{2N} \rightarrow \mathbb{R}$  be a smooth function. We say that  $g$  has order  $k$  at a point  $(p, q)$  (w.r.t system (4.2)) and write  $\text{ord}(g)(p, q) = k$  if  $L_F^j g(p, q) = 0$  for all  $0 \leq j \leq k - 1$  and  $L_F^k g(p, q) \neq 0$ . Sometimes we omit  $(p, q)$  and write  $\text{ord}(g)$  for brevity. If  $L_F^j g(p, q) = 0$  for all  $j$ , we write  $\text{ord}(g)(p, q) = \infty$ .

**Lemma 4.6.**

- if  $\text{ord}(g) > 0$ , then  $\text{ord}(L_F g) = \text{ord}(g) - 1$ ;
- $\text{ord}(g_1 g_2) = \text{ord}(g_1) + \text{ord}(g_2)$ ;
- $\text{ord}(g_1 + g_2) \geq \min\{\text{ord}(g_1), \text{ord}(g_2)\}$ ;
- if  $\text{ord}(f) \neq \text{ord}(g)$ , then  $\text{ord}(f + g) = \min\{\text{ord}(f), \text{ord}(g)\}$ .

*Proof.* Follows from the fact that  $L_F$  is a first order differential operator.  $\square$

**Lemma 4.7.** *If  $\text{ord}(H) \geq 3 \cdot 2^{N+1} - 5$  at a point  $(p, q)$ , then  $p = 0$  and  $q$  satisfies system (4.3).*

*Proof.* We again begin with the identity

$$L_F^{k+1} H = - \sum_{j=0}^k \binom{k}{j} L_F^j p_1 L_F^{k-j} p_1,$$

which implies

$$\text{ord}(p_1) = \frac{1}{2} \text{ord}(L_F H) \geq 3(2^N - 1).$$

Let us compute orders of  $p_k$  for  $k \geq 2$ . For  $p_1$  we obtain

$$L_F p_1 = -U_1' - V_1' = 0 \quad \text{and} \quad L_F^2 p_1 = -(U_1'' + V_1'')p_1 + V_1'' p_2.$$

Since  $\text{ord}(p_1) = \text{ord}(L_F^2 p_1) + 2$ , we obtain

$$\text{ord}(V_1'' p_2) = \text{ord}(L_F^2 p_1 + (U_1'' + V_1'')p_1) = \text{ord}(L_F^2 p_1).$$

Hence  $\text{ord}(p_2) \leq \text{ord}(p_1) - 2$ . If  $V_1'' \neq 0$ , then  $\text{ord}(p_2) = \text{ord}(p_1) - 2$ . On the other case  $V_1'' = 0$ , we have  $V_1''' \neq 0$  by the assumption of the theorem and  $\text{ord}(p_1) - 2 = \text{ord}(p_2) + \text{ord}(V_1''')$ . Since  $L_F V_1'' = V_1'''(p_1 - p_2)$ , we have  $\text{ord}(V_1''') = \text{ord}(p_2) + 1$ . So  $\text{ord}(p_1) - 2 = 2\text{ord}(p_2) + 1$  and  $\text{ord}(p_2) = \frac{1}{2}(\text{ord}(p_1) - 3)$ . Now we unite the both cases. Since  $\frac{1}{2}(j - 3) \leq j - 2$  for any  $j \geq 1$  and  $\text{ord}(p_1) \geq 1$ , we obtain

$$\text{ord}(p_1) - 2 \geq \text{ord}(p_2) \geq \frac{1}{2}(\text{ord}(p_1) - 3) \geq 3(2^{N-1} - 1).$$

Similarly,

$$L_F p_2 = -U_2' + V_1' - V_2' \quad \text{and} \quad L_F^2 p_2 = V_1'' p_1 - (U_2'' + V_2'' + V_1'') p_2 + V_2'' p_3$$

and

$$\text{ord}(p_2) - 2 \geq \text{ord}(p_3) \geq 3(2^{N-2} - 1).$$

Proceeding by induction, we obtain  $\text{ord}(p_k) \geq 3(2^{N-k+1} - 1)$ . In particular, since  $k \leq N$ , we have  $\text{ord}(p_k) \geq 3$ . Hence  $p_k = L_F p_k = 0$  for all  $k$ . Therefore,  $p = 0$  and  $q$  is a solution to system (4.3).  $\square$

**Lemma 4.8.** *Set of solutions to system (4.3) is bounded.*

*Proof.* Let

$$M := \max_{|x| \leq R} \max_{1 \leq k \leq N} |V_k'(x)| + 1 \geq 1.$$

Since  $V_k''(x) > 0$  for  $x \leq -R$ ,

$$(4.4) \quad x \leq R \quad \Rightarrow \quad V_k'(x) \leq M - 1 \quad \forall k.$$

Similarly, since  $V_k''(x) > 0$  for  $x \geq R$ ,

$$(4.5) \quad x \geq -R \quad \Rightarrow \quad V_k'(x) \geq -M + 1 \quad \forall k.$$

Since  $U_k'(x) \rightarrow \infty$  as  $x \rightarrow \infty$ , there is  $a > R$  such that

$$(4.6) \quad x \geq a \quad \Rightarrow \quad U_k'(x) \geq 2M \quad \forall k.$$

We claim that any solution  $q$  to system (4.3) satisfies  $q_k < a$  for all  $k$ . Indeed, let  $k$  be the smallest index for which  $q_k \geq a$ , so that in particular  $q_{k-1} < q_k$  (if  $k \neq 1$ ). Then,  $U_k'(q_k) \geq 2M$  due to (4.6), and  $V_{k-1}'(q_{k-1} - q_k) \leq M - 1$ , due to (4.4), where we set  $V_0' := 0$ . Then,

$$U_k'(q_k) - V_{k-1}'(q_{k-1} - q_k) \geq M + 1.$$

In view of the  $k$ -th equation from (4.3), this implies  $V_k'(q_k - q_{k+1}) \leq -M - 1$ . Accordingly, by (4.5),  $q_k - q_{k+1} < -R$ , so  $q_{k+1} > q_k \geq a$ . Continuing by induction,  $N - 1$ -st equation in (4.3) gives  $q_N > q_{N-1} \geq a$ , so

$$U_N'(q_N) - V_{N-1}'(q_{N-1} - q_N) \geq 2M - (M - 1) \geq 2$$

which contradicts to the last equation in (4.3).

Similarly we show that every solution  $q$  is bounded from below by  $-b < -R$  such that

$$(4.7) \quad x \leq -b \quad \Rightarrow \quad U_k^l(x) \leq -2M \quad \forall k.$$

□

*Proof of Theorem 4.3.* If  $L_F^j H(p, q) = 0$  for all  $1 \leq j \leq 3 \cdot 2^{N+1} - 5$  at a point  $(p, q)$ , then  $p = 0$  and  $q$  satisfies (4.3) by Lemma 4.7. Hence  $q$  is bounded by Lemma 4.8.

□

## 5. MIXING IN CHAINS COUPLED TO LANGEVIN THERMOSTATS

Let us consider either the chain of rotators with Hamiltonian (1.1) or the chain of oscillators with Hamiltonian (4.1). We couple the first and the last particles in the chain with Langevin thermostats of positive temperatures  $T_1$  and  $T_N$  correspondingly. That is, we consider the system of stochastic differential equations

$$(5.1) \quad \dot{q}_j = p_j, \quad \dot{p}_j = -\partial_{q_j} H + (\delta_{1j} + \delta_{Nj})(-p_j + \sqrt{2T_j} \dot{\beta}_j(t)), \quad 1 \leq j \leq N,$$

where  $\beta_j(t)$  are standard independent Brownian motions. The initial conditions  $(p_0, q_0)$  are assumed to be random and independent from the Brownian motions  $\beta_j$ .

Denote by  $\mathcal{M} \ni (p, q)$  the phase space of the system, i.e.  $\mathcal{M} = \mathbb{R}^N \times \mathbb{T}^N$  in the case of rotators and  $\mathcal{M} = \mathbb{R}^{2N}$  in the case of oscillators. Let  $\mu$  be a probability measure on  $\mathcal{M}$  and  $(p, q)(t)$  be a solution to (5.1) with initial conditions  $(p_0, q_0)$ , distributed accordingly to the measure  $\mu$ . If the distribution of solution at time  $t$ , denoted by  $\mathcal{D}(p, q)(t)$ , satisfies

$$\mathcal{D}(p, q)(t) \equiv \mu \quad \text{for all } t \geq 0,$$

then the measure  $\mu$  is called *stationary*. If equation (5.1) has a unique stationary measure  $\mu$  and distribution  $\mathcal{D}(p, q)(t)$  of *any*<sup>4</sup> its solution  $(p, q)(t)$  weakly converges to  $\mu$  as  $t \rightarrow \infty$ , then the equation is called *mixing*.

Establishing the mixing property for equation (5.1) is a problem of prime interest in non-equilibrium statistical mechanics of solids. Let us first consider the case of oscillators and assume that the potentials  $V_j$  and  $U_j$  from (4.1) behave at infinity polynomially. Then, due to our knowledge, for chains of arbitrary length  $N$  the mixing property is proven only in the case when the interaction potentials  $V_j$  are "not weaker" than the pinning potentials  $U_j$ , i.e.  $V_j$  behave at infinity as polynomials of the same or higher degree than those associated with the potentials  $U_j$  [12, 11, 19, 3, 5]. In the opposite case when the pinning dominates, the mixing is proven only for a chain of  $N = 3$  oscillators [13].

As for the chain of rotators, in [6, 4] the authors argue that it can be viewed as a system of oscillators with extremely strong pinning potentials.

---

<sup>4</sup>With "reasonable" initial conditions, e.g. with finite second moment.

They establish the mixing property for the chains of length  $N = 3, 4$  while for longer chains the problem remains open. The reason why it is so hard to prove the mixing in the chains of rotators or oscillators with "strong" pinning, is in the discussed in the Introduction decoupling of "fast" rotators (or oscillators) from their "slow" neighbours, leading to extremely slow energy transport through the chain. In particular, as a result the rate of mixing turns out to be slower than exponential, in contrast to the case of oscillators with "strong" interaction, see [8] additionally to the above mentioned works.

Let us note that for chains in which *each* particle interacts with its own Langevin thermostat, the mixing property is well-understood.

One of the most effective approaches to problems of this kind relies on the fact that to establish the mixing property it suffices to construct a strict Lyapunov function  $W : \mathcal{M} \mapsto \mathbb{R}$  outside a compact set for the generator  $\mathcal{L}^{T_1, T_2}$  of equation (5.1). More specifically, it suffices to find a sufficiently smooth proper function  $W$ , satisfying

$$\mathcal{L}^{T_1, T_2} W \leq -\psi \circ W + C,$$

where  $C$  is a constant and  $\psi$  is a positive increasing function, see for details Theorem 2.1 in [6] and references therein. We recall that action of the generator  $\mathcal{L}^{T_1, T_2}$  on a  $C^2$ -smooth function  $f(p, q)$  can be computed as

$$\mathcal{L}^{T_1, T_2} f = L_F f + T_1 \frac{\partial^2 f}{\partial p_1^2} + T_N \frac{\partial^2 f}{\partial p_N^2},$$

where  $L_F$  stands for the Lie derivative along the vector field given by the deterministic part of equation (5.1).

For the chain of oscillators with "strong" interaction which dominates the pinning, the mixing in [3, 5] is established by proving that  $\exp(\theta H)$  gives the desired Lyapunov function once  $\theta$  is small enough. In the opposite case as well as in the case of rotators the employed arguments do not work. Instead, in [6, 4] the authors construct a suitable Lyapunov functions for the chains of rotators of lengths  $N = 3$  and 4 using a method related to the discussed in the Introduction KAM-like approach subsequently employed in [7]. For the moment of writing it did not allow to prove the mixing property for longer chains since for them it is developed only for specific regime when the most part of energy is concentrated in a single rotator.

Now let us consider equation similar to (5.1), in which the Langevin thermostat is coupled only to the first oscillator, i.e. in (5.1) the factor  $\delta_{1j} + \delta_{Nj}$  is replaced by  $\delta_{1j}$ . Its generator  $\mathcal{L}^{T_1}$  acts to a  $C^2$ -smooth function  $f(p, q)$  as  $\mathcal{L}^{T_1} f = L_F f + T_1 \frac{\partial^2 f}{\partial p_1^2}$ . Then, being applied to the Lyapunov function  $W$  constructed in the Theorem 3.1, it satisfies

$$(5.2) \quad \mathcal{L}^{T_1} W \leq -H + C + T_1 \frac{\partial^2 W}{\partial p_1^2}.$$

By definition (3.4) of  $W$ ,

$$\frac{\partial^2 W}{\partial p_1^2} \sim a_0 \gamma_0 H^{\gamma_0 - 1} = a_0 \gamma_0 H^{2N - 2} \quad \text{for } H \gg 1.$$

Then, already for  $N = 2$  the r.h.s. of (5.2) growth as  $H^2$ , so  $W$  is *not* a Lyapunov function for the generator  $\mathcal{L}^{T_1}$ . Similarly one can see that our approach does not allow to prove mixing for chain (5.1) even for  $N = 3$ , in contrast to the method suggested in [6, 4, 7]. However, our construction is very explicit, simple and universal, and it allows to get the rate of energy dissipation seemingly achieved in the worst scenario. We hope that its suitable modification could help to construct the desired Lyapunov functions for generators of systems (5.1).

**Funding.** This work was supported by the Russian Science Foundation under grant no. 19-71-30012, <https://rscf.ru/en/project/23-71-33002/>.

#### REFERENCES

- [1] F. Bonetto, J. L. Lebowitz, J. Lukkarinen, *Fourier's Law for a Harmonic Crystal with Self-Consistent Stochastic Reservoirs*, Journal of Statistical Physics, **116** (2004), 783-813.
- [2] F. Bonetto, J.L. Lebowitz, L. Rey-Bellet, *Fourier's law: a challenge to theorists*, Mathematical Physics 2000, Imp. Coll. Press, London, (2000), 128-150.
- [3] P. Carmona, *Existence and uniqueness of an invariant measure for a chain of oscillators in contact with two heat baths*, Stochastic Process. Appl. **117** (2007), 1076-1092.
- [4] N. Cuneo and J.-P. Eckmann, *Non-equilibrium steady states for chains of four rotors*, Comm. Math. Phys. **345** (2016), 185-221.
- [5] N. Cuneo, J.-P. Eckmann, M. Hairer, L. Rey-Bellet, *Non-equilibrium steady states for networks of oscillators*, Electron. J. Probab. **23** (2018), 1-28.
- [6] N. Cuneo, J.-P. Eckmann, and C. Poquet, *Non-equilibrium steady state and subgeometric ergodicity for a chain of three coupled rotors*, Nonlinearity **28** (2015), 2397-2421.
- [7] N. Cuneo, J.-P. Eckmann and C. E. Wayne, *Energy Dissipation in Hamiltonian Chains of Rotators*, Nonlinearity **30** R81 (2017).
- [8] N. Cuneo and C. Poquet, *On the relaxation rate of short chains of rotors interacting with Langevin thermostats*, Electron. Commun. Probab. **22** (2017), 1-8.
- [9] D. Dolgopyat, C. Liverani, *Energy Transfer in a Fast-Slow Hamiltonian System*, Commun. Math. Phys., **308** (2011), 201-225.
- [10] A. Dymov, *Nonequilibrium Statistical Mechanics of Weakly Stochastically Perturbed System of Oscillators*, Ann. Henri Poincaré **17** (2016), 1825-1882.
- [11] J.-P. Eckmann and M. Hairer, *Non-equilibrium statistical mechanics of strongly anharmonic chains of oscillators*, Comm. Math. Phys. **212** (2000), 105-164.
- [12] J.-P. Eckmann, C.-A. Pillet, and L. Rey-Bellet, *Non-equilibrium statistical mechanics of anharmonic chains coupled to two heat baths at different temperatures*, Comm. Math. Phys. **201** (1999), 657-697.
- [13] M. Hairer and J. C. Mattingly, *Slow energy dissipation in anharmonic oscillator chains*, Comm. Pure Appl. Math. **62** (2009), 999-1032.
- [14] H.K. Khalil, *Nonlinear systems*, Prentice-Hall 2002.
- [15] S. Lepri (ed.), *Thermal Transport in Low Dimensions: From Statistical Physics to Nanoscale Heat Transfer*, Lecture Notes in Physics, **21** 2016.

- [16] C. Liverani, S. Olla, *Toward the Fourier law for a weakly interacting anharmonic crystal*, AMS, **25** (2012), 555-583.
- [17] M. Malisoff, F. Mazenc, *Construction of strict Lyapunov functions*, Springer-Verlag 2009.
- [18] J.L. Massera, *On Liapounoff's conditions of stability*, Ann. of Math. **50** (1949), 705-721.
- [19] L. Rey-Bellet and L. E. Thomas, *Exponential convergence to non-equilibrium stationary states in classical statistical mechanics*, Comm. Math. Phys. **225** (2002), 305-329.
- [20] D.A. Ruelle, *Mechanical Model for Fourier's Law of Heat Conduction*, Commun. Math. Phys., **311** (2012), 755-768.

STEKLOV MATHEMATICAL INSTITUTE OF RUSSIAN ACADEMY OF SCIENCES, MOSCOW  
119991, RUSSIA & NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS,  
MOSCOW 119048, RUSSIA

*Email address:* `dymov@mi-ras.ru`

STEKLOV MATHEMATICAL INSTITUTE OF RUSSIAN ACADEMY OF SCIENCES, MOSCOW  
119991, RUSSIA

*Email address:* `lion.lokut@gmail.com`

DIPARTIMENTO DI MATEMATICA E INFORMATICA U. DINI, UNIVERSITÀ DI FIRENZE,  
FIRENZE 50134, ITALIA

*Email address:* `andrey.sarychev@unifi.it`