

On p -torsions of geometric Brauer groups

Zhenghui Li, Yanshuai Qin

July 1, 2024

Abstract

Let X be a smooth projective integral variety over a finitely generated field k of characteristic $p > 0$. We show that the finiteness of the exponent of the p -primary part of $\mathrm{Br}(X_{k^s})^{G_k}$ is equivalent to the Tate conjecture for divisors, generalizing D’Addezio’s theorem for abelian varieties to arbitrary smooth projective varieties. As a result, we show that the cokernel of $\mathrm{Br}_{\mathrm{nr}}(K(X)) \rightarrow \mathrm{Br}(X_{k^s})^{G_k}$ is of finite exponent and complete the p -primary part of the generalization of Artin-Grothendieck’s theorem to higher relative dimensions.

1 Introduction

For any regular noetherian scheme X , the *cohomological Brauer group* $\mathrm{Br}(X)$ is defined to be the étale cohomology group $H_{\mathrm{et}}^2(X, \mathbb{G}_m)$. Let K be a finitely generated field extension. The *unramified Brauer group* $\mathrm{Br}_{\mathrm{nr}}(K)$ is defined as the intersection of Brauer groups of all DVRs with fraction field K . By the purity of Brauer groups [15], if K admits a proper regular model X , then $\mathrm{Br}_{\mathrm{nr}}(K) = \mathrm{Br}(X)$.

Theorem 1.1. *Let k be finitely generated field of characteristic $p > 0$. Let X/k be a smooth projective geometrically integral variety. Then, the finiteness of the exponent of $\mathrm{Br}(X_{k^s})^{G_k}[p^\infty]$ is equivalent to the Tate conjecture for divisors on X and the natural map $\mathrm{Br}_{\mathrm{nr}}(K(X)) \rightarrow \mathrm{Br}(X_{k^s})^{G_k}$ has a cokernel of finite exponent.*

Corollary 1.2. *Let X be an abelian variety or a K3 surface (or any smooth projective variety satisfying the Tate conjecture for divisors) over a finitely generated field k of*

Key words: Brauer group, Tate conjecture, Tate-Shafarevich group

MSC classes: 11G40, 14G17, 14J20, 11G25, 11G35

The paper is written when the first named author is in his Ph.D program in Sorbonne Université. This program has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 945332 and partially funded by the project “Group schemes, root systems, and related representations” founded by the European Union - NextGenerationEU through Romania’s National Recovery and Resilience Plan (PNRR) call no. PNRR-III-C9-2023- I8, Project CF159/31.07.2023. The second named author is supported by the DFG through CRC1085 Higher Invariants (University of Regensburg).

characteristic $p > 0$. Then the geometric Brauer group $\mathrm{Br}(X_{k^s})^{G_k}$ is of finite exponent. Moreover, it is finite if the Witt-vector cohomology $H^2(X_{\bar{k}}, W\mathcal{O}_{X_{\bar{k}}})$ is finitely generated over $W(\bar{k})$.

The prime-to- p part of the corollary was proved by Skorobogatov and Zarhin [30, 31] for abelian varieties and K3 surfaces and by Cadoret-Hui-Tamagawa [8] in general. The p -primary part for abelian varieties was prove by D'Addezio [10] by a different method which seems only working for abelian varieties. For an abelian group M and a prime ℓ , denote by $V_\ell M$ the rational Tate-module $\varprojlim_n M[\ell^n] \otimes_{\mathbb{Z}_\ell} \mathbb{Q}_\ell$.

Corollary 1.3. *Let X be a smooth projective geometrically integral variety over a finitely generated field k . Then there exists a canonical exact sequence*

$$0 \longrightarrow V_\ell \mathrm{Br}_{\mathrm{nr}}(k) \longrightarrow \mathrm{Ker}(V_\ell \mathrm{Br}_{\mathrm{nr}}(K(X))) \longrightarrow V_\ell \mathrm{Br}(X_{k^s})^{G_k} \longrightarrow V_\ell \mathrm{III}(\mathrm{Pic}_{X/k, \mathrm{red}}^0) \longrightarrow 0.$$

Moreover, the natural map

$$\mathrm{Br}_{\mathrm{nr}}(K(X)) \rightarrow \mathrm{Br}(X_{k^s})^{G_k}$$

has a cokernel of finite exponent.

This result is a higher dimensional generalization of Artin-Grothendieck's theorem [17, § 4] on the relation between Brauer groups and Tate-Shafarevich groups. The prime-to- p part ($p = \mathrm{char}(k)$) of theorem was proved by the second named author [29, 28] for k being a global field and by Ertl-Keller-Qin [13] in general.

Corollary 1.4. *Let $\mathcal{X} \rightarrow \mathcal{Y}$ be a projective surjective k -morphism between smooth projective varieties over a finitely generated field k of characteristic $p > 0$. Assume that its generic fiber X/K is smooth and geometrically integral. Then*

$$\mathrm{Br}(\mathcal{X}_{k^s})^{G_k} \longrightarrow \mathrm{Br}(X_{K^s})^{G_K}$$

has a cokernel of finite exponent.

Outline of Proofs

Firstly, by spreading out X/k , we get a smooth projective morphism $f : \mathcal{X} \rightarrow U$ between smooth integral varieties over \mathbb{F}_p . On the one hand, Pál reformulated the Tate conjecture using rigid cohomology [27, § 6]: the natural map

$$\mathrm{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_p \longrightarrow H_{\mathrm{rig}}^0(U, R^2 f_* \mathcal{O}_{\mathcal{X}}^\dagger)^{F=p}$$

is surjective. On the other hand, there is a canonical exact sequence

$$0 \longrightarrow \mathrm{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_p \longrightarrow \varprojlim_n H_{\mathrm{fppf}}^0(\mathrm{Spec}(k), R^2 f_* \mu_{p^n}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p \longrightarrow V_p \mathrm{Br}(X_{k^s})^{G_k} \longrightarrow 0$$

It remains to prove

$$\varprojlim_n H_{\mathrm{fppf}}^0(\mathrm{Spec}(k), R^2 f_* \mu_{p^n}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p \cong H_{\mathrm{rig}}^0(U, R^2 f_* \mathcal{O}_{\mathcal{X}}^\dagger)^{F=p}.$$

Secondly, we use the syntomic sheaf S_n^1 defined by Bauer in [2] which is isomorphic to the sheaf μ_{p^n} on the small syntomic site of X to relate the flat cohomology to crystalline cohomology. There is an exact sequence of sheaves on the small syntomic site [2, Thm. 3.4]

$$0 \longrightarrow \mu_{p^n} \longrightarrow \mathcal{I}_n \xrightarrow{1-p^{-1}F} \mathcal{O}_n^{\text{crys}} \longrightarrow 0.$$

It gives a natural map

$$H_{\text{syn}}^0(\text{Spec}(k), R^2 f_{\text{syn},*} \mu_{p^n}) \longrightarrow H_{\text{syn}}^0(\text{Spec}(k), R^2 f_{\text{syn},*} \mathcal{I}_n)^{p^{-1}F=1}.$$

The second term is essentially isomorphic to $H_{\text{crys}}^0(\text{Spec}(k), R^2 f_{\text{crys},*} \mathcal{O}_{X/\Sigma_n})^{F=p}$ up to groups of finite exponent (independent of n). To show the natural map is an isomorphism up to groups of finite exponent (independent of n), we use a pull-back trick developed by Colliot-Thélène and Skorobogatov in [9]. This is divided into three steps. First, we show that the claim is true for curves by a direct computation. Second, we can take the Albanese variety of X and show that the natural morphism between $R^1 f_*$ induced by the Albanese map is an isomorphism up to group of finite exponent. Third, we use the pull-back trick to show that the natural map above is an isomorphism up to groups of finite exponent (independent of n).

Notations

Fields

By a *finitely generated field*, we mean a field which is finitely generated over a prime field. For any field k , denote by k^s the separable closure. Denote by $G_k = \text{Gal}(k^s/k)$ the absolute Galois group of k .

Varieties

By a *variety* over a field k , we mean a scheme which is separated and of finite type over k . A morphism between varieties over k is a k -morphism.

For a smooth proper geometrically connected variety X over a field k , we use $\text{Pic}_{X/k}^0$ to denote the identity component of the Picard scheme $\text{Pic}_{X/k}$. Denote by $\text{Pic}_{X/k,\text{red}}^0$ the underlying reduced closed subscheme of $\text{Pic}_{X/k}^0$ (which is known to be an abelian variety cf. [16, p236-p17, Cor. 3.2.]). The Néron-Severi group $\text{NS}(X)$ of X is defined as the quotient $\text{Pic}(X)/\text{Pic}^0(X)$.

Cohomology

The default sheaves and cohomology over schemes are with respect to the small étale site. So H^i is the abbreviation of H_{et}^i . H_{fppf}^i denotes the cohomology of sheaves on the fppf site.

Brauer groups

For any noetherian scheme X , denote the *cohomological Brauer group*

$$\mathrm{Br}(X) := H^2(X, \mathbb{G}_m)_{\mathrm{tor}}.$$

Abelian group

For any abelian group M , integer m and prime ℓ , we set

$$\begin{aligned} M[m] &= \{x \in M \mid mx = 0\}, & M_{\mathrm{tor}} &= \bigcup_{m \geq 1} M[m], & M[\ell^\infty] &= \bigcup_{n \geq 1} M[\ell^n], \\ T_\ell M &= \mathrm{Hom}_{\mathbb{Z}}(\mathbb{Q}_\ell/\mathbb{Z}_\ell, M) = \varprojlim_n M[\ell^n], & V_\ell M &= T_\ell(M) \otimes_{\mathbb{Z}_\ell} \mathbb{Q}_\ell. \end{aligned}$$

For an abelian group M and a prime number p , we set

$$M(\mathrm{non-}p) = \bigcup_{p \nmid m} M[m].$$

Let $f : M \rightarrow N$ be a morphism in an abelian category, we say that f is *al. injective* (resp. *al. surjective*) if the kernel (resp. cokernel) can be killed by a positive integer. We say that f is an *al. isomorphism* iff it is al. injective and al. surjective. If f_n are a family of morphisms in an abelian category indexed by a set I , we say f is *al. injective uniformly in n* (resp. *al. surjective uniformly in n*) if the kernels (resp. cokernels) can be killed by the same positive integer simultaneously. We say that f is an *al. isomorphism uniformly in n* iff it is al. injective uniformly in n and al. surjective uniformly in n . For an object or a morphism (resp. a family of them) in an abelian category, we say that it is al. zero (resp. al. zero uniformly in n) if it can be killed by some positive integer (resp. by the same positive integer).

Crystalline sites and Topos

For the crystalline site and topos, we use the notation in [7]. Let $S = (S, I, \gamma)$ be a PD-scheme and X be an S -scheme to which γ extends, we use $\mathrm{Crys}(X/S)$ to denote the small crystalline(-Zariski) site and $(X/S)_{\mathrm{crys}}$ to denote the topos. If S is clear, we may omit it and write it as X_{crys} . For $\tau \in \{\mathrm{ZAR}, \mathrm{SYN}\}$, we use $\mathrm{CRYST}(X/S)_\tau$ to denote the big crystalline site equipped with τ -topology and $(X/S)_{\mathrm{CRYST}, \tau}$ to denote the corresponding topos.

2 Preliminaries

Let k be a field. Let $f : X \rightarrow S = \mathrm{Spec}(k)$ be a smooth projective geometrically integral variety. Let S' denote $\mathrm{Spec}(k^s)$. The goal of this section is to prove the following proposition

Proposition 2.1. *There is a canonical exact sequence*

$$0 \longrightarrow \mathrm{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_p \longrightarrow \varprojlim_n H_{\mathrm{fppf}}^0(S, R^2 f_* \mu_{p^n}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p \longrightarrow V_p \mathrm{Br}(X_{k^s})^{G_k} \longrightarrow 0,$$

where $R^2 f_* \mu_{p^n}$ is the higher direct image of the fppf sheaf μ_{p^n} on X .

Proof. Let $f : X \longrightarrow S = \mathrm{Spec}(k)$ as in the beginning. Consider the Leray spectral sequence for the fppf sheaf μ_{p^n} on X

$$E_2^{p,q} = H_{\mathrm{fppf}}^p(S, R^q f_* \mu_{p^n}) \Rightarrow H_{\mathrm{fppf}}^{p+q}(X, \mu_{p^n}).$$

It gives an edge map

$$H_{\mathrm{fppf}}^2(X, \mu_{p^n}) \longrightarrow H_{\mathrm{fppf}}^0(S, R^2 f_* \mu_{p^n}).$$

Consider the Leray spectral sequence for the fppf sheaf \mathbb{G}_m on X_{k^s}

$$E_2^{p,q} = H_{\mathrm{fppf}}^p(S', R^q f_* \mathbb{G}_m) \Rightarrow H_{\mathrm{fppf}}^{p+q}(X_{k^s}, \mathbb{G}_m).$$

Since $f_* \mathbb{G}_m = \mathbb{G}_m$, $E_2^{p,0} = H^p(S', \mathbb{G}_m) = 0$ for $p \geq 1$. There is an exact sequence

$$0 \longrightarrow H_{\mathrm{fppf}}^1(S', R^1 f_* \mathbb{G}_m) \longrightarrow H_{\mathrm{fppf}}^2(X_{k^s}, \mathbb{G}_m) \longrightarrow H_{\mathrm{fppf}}^0(S', R^2 f_* \mathbb{G}_m) \longrightarrow H_{\mathrm{fppf}}^2(S', R^1 f_* \mathbb{G}_m).$$

By Lemma 2.2 below,

$$\mathrm{Br}(X_{k^s}) = H_{\mathrm{fppf}}^2(X_{k^s}, \mathbb{G}_m) \longrightarrow H_{\mathrm{fppf}}^0(S', R^2 f_* \mathbb{G}_m)$$

is surjective and has a kernel of finite exponent. Consider the commutative diagram

$$\begin{array}{ccc} H_{\mathrm{fppf}}^2(X, \mu_{p^n}) & \longrightarrow & H_{\mathrm{fppf}}^0(S, R^2 f_* \mu_{p^n}) \\ \downarrow & & \downarrow \\ \mathrm{Br}(X_{k^s})^{G_k}[p^n] & \longrightarrow & H_{\mathrm{fppf}}^0(S', R^2 f_* \mathbb{G}_m)^{G_k}[p^n] \end{array}$$

The second row is al. isomorphism uniformly in n . By Lemma 2.5, the first column is al. surjective uniformly in n . Thus, the second column is al. surjective uniformly in n . The Kummer exact sequence on the fppf site of X

$$0 \longrightarrow \mu_p^n \longrightarrow \mathbb{G}_m \xrightarrow{p^n} \mathbb{G}_m \longrightarrow 0$$

gives an exact sequence of fppf sheaves

$$0 \longrightarrow R^1 f_* \mathbb{G}_m / p^n \longrightarrow R^2 f_* \mu_p^n \longrightarrow R^2 f_* \mathbb{G}_m.$$

Hence we get

$$0 \longrightarrow H_{\mathrm{fppf}}^0(S, R^1 f_* \mathbb{G}_m / p^n) \longrightarrow H_{\mathrm{fppf}}^0(S, R^2 f_* \mu_p^n) \longrightarrow H_{\mathrm{fppf}}^0(S, R^2 f_* \mathbb{G}_m).$$

By [10, Lem. 3.2], the natural map $H_{\text{fppf}}^0(S, R^2 f_* \mathbb{G}_m) \rightarrow H_{\text{fppf}}^0(S', R^2 f_* \mathbb{G}_m)^{G_k}$ is injective. This gives an exact sequence with the last map al. surjective uniformly in n

$$0 \rightarrow H_{\text{fppf}}^0(S, R^1 f_* \mathbb{G}_m / p^n) \rightarrow H_{\text{fppf}}^0(S, R^2 f_* \mu_p^n) \rightarrow H_{\text{fppf}}^0(S', R^2 f_* \mathbb{G}_m)^{G_k} [p^n].$$

By Lemma 2.4 below, $H_{\text{fppf}}^0(S, R^1 f_* \mathbb{G}_m / p^n)$ is al. isomorphic to $(\text{NS}(X_{\bar{k}}) / p^n)^{G_k}$ uniformly in n . Taking inverse limit and then tensoring \mathbb{Q}_p , we get

$$0 \rightarrow (\text{NS}(X_{\bar{k}}) \otimes_{\mathbb{Z}} \mathbb{Q}_p)^{G_k} \rightarrow \varprojlim_n H_{\text{fppf}}^0(S, R^2 f_* \mu_p^n) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p \rightarrow V_p H_{\text{fppf}}^0(S', R^2 f_* \mathbb{G}_m)^{G_k} \rightarrow 0.$$

This completes the proof since $V_p \text{Br}(X_{k^s})^{G_k} \cong V_p H_{\text{fppf}}^0(S', R^2 f_* \mathbb{G}_m)^{G_k}$ by arguments above and $\text{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_p \cong (\text{NS}(X_{\bar{k}}) \otimes_{\mathbb{Z}} \mathbb{Q}_p)^{G_k}$ by Lemma 2.4 below. \square

Lemma 2.2. $H_{\text{fppf}}^1(S', R^1 f_* \mathbb{G}_m)$ is of finite exponent and $H_{\text{fppf}}^2(S', R^1 f_* \mathbb{G}_m) = 0$.

Proof. $R^1 f_* \mathbb{G}_m$ is represented by the Picard scheme $P := \text{Pic}_{X/k}$. Write P^0 for $\text{Pic}_{X/k}^0$ and P_{red}^0 for $\text{Pic}_{X/k, \text{red}}^0$. By [16], there are exact sequences

$$0 \rightarrow P^0 \rightarrow P \rightarrow \text{NS}_X \rightarrow 0,$$

and

$$0 \rightarrow P_{\text{red}}^0 \rightarrow P^0 \rightarrow T \rightarrow 0,$$

where NS_X is an étale group scheme over k ($\text{NS}_X(k^s) = \text{NS}(X_{\bar{k}})$) and T is a finite group scheme over k . Since P_{red}^0 and NS_X are smooth group over k ,

$$H_{\text{fppf}}^i(S', P_{\text{red}}^0) = 0 \text{ and } H_{\text{fppf}}^i(S', \text{NS}_X) = 0$$

for $i \geq 1$. By [23, Prop. 5.1], there is an exact sequence

$$0 \rightarrow T \rightarrow G_0 \rightarrow G_1 \rightarrow 0,$$

where G_0 and G_1 are smooth group of finite type over k . Then,

$$H_{\text{fppf}}^i(S', T) = 0 \text{ for } i \geq 2.$$

The claims follow from the long exact sequences associated to two exact sequences. \square

Corollary 2.3. *The kernel of the natural map $\text{Br}(X_{k^s}) \rightarrow \text{Br}(X_{\bar{k}})$ is isomorphic to $H_{\text{fppf}}^1(S', R^1 f_* \mathbb{G}_m)$, which is of finite exponent and vanishes if $\text{Pic}_{X/k}$ is smooth over k .*

Proof.

$$0 \rightarrow H_{\text{fppf}}^1(S', R^1 f_* \mathbb{G}_m) \rightarrow H_{\text{fppf}}^2(X_{k^s}, \mathbb{G}_m) \rightarrow H_{\text{fppf}}^0(S', R^2 f_* \mathbb{G}_m)$$

is exact. By the lemma above, $H_{\text{fppf}}^1(S', R^1 f_* \mathbb{G}_m)$ is of finite exponent. It suffices to show that

$$H_{\text{fppf}}^0(S', R^2 f_* \mathbb{G}_m) \rightarrow H_{\text{fppf}}^0(\text{Spec}(\bar{k}), R^2 f_* \mathbb{G}_m) = \text{Br}(X_{\bar{k}})$$

is injective. This follows from [10, Lem. 3.2]. \square

Lemma 2.4. *There is a natural map which is al. isomorphism uniformly in n :*

$$H_{\text{fppf}}^0(S, R^1 f_* \mathbb{G}_m/p^n) \longrightarrow (\text{NS}(X_{\bar{k}})/p^n)^{G_k}.$$

Moreover, there is a canonical isomorphism

$$\text{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_p \cong (\text{NS}(X_{\bar{k}}) \otimes_{\mathbb{Z}} \mathbb{Q}_p)^{G_k}.$$

Proof. Notations as in the proof of Lemma 2.2,

$$R^1 f_* \mathbb{G}_m/p^n = P/p^n.$$

There are exact sequences of fppf sheaves

$$P^0/p^n \longrightarrow P/p^n \longrightarrow \text{NS}_X/p^n \longrightarrow 0,$$

and

$$P_{\text{red}}^0/p^n \longrightarrow P^0/p^n \longrightarrow T/p^n \longrightarrow 0.$$

$P_{\text{red}}^0/p^n = 0$ since P_{red}^0 is an abelian variety. $P^0/p^n \cong T/p^n$ is killed by the order of T . Then,

$$H_{\text{fppf}}^0(S, P/p^n) \longrightarrow H_{\text{fppf}}^0(S, \text{NS}_X/p^n)$$

is an al. isomorphism uniformly in n . Since NS_X/p^n is an étale group scheme over k ,

$$H_{\text{fppf}}^0(S, \text{NS}_X/p^n) = H_{\text{et}}^0(S', \text{NS}_X/p^n)^{G_k} = \text{NS}_X(k^s)/p^n = (\text{NS}(X_{\bar{k}})/p^n)^{G_k}.$$

For the second claim, as in the proof of Lemma 2.2, taking cohomology for the following exact sequences

$$0 \longrightarrow P^0 \longrightarrow P \longrightarrow \text{NS}_X \longrightarrow 0,$$

and

$$0 \longrightarrow P_{\text{red}}^0 \longrightarrow P^0 \longrightarrow T \longrightarrow 0,$$

we get an exact sequence

$$0 \longrightarrow P^0(k^s) \longrightarrow P(k^s) \longrightarrow \text{NS}_X(k^s) \longrightarrow H_{\text{fppf}}^1(S', P^0).$$

The last groups is of finite exponent. This shows that $\text{NS}(X_{k^s}) \longrightarrow \text{NS}(X_{\bar{k}})$ is an al. isomorphism. Thus, $\text{NS}(X_{k^s}) \otimes_{\mathbb{Z}} \mathbb{Q}_p \cong \text{NS}(X_{\bar{k}}) \otimes_{\mathbb{Z}} \mathbb{Q}_p$. By [32, §, 2.2, p10],

$$(\text{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_p) \cong (\text{NS}(X_{k^s}) \otimes_{\mathbb{Z}} \mathbb{Q}_p)^{G_k}.$$

□

Lemma 2.5. *The natural map $\text{Br}(X)[p^n] \longrightarrow \text{Br}(X_{k^s})^{G_k}[p^n]$ is al. surjective uniformly in n .*

Proof. Firstly, assume that $X(k)$ is not empty. By [9] (cf. also [32]), the natural map $\text{Br}(X) \rightarrow \text{Br}(X_{k^s})^{G_k}$ is al. surjective and there is a canonical exact sequence

$$0 \rightarrow \text{Br}(k) \rightarrow \text{Ker}(\text{Br}(X) \rightarrow \text{Br}(X_{k^s})^{G_k}) \rightarrow H^1(k, \text{Pic}_{X/k}) \rightarrow 0$$

Write $M_X =: \text{Im}(\text{Br}(X) \rightarrow \text{Br}(X_{k^s})^{G_k})$ and $N_X := \text{Ker}(\text{Br}(X) \rightarrow \text{Br}(X_{k^s})^{G_k})$. By the snake lemma,

$$0 \rightarrow N_X[p^n] \rightarrow \text{Br}(X)[p^n] \rightarrow M_X[p^n] \rightarrow N_X/p^n$$

Let $Y \subseteq X$ be a smooth projective geometrically integral curve over k obtained by taking hyperplane sections repeatedly (cf. [32]). By extending k , we assume that $Y(k)$ is not empty. Fix a k -point $P : \text{Spec}(k) \rightarrow Y$, then the first exact sequence is split. So $N_X = \text{Br}(k) \oplus H^1(k, \text{Pic}_{X/k})$. Similarly, there are two exact sequences above for Y and $N_Y = \text{Br}(k) \oplus H^1(k, \text{Pic}_{Y/k})$. By functoriality and the fact $\text{Br}(Y_{k^s}) = 0$ (so $M_Y = 0$), it suffices to prove that the natural map

$$N_X/p^n \rightarrow N_Y/p^n$$

is al. injective uniformly in n . Thus, it suffices to show that $H^1(k, \text{Pic}_{X/k})/p^n \rightarrow H^1(k, \text{Pic}_{Y/k})/p^n$ is al. injective uniformly in n . The natural map $H^1(k, \text{Pic}_{X/k}^0) \rightarrow H^1(k, \text{Pic}_{X/k})$ is al. isomorphism since $\text{NS}(X_{k^s})$ is finitely generated. Hence, it suffices to show that $H^1(k, \text{Pic}_{X/k}^0)/p^n \rightarrow H^1(k, \text{Pic}_{Y/k}^0)/p^n$ is al. injective uniformly in n . This follows from the fact that there is an abelian group D and a morphism $D \rightarrow H^1(k, \text{Pic}_{Y/k}^0)$ such that the induced map $H^1(k, \text{Pic}_{X/k}^0) \oplus D \rightarrow H^1(k, \text{Pic}_{Y/k}^0)$ is an al. isomorphism (cf. [32]). This proves the claim for X_l/l where l is some finite Galois extension of k . Let G denote the Galois group $\text{Gal}(l/k)$. Then claim for X/k follows from the fact that

$$\text{Br}(X) \rightarrow \text{Br}(X_l)^G$$

is an al. isomorphism (cf. [32] for details). \square

Lemma 2.6. *The Brauer group $\text{Br}(X_{\bar{k}})$ is an extension of a group of finite exponent by a divisible group of cofinite type. The Brauer group $\text{Br}(X_{k^s})$ is an extension of a group of cofinite type by a group of finite exponent.*

Proof. There is an exact sequence

$$0 \rightarrow \text{NS}(X_{\bar{k}}) \otimes_{\mathbb{Z}} \mathbb{Q}_p/\mathbb{Z}_p \rightarrow H_{\text{fppf}}^2(X_{\bar{k}}, \mu_{p^\infty}) \rightarrow \text{Br}(X_{\bar{k}})[p^\infty] \rightarrow 0.$$

By the proof of [18, Prop. 5.9], $H_{\text{fppf}}^2(X_{\bar{k}}, \mu_{p^\infty})$ is an extension of group of finite exponent by a group of cofinite type. Hence, the claim is true for $\text{Br}(X_{\bar{k}})[p^\infty]$. Since $\text{Br}(X_{\bar{k}})(\text{non-}p)$ is of cofinite type, this proves the first claim. The second claim follows from Corollary 2.3. \square

3 Proofs of Theorem 1.1 and its corollaries

In this section, we will prove Theorem 1.1 and all corollaries in the introduction under the assumption of Theorem 4.1 which will be proved in the last section.

Lemma 3.1. *Let $f : \mathcal{X} \rightarrow U$ be smooth projective morphism between smooth integral varieties over \mathbb{F}_p with a generic fiber X/k . Assume that X is geometrically integral over k and f has a section. Then*

$$\dim_{\mathbb{Q}_p}(H_{\text{rig}}^1(U, R^1 f_* \mathcal{O}_{\mathcal{X}}^\dagger)^{F=p}) = \dim_{\mathbb{Q}_p}(V_p \text{III}(\text{Pic}_{X/k, \text{red}}^0)) + \dim_{\mathbb{Q}_p}(\text{Pic}^0(X) \otimes_{\mathbb{Z}} \mathbb{Q}_p)$$

Proof. By the assumption, f has a section. This gives an Albanese morphism $a : X \rightarrow A$ [16, no 236, Thm. 3.3 (iii)]. By shrinking U , A/k extends to an abelian scheme $g : \mathcal{A} \rightarrow U$. By the Neron model property of g , a extends to a projective morphism $a : \mathcal{X} \rightarrow \mathcal{A}$. The induced pullback for locally constant sheaves $R^1 g_* \mu_{\ell^n} \rightarrow R^1 f_* \mu_{\ell^n}$ is an isomorphism. Thus for each $x \in U$, $a : \mathcal{X}_x \rightarrow \mathcal{A}_x$ induce an isogeny of the reduced Picard varieties. Consider the pull-back $a^* : R^1 g_* \mathcal{O}_{\mathcal{A}}^\dagger \rightarrow R^1 f_* \mathcal{O}_{\mathcal{X}}^\dagger$. This is an isomorphism. To prove this, by Kedlaya's full-faithfulness theorem [20], it suffices to show that

$$a^* : R^1 g_{\text{crys},*} \mathcal{O}_{\mathcal{A}/\mathbb{Z}_p} \otimes \mathbb{Q}_p \rightarrow R^1 f_{\text{crys},*} \mathcal{O}_{\mathcal{X}/\mathbb{Z}_p} \otimes \mathbb{Q}_p$$

is an isomorphism of F -isocrystals. This follows from [25, Prop. A.8 and Lem. A.9] since the claim holds for fibers over a closed point in U (cf. [1]).

By [27, Definition 5.16], there is a Leray spectral sequence

$$E_2^{p,q} = H_{\text{rig}}^p(U, R^q f_* \mathcal{O}_{\mathcal{X}}^\dagger) \Rightarrow H_{\text{rig}}^{p+q}(\mathcal{X}).$$

Following the pull-back trick in [29, §, 3.2], these canonical maps

$$\begin{aligned} d_2^{1,1} : E_2^{1,1} &\rightarrow E_2^{3,0} \\ d_3^{0,2} : E_3^{0,2} &\rightarrow E_3^{3,0} \\ d_2^{0,2} : E_2^{0,2} &\rightarrow E_2^{2,1} \end{aligned}$$

vanish. Thus, there are exact sequences

$$0 \rightarrow K \rightarrow H_{\text{rig}}^2(\mathcal{X}) \rightarrow H_{\text{rig}}^1(U, R^2 f_* \mathcal{O}_{\mathcal{X}}^\dagger) \rightarrow 0$$

and

$$0 \rightarrow H_{\text{rig}}^2(U) \rightarrow K \rightarrow H_{\text{rig}}^1(U, R^1 f_* \mathcal{O}_{\mathcal{X}}^\dagger) \rightarrow 0$$

Taking $F = p$ and using the pull-back trick in [29, §, 3.2] again, we get exact sequences

$$0 \rightarrow K^{F=p} \rightarrow H_{\text{rig}}^2(\mathcal{X})^{F=p} \rightarrow H_{\text{rig}}^0(U, R^2 f_* \mathcal{O}_{\mathcal{X}}^\dagger)^{F=p} \rightarrow 0$$

and

$$0 \rightarrow H_{\text{rig}}^2(U)^{F=p} \rightarrow K^{F=p} \rightarrow H_{\text{rig}}^1(U, R^1 f_* \mathcal{O}_{\mathcal{X}}^\dagger)^{F=p} \rightarrow 0$$

Since $H_{\text{rig}}^1(U, R^1 f_* \mathcal{O}_{\mathcal{X}}^\dagger)^{F=p} \cong H_{\text{rig}}^1(U, R^1 g_* \mathcal{O}_{\mathcal{A}}^\dagger)^{F=p}$ and $\text{Pic}_{X/k, \text{red}}^0 \cong \text{Pic}_{A/k}^0$, it suffices to prove the claim for \mathcal{X} being an abelian scheme. The Tate conjecture for divisor is true for abelian varieties. So $\dim_{\mathbb{Q}_p}(H_{\text{rig}}^0(U, R^2 f_* \mathcal{O}_{\mathcal{X}}^\dagger)^{F=p}) = \dim_{\mathbb{Q}_p}(\text{NS}(X)) \otimes_{\mathbb{Z}} \mathbb{Q}_p$. By [13, Prop. 2.22],

$$\dim_{\mathbb{Q}_p}(H_{\text{rig}}^2(\mathcal{X})^{F=p}) = \dim_{\mathbb{Q}_p}(\text{Pic}(\mathcal{X}) \otimes_{\mathbb{Z}} \mathbb{Q}_p) + \dim_{\mathbb{Q}_p}(V_p \text{Br}_{\text{nr}}(K(X))),$$

$$\dim_{\mathbb{Q}_p}(H_{\text{rig}}^2(U)^{F=p}) = \dim_{\mathbb{Q}_p}(\text{Pic}(U) \otimes_{\mathbb{Z}} \mathbb{Q}_p) + \dim_{\mathbb{Q}_p}(V_p \text{Br}_{\text{nr}}(k)).$$

Hence

$$\begin{aligned} \dim_{\mathbb{Q}_p}(H_{\text{rig}}^1(U, R^2 f_* \mathcal{O}_{\mathcal{X}}^\dagger)^{F=p}) - \dim_{\mathbb{Q}_p}(V_p \text{III}(\text{Pic}_{X/k}^0)) &= \dim_{\mathbb{Q}_p}(\text{Pic}(\mathcal{X}) \otimes_{\mathbb{Z}} \mathbb{Q}_p) + \dim_{\mathbb{Q}_p}(V_p \text{Br}_{\text{nr}}(K(X))) \\ &\quad - \dim_{\mathbb{Q}_p}(V_p \text{III}(\text{Pic}_{X/k}^0)) - \dim_{\mathbb{Q}_p}(\text{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_p) \\ &\quad - \dim_{\mathbb{Q}_p}(\text{Pic}(U) \otimes_{\mathbb{Z}} \mathbb{Q}_p) + \dim_{\mathbb{Q}_p}(V_p \text{Br}_{\text{nr}}(k)) \\ &= \dim_{\mathbb{Q}_p}(\text{Pic}(\mathcal{X}) \otimes_{\mathbb{Z}} \mathbb{Q}_p) - \dim_{\mathbb{Q}_p}(\text{Pic}(U) \otimes_{\mathbb{Z}} \mathbb{Q}_p) - \dim_{\mathbb{Q}_p}(\text{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_p) \\ &= \dim_{\mathbb{Q}_p}(\text{Pic}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_p) - \dim_{\mathbb{Q}_p}(\text{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_p) \\ &= \dim_{\mathbb{Q}_p}(\text{Pic}^0(X) \otimes_{\mathbb{Z}} \mathbb{Q}_p). \end{aligned}$$

The second equality follows from

$$0 \longrightarrow V_p \text{Br}_{\text{nr}}(k) \longrightarrow V_p \text{Br}_{\text{nr}}(K(X)) \longrightarrow V_p \text{III}(\text{Pic}_{X/k}^0) \longrightarrow 0,$$

which was proved in [13, Thm. (D)]. The third equality follows from the exact sequence (cf. [13, Prop. 1.3])

$$0 \longrightarrow \text{Pic}(U) \longrightarrow \text{Pic}(\mathcal{X}) \longrightarrow \text{Pic}(X) \longrightarrow 0.$$

□

Proof of Theorem 1.1

Proof. The Tate conjecture for divisors on X is equivalent to

$$\dim_{\mathbb{Q}_p}(\text{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_p) = \dim_{\mathbb{Q}_p}(H_{\text{rig}}^0(U, R^2 f_* \mathcal{O}_{\mathcal{X}}^\dagger)^{F=p}).$$

By Theorem 4.1, this is equivalent to

$$\dim_{\mathbb{Q}_p}(\text{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_p) = \dim_{\mathbb{Q}_p}(\varprojlim_n H_{\text{fppf}}^0(\text{Spec}(k), R^2 f_* \mu_{p^n}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p),$$

which is equivalent to $V_p \text{Br}(X_{k^s})^{G_k} = 0$ by Proposition 2.1. By Lemma 2.6, $\text{Br}(X_{k^s})^{G_k}[p^\infty]$ is al. isomorphic to a divisible group of cofinite type. Hence, $V_p \text{Br}(X_{k^s})^{G_k} = 0$ is equivalent to that $\text{Br}(X_{k^s})^{G_k}[p^\infty]$ is of finite exponent. By [27, Prop. 6.6] and [8], this is also equivalent to $\text{Br}(X_{k^s})^{G_k}$ is of finite exponent. This proves the first claim.

For the second claim, we will first prove it under the assumption $X(k) \neq \emptyset$. We will remove this assumption later. To prove the claim, it suffices to show that

$$V_p \text{Br}_{\text{nr}}(K(X)) \longrightarrow V_p \text{Br}(X_{k^s})^{G_k}$$

is surjective since $\text{Br}(X_{k^s})^{G_k}$ is an extension of a group of finite exponent by a group of cofinite type (cf. Lemma 2.6). Thus, $V_p \text{Br}(X_{k^s})^{G_k}$ is a finite dimensional \mathbb{Q}_p -linear space. Thus, it suffices to show

$$\dim_{\mathbb{Q}_p}(V_p \text{Br}_{\text{nr}}(K(X))) = \dim_{\mathbb{Q}_p}(\text{Ker}(V_p \text{Br}_{\text{nr}}(K(X)) \longrightarrow V_p \text{Br}(X_{k^s})^{G_k}) + \dim_{\mathbb{Q}_p}(V_p \text{Br}(X_{k^s})^{G_k}).$$

By the first part of Corollary 1.3, it suffices to show

$$\dim_{\mathbb{Q}_p}(V_p \text{Br}_{\text{nr}}(K(X))) = \dim_{\mathbb{Q}_p}(V_p \text{Br}_{\text{nr}}(k)) + \dim_{\mathbb{Q}_p}(V_p \text{III}(\text{Pic}_{X/k, \text{red}}^0)) + \dim_{\mathbb{Q}_p}(V_p \text{Br}(X_{k^s})^{G_k}).$$

By spreading out, X/k can extend to a morphism $f : \mathcal{X} \longrightarrow U$ as in Lemma 3.1. By the proof of Lemma 3.1,

$$\begin{aligned} \dim_{\mathbb{Q}_p}(H_{\text{rig}}^2(\mathcal{X})^{F=p}) &= \dim_{\mathbb{Q}_p}(H_{\text{rig}}^0(U, R^2 f_* \mathcal{O}_{\mathcal{X}}^\dagger)^{F=p}) + \dim_{\mathbb{Q}_p}(H_{\text{rig}}^2(U)^{F=p}) \\ &\quad + \dim_{\mathbb{Q}_p}(H_{\text{rig}}^1(U, R^1 f_* \mathcal{O}_{\mathcal{X}}^\dagger)^{F=p}) \end{aligned}$$

Plug the following equations into the above equation, we get the desired result.

$$\begin{aligned} \dim_{\mathbb{Q}_p}(H_{\text{rig}}^2(\mathcal{X})^{F=p}) &= \dim_{\mathbb{Q}_p}(\text{Pic}(\mathcal{X}) \otimes_{\mathbb{Z}} \mathbb{Q}_p) + \dim_{\mathbb{Q}_p}(V_p \text{Br}_{\text{nr}}(K(X))) \\ \dim_{\mathbb{Q}_p}(H_{\text{rig}}^2(U)^{F=p}) &= \dim_{\mathbb{Q}_p}(\text{Pic}(U) \otimes_{\mathbb{Z}} \mathbb{Q}_p) + \dim_{\mathbb{Q}_p}(V_p \text{Br}_{\text{nr}}(k)) \\ \dim_{\mathbb{Q}_p}(H_{\text{rig}}^0(U, R^2 f_* \mathcal{O}_{\mathcal{X}}^\dagger)^{F=p}) &= \dim_{\mathbb{Q}_p}(\text{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_p) + \dim_{\mathbb{Q}_p}(V_p \text{Br}(X_{k^s})^{G_k}) \\ \dim_{\mathbb{Q}_p}(H_{\text{rig}}^1(U, R^1 f_* \mathcal{O}_{\mathcal{X}}^\dagger)^{F=p}) &= \dim_{\mathbb{Q}_p}(\text{Pic}^0(X) \otimes_{\mathbb{Z}} \mathbb{Q}_p) + \dim_{\mathbb{Q}_p}(V_p \text{III}(\text{Pic}_{X/k, \text{red}}^0)) \end{aligned}$$

This proves the claim for the case $X(k) \neq \emptyset$. In general, there exists a finite Galois extension l/k such that $X(l) \neq \emptyset$. So $\text{Br}_{\text{nr}}(K(X_l)) \longrightarrow \text{Br}(X_{k^s})^{G_l}$ is al. surjective. Then the claim for X/k follows from the fact that $\text{Br}_{\text{nr}}(K(X)) \longrightarrow \text{Br}_{\text{nr}}(K(X_l))^{\text{Gal}(l/k)}$ is an al. isomorphism. \square

Proof of Corollary 1.4

Proof. By purity of Brauer groups [15], $\text{Br}_{\text{nr}}(K(X)) = \text{Br}(\mathcal{X})$. By Theorem 1.1,

$$\text{Br}(\mathcal{X}) \longrightarrow \text{Br}(X_{K^s})^{G_K}$$

is al. surjective. Since it factors through

$$\text{Br}(\mathcal{X}_{k^s})^{G_k} \longrightarrow \text{Br}(X_{K^s})^{G_K},$$

the claim follows. \square

4 Syntomic Cohomology and Crystalline Cohomology

Let X be a smooth projective geometrically integral variety over a field k finitely generated over \mathbb{F}_p . By spreading out X/k , we get a smooth projective morphism $f : \mathcal{X} \rightarrow U$ between smooth integral varieties over \mathbb{F}_p . Recall Pál's formulation of the Tate conjecture using rigid cohomology [27, § 6]: the natural injective map

$$\mathrm{NS}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_p \longrightarrow H_{\mathrm{rig}}^0(U, R^2 f_* \mathcal{O}_{\mathcal{X}}^\dagger)^{F=p}$$

is surjective. Pál [27, Prop. 6.6] showed that it is equivalent to the ℓ -adic Tate conjecture. The goal of this section is to prove the following theorem.

Theorem 4.1. *There is a \mathbb{Q}_p -linear isomorphism*

$$\varprojlim_n H_{\mathrm{fppf}}^0(\mathrm{Spec}(k), R^2 f_* \mu_{p^n}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p \cong H_{\mathrm{rig}}^0(U, R^2 f_* \mathcal{O}_{\mathcal{X}}^\dagger)^{F=p}.$$

This is divided into two steps, for the first we show

Proposition 4.2. *There is an al. isomorphism uniformly in n*

$$H_{\mathrm{syn}}^0(S, R^2 f_{\mathrm{syn}*} \mu_{p^n, X}) \xrightarrow{\cong} H_{\mathrm{crys}}^0(S, R^2 f_{\mathrm{crys}*} \mathcal{O}_{X/W_n})^{F=p}.$$

Then we relate crystalline cohomology of the generic fiber to rigid cohomology of \mathcal{X}

Proposition 4.3. *There is a \mathbb{Q}_p -linear isomorphism*

$$H_{\mathrm{crys}}^0(S, R^2 f_{\mathrm{crys}*} \mathcal{O}_{X/W})^{F=p}[1/p] \cong H_{\mathrm{rig}}^0(U, R^2 f_* \mathcal{O}_{\mathcal{X}}^\dagger)^{F=p}$$

4.1 Flat cohomology

For a scheme X , let $\mathrm{fppf}(X)$ (resp. $\mathrm{syn}(X)$) denote the big fppf site (resp. the small syntomic site (cf. [2, Def. 1.3])). The associated topoi are denoted by X_{fppf} (resp. X_{syn}). There is an obvious morphism of sites $v : \mathrm{fppf}(X) \rightarrow \mathrm{syn}(X)$.

Lemma 4.4. *Let G be a smooth commutative group scheme over X and $v : \mathrm{fppf}(X) \rightarrow \mathrm{syn}(X)$ be the morphism of sites. Then $R^q v_* G = 0$ for any $q > 0$.*

Proof. This follows from the the proof of [24, Chap. III, 3.9]. It is in fact the sheafication on $\mathrm{syn}(X)$ of the higher direct image from the fppf site to the category of syntomic X -scheme equipped with the étale topology. For $p > 0$, the second is 0. So it is also 0 for $p > 0$. \square

f induces a commutative diagram of topoi

$$\begin{array}{ccc} X_{\text{fppf}} & \xrightarrow{v} & X_{\text{syn}} \\ \downarrow f & & \downarrow f \\ S_{\text{fppf}} & \xrightarrow{v} & S_{\text{syn}} \end{array}$$

Consider the Kummer exact sequence $0 \rightarrow \mu_p^n \rightarrow \mathbb{G}_m \xrightarrow{p^n} \mathbb{G}_m \rightarrow 0$ on $\text{fppf}(X)$. It is still exact after taking v_* . Thus, by the Lemma above, $R^q v_* \mu_{p^n} = 0$ (also true on S) for any $q > 0$. Thus, there is a Leray spectral sequence

$$E_2^{p,q} = R^p v_* R^q f_* \mu_{p^n, X} \rightarrow R^{p+q} v_* \mu_{p^n, X}$$

Lemma 4.5. *The edge map $R^q v_* \mu_{p^n, X} \rightarrow v_* R^q f_* \mu_{p^n, X}$ is an isomorphism for $q \leq 2$.*

Proof. $f_* \mu_{p^n, X} = \mu_{p^n, S}$ Since f is geometrically connected. This shows that $E_2^{p,0} = 0$ for $p > 0$. $T := R^1 f_* \mu_{p^n, X} = \text{Pic}_{X/k}[p^n]$ is a finite flat group scheme over k . By [23, Prop. 5.1], there is an exact sequence

$$0 \rightarrow T \rightarrow G_0 \rightarrow G_1 \rightarrow 0,$$

where G_0 and G_1 are smooth group of finite type over k . $G_0 \rightarrow G_1$ is faithful flat. So it is exact on $\text{syn}(S)$ since $G_0 \rightarrow G_1$ is in fact a syntomic cover. By the Lemma above, $R^p v_* T = 0$ for $p > 0$. This proves the claim. \square

Corollary 4.6. *There is a canonical isomorphism*

$$H_{\text{fppf}}^0(S, R^2 f_* \mu_{p^n, X}) \cong H_{\text{syn}}^0(S, R^2 f_* v_* \mu_{p^n, X}).$$

In the following, we will first prove Theorem 4.1 for curves by a direct computation. Then, we prove it for abelian varieties by reducing to curves. Then, we prove the theorem for general X by reducing to curves and abelian varieties.

Lemma 4.7. *Let $f : X \rightarrow S = \text{Spec}(k)$ be smooth projective geometrically integral curve. Then*

$$H_{\text{fppf}}^0(S, R^2 f_* \mu_{p^n}) \cong H_{\text{fppf}}^0(\text{Spec}(\bar{k}), R^2 f_* \mu_{p^n}) \cong \mathbb{Z}/p^n.$$

Proof. By the proof of Proposition 2.1, there is an exact sequence

$$0 \rightarrow H_{\text{fppf}}^0(S, R^1 f_* \mathbb{G}_m/p^n) \rightarrow H_{\text{fppf}}^0(S, R^2 f_* \mu_p^n) \rightarrow H_{\text{fppf}}^0(S, R^2 f_* \mathbb{G}_m).$$

By [10, Lem. 3.2], the natural map

$$H_{\text{fppf}}^0(S, R^2 f_* \mathbb{G}_m) \rightarrow H_{\text{fppf}}^0(\text{Spec}(\bar{k}), R^2 f_* \mathbb{G}_m)$$

is injective. By the fact $H_{\text{fppf}}^0(\text{Spec}(\bar{k}), R^2 f_* \mathbb{G}_m) = \text{Br}(X_{\bar{k}}) = 0$, the second arrow in the exact sequence is an isomorphism. Since $\text{Pic}_{X/k}$ is smooth, the proof of Lemma 2.4 actually implies that

$$H_{\text{fppf}}^0(S, R^1 f_* \mathbb{G}_m/p^n) \cong ((\text{NS}(X_{\bar{k}})/p^n)^{G_k}) \cong \mathbb{Z}/p^n$$

By [10, Lem. 3.2], the natural map

$$H_{\text{fppf}}^0(S, R^2 f_* \mu_{p^n}) \rightarrow H_{\text{fppf}}^0(\text{Spec}(\bar{k}), R^2 f_* \mu_{p^n})$$

is injective. So it is an isomorphism. \square

4.2 Crystalline cohomology

Definition 4.8. 1. Let

$$(X/W_n)_{\text{CRY,SYN}} \xrightarrow{v_X} X_{\text{SYN}} \xrightarrow{p_X} X_{\text{syn}}$$

be natural morphisms of topos and $\mathcal{O}_{X,n}^{\text{crys}} := v_*(\mathcal{O}_{X/W_n})$. Since p_* is exact, we also use $\mathcal{O}_n^{\text{crys}}$ to denote $p_*\mathcal{O}_n^{\text{crys}}$ when there is no confusion.

2. Define the sheaf \mathcal{J}_n on $\text{SYN}(X)$ via the exact sequence

$$0 \longrightarrow \mathcal{J}_n \xrightarrow{j_n} \mathcal{O}_n^{\text{crys}} \longrightarrow \mathbb{G}_a \longrightarrow 0.$$

Define \mathcal{I}_n as the quotient $\mathcal{J}_{n+1}/\mathcal{J}_1$. There is an exact sequence

$$0 \longrightarrow p^n \mathcal{O}_{n+1}^{\text{crys}}/\mathcal{J}_1 \longrightarrow \mathcal{I}_n \xrightarrow{i_n} \mathcal{J}_n \longrightarrow 0.$$

Proposition 4.9 ([2, Thm. 3.4]). *There is an exact sequence of sheaves on $\text{syn}(X)$*

$$0 \longrightarrow \mu_{p^n} \longrightarrow \mathcal{I}_n \xrightarrow{1-p^{-1}F} \mathcal{O}_n^{\text{crys}} \longrightarrow 0$$

Remark 4.10. By definition, the following diagram commutes

$$\begin{array}{ccc} \mathcal{I}_n & \xrightarrow{1-p^{-1}F} & \mathcal{O}_n^{\text{crys}} \\ j_n \circ i_n \downarrow & & \downarrow p \\ \mathcal{O}_n^{\text{crys}} & \xrightarrow{p-F} & \mathcal{O}_n^{\text{crys}} \end{array}$$

such that kernel and cokernel of each vertical map is kill by p^2 . Thus the fiber of each line are al. quasi-isomorphic uniformly in n .

Take $Rf_{\text{syn}*}(-)$ of the complex in proposition 4.9 and write g_i to be the map $1-p^{-1}F$ between $R^i f_{\text{syn}*}\mathcal{J}_n$ and $R^i f_{\text{syn}*}\mathcal{O}_n^{\text{crys}}$, we get an exact sequence

$$0 \longrightarrow \text{coker}(g_1) \longrightarrow R^2 f_{\text{syn}*}\mu_{p^n} \longrightarrow \ker(g_2) \longrightarrow 0$$

which induces a long exact sequence

$$0 \rightarrow H^0(S, \text{coker}(g_1)) \rightarrow H^0(S, R^2 f_{\text{syn}*}\mu_{p^n}) \rightarrow H^0(S, \ker(g_2)) \xrightarrow{\gamma} H^1(S, \text{coker}(g_1)).$$

We need to show that $H^0(S, \text{coker}(g_1))$ is al. zero uniformly in n and the last map is al. zero uniformly in n .

4.2.1 Change of topology

Lemma 4.11. *Consider the following commutative diagram*

$$\begin{array}{ccc} X_{\text{CRY,SYN}} & \xrightarrow{f_{\text{CRY,SYN}}} & S_{\text{CRY,SYN}} \\ c_{\text{CRY,SZ}} \downarrow & & \downarrow c_{\text{CRY,SZ}} \\ X_{\text{CRY,ZAR}} & \xrightarrow{f_{\text{CRY,ZAR}}} & S_{\text{CRY,ZAR}}. \end{array}$$

Let $\mathcal{E}_{X,\text{SYN},n}^i := R^i f_{\text{CRY,SYN}*} \mathcal{O}_{X/W_n}$ and $\mathcal{E}_{X,\text{ZAR},n}^i := R^i f_{\text{CRY,ZAR}*} \mathcal{O}_{X/W_n}$ similarly, then

$$\mathcal{E}_{X,\text{SYN},n}^i \cong c_{\text{CRY,SZ}}^* \mathcal{E}_{X,\text{ZAR},n}^i.$$

Proof. We consider the presheaf \mathcal{P} on the category $\text{CRY}(S/W_n)$

$$(U, T, \delta) \rightarrow H^i((X_U/T)_{\text{CRY,SYN}}, \mathcal{O}_{X_U/T}).$$

The cohomology group is the same as $H^i((X_U/T)_{\text{CRY,ZAR}}, \mathcal{O}_{X_U/T})$ (see [5, 1.1.19]). By [5, Prop. 1.1.11], the syntomic (resp. Zariski) sheaf associated to \mathcal{P} is $\mathcal{E}_{X,\text{SYN},n}^i$ (resp. $\mathcal{E}_{X,\text{ZAR},n}^i$) and the claim follows. \square

4.2.2 Crystal property of higher direct image

Recall quasi-smoothness of morphisms between schemes is defined in [4, IV Definition 1.5.1]. Note that all smooth maps are quasi-smooth ([4, IV 1.5.5]) and for a characteristic p field k and $C(k)$ is a Cohen ring of k . Then $C_n(k) = C(k)/p^n C(k)$ is quasi-smooth over W_n because k has a p -base (see [6, Proposition 1.2.6] and [25, Remark A.4]).

Lemma 4.12. *For each $(U, T, \delta) \in \text{CRY}(S/W_n)_{\text{ZAR}}$, Zariski locally on T there exists a PD-morphism $v : T \rightarrow D_n := \text{Spec}(C_n(k))$ and*

$$(Rf_{\text{CRY}*} \mathcal{O}_{X/W_n})_T \cong Lv^* R\Gamma_{\text{crys}}(X/D_n)$$

where $R\Gamma_{\text{crys}}(X/D_n)$ is regarded as a quasi-coherent Zariski sheaf over D_n .

Proof. The sheaf $Rf_{\text{CRY}*} \mathcal{O}_{X/W_n}$ is the sheafification (in the derived ∞ -category) of $(U', T', \delta') \rightarrow R\Gamma_{\text{crys}}(X_{U'}/T')$ under the Zariski-topology. So

$$(Rf_{\text{CRY}*} \mathcal{O}_{X/W_n})_T \cong Rf_{X_U/T*} \mathcal{O}_{X_U/T}.$$

To see the existence of v , recall that the PD-envelope of $S \rightarrow D_n$ is D_n (p is the maximal ideal) so the result follows from C_n being a quasi-smooth W_n -algebra.

Assume there exists a PD-morphism $v : T \rightarrow C_n$, we apply smooth base change ([5, Corollary 7.12]) to the diagram

$$\begin{array}{ccc} X_U & \xrightarrow{g'} & X \\ f' \downarrow & & \downarrow f \\ U & \xrightarrow{g} & S \end{array}$$

and get

$$Lg_{\text{crys}}^* Rf_{\text{crys}*} \mathcal{O}_{X/W_n} \simeq Rf'_{\text{crys}*} Lg'^*_{\text{crys}} \mathcal{O}_{X/W_n}.$$

The claim follows as

$$(Lg_{\text{crys}}^* Rf_{\text{crys}*} \mathcal{O}_{X/W_n})_T \simeq Lv^*(Rf_{\text{crys}*} \mathcal{O}_{X/W_n})_{D_n} \simeq Lv^* R\Gamma_{\text{crys}}(X/D_n)$$

and

$$(Rf'_{\text{crys}*} Lg'^*_{\text{crys}} \mathcal{O}_{X/W_n})_T \simeq Rf_{X_U/T*} \mathcal{O}_{X_U/T}.$$

□

Consider the following commutative diagram of topos

$$\begin{array}{ccccc} X_{\text{CRY,SYN}} & \xrightarrow{v_X} & X_{\text{SYN}} & \xrightarrow{p_X} & X_{\text{syn}} \\ f_{\text{CRY,SYN}} \downarrow & & f_{\text{SYN}} \downarrow & & f_{\text{syn}} \downarrow \\ S_{\text{CRY,SYN}} & \xrightarrow{v_S} & S_{\text{SYN}} & \xrightarrow{p_S} & S_{\text{syn}}. \end{array}$$

By [2, Proposition 1.17], $Rp_{X*} Rv_{X,*} \mathcal{O}_{X/W_n} \cong p_{X*} v_{X,*} \mathcal{O}_{X/W_n} = \mathcal{O}_{X,n}^{\text{crys}}$. So

$$(1) \quad p_{S*} Rv_{S*} Rf_{\text{CRY,SYN}*} \mathcal{O}_{X/W_n} \simeq Rf_{\text{syn}*} \mathcal{O}_{X,n}^{\text{crys}}.$$

Proposition 4.13. *The edge map*

$$R^i f_{\text{SYN}*} \mathcal{O}_{X,n}^{\text{crys}} \rightarrow v_{S*} \mathcal{E}_{X,\text{SYN},n}^i$$

is an al. isomorphism uniformly in n .

Proof. Apply [2, Prop. 1.17] and lemma 4.11, it suffices to check there exist crystals $\mathcal{E}_{X/S,n}^i$ such that $\mathcal{E}_{X,\text{ZAR},n}^i$ is al. isomorphic to $\mathcal{E}_{X/S,n}^i$ uniformly in n . For this, we follow Morrow's method: Let C denote a Cohen ring of k , C_n denote C/p^n , $D_n := \text{Spec } C_n$ and $D := \text{Spec } C$. The same argument as in the proof of [25, Lem. A.2] provides a morphism $\psi_n^i : \mathcal{E}_{X/S,n}^i \rightarrow \mathcal{E}_{X,\text{ZAR},n}^i$ where $\mathcal{E}_{X/S,n}^i$ is the crystal constructed by Morrow in the proof of [25, Lem. A.2] whose value on affine (U, T, δ) is $g_T^* H^i(X/D_n)$ for any chosen map $g_T : T \rightarrow D_n$ lifting $U \rightarrow S$. Note that although the site considered there is the small site, [25, Lem. A.3] applies to big sites. Restricted to T , $\psi_{n,T}^i$ is given by the edge map

$$\psi_{n,T}^i : g_T^* H^i(X/D_n) \rightarrow H^i(Lg_T^* R\Gamma_{\text{crys}}(X/D_n)) \simeq (R^i f_{\text{CRY,SYN}*} \mathcal{O}_{X/W_n})_T.$$

We need to see the first map is an al. isomorphism uniform in n and g_T .

By [7, Theorem 7.24], the limit complex $R\Gamma_{\text{crys}}(X/D)$ is a perfect complex with finite Tor-amplitude and $R\Gamma_{\text{crys}}(X/D_n) \simeq R\Gamma_{\text{crys}}(X/D) \otimes_D^L D_n$. Note that D is a complete DVR thus the torsion part of $H_{\text{crys}}^i(X/D)$ is uniformly bounded and $H_{\text{crys}}^i(X/D) \otimes_D D_n$ is al. isomorphic to $H_{\text{crys}}^i(X/D_n)$ uniformly in n . So for each $i \geq 0$, the edge map

$$g_T^* H_{\text{crys}}^i(X/D_n) \rightarrow H^i(Lg_T^* R\Gamma_{\text{crys}}(X/D_n))$$

is an al. isomorphism uniformly in n and g_T . □

Corollary 4.14. *There is an al. isomorphism uniformly in n*

$$H_{\text{syn}}^0(S, R^i f_{\text{syn}*} \mathcal{O}_{X,n}^{\text{crys}}) \longrightarrow H_{\text{crys}}^0(S, R^i f_{\text{crys}*} \mathcal{O}_{X/W_n}).$$

Proof. Pushforward the edge map in proposition 4.13 along p_S and take cohomology, we get an al. isomorphism uniformly in n

$$H_{\text{syn}}^0(S, R^i f_{\text{syn}*} \mathcal{O}_{X,n}^{\text{crys}}) \longrightarrow H^0(S_{\text{CRY,SYN}}, \mathcal{E}_{X,\text{SYN},n}^i).$$

We also see in the proof that $\mathcal{E}_{X,\text{SYN},n}^i$ is al. isomorphic to $\mathcal{E}_{X/S}^i$ (coherent crystals in Zariski or syntomic topology coincide) uniformly in n . As a coherent crystal

$$H^0(S_{\text{CRY,SYN}}, \mathcal{E}_{X/S}^i) \cong H^0(S_{\text{CRY,ZAR}}, \mathcal{E}_{X/S}^i) \cong H_{\text{crys}}^0(S, \mathcal{E}_{X/S}^i).$$

Pushforward the al. isomorphism $\mathcal{E}_{X/S}^i \rightarrow \mathcal{E}_{X,\text{ZAR},n}^i$ to small crystalline site, we get an al. isomorphism $H_{\text{crys}}^0(S, \mathcal{E}_{X/S}^i) \rightarrow H_{\text{crys}}^0(S, R^i f_{\text{crys}*} \mathcal{O}_{X/W_n})$ uniformly in n . \square

Lemma 4.15. *Let $i_n : (S/W_n)_{\text{crys}} \rightarrow (S/W)_{\text{crys}}$ be the canonical morphism of topos and $\mathcal{E}_{X/S}^i$ be the crystal constructed by Morrow in the proof of [25, Lem. A.2] (recalled in proof of proposition 4.13). Then*

$$i_n^* \mathcal{E}_{X/S}^2 \longrightarrow R^2 f_{\text{crys}*} \mathcal{O}_{X/W_n}$$

is an al. isomorphism uniformly in n .

Proof. The claim follows from Morrow's proof of [25, Lemma A.2] and that

$$H_{\text{crys}}^i(X/D)/p^n \longrightarrow H_{\text{crys}}^i(X/D_n)$$

is an al. isomorphism uniformly in n for any $i \geq 0$ (cf. [7, Cor. 7.25]). \square

4.3 Relation with Albanese

Let X be a smooth projective geometrically connected variety over field k , there is an Albanese variety A associated to X which is the dual abelian variety of the reduced part of the Picard scheme $\text{Pic}_{X/k,\text{red}}^0$ (cf. [16, no 236, Thm. 3.3 (iii)]). The motivation is the following result [1]

Theorem 4.16. *Let X be a smooth and projective over a perfect field k of characteristic $p > 0$. Let $T_{\text{crys}}(-)$ denote the covariant Dieudonne module. Then, there is a canonical isomorphism*

$$T_{\text{crys}}(\text{Pic}_{X/k,\text{red}}^0) \cong H_{\text{crys}}^1(X/W(k)).$$

Assume there is a rational point $a \in X$. Let $a : X \rightarrow A$ be the Albanese morphism induced by a . By definition, a induced an isomorphism between the reduced Picard varieties [16, no 236, Thm. 3.3 (iii)]. So we have

$$\begin{array}{ccc} X & \xrightarrow{a} & A \\ & \searrow f & \swarrow g \\ & S & \end{array}$$

and a^* induces a map $\mathcal{E}_{A,\text{SYN},n}^1 \rightarrow \mathcal{E}_{X,\text{SYN},n}^1$ which is the sheafification of the corresponding map $a^* : \mathcal{E}_{A,\text{ZAR},n}^1 \rightarrow \mathcal{E}_{X,\text{ZAR},n}^1$ (Lemma 4.11). By crystalline Dieudonne theory, we know $\mathcal{E}_{A,\text{ZAR},n}^1$ is a coherent crystal (cf [5, Sec. 2.5]).

Proposition 4.17. *The morphism a^* is an al. isomorphism uniformly in n .*

Proof. Choose a Cohen ring $C = C(k)$, let $C_n := C_n(k) := C(k)/p^n$ and $D_n := \text{Spec}(C_n)$. It suffices to check the statement locally in $\text{CRYS}(S)_{\text{ZAR}}$. Let $(U, T, \delta) \in \text{CRYS}(S)_{\text{ZAR}}$ and apply lemma 4.12 we may assume there exists a PD-morphism $v : T \rightarrow D_n$. Via the proof of proposition 4.13, for each $p \geq 0$, the edge map

$$v^* H_{\text{crys}}^p(X/D_n) \rightarrow H^p(Lv^* R\Gamma_{\text{crys}}(X/D_n))$$

is an al. isomorphism uniformly in n and v . Combine it with Lemma 4.12, we get an al. isomorphism uniformly bounded in n and v

$$v^* H_{\text{crys}}^p(X/D_n) \longrightarrow \mathcal{E}_{X,\text{ZAR},n}^p|_T.$$

Similarly, it works for the Albanese variety A . Take $p = 1$, it is enough to see

$$H_{\text{crys}}^1(X/D) \rightarrow H_{\text{crys}}^1(A/D)$$

induced by a is an al. isomorphism uniformly bounded in n . To show the statement it is enough to show it after base change to $\overline{W} = W(\overline{k})$ which is faithfully flat C -algebra. Let $\overline{D} := \text{Spec } \overline{W}$, via smooth base change [7, Coro 7.12] we get

$$H_{\text{crys}}^1(X/D) \otimes_D \overline{W} \cong H_{\text{crys}}^1(X_{\overline{k}}/\overline{D}).$$

Now the result follows from Theorem 4.16 and the duality between Albanese and Picard scheme. \square

5 Proof of Proposition 4.2

5.1 Case of a curve

Assume $\dim X = 1$. Base change to $\overline{S} = \text{Spec } \overline{k}$ we have the following diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^0(S, \text{coker}(g_1)) & \longrightarrow & H^0(S, R^2 f_{\text{syn}*} \mu_{p^n}) & \xrightarrow{\beta_4} & H^0(S, \ker(g_2)) \\ & & \beta_1 \downarrow & & \beta_2 \downarrow & & \beta_5 \downarrow \\ 0 & \longrightarrow & H^0(\overline{S}, \text{coker}(g_1)_{\overline{S}}) & \longrightarrow & H^0(\overline{S}, R^2 f_{\text{syn}*} \mu_{p^n, X_{\overline{k}}}) & \xrightarrow{\beta_3} & H^0(\overline{S}, \ker(g_2)_{\overline{S}}) \end{array}$$

In the following, we show that β_5 is al. injective uniformly in n (cf. Corollary 5.4) and β_3 is an al. isomorphism uniformly in n (cf. Lemma 5.5). By Lemma 4.7, β_2 is already

an isomorphism. This shows that β_3 is an al. isomorphism uniformly in n . The digram chasing shows that $H^0(S, \text{coker}(g_1))$ and the image of

$$H^0(S, \text{coker}(g_2)) \longrightarrow H^1(S, \text{coker}(g_1))$$

can be killed by a positive integer independent of n .

Lemma 5.1. *We have $R^i f_{\text{CRYST}}^* \mathcal{O}_{X/W_n}$ ($i = 0, 1, 2$) is a coherent crystal over $\text{SYN}(S)$.*

Proof. Through the proof of proposition 4.13, the obstruction preventing $R^i f_{\text{CRYST}}^* \mathcal{O}_{X/W_n}$ from being a crystal lies in torsions in cohomology groups $H_{\text{crys}}^i(X/D_n)$. So it suffices to see $H_{\text{crys}}^i(X/D_n)$ ($i = 0, 1, 2$) are flat D_n -modules. Since \overline{W}_n is a faithfully flat C_n -algebra, apply smooth base change, it suffices to see it when k is algebraically closed. Now the claim for H^0 is clear and Poincare duality gives H^2 . For H^1 , one can apply Theorem 4.16. \square

Remark 5.2. Any smooth projective geometrically connected curve over a separable closed field k of characteristic $p > 0$ admits a lifting to $C(k)$. This follows from theory of moduli of curves.

So the edge map $R^2 f_{\text{syn}*} \mathcal{O}_n^{\text{crys}} \rightarrow v_* R^2 f_{\text{crys}*} \mathcal{O}_{X/W_n}$ is an isomorphism.

Lemma 5.3. *The natural map*

$$H^0(S, R^2 f_{\text{crys}*} \mathcal{O}_{X/W_n}) \rightarrow H^0(\overline{S}, R^2 f_{\text{crys}*} \mathcal{O}_{\overline{X}/W_n})$$

is injective.

Proof. By definition, $H^0(S, R^2 f_{\text{crys}*} \mathcal{O}_{X/W_n}) = \text{Hom}(\mathcal{O}_{S/W}, R^2 f_{\text{crys}*} \mathcal{O}_{X/W_n})$. By Lemma 5.1, $R^2 f_{\text{crys}*} \mathcal{O}_{X/W_n}$ is a coherent crystal. Since $C(k) \rightarrow W(\overline{k})$ is faithfully flat, the pull-back functor from the category of crystal on S to the one of \overline{S} is faithful. By the base change theorem of crystalline cohomology, $R^2 f_{\text{crys}*} \mathcal{O}_{\overline{X}/W_n}$ is naturally isomorphic to the pull-back of $R^2 f_{\text{crys}*} \mathcal{O}_{X/W_n}$. This proves the claim. \square

Corollary 5.4. *The natural map*

$$H_{\text{syn}}^0(S, R^2 f_{\text{syn}*} \mathcal{O}_{X,n}^{\text{crys}}) \longrightarrow H_{\text{syn}}^0(\overline{S}, R^2 f_{\text{syn}*} \mathcal{O}_{\overline{X},n}^{\text{crys}})$$

is injective. As a result, the natural map

$$H_{\text{syn}}^0(S, R^2 f_{\text{syn}*} \mathcal{I}_n)^{1=p^{-1}F} \longrightarrow H_{\text{syn}}^0(\overline{S}, R^2 f_{\text{syn}*} \mathcal{I}_n)^{1=p^{-1}F}$$

is al. injective uniformly in n .

Proof. By the proof of Proposition 4.13, the natural map

$$H_{\text{syn}}^0(S, R^2 f_{\text{syn}*} \mathcal{O}_{X,n}^{\text{crys}}) \longrightarrow H^0(S, R^2 f_{\text{crys}*} \mathcal{O}_{X/W_n})$$

is actually an isomorphism since $R^i f_{\text{CRYs}*} \mathcal{O}_{X/W_n}$ ($i = 0, 1$) is a coherent crystal over $\text{SYN}(S)$. Then, the first claim follows from Lemma 5.3. The second claim follows from the fact that the natural map

$$\mathcal{I}_n \longrightarrow \mathcal{O}_n^{\text{crys}}$$

has a kernel and cokernel killed by p . □

Lemma 5.5. *The natural map*

$$H^0(\overline{S}, R^2 f_{\text{syn}*} \mu_{p^n, X_{\overline{K}}}) \longrightarrow H_{\text{syn}}^0(\overline{S}, R^2 f_{\text{syn}*} \mathcal{I}_n)^{1=p^{-1}F}$$

is surjective and al, injective uniformly in n .

Proof. Since any syntomic cover of \overline{S} has a section, so $H_{\text{syn}}^i(\overline{S}, \mathcal{F}) = 0$ for any sheaf \mathcal{F} on $\text{syn}(\overline{S})$ and $i > 0$. By the Leray spectral sequence, for any sheaf \mathcal{G} on $\text{syn}(\overline{X})$, there is a canonical isomorphism

$$H_{\text{syn}}^i(\overline{X}, \mathcal{G}) \cong H_{\text{syn}}^0(\overline{S}, R^i f_{\text{syn}*} \mathcal{G}).$$

Take cohomology on \overline{X} for the exact sequence in Proposition 4.9. This shows the surjectivity in the claim. The natural map

$$H_{\text{syn}}^0(\overline{S}, R^2 f_{\text{syn}*} \mathcal{I}_n)^{1=p^{-1}F} \longrightarrow H^0(\overline{S}, R^2 f_{\text{crys}*} \mathcal{O}_{\overline{X}/W_n})^{F=p} = H_{\text{crys}}^2(\overline{X}/W_n)^{F=p}$$

is an al. isomorphism. Since $\text{Spec}(\overline{W})$ is a final object in $\text{CRYs}(\overline{S}/W)$ [19, II. 2.2], $H_{\text{crys}}^2(\overline{X}/W_n) = H_{\text{crys}}^2(\overline{X}/\overline{W}_n)$. By the trace map of crystalline cohomology,

$$H_{\text{crys}}^2(\overline{X}/\overline{W}_n)^{F=p} = (\overline{W}_n)^{F=id} = \mathbb{Z}/p^n.$$

The surjectivity implies that the map is al. injective uniformly in n . □

5.2 Albanese variety

Assume that X has a k -rational point a . a induces an Albanese map $a : X \rightarrow A := \text{Alb}(X)$ be the map. The prove in the last section implies a^* induces an al. isomorphism uniformly in n $R^1 f_{\text{syn}*} \mathcal{O}_{X,n}^{\text{crys}} \cong R^1 f_{\text{syn}*} \mathcal{O}_{A,n}^{\text{crys}}$ and the result holds for \mathcal{I}_n as \mathcal{I}_n is al. isomorphic to $\mathcal{O}_n^{\text{crys}}$ uniformly in n . All our construction are functorial, so $\text{coker}(g_{1,A}) \cong \text{coker}(g_{1,X})$ is al. uniformly in n .

5.3 General X

For general X , we use Grothendieck's trick: We choose a series of hyperplane sections to get $Y \subset X$ such that Y is a smooth projective geometrically connected curve. To see the al. vanishing of γ_X uniformly in n , we have the following diagram

$$\begin{array}{ccc} H^0(S, \ker(g_{2,X})) & \xrightarrow{\gamma_X} & H^1(S, \operatorname{coker}(g_{1,X})) \\ \beta_1 \downarrow & & \downarrow \beta_2 \\ H^0(S, \ker(g_{2,Y})) & \xrightarrow{\gamma_Y} & H^1(S, \operatorname{coker}(g_{1,Y})). \end{array}$$

We already know γ_Y vanishes so it is enough to see β_2 is injective. For simplicity, we assume that Y has a rational k -point. The universal property of Albanese variety induces a diagram

$$\begin{array}{ccc} Y & \longrightarrow & X \\ \downarrow & & \downarrow \\ A_Y & \longrightarrow & A_X. \end{array}$$

The natural map $A_Y \rightarrow A_X$ is surjective. This is equivalent to show that the natural map between the associated Picard varieties has a finite kernel. Since Y is obtained by taking hyperplanes section,

$$H_{et}^1(X_{k^s}, \mathbb{Q}_\ell(1)) \longrightarrow H_{et}^1(Y_{k^s}, \mathbb{Q}_\ell(1))$$

is injective. So the map between the Tate-modules of the associated Picard varieties is injective. By the Poincaré reducibility theorem, $A_Y \cong A_X \times B$ for another abelian variety B up to an isogeny. Roughly speaking, $\operatorname{coker}(g_{1,A_X})$ is almost a direct summand of the sheaf $\operatorname{coker}(g_{1,A_Y})$ uniformly in n . So the injectivity follows from the comparison between X and A_X . This proves that γ_X is al. 0 uniformly in n . The almost vanishing of $H^0(S, \operatorname{coker}(g_1))$ follows from a similar argument.

Now we need to remove the assumption of existence of a rational k -point. By taking a finite Galois extension l of k , we can assume that the claim is true for X_l/l . Then the claim for X follows from the Galois étale descent. This proves that the natural map

$$H^0(S, R^2 f_{\operatorname{syn}*} \mu_{p^n}) \rightarrow H^0(S, R^2 f_{\operatorname{syn}*} \mathcal{I}_n)^{1=p^{-1}F}$$

is an al. isomorphism uniformly in n . Composites with the natural map

$$H^0(S, R^2 f_{\operatorname{syn}*} \mathcal{I}_n)^{1=p^{-1}F} \longrightarrow H_{\operatorname{crys}}^0(S, R^2 f_{\operatorname{crys}*} \mathcal{O}_{X/W_n})^{F=p}.$$

We get an al. isomorphism uniformly in n

$$H^0(S, R^2 f_{\operatorname{syn}*} \mu_{p^n}) \rightarrow H_{\operatorname{crys}}^0(S, R^2 f_{\operatorname{crys}*} \mathcal{O}_{X/W_n})^{F=p}$$

By the Lemma below and invert p , we get an isomorphism

$$\left(\varprojlim_n H_{\operatorname{syn}}^0(S, R^2 f_{\operatorname{syn}*} \mu_{p^n}) \right) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p \xrightarrow{\cong} H_{\operatorname{crys}}^0(S, R^2 f_{\operatorname{crys}*} \mathcal{O}_{X/W})^{F=p} [1/p].$$

This completes the proof of Proposition 4.2.

Lemma 5.6. *The natural morphism*

$$H_{\text{crys}}^0(S, R^2 f_{\text{crys}*} \mathcal{O}_{X/W}) \longrightarrow \varprojlim_n H_{\text{crys}}^0(S, R^2 f_{\text{crys}*} \mathcal{O}_{X/W_n})$$

is an al. isomorphism.

Proof. By Lemma 4.15, it suffices to prove that

$$H_{\text{crys}}^0(S, \mathcal{E}_{X/S}) \longrightarrow \varprojlim_n H_{\text{crys}}^0(S, i_n^* \mathcal{E}_{X/S})$$

is an isomorphism. This follows from [7, 7.22.2] since $\mathcal{E}_{X/S}$ is a coherent crystal on S/W . \square

6 Proof of Proposition 4.3

6.1 F-crystals

We still need to see

$$H_{\text{crys}}^0(S, R^2 f_{\text{crys}*} \mathcal{O}_{X/W})^{F=p}[1/p] \cong H_{\text{rig}}^0(U, R^2 f_* \mathcal{O}_X^\dagger)$$

Let S be a Noetherian scheme of characteristic p . An F -crystal over S is an $\mathcal{O}_{S/\mathbb{Z}_p}$ -coherent crystal \mathcal{E} with a morphism $F : \mathcal{E}^{(\sigma)} \rightarrow \mathcal{E}$ such that the kernel and cokernel of F are annihilated by a power of p , where $\mathcal{O}_{S/\mathbb{Z}_p}$ is the structure sheaf of the big crystalline site S over \mathbb{Z}_p . The Morphisms are morphisms of coherent crystals compatible with F . The category of F -isocrystals is the isogeny category of F -crystals. We denote the category of F -crystals via $\text{F-Crys}(S/W)$ and F -isocrystals via $\text{F-Isoc}(S/W)$. If S is of finite type, by [3, Theorem 2.4.2], our category of F -isocrystals can be naturally seen as a full subcategory of convergent F -isocrystals and if S is smooth over \mathbb{F}_p , it is an equivalence of categories up to Tate twists.

Lemma 6.1. *If S is the spectrum of a field of characteristic $p > 0$ or an affine smooth variety over a perfect field k of characteristic $p > 0$. Then $\text{F-Crys}(S/W)$ is an abelian category and the functor from $\text{F-Crys}(S/W)$ to the category of coherent crystals on (S/W) is exact. Moreover, if S is an affine smooth variety over a perfect field k , the functor from $\text{F-Crys}(S/W)$ to the category of convergent F -isocrystals is exact.*

Proof. By [4, Prop. IV.1.7.6], the category of coherent crystals is an abelian category. It suffices to show that the kernel and cokernel inherit a non-degenerated F -structure. Let F_S denote the absolute Frobenius on S . The cokernel is clear since F_S^* is a right exact functor. Let $0 \rightarrow \mathcal{F}_1 \rightarrow \mathcal{F}_2 \rightarrow \mathcal{F}_3 \rightarrow 0$ be an exact sequence in the category of crystals. Assume that the third arrow is an morphism of F -crystals. The pull-back gives an exact sequence of crystals

$$F_S^* \mathcal{F}_1 \rightarrow F_S^* \mathcal{F}_2 \rightarrow F_S^* \mathcal{F}_3 \rightarrow 0$$

Next, we show that the surjective morphism

$$F_S^* \mathcal{F}_1 \rightarrow \ker(F_S^* \mathcal{F}_2 \rightarrow F_S^* \mathcal{F}_3)$$

is al. injective. Write $S = \text{Spec}(A)$. Let \mathcal{A} be an p -adic complete noetherian algebra such that $A = \mathcal{A}/p$ (cf. [25, Lem. A.3 and Cor. A.5] for details). Let M_i ($i = 1, 2, 3$) denote the corresponding \mathcal{A} -modules of \mathcal{F}_i . By [25, Cor. A.5], each $M_i[1/p]$ is a finite projective \mathcal{A} -module. So the exact sequence

$$0 \rightarrow M_1[1/p] \rightarrow M_2[1/p] \rightarrow M_3[1/p] \rightarrow 0$$

split as $A[1/p]$ -modules. Let $\sigma : \mathcal{A} \rightarrow \mathcal{A}$ be a lifting of the absolute Frobenius F_S . So the sequence

$$0 \rightarrow (M_1 \otimes_{\mathcal{A}, \sigma} \mathcal{A})[1/p] \rightarrow (M_2 \otimes_{\mathcal{A}, \sigma} \mathcal{A})[1/p] \rightarrow (M_3 \otimes_{\mathcal{A}, \sigma} \mathcal{A})[1/p] \rightarrow 0$$

is exact. So

$$F_S^* \mathcal{F}_1 \rightarrow \ker(F_S^* \mathcal{F}_2 \rightarrow F_S^* \mathcal{F}_3)$$

is an isomorphism after tensor \mathbb{Q}_p . Its kernel is killed by a power of p since the \mathcal{A} -module of the kernel is finitely generated. Then, by digram chasing, the induced morphism

$$F_S^* \mathcal{F}_1 \rightarrow \mathcal{F}_1$$

is an isogeny. It is easy to check by diagram chasings that \mathcal{F}_1 is the $\ker(F_S^* \mathcal{F}_2 \rightarrow F_S^* \mathcal{F}_3)$ in $\text{F-Crys}(S/W)$. This proves the first claim. For the second claim, by definition [21, Def. 2.1], the category of convergent F -isocrystals is equivalent to the category of coherent \mathcal{D} -module over $\mathcal{A}[1/p]$ with a semi-linear σ -action. The claim follows from the fact that an exact sequence of \mathcal{A} -modules is an exact sequence of $\mathcal{A}[1/p]$ -modules after tensor \mathbb{Q}_p . \square

Lemma 6.2. *Let $f : \mathcal{X} \rightarrow U$ be a smooth projective morphism with geometrically connected fibers. Assume that U is a smooth affine integral variety over \mathbb{F}_p . Then, there is an open dense subset $V \subseteq U$, a F -crystal $\mathcal{F} \in \text{F-Crys}(V/W)$ and a F -morphism $\mathcal{F} \rightarrow R^2 f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W}|_V$ whose kernel and cokernel are killed by a power of p . Moreover, the induced morphism*

$$H_{\text{crys}}^0(V, \mathcal{F})^{F=p}[p^{-1}] \cong H_{\text{crys}}^0(U, R^2 f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W})^{F=p}[p^{-1}]$$

is an isomorphism.

Proof. By [25, Lem. A.2], there is a coherent crystal $\mathcal{E}_{\mathcal{X}/U}$ and a morphism of $\mathcal{O}_{U/W}$ -module $\alpha : \mathcal{E}_{\mathcal{X}/U} \rightarrow R^2 f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W}$ whose kernel and cokernel are killed by a power of p . So, there is a morphism of $\mathcal{O}_{U/W}$ -module

$$\beta : R^2 f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W} \rightarrow \mathcal{E}_{\mathcal{X}/U}$$

such that $\alpha\beta = \beta\alpha = p^n$. Let $a : F_U^* R^2 f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W} \rightarrow R^2 f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W}$ be the morphism defined in [25, Prop. A.7 (2)]. Define $b := \beta a F_U^*(\alpha)$.

$$\alpha b = \alpha \beta a F_U^*(\alpha) = p^n a F_U^*(\alpha) = p^n a F_U^*(\alpha)$$

Let $R^2 f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W}(-n)$ denote $p^n a : F_U^* R^2 f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W} \rightarrow R^2 f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W}$. (\mathcal{E}, b) is a F -crystal isogenous to $R^2 f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W}(-n)$. By [25, Prop. A.8], all slopes of (\mathcal{E}, b) are $\geq n$. View $\mathcal{E}(n)$ as an object in the category of convergent F -isocrystals. By [22, Lem. 5.8], there is an open dense open subset V of U such that $\mathcal{E}(n)|_V$ is isomorphic to a F -crystal (\mathcal{F}, c) in the category of convergent F -isocrystals on V . So $\mathcal{E}|_V$ is isomorphic to $\mathcal{F}(-n)$ in the category of convergent F -isocrystals on V . Hence, there is an F -isogeny $\gamma : \mathcal{F}(-n) \rightarrow \mathcal{E}|_V$ in $\text{F-Crys}(V/W)$ (It is also an isogeny in the category of $\mathcal{O}_{V/W}$ -modules). By definition, $\mathcal{F}(-n) = (\mathcal{F}, p^n c)$ and $\gamma : \mathcal{F} \rightarrow \mathcal{E}|_V$ is a morphism of crystals on V . The composition $\eta := \alpha_V \gamma$ is a F -morphism $\mathcal{F}(-n) \rightarrow R^2 f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W}(-n)|_V$, where $\alpha_V := \alpha|_V$. Thus, $(p^n a|_V) F_V^*(\eta) = \eta(p^n c)$. This is same as $a|_V F_V^*(p^n \eta) = (p^n \eta) c$. Thus, $p^n \eta$ is a F -morphism $\mathcal{F} \rightarrow R^2 f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W}|_V$ and is also an isogeny in the category of $\mathcal{O}_{V/W}$ -modules since both of α_V and γ are isogenies in the category of $\mathcal{O}_{V/W}$ -modules. This proves the first claim.

Next, we prove the second claim. By definition,

$$\begin{aligned} H_{\text{crys}}^0(U, R^2 f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W})^{F=p}[p^{-1}] &= H_{\text{crys}}^0(U, R^2 f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W}(-n))^{F=p^{n+1}}[p^{-1}] \\ &= H_{\text{crys}}^0(U, \mathcal{E})^{F=p^{n+1}}[p^{-1}] = \text{Hom}_{\text{F-Isoc}}(\mathcal{O}_{U/W}(-(n+1)), \mathcal{E}). \end{aligned}$$

By [21, Thm. 5.3],

$$\begin{aligned} \text{Hom}_{\text{F-Isoc}}(\mathcal{O}_{U/W}(-(n+1)), \mathcal{E}) &\cong \text{Hom}_{\text{F-Isoc}}(\mathcal{O}_{V/W}(-(n+1)), \mathcal{E}|_V) \\ &\cong \text{Hom}_{\text{F-Isoc}}(\mathcal{O}_{V/W}(-(n+1)), \mathcal{F}(-n)) \cong \text{Hom}_{\text{F-Isoc}}(\mathcal{O}_{V/W}(-1), \mathcal{F}) \\ &\cong H_{\text{crys}}^0(V, \mathcal{F})^{F=p}[p^{-1}]. \end{aligned}$$

□

Proposition 6.3. *Let $f : \mathcal{X} \rightarrow U$ as in the previous lemma. Write $g : S = \text{Spec}(K) \rightarrow U$ for the generic point of U and X for the generic fiber. Then there is a canonical isomorphism induced by g*

$$H_{\text{crys}}^0(U, R^2 f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W})^{F=p}[p^{-1}] \cong H_{\text{crys}}^0(S, R^2 f_{\text{crys}*} \mathcal{O}_{X/W})^{F=p}[1/p].$$

Proof. By Lemma 6.4 below, the F -morphism

$$g_{\text{crys}}^* R^n f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W} \longrightarrow R^n f_{\text{crys}*} \mathcal{O}_{X/W}$$

has a kernel and a cokernel killed by a p -power. Let \mathcal{F} as in the previous lemma, $\mathcal{F}|_S \rightarrow g_{\text{crys}}^* R^n f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W}$ is an F -morphism with a kernel and a cokernel killed by a p -power. This gives

$$H_{\text{crys}}^0(S, \mathcal{F}|_S)^{F=p}[p^{-1}] \cong H_{\text{crys}}^0(S, R^2 f_{\text{crys}*} \mathcal{O}_{X/W})^{F=p}[1/p].$$

By the second claim of the previous lemma, it remains to prove that the pull-back induces a canonical isomorphism

$$H_{\text{crys}}^0(V, \mathcal{F}|_S)^{F=p}[p^{-1}] \cong H_{\text{crys}}^0(S, \mathcal{F})^{F=p}[p^{-1}].$$

This is proved in Lemma 6.5 below. \square

Lemma 6.4. *Consider the following Cartesian diagram*

$$\begin{array}{ccc} X & \xrightarrow{g'} & \mathcal{X} \\ f' \downarrow & & \downarrow f \\ S & \xrightarrow{g} & U \end{array}$$

There exists a natural F -morphism

$$g_{\text{crys}}^* R^n f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W} \longrightarrow R^n f'_{\text{crys}*} \mathcal{O}_{X/W}$$

which is an isomorphism up to a finite p -exponent.

Proof. By smooth base change, we have a map

$$g_{\text{crys}}^* R^n f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W} \rightarrow H^n(Lg_{\text{crys}}^* Rf_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W}) \xrightarrow{\cong} H^n(Rf'_{\text{crys}*} \mathcal{O}_{X/W}) = R^n f'_{\text{crys}*} \mathcal{O}_{X/W}.$$

This map is F -equivariant because both F -structures are induced from the absolute and relative Frobenius whose formation are functorial (see [25, Prop. A.7]). To see the first map is an isomorphism up to a finite p -exponent, take a p -adically complete, noetherian flat lifting R of U to W . Since $R/p^n R$ is quasi-smooth over W_n for all n , it suffices to check on each affine $(V, T = \text{Spec } B, \delta) \in \text{Crys}(S/W)$ such that there exists a map $v_T : R \rightarrow B$ lifting $V \rightarrow S$ (after a easy covering argument). We need to see for $i \geq 1$, $L^i v_T^* H_{\text{crys}}^{n+i}(\mathcal{X}/R)$ is killed by a power of p independent of v_T . This follows from [25, Coro. A.5] that $H_{\text{crys}}^{n+i}(\mathcal{X}/R)$ is a finitely generated R -module which becomes finite projective over $R[1/p]$ after inverting p (cf. [25, Proof of Lem. A.2]). \square

Lemma 6.5. *Notations as in Lemma 6.2,*

$$H_{\text{crys}}^0(V, \mathcal{F})^{F=p}[p^{-1}] \cong H_{\text{crys}}^0(S, \mathcal{F}|_S)^{F=p}[p^{-1}].$$

Proof. By shrinking V , we can assume that V is affine and the slope polygon of \mathcal{F} is constant on V (cf. [21, Thm. 3.12]). By [21, Corollary 4.2], there exists a unique filtration in the category of convergent F -isocrystals

$$0 = \mathcal{E}_0 \subset \mathcal{E}_1 \subset \cdots \mathcal{E}_l = \mathcal{F}$$

such that each successive quotient $\mathcal{E}_i/\mathcal{E}_{i-1}$ is isoclinic of slope $s_i \in \mathbb{Q}_{\geq 0}$ and $s_1 < s_2 < \cdots < s_l$. Since all slope of \mathcal{F} are nonnegative. There is a \mathcal{E}_i such that all of slopes of \mathcal{E}_i

are contained in $[0, 1]$ and all of slopes of $\mathcal{F}/\mathcal{E}_i$ are > 1 . Denote it by $\mathcal{F}_{[0,1]}$. By shrinking V further, we can assume that $\mathcal{F}_{[0,1]}$ comes from a F -crystal (cf. [22, Lem. 5.8]). There is a morphism $\mathcal{F}_{[0,1]} \rightarrow \mathcal{F}$ in $\text{F-Crys}(V/W)$ which becomes injective in $\text{F-Isoc}(V/W)$. By Lemma 6.1 replacing $\mathcal{F}_{[0,1]}$ by its image in \mathcal{F} , we can assume that $\mathcal{F}_{[0,1]} \rightarrow \mathcal{F}$ in $\text{F-Crys}(V/W)$ is injective. Let $\mathcal{F}_{>1}$ denote the cokernel which has all slope greater than 1 since restricting to a closed point is exact (cf. [25, Lem. A.9]). Let \mathcal{A} be a smooth lifting of V to W and \mathcal{B} denote the local ring of the generic point of the special fiber of $\text{Spec}(\mathcal{A})$. It's easy to see that \mathcal{B} is a DVR with the maximal ideal generated by p . Since $\mathcal{A} \rightarrow \mathcal{B}$ is flat, it is still exact after taking completion at p . By abusing notations, we still denote them by \mathcal{A} and \mathcal{B} respectively. Hence, the pull-back functor $\text{F-Crys}(V/W) \rightarrow \text{F-Crys}(S/W)$ is exact. Thus, there is an exact sequence in $\text{F-Crys}(S/W)$

$$0 \rightarrow \mathcal{F}_{[0,1]}|_S \rightarrow \mathcal{F}|_S \rightarrow \mathcal{F}_{>1}|_S \rightarrow 0.$$

This gives

$$0 \rightarrow \text{Hom}_{\text{F-Crys}}(\mathcal{O}_{S/W}(-1), \mathcal{F}_{[0,1]}|_S) \rightarrow \text{Hom}_{\text{F-Crys}}(\mathcal{O}_{S/W}(-1), \mathcal{F}|_S) \rightarrow \text{Hom}_{\text{F-Crys}}(\mathcal{O}_{S/W}(-1), \mathcal{F}_{>1}|_S)$$

The last group is of finite exponent, since the pull-back functor from $\text{F-Crys}(S/W)$ to $\text{F-Crys}(\text{Spec}(\bar{K})/W)$ is faithful and all slopes of $\mathcal{F}_{>1}|_S$ are greater than 1. Thus,

$$\text{Hom}_{\text{F-Crys}}(\mathcal{O}_{S/W}(-1), \mathcal{F}_{[0,1]}|_S)[p^{-1}] \cong \text{Hom}_{\text{F-Crys}}(\mathcal{O}_{S/W}(-1), \mathcal{F}|_S)[p^{-1}]$$

Similarly ([25, Lem. A.9]),

$$\text{Hom}_{\text{F-Crys}}(\mathcal{O}_{V/W}(-1), \mathcal{F}_{[0,1]})[p^{-1}] \cong \text{Hom}_{\text{F-Crys}}(\mathcal{O}_{V/W}(-1), \mathcal{F})[p^{-1}].$$

Thus, we may assume that $\mathcal{F}_{[0,1]} = \mathcal{F}$. By shrinking V (cf. [22, Lem. 5.8]), we may assume that \mathcal{F} is F -isogenous to a Dieudonné crystal. In fact, we can assume that \mathcal{F} is a Dieudonné crystal. Since the functor from the category of Dieudonné crystals to F-Crys on S or V is fully faithful (cf. [22, Rmk. 5.2]), it suffices to show

$$\text{Hom}_{DC(V)}(\mathcal{O}_{V/W}(-1), \mathcal{F})[p^{-1}] \cong \text{Hom}_{DC(S)}(\mathcal{O}_{S/W}(-1), \mathcal{F}|_S)[p^{-1}].$$

This follows from the equivalences between Dieudonné crystals and p -divisible groups on S and on V and the fact that the restriction functor for p -divisible groups from V to S is fully faithful (cf. [12, 11]). \square

6.2 Proof of Theorem 4.1

It remains to show that

$$H_{\text{crys}}^0(U, R^2 f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W})^{F=p}[1/p] \cong H_{\text{rig}}^0(U, R^2 f_* \mathcal{O}_{\mathcal{X}}^\dagger)^{F=p}$$

By [25, Cor. A.13], the F -isocrystal $R^2 f_{\text{crys}*} \mathcal{O}_{\mathcal{X}/W} \otimes \mathbb{Q}_p$ is naturally isomorphic to the convergent F -isocrystal $R^2 f_* \mathcal{O}_{\mathcal{X}/K}$ constructed by Ogus [26, Thm. 3.1]. By Ertl-Vezzani's recent proof of the Berthelot's conjecture [14], $R^2 f_* \mathcal{O}_{\mathcal{X}/K}$ admits an overconvergent extension $R^2 f_* \mathcal{O}_{\mathcal{X}}^\dagger$. By Kedlaya's fully faithfulness [20],

$$\text{Hom}_{\text{F-Isoc}}(\mathcal{O}_U(-1), R^2 f_* \mathcal{O}_{\mathcal{X}/K}) \cong \text{Hom}_{\text{F-Isoc}^\dagger}(\mathcal{O}_U^\dagger(-1), R^2 f_{\text{rig}} \mathcal{O}_{\mathcal{X}}^\dagger) \cong H_{\text{rig}}^0(U, R^2 f_* \mathcal{O}_{\mathcal{X}}^\dagger)^{F=p}$$

This completes the proof of Theorem 4.1.

Acknowledgements

The first named author would like to thank Prof. Matthew Morrow for important communications on F -crystals and to the second named author for bringing attention to this problem. The second named author would like to thank Veronika Ertl and Timo Keller for important discussions. Veronika Ertl and Timo Keller were in early stage of this project.

References

- [1] Fabrizio Andreatta and Luca Barbieri-Viale. Crystalline realizations of 1-motives. *Mathematische Annalen*, 331, 03 2003.
- [2] Werner Bauer. On the conjecture of Birch and Swinnerton-Dyer for abelian varieties over function fields in characteristic $p > 0$. *Inventiones mathematicae*, 108(2):263–288, 1992.
- [3] Pierre Berthelot. *Cohomologie rigide et cohomologie rigide à supports propres*. Université de Rennes 1. Institut de Recherche Mathématique de Rennes [IRMAR], 1996.
- [4] Pierre Berthelot. *Cohomologie Cristalline des Schemas de Caractéristique $p > 0$* , volume 407. Springer, 2006.
- [5] Pierre Berthelot, Lawrence Breen, and William Messing. *Théorie de Dieudonné cristalline II*, volume 930. Springer, 2006.
- [6] Pierre Berthelot and William Messing. Théorie de Dieudonné cristalline III: théorèmes d'équivalence et de pleine fidélité. *The Grothendieck Festschrift: A Collection of Articles Written in Honor of the 60th Birthday of Alexander Grothendieck*, pages 173–247, 2007.
- [7] Pierre Berthelot and Arthur Ogus. *Notes on Crystalline Cohomology*. Princeton University Press, Princeton, 1978.
- [8] Anna Cadoret, Chun Yin Hui, and Akio Tamagawa. \mathbb{Q}_ℓ -versus \mathbb{F}_ℓ -coefficients in the Grothendieck-Serre/Tate conjectures. *Israel J. Math.*, 257(1):71–101, 2023.
- [9] J.-L. Colliot-Thélène and A. N. Skorobogatov. Descente galoisienne sur le groupe de Brauer. *J. reine angew. Math.*, 682:141–165, 2013.
- [10] M. D'Addezio. Boundedness of the p -primary torsion of the Brauer group of an abelian variety. *Compositio Mathematica*, 160(2):463–480, 2024.
- [11] Aise Johan de Jong. Homomorphisms of Barsotti-Tate groups and crystals in positive characteristic. *Inventiones mathematicae*, 134:301–333, 1998.
- [12] Arthur J de Jong. Crystalline Dieudonné module theory via formal and rigid geometry. *Publications Mathématiques de l'IHÉS*, 82:5–96, 1995.

- [13] V. Ertl, T. Keller, and Y. Qin. Comparison of different Tate conjectures. Preprint at <http://arxiv.org/abs/2012.01337v2>.
- [14] Veronika Ertl and Alberto Vezzani. Berthelot’s conjecture via homotopy theory. *arXiv:2406.02182*, 2024.
- [15] K. Česnavičius. Purity for the Brauer group. *Duke Mathematical Journal*, 168(8):1461–1486, 2019.
- [16] A Grothendieck. Technique de descente et theoremes d’existence en geometrie algebrique. v, vi: Les schemas de picard. sm. bourbaki 1961/62, exp 232, 236. *Fondements de la geometrie algebrique. Extraits du Sm. Bourbaki*, 62, 1957.
- [17] A Grothendieck. Le groupe de Brauer III: exemples et compléments. *Dix Exposes sur la Cohomologie des Schemas, Masson et Cie*, 1968.
- [18] Luc Illusie. Complexe de de Rham-Witt. *Astérisque*, 63:83–112, 1979.
- [19] Nicholas Katz. Slope filtration of F-crystals. *Astérisque*, 63(1):979, 1979.
- [20] Kiran S Kedlaya. Full faithfulness for overconvergent f-isocrystals. *Geometric aspects of Dwork theory*, 2:819–835, 2004.
- [21] Kiran S Kedlaya. Notes on isocrystals. *arXiv preprint arXiv:1606.01321v6*, 2016.
- [22] Raju Krishnamoorthy and Ambrus Pál. Rank 2 local systems and abelian varieties. *Selecta Mathematica*, 27(4):51, 2021.
- [23] B Mazur and L Roberts. Local Euler characteristics. *Inventiones mathematicae*, 9:201–234, 1970.
- [24] James S Milne. *Etale cohomology (PMS-33)*. Princeton university press, 1980.
- [25] M. Morrow. A Variational Tate Conjecture in crystalline cohomology. *Journal of the European Mathematical Society*, 21(11):3467–3511, 2019.
- [26] Arthur Ogus. F-isocrystals and De Rham cohomology II–convergent isocrystals. *Duke Mathematical Journal (C)*, 1984.
- [27] Ambrus Pál. The p -adic monodromy group of abelian varieties over global function fields of characteristic p . *Documenta Mathematica*, 27:1509–1579, 2022.
- [28] Y. Qin. On the Brauer groups of fibrations II. Preprint at <https://arxiv.org/abs/2103.06910v2>.
- [29] Yanshuai Qin. On the Brauer groups of fibrations. *Mathematische Zeitschrift*, 307(1):1–20, 2024.

- [30] A.N. Skorobogatov and Y.G. Zarkhin. A finiteness theorem for the Brauer group of Abelian varieties and K3 surfaces. *Journal of Algebraic Geometry*, 17(3):481–502, 2008.
- [31] A.N. Skorobogatov and Y.G. Zarkhin. A finiteness theorem for the Brauer group of K3 surfaces in odd characteristic. *International Mathematics Research Notices*, (21):11404–11418, 2015.
- [32] X. Yuan. Comparison of arithmetic Brauer groups with geometric brauer group, 2021. Preprint at <https://arxiv.org/abs/2011.12897v2>.

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT REGENSBURG, 93040 REGENSBURG, GERMANY
E-mail address: yanshuai.qin@ur.de
IMJ-PRG, SORBONNE UNIVERSITÉ, 4 PLACE JUSSIEU, 75005 PARIS, FRANCE
E-mail address: zhenghuili@imj-prg.fr