

Existence and regularity results for a Neumann problem with mixed local and nonlocal diffusion

Craig Cowan^{a*}, Mohammad El Smaily^{b†} and Pierre Aime Feulefack^{b‡}

^a Department of Mathematics, University of Manitoba,
Winnipeg, MB, Canada

^b Department of Mathematics, University of Northern British Columbia,
Prince George, BC, Canada

Abstract

In this paper, we consider an elliptic problem driven by a mixed local-nonlocal operator with drift and subject to nonlocal Neumann condition. We prove the existence and uniqueness of a solution $u \in W^{2,p}(\Omega)$ of the considered problem with L^p -source function when p and s are in a certain range.

Keywords. Maximum principle, operators of mixed order with Drift,

2010 Mathematics Subject Classification. 35A09, 35B50, 35B65, 35R11, 35J67, 47A75.

1 Introduction and main results

In this paper, we are interested in the study of the following problem

$$\begin{cases} Lu := -\Delta u(x) + (-\Delta)^s u(x) - q(x) \cdot \nabla u(x) + a(x)u = f(x) & \text{in } \Omega \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega \\ \mathcal{N}_s u = 0 & \text{on } \mathbb{R}^N \setminus \bar{\Omega}, \end{cases} \quad (1.1)$$

with a mixed diffusion and a new type of Neumann boundary conditions. The new boundary condition is $\mathcal{N}_s u = 0$ on $\mathbb{R}^N \setminus \bar{\Omega}$, where $\mathcal{N}_s u$ — known as the nonlocal normal derivative of u is given by

$$\mathcal{N}_s u(x) = C_{N,s} \int_{\Omega} \frac{u(x) - u(y)}{|x - y|^{N+2s}} dy \quad x \in \mathbb{R}^N \setminus \bar{\Omega}. \quad (1.2)$$

The diffusion term is a superposition of the classical Laplacian (local diffusion) and the fractional Laplacian $(-\Delta)^s$ for certain values of $s \in (0, 1)$ that will be specified later. It is well known that the fractional Laplacian represents a nonlocal diffusion in the medium.

*Email: craig.cowan@umanitoba.ca

†Corresponding author. Email: mohammad.elsmaily@unbc.ca

‡Email: pierre.feulefack@aims-cameroon.org

We recall that the operator $(-\Delta)^s$, with $s \in (0, 1)$, stands for the fractional Laplacian and it is defined for compactly supported function $u : \mathbb{R}^N \rightarrow \mathbb{R}$ of class \mathcal{C}^2 by

$$(-\Delta)^s u(x) = C_{N,s} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}^N \setminus B_\varepsilon(x)} \frac{u(x) - u(y)}{|x - y|^{N+2s}} dy \quad (1.3)$$

with the same normalization constant $C_{N,s}$ as in (1.2) given by

$$C_{N,s} := \pi^{-\frac{N}{2}} 2^{2s} s \frac{\Gamma(\frac{N}{2} + s)}{\Gamma(1 - s)}. \quad (1.4)$$

We refer the reader to the results of Silvestre [10] about the spaces that $(-\Delta)^s u$ belongs to according to the regularity of the function u .

The boundary conditions in (1.1) consist of the classical Neumann boundary condition $\frac{\partial u}{\partial \nu} = 0$ on $\partial\Omega$ (ν is the outward unit normal on $\partial\Omega$) and the nonlocal boundary condition $\mathcal{N}_s u = 0$ (see [8]) on $\mathbb{R}^N \setminus \Omega$. The classical Neumann condition states that there is no flux through the boundary of the domain. On the other hand, the nonlocal boundary condition $\mathcal{N}_s u = 0$ states that if a particle is in $\mathbb{R}^N \setminus \overline{\Omega}$, it may come back to any point $y \in \Omega$ with the probability density of jumping from x to y being proportional to $|x - y|^{-N-2s}$. A detailed description of (1.1) is given in [9]. The condition $\mathcal{N}_s u = 0$ is interpreted in [9] as a condition that arises from the superposition of Brownian and Lévy processes.

The PDE

$$-\Delta u(x) + (-\Delta)^s u - q(x) \cdot \nabla u(x) + a(x)u = f(x) \quad \text{in } \Omega \quad (1.5)$$

has been extensively studied when $q \equiv 0$ and the boundary condition is of Dirichlet type. That is, $u \equiv 0$ in $\mathbb{R}^N \setminus \Omega$. Existence and regularity of solutions for (1.5), as well as maximum principles are among the results obtained in [3], [4], [5], and [11], where the advection q is absent and the boundary condition is of Dirichlet type. The authors of this paper studied (1.5) in the recent work [6], where an advection term is present and (1.5) is coupled with the Dirichlet condition $u \equiv 0$ on $\mathbb{R} \setminus \overline{\Omega}$.

The recent work [7] considers (1.5) with $q \equiv 0$ and $a \equiv 0$ to provide spectral properties of the mixed diffusion operator. The work [8] considers a purely nonlocal diffusion and provides existence results for the problem with nonlocal Neumann conditions. It is important to note that [8] does not consider a PDE with a mixed diffusion and it does not account for advection.

The domain and the coefficients. Throughout this paper, we assume that the domain Ω is an open bounded connected subset of \mathbb{R}^N with smooth boundary $\partial\Omega$. The coefficients q and a are assumed to be smooth with $a \geq 0$ and not identically zero.

The normal derivative of u on $\partial\Omega$. Our solutions will, in general, be $C^1(\overline{\Omega})$ but the extension (\tilde{u} defined later) will not be sufficiently smooth. Hence to compute $\partial_\nu u(x)$ on $\partial\Omega$, we are using

$$\partial_\nu u(x) = \lim_{t \rightarrow 0^-} \frac{u(x_0 + t\nu(x_0)) - u(x_0)}{t},$$

where $\nu(x)$ is the unit outward normal to $\partial\Omega$ at $x \in \partial\Omega$.

We prove the following results for problem (1.1).

Theorem 1.1. *Let Ω be an open bounded set of \mathbb{R}^N with smooth boundary and $f \in L^p(\Omega)$. Then,*

1. *if $\frac{N-1}{2N} < s < \frac{1}{2}$ and $N < p < \frac{1}{1-2s}$, problem (1.1) admits a unique solution $u \in W^{2,p}(\Omega)$.*
2. *if $p > N$ and $\frac{1}{2} \leq s < \frac{1}{2} + \frac{1}{2p}$, problem (1.1) admits a unique solution $u \in W^{2,p}(\Omega)$.*

2 Some functional spaces

Let $\Omega \subset \mathbb{R}^N$ be an open bounded set. For a vector field $q : \Omega \rightarrow \mathbb{R}^N$, we write $q \in L^\infty(\Omega)$ (resp. $q \in \mathcal{C}^{0,\alpha}(\overline{\Omega})$) whenever $q_j \in L^\infty(\Omega)$ (resp. $q_j \in \mathcal{C}^{0,\alpha}(\overline{\Omega})$), $j = 1, 2, \dots, N$. We denote by $\mathcal{C}^{k,\alpha}(\overline{\Omega})$, $0 < \alpha \leq 1$, the Banach space of functions $u \in \mathcal{C}^k(\overline{\Omega})$ such that derivative of order k belong to $\mathcal{C}^{0,\alpha}(\overline{\Omega})$ with the norm

$$\|u\|_{\mathcal{C}^{k,\alpha}(\overline{\Omega})} := \|u\|_{\mathcal{C}^k(\overline{\Omega})} + \sum_{|\tau|=k} [D^\tau u]_{\mathcal{C}^{0,\alpha}(\overline{\Omega})},$$

where the Hölder seminorm $[\cdot]_{\mathcal{C}^{0,\alpha}(\overline{\Omega})}$ is given by

$$[u]_{\mathcal{C}^{0,\alpha}(\overline{\Omega})} = \sup_{x,y \in \overline{\Omega}, x \neq y} \frac{|u(x) - u(y)|}{|x - y|^\alpha}$$

and $\mathcal{C}^{0,\alpha}(\overline{\Omega})$ is the Banach space of functions $u \in \mathcal{C}^0(\overline{\Omega})$ which are Hölder continuous with exponent α in Ω and the norm $\|u\|_{\mathcal{C}^{0,\alpha}(\overline{\Omega})} = \|u\|_{L^\infty(\Omega)} + [u]_{\mathcal{C}^{0,\alpha}(\overline{\Omega})}$. We refer the reader to the results of Silvestre [10] about the spaces that $(-\Delta)^s u$ belongs to according to the regularity of the function u .

If $k \in \mathbb{N}$, as usual we set

$$W^{k,p}(\Omega) := \left\{ u \in L^p(\Omega) : D^\alpha u \text{ exists for all } \alpha \in \mathbb{N}^N, |\alpha| \leq k \text{ and } D^\alpha u \in L^p(\Omega) \right\}$$

for the Banach space of k -times (weakly) differentiable functions in $L^p(\Omega)$. Moreover, in the fractional setting, for $s \in (0, 1)$ and $p \in [1, \infty)$, we set

$$W^{s,p}(\Omega) := \left\{ u \in L^p(\Omega) : \frac{u(x) - u(y)}{|x - y|^{\frac{n}{p} + s}} \in L^p(\Omega \times \Omega) \right\}.$$

The space $W^{s,p}(\Omega)$ is a Banach space with the norm

$$\|u\|_{W^{s,p}(\Omega)} = \left(\|u\|_{L^p(\Omega)}^p + \iint_{\Omega \times \Omega} \frac{|u(x) - u(y)|^p}{|x - y|^{n+sp}} dx dy \right)^{\frac{1}{p}}.$$

Next, we set $\mathcal{Q} := (\mathbb{R}^N \times \mathbb{R}^N) \setminus (\mathbb{R}^N \setminus \Omega)^2$ and define the space $\mathcal{X}^s(\Omega)$ by

$$\mathcal{X}^s(\Omega) := \left\{ u \in L^2(\Omega) : u|_\Omega \in W^{1,2}(\Omega); [u]_{\mathcal{X}^s(\Omega)} < \infty \right\}.$$

Here, the corresponding Gagliardo seminorm $[\cdot]_s$ is given by

$$([u]_s)^2 := \int_\Omega |\nabla u|^2 dx + \iint_{\mathcal{Q}} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy.$$

We also define the space

$$\mathcal{X}_K^s(\Omega) := \{u \in L^2(\Omega) : u|_\Omega \in W^{1,2}(\Omega); [u]_{s,K} < \infty\},$$

where,

$$([u]_{s,K})^2 := \int_\Omega |\nabla u|^2 dx + \int_\Omega \int_\Omega |u(x) - u(y)|^2 K_\Omega(x, y) dx dy,$$

and $K_\Omega : \Omega \times \Omega \rightarrow \mathbb{R}$ is a measurable (regional) kernel given by

$$K_\Omega(x, y) := \frac{1}{|x - y|^{N+2s}} + k_\Omega(x, y) \quad (2.1)$$

with

$$k_\Omega(x, y) := \int_{\mathbb{R}^N \setminus \Omega} \frac{1}{|x - z|^{N+2s} |y - z|^{N+2s} \int_\Omega \frac{1}{|z - z'|^{N+2s}} dz'} dz, \quad x, y \in \Omega. \quad (2.2)$$

Note that the space $\mathcal{X}^s(\Omega)$ (resp. $\mathcal{X}_K^s(\Omega)$) is a Hilbert space when furnished with the scalar product $\langle u, v \rangle_{\mathcal{X}^s(\Omega)}$ (resp. $\langle u, v \rangle_{\mathcal{X}_K^s(\Omega)}$)

$$\langle u, v \rangle_{\mathcal{X}^s(\Omega)} := \int_\Omega uv dx + \int_\Omega \nabla u \cdot \nabla v dx + \iint_\Omega \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy$$

and the corresponding norm given $\|u\|_{\mathcal{X}^s(\Omega)} = \langle u, u \rangle_{\mathcal{X}^s(\Omega)}^{\frac{1}{2}}$ (resp. $\|u\|_{\mathcal{X}_K^s(\Omega)} = \langle u, u \rangle_{\mathcal{X}_K^s(\Omega)}^{\frac{1}{2}}$). Finally, we define the space $\mathcal{L}_s^1(\mathbb{R}^N)$ by

$$\mathcal{L}_s^1(\mathbb{R}^N) := \{u : \mathbb{R}^N \rightarrow \mathbb{R}, \text{ such that } u \text{ is measurable and } \|u\|_{\mathcal{L}_s^1(\mathbb{R}^N)} < \infty\},$$

where

$$\|u\|_{\mathcal{L}_s^1(\mathbb{R}^N)} := \int_{\mathbb{R}^N} \frac{|u(y)|}{1 + |y|^{N+2s}} dx.$$

Definition 2.1. Let $u \in \mathcal{C}^{0,1}(\overline{\Omega})$ and define the function \tilde{u} on \mathbb{R}^N as

$$\tilde{u}(x) = \begin{cases} u(x) & \text{if } x \in \overline{\Omega} \\ u_1(x) & \text{if } x \in \mathbb{R}^N \setminus \overline{\Omega}, \end{cases} \quad (2.3)$$

where

$$u_1(x) := \frac{\int_\Omega \frac{u(y)}{|x - y|^{N+2s}} dy}{\int_\Omega \frac{1}{|x - y|^{N+2s}} dy}, \quad x \in \mathbb{R}^N \setminus \overline{\Omega}. \quad (2.4)$$

Remark 2.2. We note that $\mathcal{N}_s \tilde{u}(x) = 0$ for all $x \in \mathbb{R}^N \setminus \overline{\Omega}$.

We now recall the following results that lead to integration by parts in a fractional setting from [2]:

Lemma 2.3 ([2]). Let $u, v : \mathbb{R}^N \rightarrow \mathbb{R}$ be two functions such that $\mathcal{N}_s v = 0$ on $\mathbb{R}^N \setminus \overline{\Omega}$. Then

$$\begin{aligned} & \int_{\Omega} \int_{\Omega} (u(x) - u(y))(v(x) - v(y)) K_{\Omega}(x, y) \, dx dy \\ &= C_{N,s} \iint_{\mathcal{Q}} \frac{(\tilde{u}(x) - \tilde{u}(y))(v(x) - v(y))}{|x - y|^{N+2s}} \, dx dy \end{aligned} \quad (2.5)$$

From Lemma 2.3 and [8, Lemma 3.3], we deduce the integration by parts formula

$$\iint_{\mathcal{Q}} \frac{(\tilde{u}(x) - \tilde{u}(y))(v(x) - v(y))}{|x - y|^{N+2s}} \, dx dy = \int_{\Omega} v(-\Delta)^s \tilde{u} \, dx + \int_{\mathbb{R}^N \setminus \Omega} v \mathcal{N}_s \tilde{u} \, dx \quad (2.6)$$

for u and v being two \mathcal{C}^2 bounded functions in \mathbb{R}^N .

2.1 Estimates on \tilde{u}

Let $N < p < \infty$ and suppose $u \in W^{2,p}(\Omega)$ and hence u is $C^{0,1}(\overline{\Omega})$. Then note that \tilde{u} is smooth near x for any $x \notin \overline{\Omega}$. So the only real question on the smoothness of \tilde{u} is when $x \notin \Omega$ and $\delta(x) = \text{dist}(x, \partial\Omega) < 1$. For $x \notin \overline{\Omega}$, let $\hat{x} \in \partial\Omega$ be such that

$$|x - \hat{x}| = \inf_{z \in \partial\Omega} |z - x|.$$

Lemma 2.4. Let Ω be an open bounded set of \mathbb{R}^N with smooth boundary and let $N < p < \infty$ and suppose $u \in W^{2,p}(\Omega)$ with $\|u\|_{W^{2,p}} \leq 1$. The following estimates are all independent of u .

(i) For $0 < s < 1/2$ there is some C such that for all $x \in \mathbb{R}^N \setminus \overline{\Omega}$ with $\delta(x) < 1$ we have

$$|\nabla u_1(x)| \leq \frac{C}{(\delta(x))^{1-2s}}.$$

Therefore, $\tilde{u} \in W_{loc}^{1,q}(\mathbb{R}^N)$ for all $1 < q < \frac{1}{1-2s}$ — after applying the co-area formula.

(ii) If $s = 1/2$ then $\tilde{u} \in W_{loc}^{1,q}(\mathbb{R}^N)$ for all $1 \leq q < \infty$.

(iii) Let $1/2 < s < 1$. Then, there is some $C > 0$ such that $|\nabla \tilde{u}(x)| \leq C$ on \mathbb{R}^N .

Proof. Let $x \in \mathbb{R}^N \setminus \overline{\Omega}$ with $\delta(x) < 1$. For simplicity, we set

$$F(x) = \int_{\Omega} |x - y|^{-N-2s} dy.$$

Since $u \in W^{2,p}(\Omega)$ with $p > N$, we have that $u \in \mathcal{C}^{0,1}(\overline{\Omega})$. It follows from [8, Proposition 5.2] and the regularity of Ω that \tilde{u} is continuous in \mathbb{R}^N . Moreover, a direct computation shows that

$$\begin{aligned} \frac{(F(x))^2 \nabla u_1(x)}{N + 2s} &= \int_{\Omega} (u(y) - u(\hat{x})) |x - y|^{-N-2s} dy \int_{\Omega} |x - y|^{-N-2s-2} (x - y) dy \\ &+ \int_{\Omega} |x - y|^{-N-2s} dy \int_{\Omega} (u(\hat{x}) - u(y)) |x - y|^{-N-2s-2} (x - y) dy. \end{aligned}$$

We now use the bound on u given by $|u(y) - u(\hat{x})| \leq C_0|y - \hat{x}|$ to give

$$\begin{aligned} \frac{(F(x))^2 |\nabla u_1(x)|}{N+2s} &\leq C_0 \int_{\Omega} \frac{|y - \hat{x}|}{|x - y|^{N+2s}} dy \int_{\Omega} \frac{1}{|x - y|^{N+2s+1}} dy \\ &\quad + C_0 \int_{\Omega} \frac{1}{|x - y|^{N+2s}} dy \int_{\Omega} \frac{|y - \hat{x}|}{|x - y|^{N+2s+1}} dy. \end{aligned}$$

Now note that for $y \in \Omega$ we have $|x - \hat{x}| \leq |x - y|$ by the definition of \hat{x} and hence we have

$$|y - \hat{x}| \leq |y - x| + |x - \hat{x}| \leq 2|y - x|.$$

It follows using the above inequality that

$$\begin{aligned} \frac{(F(x))^2 |\nabla u_1(x)|}{N+2s} &\leq C_1 \int_{\Omega} \frac{1}{|x - y|^{N+2s-1}} dy \int_{\Omega} \frac{1}{|x - y|^{N+2s+1}} dy \\ &\quad + C_1 \int_{\Omega} \frac{1}{|x - y|^{N+2s}} dy \int_{\Omega} \frac{1}{|x - y|^{N+2s}} dy. \end{aligned}$$

Noticing that the second term in the right hand side is just $(F(x))^2$, we have

$$\frac{|\nabla u_1(x)|}{N+2s} \leq C_1 + C_2 \frac{\int_{\Omega} \frac{1}{|x - y|^{N+2s-1}} dy \int_{\Omega} \frac{1}{|x - y|^{N+2s+1}} dy}{(F(x))^2}. \quad (2.7)$$

Now, it well known from [1, Lemma 2.1] that there are constants $C_1 > 0$ and $C_2 > 0$ such that for any $x \in \mathbb{R}^N \setminus \overline{\Omega}$, we have

$$C_1 \min\{(\delta(x))^{-2s}, (\delta(x))^{-N-2s}\} \leq F(x) \leq C_2 \min\{(\delta(x))^{-2s}, (\delta(x))^{-N-2s}\}. \quad (2.8)$$

With the assumption $\delta(x) < 1$, $x \in \mathbb{R}^N \setminus \overline{\Omega}$, this reduces to

$$C_1(\delta(x))^{-2s} \leq F(x) \leq C_2(\delta(x))^{-2s}. \quad (2.9)$$

We have the more general result that for $\tau > 0$ there is some $C_1, C_2 > 0$ such that for $x \notin \overline{\Omega}$ but with $\delta(x)$ small we have

$$\frac{C_1}{(\delta(x))^\tau} \leq \int_{\Omega} \frac{1}{|x - y|^{N+\tau}} dy \leq \frac{C_2}{(\delta(x))^\tau}. \quad (2.10)$$

We now distinguish three cases: $s \in (0, 1/2)$, $s = 1/2$ or $s \in (1/2, 1)$.

Case (i): $0 < s < 1/2$. Set $R_0 = 2\text{diam}(\Omega)$, where $\text{diam}(\Omega)$ is the diameter of Ω . Since Ω is bounded and $\delta(x) < 1$, $x \in \mathbb{R}^N \setminus \overline{\Omega}$, we have that $\Omega \subset B_{2R_0+1}(x)$. It follows that

$$\int_{\Omega} \frac{1}{|x - y|^{N+2s-1}} dy \leq \int_{B_{2R_0+1}(0)} \frac{1}{|z|^{N+2s-1}} dz = C \int_0^{2R_0+1} \rho^{-2s} d\rho = C(2R_0+1)^{1-2s}. \quad (2.11)$$

Putting (2.9), (2.11) and (2.10) together, we get

$$\frac{|\nabla u_1(x)|}{N+2s} \leq C_1 + \frac{C_3}{(\delta(x))^{1-2s}}.$$

We then apply the coarea formula now to get the desired result.

Case (ii): $s = 1/2$. We know from (2.7) that

$$|\nabla u_1(x)| \leq C_1 + C_2 \frac{\int_{\Omega} \frac{1}{|x-y|^{N+2}} dy \int_{\Omega} \frac{1}{|x-y|^N} dy}{(F(x))^2},$$

and this gives combining (2.9) and (2.10),

$$|\nabla u_1(x)| \leq C_1 + C_3 \int_{\Omega} \frac{1}{|x-y|^N} dy.$$

We now estimate the last term. Since $\Omega \subset B_{2R_0+1}(x)$ and $x \notin \Omega$ with $\delta(x) < 1$. Then we have

$$G(x) = \int_{y \in \Omega} \frac{1}{|x-y|^N} dy \leq C \int_{\{z: \delta(x) \leq |z| \leq 2R_0+1\}} \frac{1}{|z|^N} dz = C \ln \left(\frac{2R_0+1}{\delta(x)} \right).$$

Then we have, after using the co-area formula,

$$\begin{aligned} \int_{\{x \notin \Omega, \delta(x) < 1\}} G(x)^q dx &\leq C^q \int_{\{x \notin \Omega, \delta(x) < 1\}} \left(\ln \left(\frac{2R_0+1}{\delta(x)} \right) \right)^q dx \\ &= C^q \int_0^1 \left(\int_{\{x \notin \Omega, \delta(x)=t\}} \left\{ \ln \left(\frac{2R_0+1}{\delta(x)} \right) \right\}^q d\sigma(x) \right) dt \\ &= C^q \int_0^1 \left(\int_{\{x \notin \Omega, \delta(x)=t\}} \left\{ \ln \left(\frac{2R_0+1}{t} \right) \right\}^q d\sigma(x) \right) dt \end{aligned}$$

There is some $C > 0$ such that $|\{x \notin \Omega : \delta(x) = t\}| \leq C$ for all $0 < t < 1$, where $|A|$ refers the the $N-1$ measure of A . From this we see (after doing a change of variables $r = 1/t$ that we have

$$\int_{\{x \notin \Omega, \delta(x) < 1\}} G(x)^q dx \leq C \int_1^{\infty} \frac{(\ln((2R_0+1)r))^q}{r^2} dr,$$

and this is finite for any $1 \leq q < \infty$. Therefore, for any $x \in \mathbb{R}^N$ and $|x| \leq R$, we have

$$\int_{B_R} |\nabla \tilde{u}(x)|^q dx \leq C_R$$

This shows that $\tilde{u} \in W_{loc}^{1,q}(\mathbb{R}^N)$ for all $1 < q < \infty$.

Case (iii): $1/2 < s < 1$. For $x \notin \bar{\Omega}$, we have from (2.7) that

$$\frac{|\nabla u_1(x)|}{N+2s} \leq C_1 + C_2 \frac{\int_{\Omega} \frac{1}{|x-y|^{N+2s-1}} dy \int_{\Omega} \frac{1}{|x-y|^{N+2s+1}} dy}{(F(x))^2}$$

Now, since $\Omega \subset \mathbb{R}^N \setminus B_{\delta(x)}(x)$ for all $x \in \Omega^c$, we compute for $s > 1/2$,

$$\int_{\Omega} \frac{1}{|x-y|^{N+2s-1}} dy \leq \int_{\mathbb{R}^n \setminus B_{d(x)}} \frac{1}{|z|^{N+2s-1}} dz = C \int_{d(x)}^{\infty} \rho^{-2s} d\rho = C \delta(x)^{1-2s}.$$

This combined with (2.9) and (2.10) yield

$$\frac{|\nabla u_1(x)|}{N+2s} \leq C_1 + C_4.$$

Hence, for all $x \in \mathbb{R}^N \setminus \overline{\Omega}$ we have

$$|\nabla u_1(x)| \leq C.$$

□

Lemma 2.5. *The following results hold:*

1. Suppose $\frac{N-1}{2N} < s < 1/2$ and $N < p < \frac{1}{1-2s}$. Then, the mapping $u \mapsto (-\Delta)^s \tilde{u}$ is continuous and compact from $W^{2,p}(\Omega)$ to $L^p(\Omega)$.
2. Suppose $p > N$ and $1/2 \leq s < \frac{1}{2} + \frac{1}{2p}$. The mapping $u \mapsto (-\Delta)^s \tilde{u}$ is continuous and compact from $W^{2,p}(\Omega)$ to $L^p(\Omega)$.

Proof. 1. For the convenience of the reader we show the continuous argument and compact argument in separate steps. Let $u \in W^{2,p}(\Omega)$ with $\|u\|_{W^{2,p}(\Omega)} \leq 1$ and let $x \in \Omega$ and then note we have

$$(-\Delta)^s \tilde{u}(x) = I + II + III$$

where

$$I(u)(x) = \int_{\{y \in \Omega: |y-x| \leq 1\}} \frac{\tilde{u}(x) - \tilde{u}(y)}{|x-y|^{N+2s}} dy,$$

$$II(u)(x) = \int_{\{y \in \mathbb{R}^N \setminus \overline{\Omega}: |y-x| \leq 1\}} \frac{\tilde{u}(x) - \tilde{u}(y)}{|x-y|^{N+2s}} dy,$$

and $III(u)(x) = \int_{\{y: |y-x| \geq 1\}} \frac{\tilde{u}(x) - \tilde{u}(y)}{|x-y|^{N+2s}} dy.$

Note that

$$|I(u)(x)| \leq \int_{\{y \notin \Omega: |y-x| \leq 1\}} \frac{C|x-y|}{|x-y|^{N+2s}} dy \leq C_2,$$

since $s < \frac{1}{2}$. Also, note that

$$|III(u)(x)| \leq \int_{\{y: |y-x| \geq 1\}} \frac{C}{|x-y|^{N+2s}} dy,$$

since \tilde{u} is bounded on \mathbb{R}^N and hence $|III(u)(x)|$ is bounded in Ω by some C . We now estimate $II(u)$. Using $z = y - x$ we have

$$|II(u)(x)| \leq \int_{|z| \leq 1} \frac{|\tilde{u}(x+z) - \tilde{u}(x)|}{|z|^{N+2s}} dz.$$

Note that

$$|\tilde{u}(x+z) - \tilde{u}(x)| \leq |z| \int_0^1 |\nabla \tilde{u}(x+tz)| dt.$$

Hence, we have

$$\begin{aligned}
\int_{\Omega} |II(u)(x)|^p dx &\leq \int_{\Omega} \left(\int_{|z|\leq 1} \frac{1}{|z|^{N+2s-1}} \int_0^1 |\nabla \tilde{u}(x+tz)| dt dz \right)^p dx \\
&\leq C \int_{\Omega} \int_{|z|\leq 1} \frac{1}{|z|^{N+2s-1}} \int_0^1 |\nabla \tilde{u}(x+tz)|^p dt dz dx \quad (\text{Jensen's inequality applied twice}) \\
&= C \int_0^1 \int_{|z|\leq 1} \frac{1}{|z|^{N+2s-1}} \left(\int_{\Omega} |\nabla \tilde{u}(x+tz)|^p dx \right) dz dt \\
&\leq C \int_0^1 \int_{|z|\leq 1} \frac{1}{|z|^{N+2s-1}} \left(\int_{|x|\leq R} |\nabla \tilde{u}(x)|^p dx \right) dz dt,
\end{aligned}$$

for some large R and hence we have

$$\int_{\Omega} |II(u)(x)|^p dx \leq C_2 \int_{|x|\leq R} |\nabla \tilde{u}(x)|^p dx.$$

Combining this with the results on $I(u)$ and $III(u)$, we see that $\Delta^s \tilde{u} \in L^p(\Omega)$ and is continuous from $W^{2,p}(\Omega)$.

We now consider the compactness. Since $W^{2,p}(\Omega)$ is a reflexive space its sufficient to show that if $u_m \rightharpoonup 0$ in $W^{2,p}(\Omega)$ then $\Delta^s \tilde{u}_m \rightarrow 0$ in $L^p(\Omega)$. Let $u_m \rightharpoonup 0$ in $W^{2,p}(\Omega)$ and hence it converges to zero in $W^{1,p}(\Omega)$ and uniformly in Ω .

$$\begin{aligned}
\frac{(F(x))^2 \nabla \tilde{u}_m(x)}{N+2s} &= \int_{\Omega} (u_m(y) - u_m(\hat{x})) |x-y|^{-N-2s} dy \int_{\Omega} |x-y|^{-N-2s-2} (x-y) dy \\
&\quad + \int_{\Omega} |x-y|^{-N-2s} dy \int_{\Omega} (u_m(\hat{x}) - u_m(y)) |x-y|^{-N-2s-2} (x-y) dy.
\end{aligned}$$

From this we see that $|\nabla \tilde{u}_m(x)| \rightarrow 0$ a.e. in \mathbb{R}^N and note that we can use the result of Lemma 2.4 with the dominated convergence theorem to see that $\tilde{u}_m \rightarrow 0$ in $W^{1,p}(B_R)$ for all $0 < R < \infty$.

Let $x \in \Omega$ and then note

$$(-\Delta)^s \tilde{u}_m(x) = J_1(u_m)(x) + III(u_m)(x)$$

where

$$J_1(u_m)(x) = I(u_m)(x) + II(u_m)(x) = \int_{\{y:|y-x|\leq 1\}} \frac{\tilde{u}_m(x) - \tilde{u}_m(y)}{|x-y|^{N+2s}} dy.$$

As before, we can write this as

$$\begin{aligned}
&\int_{\Omega} |J_1(u_m)(x)|^p dx \\
&\leq C_1 \int_0^1 \int_{\{|z|\leq 1\}} \frac{1}{|z|^{N+2s-1}} \left(\int_{x \in \Omega} |\nabla \tilde{u}_m(x+tz)|^p dx \right) dz dx \\
&\leq C_2 \int_{|x|<R} |\nabla \tilde{u}_m(x)|^p dx,
\end{aligned}$$

for some large R and we know this goes to zero from the earlier results. Let $R > 1$ be big and note we have

$$\begin{aligned} |III(u_m)(x)| &\leq \int_{\{y:1 \leq |y-x| \leq R\}} \frac{|\tilde{u}_m(x)| + |\tilde{u}_m(y)|}{|x-y|^{N+2s}} dy \\ &\quad + \int_{\{y:|y-x| \geq R\}} \frac{|\tilde{u}_m(x)| + |\tilde{u}_m(y)|}{|x-y|^{N+2s}} dy \\ &\leq \int_{\{y:1 \leq |y-x| \leq R\}} \frac{|\tilde{u}_m(x)| + |\tilde{u}_m(y)|}{|x-y|^{N+2s}} dy + CR^{-2s}, \end{aligned}$$

where C is from the fact that $|\tilde{u}_m(x)| \leq C_1$ on \mathbb{R}^N (independent of m). From this we see that

$$\begin{aligned} \int_{\Omega} |III(u_m)(x)|^p dx &\leq C_p R^{-2sp} + C_p \int_{\Omega} \left(\int_{1 \leq |y-x| \leq R} \frac{|\tilde{u}_m(x)| + |\tilde{u}_m(y)|}{|x-y|^{N+2s}} dy \right)^p dx \\ &\leq C_p R^{-2sp} + C_p \int_{\Omega} \left(2 \sup_{|\zeta| \leq R} |\tilde{u}_m(\zeta)| \int_{|z| \geq 1} \frac{1}{|z|^{N+2s}} dz \right)^p dx \end{aligned}$$

and note that the second term goes to zero when $m \rightarrow \infty$. Hence, we have

$$\limsup_m \int_{\Omega} |III(u_m)(x)|^p dx \leq C_p R^{-2sp},$$

and consequently $\int_{\Omega} |III(u_m)(x)|^p dx \rightarrow 0$ since we can set $R \rightarrow \infty$.

2. We now take $1/2 \leq s < 1$ and for these cases we split the integral in the definition of the fractional Laplacian as

$$\begin{aligned} (-\Delta)^s \tilde{u}(x) &= \frac{C_{N,s}}{2} \int_{\mathbb{R}^N} \frac{[\tilde{u}(x) - \tilde{u}(x+z)] + [\tilde{u}(x) - \tilde{u}(x-z)]}{|z|^{N+2s}} dz \\ &= \frac{C_{N,s}}{2} \sum_{i=1}^4 \int_{A_x^i} \frac{[\tilde{u}(x) - \tilde{u}(x+z)] + [\tilde{u}(x) - \tilde{u}(x-z)]}{|z|^{N+2s}} dz \\ &\quad + \frac{C_{N,s}}{2} \int_{\{z \in \mathbb{R}^N: |z| > 1\}} \frac{[\tilde{u}(x) - \tilde{u}(x+z)] + [\tilde{u}(x) - \tilde{u}(x-z)]}{|z|^{N+2s}} dz, \end{aligned}$$

where for $i = 1, \dots, 4$, the sets A_x^i are defined as

$$\begin{aligned} A_x^1 &= \{z : |z| \leq 1, x+z, x-z \in \Omega\}, \\ A_x^2 &= \{z : |z| \leq 1, x+z \notin \Omega, x-z \in \Omega\}, \\ A_x^3 &= \{z : |z| \leq 1, x+z \in \Omega, x-z \notin \Omega\}, \\ A_x^4 &= \{z : |z| \leq 1, x+z \notin \Omega, x-z \notin \Omega\}. \end{aligned}$$

We first estimate the following

$$\int_{x \in \Omega} \left(\int_{\{z \in \mathbb{R}^N: |z| > 1\}} \frac{|\tilde{u}(x+z) + \tilde{u}(x-z) - 2\tilde{u}(x)|}{|z|^{N+2s}} dz \right)^p dx.$$

First note that $\sup_{z \in \mathbb{R}^N} |\tilde{u}(z)| \leq \sup_{x \in \Omega} |u(x)|$ and hence the above quantity is bounded,

$$\int_{x \in \Omega} \left(\int_{\{z \in \mathbb{R}^N: |z| > 1\}} \frac{|\tilde{u}(x+z) + \tilde{u}(x-z) - 2\tilde{u}(x)|}{|z|^{N+2s}} dz \right)^p dx \leq C \|u\|_{W^{2,p}(\Omega)}^p.$$

We now estimate the integrals over A_x^i for $i = 1, \dots, 4$.

The A_x^1 term. Let $1/2 \leq s < 1$, $u \in W^{2,p}(\Omega)$ with $\|u\|_{W^{2,p}} \leq 1$ and let v denote is $W^{2,p}(\Omega)$ extension to all of \mathbb{R}^N which is compactly supported. Using a density we assume u, v smooth. We want to estimate

$$\int_{x \in \Omega} \left(\int_{z \in A_x^1} \frac{|\tilde{u}(x+z) + \tilde{u}(x-z) - 2\tilde{u}(x)|}{|z|^{N+2s}} dz \right)^p dx,$$

and note we can replace \tilde{u} with u in A_x^1 and then we can replace u with v . We now estimate this quantity. First note that for $|z| \leq 1$ we have

$$|v(x+z) - v(x) - \nabla v(x) \cdot z| \leq |z|^2 \int_0^1 \int_0^1 |D^2 v(x + t\tau z)| d\tau dt,$$

and from this we see that

$$|v(x+z) + v(x-z) - 2v(x)| \leq |z|^2 \int_0^1 \int_0^1 |D^2 v(x \pm t\tau z)| d\tau dt,$$

where the \pm indicates there is two terms we need to consider. Then we have

$$\int_{x \in \Omega} \left(\int_{z \in A_x^1} \frac{|\tilde{u}(x+z) + \tilde{u}(x-z) - 2\tilde{u}(x)|}{|z|^{N+2s}} dz \right)^p dx$$

is bounded above by

$$\int_{x \in \mathbb{R}^N} \left(\int_{z \in A_x^1} |z|^{-N-2s+2} \left(\int_0^1 \int_0^1 |D^2 v(x \pm t\tau z)| d\tau dt \right) dz \right)^p dx$$

and we can apply Jensen's inequality twice to get this bounded above by

$$C_1 \int_{x \in \mathbb{R}^N} \int_{z \in A_x^1} |z|^{-N-2s+2} \int_0^1 \int_0^1 |D^2 v(x \pm t\tau z)|^p d\tau dt dz dx$$

and by Fubini we see this bounded above by

$$C \int_0^1 \int_0^1 \int_{|z| \leq 1} |z|^{-N-2s+2} \left(\int_{\mathbb{R}^N} |D^2 v(x \pm t\tau z)|^p dx \right) dz dt d\tau,$$

and the Extension Theorem (see [?, Theorem 1, Page 259]) we have the term in the brackets bounded by the constant

$$\int_{\mathbb{R}^N} |D^2 v(x)|^p dx \leq C \|u\|_{W^{2,p}(\Omega)}^p$$

and this gives us the desired bound, that is

$$\int_{x \in \Omega} \left(\int_{z \in A_x^1} \frac{|\tilde{u}(x+z) + \tilde{u}(x-z) - 2\tilde{u}(x)|}{|z|^{N+2s}} dz \right)^p dx \leq C \|u\|_{W^{2,p}(\Omega)}^p.$$

The A_x^i term for $i = 2, 3, 4$. Note that if $z \in A_x^i$ for $i = 2, 3, 4$, we must have $|z| > \delta(x)$. In what follows we will estimate

$$\int_{x \in \Omega} \left(\int_{z \in A_x^i} \frac{|\tilde{u}(x+z) - \tilde{u}(x)|}{|z|^{N+2s}} dz \right)^p dx.$$

The same argument can be used to also estimate

$$\int_{x \in \Omega} \left(\int_{z \in A_x^i} \frac{|\tilde{u}(x-z) - \tilde{u}(x)|}{|z|^{N+2s}} dz \right)^p dx$$

since the only fact we will use will be that $|z| > \delta(x)$. So to estimate the full quantity we group the three terms into the following pairings

$$[\tilde{u}(x+z) - \tilde{u}(x)] + [\tilde{u}(x-z) - \tilde{u}(x)]$$

and then estimate

$$\int_{x \in \Omega} \left(\int_{z \in A_x^i} \frac{|\tilde{u}(x+z) + \tilde{u}(x-z) - 2\tilde{u}(x)|}{|z|^{N+2s}} dz \right)^p dx \quad \text{for } i = 2, \dots, 3.$$

We split the proof into two cases: $s = \frac{1}{2}$ and $s \in (1/2, 1)$.

The case $s = \frac{1}{2}$. Let $2 \leq i \leq 4$ and note that we have

$$\begin{aligned} & \int_{x \in \Omega} \left(\int_{z \in A_x^i} \frac{|\tilde{u}(x+z) - \tilde{u}(x)|}{|z|^{N+1}} dz \right)^p dx \\ & \leq \int_{x \in \Omega} \left(\int_{z \in A_x^i} \frac{\int_0^1 |\nabla \tilde{u}(x+tz)| dt}{|z|^N} dz \right)^p dx \\ & = \int_{x \in \Omega} \left(\int_{z \in A_x^i} \frac{\int_0^1 |\nabla \tilde{u}(x+tz)| dt}{|z|^{N-\alpha} |z|^\alpha} dz \right)^p dx \quad \text{for some } \alpha > 0 \text{ small, picked later} \\ & \leq \int_{x \in \Omega} \frac{1}{(\delta(x))^{\alpha p}} \left(\int_{z \in A_x^i} \frac{\int_0^1 |\nabla \tilde{u}(x+tz)| dt}{|z|^{N-\alpha}} dz \right)^p dx \\ & \leq C \int_{x \in \Omega} \frac{1}{(\delta(x))^{\alpha p}} \left(\int_{z \in A_x^i} \frac{\int_0^1 |\nabla \tilde{u}(x+tz)|^p dt}{|z|^{N-\alpha}} dz \right) dx \quad (\text{applying Jensen's inequality twice}) \\ & \leq C \int_{x \in \Omega} \frac{1}{(\delta(x))^{\alpha p}} \left(\int_{\delta(x) < |z| \leq 1} \frac{\int_0^1 |\nabla \tilde{u}(x+tz)|^p dt}{|z|^{N-\alpha}} dz \right) dx \\ & = C \int_0^1 \int_{|z| \leq 1} \frac{1}{|z|^{N-\alpha}} \left(\int_{\{x \in \Omega: \delta(x) \leq |z|\}} \frac{|\nabla \tilde{u}(x+tz)|^p}{(\delta(x))^{\alpha p}} dx \right) dz dt. \end{aligned}$$

We now fix $0 < |z| \leq 1$ and $0 < t < 1$ and note for $1 < q < \infty$ we have

$$\int_{\{x \in \Omega: \delta(x) \leq |z|\}} \frac{|\nabla \tilde{u}(x+tz)|^p}{(\delta(x))^{\alpha p}} dx \leq \left(\int_{\Omega} |\nabla \tilde{u}(x+tz)|^{pq} dx \right)^{\frac{1}{q}} \left(\int_{\Omega} \frac{1}{(\delta(x))^{\alpha pq}} dx \right)^{\frac{1}{q}}$$

and so for fixed q we can take $\alpha > 0$ small enough so that $\alpha p q' < 1$ and the internal involving the distance function is bounded and the other integral is bounded (independent of z and t) after considering the earlier gradient bound on \tilde{u} . This shows that

$$\int_{x \in \Omega} \left(\int_{z \in A_x^i} \frac{|\tilde{u}(x+z) - \tilde{u}(x)|}{|z|^{N+1}} dz \right)^p dx$$

is bounded.

The case $s \in (1/2, 1)$. Let $2 \leq i \leq 4$ and u be as above. Recalling for $z \in A_x^i$ we have $|z| > \delta(x)$ we have

$$\begin{aligned} & \int_{x \in \Omega} \left(\int_{z \in A_x^i} \frac{|\tilde{u}(x+z) - \tilde{u}(x)|}{|z|^{N+2s}} dz \right)^p dx \\ & \leq \int_{\Omega} \left(\int_{A_x^i} \frac{\int_0^1 |\nabla \tilde{u}(x+tz)| dt}{|z|^{N+2s-1}} dz \right)^p dx \\ & = \int_{\Omega} \left(\int_{A_x^i} \frac{\int_0^1 |\nabla \tilde{u}(x+tz)| dt}{|z|^{N+2s-1-\alpha} |z|^\alpha} dz \right)^p dx \quad \alpha > 0 \\ & \leq \int_{\Omega} \frac{1}{(\delta(x))^{\alpha p}} \left(\int_{A_x^i} \frac{\int_0^1 |\nabla \tilde{u}(x+tz)| dt}{|z|^{N+2s-1-\alpha}} dz \right)^p dx \\ & \leq C \int_{\Omega} \frac{1}{(\delta(x))^{\alpha p}} \left(\int_{A_x^i} \frac{\int_0^1 |\nabla \tilde{u}(x+tz)|^p dt}{|z|^{N+2s-1-\alpha}} dz \right) dx \quad (\text{Jensen's inequality applied twice}) \end{aligned}$$

If we now assume that $|\nabla \tilde{u}| \leq C$ then we get this is bounded above by

$$C \int_{x \in \Omega} \frac{1}{\delta(x)^{\alpha p}} \left(\int_{z \in A_x^i} \frac{1}{|z|^{N+2s-1-\alpha}} dz \right) dx,$$

and since $A_x^i \subset \{z : |z| \leq 1, |z| > \delta(x)\}$ then to have this bounded its sufficient that $\alpha p < 1$ and $2s - 1 - \alpha < 0$. Hence we see its sufficient that $2s - 1 < \frac{1}{p}$. The compactness proof follows the same ideas as the previous range of s . □

The following result is a maximum principle that we will use in the proof of existence of a solution.

Theorem 2.6. *Let $\Omega \subset \mathbb{R}^N$ be an open bounded set, q and a be smooth with $a(x) \geq 0$ in Ω . Suppose $u \in W^{2,p}(\Omega)$ is a solution of*

$$\begin{cases} Lu = 0 & \text{in } \Omega, \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega, \\ \mathcal{N}_s u = 0 & \text{on } \mathbb{R}^N \setminus \bar{\Omega}. \end{cases} \quad (2.12)$$

Then, $\tilde{u} \equiv 0$ in \mathbb{R}^N .

Proof. Let u denote the solution and, for notation, we also let u denote \tilde{u} outside of Ω . Suppose $x_0 \in \bar{\Omega}$ such that $u(x_0) = \inf_{\Omega} u$. We first rule out $x_0 \in \partial\Omega$. If $x_0 \in \partial\Omega$, then we take $x_m \notin \bar{\Omega}$ such that $x_m \rightarrow x_0$ as $m \rightarrow +\infty$. Using the nonlocal boundary condition we get

$$0 = \int_{\Omega} \frac{u(y) - u(x_m)}{|y - x_m|^{N+2s}} dy.$$

Passing to the limit, we see that

$$0 = \int_{\Omega} \frac{u(y) - u(x_0)}{|y - x_0|^{N+2s}} dy,$$

which shows that $u = C = \text{const.}$ is constant in Ω and hence constant in \mathbb{R}^N . Then note from the equation we have $a(x)C = f(x)$ in Ω and this shows that $u = C \geq 0$ provided $a(x)$ not identically zero which we have assumed and hence we are done. We now suppose $x_0 \in \Omega$ is such that $u(x_0) = \inf_{\Omega} u$ and we suppose $u(x_0) < 0$.

We can also suppose that $u(x_0) < \inf_{\partial\Omega} u$. Note that, from the definition of u_1 , we get $u(x) > u(x_0)$ for all $x \in \mathbb{R}^N \setminus \bar{\Omega}$.

Fix $\sigma > 0$ such that $u(x_0) + 10\sigma < \min_{\partial\Omega} u$ and we also assume $u(x_0) + 10\sigma < 0$. For $\varepsilon > 0$ small set $\Omega_\varepsilon = \{x \in \Omega : \delta(x) > \varepsilon\}$ and $\Gamma_\varepsilon = \{x \in \Omega : \delta(x) < \varepsilon\}$. By continuity there is some $\varepsilon_0 > 0$ such that

$$u(x_0) + 8\sigma < \inf_{\Gamma_{\varepsilon_0}} u.$$

For $x \notin \bar{\Omega}$ we have

$$\begin{aligned} u(x) \int_{\Omega} \frac{1}{|x - y|^{N+2s}} dy &= \int_{\Gamma_{\varepsilon_0}} \frac{u(y)}{|x - y|^{N+2s}} dy + \int_{\Omega \setminus \Gamma_{\varepsilon_0}} \frac{u(y)}{|x - y|^{N+2s}} dy \\ &\geq (u(x_0) + 8\sigma) \int_{\Gamma_{\varepsilon_0}} \frac{1}{|x - y|^{N+2s}} dy + u(x_0) \int_{\Omega \setminus \Gamma_{\varepsilon_0}} \frac{1}{|x - y|^{N+2s}} dy \\ &= u(x_0) \int_{\Omega} \frac{1}{|x - y|^{N+2s}} dy + 8\sigma \int_{\Gamma_{\varepsilon_0}} \frac{1}{|x - y|^{N+2s}} dy \end{aligned}$$

which gives

$$u(x) \geq u(x_0) + \frac{8\sigma \int_{\Gamma_{\varepsilon_0}} \frac{1}{|x - y|^{N+2s}} dy}{\int_{\Omega} \frac{1}{|x - y|^{N+2s}} dy}.$$

From this we can show there is some $c_{\varepsilon_0} > 0$ (without loss of generality we can take $c_{\varepsilon_0} < 1$) such that $u(x) \geq u(x_0) + 8\sigma c_{\varepsilon_0}$ for all $x \notin \bar{\Omega}$.

Let η denote a standard radial mollifier with η_ε the appropriately scaled function whose support is \bar{B}_ε and set $u^\varepsilon(x) = (\eta_\varepsilon * u)(x)$ for $x \in \mathbb{R}^N$. For all $0 < \varepsilon < \varepsilon_0$ we have

$$\inf_{\partial\Omega_{\frac{\varepsilon}{2}}} u^{\frac{\varepsilon}{2}} \geq \inf_{\Gamma_\varepsilon} u,$$

and also we have $\inf_{\Omega_\tau} u^\tau \rightarrow u(x_0)$ as $\tau \rightarrow 0$. From this we see there is some $0 < \varepsilon_1 < \varepsilon_0$ such that for all $0 < \varepsilon < \varepsilon_1$ we have

$$\inf_{\Omega_{\frac{\varepsilon}{2}}} u^{\frac{\varepsilon}{2}} + 6\sigma \leq u(x_0) + 8\sigma < \inf_{\Gamma_{\varepsilon_0}} u,$$

but by the monotonicity of $\varepsilon \mapsto \inf_{\Gamma_\varepsilon} u$ we have

$$\inf_{\Omega_{\frac{\varepsilon}{2}}} u^{\frac{\varepsilon}{2}} + 6\sigma < \inf_{\Gamma_{\varepsilon_0}} u \leq \inf_{\Gamma_\varepsilon} u$$

for all $0 < \varepsilon < \varepsilon_1$ and hence we have

$$\inf_{\Omega_{\frac{\varepsilon}{2}}} u^{\frac{\varepsilon}{2}} + 6\sigma < \inf_{\partial\Omega_{\frac{\varepsilon}{2}}} u^{\frac{\varepsilon}{2}},$$

for $0 < \varepsilon < \varepsilon_1$ and hence the minimum is contained in the interior of $\Omega_{\frac{\varepsilon}{2}}$. We now want to show that $\min_{\Omega_{\frac{\varepsilon}{2}}} u^{\frac{\varepsilon}{2}} \leq u^{\frac{\varepsilon}{2}}(x)$ for all $x \in \mathbb{R}^N$. Let $0 < \varepsilon < \frac{\varepsilon_1}{10}$ with:

$$u^{\frac{\varepsilon}{2}}(x_0) < u(x_0) + 8\sigma c_{\varepsilon_0}, \quad (2.13)$$

We consider the three cases:

(i) $x \in \Omega$ with $\delta(x) < \frac{\varepsilon}{2}$, (ii) $x \notin \Omega$ with $\delta(x) < \frac{\varepsilon}{2}$ and (iii) $x \notin \Omega$ with $\delta(x) > \frac{\varepsilon}{2}$.

Case (i). Here we have $x \in \Omega$ with $\delta(x) < \frac{\varepsilon}{2}$. So in this case we have

$$u^{\frac{\varepsilon}{2}}(x) = \int_{|y-x| < \frac{\varepsilon}{2}} \eta_\varepsilon(y-x)u(y)dy,$$

and note the integral can be decomposed as

$$\int_{|y-x| < \frac{\varepsilon}{2}, y \in \Omega} \eta_\varepsilon(y-x)u(y)dy + \int_{|y-x| < \frac{\varepsilon}{2}, y \notin \Omega} \eta_\varepsilon(y-x)u(y)dy$$

and from this we see that

$$\begin{aligned} u^{\frac{\varepsilon}{2}}(x) &\geq (u(x_0) + 8\sigma c_{\varepsilon_0}) \int_{|y-x| < \frac{\varepsilon}{2}} \eta_\varepsilon(y-x)dy \\ &= u(x_0) + 8\sigma c_{\varepsilon_0} \\ &> u^{\frac{\varepsilon}{2}}(x_0) \\ &\geq \inf_{\Omega_{\frac{\varepsilon}{2}}} u^{\frac{\varepsilon}{2}}. \end{aligned}$$

Case (ii). In this case we have from the definition of \tilde{u} that

$$\begin{aligned} u^{\frac{\varepsilon}{2}}(x) \int_{\Omega} \frac{1}{|x-y|^{N+2s}} dy &= \int_{\Omega} \frac{u^{\frac{\varepsilon}{2}}(y)}{|x-y|^{N+2s}} dy \\ &= \int_{\Omega} \frac{1}{|x-y|^{N+2s}} \int_{|y-z| < \frac{\varepsilon}{2}} \eta_\varepsilon(z-y)u(z)dz dy \\ &\geq \inf_{\Omega_{\frac{\varepsilon}{2}}} u^{\frac{\varepsilon}{2}} \int_{\Omega} \frac{1}{|x-y|^{N+2s}} dy, \end{aligned}$$

where we have used the computations of Case (i).

Case (iii). This follows similarly.

From the above we have, for small enough ε , that $u^{\frac{\varepsilon}{2}}(x_{\frac{\varepsilon}{2}}) = \min_{\Omega_{\frac{\varepsilon}{2}}} u^{\frac{\varepsilon}{2}}$ (some $x_{\frac{\varepsilon}{2}} \in \Omega_{\frac{\varepsilon}{2}}$) and $u^{\frac{\varepsilon}{2}}(x_{\frac{\varepsilon}{2}}) + 6\sigma < \inf_{\partial\Omega_{\frac{\varepsilon}{2}}} u^{\frac{\varepsilon}{2}}$. Also note we have (take $\tau = \varepsilon/2$) $L(u^\tau)(x) = f^\tau(x) \geq 0$ in Ω_τ and at x_τ we have

$$-\Delta u^\tau(x_\tau) + (-\Delta)^s u^\tau(x_\tau) + q(x_\tau) \cdot \nabla u^\tau(x_\tau) = f^\tau(x_\tau) - a(x_\tau)u^\tau(x_\tau) \geq 0. \quad (2.14)$$

But $-\Delta u^\tau(x_\tau) \leq 0$, $\nabla u^\tau(x_\tau) = 0$ and note that

$$(-\Delta)^s u^{\frac{\varepsilon}{2}}(x_{\frac{\varepsilon}{2}}) = \int_{y \in \mathbb{R}^N} \frac{u^{\frac{\varepsilon}{2}}(x_{\frac{\varepsilon}{2}}) - u^{\frac{\varepsilon}{2}}(y)}{|x_{\frac{\varepsilon}{2}} - y|^{N+2s}} dy,$$

and note $y \mapsto u^{\frac{\varepsilon}{2}}(x_{\frac{\varepsilon}{2}}) - u^{\frac{\varepsilon}{2}}(y)$ is continuous in y on \mathbb{R}^N , nonpositive and not identically zero. From this we see that $(-\Delta)^s u^{\frac{\varepsilon}{2}}(x_{\frac{\varepsilon}{2}}) < 0$ and this contradicts (2.14). \square

3 Proof of existence of a solution

For $u \in W^{2,p}(\Omega)$ and $\gamma \in \mathbb{R}$, we define

$$L_\gamma u(x) = -\Delta u + \gamma(-\Delta)^s \tilde{u}(x) + a(x)u(x) + q \cdot \nabla u(x), \quad x \in \Omega$$

and we consider the family of indexed problems

$$\begin{cases} L_\gamma u = f & \text{in } \Omega \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega. \end{cases} \quad (3.1)$$

Let \mathcal{A} be the set

$$\mathcal{A} := \left\{ \gamma \in [0, 1] : \exists C_\gamma > 0 \text{ such that for all } f \in L^p(\Omega), (3.1) \text{ has a solution } \right. \\ \left. u \in W^{2,p}(\Omega) \text{ such that } \|u\|_{W^{2,p}(\Omega)} \leq C_\gamma \|f\|_{L^p(\Omega)} \right\}. \quad (3.2)$$

In (3.2), we take the constant C_γ to be the smallest constant such that $\|u\|_{W^{2,p}(\Omega)} \leq C_\gamma \|f\|_{L^p(\Omega)}$ holds for all functions $f \in L^p(\Omega)$. In other words, if $C_\gamma > \varepsilon > 0$ then there exists $f_\varepsilon \in C^{0,\alpha}(\overline{\Omega})$ such that

$$\|u\|_{W^{2,p}(\Omega)} \geq (C_\gamma - \varepsilon) \|f_\varepsilon\|_{L^p(\Omega)}. \quad (3.3)$$

By classical theory, $0 \in \mathcal{A}$. Our goal is to show that \mathcal{A} is both open and closed and since $[0, 1]$ is connected we then see that $\mathcal{A} \in \{\emptyset, [0, 1]\}$, and since its non empty, we must have $\mathcal{A} = [0, 1]$ and in particular $1 \in \mathcal{A}$ which corresponds to the result we are trying to prove.

\mathcal{A} is closed. We prove \mathcal{A} is closed. Let $\gamma_m \in \mathcal{A}$ with $\gamma_m \rightarrow \gamma$ and let $C_m = C_{\gamma_m}$ denote the associated constant. We first consider the case of C_m bounded. Let $f \in L^p(\Omega)$ with $\|f\|_{L^p} = 1$. Since $\gamma_m \in \mathcal{A}$ there is some $u_m \in W^{2,p}(\Omega)$ which satisfies (3.1), with γ_m in place of γ , and $\|u_m\|_{W^{2,p}} \leq C_m \|f\|_{L^p} = C_m$. Note we can rewrite the problem as

$$-\Delta u_m + a u_m + q \cdot \nabla u_m = f + \gamma_m \Delta^s \tilde{u}_m \quad \text{in } \Omega, \quad (3.4)$$

with $\partial_\nu u_m = 0$ on $\partial\Omega$. Since C_m is bounded then we have u_m bounded in $W^{2,p}(\Omega)$ and by passing to a subsequence we can assume that $u_m \rightharpoonup u$ in $W^{2,p}(\Omega)$ and $\|u\|_{W^{2,p}} \leq \liminf_m \|u_m\|_{W^{2,p}} \leq C_1$. Also note by our earlier compactness result we have $\Delta^s \tilde{u}_m \rightarrow \Delta^s \tilde{u}$ in $L^p(\Omega)$ and this (along with the above weak $W^{2,p}$ convergence) is sufficient convergence to pass to the limit in (3.4). This shows that $\gamma \in \mathcal{A}$.

We now consider the case of $C_m \rightarrow \infty$. Then there is some $f_m \in L^p(\Omega)$ and $u_m \in W^{2,p}(\Omega)$ which solves $L_{\gamma_m} u_m = f_m$ in Ω with $\partial_\nu u_m = 0$ on $\partial\Omega$ and

$$\|u_m\|_{W^{2,p}} \geq (C_m - 1) \|f_m\|_{L^p}.$$

By normalizing we can assume $\|u_m\|_{W^{2,p}(\Omega)} = 1$ and hence $\|f_m\|_{L^p} \rightarrow 0$. By passing to subsequences we can assume that $u_m \rightharpoonup u$ in $W^{2,p}(\Omega)$ and strongly in $W^{1,p}(\Omega)$. As before we rewrite the equation for u_m by

$$-\Delta u_m + a u_m + q \cdot \nabla u_m = f_m + \gamma_m \Delta^s \tilde{u}_m \quad \text{in } \Omega, \quad (3.5)$$

with $\partial_\nu u_m = 0$ on $\partial\Omega$. If $u = 0$ then note the right hand side of (3.5) converges to zero in $L^p(\Omega)$ and by standard elliptic theory we have $u_m \rightarrow 0$ in $W^{2,p}(\Omega)$ which contradicts the normalization of u_m . We now assume $u \neq 0$. By compactness we can pass to the limit in (3.5) to see that $u \in W^{2,p}(\Omega) \setminus \{0\}$ satisfies $L_\gamma u = 0$ in Ω with $\partial_\nu u = 0$ on $\partial\Omega$ which contradicts Theorem 2.6.

\mathcal{A} is open. Let $\gamma_0 \in \mathcal{A}$ and take $|\varepsilon|$ small; when $\gamma_0 \in \{0, 1\}$ we need to restrict the sign of ε . Our goal is to show that $\gamma = \gamma_0 + \varepsilon \in \mathcal{A}$. Fix $f \in L^p(\Omega)$ with $\|f\|_{L^p} = 1$ and since $\gamma_0 \in \mathcal{A}$ there is some $v_0 \in W^{2,p}(\Omega)$ which solves $L_{\gamma_0} v_0 = f$ in Ω with $\partial_\nu v_0 = 0$ on $\partial\Omega$. We look for a solution of (3.1) of the form $u = v_0 + \phi$. Writing out the details one sees we need $\phi \in W^{2,p}(\Omega)$ to satisfy

$$L_{\gamma_0} \phi = \varepsilon \Delta^s \tilde{v}_0 + \varepsilon \Delta^s \tilde{\phi} \quad \text{in } \Omega, \quad (3.6)$$

with $\partial_\nu \phi = 0$ on $\partial\Omega$. Define the operator $J_\varepsilon(\phi) = \psi$ where ψ satisfies

$$L_{\gamma_0} \psi = \varepsilon \Delta^s \tilde{v}_0 + \varepsilon \Delta^s \tilde{\phi} \quad \text{in } \Omega, \quad (3.7)$$

with $\partial_\nu \psi = 0$ on $\partial\Omega$. We claim that for small enough ε that J_ε is a contraction mapping on $W^{2,p}(\Omega)$ and hence by the Contraction Mapping Principle there is some $\phi \in W^{2,p}(\Omega)$ with $J_\varepsilon(\phi) = \phi$. From (3.6) one would then get a $W^{2,p}(\Omega)$ bound on ϕ and hence we'd get the desired bound on u . We first note that J_ε is into $W^{2,p}(\Omega)$ after noting the right hand side of (3.6) belongs to $L^p(\Omega)$. Let $\phi_i \in W^{2,p}(\Omega)$ and $\psi_i = J_\varepsilon(\phi_i)$ and then note we have

$$L_{\gamma_0}(\psi_2 - \psi_1) = \varepsilon \Delta^s(\tilde{\phi}_2 - \tilde{\phi}_1) \quad \text{in } \Omega,$$

with the desired boundary condition. Then we have

$$\|\psi_2 - \psi_1\|_{W^{2,p}} \leq C_{\gamma_0} |\varepsilon| \|\Delta^s(\tilde{\phi}_2 - \tilde{\phi}_1)\|_{L^p} \leq C_{\gamma_0} |\varepsilon| C \|\phi_2 - \phi_1\|_{W^{2,p}},$$

and hence we see of $|\varepsilon|$ small that J_ε is a contraction on $W^{2,p}(\Omega)$.

References

- [1] N. Abatangelo, *A remark on nonlocal Neumann conditions for the fractional Laplacian*. Archiv der Mathematik 114, no. 6 (2020), pp. 699–708.
- [2] A. Audrito, J. C. Felipe-Navarro and X. Ros-Oton, *The Neumann problem for the fractional Laplacian: regularity up to the boundary*. Annali Scuola Normale Superiore-Classe di Scienze (2022).
- [3] S. Biagi, S. Dipierro, E. Valdinoci, and E. Vecchi, *Mixed local and nonlocal elliptic operators: regularity and maximum principles*, Communications in Partial Differential Equations 47, no. 3 (2022), pp. 585–629.
- [4] S. Biagi, E. Vecchi, S. Dipierro, and E. Valdinoci, *Semilinear elliptic equations involving mixed local and nonlocal operators*, Proceedings of the Royal Society of Edinburgh Section A: Mathematics 151, no. 5 (2021), pp. 1611-1641.
- [5] S. Biagi, S. Dipierro, E. Valdinoci and E. Vecchi, *A Faber-Krahn inequality for mixed local and nonlocal operators*, Journal d'Analyse Mathématique (2023): pp. 1–43.
- [6] C. Cowan, M. El Smaily, P. A. Feulefack, *The principal eigenvalue of a mixed local and nonlocal operator with drift*, preprint (2023).
- [7] S. Dipierro, E. Proietti Lippi, and E. Valdinoci, *Linear theory for a mixed operator with Neumann conditions*, Asymptotic Analysis 128, no. 4, (2022), pp. 571–594.
- [8] S. Dipierro, X. Ros-Oton, and E. Valdinoci, *Nonlocal problems with Neumann boundary conditions*. Revista Matemática Iberoamericana 33, no. 2, (2017), , pp. 377–416.
- [9] S. Dipierro and E. Valdinoci, *Description of an ecological niche for a mixed local/nonlocal dispersal: An evolution equation and a new Neumann condition arising from the superposition of Brownian and Lévy processes*, Physica A: Statistical Mechanics and its Applications, **575**, (2021), pp. 126052.
- [10] L. Silvestre, *Regularity of the obstacle problem for a fractional power of the Laplace operator*, Communications on Pure and Applied Mathematics: A Journal Issued by the Courant Institute of Mathematical Sciences 60, no. 1 (2007), pp. 67–112.
- [11] X. Su, E. Valdinoci, Y. Wei, and J. Zhang, *Regularity results for solutions of mixed local and nonlocal elliptic equations* Mathematische Zeitschrift 302, no. 3, (2022), pp. 1855–1878.