

# The principal eigenvalue of a mixed local and nonlocal operator with drift

Craig Cowan, Mohammad El Smaily and Pierre Aime Feulefack

**Abstract.** We study the eigenvalue problem involving the mixed local-nonlocal operator  $L := -\Delta + (-\Delta)^s + q \cdot \nabla$  in a bounded domain  $\Omega \subset \mathbb{R}^N$ , where a Dirichlet condition is posed on  $\mathbb{R}^N \setminus \Omega$ . The field  $q$  stands for a drift or advection in the medium. We prove the existence of a principal eigenvalue and a principal eigenfunction for  $s \in (0, 1/2]$ . Moreover, we prove  $C^{2,\alpha}$  regularity, up to the boundary, of the solution to the problem  $Lu = f$  when coupled with a Dirichlet condition and  $0 < s < 1/2$ . To prove the regularity and the existence of a principal eigenvalue, we use a continuation argument, Krein-Rutman theorem as well as a Hopf Lemma and a maximum principle for the operator  $L$ , which we derive in this paper.

*Keywords.* Fractional diffusion with drift, principal eigenvalue and eigenfunction, mixed local and nonlocal operator, regularity, non self-adjoint operators, maximum principle,

*2010 Mathematics Subject Classification.* 35A09, 35B50, 35B65, 35R11, 35J67, 47A75.

## Contents

1. Introduction and main results	2
2. Functional setting	6
3. Hopf lemma, maximum principle, and interior regularity: proof of Theorem 1.3 and Theorem 1.5	7
4. $L^p$ theory and regularity up to the boundary: proof of Theorem 1.6 and Theorem 1.7	14
5. Proof of Theorem 1.2	18
References	20

---

M. El Smaily acknowledges partial support from the Natural Sciences and Engineering Research Council of Canada through the NSERC Discovery Grant RGPIN-2017-04313. P. Feulefack is partially funded by a Research Strategic Initiatives Grant (RSIG22) from the University of Northern British Columbia. C. Cowan acknowledges support from the Natural Sciences and Engineering Research Council of Canada through the NSERC Discovery Grant.

## 1. Introduction and main results

The study of the principal eigenvalue of an operator is essential for many important results in the analysis of elliptic and parabolic PDE as well as the analysis of elliptic and parabolic integro-differential equations (IDE). For instance, the principal eigenvalue is fundamental in the study of semi-linear problems [10, 14], bifurcation theory, stability analysis of equilibrium of reaction-diffusion [7, 8], large deviation principle, and risk-sensitive control [2]. The principal eigenvalue of an operator also plays a role in determining whether the maximum principle holds or not for the operator at hand [9, 19, 26].

We are interested in the study of the principal eigenvalue for an operator involving an advection term (or drift) and a mixed local (elliptic) and nonlocal operator. We consider the following problem

$$\begin{cases} Lu = -\Delta u + (-\Delta)^s u + q \cdot \nabla u = \lambda u & \text{in } \Omega \\ u = 0 & \text{on } \mathbb{R}^N \setminus \Omega, \end{cases} \quad (1.1)$$

where

$$\Omega \text{ is an open bounded domain of } \mathbb{R}^N \text{ with } C^{2,\alpha} \text{ boundary.} \quad (1.2)$$

The operator  $L$  is an elliptic operator (non-self-adjoint) obtained by the superposition of the classical and the fractional Laplacian  $(-\Delta)^s$  where  $s \in (0, 1/2]$ . Problem (1.1) has also an advection term  $q \cdot \nabla u$ , where

$$q : \Omega \rightarrow \mathbb{R}^N \text{ is a vector field in the Hölder space } C^{0,\alpha}(\overline{\Omega}). \quad (1.3)$$

The vector field  $q$  can be viewed as a transport flow in (1.1).

We recall that the operator  $(-\Delta)^s$ ,  $s \in (0, 1)$ , stands for the fractional Laplacian and it is defined, for a compactly supported function  $u : \mathbb{R}^N \rightarrow \mathbb{R}$  of class  $C^2$ , by

$$(-\Delta)^s u(x) = C_{N,s} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}^N \setminus B_\varepsilon(x)} \frac{u(x) - u(y)}{|x - y|^{N+2s}} dy.$$

The constant  $C_{N,s}$  in the above definition is given by

$$C_{N,s} := \pi^{-\frac{N}{2}} 2^{2s} s \frac{\Gamma(\frac{N}{2} + s)}{\Gamma(1 - s)},$$

and it is chosen so that the operator  $(-\Delta)^s$  is equivalently defined by its Fourier transform

$$\mathcal{F}((-\Delta)^s u) = |\cdot|^{2s} \mathcal{F}(u).$$

It is known that we have the following limits

$$\lim_{s \rightarrow 0^+} (-\Delta)^s u = u \quad \text{and} \quad \lim_{s \rightarrow 1^-} (-\Delta)^s u = -\Delta u \quad \text{for } u \in C_c^2(\mathbb{R}^N).$$

**Definition 1.1.** *By a principal eigenvalue of  $L$ , we mean a value  $\lambda_1 \in \mathbb{R}$ , for which (1.1) admits a positive solution  $u$  ( $u > 0$ ) in  $\Omega$ . Throughout the paper, we will denote by  $\lambda_1 = \lambda_1(\Omega, q)$  the first eigenvalue of  $L$  in  $\Omega$  subject to Dirichlet condition on  $\mathbb{R}^N \setminus \Omega$  and by  $\varphi_1$  the corresponding unique (up to multiplication by a nonzero real) eigenfunction with a constant sign over  $\Omega$ .*

As in the case of a classical elliptic operator, we can adopt the following definition for the principal eigenvalue of  $L$  (see [2, 3, 5, 6, 9] and the references therein):

$$\lambda_1 := \sup \left\{ \lambda \in \mathbb{R} \text{ such that } \exists \varphi \in C^2(\Omega) \cap C^1(\overline{\Omega}) \cap C_c(\mathbb{R}^N), \varphi > 0 \text{ in } \Omega, \right. \\ \left. \varphi = 0 \text{ in } \mathbb{R}^N \setminus \Omega \text{ satisfying } L\varphi \geq \lambda\varphi \text{ in } \Omega \right\}. \quad (1.4)$$

The characterization (1.4) of  $\lambda_1$  will become clear from the proofs we provide in this paper.

The interest in the study of problems involving mixed local-nonlocal operator has been growing rapidly in recent years. This is due to their ability to describe the superposition of two stochastic processes with different scales (Brownian motion and Lévy process) [18]. The mixed local-nonlocal operator in the form (without advection)

$$L_0 := -\Delta + (-\Delta)^s, \quad s \in (0, 1),$$

has received by far great attention from different points of view. This includes existence and non-existence results [1, 4, 10, 12, 23, 25], regularity results [15, 20, 24, 29, 32], associated eigenvalue problems [11, 14, 16, 28, 30], and radial symmetry results [13].

In this paper, we consider a mixed local-nonlocal operator with the *additional advection term*  $q \cdot \nabla$ , where  $q \in L^\infty(\Omega)$  is a bounded vector field. We aim to study the existence of the principal eigenvalue and the corresponding eigenfunction in  $\Omega$  for  $L := -\Delta + (-\Delta)^s + q \cdot \nabla$  with  $s \in (0, 1/2]$ . To the best of our knowledge, the presence of an advection term has not been addressed before.

It is important to note that when the operator  $L$  does not include an advection term, that is  $L \equiv L_0$ , the operator is self-adjoint and the study of the principal eigenvalue for  $L_0$  relies on a variational characterization via the Rayleigh quotient (see [10, 11, 16]). Namely,

$$\lambda_1(\Omega) := \inf_{u \in \mathcal{X}_0^s(\Omega) \setminus \{0\}} \frac{[u]_{\mathcal{X}^s(\Omega)}}{\|u\|_{L^2(\Omega)}^2}, \quad (1.5)$$

where the space  $\mathcal{X}_0^s(\Omega)$  and the semi-norm  $[\cdot]_{\mathcal{X}^s(\Omega)}$  are defined in Section 2, below. However, in the presence of advection, the operator  $L$  is *no longer self-adjoint* and so there is no simple variational formulation for the first eigenvalue as in (1.5). We will prove the existence of such principal eigenvalue of  $L$  and the corresponding eigenfunction with the aid of the Krein-Rutman theorem (see [17]). There are many versions of the Krein-Rutman theorem in the literature. We will use that of [19, Theorem 1.2] and we recall it in Theorem A below.

Lastly, we mention that integro-differential equations arise naturally in the study of stochastic processes with jumps. They describe a biological species whose individuals diffuse either by a random walk or by a jump process according to the prescribe probabilities [32]. The generator of a Lévy process has the following

general structure

$$\mathcal{L}u := \sum_{i,j=1}^N a_{ij} D_{ij}u + \sum_{j=1}^N q_j D_j u + PV. \int_{\mathbb{R}^N} (u(x) - u(y)) K(x-y) dy, \quad (1.6)$$

where  $K$  is a measurable kernel on  $\mathbb{R}^N$  satisfying  $\int_{\mathbb{R}^N} \min\{1, |z|^2\} K(z) dz < \infty$  [16]. The first and second terms in (1.6) correspond to the diffusion and the drift respectively. The study of the operator  $\mathcal{L}$  in (1.6) with all its components (diffusion, drift and jump) appears quite intriguing. By far, there are just a few contributions in this direction. In [2], while studying the risk-sensitive control for a class of diffusion with jumps, the authors investigated the existence of the principal eigenvalue for the class of operators  $\mathcal{L}$  where the kernel is locally integrable. In [3], the authors also considered a locally integrable kernel and proved the existence of generalized principal eigenvalue in  $\mathbb{R}^N$ . We refer to [26, Chap. 3] where elliptic problems involving general second order elliptic integro-differential operator have been considered. Note that our operator  $L$  in (1.1) corresponds to  $a_{ij} = \delta_{ij}$  in (1.6). In this present work, we only consider an  $L$  where the nonlocal operator is replaced by the fractional Laplacian.

We state our first result as follows.

**Theorem 1.2.** *Let  $q \in C^{0,\alpha}(\overline{\Omega})$  for some  $0 < \alpha < 1$  and  $s \in (0, 1/2]$ . Then, there exists a principal eigenpair  $(\lambda_1(\Omega, q), \varphi_1)$  for the problem (1.1) such that*

- (a)  $\lambda_1(\Omega, q)$  is an eigenvalue of  $L$  in  $\Omega$  and the corresponding eigenfunction  $\varphi_1$  has a constant sign in  $\Omega$  and it is unique up to multiplication by a nonzero constant. Moreover, if  $s \in (0, 1/2)$ ,  $\varphi_1 \in C^{2,\alpha}(\overline{\Omega})$ . If  $s = \frac{1}{2}$ , we have  $\varphi_1 \in C^2(\Omega) \cap C^{1,\alpha}(\overline{\Omega})$ .
- (b) If  $\lambda \in \mathbb{C}$  is an eigenvalue for  $L$  in  $\Omega$ , then we have  $\lambda_1(\Omega, q) \leq |\lambda|$ .
- (c)  $\lambda_1(\Omega, q)$  can be characterized by the following min-max formula

$$\lambda_1(\Omega, q) = \max_{u \in \mathcal{V}(\Omega)} \inf_{x \in \Omega} \frac{Lu}{u}, \quad (1.7)$$

where

$$\mathcal{V}(\Omega) := \left\{ u \in C^2(\Omega) \cap C^1(\overline{\Omega}) \cap C_c(\mathbb{R}^N) : u > 0 \text{ in } \Omega \text{ and } u \equiv 0 \text{ on } \mathbb{R}^N \setminus \Omega \right\}.$$

We also prove the following Hopf Lemma for the operator  $L$ . We emphasize that the result holds for any  $s \in (0, 1)$ . We will use the following result in proving Theorem 1.2 but we will state the result in a general setting.

**Theorem 1.3 (Hopf Lemma).** *Let  $s \in (0, 1)$ . Suppose that  $\Omega \subset \mathbb{R}^N$  is a bounded  $C^2$  domain and let  $c_0 \in \mathbb{R}$ . Let  $u \in C^2(\Omega) \cap C(\mathbb{R}^N) \cap C^1(\overline{\Omega})$  such that  $u$  is bounded in  $\mathbb{R}^N$  and*

$$Lu := -\Delta u + (-\Delta)^s u + q \cdot \nabla u \geq 0 \text{ in } \Omega. \quad (1.8)$$

Let  $x_0 \in \partial\Omega$ . Assume that  $u(x) = c_0$  on  $B_{R_0}(x_0) \cap \partial\Omega$ , for some  $R_0 > 0$ , and that  $u \geq c_0$  in  $\mathbb{R}^N$ . If  $u \not\equiv c_0$  in  $\mathbb{R}^N$ , then

$$\partial_\nu u(x_0) < 0, \quad (1.9)$$

where  $\nu$  denotes the outer unit normal to  $\partial\Omega$  at  $x_0$ .

**Remark 1.4.** Observe that Theorem 1.3 holds for all  $0 < s < 1$  and that the function  $u$  is not assumed to be  $C^2(\overline{\Omega})$ . We only assume that  $u$  is differentiable at  $x_0 \in \partial\Omega$ . We will state another version of the Hopf Lemma in Theorem 3.2 below. However, the other version requires more regularity on  $u$  and its proof, which turns out to be shorter, relies on an inequality satisfied by  $(-\Delta)^s u$  in a neighbourhood of  $x_0$ . As  $C^2$  regularity, up to the boundary, is not confirmed for  $s = 1/2$ , we will see that Theorem 1.3 turns out to be more helpful, than Theorem 3.2, in proving Theorem 1.2.

The following three theorems address the regularity of solutions to the linear problem

$$\begin{cases} Lu := -\Delta u + (-\Delta)^s u + q \cdot \nabla u = f & \text{in } \Omega, \\ u = 0 & \text{on } \mathbb{R}^N \setminus \Omega. \end{cases} \quad (1.10)$$

We mention that the regularity of  $u$ —up to the boundary of  $\Omega$ , occurs for  $0 < s < 1/2$ . In the case  $s = 1/2$ , we prove an interior regularity result.

**Theorem 1.5** ( $C^{2,\alpha}$  interior regularity when  $s = 1/2$ ). *Let  $f \in C^{0,\alpha}(\overline{\Omega})$  and  $q \in C^{0,\alpha}(\overline{\Omega})$ , for some  $\alpha \in (0, 1)$ . Assume that  $s = \frac{1}{2}$ . Let  $u \in W^{2,p}(\Omega)$  be a solution of (1.10), then  $u \in C^{2,\alpha}(\overline{\Omega'})$  for every  $\Omega' \subset\subset \Omega$ . Moreover, there exists a constant  $C > 0$  such that*

$$\|u\|_{C^{2,\alpha}(\overline{\Omega'})} \leq C \|f\|_{C^{0,\alpha}(\overline{\Omega})} \quad (1.11)$$

The following theorem provides a  $W^{2,p}$  estimate for the solution to the mixed local/nonlocal problem (1.1).

**Theorem 1.6.** *Let  $\Omega \subset \mathbb{R}^N$  be an open bounded set of class  $C^{1,1}$  with  $N > 2s$  and  $s \in (0, 1/2]$ . Assume that  $f \in L^p(\Omega)$  with  $1 < p < \infty$  and  $q \in L^\infty(\Omega)$ . Then, the problem (1.10) has a unique solution  $u \in W^{2,p}(\Omega)$ . Furthermore, there exists a constant  $C := C(N, s, p, \Omega) > 0$  such that*

$$\|u\|_{W^{2,p}(\Omega)} \leq C \|f\|_{L^p(\Omega)}. \quad (1.12)$$

As a consequence of Theorem 1.6, we have the following  $C^{2,\alpha}$  regularity, up to the boundary, for (1.10) when  $0 < s < 1/2$ .

**Theorem 1.7** (Hölder regularity up to the boundary when  $s < 1/2$ ). *Let  $s \in (0, \frac{1}{2})$  and  $0 < \alpha < 1 - 2s$ . Assume that the advection term satisfies  $q \in C^{0,\alpha}(\overline{\Omega})$ . Then, there is some  $C > 0$  such that for all  $f \in C^{0,\alpha}(\overline{\Omega})$  there is some  $u \in C^{2,\alpha}(\overline{\Omega})$  that satisfies*

$$\begin{cases} Lu = f & \text{in } \Omega \\ u = 0 & \text{on } \mathbb{R}^N \setminus \Omega. \end{cases} \quad (1.13)$$

Moreover, we have

$$\|u\|_{C^{2,\alpha}(\overline{\Omega})} \leq C \|f\|_{C^{0,\alpha}(\overline{\Omega})}.$$

We briefly comment on the proof of Theorem 1.2. The main ingredient in the proof of Theorem 1.2 is the Krein-Rutman theorem, which relies on the strong maximum principle for the operator  $L$  and the  $L^p$ -theory of the problem (1.10) (see Theorem 1.6).

Indeed, with the aid of the  $L^p$ -theory of  $L_0$  developed in [32, Theorem 1.4] and for more general second order elliptic intergro-differential operators developed in [26, Theorem 3.1.23], we first prove using the method of continuity (see [27, Theorem 5.2]) that for  $f \in L^p(\Omega)$ , there exists a unique solution  $u \in W^{2,p}(\Omega)$  of problem (1.10) for any  $1 < p < \infty$ . Then, using the Sobolev (Morrey) embedding theorem we in fact have that  $u \in C^{1,\beta}(\bar{\Omega})$ , for any  $\beta \in (0, 1)$ . Moreover, the extension theorem allows us to extend  $u$  by zero to a  $C^{0,1}$ -function in  $\mathbb{R}^N$  and thanks to the regularity result of [33, Proposition 2.5], we have that  $u \in C^{2,1-2s}(\bar{\Omega})$  for  $s \in (0, 1/2)$ . This allows us to prove the uniqueness result in Lemma 4.1 and then prove Theorem 1.6 for  $s \in (0, 1/2)$ . In the case  $s = \frac{1}{2}$ , we obtain an interior  $C^{2,\alpha}$  regularity for  $u$  in Theorem 1.5, which also allows us to directly apply the strong maximum principle for  $L$  in Theorem 3.1 and thus complete the proof of Theorem 1.6 for  $s = \frac{1}{2}$ .

We point out that since our strategy of proving Theorem 1.2 relies on the  $L^p$ -theory of problem (1.10) combined with the application of the Krein-Rutmen theorem, our result holds for more general mixed local-nonlocal operators satisfying the strong maximum principle as given in (1.6) with the kernel of  $K$  satisfying

$$\frac{\kappa_1}{|x-y|^{N+2s}} \leq K(x-y) \leq \frac{\kappa_2}{|x-y|^{N+2s}}, \quad \kappa_1, \kappa_2 > 0, s \in (0, 1/2]. \quad (1.14)$$

We refer the interested reader to [26, Chap. 3] for general second order elliptic intergro-differential operators satisfying such properties. In particular, any nonlocal operator of small order will satisfy (1.14) (see [21, 22] and the references therein).

The rest of this paper is organized as follows. In Section 2, we introduce some functional spaces. In Section 3, we prove the strong maximum principle and the Hopf Lemma for  $L$ . In Section 4, we develop the  $L^p$ -theory for  $L$  and prove the existence and uniqueness of solution  $u \in W^{2,p}(\Omega)$  to problem (1.10). Section 5 is devoted to the proof of Theorem 1.2 by using most of the results in the previous sections.

## 2. Functional setting

We start this section by fixing some notation. Let  $\Omega \subset \mathbb{R}^N$  be an open bounded set. For the vector field  $q : \Omega \rightarrow \mathbb{R}^N$ , we write  $q \in L^\infty(\Omega)$  (resp.  $q \in C^{0,\alpha}(\bar{\Omega})$ ) whenever  $q_j \in L^\infty(\Omega)$  (resp.  $q_j \in C^{0,\alpha}(\bar{\Omega})$ ),  $j = 1, 2, \dots, N$ . We denote by  $C^{k,\alpha}(\bar{\Omega})$ ,  $0 < \alpha < 1$ , the Banach space of functions  $u \in C^k(\bar{\Omega})$  such that derivative of order  $k$  belong to  $C^{0,\alpha}(\bar{\Omega})$  with the norm

$$\|u\|_{C^{k,\alpha}(\bar{\Omega})} := \|u\|_{C^k(\bar{\Omega})} + \sum_{|\tau|=k} [D^\tau u]_{C^{0,\alpha}(\bar{\Omega})},$$

where

$$[u]_{C^{0,\alpha}(\bar{\Omega})} = \sup_{x,y \in \bar{\Omega}, x \neq y} \frac{|u(x) - u(y)|}{|x - y|^\alpha}$$

and  $C^{0,\alpha}(\bar{\Omega})$  is the Banach space of functions  $u \in C^0(\bar{\Omega})$  which are Hölder continuous with exponent  $\alpha$  and the norm  $\|u\|_{C^{0,\alpha}(\bar{\Omega})} = \|u\|_{L^\infty(\Omega)} + [u]_{C^{0,\alpha}(\bar{\Omega})}$ .

If  $k \in \mathbb{N}$ , as usual we set

$$W^{k,p}(\Omega) := \left\{ u \in L^p(\Omega) : D^\alpha u \text{ exists for all } \alpha \in \mathbb{N}^N, |\alpha| \leq k \text{ and } u \in L^p(\Omega) \right\}$$

for the Banach space of  $k$ -times (weakly) differentiable functions in  $L^p(\Omega)$ . Moreover, in the fractional setting, for  $s \in (0, 1)$  and  $p \in [1, \infty)$ , we set

$$W^{s,p}(\Omega) := \left\{ u \in L^p(\Omega) : \frac{u(x) - u(y)}{|x - y|^{\frac{n}{p} + s}} \in L^p(\Omega \times \Omega) \right\}.$$

The space  $W^{s,p}(\Omega)$  is a Banach space with the norm

$$\|u\|_{W^{s,p}(\Omega)} = \left( \|u\|_{L^p(\Omega)}^p + \iint_{\Omega \times \Omega} \frac{|u(x) - u(y)|^p}{|x - y|^{n+sp}} dx dy \right)^{\frac{1}{p}}.$$

We also define the space  $\mathcal{X}^s(\Omega)$  by

$$\mathcal{X}^s(\Omega) := \left\{ u \in L^2(\Omega) : u|_\Omega \in H^1(\Omega); [u]_{\mathcal{X}^s(\Omega)} < \infty \right\},$$

where the corresponding Gagliardo seminorm  $[\cdot]_{\mathcal{X}^s(\Omega)}$  is given by

$$[u]_{\mathcal{X}^s(\Omega)} := \int_\Omega |\nabla u|^2 dx + \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy.$$

Note that the space  $\mathcal{X}^s(\Omega)$  is a Hilbert space when furnished with the scalar product

$$\langle u, v \rangle_{\mathcal{X}^s(\Omega)} := \int_\Omega uv dx + \int_\Omega \nabla u \nabla v dx + \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy$$

and the corresponding norm is given by  $\|u\|_{\mathcal{X}^s(\Omega)} = \sqrt{\langle u, v \rangle_{\mathcal{X}^s(\Omega)}}$ . Define

$$\mathcal{X}_0^s(\Omega) := \left\{ u \in \mathcal{X}^s(\Omega) : u \equiv 0 \text{ on } \mathbb{R}^N \setminus \Omega \right\}.$$

Note that if  $u \in \mathcal{X}_0^s(\Omega)$  then  $u|_\Omega \in H_0^1(\Omega)$  due to the regularity assumption of  $\partial\Omega$ .

Finally, we define the space  $\mathcal{L}_s^1(\mathbb{R}^N)$  by

$$\mathcal{L}_s^1(\mathbb{R}^N) := \left\{ u : \mathbb{R}^N \rightarrow \mathbb{R}, \text{ such that } u \text{ is measurable and } \|u\|_{\mathcal{L}_s^1(\mathbb{R}^N)} < \infty \right\},$$

where

$$\|u\|_{\mathcal{L}_s^1(\mathbb{R}^N)} := \int_{\mathbb{R}^N} \frac{|u(y)|}{1 + |y|^{N+2s}} dx.$$

### 3. Hopf lemma, maximum principle, and interior regularity: proof of Theorem 1.3 and Theorem 1.5

In this section, we derive some results for the operator  $L$ . These will be important in the proof of Theorem 1.2. We start with the following result on the strong maximum principle for  $L$ .

**Theorem 3.1 (Strong Maximum Principle).** *Let  $\Omega \subset \mathbb{R}^N$  be an open bounded set and  $q \in L^\infty(\Omega)$ . Let  $s \in (0, 1)$  and  $u \in \mathcal{L}_s^1(\mathbb{R}^N)$  be a function in  $C^2(\Omega) \cap C(\mathbb{R}^N)$  that satisfies*

$$\begin{cases} Lu & \geq 0 \text{ in } \Omega \\ u & \geq 0 \text{ on } \mathbb{R}^N \setminus \Omega. \end{cases}$$

*Then  $u > 0$  in  $\Omega$  or  $u \equiv 0$  in  $\mathbb{R}^N$ .*

*Proof of Theorem 3.1.* Suppose by contradiction that  $u$  is not positive in  $\Omega$ . Since  $\Omega$  is bounded,  $\bar{\Omega}$  is compact. Since  $u$  is continuous in  $\mathbb{R}^N$  and  $u \geq 0$  in  $\mathbb{R}^N \setminus \Omega$ , there is a point  $x_0 \in \Omega$  with

$$u(x_0) = \min_{x \in \Omega} u(x) \leq 0. \quad (3.1)$$

Therefore, since  $q$  is bounded, it follows that  $q \cdot \nabla u(x_0) = 0$  and  $\Delta u(x_0) \geq 0$ . Hence, from the definition of the operator  $L$  we have that

$$(-\Delta)^s u(x_0) \geq 0.$$

Whereas by (3.1), we have that  $u(x_0) \leq u(x)$  for all  $x \in \mathbb{R}^N$ . It follows that

$$0 \leq (-\Delta)^s u(x_0) = P.V. \int_{\mathbb{R}^N} \frac{u(x_0) - u(y)}{|x_0 - y|^{N+2s}} dy = \int_{\mathbb{R}^N} \frac{u(x_0) - u(y)}{|x_0 - y|^{N+2s}} dy \leq 0.$$

Moreover, since the integrand is non-positive by assumption and (3.1), we conclude that

$$u \equiv u(x_0) \quad \text{in } \mathbb{R}^N.$$

Now, since  $u \geq 0$  in  $\mathbb{R}^N \setminus \Omega$ , it follows that  $u \equiv 0$  in  $\Omega$  and therefore  $u \equiv 0$  in  $\mathbb{R}^N$ . This leads to a contradiction and the proof is established.  $\square$

We now prove the Hopf Lemma stated in Theorem 1.3, for all  $s \in (0, 1)$ .

**Proof of Theorem 1.3.** Let be  $B_r(\bar{x})$  a ball centered at  $\bar{x} \in \Omega$  that touches  $\partial\Omega$  at  $x_0 \in \partial\Omega$ . Let  $K$  be the set defined by

$$K := B_r(\bar{x}) \cap B_{\frac{r}{2}}(x_0).$$

We introduce the auxiliary function

$$v(x) := e^{-\alpha \mathcal{K}(x)} - e^{-\alpha(1+r^2)^{\frac{1}{2}}}, \quad \text{where } \mathcal{K}(x) := (|x - \bar{x}|^2 + 1)^{\frac{1}{2}}$$

and  $\alpha$  is a positive constant to be chosen later.




FIGURE 1. The open set  $K \subset \Omega$  is the intersection of the ball centered at  $x_0$  with the ball centered at  $\bar{x}$ , which is tangent to  $\partial\Omega$  at  $x_0$ . Note that  $\bar{K} \cap \partial\Omega = \{x_0\}$ .

We have

$$v > 0 \quad \text{in } B_r(\bar{x}), \quad v = 0 \quad \text{on } \partial B_r(\bar{x}) \quad \text{and} \quad v < 0 \quad \text{on } \mathbb{R}^N \setminus B_r(\bar{x}).$$



For any  $x \in \mathbb{R}^N$ , computation shows that

$$\begin{aligned} Lv(x) &= -\Delta v(x) + (-\Delta)^s v(x) + q(x) \cdot \nabla v(x) \\ &= e^{-\alpha \mathcal{K}(x)} \left( \frac{\alpha N}{\mathcal{K}(x)} - \alpha \frac{q(x) \cdot (x - \bar{x})}{\mathcal{K}(x)} - |x - \bar{x}|^2 \left( \frac{\alpha}{\mathcal{K}^3(x)} + \frac{\alpha^2}{\mathcal{K}^2(x)} \right) \right) \\ &\quad + (-\Delta)^s v(x). \end{aligned} \quad (3.2)$$

Observe that  $1 - e^{-\rho} \leq \rho$  for all  $\rho \in \mathbb{R}$ . Therefore,

$$\begin{aligned} (-\Delta)^s v(x) &= \frac{C_{N,s}}{2} \int_{\mathbb{R}^N} \frac{2v(x) - v(x+y) - v(x-y)}{|y|^{N+2s}} dy \\ &= \frac{C_{N,s} e^{-\alpha \mathcal{K}(x)}}{2} \int_{\mathbb{R}^N} \frac{2 - e^{-\alpha(\mathcal{K}(x+y) - \mathcal{K}(x))} - e^{-\alpha(\mathcal{K}(x-y) - \mathcal{K}(x))}}{|y|^{N+2s}} dy \\ &\leq \frac{C_{N,s} \alpha e^{-\alpha \mathcal{K}(x)}}{2} \int_{\mathbb{R}^N} \frac{(\mathcal{K}(x+y) - \mathcal{K}(x)) + (\mathcal{K}(x-y) - \mathcal{K}(x))}{|y|^{N+2s}} dy \\ &= -\alpha e^{-\alpha \mathcal{K}(x)} (-\Delta)^s \mathcal{K}(x). \end{aligned} \quad (3.3)$$

Now, we compute

$$\begin{aligned} &\frac{2|(-\Delta)^s \mathcal{K}(x)|}{C_{N,s}} \\ &\leq \int_{B_1} \frac{|(\mathcal{K}(x+y) - \mathcal{K}(x)) + (\mathcal{K}(x-y) - \mathcal{K}(x))|}{|y|^{N+2s}} dy + 2\mathcal{K}(x) \int_{\mathbb{R}^N \setminus B_1} \frac{1}{|y|^{N+2s}} dy \\ &\leq \int_{B_1} \int_0^1 \frac{|\langle \nabla \mathcal{K}(x+ty) - \nabla \mathcal{K}(x-ty), y \rangle|}{|y|^{N+2s}} dt dy + \\ &\quad 2(\text{diam}(\Omega))^2 + 1 \int_{\mathbb{R}^N \setminus B_1} \frac{1}{|y|^{N+2s}} dy \\ &\leq \int_{B_1} \int_0^1 \frac{|\nabla \mathcal{K}(x+ty) - \nabla \mathcal{K}(x-ty)|}{|y|^{N+2s-1}} dt dy + \frac{2\omega_{N-1}(\text{diam}(\Omega))^2 + 1}{2s}. \end{aligned}$$

We also have

$$\begin{aligned} |\nabla \mathcal{K}(x+ty) - \nabla \mathcal{K}(x-ty)| &= \left| \frac{(x+ty)}{\mathcal{K}(x+ty)} - \frac{(x-ty)}{\mathcal{K}(x-ty)} \right| \\ &= \left| ty \left( \frac{1}{\mathcal{K}(x+ty)} + \frac{1}{\mathcal{K}(x-ty)} \right) + x \left( \frac{1}{\mathcal{K}(x+ty)} - \frac{1}{\mathcal{K}(x-ty)} \right) \right| \\ &\leq 2t|y| + \left| \frac{1}{\mathcal{K}(x+ty)} - \frac{1}{\mathcal{K}(x-ty)} \right| |x|, \text{ as } \mathcal{K} \geq 1 \\ &\leq 2t|y| \left( 1 + \left| \nabla \left( \frac{1}{\mathcal{K}(z)} \right) \right| |x| \right), \text{ for some } z \text{ in the segment } [x-ty, x+ty] \\ &\leq 2t|y| \left( 1 + \left| \frac{z - \bar{x}}{2\mathcal{K}^3(z)} \right| |x| \right) \leq 2t|y| (1 + |z - \bar{x}| |x|) \\ &\leq 2t|y| (1 + |z - x| |x| + |\bar{x}| |x|) \leq 2t|y| (1 + (3t|y| + |\bar{x}|) |x|). \end{aligned}$$

Since  $x, \bar{x} \in \Omega$  and  $\Omega$  is bounded, we have  $|x|, |\bar{x}| \leq D$ , where  $D$  is a positive constant that depends on  $\Omega$ . Also,  $|y| < 1$  for  $y \in B_1$ . Therefore,

$$\begin{aligned}
|(-\Delta)^s \mathcal{K}(x)| &\leq \frac{C_{N,s}}{2} \int_{B_1} \int_0^1 \frac{|\nabla \mathcal{K}(x+ty) - \nabla \mathcal{K}(x-ty)|}{|y|^{N+2s-1}} dt dy + \\
&\quad \frac{2\omega_{N-1}(\text{diam}(\Omega)^2 + 1)}{2s} \\
&\leq \frac{C_{N,s}}{2} \int_{B_1} 2t \int_0^1 \frac{(1 + (3t|y| + |\bar{x}|)|x|)}{|y|^{N+2s-2}} dt dy + \frac{2\omega_{N-1}(\text{diam}(\Omega)^2 + 1)}{2s} \\
&\leq \frac{C_{N,s}}{2} \int_{B_1} \frac{2(1 + (3+D)D)}{|y|^{N+2s-2}} dy + \frac{2\omega_{N-1}(\text{diam}(\Omega)^2 + 1)}{2s} \\
&\leq C_{N,s}(1 + (3+D)D) \frac{\omega_{N-1}}{2-2s} + \frac{2\omega_{N-1}(\text{diam}(\Omega)^2 + 1)}{2s} \\
&:= M.
\end{aligned} \tag{3.4}$$

We denote the constant obtained in the upper bound of  $(-\Delta)^s \mathcal{K}(x)$  by  $M$ . Thus, it follows from (3.2), (3.3) and (3.4) that, for all  $x \in \mathbb{R}^N$ ,

$$L_V(x) \leq e^{-\alpha \mathcal{K}(x)} \left( \frac{\alpha N}{\mathcal{K}(x)} - \alpha \frac{q(x) \cdot (x - \bar{x})}{\mathcal{K}(x)} - |x - \bar{x}|^2 \left( \frac{\alpha}{\mathcal{K}^3(x)} + \frac{\alpha^2}{\mathcal{K}^2(x)} \right) - \alpha M \right).$$

Now, if  $x \in K = B_r(\bar{x}) \cap B_{\frac{r}{2}}(x_0)$ , we have  $|x - \bar{x}| \geq \frac{r}{2}$  and hence

$$L_V(x) \leq e^{-\alpha \mathcal{K}(x)} \left( \frac{\alpha N}{\mathcal{K}(x)} + \alpha \frac{\|q\|_{L^\infty} |x - \bar{x}|}{\mathcal{K}(x)} - \frac{r^2}{4} \left( \frac{\alpha}{\mathcal{K}^3(x)} + \frac{\alpha^2}{\mathcal{K}^2(x)} \right) - \alpha M \right).$$

Since  $\mathcal{K}(x) \geq 1$ , for all  $x$ , we can choose  $\alpha$  large enough so that

$$L_V < 0 \quad \text{in} \quad B_{\frac{r}{2}}(x_0) \cap \Omega.$$

Consider the function  $w := -u + \varepsilon v + c_0$ , where  $\varepsilon$  is a positive constant to be chosen later. Since  $u \geq c_0$  in  $\mathbb{R}^N$ , the maximum principle in Theorem 3.1, applied to  $u - c_0$ , yields that  $u > c_0$  in  $\Omega$  (as  $u \not\equiv c_0$  in  $\mathbb{R}^N$ ). As the set  $\bar{B}_r(\bar{x}) \setminus K$  is compact, the minimum of  $u$  over  $\bar{B}_r(\bar{x}) \setminus K$  is attained. So this minimum will be strictly greater than  $c_0$ . That is,

$$\min_{\bar{B}_r(\bar{x}) \setminus K} u(x) > c_0.$$

Thus, we can find a constant  $\delta > 0$  such that

$$u \geq c_0 + \delta \quad \text{in} \quad \bar{B}_r(\bar{x}) \setminus K.$$

Then, for  $x \in \bar{B}_r(\bar{x}) \setminus K$

$$w(x) \leq -\delta + \varepsilon v = -\delta + \varepsilon (e^{-\alpha \mathcal{K}(x)} - e^{-\alpha(1+r^2)^{\frac{1}{2}}}) \leq -\delta + \varepsilon (1 - e^{-\alpha(1+r^2)^{\frac{1}{2}}}).$$

Chosen  $\varepsilon$  sufficiently small, say

$$\varepsilon < \frac{\delta}{1 - e^{-\alpha(1+r^2)^{\frac{1}{2}}}},$$

we have

$$w < 0 \quad \text{in} \quad \overline{B_r(\bar{x})} \setminus K.$$

Since  $u \geq c_0$  and  $v < 0$  in  $\mathbb{R}^N \setminus B_r(\bar{x})$ , we have that  $w < 0$  on  $\mathbb{R}^N \setminus K$ . We also have

$$w < 0 \quad \text{in} \quad \partial K \setminus \{x_0\} \quad \text{and} \quad w(x_0) = 0.$$

Moreover,

$$Lw = L(-u + \varepsilon v + c_0) \leq \varepsilon Lv < 0 \quad \text{in} \quad K.$$

From the maximum principle, applied to  $w$ , we obtain  $w < 0$  in  $K$  (since  $w \not\equiv 0$  in  $\mathbb{R}^N$ ). As  $w(x_0) = 0$ , it follows that the maximum of  $w$  over  $\overline{K}$  is attained at  $x_0$ . Therefore, the normal derivative  $\partial_\nu w$  satisfies

$$\partial_\nu w(x_0) = -\partial_\nu u(x_0) + \varepsilon \partial_\nu v(x_0) \geq 0. \quad (3.5)$$

We compute now the normal derivative of  $v$  over  $\partial B_r(\bar{x})$ . We have

$$\nabla v(x) = -\alpha \frac{(x - \bar{x})}{\mathcal{K}(x)} e^{-\alpha \mathcal{K}(x)}, \quad \text{for all } x \in \Omega.$$

Thus, for  $x \in \partial B_r(\bar{x})$ , we have

$$\partial_\nu v(x) = (x - \bar{x}) \cdot \nabla v(x) = -\alpha \frac{|x - \bar{x}|}{\mathcal{K}(x)} e^{-\alpha \mathcal{K}(x)} < 0.$$

In particular,  $\partial_\nu v(x_0) < 0$  and it follows from (3.5) that

$$\partial_\nu u(x_0) \leq \varepsilon \partial_\nu v(x_0) < 0.$$

This completes the proof.  $\square$

We state and prove another version of the Hopf Lemma in the next theorem. We refer the reader to Remark 1.4 above for more details on the difference between Theorem 3.2 and Theorem 1.3.

**Theorem 3.2 (Hopf Lemma).** *Suppose that  $\Omega \subset \mathbb{R}^N$  is a bounded  $C^2$  domain and let  $c_0 \in \mathbb{R}$ . Let  $u \in C^2(\overline{\Omega}) \cap C(\mathbb{R}^N)$  such that  $u$  is bounded in  $\mathbb{R}^N$  and*

$$Lu \geq 0 \quad \text{in} \quad \Omega. \quad (3.6)$$

*Let  $x_0 \in \partial\Omega$ . Assume that  $u(x) = c_0$  on  $B_{R_0}(x_0) \cap \partial\Omega$ , for some  $R_0 > 0$ , and that  $u \geq c_0$  in  $\mathbb{R}^N$ . If  $u \not\equiv c_0$  in  $\mathbb{R}^N$ , then*

$$\partial_\nu u(x_0) < 0, \quad (3.7)$$

*where  $\nu$  denotes the outer unit normal to  $\partial\Omega$  at  $x_0$ .*

**Proof of Theorem 3.2.** The proof mostly relies on the fact that

$$(-\Delta)^s u \leq 0 \quad \text{in} \quad B_\rho(x_0) \cap \Omega, \quad \text{for some } 0 < \rho < R_0, \quad (3.8)$$

and the Hopf lemma for elliptic operators. The proof of inequality (3.8) is done in details in [11, inequality (2.9) in the proof of Theorem 2.9] and we will omit it here.

We note that if there is a point  $y \in B_\rho(x_0)$  such that  $u(y) = c_0$ , we apply the maximum principle in Theorem 3.1 to (3.6) (knowing that  $u \geq c_0$  in  $\mathbb{R}^N$ ) and conclude that  $u \equiv c_0$  in  $\mathbb{R}^N$ , which is a contradiction.

Now, we combine (3.6) and (3.8) to deduce that there exists  $\rho > 0$  such that

$$0 \leq Lu \leq -\Delta u + q \cdot \nabla u \quad \text{in} \quad B_\rho(x_0) \cap \Omega, \quad (3.9)$$

and  $u \geq c_0$  in  $\mathbb{R}^N$ . The elliptic maximum principle (see [27, Lemma 3.4], for e.g.) implies that either  $u \equiv c_0$  in  $B_\rho(x_0) \cap \Omega$  (this cannot happen because of the note above) or  $u > c_0$  in  $B_\rho(x_0) \cap \Omega$ . Moreover, the Hopf Lemma for elliptic operators (here, we have  $-\Delta + q \cdot \nabla$ ) implies that  $\partial_\nu u(x) < 0$  for all  $x \in B_\rho(x_0) \cap \partial\Omega$ , which is part of  $\partial(B_\rho(x_0) \cap \Omega)$ . In particular, we have  $\partial_\nu u(x_0) < 0$  and this completes the proof.  $\square$

Next, we give the proof of the interior regularity when  $s = 1/2$ . We mention that, for  $0 < s < 1/2$ , we will have regularity  $C^{2,\alpha}(\bar{\Omega})$  up to the boundary. The latter is done in Theorem 1.7.

**Proof of Theorem 1.5.** Let  $f \in C^{0,\alpha}(\bar{\Omega})$ . Let  $\Omega'$  and  $\Omega_1$  be two open subsets of  $\Omega \subset \mathbb{R}^N$  such that

$$\Omega' \subset\subset \Omega_1 \subset\subset \Omega.$$

Define the cut-off function  $\eta \in C_c^\infty(\Omega_1)$  as

$$\eta(x) = 1 \quad \text{if } x \in \Omega' \quad \text{and} \quad \eta(x) = 0 \quad \text{if } x \in \mathbb{R}^N \setminus \Omega_1, \quad (3.10)$$

such that  $0 \leq \eta(x) \leq 1$  for all  $x \in \mathbb{R}^N$  and there exists  $C_1, C_2 > 0$  such that

$$|D\eta| < \frac{C_1}{\text{dist}(\Omega', \partial\Omega)} \quad \text{and} \quad |D^2\eta| < \frac{C_2}{(\text{dist}(\Omega', \partial\Omega))^2}.$$

We set

$$v := \eta u \quad \text{and} \quad w := (1 - \eta)u$$

Since  $u \in W^{2,p}(\Omega)$ , it holds that  $u \in C^{1,\alpha}(\bar{\Omega})$  applying the Bootstrap argument as in Lemma 4.1 with  $\beta = \alpha$ . Then, we compute

$$\begin{aligned} -\Delta v &= -\eta((-\Delta)^{\frac{1}{2}}u + q \cdot \nabla u - f) - 2\nabla u \cdot \nabla \eta - u\Delta\eta \\ &:= \tilde{f}. \end{aligned}$$

We note that all elements of  $\tilde{f}$  are supported in  $\Omega_1$ . We need to show that  $\tilde{f} \in C^{0,\alpha}(\mathbb{R}^N)$ . To do so, we only need to show that  $\eta(-\Delta)^{\frac{1}{2}}u \in C^{0,\alpha}(\mathbb{R}^N)$  since other terms follow easily. We write

$$\eta(-\Delta)^s u = (-\Delta)^s v - u(-\Delta)^s \eta + I(u, \eta)$$

where

$$I(u, \eta)(x) := \int_{\mathbb{R}^N} \frac{(u(x) - u(x+z))(\eta(x) - \eta(x+z))}{|z|^{N+1}} dz.$$

Since  $\text{supp } v \subset \Omega$ , we have

$$\|v\|_{C^{1,\alpha}(\mathbb{R}^N)} = \|v\|_{C^{1,\alpha}(\bar{\Omega})} \leq C\|u\|_{C^{1,\alpha}(\bar{\Omega})}. \quad (3.11)$$

Then, we use the regularity result of [33, Proposition 2.7] (with  $s = \frac{1}{2}$  and  $l = 0$ ) to get  $(-\Delta)^{\frac{1}{2}}v \in C^{0,\alpha}(\mathbb{R}^N)$  and

$$\|(-\Delta)^{\frac{1}{2}}v\|_{C^{0,\alpha}(\mathbb{R}^N)} \leq C\|v\|_{C^{1,\alpha}(\mathbb{R}^N)} \leq C\|u\|_{C^{1,\alpha}(\bar{\Omega})}.$$

Also,  $\|u(-\Delta)^{\frac{1}{2}}\eta\|_{C^{0,\alpha}(\mathbb{R}^N)} \leq C\|(-\Delta)^{\frac{1}{2}}\eta\|_{L^\infty(\mathbb{R}^N)}\|u\|_{C^{0,\alpha}(\mathbb{R}^N)} \leq C\|u\|_{C^{1,\alpha}(\bar{\Omega})}$ .

We now show that  $I(u, \eta) \in C^{0,\alpha}(\mathbb{R}^N)$ . Note that since  $\Omega_1$  is bounded, we let  $B_{R_0}$  be a ball centred at zero with radius  $R_0 > 0$  and containing  $\Omega_1$ . We set

$R := R_0 + |x| + 1$  for any fixed  $x \in \Omega_1$ . Observe that if  $|z| \geq R$ , then  $|x + z| \geq |z| - |x| \geq R - |x| = R_0 + 1 > R_0$ . Therefore,  $\eta(x + z) \equiv 0$  on  $\mathbb{R}^N \setminus B_R$ . Next, for  $x_1, x_2 \in \Omega_1$ , we write

$$|I(u, \eta)(x_1) - I(u, \eta)(x_2)| \leq |I_1| + |I_2|,$$

where

$$I_1 := \int_{B_R} \frac{\sum_{k=1}^2 (-1)^{k-1} [(u(x_k) - u(x_k + z))(\eta(x_k) - \eta(x_k + z))]}{|z|^{N+1}} dz$$

and

$$I_2 := \int_{\mathbb{R}^N \setminus B_R} \frac{\sum_{k=1}^2 (-1)^{k-1} [(u(x_k) - u(x_k + z))\eta(x_k)]}{|z|^{N+1}} dz$$

We estimate the integrand of  $I_1$  using the fundamental theorem of calculus as follows,

$$\begin{aligned} & \left| \sum_{k=1}^2 (-1)^{k-1} [(u(x_k) - u(x_k + z))(\eta(x_k) - \eta(x_k + z))] \right| \\ &= \left| \int_0^1 \langle \nabla u(x_1 + tz) - \nabla u(x_2 + tz), z \rangle dt (\eta(x_1) - \eta(x_1 + z)) \right. \\ & \quad \left. + \int_0^1 \langle \nabla \eta(x_1 + \tau z) - \nabla \eta(x_2 + \tau z), z \rangle d\tau (u(x_2) - u(x_2 + z)) \right| \\ &\leq C|z|^2 \int_0^1 |\nabla u(x_1 + tz) - \nabla u(x_2 + tz)| dt \\ & \quad + C\|u\|_{C^1(\bar{\Omega})} |z|^2 \int_0^1 |\nabla \eta(x_1 + \tau z) - \nabla \eta(x_2 + \tau z)| d\tau \\ &\leq C(\|u\|_{C^{1,\alpha}(\bar{\Omega})} + \|u\|_{C^1(\bar{\Omega})}) |z|^2 |x_1 - x_2|^\alpha \leq C\|u\|_{C^{1,\alpha}(\bar{\Omega})} |z|^2 |x_1 - x_2|^\alpha. \end{aligned}$$

Consequently,

$$|I_1| \leq C\|u\|_{C^{1,\alpha}(\bar{\Omega})} |x_1 - x_2|^\alpha \int_{B_R} \frac{|z|^2}{|z|^{N+1}} dz \leq C\|u\|_{C^{1,\alpha}(\bar{\Omega})} |x_1 - x_2|^\alpha.$$

We now estimate  $I_2$ . Observe first that

$$\begin{aligned} & \sum_{k=1}^2 (-1)^{k-1} [(u(x_k) - u(x_k + z))\eta(x_k)] \\ &= [v(x_1) - v(x_2)] - [u(x_1 + z) - u(x_2 + z)]\eta(x_1) + [\eta(x_1) - \eta(x_2)]u(x_2 + z). \end{aligned}$$

Therefore, we have

$$\begin{aligned} |I_2| &\leq C(\|v\|_{C^1(\bar{\Omega})} + \|u\|_{C^1(\bar{\Omega})} + \|u\|_{L^\infty(\Omega)}) |x_1 - x_2| \int_{\mathbb{R}^N \setminus B_R} \frac{dz}{|z|^{N+1}} \\ &\leq C\|u\|_{C^{1,\alpha}(\bar{\Omega})} |x_1 - x_2|. \end{aligned}$$

Since it not difficult to see that  $I(u, \eta) \in L^\infty(\mathbb{R}^N)$ , we conclude that  $I(u, \eta) \in C^{0,\alpha}(\mathbb{R}^N)$ . We then consider the equation

$$-\Delta v = \tilde{f} \quad \text{in } \mathbb{R}^N.$$

Since  $\tilde{f} \in C^{0,\alpha}(\mathbb{R}^N)$ , we can apply the regularity theory for classical elliptic PDEs to see that

$$v \in C^{2,\alpha}(\overline{\Omega'}).$$

Since  $v = u$  in  $\Omega'$  and since  $\Omega'$  was arbitrary, the proof of Theorem 1.5 is complete.  $\square$

#### 4. $L^p$ theory and regularity up to the boundary: proof of Theorem 1.6 and Theorem 1.7

This section is dedicated to the  $L^p$ -theory of the operator  $L$  and to the  $C^{2,\alpha}(\overline{\Omega})$  regularity. We will first prove the following problem

$$\begin{cases} Lu := -\Delta u + (-\Delta)^s u + q \cdot \nabla u = f & \text{in } \Omega, \\ u = 0 & \text{on } \mathbb{R}^N \setminus \Omega. \end{cases} \quad (4.1)$$

has a unique solution  $u \in W^{2,p}(\Omega)$  (see Theorem 1.6). This extends the  $W^{2,p}$  estimate done in [32] for  $L_0$  to the case where an advection term is present in the equation. We will need the following lemma.

**Lemma 4.1.** *Assume that  $0 < s < 1/2$ ,  $1 < p < \infty$  and that the advection term satisfies  $q \in C^{0,\alpha}(\overline{\Omega})$  for some  $0 < \alpha < 1$ . Let  $u \in W^{2,p}(\Omega)$  be a solution of*

$$\begin{cases} Lu = 0 & \text{in } \Omega \\ u = 0 & \text{on } \mathbb{R}^N \setminus \Omega. \end{cases} \quad (4.2)$$

Then,  $u = 0$ .

**Proof of Lemma 4.1.** We need to prove the solution is sufficiently regular first (so that we use the strong maximum principle, stated in Theorem 3.1).

First, consider the case of  $p \geq N$  and then note that we have  $L_0 u = -q \cdot \nabla u$ . The right hand side is in  $L^T(\Omega)$  for all  $T < \infty$  and we can then apply the  $L^p$  theory for the operator  $L_0$  in [32] to see that  $u \in W^{2,T}(\Omega)$  for all  $T < \infty$  and hence  $u \in C^{1,\beta}(\overline{\Omega})$  for all  $0 < \beta < 1$ . Now, since we have  $u \in C^{1,\beta}(\overline{\Omega})$ , and thanks to the regularity assumption on the boundary of  $\Omega$ , we can extend  $u$  by zero outside  $\Omega$  and still denote by  $u$  (see [27, Lemma 6.37]) and get that the extension is a  $C^{0,1}(\mathbb{R}^N)$  function. We can then apply the regularity result of [33, Proposition 2.5] to see that

$$g := (-\Delta)^s u \in C^{0,1-2s}(\mathbb{R}^N).$$

Now, we can write the equation as  $-\Delta u = -g - q \cdot \nabla u$  in  $\Omega$  with  $u = 0$  on  $\partial\Omega$ . Hence, as  $q \in C^{0,\alpha}(\overline{\Omega})$ , the right hand side  $-g - q \cdot \nabla u$  is a Hölder function. Thus,  $u \in C^{2,1-2s}(\overline{\Omega})$  from the classical theory of elliptic PDEs. We can then apply the maximum principle to get that  $u \equiv 0$  in  $\mathbb{R}^N$ .

Second, we suppose  $1 < p < N$  and set  $t_1 := \frac{Np}{N-p}$ . Then we have

$$u \in W^{2,p}(\Omega) \subset W^{1,t_1}(\Omega)$$

by the Sobolev embedding theorem. Hence, as  $L_0 u = -q \cdot \nabla u$ , the  $L^p$  theory for the operator  $L_0$  in [32] yields that  $u \in W^{2,t_1}(\Omega)$ . Again, the Sobolev embedding theorem implies that  $u \in W^{1,t_2}(\Omega)$ , where  $t_2 := \frac{Nt_1}{N-t_1} > t_1$ . If  $t_2 < N$ , we can do this a finite

number of times until we get  $u \in W^{2,t}(\Omega)$  for some  $t > N$ . At this stage, we become in the setting of the first case. The proof of Lemma 4.1 is complete.  $\square$

We now have all what is needed to prove Theorem 1.6, which we do as follows.

**Proof of Theorem 1.6.** We apply the method of continuity. To ease the notation, we define the operator

$$L_0 u := -\Delta u + (-\Delta)^s u,$$

and for  $\lambda \in \mathbb{R}$ , we consider the family of operators

$$L_\lambda u \equiv (1 - \lambda)L_0 u + \lambda Lu = L_0 + \lambda q \cdot \nabla u.$$

Next, for  $u \in W^{2,p}(\Omega)$  and  $\lambda \in \mathbb{R}$ , consider the problem

$$\begin{cases} L_\lambda u = f & \text{in } \Omega \\ u = 0 & \text{on } \mathbb{R}^N \setminus \Omega. \end{cases} \quad (4.3)$$

Let  $\mathcal{A}$  be the set given by

$$\mathcal{A} := \left\{ \lambda \in [0, 1] : \exists C_\lambda > 0 \text{ such that for all } f \in L^p(\Omega), (4.3) \text{ has a solution } u \text{ such that } \|u\|_{W^{2,p}(\Omega)} \leq C_\lambda \|f\|_{L^p(\Omega)} \right\}. \quad (4.4)$$

In (4.4), we take the constant  $C_\lambda$  to be the smallest constant such that  $\|u\|_{W^{2,p}(\Omega)} \leq C_\lambda \|f\|_{L^p(\Omega)}$  holds. In other words, if  $C_\lambda > \varepsilon > 0$  then there exists  $f_\varepsilon \in L^p(\Omega)$  such that

$$\|u\|_{W^{2,p}(\Omega)} \geq (C_\lambda - \varepsilon) \|f_\varepsilon\|_{L^p(\Omega)}. \quad (4.5)$$

Note that  $\mathcal{A}$  is not empty since we have that  $0 \in \mathcal{A}$  by [32, Theorem 1.4]. Therefore, we only need to show that  $1 \in \mathcal{A}$ . To do that, it suffices to prove that  $\mathcal{A}$  is both open and closed in  $[0, 1]$ . More precisely, it suffices to prove that for any fixed  $\lambda_0 \in \mathcal{A}$  and  $f \in L^p(\Omega)$ , there is an  $\varepsilon > 0$  such that  $\lambda_0 \pm \varepsilon \in \mathcal{A}$  and that any bounded sequence  $\{\lambda_n\}_n \subset \mathcal{A}$  has a convergence subsequence.

**$\mathcal{A}$  is open.** We fix  $\lambda_0 \in \mathcal{A}$ . We look for a solution  $u \in W^{2,p}(\Omega)$  of problem (4.3) in the form  $u = v_0 + \Phi$ , where  $v_0$  solves (4.3) with  $\lambda = \lambda_0$ . For  $\varepsilon \in \mathbb{R}$ , we introduce the operator  $\mathcal{N}_\varepsilon$  given by

$$\Psi = \mathcal{N}_\varepsilon(\Phi)$$

where  $\Psi$  solves the equation

$$L_{\lambda_0} \Psi = \pm \varepsilon (q \cdot \nabla v_0 + q \cdot \nabla \Phi).$$

The operator  $\mathcal{N}_\varepsilon$  maps  $W^{2,p}(\Omega)$  into itself. We claim that if  $\varepsilon$  is chosen appropriately, then  $\mathcal{N}_\varepsilon$  is a contraction in  $W^{2,p}(\Omega)$ . Indeed, since  $\lambda_0 \in \mathcal{A}$  there exists a constant  $C_{\lambda_0} > 0$  such that

$$\|\mathcal{N}_\varepsilon(\Phi)\|_{W^{2,p}(\Omega)} = \|\Psi\|_{W^{2,p}(\Omega)} \leq C_{\lambda_0} \|\pm \varepsilon (q \cdot \nabla v_0 + q \cdot \nabla \Phi)\|_{L^p(\Omega)}.$$

Now, let  $\Phi_1$  and  $\Phi_2$  be taken in  $W^{2,p}(\Omega)$ . Then

$$\begin{aligned} \|\mathcal{N}_\varepsilon(\Phi_1) - \mathcal{N}_\varepsilon(\Phi_2)\|_{W^{2,p}(\Omega)} &= \|\Psi_1 - \Psi_2\|_{W^{2,p}(\Omega)} \\ &\leq |\varepsilon| C_{\lambda_0} \|q\|_{L^\infty(\Omega)} \|\nabla(\Phi_1 - \Phi_2)\|_{L^p(\Omega)} \\ &\leq |\varepsilon| C_{\lambda_0} C \|q\|_{L^\infty(\Omega)} \|\Phi_1 - \Phi_2\|_{W^{2,p}(\Omega)}, \end{aligned}$$

where the constant  $C$  and  $C_{\lambda_0}$  are independent of  $\varepsilon$  and  $\Phi$ . Taking  $\varepsilon$  such that

$$|\varepsilon| \leq \frac{1}{2C_{\lambda_0}C\|q\|_{L^\infty(\Omega)}},$$

we get that  $\mathcal{N}_\varepsilon$  is a contraction mapping. By the fixed point theorem, for each such  $\varepsilon$  there exists a fixed point  $\Phi$  such that  $\mathcal{N}_\varepsilon(\Phi) = \Phi$ . We just showed that the equation  $L_{\lambda_0 \pm \varepsilon}u = f$  has a solution  $u \in W^{2,p}(\Omega)$ . Moreover, from the definition of  $\mathcal{N}_\varepsilon$  and the choice of  $\varepsilon$  above, it follows that

$$\begin{aligned} \|\Phi\|_{W^{2,p}(\Omega)} &= \|\mathcal{N}_\varepsilon(\Phi)\|_{W^{2,p}(\Omega)} \leq |\varepsilon|C_{\lambda_0}C\|q\|_{L^\infty(\Omega)} \left( \|\Phi\|_{W^{2,p}(\Omega)} + \|v_0\|_{W^{2,p}(\Omega)} \right) \\ &\leq \frac{1}{2} \left( \|\Phi\|_{W^{2,p}(\Omega)} + \|v_0\|_{W^{2,p}(\Omega)} \right) \end{aligned}$$

so that  $\|\Phi\|_{W^{2,p}(\Omega)} \leq 2\|v_0\|_{W^{2,p}(\Omega)}$  and the norm of  $u$  in  $W^{2,p}(\Omega)$  becomes

$$\begin{aligned} \|u\|_{W^{2,p}(\Omega)} &\leq (\|v_0\|_{W^{2,p}(\Omega)} + \|\Phi\|_{W^{2,p}(\Omega)}) \\ &\leq C_2\|v_0\|_{W^{2,p}(\Omega)} \\ &\leq C\|f\|_{L^p(\Omega)}. \end{aligned}$$

Therefore,  $\lambda_0 \pm \varepsilon \in \mathcal{A}$ .

**$\mathcal{A}$  is closed.** In order to complete the proof of theorem, we show that  $\mathcal{A}$  is closed. Let then  $\{\lambda_n\}_n \subset \mathcal{A}$  be a sequence such that  $\lambda_n \rightarrow \lambda_0 \in \mathbb{R}$  as  $n \rightarrow \infty$ . We claim that  $\lambda_0 \in \mathcal{A}$ . Let  $f \in L^p(\Omega)$ . Since  $\lambda_n \in \mathcal{A}$  for any  $n$ , there exists  $u_n$  that satisfies (4.3) with  $\lambda_n$  in place of  $\lambda$  and

$$\|u_n\|_{W^{2,p}(\Omega)} \leq C_n\|f\|_{L^p(\Omega)}$$

where  $C_n := C_{\lambda_n}$ . Therefore, if  $\{C_n\}_n$  is bounded in  $n$  then, the sequence  $\{u_n\}_n$  is also uniformly bounded in  $W^{2,p}(\Omega)$  so that passing to a subsequence, we have

$$u_n \rightharpoonup u \text{ weakly in } W^{2,p}(\Omega) \quad \text{and} \quad u_n \rightarrow u \text{ strongly in } W^{1,p}(\Omega)$$

with  $u$  satisfying the problem (4.3) with  $\lambda_0$  in place of  $\lambda$  and

$$\|u\|_{W^{2,p}(\Omega)} \leq \liminf_{n \rightarrow \infty} \|u_n\|_{W^{2,p}(\Omega)} \leq C\|f\|_{L^p(\Omega)}.$$

This shows that  $\lambda_0 \in \mathcal{A}$  and ends the proof in the case where  $\{C_n\}_n$  is bounded.

Indeed, we will show next that the only possible case. Assume to the contrary that  $\{C_n\}_n$  is unbounded. Then, passing to a subsequence, we may have that  $C_n \rightarrow \infty$  as  $n \rightarrow \infty$ . Thus, there exists a sequence  $\{f_n\}_n$  such that for large  $n$  we have

$$\|u_n\|_{W^{2,p}(\Omega)} \geq (C_n - 1)\|f_n\|_{L^p(\Omega)}.$$

Note that the above inequality holds since from (4.5) in which the constant  $C_n := C_{\lambda_n}$  is the smallest constant such that  $\|u\|_{W^{2,p}(\Omega)} \leq C_n\|f\|_{L^p(\Omega)}$  holds. Let

$$t_n := \|u_n\|_{W^{2,p}(\Omega)}, \quad v_n := \frac{u_n}{t_n}, \quad \tilde{f}_n := \frac{f_n}{t_n}.$$

Then  $\|v_n\|_{W^{2,p}(\Omega)} = 1$ ,  $\tilde{f}_n \rightarrow 0$  in  $L^p(\Omega)$  as  $n \rightarrow \infty$  and  $v_n$  satisfies the equation

$$L_0 v_n = -\lambda_n q \cdot \nabla v_n + \tilde{f}_n \quad \text{in } \Omega, \quad v_n = 0 \quad \text{on } \mathbb{R}^N \setminus \Omega. \quad (4.6)$$



It follows from [32, Theorem 1.4] that

$$\begin{aligned} \|v_n\|_{W^{2,p}(\Omega)} &\leq C \left( \|\tilde{f}_n - \lambda_n q \cdot \nabla v_n\|_{L^p(\Omega)} + \|v_n\|_{L^p(\Omega)} \right) \\ &\leq C \left( \|\tilde{f}_n\|_{L^p(\Omega)} + \|q\|_{L^\infty(\Omega)} \|v_n\|_{W^{1,p}(\Omega)} + \|v_n\|_{L^p(\Omega)} \right) \\ &\leq C (K + \|q\|_{L^\infty(\Omega)} + 1). \end{aligned}$$

This shows that the sequence  $\{v_n\}_n$  is uniformly bounded in  $W^{2,p}(\Omega)$  so that the Banach-Alaoglu Theorem implies the existence of a subsequence, which we still label as  $\{v_n\}_n$ , that converges weakly to some  $v_0 \in W^{2,p}(\Omega)$  and strongly in  $W^{1,p}(\Omega)$  thanks to the compactness of the Sobolev embedding. Hence  $\|v_0\|_{W^{2,p}(\Omega)} \leq 1$ . We now consider two possibilities.

If  $v_0 = 0$ , this will contradict the normalization  $\|v_n\|_{W^{2,p}(\Omega)} = 1$ : indeed, if  $v_0 = 0$ , then

$$\lambda_n q \cdot \nabla v_n \rightarrow 0 \quad \text{strongly in } L^p(\Omega)$$

thanks to the compactness of the Sobolev embedding. Hence, we can use the  $L^p$  theory for  $L_0$  and (4.6) to see that  $v_n \rightarrow 0$  in  $W^{2,p}(\Omega)$ , which contradicts the normalization of  $v_n$  in  $W^{2,p}(\Omega)$ .

The other possibility is that  $v_0 \neq 0$ . In such case, we can pass to the limit in (4.6) to get

$$L_{\lambda_0} v_0 = 0 \quad \text{in } \Omega, \quad v_0 = 0 \quad \text{on } \mathbb{R}^N \setminus \Omega. \quad (4.7)$$

We will discuss the consequences of (4.7) according to the values of  $s$ .

If  $0 < s < 1/2$ , then as  $v_0 \in W^{2,p}(\Omega)$  for  $1 < p < \infty$ , it follows from Lemma 4.1 that  $v_0 \equiv 0$ , which yields again a contradiction.

If  $s = 1/2$ , then we can apply the interior regularity result of Theorem 1.5 to conclude that  $v_0 \in C^2(\Omega)$ . The Sobolev embedding also tells us that  $v_0 \in C^{1,\alpha}(\bar{\Omega})$ . Thus,  $v_0 \in C^2(\Omega) \cap C(\mathbb{R}^N)$ . The maximum principle in Theorem 3.1 implies that  $v_0 \equiv 0$ , which is again a contradiction.

Therefore, for  $s \in (0, 1/2]$ , the sequence  $\{u_n\}_n$  remains uniformly bounded in  $W^{2,p}(\Omega)$  and hence  $\{C_n\}_n$  is bounded. This is a contradiction and this completes the proof of Theorem 1.6.  $\square$

**Proof of Theorem 1.7.** We use the  $L^p$  theory in Theorem 1.6. Indeed, since  $f \in C^{0,\alpha}(\bar{\Omega})$  and  $\Omega$  is bounded,  $f \in L^p(\Omega)$  for any  $p < \infty$ . It follows then from Theorem 1.6 that  $u \in C^{1,\beta}(\bar{\Omega})$  for any  $\beta \in (0, 1)$  (we choose all  $p > N$ ). Take in particular  $\beta = \alpha$ . From the uniqueness of the solution, we get that

$$\|u\|_{C^{1,\alpha}(\bar{\Omega})} \lesssim \|f\|_{L^p(\Omega)} \leq C \|f\|_{C^{0,\alpha}(\bar{\Omega})}. \quad (4.8)$$

As before, we can extend  $u$  by zero outside  $\Omega$  by a  $C^{0,1}$  function in  $\mathbb{R}^N$  and still denote by  $u$ . We apply again the regularity result of [33, Proposition 2.5] to see that  $g := (-\Delta)^s u \in C^{0,1-2s}(\mathbb{R}^N)$  with a control on the  $C^{0,\alpha}$ -norm of  $g$  as follows

$$\|g\|_{C^{0,\alpha}(\mathbb{R}^N)} \leq C \|u\|_{C^{1,\alpha}(\bar{\Omega})} \leq C \|f\|_{C^{1,\alpha}(\bar{\Omega})}. \quad (4.9)$$

Next, since  $q$  is Hölder over  $\bar{\Omega}$ , we have that  $q \cdot \nabla u \in C^{0,\alpha}(\bar{\Omega})$ . Now, we write the equation as  $-\Delta u = -g - q \cdot \nabla u$  in  $\Omega$  with  $u = 0$  on  $\partial\Omega$ . Hence, as  $\Omega$  has

smooth boundary,  $u \in C^{2,\alpha}(\overline{\Omega})$  from the classical theory of elliptic PDEs. Moreover, combining (4.8) and (4.9), there exists a constant  $C > 0$  such that

$$\|u\|_{C^{2,\alpha}(\overline{\Omega})} \leq C\|f\|_{C^{0,\alpha}(\overline{\Omega})}.$$

The proof of Theorem 1.7 is complete.  $\square$

## 5. Proof of Theorem 1.2

In this section we complete the proof of Theorem 1.2. We recall the following statement of Krein-Rutman Theorem from [19, Theorem 1.2] as we will use it in the proof of Theorem 1.2.

**Theorem A (Krein-Rutman Theorem, [19]).** *Let  $X$  be a Banach space,  $K \subset X$  a solid cone,  $T : X \rightarrow X$  a compact linear operator which satisfies  $T(K \setminus \{0\}) \subset K^\circ$ . Then,*

- (i)  $r(T) > 0$  and  $r(T)$  is a simple eigenvalue with an eigenfunction  $v \in K^\circ$ ; there is no other eigenvalue with a positive eigenfunction.
- (ii)  $|\mu| < r(T)$  for all eigenvalues  $\mu \neq r(T)$ .

We now give the proof of Theorem 1.2.

*Proof of Theorem 1.2.* Define the space

$$X := \{u \in C^{0,1}(\overline{\Omega}) : u = 0 \text{ on } \partial\Omega \text{ and } u = 0 \text{ in } \mathbb{R}^N \setminus \Omega\}$$

and the cone

$$K := \{u \in X : u \geq 0 \text{ in } \overline{\Omega}\}.$$

We will denote the interior of  $K$  by  $K^\circ$ . Indeed,

$$K^\circ = \{u \in X : \text{there is some } \varepsilon > 0 \text{ such that } u(x) \geq \varepsilon \text{dist}(x, \partial\Omega) \text{ for all } x \in \Omega\}.$$

We now define the operator

$$T : X \rightarrow X$$

by  $Tf = u$ , where  $u$  is the solution of the problem

$$\begin{cases} Lu := -\Delta u + (-\Delta)^s u + q \cdot \nabla u = f & \text{in } \Omega, \\ u = 0 & \text{on } \mathbb{R}^N \setminus \Omega. \end{cases} \quad (5.1)$$

Clearly  $T$  is a linear operator. The operator  $T$  is bounded since, by Theorem 1.6 and the Sobolev embedding, we have  $\|u\|_{C^{1,\alpha}(\overline{\Omega})} \leq C\|f\|_{C^{0,\alpha}(\overline{\Omega})}$ . Thus,

$$\|u\|_{C^{0,1}(\overline{\Omega})} \lesssim \|f\|_{C^{0,1}(\overline{\Omega})}.$$

Let us now prove that  $T(K \setminus \{0\}) \subseteq K^\circ$ . Let  $f \in K$  such that  $f \not\equiv 0$  and set  $Tf = w$ . Hence,  $Lw = f$  and  $w$  satisfies (5.1). We separate two cases according to  $s$ .

**Case 1.**  $0 < s < 1/2$ : in this case,  $w \in C^{2,\alpha}(\overline{\Omega})$  (Theorem 1.7). We can apply the strong maximum principle (Theorem 3.1) to the following problem

$$Lw = f \geq 0 \quad \text{in } \Omega, \quad w = 0 \text{ on } \mathbb{R}^N \setminus \Omega.$$

We obtain that  $w > 0$  in  $\Omega$ . Moreover, as  $w = 0$  on  $\partial\Omega$ , it follows from Theorem 3.2 that  $\partial_\nu w(x) < 0$  for all  $x \in \partial\Omega$ .

**Case 2.**  $s = 1/2$ : in this case, we have the interior regularity of  $w$  from Theorem 1.5. So we still have that  $w \in C^2(\Omega) \cap C(\mathbb{R}^n)$ , as  $w$  solves (5.1) and  $w \in C^{1,\alpha}(\bar{\Omega})$  by the Sobolev embedding. Applying the Hopf Lemma—stated in Theorem 1.3, we obtain again here that  $\partial_\nu w(x) < 0$  for all  $x \in \partial\Omega$ .

Thus, for all  $s \in (0, 1/2]$ , we have  $\partial_\nu w(x) < 0$  for all  $x \in \partial\Omega$ . As  $\partial\Omega$  is compact, then  $\max_{\partial\Omega} \partial_\nu w(x) < 0$ . This allows us find an open  $C^{0,1}$  neighbourhood  $\mathcal{O}$  of  $w$ , such that

$$\mathcal{O} \subseteq \{u \in X : \text{there is some } \varepsilon > 0 \text{ such that } u(x) \geq \varepsilon \text{dist}(x, \partial\Omega), \forall x \in \Omega\} \subseteq K^\circ.$$

Thus,  $w = Tf \in K^\circ$ .

We now verify that  $T$  is compact. Let  $\{f_n\}_n \subset X$  be a bounded sequence in  $X$ . Let us say that  $\|f_n\|_{L^\infty(\Omega)} \leq 1$ . It follows that  $f_n \in L^p(\Omega)$  for any  $1 < p < \infty$  and from the  $L^p$ -theory in Theorem 1.6, we have that  $Tf_n \in W^{2,p}(\Omega)$  for any  $1 < p < \infty$ . The Sobolev embedding implies that  $Tf_n \in C^{1,\alpha}(\bar{\Omega})$  for any  $0 < \alpha < 1$  and hence

$$\|Tf_n\|_{C^{1,\alpha}(\bar{\Omega})} \leq \|u\|_{W^{2,p}(\Omega)} \lesssim \|f_n\|_{L^p(\Omega)} \leq C,$$

where  $C$  is a constant independent of  $n$ . This implies that  $\{Tf_n\}_n$  is bounded in  $C^{1,\alpha}(\bar{\Omega})$ . By the Arzela-Ascoli theorem, the sequence  $\{Tf_n\}_n$  has a convergent subsequence (the convergence of the subsequence holds in  $C^1(\bar{\Omega})$  and hence in  $C^{0,1}(\bar{\Omega})$ ). This proves that  $T$  is compact.

Therefore, we can apply the Krein-Rutman theorem to assert that there exists a unique positive real number  $\varrho(T) > 0$  and a unique (up to multiplication by a nonzero constant) positive function  $f \in K^\circ$  such that  $Tf = \varrho(T)f$ . Therefore, the function  $\varphi_1 := Tf > 0$  satisfies the problem

$$\begin{cases} L\varphi_1 = \varrho(T)^{-1}\varphi_1 & \text{in } \Omega \\ \varphi_1 = 0 & \text{on } \mathbb{R}^N \setminus \Omega. \end{cases} \quad (5.2)$$

The function  $\varphi_1$  is in  $C^{2,\alpha}(\bar{\Omega})$  (from Theorem 1.7).

To complete the proof of Part (a) of the theorem, we set  $\lambda_1(\Omega, q) := \varrho(T)^{-1}$ . Then, from the Krein-Rutman theorem (Theorem A),  $\lambda_1(\Omega, q)$  is the principal eigenvalue for  $L$  in  $\Omega$  with the corresponding unique (up to multiplication by a nonzero constant) positive eigenfunction given by  $\varphi_1 \in C^{2,\alpha}(\bar{\Omega})$ .

From part (ii) of Theorem A, we know that  $\varrho = r(T) > 0$  satisfies: any eigenvalue  $\mu \neq \varrho$  for  $T$  satisfies  $|\mu| < \varrho$ . Now, since  $\lambda_1(\Omega, q) = \varrho(T)^{-1} > 0$ , the proof of part (b) follows.

We are left to prove the max-inf formulation (1.7) of  $\lambda_1(\Omega, q)$ —stated in part (c) of Theorem 1.2. We recall that  $\mathcal{V}(\Omega)$  is given by

$$\mathcal{V}(\Omega) := \left\{ u \in C^2(\Omega) \cap C^1(\bar{\Omega}) \cap C_c(\mathbb{R}^N) : u > 0 \text{ in } \Omega \text{ and } u \equiv 0 \text{ on } \mathbb{R}^N \setminus \Omega \right\}.$$

Since  $\varphi_1 \in \mathcal{V}(\Omega)$ , it follows that

$$\lambda_1(\Omega, q) \leq \sup_{u \in \mathcal{V}(\Omega)} \inf_{x \in \Omega} \frac{Lu(x)}{u(x)}.$$

Thus, we only need to prove that

$$\lambda_1(\Omega, q) \geq \sup_{u \in \mathcal{V}(\Omega)} \inf_{x \in \Omega} \frac{Lu(x)}{u(x)}.$$

and once we have equality then one sees we can replace the sup with a max. So we now argue by contradiction. Suppose that

$$\lambda_1(\Omega, q) < \sup_{u \in \mathcal{V}(\Omega)} \inf_{x \in \Omega} \frac{Lu(x)}{u(x)}.$$

Then, there exists  $\varepsilon > 0$  and a function  $v \in \mathcal{V}(\Omega)$  such that

$$\lambda_1(\Omega, q) + \varepsilon < \inf_{x \in \Omega} \frac{Lv(x)}{v(x)}. \quad (5.3)$$

Then note we have  $Lv > (\lambda_1(\Omega, q) + \varepsilon)v$  in  $\Omega$  with  $v = 0$  on  $\mathbb{R}^N \setminus \Omega$  and hence by Hopf's Lemma we have  $\partial_\nu v < 0$ . We now define

$$\tau^* := \sup\{\tau > 0 : v - \tau\varphi_1 \geq 0 \text{ in } \Omega\} \quad (5.4)$$

and note  $0 < \tau^* < \infty$  after noting that  $\varphi_1$  is sufficiently regular and the above Hopf result for  $v$ .

We now set  $w = v - \tau^*\varphi_1$ . First note that  $w \geq 0$  in  $\Omega$  and  $w = 0$  on  $\mathbb{R}^N \setminus \Omega$ . Since  $\varepsilon > 0$  we see that  $v$  cannot be a multiple of  $\varphi_1$  and hence  $w$  is not identically zero. Then note we have

$$Lw = Lv - \tau^*L\varphi_1 > \varepsilon v + \lambda_1(\Omega, q)w \geq 0 \quad \text{in } \Omega.$$

From the strong maximum principle we have  $w > 0$  in  $\Omega$  or  $w \equiv 0$  in  $\mathbb{R}^N$ . However,  $w \equiv 0$  contradicts that  $Lw > 0$ . Now, from Hopf Lemma (Theorem 1.3), we know that  $\partial_\nu v - \partial_\nu(\tau^*\varphi_1) = \partial_\nu w < 0$  on  $\partial\Omega$ . Thus, as  $v \geq \tau^*\varphi_1 \geq 0$  and  $\partial_\nu(\tau^*\varphi_1) > \partial_\nu v$  we can still find  $\delta > 0$  such that  $v \geq (\tau^* + \delta)\varphi_1 \geq 0$  in  $\Omega$ . This contradicts the fact that  $\tau^*$  is the largest possible in (5.4). Therefore (5.3) is false and we have

$$\lambda_1(\Omega, q) = \sup_{u \in \mathcal{V}(\Omega)} \inf_{x \in \Omega} \frac{Lu(x)}{u(x)}. \quad (5.5)$$

Furthermore, we know from part (a) of this theorem that

$$\varphi_1 \in \mathcal{V}(\Omega) \text{ and } L\varphi_1 = \lambda_1(\Omega, q)\varphi_1 \text{ in } \Omega.$$

Thus, the sup in (5.5) is indeed a max that is attained at  $\varphi_1$ . This completes the proof of (c) and hence the proof of Theorem 1.2.  $\square$

## References

1. D. Amundsen, A. Moameni and R. Y. Temgoua. *A mixed local and nonlocal supercritical Dirichlet problems*. arxiv preprint arXiv:2303.03273 (2023).
2. A. Arapostathis and A. Biswas. *Risk-sensitive control for a class of diffusions with jumps*. The Annals of Applied Probability 32, no. 6 (2022): 4106-4142.
3. A. Arapostathis, A. Biswas and P. Roychowdhury. *Generalized principal eigenvalues on  $\mathbb{R}^d$  of second order elliptic operators with rough nonlocal kernels*. Nonlinear Differential Equations and Applications NoDEA 30, no. 1 (2023): 10.

4. R. Arora, and V. D. Radulescu. *Combined effects in mixed local-nonlocal stationary problems*. arxiv preprint arXiv:2111.06701 (2021).
5. B. Barrios, L. M. Del Pezzo, J. Garcia-Melian and A. Quaas. *A priori bounds and existence of solutions for some nonlocal elliptic problems*. *Revista matemática iberoamericana* 34, no. 1 (2018): 195-220.
6. H. Berestycki, R. Ducasse and L. Rossi. *Generalized principal eigenvalues for heterogeneous road-field systems*. *Communications in Contemporary Mathematics* 22, no. 01 (2020): 1950013.
7. H. Berestycki, F. Hamel and N. Nadirashvili. *Elliptic eigenvalue problems with large drift and applications to nonlinear propagation phenomena*. *Communications in mathematical physics* 253 (2005): 451-480.
8. H. Berestycki, J. Coville and H. H. Vo. *On the definition and the properties of the principal eigenvalue of some nonlocal operators*. *Journal of Functional Analysis* 271, no. 10 (2016): 2701-2751.
9. H. Berestycki, L. Nirenberg and S. S. Varadhan. *The principal eigenvalue and maximum principle for second-order elliptic operators in general domains*. *Communications on Pure and Applied Mathematics* 47, no. 1 (1994): 47-92.
10. S. Biagi, S. Dipierro, E. Valdinoci and E. Vecchi. *A Brezis-Nirenberg type result for mixed local and nonlocal operators*. arxiv preprint arXiv:2209.07502 (2022).
11. S. Biagi, S. Dipierro, E. Valdinoci and E. Vecchi. *A Faber-Krahn inequality for mixed local and nonlocal operators*. *Journal d'Analyse Mathématique* (2023): 1-43.
12. S. Biagi, S. Dipierro, E. Valdinoci and E. Vecchi. *Mixed local and nonlocal elliptic operators: regularity and maximum principles*. *Communications in Partial Differential Equations* 47, no. 3 (2022): 585-629.
13. S. Biagi, E. Vecchi, S. Dipierro, and E. Valdinoci. *Semilinear elliptic equations involving mixed local and nonlocal operators*. *Proceedings of the Royal Society of Edinburgh Section A: Mathematics* 151, no. 5 (2021): 1611-1641.
14. A. Biswas and M. Modasiya. *Mixed local-nonlocal operators: maximum principles, eigenvalue problems and their applications*. arxiv preprint arXiv:2110.06746 (2021).
15. A. Biswas, M. Modasiya, and A. Sen. *Boundary regularity of mixed local-nonlocal operators and its application*. *Annali di Matematica Pura ed Applicata (1923-)* 202, no. 2 (2023): 679-710.
16. D. Cassani, L. Vilasi and Y. Wang. *Local versus nonlocal elliptic equations: short-long range field interactions*. *Advances in Nonlinear Analysis* 10, no. 1 (2020): 895-921.
17. K. Deimling. *Nonlinear functional analysis*. Courier Corporation, (2010).
18. S. Dipierro and E. Valdinoci. *Description of an ecological niche for a mixed local/nonlocal dispersal: an evolution equation and a new Neumann condition arising from the superposition of Brownian and Lévy processes*. *Physica A: Statistical Mechanics and its Applications* 575 (2021): 126052.
19. Y. Du. *Order structure and topological methods in nonlinear partial differential equations: Vol. 1: Maximum principles and applications*. Vol. 2. World Scientific, (2006).
20. Y. Fang, B. Shang and C. Zhang. *Regularity theory for mixed local and nonlocal parabolic  $p$ -Laplace equations*. *The Journal of Geometric Analysis* 32, no. 1 (2022): 22.
21. P. A. Feulefack. *The logarithmic Schrödinger operator and associated Dirichlet problems*. *Journal of Mathematical Analysis and Applications* 517, no. 2 (2023): 126656.

22. P. A. Feulefack and S. Jarohs. *Nonlocal operators of small order*. Annali di Matematica Pura ed Applicata (1923-) 202, no. 4 (2023): 1501-1
23. P. Garain. *On a class of mixed local and nonlocal semilinear elliptic equation with singular nonlinearity*. The Journal of Geometric Analysis 33, no. 7 (2023): 212.
24. P. Garain and J. Kinnunen. *Weak Harnack inequality for a mixed local and nonlocal parabolic equation*. Journal of Differential Equations 360 (2023): 373-406.
25. P. Garain and A. Ukhlov. *Mixed local and nonlocal Sobolev inequalities with extremal and associated quasilinear singular elliptic problems*. Nonlinear Analysis 223 (2022): 113022.
26. M.G. Garroni and J. L. Menaldi. *Second order elliptic integro-differential problems*. CRC Press, (2002).
27. D. Gilbarg and NS. Trudinger. *Elliptic partial differential equations of second order*. Vol. 224, no. 2. Berlin: springer, (1977).
28. G. Giovannardi, D. Mugnai and E. Vecchi. *An Ahmad-Lazer-Paul-type result for indefinite mixed local-nonlocal problems*. Journal of Mathematical Analysis and Applications (2023): 127442.
29. C. Mou. *Existence of  $C^\alpha$  solutions to integro-PDEs*. Calculus of Variations and Partial Differential Equations 58, no. 4 (2019): 143.
30. L. M. D. Pezzo, R. Ferreira and J. D. Rossi. *Eigenvalues for a combination between local and nonlocal  $p$ -Laplacians*. Fractional Calculus and Applied Analysis 22, no. 5 (2019): 1414-1436.
31. A. Quaas, A. Salort and A. Xia. *Principal eigenvalues of fully nonlinear integro-differential elliptic equations with a drift term*. ESAIM: Control, Optimisation and Calculus of Variations 26 (2020): 36.
32. X. Su, E. Valdinoci, Y. Wei and J. Zhang. *Regularity results for solutions of mixed local and nonlocal elliptic equations*. Mathematische Zeitschrift 302, no. 3 (2022): 1855-1878.
33. L. Silvestre. *Regularity of the obstacle problem for a fractional power of the Laplace operator*. Communications on Pure and Applied Mathematics: A Journal Issued by the Courant Institute of Mathematical Sciences 60, no. 1 (2007): 67-112.

Craig Cowan

Department of Mathematics, University of Manitoba, Winnipeg, MB, Canada

e-mail: craig.cowan@umanitoba.ca

Mohammad El Smaily

Department of Mathematics and Statistics, University of Northern British Columbia, Prince George, BC, Canada

e-mail: mohammad.elsmaily@unbc.ca

Pierre Aime Feulefack

Department of Mathematics and Statistics, University of Northern British Columbia, Prince George, BC, Canada

e-mail: PierreAime.Feulefack@unbc.ca