

# Parameter inference and nonequilibrium identification for Markovian systems based on coarse-grained observations

Bingjie Wu<sup>1</sup> and Chen Jia<sup>1\*</sup>

<sup>1</sup> Applied and Computational Mathematics Division,  
Beijing Computational Science Research Center, Beijing 100193, China.

Most experiments can only detect a set of coarse-grained clusters of a molecular system, while the internal microstates are often inaccessible. Here, based on a sufficient long coarse-grained trajectory, we obtain a set of sufficient statistics which extracts all statistic information of coarse-grained observations. Based on these sufficient statistics, we completely solve the problems of parameter inference and nonequilibrium identification for a general Markovian system with an arbitrary number of microstates and arbitrary coarse-grained partitioning. Our nonequilibrium criterion generalizes the one obtained in [J. Chem. Phys. 132:041102 (2010)] for a three-state system with two coarse-grained clusters, and is much superior to the classical criterion based on autocorrelation functions.

*Introduction* — A mesoscopic molecular system is widely modelled as a Markov process with a large number of microstates. In experiments, it often occurs that only a set of coarse-grained clusters can be detected, while the internal microstates of the system are often blurred or inaccessible. For instance, in live-cell imaging, one can obtain the time trace of the copy number of a protein in a single cell; however, it is difficult to determine whether the gene is in an active or an inactive state [1]. Due to the inability to accurately identify all microstates, the data obtained is usually the time trace of some coarse-grained states, which only retain a small degrees of freedom of the system [2–4]. Other well-known examples of partial observations include ion channel opening [5], molecular docking and undocking to a sensor [6], and flagellar motor switches [7].

Given a sufficiently long trajectory of coarse-grained states, two natural and crucial questions arise: (i) is it possible to determine the transition topology and even all transition rates between all microstates? (ii) If complete recovery of transition rates is impossible, is it still possible to determine whether the system is in equilibrium or in a nonequilibrium steady state (NESS) and even estimate the values of some macroscopic thermodynamic quantities? Over the past two decades, numerous studies have partially answered these two questions; however, there is still a lack of a unified theory for general systems.

Among these studies, some [8–10] focus on transition topology inference; some [11–13] focus on transition rate inference with a given transition topology; some [14–17] investigate nonequilibrium detection based on a two-state coarse-grained trajectory. Recently, several studies have estimated the values of some macroscopic thermodynamic quantities, such as entropy production and cycle affinities, based on partial observations [18–22]. In this study, we develop a novel approach based on sufficient statistics, and completely solve the problems of parameter inference and nonequilibrium detection for a general Markovian system with an arbitrary number of microstates and an arbitrary number of coarse-grained clusters.




FIG. 1. **Model.** (a) An  $N$ -state Markovian system with  $n$  coarse-grained states  $A_1, \dots, A_n$ , each composed of multiple microstates. (b) Illustration of an infinitely long trajectory of coarse-grained states. (c) The ladder model. (d) The cyclic model. (e) The tree model. (f) The linear model.

*Model* — We consider an  $N$ -state molecular system modeled by a continuous-time Markov chain  $(X_s)_{s \geq 0}$  with microstates  $x = 1, \dots, N$  and generator matrix  $Q = (q_{xy})$ , where  $q_{xy}$  denotes the transition rate from state  $x$  to state  $y$  whenever  $x \neq y$  and  $q_{xx} = -\sum_{y \neq x} q_{xy}$ . Recall that the transition diagram of the Markovian system is a directed graph with vertex set  $V = \{1, \dots, N\}$  and edge set  $E = \{(x, y) : q_{xy} > 0\}$ . Experimentally, it often occurs that not all microstates can be observed — what can be observed are often some coarse-grained states, each being composed of multiple microstates. Specifically, we assume that the  $N$  microstates can be divided into  $n$  coarse-grained states  $A_1, \dots, A_n$  (Fig. 1(a)). We also assume that the observation is an infinitely long trajectory of coarse-grained states (Fig. 1(b)) [8–10].

Based on the coarse-grained clusters, the generator

matrix  $Q$  can be represented by the block form

$$Q = \begin{pmatrix} Q_{A_1} & Q_{A_1 A_2} & \cdots & Q_{A_1 A_n} \\ Q_{A_2 A_1} & Q_{A_2} & \cdots & Q_{A_2 A_n} \\ \vdots & \vdots & \ddots & \vdots \\ Q_{A_n A_1} & Q_{A_n A_2} & \cdots & Q_{A_n} \end{pmatrix}.$$

The steady-state distribution  $\pi = (\pi_{A_1}, \dots, \pi_{A_n})$  of the system must satisfy  $\pi Q = \mathbf{0}$ , where  $\mathbf{0} = (0, \dots, 0)$  is the zero vector. For simplicity, we assume that each diagonal block  $Q_{A_i}$  has different eigenvalues  $-\lambda_1^i, \dots, -\lambda_{|A_i|}^i$  (any matrix can be approximated by such matrices to any degree of accuracy). Then there exists an invertible matrix  $\Phi_{A_i}$  such that  $Q_{A_i} = \Phi_{A_i}^{-1} \Lambda_{A_i} \Phi_{A_i}$ , where  $\Lambda_{A_i} = \text{diag}(-\lambda_1^i, \dots, -\lambda_{|A_i|}^i)$  is a diagonal matrix. Note that each row of  $\Phi_{A_i}$  is an eigenvector of  $Q_{A_i}$ . For convenience, the sum of elements of each eigenvector is normalized to one, i.e.,  $\Phi_{A_i} \mathbf{1}^T = \mathbf{1}^T$ , where  $\mathbf{1} = (1, \dots, 1)$ . Moreover, we set

$$\begin{aligned} \Lambda_{A_i A_j} &= \Phi_{A_i} Q_{A_i A_j} \Phi_{A_j}^{-1} = (\lambda_{kl}^{ij}), \\ \alpha_{A_i} &= \pi_{A_i} \Phi_{A_i}^{-1} = (\alpha_k^i). \end{aligned} \quad (1)$$

These quantities will play a crucial role in our analysis.

*Sufficient statistics* — In what follows, we assume that the system has reached a steady state. We next examine which information can be extracted from the infinitely long coarse-grained trajectory (Fig. 1(b)). Clearly, what can be observed from the trajectory are the jump times and the coarse-grained states before and after each jump within any finite period of time. In other words, for any given jump times  $0 \leq t_1 < \dots < t_m \leq t$  and any coarse-grained states  $A_{i_1}, \dots, A_{i_{m+1}}$ , we can estimate the following probability (density) since the trajectory is infinitely long [23]:

$$\mathbb{P}(X_s \in A_{i_1} (s < t_1), \dots, X_s \in A_{i_{m+1}} (t_m \leq s \leq t)). \quad (2)$$

These probabilities are actually all statistical information that can be extracted from coarse-grained observations.

We first consider the case where there is no jump before time  $t$ . Based on the notation in Eq. (1), it is clear that

$$\begin{aligned} \mathbb{P}(X_s \in A_i (0 \leq s \leq t)) &= \pi_{A_i} e^{Q_{A_i} t} \mathbf{1}^T \\ &= \alpha_{A_i} e^{\Lambda_{A_i} t} \mathbf{1}^T = \sum_k \alpha_k^i e^{-\lambda_k^i t}. \end{aligned} \quad (3)$$

Based on coarse-grained observations, we can estimate the probability on the left-hand side of the above equation for each time  $t$ . Since time  $t$  is arbitrary, we can estimate the values of all  $\alpha_k^i$  and  $\lambda_k^i$  and hence  $\alpha_{A_i}$  and  $\Lambda_{A_i}$  can be determined. Similarly, if there is only one jump before time  $t$ , for any  $i \neq j$ , we have

$$\begin{aligned} \mathbb{P}(X_s \in A_i (0 \leq s < t_1), X_s \in A_j (t_1 \leq s \leq t)) \\ &= \pi_{A_i} e^{Q_{A_i} t_1} Q_{A_i A_j} e^{Q_{A_j} (t-t_1)} \mathbf{1}^T \\ &= \alpha_{A_i} e^{\Lambda_{A_i} t_1} \Lambda_{A_i A_j} e^{\Lambda_{A_j} (t-t_1)} \mathbf{1}^T. \end{aligned} \quad (4)$$

Since times  $t_1$  and  $t$  are arbitrary and since we have determined  $\alpha_{A_i}$  and  $\Lambda_{A_i}$ , we can also determine  $\Lambda_{A_i A_j}$  from coarse-grained observations for any  $i \neq j$  [24].

Similarly to Eqs. (3) and (4), it can be proved that [24]

$$\begin{aligned} \mathbb{P}(X_s \in A_{i_1} (0 \leq s < t_1), \dots, X_s \in A_{i_{m+1}} (t_m \leq s \leq t)) \\ &= \pi_{A_{i_1}} e^{Q_{A_{i_1}} t_1} \cdots Q_{A_{i_m} A_{i_{m+1}}} e^{Q_{A_{i_{m+1}}} (t-t_m)} \mathbf{1}^T \\ &= \alpha_{A_{i_1}} e^{\Lambda_{A_{i_1}} t_1} \cdots \Lambda_{A_{i_m} A_{i_{m+1}}} e^{\Lambda_{A_{i_{m+1}}} (t-t_m)} \mathbf{1}^T. \end{aligned} \quad (5)$$

This shows that the probability given in Eq. (2) can be represented by  $\alpha_{A_i}$ ,  $\Lambda_{A_i}$ , and  $\Lambda_{A_i A_j}$ . Hence these three quantities contain all coarse-grained statistical information. In fact,  $\alpha_{A_i}$ ,  $\Lambda_{A_i}$ , and  $\Lambda_{A_i A_j}$  are not independent. Since  $Q \mathbf{1}^T = \mathbf{0}^T$ , it is clear that  $\Lambda \mathbf{1}^T = \mathbf{0}^T$ , where

$$\Lambda = \begin{pmatrix} \Lambda_{A_1} & \Lambda_{A_1 A_2} & \cdots & \Lambda_{A_1 A_n} \\ \vdots & \vdots & \ddots & \vdots \\ \Lambda_{A_n A_1} & \Lambda_{A_n A_2} & \cdots & \Lambda_{A_n} \end{pmatrix}.$$

This implies that  $\lambda_k^i = \sum_{j,l} \lambda_{kl}^{ij}$ , where  $\lambda_{kl}^{ij}$  are the elements of  $\Lambda_{A_i A_j}$  defined in Eq. (1). Hence  $\Lambda_{A_i}$  can be determined by all  $\Lambda_{A_i A_j}$ . Since  $\pi Q = \mathbf{0}$ , we have  $\alpha \Lambda = \mathbf{0}$ , where  $\alpha = (\alpha_{A_1}, \dots, \alpha_{A_n})$ . This equation, together with the normalization condition

$$\alpha \mathbf{1}^T = \sum_{i=1}^n \pi_{A_i} \Phi_{A_i}^{-1} \mathbf{1}^T = \sum_{i=1}^n \pi_{A_i} \mathbf{1}^T = 1,$$

shows that  $\alpha_{A_i}$  can also be determined by all  $\Lambda_{A_i A_j}$ .

Recall that a family of statistics that contain all statistical information of the observations are called sufficient statistics [25]. Since the probability in Eq. (2) can be represented by  $\alpha_{A_i}$ ,  $\Lambda_{A_i}$ , and  $\Lambda_{A_i A_j}$  and since  $\alpha_{A_i}$  and  $\Lambda_{A_i}$  can be determined by all  $\Lambda_{A_i A_j}$ , it is clear that all  $\Lambda_{A_i A_j}$  are the sufficient statistics for infinitely long coarse-grained observations. In particular, the number of these sufficient statistics is given by

$$S = \sum_{1 \leq i \neq j \leq n} |A_i| |A_j| = 2 \sum_{1 \leq i < j \leq n} |A_i| |A_j|.$$

Note that based on coarse-grained observations, two systems with different generator matrices but having the same  $\Lambda_{A_i A_j}$  are statistically indistinguishable.

*Parameter inference* — In practice, a crucial question is whether all transition rates of a Markovian system can be inferred from an infinitely long coarse-grained trajectory. Note that each transition rate  $q_{xy}$  of the system corresponds to a direct edge in the transition diagram  $(V, E)$ . Hence the system has  $|E|$  unknown parameters, where  $|E|$  denotes the number of elements in the edge set  $E$ . Note that each  $\Lambda_{A_i A_j}$  is uniquely determined by all transition rates and in the following, we rewrite it as  $\Lambda_{A_i A_j}(q_{xy})$ . Based on coarse-grained

observations, we can obtain estimates of the sufficient statistics  $\Lambda_{A_i A_j}$  and hence all transition rates should satisfy the following set of equations:

$$\Lambda_{A_i A_j}(q_{xy}) = \tilde{\Lambda}_{A_i A_j}, \quad \forall 1 \leq i \neq j \leq n, \quad (6)$$

where  $\tilde{\Lambda}_{A_i A_j}$  are the estimates of  $\Lambda_{A_i A_j}$ . Clearly, Eq. (6) has  $|E|$  unknown parameters and  $S$  equations, where  $S$  is the number of sufficient statistics. Therefore, we obtain the following criteria regarding parameter inference: (i) when  $S < |E|$ , it is impossible to determine all transition rates of the system; (ii) when  $S \geq |E|$ , all transition rates can be determined if and only if Eq. (6) has a unique solution. In general, it is very difficult to give a simple criterion for the unique solvability of Eq. (6); however, it can be checked numerically using, e.g., Gröbner basis computation [26]. Next we focus on three examples.

1) The ladder model (Fig. 1(c)). This model is widely used to model allostery of receptors in living cells with strong cooperativity and high sensitivity [27]. Consider a receptor with two conformational states and  $n$  ligand binding sites. According to the conformational state and the number of occupied binding sites, each receptor can be modeled by a Markovian system with  $N = 2n$  microstates. Due to technical limitations, we are often unable to distinguish between the two conformational states. The  $N$  microstates and  $n$  coarse-grained states are shown in Fig. 1(c). Clearly, we have  $|A_1| = \dots = |A_n| = 2$  and there are  $|E| = 6n - 4$  unknown parameters for the system. Note that for any  $n \geq 2$ , we have

$$S = 4n^2 - 4n \geq 6n - 4 = |E|.$$

Hence all transition rates of the ladder model can be inferred if and only if Eq. (6) has unique solution.

We next show that Eq. (6) is indeed uniquely solvable when  $N = 4$  and  $n = 2$ . In this case, we have  $A_1 = \{1, 2\}$  and  $A_2 = \{3, 4\}$ . It is easy to check that

$$\Phi_{A_1} = \begin{bmatrix} \frac{q_{21}}{\kappa_1 + q_{21} - q_{11}} & \frac{\kappa_1 - q_{11}}{\kappa_1 + q_{21} - q_{11}} \\ \frac{q_{21}}{\kappa_2 + q_{21} - q_{11}} & \frac{\kappa_2 - q_{11}}{\kappa_2 + q_{21} - q_{11}} \end{bmatrix},$$

$$\Phi_{A_2} = \begin{bmatrix} \frac{q_{43}}{\kappa_3 + q_{43} - q_{33}} & \frac{\kappa_3 - q_{33}}{\kappa_3 + q_{43} - q_{33}} \\ \frac{q_{43}}{\kappa_4 + q_{43} - q_{33}} & \frac{\kappa_4 - q_{33}}{\kappa_4 + q_{43} - q_{33}} \end{bmatrix},$$

where

$$\kappa_1 + \kappa_2 = q_{11} + q_{22}, \quad \kappa_1 \kappa_2 = q_{11} q_{22} - q_{12} q_{21},$$

$$\kappa_3 + \kappa_4 = q_{33} + q_{44}, \quad \kappa_3 \kappa_4 = q_{33} q_{44} - q_{34} q_{43}.$$

Then Eq. (6) reduces to

$$\Phi_{A_1} Q_{A_1 A_2} \Phi_{A_2}^{-1} = \tilde{\Lambda}_{A_1 A_2}, \quad \Phi_{A_2} Q_{A_2 A_1} \Phi_{A_1}^{-1} = \tilde{\Lambda}_{A_2 A_1}.$$

Straightforward computations show that the generator matrix  $Q$  can be estimated as

$$Q = \begin{bmatrix} \Theta_{A_1} \tilde{\Lambda}_{A_1} \Theta_{A_1}^{-1} & \Theta_{A_1} \tilde{\Lambda}_{A_1 A_2} \Theta_{A_2}^{-1} \\ \Theta_{A_2} \tilde{\Lambda}_{A_2 A_1} \Theta_{A_1}^{-1} & \Theta_{A_2} \tilde{\Lambda}_{A_2} \Theta_{A_2}^{-1} \end{bmatrix},$$

where

$$\Theta_{A_1} = \begin{bmatrix} \frac{a_{21}}{\eta_1 + a_{21} - a_{11}} & \frac{\eta_1 - a_{11}}{\eta_1 + a_{21} - a_{11}} \\ \frac{a_{21}}{\eta_2 + a_{21} - a_{11}} & \frac{\eta_2 - a_{11}}{\eta_2 + a_{21} - a_{11}} \end{bmatrix},$$

$$\Theta_{A_2} = \begin{bmatrix} \frac{b_{43}}{\eta_3 + b_{43} - b_{33}} & \frac{\eta_3 - b_{33}}{\eta_3 + b_{43} - b_{33}} \\ \frac{b_{43}}{\eta_4 + b_{43} - b_{33}} & \frac{\eta_4 - b_{33}}{\eta_4 + b_{43} - b_{33}} \end{bmatrix},$$

with

$$a_{ij} = \tilde{\lambda}_{i3}^{12} \tilde{\lambda}_{3j}^{21} + \tilde{\lambda}_{i4}^{12} \tilde{\lambda}_{4j}^{21}, \quad b_{ij} = \tilde{\lambda}_{i1}^{21} \tilde{\lambda}_{1j}^{12} + \tilde{\lambda}_{i2}^{21} \tilde{\lambda}_{2j}^{12},$$

$$\eta_1 + \eta_2 = a_{11} + a_{22}, \quad \eta_1 \eta_2 = a_{11} a_{22} - a_{12} a_{21}$$

$$\eta_3 + \eta_4 = b_{33} + b_{44}, \quad \eta_3 \eta_4 = b_{33} b_{44} - b_{34} b_{43}.$$

2) The cyclic model (Fig. 1(d)). Many crucial cellular biochemical processes can be modeled as cyclic Markovian systems such as conformational changes of enzymes and ion channels [5, 28], phosphorylation-dephosphorylation cycle [29], cell cycle progression [30], and gene state switching [31]. Note that an  $N$ -state cyclic model has  $|E| = 2N$  unknown parameters. If there are only two coarse-grained states ( $n = 2$ ), then it is impossible to infer all parameters when  $|A_1| = 1$  and  $|A_2| = N - 1$  since  $S = 2(N - 1) < |E|$ . For any other coarse-grained partitioning, we have  $S \geq 4(N - 2) \geq |E|$  and hence parameter inference is generally possible. When  $n \geq 3$ , we have  $S \geq 4N - 6 \geq |E|$ , which allows a complete parameter inference if Eq. (6) can be uniquely solved.

3) The tree and linear models (Fig. 1(e)). We finally focus on an  $N$ -state system whose transition diagram is a tree. For any coarse-grained partitioning, it is easy to check that  $S \geq 2(N - 1) = |E|$ . Hence all transition rates can be determined if Eq. (6) has unique solution. In [24], we show that Eq. (6) is uniquely solvable when  $N = 3$  and  $n = 2$ . In particular, a Markovian system with linear transitions (Fig. 1(f)) can be viewed as a tree. The linear model also widely used in biochemical studies [29].

*Nonequilibrium identification* — When the number sufficient statistics is less than the number of unknown parameters, it is impossible to determine all transition rates from coarse-grained observations. However, we may still identify whether the system is in an NESS. Our NESS criterion is based on the sufficient statistics  $\Lambda_{A_i A_j}$ . Recall that once  $\Lambda_{A_i A_j}$  are determined,  $\alpha_{A_i}$  are automatically determined by solving  $\alpha \Lambda = \mathbf{0}$  and  $\alpha \mathbf{1}^T = 1$ . In [24], we prove that if the system is in equilibrium, then the following two conditions must be satisfied:

(i) (coarse-grained probability distribution condition)

$$\alpha_k^i \geq 0. \quad (7)$$

(ii) (coarse-grained detailed balance condition)

$$\alpha_k^i \lambda_{kl}^{ij} = \alpha_l^j \lambda_{lk}^{ji}. \quad (8)$$

Recall that  $\alpha \mathbf{1}^T = 1$ . When the system is in equilibrium, we have  $\alpha_k^i \geq 0$  and thus  $\alpha = (\alpha_{A_1}, \dots, \alpha_{A_n})$  is a probability distribution. That is why Eq. (7) is called the coarse-grained probability distribution condition. On the other hand, in equilibrium, the system satisfies the detailed balance condition  $\pi_x q_{xy} = \pi_y q_{yx}$ . Eq. (8) can be viewed as the coarse-grained version of the detailed balance condition. The above result implies that if any one of Eqs. (7) and (8) is violated, then the system must be in a NESS. This gives a general criterion of nonequilibrium identification based on coarse-grained observations.

For a three-state cyclic system with two coarse-grained states ( $N = 3$  and  $n = 2$ ), we have seen that it is impossible to infer all parameters. In [16], the authors showed that this system is in a NESS when

$$TL^2 > LSM - M^2, \quad (9)$$

where  $L = q_{12} + q_{13}$ ,  $S = q_{21} + q_{23} + q_{31} + q_{32}$ ,  $T = q_{21}q_{31} + q_{21}q_{32} + q_{23}q_{31}$ , and  $M = q_{12}(q_{23} + q_{31} + q_{32}) + q_{13}(q_{21} + q_{23} + q_{32})$  are another set of sufficient statistics for the three-state system that can be determined by coarse-grained observations [16, 17]. Our result can be viewed as an extension of Eq. (9) to complex molecular systems with an arbitrary number of microstates and an arbitrary number of coarse-grained states.

In particular, when  $N = 3$  and  $n = 2$ , our criterion reduces to Eq. (9). This can be seen as follows. First, since  $\alpha \Lambda = \mathbf{0}$ , it is easy to see that

$$\alpha_1^1 \lambda_{12}^{12} = \alpha_2^2 \lambda_1^2 = \alpha_2^2 \lambda_{21}^{21}, \quad \alpha_1^1 \lambda_{13}^{12} = \alpha_3^2 \lambda_2^2 = \alpha_3^2 \lambda_{31}^{21}. \quad (10)$$

Hence the coarse-grained detailed balance condition is satisfied. On the other hand, since  $\alpha \mathbf{1}^T = \alpha_1^1 + \alpha_2^2 + \alpha_3^2 = 1$ , it follows from Eq. (10) that

$$\begin{aligned} \alpha_1^1 &= \frac{\lambda_{21}^{21} \lambda_{31}^{21}}{\lambda_{21}^{21} \lambda_{31}^{21} + \lambda_{12}^{12} \lambda_{31}^{21} + \lambda_{13}^{12} \lambda_{21}^{21}}, \\ \alpha_2^2 &= \frac{\lambda_{12}^{12} \lambda_{31}^{21}}{\lambda_{21}^{21} \lambda_{31}^{21} + \lambda_{12}^{12} \lambda_{31}^{21} + \lambda_{13}^{12} \lambda_{21}^{21}}, \\ \alpha_3^2 &= \frac{\lambda_{13}^{12} \lambda_{21}^{21}}{\lambda_{21}^{21} \lambda_{31}^{21} + \lambda_{12}^{12} \lambda_{31}^{21} + \lambda_{13}^{12} \lambda_{21}^{21}}. \end{aligned} \quad (11)$$

Furthermore, it is easy to check that [24]

$$\begin{aligned} \lambda_{21}^{21} \lambda_{31}^{21} &= T, \quad \lambda_{12}^{12} \lambda_{31}^{21} + \lambda_{13}^{12} \lambda_{21}^{21} = M, \\ \lambda_{12}^{12} \lambda_{31}^{21} \lambda_{13}^{12} \lambda_{21}^{21} &= \frac{T(LSM - M^2 - TL^2)}{(\lambda_{13}^{12} - \lambda_{12}^{12})^2}. \end{aligned} \quad (12)$$

Combining Eqs. (11) and (12), we immediately obtain

$$\begin{aligned} \alpha_1^1 &= \frac{T}{T+M} \geq 0, \quad \alpha_2^2 + \alpha_3^2 = \frac{M}{T+M} \geq 0, \\ \alpha_2^2 \alpha_3^2 &= \frac{T(LSM - M^2 - TL^2)}{(T+M)^2 (\lambda_{13}^{12} - \lambda_{12}^{12})^2}. \end{aligned}$$

Since  $\alpha_1^1 \geq 0$  and  $\alpha_2^2 + \alpha_3^2 \geq 0$ , the coarse-grained probability distribution condition is violated if and only if  $\alpha_2^2 \alpha_3^2 < 0$ , which is obviously equivalent to Eq. (9).

We now compare our NESS criterion with the classical criterion based on autocorrelation functions. For any observable  $\phi$ , recall that the autocorrelation function of the system,  $B_\phi(t) = \text{Cov}(\phi(X_0), \phi(X_t))$ , is defined as the steady-state covariance between  $\phi(X_0)$  and  $\phi(X_t)$ . For coarse-grained observations, the observable  $\phi$  is usually chosen as  $\phi(x) = i$  for all  $x \in A_i$ . It is well known [32] that if the system is in equilibrium, then

$$B_\phi(t) = \sum_{m=1}^{N-1} c_m e^{-\ell_m t}, \quad \ell_m > 0, \quad c_m \geq 0,$$

where  $-\ell_m < 0$  are all nonzero eigenvalues of the generator matrix  $Q$  and  $c_m \geq 0$  are the coefficients (note that in the case, the autocorrelation function must be monotonic). Hence if there exists some  $1 \leq m \leq N-1$  such that any one of  $\ell_m$  and  $c_m$  is negative or not real, then the system must be in a NESS.

In [24], we have proved a stronger result — if Eqs. (7) and (8) hold, then we also have  $\ell_m > 0$  and  $c_m \geq 0$ . This suggests that all NESS scenarios that can be identified by the autocorrelation criterion can definitely be identified by our criterion; in other words, our NESS criterion is superior to the autocorrelation criterion. In particular, if there are two coarse-grained states ( $n = 2$ ) with  $|A_1| = 1$  and  $|A_2| = N-1$ , then the two criteria are equivalent; in this case, the number of sufficient statistics is  $S = 2(N-1)$  and  $\ell_1, \dots, \ell_{N-1}, c_1, \dots, c_{N-1}$  is exactly a set of sufficient statistics [24]. For other coarse-grained partitionings, the two criteria are in general not equivalent and our criterion turns out to be much more powerful.

Finally, as an example, we focus on a four-state fully connected system with two different coarse-grained partitionings (Fig. 2(a),(b)). The system has 12 unknown parameters and we want to determine whether it is in a NESS. We first consider the partitioning of  $A_1 = \{1\}$  and  $A_2 = \{2, 3, 4\}$  (Fig. 2(a)). In this case, our criterion is equivalent to the autocorrelation criterion. Fig. 2(c) illustrates the NESS region in the parameter space that can be identified by our criterion, or equivalently, the autocorrelation criterion (shown in shaded blue) and the region that fails to be identified by the two criteria (shown in green). The red line shows the region of equilibrium states. In this case, NESS can only be detected in the parameter region far from the red line.

We then consider another partitioning, i.e.  $A_1 = \{1, 2\}$  and  $A_2 = \{3, 4\}$  (Fig. 2(b)). Fig. 2(c) shows the NESS regions that can be identified by our criterion (shown in blue) and by the autocorrelation criterion (shown in shaded blue). Clearly, the entire NESS region can be captured by our criterion but only a small subregion can be detected by the autocorrelation criterion. To gain






FIG. 2. **NESS identification for a four-state Markovian system.** (a) System with coarse-grained partitioning  $A_1 = \{1\}$  and  $A_2 = \{2, 3, 4\}$ . (b) System with coarse-grained partitioning  $A_1 = \{1, 2\}$  and  $A_2 = \{3, 4\}$ . (c) Phase diagram in the  $q_{14} - q_{41}$  plane for the system in (a). (d) Same as (c) but for the system in (b). In (c),(d), the red line shows the region of equilibrium states (EQ). The blue (and shaded blue) area shows the NESS region that can be identified by our criterion. The shaded blue area shows the NESS region that can be identified by the autocorrelation criterion. The green area shows the NESS region that fails to be detected by our criterion. The parameters are chosen as  $q_{12} = 1$ ,  $q_{13} = q_{21} = 2$ ,  $q_{23} = q_{24} = 4$ ,  $q_{31} = q_{32} = q_{34} = 3$ , and  $q_{42} = q_{43} = 5$ .

deeper insights, note that there are  $S = 6$  sufficient statistics for the partitioning shown in Fig. 2(a), while for the one shown in Fig. 2(b), there are  $S = 8$  sufficient statistics. More sufficient statistics implies larger NESS region that can be identified by our criterion.

*Conclusion* — Here we addressed the problems of parameter inference and nonequilibrium identification based on coarse-grained observations for a general system with an arbitrary number of microstates and arbitrary coarse-grained partitioning. Our method is based on extracting a set of sufficient statistics that incorporates all statistical information of an infinitely long coarse-grained trajectory. Using these sufficient statistics, the problems of parameter recovery and nonequilibrium detection can be brought into a unified theoretical framework.

*Acknowledgements* — C. J. acknowledges support from National Natural Science Foundation of China with grant No. U2230402 and No. 12271020.

\* Correspondence: [chenjia@csrc.ac.cn](mailto:chenjia@csrc.ac.cn)

- [1] I. Golding, J. Paulsson, S. M. Zawilski, and E. C. Cox, *Cell* **123**, 1025 (2005).  
 [2] U. Seifert, *Annual Review of Condensed Matter Physics*

- 10**, 171 (2019).  
 [3] M. Esposito, *Phys. Rev. E* **85**, 041125 (2012).  
 [4] G. Teza and A. L. Stella, *Phys. Rev. Lett.* **125**, 110601 (2020).  
 [5] B. Sakmann, *Single-channel recording* (Springer Science & Business Media, 2013).  
 [6] M. Skoge, S. Naqvi, Y. Meir, and N. S. Wingreen, *Phys. Rev. Lett.* **110**, 248102 (2013).  
 [7] H. C. Berg, *Annu. Rev. Biochem.* **72**, 19 (2003).  
 [8] O. Flomenbom, J. Klafter, and A. Szabo, *Biophys. J.* **88**, 3780 (2005).  
 [9] O. Flomenbom and R. J. Silbey, *Proc. Natl. Acad. Sci. USA* **103**, 10907 (2006).  
 [10] O. Flomenbom and R. Silbey, *J. Chem. Phys.* **128** (2008).  
 [11] Y. Deng, S. Peng, M. Qian, and J. Feng, *J. Phys. A: Math. Gen.* **36**, 1195 (2003).  
 [12] X. Xiang, X. Yang, Y. Deng, and J. Feng, *J. Phys. A: Math. Gen.* **39**, 9477 (2006).  
 [13] D. Y. Y. X. Xiang X, Zhou J, *Chaos* (2024).  
 [14] J. B. Witkoskie and J. Cao, *The Journal of Physical Chemistry B* **110**, 19009 (2006).  
 [15] Y. Tu, *Proc. Natl. Acad. Sci. USA* **105**, 11737 (2008).  
 [16] C. P. Amann, T. Schmiedl, and U. Seifert, *J. Chem. Phys.* **132**, 041102 (2010).  
 [17] C. Jia and Y. Chen, *J. Phys. A: Math. Theor.* **48**, 205001 (2015).  
 [18] D. J. Skinner and J. Dunkel, *Phys. Rev. Lett.* **127**, 198101 (2021).  
 [19] P. E. Harunari, A. Dutta, M. Polettini, and É. Roldán, *Phys. Rev. X* **12**, 041026 (2022).  
 [20] J. Van der Meer, B. Ertel, and U. Seifert, *Phys. Rev. X* **12**, 031025 (2022).  
 [21] T. Van Vu and K. Saito, *Phys. Rev. X* **13**, 011013 (2023).  
 [22] A. Ghosal and G. Bisker, *J. Phys. D: Appl. Phys.* **56**, 254001 (2023).  
 [23] D. R. Fredkin and J. A. Rice, *J. Appl. Probab.* **23**, 208 (1986).  
 [24] *Supplemental Material*.  
 [25] R. A. Fisher, *Philosophical transactions of the Royal Society of London. Series A* **222**, 309 (1922).  
 [26] V. Weispfenning, *J. Symb. Comput.* **14**, 1 (1992).  
 [27] J. Monod, J. Wyman, and J.-P. Changeux, *J. Mol. Biol.* **12**, 88 (1965).  
 [28] A. Cornish-Bowden, *Fundamentals of enzyme kinetics* (John Wiley & Sons, 2013).  
 [29] H. Qian, *Annu. Rev. Phys. Chem.* **58**, 113 (2007).  
 [30] C. Jia and R. Grima, *Phys. Rev. X* **11**, 021032 (2021).  
 [31] C. Jia and Y. Li, *SIAM J. Appl. Math.* **83**, 1572 (2023).  
 [32] M. Qian, M.-P. Qian, and X.-J. Zhang, *Phys. Lett. A* **309**, 371 (2003).