The periodic structure of local consistency^{*}

Lorenzo Ciardo University of Oxford lorenzo.ciardo@cs.ox.ac.uk Stanislav Živný University of Oxford standa.zivny@cs.ox.ac.uk

1st July 2024

Abstract

We connect the mixing behaviour of random walks over a graph to the power of the local-consistency algorithm for the solution of the corresponding constraint satisfaction problem (CSP). We extend this connection to arbitrary CSPs and their promise variant. In this way, we establish a linear-level (and, thus, optimal) lower bound against the local-consistency algorithm applied to the class of aperiodic promise CSPs. The proof is based on a combination of the probabilistic method for random Erdős-Rényi hypergraphs and a structural result on the number of fibers (i.e., long chains of hyperedges) in sparse hypergraphs of large girth. As a corollary, we completely classify the power of local consistency for the approximate graph homomorphism problem by establishing that, in the nontrivial cases, the problem has linear width.

1 Introduction

The algebraic characterisation of the power of local consistency [6,14] was one of the fundamental milestones of the algebraic theory of constraint satisfaction problems (CSPs) [15,36], and a key step towards the positive resolution of Feder–Vardi's Dichotomy Conjecture [29] obtained, independently, by Bulatov [16] and Zhuk [54]. Fix a relational structure **A** (consisting of a finite domain A and a finite list of relations over A). The constraint satisfaction problem parameterised by **A** (in short, CSP(**A**)) is the following computational problem: Given an *instance* structure **X** that admits a homomorphism (i.e., a relation-preserving map) to **A**, find an explicit such homomorphism.¹ Suitably choosing the *template* **A** allows for casting as a CSP many diverse problems, including graph colouring (if **A** is a clique \mathbf{K}_c), graph homomorphism (if **A** is an undirected graph), linear equations over a finite field (if **A** encodes the affine spaces over the field), and several variants of the Boolean satisfiability problem such as HORN SAT, k-IN-n SAT, and NOT-ALL-EQUAL SAT.

Enforcing local consistency is one of the — if not the — main algorithmic approaches developed for solving CSPs.² The κ -th level of the local-consistency algorithm searches a system of locally compatible homomorphisms from substructures of **X** of size up to κ to **A**.

^{*}This work was supported by UKRI EP/X024431/1. For the purpose of Open Access, the authors have applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission. All data is provided in full in the results section of this paper.

¹We denote the presence of a homomorphism from **X** to **A** by the expression $\mathbf{X} \to \mathbf{A}$.

²Another approach is based around the idea of finding a small generating basis of the solution space (generalising Gaussian elimination) [18,35]; there are also combinations of the two approaches, cf. [13,17,24,27].

We call such system a κ -strategy. The width of CSP(**A**) is then the minimum κ for which a κ -strategy can always be extended to a global solution. If κ is a constant, the κ -th level of the algorithm runs in time polynomial in the size of the instance. Hence, any CSP having bounded width is in P.³ Both Bulatov's [16] and Zhuk's [54] algorithms use local-consistency checking as a subroutine. Indeed, the ability to capture the power of local consistency [6,14] via tame congruence theory [33] was a key step towards the resolution of the Dichotomy Conjecture. In particular, in [42], it was established that the class of bounded-width CSPs admits a natural characterisation in terms of universal-algebraic objects known as polymorphisms — i.e., homomorphisms $f: \mathbf{A}^n \to \mathbf{A}$, where \mathbf{A}^n is the *n*-th direct power of \mathbf{A} . More precisely, it was shown that CSP(\mathbf{A}) has bounded width precisely when it admits weak near-unanimity (WNU) polymorphisms of all arities — where f is a WNU operation when it satisfies the identities $f(a, b, b, \dots, b, b) = f(b, a, b, \dots, b, b) = \dots = f(b, b, b, \dots, a, b) = f(b, b, b, \dots, b, a)$.

Despite its success in the realm of CSPs, the algebraic description of the power of local consistency has a limitation — it does not extend to the *promise* variant of CSPs. This can be formulated as follows: Fix two structures A and B such that $A \rightarrow B$; given some structure **X** such that $\mathbf{X} \to \mathbf{A}$ as input, find an explicit homomorphism from **X** to **B**. Here, **A** and **B** should be interpreted as the *strong* and the *weak* versions of the template, respectively. Hence, the promise CSP parameterised by \mathbf{A} and \mathbf{B} (in short, $PCSP(\mathbf{A}, \mathbf{B})$) asks to find an explicit, weakly satisfying assignment for an instance that is promised to be strongly satisfiable; in this sense, it captures approximability of the CSP [3, 5, 12]. Many fundamental computational questions that are inexpressible as standard CSPs admit a formulation as PCSPs. Primary examples are the approximate graph colouring problem and its natural generalisation, the approximate graph homomorphism problem. The former corresponds to letting \mathbf{A} and \mathbf{B} be two cliques \mathbf{K}_c and \mathbf{K}_d , respectively, with $c \leq d$; the latter corresponds to letting \mathbf{A} and \mathbf{B} be two undirected graphs. The complexity of both these problems is notoriously open. This is in striking contrast with their non-promise variants — the graph colouring and the graph homomorphism problems — whose complexity was classified already in 1972 by Karp [37] and in 1990 by Hell–Nešetřil [32], respectively.

Part of the algebraic machinery developed in the decades of research on CSPs can be transferred to PCSPs. In particular, polymorphisms have a natural promise version (homomorphisms $\mathbf{A}^n \to \mathbf{B}$) and, just like for CSPs, they completely determine the complexity of PCSPs, as established in [5, 12]. Yet, the CSP tools appear to lose a substantial part of their sharpness when they are adapted to the promise world. The ultimate reason for this weakening is that the common, universal-algebraic core underpinning many distinct CSP phenomena is less structured for PCSPs. In particular, PCSP polymorphisms do not compose, which makes it impossible to obtain new polymorphic identities from old ones. A clear example of this phenomenon is the break of the "WNU – local consistency" bond: While the local-consistency algorithm can be applied to solve PCSPs and, thus, the notion of width is well defined in the promise setting, the presence of WNU polymorphisms of all arities does not guarantee bounded width for PCSPs, as established in [1]. For example, the PCSP parameterised by the structures encoding 2-IN-4 SAT and NOT-ALL-EQUAL SAT has WNU polymorphisms of all arities although its width is unbounded — and, in fact, linear; see later.⁴ In particular, this

³For CSPs, the notion of bounded width admits various equivalent descriptions, in terms of expressibility via Datalog programs [40], pebble games [29], tree-width duality [19], and robust tractability [7], as well as solvability via different algorithmic techniques including linear programming [50] and semidefinite programming [2, 51].

⁴Unless P=NP, WNU polymorphisms do not even guarantee polynomial-time tractability (via any algorithm): For instance, PCSP($\mathbf{K}_c, \mathbf{K}_{2c}$) has WNUs of all arities; yet, it was shown in [43] that the problem is NP-hard

makes it infeasible to investigate the width of promise problems such as approximate graph homomorphism using the polymorphic tools currently available.

Contributions We adopt a different, *structural* approach to study the power of local consistency. To illustrate it, we start with a simple example — $\text{CSP}(\mathbf{C}_n)$, where \mathbf{C}_n is the undirected cycle of length n. In this case, the parity of n determines the width of the problem: It is well known and not hard to check that $\text{CSP}(\mathbf{C}_n)$ has width 3 if n is even, while it has unbounded width if n is odd. Can we link these two different width behaviours to a separating *structural* property depending on the parity of the cycle length?

Consider the setting of a random walker starting from some vertex v_0 of \mathbf{C}_n and moving, at each discrete time step, to a vertex adjacent to her current location, picking the left or the right one with probability $\frac{1}{2}$, independently. After a number τ of time steps, suppose that the walker ends up on vertex v_{τ} . How much information on the initial position v_0 can she deduce from knowing the final position v_{τ} as well as the elapsed time τ ? If n is odd, all information gets lost after a sufficiently long time. Indeed, in this case, any vertex can be reached from any other vertex in precisely τ steps, for any $\tau \geq n-1$. On the other hand, if n is even, while the initial vertex cannot be determined with certainty, its parity may be recovered by summing v_{τ} and τ (mod 2). This behaviour of random walks is known as *periodicity*:⁵ Even cycles are periodic, odd cycles are aperiodic. Thus, in the very special case of CSPs over undirected cycles, the different width of odd and even cycles corresponds to the periodicity property of the corresponding random walk.

The main conceptual contribution of this work is establishing that the power of local consistency is deeply linked to the periodic behaviour of the random walk described above. Moreover, this connection holds in a surprisingly broader context — well beyond the state-of-the-art knowledge on the complexity of the corresponding computational problems. For undirected graphs, aperiodicity corresponds to the existence of some integer τ (the mixing time) such that any two vertices of the graph are connected by a walk of length τ .⁶ In the case of directed graphs (digraphs), the definition is similar, the only difference being that the presence of both a directed and an alternating walks of length τ is required. In both cases, aperiodicity can be alternatively, naturally expressed in terms of the matrix-theoretic properties of primitivity and irreducibility (or ergodicity) applied to the corresponding adjacency matrices. Considering walks (and orientations thereof) in hypergraphs, the notion of aperiodicity is naturally extended to arbitrary relational structures — thus covering the whole class of CSPs. Lastly, and most significantly, we lift the link between aperiodicity and width to the promise regime. Hence, unlike the WNU polymorphic description, this approach is applicable to PCSPs.

We have seen that a (P)CSP has bounded width if it is solved by a constant level of the local-consistency algorithm — in which case, the running time of the algorithm is polynomial in the size of the instance. (P)CSPs of *linear* width lie at the opposite end of the width spectrum: Not even ϵn levels of the local-consistency algorithm are enough to solve these

for $c \ge 6$.

⁵See Remark 22 for a further discussion on this point.

⁶In the literature on Markov chains, there exist several different formalisations of the concept of "mixing time". One primary example is *Kemeny's constant* — the expected number of time steps needed to connect two vertices, independently sampled according to the stationary distribution of the chain [38,45]. The definition used in this work is a combinatorial version of the latter (generalised to arbitrary relational structures), in that it minimises the number of steps needed for connecting *every* pair of vertices; see Definition 3.

problems (where n is the size of the instance and ϵ is a positive constant).⁷ Classic examples of CSPs of linear width are systems of linear equations [2,29] (see also [20,30,52] for other lower bounds on systems of linear equations). Our main result identifies aperiodicity as a sufficient condition for having linear — and, thus, unbounded — width.⁸

Theorem 1. Let $\mathbf{A} \to \mathbf{B}$ be relational structures such that \mathbf{A} is aperiodic and \mathbf{B} is loopless.⁹ Then, PCSP(\mathbf{A}, \mathbf{B}) has linear width.¹⁰

Observe that n levels of the local-consistency algorithm always suffice for instances of size n. Hence, the linear-level lower bound in Theorem 1 is tight (up to optimising the constant ϵ). An immediate consequence of the result above is the complete characterisation of the power of local consistency for the approximate graph homomorphism problem.

Corollary 2. Let $\mathbf{A} \to \mathbf{B}$ be undirected graphs, and suppose that \mathbf{A} is non-bipartite and \mathbf{B} is loopless. Then, $PCSP(\mathbf{A}, \mathbf{B})$ has linear width.¹¹

Related work Our work is inspired by the recent paper [1] by Atserias–Dalmau, which gives a sufficient condition for a PCSP to have linear width.¹² In the language of the current paper, that condition requires that any pair of vertices should be connected by oriented walks of length 2; thus, it corresponds to the case $\tau = 2$ of our aperiodicity notion. Atserias–Dalmau's condition is satisfied, in particular, by cliques, thus resulting in a linear-level lower bound against local consistency for the solution of approximate graph *colouring*. Extending the lower bound to approximate graph *homomorphism* requires dealing with graphs having arbitrarily high mixing time and, thus, it needs the full power of our Theorem 1. Some of the machinery of [1] is used in this paper as a black box (in particular, a result on the sparsity of random hypergraphs obtained via Chernoff bounds). However, the transition from $\tau = 2$ to arbitrarily high τ requires rather different proof ideas. In particular, the concept of τ -fibrosity that we introduce in this work to study the number of long, isolated chains of hyperedges in sparse hypergraphs turns out to behave differently for $\tau \leq 3$ and for $\tau \geq 4$: In the latter case, large fibrosity is not implied by sparsity alone, and bounds on the length of the shortest cycles in hypergraphs need to be established via a probabilistic analysis.

A recent line of work established lower bounds against relaxations of approximate graph colouring [22,23] and homomorphism [25] based on linear and semidefinite programming as well as linear-equation solvers. NP-hardness of the former problem was proved in [31] conditional to Khot's *d*-to-1 Conjecture [39]. Brakensiek–Guruswami conjectured that approximate graph homomorphism should be tractable in polynomial time if **A** is bipartite or **B** has a loop, and

⁷In particular, every PCSP of linear width has unbounded width. For non-promise CSPs, the converse also holds: Every CSP of unbounded width has linear width [2]. For PCSPs, the analogous statement is not known to hold.

⁸Put differently, Theorem 1 gives a necessary condition for not having linear width, and thus in particular a necessary condition for having bounded width.

⁹A structure **B** contains a *loop* if there exists an element $b \in B$ such that every relation of **B** contains the constant tuple (b, \ldots, b) of suitable arity. Otherwise, we say that **B** is *loopless*.

¹⁰Our result holds both for the *search* version of $PCSP(\mathbf{A}, \mathbf{B})$ defined above and for its *decision* version — distinguish whether an instance **X** is homomorphic to **A** or is not even homomorphic to **B**.

¹¹It is well known that $PCSP(\mathbf{A}, \mathbf{B})$ has bounded width if \mathbf{A} is a bipartite graph or \mathbf{B} has a loop [29]. Note that, in the statement of Corollary 2, we are implicitly identifying an undirected graph with a symmetric digraph containing both orientations of each edge.

¹²While the result in [1] is stated in terms of *sublinear*, as opposed to linear, width, the proof yields a linear-level, optimal lower bound.

NP-hard otherwise [12]. Therefore, our Corollary 2 is consistent with — and supports — the conjectured complexity dichotomy for this problem.

Notation We let \mathbb{N} denote the set of positive integer numbers. For $n \in \mathbb{N}$, we let $[n] = \{1, \ldots, n\}$, while $[0] = \emptyset$. We usually write tuples in boldface, and their entries in normal font; for example, the *i*-th entry of a tuple **a** shall be denoted by a_i . Also, we denote by set(**a**) the set of all entries of **a**. Given tuples **a**, **b**, **c**, ..., we let $\langle \mathbf{a}, \mathbf{b}, \mathbf{c}, \ldots \rangle$ denote their concatenation. Let σ be a signature; i.e., a finite list of relation symbols R, each with an assigned arity $\operatorname{ar}(R) \in \mathbb{N}$. A relational structure **A** over the signature σ consists of a finite, nonempty domain A and a relation $R^{\mathbf{A}} \subseteq A^{\operatorname{ar}(R)}$ for each symbol $R \in \sigma$. We denote by $n_{\mathbf{A}}$ the size of the domain of **A**; i.e., $n_{\mathbf{A}} = |A|$. Given two structures **A** and **A'**, both over the signature σ , a homomorphism from **A** to **A'** is a function $f : A \to A'$ such that, for each $R \in \sigma$ and each $\mathbf{a} \in R^{\mathbf{A}}$, $f(\mathbf{a}) \in R^{\mathbf{A'}}$ (where $f(\mathbf{a}) = (f(a_1), \ldots, f(a_{\operatorname{ar}(R)}))$) is the entrywise application of f to the entries of **a**).

2 Overview of results and techniques

To prove Theorem 1, we follow the intuition that the power of the local-consistency algorithm corresponds to the amount of information retained in a random walk.¹³ Suppose that we have two structures \mathbf{X} (the *instance*) and \mathbf{A} (the *template*), and assume, for simplicity, that they are digraphs. The goal is to show that some high level κ of the local-consistency algorithm applied to X and A accepts (in symbols, $LC^{\kappa}(X, A) = YES^{14}$) but $X \not\rightarrow B$. This would mean that \mathbf{X} is a *fooling instance* for the algorithm applied to the decision version of PCSP(\mathbf{A}, \mathbf{B}); as the latter reduces to the search version [5], this would be sufficient to prove Theorem 1 for both versions. Consider two random walkers, Xerxes and Alice, walking on \mathbf{X} and on \mathbf{A} , respectively. Let x_0, x_1, x_2, \ldots and a_0, a_1, a_2, \ldots be the two sequences of vertices resulting from their walks. In order for the algorithm to accept, we want that the assignment $x_i \mapsto a_i$ should locally look like a homomorphism. As discussed in the Introduction, if **A** is aperiodic, Alice has a limited memory: She is not able to retain any local information on her walk after some number τ of steps. This can be exploited by Xerxes. Suppose that X contains a path of length τ isolated from the rest of the structure. Xerxes enters the path at some time t and reaches the other end at time $t + \tau$. By that time, Alice will have no memory of her location at time t. Hence, she will not be able to spot any inconsistency involving the vertices of the path. If many of these deceptive paths exist in **X**, chances are that Xerxes will always manage to make his walk locally consistent with Alice's. Now, the longer time τ is needed for Alice to lose here memory, the longer Xerxes' paths need to be, in order to be effective. The risk, of course, is that if too many long paths are required, it could become easy to build a global homomorphism from \mathbf{X} to any other structure \mathbf{B} , thus preventing it from being a fooling instance.

Our proof of Theorem 1 consists of four stages, discussed in the next four subsections of this overview. The first stage (*aperiodicity*) concerns the template; the second and third stages (*fibrosity* and *sparsity*) concern the instance; the final stage (*consistency*) concerns both the instance and the template. Further details and complete proofs are given in the body of the

¹³The informal description below should be compared to the *pebble-game* interpretation of local consistency [2]. ¹⁴The formal definition of the algorithm is given in Section 3.

paper. The last subsection of this overview contains a discussion on some consequences of our main result.

2.1 Aperiodicity

First of all, we need to formally define aperiodicity for arbitrary relational structures. If the structure has a unique, binary, symmetric relation (i.e., it is a graph), aperiodicity means that any pair of vertices is connected by a walk of some fixed length. If the symmetry constraint is relaxed (thus resulting in a digraph), we require the presence of both a directed and an alternating walks of fixed length. When lifting the notion of aperiodicity to arbitrary structures, it is necessary to extend the definition of a walk, and decide what it means for it to be "directed" or "alternating". We first consider the case that **A** is *monic* — i.e., its signature consists of a unique symbol.

Definition 3 (Aperiodicity, monic case). Let \mathbf{A} be a monic relational structure and let R be the unique symbol in its signature. Given $\tau \in \mathbb{N}$, we let a τ -pattern λ be a $\tau \times 2$ matrix having entries in [ar(R)] such that the two entries in each row are different. Fix a τ -pattern λ , and denote its entries by $\lambda_{i,j}$. A λ -walk in \mathbf{A} consists of a list a_0, \ldots, a_{τ} of vertices in A and a list $\mathbf{a}^{(1)}, \ldots, \mathbf{a}^{(\tau)}$ of tuples in $R^{\mathbf{A}}$ satisfying the following requirement: For each $i \in [\tau]$, $a_{\lambda_{i,1}}^{(i)} = a_{i-1}$ and $a_{\lambda_{i,2}}^{(i)} = a_i$. In this case, we say that the λ -walk connects a_0 to a_{τ} . We say that \mathbf{A} is aperiodic if there exists $\tau \in \mathbb{N}$ such that, for each $a, b \in A$ and each τ -pattern λ , there exists a λ -walk connecting a to b.¹⁵ In this case, the mixing time of \mathbf{A} is the minimum $\tau \in \mathbb{N}$ for which the condition holds.

In the definition above, a λ -walk should be viewed as an "oriented" walk, where the orientation is prescribed by the τ -pattern λ . If **A** is not monic, we turn it into a monic structure \mathbf{A}^{mon} by taking the product of all relations in **A**. Formally, let $\sigma = \{R_1, \ldots, R_\ell\}$ be the signature of **A**, and consider the signature $\tilde{\sigma}$ having a unique symbol of arity $\operatorname{ar}(R_1) + \ldots + \operatorname{ar}(R_\ell)$. We let \mathbf{A}^{mon} be the monic structure over the signature $\tilde{\sigma}$, whose domain is A and whose unique relation is the set $\{\langle \mathbf{a}^{(1)}, \ldots, \mathbf{a}^{(\ell)} \rangle : \mathbf{a}^{(i)} \in R_i^{\mathbf{A}} \,\forall i \in [\ell]\}$.

Definition 4 (Aperiodicity, general case). A relational structure **A** is *aperiodic* if \mathbf{A}^{mon} is aperiodic; in this case, the *mixing time* of **A** is defined to be the mixing time of \mathbf{A}^{mon} .¹⁶

If **A** is a digraph, each row of a τ -pattern λ is either the pair (1, 2) (the "forward" direction) or the pair (2, 1) (the "backward" direction). Thus, in this case, Definition 3 asks for the existence of some τ such that every two vertices a, b are connected by a walk of length τ , oriented in *every* possible way. It turns out that considering two orientations only ("directed" and "alternating") is enough — thus recovering the definition encountered in the Introduction for the digraph case. In fact, via matrix-theoretic results (in particular, a classical theorem by Wielandt on the maximum index of primitivity of primitive matrices), we can make this condition even weaker. This also allows upper bounding the mixing time of an aperiodic digraph.

Theorem 5. Let A be a digraph. The following are equivalent:

 $^{^{15}}$ To avoid cumbersome assumptions in the statements of our results, it shall be convenient to consider a monic structure having an empty relation of arity 1 non-aperiodic.

¹⁶As mentioned in the Introduction, the condition for linear width in [1, Thm. 3] is equivalent to the structure being aperiodic with mixing time 2.

- (*i*) **A** is aperiodic;
- (ii) there exists $t \in \mathbb{N}$ such that, for each $a, b \in A$, there exist a directed walk and an alternating walk from a to b, both of length t;
- (iii) for each $a, b \in A$, there exist a directed walk of length $n_{\mathbf{A}}^2 2n_{\mathbf{A}} + 2$ and an alternating walk of some even length from a to b.

If any of the equivalent conditions above holds, the mixing time of A is at most $n_{\rm A}^4 - 2n_{\rm A}^3 + 2n_{\rm A}^2$.

We proceed with the proof of the main theorem, whose statement, having formally introduced aperiodicity, is now well defined.

2.2 Fibrosity

As we have seen — at least at the intuitive level — the key to making an instance \mathbf{X} accepted by the local-consistency algorithm applied to an aperiodic template \mathbf{A} is to require the presence of long paths (which we call *fibers*) in \mathbf{X} , isolated from the rest of the instance (except for the endpoints). In particular, we shall require linearly many long fibers in all portions of the instance of some bounded size. As illustrated by the game of Xerxes and Alice, the length of these fibers should be at least equal to the mixing time of \mathbf{A} . If, in addition to being highly fibrous, the instance has a large chromatic number, it is a good candidate for fooling local consistency. Instead of working with relational structures, it shall be easier, for now, to deal with hypergraphs. Our strategy for achieving the goal is to split it into two subtasks: First, we show that a hypergraph meets the necessary fibrosity requirements provided that it is sparse and has large girth. Then, we prove that sparse hypergraphs having large girth and chromatic number do exist. In this subsection, we discuss the first of the two subtasks.

For an integer $r \geq 2$, an *r*-uniform hypergraph $\mathcal{H} = (V(\mathcal{H}), E(\mathcal{H}))$ consists of a nonempty set $V(\mathcal{H})$ of vertices and a set $E(\mathcal{H})$ of hyperedges, where a hyperedge is a set of vertices of cardinality exactly r. Henceforth, we shall let r be fixed, and we shall refer to r-uniform hypergraphs simply as hypergraphs. We denote the number of vertices and the number of hyperedges of \mathcal{H} by $n_{\mathcal{H}}$ and $m_{\mathcal{H}}$, respectively. Two hyperedges are *adjacent* if they have a nonempty intersection; a vertex is *incident* to a hyperedge if it belongs to it. For $v \in V(\mathcal{H})$, $\deg_{\mathcal{H}}(v)$ is the *degree* of v; i.e., the number of hyperedges of \mathcal{H} to which v is incident. We say that a hyperedge e is a *link* if exactly two vertices in e have degree 2, and all of the other r-2vertices have degree 1. A set f of distinct links is called a *fiber* if the elements of f can be labelled as e_1, \ldots, e_{τ} , for some $\tau \in \mathbb{N}$, in a way that e_i is adjacent to e_{i+1} for each $i \in [\tau - 1]$. If we need to stress that f has cardinality τ , we refer to it as a τ -fiber. The τ -fibersity of \mathcal{H} , which we denote by $\varphi_{\mathcal{H}}^{\tau}$, is the maximum cardinality of a set of mutually disjoint τ -fibers in \mathcal{H} .

How to enforce large fibrosity in a hypergraph? Intuitively, if a hypergraph is dense, it is unlikely to contain many long fibers. Hence, we shall require that the hypergraph should be sparse. More precisely, given a real number $\beta > 1$, we say that \mathcal{H} is β -sparse if $m_{\mathcal{H}} < \frac{\beta}{r-1}n_{\mathcal{H}}$. Furthermore, we say that \mathcal{H} is *hereditarily* β -sparse if every subhypergraph of \mathcal{H} (including \mathcal{H} itself) is β -sparse. Unfortunately, sparsity is not enough for guaranteeing large fibrosity, even in the binary case. For example, the τ -fibrosity of a star (i.e., a tree having all but one vertex of degree 1) is 0 while, like all acyclic graphs, stars are hereditarily β -sparse for any $\beta > 1$. Given a hyperedge $e \in E(\mathcal{H})$, we say that e is *pendent* if at most one vertex of e has degree at least 2; the *pendency* of \mathcal{H} , denoted by $\pi_{\mathcal{H}}$, is the number of pendent hyperedges in \mathcal{H} . In the star case, the obstacle to large fibrosity is the presence of (linearly) many pendent edges. As we shall see, these do not affect acceptance by the local-consistency algorithm. This suggests that the quantity to be lower-bounded should be the sum of the τ -fibrosity and the pendency. While sparsity is enough for guaranteeing large τ -fibrosity + pendency if $\tau \leq 3$, for larger values of τ it turns out to be insufficient.¹⁷ For example, the graph obtained as the disjoint union of many triangles is hereditarily β -sparse for any $\beta > 1$, but it has no pendent edges nor τ -fibers for any $\tau > 3$ — while it has linearly many mutually disjoint 1-fibers, 2-fibers, and 3-fibers. In this specific case, the problem is generated by the presence of short cycles — which we now define in the hypergraph case, following [9]. For $2 \leq \ell \in \mathbb{N}$, a (Berge) cycle of length ℓ in \mathcal{H} is a sequence $v_0, e_0, v_1, e_1, \ldots, v_{\ell-1}, e_{\ell-1}, v_0$ such that $v_i \in V(\mathcal{H})$ and $e_i \in E(\mathcal{H})$ for each i, all v_i 's and e_i 's are distinct, and $v_i \in e_{i-1} \cap e_i \pmod{\ell}$. The girth of \mathcal{H} (in symbols, gir(\mathcal{H})) is the smallest length of a cycle in \mathcal{H} . Note that two distinct hyperedges of a hypergraph of girth at least 3 cannot intersect in more than one vertices, as this would create a cycle of length 2.

The main result of this subsection is that short cycles are in fact the only obstacle: Sparsity and large girth are enough to guarantee linear fibrosity + pendency.

Theorem 6. For $\tau \in \mathbb{N}$ and $1 < \beta \in \mathbb{R}$, let \mathcal{H} be a hereditarily β -sparse hypergraph of girth at least τ having no isolated vertices.¹⁸ Then, $\varphi_{\mathcal{H}}^{\tau} + \pi_{\mathcal{H}} > \left(\frac{1}{10r\tau} - \beta + 1\right) n_{\mathcal{H}}$.

If $\beta < 1 + \frac{1}{10r\tau}$, the inequality above yields a linear lower bound on the sum of τ -fibrosity and pendency. We stress that this condition on β involves r and τ , which are fixed constants of the template, but is independent of the number of vertices in \mathcal{H} . As will become clear later, this is the first of two crucial details that are needed in order to make the instance able to fool linear — as opposed to only constant — levels of the local-consistency algorithm.

In order to prove Theorem 6, it shall be useful to introduce a more expressive nomenclature for fibers. Given a fiber f in \mathcal{H} , let \mathcal{H}_f be the subhypergraph of \mathcal{H} induced by the vertices contained in the hyperedges of f. We say that f is *degenerate* if \mathcal{H}_f is a cycle; otherwise, we say that f is *non-degenerate*. We say that a fiber f is *maximal* if it is not properly included in any other fiber. We denote by $\varphi_{\mathcal{H}}^{\max}$ the number of maximal fibers of \mathcal{H} . Also, we denote by $\lambda_{\mathcal{H}}$ the number of links of \mathcal{H} . Notice that two distinct maximal fibers are necessarily disjoint; otherwise, their union would yield a longer fiber. Hence, the set of maximal fibers partitions the set of links. The proof of Theorem 6 is obtained by showing that hereditarily sparse hypergraphs have a small number of maximal fibers (provided that no fiber is degenerate), while having a large number of links, as stated in the two results below.

Proposition 7. For $\beta > 1$, let \mathcal{H} be a hereditarily β -sparse hypergraph all of whose fibers are non-degenerate. Then, $\varphi_{\mathcal{H}}^{\max} < 3(\beta - 1)n_{\mathcal{H}} + 3\pi_{\mathcal{H}}$.

Proposition 8. For $\beta > 1$, let \mathcal{H} be a hereditarily β -sparse hypergraph having no isolated vertices. Then, $\lambda_{\mathcal{H}} > (\frac{1}{r} + 6 - 6\beta) n_{\mathcal{H}} - 7\pi_{\mathcal{H}}$.

The numbers of links, maximal fibers, and τ -fibers is constrained by the following relation.

Proposition 9. For any hypergraph \mathcal{H} and any $\tau \in \mathbb{N}$, $\varphi_{\mathcal{H}}^{\tau} + \varphi_{\mathcal{H}}^{\max} > \frac{\lambda_{\mathcal{H}}}{\tau}$.

Then, the claimed lower bound on τ -fibrosity + pendency easily follows.

Proof of Theorem 6. Observe that τ -fibrosity and pendency are additive with respect to disjoint unions; i.e., if \mathcal{H} and \mathcal{H}' are disjoint hypergraphs, $\varphi_{\mathcal{H}\cup\mathcal{H}'}^{\tau} = \varphi_{\mathcal{H}}^{\tau} + \varphi_{\mathcal{H}'}^{\tau}$ and $\pi_{\mathcal{H}\cup\mathcal{H}'} = \varphi_{\mathcal{H}}^{\tau} + \varphi_{\mathcal{H}'}^{\tau}$

¹⁷We point out that this issue does not occur in [1], which corresponds to the case $\tau = 2$ of our result.

 $^{^{18}\}mathrm{An}$ isolated vertex is a vertex of degree 0.

 $\pi_{\mathcal{H}} + \pi_{\mathcal{H}'}$. Clearly, the same holds for the number $n_{\mathcal{H}}$ of vertices. Hence, we can assume without loss of generality that \mathcal{H} is connected. Suppose first that all fibers of \mathcal{H} are non-degenerate. The combination of Propositions 7, 8, and 9 yields

$$\varphi_{\mathcal{H}}^{\tau} > \frac{\lambda_{\mathcal{H}}}{\tau} - \varphi_{\mathcal{H}}^{\max} > \left(\frac{1}{r\tau} + \frac{6}{\tau} - \frac{6\beta}{\tau}\right) n_{\mathcal{H}} - \frac{7}{\tau}\pi_{\mathcal{H}} - 3(\beta - 1)n_{\mathcal{H}} - 3\pi_{\mathcal{H}}$$
$$= \left(\frac{1}{r\tau} + \frac{6}{\tau} - \frac{6\beta}{\tau} - 3(\beta - 1)\right) n_{\mathcal{H}} - \left(\frac{7}{\tau} + 3\right) \pi_{\mathcal{H}} > \left(\frac{1}{r\tau} - 10\beta + 10\right) n_{\mathcal{H}} - 10\pi_{\mathcal{H}},$$

whence it follows that

$$\varphi_{\mathcal{H}}^{\tau} + \pi_{\mathcal{H}} \ge \frac{1}{10} (\varphi_{\mathcal{H}}^{\tau} + 10\pi_{\mathcal{H}}) > \left(\frac{1}{10r\tau} - \beta + 1\right) n_{\mathcal{H}}$$

as required. Suppose now that at least one fiber of \mathcal{H} is degenerate. Since \mathcal{H} is connected and has girth at least τ , this implies that \mathcal{H} is a cycle of length at least τ . Hence, $\tau \leq m_{\mathcal{H}} = \frac{n_{\mathcal{H}}}{r-1}$. By decomposing the cycle into mutually disjoint τ -fibers, we deduce that, in this case,

$$\varphi_{\mathcal{H}}^{\tau} + \pi_{\mathcal{H}} = \varphi_{\mathcal{H}}^{\tau} = \left\lfloor \frac{m_{\mathcal{H}}}{\tau} \right\rfloor > \frac{m_{\mathcal{H}}}{2\tau} = \frac{n_{\mathcal{H}}}{2(r-1)\tau} \ge \frac{n_{\mathcal{H}}}{10r\tau}$$

and the claimed inequality follows since $\beta > 1$.

2.3 Sparsity

We now know how to translate hereditary sparsity and large girth into linear fibrosity + pendency — which, in turn, shall be the key to making the instance locally consistent. The next step is to show that there exist hereditarily sparse hypergraphs of large girth that, at the same time, are highly chromatic. This last property will guarantee that the corresponding instance is a No-instance (i.e., it is not homomorphic to **B**), thus making it able to fool the local-consistency algorithm.

We will need a slightly more general version of sparsity. Given two real numbers $\beta > 1$ and $\gamma > 0$, we say that a hypergraph \mathcal{H} is (γ, β) -threshold-sparse if every subhypergraph \mathcal{H}' of \mathcal{H} such that $m_{\mathcal{H}'} \leq \gamma$ is β -sparse. Observe that \mathcal{H} is hereditarily β -sparse precisely when it is $(m_{\mathcal{H}}, \beta)$ -threshold-sparse. Also, the chromatic number of \mathcal{H} — in symbols, $chr(\mathcal{H})$ — is the smallest integer c for which the vertices of \mathcal{H} can be partitioned into c classes such that every hyperedge intersects at least two classes of the partition. We shall prove the following result.

Theorem 10. Take two positive integer numbers g and h and a real number $\beta > 1$. There exists a positive real number $\delta = \delta(g, h, \beta)$ and a positive integer number $n_0 = n_0(g, h, \beta)$ such that, for each $n \ge n_0$, there exists a hypergraph \mathcal{H} with the following properties:

1. \mathcal{H} has n vertices;	3. $\operatorname{chr}(\mathcal{H}) \ge h;$
2. $\operatorname{gir}(\mathcal{H}) \ge g;$	4. \mathcal{H} is $(\delta n, \beta)$ -threshold-sparse

The proof of Theorem 10 is probabilistic. Specifically, we sample a random (*r*-uniform) Erdős-Rényi hypergraph with *n* vertices and hyperedge probability *p* carefully chosen. For suitable values of *p*, one can show that \mathcal{H} , with a nonzero probability, has the following properties: (*i*) it contains a small number (less than, say, $\frac{n}{2}$) of short cycles; (*ii*) it has a small independence number (less than $\frac{n}{2h}$); and (*iii*) it is threshold-sparse. At this point, to

conclude, one simply needs to break all short cycles and check that the resulting hypergraph meets all the requirements listed in the theorem. The proof of (i) and (ii) goes along the lines of the classic result on the existence of hypergraphs with arbitrarily large girth and chromatic number (obtained in [28] as an extension of the analogous result for graphs). To prove (iii), we use as a black box a sparsity result for random hypergraphs established in [1] via Chernoff bounds (see also [21]).

It should be observed that, while the minimum required girth and chromatic number of the hypergraph obtained via the probabilistic construction above are fixed constants and, thus, can be arbitrarily small compared to the number n of vertices, the first parameter of the threshold-sparsity is required to be linear in n. This is the second crucial detail that allows our argument to push the lower bound of Theorem 1 up to a linear level of consistency. Indeed, unfolding the definition of threshold-sparsity, part 4. of Theorem 10 asks that every subhypergraph of \mathcal{H} with at most δn hyperedges should be (hereditarily) β -sparse. This allows calling Theorem 6 for subhypergraphs of linear size — precisely what is needed in order to obtain consistency up to a linear level.

2.4 Consistency

In this final stage of our proof of Theorem 1, we study the interaction between the template property of Subsection 2.1 and the instance properties of Subsections 2.2 and 2.3. It shall be useful to select a set of one or two distinguished vertices (we call this set a *joint*) from any pendent hyperedge and any fiber of a hypergraph \mathcal{H} . If e is a pendent hyperedge of \mathcal{H} , a set $J = \{v\}$ is a joint of e if $v \in e$ and v has maximum degree among the vertices of e. Let now f be a fiber of \mathcal{H} , and recall that \mathcal{H}_f denotes the subhypergraph of \mathcal{H} induced by the vertices contained in the hyperedges of f. If f is non-degenerate, there exist precisely two vertices $u \neq v$ of \mathcal{H}_f that are incident to edges not in f. In this case, the joint of f is the set $J = \{v\}$ where v is a vertex of \mathcal{H}_f such that $\deg_{\mathcal{H}}(v) = 2$. Notice that joints are not uniquely determined, in general. Given a monic structure \mathbf{X} whose symbol R has arity $r \geq 2$, we say that \mathbf{X} is oriented if $|\operatorname{set}(\mathbf{x})| = r$ for each $\mathbf{x} \in R^{\mathbf{X}}$, and $\operatorname{set}(\mathbf{x}) = \operatorname{set}(\mathbf{x}')$ implies $\mathbf{x} = \mathbf{x}'$ for each $\mathbf{x}, \mathbf{x}' \in R^{\mathbf{X}}$. Given an oriented monic structure \mathbf{X} , we let $\mathbf{X}^{\operatorname{sym}}$ be the (r-uniform) hypergraph having vertex set X and hyperedge set $\{\operatorname{set}(\mathbf{x}) : \mathbf{x} \in R^{\mathbf{X}}\}$.

The next two results argue that, if \mathbf{A} is aperiodic, a partial homomorphism to \mathbf{A} can always by extended to include a fiber or a pendent hyperedge, respectively.¹⁹

Proposition 11. Let \mathbf{X} , \mathbf{A} be monic structures such that \mathbf{X} is oriented and \mathbf{A} is aperiodic with mixing time τ , let f be a τ -fiber of \mathbf{X}^{sym} , let J be a joint of f, and let \mathbf{Y} be the substructure of \mathbf{X} induced by $(X \setminus \bigcup_{e \in f} e) \cup J$. Then, any homomorphism $h : \mathbf{Y} \to \mathbf{A}$ can be extended to a homomorphism $h' : \mathbf{X} \to \mathbf{A}$.

Proposition 12. Let \mathbf{X} , \mathbf{A} be monic structures such that \mathbf{X} is oriented and \mathbf{A} is aperiodic, let e be a pendent hyperedge for \mathbf{X}^{sym} , let J be a joint of e, and let \mathbf{Y} be the substructure of \mathbf{X} induced by $(X \setminus e) \cup J$. Then, any homomorphism $h : \mathbf{Y} \to \mathbf{A}$ can be extended to a homomorphism $h' : \mathbf{X} \to \mathbf{A}$.

When \mathbf{A} is aperiodic and \mathbf{X} is sparse and has a large girth, the two propositions above allow for showing that acceptance by the local-consistency algorithm can be derived from the presence of a suitably large consistency gap, which we define next (following [1]).

¹⁹In the terminology of [1], this means that fibers and pendent hyperedges yield *boundary sets* (see also [47]).

Definition 13. Let **X** and **A** be monic structures, let **Y** be a substructure of **X**, and let X' be a subset of X. We say that a function $f : X' \to A$ is *consistent with* **Y** if there is a homomorphism $h : \mathbf{Y} \to \mathbf{A}$ such that f and h agree on $X' \cap Y$. Moreover, for $\kappa \in \mathbb{N}$ and $\gamma \in \mathbb{R}$, we say that the pair (\mathbf{X}, \mathbf{A}) has a (κ, γ) -consistency gap if, for each substructure **W** of **X** such that $n_{\mathbf{W}} \leq \kappa$ and each homomorphism $f : \mathbf{W} \to \mathbf{A}$, one of the following conditions holds:

- f is consistent with every substructure of X whose relation contains at most γ tuples, or
- f is not consistent with some substructure of **X** whose relation contains at most $\frac{\gamma}{n_{\mathbf{A}}}$ tuples.

Theorem 14. Let **X** and **A** be monic structures, and take two numbers $\kappa \in \mathbb{N}$ and $\gamma \in \mathbb{R}$ such that $\gamma \geq n_{\mathbf{A}}$. Suppose that

- A is aperiodic with mixing time τ ;
- X is oriented, and X^{sym} has girth at least τ and is (γ, β)-threshold-sparse for some real number 1 < β < 1 + ¹/_{10rτ};
- (\mathbf{X}, \mathbf{A}) has a (κ, γ) -consistency gap.

Then, $LC^{\kappa}(\mathbf{X}, \mathbf{A}) = YES.$

To finalise the proof of Theorem 1, we just need to put all the pieces together: We construct a sparse, highly chromatic hypergraph \mathcal{H} of large girth via Theorem 10; we use Theorem 6 to argue that all subhypergraphs of \mathcal{H} (up to linear size) have linear fibrosity + pendency; we turn \mathcal{H} into an oriented instance **X** by choosing an orientation for each hyperedge; we show that the pair (**X**, **A**) has a ($\lfloor \epsilon n \rfloor, \delta n$)-consistency gap; we deduce via Theorem 14 that $\mathrm{LC}^{\lfloor \epsilon n \rfloor}(\mathbf{X}, \mathbf{A}) = \mathrm{YES}$; finally, we use the facts that \mathcal{H} has large chromatic number and **B** is loopless to conclude that **X** fools linear levels of the local-consistency algorithm.

2.5 Epilogue

Theorem 1 gives a sufficient condition for a PCSP to have linear width. In fact, we can effortlessly strengthen it by using the fact that, if $\mathbf{A}' \to \mathbf{A} \to \mathbf{B}$ and $\text{PCSP}(\mathbf{A}', \mathbf{B})$ has linear width, the same is true for $\text{PCSP}(\mathbf{A}, \mathbf{B})$ — as is easily derived from Lemma 29. We obtain the following result.

Corollary 15. Let $\mathbf{A} \to \mathbf{B}$ be relational structures, and suppose that \mathbf{B} is loopless and there exists some aperiodic relational structure \mathbf{A}' such that $\mathbf{A}' \to \mathbf{A}$. Then, $\mathrm{PCSP}(\mathbf{A}, \mathbf{B})$ has linear width.

As an immediate consequence, we can completely characterise the power of local consistency applied to the approximate graph homomorphism problem.

Proof of Corollary 2. Since **A** is non-bipartite, it contains an odd undirected cycle \mathbf{C}_p ; thus, $\mathbf{C}_p \to \mathbf{A}$. The result then follows from Corollary 15 by observing that every odd cycle is aperiodic.

A natural question, at this point, is whether the sufficient condition for linear width in Corollary 15 is also necessary. The answer turns out to be negative, even in the CSP setting. Indeed, it is known that there exists an oriented tree \mathbf{T} whose corresponding CSP has unbounded width [11]. By [2] (see Footnote 7), $\text{CSP}(\mathbf{T})$ has in fact linear width. Suppose that there exists some aperiodic digraph \mathbf{A}' such that $\mathbf{A}' \to \mathbf{T}$. It is not hard to check (see Proposition 25) that this implies the existence of some aperiodic induced substructure \mathbf{T}' of \mathbf{T} . This is a contradiction, as \mathbf{T}' is an oriented forest and, thus, no directed walk in \mathbf{T}' connects a vertex to itself. Another natural question is whether there exist PCSPs having *intermediate* width; i.e., super-constant but nonlinear. In the two cases of non-promise CSPs and binary symmetric PCSPs no such problems exist, as follows from [2] and from Corollary 2, respectively. Does the width dichotomy extend to the entire class of PCSPs?

Finally, we observe that, in the binary case, aperiodicity is formulated in terms of the matrix-theoretic notion of *primitivity* applied to the adjacency matrix of \mathbf{A} (see Section 4). In turn, the nature of primitivity is ultimately spectral: The Perron–Frobenius Theorem implies that an irreducible matrix is primitive precisely when it has a unique eigenvalue having maximum modulus [46]. In other words, the necessary condition for nonlinear width given by the contrapositive of Theorem 1 involves the requirement that the adjacency spectrum of \mathbf{A} should collapse, in the sense that at least two eigenvalues should have the same modulus. Now, a trivial topological consideration shows that the collapse of the spectrum of a square matrix is a *singularity*, in the sense that the spectrum of a random matrix does not collapse with high probability. Resolving a 30-year-old conjecture of Babai, it was recently shown by Tao and Vu [49] that the discrete counterpart of this fact also holds: The adjacency spectrum of a random graph does not collapse with high probability.²⁰ This suggests that, on a high level, PCSPs of nonlinear width should be regarded as a singularity within the class of PCSPs. While making this statement mathematically formal goes beyond the scope of the current paper, we find this phenomenon worth of further consideration.

3 The local-consistency algorithm

In this preliminary section, we formally describe the local-consistency algorithm, following [8,41]. Let **X** and **A** be two relational structures having the same signature σ . We say that a structure **Y** of signature σ is a substructure of **X** if $Y \subseteq X$ and $R^{\mathbf{Y}} \subseteq R^{\mathbf{X}}$ for each $R \in \sigma$. If, in particular, $R^{\mathbf{Y}} = R^{\mathbf{X}} \cap Y^{\operatorname{ar}(R)}$ for each $R \in \sigma$, we say that \mathbf{Y} is an *induced substructure* of \mathbf{X} . (We define subhypergraphs and induced subhypergraphs analogously.) A partial homomorphism from \mathbf{X} to \mathbf{A} is either a homomorphism from some induced substructure of \mathbf{X} to \mathbf{A} , or the empty mapping from \emptyset to A. Take an integer $\kappa \in \mathbb{N}$, and consider a nonempty family \mathscr{F} of partial homomorphisms from X to A. Given a function f, let dom(f) denote its domain. We say that \mathscr{F} is a κ -strategy for **X** and **A** if it is closed under restrictions (i.e., for every $f \in \mathscr{F}$ and $Y \subseteq \text{dom}(f)$, the restriction of f to Y is in \mathscr{F}) and has the extension property up to κ (i.e., for every $f \in \mathscr{F}$ with $|\operatorname{dom}(f)| < \kappa$ and every $x \in X \setminus \operatorname{dom}(f)$, there exists some $f' \in \mathscr{F}$ that extends f and has domain dom $(f) \cup \{x\}$). Testing for the existence of a κ -strategy for **X** and **A** can be performed in time polynomial in $(n_{\mathbf{X}} + n_{\mathbf{A}})^{\kappa}$ through the so-called *local-consistency* algorithm, which starts with all partial homomorphisms from \mathbf{X} to \mathbf{A} with domain size at most κ , and iteratively discards those that do not satisfy the two conditions above, until a fixed point is reached. If the resulting family of partial homomorphisms is nonempty, it must

²⁰Babai's conjecture was motivated by his work with Grigoryev and Mount [4], proving that the graph *isomorphism* problem is in P when restricted to the class of graphs having distinct adjacency eigenvalues. Interestingly, the same condition makes the corresponding *homomorphism* problem consistency-hard, as follows from the current paper.

be a κ -strategy; in this case, we write $LC^{\kappa}(\mathbf{X}, \mathbf{A}) = YES$. Otherwise, we are guaranteed that no κ -strategy exists, and we write $LC^{\kappa}(\mathbf{X}, \mathbf{A}) = NO.^{21}$

Let now A and B be two relational structures such that $A \rightarrow B$. Informally, the width of $PCSP(\mathbf{A}, \mathbf{B})$ measures the power of the local-consistency algorithm for its solution. More precisely, observe that, given an instance \mathbf{X} , if h is a homomorphism from \mathbf{X} to \mathbf{A} , restricting h to all subsets of X of size at most κ yields a proper κ -strategy for X and A. Hence, if $\mathbf{X} \to \mathbf{A}, \operatorname{LC}^{\kappa}(\mathbf{X}, \mathbf{A}) = \operatorname{YES}$. If, on the other hand, $\mathbf{X} \to \mathbf{B}$ whenever $\operatorname{LC}^{\kappa}(\mathbf{X}, \mathbf{A}) = \operatorname{YES}$, the κ -consistency algorithm effectively solves $PCSP(\mathbf{A}, \mathbf{B})$. Given $n \in \mathbb{N}$, let $\kappa(n)$ be the minimum nonnegative integer such that $LC^{\kappa(n)}(\mathbf{X}, \mathbf{A}) = YES$ implies $\mathbf{X} \to \mathbf{B}$ for every instance \mathbf{X} having n vertices. (Note that the minimum is well defined as, if **X** has n vertices, it always holds that $LC^n(\mathbf{X}, \mathbf{A}) = YES$ if and only if $\mathbf{X} \to \mathbf{A}$; hence, in particular, $\kappa(n) \leq n$ for any PCSP.) The resulting function $\kappa : \mathbb{N} \to \mathbb{N} \cup \{0\}$ is called the *width* of PCSP(A, B). Observe that κ is necessarily a non-decreasing function. Indeed, letting **X** be an instance on *n* vertices such that $LC^{\kappa(n)-1}(\mathbf{X}, \mathbf{A}) = YES$ and $\mathbf{X} \not\to \mathbf{B}$, adding to \mathbf{X} any number c of isolated vertices results in an instance \mathbf{X}' on n + c vertices that satisfies $\mathrm{LC}^{\kappa(n)-1}(\mathbf{X}', \mathbf{A}) = \mathrm{YES}$ and $\mathbf{X}' \neq \mathbf{B}$, thus witnessing that $\kappa(n+c) \geq \kappa(n)$. If PCSP(A, B) has bounded width (meaning that $\kappa(n) \leq \kappa^*$ for some fixed κ^* independent of n, the local-consistency algorithm certifies that the problem is solvable in polynomial time. At the opposite extreme, we say that $PCSP(\mathbf{A}, \mathbf{B})$ has *linear* width if there exists a positive constant ϵ for which $\kappa(n) > \epsilon n$ for any $n \in \mathbb{N}$; i.e., for any $n \in \mathbb{N}$ there exists an instance **X** on *n* vertices such that $LC^{\lfloor \epsilon n \rfloor}(\mathbf{X}, \mathbf{A}) = YES$ but $\mathbf{X}\not\rightarrow \mathbf{B}.$

4 Aperiodicity

Let **A** be a digraph and let w consist of a list a_0, \ldots, a_t of vertices of **A** and a list $\mathbf{a}^{(1)}, \ldots, \mathbf{a}^{(t)}$ of directed edges of **A**. If $\mathbf{a}^{(i)} = (a_{i-1}, a_i)$ for each $i \in [t]$, we say that w is a *directed* walk of length t from a_0 to a_t ; if $\mathbf{a}^{(i)} = (a_{i-1}, a_i)$ for each odd $i \in [t]$ and $\mathbf{a}^{(i)} = (a_i, a_{i-1})$ for each even $i \in [t]$, we say that w is an *alternating* walk of length t from a_0 to a_t . The goal of this section is to prove the following result.

Theorem (Theorem 5 restated). Let A be a digraph. The following are equivalent:

- (i) \mathbf{A} is aperiodic;
- (ii) there exists $t \in \mathbb{N}$ such that, for each $a, b \in A$, there exist a directed walk and an alternating walk from a to b, both of length t;
- (iii) for each $a, b \in A$, there exist a directed walk of length $n_{\mathbf{A}}^2 2n_{\mathbf{A}} + 2$ and an alternating walk of some even length from a to b.

If any of the equivalent conditions above holds, the mixing time of **A** is at most $n_{\mathbf{A}}^4 - 2n_{\mathbf{A}}^3 + 2n_{\mathbf{A}}^2$.

The conditions (*ii*) and (*iii*) of Theorem 5 can be conveniently reformulated in terms of the adjacency matrix $\mathscr{A}(\mathbf{A})$ of \mathbf{A} — as we shall see after introducing the necessary terminology. By \mathbf{e}_i , we indicate the *i*-th standard unit vector (whose size shall be clear from the context). We say that a matrix is *positive* (resp. *nonnegative*) if all of its entries are positive (resp. nonnegative) real numbers. A nonnegative square matrix M is *primitive* if M^t is positive for some $t \in \mathbb{N}$, and it is *irreducible* if for each pair (i, j) of indices there exists $t \in \mathbb{N}$ such that

²¹It shall be convenient to let $LC^{0}(\mathbf{X}, \mathbf{A}) = YES$ for each \mathbf{X} and \mathbf{A} .

 $\mathbf{e}_i^{\top} M^t \mathbf{e}_j$ (i.e., the (i, j)-th entry of M^t) is positive. Clearly, any primitive matrix is irreducible, while the converse is false. See also Remark 22 for a more detailed discussion on irreducible and primitive matrices.

For two vertices $a, b \in A$, a directed walk of length t from a to b exists precisely when $\mathbf{e}_a^{\top} \mathscr{A}(\mathbf{A})^t \mathbf{e}_b > 0$. Moreover, if t is even, an alternating walk of length t from a to b exists precisely when $\mathbf{e}_a^{\top} (\mathscr{A}(\mathbf{A}) \mathscr{A}(\mathbf{A})^{\top})^{\frac{t}{2}} \mathbf{e}_b > 0$. Note that multiplying a positive and a primitive matrices yields a primitive matrix; moreover, M^{t+1} is positive if M^t is positive. Thus, condition (ii) in Theorem 5 is equivalent to requiring that both $\mathscr{A}(\mathbf{A})$ and $\mathscr{A}(\mathbf{A}) \mathscr{A}(\mathbf{A})^{\top}$ should be primitive matrices. Moreover, condition (iii) is equivalent to requiring that $\mathscr{A}(\mathbf{A})$ should be primitive with index of primitivity²² at most $n_{\mathbf{A}}^2 - 2n_{\mathbf{A}} + 2$, and $\mathscr{A}(\mathbf{A}) \mathscr{A}(\mathbf{A})^{\top}$ should be irreducible.

We shall make use of the following two matrix-theoretic results. The first is a classic theorem by Wielandt, giving a (sharp) upper bound on the index of primitivity of primitive matrices. The second is folklore; the proof can be found, for example, in [10, Ch. 3].

Theorem 16 ([48,53]). An $n \times n$ nonnegative matrix is primitive if and only if its (n^2-2n+2) -th power is positive.

Theorem 17 ([10]). Any irreducible matrix with positive trace is primitive.

By $\operatorname{supp}(M)$ we denote the *support* of M; i.e., the set of pairs (i, j) of indices such that $\mathbf{e}_i^{\top} M \mathbf{e}_j \neq 0$. Given two matrices M_1 and M_2 of equal size, we write $M_1 \leq M_2$ (resp. $M_1 \leq M_2$, $M_1 \sim M_2$) to indicate that $\operatorname{supp}(M_1) \subseteq \operatorname{supp}(M_2)$ (resp. $\operatorname{supp}(M_1) \subseteq \operatorname{supp}(M_2)$, $\operatorname{supp}(M_1) = \operatorname{supp}(M_2)$). The following fact is trivially proved.

Lemma 18. Let M_1, M_2, M_3, M_4 be nonnegative square matrices, and suppose that $M_1 \leq M_2$ and $M_3 \leq M_4$. Then, $M_1M_3 \leq M_2M_4$.

We denote by **0**, *I*, and *J* the all-zero vector, the identity matrix, and the all-one matrix, respectively; their sizes shall be clear from the context. Given a nonnegative square matrix M, a nonnegative integer t, and a tuple $\mathbf{x} \in \{-1, 1\}^t$ (which we shall refer to as an *indicator tuple*), we consider the matrix $M^{\mathbf{x}} = \prod_{i=1}^{t} N_i$, where $N_i = M$ if $x_i = 1$, and $N_i = M^{\top}$ if $x_i = -1$. If t = 0 (and, thus, \mathbf{x} is the empty tuple), we define $M^{\mathbf{x}} = I$. Recall that the concatenation of the tuples $\mathbf{x}, \mathbf{y}, \mathbf{z}, \ldots$ is denoted by $\langle \mathbf{x}, \mathbf{y}, \mathbf{z}, \ldots \rangle$.

Lemma 19. Let M be an irreducible matrix and let $\mathbf{x}, \mathbf{y}, \mathbf{z}$ be indicator tuples such that $M^{\mathbf{y}} \sim J$. Then, $M^{\langle \mathbf{x}, \mathbf{y}, \mathbf{z} \rangle} \sim J$.

Proof. It easily follows from the definition of an irreducible matrix that no row or column of M is the all-zero vector. We deduce that $M^{\mathbf{w}}\mathbf{e}_i \neq \mathbf{0}$ for any indicator tuple \mathbf{w} and any index i. Take now two indices j, ℓ , and observe that

$$\mathbf{e}_{i}^{\top} M^{\langle \mathbf{x}, \mathbf{y}, \mathbf{z} \rangle} \mathbf{e}_{\ell} = \mathbf{e}_{i}^{\top} M^{\mathbf{x}} M^{\mathbf{y}} M^{\mathbf{z}} \mathbf{e}_{\ell}.$$

Since both $\mathbf{e}_j^{\top} M^{\mathbf{x}}$ and $M^{\mathbf{z}} \mathbf{e}_{\ell}$ are nonzero vectors and, by assumption, $M^{\mathbf{y}} \sim J$, we deduce that $\mathbf{e}_j^{\top} M^{\langle \mathbf{x}, \mathbf{y}, \mathbf{z} \rangle} \mathbf{e}_{\ell} \neq 0$, which concludes the proof.

We say that an indicator tuple is *balanced* if the sum of its entries is 0. Also, we indicate by ω_i the tuple (1, -1, 1, -1, ...) of length *i*. Henceforth in this section, we fix a digraph **A** and let *M* be its $n_{\mathbf{A}} \times n_{\mathbf{A}}$ adjacency matrix.

 $^{^{22}\}mbox{I.e.},$ the smallest t for which the $t\mbox{-th}$ power of the matrix is positive.

Lemma 20. Let A be a digraph satisfying part (iii) of Theorem 5, and let \mathbf{x} be a balanced indicator tuple. Then, $I \leq M^{\mathbf{x}}$.

Proof. We use induction over the length ℓ of \mathbf{x} (which must be even, since \mathbf{x} is balanced). If $\ell = 0$, $M^{\mathbf{x}} = I$ and the result is clear. If $\ell = 2$, \mathbf{x} is either ω_2 or $-\omega_2$. Since M is primitive, it is irreducible, thus it has no all-zero row or column. Hence, the diagonals of the matrices $M^{\omega_2} = MM^{\top}$ and $M^{-\omega_2} = M^{\top}M$ are strictly positive, which means that $I \leq M^{\mathbf{x}}$, as needed. For the inductive step, suppose that $\ell \geq 4$ and notice that \mathbf{x} can be written as $\mathbf{x} = \langle \mathbf{w}, \mathbf{z}, \tilde{\mathbf{w}} \rangle$ with \mathbf{z} balanced of length 2 and $\langle \mathbf{w}, \tilde{\mathbf{w}} \rangle$ balanced of length $\ell - 2$. Using Lemma 18 and the inductive hypothesis, we deduce that

$$I \leq M^{\langle \mathbf{w}, \tilde{\mathbf{w}} \rangle} = M^{\mathbf{w}} M^{\tilde{\mathbf{w}}} = M^{\mathbf{w}} I M^{\tilde{\mathbf{w}}} \leq M^{\mathbf{w}} M^{\mathbf{z}} M^{\tilde{\mathbf{w}}} = M^{\langle \mathbf{w}, \mathbf{z}, \tilde{\mathbf{w}} \rangle} = M^{\mathbf{x}},$$

as needed.

Lemma 21. Let **A** be a digraph satisfying part (iii) of Theorem 5, let \mathbf{x}, \mathbf{y} be indicator tuples, and suppose that \mathbf{y} is balanced and has a positive length. Then, either $J \sim M^{\mathbf{x}}$ or $M^{\mathbf{x}} \leq M^{\langle \mathbf{x}, \mathbf{y} \rangle}$.

Proof. If $J \sim M^{\mathbf{x}}$ there is nothing to prove, so we assume that this is not the case. Note that combining Lemma 18 and Lemma 20 yields $M^{\mathbf{x}} \leq M^{\langle \mathbf{x}, \mathbf{y} \rangle}$, so we only need to exclude that $M^{\mathbf{x}} \sim M^{\langle \mathbf{x}, \mathbf{y} \rangle}$. Assume, for the sake of contradiction, that $M^{\mathbf{x}} \sim M^{\langle \mathbf{x}, \mathbf{y} \rangle}$. By assumption, the matrix MM^{\top} is irreducible. Moreover, applying Lemma 20 to the balanced indicator tuple $\boldsymbol{\omega}_2$ yields $I \leq MM^{\top}$. We can then invoke Theorem 17 to deduce that MM^{\top} is in fact primitive and, thus, $J \sim M^{\boldsymbol{\omega}_t}$ for some $t \in \mathbb{N}$. We now proceed by induction on the length ℓ of \mathbf{y} .

If $\ell = 2$, suppose first that $\mathbf{y} = \boldsymbol{\omega}_2$. Take an even integer z. Since $\boldsymbol{\omega}_z$ is balanced, Lemma 18 and Lemma 20 imply that $M^{\mathbf{x}} \leq M^{\langle \mathbf{x}, \boldsymbol{\omega}_z \rangle}$. Moreover, using the assumption that $M^{\mathbf{x}} \sim M^{\langle \mathbf{x}, \boldsymbol{\omega}_z \rangle}$, a repeated application of Lemma 18 yields $M^{\langle \mathbf{x}, \boldsymbol{\omega}_z \rangle} \leq M^{\mathbf{x}}$. It follows that $M^{\mathbf{x}} \sim M^{\langle \mathbf{x}, \boldsymbol{\omega}_z \rangle}$. Choosing z so that $z \geq t$ and applying Lemma 19 yields

$$J \sim M^{\langle \mathbf{x}, \boldsymbol{\omega}_z \rangle} \sim M^{\mathbf{x}},$$

a contradiction. Similarly, if $\mathbf{y} = -\boldsymbol{\omega}_2$, we find $M^{\mathbf{x}} \sim M^{\langle \mathbf{x}, -1, \boldsymbol{\omega}_z \rangle}$ for any odd z, and the same argument as above yields the contradiction $J \sim M^{\mathbf{x}}$.

For the inductive step, suppose that $\ell \geq 4$. We can write \mathbf{y} as the concatenation $\mathbf{y} = \langle \mathbf{w}, \mathbf{z}, \tilde{\mathbf{w}} \rangle$, where \mathbf{z} is balanced and has length 2, and $\langle \mathbf{w}, \tilde{\mathbf{w}} \rangle$ is balanced and has length $\ell - 2 > 0$. Lemma 18 and Lemma 20 yield $M^{\langle \mathbf{x}, \mathbf{w}, \tilde{\mathbf{w}} \rangle} \leq M^{\langle \mathbf{x}, \mathbf{w}, \mathbf{z}, \tilde{\mathbf{w}} \rangle} = M^{\langle \mathbf{x}, \mathbf{y} \rangle}$, while the inductive hypothesis gives $M^{\mathbf{x}} \leq M^{\langle \mathbf{x}, \mathbf{w}, \tilde{\mathbf{w}} \rangle}$. As a consequence,

$$M^{\mathbf{x}} \triangleleft M^{\langle \mathbf{x}, \mathbf{w}, \widetilde{\mathbf{w}} \rangle} \trianglelefteq M^{\langle \mathbf{x}, \mathbf{y} \rangle} \sim M^{\mathbf{x}},$$

a contradiction.

Given a tuple **x** of length t and two indices $i \leq j \in [t]$, we denote by $\mathbf{x}_{[i:j]}$ the tuple $(x_i, x_{i+1}, \ldots, x_j)$.

Proof of Theorem 5. To prove the implication $(i) \Rightarrow (ii)$, suppose that **A** is aperiodic and let τ be its mixing time. Consider the τ -pattern λ consisting in the $\tau \times 2$ matrix whose first column is identically 1 and whose second column is identically 2. Observe that a λ -walk is

precisely a directed walk of length τ . Similarly, an alternating walk of length τ is the same as a $\tilde{\lambda}$ -walk corresponding to the τ -pattern

$$\tilde{\lambda} = \begin{bmatrix} 1 & 2 \\ 2 & 1 \\ 1 & 2 \\ 2 & 1 \\ \vdots & \vdots \end{bmatrix}$$

Hence, (ii) follows from (i) by the definition of aperiodic structures.

Observe now that the condition (ii) is equivalent to the fact that $M^t \sim J$ and $M^{\omega_t} \sim J$ for some $t \in \mathbb{N}$. By Lemma 19, t can be assumed to be even. Hence, the implication $(ii) \Rightarrow (iii)$ directly follows by applying Theorem 16 to M.

We now establish the implication $(iii) \Rightarrow (i)$. Let $q = n_{\mathbf{A}}^2 - 2n_{\mathbf{A}} + 2$, and observe that (iii) implies that $M^q \sim J$. Let $\tau = n_{\mathbf{A}}^4 - 2n_{\mathbf{A}}^3 + 2n_{\mathbf{A}}^2$. We claim that \mathbf{A} is aperiodic with mixing time at most τ . Given a τ -pattern λ and two vertices $a, b \in A$, observe that a λ -walk connecting a and b exists if and only if the (a, b)-th entry of the matrix $M^{\mathbf{x}}$ is positive, where \mathbf{x} is the indicator tuple obtained by taking the first column of λ and replacing all occurrences of "2" with "-1". Hence, the claim is equivalent to the fact that $J \sim M^{\mathbf{x}}$ for any indicator tuple \mathbf{x} of length τ . Pick one such tuple \mathbf{x} . Define integer numbers $y_0 = 0$ and, for $i \in [\tau]$, $y_i = y_{i-1} + x_i$. Let m (resp. \tilde{m}) be the maximum (resp. the minimum) of the $y'_i s$, and define $\mu = m - \tilde{m}$. We now consider two cases.

Suppose first that $\mu \ge q$. Let $i, j \in \{0, ..., \tau\}$ be such that $y_i = \tilde{m}$ and $y_j = m$. Without loss of generality, we can assume that i < j (otherwise, we proceed analogously reasoning on M^{\top} instead of M). Consider the subtuple $\tilde{\mathbf{x}} = \mathbf{x}_{[i:j]}$, and observe that it can be written as

$$\tilde{\mathbf{x}} = \langle 1, \mathbf{w}_1, 1, \mathbf{w}_2, 1, \mathbf{w}_3, 1, \dots, \mathbf{w}_{\mu-1}, 1 \rangle,$$

where $\mathbf{w}_1, \ldots, \mathbf{w}_{\mu-1}$ are balanced indicator tuples. Using Lemma 18 and Lemma 20, we deduce that

$$M^{\mu} \leq M M^{\mathbf{w}_1} M M^{\mathbf{w}_2} M M^{\mathbf{w}_3} M \dots M^{\mathbf{w}_{\mu-1}} M = M^{\tilde{\mathbf{x}}}.$$

Since $\mu \ge q$ and $J \sim M^q$, we deduce from Lemma 19 that $J \sim M^{\mu} \sim M^{\tilde{\mathbf{x}}} \sim M^{\mathbf{x}}$, as required.

Suppose now that $\mu < q$. Since $\tau = q n_{\mathbf{A}}^2$, by the pigeonhole principle, there exists some integer ℓ such that $\tilde{m} \leq \ell \leq m$ and, letting $K = \{i \in \{0, \ldots, \tau\} : y_i = \ell\}, |K| \geq \frac{\tau+1}{q} > \frac{\tau}{q} = n_{\mathbf{A}}^2$. Label the elements of K as k_1, \ldots, k_z in increasing order (where z = |K|), and consider the subtuples $\mathbf{w}_0 = \mathbf{x}_{[1:k_1]}, \mathbf{w}_i = \mathbf{x}_{[k_i+1:k_{i+1}]}$ for $i \in [z-1]$, and $\mathbf{w}_z = \mathbf{x}_{[k_z+1:\tau]}$. (If $k_1 = 0$, we let \mathbf{w}_0 be the empty tuple; similarly, if $k_z = \tau$, we let \mathbf{w}_z be the empty tuple). Observe that $\mathbf{x} = \langle \mathbf{w}_0, \mathbf{w}_1, \ldots, \mathbf{w}_{z-1}, \mathbf{w}_z \rangle$. Moreover, for each $i \in [z-1], \mathbf{w}_i$ is a balanced indicator tuple of length at least 2. If $J \sim M^{\langle \mathbf{w}_0, \mathbf{w}_1, \ldots, \mathbf{w}_i \rangle}$ for some $i \in \{0, \ldots, z\}$, we can conclude through Lemma 19 that $J \sim M^{\mathbf{x}}$, as needed. Otherwise, Lemma 21 yields

$$M^{\mathbf{w}_0} \triangleleft M^{\langle \mathbf{w}_0, \mathbf{w}_1 \rangle} \triangleleft M^{\langle \mathbf{w}_0, \mathbf{w}_1, \mathbf{w}_2 \rangle} \triangleleft \dots \triangleleft M^{\langle \mathbf{w}_0, \dots, \mathbf{w}_{z-1}
angle} \triangleleft J,$$

which is impossible as $z > n_{\mathbf{A}}^2$.

The next three remarks contain some further observations on aperiodicity.

Remark 22. As we have discussed in the Overview at the intuitive level, our proof of Theorem 1 works by exploiting the fact that randomly walking over an aperiodic \mathbf{A} for a sufficiently long time makes Alice oblivious of her past locations. Using standard theory of discrete-time Markov chains,²³ we can give this statement a more rigorous meaning. Suppose, for the sake of simplicity, that **A** is strongly connected. Alice starts her walk from a vertex a sampled uniformaly at random among all vertices of **A**. After a number i of time steps, she ascertains her new location b and tries to guess her starting point a based on the known data b and i. Clearly, the trivial strategy of choosing at random yields a winning chance of $\frac{1}{n_{\mathbf{A}}}$. Can she do better? Let T be the transition matrix of **A**; i.e., the row-stochastic matrix $T = D^{-1}M$, where D is the diagonal matrix of the outdegrees of the vertices and M is the adjacency matrix. Since A is strongly connected, the Markov chain is ergodic, which means that there exists a unique stationary probability distribution \mathbf{p} satisfying $\mathbf{p}^{\top}T = \mathbf{p}^{\top}$. The probability of Alice ending up in b after i time steps starting from a is $\mathbf{e}_a^{\top} T^i \mathbf{e}_b$ (where \mathbf{e}_a is the probability distribution assigning all the weight to a). By Bayes' Theorem, to maximise her winning chances, Alice should guess a vertex that maximises the corresponding entry of the vector $T^i \mathbf{e}_b$ — which is a deterministic vector, as Alice is aware of both i and b. Now, it is easy to show that T is primitive if and only if M is primitive. If this is the case, T^i converges to the rank-one matrix $\mathbf{1p}^{\top}$ for $i \to \infty$ (where **1** is the all-one vector). Hence, $T^i \mathbf{e}_b$ converges to the constant vector $p_b \mathbf{1}$ — meaning that, after a large-enough number of steps, all choices will roughly yield, for Alice, a disappointing $\frac{1}{n_{A}}$ winning chance. On the other hand, if the primitivity requirement is relaxed, the Markov chain is not guaranteed to converge to the stationary distribution. In fact, it follows from the Perron–Frobenius Theorem that there exist subsequences of the sequence $(T^i)_{i\in\mathbb{N}}$ converging to precisely c different limits, where c is the number of eigenvalues of M having maximum modulus [34, § 9.2]; note that M is primitive precisely when c = 1. In particular, $T^i \mathbf{e}_b$ may not converge to a constant vector, which makes it possible for Alice to guess correctly with probability strictly larger than $\frac{1}{n_A}$.



(a) The adjacency matrix M of the digraph above satisfies $M^5 \sim J$, so M is primitive. However, MM^{\top} is not irreducible.



(b) The adjacency matrix M of the digraph above satisfies $(MM^{\top})^2 \sim J$, so MM^{\top} is primitive. However, M is not irreducible.

Figure 1: Digraphs from Remark 23.

Remark 23. While, in the case of arbitrary structures, the definition of aperiodicity requires that any pair of vertices should be connected by a walk of some fixed length τ oriented in all possible ways (more precisely, by a λ -walk for all possible τ -patterns λ), Theorem 5 shows that, in the binary case, it is enough to consider two orientations only — directed and alternating. The condition cannot be further simplified: Figure 1 illustrates two examples of digraphs

 $^{^{23}}$ For example, see [38].

having the property that directed (resp. alternating) walks of some fixed length connecting any pair of vertices exist, but there are pairs of vertices not connected by any alternating (resp. directed) walk.

Remark 24. In order to obtain more intuition of the definition of aperiodicity in the nonbinary case, we now give a simple example of an aperiodic structure capturing equations over groups. Let G be a finite group with at least two elements, and let **G** be the relational structure with domain G, having a ternary relation $R_g^{\mathbf{G}} = \{(h_1, h_2, h_3) : h_1h_2h_3 = g\}$ for each $g \in G$. Consider the structure \mathbf{G}^{mon} , take $g, g' \in G$, and let $\lambda = (\lambda_1, \lambda_2)$ be a 1-pattern, where $\lambda_1 \neq \lambda_2 \in [3 \cdot |G|]$. A λ -walk in \mathbf{G}^{mon} connecting g and g' is simply given by a tuple **g** in the unique relation of \mathbf{G}^{mon} satisfying $g_{\lambda_1} = g$ and $g_{\lambda_2} = g'$. Using the fact that g and g' have inverses in G, it is straightforward to check that such **g** must exist. It follows that $\mathbf{G}^{\text{mon}} \longrightarrow$ and, thus, \mathbf{G} — is aperiodic with mixing time $\tau = 1$. Since, in addition, **G** is loopless when $|G| \geq 2$, Theorem 1 can be applied to $\text{CSP}(\mathbf{G}) = \text{PCSP}(\mathbf{G}, \mathbf{G})$, and it yields the well-known fact that equations over groups have linear width [29]. We note that this argument also yields linear width for promise equations over groups, as recently studied in [44].

We conclude this section by proving the following monotonicity property of aperiodic structures.

Proposition 25. Let $\mathbf{A}' \to \mathbf{A}$ be relational structures such that \mathbf{A}' is aperiodic. Then, some induced substructure of \mathbf{A} is aperiodic.

Proof. Suppose first that \mathbf{A}' and \mathbf{A} are monic. Take a homomorphism $f : \mathbf{A}' \to \mathbf{A}$, and let $\tilde{\mathbf{A}}$ be the substructure of \mathbf{A} induced by the range of f. Let τ be the mixing time of \mathbf{A}' , and choose two vertices $a, b \in \tilde{\mathbf{A}}$ and a τ -pattern λ . Since f is surjective onto $\tilde{\mathbf{A}}$, there exist two vertices $a', b' \in A'$ such that f(a') = a and f(b') = b. Using that \mathbf{A}' has mixing time τ , we can find a λ -walk in \mathbf{A}' consisting of vertices $a' = a_0, \ldots, a_\tau = b'$ and tuples $\mathbf{a}^{(1)}, \ldots, \mathbf{a}^{(\tau)}$ in $R^{\mathbf{A}'}$. Then, one easily checks that the vertices $f(a_0), \ldots, f(a_\tau)$ and the tuples $f(\mathbf{a}^{(1)}), \ldots, f(\mathbf{a}^{(\tau)})$ in $R^{\tilde{\mathbf{A}}}$ yield a λ -walk in $\tilde{\mathbf{A}}$ connecting $a = f(a_0)$ to $b = f(a_\tau)$. This shows that $\tilde{\mathbf{A}}$ is aperiodic, as required.

Suppose now that \mathbf{A}' and \mathbf{A} are not monic. Observe that $\mathbf{A}'^{\text{mon}} \to \mathbf{A}^{\text{mon}}$ and, by Definition 4, \mathbf{A}'^{mon} is aperiodic. We deduce from the first part of the proof that there exists some nonempty subset $S \subseteq A$ such that the substructure of \mathbf{A}^{mon} induced by S — in symbols, $\mathbf{A}^{\text{mon}}|_{S}$ — is aperiodic. It is not hard to check that $\mathbf{A}^{\text{mon}}|_{S} = (\mathbf{A}|_{S})^{\text{mon}}$. We deduce that $\mathbf{A}|_{S}$ is aperiodic, thus concluding the proof.

5 Fibrosity

Recall that, given an (*r*-uniform) hypergraph \mathcal{H} , $\varphi_{\mathcal{H}}^{\tau}$ denotes the maximum cardinality of a set of mutually disjoint τ -fibers²⁴ in \mathcal{H} , while $\varphi_{\mathcal{H}}^{\max}$, $\pi_{\mathcal{H}}$, and $\lambda_{\mathcal{H}}$ denote the number of maximal fibers, pendent hyperedges, and links in \mathcal{H} , respectively. The goal of this section is to prove the following result.

²⁴We point out that, if f and f' are two disjoint τ -fibers in \mathcal{H} consisting of the hyperedges e_1, \ldots, e_{τ} and e'_1, \ldots, e'_{τ} , respectively, it is not forbidden that, for some $i, j \in [\tau]$, the hyperedges e_i and e'_j are adjacent; however, they cannot be equal.

Theorem (Theorem 6 restated). For $\tau \in \mathbb{N}$ and $1 < \beta \in \mathbb{R}$, let \mathcal{H} be a hereditarily β -sparse hypergraph of girth at least τ having no isolated vertices. Then,

$$\varphi_{\mathcal{H}}^{\tau} + \pi_{\mathcal{H}} > \left(\frac{1}{10r\tau} - \beta + 1\right) n_{\mathcal{H}}.$$

Given a hyperedge e, we let $\operatorname{sdr}(e)$ be the sum of the degree reciprocals of e; i.e., $\operatorname{sdr}(e) = \sum_{v \in e} \frac{1}{\operatorname{deg}_{\mathcal{H}}(v)}$. One easily checks that the equality

$$n_{\mathcal{H}} = \sum_{e \in E(\mathcal{H})} \operatorname{sdr}(e) \tag{1}$$

holds, provided that \mathcal{H} has no isolated vertices.

Proposition (Proposition 7 restated). For $\beta > 1$, let \mathcal{H} be a hereditarily β -sparse hypergraph all of whose fibers are non-degenerate. Then,

$$\varphi_{\mathcal{H}}^{\max} < 3(\beta - 1)n_{\mathcal{H}} + 3\pi_{\mathcal{H}}.$$

Proof. The result is clear if \mathcal{H} consists in an isolated vertex or a single hyperedge, so we assume that this is not the case. Furthermore, since the quantities $\varphi_{\mathcal{H}}^{\max}$, $n_{\mathcal{H}}$, and $\pi_{\mathcal{H}}$ are all additive with respect to disjoint unions, and since a subhypergraph of a hereditarily β -sparse hypergraph is hereditarily β -sparse itself, we can assume without loss of generality that \mathcal{H} is a connected hypergraph.

Let \mathscr{W} be the set of pendent hyperedges of \mathcal{H} that are adjacent to some link. Moreover, let \mathscr{Z} be the set containing every hyperedge of \mathcal{H} that is not pendent, is not a link, and is adjacent to some link. It shall be useful to label the hyperedges in \mathscr{Z} according to the number of links adjacent to them. Note that this number is at least 1 (by definition of \mathscr{Z}) and at most r (since, by definition of a link, any vertex $v \in e \in \mathscr{Z}$ is incident to at most one link). Thus, for $i \in [r]$, we let \mathscr{Z}_i be the set of hyperedges in \mathscr{Z} that are adjacent to exactly *i*-many links. We also let w, z, and z_i denote the cardinalities of the sets \mathscr{W}, \mathscr{Z} , and \mathscr{Z}_i , respectively. The result is obtained by estimating the total sdr of \mathscr{Z} , and comparing it to the total sdr of all other hyperedges.

Take $i \in [r]$ and $e \in \mathscr{Z}_i$, and observe that e intersects exactly i links in exactly i different vertices. Since the maximum degree of a vertex in a link is 2, we deduce that $\operatorname{sdr}(e) \leq r - i + \frac{i}{2} = r - \frac{i}{2}$. However, we can improve this bound in the case that i is 1 or 2. In the latter case, if $\operatorname{sdr}(e)$ were exactly $r - \frac{i}{2} = r - 1$, it would follow that e has exactly two vertices of degree 2 and r - 2 vertices of degree 1; i.e., e would need to be a link, which is impossible by the definition of \mathscr{Z} . As a consequence, at least one of the r - 2 vertices of e that are not incident to a link has degree at least 2. This yields $\operatorname{sdr}(e) \leq r - 3 + 1 + \frac{1}{2} = r - \frac{i}{2} - \frac{1}{2}$. If i = 1, one can do even better. Let V be the set of r - 1 vertices of e that are not incident to a link. If all of them had degree 1, e would be a pendent hyperedge, against our assumption. Furthermore, if one of the vertices of V had degree 2 and all the others had degree 1, e would be a link, again contradicting the hypothesis. It follows that either two vertices of V have degree at least 2, or a vertex of V has degree at least 3. The latter choice yields the highest sdr, and thus we can use it to give an upper bound: $\operatorname{sdr}(e) \leq r - 2 + \frac{1}{2} + \frac{1}{3} = r - \frac{i}{2} - \frac{2}{3}$. As a

consequence, we bound the total sdr of ${\mathscr Z}$ as follows:

$$\begin{split} \sum_{e \in \mathscr{X}} \mathrm{sdr}(e) &= \sum_{e \in \mathscr{X}_1} \mathrm{sdr}(e) + \sum_{e \in \mathscr{X}_2} \mathrm{sdr}(e) + \sum_{i \ge 3} \sum_{e \in \mathscr{X}_i} \mathrm{sdr}(e) \\ &\leq z_1 \left(r - \frac{1}{2} - \frac{2}{3} \right) + z_2 \left(r - 1 - \frac{1}{2} \right) + \sum_{i \ge 3} z_i \left(r - \frac{i}{2} \right) \\ &= r \sum_{i \in [r]} z_i - \frac{1}{2} \sum_{i \in [r]} i z_i - \frac{2}{3} z_1 - \frac{1}{2} z_2 \\ &= (r - 1)z + z - \frac{1}{2} \sum_{i \in [r]} i z_i - \frac{2}{3} z_1 - \frac{1}{2} z_2, \end{split}$$

where we have used that $\sum_{i \in [r]} z_i = z$. Note that

$$\begin{aligned} z - \frac{1}{2} \sum_{i \in [r]} iz_i - \frac{2}{3} z_1 - \frac{1}{2} z_2 &= \sum_{i \in [r]} \left(z_i - \frac{iz_i}{2} \right) - \frac{2}{3} z_1 - \frac{1}{2} z_2 = -\frac{1}{6} z_1 - \frac{1}{2} z_2 - \sum_{i \ge 3} \left(\frac{i}{2} - 1 \right) z_i \\ &\leq -\frac{1}{6} \sum_{i \in [r]} iz_i. \end{aligned}$$

Therefore, we obtain

$$\sum_{e \in \mathscr{Z}} \operatorname{sdr}(e) \le (r-1)z - \frac{1}{6} \sum_{i \in [r]} iz_i.$$
⁽²⁾

We say that a fiber f connects two (possibly equal) hyperedges $e, e' \in E(\mathcal{H}) \setminus f$ if some hyperedge in f is adjacent to e and some hyperedge in f is adjacent to e'. Consider the multigraph (with loops allowed) \mathcal{G} whose vertex set is $\mathscr{Z} \cup \mathscr{W}$ and whose edge multiset is defined as follows: For $e, e' \in \mathscr{Z} \cup \mathscr{W}$ (possibly equal), we insert one edge in \mathcal{G} joining e and e'for each maximal fiber of \mathcal{H} connecting e and e'. We claim that $m_{\mathcal{G}} = \varphi_{\mathcal{H}}^{\max}$ (where $m_{\mathcal{G}}$ is the number of edges of the multigraph \mathcal{G} , counted with multiplicity). First, since any maximal fiber yields at most one edge in \mathcal{G} , it is clear that $m_{\mathcal{G}} \leq \varphi_{\mathcal{H}}^{\max}$. Suppose that $m_{\mathcal{G}} < \varphi_{\mathcal{H}}^{\max}$. This would mean that there exists a maximal fiber f such that, for each $e \in f$, e is only adjacent to other links. Since f is maximal, e is in fact only adjacent to other hyperedges in f. In other words, f induces a cycle, which is impossible as f is non-degenerate by assumption. It follows that the claim is true. For each $e \in \mathscr{Z} \cup \mathscr{W}$, let $\deg_{\mathcal{G}}(e)$ be the degree of e in \mathcal{G} ; i.e., the number of edges of \mathcal{G} to which e is incident (where a loop contributes 2). Note that, if $e \in \mathscr{Z}_i$ for some $i \in [r]$, $\deg_{\mathcal{G}}(e) = i$. Moreover, if $e \in \mathscr{W}$, $\deg_{\mathcal{G}}(e) = 1$. Via the handshaking lemma, we deduce that

$$2\varphi_{\mathcal{H}}^{\max} = 2m_{\mathcal{G}} = \sum_{e \in \mathscr{Z}} \deg_{\mathcal{G}}(e) + \sum_{e \in \mathscr{W}} \deg_{\mathcal{G}}(e) = \sum_{i \in [r]} iz_i + w.$$

Plugging this information into (2) yields

$$\sum_{e \in \mathscr{Z}} \operatorname{sdr}(e) \le (r-1)z - \frac{1}{3}\varphi_{\mathcal{H}}^{\max} + \frac{1}{6}w$$

Let \mathscr{P} be the set of pendent hyperedges of \mathcal{H} . Since \mathcal{H} is connected and does not consist in a single hyperedge, $\operatorname{sdr}(e) \leq r - \frac{1}{2}$ for each $e \in E(\mathcal{H})$. Moreover, $\operatorname{sdr}(e) \leq r - 1$ if $e \notin \mathscr{P}$. Noting also that, by assumption, \mathcal{H} does not contain isolated vertices, (1) yields

$$n_{\mathcal{H}} = \sum_{e \in E(\mathcal{H})} \operatorname{sdr}(e) = \sum_{e \in \mathscr{P}} \operatorname{sdr}(e) + \sum_{e \in \mathscr{Z}} \operatorname{sdr}(e) + \sum_{e \in \mathcal{H} \setminus (\mathscr{P} \cup \mathscr{Z})} \operatorname{sdr}(e)$$
$$\leq \left(r - \frac{1}{2}\right) \pi_{\mathcal{H}} + (r - 1)z - \frac{1}{3}\varphi_{\mathcal{H}}^{\max} + \frac{1}{6}w + (m_{\mathcal{H}} - \pi_{\mathcal{H}} - z)(r - 1)$$
$$\leq -\frac{1}{3}\varphi_{\mathcal{H}}^{\max} + \pi_{\mathcal{H}} + m_{\mathcal{H}}(r - 1),$$

where we have used that $w \leq \pi_{\mathcal{H}}$. We now make use of the fact that \mathcal{H} is hereditarily β -sparse and thus, in particular, it is β -sparse. This gives

$$n_{\mathcal{H}} < -\frac{1}{3}\varphi_{\mathcal{H}}^{\max} + \pi_{\mathcal{H}} + \beta n_{\mathcal{H}}$$

which concludes the proof.

Proposition (Proposition 8 restated). For $\beta > 1$, let \mathcal{H} be a hereditarily β -sparse hypergraph having no isolated vertices. Then,

$$\lambda_{\mathcal{H}} > \left(\frac{1}{r} + 6 - 6\beta\right) n_{\mathcal{H}} - 7\pi_{\mathcal{H}}.$$

Proof. We can assume without loss of generality that \mathcal{H} is connected. Let \mathscr{P} and \mathscr{L} denote the set of pendent hyperedges and the set of links of \mathcal{H} , respectively. Notice that $\operatorname{sdr}(e) = r - 1$ if $e \in \mathscr{L}$, $\operatorname{sdr}(e) \leq r - \frac{7}{6}$ if $e \notin \mathscr{P} \cup \mathscr{L}$, and, clearly, $\operatorname{sdr}(e) \leq r$ for any hyperedge $e \in E(\mathcal{H})$. Similarly to Proposition 7, the result is obtained by comparing the total sdr of a certain set of hyperedges to that of all other hyperedges. In this case, instead of the set \mathscr{L} , we consider the set \mathscr{L} . Since, by assumption, \mathcal{H} has no isolated vertices, (1) gives

$$n_{\mathcal{H}} = \sum_{e \in E(\mathcal{H})} \operatorname{sdr}(e) = \sum_{e \in \mathscr{P}} \operatorname{sdr}(e) + \sum_{e \in \mathscr{L}} \operatorname{sdr}(e) + \sum_{e \in E(\mathcal{H}) \setminus (\mathscr{P} \cup \mathscr{L})} \operatorname{sdr}(e)$$
$$\leq \pi_{\mathcal{H}}r + \lambda_{\mathcal{H}}(r-1) + (m_{\mathcal{H}} - \pi_{\mathcal{H}} - \lambda_{\mathcal{H}}) \left(r - \frac{7}{6}\right) = \frac{1}{6}\lambda_{\mathcal{H}} + \frac{7}{6}\pi_{\mathcal{H}} + m_{\mathcal{H}}(r-1) - \frac{1}{6}m_{\mathcal{H}}.$$

Note that $m_{\mathcal{H}}$ appears twice, with opposite signs, in the right-hand side of the expression above. Hence, to continue the chain of inequalities, we need both an upper and a lower bound on $m_{\mathcal{H}}$. The upper bound is the one dictated by the fact that \mathcal{H} is β -sparse; i.e., $m_{\mathcal{H}} < \frac{\beta}{r-1}n_{\mathcal{H}}$. The lower bound comes from the fact that \mathcal{H} is connected and, thus, $m_{\mathcal{H}} \geq \frac{n_{\mathcal{H}}-1}{r-1}$ (as can be easily shown by induction on $m_{\mathcal{H}}$). Moreover, since \mathcal{H} contains no isolated vertices, $n_{\mathcal{H}} \geq r$, and it follows that $\frac{n_{\mathcal{H}}-1}{r-1} \geq \frac{n_{\mathcal{H}}}{r}$. Therefore, we obtain

$$n_{\mathcal{H}} < \frac{1}{6}\lambda_{\mathcal{H}} + \frac{7}{6}\pi_{\mathcal{H}} + \beta n_{\mathcal{H}} - \frac{1}{6r}n_{\mathcal{H}}$$

which directly yields the claimed inequality.

Proposition (Proposition 9 restated). For any hypergraph \mathcal{H} and any $\tau \in \mathbb{N}$,

$$\varphi_{\mathcal{H}}^{\tau} + \varphi_{\mathcal{H}}^{\max} > \frac{\lambda_{\mathcal{H}}}{\tau}.$$

Proof. For $i \in \mathbb{N}$, let p_i be the number of maximal fibers of \mathcal{H} of size exactly *i*. Note that

$$\sum_{i\in\mathbb{N}}p_i=\varphi_{\mathcal{H}}^{\max}$$

Moreover, since the maximal fibers of \mathcal{H} partition the set of links of \mathcal{H} , we have that

$$\sum_{i\in\mathbb{N}}ip_i=\lambda_{\mathcal{H}}.$$

Observe now that a maximal fiber of size *i* can be split into a set of $\lfloor \frac{i}{\tau} \rfloor$ mutually disjoint τ -fibers (plus an additional $(i - \tau \lfloor \frac{i}{\tau} \rfloor)$ -fiber). As a consequence, we find that

$$\varphi_{\mathcal{H}}^{\tau} = \sum_{i \in \mathbb{N}} p_i \left\lfloor \frac{i}{\tau} \right\rfloor > \sum_{i \in \mathbb{N}} p_i \left(\frac{i}{\tau} - 1 \right) = \frac{\lambda_{\mathcal{H}}}{\tau} - \varphi_{\mathcal{H}}^{\max},$$

and the conclusion follows.

Theorem 6 follows by combining Propositions 7, 8, and 9, as detailed in Subsection 2.2.

6 Sparsity

In this section, we prove the existence of sparse and highly chromatic hypergraphs of large girth, as stated next.

Theorem (Theorem 10 restated). Take two positive integer numbers g and h and a real number $\beta > 1$. There exists a positive real number $\delta = \delta(g, h, \beta)$ and a positive integer number $n_0 = n_0(g, h, \beta)$ such that, for each $n \ge n_0$, there exists a hypergraph \mathcal{H} with the following properties:

- 1. \mathcal{H} has n vertices;
- 2. $\operatorname{gir}(\mathcal{H}) \geq g;$
- 3. $\operatorname{chr}(\mathcal{H}) \ge h;$
- 4. \mathcal{H} is $(\delta n, \beta)$ -threshold-sparse.

As discussed in the Overview, our proof of Theorem 10 is probabilistic, and it combines Erdős–Hajnal's proof of the existence of hypergraphs with arbitrarily large girth and chromatic number [28] with Atserias–Dalmau's result on the sparsity of random Erdős-Rényi hypergraphs [1]. The latter result is stated in terms of a different version of threshold-sparsity, which we now define.²⁵

For μ and ν positive real numbers, we say that a hypergraph \mathcal{H} is (μ, ν) -vertex-thresholdsparse if any subhypergraph \mathcal{H}' of \mathcal{H} such that $n_{\mathcal{H}'} \leq \mu \cdot n_{\mathcal{H}}$ satisfies $m_{\mathcal{H}'} < \nu \cdot n_{\mathcal{H}'}$. The following simple result connects the two notions of threshold-sparsity.

Lemma 26. Let $\beta > 1, \gamma, \mu, \nu$ be positive real numbers, let \mathcal{H} be a (μ, ν) -vertex-threshold-sparse hypergraph, and suppose that $\frac{\beta}{\nu} \ge r-1$ and $n_{\mathcal{H}} \ge (r-1)\frac{\gamma}{\beta\mu}$. Then, \mathcal{H} is (γ, β) -threshold-sparse.

 $^{^{25}}$ We remark that the notion of threshold-sparsity given in Subsection 2.3 is also implicit in [1].

Proof. Let \mathcal{H}' be a subhypergraph of \mathcal{H} satisfying $m_{\mathcal{H}'} \leq \gamma$, and suppose that \mathcal{H}' is not β -sparse. We obtain

$$n_{\mathcal{H}'} \leq (r-1)\frac{m_{\mathcal{H}'}}{\beta} \leq (r-1)\frac{\gamma}{\beta} \leq \mu \cdot n_{\mathcal{H}}.$$

Since \mathcal{H} is (μ, ν) -vertex-threshold-sparse, we deduce that

$$m_{\mathcal{H}'} < \nu \cdot n_{\mathcal{H}'} \le \frac{\beta}{r-1} n_{\mathcal{H}'}$$

which contradicts the fact that \mathcal{H}' is not β -sparse.

For $n \in \mathbb{N}$ and $p \in \mathbb{R}$, $0 \leq p \leq 1$, the expression $\mathcal{H} \sim \mathscr{H}(n, p)$ shall denote that \mathcal{H} is a random (*r*-uniform) Erdős-Rényi hypergraph with *n* vertices, where each *r*-element set of vertices forms a hyperedge with probability *p*, independently. The next result gives an upper bound on the probability that a random Erdős-Rényi hypergraph is not vertex-threshold-sparse.

Proposition 27 ([1]). Let μ, ν, ℓ be positive real numbers and let n be a positive integer number. If $1 \leq \ell \leq n^{r-1}$ and $\mu \leq (\frac{\nu}{\ell})^{\frac{1}{r-1}}(\frac{r}{e})^{\frac{r}{r-1}}$, the probability that a random hypergraph $\mathcal{H} \sim \mathscr{H}(n, \ell n^{1-r})$ is not (μ, ν) -vertex-threshold-sparse is at most

$$\pi(r,\ell,n,\mu,\nu) = \sum_{i=1}^{\lfloor \mu n \rfloor} \left(\left(\frac{n}{i}\right)^{1-(r-1)\nu} \ell^{\nu} e^{1+(r+1)\nu} r^{-r\nu} \nu^{-\nu} \right)^i.$$
(3)

,

Proof of Theorem 10. Consider the following parameters:

$$\ell = \frac{(3hr)^r}{2h} \log(3eh) + 1,$$

$$\nu = \frac{\beta}{r-1},$$

$$\vartheta = \ell^{\nu} e^{1+(r+1)\nu} r^{-r\nu} \nu^{-\nu},$$

$$\mu = \min\left(\left(\frac{\nu}{\ell}\right)^{\frac{1}{r-1}} \left(\frac{r}{e}\right)^{\frac{r}{r-1}}, (3\vartheta)^{-\frac{1}{\beta-1}}\right),$$

$$\delta = \frac{\beta\mu}{r-1}.$$
(4)

Notice that all of them are positive real numbers. Let n be an even positive integer number to be determined later, and suppose that $n \ge \ell^{\frac{1}{r-1}}$. Let $p = \ell n^{1-r}$, and sample a hypergraph $\mathcal{H} \sim \mathscr{H}(n,p)$.

Fix $w = \lfloor \frac{n}{2h} \rfloor$, and let W be the random variable counting the number of independent sets in \mathcal{H} of size w (where an independent set is a set S of vertices such that $e \not\subseteq S$ for any $e \in E(\mathcal{H})$). Suppose that $n \ge 2hr$ (which implies that $w \ge r$) and that $n \ge 6h$ (which implies that $w \ge \frac{n}{2h} - 1 \ge \frac{n}{3h}$). It is well known that, for $1 \le x \le y$, the inequalities

$$\left(\frac{y}{x}\right)^x \le \binom{y}{x} \le \left(\frac{ey}{x}\right)^x$$

hold (for example, see [26]). As a consequence, using the linearity of expectation, we obtain the following upper bound for the expected value of W:

$$\mathbb{E}[W] = \binom{n}{w} (1-p)^{\binom{w}{r}} \le \binom{n}{w} e^{-p\binom{w}{r}} \le \left(\frac{en}{w}\right)^w e^{-p\left(\frac{w}{r}\right)^r} \le (3eh)^{\frac{n}{2h}} e^{-\ell n^{1-r} \frac{n^r}{(3hr)^r}} = \left(\frac{(3eh)^{\frac{1}{2h}}}{e^{\frac{\ell}{(3hr)^r}}}\right)^n.$$

The choice of ℓ guarantees that $(3eh)^{\frac{1}{2h}} < e^{\frac{\ell}{(3hr)^r}}$, whence it follows that $\mathbb{E}[W]$ approaches 0 as n tends to infinity. Recall that the independence number of \mathcal{H} (in symbols, $\operatorname{ind}(\mathcal{H})$) is the maximum cardinality of an independent set in \mathcal{H} . By Markov's inequality, the probability that W is at least 1 - i.e., that $\operatorname{ind}(\mathcal{H}) \geq w$ — is at most $\mathbb{E}[W]$ and, thus, it tends to 0 as n tends to infinity. In particular, for n large enough,

$$\mathbb{P}\left(W \ge 1\right) < \frac{1}{4}.\tag{5}$$

It is well known [9] that a cycle of length $j \ge 2$ in a hypergraph yields a collection of distinct hyperedges e_1, \ldots, e_j such that $|\bigcup_{i \in [j]} e_i| \le j(r-1)$ and, vice versa, a collection of j distinct hyperedges satisfying this condition contains a cycle of length at most j. The number of such collections is at most

$$\binom{n}{j(r-1)}\binom{j(r-1)}{r}^j,$$

which is upper-bounded by $x_j n^{j(r-1)}$ for some constant x_j independent of n. Let a *short cycle* indicate a cycle of length at most g-1, and let V be the random variable counting the number of short cycles in \mathcal{H} . By linearity of expectation, we find

$$\mathbb{E}[V] \le \sum_{j=2}^{g-1} x_j n^{j(r-1)} p^j = \sum_{j=2}^{g-1} \ell^j x_j.$$

Hence, using Markov's inequality, we deduce that the probability that V is at least $\frac{n}{2}$ is at most

$$\frac{2\mathbb{E}[V]}{n} \le \frac{2\sum_{j=2}^{g-1} \ell^j x_j}{n}.$$

Since the numerator in the expression above is independent of n, the fraction approaches 0 as n tends to infinity. In particular,

$$\mathbb{P}\left(V \ge \frac{n}{2}\right) < \frac{1}{4} \tag{6}$$

for n large enough.

Let A be the event that \mathcal{H} is not (μ, ν) -vertex-threshold-sparse. The choice of μ guarantees that Proposition 27 applies. Hence, the probability of A is upper bounded by the quantity $\pi = \pi(r, \ell, n, \mu, \nu)$ in (3). From the fact that $\mu \leq (3\vartheta)^{-\frac{1}{\beta-1}}$, we deduce that $\mu^{\beta-1}\vartheta \leq \frac{1}{3}$. Therefore,

$$\mathbb{P}(A) \le \pi = \sum_{i=1}^{\lfloor \mu n \rfloor} \left(\left(\frac{i}{n}\right)^{\beta-1} \vartheta \right)^i \le \sum_{i=1}^{\lfloor \mu n \rfloor} \left(\mu^{\beta-1} \vartheta \right)^i \le \sum_{i=1}^{\lfloor \mu n \rfloor} \frac{1}{3^i} < \sum_{i=1}^{\infty} \frac{1}{3^i} = \frac{1}{2}.$$
 (7)

By the union bound, we deduce from (5), (6), and (7) that, if n is large enough, there exists a hypergraph \mathcal{H} having n vertices such that W = 0, $V < \frac{n}{2}$, and A does not hold. In other words, $\operatorname{ind}(\mathcal{H}) < w$, \mathcal{H} has fewer than $\frac{n}{2}$ short cycles, and \mathcal{H} is (μ, ν) -vertex-threshold-sparse.

Select a set S of vertices of \mathcal{H} of cardinality $\frac{n}{2}$ in a way that each short cycle of \mathcal{H} contains a vertex of S, and let $\tilde{\mathcal{H}}$ be the subhypergraph of \mathcal{H} induced by $V(\mathcal{H}) \setminus S$. Note that $n_{\tilde{\mathcal{H}}} = \frac{n}{2}$. Furthermore, $\operatorname{ind}(\tilde{\mathcal{H}}) \leq \operatorname{ind}(\mathcal{H}) < w$, while $\operatorname{gir}(\tilde{\mathcal{H}}) \geq g$ since all short cycles have been broken. Since the product of the chromatic and the independence numbers of a hypergraph is at least the number of its vertices (see [9]), we find

$$\operatorname{chr}(\tilde{\mathcal{H}}) \ge \frac{n_{\tilde{\mathcal{H}}}}{\operatorname{ind}(\tilde{\mathcal{H}})} \ge \frac{n}{2w} \ge h.$$

Finally, observe that $\hat{\mathcal{H}}$ is (μ, ν) -vertex-threshold-sparse by monotonicity of vertex-threshold-sparsity. The choice of δ guarantees that

$$(r-1)\frac{\delta n_{\tilde{\mathcal{H}}}}{\beta\mu} = n_{\tilde{\mathcal{H}}}$$

Therefore, Lemma 26 implies that \mathcal{H} is $(\delta n_{\mathcal{H}}, \beta)$ -threshold-sparse, thus concluding the proof of Theorem 10.

7 Consistency

The goal of this section is to establish that a (κ, γ) -consistency gap — for some suitable parameter γ — can be used to recover a κ -strategy and, thus, to certify acceptance by the κ -th level of the local-consistency algorithm applied to an aperiodic template and a sparse instance having large girth.

Theorem (Theorem 14 restated). Let **X** and **A** be monic structures, and take two numbers $\kappa \in \mathbb{N}$ and $\gamma \in \mathbb{R}$ such that $\gamma \geq n_{\mathbf{A}}$. Suppose that

- A is aperiodic with mixing time τ ;
- **X** is oriented, and **X**^{sym} has girth at least τ and is (γ, β) -threshold-sparse for some real number $1 < \beta < 1 + \frac{1}{10r\tau}$;
- (\mathbf{X}, \mathbf{A}) has a (κ, γ) -consistency gap.

Then, $LC^{\kappa}(\mathbf{X}, \mathbf{A}) = YES.$

The key to proving Theorem 14 is to show that τ -fibers and pendent hyperedges yield boundary sets (as considered in [1,47]) for aperiodic templates having mixing time τ , in the sense that any partial homomorphism to such templates can be extended to include τ -fibers and pendent hyperedges. This is done in the next two propositions.

Proposition (Proposition 11 restated). Let \mathbf{X} , \mathbf{A} be monic structures such that \mathbf{X} is oriented and \mathbf{A} is aperiodic with mixing time τ , let f be a τ -fiber of \mathbf{X}^{sym} , let J be a joint²⁶ of f, and let \mathbf{Y} be the substructure of \mathbf{X} induced by $(X \setminus \bigcup_{e \in f} e) \cup J$. Then, any homomorphism $h: \mathbf{Y} \to \mathbf{A}$ can be extended to a homomorphism $h': \mathbf{X} \to \mathbf{A}$.

Proof. Let $J = \{u, v\}$, where u and v are possibly equal vertices of \mathcal{H}_f . Label the links in f as e_1, \ldots, e_τ , where e_i is adjacent to e_{i+1} for each $i \in [\tau - 1]$. Assume, without loss of generality, that $u \in e_1$ and $v \in e_\tau$. Set $w_0 = u$ and define iteratively, for each $i \in [\tau]$, the vertex w_i as the only vertex of $e_i \setminus \{w_{i-1}\}$ having degree 2. Note that this implies $w_\tau = v$. Recall that R denotes the unique relation symbol in the signature of \mathbf{X} and \mathbf{A} , and $r = \operatorname{ar}(R)$. Since \mathbf{X} is oriented,

 $^{^{26}\}mathrm{Recall}$ the definition of joints given in Subsection 2.4.

by the definition of \mathbf{X}^{sym} , there exist tuples $\mathbf{x}^{(1)}, \ldots, \mathbf{x}^{(\tau)} \in R^{\mathbf{X}}$ such that $e_i = \text{set}(\mathbf{x}^{(i)})$ for each $i \in [\tau]$, and the $\mathbf{x}^{(i)}$'s are uniquely determined. Hence, there exist two tuples $\mathbf{p}, \mathbf{q} \in [r]^{\tau}$ such that, for each $i \in [\tau]$, $x_{p_i}^{(i)} = w_{i-1}$ and $x_{q_i}^{(i)} = w_i$. From the way we chose the vertices w_i , it follows that $w_{i-1} \neq w_i$ for each $i \in [\tau]$. Therefore, $p_i \neq q_i$ for each $i \in [\tau]$. Hence, the matrix $\lambda = [\mathbf{p} \ \mathbf{q}]$ is a τ -pattern. Observe that $w_0 = u$ and $w_{\tau} = v$ are both vertices of \mathbf{Y} . Using the assumption that \mathbf{A} has mixing time τ , we can then find a λ -walk connecting $h(w_0)$ to $h(w_{\tau})$. Suppose that the λ -walk consists of vertices $a_0 = h(w_0), a_1, \ldots, a_{\tau} = h(w_{\tau})$ in Aand tuples $\mathbf{a}^{(1)}, \ldots, \mathbf{a}^{(\tau)}$ in $R^{\mathbf{A}}$. Consider the map $h' : X \to A$ defined as follows:

(I) $h'(x_j^{(i)}) = a_j^{(i)}$ for each $i \in [\tau]$ and each $j \in [r]$; (II) h'(x) = h(x) for each $x \in Y$.

First, we show that h' is well defined. The two cases above cover all vertices of X, and the only vertices covered multiple times are the vertices w_i for $i \in [\tau]$. Indeed, $w_0 = x_{p_1}^{(1)}$, so we need to check that $h(w_0) = a_{p_1}^{(1)}$. This is true since, using the definition of λ -walk, we have

$$a_{p_1}^{(1)} = a_{\lambda_{1,1}}^{(1)} = a_0 = h(w_0).$$

Similarly, the homonymy $w_{\tau} = x_{q_{\tau}}^{(\tau)}$ does not yield problems as

$$a_{q_{\tau}}^{(\tau)} = a_{\lambda_{\tau,2}}^{(\tau)} = a_{\tau} = h(w_{\tau}).$$

Furthermore, for $i \in [\tau - 1]$, we have $w_i = x_{p_{i+1}}^{(i+1)} = x_{q_i}^{(i)}$. Also this homonymy is not problematic, as

$$a_{p_{i+1}}^{(i+1)} = a_{\lambda_{i+1,1}}^{(i+1)} = a_i = a_{\lambda_{i,2}}^{(i)} = a_{q_i}^{(i)}.$$

Therefore, h' is well defined. We now claim that it is a homomorphism from **X** to **A**. Since h' clearly extends h, this would be enough to conclude the proof. Take $\mathbf{x} \in R^{\mathbf{X}}$; we need to show that $h'(\mathbf{x}) \in R^{\mathbf{A}}$. If $\mathbf{x} \in R^{\mathbf{Y}}$, the claim is clear as, in this case, $h'(\mathbf{x}) = h(\mathbf{x}) \in R^{\mathbf{A}}$ since h is a homomorphism. Otherwise, by definition of fiber, it must be the case that $\operatorname{set}(\mathbf{x}) \in f$. Thus, since **X** is oriented, we deduce that $\mathbf{x} = \mathbf{x}^{(i)}$ for some $i \in [\tau]$. Using the rule (I) above, we conclude that $h'(\mathbf{x}) = h'(\mathbf{x}^{(i)}) = \mathbf{a}^{(i)} \in R^{\mathbf{A}}$, as required.

Proposition (Proposition 12 restated). Let \mathbf{X} , \mathbf{A} be monic structures such that \mathbf{X} is oriented and \mathbf{A} is aperiodic, let e be a pendent hyperedge for \mathbf{X}^{sym} , let J be a joint of e, and let \mathbf{Y} be the substructure of \mathbf{X} induced by $(X \setminus e) \cup J$. Then, any homomorphism $h : \mathbf{Y} \to \mathbf{A}$ can be extended to a homomorphism $h' : \mathbf{X} \to \mathbf{A}$.

Proof. Call v the unique vertex in J, let $\mathbf{x} \in R^{\mathbf{X}}$ be such that $e = \operatorname{set}(\mathbf{x})$, and let $j \in [r]$ be such that $x_j = v$, where r is the arity of R. Let τ be the mixing time of \mathbf{A} . The definition of oriented structure guarantees that $r \geq 2$; hence, there exists some τ -pattern λ such that $\lambda_{1,1} = j$. Using the aperiodicity of \mathbf{A} , we find a λ -walk connecting h(v) to itself. In particular, this means that there exists a tuple $\mathbf{a} \in R^{\mathbf{A}}$ such that $h(v) = a_{\lambda_{1,1}} = a_j$. Since all vertices in $e \setminus \{v\}$ have degree 1 in $\mathbf{X}^{\operatorname{sym}}$, the map $h' : X \to A$ given by $h'(x_i) = a_i$ for each $i \in [r]$ and h'(x) = h(x) for all other vertices of \mathbf{X} is easily seen to yield a homomorphism from \mathbf{X} to \mathbf{A} extending h. We can now prove Theorem 14. This result extends [1, Lemmas 8,10], and the proof below is an adaptation of the one in [1]. Given a monic structure \mathbf{Y} whose unique relation symbol is R, we let $m_{\mathbf{Y}} = |R^{\mathbf{Y}}|$.

Proof of Theorem 14. If r = 2 and $\tau = 1$, it follows from Definition 3 that \mathbf{A} has a loop, which implies that $\mathbf{X} \to \mathbf{A}$ and, thus, that $\mathrm{LC}^{\kappa}(\mathbf{X}, \mathbf{A}) = \mathrm{YES}$, as needed; hence, we assume that $r \geq 3$ or $\tau \geq 2$ (note that the case r = 1 is forbidden by the definition of oriented structure). Similarly, we can assume that $m_{\mathbf{X}} \neq 0$. Let $\hat{\mathbf{X}}$ be the substructure of \mathbf{X} obtained by removing all vertices that do not belong to any tuple in $R^{\mathbf{X}}$. It is easy to check from Definition 13 that a (κ, γ) -consistency gap for (\mathbf{X}, \mathbf{A}) implies a (κ, γ) -consistency gap for $(\hat{\mathbf{X}}, \mathbf{A})$. Moreover, any κ -strategy $\hat{\mathscr{F}}$ for $\hat{\mathbf{X}}$ and \mathbf{A} can be straightforwardly turned into a κ -strategy for \mathbf{X} and \mathbf{A} by extending the partial homomorphisms in $\hat{\mathscr{F}}$ in an arbitrary way to cover the vertices of $X \setminus \hat{X}$. This means that we can assume, without loss of generality, that \mathbf{X}^{sym} has no isolated vertices. Let \mathscr{F} be the set of all functions from a subset of X of size at most κ to A, that are consistent with every substructure of \mathbf{X} whose relation has at most γ tuples. We claim that \mathscr{F} is a κ -strategy for \mathbf{X} and \mathbf{A} .

Take a nonempty map $f \in \mathscr{F}$ having domain $X' \subseteq X$, let \mathbf{X}' be the substructure of \mathbf{X} induced by X', let $\mathbf{x} \in R^{\mathbf{X}'}$, and let \mathbf{Y} be the substructure of \mathbf{X} whose vertex set is $Y = \operatorname{set}(\mathbf{x})$ and whose relation is $R^{\mathbf{Y}} = \{\mathbf{x}\}$. Since $m_{\mathbf{Y}} = 1 \leq n_{\mathbf{A}} \leq \gamma$, f is consistent with \mathbf{Y} by definition of \mathscr{F} , so there exists a homomorphism $h : \mathbf{Y} \to \mathbf{A}$ that agrees with f on $X' \cap Y = Y$. It follows that $f(\mathbf{x}) = h(\mathbf{x}) \in R^{\mathbf{A}}$. Therefore, f is a homomorphism from \mathbf{X}' to \mathbf{A} (and, thus, a partial homomorphism from \mathbf{X} to \mathbf{A}).

We now claim that all substructures \mathbf{Y} of \mathbf{X} such that $m_{\mathbf{Y}} \leq \gamma$ are homomorphic to \mathbf{A} . Otherwise, let \mathbf{Y} be a witness of the contrary; we may assume without loss of generality that any proper substructure of \mathbf{Y} is homomorphic to \mathbf{A} and that $m_{\mathbf{Y}} \geq 1$. Note that \mathbf{Y} is oriented and \mathbf{Y}^{sym} is hereditarily β -sparse. Applying Theorem 6, we deduce that

$$\varphi_{\mathbf{Y}^{\text{sym}}}^{\tau} + \pi_{\mathbf{Y}^{\text{sym}}} > \left(\frac{1}{10r\tau} - \beta + 1\right) n_{\mathbf{Y}^{\text{sym}}} > 0,$$

where we have used that $\beta < 1 + \frac{1}{10\tau\tau}$. Hence, \mathbf{Y}^{sym} has at least one τ -fiber or pendent hyperedge. In the former case, let $S = \bigcup_{e \in f} e$ where f is a τ -fiber of \mathbf{Y}^{sym} , and let J be a joint of f; in the latter case, let S = e where e is a pendent hyperedge of \mathbf{Y}^{sym} , and let Jbe a joint of e. Let \mathbf{Y}' be the substructure of \mathbf{Y} induced by $(Y \setminus S) \cup J$. Since at least one of the conditions $r \geq 3$ and $\tau \geq 2$ holds — as assumed at the beginning of the proof — we observe that $S \setminus J \neq \emptyset$. Thus, $n_{\mathbf{Y}'} < n_{\mathbf{Y}}$. By the minimality of \mathbf{Y} , we have that $\mathbf{Y}' \to \mathbf{A}$. It follows from Proposition 11 or from Proposition 12 that $\mathbf{Y} \to \mathbf{A}$, a contradiction. Therefore, the claim is true. In particular, we deduce that the empty function from \emptyset to A belongs to \mathscr{F} , thus yielding $\mathscr{F} \neq \emptyset$.

The fact that \mathscr{F} is closed under restrictions directly follows from the definition of consistent functions. To show that \mathscr{F} has the extension property up to κ , take $\mathscr{F} \ni f: X' \to A$ with $|X'| < \kappa$ and pick $x \in X \setminus X'$. For $a \in A$, consider the map $f_a: X' \cup \{x\} \to A$ mapping x to aand x' to f(x') for each $x' \in X'$. Suppose that for each $a \in A$ there exists some substructure \mathbf{X}_a of \mathbf{X} whose relation has at most $\frac{\gamma}{n_{\mathbf{A}}}$ tuples such that f_a is not consistent with \mathbf{X}_a , and let $\mathbf{Z} = \bigcup_{a \in A} \mathbf{X}_a$ (i.e., $Z = \bigcup_{a \in A} X_a$ and $R^{\mathbf{Z}} = \bigcup_{a \in A} R^{\mathbf{X}_a}$). Note that, for each $a \in A$, f is consistent with \mathbf{X}_a since $f \in \mathscr{F}$; hence, $x \in \bigcap_{a \in A} X_a \subseteq Z$. If f is consistent with \mathbf{Z} , there exists a homomorphism $h: \mathbf{Z} \to \mathbf{A}$ such that f and h agree on $X' \cap Z$. Using that $x \in Z$, observe that the restriction of h to $\mathbf{X}_{h(x)}$ is a homomorphism from $\mathbf{X}_{h(x)}$ to \mathbf{A} that agrees with $f_{h(x)}$ on $(X' \cup \{x\}) \cap X_{h(x)}$. This means that $f_{h(x)}$ is consistent with $\mathbf{X}_{h(x)}$, a contradiction. As a consequence, f is not consistent with \mathbf{Z} . But this contradicts the fact that $f \in \mathscr{F}$, as

$$m_{\mathbf{Z}} \leq \sum_{a \in A} m_{\mathbf{X}_a} \leq \sum_{a \in A} \frac{\gamma}{n_{\mathbf{A}}} = \gamma.$$

It follows that there exists some $a' \in A$ such that $f_{a'}$ is consistent with every substructure of **X** whose relation has at most $\frac{\gamma}{n_{\mathbf{A}}}$ tuples. Let **W** be the substructure of **X** induced by $X' \cup \{x\}$. Since, by assumption, $\frac{\gamma}{n_{\mathbf{A}}} \geq 1$, the same argument as in the beginning of the proof shows that $f_{a'}$ is a homomorphism from **W** to **A**. Moreover, $n_{\mathbf{W}} \leq \kappa$. Using the fact that (\mathbf{X}, \mathbf{A}) has a (κ, γ) -consistency gap, we deduce that $f_{a'}$ is consistent with every substructure of **X** whose relation has at most γ tuples; i.e., $f_{a'} \in \mathscr{F}$. Since, clearly, $f_{a'}$ extends f, this means that \mathscr{F} has the extension property up to κ , as required.

In conclusion, we have shown that \mathscr{F} is a κ -strategy for (\mathbf{X}, \mathbf{A}) and, thus, $LC^{\kappa}(\mathbf{X}, \mathbf{A}) = YES$.

8 Everything together

In this section, we put together all the pieces of the puzzle and prove our main result.

Theorem (Theorem 1 restated). Let $\mathbf{A} \to \mathbf{B}$ be relational structures such that \mathbf{A} is aperiodic and \mathbf{B} is loopless. Then, PCSP(\mathbf{A}, \mathbf{B}) has linear width.

It shall be convenient to only deal with monic structures. The next result shows that this assumption does not yield a loss of generality. While it is essentially folklore, we give a self-contained proof in Appendix A for completeness.

Lemma 28. Let $\mathbf{A} \to \mathbf{B}$ be relational structures, and suppose that $PCSP(\mathbf{A}^{mon}, \mathbf{B}^{mon})$ has linear width. Then, $PCSP(\mathbf{A}, \mathbf{B})$ has linear width, too.

Let **Y** be a monic structure whose relation has arity r. For $c \in \mathbb{N}$, let $\mathbf{K}_{r,c}$ be the monic structure with domain [c] whose unique, r-ary relation is the subset of $[c]^r$ obtained by removing all constant tuples. The *chromatic number* of **Y** is the minimum c such that $\mathbf{Y} \to \mathbf{K}_{r,c}$. Note that **Y** has a finite chromatic number if and only if it is loopless.

Proof of Theorem 1. Observe that a structure \mathbf{Y} is aperiodic (resp. loopless) if and only if \mathbf{Y}^{mon} is aperiodic (resp. loopless). Hence, by virtue of Lemma 28, it is enough to prove the result in the case that \mathbf{A} and \mathbf{B} are monic structures. Let τ be the mixing time of \mathbf{A} and let c be the chromatic number of \mathbf{B} ; since \mathbf{A} is aperiodic and \mathbf{B} is loopless, these numbers are both finite. Also, let R be the unique symbol in their signature and let r be the arity of R. If r = 1 or if r = 2 and $\tau = 1$, Definition 3 (and Footnote 15) would imply that \mathbf{A} contains a loop, which is impossible since $\mathbf{A} \to \mathbf{B}$ and \mathbf{B} is loopless.

Pick a real number β such that $1 < \beta < 1 + \frac{1}{10\tau\tau}$, and apply Theorem 10 to the parameters $g = \tau$, h = c + 1, and β . Let $\delta = \delta(\tau, c + 1, \beta)$ and $n_0 = n_0(\tau, c + 1, \beta)$ be as in the statement of Theorem 10, and set

$$\epsilon = \min\left(\frac{1}{n_0}, \left(\frac{1}{10r\tau} - \beta + 1\right)\frac{(r-1)\delta}{\beta n_{\mathbf{A}}}\right).$$
(8)

Observe that $\epsilon > 0$. We claim that ϵ is a constant witnessing that $PCSP(\mathbf{A}, \mathbf{B})$ has linear width (see Section 3).

Take $n \in \mathbb{N}$. We can assume without loss of generality that $n \geq \frac{1}{\epsilon}$. Otherwise, we might simply take as **X** any instance on *n* vertices having a loop. Since **B** is loopless, $\mathbf{X} \neq \mathbf{B}$. Moreover, $LC^{\lfloor \epsilon n \rfloor}(\mathbf{X}, \mathbf{A}) = LC^0(\mathbf{X}, \mathbf{A}) = YES$, as needed. The fact that $n \geq \frac{1}{\epsilon}$ implies that $n \ge n_0$, via (8). Hence, invoking Theorem 10, we find a $(\delta n, \beta)$ -threshold-sparse r-uniform hypergraph \mathcal{H} such that $n_{\mathcal{H}} = n$, $gir(\mathcal{H}) \geq \tau$, and $chr(\mathcal{H}) \geq c+1$. Consider now a structure **X** obtained from \mathcal{H} by choosing an arbitrary orientation for each hyperedge; i.e., X has domain $V(\mathcal{H})$ and its unique relation $R^{\mathbf{X}}$ (of arity r) contains, for each $e \in E(\mathcal{H})$, precisely one r-tuple **x** such that $set(\mathbf{x}) = e$. Note that **X** is an oriented monic structure, and $\mathbf{X}^{sym} = \mathcal{H}$. We claim that the pair (\mathbf{X}, \mathbf{A}) has a $(|\epsilon n|, \delta n)$ -consistency gap. Take a homomorphism $f : \mathbf{W} \to \mathbf{A}$, where **W** is a substructure of **X** with $n_{\mathbf{W}} \leq \lfloor \epsilon n \rfloor$. Suppose, for the sake of contradiction, that there exists a substructure **Y** of **X** such that f is not consistent with **Y** and $\frac{\delta n}{n_A} < m_Y \leq \delta n$, and assume without loss of generality that \mathbf{Y} is minimal, in the sense that f is consistent with any proper substructure of **Y**. Let $\mathcal{G} = \mathbf{Y}^{\text{sym}}$, and let \mathcal{G}' be the subhypergraph of \mathcal{G} obtained by removing all isolated vertices of \mathcal{G} . By the monotonicity of threshold-sparsity, we deduce that \mathcal{G}' is hereditarily β -sparse. Moreover, its girth is at least τ . It follows from Theorem 6 that

$$\varphi_{\mathcal{G}'}^{\tau} + \pi_{\mathcal{G}'} > \left(\frac{1}{10r\tau} - \beta + 1\right) n_{\mathcal{G}'}.$$
(9)

Notice that $\varphi_{\mathcal{G}}^{\tau} = \varphi_{\mathcal{G}'}^{\tau}$ and $\pi_{\mathcal{G}} = \pi_{\mathcal{G}'}$. Moreover, by sparsity, we have that $m_{\mathcal{G}} = m_{\mathcal{G}'} < \frac{\beta}{r-1}n_{\mathcal{G}'}$. Plugging this into (9), using (8), and recalling that $\frac{\delta n}{n_{\mathbf{A}}} < m_{\mathbf{Y}} = m_{\mathcal{G}}$, we find

$$\varphi_{\mathcal{G}}^{\tau} + \pi_{\mathcal{G}} > \left(\frac{1}{10r\tau} - \beta + 1\right) \frac{r-1}{\beta} m_{\mathcal{G}} \ge \frac{\epsilon n_{\mathbf{A}} m_{\mathcal{G}}}{\delta} > \epsilon n \ge \lfloor \epsilon n \rfloor.$$

Let Ω be the union of the set of pendent hyperedges of \mathcal{G} and a maximum set of mutually disjoint τ -fibers of \mathcal{G} . Label the elements of Ω as $\omega_1, \ldots, \omega_\ell$, where $\ell = \varphi_{\mathcal{G}}^{\tau} + \pi_{\mathcal{G}}$. For $i \in [\ell]$, let J_i be a joint of ω_i . Furthermore, consider the set U_i given by $U_i = \omega_i \setminus J_i$ if ω_i is a pendent hyperedge, and $U_i = \bigcup_{e \in \omega_i} e \setminus J_i$ if ω_i is a fiber. Since at least one of $r \geq 3$ and $\tau \geq 2$ holds, the sets U_i are nonempty. Moreover, they are mutually disjoint. By the minimality of \mathbf{Y} , for each $i \in [\ell]$, f is consistent with the substructure \mathbf{Y}_i of \mathbf{Y} induced by $Y \setminus U_i$. Thus, let $h_i : \mathbf{Y}_i \to \mathbf{A}$ be a homomorphism that agrees with f on $W \cap Y_i$. Via Proposition 11 or Proposition 12, we can extend each h_i to a homomorphism $h'_i : \mathbf{Y} \to \mathbf{A}$. Since f is not consistent with \mathbf{Y} , for each $i \in [\ell]$ there exists some vertex $u_i \in W \cap U_i$ such that $f(u_i) \neq h'_i(u_i)$. In particular, this means that $n_{\mathbf{W}} \geq \ell$, as the sets U_i are mutually disjoint. We deduce that

$$n_{\mathbf{W}} \ge \ell = \varphi_{\mathcal{G}}^{\tau} + \pi_{\mathcal{G}} > \lfloor \epsilon n \rfloor,$$

a contradiction. It follows that the pair (\mathbf{X}, \mathbf{A}) has a $(\lfloor \epsilon n \rfloor, \delta n)$ -consistency gap, as claimed. Observe now that

$$\left(\frac{1}{10r\tau} - \beta + 1\right)\frac{r-1}{\beta} = \frac{1}{10\tau} \cdot \frac{r-1}{r} \cdot \frac{1 - 10r\tau(\beta - 1)}{\beta} \le 1.$$

Hence, using (8) and the assumption that $n \geq \frac{1}{\epsilon}$, we deduce that

$$n_{\mathbf{A}} \le \epsilon n n_{\mathbf{A}} \le \left(\frac{1}{10r\tau} - \beta + 1\right) \frac{(r-1)\delta}{\beta} n \le \delta n.$$

It follows from Theorem 14 that $LC^{\lfloor \epsilon n \rfloor}(\mathbf{X}, \mathbf{A}) = YES$. Finally, from $chr(\mathcal{H}) \ge c+1$ we deduce that $\mathbf{X} \not\to \mathbf{K}_{r,c}$ and, thus, $\mathbf{X} \not\to \mathbf{B}$. The proof is concluded. \Box

A Proof of Lemma 28

We prove the following result.

Lemma (Lemma 28 restated). Let $\mathbf{A} \to \mathbf{B}$ be relational structures, and suppose that $PCSP(\mathbf{A}^{mon}, \mathbf{B}^{mon})$ has linear width. Then, $PCSP(\mathbf{A}, \mathbf{B})$ has linear width, too.

We shall make use of a result from [5] based on the algebraic approach to PCSPs. First, we introduce the necessary terminology. Let σ be the common signature of **A** and **B**. For $d \in \mathbb{N}$, let \mathbf{A}^d be the *d*-th direct power of **A**; i.e., \mathbf{A}^d is the relational structure on the signature σ whose domain is A^d and whose relations are defined as follows: For $R \in \sigma$ and any $d \times \operatorname{ar}(R)$ matrix M whose rows are tuples in $R^{\mathbf{A}}$, the columns of M form a tuple in $R^{\mathbf{A}^d}$. We let $\operatorname{Pol}^{(d)}(\mathbf{A}, \mathbf{B})$ be the set of homomorphisms $\mathbf{A}^d \to \mathbf{B}$, and we let $\operatorname{Pol}(\mathbf{A}, \mathbf{B})$ (the polymorphism set of PCSP(\mathbf{A}, \mathbf{B})) be the disjoint union of $\operatorname{Pol}^{(d)}(\mathbf{A}, \mathbf{B})$ for $d \in \mathbb{N}$. Given $d, d' \in \mathbb{N}, f \in \operatorname{Pol}^{(d)}(\mathbf{A}, \mathbf{B})$, and a function $\pi : [d] \to [d']$, we define $f_{/\pi}$ (the "minor of f under π ") as the function from $A^{d'}$ to B given by $(a_1, \ldots, a_{d'}) \mapsto f(a_{\pi(1)}, \ldots, a_{\pi(d)})$. It is not hard to check that $f_{/\pi}$ yields a homomorphism $\mathbf{A}^{d'} \to \mathbf{B}$ and, thus, $f_{/\pi} \in \operatorname{Pol}^{(d')}(\mathbf{A}, \mathbf{B})$. For $\mathbf{A} \to \mathbf{B}$ and $\mathbf{A}' \to \mathbf{B}'$, a map $\xi : \operatorname{Pol}(\mathbf{A}, \mathbf{B}) \to \operatorname{Pol}(\mathbf{A}', \mathbf{B}')$ is a minion homomorphism²⁷ if it preserves arities (i.e., the range of the restriction of ξ to $\operatorname{Pol}^{(d)}(\mathbf{A}, \mathbf{B})$ is included in $\operatorname{Pol}^{(d)}(\mathbf{A}', \mathbf{B}')$) and it preserves minors (i.e., $\xi(f_{/\pi}) = \xi(f)_{/\pi}$ for each compatible f and π). Observe that the signature σ of \mathbf{A} and \mathbf{B} and the signature σ' of \mathbf{A}' and \mathbf{B}' are not required to be equal for a minion homomorphism ξ to exist.

It was shown in [5] that the property of having bounded width is preserved under minion homomorphisms. The same proof also works in the linear-width regime (as was noted in [1] for the case of sublinear width) and it yields the following statement.

Lemma 29 ([5]). Let $\mathbf{A} \to \mathbf{B}$ and $\mathbf{A}' \to \mathbf{B}'$ be relational structures such that there exists a minion homomorphism $\operatorname{Pol}(\mathbf{A}, \mathbf{B}) \to \operatorname{Pol}(\mathbf{A}', \mathbf{B}')$, and suppose that $\operatorname{PCSP}(\mathbf{A}', \mathbf{B}')$ has linear width. Then, $\operatorname{PCSP}(\mathbf{A}, \mathbf{B})$ has linear width, too.

Lemma 28 follows by comparing the polymorphisms of $PCSP(\mathbf{A}^{mon}, \mathbf{B}^{mon})$ to those of $PCSP(\mathbf{A}, \mathbf{B})$.

Proof of Lemma 28. Observe that any homomorphism f from \mathbf{A} to \mathbf{B} is also a homomorphism from \mathbf{A}^{mon} to \mathbf{B}^{mon} , as easily follows from the definition of \mathbf{A}^{mon} and \mathbf{B}^{mon} . Observe also that the structures $(\mathbf{A}^d)^{\text{mon}}$ and $(\mathbf{A}^{\text{mon}})^d$ are homomorphically equivalent for each $d \in \mathbb{N}$. Take a homomorphism $h_d : (\mathbf{A}^{\text{mon}})^d \to (\mathbf{A}^d)^{\text{mon}}$. We deduce that $f \circ h_d \in \text{Pol}^{(d)}(\mathbf{A}^{\text{mon}}, \mathbf{B}^{\text{mon}})$ for each $f \in \text{Pol}^{(d)}(\mathbf{A}, \mathbf{B})$. Moreover, it is immediate to check that the assignment $f \mapsto f \circ h_d$ preserves arities and minors, and it thus yields a minion homomorphism from Pol(\mathbf{A}, \mathbf{B}) to Pol($\mathbf{A}^{\text{mon}}, \mathbf{B}^{\text{mon}}$). The result then follows from Lemma 29.

We point out that the converse of Lemma 28 also holds, provided that all relations of \mathbf{A} are nonempty. Indeed, in that case, any homomorphism from \mathbf{A}^{mon} to \mathbf{B}^{mon} is also a homomorphism from \mathbf{A} to \mathbf{B} , and a similar argument as in the proof above shows that there exists a minion homomorphism from $\text{Pol}(\mathbf{A}^{\text{mon}}, \mathbf{B}^{\text{mon}})$ to $\text{Pol}(\mathbf{A}, \mathbf{B})$.

 $^{^{27}}$ A minion is the algebraic structure whose operations are all minor maps.

References

- Albert Atserias and Víctor Dalmau. Promise constraint satisfaction and width. In Proc. 2022 ACM-SIAM Symposium on Discrete Algorithms (SODA'22), pages 1129–1153, 2022. arXiv:2107.05886, doi:10.1137/1.9781611977073.48.
- [2] Albert Atserias and Joanna Ochremiak. Proof complexity meets algebra. ACM Trans. Comput. Log., 20(1):1:1–1:46, 2019. arXiv:1711.07320, doi:10.1145/3265985.
- [3] Per Austrin, Venkatesan Guruswami, and Johan Håstad. (2+ε)-Sat is NP-hard. SIAM J. Comput., 46(5):1554–1573, 2017. doi:10.1137/15M1006507.
- [4] László Babai, D. Yu. Grigoryev, and David M. Mount. Isomorphism of graphs with bounded eigenvalue multiplicity. In Proc. 14th Annual ACM Symposium on Theory of Computing (STOC'82), pages 310–324, 1982. doi:10.1145/800070.802206.
- [5] Libor Barto, Jakub Bulín, Andrei A. Krokhin, and Jakub Opršal. Algebraic approach to promise constraint satisfaction. J. ACM, 68(4):28:1-28:66, 2021. arXiv:1811.00970, doi:10.1145/ 3457606.
- [6] Libor Barto and Marcin Kozik. Constraint satisfaction problems solvable by local consistency methods. J. ACM, 61(1), 2014. Article No. 3. doi:10.1145/2556646.
- [7] Libor Barto and Marcin Kozik. Robustly solvable constraint satisfaction problems. SIAM J. Comput., 45(4):1646-1669, 2016. arXiv:1512.01157, doi:10.1137/130915479.
- [8] Libor Barto, Andrei Krokhin, and Ross Willard. Polymorphisms, and how to use them. In Andrei Krokhin and Stanislav Živný, editors, *The Constraint Satisfaction Problem: Complexity* and Approximability, volume 7 of Dagstuhl Follow-Ups, pages 1–44. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl, Germany, 2017. doi:10.4230/DFU.Vol7.15301.1.
- [9] Claude Berge. Hypergraphs, volume 45 of North-Holland mathematical library. North-Holland, 1989.
- [10] Abraham Berman and Robert J Plemmons. Nonnegative matrices in the mathematical sciences. SIAM, 1994.
- [11] Manuel Bodirsky, Jakub Bulín, Florian Starke, and Michael Wernthaler. The smallest hard trees. Constraints, 28(2):105–137, 2023. arXiv:2205.07528, doi:10.1007/S10601-023-09341-8.
- [12] Joshua Brakensiek and Venkatesan Guruswami. Promise constraint satisfaction: Algebraic structure and a symmetric Boolean dichotomy. SIAM J. Comput., 50(6):1663-1700, 2021. arXiv: 1704.01937, doi:10.1137/19M128212X.
- [13] Joshua Brakensiek, Venkatesan Guruswami, Marcin Wrochna, and Stanislav Zivný. The power of the combined basic LP and affine relaxation for promise CSPs. SIAM J. Comput., 49:1232–1248, 2020. arXiv:1907.04383, doi:10.1137/20M1312745.
- [14] Andrei Bulatov. Bounded relational width. Unpublished manuscript, 2009. URL: https://www2. cs.sfu.ca/~abulatov/papers/relwidth.pdf.
- [15] Andrei Bulatov, Peter Jeavons, and Andrei Krokhin. Classifying the complexity of constraints using finite algebras. SIAM J. Comput., 34(3):720–742, 2005. doi:10.1137/S0097539700376676.
- [16] Andrei A. Bulatov. A dichotomy theorem for nonuniform CSPs. In Proc. 58th Annual IEEE Symposium on Foundations of Computer Science (FOCS'17), pages 319–330, 2017. arXiv: 1703.03021, doi:10.1109/F0CS.2017.37.
- [17] Andrei A. Bulatov. Constraint Satisfaction Problems over semilattice block Mal'tsev algebras. Inf. Comput., 268, 2019. arXiv:1701.02623, doi:10.1016/J.IC.2019.104437.

- [18] Andrei A. Bulatov and Víctor Dalmau. A simple algorithm for Mal'tsev constraints. SIAM J. Comput., 36(1):16–27, 2006. doi:10.1137/050628957.
- [19] Andrei A. Bulatov, Andrei Krokhin, and Benoit Larose. Dualities for Constraint Satisfaction Problems. In Complexity of Constraints, volume 5250 of Lecture Notes in Computer Science, pages 93–124. Springer, 2008. doi:10.1007/978-3-540-92800-3_5.
- [20] Siu On Chan. Approximation resistance from pairwise-independent subgroups. J. ACM, 63(3):27:1– 27:32, 2016. doi:10.1145/2873054.
- [21] Vašek Chvátal and Endre Szemerédi. Many hard examples for resolution. J. ACM, 35(4):759–768, 1988. doi:10.1145/48014.48016.
- [22] Lorenzo Ciardo and Stanislav Živný. Approximate graph colouring and crystals. In Proc. 2023 ACM-SIAM Symposium on Discrete Algorithms (SODA'23), pages 2256–2267, 2023. arXiv: 2210.08293, doi:10.1137/1.9781611977554.ch86.
- [23] Lorenzo Ciardo and Stanislav Živný. Approximate graph colouring and the hollow shadow. In Proc. 55th Annual ACM Symposium on Theory of Computing (STOC'23), pages 623–631. ACM, 2023. arXiv:2211.03168, doi:10.1145/3564246.3585112.
- [24] Lorenzo Ciardo and Stanislav Živný. CLAP: A new algorithm for promise CSPs. SIAM J. Comput., 52(1):1–37, 2023. arXiv:2107.05018, doi:10.1137/22M1476435.
- [25] Lorenzo Ciardo and Stanislav Zivný. Semidefinite programming and linear equations vs. homomorphism problems. In Proc. 56th Annual ACM Symposium on Theory of Computing (STOC'24), pages 1935–1943. ACM, 2024. arXiv:2311.00882, doi:10.1145/3618260.3649635.
- [26] Thomas H Cormen, Charles E Leiserson, Ronald L Rivest, and Clifford Stein. Introduction to Algorithms. MIT press, 2022.
- [27] Victor Dalmau and Jakub Opršal. Local consistency as a reduction between constraint satisfaction problems. In Proc. 39th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS'24), pages 29:1–29:15. ACM, 2024. arXiv:2301.05084, doi:10.1145/3661814.3662068.
- [28] Paul Erdős and András Hajnal. On chromatic number of graphs and set-systems. Acta Math. Hung., 17(1-2):61–99, 1966. doi:10.1007/bf02020444.
- [29] Tomás Feder and Moshe Y. Vardi. The computational structure of monotone monadic SNP and constraint satisfaction: A study through Datalog and group theory. SIAM J. Comput., 28(1):57–104, 1998. doi:10.1137/S0097539794266766.
- [30] Dima Grigoriev. Linear lower bound on degrees of Positivstellensatz calculus proofs for the parity. Theor. Comput. Sci., 259(1-2):613-622, 2001. doi:10.1016/S0304-3975(00)00157-2.
- [31] Venkatesan Guruswami and Sai Sandeep. d-to-1 hardness of coloring 3-colorable graphs with O(1) colors. In Proc. 47th International Colloquium on Automata, Languages, and Programming (ICALP'20), volume 168 of LIPIcs, pages 62:1–62:12. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2020. doi:10.4230/LIPIcs.ICALP.2020.62.
- [32] Pavol Hell and Jaroslav Nešetřil. On the complexity of H-coloring. J. Comb. Theory, Ser. B, 48(1):92–110, 1990. doi:10.1016/0095-8956(90)90132-J.
- [33] D. Hobby and R.N. McKenzie. The Structure of Finite Algebras, volume 76 of Contemporary Mathematics. American Mathematical Society, Providence, R.I., 1988.
- [34] Leslie Hogben. Handbook of Linear Algebra. CRC Press, 2013.
- [35] Pawel M. Idziak, Petar Markovic, Ralph McKenzie, Matthew Valeriote, and Ross Willard. Tractability and learnability arising from algebras with few subpowers. SIAM J. Comput., 39(7):3023–3037, 2010. doi:10.1137/090775646.

- [36] Peter G. Jeavons, David A. Cohen, and Marc Gyssens. Closure properties of constraints. J. ACM, 44(4):527–548, 1997. doi:10.1145/263867.263489.
- [37] Richard M. Karp. Reducibility among combinatorial problems. In Proc. Complexity of Computer Computations, pages 85–103, 1972. doi:10.1007/978-1-4684-2001-2_9.
- [38] John G. Kemeny and Laurie J. Snell. *Finite Markov Chains*, volume 26. van Nostrand Princeton, NJ, 1969.
- [39] Subhash Khot. On the power of unique 2-prover 1-round games. In Proc. 34th Annual ACM Symposium on Theory of Computing (STOC'02), pages 767–775. ACM, 2002. doi:10.1145/ 509907.510017.
- [40] Phokion G. Kolaitis and Moshe Y. Vardi. Conjunctive-query containment and constraint satisfaction. J. Comput. Syst. Sci., 61(2):302–332, 2000. doi:10.1006/jcss.2000.1713.
- [41] Phokion G. Kolaitis and Moshe Y. Vardi. A logical approach to constraint satisfaction. In Nadia Creignou, Phokion G. Kolaitis, and Heribert Vollmer, editors, Complexity of Constraints - An Overview of Current Research Themes [Result of a Dagstuhl Seminar], volume 5250 of Lecture Notes in Computer Science, pages 125–155. Springer, 2008. doi:10.1007/978-3-540-92800-3_6.
- [42] Marcin Kozik, Andrei Krokhin, Matt Valeriote, and Ross Willard. Characterizations of several Maltsev conditions. Algebra Univers., 73(3):205–224, 2015. doi:10.1007/s00012-015-0327-2.
- [43] Andrei Krokhin, Jakub Opršal, Marcin Wrochna, and Stanislav Živný. Topology and adjunction in promise constraint satisfaction. SIAM J. Comput., 52(1):37–79, 2023. arXiv:2003.11351, doi:10.1137/20M1378223.
- [44] Alberto Larrauri and Stanislav Zivný. Solving promise equations over monoids and groups. In Proc. 51st International Colloquium on Automata, Languages, and Programming (ICALP'24), 2024. arXiv:2402.08434.
- [45] Mark Levene and George Loizou. Kemeny's constant and the random surfer. Am. Math. Mon., 109(8):741–745, 2002.
- [46] Henryk Minc. Nonnegative Matrices, volume 170. Wiley New York, 1988.
- [47] Michael Molloy and Mohammad R Salavatipour. The resolution complexity of random constraint satisfaction problems. SIAM J. Comput., 37(3):895–922, 2007. doi:10.1137/S0097539703436485.
- [48] Hans Schneider. Wielandt's proof of the exponent inequality for primitive nonnegative matrices. Linear Algebra Its Appl., 353(1-3):5–10, 2002. doi:10.1016/S0024-3795(02)00414-7.
- [49] Terence Tao and Van Vu. Random matrices have simple spectrum. Comb., 37:539–553, 2017. arXiv:1412.1438, doi:10.1007/s00493-016-3363-4.
- [50] Johan Thapper and Stanislav Živný. The power of Sherali-Adams relaxations for general-valued CSPs. SIAM J. Comput., 46(4):1241–1279, 2017. arXiv:1606.02577, doi:10.1137/16M1079245.
- [51] Johan Thapper and Stanislav Živný. The limits of SDP relaxations for general-valued CSPs. ACM Trans. Comput. Theory, 10(3):12:1–12:22, 2018. arXiv:1612.01147, doi:10.1145/3201777.
- [52] Madhur Tulsiani. CSP gaps and reductions in the Lasserre hierarchy. In Proc. 41st Annual ACM Symposium on Theory of Computing (STOC'09), pages 303–312. ACM, 2009. doi:10.1145/ 1536414.1536457.
- [53] Helmut Wielandt. Unzerlegbare, nicht negative Matrizen. Math. Z., 52(1):642-648, 1950. doi: 10.1007/BF02230720.
- [54] Dmitriy Zhuk. A proof of the CSP dichotomy conjecture. J. ACM, 67(5):30:1-30:78, 2020. arXiv:1704.01914, doi:10.1145/3402029.