

AN ENTIRE FUNCTION DEFINED BY RIEMANN

J. ARIAS DE REYNA

ABSTRACT. In one of the sheets in Riemann's Nachlass he defines an entire function and connect it with his zeta function. As in many pages in his Nachlass, Riemann is not giving complete proofs.

He obtains an L^∞ function whose Fourier transform vanish at the real values γ with $\zeta(\frac{1}{2} + i\gamma) = 0$. We give proofs of Riemann formulas.

This is an integral representation of the zeta function different from the known ones. I believe this is the first time it has been published.

1. INTRODUCTION

After the publication of Riemann's complete works, it was suspected that his posthumous writings contained an expression of $\Xi(t)$. Thanks to the contributions of the librarian Distel and of Bessel Hagen, the Riemann-Siegel formula and the integral expression $\mathcal{R}(s)$ were discovered. They were studied by Siegel in [2]. Here, we consider a different integral representation that we have found in Riemann's posthumous papers.¹

As usual in Riemann we only find a set of equations. I give a proof of his formulas and some notice of my interpretation of these formulas. Riemann defines a bounded function on \mathbf{R} whose Fourier transform vanishes at the real values γ with $\zeta(\frac{1}{2} + i\gamma) = 0$. Riemann defines an entire function

$$(1) \quad \chi(z) := \int_L \frac{e^{izw}}{e^{2\pi w^2} - 1} dw,$$

where L is a line parallel to the real axis at a height $0 < b < 2^{-1/2}$. It has the power series expansion

$$(2) \quad \chi(z) = \frac{z}{2} + \frac{1}{\sqrt{2}} \sum_{n=0}^{\infty} \frac{\zeta(n + \frac{1}{2})}{n!} \left(-\frac{z^2}{8\pi}\right)^n,$$

and satisfies the functional equation $\chi(-z) = \chi(z) - z$. And is connected with $\zeta(s)$ by

$$(3) \quad \int_0^{\infty} \chi(x) x^{2s-1} dx = \Gamma(2s) \zeta(s + \frac{1}{2}) \cos \frac{\pi(2s+1)}{4}, \quad \operatorname{Re}(s) > 0.$$

2. DEFINITION OF RIEMANN'S CHI FUNCTION.

Proposition 1. *The integral*

$$(4) \quad \chi(z) := \int_L \frac{e^{izw}}{e^{2\pi w^2} - 1} dw,$$

2020 *Mathematics Subject Classification.* Primary 11M06; Secondary 30D99.

Key words and phrases. zeta function, Riemann's auxiliar function.

¹I have been able to see Riemann's posthumous writings on number theory thanks to W. Gabcke who sent them to me with permission from the library in Gotingen. The integral expression under consideration is contained in page 00000093.jpg

where L is a line parallel to the real axis at a height $0 < b < 2^{-1/2}$ define an entire function.

Proof. For a fixed z the integrand is meromorphic in w with simple poles at $\pm\sqrt{n} e^{\pm\frac{\pi i}{4}}$ with $n \in \mathbf{N}$ and a double pole at $w = 0$.

Let $R > 0$ and $0 < \varepsilon < 2^{-1/2}$ be positive real numbers. For $|z| < R$ and $w = x + ib$ with $\varepsilon < b < 2^{-1/2} - \varepsilon$ and x real, we have

$$\left| \frac{e^{izw}}{e^{2\pi w^2} - 1} \right| \leq C e^{-\pi x^2}$$

for an adequate constant $C = C(R, \varepsilon)$. Therefore $\chi(z)$ is well defined and holomorphic in $|z| < R$. \square

3. CALCULATING THE INTEGRAL BY RESIDUES.

Riemann first obtains a series for his function $\chi(z)$.

Proposition 2. For $|\arg(z)| < \frac{\pi}{4}$ we have

$$(5) \quad \chi(z) = \frac{1}{2} \sum_{n=1}^{\infty} \left(\frac{e^{-z\sqrt{n}e^{-\pi i/4}}}{\sqrt{n}} e^{\pi i/4} + \frac{e^{-z\sqrt{n}e^{\pi i/4}}}{\sqrt{n}} e^{-\pi i/4} \right).$$

In particular, for $z = x > 0$ real

$$(6) \quad \chi(x) = \sum_{n=1}^{\infty} \frac{e^{-x\sqrt{n/2}}}{\sqrt{n}} \sin\left(\frac{\pi}{4} - x\sqrt{\frac{n}{2}}\right).$$

Proof. Let N be a non-negative integer and define b by $b\sqrt{2} = N + \frac{1}{2}$. Let L_N be the path consisting of the line $x + ib$. This line is chosen so that it passes between successive poles of the integrand defining $\chi(z)$.

By Cauchy's Theorem

$$\begin{aligned} \int_{L_0} \frac{e^{izw}}{e^{2\pi w^2} - 1} dw - \int_{L_N} \frac{e^{izw}}{e^{2\pi w^2} - 1} dw \\ = 2\pi i \sum_{n=1}^N \left(\operatorname{Res}_{w=-\sqrt{n}e^{-\pi i/4}} \frac{e^{izw}}{e^{2\pi w^2} - 1} + \operatorname{Res}_{w=\sqrt{n}e^{\pi i/4}} \frac{e^{izw}}{e^{2\pi w^2} - 1} \right). \end{aligned}$$

After some computations

$$\chi(z) = \frac{1}{2} \left(\sum_{n=1}^N \frac{e^{-z\sqrt{n}e^{-\pi i/4}}}{\sqrt{n}} e^{-\pi i/4} + \frac{e^{-z\sqrt{n}e^{\pi i/4}}}{\sqrt{n}} e^{\pi i/4} \right) + \int_{L_N} \frac{e^{izw}}{e^{2\pi w^2} - 1} dw.$$

For a fixed $z = a > 0$ we show that the integral in L_N converges to 0 when $N \rightarrow \infty$. In fact, we have

$$U_N := \int_{L_N} \frac{e^{iaw}}{e^{2\pi w^2} - 1} dw = \int_{-\infty}^{+\infty} \frac{e^{-ab} e^{iax} dx}{e^{2\pi(x^2 - b^2) + 4\pi ibx} - 1}.$$

Therefore,

$$\begin{aligned} |U_N| &\leq e^{-ab} \int_{-\infty}^{+\infty} \frac{dx}{|e^{2\pi(x^2-b^2)} - e^{-4\pi ibx}|} \\ &= be^{-ab} \int_{-\infty}^{+\infty} \frac{dx}{|e^{2\pi b^2(x^2-1)} - e^{-4\pi ib^2x}|}. \end{aligned}$$

Recall that $2b^2 = (N + \frac{1}{2})^2$. It can be shown (we will show this a little further on) that there is a constant $C > 0$ and an integer N_0 such that for $N \geq N_0$ and all x

$$(7) \quad |e^{2\pi b^2(x^2-1)} - e^{-4\pi ib^2x}| > Cb^{-2}.$$

Assuming this, we have for $N \geq N_0$

$$\begin{aligned} |U_N| &\leq be^{-ab} \frac{4}{C} b^2 + be^{-ab} \int_{|x|>2} \frac{dx}{|e^{2\pi b^2(x^2-1)} - e^{-4\pi ib^2x}|} \\ &\ll b^3 e^{-ab} + be^{-ab} \int_2^{+\infty} e^{-2\pi b^2(x^2-1)} dx. \end{aligned}$$

Since this tends to 0 for $b \rightarrow +\infty$, we have $\lim_{N \rightarrow \infty} U_N = 0$.

To prove (7) observe that (7) is true, with $C = 1$ when $|e^{2\pi b^2(x^2-1)} - 1| \geq b^{-2}$. Therefore, we may assume that $|e^{2\pi b^2(x^2-1)} - 1| < b^{-2}$. In this case, for $n \geq N_0$, we have

$$-\frac{1}{2}b^{-2} < \log(1 - b^{-2}) < 2\pi b^2(x^2 - 1) < \log(1 + b^{-2}) < 2b^{-2}.$$

This implies that $|x - 1| \leq cb^{-4}$ or $|x + 1| \leq cb^{-4}$ for some absolute constant $c > 0$. This implies that $e^{4\pi ib^2x}$ is very near i or $-i$, for example, in the first case $x = 1 + \theta b^{-4}$, for some real θ with $|\theta| \leq c$

$$e^{4\pi ib^2x} = e^{2\pi i(N^2 + N + \frac{1}{4})(1 + \theta b^{-4})} = e^{\frac{\pi i}{2} + 4\pi i\theta b^{-2}},$$

so that for a new absolute constant $c > 0$ we have $|e^{4\pi ib^2x} - i| \leq cb^{-2}$. It follows that

$$\begin{aligned} |e^{2\pi b^2(x^2-1)} - e^{-4\pi ib^2x}| &= |1 - i + (e^{2\pi b^2(x^2-1)} - 1) - (e^{-4\pi ib^2x} - i)| \\ &\geq \sqrt{2} - b^{-2} - cb^{-2} \geq \frac{1}{2} > Cb^{-2}. \end{aligned}$$

This proof (5) when z is real and positive. But it is easy to see that both members are well defined and holomorphic in $|\arg(z)| < \pi/4$. So the equality extends to this range. Equation (6) is obtained from (5) by using the definition of the sin function. It is also valid in the same range, although we state it only for $x > 0$ real. \square

Proposition 3. For $x > 0$ and any natural number N we have

$$(8) \quad \chi(x) = \sum_{n=1}^N \frac{e^{-x\sqrt{n/2}}}{\sqrt{n}} \sin\left(\frac{\pi}{4} - x\sqrt{\frac{n}{2}}\right) + R_N,$$

with

$$|R_N| \leq \left(\frac{1}{\sqrt{N+1}} + \frac{2\sqrt{2}}{x}\right) e^{-x\sqrt{(N+1)/2}}.$$

Proof. By (6) we have

$$|R_n| = \left| \sum_{n=N+1}^{\infty} \frac{e^{-x\sqrt{n/2}}}{\sqrt{n}} \sin\left(\frac{\pi}{4} - x\sqrt{\frac{n}{2}}\right) \right| \leq \sum_{n=N+1}^{\infty} \frac{e^{-x\sqrt{n/2}}}{\sqrt{n}}.$$

Since the function $t \mapsto \frac{e^{-x\sqrt{t/2}}}{\sqrt{t}}$ is decreasing, we have

$$|R_N| \leq \frac{e^{-x\sqrt{(N+1)/2}}}{\sqrt{N+1}} + \int_{N+1}^{\infty} \frac{e^{-x\sqrt{t/2}}}{\sqrt{t}} dt = \frac{e^{-x\sqrt{(N+1)/2}}}{\sqrt{N+1}} + \frac{2\sqrt{2}}{x} e^{-x\sqrt{(N+1)/2}}.$$

□

4. AN ALTERNATIVE INTEGRAL REPRESENTATION.

Proposition 4. *For any $z \in \mathbf{C}$ we have*

$$(9) \quad \chi(z) = \frac{\zeta(1/2)}{\sqrt{2}} + \frac{z}{2} + 2 \int_0^{\infty} \frac{\cos(xz) - 1}{e^{2\pi x^2} - 1} dx.$$

Proof. Let $0 < \varepsilon < 1$ be a given real number. By Cauchy's Theorem, the integration path in the definition (4) of $\chi(z)$ can be transformed into the path consisting of the portion $(-\infty, -\varepsilon)$ of the negative real axis, followed by the semicircle of center 0 and radius ε above the real axis, and the portion (ε, ∞) of the real axis. This gives us the equality

$$\chi(z) = \int_{\varepsilon}^{\infty} \frac{e^{-zxi}}{e^{2\pi x^2} - 1} dx - \int_{C_{\varepsilon}} \frac{e^{zwi}}{e^{2\pi w^2} - 1} dw + \int_{\varepsilon}^{\infty} \frac{e^{zxi}}{e^{2\pi x^2} - 1} dx.$$

For $|w| < 1$ we have

$$\frac{e^{zwi}}{e^{2\pi w^2} - 1} = \frac{1}{2\pi w^2} + \frac{iz}{2\pi w} + g(w),$$

where $g(w)$ is holomorphic. Therefore,

$$\int_{C_{\varepsilon}} \frac{e^{zwi}}{e^{2\pi w^2} - 1} dw = \frac{1}{\pi\varepsilon} - \frac{z}{2} + G(\varepsilon),$$

with $\lim_{\varepsilon \rightarrow 0} G(\varepsilon) = 0$. So, we obtain

$$\chi(z) = 2 \int_{\varepsilon}^{\infty} \frac{\cos(xz)}{e^{2\pi x^2} - 1} dx - \frac{1}{\pi\varepsilon} + \frac{z}{2} - G(\varepsilon).$$

We can write this in the form

$$\chi(z) = 2 \int_{\varepsilon}^{\infty} \frac{\cos(xz) - 1}{e^{2\pi x^2} - 1} dx + 2 \int_{\varepsilon}^{\infty} \left(\frac{1}{e^{2\pi x^2} - 1} - \frac{1}{2\pi x^2} \right) dx + \frac{z}{2} - G(\varepsilon).$$

Taking the limits when $\varepsilon \rightarrow 0^+$ we obtain

$$\chi(z) = 2 \int_0^{\infty} \frac{\cos(xz) - 1}{e^{2\pi x^2} - 1} dx + 2 \int_0^{\infty} \left(\frac{1}{e^{2\pi x^2} - 1} - \frac{1}{2\pi x^2} \right) dx + \frac{z}{2}.$$

The second integral is transformed by a change of variables, and the resulting integral is found in Titchmarsh [3, eq. (2.7.1)]

$$2 \int_0^{\infty} \left(\frac{1}{e^{2\pi x^2} - 1} - \frac{1}{2\pi x^2} \right) dx = \frac{1}{\sqrt{2\pi}} \int_0^{\infty} \left(\frac{1}{e^y - 1} - \frac{1}{y} \right) \frac{dy}{\sqrt{y}} = \frac{\zeta(1/2)\Gamma(1/2)}{\sqrt{2\pi}}.$$

□

Proposition 5. *The function $\chi(z)$ has the following power series expansion*

$$(10) \quad \chi(z) = \frac{z}{2} + \frac{1}{\sqrt{2}} \sum_{n=0}^{\infty} \frac{\zeta(n + \frac{1}{2})}{n!} \left(-\frac{z^2}{8\pi}\right)^n.$$

Proof. In the integral in equation (9) expand the cosine in power series. It is easy to justify integration term by term. We obtain

$$\chi(z) = \frac{\zeta(1/2)}{\sqrt{2}} + \frac{z}{2} + \sum_{n=1}^{\infty} (-1)^n \frac{z^{2n}}{(2n)!} \cdot 2 \int_0^{\infty} \frac{x^{2n}}{e^{2\pi x^2} - 1} dx.$$

The integral can be computed by transforming it by a change of variables to the integral in Titchmarsh [3, eq. (2.4.1)], and then substituting the known value of $\Gamma(n + \frac{1}{2})$

$$\begin{aligned} 2 \int_0^{\infty} \frac{x^{2n}}{e^{2\pi x^2} - 1} dx &= \frac{2}{(2\pi)^{n+1/2}} \int_0^{\infty} \frac{y^n}{e^y - 1} \frac{dy}{2\sqrt{y}} = \frac{1}{(2\pi)^{n+1/2}} \int_0^{\infty} \frac{y^{n+\frac{1}{2}}}{e^y - 1} \frac{dy}{y} \\ &= \frac{\Gamma(n + \frac{1}{2})\zeta(n + \frac{1}{2})}{(2\pi)^{n+1/2}} = \frac{\zeta(n + \frac{1}{2})}{(2\pi)^{n+1/2}} \frac{(2n)! \sqrt{\pi}}{n! 2^{2n}}. \end{aligned}$$

Therefore,

$$\begin{aligned} \chi(z) &= \frac{\zeta(1/2)}{\sqrt{2}} + \frac{z}{2} + \sum_{n=1}^{\infty} (-1)^n \frac{z^{2n}}{(2n)!} \cdot \frac{\zeta(n + \frac{1}{2})}{(2\pi)^{n+1/2}} \frac{(2n)! \sqrt{\pi}}{n! 2^{2n}} \\ &= \frac{\zeta(1/2)}{\sqrt{2}} + \frac{z}{2} + \frac{1}{\sqrt{2}} \sum_{n=1}^{\infty} \frac{\zeta(n + \frac{1}{2})}{n!} \left(-\frac{z^2}{8\pi}\right)^n \\ &= \frac{z}{2} + \frac{1}{\sqrt{2}} \sum_{n=0}^{\infty} \frac{\zeta(n + \frac{1}{2})}{n!} \left(-\frac{z^2}{8\pi}\right)^n. \end{aligned} \quad \square$$

Remark 6. Riemann ‘derives’ the power series from (5) by means of an ‘Eulerian’ computation, that I cannot resist duplicating:

$$\begin{aligned} \chi(z) &= \frac{1}{2} \sum_{n=1}^{\infty} \left(\frac{e^{z\sqrt{n}e^{3\pi i/4}}}{\sqrt{n}} e^{\pi i/4} + \frac{e^{z\sqrt{n}e^{\pi i/4}}}{\sqrt{n}} e^{-3\pi i/4} \right) \\ &= \frac{1}{2} \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \frac{z^m n^{\frac{m-1}{2}}}{m!} \left(e^{\frac{3\pi i}{4}m + \frac{\pi i}{4}} + e^{-\frac{3\pi i}{4}m - \frac{\pi i}{4}} \right) \\ &= \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \frac{z^m n^{\frac{m-1}{2}}}{m!} \cos\left(\frac{3m+1}{4}\pi\right) \\ &= \sum_{m=0}^{\infty} \frac{z^m \zeta\left(\frac{1-m}{2}\right)}{m!} \cos\left(\frac{3m+1}{4}\pi\right). \end{aligned}$$

Naturally, this is not correct, but similar calculations lead to good results on many occasions. We should not underestimate Riemann, it is quite possible that he knew a general method to justify this type of reasoning. The power series obtained by Riemann is equivalent to (10).

$$(11) \quad \chi(z) = \frac{z}{2} + \frac{1}{\sqrt{2}} \sum_{n=0}^{\infty} \frac{\zeta(n + \frac{1}{2})}{n!} \left(-\frac{z^2}{8\pi}\right)^n = \sum_{m=0}^{\infty} \frac{z^m \zeta\left(\frac{1-m}{2}\right)}{m!} \cos\left(\frac{3m+1}{4}\pi\right),$$

as can be checked easily by means of the functional equation.

Notice that we also obtain the curious equality

$$(12) \quad \chi^{(m)}(0) = \zeta\left(\frac{1-m}{2}\right) \cos\left(\frac{3m+1}{4}\pi\right).$$

Corollary 7. *The function $\chi(z) - \frac{z}{2}$ is even so that the function $\chi(z)$ satisfies the functional equation*

$$(13) \quad \chi(-z) = \chi(z) - z.$$

5. CONNECTION TO THE ZETA FUNCTION.

Proposition 8. *We have*

$$(14) \quad \int_0^\infty \chi(x)x^{2s-1} dx = \Gamma(2s)\zeta\left(s + \frac{1}{2}\right) \cos\frac{\pi(2s+1)}{4}, \quad \operatorname{Re}(s) > 0.$$

Proof. By (8) with $N = 1$ we have $|\chi(x)| \leq (1 + \frac{4}{x})e^{-x/\sqrt{2}}$ for $x > 1$ and there is a constant M such that $|\chi(x)| \leq M$ for $0 < x < 1$. This implies that the left member of (14) defines an analytic function for $\sigma > 0$. Therefore, we only have to show the equality for $\sigma = \operatorname{Re}(s) > 1/2$. By (5)

$$\int_0^\infty \chi(x)x^{2s} \frac{dx}{x} = \int_0^\infty \frac{1}{2} \sum_{n=1}^\infty \left(\frac{e^{-x\sqrt{n}e^{-\pi i/4}}}{\sqrt{n}} e^{\pi i/4} + \frac{e^{-x\sqrt{n}e^{\pi i/4}}}{\sqrt{n}} e^{-\pi i/4} \right) x^{2s} \frac{dx}{x}.$$

We may interchange sum and integral (for example, taking in absolute value all the terms and integrating we obtain a convergent series), therefore, we obtain

$$\begin{aligned} \int_0^\infty \chi(x)x^{2s-1} dx &= \frac{1}{2} \sum_{n=1}^\infty \left(\frac{e^{\pi i/4}}{\sqrt{n}} \int_0^\infty e^{-x\sqrt{n}e^{-\pi i/4}} x^{2s-1} dx + \frac{e^{-\pi i/4}}{\sqrt{n}} \int_0^\infty e^{-x\sqrt{n}e^{\pi i/4}} x^{2s-1} dx \right). \end{aligned}$$

These integrals are elementary, and we obtain for $\sigma > 1/2$

$$\begin{aligned} \int_0^\infty \chi(x)x^{2s-1} dx &= \frac{1}{2} \sum_{n=1}^\infty \left(\frac{e^{\pi i/4}}{\sqrt{n}} \frac{\Gamma(2s)}{n^s} e^{\frac{\pi i s}{2}} + \frac{e^{-\pi i/4}}{\sqrt{n}} \frac{\Gamma(2s)}{n^s} e^{-\frac{\pi i s}{2}} \right) \\ &= \Gamma(2s)\zeta\left(s + \frac{1}{2}\right) \cos\left(\frac{\pi}{4} + \frac{\pi s}{2}\right). \quad \square \end{aligned}$$

6. APPLICATION OF EULER-MACLAURIN EXPANSION.

Equation (5) is well suited to apply the Euler-Maclaurin expansion, which we quote from [1, p. 104]:

Proposition 9. *Suppose that M is a positive integer and that f has continuous derivatives through the $2M$ th order in the interval $[a, b]$ where a and b are integers with $a < b$. Then,*

$$(15) \quad \sum_{n=a}^b f(n) = \int_a^b f(x) dx + \frac{f(a) + f(b)}{2} + \sum_{m=1}^M \frac{B_{2m}}{(2m)!} (f^{(2m-1)}(b) - f^{(2m-1)}(a)) \\ - \frac{1}{(2M)!} \int_a^b f^{(2M)}(x) \tilde{B}_{2M}(x) dx.$$

Here, B_m are the Bernoulli numbers

$$B_0 = 1, \quad B_1 = -\frac{1}{2}, \quad B_2 = \frac{1}{6}, \quad B_4 = -\frac{1}{30}, \quad \dots, \quad B_{2n+1} = 0, \quad n \geq 1$$

We also notice the inequality

$$\max_x |\tilde{B}_{2n}(x)| = |B_{2n}|.$$

Proposition 10. For $x > 0$ real and any natural number N we have

$$(16) \quad \chi(x) = \sum_{n=1}^N \frac{e^{-x\sqrt{n/2}}}{\sqrt{n}} \sin\left(\frac{\pi}{4} - x\sqrt{n/2}\right) - \frac{1}{2} \frac{e^{-x\sqrt{N/2}}}{\sqrt{N}} \sin\left(\frac{\pi}{4} - x\sqrt{N/2}\right) \\ - \frac{2}{x} e^{-x\sqrt{N/2}} \sin(x\sqrt{N/2}) + \frac{x}{24N} e^{-x\sqrt{N/2}} \cos(x\sqrt{N/2}) \\ + \frac{1}{24N^{3/3}} e^{-x\sqrt{N/2}} \cos\left(x\sqrt{N/2} + \frac{\pi}{4}\right) + R(x, N),$$

with

$$|R(x, N)| \leq \frac{e^{-x\sqrt{N/2}}}{24\sqrt{N}} (x + N^{-1/2})^2.$$

Proof. In Proposition 9 take

$$f(t) = \frac{1}{2} \left(\frac{e^{-x\sqrt{t}e^{-\pi i/4}}}{\sqrt{t}} e^{\pi i/4} + \frac{e^{-x\sqrt{t}e^{\pi i/4}}}{\sqrt{t}} e^{-\pi i/4} \right) = \operatorname{Re} \left\{ \frac{e^{-x\sqrt{t}e^{-\pi i/4}}}{\sqrt{t}} e^{\pi i/4} \right\}.$$

For $x > 0$ we have

$$\sum_{n=N+1}^{\infty} f(n) = \chi(x) - \sum_{n=1}^N f(n).$$

By Euler-Maclaurin (taking $M = 1$, $a = N$ and $b = L$)

$$\sum_{n=N}^L f(n) = \int_N^L f(t) dt + \frac{f(N) + f(L)}{2} + \frac{B_2}{2} (f'(L) - f'(N)) - R(x, N, L),$$

where

$$R(x, N, L) = \frac{1}{2} \int_N^L f''(t) \tilde{B}_2(t) dt.$$

Taking the limit for $L \rightarrow \infty$ yields

$$\chi(x) - \sum_{n=1}^N f(n) = \int_N^{\infty} f(t) dt - \frac{f(N)}{2} - \frac{f'(N)}{12} - \frac{1}{2} \int_N^{\infty} f''(t) \tilde{B}_2(t) dt.$$

Since

$$f''(t) = \operatorname{Re} \left\{ \frac{e^{x\sqrt{\frac{t}{2}}(-1+i)}}{4t^{3/2}} \left(\frac{3e^{\frac{\pi}{4}i}}{t} + \frac{3x}{\sqrt{t}} + x^2 e^{\frac{7\pi}{4}i} \right) \right\},$$

we have

$$\begin{aligned} |R(x, N)| &\leq \frac{1}{12} \int_N^\infty e^{-x\sqrt{t/2}} \left(\frac{3}{4}t^{-5/2} + \frac{3x}{4}t^{-2} + \frac{x^2}{4}t^{-3/2} \right) dt \\ &\leq \frac{1}{48} e^{-x\sqrt{N/2}} (2N^{-3/2} + 3xN^{-1} + 2x^2N^{-1/2}) \\ &\leq \frac{1}{24} e^{-x\sqrt{N/2}} (N^{-3/4} + xN^{-1/4})^2. \end{aligned}$$

The integral of $f(t)$ can be computed and we obtain

$$\begin{aligned} \chi(x) &= \frac{1}{2} \sum_{n=1}^N \left(\frac{e^{-x\sqrt{n}e^{-\pi i/4}}}{\sqrt{n}} e^{\pi i/4} + \frac{e^{-x\sqrt{n}e^{\pi i/4}}}{\sqrt{n}} e^{-\pi i/4} \right) \\ &\quad + \frac{e^{-x\sqrt{N}e^{-\pi i/4}}}{x} i - \frac{e^{-x\sqrt{N}e^{\pi i/4}}}{x} i - \frac{1}{4} \left(\frac{e^{-x\sqrt{N}e^{-\pi i/4}}}{\sqrt{N}} e^{\pi i/4} + \frac{e^{-x\sqrt{N}e^{\pi i/4}}}{\sqrt{N}} e^{-\pi i/4} \right) \\ &\quad + \frac{1}{12} \operatorname{Re} \left\{ \frac{e^{-x\sqrt{N}e^{-\pi i/4}}}{2N} \left(\frac{e^{\frac{\pi}{4}i}}{\sqrt{N}} + x \right) \right\} - R(x, N). \quad \square \end{aligned}$$

Remark 11. Riemann substitutes in (14) the asymptotic expansion for $\chi(s)$ obtained in Proposition 10, integrates term by term to get an asymptotic expansion for $\zeta(s + \frac{1}{2})$. But he only gets the Euler-Maclaurin expansion of $\zeta(s + \frac{1}{2})$.

Riemann uses the Euler Maclaurin expansion to compute $\chi(0) = \zeta(1/2)/\sqrt{2}$. Which we have computed in Section 4 by other means.

Taking the limits when $x \rightarrow 0^+$ in (16) he obtains:

$$\chi(0) = \sum_{n=1}^N \frac{1}{\sqrt{n}} \frac{1}{\sqrt{2}} - \frac{1}{2\sqrt{2N}} - \sqrt{2N} + \frac{1}{24\sqrt{2}N^{3/2}} + R(0, N).$$

Therefore,

$$\chi(0) = \lim_{N \rightarrow \infty} \left(\frac{1}{\sqrt{2}} \sum_{n=1}^N \frac{1}{\sqrt{n}} - \sqrt{2N} \right).$$

Riemann recognizes the limit as $\zeta(1/2)/\sqrt{2}$.

7. A LAST REPRESENTATION OF χ .

Proposition 12. For any $z \in \mathbf{C}$ we have

$$(17) \quad \chi(z) = \frac{\zeta(\frac{1}{2})}{\sqrt{2}} + \frac{z}{2} + \sum_{n=1}^{\infty} \frac{e^{-\frac{z^2}{8\pi n}} - 1}{\sqrt{2n}}.$$

Proof. This can be obtained directly from the power series (10). Riemann does not write (10) only his equivalent form in (11), therefore, I try to follow what appear to be Riemann path to this series expansion.

Differentiating (4) we obtain

$$(18) \quad \chi''(z) = - \int_L \frac{w^2 e^{izw}}{e^{2\pi w^2} - 1} dw.$$

The integrand has no singularity at $w = 0$, therefore we may change the line of integration to the real line,

$$\chi''(z) = - \int_{-\infty}^{\infty} \frac{x^2 e^{izx}}{e^{2\pi x^2} - 1} dx.$$

From which we obtain easily

$$\chi''(z) = - \sum_{n=1}^{\infty} \int_{-\infty}^{\infty} x^2 e^{izx} e^{-2\pi n x^2} dx = - \sum_{n=1}^{\infty} \frac{e^{-\frac{z^2}{8\pi n}} (4\pi n - z^2)}{16\pi^2 \sqrt{2n^{5/2}}}.$$




FIGURE 1. X-ray of $\chi(s)$ on $(-50, 50)^2$

We have

$$\frac{\partial^2}{\partial z^2} \frac{e^{-\frac{z^2}{8\pi n}} - 1}{\sqrt{2n}} = - \frac{e^{-\frac{z^2}{8\pi n}} (4\pi n - z^2)}{16\pi^2 \sqrt{2n^{5/2}}}$$

Since the series in (17) converges uniformly in compact sets, we derive from this that

$$\chi(z) = az + b + \sum_{n=1}^{\infty} \frac{e^{-\frac{z^2}{8\pi n}} - 1}{\sqrt{2n}},$$

for some adequate constants a and b . Taking $z = 0$, we obtain $b = \chi(0)$ and it is also clear that $\chi'(0) = a$. This proves (17). \square

8. X-RAY AND ZEROS.

We may use the power series to compute $\chi(z)$ for $|z| \leq r$. Since there is a large cancellation between the terms of the power series, we have to use large precision in computing the terms. Thus, we have been able to compute $\chi(z)$ in the square $(-50, 50)^2$. This allows us to compute the x-ray of this function in this square.

Taking $N = 1$ in Proposition 3 we have for $x > 0$

$$\chi(x) \sim e^{-x/\sqrt{2}} \sin\left(\frac{\pi}{4} - \frac{x}{\sqrt{2}}\right) + O(e^{-x}).$$

Therefore, the real zeros are approximately equal to $a_n \sim \pi\sqrt{2}(n + \frac{1}{4})$. In practice, this is a very good approximation; for example, for $n = 10$ the difference a_{10} and its approximation is 1.12×10^{-6} .

The zeros in the direction $e^{3\pi i/4}$ are approximately situated at $4\pi\sqrt{n}e^{3\pi i/4}$. But this is only a guess founded on the numbers, and it is not so good as in the real case.

The zeros in the direction $e^{\pi i/4}$ are more difficult to predict because some of them *are missing* because they are *migrated* to the real line.

REFERENCES

- [1] H. M. EDWARDS, *Riemann's Zeta function*, Academic Press, New York, 1974. (Reprinted by Dover Publications, Mineola, N.Y. 2001.)
- [2] C. L. SIEGEL, *Über Riemann's Nachlaß zur analytischen Zahlentheorie*, *Quellen und Studien zur Geschichte der Mathematik, Astronomie und Physik* **2** (1932), 45–80. [English version](#).
- [3] E. C. Titchmarsh, *The theory of the Riemann zeta-function*, 2nd ed. with a preface by D. R. Heath-Brown, Oxford University Press, New York, 1986.

UNIVERSIDAD DE SEVILLA, FACULTAD DE MATEMÁTICAS, C/TARFIA, SN, 41012-SEVILLA, SPAIN.
Email address: arias@us.es, ariasdereyna1947@gmail.com