

CANONICAL HEIGHTS, PERIODS AND THE HURWITZ ZETA FUNCTION

ROLF ANDREASSON, ROBERT J. BERMAN

ABSTRACT. Let $(\mathcal{X}, \mathcal{D})$ be a projective log pair over the ring of integers of a number field such that the log canonical line bundle $\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$, or its dual $-\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$, is relatively ample. We introduce a canonical height of $\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$, which is finite precisely when the complexifications of $(\mathcal{X}, \mathcal{D})$ are K-semistable. When the complexifications of $(\mathcal{X}, \mathcal{D})$ are K-polystable, the canonical height is the height of $\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ wrt any volume-normalized Kähler-Einstein metric on the complexifications of $\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$. The canonical height is shown to have a number of useful variational properties. Moreover, it may be expressed as a limit of periods on the N -fold products of the complexifications of \mathcal{X} , as N tends to infinity. In particular, using this limit formula, the canonical height for the arithmetic log surfaces $(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D})$, where \mathcal{D} has at most three components, is computed explicitly in terms of the Hurwitz zeta function and its derivative at $s = -1$. Combining this explicit formula with previous height formulas for quaternionic Shimura curves yields a procedure for extracting information about the canonical integral models of some Shimura curves, such as wild ramification. Furthermore, explicit formulas for the canonical height of twisted Fermat curves are obtained, implying explicit Parshin type bounds for the Arakelov metric.

1. INTRODUCTION

Let $(\mathcal{X}, \mathcal{L})$ be a polarized arithmetic variety over the ring of integers $\mathcal{O}_{\mathbb{F}}$ of a number field \mathbb{F} , i.e. a projective flat scheme \mathcal{X} over $\mathcal{O}_{\mathbb{F}}$ of relative dimension n , endowed with a relatively ample line bundle \mathcal{L} . A key role in arithmetic geometry is played by the *height* $h_{\|\cdot\|}(\mathcal{X}, \mathcal{L})$ of $(\mathcal{X}, \mathcal{L})$, which is defined with respect to a given metric $\|\cdot\|$ on the \mathbb{C} -points of $\mathcal{L} \rightarrow \mathcal{X}$. The height is a measure of the arithmetic complexity of $(\mathcal{X}, \mathcal{L})$ and may be defined in terms of arithmetic intersection theory in the context of Arakelov geometry [52, 45, 20]:

$$h_{\|\cdot\|}(\mathcal{X}, \mathcal{L}) := \overline{\mathcal{L}}^{n+1}, \quad \hat{h}_{\|\cdot\|}(\mathcal{X}, \mathcal{L}) := \overline{\mathcal{L}}^{n+1} / ([\mathbb{F} : \mathbb{Q}] L_{\mathbb{F}}^n(n+1))$$

where $\overline{\mathcal{L}}$ is a shorthand for the metrized line bundle $(\mathcal{L}, \|\cdot\|)$ and $\hat{h}_{\|\cdot\|}(\mathcal{X}, \mathcal{L})$ is called the *normalized height* (which is invariant under base change). In the classical “geometric analog” - where the scheme \mathcal{X} over \mathbb{Z} corresponds to a fibration $\mathcal{X} \rightarrow \mathbb{A}_{\mathbb{C}}^1$ over the complex affine line - the role of the metric $\|\cdot\|$ is played by an extension $(\overline{\mathcal{X}}, \overline{\mathcal{L}})$ of $(\mathcal{X}, \mathcal{L})$ over the compactification $\mathbb{P}_{\mathbb{C}}^1$ of $\mathbb{A}_{\mathbb{C}}^1$. Accordingly, the role of the height is played by the degree of $(\overline{\mathcal{X}}, \overline{\mathcal{L}})$.

From an adelic perspective a metrized polarized arithmetic variety $(\mathcal{X}, \mathcal{L}, \|\cdot\|)$ over $\mathcal{O}_{\mathbb{F}}$ induces an adelic metric on the adelic extension of the line bundle $L_{\mathbb{F}} \rightarrow X_{\mathbb{F}}$ (determined

by $(\mathcal{X}, \mathcal{L})$ and $\|\cdot\|$ over the finite and infinite places of \mathbb{F} , respectively) [106]. In the case when $X_{\mathbb{F}}$ is a curve and $K_{X_{\mathbb{F}}}$ is ample there is - after performing a base change - a canonical relatively ample model $(\mathcal{X}, \mathcal{K}_{\mathcal{X}})$ of (X, K_X) over $\mathcal{O}_{\mathbb{F}}$, which is *stable* (in the sense of Deligne-Mumford [38]). It was proposed by Manin [81], and along similar lines by Bost [21] and Zhang [107], that the analog, over the infinite places of \mathbb{F} of such a canonical model is played by a *Kähler-Einstein metric*, i.e. a Kähler metric ω on the complex points $X(\mathbb{C})$ with constant Ricci curvature (which is negative when $K_{X_{\mathbb{F}}}$ is ample). Manin’s proposal can be made precise using work of Odaka [86], that will be further developed here in the more general context of log pairs (see Remark 3.11).

We will be mainly concerned with the case where \mathcal{L} is the *log canonical line bundle* $\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ or its dual $-\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ of a *log pair* $(\mathcal{X}, \mathcal{D})$ in the usual sense of the Minimal Model Program in birational geometry (recalled in Section 2.1). This means that \mathcal{D} is an effective \mathbb{R} -divisor on \mathcal{X}

$$\mathcal{D} = \sum w_i \mathcal{D}_i, \text{ and } \mathcal{K}_{(\mathcal{X}, \mathcal{D})} := \mathcal{K}_{\mathcal{X}} + \mathcal{D},$$

where \mathcal{D}_i are irreducible effective divisors and w_i are non-negative real numbers. The main arithmetic applications concern the “orbifold/cusp” case, where \mathcal{D} has simple normal crossings and $w_i = 1 - 1/m_i$ for “ramification indices” $m_i \in \mathbb{N} \cup \{\infty\}$. However, allowing general non-negative coefficients w_i will be important in order to apply variational arguments.

1.1. Motivation. In contrast to algebraic degrees, heights can rarely be computed explicitly. But for special arithmetic varieties \mathcal{X} and metrics - typically admitting a modular interpretation - it has been conjectured that the corresponding normalized height can be computed explicitly in terms of special values of logarithmic derivatives of Dedekind zeta functions $\zeta_{\mathbb{F}}$ (and, more generally, of Artin L-functions). [79, 80, 67]. Very recently, the case of quaternionic Shimura curves over a totally real field \mathbb{F} was settled by Yuan [103]. The proof, that builds on [101, 102], uses automorphic forms over the adelic group $GL_2(\mathbb{A}_{\mathbb{F}})$ (the case of the classical modular curve was first shown by Bost and Kuhn [72], using modular forms, and the case when $\mathbb{F} = \mathbb{Q}$ was established by Kudla-Rapoport-Yang [68, Thm 1.0.5] in the context of Kudla’s program). Such a Shimura curve defines a log pair (X, Δ) over \mathbb{F} , whose log canonical line bundle $K_{(X, \Delta)}$ is ample. Moreover, by [102], X admits a canonical model \mathcal{X} over $\mathcal{O}_{\mathbb{F}}$. Likewise, $K_{(X, \Delta)}$ admits a canonical model over $\mathcal{O}_{\mathbb{F}}$ called the *Hodge bundle* [102], which is endowed with the *Petersson metric*. The Hodge bundle may be identified with the log canonical line bundle $\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ for a canonical effective \mathbb{Q} -divisor \mathcal{D} on \mathcal{X} . Moreover, the Petersson metric can be characterized as the unique Kähler-Einstein metric on $K_{(X, \Delta)}$ with volume $\pi \deg K_{(X, \Delta)}/2$ (Lemma 9.2). Yuan’s formula [103] for the normalized height of $(\mathcal{X}, \overline{\mathcal{K}_{(\mathcal{X}, \mathcal{D})}})$ reads as follows:

$$(1.1) \quad \hat{h}_{\text{Pet}}(\overline{\mathcal{K}_{(\mathcal{X}, \mathcal{D})}}) = -\frac{1}{2} - \frac{1}{[\mathbb{F} : \mathbb{Q}]} \frac{\zeta'_{\mathbb{F}}(-1)}{\zeta_{\mathbb{F}}(-1)} + \frac{1}{[\mathbb{F} : \mathbb{Q}]} \frac{3N(\mathfrak{p}) - 1}{4(N(\mathfrak{p}) - 1)} \sum_{\mathfrak{p}} \log N(\mathfrak{p})$$

where \mathfrak{p} ranges over the prime ideals in $\mathcal{O}_{\mathbb{F}}$ which are in the ramification locus of the corresponding quaternion algebra and $N(\mathfrak{p}) := \sharp(\mathcal{O}_{\mathbb{F}}/\mathfrak{p})$. When $\mathbb{F} = \mathbb{Q}$, \mathcal{X} is the coarse

moduli scheme parametrizing all Abelian surfaces over \mathbb{Z} with quaternion multiplication [104] and the divisor \mathcal{D} on \mathcal{X} is the Zariski closure of Δ (see Example 9.1).

For general heights, all one can hope for is to obtain explicit bounds. For example, as shown in the early days of Arakelov geometry by Parshin [88] (and further developed in [99, 83]) an effective version of the Mordell conjecture (i.e. an effective bound on the height of \mathbb{F} -points of a given curve $X_{\mathbb{F}}$ with $K_{X_{\mathbb{F}}}$ ample) would follow from a height inequality for stable arithmetic surfaces \mathcal{X} (in the sense of Deligne-Mumford [38]), when $K_{X(\mathbb{C})}$ is endowed with the *Arakelov metric*

$$(1.2) \quad \hat{h}_{\text{Ar}}(\overline{\mathcal{K}_{\mathcal{X}}}) \leq c_0 + c_1 \sum_{\mathfrak{p} \text{ bad}} \log N(\mathfrak{p}) + c_2 \log |D_{\mathbb{F}}|$$

for constants c_1, c_2 independent of X and a constant c_0 depending on X , summing over closed points $\mathfrak{p} \in \text{Spec } \mathcal{O}_{\mathbb{F}}$ of bad reduction and $D_{\mathbb{F}}$ denotes the discriminant of \mathbb{F} (in particular, $\log |D_{\mathbb{F}}| \geq 0$ with equality when $\mathbb{F} = \mathbb{Q}$). The geometric analog of this inequality follows, with explicit constants, from the Miyaoka–Yau inequality. However, the direct arithmetic analog of Miyaoka–Yau fails for the Arakelov metric, as shown in [19] (by explicit computations on curves of genus two). As discussed in [99], there seems to be no clear understanding of what the constants c_i in the inequality 1.2 should be (but, necessarily, $c_2 \geq 0$, as pointed out in [83]). A suggestion is put forth in Section 10.3.

Diophantine aspects of orbifolds (X, Δ) have also recently been explored in [29, 1]. Moreover, in the opposite situation where $X_{\mathbb{F}}$ is a Fano variety, i.e. $-K_{X_{\mathbb{F}}}$ is ample, some intriguing conjectural relations between the density of \mathbb{F} -points on $X_{\mathbb{F}}$ and bounds on the height of $-\mathcal{K}_{\mathcal{X}}$ are discussed in [17] (for appropriate models \mathcal{X} of $X_{\mathbb{F}}$ over $\mathcal{O}_{\mathbb{F}}$).

1.2. Main results. Let $(\mathcal{X}, \mathcal{D})$ be a log pair over $\mathcal{O}_{\mathbb{F}}$ and consider, for simplicity, first the case when $\mathcal{O}_{\mathbb{F}} = \mathbb{Z}$. Assume that $\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is relatively ample, i.e. either the log canonical line bundle $\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ or its dual $-\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is relatively ample. Then the complexification $\pm K_{(X, \Delta)}$ of $\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ admits a Kähler-Einstein metric if and only if $\pm K_{(X, \Delta)}$ is *K-polystable* (as recalled in Section 2.3). For example, when $K_{(X, \Delta)} > 0$ any orbifold/cusp pair (X, Δ) is K-polystable. Accordingly, when $\pm K_{(X, \Delta)}$ is K-polystable we define the *canonical height* $h_{\text{can}}(\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})})$ of $\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ as the height of $\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ wrt to any volume-normalized Kähler-Einstein metric on $\pm K_{(X, \Delta)}$.

1.2.1. General results. We show that the canonical height has a number of useful properties, as the coefficients of $\mathbf{w} \in \mathbb{R}^m$ of \mathcal{D} are varied, assuming that $\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ stays proportional to one and the same \mathbb{R} -line bundle (see Prop 4.1 and Prop 4.3 respectively):

- $\pm\hat{h}_{\text{can}}(\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})})$ is *concave* and *continuous* wrt \mathbf{w} up to the boundary of the convex domain in \mathbb{R}^m where $\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is K-semistable.
- $\pm\hat{h}_{\text{can}}(\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})})$ is *real-analytic* wrt \mathbf{w} in the region where (X, Δ) is K-stable, if (X, Δ) is log smooth.

This last statement requires some explanation. So far, we have not defined $\hat{h}_{\text{can}}(\pm\mathcal{K}_{(\mathcal{X},\mathcal{D})})$ in the “log Calabi-Yau case” when $\mathcal{K}_{(\mathcal{X},\mathcal{D})}$ is trivial. But $\hat{h}_{\text{can}}(\pm\mathcal{K}_{(\mathcal{X},\mathcal{D})})$ extends real-analytically over this region and moreover, coincides with a Faltings type height of $(\mathcal{X},\mathcal{D})$ there, defined in terms of the period of the trivializing section of $\mathcal{K}_{(\mathcal{X},\mathcal{D})}$. Likewise, we show that when $\mathcal{K}_{(\mathcal{X},\mathcal{D})}$ is relatively ample, the canonical height of $\mathcal{K}_{(\mathcal{X},\mathcal{D})}$ may be expressed as a limit of periods on large N -fold powers X^N of a canonical algebraic top form α (see Theorem 5.2):

$$(1.3) \quad \hat{h}_{\text{can}}(\mathcal{K}_{(\mathcal{X},\mathcal{D})}) = - \lim_{N \rightarrow \infty} \left(\frac{i}{2}\right)^{(Nn)^2} \log \int_{X^N} \alpha \wedge \bar{\alpha},$$

assuming that $(\mathcal{X},\mathcal{D})$ is klt (which is a generic condition). The proof leverages the probabilistic approach to Kähler-Einstein metrics introduced in [5, 6, 9]. A similar limit formula is established for $-\hat{h}_{\text{can}}(-\mathcal{K}_{(\mathcal{X},\mathcal{D})})$, conditioned on some conjectures in [6, 9], which hold when $n = 1$. Anyhow, in practice, $-\hat{h}_{\text{can}}(-\mathcal{K}_{(\mathcal{X},\mathcal{D})})$ may be obtained by analytic continuation of the formula for the canonical height of relatively ample log canonical line bundles.

The results above naturally extend to any number field \mathbb{F} , by taking all the complexifications X_σ of \mathcal{X} into account (labeled by the embeddings σ of \mathbb{F} in \mathbb{C}). The real-analyticity properties above indicate that when switching the sign of $\mathcal{K}_{(\mathcal{X},\mathcal{D})}$ the role of $h_{\text{can}}(\overline{\mathcal{K}_{(\mathcal{X},\mathcal{D})}})$ is played by the *negative* of $h_{\text{can}}(\overline{-\mathcal{K}_{(\mathcal{X},\mathcal{D})}})$. In Prop 3.3 this phenomenon is illuminated by establishing a unified variational principle for the invariant $\pm h_{\text{can}}(\pm\mathcal{K}_{(\mathcal{X},\mathcal{D})})$. It shows, in particular, that the Kähler-Einstein metrics minimize a logarithmic generalization of the arithmetic Mabuchi functional introduced by Odaka [86]. Furthermore, we introduce a notion of optimality of models: a model $(\mathcal{X}^o, \mathcal{D}^o)$ over $\mathcal{O}_{\mathbb{F}}$ for a log pair $(X_{\mathbb{F}}, \Delta_{\mathbb{F}})$ is called *optimal* if $\pm\mathcal{K}_{(\mathcal{X}^o, \mathcal{D}^o)}$ is relatively ample (for some sign) and $(\mathcal{X}^o, \mathcal{D}^o)$ minimizes $\pm h(\overline{\pm\mathcal{K}_{(\mathcal{X},\mathcal{D})}})$:

$$(1.4) \quad \pm h(\overline{\pm\mathcal{K}_{(\mathcal{X}^o, \mathcal{D}^o)}}) = \min_{(\mathcal{X}, \mathcal{D})} \pm h(\overline{\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})}}),$$

for any fixed metric on $\pm K_{(X, \Delta)(\mathbb{C})}$, where $(\mathcal{X}, \mathcal{D})$ ranges over all models over $\mathcal{O}_{\mathbb{F}}$ for $(X_{\mathbb{F}}, \Delta_{\mathbb{F}})$ such that $\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is relatively ample. For example, when K_X is a curve with $K_X > 0$, a stable model of X over $\mathcal{O}_{\mathbb{F}}$ (in the sense of Deligne-Mumford [38]) is an optimal model for $(X, 0)$. This follows from work of Odaka [85] (see Cor 3.10). Conjecturally, in general, our notion of optimality is related to Odaka’s notion of global K-semistability [85] (see Remark 3.17). It should be stressed that the optimality condition 1.4 is independent of the choice of metric (see Lemma 2.4).

1.2.2. *Explicit formulae for log pairs on $\mathbb{P}_{\mathbb{Z}}^1$ with three components.* Consider an effective divisor $\Delta_{\mathbb{Q}}$ on $\mathbb{P}_{\mathbb{Q}}^1$ such that the complexification of $(\mathbb{P}_{\mathbb{Q}}^1, \Delta_{\mathbb{Q}})$ is K-semistable. As is well-known [51, 9], this amounts to the following condition on the weights w_i of $\Delta_{\mathbb{Q}}$:

$$(1.5) \quad 0 \leq w_i \leq 1, \quad w_i \leq \frac{1}{2}V + 1, \quad V := \sum_j w_j - 2$$

(note that, since V is the volume/degree of $K_{(\mathbb{P}_{\mathbb{Q}}^1, \Delta_{\mathbb{Q}})}$, the second condition is automatic when $K_{(\mathbb{P}_{\mathbb{Q}}^1, \Delta_{\mathbb{Q}})}$ is semi-ample, i.e. when $V \geq 0$). We show (Prop 3.14) that when $\Delta_{\mathbb{Q}}$ is supported on *three* points, the optimal model of $(\mathbb{P}_{\mathbb{Q}}^1, \Delta_{\mathbb{Q}})$ over \mathbb{Z} is $(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)$, where \mathcal{D}^o is the Zariski closure of the divisor on $\mathbb{P}_{\mathbb{Q}}^1$ supported on $\{0, 1, \infty\}$ having the same coefficients as $\Delta_{\mathbb{Q}}$. Using the period formula 1.3 the canonical height of $(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)$ is expressed explicitly in terms of the Hurwitz zeta function $\zeta(s, x)$ and its derivative $\zeta'(s, x)$ wrt s . More precisely, setting

$$(1.6) \quad \gamma(a, b) := F(b) + F(1 - b) - F(a) - F(1 - a), \quad F(x) := \zeta(-1, x) + \zeta'(-1, x),$$

for $a, b \in [0, 1]$ we show:

Theorem 1.1. *Let $(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)$ be as above. When $K_{(\mathbb{P}_{\mathbb{Q}}^1, \Delta_{\mathbb{Q}})}$ is semi-ample (i.e. $V \geq 0$)*

$$(1.7) \quad \hat{h}_{can}(\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)}) = f(\mathbf{w}) := \frac{1}{2}(1 - \log(\pi \frac{V}{2})) - \frac{\gamma(0, \frac{V}{2}) - \sum_{i=1}^3 \gamma(w_i - \frac{V}{2}, w_i)}{V}$$

and when $-K_{(\mathbb{P}_{\mathbb{Q}}^1, \Delta_{\mathbb{Q}})}$ is ample (i.e. $V < 0$)

$$\hat{h}_{can}(-\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)}) = \frac{1}{2}(1 + \log \frac{\pi}{-V/2}) + \frac{\gamma(0, \frac{-V}{2}) + \sum_{i=1}^3 \gamma(w_i, w_i - \frac{V}{2})}{V}.$$

The theorem applies, in particular, when Δ is an orbifold/cusp divisor, i.e. when $w_i = 1 - 1/m_i$ for $m_i \in \mathbb{N} \cup \{\infty\}$, $i \leq 3$. Indeed, in this case $(\mathbb{P}_{\mathbb{Q}}^1, \Delta_{\mathbb{Q}})$ is always K-polystable - the corresponding Kähler-Einstein metric is the one induced by uniformization. In fact, *any* orbifold $(\mathbb{P}_{\mathbb{Q}}^1, \Delta_{\mathbb{Q}})$ such that $-K_{(\mathbb{P}_{\mathbb{Q}}^1, \Delta_{\mathbb{Q}})} > 0$ has the property that $\Delta_{\mathbb{Q}}$ is supported on three points and $(\mathbb{P}_{\mathbb{Q}}^1, \Delta_{\mathbb{Q}})$ is isomorphic to the quotient \mathbb{P}^1/G , where G is a finite subgroup of $SU(2)$ [70, Chapter 8]. Moreover, when $K_{(\mathbb{P}_{\mathbb{Q}}^1, \Delta_{\mathbb{Q}})} > 0$ and Δ is an orbifold/cusp divisor supported on three points the Kähler-Einstein metric for (\mathbb{P}^1, Δ) is the one induced by the action on the upper-half plane \mathbb{H} by a discrete subgroup Γ of $SL(2, \mathbb{R})$ such that $\overline{\mathbb{H}/\Gamma} \simeq \mathbb{P}_{\mathbb{C}}^1$ (known as a triangle group [92, 31, Prop 1]). In this case $\hat{h}_{can}(\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)})$ appears in the arithmetic Riemann-Roch formula established in [47] (using a different volume-normalization). Combining the previous theorem with [47, Thm 10.1] thus yields an explicit formula for the derivative at $s = 1$ of the corresponding Selberg zeta function $Z(s, \Gamma)$ in terms of the arithmetic degree of the line bundle ψ_W defined in [47] (generalizing the case when $\Gamma = SL(2, \mathbb{Z})$, established in [47, Thm 10.2]).

In another direction, the previous theorem yields explicit expressions for Odaka's modular invariant of any polarized log pair $(\mathbb{P}_{\mathbb{Q}}^1, \Delta_{\mathbb{Q}})$ such that $\Delta_{\mathbb{Q}}$ is supported on three points (see formula 7.1).

1.2.3. *Sharp bounds for log pairs on $\mathbb{P}_{\mathbb{Z}}^1$.* In [3] a logarithmic arithmetic analog of Fujita's sharp bound [50] on the degree of a K-semistable Fano variety was proposed, which may be formulated as the following bound

$$(1.8) \quad h_{\text{can}}(\mathcal{X}, \mathcal{D}) \leq h_{\text{can}}(\mathbb{P}_{\mathbb{Z}}^n, 0) \quad \left(= \frac{1}{2}(n+1)^{n+1} \left((n+1) \sum_{k=1}^n k^{-1} - n + \log\left(\frac{\pi^n}{n!}\right) \right) \right)$$

assuming that \mathcal{X} is a projective scheme over \mathbb{Z} such that $-\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is a relatively ample \mathbb{Q} -line bundle and its complexification is K-semistable. Moreover, for \mathcal{X} normal equality should hold only for $(\mathbb{P}_{\mathbb{Z}}^n, 0)$. The conjecture was, in particular, settled for $n = 1$ when $\mathcal{D} \otimes \mathbb{Q}$ is supported at three points. It should, however, be stressed that the conjectured inequality 1.8 does *not* hold when the canonical height is replaced by its normalization \hat{h}_{can} . In fact, for all we know it could actually be that $\hat{h}_{\text{can}}(\mathcal{X}, \mathcal{D})$ is *minimal* for $(\mathbb{P}_{\mathbb{Z}}^n, 0)$ among all arithmetic log Fano varieties $(\mathcal{X}, \mathcal{D})$ over \mathbb{Z} , if $(\mathcal{X}, \mathcal{D})$ is taken to be an optimal model over \mathbb{Z} . Here we show that this is, indeed, the case for any arithmetic log Fano surface $(\mathcal{X}, \mathcal{D})$ such that $\mathcal{D} \otimes \mathbb{Q}$ is supported at three points. In this case it was shown in [3, Section 6] that the optimal model of $(\mathcal{X}, \mathcal{D})$ is of the form $(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)$, as appearing in the previous section. The minimality in question thus follows from the first inequality in the following theorem:

Theorem 1.2. *Let $(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)$ be as in the previous section. Then*

$$\pm \hat{h}_{\text{can}}(\pm \mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)}) \leq -\hat{h}_{\text{can}}(-\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, 0)}) \quad \left(= -\frac{1}{2}(1 + \log \pi) \right)$$

with equality iff $\mathcal{D}^o = 0$. Furthermore, if $K_{(\mathbb{P}^1, \Delta)}$ is semi-ample, then the following more precise inequality holds:

$$\hat{h}_{\text{can}}(\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)}) \leq -\frac{1}{2} \log(\pi) + \frac{3}{2} \log \frac{\Gamma(\frac{2}{3})}{\Gamma(\frac{1}{3})} (< 0)$$

where equality holds iff $K_{(\mathbb{P}^1, \Delta)}$ is trivial and all the coefficients of \mathcal{D}^o equal $2/3$. More generally, the latter inequality holds when the Kähler-Einstein metric is replaced by any volume-normalized continuous metric on $K_{(\mathbb{P}^1, \Delta)}$.

The proof combines the explicit formula in Theorem 1.1 with the concavity of $\pm \hat{h}_{\text{can}}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})})$. Interestingly, the second inequality in the previous theorem is reminiscent of the result in [37, page 29] saying that the maximal value of Faltings' stable height of elliptic curves is attained at the semistable reduction of the Néron model \mathcal{X}_0 of any elliptic curve X with vanishing j -invariant. Incidentally, after a base change, such an elliptic curve is a Galois cover of the log pair (\mathbb{P}^1, Δ) saturating the second inequality in the previous theorem.

1.2.4. Specific values of canonical heights. In some orbifold cases the terms involving the Hurwitz zeta function in Theorem 1.1 can be eliminated in favor of a single logarithmic derivative of a Dedekind zeta function $\zeta_{\mathbb{F}}(s)$ of an appropriate totally real number field \mathbb{F} . More precisely, when the degree V of $K_{(\mathbb{P}_{\mathbb{Q}}^1, \Delta_{\mathbb{Q}})}$ is positive, using the classical multiplication theorem for the Hurwitz zeta function leads - after some computations - to the results in Table 1, where (m_1, m_2, m_3) denotes the ramification indices of $\mathcal{D}^o \otimes \mathbb{Q}$ and $\hat{h}_{\text{Pet}} := \hat{h}_{\text{can}}(\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)}) - \log \frac{\pi V}{2}$ (which is the height wrt the Peterson metric). In

some of these cases the coefficients in front of the $\log p$ - terms carry information about the canonical integral model of the Shimura curve attached to \mathbb{F} , as explained in the following section.

(m_1, m_2, m_3)	$\hat{h}_{\text{Pet}} + \frac{1}{2} + \frac{1}{[\mathbb{F}:\mathbb{Q}]} \frac{\zeta_{\mathbb{F}}'(-1)}{\zeta_{\mathbb{F}}(-1)}$
$(2, 3, \infty)$	$-\frac{1}{2} \log 2 - \frac{1}{4} \log 3, \mathbb{F} = \mathbb{Q}$
$(6, 2, 6)$	$-\frac{1}{6} \log 2 + \frac{1}{8} \log 3, \mathbb{F} = \mathbb{Q}$
$(4, 4, 4)$	$-\frac{19}{12} \log 2, \mathbb{F} = \mathbb{Q}(\sqrt{2})$
$(3, 3, 6)$	$-\frac{5}{6} \log 2 - \frac{13}{16} \log 3, \mathbb{F} = \mathbb{Q}(\sqrt{3})$
$(2, 4, 12)$	$-\frac{5}{3} \log 2 - \frac{7}{16} \log 3, \mathbb{F} = \mathbb{Q}(\sqrt{3})$
$(6, 6, 6)$	$-\frac{5}{6} \log 2 - \frac{7}{16} \log 3, \mathbb{F} = \mathbb{Q}(\sqrt{3})$
$(5, 5, 5)$	$\frac{25}{48} \log 5, \mathbb{F} = \mathbb{Q}(\sqrt{5})$
$(3, 4, 6)$	$-\frac{9}{16} \log 3 - \frac{11}{12} \log 2, \mathbb{F} = \mathbb{Q}(\sqrt{6})$
$(7, 7, 7)$	$-\frac{95}{12^2} \log 7, \mathbb{F} = \mathbb{Q}(\cos(\pi/7))$
$(9, 9, 9)$	$-\frac{31}{24} \log 3, \mathbb{F} = \mathbb{Q}(\cos(\pi/9))$

TABLE 1. Ramification indices and the corresponding normalized height for some log canonically polarized orbifolds.

In the opposite case of Fano orbifold curves (which are automatically K-polystable) further cancellations take place (see Section 8.2).

1.2.5. *Application to the canonical integral models of some quaternionic Shimura curves.* Given a quaternionic Shimura curve X over a totally real field \mathbb{F} , consider the corresponding canonical model $(\mathcal{X}, \mathcal{D})$ of (X, Δ) over $\mathcal{O}_{\mathbb{F}}$, appearing in formula 1.1. The log pair (X, Δ) over \mathbb{F} is *stable*, in the sense of the Minimal Model Program (MMP) in birational geometry (generalizing Deligne-Mumford's notion of stability to log pairs; see Section 2.4.2). However, in general, there exist prime ideals \mathfrak{p} such that the corresponding log pair $(\mathcal{X}, \mathcal{D})$ over the field $(\mathcal{O}_{\mathbb{F}}/\mathfrak{p})$ is *not* stable (even when \mathcal{D} is horizontal). That is to say that the fibers of the log pair $(\mathcal{X}, \mathcal{D})$ over the closed fibers over $\text{Spec } \mathcal{O}_{\mathbb{F}}$ are not stable, in general. However, there exists, after perhaps performing a finite base change $\mathbb{F} \subset \mathbb{F}'$, a unique model $(\mathcal{X}^{\circ}, \mathcal{D}^{\circ})$ for (X, Δ) over $\mathcal{O}_{\mathbb{F}'}$ all of whose fibers are stable (see Section 3.3). Such a model is, in fact, an optimal model in the sense of formula 1.4. In order to get a measure of how much $(\mathcal{X}, \mathcal{D})$ differs from $(\mathcal{X}^{\circ}, \mathcal{D}^{\circ})$ one can fix a metric on $K_{(X, \Delta)(\mathbb{C})}$ and consider the corresponding height difference

$$(1.9) \quad h(\overline{\mathcal{K}_{(\mathcal{X}, \mathcal{D})} \otimes_{\mathcal{O}_{\mathbb{F}}} \mathcal{O}_{\mathbb{F}'}}}) - h(\overline{\mathcal{K}_{(\mathcal{X}^{\circ}, \mathcal{D}^{\circ})}}}) = \sum_{\mathfrak{p}} h(\mathfrak{p}) \log N(\mathfrak{p})$$

for a finite set \mathfrak{p} of closed points in $\text{Spec } \mathcal{O}_{\mathbb{F}'}$. The numbers $h(\mathfrak{p})$ are independent of the choice of metric. Indeed, they may be expressed as algebro-geometric intersection numbers on the fiber $\mathcal{Y}_{\mathfrak{p}}$ over \mathfrak{p} of any given normal model \mathcal{Y} of X over $\mathcal{O}_{\mathbb{F}'}$, dominating $\mathcal{X} \otimes_{\mathcal{O}_{\mathbb{F}}} \mathcal{O}_{\mathbb{F}'}$ and \mathcal{X}^o (see Lemma 2.4). Moreover, $h(\mathfrak{p})$ is non-negative and vanishes iff $\mathcal{X} \otimes_{\mathcal{O}_{\mathbb{F}}} \mathcal{O}_{\mathbb{F}'}$ is isomorphic to \mathcal{X}^o locally around \mathfrak{p} (see Remark 3.15).

We will consider some cases where $(\mathcal{X}^o, \mathcal{D}^o)$ is of the form $(\mathbb{P}_{\mathcal{O}_{\mathbb{F}'}}^1, \mathcal{D}^o)$ for a horizontal divisor \mathcal{D}^o on $\mathbb{P}_{\mathcal{O}_{\mathbb{F}'}}^1$. Given a prime number p , we will compute the rational numbers

$$h(p) := \frac{1}{[\mathbb{F}' : \mathbb{Q}]} \sum_{\mathfrak{p} \in \mathfrak{p}_i} f_i h(\mathfrak{p}_i), \quad N(\mathfrak{p}_i) := p^{f_i},$$

that are invariant under base change. To the best of our knowledge these numbers have not been computed before, except for the indefinite quaternion algebra over \mathbb{Q} with the smallest discriminant ($= 1$). In this case, $\mathcal{X} = \mathbb{P}_{\mathbb{Z}}^1$ and \mathcal{D} is an explicit horizontal divisor supported at three \mathbb{Z} -points, as follows from classical results for the j -invariant of elliptic curves. Accordingly, the numbers $h(\mathfrak{p})$ may be computed directly from the intersection-theoretic formula in Lemma 2.4. The result is that $h(p)$ vanishes unless $p = 2$ or $p = 3$ and

$$(1.10) \quad h(2) = 1/2, \quad h(3) = 1/4.$$

In particular, the log pairs $(\mathcal{X}, \mathcal{D}) \otimes \mathbb{Z}/(p)$ are not stable when $p = 2$ or $p = 3$, although \mathcal{X} is smooth over \mathbb{Z} (see Remark 9.4). It may also be worth pointing out that in this case Theorem 1.1 yields a new proof of the height formula 1.1, that does not use automorphic (or modular) forms, nor uniformization (see Section 9.2.2).

For the indefinite quaternion algebra over \mathbb{Q} with the next to smallest discriminant ($= 6$), we compute the corresponding rational numbers $h(p)$ indirectly, by combining Theorem 1.1 with formula 1.1. In this case, by a result of Ihara, the canonical model of the corresponding quaternionic Shimura curve $X_{\mathbb{Q}}$ is isomorphic to \mathbb{P}^1 over $\mathbb{Q}(\sqrt{3}, i)$. Moreover, Δ is supported on four $\mathbb{Q}(\sqrt{3}, i)$ -points with ramification indices $(3, 3, 2, 2)$ and cross ratio -1 [42]. We show that the corresponding unique optimal model of $(X_{\mathbb{Q}}, \Delta_{\mathbb{Q}}) \otimes \mathbb{Q}(\sqrt{3}, i)$ over $\mathcal{O}_{\mathbb{Q}(\sqrt{3}, i)}$ is given by $(\mathbb{P}_{\mathcal{O}_{\mathbb{Q}(\sqrt{3}, i)}}^1, \mathcal{D}^o)$, where \mathcal{D}^o denotes the Zariski closure of the divisor on $\mathbb{P}_{\mathbb{Q}(\sqrt{3}, i)}^1$ supported on the pair of two points $\{\infty, 0; 1, -1\}$, having the same ramification indices $(3, 3; 2, 2)$ as the divisor Δ under the action of an automorphism of \mathbb{P}^1 .

Theorem 1.3. *Consider the quaternion algebra over \mathbb{Q} with discriminant 6 and denote by $(\mathcal{X}, \mathcal{D})$ the canonical model over \mathbb{Z} of the corresponding Shimura curve $(X_{\mathbb{Q}}, \Delta_{\mathbb{Q}})$. Then $h(p) = 0$ unless $p = 2$ or $p = 3$ and*

$$h(2) = 11/18, \quad h(3) = 7/12.$$

It follows that $(\mathcal{X} \otimes_{\mathbb{Z}} \mathcal{O}_{\mathbb{Q}(\sqrt{3}, i)}, \mathcal{D})$ is isomorphic to $(\mathbb{P}_{\mathcal{O}_{\mathbb{Q}(\sqrt{3}, i)}}^1, \mathcal{D}^o)$ away from the fibers over $p = 2$ and $p = 3$ (see Remark 3.15). This isomorphism also follows from the explicit model for $(\mathcal{X}, \mathcal{D})$ over $\mathbb{Z}[1/6]$ established in [60]. Since there is a single prime ideal \mathfrak{p}_2 in $\mathcal{O}_{\mathbb{Q}(\sqrt{3}, i)}$ over 2 and a single one \mathfrak{p}_3 over 3 the previous theorem also allows one

to compute $h(\mathfrak{p}_2)$ and $h(\mathfrak{p}_3)$. It also follows from the previous theorem (together with Prop 3.14) that the fibers of the log arithmetic surface $(\mathcal{X}, \mathcal{D}) \otimes_{\mathbb{Z}} \mathcal{O}_{\mathbb{Q}(\sqrt{3}, i)}$ are not stable over \mathfrak{p}_2 and \mathfrak{p}_3 . In the proof of the theorem a covering argument is used to replace the divisor \mathcal{D}^o with the Zariski closure of the divisor supported on the points $\{0, 1, \infty\}$ with ramification indices $(6, 2, 6)$. For the later divisor the corresponding the explicit formula in Table 1 can be employed.

More generally, by the classification results in [93, Table 3], there are 19 Shimura curves $X_{\mathbb{F}}$ that are isomorphic to $\mathbb{P}_{\bar{\mathbb{Q}}}^1$ over $\bar{\mathbb{Q}}$ and such that the corresponding divisor $\Delta_{\bar{\mathbb{Q}}}$ is supported on at most three $\bar{\mathbb{Q}}$ -points, up to taking finite covers. The cases when $\mathbb{F} = \mathbb{Q}$ are precisely the two cases discussed above. Another case is considered in the following result (see also Theorem 9.6 for one more case).

Theorem 1.4. *Consider the quaternion algebra over $\mathbb{Q}(\sqrt{3})$ that is ramified precisely over the unique prime ideal in $\mathcal{O}_{\mathbb{Q}(\sqrt{3})}$ containing 3 and denote by $(\mathcal{X}, \mathcal{D})$ the canonical model over $\mathcal{O}_{\mathbb{Q}(\sqrt{3})}$ of the corresponding Shimura curve $(X_{\mathbb{Q}(\sqrt{3})}, \Delta_{\mathbb{Q}(\sqrt{3})})$. Fix a finite field extension \mathbb{F} of $\mathbb{Q}(\sqrt{3})$ such that $X_{\mathbb{Q}(\sqrt{3})} \otimes \mathbb{F}$ is isomorphic to $\mathbb{P}_{\mathbb{F}}^1$ and $\Delta_{\mathbb{F}}$ is supported on three \mathbb{F} -points. Then the optimal model of $(X_{\mathbb{Q}(\sqrt{3})}, \Delta_{\mathbb{Q}(\sqrt{3})}) \otimes \mathbb{F}$ over $\mathcal{O}_{\mathbb{Q}(\sqrt{3})}$ is given by $(\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1, \mathcal{D}^o)$, where \mathcal{D}^o denotes the Zariski closure of the divisor on $\mathbb{P}_{\mathbb{F}}^1$ supported on $\{0, 1, \infty\}$ having the same ramification indices $(2, 4, 12)$ as the divisor $\Delta_{\mathbb{F}}$. Moreover $h(p) = 0$ unless $p = 2$ or $p = 3$ and*

$$h(2) = \frac{5}{9}, \quad h(3) = \frac{15}{48}.$$

It follows from the vanishing of $h(p)$ for $p \neq 2, 3$ in the previous theorem that the base change of $(\mathcal{X}, \mathcal{D})$ to $\mathcal{O}_{\mathbb{F}}$ is isomorphic to $(\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1, \mathcal{D}^o)$ away from the fibers of \mathcal{X} over $\{(2), (3)\} \in \text{Spec} \mathbb{Z}$ (see Remark 3.15). As a consequence, over this Zariski open subset the following three properties hold: (i) \mathcal{X} is (geometrically) smooth, (ii) \mathcal{D} is horizontal and (iii) the irreducible components of \mathcal{D} are mutually non-intersecting. In fact, (i) and (ii) hold over any prime ideal \mathfrak{p} of $\mathcal{O}_{\mathbb{Q}(\sqrt{3})}$ which is not in the ramification locus of the corresponding quaternion algebra, i.e. when $\mathfrak{p} \neq \mathfrak{p}_3$. This follows from general results in [103] (see the beginning of Section 9.3). In particular, they also hold for $\mathfrak{p} = \mathfrak{p}_2$. However, the non-vanishing of $h(2)$, in the previous theorem, implies that some of the components of \mathcal{D} must intersect over \mathfrak{p}_2 . Furthermore, in Section 9.3, we show, building on [103], that there is wild ramification over $\mathcal{X}_{\mathfrak{p}_2}$ (in the sense of stacks).

Remark 1.5. In order to extend Theorem 1.3 to further Shimura curves in [93, Table 3], one needs to extend Table 1 to the other ramification indices in column 5 in [93, Table 3]. See the discussion in Section 8.3.

1.2.6. *Application to twisted Fermat curves.* Given integers a_0, a_1 and a_2 , consider the corresponding twisted Fermat curve $X_a^{(m)}$ of degree m in $\mathbb{P}_{\mathbb{Q}}^2$, cut out by the polynomial $a_0 x_0^m + a_1 x_1^m + a_2 x_2^m$, for $m \geq 3$. When $a_1 = a_2 = 1$ and $a_0 = -1$ this is the classical Fermat curve that we shall denote by $X^{(m)}$. Using that $X^{(m)}$ is a Galois cover of \mathbb{P}^1 of degree m with branching divisor Δ supported at $\{0, 1, \infty\}$ with ramification indices (m, m, m) , we deduce the following result from Theorems 1.1, 1.2:

Theorem 1.6. Denoting by $\mathcal{X}_a^{(m)}$ the Zariski closure of $X_a^{(m)}$ in $\mathbb{P}_{\mathbb{Z}}^2$,

$$\hat{h}_{\text{can}}(\mathcal{K}_{\mathcal{X}_a^{(m)}}) = f(1 - 1/m, 1 - 1/m, 1 - 1/m) + \log m + \left(\frac{(m-3)}{2} + 1 \right) \frac{1}{m} \sum_i \log |a_i|,$$

where f is defined in 1.7. As a consequence, for any stable model $\mathcal{Y}_a^{(m)}$ of $X_a^{(m)}$ over $\mathcal{O}_{\mathbb{F}}$ and any volume-normalized metric on $K_{\mathcal{X}_a^{(m)}(\mathbb{C})}$

$$(1.11) \quad \hat{h}(\overline{\mathcal{K}_{\mathcal{Y}_a^{(m)}}}) \leq f(1 - 1/m, 1 - 1/m, 1 - 1/m) + \log m (\leq \log m).$$

Consider in particular the Zariski closure in $\mathcal{X}^{(m)} \subset \mathbb{P}_{\mathbb{Z}}^2$ of the Fermat curve of degree $m (\geq 4)$. For $m \in \{4, 5, 6, 7, 9\}$ Table 1 yields an explicit expression for $\hat{h}_{\text{can}}(\mathcal{K}_{\mathcal{X}^{(m)}})$ in terms of the logarithmic derivatives of Dedekind zeta functions of the following form:

$$(1.12) \quad \hat{h}_{\text{can}}(\mathcal{K}_{\mathcal{X}^{(m)}}) = \log \frac{\pi(1 - \frac{3}{m})}{2} - \frac{1}{2} - \frac{1}{[\mathbb{F}_m : \mathbb{Q}]} \frac{\zeta'_{\mathbb{F}_m}(-1)}{\zeta_{\mathbb{F}_m}(-1)} - c_m \log m,$$

where $\mathbb{F}_m := \mathbb{Q}(\cos(\pi/m))$, $c_m \in \mathbb{Q}$ and the sum ranges over all primes p dividing m . However, there are reasons to doubt that this formula holds for *all* $m \geq 4$ - even for $m = 8$ (see the discussion in Section 8.3).

In another direction, using the relations between the Kähler-Einstein metric and the Arakelov metric in [62, 63] we deduce the following general inequalities from the previous theorem:

Corollary 1.7. Denoting by $\mathcal{X}^{(m)}$ the Zariski closure in $\mathbb{P}_{\mathbb{Z}}^2$ of the Fermat curve of degree $m (\geq 4)$,

$$(1.13) \quad \begin{aligned} \hat{h}_{\text{Ar}}(\overline{\mathcal{K}_{\mathcal{X}^{(m)}}}) &\leq \frac{1}{2} (1 - \log(\frac{1-3/m}{2})) - \frac{\gamma(0, \frac{1-3/m}{2}) - 3\gamma(\frac{1}{2} + \frac{1}{2m}, 1 - \frac{1}{m})}{1 - 3/m} + 2 \log m + \epsilon_m \leq \\ &\quad - \frac{1}{2} - \frac{13}{12} \log 2 - \frac{1}{2} \frac{\zeta'_{\mathbb{Q}(\sqrt{2})}(-1)}{\zeta_{\mathbb{Q}(\sqrt{2})}(-1)} + \epsilon_m + 2 \log m, \end{aligned}$$

where ϵ_m is the following number, decreasing to 0, as $m \rightarrow \infty$,

$$\epsilon_m := \frac{1}{2} \frac{4 \log((m-1)(m-2) - 2) + 1}{(m-1)(m-2)/2 - 1} + \frac{1}{2} \log\left(\frac{(m-1)(m-2) - 2}{m^2}\right).$$

In general, $\hat{h}_{\text{Ar}}(\overline{\mathcal{K}_{\mathcal{X}}}) \geq 0$, when $\mathcal{K}_{\mathcal{X}}$ is relatively ample, [44]. Accordingly, the previous corollary gives (using that $\epsilon_m \leq \epsilon_4$ and by evaluating the constants in question) that

$$(1.14) \quad 0 \leq \hat{h}_{\text{Ar}}(\overline{\mathcal{K}_{\mathcal{X}^{(m)}}}) \leq -0.88\dots + 2 \log(m).$$

Since $\hat{h}(\overline{\mathcal{K}_{\mathcal{Y}^{(m)}}}) \leq \hat{h}(\overline{\mathcal{K}_{\mathcal{X}^{(m)}}})$ for any stable model $\mathcal{Y}^{(m)}$ of $X^{(m)}$ we thus deduce an explicit Parshin inequality for $\mathcal{Y}^{(m)}$ (as in inequality 1.2). The inequality 1.14 also holds when $\mathcal{X}^{(m)}$ is replaced by the minimal regular model $\mathcal{X}_{\text{min}}^{(m)}$ attached to any given finite field extension \mathbb{F} of \mathbb{Q} (by Prop 3.18). Upper bounds on $\hat{h}_{\text{Ar}}(\mathcal{K}_{\mathcal{X}_{\text{min}}^{(m)}})$ have previously been obtained in [73, 34, 35], when $\mathbb{F} = \mathbb{Q}(\zeta_m)$ where ζ_m denotes an m -th root of

unity (assuming that m is a prime number or square-free). However, the bounds in [73, 34, 35] involve two non-explicit constants κ_1 and κ_2 , appearing in the analytic contribution $\kappa_1 \log m + \kappa_2$ to the bounds in [73, 34, 35] (originating in [73, Thm 2.10]). Explicit bounds on $\hat{h}_{\psi_{\Delta r}}(\overline{\mathcal{K}_{\mathcal{Y}(m)}})$ that are polynomial in m are contained in [61, Cor 1.5.1].

1.3. Acknowledgments. We are deeply grateful to Noam Elkies, Dennis Eriksson, Gerard Freixas i Montplet, Christian Johansson, John Voight and Xinyi Yuan for very helpful discussions and feedback. This work was supported by a Wallenberg Scholar grant from the Knut and Alice Wallenberg foundation.

2. SETUP

Henceforth, \mathcal{X} will denote an *arithmetic variety* (over $\mathcal{O}_{\mathbb{F}}$), i.e. a projective flat scheme \mathcal{X} over the ring of integers $\mathcal{O}_{\mathbb{F}}$ of a number field \mathbb{F} of relative dimension n such that \mathcal{X} is reduced, pure dimensional, satisfies Serre's conditions S_2 and has a relative canonical sheaf $\omega_{\mathcal{X}/B}$ [71, Condition 1.6.1] - the \mathbb{Q} -divisor corresponding to $\omega_{\mathcal{X}/B}$ will be denoted by $\mathcal{K}_{\mathcal{X}}$. For example, these conditions are satisfied if \mathcal{X} is normal. We will denote by π the corresponding structure morphism from \mathcal{X} to $\text{Spec } \mathcal{O}_{\mathbb{F}}$. The corresponding scheme over \mathbb{F} , $\mathcal{X} \otimes_{\mathcal{O}_{\mathbb{F}}} \mathbb{F}$, will be denoted by $X_{\mathbb{F}}$. Furthermore, we will denote by X_{σ} the complex varieties corresponding to \mathcal{X} , labeled by the embeddings $\sigma : \mathbb{F} \hookrightarrow \mathbb{C}$. The \mathbb{C} -points of \mathcal{X} will be denoted by $X(\mathbb{C})$:

$$X(\mathbb{C}) = \bigsqcup_{\sigma} X_{\sigma}, \quad X_{\sigma} := \mathcal{X} \otimes_{\sigma} \mathbb{C}.$$

Throughout the paper we will assume that X_{σ} is normal. Given a line bundle $\mathcal{L} \rightarrow \mathcal{X}$ we will denote by L^n the corresponding algebraic top intersection over the generic fiber of \mathcal{X} (or, equivalently, over the complexifications X_{σ} for any σ). We will use additive notation for tensor products of line bundles and say that $\pm \mathcal{L}$ is relatively ample if either \mathcal{L} or its dual $-\mathcal{L}$ is relatively ample.

2.1. Log pairs and models. A *log pair* $(\mathcal{X}, \mathcal{D})$ over $\mathcal{O}_{\mathbb{F}}$ (also called an *arithmetic log variety*) of relative dimension n is an arithmetic variety \mathcal{X} endowed with an effective \mathbb{R} -divisor \mathcal{D} on \mathcal{X} , not contained in the singular locus of \mathcal{X} , such that $\mathcal{K}_{\mathcal{X}} + \mathcal{D}$ is \mathbb{R} -Cartier (i.e. a real multiple of $\mathcal{K}_{\mathcal{X}} + \mathcal{D}$ defines a line bundle). See [71, Section 1.1] where log pairs are defined over any excellent ring for \mathbb{Q} -divisors and the same setup applies to \mathbb{R} -divisors [71, Remark 2.20]. The complexifications of $(\mathcal{X}, \mathcal{D})$ will be denoted by $(X_{\sigma}, \Delta_{\sigma})$. A triple $(\mathcal{X}, \mathcal{D}; \mathcal{L})$ consisting of a log pair $(\mathcal{X}, \mathcal{D})$ over $\mathcal{O}_{\mathbb{F}}$ and a relatively ample \mathbb{Q} -line bundle \mathcal{L} on \mathcal{X} will be called a *polarized log pair over $\mathcal{O}_{\mathbb{F}}$* .

Given a polarized log pair $(X, \Delta; L)$ over \mathbb{F} , a *model for $(X, \Delta; L)$ over $\mathcal{O}_{\mathbb{F}}$* consists, by definition, of a polarized log pair $(\mathcal{X}, \mathcal{D}; \mathcal{L})$ over $\mathcal{O}_{\mathbb{F}}$ and an isomorphism between $(\mathcal{X}, \mathcal{D}; \mathcal{L}) \otimes_{\mathcal{O}_{\mathbb{F}}} \mathbb{F}$ and $(X, \Delta; L)$.

2.1.1. *Singularities of log pairs.* Given a log pair $(\mathcal{X}, \mathcal{D})$ over an excellent ring, with \mathcal{X} normal, consider a blow-up morphism $p : \mathcal{Y} \rightarrow \mathcal{X}$ from a normal scheme \mathcal{Y} to \mathcal{X} and decompose

$$\mathcal{K}_{\mathcal{Y}/\mathcal{X}} - p^*\mathcal{D} = \sum_i a_i E_i, \quad a_i \geq -1, \quad \mathcal{K}_{\mathcal{Y}/\mathcal{X}} := \mathcal{K}_{\mathcal{Y}} - p^*\mathcal{K}_{\mathcal{X}},$$

where the prime divisor E_i is either an exceptional divisor of p or the proper transform of a component of \mathcal{D} . Following [71, Section 2] $(\mathcal{X}, \mathcal{D})$ is said to be *log canonical (lc)* if $a_i \geq -1$ for any such $p : \mathcal{Y} \rightarrow \mathcal{X}$ and *Kawamata Log Terminal (klt)* if $a_i > 1$. Without assuming that \mathcal{X} is normal there is also a notion of *semi-log canonical* pairs $(\mathcal{X}, \mathcal{D})$ (coinciding with lc pairs when \mathcal{X} is normal)[71]. When $(\mathcal{X}, 0)$ is lc (klt) \mathcal{X} is said to have lc (klt) singularities.

For example, when $(\mathcal{X}, \mathcal{D})$ is *log smooth*, i.e. \mathcal{X} is regular and \mathcal{D} has simple normal crossings, $(\mathcal{X}, \mathcal{D})$ is lc if $w_i \leq 1$ for all coefficients w_i of \mathcal{D} and klt if $w_i < 1$ (by [71, Cor 2.13]). Moreover, in general, if \mathcal{X} is a normal scheme of dimension one over a perfect field, then $(\mathcal{X}, \mathcal{D})$ is lc (klt) iff $w_i \leq 1$ ($w < 1$) for all coefficients w_i of \mathcal{D} [71, page 43].

2.2. **Metrics and measures.** In this section X will denote a compact complex manifold.

2.2.1. *Local representations of metrics and measures.* As in [2, 3] we will use additive notation for metrics on holomorphic line bundles $L \rightarrow X$. This means that we identify a continuous Hermitian metric $\|\cdot\|$ on L with a collection of continuous local functions ϕ_U associated to a given covering of X by open subsets U and trivializing holomorphic sections e_U of $L \rightarrow U$, $\phi_U := -\log(\|e_U\|^2)$. The curvature current of the metric may then, locally, be expressed as

$$dd^c\phi_U := \frac{i}{2\pi}\partial\bar{\partial}\phi_U.$$

Accordingly, as is customary, we will symbolically denote by ϕ a given continuous Hermitian metric on L and by $dd^c\phi$ its curvature current. More generally, a *singular metric* ϕ on L is defined by the condition that $\phi_U \in L^1_{\text{loc}}$. When $dd^c\phi_U \geq 0$ such a metric is called a *psh metric* (when ϕ_U is taken to be strongly upper semi-continuous).

To a log pair (X, Δ) together with a (multi-valued) section s_Δ cutting out Δ and a continuous metric ϕ on $\pm K_{(X, \Delta)}$ we attach a measure μ_ϕ on X , in the following standard way. First, by definition this measure puts no mass on $X - X_{\text{reg}}$, where X_{reg} denotes the regular locus of X . Next, locally on X_{reg} the measure μ_ϕ is defined by

$$\mu_\phi = e^{\pm\phi_U} |s_U|^{-2} \left(\frac{i}{2}\right)^{n^2} dz \wedge d\bar{z}, \quad dz := dz_1 \wedge \cdots \wedge dz_n$$

by taking $e_U = \partial/\partial z_1 \wedge \cdots \wedge \partial/\partial z_n \otimes e_\Delta$ where e_Δ is a local trivialization of the \mathbb{Q} -line bundle over X_{reg} corresponding to the divisor Δ and $s_U e_\Delta$ is the (multi-valued) holomorphic section cutting out Δ . This measure is globally well-defined and gives finite mass to X iff (X, Δ) is klt [16, Section 3.1]. Accordingly, a metric ϕ on $\pm K_{(X, \Delta)}$ is called *volume-normalized* if $\int_X \mu_\phi = 1$. More generally, if X has several components X_σ , then ϕ is called volume-normalized if $\int_{X_\sigma} \mu_\phi = 1$ for all components.

2.2.2. *The complex Monge-Ampère measure and finite energy metrics.* Let L be a semi-ample line bundle over X and fix a continuous metric ϕ_0 on L with positive curvature current. We define the complex *Monge-Ampère measure* of a singular metric ϕ on L as the n -fold product

$$\text{MA}(\phi) := (dd^c \phi)^n / L^n,$$

using the notion of non-pluripolar products of positive currents, introduced in [23]. A psh metric ϕ on L is said to have *finite energy*, if $\text{MA}(\phi)$ is a probability measure and $\int_X (\phi - \phi_0) \text{MA}(\phi) < \infty$ (see [23, 16]). For any such metric

$$(2.1) \quad \mathcal{E}(\phi) := \mathcal{E}_{\phi_0}(\phi) := \int_X \sum_{j=0}^n (\phi - \phi_0) (dd^c \phi)^j \wedge (dd^c \phi_0)^{n-j} < \infty.$$

Remark 2.1. When $n = 1$ a psh metric ϕ has finite energy iff, locally, the gradient $\nabla \phi \in L^2_{\text{loc}}$.

Given a log pair (X, Δ) and a psh metric ϕ on $\pm K_{(X, \Delta)}$ of finite energy we obtain, just as when ϕ is continuous, a measure μ_ϕ on X . If (X, Δ) is klt then the measure μ_ϕ still gives finite total mass to X [16, 14].

Example 2.2. Let (X, Δ) be a log smooth lc pair such that $K_{(X, \Delta)}$ is ample and denote by s_i the holomorphic sections cutting out the components Δ_i of Δ with coefficient $w_i = 1$. A psh metric ϕ on $K_{(X, \Delta)}$ is said to have log-log singularities if ϕ is locally of the form $-\sum_i \log(\log |s_i|^{-2}) + O(1)$. Such a psh metric ϕ has finite energy [54, Prop 2.3]. The corresponding measure μ_ϕ gives finite total mass to X . In contrast, if ϕ is locally bounded then $\int_X \mu_\phi < \infty \iff \Delta$ is klt.

2.2.3. *Kähler-Einstein metrics.* Given a projective log pair (X, Δ) over \mathbb{C} a metric ϕ on $\pm K_{(X, \Delta)}$ is said to be a *Kähler-Einstein metric*, if ϕ has finite energy and its curvature current $dd^c \phi$ induces a Kähler metric with constant positive Ricci curvature on the complement of Δ in X_{reg} [16, 14]. In particular, by [16, 14], a Kähler-Einstein metric ϕ on $\pm K_{(X, \Delta)}$ is volume-normalized iff

$$(2.2) \quad \text{MA}(\phi) = \mu_\phi$$

By the resolution of the Yau-Tian-Donaldson conjecture (X, Δ) admits a Kähler-Einstein metric iff (X, Δ) is *K-polystable* (as defined in the following section). When $K_{(X, \Delta)} > 0$ this follows from combining the characterization of K-stability in [85, 24] with [14] and when $-K_{(X, \Delta)} > 0$ it follows - in the general singular setup - from the combination of [75] and [77].

Example 2.3. Let (X, Δ) be a log smooth lc pair such that $K_{(X, \Delta)} > 0$. Then $K_{(X, \Delta)}$ admits a Kähler-Einstein metric ϕ (unique up to scalings) and ϕ has log-log singularities (see Example 2.2).

Given a variety $X_{\mathbb{F}}$ defined over \mathbb{F} we will say that a metric ϕ on $\pm K_{(X, \Delta)(\mathbb{C})}$ is Kähler-Einstein if the restriction of ϕ to each component X_σ is a Kähler-Einstein metric on $\pm K_{(X_\sigma, \Delta_\sigma)}$.

2.3. K-stability. We next recall the definition of K-stability in terms of intersection numbers (see the survey [100] for more background). Let (X, Δ) be a log pair over \mathbb{C} and L an ample line bundle over X . A *test configuration* for a polarized log pair (X, L) is a \mathbb{C}^* -equivariant normal model $(\mathcal{X}, \mathcal{L})$ for (X, L) over the complex affine line $\mathbb{A}_{\mathbb{C}}^1$. More precisely, \mathcal{X} is a normal complex variety endowed with a \mathbb{C}^* -action ρ , a \mathbb{C}^* -equivariant holomorphic surjection π to $\mathbb{A}_{\mathbb{C}}^1$ and a relatively ample \mathbb{C}^* -equivariant \mathbb{Q} -line bundle \mathcal{L} (endowed with a lift of ρ):

$$(2.3) \quad \pi : \mathcal{X} \rightarrow \mathbb{A}_{\mathbb{C}}^1, \quad \mathcal{L} \rightarrow \mathcal{X}, \quad \rho : \mathcal{X} \times \mathbb{C}^* \rightarrow \mathcal{X}$$

such that the fiber of \mathcal{X} over $1 \in \mathbb{A}_{\mathbb{C}}^1$ is equal to (X, L) . A log pair (X, Δ) is said to be *K-semistable* if $\text{DF}_{\Delta}(\mathcal{X}, \mathcal{L}) \geq 0$ for any test configuration $(\mathcal{X}, \mathcal{L})$, where $\text{DF}_{\Delta}(\mathcal{X}, \mathcal{L})$ is the *Donaldson-Futaki invariant*:

$$(2.4) \quad n! \text{DF}_{\Delta}(\mathcal{X}, \mathcal{L}) := \frac{a}{(n+1)!} \overline{\mathcal{L}}^{n+1} + \mathcal{K}_{(\overline{\mathcal{X}}, \mathcal{D})/\mathbb{P}_{\mathbb{C}}^1} \cdot \overline{\mathcal{L}}^n, \quad a = -n(K_{(X, \Delta)} \cdot L^{n-1})/L^n$$

where $\overline{\mathcal{L}}$ denotes the \mathbb{C}^* -equivariant extension of \mathcal{L} to the \mathbb{C}^* -equivariant compactification $\overline{\mathcal{X}}$ of \mathcal{X} over $\mathbb{P}_{\mathbb{C}}^1$ and $\mathcal{K}_{(\overline{\mathcal{X}}, \mathcal{D})/\mathbb{P}_{\mathbb{C}}^1}$ denotes the relative log canonical divisor of the pair $(\overline{\mathcal{X}}, \mathcal{D})$ with \mathcal{D} denoting the Zariski closure in $\overline{\mathcal{X}}$ of the \mathbb{C}^* -orbit of the divisor Δ . Furthermore, $(X, \Delta; L)$ is said to be *K-polystable* if $\text{DF}_{\Delta}(\mathcal{X}, \mathcal{L}) \geq 0$ with equality iff $\mathcal{X} \simeq X \times \mathbb{A}_{\mathbb{C}}^1$ and *K-stable* if equality only holds when $\mathcal{X} \simeq X \times \mathbb{A}_{\mathbb{C}}^1$ for a \mathbb{C}^* -equivariant isomorphism.

In the case that $\pm K_{(X, \Delta)} > 0$ we will say that (X, Δ) is K-polystable if $(X, \Delta; \pm K_{(X, \Delta)})$ is K-polystable (and likewise for K-semistability). We recall the following results from [85, 87, 24]:

- When $K_X > 0$ (X, Δ) is K-polystable iff it is K-semistable iff (X, Δ) is log canonical
- When $kK_{(X, \Delta)}$ is trivial for some k , $(X, \Delta; L)$ is K-polystable for any L iff (X, Δ) is klt and K-semistable for any L iff (X, Δ) is lc.
- When $-K_X > 0$ the K-semistability of (X, Δ) implies that (X, Δ) is klt (however, the converse does not hold, in general).

When X is defined over \mathbb{F} we will say that $X(\mathbb{C})$ is K-polystable (etc) if X_{σ} is K-polystable (etc) for all complexifications X_{σ} .

2.4. Canonical heights and optimal models.

2.4.1. Canonical heights. A *metrized line bundle* $\overline{\mathcal{L}}$ is a line bundle $\mathcal{L} \rightarrow \mathcal{X}$ over an arithmetic variety \mathcal{X} such that the corresponding line bundle $L(\mathbb{C}) \rightarrow X(\mathbb{C})$ is endowed with a metric, that we shall denote by ϕ (as in Section 2.2.1); $\overline{\mathcal{L}} := (\mathcal{L}, \phi)$. We will assume that ϕ has finite energy. When ϕ is continuous the height $h_{\phi}(\mathcal{X}, \mathcal{L})$ and the normalized height $\hat{h}_{\phi}(\mathcal{X}, \mathcal{L})$ are defined by

$$h_{\phi}(\mathcal{X}, \mathcal{L}) := \overline{\mathcal{L}}^{n+1}, \quad \hat{h}_{\phi}(\mathcal{X}, \mathcal{L}) := \frac{\overline{\mathcal{L}}^{n+1}}{[\mathbb{F} : \mathbb{Q}] L^n (n+1)},$$

expressed in terms of the arithmetic top intersection numbers of $\overline{\mathcal{L}}$ [52, 45, 20, 106]. The normalized height is equivariant under scalings of the metric,

$$(2.5) \quad \hat{h}_{\phi+c} = \hat{h}_{\phi} + c/2, \quad \forall c \in \mathbb{R}.$$

and invariant under base change, induced by finite extensions of \mathbb{F} [20, Section 3.1.4]. The definition of $h_{\phi}(\mathcal{X}, \mathcal{L})$ extends naturally to any metrized \mathbb{R} -line bundle \mathcal{L} , by imposing homogeneity.

Lemma 2.4. *Let (X, L) be a polarized projective normal scheme over \mathbb{F} . Consider two metrized models $(\mathcal{X}, \overline{\mathcal{L}})$ and $(\mathcal{X}', \overline{\mathcal{L}'})$ of (X, L) over $\mathcal{O}_{\mathbb{F}}$. Assume that the induced isomorphism between $(\mathcal{X}, \mathcal{L})$ and $(\mathcal{X}', \mathcal{L}')$ yields an isometry between $\overline{\mathcal{L}}(\mathbb{C})$ and $\overline{\mathcal{L}'}(\mathbb{C})$. Then there exist integers $h(\mathfrak{p})$, where \mathfrak{p} ranges over a finite number of closed points of $\text{Spec } \mathcal{O}_{\mathbb{F}}$, such that*

$$h_{\phi'}(\mathcal{X}', \mathcal{L}') - h_{\phi}(\mathcal{X}, \mathcal{L}) = \sum_{\mathfrak{p}} h(\mathfrak{p}) \log N(\mathfrak{p})$$

Moreover, fixing a model \mathcal{Y} of $X_{\mathbb{F}}$ over $\mathcal{O}_{\mathbb{F}}$ dominating both \mathcal{X} and \mathcal{X}' and identifying \mathcal{L} and \mathcal{L}' with their pull-backs to \mathcal{Y} ,

$$h(\mathfrak{p}) = \sum_{0 \leq j \leq n} (\mathcal{L}'^j_{|\mathcal{Y}_{\mathfrak{p}}} \cdot \mathcal{L}_{|\mathcal{Y}_{\mathfrak{p}}}^{n-j}) \cdot E_{\mathfrak{p}}, \quad \sum_{\mathfrak{p}} E_{\mathfrak{p}} := (s = 0)$$

where $(s = 0)$ denotes the zero-divisor on \mathcal{Y} of the rational section s of $\overline{\mathcal{L}'} - \overline{\mathcal{L}}$ whose restriction to the generic fiber of \mathcal{X} equals $1 \in H^0(X_{\mathbb{F}}, \mathcal{O}_{X_{\mathbb{F}}}) (= \mathbb{F})$ and the intersection numbers are computed on the projective scheme $\mathcal{Y}_{\mathfrak{p}}$ over the residue field of \mathfrak{p} . More generally, the formulas above extend, by homogeneity, to the case when $L_{\mathbb{F}}$ is an \mathbb{R} -line bundle.

Proof. This follows from basic properties of arithmetic intersection numbers. For future reference we provide a proof. Using the multilinearity of arithmetic intersection numbers,

$$h(\overline{\mathcal{L}'}) - h(\overline{\mathcal{L}}) = \left(\sum_{j=0}^n \overline{\mathcal{L}'^j} \cdot \overline{\mathcal{L}^{n-j}} \right) \cdot (\overline{\mathcal{L}'} - \overline{\mathcal{L}}).$$

Now pull back \mathcal{L} and \mathcal{L}' to a model \mathcal{Y} as described in the lemma. By assumption, the restriction of $(\overline{\mathcal{L}'} - \overline{\mathcal{L}})$ to the generic fiber $X_{\mathbb{F}}$ of $\mathcal{Y} \rightarrow \text{Spec } \mathcal{O}_{\mathbb{F}}$ may be identified with the trivial line bundle $\mathcal{O}_{X_{\mathbb{F}}} \rightarrow X_{\mathbb{F}}$ endowed with its standard metric. The restriction formula for (generalized) arithmetic intersection numbers [20, Prop 2.3.1] ([46, Prop 6.3]) thus gives

$$\overline{\mathcal{L}'^j} \cdot \overline{\mathcal{L}^{n-j}} \cdot (\overline{\mathcal{L}'} - \overline{\mathcal{L}}) = (\overline{\mathcal{L}'^j} \cdot \overline{\mathcal{L}^{n-j}}) \cdot (s = 0) - \int_{X(\mathbb{C})} \log \|s\| (dd^c \phi)^{n-j} \wedge (dd^c \phi')^j.$$

Since $\|s\| = 1$ on $X(\mathbb{C})$ and $(s = 0)$ is a vertical divisor on \mathcal{Y} this concludes the proof. \square

Following [15], the functional $\phi \mapsto h_{\phi}(\mathcal{X}, \mathcal{L})$ admits a canonical extension to a functional on the space of all singular metrics ψ on \mathcal{L} with positive curvature current (using that $h_{\phi}(\mathcal{X}, \mathcal{L})$ is increasing in ϕ):

$$h_{\psi}(\mathcal{X}, \mathcal{L}) := \sup_{\phi \leq \psi} h_{\phi}(\mathcal{X}, \mathcal{L}),$$

where ϕ is assumed to be a continuous metric on $L(\mathbb{C})$ with positive curvature current. As observed in [15] $h_\psi(\mathcal{X}, \mathcal{L})$ is finite iff ψ has finite energy and then, for any fixed continuous metric ϕ_0 on $L(\mathbb{C})$,

$$(2.6) \quad h_\psi(\mathcal{X}, \mathcal{L}) = h_{\phi_0}(\mathcal{X}, \mathcal{L}) + \frac{1}{2} \mathcal{E}_{\phi_0}(\psi), \quad \mathcal{E}_{\phi_0}(\phi) = \sum_{\sigma} \mathcal{E}_{\phi_0^\sigma}(\phi^\sigma),$$

where ϕ^σ and ϕ_0^σ denote the restrictions of the metrics ϕ and ϕ_0 to $L_\sigma \rightarrow X_\sigma$ and \mathcal{E} denotes the functional 2.1. When $n = 1$ and $\nabla\psi \in L_{\text{loc}}^2$ this shows that the height $h_\psi(\mathcal{X}, \mathcal{L})$ coincides with the height defined wrt the generalized arithmetic intersection theory in [22].

Let now $(\mathcal{X}, \mathcal{D})$ be an arithmetic log pair such that $\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is relatively ample. We define the canonical height of $\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ as

$$(2.7) \quad h_{\text{can}}(\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})}) := \sup_{\phi} h_{\phi}(\mathcal{X}, \pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})}),$$

where the sup ranges over all volume-normalized psh metrics ϕ on $\pm K_{(X(\mathbb{C}), \Delta(\mathbb{C}))}$ of finite energy.

2.4.2. *Optimal models and the canonical height over \mathbb{F} .* A model $(\mathcal{X}^\circ, \mathcal{D}^\circ)$ over $\mathcal{O}_{\mathbb{F}}$ for a log pair $(X_{\mathbb{F}}, \Delta_{\mathbb{F}})$ will said to be *optimal* if $\pm\mathcal{K}_{(\mathcal{X}^\circ, \mathcal{D}^\circ)}$ is relatively ample (for some sign) and

$$\pm h(\overline{\pm\mathcal{K}_{(\mathcal{X}^\circ, \mathcal{D}^\circ)}}) = \min_{(\mathcal{X}, \mathcal{D})} \pm h(\overline{\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})}}),$$

for any fixed metric on $\pm K_{(X, \Delta)}$, where $(\mathcal{X}, \mathcal{D})$ ranges over all models over $\mathcal{O}_{\mathbb{F}}$ for $(X_{\mathbb{F}}, \Delta_{\mathbb{F}})$ such that $\pm\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is relatively ample. This definition is independent of the choice of metric, by Lemma 2.4.

2.5. The arithmetic Mabuchi functional and Odaka's modular invariant. Let $\overline{\mathcal{L}} \rightarrow \mathcal{X}$ be a metrized relatively ample line bundle over an arithmetic variety \mathcal{X} over $\mathcal{O}_{\mathbb{F}}$. When $X_{\mathbb{F}}$ is non-singular and the metric on $L(\mathbb{C})$ is smooth, then the corresponding *arithmetic Mabuchi (K-energy) functional* is defined as follows (in terms of Gillet-Soulé's arithmetic intersection numbers [52]):

$$(2.8) \quad \mathcal{M}_{\mathcal{X}}(\overline{\mathcal{L}}) := \frac{a}{(n+1)!} \overline{\mathcal{L}}^{n+1} + \frac{1}{n!} \overline{\mathcal{K}}_{\mathcal{X}} \cdot \overline{\mathcal{L}}^n, \quad a = -n(K_{X_{\mathbb{F}}} \cdot L_{\mathbb{F}}^{n-1})/L_{\mathbb{F}}^n,$$

where $K_{X(\mathbb{C})}$ is endowed with the metric induced by the normalized volume form ω^n/L^n of the curvature form ω of $\overline{\mathcal{L}}$ (giving total volume one to X).

Remark 2.5. We have followed the normalizations adopted in [2], which differ from Odaka's arithmetic Mabuchi functional [86] which uses the metric on K_X induced by the non-normalized volume form $\omega^n/n!$ (as explained in [2, Section 6.4], when X is Fano, and further discussed in Remark 4.2).

Let now $(\mathcal{X}, \mathcal{D})$ be a log pair over $\mathcal{O}_{\mathbb{F}}$ and $\overline{\mathcal{L}} \rightarrow \mathcal{X}$ a metrized relatively ample line bundle over \mathcal{X} . When $(X_{\mathbb{F}}, \Delta_{\mathbb{F}})$ is log smooth, log canonical and the metric ϕ has pre-log-log singularities in the sense of [28] (along the non-klt components of Δ) we define the

arithmetic log Mabuchi functional as follows, using the arithmetic intersection theory in [28, 27] (see also [72] for the case $n = 1$):

$$(2.9) \quad \mathcal{M}_{(\mathcal{X}, \mathcal{D})}(\overline{\mathcal{L}}) := \frac{a}{(n+1)!} \overline{\mathcal{L}}^{n+1} + \frac{1}{n!} (\overline{\mathcal{K}}_{(\mathcal{X}, \mathcal{D})}) \cdot \overline{\mathcal{L}}^n, \quad a = -n(K_{(X, \Delta)} \cdot L^{n-1})/L^n,$$

where $K_{(X, \Delta)(\mathbb{C})}$ is endowed with the normalized volume form ω^n/L^n of the curvature form ω of $\overline{\mathcal{L}}$, tensored with the singular metric on the \mathbb{Q} -line bundle Δ , induced by the (multivalued) holomorphic section cutting out Δ . The definition of $\mathcal{M}_{(\mathcal{X}, \mathcal{D})}(\overline{\mathcal{L}})$ mimics the definition of the Donaldson-Futaki invariant 2.4.

Note that in the case that $\mathcal{L} = \pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}$, that we shall focus on here,

$$(2.10) \quad \mathcal{M}_{(\mathcal{X}, \mathcal{D})}(\pm \overline{\mathcal{K}}_{(\mathcal{X}, \mathcal{D})}) = -\pm \frac{n}{(n+1)!} \overline{\mathcal{L}}^{n+1} + \frac{1}{n!} (\overline{\mathcal{K}}_{(\mathcal{X}, \mathcal{D})}) \cdot \overline{\mathcal{L}}^n$$

The *normalized arithmetic log Mabuchi functional* is defined by

$$\hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{D})}(\overline{\mathcal{L}}) := \frac{\mathcal{M}_{(\mathcal{X}, \mathcal{D})}(\overline{\mathcal{L}})}{[\mathbb{F} : \mathbb{Q}] L^n / n!}$$

It follows readily from the definition that if $q : \mathcal{Y} \rightarrow \mathcal{X}$ is a birational morphism over $\mathcal{O}_{\mathbb{F}}$ with \mathcal{Y} and \mathcal{X} normal, then

$$(2.11) \quad \mathcal{M}_{(\mathcal{X}, \mathcal{D})}(\overline{\mathcal{L}}) = \mathcal{M}_{(\mathcal{Y}, q^* \mathcal{D})}(\overline{q^* \mathcal{L}}).$$

2.5.1. Odaka's modular invariant. Consider now a polarized log pair $(X_{\mathbb{F}}, D_{\mathbb{F}}; L_{\mathbb{F}})$ over a number field \mathbb{F} . Following [86] (but using our different normalizations) we define its *normalized modular invariant* by

$$\hat{\mathcal{M}}(X_{\mathbb{F}}, D_{\mathbb{F}}; L_{\mathbb{F}}) := \inf \hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{D})}(\overline{\mathcal{L}}) \in]-\infty, \infty[$$

where the infimum runs over all metrized polarized log pairs $(\mathcal{X}, \mathcal{D}; \overline{\mathcal{L}})$ over $\mathcal{O}_{\mathbb{F}'}$ where \mathbb{F}' is a finite field extension of \mathbb{F} . The (non-normalized) *modular invariant* $\mathcal{M}(X_{\mathbb{F}}, D_{\mathbb{F}}; L_{\mathbb{F}})$ is defined by $\hat{\mathcal{M}}(X_{\mathbb{F}}, D_{\mathbb{F}}; L_{\mathbb{F}}) [\mathbb{F} : \mathbb{Q}] L^n / n!$

Example 2.6. When $X_{\mathbb{F}}$ is an abelian variety and $D_{\mathbb{F}} = 0$, our normalizations ensure that $\hat{\mathcal{M}}(X_{\mathbb{F}}, D_{\mathbb{F}}; L_{\mathbb{F}})$ is precisely Faltings' height [45] of $X_{\mathbb{F}}$, as follows from 3.7, combined with [86, Thm 2.14].

3. VARIATIONAL PRINCIPLES

3.1. The arithmetic Mabuchi functional when $L_{\mathbb{F}} = \pm K_{(X_{\mathbb{F}}, \Delta_{\mathbb{F}})}$ and finite energy metrics. Consider a general polarized log pair $(\mathcal{X}, \mathcal{D}; \mathcal{L})$ such that $L_{\mathbb{F}} = \pm K_{(X_{\mathbb{F}}, \Delta_{\mathbb{F}})}$. Denote by E_{\pm} the vertical divisor on \mathcal{X} cut out by the rational section of $\mathcal{L} - \pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ whose restriction to the generic fiber $X_{\mathbb{F}}$ of \mathcal{X} coincides with $1 \in H^0(X_{\mathbb{F}}, \mathcal{O}_{\mathbb{F}})$. Given a finite energy metric ϕ on $L(\mathbb{C})$ we then define $\hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{D})}(\overline{\mathcal{L}})$ as follows,

$$(3.1) \quad \hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{D})}(\mathcal{L}, \phi) := \pm \hat{h}_{\phi}(\mathcal{X}, \mathcal{L}) + \frac{1}{2} \text{Ent}(\text{MA}(\phi) | \mu_{\phi}) - \frac{E_{\pm} \cdot \mathcal{L}^n}{L_{\mathbb{F}}^n}$$

where $\text{Ent}(\mu|\mu_0)$ denotes the *entropy* of a measure μ relative to a measure μ_0 :

$$(3.2) \quad \text{Ent}(\mu|\mu_0) := \int_{X(\mathbb{C})} \log \frac{\mu}{\mu_0} \mu \left(= \sum_{\sigma} \int_{X_{\sigma}} \log \frac{\mu}{\mu_0} \mu \right),$$

if μ is absolutely continuous wrt μ_0 and $\text{Ent}(\mu|\mu_0) := \infty$, otherwise. Note that since ϕ has finite energy the height term is always finite. Moreover, since E_{\pm} is a vertical divisor the last term in formula 3.1 is independent of the metric on \mathcal{L} .

Lemma 3.1. *Assume that $(X_{\mathbb{F}}, \Delta_{\mathbb{F}})$ is log smooth and log canonical, that $L_{\mathbb{F}} = \pm K_{(X_{\mathbb{F}}, \Delta_{\mathbb{F}})}$ and that ϕ is a metric on $L(\mathbb{C})$ with pre-log-log singularities. Then the definitions 2.9 and 2.9 are compatible*

Proof. Let ϕ be a psh metric with pre-log-log singularities. Then ϕ has finite energy (see Example 2.2) and, as a consequence, μ_{ϕ} has total mass. We rewrite the definition 2.10 of $\mathcal{M}_{(\mathcal{X}, \mathcal{D})}(\overline{\mathcal{L}})$, where $\overline{\mathcal{L}} = (\mathcal{L}, \phi)$, as

$$\begin{aligned} \mathcal{M}_{(\mathcal{X}, \mathcal{D})}(\overline{\mathcal{L}}) &= -\pm \frac{n}{(n+1)n!} \overline{\mathcal{L}}^{n+1} + \frac{\pm}{n!} \overline{\mathcal{L}}^n + \frac{1}{n!} (\overline{\mathcal{K}}_{(\mathcal{X}, \mathcal{D})} - \pm \overline{\mathcal{L}}) \cdot \overline{\mathcal{L}}^n = \\ &\pm \frac{1}{(n+1)!} \overline{\mathcal{L}}^n + \frac{1}{n!} (\overline{\mathcal{K}}_{(\mathcal{X}, \mathcal{D})} - \pm \overline{\mathcal{L}}) \cdot \overline{\mathcal{L}}^n. \end{aligned}$$

Denote by ψ the induced metric on $K_{(X, \Delta)(\mathbb{C})}$. Since $\overline{\mathcal{K}}_{(\mathcal{X}, \mathcal{D})} - \pm \overline{\mathcal{L}} =: -E_{\pm}$ is a vertical divisor the restriction formula [46, Prop 6.3] yields

$$(\overline{\mathcal{K}}_{(\mathcal{X}, \mathcal{D})} - \pm \overline{\mathcal{L}}) \cdot \overline{\mathcal{L}}^n = \int_{X(\mathbb{C})} (\psi - \pm \phi)(dd^c \phi)^n - E_{\pm} \cdot \mathcal{L}^n.$$

Moreover, since the measure $(dd^c \phi)^n$ does not charge $X(\mathbb{C}) - \text{supp}(\Delta(\mathbb{C}))$ and ψ is represented by $\log \text{MA}(\phi)$ on $X - \text{supp}(\Delta)$ it follows that

$$\frac{1}{L_{\mathbb{F}}^n} (\overline{\mathcal{K}}_{(\mathcal{X}, \mathcal{D})} - \pm \overline{\mathcal{L}}) \cdot \overline{\mathcal{L}}^n = \frac{1}{2} \int_{X(\mathbb{C})} \log \frac{\text{MA}(\phi)}{\mu_{\phi}} \text{MA}(\phi) - E_{\pm} \cdot \mathcal{L}^n,$$

which concludes the proof. \square

3.2. Variational principles for metrics.

Lemma 3.2. *Let $(\mathcal{X}, \mathcal{D})$ be a log pair such that $\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is relatively ample and ϕ a volume-normalized psh metric on $\pm K_{(X, \Delta)(\mathbb{C})}$ with finite energy. Then*

$$\hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{D})}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}, \phi) \geq \pm \hat{h}_{\phi}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})})$$

with equality iff ϕ is a Kähler-Einstein metric.

Proof. When $\mathcal{L} = \pm \mathcal{K}_{\mathcal{X}}$ we have that $E_{\pm} = 0$ in formula 3.1. Indeed, $\mathcal{L} \pm \mathcal{K}_{\mathcal{X}}$ is the trivial line bundle and $1 \in H^0(\mathcal{X}, \mathcal{O}_{\mathcal{X}})$ has no zeroes on \mathcal{X} under our assumptions on \mathcal{X} (as shown precisely as in the case $\mathbb{F} = \mathbb{Q}$ considered in [2, Lemma 2.3]). The lemma thus follows from combining the expression 3.1 for $\hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{D})}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}, \phi)$ with the Kähler-Einstein equation 2.2, using that, for any given probability measures μ and μ_0 , $\text{Ent}(\mu|\mu_0) \geq 0$ with equality iff $\mu = \mu_0$ (by Jensen's inequality). \square

Proposition 3.3. *Let $(\mathcal{X}, \mathcal{D})$ be a log pair, whose complexification is klt, such that either $\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ or $-\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is relatively ample. Then*

$$(3.3) \quad \inf_{\phi} \hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{D})}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}, \phi) = \pm \sup_{\phi} \left(\hat{h}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}, \phi) : \phi \text{ vol-normalized} \right) := \pm \hat{h}_{can}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}),$$

where ϕ ranges over all finite energy psh metrics on $\pm K_{(X, \Delta)(\mathbb{C})}$. Moreover, the inf and sup above are attained iff ϕ is a Kähler-Einstein metric. In particular,

$$(3.4) \quad \inf_{\phi} \hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{D})}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}, \phi) = \pm \hat{h}_{can}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}) = \pm \hat{h}_{\phi_{KE}}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})})$$

for any volume-normalized Kähler-Einstein metric ϕ_{KE} , if such a metric exists (i.e. if $\pm K_{(X, \Delta)}$ is K -polystable). More generally, if $\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is relatively ample and $(\mathcal{X}, \mathcal{D})$ is log canonical (equivalently, (X, Δ) is K -stable and $K_{(X, \Delta)}$ admits a Kähler-Einstein metric) then the identities 3.4 still hold.

Proof. Introducing the normalized arithmetic log Ding functional defined by

$$(3.5) \quad \hat{D}_{(\mathcal{X}, \mathcal{D})}((\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}, \phi) = -\hat{h}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}, \phi) \pm \sum_{\sigma} \frac{1}{2[\mathbb{F} : \mathbb{C}]} \log \int_{X_{\sigma}} \mu_{\phi},$$

it is equivalent (by scaling the restrictions of ϕ to X_{σ}) to prove that

$$\inf_{\phi} \hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{D})}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}, \phi) = -\pm \inf_{\phi} \hat{D}((\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}, \phi),$$

where ϕ ranges over all finite energy metrics on $\pm K_{(X, \Delta)(\mathbb{C})}$. Now set $\mathcal{L} = \pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ and fix a reference metric ϕ_0 of finite energy on $L(\mathbb{C})$ (for example a continuous psh metric). We can then, using formula 3.1, rewrite

$$(3.6) \quad \hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{D})}(\mathcal{L}, \phi) := \pm \frac{1}{2} \left(2\hat{h}(\mathcal{L}, \phi) - \int_X (\phi - \phi_0) \text{MA}(\phi) \right) + \frac{1}{2} \text{Ent}(\text{MA}(\phi) | \mu_{\phi_0})$$

(the klt assumption ensures that μ_{ϕ_0} has finite total mass). Hence,

$$(3.7) \quad \hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{D})}(\mathcal{L}, \phi) = \frac{1}{2} \hat{\mathcal{M}}_{\phi_0}(\phi) \pm \hat{h}(\mathcal{L}, \phi_0)$$

where $\hat{\mathcal{M}}_{\phi_0}(\phi)$ is defined by replacing $(\mathcal{L}, \phi)^{n+1}$ in formula 3.6 with $\mathcal{E}_{\phi_0}(\phi)/2$. Likewise,

$$(3.8) \quad \hat{D}_{(\mathcal{X}, \mathcal{D})}((\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}, \phi) = \frac{1}{2} \hat{D}_{\phi_0}(\phi) - \hat{h}(\mathcal{L}, \phi_0),$$

where $\hat{D}_{\phi_0}(\phi)$ is defined by replacing $(\mathcal{L}, \phi)^{n+1}$ in formula 3.5 with $\mathcal{E}_{\phi_0}(\phi)/2$. All in all, by decomposing

$$\hat{\mathcal{M}}_{\phi_0}(\phi) = \sum_{\sigma} \hat{\mathcal{M}}_{\phi_0^{\sigma}}(\phi^{\sigma}), \quad \hat{D}_{\phi_0}(\phi) = \sum_{\sigma} \hat{D}_{\phi_0^{\sigma}}(\phi^{\sigma}),$$

where ϕ^{σ} is the restriction of ϕ to X_{σ} and $\hat{\mathcal{M}}_{\phi_0^{\sigma}}(\phi^{\sigma})$ and $\hat{D}_{\phi_0^{\sigma}}(\phi^{\sigma})$ are defined by decomposing both terms appearing in the definitions of $\hat{\mathcal{M}}_{\phi_0}(\phi)$ and $\hat{D}_{\phi_0}(\phi)$ wrt σ . All in all, this means that it is equivalent to prove the following identity:

$$(3.9) \quad \inf_{\phi^{\sigma}} \hat{\mathcal{M}}_{\phi_0^{\sigma}}(\phi^{\sigma}) = -\pm \inf_{\phi^{\sigma}} \hat{D}_{\phi_0^{\sigma}}(\phi^{\sigma}),$$

where ϕ_0^σ ranges over all psh metrics on L_σ with finite energy. But this identity follows from results in [4, 16]. For future reference we recall the reduction to [4, 16], which uses the thermodynamical formalism introduced in [4]. Let X be a complex projective variety and assume that $L = \pm K_{(X,\Delta)}$ is ample. Given a reference metric ϕ_0 on L consider the functional E on the space $\mathcal{P}(X)$ of all probability measures μ on X defined by

$$E(\mu) = \sup_{\phi} \left(\frac{\mathcal{E}_{\phi_0}(\phi)}{(n+1)L^n} - \int_X (\phi - \phi_0)\mu \right),$$

where the sup ranges over all psh metrics ϕ on L with finite energy. In the terminology introduced in [11] $E(\mu)$ is the *pluricomplex energy* of μ (relative to $dd^c\phi_0$).¹ Next, given $\beta \in \mathbb{R}$ the corresponding *free energy functional* F_β on $\mathcal{P}(X)$ is defined by

$$(3.10) \quad F_\beta(\mu) = \beta E(\mu) + \text{Ent}(\mu|\mu_{\phi_0}),$$

if $E(\mu) < \infty$. Otherwise, $F_\beta(\mu) := \infty$.² Recall that $\text{Ent}(\mu|\mu_0)$ is the relative entropy defined in formula 3.2. By [11, Thm A], $E(\mu) < \infty$ iff there exists a finite energy psh metric ϕ_μ solving $\text{MA}(\phi_\mu) = \mu$. Moreover, the sup defining $E(\mu)$ is then attained at ϕ_μ . Hence, if ϕ has finite energy, we can express

$$(3.11) \quad \hat{\mathcal{M}}_{\phi_0}(\phi) = F_{\pm 1}(\text{MA}(\phi)).$$

The identity 3.9 thus follows from the following identity, applied to $\beta = \pm 1$:

$$(3.12) \quad \inf_{\mu \in \mathcal{P}(X)} F_\beta(\mu) = -\beta \inf_{\phi} \hat{D}_{\phi_0}(\phi).$$

When $\beta = -1$ this identity follows from [4, Thm 1.1] for X non-singular and the same argument applies in general (see [16, Lemma 4.4]). When $\beta = 1$ the identity 3.12 follows from [4, Thm 3.3] when X is non-singular and, again, the same argument applies in general. Anyhow, we will prove the case $\beta = 1$ directly in the more general setup of log canonical pairs. But we first note that the statement in the proposition about Kähler-Einstein metrics follows from the well-known fact that the optimizers of both the Mabuchi functional $\hat{\mathcal{M}}_{\phi_0}$ and the Ding functional \hat{D}_{ϕ_0} (relative to ϕ_0) are precisely the Kähler-Einstein metrics on $\pm K_{(X,\Delta)}$. See [4, Thm 3.3] for the case $\beta = -1$ and [11, Thm C] and for the case $\beta = 1$.

Finally, assume that $\mathcal{K}_{(\mathcal{X},\mathcal{D})}$ is relatively ample and $(\mathcal{X}, \mathcal{D})$ is log canonical. By [14], $K_{(X,\Delta)(\mathbb{C})}$ admits a unique Kähler-Einstein metric ϕ_{KE} with finite energy. In particular, the corresponding measure $\mu_{\phi_{\text{KE}}}$ has finite total mass (as follows from the Kähler-Einstein equation 2.2). In fact, as shown in [14], ϕ_{KE} minimizes the corresponding Ding functional \hat{D}_{ϕ_0} , appearing in formula 3.8. It will thus be enough to show that ϕ_{KE} also minimizes $\hat{\mathcal{M}}_{\phi_0}$, or equivalently: that $\text{MA}(\phi_{\text{KE}})$ minimizes the corresponding free energy functionals F_1 for any X_σ . To this end restrict to X_σ and set $\phi_0 := \phi_{\text{KE}}$, assuming that ϕ_{KE} is volume-normalized, i.e. that μ_{ϕ_0} is a probability measure. This implies (by

¹we have followed the notation in [4], which differs from the notation in [11] where the pluricomplex energy is denoted $E^*(\mu)$.

²we have followed the notation in [7] which differs from the notation in [4] where the role of F_β is played by βF_β .

Jensen's inequality) that $\text{Ent}(\mu|\mu_{\phi_0}) \geq 0$ iff $\mu = \mu_{\phi_0}$. But, in general, we also have $E(\mu) = 0$ iff $\mu = \text{MA}(\phi_0)$ [11, 16]. Hence, $F(\mu) \geq 0$ with equality iff $\mu = \mu_{\phi_0}$. Since ϕ_0 is assumed to be Kähler-Einstein metric this concludes the proof. \square

We also note the following

Lemma 3.4. *The sup defining $\hat{h}_{\text{can}}(\pm\mathcal{K}_{(\mathcal{X},\mathcal{D})})$ may, equivalently, be taken over all continuous psh metrics on $\pm K_{(X,\Delta)(\mathbb{C})}$ and when $n = 1$ the sup may be taken over all continuous metrics. Moreover, if $(X, \Delta)(\mathbb{C})$ is log smooth and klt then both the inf and the sup in formula 3.3 may, equivalently, be taken over all log smooth psh metrics ϕ (i.e. such that the curvature form ω_ϕ of ϕ has conical singularities along Δ).*

Proof. To prove the first result recall that when L is an ample line bundle over a normal complex projective variety X any psh metric ψ on L is the decreasing limit of continuous (and even smooth) psh metrics ψ_j [33, Cor C]. Hence, the first statement of the lemma follows from the fact that the Ding function \hat{D}_{ϕ_0} , appearing in formula 3.8, is continuous under decreasing limits (indeed, for the integral term this follows from the monotone convergence theorem in integration theory and for the term $\mathcal{E}_{\phi_0}(\phi)$ this follows from [23, Thm 2.17]). Next consider the case when $n = 1$. Following [10], given a continuous psh metric ϕ on $L := \pm K_{(X,\Delta)(\mathbb{C})}$ denote by $P_X\phi$ the continuous psh metric on L defined as the sup of all continuous psh metrics ψ on L satisfying $\psi \leq \phi$. Then $P_X\phi \leq \phi$, giving $\pm \log \mu_{P_X\phi}(X_\sigma) \leq \pm \log \mu_\phi(X_\sigma)$. Hence, by formula 3.5, it is enough to show that $h_\phi(\mathcal{L}) \leq h_{P_X\phi}(\mathcal{L})$. But, by formula 2.6, this follows from

$$\int_{X(\mathbb{C})} (P_X\phi - \phi)(dd^c P_X\phi + dd^c \phi) = \int_{X(\mathbb{C})} (P_X\phi - \phi)(-dd^c P_X\phi + dd^c \phi) \geq 0,$$

using in the first equality that $\int_{X(\mathbb{C})} (P_X\phi - \phi)(dd^c P_X\phi) = 0$ (by [10, Prop 2.10]). The inequality then follows by integrating by parts to get $\int_{X(\mathbb{C})} d(P_X\phi - \phi) \wedge d^c(P_X\phi - \phi)$, which is an L^2 -norm and thus non-negative. Finally, to prove the statement concerning pairs $(X, \Delta)(\mathbb{C})$ that are log smooth and klt first note that, as in the proof of the previous proposition, it is enough to prove the corresponding statement for the log Mabuchi functional $\hat{\mathcal{M}}_{\phi_0}$ and log Ding functional \hat{D}_{ϕ_0} . But the latter property follows from essentially well-known regularization results for \hat{D}_{ϕ_0} and $\hat{\mathcal{M}}_{\phi_0}$. For example, when $\Delta = 0$, the regularization result in question for $\hat{\mathcal{M}}_{\phi_0}$ appears in [13, Lemma 3.1] and the case when $\Delta \neq 0$ is shown in precisely the same way, but replacing the use of the Calabi theorem in the proof of [13, Lemma 3.1] with [55, Thm A] (with $\mu = 0$). \square

It should be stressed that, in general, the finiteness of $\hat{h}_{\text{can}}(-\mathcal{K}_{(\mathcal{X},\mathcal{D})})$ does not imply that $-K_{(X,\Delta)}$ admits a Kähler-Einstein metric, or equivalently, that (X, Δ) is K-polystable. For example, when $\Delta = 0$, it was shown in [2, Thm 2.4] that the finiteness in question is equivalent to the K-semistability of X (which, in general, is weaker than K-polystability). More generally, we have:

Theorem 3.5. *Let $(\mathcal{X}, \mathcal{D})$ be a log pair such that $\pm\mathcal{K}_{(\mathcal{X},\mathcal{D})}$ is relatively ample. Then $\hat{h}_{\text{can}}(\pm\mathcal{K}_{(\mathcal{X},\mathcal{D})}) < \infty$ if and only if (X, Δ) is K-semistable.*

Proof. When $-\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is relatively ample this is shown in, essentially, the same way as in case $\Delta = 0$, considered in [2, Thm 2.4]. Next, for log pairs over \mathbb{C} such that $K_{(X, \Delta)}$ is ample it is shown in [14, Thm 2.4] that the inf of the corresponding Ding functional $\mathcal{D}_{\phi_0}(\phi)$ of all psh metrics ϕ of finite energy is finite iff (X, Δ) is lc, which concludes the proof using formula 3.8 and the results described in Section 2.3. \square

Corollary 3.6. *Let $(\mathcal{X}, \mathcal{D})$ be a log pair. If $-\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is relatively ample and $\hat{h}_{can}(-\mathcal{K}_{(\mathcal{X}, \mathcal{D})})$ is finite, then (X, Δ) is klt. If $\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is relatively ample and $\hat{h}_{can}(\mathcal{K}_{(\mathcal{X}, \mathcal{D})})$ is finite, then (X, Δ) is lc.*

Proof. This follows from the previous theorem, using the relations between the K-semistability of (X, Δ) and the singularities of (X, Δ) recalled in Section 2.3. Alternatively, a direct analytic proof can be given using that for any given finite energy psh metric ϕ on $-K_{(X, \Delta)}$ ($K_{(X, \Delta)}$) the total mass $\mu_\phi(X)$ is finite iff (X, Δ) is klt (lc) [14]. \square

3.2.1. *Intermezzo: the log Calabi-Yau case and Faltings' height.* Next, assume that $(\mathcal{X}, \mathcal{D})$ is a log Calabi-Yau pair, in the sense that there exists a positive integer k such that $k\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is trivial. Denote by α the multivalued meromorphic top form on X_σ defined as the tensor product of the k :th root of a generator of $H^0(\mathcal{X}, k\mathcal{K}_{(\mathcal{X}, \mathcal{D})})$ with the inverse of the (multivalued) section $s_{\mathcal{D}}$ cutting out \mathcal{D} . Then one can define a *Faltings' height* of $(\mathcal{X}, \mathcal{D})$ by

$$(3.13) \quad h_{\text{Falt}}(\mathcal{X}, \mathcal{D}) := -\frac{1}{2[\mathbb{F} : \mathbb{Q}]} \log \prod_{\sigma} \left(\frac{i}{2}\right)^{n^2} \int_{X_\sigma} \alpha_\sigma \wedge \bar{\alpha}_\sigma \in [\infty, \infty[,$$

which is finite iff $(X, \Delta)(\mathbb{C})$ is klt (as follows directly from the analytic characterization of klt pairs). When \mathcal{X} is an abelian variety and $\mathcal{D} = 0$ this is the usual definition of the Faltings height [45] (see also [37], where a different normalization is adopted).

Proposition 3.7. *Assume that some tensor power of $\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is trivial. Then, for any relatively ample line bundle \mathcal{L} over \mathcal{X} ,*

$$(3.14) \quad \inf_{\psi} \hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{L})}(\psi) = h_{\text{Falt}}(\mathcal{X}, \mathcal{D}),$$

where the inf ranges over all psh metrics on $L(\mathbb{C})$ of finite energy. In particular, the inf above is finite iff $(X, \Delta)(\mathbb{C})$ is klt.

Proof. In the case that (X, Δ) is klt the proof proceeds as in the case when $\Delta = 0$, considered in [3, Prop 6.5]. Next, when (X, Δ) is not klt we need to prove that the inf in the lemma equals $-\infty$. To this end fix a sequence of increasing compact sets C_j exhausting the complement in X of the support of Δ . We can take C_j to be the closure of open domains in X and consider the probability measures

$$\mu_j := 1_{C_j} (i/2)^{n^2} \alpha \wedge \bar{\alpha} / \int_{C_j} (i/2)^{n^2} \alpha \wedge \bar{\alpha},$$

where 1_{C_j} denotes the characteristic function of C_j . By [23, Thm B] there exists psh metrics ϕ_j on L of finite energy such that $\text{MA}(\phi_j) = \mu_j$. Indeed, ϕ_j is even locally

bounded. A slight variant of Lemma 3.1 gives

$$\mathcal{M}_{(\mathcal{X}, \mathcal{L})}(\phi_j) = \frac{1}{2} \text{Ent}(\text{MA}(\phi_j) | (i/2)^{n^2} \alpha \wedge \bar{\alpha}) = -\log \int_{C_j} (i/2)^{n^2} \alpha \wedge \bar{\alpha},$$

which converges to $-\log \int_X (i/2)^{n^2} \alpha \wedge \bar{\alpha}$, as $j \rightarrow \infty$, by the monotone convergence theorem. Finally, since, by the analytic characterization of klt pairs, $\int_X i^{n^2} \alpha \wedge \bar{\alpha}$ is finite iff (X, Δ) is klt, this concludes the proof. \square

In the light of the variational principles in Prop 3.3 and Lemma 3.7 it is thus natural to define

$$\pm \hat{h}_{\text{can}}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}) := h_{\text{Falt}}(\mathcal{X}, \mathcal{D})$$

when $(\mathcal{X}, \mathcal{D})$ is log Calabi-Yau.

Remark 3.8. As recalled in Section 2.3, a polarized log Calabi-Yau pair (X, Δ, L) is K-semistable iff (X, Δ) is lc. Hence, the previous lemma reveals that - in contrast to the case when $L = \pm K_{(X, \Delta)}$ - K-semistability is *not* equivalent to the finiteness of the inf of $\hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{L})}(\phi)$ over all metrics on L of finite energy, in the log Calabi-Yau case (only K-polystability is).

3.3. Variational principles for models. In this section we will, for simplicity, assume that all arithmetic varieties are *normal*. Consider two metrized models $(\mathcal{X}, \mathcal{D}; \bar{\mathcal{L}})$ and $(\mathcal{X}', \mathcal{D}'; \bar{\mathcal{L}}')$ for $(X_{\mathbb{F}}, \Delta_{\mathbb{F}}, L_{\mathbb{F}})$ over $\mathcal{O}_{\mathbb{F}}$. Assume that the induced isomorphism between $\bar{\mathcal{L}}(\mathbb{C})$ and $\bar{\mathcal{L}}'(\mathbb{C})$ is an isometry. Then the difference $\mathcal{M}_{(\mathcal{X}, \mathcal{D})}(\bar{\mathcal{L}}) - \mathcal{M}_{(\mathcal{X}', \mathcal{D}')}(\bar{\mathcal{L}}')$ is independent of the induced metric on $L_{\mathbb{F}}(\mathbb{C})$:

$$(3.15) \quad \mathcal{M}_{(\mathcal{X}', \mathcal{D}')}(\bar{\mathcal{L}}') - \mathcal{M}_{(\mathcal{X}, \mathcal{D})}(\bar{\mathcal{L}}) = \sum_b m(\mathfrak{b}) \log N(\mathfrak{b})$$

for a finite number of closed points $b \in \text{Spec } \mathcal{O}_{\mathbb{F}}$, where $N(b)$ denotes the cardinality of the residue field of b and $m(\mathfrak{b})$ may be expressed in terms of intersection numbers over the fiber \mathcal{Y}_b of any fixed model \mathcal{Y} dominating both \mathcal{X} and \mathcal{X}' . This is shown precisely as in the proof of Lemma 2.4. In fact, the difference 3.15 is even independent of the choice of a fixed metric on $K_{(X, \Delta)}(\mathbb{C})$ in formula 2.9. Accordingly, in this section we shall fix any pair of metrics on $L(\mathbb{C})$ and $K_{(X, \Delta)}(\mathbb{C})$ and denote by $\mathcal{M}_{(\mathcal{X}, \mathcal{D})}(\mathcal{L})$ the corresponding (generalized) arithmetic Mabuchi functional.

We recall the following result from [86, Thm 2.14]:

Theorem 3.9. (*Odaka*). *Given a projective scheme $(X_{\mathbb{F}}, K_{X_{\mathbb{F}}})$ such that $K_{X_{\mathbb{F}}}$ defines an ample \mathbb{Q} -line bundle, assume that \mathcal{X} is a model of $X_{\mathbb{F}}$ over $\mathcal{O}_{\mathbb{F}}$ such that \mathcal{X} is normal, $\mathcal{K}_{\mathcal{X}}$ is relatively ample and $(\mathcal{X}, \mathcal{X}_b)$ is log canonical for any closed point b . Then*

$$\mathcal{M}_{\mathcal{X}^\circ}(\mathcal{K}_{\mathcal{X}^\circ}) \leq \mathcal{M}_{\mathcal{X}'}(\mathcal{L}')$$

for any relatively ample model $(\mathcal{X}', \mathcal{L}')$ of $(X_{\mathbb{F}}, K_{X_{\mathbb{F}}})$ over $\text{Spec } \mathcal{O}_{\mathbb{F}}$ and metric on $K_X(\mathbb{C})$.

It follows from the previous theorem (applied to \mathcal{L}' of the form $\mathcal{K}_{\mathcal{X}'}$) that \mathcal{X}° is an optimal integral model for $X_{\mathbb{F}}$ (in the sense of 2.4.2). Using inversion of adjunction we

also deduce the following corollary, where \mathcal{X}_s is a stable model in the sense of Deligne-Mumford [38] (which always exists, after a base change):

Corollary 3.10. *Let $X_{\mathbb{F}}$ be a non-singular projective curve over \mathbb{F} such that $K_{X_{\mathbb{F}}} > 0$ and \mathcal{X}^s a stable model of $X_{\mathbb{F}}$ over $\text{Spec } \mathcal{O}_{\mathbb{F}}$. Then*

$$\mathcal{M}_{\mathcal{X}^s}(\mathcal{K}_{\mathcal{X}^s}) \leq \mathcal{M}_{\mathcal{X}'}(\mathcal{L}')$$

for any relatively ample model $(\mathcal{X}', \mathcal{L}')$ of $(X_{\mathbb{F}}, K_{X_{\mathbb{F}}})$ over $\text{Spec } \mathcal{O}_{\mathbb{F}}$. In particular, \mathcal{X}^s is an optimal model for $X_{\mathbb{F}}$ (in the sense of section 2.4.2).

Proof. Recall that $\mathcal{K}_{\mathcal{X}^s}$ is relatively ample [38, page 78]. Next, by inversion of adjunction for surfaces X over excellent rings [98, Thm 5.1] a log pair (X, C) , where C is assumed to be a reduced divisor, is log canonical iff the scheme C has semi-log canonical singularities, i.e. iff the log pair (C^ν, D^ν) is log canonical, where C^ν denotes the normalization of C and D^ν denotes the reduced divisor on C^ν defined by the conductor. Now, by the very definition of stable models in [38] the scheme \mathcal{X}_b^s is geometrically reduced and thus, in particular, reduced. Moreover, since the scheme \mathcal{X}_b^s has only ordinary double points (by definition), its normalization is regular. C^ν being reduced, it thus follows that $((\mathcal{X}_b^s)^\nu, D^\nu)$ is log canonical, as desired. \square

Remark 3.11. Let $X_{\mathbb{F}}$ be as in the previous corollary. Combining the previous corollary with Prop 3.3 reveals that $\mathcal{M}_{\mathcal{X}}(\mathcal{L}, \phi)$ is minimal when $(\mathcal{X}, \mathcal{L}) = (\mathcal{X}_s, \mathcal{K}_{\mathcal{X}^s})$ and ϕ is a Kähler-Einstein metric on $K_{X(\mathbb{C})}$. Likewise, the minimum of $h(\mathcal{K}_{\mathcal{X}})$ over all models \mathcal{X} of $X_{\mathbb{F}}$ with relatively ample $\mathcal{K}_{\mathcal{X}}$ and volume-normalized continuous psh metrics ϕ is attained for $\mathcal{X} = \mathcal{X}^s$ and ϕ the unique volume-normalized Kähler-Einstein metric on $K_{X(\mathbb{C})}$. This is in line with the suggestion put forth in [81, Section 3.1].

The previous theorem can be generalized to log pairs $(\mathcal{X}, \mathcal{D})$ such that:

$$(3.16) \quad (1) (\mathcal{X}, \mathcal{D} + \mathcal{X}_b) \text{ is lc for any closed } b \in \text{Spec } \mathcal{O}_{\mathbb{F}}, \quad (2) \mathcal{K}_{(\mathcal{X}, \mathcal{D})} \text{ is relatively ample}$$

But for our purposes it will be enough to consider the case of arithmetic surfaces:

Proposition 3.12. *Let $(\mathcal{X}, \mathcal{D})$ be an arithmetic log pair over $\mathcal{O}_{\mathbb{F}}$ satisfying conditions 1 and 2 above. Then*

$$\mathcal{M}_{(\mathcal{X}, \mathcal{D})}(\mathcal{K}_{(\mathcal{X}, \mathcal{D})}) \leq \mathcal{M}_{(\mathcal{X}', \mathcal{D}')}(\mathcal{L}')$$

for any relatively ample model $(\mathcal{X}', \mathcal{D}'; \mathcal{L}')$ of $(X_{\mathbb{F}}, \Delta_{\mathbb{F}}; K_{X_{\mathbb{F}}})$ over $\text{Spec } \mathcal{O}_{\mathbb{F}}$. More precisely, $m(b) \geq 0$ for any closed point $b \in \text{Spec } \mathcal{O}_{\mathbb{F}}$ (where $m(b)$ is the number appearing in formula 3.17). In particular, if $\mathcal{L}' = \mathcal{K}_{(\mathcal{X}', \mathcal{D}')}$, then $h(b) \geq 0$ for any closed point b and, as a consequence, $(\mathcal{X}, \mathcal{D})$ is an optimal model for $(X_{\mathbb{F}}, \Delta_{\mathbb{F}})$.

Proof. We will generalize the proof of Cor 3.10 and Thm 3.9, following the argument in [3, Section 6]. Set $\mathcal{L} := \mathcal{K}_{(\mathcal{X}, \mathcal{D})}$. By Step 1 in [3, Sections 6.2, 6.3.1], there exists a regular arithmetic surface \mathcal{Y} with birational morphisms p and q to \mathcal{X} and \mathcal{X}' , respectively (which are isomorphisms over the generic point of $\text{Spec } \mathcal{O}_{\mathbb{F}}$) such that

$$q^* \mathcal{L}' = p^* \mathcal{L} - E, \quad (\implies \mathcal{M}_{(\mathcal{X}', \mathcal{D}')}(\mathcal{L}') = \mathcal{M}_{(\mathcal{Y}, q^* \mathcal{D}')} (p^* \mathcal{L} - E))$$

for a p -exceptional effective \mathbb{Q} -divisor E on \mathcal{Y} , which vanishes iff p is an isomorphism 2.11 (using the pull-back formula 2.11 for the implication). A direct computation gives

$$(3.17) \quad \mathcal{M}_{(\mathcal{X}', \mathcal{D}')}(\mathcal{L}') - \mathcal{M}_{(\mathcal{X}, \mathcal{D})}(\mathcal{L}) = \frac{1}{2}q^*\mathcal{L}' \cdot E + q^*\mathcal{L}' \cdot (\mathcal{K}_{\mathcal{Y}/\mathcal{X}} - p^*\mathcal{D} + q^*\mathcal{D}').$$

The first term above is non-negative, since E is effective and $q^*\mathcal{L}'$ is semi-ample. Thus all that remains is to verify that $\mathcal{K}_{\mathcal{X}'/\mathcal{X}} - p^*\mathcal{D} + q^*\mathcal{D}'$ is effective, under the assumptions on \mathcal{D} . The condition 1 is, by inversion of adjunction on excellent surfaces [98, Thm 5.1], equivalent (since \mathcal{X} is normal) to the following property: $(\mathcal{X}, \mathcal{X}_b + \mathcal{D})$ is log canonical for any b . But then it follows from [66, Lemma 7.2 (4)] that $\mathcal{K}_{\mathcal{X}'/\mathcal{X}} - p^*\mathcal{D} + q^*\mathcal{D}'$ is effective. This proves the inequality 3.18. \square

In general, morphisms $(\mathcal{X}, \mathcal{D}) \rightarrow B$ satisfying the condition 1 are called *log canonical (lc)* in the context of the Minimal Model Program (MMP) [66]. The fibers \mathcal{X}_b are automatically reduced and if condition 2 also holds, then the restricted log pair $(\mathcal{X}_b, \mathcal{D}_b)$ is a *stable pair* in the sense of the MMP (i.e. $(\mathcal{X}_b, \mathcal{D}_b)$ is semi-log canonical and $\mathcal{K}_{(\mathcal{X}_b, \mathcal{D}_b)}$ is ample [65]). When $B = \text{Spec } \mathcal{O}_{\mathbb{F}}$ and $n = 1$ the existence of a model satisfying 1 and 2 above, after a base change, follows from [58, Prop 3.7] (under some regularity assumptions). The existence in any dimension is shown in [57, Cor 1.5], when the ground field is \mathbb{C} . Here we will focus on the following simple case:

Lemma 3.13. *Consider a log canonical pair $(\mathbb{P}_{\mathbb{F}}^1, \Delta_{\mathbb{F}})$ (i.e. the coefficients w_i of $\Delta_{\mathbb{F}}$ are in $[0, 1]$) such that $K_{(\mathbb{P}_{\mathbb{F}}^1, \Delta_{\mathbb{F}})} > 0$. Assume that either, (a) $\Delta_{\mathbb{F}}$ is supported on three \mathbb{F} -points (p_0, p_1, p_{∞}) in $\mathbb{P}_{\mathbb{F}}^1$, or, (b) $\Delta_{\mathbb{F}}$ is supported on four \mathbb{F} -points $(p_{\infty}, p_0; p_1, p_{-1})$ in $\mathbb{P}_{\mathbb{F}}^1$ with cross ratio -1 , such that $w_1 + w_{-1} \leq 1$. Then there exists an automorphism f of $\mathbb{P}_{\mathbb{F}}^1$ mapping (p_0, p_1, p_{∞}) to $(0, 1, \infty)$ and $(p_{\infty}, p_0; p_1, p_{-1})$ to $(\infty, 0; 1, -1)$ respectively and the Zariski closure \mathcal{D} of $f_*(\Delta_{\mathbb{F}})$ in $\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1$ has the property that $(\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1, \mathcal{D})$ satisfies conditions 1 and 2 above.*

Proof. The existence of f is a classical fact. By inversion of adjunction for excellent surfaces (see the proof of the second point below) it is enough to show that \mathcal{D}_b is log canonical for any b in $\text{Spec } \mathcal{O}_{\mathbb{F}}$, i.e. that $\mathcal{D} \otimes_{\mathcal{O}_{\mathbb{F}}} \mathbb{F}_b$ has coefficients in $[0, 1]$, where \mathbb{F}_b denotes the residue field of b . But this is immediate, since $\{0, 1, \infty\}$ corresponds to three distinct points in $\mathbb{P}_{\mathbb{F}_b}^1$ for any b . When the divisor $\mathcal{D} \otimes_{\mathcal{O}_{\mathbb{F}}} \mathbb{F}$ on $\mathbb{P}_{\mathbb{F}}^1$ is supported on $\{0, 1, -1, \infty\}$ the only case where $\{0, 1, -1, \infty\}$ does not correspond to four distinct points in $\mathbb{P}_{\mathbb{F}_b}^1$ is when 1 and -1 correspond to the same point in $\mathbb{P}_{\mathbb{F}_b}^1$ which thus acquires the weight $w_{-1} + w_1$. Hence, if $w_{-1} + w_1 \leq 1$, then $\mathcal{D}|_{\mathbb{P}_{\mathbb{F}_b}^1}$ has coefficients in $[0, 1]$, as desired. \square

The following proposition shows, in particular, that the log pair $(\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1, \mathcal{D})$ featuring in the previous lemma is the unique optimal model.

Proposition 3.14. *Consider a log pair $(\mathbb{P}_{\mathbb{F}}^1, \Delta_{\mathbb{F}})$ over a number field \mathbb{F} with coefficients $w_i \in [0, 1]$ such that $\pm K_{(\mathbb{P}_{\mathbb{F}}^1, \Delta_{\mathbb{F}})} > 0$. Let $(\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1, \mathcal{D}; \pm \mathcal{K}_{(\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1, \mathcal{D})})$ be a relatively ample model for $(\mathbb{P}_{\mathbb{F}}^1, \Delta_{\mathbb{F}}; \pm K_{(\mathbb{P}_{\mathbb{F}}^1, \Delta_{\mathbb{F}})})$ over $\mathcal{O}_{\mathbb{F}}$ satisfying the following conditions:*

- When $K_{(\mathbb{P}_{\mathbb{F}}^1, \Delta_{\mathbb{F}})} > 0$: $(\mathbb{P}_{\mathbb{F}_b}^1, \mathcal{D}_{|\mathbb{P}_{\mathbb{F}_b}^1})$ is log canonical for any closed point $b \in \text{Spec} \mathcal{O}_{\mathbb{F}}$, where \mathbb{F}_b denotes the residue field of b (i.e. the coefficients of $\mathcal{D}_{|\mathbb{P}_{\mathbb{F}_b}^1}$ are in $[0, 1]$) or equivalently: 1 and 2 in formula 3.16 holds.
- When $-K_{(\mathbb{P}_{\mathbb{F}}^1, \Delta_{\mathbb{F}})} > 0$: \mathcal{D} is the Zariski closure of the divisor on $\mathbb{P}_{\mathbb{F}}^1$ supported on $\{0, 1, \infty\}$ having the same coefficients as $\Delta_{\mathbb{Q}}$ and $(\mathbb{P}^1, \Delta)(\mathbb{C})$ is K -semistable (i.e. the weight conditions 1.5 hold)

Then

$$(3.18) \quad \mathcal{M}_{(\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1, \mathcal{D})}(\pm \mathcal{K}_{(\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1, \mathcal{D})}) \leq \mathcal{M}_{(\mathcal{X}', \mathcal{D}')}(\mathcal{L}')$$

for any model $(\mathcal{X}', \mathcal{D}'; \mathcal{L}')$ of $(\mathbb{P}_{\mathbb{F}}^1, \Delta_{\mathbb{F}}; \pm \mathcal{K}_{(\mathbb{P}_{\mathbb{F}}^1, \Delta_{\mathbb{F}})})$ over $\mathcal{O}_{\mathbb{F}}$ such that \mathcal{L}' is relatively ample. When $(\mathbb{P}_{\mathbb{C}}^1, \Delta_{\mathbb{C}})$ is K -stable (which is automatic if $K_{(\mathbb{P}_{\mathbb{F}}^1, \Delta_{\mathbb{F}})} > 0$) equality holds in 3.18 iff $(\mathcal{X}', \mathcal{D}') = (\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1, \mathcal{D})$ and $\mathcal{L}' = \pm \mathcal{K}_{(\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1, \mathcal{D})} + \pi^* M$ for some line bundle $M \rightarrow \text{Spec} \mathcal{O}_{\mathbb{F}}$.

Proof. The case when $-K_{(\mathbb{P}_{\mathbb{F}}^1, \Delta_{\mathbb{F}})} > 0$ is shown in [3, Section 6] when $\mathbb{F} = \mathbb{Q}$ (see [3, Remark 6.5] for the equality case) and the proof in the general case is essentially the same. In the case $K_{(\mathbb{P}_{\mathbb{F}}^1, \Delta_{\mathbb{F}})} > 0$ the inequality follows from proposition 3.12. We thus assume that equality holds. Then it follows from the proof of proposition 3.12 that $q^* \mathcal{L}' \cdot E = 0$, which means that $E^2 = 0$. Since E is a vertical divisor and \mathcal{X} is regular this can only happen if $E = \sum_b \lambda_b \mathcal{Y}_b$ for some $\lambda_b \in \mathbb{R}$. It follows that $E = 0$, since E is p -exceptional. This means that p is an isomorphism and thus $\mathcal{Y} \simeq \mathcal{X}$. Next, since all the fibers of $\mathcal{X} (= \mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1)$ over $\text{Spec} \mathcal{O}_{\mathbb{F}}$ are reduced and irreducible it then follows that q is also an isomorphism, $\mathcal{X}' \simeq \mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1$ and $\mathcal{L}' \simeq \mathcal{L} + \pi^* M$, if \mathcal{X}' is normal. The vanishing of the right hand side in formula 3.17 then forces $\mathcal{D}' = \mathcal{D}$, as desired (since $\mathcal{K}_{\mathcal{X}'/\mathcal{X}} - p^* \mathcal{D} + q^* \mathcal{D}'$ is effective, as shown in the proof of proposition 3.12). \square

Remark 3.15. The proof of the previous proposition shows that, in fact, $m(b) \geq 0$ for any closed point b , where $m(b)$ is the number appearing in formula 3.17. Moreover, if $m(b) = 0$ for all b in a open subset U of $\text{Spec} \mathcal{O}_{\mathbb{F}}$, then \mathcal{X}' is isomorphic to $\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1$ over U and, under such an isomorphism, $\mathcal{D}' = \mathcal{D}$ over U .

In the light of the discussion following Remark 3.11 it seems natural to pose the following conjecture:

Conjecture 3.16. *Given a number field \mathbb{F} and a log pair $(X_{\mathbb{F}}, D_{\mathbb{F}})$ such that $\pm K_{(X_{\mathbb{F}}, \Delta_{\mathbb{F}})}$ is ample*

$$\inf_{(\mathcal{X}, \mathcal{D}; \mathcal{L})} \hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{D})}(\mathcal{L}) = \inf_{(\mathcal{X}, \mathcal{D})} \hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{D})}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})})$$

where $(\mathcal{X}, \mathcal{D}; \mathcal{L})$ ranges over all polarized models of $(X_{\mathbb{F}}, \Delta_{\mathbb{F}}; \pm K_{(X_{\mathbb{F}}, \Delta_{\mathbb{F}})})$ over $\mathcal{O}_{\mathbb{F}}$ and $(\mathcal{X}, \mathcal{D})$ ranges over all models of $(X_{\mathbb{F}}, \Delta_{\mathbb{F}})$ such that $\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is relatively ample.

Assuming the validity of this conjecture and the existence of optimal models $(\mathcal{X}^{\circ}, \mathcal{D}^{\circ})$ (defined in section 2.4.2) for sufficiently large field extensions, it follows that the normalized modular invariant $\hat{\mathcal{M}}(X_{\mathbb{F}}, D_{\mathbb{F}}; \pm K_{(X_{\mathbb{F}}, \Delta_{\mathbb{F}})})$ coincides with $\pm h_{\text{can}}(\mathcal{X}^{\circ}, \mathcal{D}^{\circ})$. For

example, the previous conjecture holds for the log pairs $(\mathbb{P}_{\mathbb{F}}^1, \Delta_{\mathbb{F}})$ appearing in the previous proposition.

Remark 3.17. According to a conjecture of Odaka [86], the infimum of $\hat{\mathcal{M}}_{\mathcal{X}}(\mathcal{L})$ over $(\mathcal{X}, \mathcal{L})$ is attained at any globally K-semistable model, i.e. $(\mathcal{X}, \mathcal{L})$, i.e. a model all whose fibers over $\text{Spec } \mathcal{O}_{\mathbb{F}}$ are K-semistable (see [59] for recent progress on this conjecture). In particular, if this conjecture holds, then any globally K-semistable model of the form $(\mathcal{X}, \pm \mathcal{K}_{\mathcal{X}})$ is optimal in the sense of Section 2.4.2.

3.3.1. *Relatively minimal models.* Consider a non-singular projective curve X over a number field \mathbb{F} such that $K_X > 0$. It admits a unique regular model \mathcal{X}_{\min} over $\mathcal{O}_{\mathbb{F}}$ which is *minimal*, or equivalently: *relatively minimal* (obtained by repeatedly blowing down vertical (-1) -curves). $\mathcal{K}_{\mathcal{X}_{\min}}$ is nef. Contracting the vertical (-2) -curves in \mathcal{X}_{\min} yields a birational morphism from \mathcal{X}_{\min} to a projective normal scheme over $\mathcal{O}_{\mathbb{F}}$, called the *canonical model* \mathcal{X}_{can} of X (not to be confused with the canonical model of a Shimura curve). \mathcal{X}_{can} is Gorenstein and $\mathcal{K}_{\mathcal{X}_{\text{can}}}$ is relatively ample [76, Cor 4.18].

Proposition 3.18. *Let $X_{\mathbb{F}}$ be a non-singular projective curve over \mathbb{F} such that $K_{X_{\mathbb{F}}} > 0$. Then*

$$\mathcal{M}_{\mathcal{X}_{\text{can}}}(\mathcal{K}_{\mathcal{X}_{\text{can}}}) = \mathcal{M}_{\mathcal{X}_{\min}}(\mathcal{K}_{\mathcal{X}_{\min}}) \leq \mathcal{M}_{\mathcal{X}'}(\mathcal{L}')$$

for any relatively ample model $(\mathcal{X}', \mathcal{L}')$ of $(X_{\mathbb{F}}, K_{X_{\mathbb{F}}})$ over $\text{Spec } \mathcal{O}_{\mathbb{F}}$. In particular, \mathcal{X}_{can} is an optimal model for $X_{\mathbb{F}}$ (in the sense of section 2.4.2).

Proof. This follows from results outlined in [86]. For completeness we provide a proof. Denoting by $\nu : \mathcal{X}'_{\nu} \rightarrow \mathcal{X}'$ the normalization of \mathcal{X}' , one first observes that $\mathcal{M}_{\mathcal{X}'_{\nu}}(\nu^* \mathcal{K}_{\mathcal{X}'}) \leq \mathcal{M}_{\mathcal{X}'}(\mathcal{K}_{\mathcal{X}'})$, using that the Weil divisor $\mathcal{K}_{\mathcal{X}'_{\nu}/\mathcal{X}'}$ is anti-effective (just as in the proof of [86, Prop 2.17]). Next, fixing a birational morphism $f : \mathcal{X} \rightarrow \mathcal{X}'_{\nu}$ from a regular arithmetic surface \mathcal{X} , yields, by 2.11, $\mathcal{M}_{\mathcal{X}'_{\nu}}(\nu^* \mathcal{K}_{\mathcal{X}'}) = \mathcal{M}_{\mathcal{X}}(\mathcal{L})$, where \mathcal{L} is the relatively semi-ample line bundle on \mathcal{X} obtained by pulling back $\nu^* \mathcal{K}_{\mathcal{X}'}$ to \mathcal{X} . Finally, it follows from [86, Thm 2.20] that

$$(3.19) \quad \mathcal{M}_{\mathcal{X}_{\min}}(\mathcal{K}_{\mathcal{X}_{\min}}) \leq \mathcal{M}_{\mathcal{X}}(\mathcal{L}).$$

by running the Minimal Model Program with scaling [86, Thm 2.20]. For completeness we detail the proof of the inequality 3.19 in the present setup. Let $\mathcal{L}^{(0)}$ be a relatively nef line bundle over a regular model $\mathcal{X}^{(0)}$ of X and set $\mathcal{L}_t^{(0)} := \mathcal{L}^{(0)} + t(\mathcal{K}_{\mathcal{X}^{(0)}} - \mathcal{L}^{(0)})$. A direct computation reveals that

$$d\mathcal{M}_{\mathcal{X}^{(0)}}(\mathcal{L}_t^{(0)})/dt = (\mathcal{K}_{\mathcal{X}^{(0)}} - \mathcal{L}^{(0)}) \cdot (\mathcal{K}_{\mathcal{X}^{(0)}} - \mathcal{L}^{(0)}) \leq 0,$$

using in the last step that $\mathcal{K}_{\mathcal{X}^{(0)}} - \mathcal{L}^{(0)}$ is a vertical divisor on a regular arithmetic surface. In particular, if $\mathcal{K}_{\mathcal{X}^{(0)}}$ is relatively nef, then we are done, since then $\mathcal{K}_{\mathcal{X}^{(0)}}$ is the unique relative minimal model ($\mathcal{X}^{(0)}$ cannot contain any (-1) -curves, by adjunction).

Otherwise, denote by t_0 the sup over all $t \in [0, 1]$ such that $\mathcal{L}_t^{(0)}$ is relatively nef. By assumption, $t_0 \in [0, 1[$. Note that the cone of all effective vertical divisors on $\mathcal{X}^{(0)}$, modulo numerical equivalence, is generated by a finite number of extremal effective divisors C_i (for elementary reasons; see [98, Lemma 2.13] for a more general statement).

As a consequence, t_0 is rational and there exists an extremal effective divisor C such that $\mathcal{L}_{t_0}^{(0)} \cdot C = 0$ and $\mathcal{K}_{\mathcal{X}^{(0)}} \cdot C < 0$ (just as in [18, Lemma 3.10.8]). It follows that there exists a birational morphism

$$f_0 : \mathcal{X}^{(0)} \rightarrow \mathcal{X}^{(1)}$$

to a regular model $\mathcal{X}^{(1)}$ of X that contracts precisely C . Indeed, in general, C induces an extremal contraction $f : \mathcal{X} \rightarrow \mathcal{Z}$ and since $K_X > 0$ this contraction is birational and $f(C)$ is 0-dimensional (see [98, Thm 4.4] and [18, Remark 3.10.9]). Thus, by [71, Thm 1.5], there exists a birational morphism $f_0 : \mathcal{X}^{(0)} \rightarrow \mathcal{X}^{(1)}$ with $\mathcal{X}^{(1)}$ regular, mapping C to closed point x on $\mathcal{X}^{(1)}$ and such that f_0 restricts to an isomorphism from $\mathcal{X}^{(0)} - C$ to $\mathcal{X}^{(1)} - \{x\}$. Denote by $\mathcal{L}^{(1)}$ the \mathbb{Q} -line bundle on $\mathcal{X}^{(1)}$ defined as the push-forward of $\mathcal{L}_{t_0}^{(0)}$ under f_0 . It satisfies $f_0^* \mathcal{L}^{(1)} = \mathcal{L}_{t_0}^{(0)}$ and thus $\mathcal{M}_{\mathcal{X}^{(0)}}(\mathcal{L}_{t_0}^{(0)}) = \mathcal{M}_{\mathcal{X}^{(1)}}(\mathcal{L}^{(1)})$, by 2.11. Since $\mathcal{L}^{(1)}$ is relatively nef we can repeat this procedure in a finite number of steps until $\mathcal{K}_{\mathcal{X}^{(j)}}$ is nef (using that the Picard number decreases at each step) and thus $\mathcal{X}^{(j)}$ is the unique relative minimal model of X . This proves the inequality 3.19. Finally, since the canonical morphism $F : \mathcal{X}_{\min} \rightarrow \mathcal{X}_{\text{can}}$ satisfies $F^* \mathcal{K}_{\mathcal{X}_{\text{can}}} = \mathcal{K}_{\mathcal{X}_{\min}}$ [76, Cor 4.18] this concludes the proof of the proposition. \square

Remark 3.19. The previous proposition yields an alternative proof of Cor 3.10. Indeed, if X admits a semistable model over $\mathcal{O}_{\mathbb{F}}$ (in the sense of Deligne-Mumford), then the regular minimal model \mathcal{X}_{\min} of X over $\mathcal{O}_{\mathbb{F}}$ is also semistable and \mathcal{X}_{can} is the stable model [76, Thm 10.3.34].

4. VARIATIONS OF THE CANONICAL HEIGHT WITH RESPECT TO THE COEFFICIENTS OF \mathcal{D}

We will say that a set of log pairs $\{(\mathcal{X}, \mathcal{D})\}$ is a *linear family* if \mathcal{X} and the irreducible components of \mathcal{D} are fixed and $\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is proportional to one and the same relatively ample line bundle, i.e. $\mathcal{K}_{(\mathcal{X}, \mathcal{D})} \cong s\mathcal{L}_0$ for some $s \in \mathbb{R}$ (depending on the coefficients w_i of \mathcal{D}) and some relatively ample line bundle \mathcal{L}_0 (independent of w_i).

Proposition 4.1. *Let $\{(\mathcal{X}, \mathcal{D})\}$ be a linear family of log canonical (lc) pairs. Then $\pm \hat{h}_{\text{can}}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})})$ is concave wrt the coefficients $\mathbf{w} \in \mathbb{R}^m$ of Δ (assuming that the sign is chosen so that $\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is relatively ample). In particular, $\pm \hat{h}_{\text{can}}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})})$ is continuous wrt \mathbf{w} in the interior \dot{C} of the convex set $C \subseteq \mathbb{R}^m$ of all \mathbf{w} for which $\pm \hat{h}_{\text{can}}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})})$ is finite. Furthermore, $\pm \hat{h}_{\text{can}}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})})$ is continuous along any affine segment I in C , homeomorphic to $]0, 1]$, if the interior of I is contained in \dot{C} .*

Proof. By the variational principles in Prop 3.3 and Prop 3.7 we can, express

$$(4.1) \quad \pm \hat{h}_{\text{can}}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}) = \inf_{\psi_0} \hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{D})}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}, s\psi_0),$$

where ψ_0 ranges over all psh metrics on $L_0(\mathbb{C})$ of finite energy. Indeed, since we have assumed $\mathcal{K}_{(\mathcal{X}, \mathcal{D})} \cong s\mathcal{L}_0$, any psh metric ψ on $\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}(\mathbb{C})$ may be expressed as $\psi = s\psi_0$ for some psh metric ψ_0 on $L_0(\mathbb{C})$ (namely, $\psi_0 := \psi/s$). Moreover, since we are assuming that the infimum is finite we may as well assume that $\text{Ent}(\text{MA}(\psi_0)|_{\mu_{s\psi_0}}) < \infty$. To

prove the concavity of $\pm \hat{h}_{\text{can}}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})})$ wrt \mathbf{w} it will thus be enough to show that for a fixed such psh metric ψ_0 on L_0 , $\hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{D})}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}, \pm s\psi_0)$ is affine with respect to \mathbf{w} . To this end we will exploit the expression 3.1. Setting $\phi = \pm s\psi_0$ we have

$$(4.2) \quad \text{MA}(\pm s\psi_0) = \text{MA}(\psi_0), \quad \mu_\phi = e^{s\psi_0} |s_1|^{-2w_1} \dots \left(\frac{i}{2}\right)^{n^2} dz \wedge d\bar{z}$$

using homogeneity in the first equality and, in the second equality, a local representation as in section, where s_i are the sections cutting out the irreducible components of \mathcal{D} . Since since s is affine in \mathbf{w} it we deduce that $\log \frac{\text{MA}(s\psi_0)}{\mu_{s\psi_0}}$ is affine in \mathbf{w} , which implies that $\text{Ent}(\text{MA}(s\psi_0)|\mu_{s\psi_0})$ is affine in \mathbf{w} . Finally, by homogeneity,

$$\pm h_{\pm s\psi_0}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}) = s h_{\psi_0}(\mathcal{L}_0).$$

which concludes the proof that $\hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{D})}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}, \pm s\psi_0)$ is affine, showing that $\hat{h}_{\text{can}}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})})$ is concave. Since any convex functions is continuous on a open subset where it is finite it follows that $\hat{h}_{\text{can}}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})})$ is continuous in \dot{C} . The last continuity statement in the proposition also follows from elementary properties of convex functions (see [2, Lemma 2.10]). \square

Remark 4.2. If one were to instead metrize $K_{(X, \Delta)}$ with the metric induced by the volume form $\omega_{\text{KE}}^n/n!$ of the Kähler-Einstein metric ω_{KE} (without normalizing the volume), then the corresponding normalized height $\pm \hat{h}(\pm \mathcal{K}_{(X, \Delta)})$ would always diverge as $K_{(X, \Delta)}$ approaches the trivial line bundle. Indeed, by the scaling relation 2.5

$$\pm \hat{h}(\pm \mathcal{K}_{(X, \Delta)}) = \pm \hat{h}_{\text{can}}(\pm \mathcal{K}_{(X, \Delta)}) \pm \frac{1}{2} \log \frac{c_1(\pm K_{(X, \Delta)})^n}{n!},$$

where the second term diverges as $K_{(X, \Delta)}$ approaches the trivial line bundle.

Proposition 4.3. *Let $\{(\mathcal{X}, \mathcal{D}_{\mathbf{w}})\}$ be a linear family of log smooth klt pairs such that $\pm K_{(X, \Delta_{\mathbf{w}})}$ is K-stable and assume that the coefficients \mathbf{w} range over an open subset G of \mathbb{R}^m . Then $\pm \hat{h}_{\text{can}}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D}_{\mathbf{w}})})$ is real-analytic wrt \mathbf{w} in G .*

Proof. By Hartog's classical theorem on separate holomorphicity it is enough to consider the case when $\{(\mathcal{X}, \mathcal{D}_{\mathbf{w}})\}$ is a one-parameter family: i.e. \mathbf{w} depends linearly on a parameter $t \in]0, 1[$. To simplify the notation we assume that $\mathbb{F} = \mathbb{Q}$ so that there is only one complex embedding σ of \mathbb{F} . But the proof in the general case is essentially the same. We will write $X = X(\mathbb{C})$ and $L = L(\mathbb{C})$. As recalled in Section 2.3, the K-stability assumption is equivalent to the existence of a unique volume-normalized Kähler-Einstein metric ϕ_t on $\pm K_{(X, \Delta_{\mathbf{w}(t)})}$ (which, by [36] and [12, Thm 2.19], is equivalent to the properness of the Mabuchi functional appearing in formula 3.7). Expressing $\phi_t = \pm s\psi$ for ψ a metric on L_0 , just as in the proof of the previous proposition (where ψ was denoted ψ_0), the Kähler-Einstein equation 2.2 for ϕ_t translates (using 4.2) into

$$(4.3) \quad \text{MA}(\psi) = e^{s(t)\psi} |s_1|^{-2w_1(t)} \dots \left(\frac{i}{2}\right)^{n^2} dz \wedge d\bar{z}.$$

Note that the right hand side of this equation depends real-analytically on (t, ψ) , since $s(t)$ and $w_i(t)$ depend linearly on t . Hence, assuming that one can apply the implicit function theorem in an appropriate Banach space, the real-analyticity of $\pm \hat{h}_{\text{can}} (\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})})$ wrt t then follows precisely as in the proof of [7, Thm 7.9] (where a different family of twisted Kähler-Einstein equations was considered, where the role s is played by β and $\Delta = 0$). Finally, as explained in [9, Section 2.4.3], the implicit function theorem can indeed be applied under the assumptions of the proposition, using the theory for linearizations of equations of the form 4.3, established in [40, 64] in the case that the components of Δ do not intersect and announced in [82] in the log smooth case. \square

Example 4.4. Let \mathcal{X} be an arithmetic Fano variety and \mathcal{D}_1 a divisor cut out by an element in $H^0(\mathcal{X}, -\mathcal{K}_{\mathcal{X}})$. Then, for $w \in \mathbb{R}$, $(\mathcal{X}, w\mathcal{D}_1)$ is a linear family as above with $s = 1 - w$ and $\mathcal{L}_0 = -\mathcal{K}$. When Δ_1 is defined by an irreducible non-singular hypersurface it follows from a conjecture of Donaldson, established in [4], that the log Fano variety $(X, (1 - s)\Delta_1)$ admits a unique Kähler-Einstein metric ω_s for any sufficiently small positive number s (corresponding to a psh metric ϕ_s on \mathcal{L} of finite energy). As a consequence, by the previous two propositions $\hat{h}_{\text{can}} (-\mathcal{K}_{(\mathcal{X}, (1-s)\mathcal{D})})$ is real-analytic and concave for s sufficiently small. As another example (where \mathcal{D}_1 is not irreducible) let \mathcal{X} be the canonical model over \mathbb{Z} of a toric Fano variety $X_{\mathbb{Q}}$ and let \mathcal{D}_1 be the standard torus invariant anti-canonical divisor on \mathcal{X} . Then $(X, (1 - s)\Delta_1)$ is K-polystable for any $s \in [0, 1]$ and, by [3, Lemma 3.2], $-\hat{h}_{\text{can}} (-\mathcal{K}_{(\mathcal{X}, (1-s)\mathcal{D})}) = -\hat{h}_{\text{can}} (-\mathcal{K}_{\mathcal{X}}) + \frac{n}{2} \log s$, which is, indeed concave wrt s and continuous as $s \rightarrow 1$ and $s \rightarrow 0$ (since the log Calabi-Yau $(\mathcal{X}, \mathcal{D})$ is not klt) in accordance with Prop 4.1. However, while $-\hat{h}_{\text{can}} (-\mathcal{K}_{(\mathcal{X}, (1-s)\mathcal{D})})$ is real-analytic wrt s , this does not follow from Prop 4.3, since $(X, (1 - s)\Delta)$ is not K-stable in this case (but it seems likely that the real-analyticity could be deduced from a generalization of Prop 4.3 taking a maximal compact subgroup of the automorphism group of X into account).

5. CANONICAL HEIGHTS IN TERMS OF PERIODS

We start with some notation. Given a \mathbb{Q} -divisor Δ on a complex projective variety X and a positive integer N we will use the same notation Δ for the divisor on the N -fold product X^N of X defined as the sum of the N :th pull backs of the divisor Δ on X under the N projections onto the different factors of X^N . We will denote by s_{Δ} the corresponding (multi-valued) holomorphic section over X_{reg}^N cutting out the restriction of Δ to X_{reg}^N (where X_{reg} denotes the regular locus of X).

5.1. The case $K_{(X, \Delta)} > 0$. Let $(\mathcal{X}, \mathcal{D})$ be an arithmetic log pair over $\mathcal{O}_{\mathbb{F}}$ such that that $\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is a relatively ample \mathbb{Q} -line bundle over \mathcal{X} and assume that $(X_{\mathbb{Q}}, \Delta_{\mathbb{Q}})$ is klt. Given a positive real number k such that $k\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is a line bundle (i.e. Cartier) denote by N_k the rank of the $\mathcal{O}_{\mathbb{F}}$ -module $H^0(\mathcal{X}, k\mathcal{K}_{(\mathcal{X}, \mathcal{D})})$:

$$N_k := \dim_{\mathbb{C}} (H^0(\mathcal{X}, k\mathcal{K}_{(\mathcal{X}, \mathcal{D})}) \otimes_{\sigma} \mathbb{C})$$

for any embedding σ of \mathbb{F} into \mathbb{C} (the subscript k will occasionally be omitted to simplify the notation). The exterior power $\Lambda^{N_k} (H^0(\mathcal{X}, k\mathcal{K}_{(\mathcal{X}, \mathcal{D})}))$ thus has rank one. We fix

a non-trivial element in $\Lambda^{N_k} (H^0(\mathcal{X}, k\mathcal{K}_{(\mathcal{X}, \mathcal{D})}))$ that we note by $\det S^{(k)}$. For example, $\det S^{(k)}$ can be taken to be the N_k -fold exterior product of any N_k elements $s_1^{(k)}, \dots, s_{N_k}^{(k)}$ in $H^0(\mathcal{X}, k\mathcal{K}_{(\mathcal{X}, \mathcal{D})})$ that define a basis in $H^0(\mathcal{X}, k\mathcal{K}_{(\mathcal{X}, \mathcal{D})}) \otimes_{\sigma} \mathbb{C}$. Under the standard natural embedding of $\Lambda^{N_k} (H^0(X, k\mathcal{K}_{(X, \Delta)_{\sigma}}))$ into $H^0(\mathcal{X}^N, k\mathcal{K}_{(X^N, \Delta)_{\sigma}})$ we can identify the complexifications of $\det S^{(k)}$ with a holomorphic section of $kK_{(X^{N_k}, \Delta_k)}$:

$$(5.1) \quad (\det S_{\sigma}^{(k)})(x_1, x_2, \dots, x_{N_k}) = \det(s_i^{(k)}(x_j)),$$

Thus

$$\alpha_{\sigma}^{(k)} := \left(\det S_{\sigma}^{(k)} \right)^{1/k} \otimes s_{\Delta}^{-1}$$

defines a multivalued meromorphic top form on X^{N_k} (i.e. a multivalued meromorphic section of $K_{X^{N_k}}$). Set

$$\mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D})_{\sigma} := \left(\frac{i}{2} \right)^{(N_k n)^2} \int_{X_{\sigma}^{N_k}} \alpha_{\sigma}^{(k)} \wedge \overline{\alpha_{\sigma}^{(k)}} \in \mathbb{R}, \quad \mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D}) := \prod_{\sigma} \mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D})_{\sigma}$$

(the klt assumption ensures that $\mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D})_{\sigma} < \infty$). The product

$$(5.2) \quad \left(\frac{\Lambda^{N_k} (H^0(\mathcal{X}, k\mathcal{K}_{(\mathcal{X}, \mathcal{D})}))}{\mathcal{O}_{\mathbb{F}}(\det S^{(k)})} \right)^{1/k} \cdot \mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D})$$

is, for any given k , an invariant of $(\mathcal{X}, \mathcal{D})$, as follows directly from the product formula in \mathbb{F} .

Remark 5.1. If $\mathbb{F} = \mathbb{Q}$ then $\Lambda^{N_k} (H^0(\mathcal{X}, k\mathcal{K}_{(\mathcal{X}, \mathcal{D})}))$ is a free \mathbb{Z} -module of rank one and thus taking $\det S^{(k)}$ to be a generator of $\Lambda^{N_k} (H^0(\mathcal{X}, k\mathcal{K}_{(\mathcal{X}, \mathcal{D})}))$ eliminates the first factor in the product

Theorem 5.2. *Let $(\mathcal{X}, \mathcal{D})$ be an arithmetic log pair over $\mathcal{O}_{\mathbb{F}}$ such that $\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is a relatively ample \mathbb{Q} -line bundle over \mathcal{X} and assume that $(X_{\mathbb{Q}}, \Delta_{\mathbb{Q}})$ is klt. Then*

$$\hat{h}_{can}(\mathcal{K}_{(\mathcal{X}, \mathcal{D})}) = - \lim_{k \rightarrow \infty} \frac{1}{2N_k} \log \left(\left(\frac{\Lambda^{N_k} (H^0(\mathcal{X}, k\mathcal{K}_{(\mathcal{X}, \mathcal{D})}))}{\mathcal{O}_{\mathbb{F}}(\det S^{(k)})} \right)^{-2/k} \cdot \mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D}) \right).$$

Proof. Fix σ and write $X = X_{\sigma}$. For any fixed continuous metric $\|\cdot\|$ on $K_{(X, \Delta)}$ with positive curvature current we can express

$$\mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D})_{\sigma} = \int_{X^{N_k}} \left\| \det S^{(k)} \right\|^{2/k} dV^{\otimes N_k}.$$

where dV denotes the measure on X corresponding to the metric $\|\cdot\|$ (using the additive notation ϕ_0 for the metric $\|\cdot\|$ this means that $dV = \mu_{\phi_0}$ in the notation of Section 2.2.1). Indeed, in general, given $s \in H^0(X, kK_{(Y, \Delta)})$ and a volume form dV on Y we can, locally on Y , express

$$\|s\|^{2/k} dV := |s|^{2/k} e^{-\phi_0} \cdot \left(\frac{i}{2} \right)^{(\dim Y)^2} e^{\phi_0} dz \wedge d\bar{z} = \left(\frac{i}{2} \right)^{(\dim Y)^2} (s^{1/k} dz) \wedge (\overline{s^{1/k} dz}).$$

Next, fix a basis in $H^0(X, k(K_X + \Delta))$ which is orthonormal wrt the scalar product $\langle \cdot, \cdot \rangle$ on $H^0(X, k(K_X + \Delta))$ induced by $(\|\cdot\|, dV)$ and denote by $\det S_0^{(k)}$ the corresponding section of $K_{(X^{N_k}, \Delta_k)}$, defined as in formula 5.1. By basic linear algebra $\det S^{(k)} = \det_{i,j \leq N_k} \langle s_i^{(k)}, s_j^{(k)} \rangle \det S_0^{(k)}$. Hence,

$$\frac{1}{N_k} \log \mathcal{Z}_{N_k} = \sum_{\sigma} \frac{1}{kN_k} \log \det_{i,j \leq N_k} \langle s_i, s_j \rangle_{\sigma} + \int_{X^{N_k}} \left\| \det S_0^{(k)} \right\|^{2/k} dV^{\otimes N_k}.$$

By the arithmetic Hilbert-Samuel formula [53, 105],

$$(5.3) \quad \frac{1}{kN_k} \log \left(\prod_{\sigma} \det_{i,j \leq N_k} \langle s_i, s_j \rangle_{\sigma} \# \left(\frac{\Lambda^{N_k}(H^0(\mathcal{X}, k\mathcal{K}(\mathcal{X}, \mathcal{D})))}{\mathcal{O}_{\mathbb{F}}(\det S^{(k)})} \right)^{-2} \right) \rightarrow -2\hat{h}(\mathcal{K}(\mathcal{X}, \mathcal{D}), \|\cdot\|)$$

as $k \rightarrow \infty$. Next, by the large deviation principle in [5, Thm 1.1] for X non-singular and $\Delta = 0$ and [6, Thm 4.3], in general: for any given metric $\|\cdot\|$ on $K_{(X, \Delta)}$,

$$(5.4) \quad -\frac{1}{N_k} \log \int_{X^{N_k}} \left\| \det S_0^{(k)} \right\|^{2/k} dV^{\otimes N_k} \rightarrow \inf_{\mu \in \mathcal{P}(X)} F_1(\mu).$$

where $F_1(\mu)$ is the free energy type functional defined in formula 3.10. Hence, combining 5.3 and 5.4 gives, using the identities 3.7 and 3.11,

$$-\lim_{k \rightarrow \infty} \frac{1}{2N_k} \log \left(\left(\# \frac{\Lambda^{N_k}(H^0(\mathcal{X}, k\mathcal{K}(\mathcal{X}, \mathcal{D})))}{\mathcal{O}_{\mathbb{F}}(\det S^{(k)})} \right)^{-2/k} \cdot \mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D}) \right) = \inf_{\phi} \mathcal{M}_{\phi}(\mathcal{K}(\mathcal{X}, \mathcal{D}), \phi),$$

where the inf ranges over all finite energy metrics ϕ on $K_{(X, \Delta)(\mathbb{C})}$. Invoking the variational principle in Prop 3.3 thus concludes the proof. \square

5.1.1. *Intermezzo: the case when $\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is semi-ample and Faltings' height.* Before moving on to the log Fano case we note that $\mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D})$ is well-defined as soon as $k\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is effective, i.e. $N_k \geq 1$. In particular, if $\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is semi-ample, then $\mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D})$ is well-defined for k sufficiently divisible. For example, when $k\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is trivial $-\frac{1}{2N_k} \log \mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D})$ coincides, by definition, with Faltings' height 3.13. There is thus no need to let k tend to infinity in this case. In general, when $\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is semi-ample the proof of Theorem 5.2 reveals, together with the results described in [6, Section 5.2], that

$$-\lim_{k \rightarrow \infty} \frac{1}{2N_k} \log \left(\left(\# \frac{\Lambda^{N_k}(H^0(\mathcal{X}, k\mathcal{K}(\mathcal{X}, \mathcal{D})))}{\mathcal{O}_{\mathbb{F}}(\det S^{(k)})} \right)^{-2/k} \cdot \mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D}) \right) = h_{\phi_{\text{can}}}(\mathcal{K}_{(\mathcal{X}, \mathcal{D})}),$$

where ϕ_{can} is the volume-normalized metric on $\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ introduced in [91, 90], whose curvature form is the pull-back to X of a canonical twisted Kähler-Einstein metric on the canonical model of X over \mathbb{C} (i.e. the Proj of the canonical ring of X).

5.2. **The case** $-K_{(X,\Delta)} > 0$. Let now $(\mathcal{X}, \mathcal{D})$ be an arithmetic log pair over $\mathcal{O}_{\mathbb{F}}$ such that $-\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is a relatively ample \mathbb{Q} -line bundle over \mathcal{X} . Given a positive real number k such that $-k\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is a bona fide line bundle (i.e. Cartier) we can, after replacing k with $-k$, proceed as before. More precisely, we set

$$N_k := \dim_{\mathbb{R}} (H^0(\mathcal{X}, -k\mathcal{K}_{(\mathcal{X}, \mathcal{D})}) \otimes \mathbb{R})$$

and define

$$\mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D})_{\sigma} := \left(\frac{i}{2}\right)^{(N_k n)^2} \int_{X_{\sigma}^{N_k}} \alpha_k \wedge \overline{\alpha_k}, \quad \alpha_k := \left(\det S^{(k)}\right)^{-1/k} \otimes s_{\Delta}$$

where α_k still defines a meromorphic top form on a Zariski open subset of X^{N_k} . We then define $\mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D})$ as the product over σ of $\mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D})_{\sigma}$. However, in this case $\mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D})_{\sigma}$ may diverge (even if $\mathcal{D} = 0$).

Theorem 5.3. *Assume that $-K_{(X,\Delta)} > 0$ and that $n = 1$. Then $\mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D})$ is finite for k sufficiently large iff (X, Δ) is K -stable. Moreover,*

$$\hat{h}_{can}(-\mathcal{K}_{(\mathcal{X}, \mathcal{D})}) = \lim_{k \rightarrow \infty} \frac{1}{2N_k} \log \left(\mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D}) \left(\frac{\Lambda^{N_k}(H^0(\mathcal{X}, k\mathcal{K}_{(\mathcal{X}, \mathcal{D})}))}{\mathcal{O}_{\mathbb{F}}(\det S^{(k)})} \right)^{2/k} \right).$$

Proof. Proceeding as in the proof of Theorem 5.2, but replacing k with $-k$ yields, if $\mathcal{Z}_{N_k} < \infty$,

$$\frac{1}{N_k} \log \mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D})_{\sigma} = -\frac{1}{kN_k} \log \det_{i,j \leq N_k} \langle s_i, s_j \rangle + \int_{X_{\sigma}^{N_k}} \left\| \det S_0^{(k)} \right\|^{-2/k} dV^{\otimes N_k}$$

for any given metric $\|\cdot\|$ on $-K_{(X,\Delta)}$, where dV denotes the corresponding measure on X . Now assume that $n = 1$ and (X, Δ) is K -stable. By [9, Thm 4.1] this equivalently means that $\mathcal{Z}_{N_k} < \infty$ for k sufficiently large. Moreover, by [9, Thm 4.4],

$$-\frac{1}{N_k} \log \int_{X^{N_k}} \left\| \det S_0^{(k)} \right\|^{2/k} dV^{\otimes N_k} \rightarrow \inf_{\mu \in \mathcal{P}(X)} F_{-1}(\mu)$$

Hence, invoking the Hilbert-Samuel formula and the identities 3.7 concludes the proof, precisely as in the case $K_{(X,\Delta)} > 0$. \square

For a general relative dimension n a notion of *Gibbs stability* is introduced in [6], which - in the arithmetic present setup - amounts to the finiteness of $\mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D})$ for k sufficiently large. It is conjectured in [6, 9] that $(\mathcal{X}, \mathcal{D})$ is Gibbs stable iff (X, Δ) is K -stable (the “only if” direction is established in [49]). Moreover, under the following (a priori) stronger assumption:

$$(5.5) \quad \left(\frac{\Lambda^{N_k}(H^0(\mathcal{X}, k\mathcal{K}_{(\mathcal{X}, \mathcal{D})}))}{\mathcal{O}_{\mathbb{F}}(\det S^{(k)})} \right)^{-\pm 2/k} \cdot \mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D}) \leq C^{N_k}$$

it was pointed out in [9] that the convergence in Theorem 5.3 holds under a certain zero-free hypothesis, discussed in the following section.

Remark 5.4. It is sometimes convenient to use a different normalization, where $\alpha_k \wedge \overline{\alpha_k}$ is replaced by $\pi^{-nN} \alpha_k \wedge \overline{\alpha_k}$. These two different normalizations are analogous to the two different normalizations for Faltings' height of abelian varieties appearing in the literature ([43] vs. [37]). Then the right hand side in Theorems 5.2, 5.3 gets replaced by $\pm \hat{h}_{\text{can}}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}) + \frac{n}{2} \log \pi$. This is the height of $\pm \hat{h}_{\text{can}}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})})$ computed wrt the Kähler-Einstein metric on $\pm K_{(X, \Delta)}$ giving volume π^n to X . In all the explicit formulas that we have been able to compute (e.g. Theorem 1.1) this normalization has the effect of removing π from the explicit formulas.

5.3. Real-analyticity and the zero-free hypothesis. Consider a linear family $(\mathcal{X}, \mathcal{D}_{\mathbf{w}})$ of log pairs with coefficients $\mathbf{w} \in \mathbb{R}^m$, as defined in Section 4. This means that $\mathcal{K}_{(\mathcal{X}, \mathcal{D}_{\mathbf{w}})} \cong s(\mathbf{w})\mathcal{L}_0$, where s is an affine function of \mathbf{w} . Given a positive integer l set $k := ls^{-1}$ which is thus negative when $s < 0$. By definition, $k\mathcal{K}_{(\mathcal{X}, \mathcal{D}_{\mathbf{w}})} \cong l\mathcal{L}_0$, giving

$$H^0(\mathcal{X}, k\mathcal{K}_{(\mathcal{X}, \mathcal{D}_{\mathbf{w}})}) \cong H^0(\mathcal{X}, l\mathcal{L}_0).$$

Hence, denoting by N the dimension of $H^0(\mathcal{X}, l\mathcal{L}_0) \otimes \mathbb{R}$ and by $\det S$ the corresponding section over X^N (both depending only l) we can express

$$\mathcal{Z}_N(\mathcal{X}, \mathcal{D}_{\mathbf{w}})_{\sigma} := \left(\frac{1}{2}\right)^{(Nn)^2} \int_{X^N} \left| (\det S_{\sigma})^{s(\mathbf{w})/l} \otimes s_1^{w_1} \otimes \dots \otimes s_m^{w_m} \right|^2.$$

For a fixed positive integer l this function is manifestly real-analytic (and log convex) wrt $\mathbf{w} \in \mathbb{R}^m$ in the open region where $\mathcal{Z}_N(\mathcal{X}, \mathcal{D}) < \infty$. More precisely, allowing complex coefficients, $\mathbf{w} \in \mathbb{C}^m$, the corresponding function $\mathcal{Z}_N(\mathcal{X}, \mathcal{D}_{\mathbf{w}})$ is holomorphic in the tube domain in \mathbb{C}^m over the open subset $\{\mathcal{Z}_N(\mathcal{X}, \mathcal{D}_{\mathbf{w}}) < \infty\} \subseteq \mathbb{R}^m$. In [9] a “zero-free hypothesis” is introduced, which in the present arithmetic setup may be formulated as follows:

$$(5.6) \quad \exists \Omega \subset \mathbb{C}^m: \mathcal{Z}_N(\mathcal{X}, \mathcal{D}_{\mathbf{w}}) \neq 0 \text{ in } \Omega,$$

where Ω is assumed to be a connected open subset of \mathbb{C}^m independent of N (i.e on l) and contained in the tube-domain $\{\mathcal{Z}_N(\mathcal{X}, \mathcal{D}_{\mathbf{w}}) < \infty\}$.

Proposition 5.5. *Assume that $-\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is relatively ample and that the uniform bound 5.5 holds. If $(\mathcal{X}, \mathcal{D})$ contained in a linear family $(\mathcal{X}, \mathcal{D}_{\mathbf{w}})$ containing some log pair $(\mathcal{X}, \mathcal{D}_{\mathbf{w}_0})$ such that $\mathcal{K}_{(\mathcal{X}, \mathcal{D}_{\mathbf{w}_0})}$ is relatively ample, then the convergence in Theorem 5.3 holds under the condition that the zero-free hypothesis 5.6 holds.*

Proof. This follows from arguments in [9], which go as follows. First, using basic properties of holomorphic functions and convexity, after passing to a subsequence, the following limit holds uniformly on compact subsets of Ω , as $l \rightarrow \infty$:

$$-\frac{1}{2N_k} \log \left(\left(\frac{\Lambda^{N_k}(H^0(\mathcal{X}, k\mathcal{K}_{(\mathcal{X}, \mathcal{D}))))}{\mathcal{O}_{\mathbb{F}}(\det S^{(k)})} \right)^{-2s/l} \cdot \mathcal{Z}_N(\mathcal{X}, \mathcal{D}) \right) \rightarrow g(\mathbf{w})$$

for some holomorphic function g on Ω (indeed, by assumption, g is a uniform limit of uniformly bounded holomorphic functions on Ω). But Theorem 5.2 implies that

$g = \pm \hat{h}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D}_w)})$ on $\Omega \cap \mathbb{R}^m \cap \{s > 0\}$. Hence, by uniqueness of real-analytic extensions, it follows from Prop 4.3 that $g = \pm \hat{h}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D}_w)}) = f$ on all of $\Omega \cap \mathbb{R}^m$, which concludes the proof. \square

5.4. Synthesis on arithmetic log surfaces. When $n = 1$ combining Theorems 5.2, 5.3 yields:

Theorem 5.6. *Let $(\mathcal{X}, \mathcal{D})$ be an arithmetic log pair over $\mathcal{O}_{\mathbb{F}}$ of relative dimension one such that $\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is a relatively ample \mathbb{Q} -line bundle over \mathcal{X} and assume that $(X_{\mathbb{Q}}, \Delta_{\mathbb{Q}})$ is klt. Then*

$$\pm \hat{h}_{can}(\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}) = - \lim_{k \rightarrow \infty} \frac{1}{2N_k} \log \left(\left(\frac{\Lambda^{N_k}(H^0(\mathcal{X}, \pm k \mathcal{K}_{(\mathcal{X}, \mathcal{D})}))}{\mathcal{O}_{\mathbb{F}}(\det S^{(k)})} \right)^{-\pm 2/k} \cdot \mathcal{Z}_{N_k}(\mathcal{X}, \mathcal{D}) \right).$$

6. THE CANONICAL HEIGHT OF LOG PAIRS ON $\mathbb{P}_{\mathbb{Z}}^1$ AND THE HURWITZ ZETA FUNCTION

In this section we will, in particular, prove Theorem 1.1. Recall that \mathcal{D}^o denotes the divisor on $\mathbb{P}_{\mathbb{Z}}^1$ defined as the Zariski closure of the divisor $\Delta_{\mathbb{Q}}$ on $\mathbb{P}_{\mathbb{Q}}^1$ supported at $\{0, 1, \infty\}$ with coefficients $\mathbf{w} = (w_1, w_2, w_3)$ contained in the convex domain $C \subset \mathbb{R}^3$ defined by the weight conditions 1.5 (i.e. $(\mathbb{P}^1, \Delta_{\mathbf{w}})$ is K-semistable). Denote by $f(\mathbf{w})$ the function

$$(6.1) \quad f(\mathbf{w}) := \frac{1 - \log(\pi \frac{V}{2})}{2} - \frac{\gamma(0, \frac{V}{2}) - \sum_{i=1}^3 \gamma(w_i - \frac{V}{2}, w_i)}{V}, \quad V := -2 + \sum_{i=1}^3 w_i$$

defined in the interior of C when $V > 0$, where $\gamma(a, b)$ is defined by formula 1.6 (note that V is the degree of $K_{(\mathbb{P}^1, \Delta)}$.)

Lemma 6.1. *Given $a, b \in]0, 1[$,*

$$\gamma(a, b) = \int_a^b \log l(x) dx, \quad l(x) := \frac{\Gamma(x)}{\Gamma(1-x)}, \quad \Gamma(x) := \int_0^{\infty} t^{x-1} e^{-t} dt.$$

Proof. The formula follows directly from the well-known fact that $\zeta(-1, t) + \zeta'(-1, t) + \frac{(t-1)}{2} \log(2\pi)$ is a primitive of $\log(\Gamma(t))$ on $]0, 1[$ [32, formula 3.11]. \square

The previous lemma reveals that f is real-analytic when $V > 0$. Furthermore, Theorem 5.6 will imply that f extends real-analytically to all of the interior of C . We extend f to a finite function on the subset of the boundary of C where $V \neq 0$, by declaring its value to be the limit of its values along any affine segment I in the interior of C reaching the boundary.

6.1. **The case** $K_{(\mathbb{P}^1, \Delta)} > 0$. It is enough to consider case when Δ is klt, i.e. $w_i < 1$, by the continuity in Prop 4.1. Since

$$k\mathcal{K}_{(\mathbb{P}^1_{\mathbb{Z}}, \mathcal{D}^o)} \simeq kV\mathcal{O}(1), \quad V := V(K_{(\mathbb{P}^1, \Delta)})$$

the free \mathbb{Z} -module $H^0(\mathcal{X}, k\mathcal{K}_{(\mathbb{P}^1_{\mathbb{Z}}, \mathcal{D}^o)})$ may be identified with the space of all homogeneous polynomials of degree kV on \mathbb{C}^2 with integer coefficients. We fix the standard basis s_1, \dots, s_{N_k} of monomials in the latter free \mathbb{Z} -module and denote by $\det S^{(k)}$ the corresponding generator of $\Lambda^{N_k}(H^0(\mathcal{X}, k\mathcal{K}_{(\mathcal{X}, \mathcal{D})}))$, as in Remark 5.1. Denote by p_1, \dots, p_m the irreducible components of Δ and assume that $z = \infty$ at p_m , where z denotes the affine coordinate on the standard affine piece \mathbb{C} of $\mathbb{P}^1_{\mathbb{C}}$. Identifying $\det S^{(k)}$ with the Vandermonde determinant $\prod_{i < j \leq N_k} (z_i - z_j)$ (where $N_k = kV + 1$) we can thus express

$$(6.2) \quad \mathcal{Z}_N = \int_{\mathbb{C}^N} \left(\prod_{i \neq j} |z_i - z_j| \right)^{\frac{V}{N-1}} \prod_{i \leq N, j \leq m-1} |z_i - p_j|^{-2w_i} \prod_i \frac{i}{2} dz_i \wedge d\bar{z}_i,$$

where we have, for simplicity, dropped the subindex k in the notation N_k ([9, Lemma 4.3]). For $m = 3$ and $(p_1, p_2) = (0, 1)$ the integral appearing in the right hand side of the previous formula is known as the Dotsenko-Fateev integral (and can be viewed as a Selberg integral over the field \mathbb{C} [48]). By [41, Formula B.9] (and [48, formula 3.1]) it may be explicitly computed in terms of the function $l(x)$ appearing in Lemma 6.1:

$$(6.3) \quad \mathcal{Z}_N = N! \left(\frac{\pi}{l(\frac{1}{2} \frac{V}{N-1})} \right)^N \prod_{j=0}^{N-1} \frac{l(\frac{j+1}{2} \frac{V}{N-1})}{l(w_1 - \frac{j}{2} \frac{V}{N-1}) l(w_2 - \frac{j}{2} \frac{V}{N-1}) l(w_3 - \frac{j}{2} \frac{V}{N-1})},$$

Hence, by Theorem 5.6,

$$(6.4) \quad \begin{aligned} -\hat{h}_{\text{can}}(\mathcal{K}_{(\mathcal{X}, \Delta)}) &= \lim_{N \rightarrow \infty} \frac{1}{2N} \log \mathcal{Z}_N \\ &= \lim_{N \rightarrow \infty} \frac{1}{2N} \left(\log N! + N \log(\pi) - N \log(l(\frac{1}{2} \frac{V}{N-1})) \right) + \\ &\quad \frac{1}{V} \sum_{j=0}^{N-1} \log(l(\frac{j+1}{2} \frac{V}{N-1})) \frac{V}{2N} - \frac{1}{V} \sum_{k=1}^3 \sum_{j=0}^{N-1} \log l(w_k - \frac{j}{2} \frac{V}{N-1}) \frac{V}{2N} \end{aligned}$$

Using Stirling's approximation and the fact that the gamma function has a simple pole with residue 1 at 0 gives

$$\begin{aligned} &\frac{1}{2N} (\log N! - N \log(l(\frac{1}{2} \frac{V}{N-1}))) = \\ &\frac{1}{2} \left(\log N - 1 - \log(\Gamma(\frac{V}{2} \frac{1}{N-1})) + \log(\Gamma(1 - \frac{V}{2} \frac{1}{N-1})) \right) + O(N^{-1} \log N) \\ &\frac{1}{2} \left(\log N - 1 - \log(\frac{2}{V}(N-1) + O(1)) + \log(\Gamma(1 - \frac{V}{2} \frac{1}{N-1})) \right) + O(N^{-1} \log(N)) \\ &\quad \rightarrow_{N \rightarrow \infty} \frac{1}{2} (\log \frac{V}{2} - 1). \end{aligned}$$

All in all, recognizing the sums over j in 6.4 as either right or left Riemann sums, this proves $\hat{h}_{\text{can}}(\mathcal{K}_{(\mathbb{P}^1_{\mathbb{Z}}, \mathcal{D}^o)}) = f(\mathbf{w})$.

6.2. The case $-K_{(\mathbb{P}^1, \Delta)} > 0$. It is enough to consider the case when (\mathbb{P}^1, Δ) is K -stable (i.e. the case when \mathbf{w} is contained in the interior of C), by the continuity in Prop 4.1. Since $|V|$ is the volume (degree) of $-K_{(\mathbb{P}^1, \Delta)}$,

$$-k\mathcal{K}_{(\mathbb{P}^1_{\mathbb{Z}}, \mathcal{D}^o)} \simeq k|V|\mathcal{O}(1), \quad |V| = 2 - \sum_{i=1}^m w_i.$$

Using that $|V| = -V$ formula 6.3 now yields

$$\mathcal{Z}_N = N! \left(\frac{\pi}{-l(-\frac{1}{2}\frac{|V|}{N-1})} \right)^N \prod_{j=0}^{N-1} \frac{-l(-\frac{(j+1)}{2}\frac{|V|}{N-1})}{l(w_1 + \frac{j}{2}\frac{|V|}{N-1})l(w_2 + \frac{j}{2}\frac{|V|}{N-1})l(w_3 + \frac{j}{2}\frac{|V|}{N-1})}.$$

Hence, proceeding precisely as before, gives

$$\frac{1}{2}h_{\text{can}}(-\mathcal{K}_{(\mathbb{P}^1_{\mathbb{Z}}, \mathcal{D}^o)}) = \frac{|V|}{2}(\log \frac{|V|}{2} + \log \pi - 1) + \int_{-\frac{|V|}{2}}^0 \log(-l(x))dx - \sum_{k=1}^3 \int_{w_k}^{w_k + \frac{|V|}{2}} \log l(x)dx.$$

Finally, exploiting that $\Gamma(x+1) = x\Gamma(x)$ the integral over $[-|V|/2, 0]$ may be rewritten as

$$- \int_0^{\frac{|V|}{2}} (\log l(x) - 2 \log(x)) dx = - \int_0^{\frac{|V|}{2}} \log l(x)dx - |V| \log(\frac{|V|}{2}) + |V|$$

so that in total

$$\hat{h}_{\text{can}}(-\mathcal{K}_{(\mathbb{P}^1_{\mathbb{Z}}, \mathcal{D}^o)}) = \frac{1}{2}(-\log \frac{|V|}{2} + \log \pi + 1) - \frac{1}{|V|} \int_0^{\frac{|V|}{2}} \log(l(x))dx - \sum_{k=1}^3 \frac{1}{|V|} \int_{w_k}^{w_k + \frac{|V|}{2}} \log l(x)dx.$$

6.2.1. Real-analyticity. Let us give three different proofs that $\pm \hat{h}_{\text{can}}(\pm \mathcal{K}_{(\mathbb{P}^1_{\mathbb{Z}}, \mathcal{D}^o)})$ is real-analytic in the interior of C . First, this is a special case of Prop 4.3. Secondly, let us show directly from Theorem 1.1 that $\pm \hat{h}_{\text{can}}(\pm \mathcal{K}_{(\mathbb{P}^1_{\mathbb{Z}}, \mathcal{D}^o)})$ is real-analytic. By Theorem 1.1

$$\pm \hat{h}_{\text{can}}(\pm \mathcal{K}_{(\mathbb{P}^1_{\mathbb{Z}}, \mathcal{D}^o)}) = \frac{1}{2}(-\log(\pi) - \int_0^1 \log(\frac{Vt}{2}l(\frac{Vt}{2}))dt + \sum_{k=1}^3 \int_0^1 \log l(w_k - \frac{Vt}{2})dt).$$

Recalling that the gamma function has a simple pole at 0, so that $\Gamma(x)x$ is real-analytic and positive on $(-1, \infty)$, it is not to hard to see that the expression above is a real-analytic function of the weights w_i in the interior of C . The third proof of the real-analyticity exploits that the assumptions in Prop 5.5 are satisfied in this case, since \mathcal{Z}_N is a product of Gamma-functions (see the end of 4.3). As a consequence, $\pm \hat{h}_{\text{can}}(\pm \mathcal{K}_{(\mathbb{P}^1_{\mathbb{Z}}, \mathcal{D}^o)})$ is the restriction of a uniform limit of holomorphic functions on Ω and thus real-analytic.

6.3. A case with more than three points. Assume given $(w_0, w_1, w_\infty) \in]0, 1[^3$. Consider the divisor $\tilde{\Delta}$ on \mathbb{P}^1 supported at the points $\{0, 1, -1, \infty\}$ with weights $(w_0, w_1, w_{-1}, w_\infty)$. Assume that $K_{(\mathbb{P}^1, \tilde{\Delta})} > 0$. Denote by Δ the divisor on \mathbb{P}^1 supported at the points $\{0, 1, \infty\}$ with weights $(1 + \frac{w_0-1}{2}, w_1, 1 + \frac{w_\infty-1}{2})$ (in particular, if Δ has ramification m at p_0 , then $\tilde{\Delta}$ has ramification $2m$). Denote by $\tilde{\mathcal{D}}$ and \mathcal{D} the Zariski closures in $\mathbb{P}_{\mathbb{Z}}^1$ of $\tilde{\Delta}$ and Δ , respectively.

Proposition 6.2. *The following formula holds*

$$\hat{h}_{can}(\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \tilde{\mathcal{D}})}) = \hat{h}_{can}(\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D})}) + \frac{1}{2} \log 2 = f\left(1 + \frac{w_0-1}{2}, w_1, 1 + \frac{w_\infty-1}{2}\right) + \frac{1}{2} \log 2$$

a

Proof. Using the standard isomorphism $\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D})} \simeq V(K_{(\mathbb{P}^1, \Delta)})\mathcal{O}(1)$ over \mathbb{Z} we can identify a given metric ϕ on $K_{(\mathbb{P}^1, \Delta)}$ with a metric on $(K_{(\mathbb{P}^1, \Delta)})\mathcal{O}(1)$. Consider the standard map $F : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ of degree 2, which in the standard affine coordinate is given by $y = x^2$. We will use the same notation F for its standard lift satisfying $F^*\mathcal{O}(1) \simeq 2\mathcal{O}(1)$. Then $F^*\phi$ defines a metric on $2V(K_{(\mathbb{P}^1, \Delta)})\mathcal{O}(1)$ which (as before) may be identified with a metric on $\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \tilde{\mathcal{D}})}$ (using that $2V(K_{(\mathbb{P}^1, \Delta)}) = V(K_{(\mathbb{P}^1, \tilde{\Delta})})$). By basic functoriality of normalized heights $\hat{h}(\mathcal{O}(1), \psi) = \hat{h}(F^*\mathcal{O}(1), F^*\psi)$ for any metric ψ on $\mathcal{O}(1)$. In particular, $\hat{h}(V\mathcal{O}(1), \phi) = \hat{h}(F^*(V\mathcal{O}(1)), F^*\phi)$. Hence, it will be enough to show that

$$\int_{\mathbb{P}^1} \mu_{F^*\phi} = 2^{-1} \int_{\mathbb{P}^1} \mu_\phi.$$

To this end first observe that

$$dx = 2^{-1}y^{-1/2}dy, \quad (y-1)(y+1) = (x-1).$$

Hence,

$$\mu_{F^*\phi} = 2^{-2}F^*\mu_\phi.$$

Since the map F has degree 2 it follows that

$$\int_{\mathbb{P}^1} \mu_{F^*\phi} = \int_{\mathbb{P}^1} 2^{-2}F^*\mu_\phi = 2 \int_{\mathbb{P}^1} 2^{-2}\mu_\phi = 2^{-1} \int_{\mathbb{P}^1} \mu_\phi,$$

as desired. \square

Remark 6.3. A similar formula holds when the two points $\{1, 1\}$ in the support of $\tilde{\Delta}$ are replaced by the d points defined as the d roots of unity (by taking the map F in the proof above to be defined by $y = x^d$).

7. SHARP BOUNDS ON $\mathbb{P}_{\mathbb{Z}}^1$

In this section we will, in particular, prove Theorem 1.2. We continue with the notations from Section 6. But we start with the following refinement of the conjectural Fujita type inequality 1.8 in the present case:

Theorem 7.1. *Let $(\mathcal{X}, \mathcal{D}; \mathcal{L})$ be a polarized arithmetic log surface $(\mathcal{X}, \mathcal{D}; \mathcal{L})$ over \mathbb{Z} with \mathcal{X} normal such that the complexification X of \mathcal{X} equals \mathbb{P}^1 and the complexification L of \mathcal{L} equals either $K_{(\mathbb{P}^1, \Delta)}$ or $-K_{(\mathbb{P}^1, \Delta)}$. Assume that Δ is supported on at most three points and that (X, Δ) is K -semistable (i.e. the weight conditions 1.5 hold). Then*

$$\hat{\mathcal{M}}_{(\mathcal{X}, \mathcal{D})}(\overline{\mathcal{L}}) \geq f(\mathbf{w})$$

and if (X, Δ) is K -stable, then equality holds iff $(\mathcal{X}, \mathcal{D}) = (\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)$ and $\mathcal{L} = \pm \mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)}$. As a consequence, if $\pm \mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is relatively ample, then

$$\pm \hat{h}_{can}(\overline{\pm \mathcal{K}}_{(\mathcal{X}, \mathcal{D})}) \geq f(\mathbf{w})$$

and if $-\mathcal{K}_{(\mathcal{X}, \mathcal{D})}$ is relatively ample, then

$$-\hat{h}(\overline{-\mathcal{K}}_{(\mathcal{X}, \mathcal{D})}) \geq f(\mathbf{w})$$

for any volume-normalized continuous metric on $-K_{(\mathbb{P}^1, \Delta)}$.

Proof. The first inequality follows directly from combining the variational principles for metrics and models in Prop 3.3 and Prop 3.14, respectively, with Theorem 1.1. In particular, taking the inf over all psh metrics of finite energy yields the second inequality. The third inequality then follows from Lemma 3.4, since $n = 1$. \square

In fact, the inequalities in the previous theorem hold more generally when \mathbb{Z} is replaced by $\mathcal{O}_{\mathbb{F}}$ (using Prop 3.14 and the fact that the normalized height is invariant under base-change). In particular, the first inequality yields the following explicit expression for the normalized modular invariant, defined in Section 2.5.1,

$$(7.1) \quad \hat{\mathcal{M}}(\mathbb{P}_{\mathbb{F}}^1, \Delta_{\mathbb{F}}; \pm K_{(\mathbb{P}_{\mathbb{Q}}^1, \Delta_{\mathbb{Q}})}) = f(\mathbf{w}),$$

when $\Delta_{\mathbb{F}}$ is supported on three points in $\mathbb{P}_{\mathbb{F}}^1$.

7.1. Proof of Theorem 1.2. We first establish the following refinement of the first inequality in Theorem 1.2:

Theorem 7.2. *The following inequality holds*

$$\hat{h}(-\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)}) \geq \frac{1}{2}(1 + \log \pi) + \frac{1}{4}(1 + \log \frac{3}{4}) \sum_{k=1}^3 w_k$$

In particular, $\hat{h}(-\mathcal{K}_{(\mathcal{X}, \Delta)}) \geq \hat{h}(-\mathcal{K}_{\mathbb{P}_{\mathbb{Z}}^1}) > 0$.

Proof. Set $g(\mathbf{w}) = -\pm \hat{h}(\pm \mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)})$ and first consider the case when $-K_{(\mathbb{P}^1, \Delta)} > 0$, so that $|V|$ is the volume (degree) of $-K_{(\mathbb{P}^1, \Delta)}$. In this case, $g = \hat{h}(\pm \mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)})$ and, using

Theorem 1.1,

$$\begin{aligned}\frac{\partial g}{\partial w_i} &= \frac{1}{2V} - \frac{1}{|V|} \int_0^{\frac{|V|}{2}} \log l(x) dx - \sum_{k=1}^3 \frac{1}{|V|} \int_{w_k}^{w_k + \frac{|V|}{2}} \log l(x) dx \\ &+ \frac{1}{2|V|} \log l\left(\frac{|V|}{2}\right) - \frac{1}{2|V|} \log l\left(w_i + \frac{|V|}{2}\right) + \frac{1}{|V|} \log l(w_i) \\ &+ \sum_{k=1, k \neq i}^3 \frac{1}{2|V|} \log l\left(w_k + \frac{|V|}{2}\right).\end{aligned}$$

Next we compute the limit of the gradient at zero along the curve $w(t) = (t, t, t)$, i.e.

$$\lim_{t \rightarrow 0} \frac{d}{dt} g(w(t)) = 3 \lim_{t \rightarrow 0} \frac{d}{dw_i} g(w)|_{w_t} = \frac{3}{4} \left(1 + \lim_{t \rightarrow 0} \log\left(l\left(1 - \frac{3t}{2}\right) l(t)^2 l\left(1 - \frac{t}{2}\right)\right)\right) = \frac{3}{4} \left(1 + \log \frac{3}{4}\right).$$

By Prop 4.1, $g(w)$ is, in general, convex. Hence, along the curve $w(t)$ we have $g(w(t)) \geq g(0) + t \lim_{t \rightarrow 0} \frac{d}{dt} g(w(t))$. Furthermore, as g is symmetric in the weights and convex, we have $g(w) \geq g(w(t))$ for any w where $\sum_{k=1}^3 w_k = \sum_{k=1}^3 w_k(t) = 3t$. Putting it all together we have shown,

$$g(w) \geq \frac{1}{2} (1 + \log \pi) + \frac{1}{4} \left(1 + \log \frac{3}{4}\right) \sum_{k=1}^3 w_k,$$

as desired. \square

Finally, we establish the following refinement of the second inequality in Theorem 1.2:

Theorem 7.3. *The following inequality holds when $K_{(\mathbb{P}^1, \Delta)}$ is semi-ample:*

$$\hat{h}(\mathcal{K}_{(\mathbb{P}^1, \mathcal{D}^\circ)}) \leq -\frac{1}{2} \log(\pi) + \frac{3}{2} \log \frac{\Gamma(\frac{2}{3})}{\Gamma(\frac{1}{3})} + \frac{3}{4} \left(\gamma + \frac{1}{2} \left(\frac{\Gamma'(2/3)}{\Gamma(2/3)} + \frac{\Gamma'(1/3)}{\Gamma(2/3)}\right)\right) \left(\sum_{k=1}^3 w - 2\right)$$

equality holds for the weights $(2/3, 2/3, 2/3)$. In particular, $\hat{h}(\mathcal{K}_{(\mathbb{P}^1, \mathcal{D}^\circ)}) < 0$.

Proof. We have for the gradient of $\hat{h} := \hat{h}(\mathcal{K}_{(\mathcal{X}, \Delta)})$, with V denoting the degree of $K_{(\mathbb{P}^1, \Delta)}$, which, by assumption is non-negative,

$$\begin{aligned}\frac{d}{dw_i} \hat{h}(w) &= -\frac{1}{2V} + \frac{1}{V^2} \int_0^{\frac{V}{2}} \log l(x) dx - \frac{1}{V^2} \sum_{k=1}^3 \int_{w_k - \frac{V}{2}}^{w_k} \log l(x) dx \\ &- \frac{1}{2V} \log l\left(\frac{V}{2}\right) + \frac{1}{V} \log l(w_i) - \frac{1}{2V} \log l\left(w_i - \frac{V}{2}\right) \\ &+ \sum_{k=1, k \neq i}^3 \frac{1}{2V} \log l\left(w_k - \frac{V}{2}\right).\end{aligned}$$

We will compute the limit of the gradient as $t \rightarrow \frac{2}{3}_+$, along the curve $w(t) := (t, t, t)$. We begin by making a few preparatory calculations. First, as $V \rightarrow 0$,

$$\begin{aligned} \frac{1}{V^2} \int_0^{\frac{V}{2}} \log l(x) dx &= \frac{1}{V^2} \int_0^{\frac{V}{2}} \log\left(\frac{1}{x} - \gamma + \mathcal{O}(x)\right) + \gamma x + \mathcal{O}(x^2) dx \\ &= \frac{1}{V^2} \int_0^{\frac{V}{2}} -\log(x) - 2\gamma x + \mathcal{O}(x^2) dx = -\frac{1}{2V} \log \frac{V}{2} + \frac{1}{2V} - \frac{\gamma}{4} \end{aligned}$$

where we have used the Laurent series of Γ around 0 and $\log \Gamma$ around 1. Next, as $t \rightarrow \frac{2}{3}$,

$$\begin{aligned} -\frac{3}{(3t-2)^2} \int_{t-(3t-2)/2}^t \log l(x) dx &= -\frac{3}{(3t-2)^2} \int_{t-(3t-2)/2}^t \log l\left(\frac{2}{3}\right) + (\log l)'\left(\frac{2}{3}\right)\left(x - \frac{2}{3}\right) + \mathcal{O}\left(\left(x - \frac{2}{3}\right)^2\right) dx \\ &= -\frac{3}{2(3t-2)} \log l\left(\frac{2}{3}\right) - \frac{1}{8} (\log l)'\left(\frac{2}{3}\right) + \mathcal{O}\left(t - \frac{2}{3}\right) \end{aligned}$$

and

$$\begin{aligned} -\frac{1}{2V} \log l\left(\frac{V}{2}\right) &= -\frac{1}{2V} \log\left(\frac{2}{V} - \gamma + \mathcal{O}(V)\right) + \frac{\gamma}{4} + \mathcal{O}(V) \\ &= \frac{1}{2V} \log \frac{V}{2} + \frac{\gamma}{2} + \mathcal{O}(V) \end{aligned}$$

Thus, in total

$$\begin{aligned} \lim_{t \rightarrow \frac{2}{3}_+} \frac{d}{dt} \hat{h}(w(t)) &= 3 \lim_{t \rightarrow \frac{2}{3}_+} \frac{d}{dw_i} \hat{h}(w)|_{w_t} \\ &= 3 \lim_{t \rightarrow \frac{2}{3}_+} \frac{\gamma}{4} - \frac{3}{2(3t-2)} \log l\left(\frac{2}{3}\right) - \frac{1}{8} (\log l)'\left(\frac{2}{3}\right) \\ &\quad + \frac{1}{3t-2} \log l(t) + \frac{1}{2(3t-2)} \log l\left(t - \frac{3t-2}{2}\right) \\ &= 3 \lim_{s \rightarrow 0_+} -\frac{3}{2s} \log l\left(\frac{2}{3}\right) - \frac{1}{8} (\log l)'\left(\frac{2}{3}\right) \\ &\quad + \frac{1}{s} \log l\left(\frac{s}{3} + \frac{2}{3}\right) + \frac{1}{2s} \log l\left(-\frac{s}{6} + \frac{2}{3}\right) \\ &= \frac{3\gamma}{4} + \frac{3}{8} (\log l)'\left(\frac{2}{3}\right) \\ &= \frac{3}{4} \left(\gamma + \frac{1}{2} \left(\frac{\Gamma'(2/3)}{\Gamma(2/3)} + \frac{\Gamma'(1/3)}{\Gamma(2/3)} \right) \right). \end{aligned}$$

We also want to evaluate the height (or rather take the limit) for the weights $(2/3, 2/3, 2/3)$, which is an easy variation of the above calculation,

$$\lim_{t \rightarrow \frac{2}{3}_+} \hat{h}(w(t)) = -\frac{1}{2} \log(\pi) + \frac{3}{2} \log l\left(\frac{2}{3}\right)$$

By a similar argument as in Theorem 7.2, using the concavity of the height, we have thus shown

$$\hat{h} \leq -\frac{1}{2} \log(\pi) + \frac{3}{2} \log \frac{\Gamma(\frac{2}{3})}{\Gamma(\frac{1}{3})} + \frac{3}{4} \left(\gamma + \frac{1}{2} \left(\frac{\Gamma'(2/3)}{\Gamma(2/3)} + \frac{\Gamma'(1/3)}{\Gamma(2/3)} \right) \right) \left(\sum_{k=1}^3 w - 2 \right).$$

□

8. SPECIFIC VALUES OF CANONICAL HEIGHTS

In this section we continue with the case when $\mathcal{X} = \mathbb{P}_{\mathbb{Z}}^1$ and \mathcal{D}^o is the Zariski closure of the divisor Δ on $\mathbb{P}_{\mathbb{Q}}^1$ supported on the three points $\{0, 1, \infty\}$. We consider only the “orbifold/cusp case” where the coefficients w_i of Δ are of the form $w_i = 1 - 1/m_i$ for $m_i \in \mathbb{N} \cup \{\infty\}$, where m_i are called *ramification indices*. The formulas in Table 1 (Section 1.2.4) are obtained by simplifying the explicit expression $f(\mathbf{w})$ appearing in Theorem 1.1. Moreover, we also compute the canonical height in some log Fano cases. We will provide the complete calculation only for the simplest cases. The calculations in the remaining cases are similar, but since they are somewhat lengthy they are merely outlined (in order to avoid computational mistakes, we have numerically verified the end results to machine precision, using standard implementations of the expression $f(\mathbf{w})$).

8.1. The case when $K_{(\mathbb{P}^1, \Delta)} > 0$. Set

$$\hat{h} := \hat{h}_{\text{can}}(\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)}) + \log \frac{\pi V}{2} = \frac{1}{2} - \frac{\gamma(0, \frac{V}{2}) - \sum_{i=1}^3 \gamma(w_i - \frac{V}{2}, w_i)}{V},$$

using, in the second equality, Theorem 1.1. In the application to Shimura curves, considered in Section 9, \hat{h} is the normalized height of $\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D})}$ with respect to the Petersson metric.

Proposition 8.1. *For the ramification indices $(2, 3, \infty)$*

$$\hat{h} = -\frac{\zeta'(-1)}{\zeta(-1)} - \frac{1}{2} - \frac{1}{4} \log(12).$$

Proof. Denote as before $F(x) = \zeta(-1, x) + \zeta'(-1, x)$. By Theorem 1.1 and 1.6 we have (using also that $F(0) = F(1)$ interpreted correctly)

$$\begin{aligned} \hat{h} &= \frac{1}{2} + \frac{1}{V} (-F(1/12) - F(11/12) + F(0) + F(12/12) \\ &\quad + F(6/12) + F(6/12) - F(5/12) - F(7/12) \\ &\quad + F(8/12) + F(4/12) - F(7/12) - F(5/12) \\ &\quad + F(12/12) + F(0) - F(11/12) - F(1/12)) \\ &= 1/2 + \frac{1}{V} (-2F(1/12) + F(4/12) - 2F(5/12) + 2F(6/12) \\ &\quad - 2F(7/12) + F(8/12) - 2F(11/12) + F(12/12)) \end{aligned}$$

To relate the linear combination of Hurwitz zeta functions to the Riemann zeta function, we use the *multiplication theorem*

$$(8.1) \quad k^s \zeta(s) = \sum_{i=1}^k \zeta(s, i/k)$$

repeatedly for appropriate values of k and end up with

$$\begin{aligned} & -2\zeta(s, 1/12) + \zeta(s, 4/12) - 2\zeta(s, 5/12) + 2\zeta(6/12) \\ & -2\zeta(s, 7/12) + \zeta(s, 8/12) - 2\zeta(s, 11/12) + \zeta(s, 12/12) \\ & = (-2 \cdot 12^s + 2 \cdot 6^s + 2 \cdot 4^s + 3^s + 1)\zeta(s) \end{aligned}$$

Using this together with the definition 1.6 of F leads to

$$\begin{aligned} \hat{h} &= \frac{1}{2} + \frac{1}{V}((-2 \cdot 12^{-1} + 2 \cdot 6^{-1} + 2 \cdot 4^{-1} + 3^{-1} + 1)\zeta(-1) \\ & \quad + (-2 \log(12)12^{-1} + 2 \log(6)6^{-1} + 2 \log(4)4^{-1} + \log(3)3^{-1})\zeta(-1) \\ & \quad + (-2 \cdot 12^{-1} + 2 \cdot 6^{-1} + 2 \cdot 4^{-1} + 3^{-1} + 1)\zeta'(-1) \\ &= \frac{1}{2} + \frac{1}{V}\left(-\frac{2}{12} - \frac{\log(12)}{24} + 2\zeta'(-1)\right) \\ &= -\frac{\zeta'(-1)}{\zeta(-1)} - \frac{1}{2} - \frac{1}{4}\log(12). \end{aligned}$$

□

For the ramification indices $(6, 2, 6)$ the calculation is similar to the previous case. Next, for the case of ramification indices $(4, 4, 4)$ the calculation also proceeds in a similar way, but now using that

$$\zeta_{\mathbb{Q}(\sqrt{2})}(s) = \zeta(s)8^{-s}\left(\zeta\left(s, \frac{1}{8}\right) - \zeta\left(s, \frac{3}{8}\right) - \zeta\left(s, \frac{5}{8}\right) + \zeta\left(s, \frac{7}{8}\right)\right),$$

by the standard factorization formula for $\zeta_{\mathbb{F}}(s)$, when \mathbb{F} is an abelian Galois extension of \mathbb{Q} . From the explicit formula in Theorem 1.1 for \hat{h} we get a linear combination of Hurwitz zeta functions rather than a linear combinations of products of Hurwitz zeta functions as in the equation above. But, after differentiating and evaluating at -1 , we can still use the multiplication theorem for the Hurwitz zeta function a number of times on the derivative terms, while evaluating the rest in terms of explicit rational numbers. This uses the well known relation between values of the Hurwitz zeta function at -1 and the second Bernoulli polynomial. Indeed,

$$\zeta(-1, a) = -\frac{B_2(a)}{2}.$$

Finally, for the rest of the cases, i.e. $(3, 3, 6)$, $(5, 5, 5)$, $(6, 6, 6)$, $(2, 4, 12)$, $(7, 7, 7)$ and $(9, 9, 9)$, the strategy is the same as for the case of $(4, 4, 4)$, noting that the respective number fields are all abelian Galois extensions.

8.2. **The case when $-K_{(\mathbb{P}^1, \Delta)} > 0$.** In this section we will use the shorthand $\hat{h}_{\text{can}} := \hat{h}_{\text{can}}(-\mathcal{K}_{(\mathbb{P}^1_{\mathbb{Z}}, \mathcal{D})})$. We start by verifying that Theorem 1.1 recovers some simple cases of Fano orbifolds, where the canonical height has previously been computed.

8.2.1. *The case when Δ is supported on two points.* In this case we may, by symmetry, assume that $w_2 = 0$. In this case the K-semistability assumption implies that $w_1 = w_3$. By [3, Lemma 3.2],

$$\hat{h}_{\text{can}} = \frac{1}{2}(1 + \log \pi - \log \frac{V}{2})$$

(when all weights vanish this specializes to the well-known formula for the canonical height of $\mathbb{P}^1_{\mathbb{Z}}$). In order for the previous formula to be consistent with the previous theorem it must be that

$$\gamma(0, \frac{V}{2}) + \gamma(1 - \frac{V}{2}, 1) = 0.$$

Let us give a direct proof of this vanishing, using a symmetry argument. Setting $\lambda = V/2$ and $g(t) = \log \Gamma(x)$ the left hand side in the previous formula may be expressed as the integral over $[-\lambda/2, \lambda/2]$ of the function

$$(g(x + \lambda/2) - g(-x + \lambda/2)) + (g(x + 1 - \lambda/2) - g(-x + 1 - \lambda/2)),$$

which is odd (since both terms are). Hence, the integral over $[-\lambda/2, \lambda/2]$ indeed vanishes.

8.2.2. *The case of ramification indices (2, 2, 2) and the Fermat curve \mathcal{X}_2 of degree two.* Let us next show that when all weights w_i equal 1/2

$$(8.2) \quad \hat{h}_{\text{can}} = \frac{1}{2}(1 + \log \pi) + \frac{1}{2} \log 2$$

by expressing $\hat{h}_{\text{can}}(\mathbb{P}^1_{\mathbb{Z}}, \mathcal{D})$ in terms of the canonical height of the Zariski closure \mathcal{X}_2 in $\mathbb{P}^2_{\mathbb{Z}}$ of the Fermat curve of degree two:

$$\hat{h}_{\text{can}} = \hat{h}_{\text{can}}(-\mathcal{K}_{\mathcal{X}_2}) + \log 2$$

(as follows from realizing $(\mathbb{P}^1_{\mathbb{Z}}, \mathcal{D})$ as a Galois cover of \mathcal{X}_2 as in [3, Section 5.2]). Formula 8.2 thus follows from

$$2\hat{h}_{\text{can}}(-\mathcal{K}_{\mathcal{X}_2}) = 1 + \log \pi - \log 2,$$

which can be deduced from the height formula for quadrics in [30] (or by noting that \mathcal{X}_2 is the blow-up of $\mathbb{P}^1_{\mathbb{Z}}$ of a closed point on the fiber over the prime (2)). Since, in this case, $V = 1/2$, formula 8.2 is, by Theorem 1.1, equivalent to the identity

$$\gamma(0, \frac{1}{4}) + 3\gamma(\frac{1}{2}, \frac{3}{4}) = \frac{1}{4} \log 2.$$

This identity can, indeed, be verified using the multiplication formula 8.1. Indeed, applying this formula for $k = 2, 4$ one easily finds the above relation.

(m_1, m_2, m_3)	$\hat{h}_{\text{can}} - \frac{1}{2}(1 + \log \pi)$
$(2, 2, 3)$	$-\frac{1}{6} \log 2 + \frac{2}{3} \log 3$
$(2, 2, 4)$	$\frac{3}{4} \log 2$
$(2, 3, 3)$	$\frac{1}{2} \log 2 + \frac{1}{8} \log 3$
$(2, 3, 4)$	$\frac{7}{12} \log 2 + \frac{1}{8} \log 3$

TABLE 2. Ramification indices and the corresponding normalized height for some log Fano orbifolds.

8.2.3. *New cases.* We next consider some cases where the canonical height has not been computed before:

In general, any K-stable Fano orbifold curve can have ramification indices (m_1, m_2, m_3) from either the infinite list $(2, 2, r), r \geq 2$ or the exceptional list $(2, 3, 3), (2, 3, 4), (2, 3, 5)$.

Proposition 8.2. *For the ramification indices $(2, 3, 3)$*

$$\hat{h}_{\text{can}} = \frac{1}{2} + \frac{1}{2} \log \pi + \frac{1}{8} \log 48.$$

Proof. Denote as before $F(x) = \zeta(-1, x) + \zeta'(-1, x)$. Then

$$\begin{aligned} \hat{h}_{\text{can}} = & \frac{1}{2} \left(-\log \frac{V}{2} + \log \pi + 1 \right) - \frac{1}{V} \left(\right. \\ & F(1/12) + F(11/12) - F(0) - F(1) \\ & + F(7/12) + F(5/12) - 2F(6/12) \\ & \left. + 2F(9/12) + 2F(3/12) - 2F(8/12) - 2F(4/12) \right). \end{aligned}$$

By repeatedly using the multiplication theorem 8.1 we find the following identity for the relevant linear combination of Hurwitz zeta functions.

$$\begin{aligned} & -\zeta(s, 1/12) - 2\zeta(s, 3/12) + 2\zeta(s, 4/12) \\ & -\zeta(s, 5/12) + 2\zeta(s, 6/12) - \zeta(s, 7/12) \\ & + 2\zeta(s, 8/12) - 2\zeta(s, 9/12) - \zeta(s, 11/12) \\ & = (-12^s + 6^s - 4^s + 2 \cdot 3^s + 3 \cdot 2^s - 2)\zeta(s). \end{aligned}$$

This allows us to compute \hat{h} , and quite remarkably, since

$$-12^{-1} + 6^{-1} - 4^{-1} + 2 \cdot 3^{-1} + 3 \cdot 2^{-1} - 2 = 0,$$

there is no $\zeta'(-1)$ appearing in the expression, which is in total

$$\hat{h}_{\text{can}} = \frac{1}{2} + \frac{1}{2} \log \pi + \frac{1}{8} \log 48.$$

The rest of the cases in Table 2 are computed in a similar manner.

In all cases considered above, $\hat{h}_{\text{can}}(-\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)}) - \hat{h}_{\text{can}}(-\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1)})$ is a sum of terms $q(p) \log p$ for primes p and $q(p) \in \mathbb{Q}$. As next shown, this is always the case: \square

Proposition 8.3. *For any ramification indices (m_1, m_2, m_3) such that the corresponding log pair $(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)$ is Fano, i.e. $-\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)}$ is relatively ample,*

$$\hat{h}_{\text{can}}(-\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)}) = \hat{h}_{\text{can}}(-\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1)}) + \sum_p q(p) \log p,$$

where p ranges over a finite number of primes and $q(p) \in \mathbb{Q}$.

Proof. By the ADE-classification of log Fano orbifolds over \mathbb{C} , the orbifold $(\mathbb{P}_{\mathbb{C}}^1, \Delta)$ over \mathbb{C} , induced by $(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)$, coincides with the orbifold induced from an action on \mathbb{P}^1 by a finite group $G \subset SU(2)$ [70, Chapter 8]. By Hurwitz formula, $f^*(-K_{(\mathbb{P}_{\mathbb{C}}^1, \Delta)}) = -K_{\mathbb{P}_{\mathbb{C}}^1}$. Moreover, since G preserves the Fubini-Study metric ω_{FS} on \mathbb{P}^1 its push forward $f_*\omega_{\text{FS}}$ is a Kähler-Einstein metric $(\mathbb{P}_{\mathbb{C}}^1, \Delta)$. Fixing a Kähler-Einstein metric ϕ on $-K_{(\mathbb{P}_{\mathbb{C}}^1, \Delta)}$ this means that $f^*\phi$ defines a Kähler-Einstein metric on $-K_{\mathbb{P}_{\mathbb{C}}^1}$. Next, since the branching locus of the corresponding quotient morphism $f : \mathbb{P}^1 \rightarrow \mathbb{P}^1/G$ is contained in $\{0, 1, \infty\}$ (i.e. it is a Belyi function), f is defined over a number field \mathbb{F} (in fact, there is an explicit formula for f going back to Klein [74, Section 4.1]). As a consequence, there exists a regular projective model \mathcal{Y} of $\mathbb{P}_{\mathbb{F}}^1$ over $\mathcal{O}_{\mathbb{F}}$ and generically finite morphisms g_1 and g_2 from \mathcal{Y} to $\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1$ such that, on the generic fiber $f = g_2 \circ (g_1)^{-1}$ and

$$g_1^*(-\mathcal{K}_{\mathbb{P}_{\mathbb{F}}^1}) = -(\mathcal{K}_{\mathcal{Y}} + \mathcal{R}_1), \quad g_2^*(-\mathcal{K}_{\mathbb{P}_{\mathbb{F}}^1}) = -(\mathcal{K}_{\mathcal{Y}} + \mathcal{R}_2)$$

for \mathbb{Q} -divisors \mathcal{R}_i on \mathcal{Y} . It thus follows from the proof of Lemma 2.4 that

$$\hat{h}_{\phi}(-\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)}) - \hat{h}_{f^*\phi}(-\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1)}) = \sum_p a(p) \log p$$

for a finite number of primes p and $a(p) \in \mathbb{Q}$. Finally, since the integral of the measure attached to ϕ coincides with the integral of the measure attached to $f^*\phi$, up to multiplication by the degree of f , this proves the proposition. \square

This leads one to wonder if the canonical height of any K-semistable Fano orbifold is always a rational number mod $\log(\pi^{\mathbb{Q}}\mathbb{N}^{\mathbb{Q}})$? For example, this is in line with previous explicit height formulas on Fano varieties wrt Kähler-Einstein metrics, which - as far as we know - all concern *homogeneous* Fano varieties [78, 30, 69, 94, 95, 96].

8.3. Extension to other arithmetic triangle groups (?) Let us come back to the case of log pairs satisfying $\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)}(\mathbb{C}) > 0$, for given ramification indices (m_1, m_2, m_3) . Assume that $\Gamma(m_1, m_2, m_3)$ satisfies the following ‘‘arithmetic’’ condition: $\Gamma(m_1, m_2, m_3)$ is a subgroup of $\Gamma^{(+)}(B, \mathcal{O}_B)$ of finite index, where $\Gamma^{(+)}(B, \mathcal{O}_B)$ is a triangle group corresponding to a quaternion algebra B over a totally real number field \mathbb{F} . This means that $\Gamma^{(+)}(B, \mathcal{O}_B)$ is also of the form $\Gamma(n_1, n_2, n_3)$ and that $\mathbb{H}/\Gamma^{(+)}(B, \mathcal{O}_B)$ is one of the components of the complex points of a Shimura curve $(X_{\mathbb{F}}, \Delta_{\mathbb{F}})$ (see Section 9.1.1). In

this case it follows from Yuan's formula 1.1 that

$$(8.3) \quad \hat{h}_{\text{Pet}} \left(\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)} \right) = -\frac{1}{2} - \frac{1}{[\mathbb{F} : \mathbb{Q}]} \frac{\zeta'_{\mathbb{F}}(-1)}{\zeta_{\mathbb{F}}(-1)} + \sum_p q(p) \log p,$$

where p ranges over a finite number of primes and $q(p) \in \mathbb{Q}$. Indeed, the quotient map $f : \mathbb{H}/\Gamma(m_1, m_2, m_3) \rightarrow \mathbb{H}/\Gamma(n_1, n_2, n_3)$ preserves the Peterson metric (see formula 9.2). Moreover, since f is a Belyi function, it is defined over $\bar{\mathbb{Q}}$. Hence, formula 8.3 follows from combining the argument in the proof of Proposition 8.3 with Lemma 2.4 and formula 1.1.

However, computing the numbers $q(p)$ explicitly would require an explicit knowledge of the arithmetic geometry of the canonical model $(\mathcal{X}, \mathcal{D})$, as well as the corresponding Belyi function. In contrast, for the cases in Table 1 (which all satisfy the arithmetic condition above) $q(p)$ is computed explicitly using the formula in Theorem 1.1. Accordingly, it would be interesting to know if Theorem 1.1 could be used to compute $q(p)$ explicitly in *all* these arithmetic cases. While there are probably a few remaining cases where the procedure in Section 8.1 works, we found several cases indicating that the relations coming from the multiplication identity for the Hurwitz zeta function are not enough to establish an explicit formula of the form 8.3. It may still be that some more complicated identities could be leveraged. For instance, $\Gamma(m, m, m)$ satisfies the arithmetic condition in question precisely when $m \in \{4, 5, 6, 7, 8, 9, 12, 15\}$ and then $\mathbb{F} = \mathbb{Q}(\cos(\pi/m))$ [93]. If it would be enough to apply the multiplication identity for the Hurwitz zeta function to extend Table 1 to the case $m = 8$, then the resulting formula would only involve a $\log p$ -term for $p = 2$ (as in formula 1.12). However, numerical investigations indicate that the naive height of the rational coefficient c_2 in front of $\log 2$ would have to be very large ($\gtrsim 30000$).

9. APPLICATIONS TO SHIMURA CURVES

9.1. Setup. We start by recalling the setup in [103, 102]. Let \mathbb{F} be a totally real number field and Σ a finite set of places of \mathbb{F} of odd cardinality, containing all the infinite places of \mathbb{F} . Let \mathbb{B} be a totally definite incoherent quaternion algebra over the adèle ring \mathbb{A} associated to \mathbb{F} . To a compact open subgroup $U \subseteq \mathbb{B}_f^\times$ is attached a *Shimura curve* X_U over \mathbb{F} . This is a non-singular projective curve over \mathbb{F} , which may be defined as a coarse moduli scheme [102, Section 1.2.1]. Its complex points $X_U(\mathbb{C})$ may be represented as follows. Fix an infinite place σ of \mathbb{F} and denote by B the indefinite quaternion algebra over \mathbb{F} with ramification locus $\Sigma - \{\sigma\}$ (denoted by Σ_f). Then $X_{\mathbb{F}}(\mathbb{C})$ is the compactification of

$$B^\times \backslash \mathbb{H}^\pm \times \mathbb{B}_f^\times / U \left(\simeq \bigsqcup_{\sigma} \mathbb{H} / \Gamma_{\sigma} \right)$$

for a finite number of appropriate discrete subgroups Γ_{σ} of $SL(2, \mathbb{R})$. More precisely, this quotient construction induces a log pair (X_U, Δ) such that $K_{(X, \Delta)}$ is ample, where $\Delta_{\mathbb{F}}$ is the orbifold/cusp divisor appearing as the branching divisor, plus the cusps. Denote by X the Shimura curve corresponding to a *maximal* compact open subgroup $U \subseteq \mathbb{B}_f^\times$. By

[102, Section 4.2], it has a canonical integral model \mathcal{X} over $\mathcal{O}_{\mathbb{F}}$, which is a projective flat, normal and \mathbb{Q} -factorial arithmetic surface over $\mathcal{O}_{\mathbb{F}}$. Briefly, the model \mathcal{X} is defined as follows, locally over the base $\text{Spec}\mathcal{O}_{\mathbb{F}}$. First, to an appropriate compact open subgroup U' of \mathbb{B}_f^{\times} is attached a regular model $\mathcal{X}_{U'}$ of $X_{U'}$ that is stable over $\mathcal{O}_{\mathbb{F}}$, in the sense of Deligne-Mumford. Then the scheme \mathcal{X} is defined as the quotient of $\mathcal{X}_{U'}$ by the finite group U'/U . In particular, there is a finite morphism

$$(9.1) \quad g : \mathcal{X}_{U'} \rightarrow \mathcal{X}.$$

Moreover, as shown in [102, Section 4.2], $K_{(X,\Delta)}$ admits a canonical relatively ample model \mathcal{L} over $\mathcal{O}_{\mathbb{F}}$, dubbed the *Hodge bundle*. It may be defined as the norm $N_g(\mathcal{K}_{\mathcal{X}_{U'}})$ of the relative canonical line bundle of $\mathcal{X}_{U'}$ under g , divided by the degree of g . As a consequence, we can express \mathcal{L} as the log canonical line bundle of a canonical effective divisor \mathcal{D} on \mathcal{X} :

$$\mathcal{L} = \mathcal{K}_{(\mathcal{X},\mathcal{D})}, \quad \mathcal{D} := (\deg g)^{-1} N_g(\mathcal{R}), \quad \mathcal{R} := (\mathcal{K}_{\mathcal{X}_{U'}} - g^* \mathcal{K}_{\mathcal{X}}),$$

where \mathcal{R} is an effective divisor on $\mathcal{X}_{U'}$ (using that g is a ramified cover in the sense of [71, Def 2.39]).

The complex points of \mathcal{L} may be identified with $K_{(X,\Delta)}(\mathbb{C})$, which is endowed with the *Petersson metric*. Using the normalization adopted in [103], this is the metric ϕ_{Pet} on $K_{(X,\Delta)}(\mathbb{C})$ which pulls back to the Poincaré metric on the canonical line bundle of the upper half-plane \mathbb{H} under the uniformization maps $\mathbb{H} \rightarrow (X_{\sigma}(\mathbb{C}), \Delta_{\sigma})$ (ramified along Δ_{σ}), where the Poincaré metric is defined by

$$(9.2) \quad \|d\tau\|_{\text{Pet}} := 2\text{Im}(\tau), \quad \mathbb{H} = \{\text{Im}(\tau) > 0\} \Subset \mathbb{C}.$$

Example 9.1. When $\mathbb{F} = \mathbb{Q}$ it is shown in [104, Lemma 2.1] that \mathcal{D} is the Zariski closure of Δ . Moreover, the scheme \mathcal{X} is the coarse moduli scheme of the moduli stack parametrizing all Abelian schemes over \mathbb{Z} of relative dimension 2 with a special action by a maximal order \mathcal{O}_B in B (see [68] and [104, Sections 2.1, 2.2] with $n(U) = 1$). More precisely, when $B = M_2(\mathbb{Q})$ the coarse moduli space has to be “compactified” and then $X_{\mathbb{Q}}$ is the classical modular curve over \mathbb{Q} [39]. In general, when $\mathbb{F} = \mathbb{Q}$, there is a universal abelian scheme $\pi : \mathcal{A} \rightarrow \mathcal{X}$ and, by [104, Thm 1.1], the normalized height of $\overline{\mathcal{L}}$ coincides with normalized height of the Hodge bundle $\pi_* \mathcal{K}_{\mathcal{A}/\mathcal{X}} \rightarrow \mathcal{X}$ plus $2^{-1} \log d_B$ (where d_B denotes the discriminant of B), when the Hodge bundle is endowed with the Faltings metric, normalized in the following way: $\|\alpha\|_x^2 := (2\pi)^{-2} \left| \int_{\pi^{-1}(x)(\mathbb{C})} \alpha \wedge \bar{\alpha} \right|$.

9.1.1. *The case when X has geometric genus zero.* Now specialize to the case when $X_{\mathbb{Q}} \cong \mathbb{P}_{\mathbb{Q}}^1$ and the corresponding divisor Δ is supported at most three points, up to taking finite covers. By the classification result in [93], there are 19 different classes of such quaternionic Shimura curves, corresponding to 13 different totally real fields \mathbb{F} . The ramification indices of the corresponding divisors are explicitly given in [93, Table 3]. Let us recall the classical terminology in [93], since it different than the one in [103]. There is an isomorphism ρ_1 of the quaternion algebra B into $M_2(\mathbb{R})$. The image in $M_2(\mathbb{R})$ of the group of all elements in a maximal order \mathcal{O}_B in A whose reduced norm is a totally positive element in \mathbb{F}^{\times} is denoted by $\Gamma^{(+)}(B, \mathcal{O}_B)$. In the

terminology of [103] the quotient $\mathbb{H}/\Gamma^{(+)}(B, \mathcal{O})$ coincides with one connected component of $X_{\mathbb{F}}(\mathbb{C})$. Indeed, as shown in [101, Section 3.1.1] the complex points $X(\mathbb{C})$ may (up to compactifying the cusps) be decomposed in connected components of the form \mathbb{H}/Γ_h where $\Gamma_h := B_+^{\times} \cap hUh^{-1}$, where B_+ denotes the elements in B with positive reduced norm and h ranges over a some elements including the identity e . Since U is assumed maximal it can be taken to be $\hat{\mathcal{O}}_B := \mathcal{O}_B \times \hat{\mathcal{O}}_{\mathbb{F}}$. Thus, when h is the identity e we get $\Gamma_e = B_+^{\times} \cap \mathcal{O}_B^{\times} = \Gamma^{(+)}(B, \mathcal{O}_B)$.

9.2. Proofs of Theorem 1.3, 1.4. We continue with the case when the Shimura curve $X_{\mathbb{F}}$ has geometric genus zero. Equivalently, there is a finite base change $\mathbb{F} \hookrightarrow \mathbb{F}'$ such that $X_{\mathbb{F}'} \cong \mathbb{P}_{\mathbb{F}'}^1$. Moreover, after perhaps increasing \mathbb{F}' , we may assume that the irreducible components of $\Delta_{\mathbb{F}'} := \Delta_{\mathbb{F}} \otimes \mathbb{F}'$ are defined by \mathbb{F}' -points. The scheme $\mathcal{X} \otimes_{\mathcal{O}_{\mathbb{F}}} \mathcal{O}_{\mathbb{F}'}$ is still normal and \mathbb{Q} -Gorenstein (see [103, Section 4.1]).

In the cases considered below we will show that the optimal model of $(X_{\mathbb{F}'}, \Delta_{\mathbb{F}'})$ over $\mathcal{O}_{\mathbb{F}'}$ is of the form $(\mathbb{P}_{\mathcal{O}_{\mathbb{F}'}}^1, \mathcal{D}^o)$ for a divisor \mathcal{D}^o on $\mathbb{P}_{\mathcal{O}_{\mathbb{F}'}}^1$. It follows from Lemma 2.4 that, for any fixed metric on $K_{(X, \Delta)(\mathbb{C})}$,

$$(9.3) \quad \frac{1}{[\mathbb{F}' : \mathbb{Q}]} \left(h(\overline{\mathcal{K}_{(\mathcal{X}, \mathcal{D})} \otimes_{\mathcal{O}_{\mathbb{F}}} \mathcal{O}_{\mathbb{F}'}}) - h(\overline{\mathcal{K}_{(\mathbb{P}_{\mathcal{O}_{\mathbb{F}'}}^1, \mathcal{D}^o)}}) \right) = \sum_p h(p) \log p,$$

for a finite number of prime numbers p and rational numbers $h(p)$, independent of the choice of \mathbb{F}' . The number $h(p)$ may be geometrically expressed as follows. Fix any normal model \mathcal{Y} of $X_{\mathbb{F}'}$ over $\mathcal{O}_{\mathbb{F}'}$, dominating both $\mathcal{X} \otimes_{\mathcal{O}_{\mathbb{F}}} \mathcal{O}_{\mathbb{F}'}$ and $\mathbb{P}_{\mathcal{O}_{\mathbb{F}'}}^1$. Denote by \mathfrak{p}_i the prime ideals in $\mathcal{O}_{\mathbb{F}'}$ over p . Then

$$h(p) = \frac{1}{[\mathbb{F}' : \mathbb{Q}]} \sum_i h(\mathfrak{p}_i) f_i, \quad N(\mathfrak{p}_i) =: p^{f_i}$$

where $h(\mathfrak{p}_i)$ is the sum of intersection numbers on the fiber $\mathcal{Y}_{\mathfrak{p}_i}$ defined in Lemma 2.4, for $\mathcal{L}' = \mathcal{K}_{(\mathcal{X}, \mathcal{D})} \otimes_{\mathcal{O}_{\mathbb{F}}} \mathcal{O}_{\mathbb{F}'}$ and $\mathcal{L} = \mathcal{K}_{(\mathbb{P}_{\mathcal{O}_{\mathbb{F}'}}^1, \mathcal{D}^o)}$. Note that $h(\mathfrak{p}_i) \geq 0$. Indeed, since $\mathcal{K}_{(\mathcal{X}, \mathcal{D})} \otimes_{\mathcal{O}_{\mathbb{F}}} \mathcal{O}_{\mathbb{F}'}$ is the log canonical line bundle of the log pair $(\mathcal{X}, \mathcal{D}) \otimes_{\mathcal{O}_{\mathbb{F}}} \mathcal{O}_{\mathbb{F}'}$ (using that $\mathcal{K}_{\mathcal{X}} \otimes_{\mathcal{O}_{\mathbb{F}}} \mathcal{O}_{\mathbb{F}'} = \mathcal{K}_{\mathcal{X} \otimes_{\mathcal{O}_{\mathbb{F}}} \mathcal{O}_{\mathbb{F}'}}$), the non-negativity of $h(\mathfrak{p}_i)$ follows from Remark 3.15.

We will compute $h(p)$ for some Shimura curves. By the uniqueness of prime factorization it will be enough to compute the left hand side in formula 9.3.

9.2.1. Height formulas. All heights will be computed wrt the Petersson metric on $K_{(X, \Delta)(\mathbb{C})}$, denoted by h_{Pet} .

Lemma 9.2. *The volume of the measure μ corresponding to the Petersson metric on $K_{(X, \Delta)}$ is equal to $\pi V/2$:*

$$\int_X \mu = \pi V/2, \quad V := V(K_{(X, \Delta)}).$$

In particular, $\hat{h}_{\text{Pet}}(\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)}) = \hat{h}_{\text{can}}(\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)}) + \frac{1}{2} \log(\pi V/2)$.

Proof. First observe that, in general, if μ is the measure corresponding to a finite energy metric ϕ on $K_{(X,\Delta)}$, then

$$\int_{X-\Delta} dd^c \phi = V(K_{(X,\Delta)})$$

(using that $dd^c \phi$ does not charge finite subsets). Now, let μ be the measure induced by Petersson metric on $K_{(X,\Delta)}$. As recalled above this means that $\mu = \mu_\phi$ where, locally, $\phi := -\log(\|d\tau\|^2) := -\log((2y)^2)$. Note that $dd^c \phi = \frac{1}{\pi} \mu$ on $X - \Delta$. Indeed,

$$dd^c \phi := \frac{1}{\pi} i \partial \bar{\partial} \phi := \frac{1}{\pi} \left(\frac{\partial}{\partial z} \frac{\partial}{\partial \bar{z}} \phi \right) dx dy = \frac{2}{\pi} \frac{1}{(2y)^2} dx dy =: \frac{2}{\pi} e^\phi dx dy =: \frac{2}{\pi} \mu$$

All in all this means that $V(K_{(X,\Delta)}) = \int_{X-\Delta} dd^c \phi = \int_{X-\Delta} \frac{2}{\pi} \mu$, proving the desired formula. \square

Theorem 1.1 thus implies the following corollary, where \mathcal{D}° denotes the divisor on $\mathbb{P}_{\mathbb{Z}}^1$ defined as the Zariski closure of the divisor $\Delta_{\mathbb{Q}}$ on $\mathbb{P}_{\mathbb{Q}}^1$ supported on $\{0, 1, \infty\}$ with weights $w_i \in [0, 1]$.

Corollary 9.3. *The following formula holds when $K_{(\mathbb{P}^1, \Delta)}$ is ample:*

$$\hat{h}_{\text{Pet}}(\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^\circ)}) = \frac{1}{2} - \frac{\gamma(0, \frac{V}{2}) - \sum_{i=1}^3 \gamma(w_i - \frac{V}{2}, w_i)}{V}$$

9.2.2. *The case $\mathbb{F} = \mathbb{Q}$, $\Sigma_f = \emptyset$.* Let us show how to recover height formula 1.1 in the case $\Sigma_f = \emptyset$ from Theorem 1.1. By [39] the corresponding canonical model \mathcal{X} is isomorphic to $\mathbb{P}_{\mathbb{Z}}^1$ over \mathbb{Z} (under the morphism defined by the j -invariant) and $\Delta_{\mathbb{Q}} (:= \mathcal{D} \otimes_{\mathbb{Z}} \mathbb{Q})$ is supported on the three points $0, 1728$ and ∞ in $\mathbb{P}_{\mathbb{Q}}^1$ with ramification indices $2, 3$ and ∞ , respectively. In general, if $a \in \mathbb{Z}$ the divisor \mathcal{D}_a on $\mathbb{P}_{\mathbb{Z}}^1$ defined as the Zariski closure of the divisor on $\mathbb{P}_{\mathbb{Q}}^1$ supported at the points $0, a$ and ∞ (with given weights w_0, w_1 and w_∞) satisfies

$$\hat{h}_{\text{Pet}}(\mathcal{K}_{(\mathcal{X}, \mathcal{D}_a)}) = \hat{h}_{\text{Pet}}(\mathcal{K}_{(\mathcal{X}, \mathcal{D}_1)}) - \left(\frac{\sum_{i \leq \infty} w_i}{2} - \sum_{i < \infty} w_i \right) \log a.$$

This follows Theorem 5.6, using the change of variables $z_i = a\zeta_i$ in the integral formula 6.2 (but it can also be shown directly using scheme theory). In the present case $\mathcal{D} = \mathcal{D}_{1728}$, i.e. $a = 1728$. The bracket above thus becomes $-\frac{1}{12}$. Since $1728 = 12^3 (= 2^6 3^3)$ it follows that

$$\hat{h}_{\text{Pet}}(\mathcal{K}_{(\mathcal{X}, \mathcal{D}_a)}) = \hat{h}_{\text{Pet}}(\mathcal{X}, \mathcal{D}_1) + \frac{1}{12} \log(12^3) (\implies h(2) = \frac{1}{2}, h(3) = \frac{1}{4}).$$

A new proof of formula 1.1 is thus obtained by invoking the formula for ramification indices $(2, 3, \infty)$ in Table 1.

Remark 9.4. From the identity $\mathcal{D} = \mathcal{D}_{1728}$ one sees directly that the reduction mod p of $(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D})$ is log canonical iff the prime p is not in $\{2, 3\}$ and that $\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D})}$ is isomorphic to $\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^\circ)}$ precisely over the complement in $\mathbb{P}_{\mathbb{Z}}^1$ of the fibers over (2) and (3). This is

consistent (as it must) with the fact, shown above, that $h(p)$ vanishes iff p is not in $\{2, 3\}$ (see Remark 3.15).

9.2.3. *The case $\mathbb{F} = \mathbb{Q}, \Sigma_f = \{2, 3\}$ (proof of Theorem 1.3).* Now consider the case when the indefinite quaternion algebra B over \mathbb{Q} has discriminant 6, i.e. it is ramified at 2 and 3. According to a result attributed to Ihara, the corresponding Shimura curve $X_{\mathbb{Q}}$ is the subscheme of $\mathbb{P}_{\mathbb{Q}}^2$ cut out by $x_0^2 + 3x_1^2 + x_2^2$ (see [42, Section 3.1] for a proof). In particular, $X_{\mathbb{Q}}$ admits a $\mathbb{Q}(\sqrt{-3})$ -point, e.g. $[1 : (\sqrt{-3})^{-1} : 0]$. It follows that $X_{\mathbb{Q}} \otimes \mathbb{Q}(\sqrt{-3})$ is isomorphic to $\mathbb{P}_{\mathbb{Q}(\sqrt{-3})}^1$ (by stereographic projection through any \mathbb{F} -point). Furthermore, by [42, Section 3.1] and [93, Table 3], setting $\mathbb{F} := \mathbb{Q}(\sqrt{3}, i)$ the corresponding divisor $\Delta_{\mathbb{Q}} \otimes \mathbb{F}$ is supported at four \mathbb{F} -points with ramification indices $(3, 3; 2, 2)$. Moreover, as explained in [42, Section 3.1], the cross ratio of the corresponding pair of two points is -1 . Denote by $(\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1, \mathcal{D}^o)$ the corresponding unique optimal model over $\mathcal{O}_{\mathbb{F}}$, furnished by Lemma 3.13 and Prop 3.14.

We will compute the left hand side in formula 9.3 wrt the Petersson metric.

Lemma 9.5. *The following formula holds,*

$$\hat{h}_{\text{Pet}}(\overline{\mathcal{K}_{(\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1, \mathcal{D}^o)}}) = -\frac{\zeta'(-1)}{\zeta(-1)} - \frac{1}{2} - \left(\left(\frac{1}{6} - \frac{1}{2} \right) \log 2 - \frac{1}{8} \log 3 \right),$$

which, combined with formula 1.1, gives

$$\hat{h}(\overline{\mathcal{K}_{(\mathcal{X}, \mathcal{D})}}) - \hat{h}(\overline{\mathcal{K}_{(\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1, \mathcal{D}^o)}}) = \frac{11}{12} \log 2 + \frac{7}{8} \log 3$$

Proof. By Prop 6.2

$$\hat{h}_{\text{Pet}}(\overline{\mathcal{K}_{(\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1, \mathcal{D}^o)}}) = \hat{h}_{\text{Pet}}(\overline{\mathcal{K}_{(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}')}}) + \frac{1}{2} \log 2,$$

where \mathcal{D}' is the Zariski closure of the divisor supported at $(0, 1, \infty)$ with ramification indices $(6, 2, 6)$. Hence, by Table 1,

$$\hat{h}_{\text{Pet}}(\overline{\mathcal{K}_{(\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1, \mathcal{D}^o)}}) = -\frac{\zeta'(-1)}{\zeta(-1)} - \frac{1}{2} - \left(\frac{1}{6} \log 2 - \frac{1}{8} \log 3 \right) + \frac{1}{2} \log 2,$$

Combining this result with Yuan's formula 1.1 for $\mathfrak{p} = (2)$ and $\mathfrak{p} = (3)$ in \mathbb{Z} reveals that

$$\begin{aligned} \hat{h}(\overline{\mathcal{K}_{(\mathcal{X}, \mathcal{D})}}) - \hat{h}(\overline{\mathcal{K}_{(\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1, \mathcal{D}^o)}}) &= \frac{3 \cdot 2 - 1}{4} \log 2 + \frac{3 \cdot 3 - 1}{4(3-1)} \log 3 + \left(\left(\frac{1}{6} - \frac{1}{2} \right) \log 2 - \frac{1}{8} \log 3 \right) = \\ &= \left(\frac{3 \cdot 2 - 1}{4} + \frac{1}{6} - \frac{1}{2} \right) \log 2 + \left(\frac{3 \cdot 3 - 1}{4(3-1)} - \frac{1}{8} \right) \log 3 = \frac{11}{12} \log 2 + \frac{7}{8} \log 3 \end{aligned}$$

□

Since the normalized height is invariant under base change we have $\hat{h}(\overline{\mathcal{K}_{(\mathcal{X}, \mathcal{D})}} \otimes_{\mathbb{Z}} \mathcal{O}_{\mathbb{F}}) = \hat{h}(\overline{\mathcal{K}_{(\mathcal{X}, \mathcal{D})}})$. Hence, setting $\hat{h}(p) := h(p)/(2K_{(X_{\mathbb{F}}, \Delta_{\mathbb{F}})} \cdot X_{\mathbb{F}})$ the previous lemma gives $\hat{h}(2) = \frac{11}{12}$ and $\hat{h}(3) = \frac{7}{8}$ (using uniqueness of prime factorization in \mathbb{Z}). Since $h(p) = \hat{h}(p) \cdot 2/3$ this means that $h(2) = 11/18$ and $h(3) = 7/12$.

9.2.4. *The quaternion algebra over $\mathbb{Q}(\sqrt{3})$ ramified over 3 (proof of Theorem 1.4).* Now consider the quaternion algebra over $\mathbb{Q}(\sqrt{3})$ that is only ramified at the unique prime ideal \mathfrak{p}_3 in $\mathcal{O}_{\mathbb{Q}(\sqrt{3})}$ containing 3. In fact, $\mathfrak{p}_3 = (\sqrt{3})$. Indeed, (3) is the square of the ideal $(\sqrt{3})$, which has norm $N(\mathfrak{p}_3) = 3$. As a consequence, the contribution from prime ideals in formula 1.1 for $\hat{h}(\overline{\mathcal{K}(\mathcal{X}, \mathcal{D})})$ is

$$\frac{3N(\mathfrak{p}) - 1}{4(N(\mathfrak{p}) - 1)} \sum_{\mathfrak{p}} \log N(\mathfrak{p}) = \log 3,$$

Moreover, by [93, Table 3], $(\mathcal{X}, \mathcal{D}) \otimes \bar{\mathbb{Q}}$ is isomorphic to $(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o) \otimes \bar{\mathbb{Q}}$, where \mathcal{D}^o is the divisor appearing in Theorem 1.1 with ramification indices (2, 4, 12). Fix a finite field extension \mathbb{F} of $\mathbb{Q}(\sqrt{3})$ such that $(\mathcal{X}, \mathcal{D}) \otimes \mathbb{F}$ is isomorphic to $(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o) \otimes \mathbb{F}$. Combining formula 1.1 with Theorem 1.1 thus yields, using Table 1,

$$\hat{h}(\overline{\mathcal{K}(\mathcal{X}, \mathcal{D})}) - \hat{h}(\overline{\mathcal{K}(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)}) = \frac{1}{2} \log 3 + \frac{5}{3} \log 2 + \frac{7}{16} \log 3.$$

Note that $1/2 + 7/16 = 15/16$. Since the normalized height is invariant under base change, it follows that

$$\hat{h}(\overline{\mathcal{K}(\mathcal{X}, \mathcal{D}) \otimes \mathcal{O}_{\mathbb{F}}}) - \hat{h}(\overline{\mathcal{K}(\mathbb{P}_{\mathbb{F}}^1, \mathcal{D}^o)}) = \hat{h}_2 \log 2 + \hat{h}_3 \log 3, \quad \hat{h}_2 = \frac{5}{3}, \quad \hat{h}_3 = \frac{15}{16}.$$

Since $h(p) = \hat{h}(p) \cdot (2K_{(X_{\mathbb{F}}, \Delta_{\mathbb{F}})} \cdot X_{\mathbb{F}})$ and $K_{(X_{\mathbb{F}}, \Delta_{\mathbb{F}})} \cdot X_{\mathbb{F}} = 1/2 + 3/4 + 1/12 - 2 = 1/6$ we deduce that

$$h(2) = 2 \cdot (1/6) \cdot (5/3) = \frac{5}{9}, \quad h(3) = 2 \cdot (1/6) \cdot (15/16) = \frac{15}{48}.$$

9.2.5. *The quaternion algebra over $\mathbb{Q}(\sqrt{6})$ ramified over 2.*

Theorem 9.6. *Consider the quaternion algebra over $\mathbb{Q}(\sqrt{6})$ ramified over the unique prime ideal \mathfrak{p}_2 containing 2 and denote by $(\mathcal{X}, \mathcal{D})$ the canonical model over $\mathcal{O}_{\mathbb{Q}(\sqrt{6})}$ of the corresponding Shimura curve $(X_{\mathbb{Q}(\sqrt{6})}, \Delta_{\mathbb{Q}(\sqrt{6})})$. Fix a finite field extension \mathbb{F} of $\mathbb{Q}(\sqrt{6})$ such that $X_{\mathbb{Q}(\sqrt{6})} \otimes \mathbb{F}$ is isomorphic to $\mathbb{P}_{\mathbb{F}}^1$ and $\Delta_{\mathbb{F}}$ is supported on three \mathbb{F} -points. Then the optimal model of $(X_{\mathbb{Q}(\sqrt{6})}, \Delta_{\mathbb{Q}(\sqrt{6})}) \otimes \mathbb{F}$ over $\mathcal{O}_{\mathbb{Q}(\sqrt{6})}$ is given by $(\mathbb{P}_{\mathbb{F}}^1, \mathcal{D}^o)$, where \mathcal{D}^o denotes the Zariski closure of the divisor on $\mathbb{P}_{\mathbb{F}}^1$ supported on $\{0, 1, \infty\}$ having the same ramification indices (3, 4, 6) as the divisor $\Delta_{\mathbb{F}}$. Moreover $h(p) = 0$ unless $p = 2$ or $p = 3$ and*

$$h(2) = \frac{43}{144}, \quad h(3) = \frac{3}{32}.$$

Proof. The unique prime ideal \mathfrak{p}_2 of $\mathbb{Q}(\sqrt{6})$ containing 2 is given by $(2 + \sqrt{6})$, which has norm $N(\mathfrak{p}_2) = 2$. As such, the contribution coming from the prime ideals in 1.1 is given by

$$\frac{3N(\mathfrak{p}) - 1}{4(N(\mathfrak{p}) - 1)} \sum_{\mathfrak{p}} \log N(\mathfrak{p}) = \frac{7}{4} \log 2.$$

By [93, Table 3], $(\mathcal{X}, \mathcal{D}) \otimes \bar{\mathbb{Q}}$ is isomorphic to $(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o) \otimes \bar{\mathbb{Q}}$, where \mathcal{D}^o is the divisor appearing in Theorem 1.1 with ramification indices $(3, 4, 6)$. Combining formula 1.1 and row 7 in Table 1 yields,

$$\hat{h}(\overline{\mathcal{K}(\mathcal{X}, \mathcal{D})}) - \hat{h}(\overline{\mathcal{K}(\mathbb{P}_{\mathbb{Z}}^1, \mathcal{D}^o)}) = \frac{1}{2} \frac{7}{4} \log 2 + \frac{9}{16} \log 3 + \frac{11}{12} \log 2.$$

□

Since the normalized height is invariant under base change, it follows that

$$\hat{h}(\overline{\mathcal{K}(\mathcal{X}, \mathcal{D}) \otimes \mathcal{O}_{\mathbb{F}}}) - \hat{h}(\overline{\mathcal{K}(\mathbb{P}_{\mathcal{O}_{\mathbb{F}}}^1, \mathcal{D}^o)}) = \hat{h}_2 \log 2 + \hat{h}_3 \log 3, \quad \hat{h}_2 = \frac{43}{24}, \quad \hat{h}_3 = \frac{9}{16}.$$

Since $h(p) = \hat{h}(p) \cdot (2K_{(X_{\mathbb{F}}, \Delta_{\mathbb{F}})} \cdot X_{\mathbb{F}})$ and $K_{(X_{\mathbb{F}}, \Delta_{\mathbb{F}})} \cdot X_{\mathbb{F}} = 2/3 + 3/4 + 5/6 - 2 = 1/12$ we deduce that

$$h(2) = 2 \cdot (1/12) \cdot (43/24) = \frac{43}{144}, \quad h(3) = 2 \cdot (1/12) \cdot (9/16) = \frac{3}{32}.$$

9.3. Implications for wild ramification and intersections over special places.

As recalled in Section 9.1, the canonical integral model \mathcal{X} of a quaternionic Shimura curve, comes, locally over the base $\text{Spec } \mathcal{O}_{\mathbb{F}}$, with a finite morphism from a regular scheme \mathcal{X}' to \mathcal{X} (formula 9.1), induced by the action of a finite group G on \mathcal{X}' . The morphism induces an effective divisor \mathcal{D} on \mathcal{X} . Consider now $\mathfrak{p} \in \text{Spec } \mathcal{O}_{\mathbb{F}}$ which is *split*, i.e. \mathfrak{p} is not in the ramification locus of the quaternion algebra B . Denote by κ the residue field of \mathfrak{p} and by $\bar{\kappa}$ its algebraic closure. Both $\mathcal{X}'_{\mathfrak{p}} \otimes_{\kappa} \bar{\kappa}$ and $\mathcal{X}_{\mathfrak{p}} \otimes_{\kappa} \bar{\kappa}$ are smooth [103, Section 4.1]. Moreover, by [103, Prop 4.1], the restricted finite morphism

$$(9.4) \quad g : \mathcal{X}'_{\mathfrak{p}} \rightarrow \mathcal{X}_{\mathfrak{p}}$$

is unramified at the generic points of $\mathcal{X}'_{\mathfrak{p}}$. This means that the restricted finite morphism 9.4 is a ramified cover in the sense of [71, Def 2.39]. Accordingly, the restriction of \mathcal{D} to $\mathcal{X}_{\mathfrak{p}}$ defines a divisor on $\mathcal{X}_{\mathfrak{p}}$ that we shall denote by $\mathcal{D}_{\mathfrak{p}}$. In general, a ramified cover is called *tame* at a given prime divisor P' on $\mathcal{X}'_{\mathfrak{p}}$ if the characteristic of the residue field of P' does not divide the ramification index of g along P' . We will say that g has *wild ramification* if the ramification is not tame at all prime divisors P' on $\mathcal{X}'_{\mathfrak{p}}$. Theorems 1.4, 9.6 imply the following

Corollary 9.7. *When \mathcal{X} is the canonical model in Theorem 1.4 the log pair $(\mathcal{X}_{\mathfrak{p}_2}, \mathcal{D}_{\mathfrak{p}_2})$ is not log stable, i.e. it is not log canonical (lc). As a consequence, the ramified cover 9.4 has wild ramification over \mathfrak{p}_2 and some of the irreducible components of the divisor \mathcal{D} on \mathcal{X} coincide, when restricted to the fiber of \mathcal{X} over \mathfrak{p}_2 . Moreover, when \mathcal{X} is the canonical model in Theorem 9.6, the corresponding result holds over \mathfrak{p}_3 .*

Proof. Denote by \mathfrak{p} a prime ideal appearing in the statement of the corollary and by \mathbb{F} the totally real field in question. Since \mathfrak{p} is split [103, Prop 4.1] shows, as recalled above, that 9.4 is a ramified cover. Let us first show that the log pair $(\mathcal{X}_{\mathfrak{p}}, \mathcal{D}_{\mathfrak{p}})$ is not lc. Assume, in order to get a contradiction, that $(\mathcal{X}_{\mathfrak{p}}, \mathcal{D}_{\mathfrak{p}})$ is lc. Take a finite field extension \mathbb{F}' of \mathbb{F} to which Theorem 1.4 applies and fix a prime ideal \mathfrak{p}' in $\mathcal{O}_{\mathbb{F}'}$ over \mathfrak{p} . Then the restriction

of $(\mathcal{X}, \mathcal{D}) \otimes_{\mathcal{O}_{\mathbb{F}'}} \mathcal{O}_{\mathbb{F}'}$ to the fiber over \mathfrak{p}' , that we denote by $(\mathcal{X}_{\mathfrak{p}'}, \mathcal{D}_{\mathfrak{p}'})$, is also lc. Indeed, in general, as recalled in Section 2.1.1, if X is a normal scheme of dimension one over a perfect field, then (X, \mathcal{D}) is lc iff $w_i \leq 1$ for all coefficients w_i of \mathcal{D} . Since any finite field is perfect this applies to $(\mathcal{X}_{\mathfrak{p}}, \mathcal{D}_{\mathfrak{p}})$. Hence, decomposing $\mathcal{D}_{\mathfrak{p}} = \sum w_i D_{\mathfrak{p}}^{(i)}$ where $D_{\mathfrak{p}}^{(i)}$ is a prime divisor on X we have $w_i \leq 1$. Next, since the residue field $\mathbb{F}_{\mathfrak{p}}$ is perfect, $D_{\mathfrak{p}}^{(i)} \otimes_{\mathbb{F}_{\mathfrak{p}}} \mathbb{F}_{\mathfrak{p}'}$ is a sum of *distinct* irreducible divisors $D_{\mathfrak{p}'}^{(i,j)}$. As a consequence, the coefficients of $\mathcal{D}_{\mathfrak{p}'}$ are at most 1, showing that $(\mathcal{X}_{\mathfrak{p}'}, \mathcal{D}_{\mathfrak{p}'})$ is indeed lc. But this implies that $h(\mathfrak{p}') = 0$, where $h(\mathfrak{p}')$ is defined in formula 1.9, comparing $(\mathcal{X}, \mathcal{D}) \otimes_{\mathcal{O}_{\mathbb{F}'}} \mathcal{O}_{\mathbb{F}'}$ with the optimal model appearing in Theorem 1.4. Indeed, since $(\mathcal{X}_{\mathfrak{p}'}, \mathcal{D}_{\mathfrak{p}'})$ is lc the vanishing $h(\mathfrak{p}') = 0$ follows from Prop 3.12. Finally, the vanishing of $h(\mathfrak{p}')$ for all prime ideals \mathfrak{p}' over \mathfrak{p} implies that $h(\mathfrak{p}) = 0$, which contradicts Theorem 1.4. Next, to show the statement about wild ramification, first observe that, since 9.4 is a ramified cover we have

$$g^*(\mathcal{K}_{\mathcal{X}_{\mathfrak{p}}} + \mathcal{D}_{\mathfrak{p}}) = \mathcal{K}_{\mathcal{X}'_{\mathfrak{p}}}.$$

Assume, to get a contradiction, that g does not have wild ramification over \mathfrak{p} . This implies, since $(\mathcal{X}'_{\mathfrak{p}}, 0)$ is lc (and even klt) that $(\mathcal{X}_{\mathfrak{p}}, \mathcal{D}_{\mathfrak{p}})$ is lc, by a Hurwitz type formula (see [71, Cor 2.43]). This is a contradiction. Likewise, if the irreducible components of $\mathcal{D}_{\mathfrak{p}}$ were all distinct, then the coefficients of $\mathcal{D}_{\mathfrak{p}}$ would all be of the form $1 - 1/m_i$ for positive integers m_i (since \mathcal{D} is the Zariski closure of an orbifold divisor on the generic fiber). Thus $(\mathcal{X}_{\mathfrak{p}}, \mathcal{D}_{\mathfrak{p}})$ would be klt, contradicting that it is not even lc. \square

In general, when \mathfrak{p} is split, $\mathcal{X}_{\mathfrak{p}}$ is isomorphic to $\mathcal{X}'_{\mathfrak{p}}/G$, by [103, Prop 4.1]. The previous corollary also applies to the classical case when $B = M_2(\mathbb{Q})$, where \mathcal{X} is the compactification of the coarse moduli space of elliptic curves over \mathbb{Z} . In this case all \mathfrak{p} are split. The non-vanishing 1.10 thus implies that 9.4 has wild ramification over $\mathfrak{p} = (2)$ and $\mathfrak{p} = (3)$. This also follows from classical results about elliptic curves. Indeed, for $p = 2$ and $p = 3$ there exist elliptic curves E over $\mathbb{Z}/(p)$ such that $\sharp(\text{Aut}(E)/\{\pm 1\})$ is 12 or 6, respectively. These elliptic curves give rise to local ramification indices for the morphism 9.4 of order 12 and 6, respectively, which are thus divided by p .

10. APPLICATION TO TWISTED FERMAT CURVES

In this Section we will, in particular, prove Theorem 1.6. Given integers a_i consider the subscheme \mathcal{X}_a of $\mathbb{P}_{\mathbb{Z}}^{n+1}$ cut out by the homogeneous polynomial $\sum_{i=0}^{n+1} a_i x_i^d$. This scheme will be denoted by \mathcal{X}_1 in the case $a_i = 1$.

Proposition 10.1. *The following formula holds when $\pm K_{\mathcal{X}_a}$ is ample (i.e when $\pm(d - (n + 2)) > 0$)*

$$h_{can}(\mathcal{K}_{\mathcal{X}_a}) = h_{can}(\mathcal{K}_{\mathcal{X}_1}) + \left(\frac{|n+2-d|}{n+1} \pm 1\right) d^{-1} \sum_i \log(|a_i|).$$

Proof. The case when $-K_{\mathcal{X}} > 0$ is the content of [3, formula 5.5] (applied to $k = n + 2 - d$). The proof in the case when $K_{\mathcal{X}} > 0$ is essentially the same, but then k is

[3, Lemma 5.3] is taken as $d - (n + 2)$ (by adjunction) and the minus sign in \pm results from change in sign in front of $\log \int \mu_\phi$ (see [3, Lemma 5.4]). \square

It follows that

$$h_{\text{can}}(\mathcal{X}_a) \leq h_{\text{can}}(\mathcal{X}_1) \text{ when } -K_{X_a} > 0, \quad h_{\text{can}}(\mathcal{X}_1) \leq h_{\text{can}}(\mathcal{X}_a), \text{ when } K_{X_a} > 0$$

Equivalently, by Prop 3.3, this means that

$$\inf_{\psi} \mathcal{M}_{\mathcal{X}_1}(\pm \mathcal{K}_{\mathcal{X}_1}, \psi) \leq \inf_{\psi} \mathcal{M}_{\mathcal{X}_a}(\pm \mathcal{K}_{\mathcal{X}_a}, \psi).$$

Now we specialize to $n = 1$. Given a positive integer m consider the divisor \mathcal{D} on \mathbb{P}^1 supported on $\{0, 1, \infty\}$ with coefficients $(1 - 1/m)$.

Lemma 10.2. *Denote by \mathcal{X} the Fermat hypersurface of a given degree $m (> 2)$. Then,*

$$\hat{h}_{\text{can}}(\mathcal{K}_{\mathcal{X}}) = \hat{h}_{\text{can}}(\mathcal{K}_{(\mathbb{P}^1, \mathcal{D})}) + \frac{1}{2} \log \frac{V(X)}{V(\mathbb{P}^1, \Delta)} \quad \left(\frac{V(X)}{V(\mathbb{P}^1, \Delta)} = m^2 \right)$$

Proof. This is shown exactly as in the Fano case in [3, Prop 5.6], but now the last term comes with a different sign (due to the sign difference in the definition of $\hat{h}_{\text{can}}(\mathcal{X})$). \square

10.1. Proof of Theorem 1.6. The first formula in Theorem 1.6 follows directly from combining the previous proposition and lemma with Theorem 1.1. Next, fix m and a and set $\mathcal{X}_a = \mathcal{X}_a^{(m)}$. By [38], there exists a stable model \mathcal{X}^s for $\mathcal{X}_a \otimes_{\mathbb{Z}} \mathcal{O}_{\mathbb{F}}$ over $\mathcal{O}_{\mathbb{F}}$ for some number field \mathbb{F} . Since the base change of a stable model is still a stable model [?, Section 1.5], we may as well assume that \mathbb{F} contains all $a_i^{1/m}$. Thus $X_a \otimes_{\mathbb{Q}} \mathbb{F}$ is isomorphic to $X_1 \otimes_{\mathbb{Q}} \mathbb{F}$ over \mathbb{F} , showing that \mathcal{X}^s is also a stable model for $\mathcal{X}_1 \otimes_{\mathbb{Z}} \mathcal{O}_{\mathbb{F}}$. Hence, to prove the inequalities in Theorem 1.6, it will - by the first formula in Theorem 1.6 (combined with Theorem 1.2 and Lemma 3.4) - be enough to show that

$$(10.1) \quad \hat{h}_{\text{can}}(\mathcal{K}_{\mathcal{X}^s}) \leq \hat{h}_{\text{can}}(\mathcal{K}_{\mathcal{X}_1}).$$

But, by Cor 3.10, $\hat{h}_{\text{can}}(\mathcal{K}_{\mathcal{X}^s}) \leq \hat{h}_{\text{can}}(\mathcal{K}_{\mathcal{X}_1 \otimes_{\mathbb{Z}} \mathcal{O}_{\mathbb{F}}})$. Since $\mathcal{K}_{\mathcal{X}_1 \otimes_{\mathbb{Z}} \mathcal{O}_{\mathbb{F}}}$ is isomorphic to $\mathcal{K}_{\mathcal{X}_1} \otimes_{\mathbb{Z}} \mathcal{O}_{\mathbb{F}}$ (by the adjunction formula) and the normalized height is invariant under base change this proves the inequality 10.1.

10.2. The Arakelov vs the Kähler-Einstein metric (proof of Cor 1.7). Let $X_{\mathbb{Q}}$ be a non-singular projective curve of degree m in $\mathbb{P}_{\mathbb{Q}}^2$ and denote by X its complex points. Assume that K_X is ample, i.e. $m \geq 4$. Denote by g_X the genus of X . The Arakelov metric on K_X may be defined as the metric which turns the adjunction formula into an isometry [44].

The first inequality in Cor 1.7 follows directly from combining Theorem 1.6 with the following bound (using that $V = 1 - 3/m$):

Lemma 10.3. *For any given model \mathcal{X} of X over \mathbb{Z}*

$$\hat{h}_{\text{Ar}}(\mathcal{X}) \leq \hat{h}_{\text{can}}(\mathcal{X}) + \frac{1}{2} \log \pi + \frac{1}{2} \frac{4 \log((m-1)(m-2)-2) + 1}{(m-1)(m-2)/2-1} + \frac{1}{2} \log((m-1)(m-2)/2-1).$$

This bound follows from results in [62, 63], as next explained. First recall that, by 2.6,

$$2\hat{h}_{\psi_{\text{Ar}}}(\mathcal{X}) - 2\hat{h}_{\text{can}}(\mathcal{X}) = \frac{\mathcal{E}(\psi_{\text{Ar}}, \psi_{\text{KE}})}{2V(K_X)},$$

where ψ_{KE} denotes the unique volume-normalized Kähler-Einstein metric on X . Comparing with the notation in [63, Section 2.1], $\mu_{\text{hyp}} := 4\pi(g_X - 1)\mu_{\psi_{\text{KE}}}$, where $\mu_{\psi_{\text{KE}}}$ denotes the measure on X corresponding to ψ_{KE} . Denoting by ψ_{hyp} the Kähler-Einstein metric on K_X corresponding to μ_{hyp} , [63, Prop 4.5] thus yields the following bound:

$$\frac{\mathcal{E}(\psi_{\text{Ar}}, \psi_{\text{hyp}})}{2V(K_X)} := \int_X (\psi_{\text{Ar}} - \psi_{\text{hyp}}) \frac{(dd^c \psi_{\text{hyp}} + dd^c \psi_{\text{Ar}})}{2V(K_X)} \leq -\frac{c_X - 1}{g_X - 1} - \log 4,$$

where c_X is the finite part of the logarithmic derivative of the Selberg zeta function at $s = 1$, defined before [63, formula 2.8]. Since $\psi_{\text{hyp}} := \psi_{\text{KE}} + \log(4\pi(g_X - 1))$ this means that

$$\frac{\mathcal{E}(\psi_{\text{Ar}}, \psi_{\text{KE}})}{2V(K_X)} \leq \frac{-c_X + 1}{g_X - 1} - \log 4 + \log(4\pi(g_X - 1)).$$

Next, we recall that, by [62, Thm 3.3],

$$-c_X \leq 4\log(2g_X - 2)$$

(since X has no cusps, nor elliptic points; compare [62, Section 2.1]). Hence,

$$(10.2) \quad 2\hat{h}_{\psi_{\text{Ar}}}(\mathcal{X}) - 2\hat{h}_{\text{can}}(\mathcal{X}) \leq \frac{4\log(2g_X - 2) + 1}{g_X - 1} + \log(\pi(g_X - 1))$$

and since $2g_X = (m - 1)(m - 2)$ this proves Lemma 10.3.

Finally, to prove the second inequality in Cor 1.7 recall that it was shown in the course of the proof of Theorem 1.2 that $f(t, t, t)$ is decreasing in t . In the present case $t = 1 - 1/m$ where $m \geq 4$. Moreover, since $f(1 - 1/4, 1 - 1/4, 1 - 1/4)$ is expressed in Table 1 (for the ramification indices $(4, 4, 4)$) this concludes the proof (the fact that ϵ_m is decreasing can be shown by elementary methods).

10.3. Comparison with Parshin's inequality in the geometric case. Let \mathcal{B} be a complex projective curve of genus $g_{\mathcal{B}}$ and \mathcal{X} a complex projective surface with a morphism $\mathcal{X} \rightarrow \mathcal{B}$ such that the relative canonical line bundle $\mathcal{K}_{\mathcal{X}/\mathcal{B}}$ is relatively ample. Assume that \mathcal{X}, \mathcal{B} and the generic fiber X of \mathcal{B} are regular and denote by s the number of singular fibers of $\mathcal{X} \rightarrow \mathcal{B}$. By [88, 99], the following geometric analog of Parshin's proposed arithmetic inequality 1.2 holds:

$$(10.3) \quad \hat{h}(\mathcal{K}_{\mathcal{X}/\mathcal{B}}) := \frac{\mathcal{K}_{\mathcal{X}/\mathcal{B}} \cdot \mathcal{K}_{\mathcal{X}/\mathcal{B}}}{2 \deg(K_X)} \leq \max(0, g_{\mathcal{B}} - 1) + \frac{1}{2}s,$$

when $\mathcal{X} \rightarrow \mathcal{B}$ is *(semi)-stable*. This is a consequence of the Miyaoka–Yau inequality for \mathcal{X} . Moreover, by [97],

$$(10.4) \quad \hat{h}(\mathcal{K}_{\mathcal{X}/\mathcal{B}}) \leq \max(0, g_{\mathcal{B}} - 1) + \frac{3}{2}s,$$

when $\mathcal{X} \rightarrow \mathcal{B}$ is merely *relatively minimal*.

In the arithmetic case the role of $g_{\mathcal{B}}$ is played by $\log |D_{\mathbb{F}}|$ and the role of s is played by $\sum_{\mathfrak{p} \text{ bad}} \log N(\mathfrak{p})$ (see the discussion in [99]). In particular, the case $g_{\mathcal{B}} = 0$ corresponds to the case when $\mathbb{F} = \mathbb{Q}$. Coming back to case of the Zariski closure $\mathcal{X}^{(m)}$ in $\mathbb{P}_{\mathbb{Z}}^2$ of the Fermat curve $X^{(m)}$ over \mathbb{Q} of degree m , recall that, by Cor 1.7,

$$\hat{h}_{\text{can}}(\overline{\mathcal{K}_{\mathcal{X}^{(m)}}}) < 0 + \log m, \quad \hat{h}_{\text{Ar}}(\overline{\mathcal{K}_{\mathcal{X}^{(m)}}}) < 0 + 2 \log m.$$

As a consequence, the corresponding inequalities also hold when $\mathcal{X}^{(m)}$ is replaced by a stable model or a relatively minimal model (By Cor 3.10 and Prop 3.18). Now specialize to the case when m is square-free. Then the role of $\log m$ is played by s in the geometric case. Thus, the inequality for $\hat{h}_{\text{can}}(\overline{\mathcal{K}_{\mathcal{X}_{\min}^{(m)}}})$ is actually better than the inequality one would obtain from the geometric inequality 10.4, would it translate to the arithmetic setup (since $1 < 3/2$).

In view of the previous discussion it seems natural to ask if, in general, the direct analog of the geometric inequality 10.3 holds for the volume-normalized Kähler-Einstein metric on K_X ? This would imply Parshin's inequality 1.2 for the Arakelov metric on K_X with explicit constants (using the inequality 10.2). For example, when $\mathbb{Q} = \mathbb{F}$ one would get $c_1 = 1/2, c_2 = 0$ and c_0 explicitly bounded from above by $\log \deg K_X$. More generally, consider a projective regular curve X over a number field \mathbb{F} , endowed with a divisor Δ such that $K_{(X, \Delta)} > 0$ and fix a finite field extension \mathbb{F}' of \mathbb{F} such that $(X, \Delta)_{\mathbb{F}} \otimes \mathbb{F}'$ admits a relatively stable model $(\mathcal{X}, \mathcal{D})$ over $\mathcal{O}_{\mathbb{F}'}$ (as discussed before the statement of Lemma 3.13). Does the following inequality hold,

$$\hat{h}(\mathcal{K}_{(\mathcal{X}, \mathcal{D})}) \leq \max(0, \log |D_{\mathbb{F}}| - 1) + \frac{1}{2} \sum_{\mathfrak{p} \text{ bad}} \log N(\mathfrak{p})?$$

The case when $\mathbb{F} = \mathbb{F}' = \mathbb{Q}$, $X = \mathbb{P}_{\mathbb{Q}}^1$ and Δ has three irreducible components follows from the second inequality in Theorem 7.1 (there are no bad \mathfrak{p} in this case).

REFERENCES

- [1] Dan Abramovich. Birational geometry for number theorists. In Arithmetic geometry, volume 8 of Clay Math. Proc., pages 335–373. Amer. Math. Soc., Providence, RI, 2009.
- [2] R.Andreasson; R J. Berman: Sharp bounds on the height of K-semistable Fano varieties I, the toric case. arXiv:2205.00730 (2022)
- [3] R.Andreasson; R.J. Berman: Sharp bounds on the height of K-semistable Fano varieties II, The log case. arXiv:2312.05064, 2023 - arxiv.org
- [4] R. J. Berman: A thermodynamical formalism for Monge-Ampère equations, Moser-Trudinger inequalities and Kähler-Einstein metrics. Adv. Math. 248 (2013), 1254-1297
- [5] Berman, R. J.: Large deviations for Gibbs measures with singular Hamiltonians and emergence of Kähler-Einstein metrics. Comm. Math. Phys. 354, 1133{1172 (2017).
- [6] Berman, R.J: Kähler-Einstein metrics, canonical random point processes and birational geometry. In Proceedings of Symposia in Pure Mathematics. Volume 97.1 : Algebraic Geometry Salt Lake City 2015 (Part 1). pp 29-74
- [7] Berman, R. J., An invitation to Kähler-Einstein metrics and random point processes. In Surveys in Differential Geometry Volume 23 (2018) Pages: 35 – 87
- [8] Berman, R.J: K-polystability of Q-Fano varieties admitting Kähler-Einstein metrics. Inventiones Math. March 2016, Volume 203, Issue 3, pp 973-1025

- [9] Berman, R.J: Kähler-Einstein metrics and Archimedean zeta functions. European Congress of Mathematics Portorož, 20–26 June, 2021 pp. 199–251. EMS press (2023)
- [10] Berman, R.J.; Boucksom, S: Growth of balls of holomorphic sections and energy at equilibrium. *Invent. Math.* Vol. 181, Issue 2 (2010), p. 337
- [11] R. J. Berman, S. Boucksom, V. Guedj and A. Zeriahi, A variational approach to complex Monge–Ampère equations, *Publ. Math. Inst. Hautes Études Sci.* 117 (2013), 179–245.
- [12] R. J. Berman, S. Boucksom, and M. Jonsson, A variational approach to the Yau-Tian-Donaldson conjecture, *J. Amer. Math. Soc.* 34 no. 3 (2021), 605–652
- [13] R. J. Berman; T. Darvas; C.H. Lu: Convexity of the extended K-energy and the large time behavior of the weak Calabi flow. *Geom. and Topol.* 21(2017), no. 5, 2945–2988
- [14] R J Berman, H Guenancia: Kähler–Einstein metrics on stable varieties and log canonical pairs, *Geom. Funct. Anal.* 24 (2014) 1683–1730
- [15] Berman, R.J; Freixas i Montplet: G: An arithmetic Hilbert-Samuel theorem for singular hermitian line bundles and cusp forms. *Compositio Mathematica.* Vol. 150/Issue 10 (2014)
- [16] R.J.Berman; Eyssidieux, P: S. Boucksom, V. Guedj, A. Zeriahi: Kähler-Einstein metrics and the Kähler-Ricci flow on log Fano varieties. *Journal für die Reine und Angewandte Mathematik* (published on-line 2016).
- [17] R.J. Berman: On K-stability, height bounds and the Manin-Peyre conjecture. Preprint at <https://arxiv.org/abs/2305.07272>.
- [18] C. Birkar, P. Cascini, C. Hacon and J. McKernan; Existence of minimal models for varieties of log general type, *J. Amer. Math. Soc.* 23 (2010), no. 2, 405–468.
- [19] Bost, J.-B., Mestre, J.-F., Moret-Bailly, L.: Sur le calcul explicite des “classes de Chern” des surfaces arithmétiques de genre 2. *Astérisque*, no. 183, pp. 69–105 (1990). *Séminaire sur les Pinceaux de Courbes Elliptiques* (Paris, 1988)
- [20] Bost, J.-B.; Gillet, H.; Soulé, C.: Heights of projective varieties and positive Green forms. *J. Amer. Math. Soc.* 7 (1994), no. 4, 903–102
- [21] Bost, J.-B.: Intrinsic heights of stable varieties and abelian varieties. *Duke Math. J.* 82 (1996), no. 1, 21–70
- [22] J.-B. Bost: Potential theory and Lefschetz theorems for arithmetic surfaces. *Ann. Sci. École Norm. Sup.* (4) 32 (1999), no. 2, 241–312.
- [23] S.Boucksom, P.Eyssidieux, V.Guedj; A.Zeriahi: Monge- Ampere equations in big cohomology classes, *Acta Math.* 205 (2010), no. 2, 199–262
- [24] S.Boucksom, T.Hisamoto, M.Jonsson: Uniform K-stability, Duistermaat-Heckman measures and singularities of pairs, *Ann. Inst. Fourier (Grenoble)* 67 (2017), no. 2, 743–841.
- [25] J.-F. Boutot and H. Carayol. Uniformisation p-adique des courbes de Shimura: les théorèmes de Cerednik et de Drinfeld. *Courbes modulaires et courbes de Shimura*, *Astérisque*, tome 196–197, 1991.
- [26] JF Boutot, T Zink: On the p-adic uniformization of quaternionic Shimura curves. arXiv preprint [arXiv:2212.06886](https://arxiv.org/abs/2212.06886), 2022 - arxiv.org
- [27] Bruinier, J.H., Burgos Gil, J.-I., Kühn, U.: Borcherds Products and Arithmetic Intersection Theory on Hilbert Modular Surfaces. *Duke Math J.* 139(1), 1–88 (2007)
- [28] I. Burgos Gil, J. Kramer, and U. Kuhn, Cohomological arithmetic Chow rings, *J. Inst. Math. Jussieu* 6 (2007), 1–172.
- [29] F. Campana: Special manifolds, arithmetic and hyperbolic aspects: a short survey, *Rational points, rational curves, and entire holomorphic curves on projective varieties*, 2015, pp. 23–52
- [30] J.Cassaigne;V.Maillot. Hauteur des hypersurfaces et fonctions zeta d’Igusa. *Journal of Number Theory*, 83:226–255, 2000.
- [31] T. C. Chau; A note concerning Fox’s paper on Fenchel’s conjecture, *Proc. Amer. Math. Soc.* 88 (1983), no. 4, 584–586.
- [32] Choi, J; Srivastava, H.M.: A family of log-gamma integrals and associated results, *Journal 70 of Mathematical Analysis and Applications*, 303 (2005) 436–449, Science Direct, Elsevier.

- [33] Coman, D., Guedj, V., Zeriahi, A.: Extension of plurisubharmonic functions with growth control. *J. Reine Angew. Math.* 676, 33–49 (2013)
- [34] C. Curilla; U. Kühn. On the arithmetic self-intersection numbers of the dualizing sheaf for Fermat curves of prime exponent. arXiv:0906.3891.
- [35] Christian Curilla, J. Steffen Müller: The minimal regular model of a Fermat curve of odd square-free exponent and its dualizing sheaf. *Kyoto J. Math.* 60, no. 1 (2020), 219–268
- [36] Darvas, T., Rubinstein, Y.: Tian’s properness conjectures and Finsler geometry of the space of Kähler metrics. *J. Am. Math. Soc.* 30, 347–387 (2017)
- [37] Deligne, P: Preuve des conjectures de Tate et de Shafarevitch. Séminaire Bourbaki : volume 1983/84, exposés 615–632, Astérisque, no. 121–122 (1985), Talk no. 616, 17 p.
- [38] Deligne, P; Mumford, D.: The irreducibility of the space of curves of given genus, *Publications Mathématiques I.H.E.S.*, 36 (1969), 75–109
- [39] P. Deligne, M. Rapoport, Les schémas de modules de courbes elliptiques. *Modular Functions of One Variable II*, Proc. Internat. Summer School, Univ. Antwerp 1972, Lect. Notes Math. 349 (1973), 143–316
- [40] S.K. Donaldson Kähler metrics with cone singularities along a divisor. *Essays in mathematics and its applications*, Springer, Heidelberg, 2012, pp. 49–79,
- [41] V. S. Dotsenko and V. A. Fateev: Four-point correlation functions and the operator algebra in 2D conformal invariant theories with central charge $C \leq 1$. *Nucl. Phys.B* 251(1985)
- [42] Elkies, N.D.: Shimura curve computations. Pages 1–47 in *Algorithmic Number Theory (Proceedings of ANTS-III; J. P. Buhler, ed.; Berlin: Springer, 1998; Lecture Notes in Computer Science 1423)*.
- [43] Faltings, G: Endlichkeitssätze für abelsche Varietäten über Zahlkörpern, *Invent. Math.* (1983)
- [44] Faltings, G: Calculus on arithmetic surfaces, *Ann. of Math.* (2), 119 (1984), 387–424.
- [45] Faltings, G.: Diophantine approximation on abelian varieties, *Ann. of Math.* 119’ (1991) 549– 576
- [46] Freixas i Montplet, G: Heights and metrics with logarithmic singularities. *Journal für die Reine und Angewandte Mathematik*, 627 (2009), 97–153.
- [47] Freixas i Montplet, G., von Pippich, A.: Riemann-Roch isometries in the non-compact orbifold setting. *J. Eur. Math. Soc.* (2018). <https://doi.org/10.4171/JEMS/992>
- [48] Z.Fu; Y.Zhu: Selberg integral over local fields. *Forum Math.* 2019; 31(5): 1085–1095
- [49] Fujita, Kento .J; Odaka, Y: On the K-stability of Fano varieties and anticanonical divisors. *Tohoku Math. J.* (2) 70 (2018), no. 4, 511–521.
- [50] K. Fujita. Optimal bounds for the volumes of Kähler-Einstein Fano Manifolds. *Amer. J. Math.* 140 (2018), 391–414
- [51] K.Fujita. K-stability of log Fano hyperplane arrangements. *J. Algebraic Geom.*, 30(4):603– 630, 2021.
- [52] Gillet, H.; Soulé, C: Arithmetic intersection theory, *Inst. Hautes Etudes Sci. Publ. Math.* 72 (1990), 94–174.
- [53] H. Gillet and C. Soulé: An arithmetic Riemann-Roch theorem, *Invent. Math.* 110 (1992), 473–54
- [54] H. Guenancia: Kähler-Einstein metrics with mixed Poincaré and cone singularities along a normal crossing divisor”, *Ann. Inst. Fourier* 64 (2014), no. 6, p. 1291–1330.
- [55] Guenancia, H; Păun, M: Conic singularities metrics with prescribed Ricci curvature: general cone angles along normal crossing divisors. *J. Differential Geom.* 103 (2016), no. 1, 15–57
- [56] H. Guenancia; D. Wu: On the boundary behavior of Kähler-Einstein metrics on log canonical pairs. *Math. Annalen* 366 (2016), no. 1, p. 101–120
- [57] Hacon, C.D., Xu, C. Existence of log canonical closures. *Invent. math.* 192, 161–195 (2013).
- [58] B. Hassett, Moduli spaces of weighted pointed stable curves, *Adv. Math.* 173 (2003), no. 2, 316–352
- [59] M. Hattori, Y. Odaka: Minimization of Arakelov K-energy for many cases. Preprint at <https://arxiv.org/abs/2211.03415>.
- [60] M.Hill;T.Lawson, Automorphic forms and cohomology theories on Shimura curves of small discriminant, *Adv. Math.* 225 (2010), no. 2, 1013–1045.

- [61] Javanpeykar, A.: Polynomial bounds for Arakelov invariants of Belyi curves. With an appendix by Peter Bruin. *Algebra Number Theory* 8(1), 89–140 (2014)
- [62] J. Jorgenson and J. Kramer, Bounds for special values of Selberg zeta functions of Riemann surfaces, *J. Reine Angew. Math.* 541 (2001)
- [63] J. Jorgenson, J. Kramer, Expressing Arakelov invariants using hyperbolic heat kernels. In: *The Ubiquitous Heat Kernel*. J. Jorgenson, L. Walling (eds.), *Contemp. Math.* AMS 398 (2006), 295–309.
- [64] Jeffres, T; Mazzeo, R; Rubinstein, Y.A.; Kähler-Einstein metrics with edge singularities. *Ann. of Math.* (2) 183 (2016), no. 1, 95–176.
- [65] J. Kollár, Moduli of varieties of general type, *Handbook of moduli*. Vol. II, *Adv. Lect. Math.* (ALM), vol. 25, Int. Press, Somerville, MA, 2013, pp. 131–157.
- [66] J. Kollár, S. Mori, *Birational geometry of algebraic varieties*. With the collaboration of C. H. Clemens and A. Corti. Translated from the 1998 Japanese original. *Cambridge Tracts in Mathematics*, 134. Cambridge University Press, Cambridge, 1998.
- [67] Stephen S. Kudla: Special cycles and derivatives of Eisenstein series, *Heegner Points and Rankin L-Series*, *Math. Sci. Res. Inst. Publ.*, vol. 49, Cambridge University Press, Cambridge, 2004, pp. 243–270.
- [68] Stephen S. Kudla, Michael Rapoport, and Tonghai Yang, *Modular forms and special cycles on Shimura curves*, *Annals of Mathematics Studies*, vol. 161, Princeton University Press, Princeton, NJ, 2006
- [69] C. Kaiser; K. Köhler : A fixed point formula of Lefschetz type in Arakelov geometry III: representations of Chevalley schemes and heights of flag varieties, *Invent. Math.* 147 (2002), 633–669.
- [70] A.Kirillov Jr; *Quiver Representations and Quiver Varieties*. AMS. 2016, Graduate studies in Math. Vol 174
- [71] J. Kollár: *Singularities of the minimal model program*, *Cambridge Tracts in Mathematics*, vol. 200, Cambridge University Press, Cambridge, 2013, With the collaboration of Sándor Kovács.
- [72] Ulf Kuhn, Generalized arithmetic intersection numbers, *J. Reine Angew. Math.* 534 (2001), 209–236
- [73] U. Kuhn: On the arithmetic self-intersection number of the dualizing sheaf on arithmetic surfaces, 2009, <http://arxiv.org/abs/0906.2056>.
- [74] C.Kurkoglu: Rational Exceptional Belyi Coverings. <https://arxiv.org/abs/2404.14675>
- [75] Li, C: G-uniform stability and Kähler-Einstein metrics on Fano varieties. *Invent. Math.* 227 (2022), no. 2, 661–744.
- [76] Liu, Q; and R.Erne: *Algebraic Geometry and Arithmetic Curves*, Oxford University Press, Incorporated, 2002.
- [77] Y Liu, C Xu, Z Zhuang: Finite generation for valuations computing stability thresholds and applications to K-stability. *Annals of Mathematics* (2), 196(2):507–566, 2022
- [78] Maillot, V.: Un calcul de Schubert arithmétique. *Duke Math. J.* 80, 195–221 (1995)
- [79] Vincent Maillot and Damien Roessler, Conjectures sur les dérivées logarithmiques des fonctions L d’Artin aux entiers négatifs, *Math. Res. Lett.* 9 (2002), no. 5-6, 715–72
- [80] V Maillot, D Rössler: Conjectures on the logarithmic derivatives of Artin L-functions II. To appear in the proceedings of the summer school "Motives and Complex Multiplication" (14-19 August 2016, Monte Verita, Ascona, Switzerland, edited by Javier Fréсан and Peter Jossen). arXiv preprint arXiv:1808.03068
- [81] Y. Manin, *New dimensions in geometry*, Workshop at Bonn 1984 (Bonn, 1984), 59-101. *Lecture Notes in Mathematics* vol. 1111, Springer (1985)
- [82] R.Mazzeo; Y.A.Rubinstein: The Ricci continuity method for the complex Monge–Ampère equation, with applications to Kähler–Einstein edge metrics. *Comptes Rendus Mathématique* Volume 350, Issues 13–14, July 2012, Pages 693–697
- [83] L. Moret-Bailly, *Hauteurs et classes de Chern sur les surfaces arithmétiques*, *Asterisque* 183 (1990), 37–58.

- [84] Mustata, M: Zeta functions in algebraic geometry. Lecture notes available at <https://public.websites.umich.edu/~mmustata/>.
- [85] Odaka, Y: The GIT stability of polarized varieties via discrepancy. *Ann. of Math.* Pages 645-661, Vol. 177 (2013), Issue 2
- [86] Odaka, Y: Canonical Kähler metrics and arithmetics: Generalizing Faltings heights. *Kyoto J. Math.* 58(2) (2018) 243-288
- [87] Y.Odaka, S.Sun, Testing Log K-stability by blowing up formalism, *Annales de la Faculté des sciences de Toulouse : Mathématiques, Série 6, Tome 24* (2015) no. 3, p. 505-522
- [88] Parshin, A.N.: The Bogomolov-Yau-Miyaoka inequality for arithmetical surfaces and its applications. In *Seminaire de theorie des nombres de Paris. Progr. Maths.* 75 1988 Basel Boston: Birkhauser, pp. 299 312
- [89] Siegel, C. L.: Berechnung von Zetafunktionen an ganzzahligen Stellen. *Nachr. Akad. Wiss. Göttingen, Math.-Phys. Klasse 10* (1969), pp. 87-102.
- [90] Song, Y; Tian, G: Canonical measures and Kähler-Ricci flow. *J. Amer. Math. Soc.* 25 (2012), no. 2, 303-353.
- [91] Tsuji,H: Canonical measures and the dynamical systems of Bergman kernels. Preprint arXiv:0805.1829, 2008
- [92] Takeuchi, K: Arithmetic triangle groups, *J. Math. Soc. Japan* 29 (1977), 91-10
- [93] Takeuchi, K.: Commensurability classes of arithmetic triangle groups, *J. Fac. Sci. Univ. Tokyo* 24 (1977), 201-212
- [94] H. Tamvakis : Schubert calculus on the arithmetic Grassmannian, *Duke Math. J.* 98 (1999), 421-443.
- [95] H. Tamvakis : Arakelov theory of the Lagrangian Grassmannian, *J. reine angew. Math.* 516 (1999),207-223.
- [96] H. Tamvakis : Height formulas for homogeneous varieties, *Michigan Math. J.* 48 (2000), 593-610.
- [97] Tan, SL. On the invariants of base changes of pencils of curves, II. *Math Z* 222, 655-676 (1996)
- [98] H. Tanaka, Minimal model program for excellent surfaces, *Ann. Inst. Fourier (Grenoble)*, 68, no. 1, 345-376, (2018).
- [99] Vojta, P: Diophantine inequalities and Arakelov theory In: S. Lang, *Introduction to Arakelov Theory*, Springer, 1988, pp. 155-178.
- [100] Xu, C: K-stability of Fano varieties: an algebro-geometric approach. *EMS Surv. Math. Sci.* 8 (2021), no. 1-2, 265-354.
- [101] X. Yuan, S. Zhang, W. Zhang, *The Gross-Zagier formula on Shimura curves*, *Annals of Math. Studies*, No. 184, Princeton University Press, 2012
- [102] X. Yuan, S. Zhang, On the averaged Colmez conjecture, *Ann. of Math.* (2) 187 (2018), no. 2, 533-638.
- [103] X. Yuan: Modular heights of quaternionic Shimura curves. preprint arXiv:2205.13995 (2022)
- [104] X. Yuan: Explicit Kodaira-Spencer map over Shimura Curves. *Acta Math. Sinica (Chinese Ser.)* 67 (2024), no. 2, 227-249. arXiv: 2205.11334
- [105] Zhang, S: Positive line bundles on arithmetic varieties. *J. Amer. Math. Soc.* 8 (1995), 187-221
- [106] S. Zhang, Small points and adelic metrics, *J. Alg. Geometry* 4 (1995), 281-300
- [107] S. Zhang: Heights and reductions of semi-stable varieties, *Compositio Mathematica*. Vol 104, no. 1 (1996), p.77-105

Email address: rolfan@chalmers.se, robertb@chalmers.se