

ON MULTIDIMENSIONAL BOHR RADII FOR BANACH SPACES

VASUDEVARAO ALLU AND SUBHADIP PAL

ABSTRACT. In this paper, we study a more general version of multidimensional Bohr radii for the holomorphic functions defined on unit ball of ℓ_q^n ($1 \leq q \leq \infty$) spaces with values in arbitrary complex Banach spaces. More precisely, we study the multidimensional Bohr radii for bounded linear operators between complex Banach spaces, primarily motivated by the work of A. Defant, M. Maestre, and U. Schwaning [Adv. Math. 231 (2012), pp. 2837–2857]. We obtain the exact asymptotic estimates of multidimensional Bohr radius for both finite and infinite dimensional Banach spaces. As an application, we find the lower bound of arithmetic Bohr radius.

1. INTRODUCTION

In the beginning of the 20th century while studying the well-known Riemann ζ -function, a Danish mathematician Harald Bohr made significant developments to the general theory of Dirichlet series. A Dirichlet series is an summation expression of the form

$$(1.1) \quad D(s) := \sum_{n \geq 1} \frac{a_n}{n^s},$$

where $a_n \in \mathbb{C}$ and $s = \sigma + it$ is a complex variable. It is well-known that the region of convergence, absolute convergence, and uniform convergence of a Dirichlet series define to be the half planes of the form $\{s \in \mathbb{C} : \operatorname{Re}(s) > \sigma_0\}$ in the complex plane \mathbb{C} . Bohr was mainly interested in tackling the region of convergence of this Dirichlet series. More precisely, Bohr was investigating the width of the greatest strip for which a Dirichlet series converges uniformly but not absolutely. This problem is known as *absolute convergence problem* of Bohr in the theory of Dirichlet series. In order to tackle this problem, Bohr has obtained a nice connection between a Dirichlet series and a power series of infinite variables. Indeed, given a Dirichlet series of the form (1.1), Bohr considered for each $n \in \mathbb{N}$ the prime decomposition of $n = p_1^{\alpha_1} \cdots p_r^{\alpha_r}$ (where $(p_m)_{m \in \mathbb{N}}$ denotes the sequence of order prime numbers) and defined $z = (p_1^{-s}, \dots, p_r^{-s})$. Therefore, Bohr has established the following identity:

$$(1.2) \quad D(s) = \sum_{n \geq 1} a_n (p_1^{-s})^{\alpha_1} \cdots (p_r^{-s})^{\alpha_r} = \sum a_n z_1^{\alpha_1} \cdots z_r^{\alpha_r}.$$

The above correspondence (1.2) is known as the *Bohr transform*. It helps us to translate the problem on Dirichlet series in terms of power series and thus, one can apply complex analysis techniques to solve this problem. As a part of these studies, Bohr once asked whether is it possible to compare the absolute value of a power series in one complex variable with the sum of the absolute values of its coefficients. In 1914, Bohr [12] proved the following remarkable result which nowadays known as *Bohr's inequality*:

File: AP-Bohr-1-2024.tex, printed: 2024-7-1, 0.31

2020 *Mathematics Subject Classification.* Primary 32A05, 32A10, 46B07; Secondary 46B09, 46G20.

Key words and phrases. Bohr radius, Power series, Holomorphic functions, Homogeneous polynomials, Banach spaces.

Theorem A. [12] If $f(z) = \sum_{n=0}^{\infty} a_n z^n$ is a holomorphic function in unit disk such that $\sup_{z \in \mathbb{D}} |f(z)| < \infty$, then

$$(1.3) \quad \sum_{n=0}^{\infty} |a_n| r^n \leq \sup_{z \in \mathbb{D}} |f(z)|$$

for $|z| = r \leq 1/3$, and the constant $1/3$, referred to as the Bohr radius, which is sharp.

In fact, Bohr himself obtained the radius as $1/6$. Later, this radius was further improved to $1/3$ by M. Riesz, I. Schur, and N. Wiener [37, 38], independently. This significant inequality was forgotten for long eighty years until Dixon [27] used the inequality in his paper to disprove a long-standing conjecture in Operator Algebras: If a Banach algebra satisfies the von Neumann inequality then it is necessarily an operator algebra, *i.e.*, which is isometrically isomorphic to a closed subalgebra of $\mathcal{L}(H)$, the algebra of all bounded operators for some Hilbert space H . After that, several researchers like Aizenberg [1, 2, 3], Boas and Khavinson [10], Boas [11], Bombieri and Bourgain [15], Galicer, Mansilla and Muro [30], Hamada, Honda and Kohr [31], Paulsen and Singh [35], Popescu [36] have paid much attention to this result to use it in different contexts of mathematics and generalized it. In this paper, we mostly focus on the unit ball of Minkowski space ℓ_q^n , where $q \in [1, \infty]$. For $q \in [1, \infty)$, we denote

$$B_{\ell_q^n} = \left\{ z \in \mathbb{C}^n : \|z\|_q = \left(\sum_{i=1}^n |z_i|^q \right)^{1/q} < 1 \right\},$$

and

$$B_{\ell_\infty^n} = \left\{ z \in \mathbb{C}^n : \|z\|_\infty = \sup_{1 \leq i \leq n} |z_i| < 1 \right\}.$$

Before proceeding for further discussions, let us first introduce the following notion of multidimensional Bohr radius for holomorphic functions defined on $B_{\ell_q^n}$, $1 \leq q \leq \infty$ with values in a complex Banach space X , primarily motivated by the work of Defant *et. al.* [25].

Definition 1.1. Let $1 \leq q \leq \infty$, $\lambda \geq 1$, $n \in \mathbb{N}$, and let $T : X \rightarrow Y$ be a bounded linear operator between complex Banach spaces X and Y such that $\|T\| \leq \lambda^{1/p}$. For $1 \leq p < \infty$, the λ_p -Bohr radius of T , denoted by $K^p(B_{\ell_q^n}, T, \lambda)$, is the supremum of all $r \geq 0$ such that for all holomorphic functions $f(z) = \sum_{\alpha \in \mathbb{N}_0^n} x_\alpha z^\alpha$ on $B_{\ell_q^n}$ we have

$$(1.4) \quad \sup_{z \in r B_{\ell_q^n}} \sum_{\alpha \in \mathbb{N}_0^n} \|T(x_\alpha) z^\alpha\|_Y^p \leq \lambda \left\| \sum_{\alpha \in \mathbb{N}_0^n} x_\alpha z^\alpha \right\|_X^p.$$

We write $K^p(B_{\ell_q^n}, T)$ for $\lambda = 1$. If T is the identity map on X then we use the notation $K^p(B_{\ell_q^n}, X, \lambda)$ and $K^p(B_{\ell_q^n}, X)$. For $X = \mathbb{C}$, we write $K^p(B_{\ell_q^n}, \lambda)$ for $K^p(B_{\ell_q^n}, \mathbb{C}, \lambda)$ and $K(B_{\ell_q^n})$ whenever $p = 1$ and $\lambda = 1$.

We observe that if f is an unbounded holomorphic function on $B_{\ell_q^n}$ then the inequality (1.4) holds trivially. Therefore, throughout our discussion, we assume the holomorphic functions to be bounded on $B_{\ell_q^n}$. Clearly, with the above notation, Theorem A reduces to $K(\mathbb{D}) = 1/3$.

In the classical sense, Bombieri and Bourgain [14, 15] have studied the constant $K(\mathbb{D}, \lambda)$ whenever $\lambda \geq 1$. In fact, Bombieri [14] has obtained the exact value

$$K(\mathbb{D}, \lambda) = \frac{1}{3\lambda - 2\sqrt{2(\lambda^2 - 1)}},$$

where $\lambda \in [1, \sqrt{2}]$, and Bombieri and Bourgain [15] have determined the exact asymptotic behavior of $K(\mathbb{D}, \lambda)$ as $\sqrt{\lambda^2 - 1}/\lambda$ whenever $\lambda \rightarrow \infty$. In 2010, Blasco [8] established that $K(\ell_q^2) = 0$ for $1 \leq q \leq \infty$. Thus, the study of Bohr radius problem becomes irrelevant for $\dim(X) > 1$. Hence, it seems reasonable to modify the Bohr inequality to obtain more interesting result. That is why the authors [25] have introduced λ in the definition of Bohr radius $K(B_{\ell_\infty^n}, T, \lambda)$ of bounded linear operator T . In [25], the authors have established that $K(B_{\ell_\infty^n}, T, \lambda) > 0$ for every bounded linear operator $T : X \rightarrow Y$ between complex Banach spaces X and Y . Later, in this direction, the work was extended by Kumar *et al.* [32] and studied the Bohr radius $K(B_{\ell_q^n}, T, \lambda)$ for any bounded operator T , $\lambda \geq 1$, and $1 \leq q \leq \infty$.

In 1989, Dineen and Timoney [29] extensively studied the constant $K(B_{\ell_\infty^n})$ and later, their result has been quickly refined in [10]. In 1997, Boas and Khavinson [10] proved the following estimate for each $n \in \mathbb{N}$ with $n \geq 2$,

$$\frac{1}{3\sqrt{n}} \leq K(B_{\ell_\infty^n}) \leq 2\sqrt{\frac{\log n}{n}}.$$

In view of results of Aizenberg, Boas, Dineen, Khavinson and Timoney (see [1],[10], [11], [29]), estimating the multidimensional Bohr radii, for every $1 \leq q \leq \infty$ and all n we have

$$\frac{1}{C} \left(\frac{1}{n}\right)^{1 - \frac{1}{\min\{q, 2\}}} \leq K(B_{\ell_q^n}) \leq C \left(\frac{\log n}{n}\right)^{1 - \frac{1}{\min\{q, 2\}}},$$

where C is a constant independent of q and n . One of the main reasons behind the occurrence of log-term in the upper estimates of all known non-trivial results on multidimensional Bohr radii is the probabilistic methods used by the authors. In [11, p. 326] Boas conjectured that *“This logarithmic factor, an artifact of the proof, presumably should not really be present”*. In 2006, Dafant and Frerick [18] disproved the conjecture by obtaining a logarithmic lower bound which is almost correct asymptotic estimate for the Bohr radius $K(B_{\ell_q^n})$ $1 \leq q \leq \infty$. Using the methods developed in [21], Defant and Frerick [18] have proved that there is a constant $C > 0$ such that for every $1 \leq q \leq \infty$ and n we have

$$(1.5) \quad \frac{1}{C} \left(\frac{\log n / \log \log n}{n}\right)^{1 - \frac{1}{\min\{q, 2\}}} \leq K(B_{\ell_q^n}).$$

Later, (1.5) was improved by the same authors in [19]. The case for $q = \infty$ was settled by Defant and his co-authors [20] and they showed that $K(B_{\ell_\infty^n})$ behaves asymptotically as $\sqrt{\log n/n}$ as a consequence of hypercontractivity of Bohnenblust-Hille inequality. Further, Bayart *et al.* [5] have extended the techniques and obtained the following:

$$\lim_{n \rightarrow \infty} \frac{K(B_{\ell_\infty^n})}{\sqrt{\frac{\log n}{n}}} = 1.$$

The powered versions of the Bohr radii $K^p(\mathbb{D})$ and $K^p(B_{\ell_\infty^n})$ for scalar-valued holomorphic functions were first studied by Djakov and Ramanujan [28]. Later, Bénéteau *et al.* [7]

explored the analogous radii in the context of Hardy spaces. Apart from the scalar valued functions, Blasco [9] has investigated the radius $K^p(\mathbb{D}, X)$ for arbitrary complex Banach spaces. In fact, Blasco [9, Theorem 1.10] has proved that the Bohr radius $K^p(\mathbb{D}, X) > 0$ if, and only if, X is p -uniformly \mathbb{C} -convex, where $2 \leq p < \infty$. Recently, Das [16, 17] has showed the precise asymptotic value of $K^p(B_{\ell_q^n})$ and investigated p -Bohr radius for Hardy space of X -valued holomorphic functions defined on complete Reinhardt domain.

To explore more general aspects of multidimensional Bohr radii, we introduce λ in our Definition 1.1, in more general sense. Indeed, we shall prove that for every $1 \leq p < \infty$ and bounded linear operator T with $\|T\| < \lambda^{1/p}$, the multidimensional Bohr radius $K^p(B_{\ell_q^n}, T, \lambda)$ is nonzero. Further, our aim is to study the correct asymptotic estimates for the Bohr radii $K^p(B_{\ell_q^n}, T, \lambda)$ for arbitrary Banach spaces. As a consequence, we obtain lower bound of arithmetic Bohr radius for Banach spaces. The organization of this paper is as follows. After the preliminaries, Section 3 accumulates all the main results of this paper. In this section, Proposition 3.1 shows the for every bounded operator T with $\|T\| < \lambda^{1/p}$ the Bohr radius $K^p(B_{\ell_q^n}, T, \lambda)$ is positive. Further, Theorem 3.1 and Theorem 3.2 give the asymptotic behavior of $K^p(B_{\ell_q^n}, X, \lambda)$ for finite and infinite dimensional Banach space X respectively, in case of identity operator on X . Finally, in Theorem 3.4, we obtain lower estimates for $K^p(B_{\ell_q^n}, T, \lambda)$. Section 4 keeps two important lemmas which play vital role to prove the main results. Section 5 contains the proof of main results.

2. PRELIMINARIES

We employ standard terminologies and notions from classical theory of Banach spaces (see e.g. [33, 34]). We denote \mathbb{D} to be the open unit disk in complex plane \mathbb{C} . Throughout the paper, it has been assumed that every Banach space X is complex. X^* will denote the topological dual of Banach space X and we write B_X for the open unit ball in X . As usual, we denote ℓ_q^n for the Banach space of all n -tuples $z = (z_1, \dots, z_n) \in \mathbb{C}^n$ endowed with the norm

$$\|z\|_q := \left(\sum_{i=1}^n |z_i|^q \right)^{\frac{1}{q}} \quad \text{for } 1 \leq q < \infty$$

and $\|z\|_\infty := \max_{1 \leq i \leq n} |z_i|$ for $q = \infty$. For $1 \leq q \leq \infty$, the conjugate exponent q' for q is defined by $1/q + 1/q' = 1$. For $x \in X$, the absolute value of x is defined by $|x| = x \vee (-x)$. Recall that a Banach lattice is a Banach space X which is a vector lattice with $|x| \leq |y|$ implies $\|x\| \leq \|y\|$ for all $x, y \in X$. A Banach lattice is said to be r -concave, where $1 \leq r < \infty$, if there is a constant $M > 0$ such that

$$(2.1) \quad \left(\sum_{j=1}^n \|x_j\|^r \right)^{\frac{1}{r}} \leq M \left\| \left(\sum_{j=1}^n |x_j|^r \right)^{\frac{1}{r}} \right\|$$

for every arbitrarily chosen finite elements $x_1, \dots, x_n \in X$. The best such constant satisfying (2.1) is as usual denoted by $M_r(X)$. For $2 \leq r \leq \infty$, a Banach space X is said to have cotype r , if there is a constant $C > 0$ such that for every choice of finitely many vectors $x_1, \dots, x_n \in X$, we have

$$(2.2) \quad \left(\sum_{j=1}^n \|x_j\|^r \right)^{\frac{1}{r}} \leq C \left(\int_0^1 \left\| \sum_{j=1}^n r_j(t) x_j \right\|^2 dt \right)^{\frac{1}{2}},$$

where r_j is the j th Rademacher function on $[0, 1]$. Further, the best such constant C satisfying (2.2) is usually denoted by $C_r(X)$. We write

$$\text{Cot}(X) := \inf \{2 \leq r \leq \infty : X \text{ has cotype } r\}.$$

For a Banach space with cotype ∞ *i.e.*, $\text{Cot}(X) = \infty$, we denote $\frac{1}{\text{Cot}(X)} = 0$. It is to be noted that the notions of concavity and cotype are closely related concepts in the context of Banach lattices. In fact, a r -concave Banach lattice with $r \geq 2$ is of cotype r . On the other way, each Banach lattice of cotype 2 is 2-concave, and a Banach lattice of cotype $r > 2$ is s -concave for all $s > r$.

Given $m, n \in \mathbb{N}$, we consider the following sets of indices:

$$\begin{aligned} \mathcal{M}(m, n) &= \{\mathbf{j} = (j_1, \dots, j_m) : 1 \leq j_1, \dots, j_m \leq n\} \\ &= \{1, 2, \dots, n\}^m, \end{aligned}$$

$$\mathcal{J}(m, n) = \{\mathbf{j} \in \mathcal{M}(m, n) : 1 \leq j_1 \leq \dots \leq j_m \leq n\}.$$

Furthermore, we write another closely related multi-index set

$$\Lambda(m, n) = \{\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}_0^n : |\alpha| = m\},$$

where $|\alpha| = \alpha_1 + \dots + \alpha_n$. We may identify $\Lambda(m, n)$ with $\mathcal{J}(m, n)$ and vice-versa. For $\alpha = (\alpha_1, \dots, \alpha_n) \in \Lambda(m, n)$, the associated index $\mathbf{j}_\alpha \in \mathcal{J}(m, n)$ is given by $\mathbf{j}_\alpha = (1, \overset{\alpha_1}{\cdot}, 1, 2, \overset{\alpha_2}{\cdot}, 2, \dots, n, \overset{\alpha_n}{\cdot}, n)$. Moreover, for $\mathbf{j} \in \mathcal{J}(m, n)$ the associated multi-index $\mathbf{j}_\alpha \in \Lambda(m, n)$ is given by $\mathbf{j}_{\alpha_r} = |\{k : j_k = r\}|$. This gives a one-to-one correspondence between $\Lambda(m, n)$ and $\mathcal{J}(m, n)$. To identify $\mathcal{M}(m, n)$, we define the following equivalence relation on $\mathcal{M}(m, n)$: for $\mathbf{i}, \mathbf{j} \in \mathcal{M}(m, n)$, we denote $\mathbf{i} \sim \mathbf{j}$ if, and only if, there is a permutation $\sigma \in S_m$ such that $\mathbf{j} = (i_{\sigma(1)}, \dots, i_{\sigma(m)})$. We observe that

$$\mathcal{M}(m, n) = \cup_{\mathbf{j} \in \mathcal{J}(m, n)} [\mathbf{j}].$$

Moreover, it is worth noting that for each $\alpha \in \Lambda(m, n)$ we have

$$\text{card}[\mathbf{j}_\alpha] = \frac{m!}{\alpha!},$$

where $\alpha! = \alpha_1! \cdots \alpha_n!$. Further, we consider the following notation from [6]: given a subset $\mathcal{J} \subset \mathcal{J}(m, n)$, we denote

$$\mathcal{J}^* = \{\mathbf{j} \in \mathcal{J}(m-1, n) : \text{there is } k \geq 1, (\mathbf{j}, k) \in \mathcal{J}\}.$$

For $1 \leq q \leq \infty$, let $\mathcal{P}({}^m\ell_q^n, X)$ be the linear space of all m -homogeneous polynomial $Q : \ell_q^n \rightarrow X$ of n complex variables with values in X , equipped with uniform (or sup) norm

$$\|Q\|_{\ell_q^n} := \sup_{z \in B_{\ell_q^n}} \|Q(z)\|_X.$$

Using the above multi-index notations, a polynomial $Q \in \mathcal{P}({}^m\ell_q^n, X)$ can be expressed in the form

$$Q(z) = \sum_{\alpha \in \Lambda(m, n)} x_\alpha z^\alpha = \sum_{\mathbf{j} \in \mathcal{J}(m, n)} c_{\mathbf{j}} z_{\mathbf{j}},$$

where $x_\alpha, c_{\mathbf{j}} \in X$ and $x_\alpha = c_{\mathbf{j}}$ with $\mathbf{j} = (1, \overset{\alpha_1}{\cdot}, 1, 2, \overset{\alpha_2}{\cdot}, 2, \dots, n, \overset{\alpha_n}{\cdot}, n)$. If $f : B_{\ell_q^n} \rightarrow X$ is an X -valued holomorphic function then we write $f(z) = \sum_{\alpha \in \mathbb{N}_0^n} x_\alpha(f) z^\alpha$ as the monomial series expansion of f , where $x_\alpha(f) = (\partial^\alpha f(0))/\alpha!$ is the α th coefficient of the expansion.

A bounded linear operator $T : X \rightarrow Y$ between two Banach spaces X and Y is called (r, s) -summing, $1 \leq r, s < \infty$, if there is a constant $\tilde{C} > 0$ such that for each choice of finitely many vectors $x_1, \dots, x_n \in X$, we have

$$(2.3) \quad \left(\sum_{j=1}^n \|T(x_j)\|^r \right)^{\frac{1}{r}} \leq \tilde{C} \sup_{x^* \in B_{X^*}} \left(\sum_{j=1}^n |x^*(x_j)|^s \right)^{\frac{1}{s}}.$$

Also, the best such constant \tilde{C} satisfying (2.3) is as usual denoted by $\pi_{r,s}(T)$. In particular, if $r = s$, we call T is r -summing and we write the best constant as $\pi_r(T)$.

3. MAIN RESULTS

As a crucial property, we ensure that for every bounded operator T with $\|T\| < \lambda^{1/p}$, the λ_p -Bohr radius of T is non-zero.

Proposition 3.1. *Let $1 \leq p < \infty$ and $\lambda > 1$. Suppose $T : X \rightarrow Y$ is a bounded operator between complex Banach spaces X and Y such that $\|T\| < \lambda^{1/p}$. Then there exists a constant $M > 0$ such that for all $1 \leq q \leq \infty$ and $n \in \mathbb{N}$ we have*

$$K^p(B_{\ell_q^n}, T, \lambda) \geq M \frac{1}{n^{\frac{1}{q} - \frac{1}{pq}}},$$

where

$$M = \begin{cases} \max \left\{ \left(\frac{\lambda - \|T\|^p}{2\lambda - \|T\|^p} \right)^{1/p}, \left(\frac{\lambda - \|T\|^p}{(\lambda - \|T\|^p + 1)\|T\|} \right)^{1/p} \right\}, & \text{if } \|T\| \geq 1, \\ \max \left\{ \left(\frac{\lambda - \|T\|^p}{2\lambda - \|T\|^p} \right)^{1/p}, \left(\frac{\lambda - \|T\|^p}{\lambda - \|T\|^p + 1} \right)^{1/p} \right\}, & \text{if } 0 < \|T\| < 1. \end{cases}$$

As a consequence, whenever we consider the identity operator on X , Proposition 3.1 demonstrates the lower bound for the classical p -Bohr radius of Banach spaces.

Corollary 3.2. *Let X be a complex Banach space and $\lambda > 1$, $1 \leq p < \infty$. Then for all $1 \leq q \leq \infty$ and $n \in \mathbb{N}$ we have*

$$K^p(B_{\ell_q^n}, X, \lambda) \geq \left(\frac{\lambda - 1}{\lambda} \right)^{\frac{1}{p}} \frac{1}{n^{\frac{1}{p} - \frac{1}{pq}}}.$$

Remark 3.1. It is worth noting that for $\lambda = 1$ and $p = 1$, we actually obtain the trivial lower bound $K(B_{\ell_q^n}, X) \geq 0$. Indeed, as we have discussed before, Blasco [8, Theorem 1.2] has also observed that $K(\ell_q^2) = 0$ for every $1 \leq q \leq \infty$.

Our main aim of this paper is to investigate precise asymptotic behavior of λ_p -Bohr radius $K^p(B_{\ell_q^n}, T, \lambda)$, where T is a bounded operator between two complex Banach spaces. Motivated by the work of Defant *et. al.* [25], one of our contributions in this paper is the following theorem, which provides the correct asymptotic estimates for λ_p -Bohr radius of identity operator on Banach spaces, where $1 \leq p < \infty$. Let us first concentrate on finite dimensional Banach spaces.

Theorem 3.1. *Let X be a finite dimensional complex Banach space and $\lambda > 1$, $1 \leq p < \infty$. Then there are constants B and C such that for each $1 \leq q \leq \infty$ and $n \in \mathbb{N}$ we have*

$$K^p(B_{\ell_q^n}, X, \lambda) \leq B \lambda^{2/p} n^{1-1/p} \left(\frac{\log n}{n} \right)^{1 - \frac{1}{\min\{q, 2\}}}$$

and

$$K^p(B_{\ell_q^n}, X, \lambda) \geq \begin{cases} C \left(\frac{\lambda - \|T\|^p}{2\lambda - \|T\|^p} \right)^{1/p} \left(\frac{\log n}{n} \right)^{1 - \frac{1}{r}}, & \text{for } q \leq r \\ C \left(\frac{\lambda - \|T\|^p}{2\lambda - \|T\|^p} \right)^{1/p} n^{\frac{1}{q} - \frac{1}{r}} \left(\frac{\log n}{n} \right)^{1 - \frac{1}{r}}, & \text{for } r < q \end{cases}$$

where $r = pq'/(pq' - 1)$. Here B is a universal constant and C depends only on X .

Now, we want to study the Bohr radius $K^p(B_{\ell_q^n}, X, \lambda)$ for the infinite dimensional Banach spaces. In this cases, we observe that the log term is missing from the estimates of $K^p(B_{\ell_q^n}, X, \lambda)$. In fact, the asymptotic decay is very much influenced by geometry of the Banach space X i.e., by its optimal cotype $\text{Cot}(X)$.

Theorem 3.2. *Let $1 \leq p < \infty$ and $1 \leq q \leq \infty$. If X is an infinite dimensional complex Banach space of cotype r with $2 \leq r \leq \infty$, then we have*

$$K^p(B_{\ell_q^n}, X, \lambda) \leq \begin{cases} \frac{\lambda^{1/p}}{n^{\frac{1}{p} - \frac{1}{q}}}, & \text{for } q \leq \text{Cot}(X) = r \\ \frac{\lambda^{1/p}}{n^{\frac{1}{p}(1 - \frac{1}{\text{Cot}(X)}) - \frac{1}{q}(1 - \frac{1}{p})}}, & \text{for } q > \text{Cot}(X) = r \end{cases}$$

and

$$K^p(B_{\ell_q^n}, X, \lambda) \geq \begin{cases} \left(\frac{\lambda - 1}{\lambda} \right)^{1/p} \frac{1}{e^{C_r(X)}}, & \text{for } q \leq s \\ \left(\frac{\lambda - 1}{\lambda} \right)^{1/p} \frac{1}{e^{C_r(X)} n^{\frac{1}{s} - \frac{1}{q}}}, & \text{for } q > s, \end{cases}$$

where $s = r/(r - 1)$.

Defant *et. al.* [25, Theorem 4.1] have used the hypercontractivity of the Bohnenblust-Hille inequality to establish the asymptotic behavior of of the Bohr radii $K^p(B_{\ell_\infty^n}, X, \lambda)$ for $p = 1$. Moreover, as a technical tool, Defant *et. al.* [25, Theorem 5.3] have shown Bohnenblust-Hille type inequality for $(s, 1)$ -summing operator to estimate the multidimensional Bohr radii for operators: let Y be a r -concave Banach lattice, with $2 \leq r < \infty$, and $T : X \rightarrow Y$ an $(s, 1)$ -summing operator with $1 \leq s \leq r$. Then there is a constant $C > 0$ such that for every m -homogeneous polynomial $Q : \ell_\infty^n \rightarrow X$, $Q(z) = \sum_{\alpha \in \Lambda(m, n)} x_\alpha z^\alpha$

$$\left(\sum_{\alpha \in \Lambda(m, n)} \|T(x_\alpha)\|_Y^\rho \right)^{\frac{1}{p}} \leq C^m \|Q\|_{B_{\ell_\infty^n}},$$

where

$$\rho := \frac{rsm}{r + (m - 1)s}.$$

Recently, by considering the diagonal operator $D_z : \ell_\infty^n \rightarrow \ell_q^n$, Kumar and Manna [32] have obtained an extended version of the above result for m -homogeneous polynomials $Q \in \mathcal{P}(^m \ell_q^n, X)$, $1 \leq q < \infty$.

Theorem 3.3. [32] *Let $2 \leq r < \infty$. Suppose Y is a r -concave Banach lattice, and the operator $T : X \rightarrow Y$ is an $(s, 1)$ -summing operator with $1 \leq s \leq r$. Then there is a constant $\tilde{C} > 0$ such that the following holds*

$$\left(\sum_{\alpha \in \Lambda(m, n)} \|T(x_\alpha)z^\alpha\|_Y^\rho \right)^{\frac{1}{p}} \leq \tilde{C}^m \|Q\|_{B_{\ell_q^n}}$$

for every polynomial $Q(z) = \sum_{\alpha \in \Lambda(m,n)} x_\alpha z^\alpha \in \mathcal{P}(m\ell_q^n)$, $1 \leq q < \infty$.

We are now ready to state our next main result. As a consequence of the above technical discussions, Theorem 3.4 provides the lower estimates of the Bohr radius $K^p(B_{\ell_q^n}, T, \lambda)$ for any bounded operator T .

Theorem 3.4. *Let $1 \leq p < \infty$ and $T : X \rightarrow Y$ be a bounded operator between Banach spaces X and Y .*

- (1) *Assume that X or Y is of cotype r with $2 \leq r \leq \infty$. Then there is a constant $D > 0$ such that for every $\|T\| < \lambda^{1/p}$, $1 \leq q \leq \infty$, and n*

$$K^p(B_{\ell_q^n}, T, \lambda) \geq \begin{cases} D \left(\frac{\lambda - \|T\|^p}{\lambda} \right)^{\frac{1}{p}}, & \text{for } q \leq s \\ D \left(\frac{\lambda - \|T\|^p}{\lambda} \right)^{\frac{1}{p}} \left(\frac{1}{n} \right)^{\frac{1}{s} - \frac{1}{q}}, & \text{for } q > s, \end{cases}$$

where $s = r/(r-1)$.

- (2) *Assume that Y is a r -concave Banach lattice with $2 \leq r < \infty$ and there is a $1 \leq s < r$ such that the operator T is $(s, 1)$ -summing. Then there is a constant $D > 0$ such that*

$$K^p(B_{\ell_q^n}, T, \lambda) \geq D \left(\frac{\lambda - \|T\|^p}{2\lambda - \|T\|^p} \right)^{\frac{1}{p}} \left(\frac{\log n}{n} \right)^{1 - \frac{1}{r}}$$

for every $\|T\| < \lambda^{1/p}$, $1 \leq q \leq \infty$, and $n \in \mathbb{N}$.

We now move our focus for a while on arithmetic Bohr radius of Banach spaces, which is another interesting multidimensional variant of Bohr radius. In fact, it has very close relationship with classical multidimensional Bohr radii. In order to study the upper inclusion of domains of convergence of monomial expansions, as a technical tool, Defant *et. al.* [24] introduced the arithmetic Bohr radius for scalar-valued holomorphic functions. Later, it was studied independently by the same authors in [23]. Recently, the study of arithmetic Bohr radius has been extended for vector-valued holomorphic functions (see [4]). Indeed, the authors [4] have introduced the following notion of arithmetic Bohr radius:

Definition 3.1. [4] Let $\mathcal{F}(B_{\ell_q^n}, X)$ be set of all holomorphic functions on $B_{\ell_q^n}$ into a complex Banach space X . For each $1 \leq p < \infty$ and $\lambda \geq 1$, the λ_p -arithmetic Bohr radius of $B_{\ell_q^n}$ with respect to $\mathcal{F}(B_{\ell_q^n}, X)$ is defined as

$$A_p(\mathcal{F}(B_{\ell_q^n}, X), \lambda) := \sup \left\{ \frac{1}{n} \sum_{i=1}^n r_i \mid r \in \mathbb{R}_{\geq 0}^n, \forall f \in \mathcal{F}(B_{\ell_q^n}, X) : \sum_{\alpha \in \mathbb{N}_0^n} \|x_\alpha(f)\|^p r^{p\alpha} \leq \lambda \|f\|_{B_{\ell_q^n}}^p \right\},$$

where $\mathbb{R}_{\geq 0}^n = \{r = (r_1, \dots, r_n) \in \mathbb{R}^n : r_i \geq 0, 1 \leq i \leq n\}$. We write $A_p(B_{\ell_q^n}, X, \lambda)$ for $A_p(H_\infty(B_{\ell_q^n}, X), \lambda)$ and $A_p(B_{\ell_q^n}, X)$ for $A_p(B_{\ell_q^n}, X, 1)$.

In view of [4, Theorem 2.2] and Theorem 3.1, as an application, we obtain the following lower estimate for the arithmetic Bohr radius $A_p(B_{\ell_q^n}, X, \lambda)$ in case of X is finite dimensional.

Theorem 3.5. *Let $1 \leq p < \infty$, $\lambda \geq 1$, and $1 \leq q \leq \infty$. Suppose X is a finite dimensional Banach space. Then there is a constant $C > 0$ such that for every $n > 1$ we have*

$$A_p(B_{\ell_q^n}, X, \lambda) \geq \begin{cases} C \left(\frac{\lambda-1}{2\lambda-1}\right)^{1/p} n^{-\frac{1}{q}} \left(\frac{\log n}{n}\right)^{1-\frac{1}{r}}, & \text{for } q \leq r \\ C \left(\frac{\lambda-1}{2\lambda-1}\right)^{1/p} n^{-\frac{1}{r}} \left(\frac{\log n}{n}\right)^{1-\frac{1}{r}}, & \text{for } r < q, \end{cases}$$

where $r = pq'/(pq' - 1)$, and the constant C depends only on X .

4. KEY LEMMAS

As demonstrated by the scalar case, studying the m -homogeneous polynomial first is typically a beneficial approach for obtaining nontrivial estimates for Bohr radii. The following definition is the m -homogeneous analogue of Definition 1.1 for $m \in \mathbb{N}$.

Definition 4.1. Let $1 \leq p < \infty$, $1 \leq q \leq \infty$, $\lambda \geq 1$, $m, n \in \mathbb{N}$, and $T : X \rightarrow Y$ be a bounded operator in complex Banach spaces such that $\|T\| \leq \lambda^{1/p}$. Then $K_m^p(B_{\ell_q^n}, X, \lambda)$ is defined to be the supremum of all $r \geq 0$ such that for every m -homogeneous polynomials $Q \in \mathcal{P}(^m \ell_q^n, X)$, $Q(z) = \sum_{\alpha \in \Lambda(m, n)} x_\alpha z^\alpha$, we have

$$\sup_{z \in r B_{\ell_q^n}} \sum_{\alpha \in \Lambda(m, n)} \|T(x_\alpha) z^\alpha\|_Y^p \leq \lambda \left\| \sum_{\alpha \in \Lambda(m, n)} x_\alpha z^\alpha \right\|_X^p.$$

It is worth mentioning that if T is a null operator, then the above supremum is ∞ . Thus, by assuming that $T \neq 0$ throughout the paper, we are avoiding this trivial scenario. The constant $K_m^p(B_{\ell_q^n}, X, \lambda)$ in the Definition 4.1 can be rephrased as

$$(4.1) \quad K_m^p(B_{\ell_q^n}, X, \lambda) := \sup \left\{ r \geq 0 : \sup_{z \in B_{\ell_q^n}} \sum_{\alpha \in \Lambda(m, n)} \|T(x_\alpha) z^\alpha\|_Y^p \leq \frac{\lambda}{r^{mp}} \sup_{z \in B_{\ell_q^n}} \|Q(z)\|_X^p \right. \\ \left. \text{for all } Q \in \mathcal{P}(^m \ell_q^n, X) \right\}.$$

Further, it is worth noting that

$$(4.2) \quad K_m^p(B_{\ell_q^n}, T, \lambda) = \sqrt[m]{\lambda} K_m^p(B_{\ell_q^n}, T)$$

and

$$(4.3) \quad K^p(B_{\ell_q^n}, T, \lambda) \geq \max \left\{ K^p(B_{\ell_q^n}, X, \lambda / \|T\|^p), K^p(B_{\ell_q^n}, Y, \lambda / \|T\|^p) \right\}.$$

The following lemma provides a nice connection between Definitions 1.1 and 4.1, which will be one of the most important technical tools to prove our main results in this paper.

Lemma 4.4. *Let $1 \leq p < \infty$ and $T : X \rightarrow Y$ be a bounded operator between complex Banach spaces X and Y with $\|T\| < \lambda^{1/p}$. Then we have*

$$(a) \quad \left(\frac{\lambda - \|T\|^p}{2\lambda - \|T\|^p} \right)^{1/p} \inf_{m \in \mathbb{N}} K_m^p(B_{\ell_q^n}, T, \lambda) \leq K^p(B_{\ell_q^n}, T, \lambda) \leq \inf_{m \in \mathbb{N}} K_m^p(B_{\ell_q^n}, T, \lambda) \\ (b) \quad \left(\frac{\lambda - \|T\|^p}{\lambda - \|T\|^p + 1} \right)^{1/p} \inf_{m \in \mathbb{N}} K_m^p(B_{\ell_q^n}, T) \leq K^p(B_{\ell_q^n}, T, \lambda) \leq \lambda^{1/p} \inf_{m \in \mathbb{N}} K_m^p(B_{\ell_q^n}, T).$$

Proof. The right hand side inequality of (a) is clear from the fact that $K^p(B_{\ell_q^n}, T, \lambda) \leq K_m^p(B_{\ell_q^n}, T, \lambda)$ for all $m \geq 1$. So, we now proceed for the left hand side inequality of (a).

Let $f(z) = \sum_{\alpha \in \mathbb{N}_0^n} x_\alpha z^\alpha$ be a holomorphic function on $B_{\ell_q^n}$ with values in the complex Banach space X . Further, suppose $v \in \mathbb{C}^n$ such that $\|v\| \leq \inf_{m \in \mathbb{N}} \{K_m^p(B_{\ell_q^n}, T, \lambda)\}$. It is worth noting that for $1 \leq p < \infty$

$$0 < \left(\frac{\lambda - \|T\|^p}{2\lambda - \|T\|^p} \right)^{1/p} < 1.$$

Since $v \in K_m^p(B_{\ell_q^n}, T, \lambda) \overline{B_{\ell_q^n}}$ for all $m \in \mathbb{N}$ and using the Cauchy-Riemann inequalities we have that

$$\begin{aligned} & \sum_{\alpha \in \mathbb{N}_0^n} \left\| T(x_\alpha) \left(\left(\frac{\lambda - \|T\|^p}{2\lambda - \|T\|^p} \right)^{1/p} v \right)^\alpha \right\|_Y^p \\ &= \|T(x_0)\|^p + \sum_{m=1}^{\infty} \left(\frac{\lambda - \|T\|^p}{2\lambda - \|T\|^p} \right)^m \sum_{\alpha \in \Lambda(m,n)} \|T(x_\alpha)\|^p |v|^{p\alpha} \\ &\leq \|T\|^p \|x_0\|^p + \sum_{m=1}^{\infty} \left(\frac{\lambda - \|T\|^p}{2\lambda - \|T\|^p} \right)^m \lambda \left\| \sum_{\alpha \in \Lambda(m,n)} x_\alpha z^\alpha \right\|_{B_{\ell_q^n}}^p \\ &\leq \left(\|T\|^p + \lambda \sum_{m=1}^{\infty} \left(\frac{\lambda - \|T\|^p}{2\lambda - \|T\|^p} \right)^m \right) \|f\|_{B_{\ell_q^n}}^p = \lambda \|f\|_{B_{\ell_q^n}}^p. \end{aligned}$$

We shall prove (b) in the similar way. Since $K_m^p(B_{\ell_q^n}, T, \lambda) = \lambda^{1/pm} K_m^p(B_{\ell_q^n}, T)$, we obtain

$$K^p(B_{\ell_q^n}, T, \lambda) \leq \inf_{m \in \mathbb{N}} \left\{ \lambda^{\frac{1}{pm}} K_m^p(B_{\ell_q^n}, T) \right\} \leq \lambda^{1/p} \inf_{m \in \mathbb{N}} \{K_m^p(B_{\ell_q^n}, T)\}.$$

In this case, we suppose again that $v \in \mathbb{C}^n$ as above but here we proceed by considering $\|v\| \leq \inf_{m \in \mathbb{N}} \{K_m^p(B_{\ell_q^n}, T)\}$. Thus, for $v \in K_m^p(B_{\ell_q^n}, T) \overline{B_{\ell_q^n}}$, we obtain

$$\begin{aligned} & \sum_{\alpha \in \mathbb{N}_0^n} \left\| T(x_\alpha) \left(\left(\frac{\lambda - \|T\|^p}{\lambda - \|T\|^p + 1} \right)^{1/p} v \right)^\alpha \right\|_Y^p \\ &= \|T(x_0)\|^p + \sum_{m=1}^{\infty} \left(\frac{\lambda - \|T\|^p}{\lambda - \|T\|^p + 1} \right)^m \sum_{\alpha \in \Lambda(m,n)} \|T(x_\alpha)\|^p |v|^{p\alpha} \\ &\leq \|T\|^p \|x_0\|^p + \sum_{m=1}^{\infty} \left(\frac{\lambda - \|T\|^p}{\lambda - \|T\|^p + 1} \right)^m \left\| \sum_{\alpha \in \Lambda(m,n)} x_\alpha z^\alpha \right\|_{B_{\ell_q^n}}^p \\ &\leq \left(\|T\|^p + \sum_{m=1}^{\infty} \left(\frac{\lambda - \|T\|^p}{\lambda - \|T\|^p + 1} \right)^m \right) \|f\|_{B_{\ell_q^n}}^p = \lambda \|f\|_{B_{\ell_q^n}}^p. \end{aligned}$$

This completes the proof. \square

The following lemma of Bayart *et. al.* [6, Lemma 3.5] is a key tool to prove Theorem 3.1. For a m -homogeneous polynomial $Q \in \mathcal{P}(^m \ell_q^n)$, ($1 \leq q \leq \infty$), it guarantees the ℓ_q -summability of certain coefficients of Q in terms of its uniform norm on ℓ_q^n .

Lemma 4.5. [6] *Let $1 \leq t \leq \infty$ and P be an m -homogeneous polynomial in n variables. Then for any $\mathbf{j} \in \mathcal{J}(m-1, n)$,*

$$\left(\sum_{k=j_{m-1}}^n |c_{(\mathbf{j},k)}(P)|^{t'} \right)^{1/t'} \leq m e^{1+\frac{m-1}{t}} |\mathbf{j}|^{1/t} \|P\|_{\mathcal{P}(m, \ell_t^n)}.$$

5. PROOF OF THE MAIN RESULTS

Proof of Proposition 3.1. Let $Q(z) = \sum_{\alpha \in \Lambda(m,n)} x_\alpha z^\alpha \in \mathcal{P}(m, \ell_q^n, X)$ be an m -homogeneous polynomial. By using Hölder's inequality and for all $z \in \mathbb{C}^n$, we have

$$\begin{aligned} \sum_{\alpha \in \Lambda(m,n)} \|T(x_\alpha) z^\alpha\|^p &\leq \max_{\alpha \in \Lambda(m,n)} \|T(x_\alpha)\|^p \sum_{\alpha \in \Lambda(m,n)} |z^\alpha|^p \\ &\leq \|T\|^p \max_{\alpha \in \Lambda(m,n)} \|x_\alpha\|^p (|z_1|^p + \cdots + |z_n|^p)^m \\ &\leq \lambda \max_{\alpha \in \Lambda(m,n)} \|x_\alpha\|^p (|z_1|^{pq} + \cdots + |z_n|^{pq})^{\frac{m}{q}} n^{(1-\frac{1}{q})m} \\ &\leq \lambda \|Q\|_{\ell_q^n}^p \|z\|_{pq}^{mp} n^{mp(\frac{1}{p}-\frac{1}{pq})} \\ &\leq \lambda \|Q\|_{\ell_q^n}^p \|z\|_q^{mp} n^{mp(\frac{1}{p}-\frac{1}{pq})}. \end{aligned}$$

Therefore, for all $z \in (1/n^{(1/p-1/pq)}) B_{\ell_q^n}$, we obtain

$$\sum_{\alpha \in \Lambda(m,n)} \|T(x_\alpha) z^\alpha\|^p \leq \lambda \|Q\|^p.$$

Thus, for every m -homogeneous polynomial $Q \in \mathcal{P}(m, \ell_q^n)$, we obtain

$$\sup \left\{ \sum_{\alpha \in \Lambda(m,n)} \|T(x_\alpha) z^\alpha\|^p \mid z \in \frac{1}{n^{\frac{1}{p}-\frac{1}{pq}}} B_{\ell_q^n} \right\} \leq \lambda \|Q\|^p,$$

which yields

$$K_m^p(B_{\ell_q^n}, T, \lambda) \geq \frac{1}{n^{\frac{1}{p}-\frac{1}{pq}}}.$$

In view of Lemma 4.4(a), for each $1 \leq p < \infty$, we have

$$K^p(B_{\ell_q^n}, T, \lambda) \geq \frac{1}{n^{\frac{1}{p}-\frac{1}{pq}}} \left(\frac{\lambda - \|T\|^p}{2\lambda - \|T\|^p} \right)^{1/p}.$$

By going similar lines of arguments as above for every m -homogeneous polynomial $Q(z) = \sum_{|\alpha|=m} x_\alpha z^\alpha$, we obtain that

$$z \in \left(\frac{1}{m \sqrt{\|T\|} n^{\frac{1}{p}-\frac{1}{pq}}} \right) B_{\ell_q^n} \sum_{\alpha \in \Lambda(m,n)} \|T(x_\alpha) z^\alpha\|^p = \frac{1}{\|T\|^p} \sup_{z \in \left(\frac{1}{n^{\frac{1}{p}-\frac{1}{pq}}} \right) B_{\ell_q^n}} \sum_{\alpha \in \Lambda(m,n)} \|T(x_\alpha) z^\alpha\|^p \leq \|Q\|_{\mathcal{P}(m, \ell_q^n)}^p.$$

Hence, for every $m \in \mathbb{N}$, we have

$$K_m^p(B_{\ell_q^n}, T) \geq \frac{1}{m \sqrt{\|T\|} n^{\frac{1}{p}-\frac{1}{pq}}}.$$

Now, by the virtue of Lemma 4.4(b), we have

$$K^p(B_{\ell_q^n}, T, \lambda) \geq \left(\frac{\lambda - \|T\|^p}{\lambda - \|T\|^p + 1} \right)^{1/p} \frac{1}{\|T\| n^{\frac{1}{p} - \frac{1}{pq}}} \quad \text{if } \|T\| \geq 1$$

and

$$K^p(B_{\ell_q^n}, T, \lambda) \geq \left(\frac{\lambda - \|T\|^p}{\lambda - \|T\|^p + 1} \right)^{1/p} \frac{1}{n^{\frac{1}{p} - \frac{1}{pq}}} \quad \text{if } 0 < \|T\| < 1.$$

This completes the proof. \square

Proof of Theorem 3.1. We first check that the upper bound of $K^p(B_{\ell_q^n}, X, \lambda)$ is true. In view of [4, Lemma 2.3], we have

$$(5.1) \quad K^p(B_{\ell_q^n}, X, \lambda) \leq n^{1/q} A_p(B_{\ell_q^n}, X, \lambda), \quad \lambda \geq 1$$

for any $1 \leq p < \infty$ and $n \in \mathbb{N}$, where $A_p(B_{\ell_q^n}, X, \lambda)$ is the λ_p -arithmetic Bohr radius for the unit ball of ℓ_q^n spaces. In the same paper [4, Theorem 2.2], the authors have established that there exists a constant $B > 0$ such that

$$(5.2) \quad A_p(B_{\ell_q^n}, X, \lambda) \leq B \lambda^{\frac{2}{p \log n}} \frac{(\log n)^{1 - (1/\min\{q, 2\})}}{n^{\frac{1}{p} + \frac{1}{\max\{2, q\}} - \frac{1}{2}}}.$$

Using (5.1) and (5.2) and a simple computation yields

$$K^p(B_{\ell_q^n}, X, \lambda) \leq B \lambda^{\frac{2}{p \log n}} \frac{(\log n)^{1 - \frac{1}{\min\{q, 2\}}}}{n^{\frac{1}{p} - \frac{1}{\min\{q, 2\}}}} = B \lambda^{\frac{2}{p \log n}} n^{1 - \frac{1}{p}} \left(\frac{\log n}{n} \right)^{1 - \frac{1}{\min\{q, 2\}}}.$$

Now we shall prove the lower bound for $K^p(B_{\ell_q^n}, X, \lambda)$. By the virtue of Lemma 4.4, it is sufficient to show the existence of a constant $C(X) > 0$ such that

$$\inf_{m \in \mathbb{N}} K_m^p(B_{\ell_q^n}, X, \lambda) \geq \begin{cases} C \left(\frac{\log n}{n} \right)^{1 - \frac{1}{r}}, & \text{for } q \leq r \\ C n^{\frac{1}{q} - \frac{1}{r}} \left(\frac{\log n}{n} \right)^{1 - \frac{1}{r}}, & \text{for } r < q \end{cases}$$

for every $1 \leq p < \infty$. Let us fix $Q \in \mathcal{P}(^m \ell_q^n, X)$ and $u \in \ell_q^n$. Then, by Lemma 4.5, it is worth noting that

$$(5.3) \quad \left(\sum_{k: (\mathbf{j}, k) \in \mathcal{J}(m, n)} |c_{(\mathbf{j}, k)}(P)|^{t'} \right)^{1/t'} = \left(\sum_{k=j_{m-1}}^n |c_{(\mathbf{j}, k)}(P)|^{t'} \right)^{1/t'} \leq m e^{1 + \frac{m-1}{t}} |\mathbf{j}|^{1/t} \|P\|_{\mathcal{P}(^m \ell_q^n)}$$

holds for any m -homogeneous polynomial $P \in \mathcal{P}(^m \ell_q^n)$, $1 \leq t \leq \infty$. Since X is a finite dimensional Banach space, the identity map Id_X on X is absolutely summing *i.e.*, 1-summing and hence the operator Id_X is s -summing for every $s \geq 1$ with $\pi_s(\text{Id}_X) \leq \pi_1(\text{Id}_X)$ (see [26, Theorem 10.4]). In other words, there exists a constant $d > 0$ such that for every $s \geq 1$ and for every finitely many $x_1, \dots, x_k \in X$,

$$\left(\sum_{i=1}^k \|x_i\|^s \right)^{1/s} \leq d \sup_{x^* \in B_{X^*}} \left(\sum_{i=1}^k |x^*(x_i)|^s \right)^{1/s}.$$

Hence, if $Q(z) = \sum_{|\alpha|=m} x_\alpha(Q) z^\alpha$, then using the above inequality (5.3), Hölder's inequality (two times), and the multinomial formula, for every $1 \leq p < \infty$ we obtain

$$\begin{aligned}
\sum_{\mathbf{j} \in \mathcal{J}(m,n)} \|x_{\mathbf{j}}(Q)\|^p |u_{\mathbf{j}}|^p &= \sum_{\mathbf{j} \in \mathcal{J}(m,n)^*} \left(\sum_{k: (\mathbf{j},k) \in \mathcal{J}(m,n)} \|x_{\mathbf{j}}(Q)\|^p |u_{\mathbf{j}}|^p |u_k|^p \right) \\
&\leq \sum_{\mathbf{j} \in \mathcal{J}(m,n)^*} |u_{\mathbf{j}}|^p \left(\sum_{k: (\mathbf{j},k) \in \mathcal{J}(m,n)} \|x_{\mathbf{j}}(Q)\|^{pq'} \right)^{1/q'} \left(\sum_k |u_k|^{pq} \right)^{1/q} \\
&= \sum_{\mathbf{j} \in \mathcal{J}(m,n)^*} |u_{\mathbf{j}}|^p \left(\left(\sum_{k: (\mathbf{j},k) \in \mathcal{J}(m,n)} \|x_{\mathbf{j}}(Q)\|^{pq'} \right)^{1/pq'} \|u\|_{pq} \right)^p \\
&\leq \pi_1^p(\text{Id}_X) \sum_{\mathbf{j} \in \mathcal{J}(m,n)^*} |u_{\mathbf{j}}|^p \left(\sup_{x^* \in B_{X^*}} \left(\sum_{k: (\mathbf{j},k) \in \mathcal{J}(m,n)} |x^*(x_{(\mathbf{j},k)}(Q))|^{pq'} \right)^{1/pq'} \right)^p \|u\|_{pq}^p \\
&\leq \pi_1^p(\text{Id}_X) \left(me^{1+\frac{m-1}{r}} \right)^p \|u\|_{pq}^p \sup_{x^* \in B_{X^*}} \sup_{z \in \ell_q^n} \left| x^* \left(\sum_{\mathbf{j} \in \mathcal{J}(m,n)} x_\alpha(Q) z^\alpha \right) \right|^p \sum_{\mathbf{j} \in \mathcal{J}(m,n)^*} |\mathbf{j}|^{p/r} |u_{\mathbf{j}}|^p \\
&\leq \pi_1^p(\text{Id}_X) \left(me^{1+\frac{m-1}{r}} \right)^p \|u\|_q^p \|Q\|_{\mathcal{P}(m, \ell_q^n)}^p \left(\sum_{\mathbf{j} \in \mathcal{J}(m,n)^*} |\mathbf{j}|^{1/r} |u_{\mathbf{j}}| \right)^p \\
&\leq \pi_1^p(\text{Id}_X) \left(me^{1+\frac{m-1}{r}} \right)^p \|u\|_q^p \|Q\|_{\mathcal{P}(m, \ell_q^n)}^p \left(\left(\sum_{\mathbf{j} \in \mathcal{J}(m,n)^*} |\mathbf{j}| |u_{\mathbf{j}}|^r \right)^{1/r} \left(\sum_{\mathbf{j} \in \mathcal{J}(m,n)^*} 1 \right)^{1-1/r} \right)^p.
\end{aligned}$$

It is worth noting that

$$\left(\sum_{\mathbf{j} \in \mathcal{J}(m,n)^*} |\mathbf{j}| |u_{\mathbf{j}}|^r \right)^{1/r} = \|u\|_r^{m-1}.$$

Hence, we obtain

$$\sum_{\mathbf{j} \in \mathcal{J}(m,n)} \|x_{\mathbf{j}}(Q)\|^p |u_{\mathbf{j}}|^p \leq \pi_1^p(\text{Id}_X) \left(me^{1+\frac{m-1}{r}} \right)^p \|u\|_r^{p(m-1)} \|u\|_q^p \|Q\|_{\mathcal{P}(m, \ell_q^n)}^p \left(\sum_{\mathbf{j} \in \mathcal{J}(m,n)^*} 1 \right)^{p(1-\frac{1}{r})}.$$

Using the facts that

$$\text{card} \{ \mathbf{j} = (j_1, \dots, j_{m-1}) : 1 \leq j_1 \leq \dots \leq j_{m-1} \leq n \} = \binom{n+m-2}{n-1}$$

and

$$\binom{n+m-2}{n-1} \leq e^m \left(1 + \frac{n}{m} \right)^{m-1} \text{ for all } n, m \in \mathbb{N},$$

finally we have the following inequality

$$\sum_{\mathbf{j} \in \mathcal{J}(m,n)} \|x_{\mathbf{j}}(Q)\|^p |u_{\mathbf{j}}|^p \leq \pi_1^p(\text{Id}_X) \left(m e^{m+\frac{1}{r'}} \right)^p \left(1 + \frac{n}{m} \right)^{\frac{p(m-1)}{r'}} \|u\|_r^{p(m-1)} \|u\|_q^p \|Q\|_{\mathcal{P}(m\ell_q^n)}^p.$$

For $q \leq r$, we have $\|u\|_r \leq \|u\|_q$. Then, we obtain

$$(5.4) \quad \sum_{\mathbf{j} \in \mathcal{J}(m,n)} \|x_{\mathbf{j}}(Q)\|^p |u_{\mathbf{j}}|^p \leq \pi_1^p(\text{Id}_X) \left(m e^{m+\frac{1}{r'}} \right)^p \left(1 + \frac{n}{m} \right)^{\frac{p(m-1)}{r'}} \|u\|_q^{mp} \|Q\|_{\mathcal{P}(m\ell_q^n)}^p$$

and hence, in view of (4.1) and (5.4), it follows that

$$(5.5) \quad K_m^p(B_{\ell_q^n}, X, \lambda) \geq \sqrt[p]{\lambda} \left(\pi_1(\text{Id}_X) m e^{m+\frac{1}{r'}} \right)^{-1/m} \left(1 + \frac{n}{m} \right)^{-\frac{m-1}{mr'}}.$$

On the other hand, we have

$$\|u\|_r \leq n^{\frac{1}{r}-\frac{1}{q}} \|u\|_q \quad \text{for } r < q.$$

Thus, we obtain

$$\sum_{\mathbf{j} \in \mathcal{J}(m,n)} \|x_{\mathbf{j}}(Q)\|^p |u_{\mathbf{j}}|^p \leq \pi_1^p(\text{Id}_X) \left(m e^{m+\frac{1}{r'}} \right)^p \left(1 + \frac{n}{m} \right)^{\frac{p(m-1)}{r'}} \left(n^{\frac{1}{r}-\frac{1}{q}} \right)^{p(m-1)} \|u\|_q^{mp} \|Q\|_{\mathcal{P}(m\ell_q^n)}^p$$

and hence (4.1) gives

$$(5.6) \quad K_m^p(B_{\ell_q^n}, X, \lambda) \geq \sqrt[p]{\lambda} \left(\pi_1(\text{Id}_X) m e^{m+\frac{1}{r'}} \right)^{-1/m} \left(n^{\frac{1}{r}-\frac{1}{q}} \right)^{-\frac{m-1}{m}} \left(1 + \frac{n}{m} \right)^{-\frac{m-1}{mr'}}.$$

From (5.5) and (5.6), we observe that there exists a constant $E(X)$, depending on X , such that

$$K_m^p(B_{\ell_q^n}, X, \lambda) \geq \begin{cases} E(X) \sqrt[p]{\lambda} \left(1 + \frac{n}{m} \right)^{-\frac{m-1}{mr'}}, & \text{for } q \leq r \\ E(X) \sqrt[p]{\lambda} \left(n^{\frac{1}{r}-\frac{1}{q}} \right)^{-\frac{m-1}{m}} \left(1 + \frac{n}{m} \right)^{-\frac{m-1}{mr'}}, & \text{for } r < q \end{cases}$$

for all $1 \leq p < \infty$ and $m \in \mathbb{N}$. The concluding part of this proof lies on the minimizing the right side of this inequality in m . We shall use the following important estimate

$$\left(1 + \frac{n}{m} \right)^{-\frac{m-1}{m}} \geq 2^{-\frac{m-1}{m}} \min \left\{ 1, \left(\frac{n}{m} \right)^{-\frac{m-1}{m}} \right\} \geq \frac{1}{2} \min \left\{ 1, \frac{mn^{1/m}}{nm^{1/m}} \right\}.$$

Now, we observe that the function $h : (0, \infty) \rightarrow \mathbb{R}$ defined by $h(x) = xn^{1/x}$ attains a minimum value $e \log n$ at $x = \log n$. This implies that

$$\left(1 + \frac{n}{m} \right)^{-\frac{m-1}{m}} \geq \frac{1}{2} \left(\frac{\log n}{n} \right).$$

Therefore, there exists a constant $C(X)$ such that

$$\inf_{m \in \mathbb{N}} K_m^p(B_{\ell_q^n}, X, \lambda) \geq \begin{cases} C(X) \left(\frac{\log n}{n} \right)^{\frac{1}{r'}}, & \text{for } q \leq r \\ C(X) n^{\frac{1}{q}-\frac{1}{r'}} \left(\frac{\log n}{n} \right)^{\frac{1}{r'}}, & \text{for } r < q. \end{cases}$$

This completes the proof. \square

Proof of Theorem 3.2. At first, we establish the claim of the upper bound for $K^p(B_{\ell_q^n}, X, \lambda)$. To prove this, we need an important result by Maurey and Pisier [26, Theorem 14.5]: For every infinite dimensional Banach space X ,

$$\inf \{2 \leq r \leq \infty : X \text{ has cotype } r\} = \sup \{2 \leq r \leq \infty : X \text{ finitely factors } \ell_r \hookrightarrow \ell_\infty\}.$$

We say that X finitely factors $\ell_r \hookrightarrow \ell_\infty$ if for every $\epsilon > 0$ and $n \in \mathbb{N}$, there exist $x_1, \dots, x_n \in X$ such that

$$(5.7) \quad \frac{1}{1+\epsilon} \|z\|_\infty \leq \left\| \sum_{j=1}^n z_j x_j \right\| \leq \|z\|_r$$

for all $z = (z_1, \dots, z_n) \in \mathbb{C}^n$. In particular, by considering $z = e_j = (0, \dots, 0, 1, 0, \dots, 0)$, (5.7) reduces to

$$\frac{1}{1+\epsilon} = \frac{1}{1+\epsilon} \|e_j\|_\infty \leq \|x_j\|, \quad 1 \leq j \leq n.$$

Hence, for a given $\epsilon > 0$, there exist $x_1, \dots, x_n \in X$ such that

$$(5.8) \quad \frac{n}{(1+\epsilon)^p} \leq \sum_{j=1}^n \|x_j\|^p, \quad 1 \leq p < \infty.$$

In view of (5.7) and the definition of $K_1^p(B_{\ell_q^n}, X)$ for 1-homogeneous polynomial in X , for any $1 \leq p < \infty$ we obtain

$$(5.9) \quad \begin{aligned} \sum_{j=1}^n \|x_j\|^p \frac{1}{n^{p/q}} &\leq \frac{1}{(K_1^p(B_{\ell_q^n}, X))^p} \sup_{z \in B_{\ell_q^n}} \left\| \sum_{j=1}^n z_j x_j \right\|^p \\ &\leq \frac{1}{(K_1^p(B_{\ell_q^n}, X))^p} \sup_{z \in B_{\ell_q^n}} \|z\|_{\text{Cot}(X)}^p. \end{aligned}$$

Now, we note that if $q \leq \text{Cot}(X)$ then $\sup_{z \in B_{\ell_q^n}} \|z\|_{\text{Cot}(X)} \leq 1$. Then, using (5.8) and (5.9), we have that

$$(K_1^p(B_{\ell_q^n}, X))^p \leq \frac{(1+\epsilon)^p}{n^{1-\frac{p}{q}}}$$

for every $1 \leq p < \infty$ and for all $\epsilon > 0$. Thus, we have

$$K_1^p(B_{\ell_q^n}, X) \leq \frac{1}{n^{\frac{1}{p}-\frac{1}{q}}}, \quad 1 \leq p < \infty.$$

On the other hand, if $q > \text{Cot}(X) = r$ then we have

$$\|z\|_{\text{Cot}(X)} \leq n^{\frac{1}{\text{Cot}(X)}-\frac{1}{q}} \|z\|_q. \quad (\text{We assume the convention } n^{\frac{1}{\infty}} = 1)$$

In view of (5.8) and (5.9), for any $1 \leq p < \infty$ it is easy to see that

$$(K_1^p(B_{\ell_q^n}, X))^p \leq \frac{(1+\epsilon)^p}{n^{1-\frac{p}{q}-\frac{1}{\text{Cot}(X)}+\frac{1}{q}}}$$

holds for all $\epsilon > 0$. Consequently, it follows that

$$K_1^p(B_{\ell_q^n}, X) \leq \frac{1}{n^{\frac{1}{p}(1-\frac{1}{\text{Cot}(X)})-\frac{1}{q}(1-\frac{1}{p})}}.$$

Now, the conclusion follows immediately from the following observation:

$$K^p(B_{\ell_q^n}, X, \lambda) \leq K_1^p(B_{\ell_q^n}, X, \lambda) = \lambda^{1/p} K_1^p(B_{\ell_q^n}, X).$$

By the virtue of Corollary 3.2, the lower bound for $K^p(B_{\ell_q^n}, X, \lambda)$ has already been discussed for complex Banach spaces with no finite cotype. Suppose that X has cotype r . To prove the lower bound of $K^p(B_{\ell_q^n}, X, \lambda)$ in this case, we use the following technical result by Bombal [13, Theorem 3.2]: Given an m -homogeneous polynomial $Q : \mathbb{C}^n \rightarrow X$, $Q(z) = \sum_{|\alpha|=m} x_\alpha z^\alpha$, we denote by $H : \mathbb{C}^n \times \cdots \times \mathbb{C}^n \rightarrow X$ the unique symmetric m -linear mapping associated to Q . Then we have

$$\left(\sum_{i \in \mathcal{M}(m,n)} \|H(e_{i_1}, \dots, e_{i_m})\|^r \right)^{\frac{1}{r}} \leq C_r(X)^m \|H\|_{B_{\ell_q^n}}.$$

Let us consider $s > 1$ be such that $1/r + 1/s = 1$. Then, by applying Hölder's inequality, for all $z \in \mathbb{C}^n$, we obtain

$$\begin{aligned} \sum_{\alpha \in \Lambda(m,n)} \|x_\alpha z^\alpha\|^p &= \sum_{\mathbf{j} \in \mathcal{J}(m,n)} \text{card}[\mathbf{j}]^p \|H(e_{j_1}, \dots, e_{j_m})\|^p |z_{j_1} \cdots z_{j_m}|^p \\ &= \sum_{i \in \mathcal{M}(m,n)} \|H(e_{i_1}, \dots, e_{i_m})\|^p |z_{i_1} \cdots z_{i_m}|^p \\ &\leq \left(\sum_{i \in \mathcal{M}(m,n)} \|H(e_{i_1}, \dots, e_{i_m})\| |z_{i_1} \cdots z_{i_m}| \right)^p \\ &\leq \left[\left(\sum_{i \in \mathcal{M}(m,n)} \|H(e_{i_1}, \dots, e_{i_m})\|^r \right)^{\frac{1}{r}} \left(\sum_{i \in \mathcal{M}(m,n)} |z_{i_1} \cdots z_{i_m}|^s \right)^{\frac{1}{s}} \right]^p \\ &\leq C_r(X)^{mp} \|H\|_{B_{\ell_q^n}}^p \left(\sum_{i \in \mathcal{M}(m,n)} |z_{i_1} \cdots z_{i_m}|^s \right)^{\frac{p}{s}} \\ &= C_r(X)^{mp} \|H\|_{B_{\ell_q^n}}^p (|z_1|^s + \cdots + |z_n|^s)^{\frac{mp}{s}}. \end{aligned}$$

Now, for $q \leq s$, we have $\|z\|_s \leq \|z\|_q$, whereas

$$\|z\|_s \leq n^{\frac{1}{s}-\frac{1}{q}} \|z\|_q \quad \text{for } s < q.$$

Hence, we obtain

$$\sum_{\alpha \in \Lambda(m,n)} \|x_\alpha z^\alpha\|^p \leq \begin{cases} C_r(X)^{mp} \|H\|_{B_{\ell_q^n}}^p \|z\|_q^{mp}, & \text{for } q \leq s \\ C_r(X)^{mp} \left(n^{\frac{1}{s}-\frac{1}{q}}\right)^{mp} \|H\|_{B_{\ell_q^n}}^p \|z\|_q^{mp}, & \text{for } q > s. \end{cases}$$

Therefore, by the virtue of polarization formula for multilinear forms (see e.g. [22, Section 2.6]), for all $z \in B_{\ell_q^n}$ we obtain

$$\sum_{\alpha \in \Lambda(m,n)} \|x_\alpha z^\alpha\|^p \leq \begin{cases} C_r(X)^{mp} \left(\frac{m^m}{m!}\right)^p \|Q\|_{B_{\ell_q^n}}^p, & \text{for } q \leq s \\ C_r(X)^{mp} \left(n^{\frac{1}{s}-\frac{1}{q}}\right)^{mp} \left(\frac{m^m}{m!}\right)^p \|Q\|_{B_{\ell_q^n}}^p, & \text{for } q > s \end{cases}$$

and whence,

$$\sum_{\alpha \in \Lambda(m,n)} \|x_\alpha z^\alpha\|^p \leq \begin{cases} C_r(X)^{mp} e^{mp} \|Q\|_{B_{\ell_q^n}^p}^p, & \text{for } q \leq s \\ C_r(X)^{mp} \left(n^{\frac{1}{s}-\frac{1}{q}}\right)^{mp} e^{mp} \|Q\|_{B_{\ell_q^n}^p}^p, & \text{for } q > s. \end{cases}$$

In view of Definition 4.1, for every $m \in \mathbb{N}$ and $1 \leq p < \infty$ we have

$$K_m^p(B_{\ell_q^n}, X) \leq \begin{cases} \frac{1}{e^{C_r(X)}}, & \text{for } q \leq s \\ \frac{1}{e^{C_r(X)} n^{\frac{1}{s}-\frac{1}{q}}}, & \text{for } q > s. \end{cases}$$

Finally, by applying Lemma 4.4(b), for every $\lambda > 1$, $1 \leq p < \infty$, and $1 \leq q \leq \infty$, we obtain

$$K^p(B_{\ell_q^n}, X, \lambda) \leq \begin{cases} \left(\frac{\lambda-1}{\lambda}\right)^{1/p} \frac{1}{e^{C_r(X)}}, & \text{for } q \leq s \\ \left(\frac{\lambda-1}{\lambda}\right)^{1/p} \frac{1}{e^{C_r(X)} n^{\frac{1}{s}-\frac{1}{q}}}, & \text{for } q > s. \end{cases}$$

This completes the proof. \square

Proof of Theorem 3.4. We note that the proof of (1) is just a consequence of (4.3) and Theorem 3.2. With the help of these observations, we obtain that

$$\begin{aligned} K^p(B_{\ell_q^n}, T, \lambda) &\geq \max \left\{ K^p(B_{\ell_q^n}, X, \lambda / \|T\|^p), K^p(B_{\ell_q^n}, Y, \lambda / \|T\|^p) \right\} \\ &\geq \begin{cases} \left(\frac{\lambda - \|T\|^p}{\lambda}\right)^{\frac{1}{p}} \frac{1}{e^{\min\{C_r(X), C_r(Y)\}}} & \text{if } q \leq s \\ \left(\frac{\lambda - \|T\|^p}{\lambda}\right)^{\frac{1}{p}} \frac{1}{e^{\min\{C_r(X), C_r(Y)\}}} \left(\frac{1}{n}\right)^{\frac{1}{s}-\frac{1}{q}} & \text{if } q > s, \end{cases} \end{aligned}$$

where $s = r/(r-1)$. Hence, the result follows.

Next, for the proof of (2), we partly follow the techniques of the proof of Theorem 3.1 and hypercontractivity of vector-valued Bohnenblust-Hille type inequality. Applying Theorem 3.3 and Hölder's inequality, there is a constant $\tilde{C} > 0$ such that for every m -homogeneous polynomial $Q(z) = \sum_{\alpha \in \Lambda(m,n)} x_\alpha z^\alpha \in \mathcal{P}(m\mathbb{C}^n, X)$, we have

$$\begin{aligned} \sum_{\alpha \in \Lambda(m,n)} \|T(x_\alpha) z^\alpha\|^p &\leq \left[\left(\sum_{\alpha \in \Lambda(m,n)} 1 \right)^{\frac{(r-1)ms-r+s}{rms}} \times \left(\sum_{\alpha \in \Lambda(m,n)} \|T(x_\alpha) z^\alpha\|^{\frac{rsm}{r+(m-1)s}} \right)^{\frac{r+(m-1)s}{rsm}} \right]^p \\ &\leq \left(\sum_{\alpha \in \Lambda(m,n)} 1 \right)^{\frac{((r-1)ms-r+s)p}{rms}} \tilde{C}^{mp} \|Q\|_{\ell_q^n}^p. \end{aligned}$$

We observe that

$$(5.10) \quad \sum_{\alpha \in \Lambda(m,n)} 1 = \binom{n+m-1}{m} \leq e^m \left(1 + \frac{n}{m}\right)^m.$$

By the virtue of (4.1) and (4.2) and using the fact (5.10), there is a constant $E > 0$ such that

$$(5.11) \quad K_m^p(B_{\ell_q^n}, T, \lambda) \geq E \lambda^{\frac{1}{mp}} \left(1 + \frac{n}{m}\right)^{-\frac{(r-1)ms-r+s}{rms}}$$

for every m , $1 \leq p < \infty$, and $\|T\| < \lambda^{1/p}$. By minimizing the right side of (5.11), as demonstrated in the proof of Theorem 3.1, we observe that there is a constant $D > 0$ such that for $1 \leq p < \infty$ and $\|T\| < \lambda^{1/p}$

$$\inf_{m \in \mathbb{N}} \left\{ K_m^p(B_{\ell_q^n}, T, \lambda) \geq C \left(\frac{\log n}{n}\right)^{1-\frac{1}{r}} \right\}.$$

Finally, using Lemma 4.4(a), we obtain desired conclusion. This completes the proof. \square

Acknowledgment: The research of the second named author is supported by DST-INSPIRE Fellowship (IF 190721), Department of Science and Technology, New Delhi, India.

REFERENCES

- [1] L. AIZENBERG, Multidimensional analogues of Bohr's theorem on power series, *Proc. Amer. Math. Soc.* **128** (2000), 1147–1155.
- [2] L. AIZENBERG, A. AYTUNA, and P. DJAKOV, An abstract approach to Bohr's phenomenon, *Proc. Amer. Math. Soc.* **128** (2000), 2611–2619.
- [3] L. AIZENBERG, A. AYTUNA and P. DJAKOV, Generalization of theorem on Bohr for bases in spaces of holomorphic functions of several complex variables, *J. Math. Anal. Appl.* **258** (2001), 429–447.
- [4] V. ALLU, H. HALDER, and S. PAL, Multidimensional Bohr radii for vector-valued holomorphic functions, see <https://arxiv.org/abs/2308.07825> (2023).
- [5] F. BAYART, D. PELLEGRINO, and J. B. SEOANE-SEPÚLVEDA, The Bohr radius of the n -dimensional polydisk is equivalent to $\sqrt{(\log n)/n}$, *Adv. Math.* **264** (2014), 726–746.
- [6] F. BAYART, A. DEFANT, and S. SCHLÜTERS, Monomial convergence for holomorphic functions on ℓ_r , *J. Anal. Math.* **138** (2019), 107–134.
- [7] C. BÉNÉTEAU, A. DAHLNER and D. KHAVINSON, Remarks on the Bohr phenomenon, *Comput. Methods Funct. Theory* **4**(1) (2004), 1–19.
- [8] O. BLASCO, The Bohr radius of a Banach space, *In Vector measures, integration and related topics, 5964, Oper. Theory Adv. Appl., 201, Birkhäuser Verlag, Basel, 2010*.
- [9] O. BLASCO, The p -Bohr radius of a Banach space, *Collect. Math.* **68** (2017), 87–100.
- [10] H. P. BOAS and D. KHAVINSON, Bohr's power series theorem in several variables, *Proc. Amer. Math. Soc.* **125** (1997), 2975–2979.
- [11] H. P. BOAS, Majorant Series, *J. Korean Math. Soc.* **37** (2000), 321–337.
- [12] H. BOHR, A theorem concerning power series, *Proc. Lond. Math. Soc.* s2-13 (1914), 1–5.
- [13] F. BOMBAL, D. PÉREZ-GARCÍA and I. VILLANUEVA, Multilinear extensions of Grothendieck's theorem, *Q. J. Math.* **55** (2004), 441–450.
- [14] E. BOMBIERI, Sopra un teorema di H. Bohr e G. Ricci sulle funzioni maggioranti delle serie di potenze, *Boll. Un. Mat. Ital.* **17** (1962), 276–282.
- [15] E. BOMBIERI and J. BOURGAIN, A remark on Bohr's inequality, *Internat. Math. Res. Notices* **80** (2004), 4307–4330.
- [16] N. DAS, Estimates for generalized Bohr radii in one and higher dimensions, *Canad. Math. Bull.* **66** (2023), 682–699.
- [17] N. DAS, The p -Bohr radius for vector-valued holomorphic and pluriharmonic functions, *Forum Math.* **36** (2024), 765–782.
- [18] A. DEFANT and L. FRERICK, A logarithmic lower bound for multi-dimensional Bohr radii, *Israel J. Math.* **152** (2006), 17–28.
- [19] A. DEFANT and L. FRERICK, The Bohr radius of the unit ball of ℓ_p^n , *J. Reine Angew. Math.* **660** (2011), 131–147.

- [20] A. DEFANT, L. FRERICK, J. ORTEGA-CERDÀ, M. OUNAÏES, and K. SEIP, The Bohnenblust-Hille inequality for homogeneous polynomials in hypercontractive, *Ann. of Math.* **174** (2011), 512–517.
- [21] A. DEFANT, D. GARCÍA, and M. MAESTRE, Bohr power series theorem and local Banach space theory, *J. Reine Angew. Math.* **557** (2003), 173–197.
- [22] A. DEFANT, D. GARCÍA, M. MAESTRE, and P. SEVILLA-PERIS, *Dirichlet Series and Holomorphic Functions in High Dimensions*, New Mathematical Monographs: **37**, Cambridge University Press, Cambridge, (2019).
- [23] A. DEFANT, M. MAESTRE, and C. PRENGEL, The arithmetic Bohr radius, *Q. J. Math.* **59** (2008), 189–205.
- [24] A. DEFANT, M. MAESTRE, and C. PRENGEL, Domains of convergence for monomial expansions of holomorphic functions in infinitely many variables, *J. Reine Angew. Math.* **634** (2009), 13–49.
- [25] A. DEFANT, M. MAESTRE and U. SCHWARTING, Bohr radii of vector valued holomorphic functions, *Adv. Math.* **231** (2012), 2837–2857.
- [26] J. DIESTEL, H. JARCHOW, and A. TONGE, *Absolutely Summing Operators*, in: Cambridge Studies in Advance Mathematics **43**, Cambridge University Press, Cambridge, (1995).
- [27] P. G. DIXON, Banach algebras satisfying the non-unital von Neumann inequality, *Bull. Lond. Math. Soc.* **27** (4) (1995), 359–362.
- [28] P. B. DJAKOV and M. S. RAMANUJAN, A remark on Bohr’s theorem and its generalizations, *J. Anal.* **8** (2000), 65–77.
- [29] S. DINEEN and R. M. TIMONEY, Absolute bases, tensor products and a theorem of Bohr, *Studia Math.* **94** (1989), 227–234.
- [30] D. GALICER, M. MANSILLA and S. MURO, Mixed Bohr radius in several variables, *Trans. Amer. Math. Soc.* **373** (2020), 777–796.
- [31] H. HAMADA, T. HONDA and G. KOHR, Bohr’s theorem for holomorphic mappings with values in homogeneous balls, *Israel J. Math.* **173** (2009), 177–187.
- [32] S. KUMAR and R. MANNA, Multi-dimensional Bohr radii of Banach space valued holomorphic functions, see <https://arxiv.org/pdf/2303.17416> (2023).
- [33] J. LINDENSTRAUSS and L. TZAFRIRI, *Classical Banach Spaces I: Sequence Spaces*, in: Ergebnisse der Mathematik und ihrer Grenzgebiete, **92**, Springer-Verlag, Berlin-Heidelberg-New York, (1977).
- [34] J. LINDENSTRAUSS and L. TZAFRIRI, *Classical Banach Spaces II: Function Spaces*, in: Ergebnisse der Mathematik und ihrer Grenzgebiete, **97**, Springer-Verlag, Berlin-Heidelberg-New York, (1979).
- [35] V. I. PAULSEN, G. POPESCU and DINESH SINGH, On Bohr’s inequality, *Proc. Lond. Math. Soc.* s3-85 (2002), 493–512.
- [36] G. POPESCU, Bohr inequalities for free holomorphic functions on polyballs, *Adv. Math.* **347** (2019), 1002–1053.
- [37] S. SIDON, Über einen satz von Herrn Bohr, *Math. Zeit.* **26** (1927), 731–732.
- [38] M. TOMIC, Sur un theoreme de H. Bohr, *Math. Scand.* **11** (1962), 103–106.

VASUDEVARAO ALLU, SCHOOL OF BASIC SCIENCES, INDIAN INSTITUTE OF TECHNOLOGY BHUBANESWAR, BHUBANESWAR-752050, ODISHA, INDIA.

Email address: avrao@iitbbs.ac.in

SUBHADIP PAL, SCHOOL OF BASIC SCIENCES, INDIAN INSTITUTE OF TECHNOLOGY BHUBANESWAR, BHUBANESWAR-752050, ODISHA, INDIA.

Email address: subhadippal33@gmail.com