(Dated: 2 July 2024)

The (pseudo-)Riemann-metrizability and Ricci-flatness of Finsler spaces with m-Kropina metric $F = \alpha^{1+m}\beta^{-m}$ of Berwald type are investigated. We prove that the affine connection on F can locally be understood as the Levi-Civita connection of some (pseudo-)Riemannian metric if and only if the Ricci tensor of the canonical affine connection is symmetric. We also obtain a third equivalent characterization in terms of the covariant derivative of the 1-form β . We use these results to classify all locally metrizable m-Kropina spaces whose 1-forms have a constant causal character. In the special case where the first de Rahm cohomology group of the underlying manifold is trivial (which is true of simply connected manifolds, for instance), we show that global metrizability is equivalent to local metrizability and hence, in that case, our necessary and sufficient conditions also characterize global metrizability.

Berwald *m*-Kropina Spaces of Arbitrary Signature: Metrizability and Ricci-Flatness

In addition, we further obtain explicitly all Ricci-flat, locally metrizable *m*-Kropina metrics in (3+1)D whose 1-forms have a constant causal character. In fact, the only possibilities are essentially the following two: either α is flat and β is α -parallel, or α is a pp-wave and β is α -parallel.

^{a)}Electronic mail: s.j.heefer@tue.nl

CONTENTS

I. Introduction	2
II. Prerequisites	4
A. Some notational conventions	4
B. Basic notions in Finsler geometry	5
C. Berwald spaces	7
D. Metrizability	7
III. m-Kropina metrics	8
A. The Berwald condition	9
IV. Characterization of metrizability	9
A. The main theorem for local metrizability	9
B. Special types of 1-forms b	12
1. b is nowhere null	12
2. b is null and closed	13
3. b is null but not necessarily closed	13
C. Global metrizability	15
V. Ricci-flatness of <i>m</i> -Kropina spaces in $(1+3)D$	17
A. Ricci-flat nowhere-null m -Kropina metrics in $(1+3)D$	17
B. Ricci-flat null <i>m</i> -Kropina metrics in $(1+3)D$	19
VI. Discussion	21
VII. Acknowledgements	22
References	22

I. INTRODUCTION

A Berwald space is a Finsler space for which the canonical nonlinear connection reduces to a linear connection on the tangent bundle TM. This affine connection is necessarily torsionfree. The natural question thus arises whether this affine connection arises as the Levi-Civita connection of some pseudo-Riemannian metric. Simply put, is (the affine connection on) every Berwald space metrizable? In the classical positive definite setting, this is indeed the case, as was first shown in (Szabó, 1981)¹.

The proofs of this result^{1,2} rely on procedures such as averaging³ over the indicatrix, for which it is essential that the Finsler metric F is sufficiently smooth and defined everywhere on the slit tangent bundle TM_0 . In the case of Finsler metrics of indefinite signature, however, the domain $\mathcal{A} \subset TM$ where F is defined is typically only a proper subset of TM_0 and hence the proofs do not extend to this case. And even in the special case that $\mathcal{A} = TM_0$, the fact that the indicatrix is not compact in indefinite signatures poses problems.

It was shown in (Fuster et al., 2020)⁴ that Szabó's metrization theorem indeed does not generalize to the more general setting. In (Heefer et al., 2023)⁵, the situation was further investigated and necessary and sufficient conditions for local metrizability were obtained in the specific case of *m*-Kropina metrics with closed null 1-form. More recently, the metrizability of spherically symmetric Berwald spaces was investigated in (Voicu et al., 2024)⁶.

Here we extend the analysis of (Heefer et al., 2023)⁵ to arbitrary *m*-Kropina metrics, i.e. Finsler metrics of the form $F = \alpha^{1+m}\beta^{-m}$, with no restriction on the 1-form. The culprit behind all counterexamples to Szabó's theorem obtained in (Heefer et al., 2023)⁵ was the fact that the Ricci tensor constructed from the affine connection—the 'affine Ricci tensor'—is in general not symmetric. This property—symmetry of the affine Ricci tensor—is in general a necessary condition for local metrizability, and for *m*-Kropina metrics with closed null 1-form it was shown to be a necessary condition as well⁵.

Here we show that this results extends to all *m*-Kropina metrics, i.e. we prove that any *m*-Kropina metric is locally metrizable if and only if its affine Ricci tensor is symmetric. Additionally, we further obtain a third necessary and sufficient condition for local metrizability in terms of the covariant derivative of the 1-form β . We use these results to classify all locally metrizable *m*-Kropina spaces whose 1-forms have a constant causal character.

We also briefly touch upon the issue of global metrizability. We do not obtain a conclusive characterization of global metrizability in general, but in the special case where the first de Rahm cohomology group of the underlying manifold is trivial (which is true of simply connected manifolds, for instance), we show that global metrizability is equivalent to local metrizability. Hence, in this case, our necessary and sufficient conditions for local metrizability also characterize global metrizability.

Finally, as an application of our results established thus far, we obtain explicitly all Ricci-flat, locally metrizable *m*-Kropina metrics in (3 + 1)D whose 1-forms have a constant causal character. Here we note that an *m*-Kropina space is locally metrizable and Ricci-flat if and only if it is affinely Ricci flat—meaning that its affine Ricci tensor vanishes. Affine Ricci-flatness turns out to be rather restrictive property. In fact, the only possibilities are essentially the following two: either α is flat and β is α -parallel, or α is a pp-wave and β is α -parallel.

In the literature, *m*-Kropina metrics have been used for modeling the geometry of spacetime in Finsler extensions of Einstein's general theory of relativity.^{7–12}. Indeed *m*-Kropina metrics have been found to yield exact solutions^{8,9,13} to the Pfeifer and Wohlfarth's Finslerian extension of Einstein's field equation^{14–16}; see (Heefer, 2024)⁷ for a recent review. Our results put strict bounds on the remaining possibilities for solutions of Berwald *m*-Kropina type to this field equation.

Many of the results obtained in this article were first described in the PhD thesis (Heefer, $2024)^7$.

II. PREREQUISITES

Below we introduce our notational conventions and we briefly recall several relevant notions in Finsler geometry. For a more in-depth treatment, the reader is referred to $(\text{Heefer}, 2024)^7$.

A. Some notational conventions

In what follows, we will often work in local coordinates. Given a smooth manifold M of dimension n we will typically assume that some chart $\varphi : U \subset M \to \mathbb{R}^n$ is provided, and we will effectively identify any $p \in U$ with its image $x = (x^1, \ldots, x^n) \coloneqq \varphi(p) \in \mathbb{R}^n$ under φ . If $p \in U$ then each $X_p \in T_p M$ can be written as $X_p = y^i \partial_i |_p \rightleftharpoons y^i \partial_i$ (we will often be sloppy and suppress the dependence on the point p), where the tangent vectors $\partial_i \coloneqq \frac{\partial}{\partial x_i}$ make up the coordinate basis of $T_p M$. This decomposition provides natural local coordinates on the tangent bundle TM via the chart

$$\tilde{\varphi}: TU \to \mathbb{R}^n \times \mathbb{R}^n, \qquad \tilde{\varphi}(p, Y) = (\varphi(p), y^1, \dots, y^n) \rightleftharpoons (x, y),$$
(1)

where TU is given by

$$TU = \bigcup_{p \in U} \{p\} \times T_p M \subset TM.$$
⁽²⁾

These local coordinates on TM in turn provide a natural basis $\{\partial_i, \bar{\partial}_i\}_{i=1,...,n}$ of its tangent space $T_{(x,y)}TM$ at (x, y), where we define

$$\partial_i \coloneqq \frac{\partial}{\partial x^i}, \qquad \bar{\partial}_i \coloneqq \frac{\partial}{\partial y^i}.$$
 (3)

Finally, we will use the notation $dx^i dx^j$ for the symmetrized tensor product of 1-forms, i.e. $dx^i dx^j \equiv \frac{1}{2} (dx^j \otimes dx^j + dx^j \otimes dx^j)$, and whenever we work specifically in Lorentzian signature we will adhere to the sign convention (-, +, ..., +).

B. Basic notions in Finsler geometry

For our purposes, a Finsler space will be a triple (M, \mathcal{A}, F) , where M is a smooth manifold, \mathcal{A} a conic subbundle of TM, and F is a Finsler metric on \mathcal{A} , defined as follows.

Definition 1. A conic subbundle of TM is an open subset $\mathcal{A} \subset TM$ which is conic in the sense that $(x, \lambda y) \in \mathcal{A}$ for any $(x, y) \in \mathcal{A}$ and $\lambda > 0$, and which satisfies $\pi(\mathcal{A}) = M$, where $\pi : TM \to M$ is the canonical projection of the tangent bundle. The latter property says that the fibers \mathcal{A}_x of \mathcal{A} are nonempty.

Definition 2. A Finsler metric on a conic subbundle \mathcal{A} is a smooth map $F : \mathcal{A} \to [0, \infty)$ such that

• F is positively homogeneous of degree one:

$$F(x,\lambda y) = \lambda F(x,y), \quad \forall \lambda > 0;$$
(4)

• The fundamental tensor, with components $g_{ij} = \bar{\partial}_i \bar{\partial}_j \left(\frac{1}{2}F^2\right)$, is nondegenerate.

The length of a curve $\gamma: (a, b) \to M$ (whose lift stays within \mathcal{A}) is then defined as

$$\ell(\gamma) = \int_{a}^{b} F(\gamma(\lambda), \dot{\gamma}(\lambda)) \,\mathrm{d}\lambda, \qquad \dot{\gamma} = \frac{\mathrm{d}\gamma}{\mathrm{d}\lambda}.$$
(5)

We will sometimes omit the specification of \mathcal{A} and say, loosely speaking, that F is a Finsler metric on M and L a Finsler Lagrangian on M. We say that F or L is positive definite if

 g_{ij} is positive definite, and more generally that F or L has a certain signature if g_{ij} has that signature.

The connection coefficients of the canonical torsion-free, metric-compatible homogeneous nonlinear connection are given by

$$N_i^j = \frac{1}{4}\bar{\partial}_i \left[g^{jk} \left(y^m \partial_m \bar{\partial}_k L - \partial_k L \right) \right].$$
(6)

This nonlinear connection yields a smooth decomposition of the tangent bundle $T\mathcal{A}$ into a horizontal and a vertical subbundle¹⁷,

$$T\mathcal{A} = H\mathcal{A} \oplus V\mathcal{A},\tag{7}$$

where \oplus denotes the Whitney sum of vector bundles. The vertical subbundle $V\mathcal{A}$ is canonically defined as

$$V\mathcal{A} = \ker(\mathrm{d}\pi) = \operatorname{span}\left\{\bar{\partial}_i\right\}.$$
(8)

The horizontal subbundle, on the other hand, is defined by the nonlinear connection as

$$H\mathcal{A} = \operatorname{span}\left\{\delta_i\right\}, \qquad \delta_i \equiv \partial_i - N^j{}_i\bar{\partial}_j, \tag{9}$$

where we have also introduced the horizontal derivatives δ_i .

The curvature tensor of the nonlinear connection has components

$$R_{ij}^k = \delta_i N_j^k - \delta_j N_i^k, \tag{10}$$

and from this we can further define the following two quantities:

$$\operatorname{Ric} = R^{i}{}_{ij}y^{j}, \qquad R_{ij} = \frac{1}{2}\bar{\partial}_{i}\bar{\partial}_{j}\operatorname{Ric}.$$
(11)

In the literature, R_{ij} is often just referred to as the Ricci tensor, but we will give it the name *Finsler-Ricci tensor* to distinguish it from the *affine Ricci tensor* that will be introduced momentarily for Berwald spaces. Similarly, we will refer to Ric as the *Finsler-Ricci scalar*. From homogeneity and Euler's theorem, it follows that the second relation in (11) can be inverted as

$$\operatorname{Ric} = R_{ij} y^i y^j. \tag{12}$$

Ric and R_{ij} thus contain the same information, and in particular, we have Ric = 0 if and only if $R_{ij} = 0$, in which the Finsler space is said to be *Ricci-flat*.

C. Berwald spaces

A Finsler space is said to be of *Berwald* type, or simply Berwald, if the canonical nonlinear connection reduces to a (necessarily smooth) linear connection on TM, or in other words, an affine connection on the base manifold M, meaning that the connection coefficients are of the form $N_i^k = \Gamma_{ij}^k(x)y^j$ for certain coefficients $\Gamma_{ij}^k(x)$. In this case, the functions Γ_{jk}^i are precisely the Christoffel symbols of said affine connection on M. We will refer to this affine connection as the associated affine connection, or simply the affine connection on the Berwald space. It follows from the torsion-freeness of the nonlinear connection that the affine connection is torsion-free as well.

The curvature tensors (10)-(11) of a Berwald space can be written as

$$R_{ij}^{k} = \bar{R}_{\ell ij}^{k}(x)y^{\ell}, \quad \text{Ric} = \bar{R}_{ij}(x)y^{i}y^{j}, \quad R_{ij} = \frac{1}{2}\left(\bar{R}_{ij}(x) + \bar{R}_{ji}(x)\right)$$
(13)

in terms of the curvature tensor and Ricci tensor of the affine connection, given by

$$\bar{R}^{k}{}_{\ell i j} = \partial_{i}\Gamma^{k}_{j\ell} - \partial_{j}\Gamma^{k}_{i\ell} + \Gamma^{k}_{im}\Gamma^{m}_{j\ell} - \Gamma^{k}_{jm}\Gamma^{m}_{i\ell}, \qquad \bar{R}_{lk} = \bar{R}^{i}{}_{lik}, \tag{14}$$

respectively. We will refer to \bar{R}^{k}_{lij} and \bar{R}_{lk} as (the components of) the affine curvature tensor and affine Ricci tensor, respectively. Since the last identity in (13) will prove to be of particular interest later, we restate it as a lemma.

Lemma 3. The Finsler-Ricci tensor of a Berwald space is the symmetrization of the affine Ricci tensor.

We say that a Berwald space is affinely Ricci-flat if $\bar{R}_{ij} = 0$. Note that this is a stronger condition than (standard) Ricci-flatness, $R_{ij} = 0$.

D. Metrizability

Given a Finsler space of Berwald type, the canonical nonlinear connection reduces to a linear connection on TM that is torsion-free. This leads to the natural question of whether this linear connection is 'metrizable'.

Definition 4. Let F be a Berwald metric with underlying manifold M. We say that (the canonical affine connection of) F is metrizable if it arises as the Levi-Civita connection of some pseudo-Riemannian metric on M.

For readability, we usually omit the phrase 'the canonical affine connection of' in between brackets and we simply say that F is metrizable. A slightly weaker property is that of local metrizability.

Definition 5. A Berwald space with underlying manifold M is said to be locally metrizable if each point in M has a neighborhood that admits a pseudo-Riemannian metric whose Levi-Civita connection coincides with the affine connection restricted to that neighborhood.

III. *m*-KROPINA METRICS

An *m*-Kropina metric (also called generalized Kropina metric, Bogoslovsky metric or Bogoslovsky-Kropina metric) is a Finsler metric of the form

$$F = \alpha^{1+m} \beta^{-m},\tag{15}$$

where the building blocks

$$\alpha = \sqrt{|a_{ij}y^i y^j|}, \qquad \beta = b_i y^i \tag{16}$$

are constructed from a (pseudo-)Riemannian metric $a = a_{ij}(x)dx^i dx^j$ and a 1-form $b = b_i(x)dx^i$ and m, respectively, and where m is a real parameter. By a slight abuse of terminology one also refers to α and β simply as the (pseudo-)Riemannian metric and the 1-form, respectively. We remark that it follows from our definition of a Finsler metric that the 1-form b is nowhere vanishing, which we will henceforth take for granted. When m = 1 the m-Kropina metric reduces to the standard Kropina metric¹⁸ $F = \alpha^2/\beta$.

In the physics literature, spacetimes with a metric of *m*-Kropina type have also been dubbed Very General Relativity (VGR) spacetimes⁸ or General Very Special Relativity (GVSR) spacetimes¹⁰, introduced as generalizations of Very Special Relativity (VSR)^{11,12} spacetimes. Indeed, as an important application, *m*-Kropina metrics have been found to yield exact solutions^{8,9,13} to the Pfeifer and Wohlfarth's Finslerian extension of Einstein's field equation^{14–16}; see (Heefer, 2024)⁷ for a recent review.

We further introduce the notation

$$|b|^2 \equiv a_{ij}b^i b^j \tag{17}$$

for the squared norm of β with respect to α . Also, throughout the remainder of this article, all indices are raised and lowered with a_{ij} , and $\overset{\circ}{\nabla}$ denotes the Levi-Civita connection of a_{ij} .

A. The Berwald condition

As shown recently⁵ (see also (Heefer, 2024)⁷ for a more in-depth treatment and an alternative proof), an m-Kropina metric $F = \alpha^{1+m}\beta^{-m}$ with $n = \dim M > 2$ is of Berwald type if and only if there exists a smooth vector field with components f^i on M such that

$$\overset{\circ}{\nabla}_j b_i = m(f_k b^k) a_{ij} + b_i f_j - m f_i b_j.$$
(18)

Here and throughout the remainder of the article, $\mathring{\nabla}$ denotes the Levi-Civita connection of a_{ij} . This Berwald condition agrees with the one obtained by Matsumuto¹⁹, which was proven under slightly different assumptions.

Whenever the condition (18) is satisfied, the affine connection coefficients of the Berwald m-Kropina space can be expressed in terms of the Christoffel symbols $\mathring{\Gamma}_{ij}^k$ of the Levi-Civita connection corresponding to α as follows^{5,7,19}

$$\Gamma_{ij}^{\ell} = \mathring{\Gamma}_{ij}^{\ell} + \Delta \Gamma_{ij}^{k}, \qquad \Delta \Gamma_{ij}^{k} = ma^{\ell k} \left(a_{ij} f_k - a_{jk} f_i - a_{ki} f_j \right)$$
(19)

This implies in particular that for any given data (α, β) there can be at most one vector field f^k satisfying (18). (This can be seen for instance by considering $\Delta\Gamma_{ij}^{\ell}$ with $i = j \neq \ell$.)

IV. CHARACTERIZATION OF METRIZABILITY

From here onwards we will focus on *m*-Kropina metrics of Berwald type with $n = \dim M > 2$. The main aim in this section is answer the following question:

Q: Under what conditions can the affine connection on a Berwald m-Kropina space be understood, locally, as the Levi-Civita connection of a pseudo-Riemannian metric?

In §IV A, we obtain necessary and sufficient conditions for local metrizability. Then we discuss several special cases in §IV B. And finally, in §IV C, we have a brief discussion on global metrizability.

A. The main theorem for local metrizability

It is a standard result that the Ricci tensor of any Levi-Civita connection is symmetric. This immediately implies that symmetry of the Ricci tensor constructed from the affine connection of a Berwald space—that is, symmetry of the affine Ricci tensor (see Section II C)—is a necessary condition for metrizability. To characterize this condition, we will first derive a simple formula for the skew-symmetric part of the affine Ricci tensor, denoted by $\mathcal{A}(\bar{R})$ and having components $\mathcal{A}(\bar{R})_{ij} = \bar{R}_{[ij]} = \frac{1}{2} (\bar{R}_{ij} - \bar{R}_{ji})$. Moreover, to prevent confusion, we remark that the expression df below denotes the exterior derivative of the 1-form f with components $f_i = a_{ij}f^j$ appearing in the Berwald condition (18). We also recall that $n = \dim M$.

Lemma 6. The skew-symmetric part of the affine Ricci tensor is given by

$$\mathcal{A}(\bar{R}) = -\frac{mn}{2}df \tag{20}$$

Proof. From the definition (14) of the affine Ricci tensor of a Berwald space and the fact that $\Gamma_{ij}^k = \Gamma_{ji}^k$ it follows that the skew-symmetric part of the affine Ricci tensor can be written as

$$\bar{R}_{[ij]} \equiv \frac{1}{2} \left(\bar{R}_{ij} - \bar{R}_{ji} \right) = \partial_{[i} \Gamma^k_{j]k}.$$
⁽²¹⁾

Since the Levi-Civita connection $\mathring{\Gamma}_{ij}^k$ has a symmetric Ricci tensor, it follows that $\partial_{[i}\mathring{\Gamma}_{j]k}^k = 0$ and hence we have

$$\bar{R}_{[ij]} = \partial_{[i}\Gamma^k_{j]k} = \partial_{[i}\Delta\Gamma^k_{j]k} = \frac{1}{2} \left(\partial_i\Delta\Gamma^k_{kj} - \partial_j\Delta\Gamma^k_{ki} \right).$$
(22)

Employing the expression (19) for $\Delta\Gamma_{ij}^k$, we find that

$$\Delta\Gamma_{kj}^k = -mnf_j \tag{23}$$

and this yields

$$\bar{R}_{[ij]} = \frac{mn}{2} (\partial_j f_i - \partial_i f_j), \qquad (24)$$

which is precisely the coordinate expression of (20), as desired.

We then have the following characterization of local metrizability.

Theorem 7. For an *m*-Kropina metric $F = \alpha^{1+m}\beta^{-m}$ of Berwald type with dim M > 2, the following are equivalent:

(i) F is locally metrizable;

(ii) The affine Ricci tensor is symmetric, $\bar{R}_{ij} = \bar{R}_{ji}$;

(iii) f_i is a closed 1-form.

In this case we can locally write $f_i = \partial_i \psi$ for some $\psi \in C^{\infty}(M)$ and we may define $\tilde{a}_{ij} = e^{-2m\psi}a_{ij}$ and $\tilde{b}_i = e^{-(1+m)\psi}b_i$ and define $\tilde{\alpha}$ and $\tilde{\beta}$ accordingly. Then the following statements hold:

- F can be written as $F = \tilde{\alpha}^{1+m} \tilde{\beta}^{-m}$;
- $\tilde{\beta}$ is parallel with respect to $\tilde{\alpha}$;
- \tilde{a}_{ij} provides a local metrization of F.

Proof. We will prove that $(i) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (i)$. The first two implications are rather straightforward.

(i) \Rightarrow (ii): If F is locally metrizable then it follows trivially that the Ricci tensor is symmetric.

(ii) \Rightarrow (iii): If the Ricci tensor is symmetric then Lemma 6 shows that df = 0.

The remaining nontrivial part of the proof amounts to showing that (iii) implies (i).

(iii) \Rightarrow (i): If f_i is closed then f is locally exact, i.e. $f_i = \partial_i \psi$ for some (locally defined) smooth function ψ on M. Consider the following transformed metric and 1-form, $\tilde{a}_{ij} = e^{-2m\psi}a_{ij}$ and $\tilde{b}_i = e^{-(1+m)\psi}$ and define $\tilde{\alpha}$ and $\tilde{\beta}$ accordingly. Then $F = \alpha^{1+m}\beta^{-m} = \tilde{\alpha}^{1+m}\tilde{\beta}^{-m}$ and it is straightforward to show that the Christoffel symbols of \tilde{a}_{ij} are given by

$$\tilde{\Gamma}_{ij}^{k} = \mathring{\Gamma}_{ij}^{k} - m\left(f_{i}\delta_{j}^{k} + f_{j}\delta_{i}^{k} - f^{k}a_{ij}\right)$$
(25)

in terms of those of a_{ij} . Hence if we denote by $\tilde{\nabla}$ the Levi-Civita connection of $\tilde{\alpha}$ then we find that

$$\tilde{\nabla}_j \tilde{b}_i = e^{-(1+m)\psi} \left[\mathring{\nabla}_j b_i - (m(f_k b^k) a_{ij} + b_i f_j - m f_i b_j) \right] = 0,$$
(26)

which vanishes, by (18). In other words, \tilde{b}_i is parallel w.r.t. \tilde{a}_{ij} . Since we can write F as $F = \tilde{\alpha}^{1+m}\tilde{\beta}^{-m}$, it follows from Prop. 8 below that the affine connection on F is just the Levi-Civita connection of \tilde{a}_{ij} . Hence F is locally metrizable, namely by \tilde{a}_{ij} , completing the proof.

The mapping $(\alpha, \beta) \mapsto (\tilde{\alpha}, \tilde{\beta})$ is a generalization of what is called the *f*-change in (Matsumoto, 2003)¹⁹. In the proof we have made use of the following well-known result; see e.g. (Heefer, 2024)⁷.

Proposition 8. Let $F = \alpha \phi(\beta/\alpha)$ be an (α, β) -metric and suppose that $\mathring{\nabla}_i b_j = 0$. Then F is Berwald and the affine connection of F coincides with that of α .

B. Special types of 1-forms b

Having established the general criteria for local metrizability in Theorem 7, we now turn to several specific possibilities with respect to the causal character of the 1-form b:

- 1. b is nowhere null;
- 2. b is null and closed;
- 3. b is null but not necessarily closed.

1. b is nowhere null

In this case, we can show that f_i is necessarily closed¹⁹, and hence our theorem guarantees metrizability. This case was already known to Matsumoto¹⁹. To see that f_i is closed, note that contracting (18) with b^i leads to

$$\frac{1}{2} \mathring{\nabla}_j \left(|b|^2 \right) = |b|^2 f_j, \tag{27}$$

from which it follows, as long as b is nowhere null, that

$$f_j = \partial_j \ln\left(\sqrt{||b|^2|}\right). \tag{28}$$

Hence we have the following corollaries.

Corollary 9. An *m*-Kropina space with $|b|^2 \neq 0$ is locally metrizable.

Corollary 10. An *m*-Kropina space with positive definite α is locally metrizable.

2. b is null and closed

This case was the subject of our earlier paper⁵. We restate the main relevant result here.

Proposition 11. If b is null and closed then F is locally metrizable if and only if around each point in M there exist coordinates $(u, v, x^3, ..., x^n)$ such that b = du and

$$a = -2 \, du \, dv + \left[\tilde{H}(u, x) + \rho(u) v \right] \, du^2 + 2 W_a(u, x) \, du \, dx^a + h_{ab}(u, x) \, dx^a \, dx^b, \tag{29}$$

where $a, b = 3, \ldots n$ and $\tilde{H}, \rho, W_a, h_{ab}$ are smooth functions.

In this case, the affine connection restricted to the chart corresponding to the coordinates $(u, v, x^3, \ldots x^n)$ is metrizable by the following pseudo-Riemannian metric:

$$\tilde{a} = e^{\frac{m}{1-m}\int^{u}\rho(\tilde{u})\mathrm{d}\tilde{u}}a\tag{30}$$

3. b is null but not necessarily closed

Finally, if b is null but not necessarily closed then the local metrizability of F can be characterized in a way similar to that of Prop. 11, as we will show next.

Proposition 12. If $|b|^2 = 0$ then F is locally metrizable if and only if around each point in M there exist coordinates $(u, v, x^3, ..., x^n)$ such that

$$a = e^{2m\psi} \left(-2\,du\,dv + H\,du^2 + 2W_a\,du\,dx^a + h_{ab}\,dx^a\,dx^b \right),\tag{31}$$

$$b = e^{(1+m)\psi} du,\tag{32}$$

where ψ is any smooth function on M, $a, b = 3, \ldots n$ and $\tilde{H}, \rho, W_a, h_{ab}$ are smooth functions depending only on u and x^a .

In this case, F can be written as

$$F = \left| -2 \, du \, dv + H \, du^2 + 2 W_a \, du \, dx^a + h_{ab} \, dx^a \, dx^b \right|^{(1+m)/2} (du)^{-m} \tag{33}$$

and the metric $-2 du dv + H du^2 + 2W_a du dx^a + h_{ab} dx^a dx^b$ provides a local metrization of F.

Proof. We first prove the 'only if' part. By Theorem 7, if F is locally metrizable then there exist $\tilde{a}_{ij} = e^{-2m\psi}a_{ij}$ and $\tilde{b}_i = e^{-(1+m)\psi}b_i$ with corresponding $\tilde{\alpha}$ and $\tilde{\beta}$ such that $F = \tilde{\alpha}^{1+m}\tilde{\beta}^{-m}$ and such that $\tilde{\beta}$ is parallel with respect to \tilde{a}_{ij} . It is a well-known result that any pseudo-Riemannian metric \tilde{a}_{ij} admitting a nowhere vanishing parallel null 1-form \tilde{b} , admits coordinates $(u, v, x^3, \dots x^n)$ around any point such that

$$\tilde{a} = -2\mathrm{d}u\mathrm{d}v + H\mathrm{d}u^2 + 2W_a\mathrm{d}u\mathrm{d}x^a + h_{ab}\mathrm{d}x^a\mathrm{d}x^b,\tag{34}$$

$$\tilde{b} = \mathrm{d}u,\tag{35}$$

where the functions H, W_a, h_{ab} depend only on u and x^a , and where $a = 3, 4, \ldots n$. It is immediately verified that a, b, and F now attain the desired form, completing the 'only if' direction of the proof.

Conversely, if a and b have the stated form, then F is given by (33) in terms of \tilde{a} and \tilde{b} , and since $\tilde{b} = du$ is parallel with respect to $\tilde{\alpha}$, we must have $f_k = 0$ in the Berwald condition (18). It thus follows from (19) that $\tilde{\alpha}$ provides a local metrization of F, completing the proof.

To end this section, we note that as a special case of Prop. 12 we recover Prop. 11, thus providing an alternative proof of the result.

Proof of Prop. 11 from Prop. 12. Suppose first that F is locally metrizable, so that a and b are given by (31) and (32), where the metric functions do not depend on the coordinate v. The fact that the 1-form $b = e^{(1+m)\psi} du$ is closed implies that $\psi = \psi(u)$. In this case, it can be verified that the coordinate transformation given by $\bar{u} = \int^{u} e^{(1+m)\psi(u)} du$ and $\bar{v} = e^{(m-1)\psi(u)}v$ brings the 1-form into the form $b = d\bar{u}$ (as desired) and the metric into the form

$$a = -2\mathrm{d}\bar{u}\mathrm{d}\bar{v} + \bar{H}\mathrm{d}\bar{u}^2 + 2\bar{W}_a\mathrm{d}\bar{u}\mathrm{d}x^a + \bar{h}_{ab}\mathrm{d}x^a\mathrm{d}x^b,\tag{36}$$

where

$$\bar{H} = e^{-2\psi} \left[H + 2(m-1)\psi'v \right], \quad \bar{W}_a = e^{(m-1)\psi}W_a, \quad \bar{h}_{ab} = e^{2m\psi}h_{ab}.$$
(37)

Identifying $\tilde{H}(\bar{u}, x^a) = e^{-2\psi(u(\bar{u}))}H$ and $\rho(\bar{u}) = 2e^{-(1+m)\psi(u(\bar{u}))}(m-1)\psi'(u(\bar{u}))$ it is easily verified that *a* also attains the desired form (29) in the new coordinates. This proves the implication to the right.

Conversely, if a and b have the stated form, then we may define $\psi(u)$ such that $\rho(u) = 2e^{-(1+m)\psi}(m-1)\psi' = 2(m-1)d\psi/d\bar{u}$, by setting $\psi(u) = \frac{1}{2(m-1)}\int^{\bar{u}(u)}\rho(u)du$, and then we can simply perform the same coordinate transformation backward. It is clear that this results in a metric and 1 form of the form (31) and (32) and hence, by Prop. 12, F is locally

metrizable, proving the implication to the left. Furthermore, the local metrization of Prop. 12 given by $-2dudv + Hdu^2 + 2W_a dudx^a + h_{ab} dx^a dx^b$ turns under this transformation into (30), showing that (30) provides a local metrization for F, completing the proof.

C. Global metrizability

Note that in the proof of Theorem 7 we made use of the fact that df = 0 implies that f is a locally exact 1-form. Here f is the 1-form appearing in the Berwald condition (18). In the special case where f is (globally) exact, the proof guarantees that the metric \tilde{a} providing the metrization is in fact globally defined, implying that the Finsler space under consideration is globally metrizable. Hence we have the following.

Proposition 13. If f is an exact 1-form then F is (globally) metrizable.

In general, not every closed 1-form is (globally) exact, and hence the proof of the theorem does not tell us whether local metrizability is equivalent to global metrizability. In some cases it is, however.

Theorem 14. Suppose the first de Rahm cohomology group of M is trivial. Then following are equivalent:

- (i) F is metrizable;
- (ii) F is locally metrizable;
- (iii) The affine Ricci tensor is symmetric;
- (iv) f_i is a closed 1-form.

In that case, writing $f_i = \partial_i \psi$, a global metrization of F is given by $\tilde{a}_{ij} = e^{-2m\psi}a_{ij}$ and moreover, $\tilde{b}_i = e^{-(1+m)\psi}b_i$ is parallel with respect to \tilde{a}_{ij} .

Proof. We only need to prove the equivalence of global metrizability with the other three (already equivalent) statements. Since global metrizability trivially implies local metrizability, it suffices to show that f being closed implies that F is metrizable. So suppose that f is closed. Then since the first de Rahm cohomology group is trivial, f is exact, $f = d\phi$, with ϕ defined globally. The proof of Theorem 7 shows that $\tilde{a} = e^{2\phi}a$ provides a global metrization. Hence F is globally metrizable.

Since any simply connected manifold has a trivial first de Rahm cohomology group, we have, in particular, the following result.

Corollary 15. If M is simply connected then (i)-(iv) are equivalent.

The remaining question is the following one:

Q: Do locally but not globally metrizable m-Kropina spaces exist at all?

This is still an open question and we defer its analysis to future work, even though it seems likely that the answer should be affirmative. It is clear, at least, that such a space would need to have a 1-form f appearing in the Berwald condition that is **closed but not exact**. For if it were not closed, the space would not be locally metrizable and hence not metrizable; and if it were exact, then it would be globally metrizable. A straightforward method for trying to produce a counterexample is thus given as follows.

- Choose your favorite not simply connected manifold M (or more generally any manifold M with nontrivial first de Rahm cohomology group).
- 2. Pick a 1-form f that is closed but not exact.
- 3. Pick a pseudo-Riemannian metric α .
- 4. Try to solve the Berwald condition for β .

An obvious first guess in dimension 3 would be to let $M = \mathbb{R}^2 \times S^1$ with the metric $ds^2 = dx^2 + dy^2 + d\theta^2$ and $f = d\theta$, where (x, y) are the standard coordinates on \mathbb{R}^2 and θ is the standard angle coordinate on the circle S^1 . The 1-form $f = d\theta$ is the prime example of a closed but not exact 1-form, of course. In this case, which is relatively computable, the system defined by the Berwald condition turns out not to be solvable, however. That is, there is no 1-form β that leads to the prescribed $f = d\theta$. The same is true if one instead uses the metric $ds^2 = dx^2 + dy^2 + y^2 d\theta^2$, which corresponds to the standard flat metric $ds^2 = dx^2 + d\tilde{y}^2 + dz^2$ on $\mathbb{R}^3 \setminus \{0\} \cong \mathbb{R}^2 \times S^1$, where y is basically the radial coordinate in the $\tilde{y}z$ -plane. Hence, in order to produce counterexamples—if these exist (which we do expect)—one needs to consider more elaborate examples of manifolds with closed but not exact 1-forms. We leave this to future study.

V. RICCI-FLATNESS OF *m*-KROPINA SPACES IN (1+3)D

We now turn to Ricci-flatness. Recall that a Finsler space is said to be Ricci-flat if the Finsler-Ricci tensor vanishes, $R_{ij} = 0$, or equivalently, if the Ricci scalar vanishes, Ric = 0. We restrict ourselves in this section to the case where M is four-dimensional with a_{ij} having Lorentzian signature. This is the case that is most relevant to applications in gravitational physics. We'll refer to such metrics as (1+3)D *m*-Kropina metrics.

A. Ricci-flat nowhere-null *m*-Kropina metrics in (1+3)D

First, we characterize Ricci-flatness of *m*-Kropina metrics in (1+3)D with nowhere-null 1-form, $|b|^2 \neq 0$.

Proposition 16. A (1+3)D m-Kropina space with $|b|^2 \neq 0$ is Ricci-flat if and only if it is locally of the form

$$F = \left| \eta_{ij} dx^{i} dx^{j} \right|^{(1+m)/2} \left(c_{i} dx^{i} \right)^{-m}$$
(38)

with $[\eta_{ij}] = diag(-1, 1, 1, 1)$ and $c_i = const.$

Proof. We first prove the 'if' direction, i.e. we assume F is of the form (38) and prove that F is Ricci-flat. The 1-form $c_i dx^i$ is parallel with respect to η_{ij} , and hence it follows by Prop. 8 that F has the same affine connection as η , which is Ricci-flat. Hence the Ricci tensor constructed from the affine connection of F—the affine Ricci tensor—vanishes as well, i.e. F is affinely Ricci-flat. This implies, in particular, that F is Ricci-flat, by Lemma 3.

Conversely, to prove the 'only if' direction, suppose that F is Ricci-flat. Then, defining $\tilde{\alpha}, \tilde{\beta}$ as in Theorem 7, we may write $F = \tilde{\alpha}^{1+m}\tilde{\beta}^{-m}$ and in this case F has the same affine connection as $\tilde{\alpha}$, so since F is Ricci-flat, $\tilde{\alpha}$ is Ricci-flat. But by Theorem 7, $\tilde{\alpha}$ admits a non-null parallel 1-form, namely \tilde{b} , so if $\tilde{\alpha}$ is Ricci-flat it must actually be flat, by Lemma 17 below. Hence, in suitable coordinates, it can be written as $\tilde{a} = \eta_{ij} dx^i dx^j$. Since $\tilde{\beta}$ is parallel with respect to this metric, it must have constant coefficients in these coordinates, and (38) follows, completing the proof.

In the proof we have made use of the following well-known result; see e.g. (Heefer, 2024)⁷ for a self-contained proof.

Lemma 17. Any Ricci-flat Lorentzian metric admitting a non-null parallel vector field or 1-form is flat.

We remark that the *representation* of F in terms of a specific choice of α and β is, of course, not unique (we can multiply both α and β by suitably related conformal factors without changing F), and the proposition does not imply that α has to be equal to the Minkowski metric η . However, since any representative α is related to the $\tilde{\alpha}$ from the proposition by a conformal transformation, we have the following two corollaries.

Corollary 18. If F is Ricci-flat and $|b|^2 \neq 0$ then α is conformally flat.

By 'flat coordinates' below we mean coordinates in which the metric takes the form of the Minkowski metric.

Corollary 19. If α is Ricci-flat and $|b|^2 \neq 0$ then F is Ricci-flat only if α is flat and β has constant components in flat coordinates.

Proof. If F is Ricci-flat and $|b|^2 \neq 0$ then, by the previous corollary, α is conformally flat as well as Ricci-flat. It is thus a simple consequence of the Ricci-decomposition of the Riemann tensor that α must be flat. From Prop. 16 it then follows that β must have constant coefficients in flat coordinates.

Prop. 16 shows that any Ricci-flat *m*-Kropina metric $|b|^2 \neq 0$ must be of the form (38). We end this section with an example of a nowhere-null *m*-Kropina space from the literature and show that it is indeed the trivial Finsler metric (38) in disguise.

Example 20. In (Fuster et al., 2018)⁸ the following m-Kropina metric was considered as a cosmological model of spacetime in the context of Finsler gravity—an extension of Einstein's general theory of relativity that allows for Finsler spacetime geometries:

$$F = \sqrt{\left| dt^2 - a(t)^2 \left(dx^2 + dy^2 + dz^2 \right) \right|^{1+m}} \left(c \, a(t)^{1/m} dt \right)^{-m}, \tag{39}$$

The metric is written in terms of the coordinates (t, x, y, z), a(t) is an arbitrary function, and c is a positive constant. We remark that the notation in the original article was somewhat different.²⁰ This metric is Ricci-flat (and hence a solution to most candidate field equations in Finsler gravity).

Since the 1-form $b = c a(t)^{1/m} dt$ is nowhere null, Prop. 16 dictates that the metric must be of the trivial form (38) in suitable coordinates. It indeed turns out (and it is easy to check) that the coordinate transformation

$$\tilde{t} = \int^{t} \frac{c^{-m}}{a(t)} dt, \qquad \tilde{x} = c^{-m} x, \qquad \tilde{y} = c^{-m} y, \qquad \tilde{z} = c^{-m} z,$$
(40)

brings the metric into the trivial form

$$F = \sqrt{\left| d\tilde{t}^2 - d\tilde{x}^2 - d\tilde{y}^2 - d\tilde{z}^2 \right|^{1+m}} \left(d\tilde{t} \right)^m.$$
(41)

B. Ricci-flat null *m*-Kropina metrics in (1+3)D

While an *m*-Kropina space with $|b|^2 \neq 0$ is always locally metrizable, as we have seen in Prop. 9, this is not always the case when the 1-form is null. With regard to Ricciflatness, it turns out that in the $|b|^2 = 0$ case we have to add the additional assumption that *F* is locally metrizable in order to obtain nice results. Alternatively, one may choose to replace everywhere the two conditions *locally metrizable and Ricci-flat* by the single condition *affinely Ricci-flat*, the latter meaning that the *affine* Ricci tensor vanishes (rather than the Finsler-Ricci tensor, which is its symmetrization, by Lemma 3). Affine Ricci-flatness is not a very common notion in the literature, though, so we will stick to locally metrizable and Ricci-flat throughout the article.

Proposition 21. Let F be an m-Kropina metric of Berwald type. Then

F is affinely Ricci-flat \Leftrightarrow F is locally metrizable and Ricci-flat

Proof. If F is affinely Ricci-flat then it is trivially Ricci-flat, since the Finsler-Ricci tensor is the symmetrization of the affine Ricci tensor. In that case, F also locally metrizable, by Theorem 7, since the (vanishing) affine Ricci tensor is, in particular, symmetric. For the converse implication, if F is locally metrizable then, by Theorem 7, its affine Ricci tensor is symmetric, and hence the affine Ricci tensor coincides with the Finsler-Ricci tensor. The vanishing of the Finsler-Ricci tensor thus implies, in this case, the vanishing of the affine Ricci tensor.

To obtain the $|b|^2 = 0$ analog of Prop. 16, characterizing Ricci-flatness, we will employ the following result proven in (Heefer et al., 2023)¹³, an alternative account of which—more in the classical *definition-theorem-proof* style—can be found in (Heefer, 2024)⁷. **Lemma 22.** Let a Lorentzian metric a and 1-form b on a 4-dimensional manifold be locally given by

$$a = -2 du dv + H(u, x) du^{2} + 2W_{a}(u, x) dx^{a} du + h_{ab}(u, x) dx^{a} dx^{b},$$
(42)

$$b = du. (43)$$

If a is Ricci-flat then a and b can be expressed locally, in suitable coordinates, as

$$a = -2d\bar{u}d\bar{v} + \bar{H}(\bar{u},\bar{x},\bar{y})d\bar{u}^2 + d\bar{x}^2 + d\bar{y}^2,$$
(44)

$$b = d\bar{u},\tag{45}$$

such that \bar{H} satisfies $\partial_{\bar{x}}^2 \bar{H} + \partial_{\bar{y}}^2 \bar{H} = 0$.

Below δ_{ab} denotes the Kronecker symbol: $\delta_{ab} = 1$ iff a = b and otherwise $\delta_{ab} = 0$.

Proposition 23. Any locally metrizable m-Kropina space in (1 + 3)D with $|b|^2 = 0$ is Ricci-flat if and only if it is locally of the form

$$F = \left| -2 du dv + H(u, x) du^2 + \delta_{ab} dx^a dx^b \right|^{(1+m)/2} (du)^{-m}$$
(46)

such that H satisfies $\delta^{ab}\partial_a\partial_b H = 0$.

Proof. First of all, since du is parallel, it follows by Prop. 8 that any F defined by (46) is Ricci-flat, since the Ricci tensor of the Lorentzian metric (which vanishes, as can be checked explicitly) must coincide with the affine Ricci tensor of F, which therefore also vanishes, implying that also the Finsler-Ricci tensor vanishes, by Lemma 3.

Conversely, suppose that F is Ricci-flat. Since F is locally metrizable it can be written in the standard form (33) with a and b given by (42) and (43), respectively. Since the 1-form b = du is parallel with respect to the corresponding metric a, it follows that the affine connection on F coincides with the affine connection on a. So since F is Ricci-flat, a is Ricci-flat as well and hence it follows from Lemma 22 that F can be written locally in the form (46), as desired.

With this, we have classified (locally, at least) all locally metrizable, Ricci-flat *m*-Kropina metrics in (1 + 3)D with constant causal character.

VI. DISCUSSION

In this paper, we have studied the local and global metrizability as well as Ricci-flatness of *m*-Kropina spaces of Berwald type. First of all, we obtained necessary and sufficient conditions for the metrizability of Berwald *m*-Kropina metrics $F = \alpha^{1+m}\beta^{-m}$ of arbitrary signature. More precisely, Theorem 7 shows that the canonical affine connection of F can locally be obtained as the Levi-Civita connection of some pseudo-Riemannian metric if and only if the Ricci tensor constructed from the affine connection is symmetric. In addition, the theorem gives a third equivalent characterization of local metrizability in terms of the covariant derivative of the defining 1-form β with respect to the defining pseudo-Riemannian metric α . Using these characterizations, we have classified all locally metrizable *m*-Kropina metrics whose 1-forms have a constant causal character. These results are consistent with, and extend, the findings obtained in (Heefer et al., 2023)⁵, where only closed null 1-forms β were considered.

In general, global metrizability is a stronger notion than local metrizability. Nevertheless, in the special case where the first de Rahm cohomology group of the underlying manifold is trivial (which is true of simply connected manifolds, for instance), we have shown that global metrizability is equivalent to local metrizability. Hence, in this case, our necessary and sufficient conditions for local metrizability also characterize global metrizability.

Finally, we have classified all Ricci-flat, locally metrizable m-Kropina metrics in (3+1)D whose 1-forms have a constant causal character. As it turns out, an m-Kropina space is locally metrizable and Ricci-flat if and only if it is affinely Ricci flat—meaning that the Ricci tensor constructed from the canonical affine connection vanishes. We have explicitly constructed all affinely Ricci-flat m-Kropina spaces of Berwald type.

In part, the results obtained in this paper should be viewed as a small step in furthering our understanding of the pseudo-Riemann metrizability of Berwald spaces in general.^{4–6} On the other hand, our results have consequences for Finsler extensions of Einstein's general theory of relativity. More precisely, they constrain the possible types of vacuum solutions of Berwald *m*-Kropina type to Pfeifer and Wohlfarth's field equation in Finsler gravity^{14–16}, as these are typically Ricci-flat.²¹ In particular, our classification implies that any Ricci-flat vacuum solution of Berwald *m*-Kropina type with non-null 1-form must be trivial. Indeed, as an application, we considered an explicit *m*-Kropina metric from the literature used in cosmology that looks non-trivial at first sight, yet we showed that it is trivial after all. Thus, our results may also be seen as a step toward the complete classification of vacuum solutions of Berwald *m*-Kropina type to Pfeifer and Wohlfarth's field equation.

VII. ACKNOWLEDGEMENTS

S. Heefer wants to thank M. Sánchez for asking several inspiring questions regarding the difference between local and global metrizability.

REFERENCES

- ¹Z. Szabó, "Positive definite Berwald spaces," Tensor **35**, 25–39 (1981).
- ²C. Vincze, "A new proof of Szabó's theorem on the Riemann-metrizability of Berwald manifolds," Acta Math. Paedagog. Nyházi. **21**, 199–204 (2005).
- ³M. Crampin, "On the construction of Riemannian metrics for Berwald spaces by averaging," Houston journal of mathematics **40**, 737–750 (2014).
- ⁴A. Fuster, S. Heefer, C. Pfeifer, and N. Voicu, "On the non metrizability of Berwald Finsler spacetimes," Universe **6**, 64 (2020), arXiv:2003.02300.
- ⁵S. Heefer, C. Pfeifer, J. van Voorthuizen, and A. Fuster, "On the metrizability of m-Kropina spaces with closed null one-form," J. Math. Phys. **64**, 022502 (2023), arXiv:2210.02718.
- ⁶N. Voicu and S. G. Elgendi, "Metrizability of so(3)-invariant connections: Riemann versus Finsler," (2024), arXiv:2404.02980.
- ⁷S. Heefer, *Finsler Geometry, Spacetime, & Gravity*, Ph.D. thesis, Eindhoven University of Technology (2024), arXiv:2404.09858.
- ⁸A. Fuster, C. Pabst, and C. Pfeifer, "Berwald spacetimes and very special relativity," Phys. Rev. D **98**, 084062 (2018), arXiv:1804.09727.
- ⁹A. Fuster and C. Pabst, "Finsler *pp*-waves," Phys. Rev. D **94**, 104072 (2016), arXiv:1510.03058.
- ¹⁰A. P. Kouretsis, M. Stathakopoulos, and P. C. Stavrinos, "The general very special relativity in Finsler cosmology," Phys. Rev. D **79**, 104011 (2009), arXiv:0810.3267.

- ¹¹A. G. Cohen and S. L. Glashow, "Very special relativity," Phys. Rev. Lett. **97**, 021601 (2006), arXiv:hep-ph/0601236.
- ¹²G. Gibbons, J. Gomis, and C. Pope, "General very special relativity is Finsler geometry," Phys. Rev. D **76**, 081701 (2007), arXiv:0707.2174.
- ¹³S. Heefer and A. Fuster, "Finsler gravitational waves of (α, β) -type and their observational signature," Class. Quantum Gravity **40**, 184002 (2023), arXiv:2302.08334.
- ¹⁴C. Pfeifer and M. N. R. Wohlfarth, "Finsler geometric extension of Einstein gravity," Phys. Rev. D 85, 064009 (2012), arXiv:1112.5641.
- ¹⁵C. Pfeifer, *The Finsler spacetime framework: backgrounds for physics beyond metric geometry*, Ph.D. thesis, University of Hamburg (2013).
- ¹⁶M. Hohmann, C. Pfeifer, and N. Voicu, "Finsler gravity action from variational completion," Phys. Rev. D **100**, 064035 (2019), arXiv:1812.11161.
- 17 For alternative equivalent definitions, see (Javaloyes et al., 2022)²² and references therein.
- ¹⁸V. Kropina, "On projective two-dimensional Finsler spaces with special metric," Trudy Sem. Vektor. Tenzor. Anal. **11**, 277–292 (2006), arXiv:math/0605759.
- ¹⁹M. Matsumoto, Finsler Geometry in the 20th-Century (Published in Handbook of Finsler Geometry, edited by P. L. Antonelli) (Kluwer Academic Publishers, Dordrecht, 2003).
- ²⁰A slightly different framework is used in (Fuster et al., 2018)⁸, where the geometry is in fact characterized by an *r*-homogeneous Lagrangian (for arbitrary *r*) rather than a Finsler metric. But the metric (39) displayed here represents the same geometry. It is related to the Lagrangian in (Fuster et al., 2018)⁸ by $L = F^{\frac{2}{1+m}}$. Similarly, our parameter *m* is related to the parameter *m* in (Fuster et al., 2018)⁸ by $n = \frac{-2m}{1+m}$.
- ²¹In fact, we expect that any vacuum solution of Berwald *m*-Kropina metric is Ricci-flat, although this has only been proven in the case with non-null 1-form⁸.
- ²²M. Á. Javaloyes, M. Sánchez, and F. F. Villaseñor, "The Einstein-Hilbert-Palatini formalism in pseudo-Finsler geometry," Adv. Theor. Math. Phys. 26 (2022), arXiv:2108.03197.