ON RELATIVE COHOMOLOGY IN LIE THEORY

MARÍA AMELIA SALAZAR

ABSTRACT. Motivated by our attempt to understand characteristic classes of Lie groupoids and geometric structures, we are brought back to the fundamentals of the cohomology theories of Lie groupoids and algebroids. One element that was missing in the literature was the notion of *relative cohomology* in this setting. The main aim of this paper is to develop the structural theory of this notion, the relation between the relative cohomology of groupoids and that of algebroids via van Est maps, and to indicate how it can be used to provide an intrinsic definition of characteristic classes.

Contents

1.	Introduction	1
2.	Relative cohomology of Lie groupoids	4
3.	Relative cohomology of Lie algebroids	8
4.	The transitive case	12
5.	Van Est maps	14
6.	Characteristic classes	27
Ap	34	
Re	36	

1. INTRODUCTION

Lie group and Lie algebra cohomology have been used successfully to study the interaction between the algebraic and geometric properties of these objects. Lie algebra cohomology was first introduced by Élie Cartan in [4], and later extended by Chevalley and Eilenberg in [5]. It was originally used to study the topology of Lie groups and homogeneous spaces by connecting de Rham cohomology with the properties of the Lie algebra. On the other hand, Lie group cohomology is closely related to the cohomology of the classifying space (see the work of Stasheff [21]). Its relation to Lie algebra cohomology has been extensively studied and produced many classical results. In particular, van Est [24, 25, 26] in one of his results produces a map VE_G from the cochain complex that computes the differentiable cohomology of a Lie group G to the Chevalley-Eilenberg complex of its Lie algebra; under some connectivity assumptions on G, this induces an isomorphism in cohomology.

Lie groupoids and Lie algebroids are a far reaching generalization of Lie groups and Lie algebras respectively, which have been used to codify geometric structures and their symmetries (foliations, Poisson manifolds and related geometries, PDEs, etc). Their cohomologies are natural extensions of those for Lie groups and Lie algebras, and have been very relevant when applied to geometric structures such as de Rham cohomology, foliated cohomology, Poisson cohomology, etc. (see, among others, [6, 8, 9, 18]). The relation between the cohomology of Lie groupoids and Lie algebroids is a flourishing theory that started with the work of Weinstein and Xu in [27], where they first extended the van Est map to a map $VE_G : C(G) \to C(A)$ from the cochain complex that computes the differentiable cohomology of a Lie groupoid Gto the Chevalley-Eilenberg complex of its Lie algebroid A. Later Crainic [6] produced other versions of van Est results; in *op. cit.* he also defined characteristic classes of representations of Lie algebroids, extending the classical characteristic classes of flat vector bundles [1, 15], and the modular class of Poisson manifolds [11]. (See also [7, 12] for characteristic classes of Lie algebroids.)

Motivated by understanding characteristic classes of representations of Lie algebroids intrinsically and by our attempt to develop a parallel theory of characteristic classes of representations of Lie groupoids, we are brought back to understanding the fundamentals of the cohomology theories of Lie groupoids and algebroids. One element that was notably missing in the literature was the notion of *relative cohomology* for Lie groupoids and Lie algebroids. The main aim of this paper is to introduce and study this notion, as well as to indicate how it can be used to provide an intrinsic definition of characteristic classes for both Lie groupoids and Lie algebroids. (These will be studied in more detail in future work.) Throughout we pay special attention to "van Est type" results, which illustrate the relation between the notions at the global (Lie groupoid) and infinitesimal (Lie algebroid) level. Our point of view is to find intrinsic and simple notions, as well as explicit constructions. In what follows we illustrate the structure of the paper and the main results.

Relative cohomology of Lie groupoids (Section 2). We begin by revisiting the cochain complex C(G) of a Lie groupoid G computing its differentiable cohomology. Associated to any (wide) Lie subgroupoid K, we define a cochain subcomplex $C(G)^K$ (and its normalized version) which at degree p consists of cochains invariant under p + 1 natural K-actions. For K proper, one can take the average of the actions and produce a cochain map

(1)
$$\operatorname{Av}: \mathsf{C}(G) \to \mathsf{C}(G)^K$$

which turns out to be a homotopy equivalence with the inclusion $\iota: C(G)^K \to C(G)$ its homotopy inverse (see Theorem 2.5). Our definitions and results are inspired by the work of Guichardet [14] in the context of Lie groups, where he obtains a similar (but more involved) map to (1) which he proves is a homotopy equivalence when K is a finite subgroup of G. Hence, to the best of our knowledge, our approach provides a new insight already for Lie groups.

Relative cohomology of Lie algebroids (Section 3). In order to define relative cohomology in the context of Lie algebras and Lie groups, one deals with the \mathfrak{k} -basic subcomplex $C(\mathfrak{g})_{\mathfrak{k}-\text{basic}}$ of the Chevalley-Eilenberg complex $C(\mathfrak{g})$ of a Lie algebra \mathfrak{g} and a Lie subalgebra \mathfrak{k} , consisting of cochains that are \mathfrak{k} -horizontal and \mathfrak{k} -invariant under the restriction to \mathfrak{k} of coadjoint action on \mathfrak{g}^* . Similarly, when \mathfrak{g} and \mathfrak{k} are the Lie algebras of a Lie group G and a Lie subgroup K respectively, one considers the K-basic subcomplex $C(\mathfrak{g})_{K-\text{basic}}$; in this case, the relevant action is the restriction to K of the coadjoint action on \mathfrak{g}^* . Generalizing these notions to Lie algebroids and Lie groupoids is not obvious as, in general, a Lie algebroid A does not act on itself, nor does a Lie groupoid G of A act on A. To solve this issue, we use the key observation that when $B \subseteq A$ is a Lie subalgebroid, there is a natural action of B on the quotient A/B; similarly, if $K \subseteq G$ is a Lie subgroupoid integrating B, then K acts naturally on A/B. This allows us to define and give the main properties of the B-basic subcomplex and the K-basic subcomplex respectively

$$C(A)_{B-\text{basic}} \subseteq C(A), \quad C(A)_{K-\text{basic}} \subseteq C(A),$$

which coincide with the standard notions in the context of Lie algebras and Lie groups.

The transitive case (Section 4). Here we assume the existence of a wide transitive Lie subgroupoid $K \subseteq G$. We show that there is a natural extension map

$$\operatorname{ext}: \mathsf{C}^{\bullet}(G_z)^{K_z} \to \mathsf{C}^{\bullet}(G)^K,$$

where K_z , G_z are the isotropy Lie groups of K and G respectively, over a point z of the base. This map is an isomorphism of chain complexes with inverse given by the restriction (see Theorem 4.1). An analogous result holds true (see Theorem 4.3) when looking at the infinitesimal counterpart:

ext:
$$\mathsf{C}^{\bullet}(\mathfrak{g}_z)_{K_z-\text{basic}} \to \mathsf{C}^{\bullet}(A)_{K-\text{basic}};$$

here A is the Lie algebroid of G and \mathfrak{g}_z is the isotropy Lie algebra. Our theory shows that, in this case, the relative cohomology of Lie groupoids and Lie algebroids is the same as the relative cohomology of their isotropy Lie groups and Lie algebras.

Van Est maps (Section 5). In this section we study the van Est maps in the context of relative cohomology, and pay special attention to the case where the Lie subgroupoid K is proper. We make use of tubular structures and Cartan decompositions of (G, K) (see Definitions 5.1 and 5.2) to obtain van Est integration maps (i.e., cochain maps going from the relative cohomology of the Lie algebroid to the relative cohomology of the Lie groupoid) and as a consequence, conclude that in this case the van Est differentiation map (see equation (33))

$$VE_{G/K} \colon \mathsf{C}^{\bullet}(G) \to \mathsf{C}^{\bullet}(A)_{K-\text{basic}}$$

is a homotopy equivalence when K is a proper subgroupoid (see Theorem 5.12). For example, for a Lie group G and $K \subseteq G$ a maximal compact subgroup, tubular structures always exist (see [22]). To prove the results we use the Perturbation Lemma, treated in the Appendix. We apply the previous results to conclude that in the transitive case, the differentiable cohomology of the Lie groupoid is homotopy equivalent to the cohomology of its isotropy Lie algebra relative to the maximal compact subgroup of its isotropy Lie group (see Theorem 5.17).

Characteristic classes (Section 6). As an application of our theory we introduce the definition of characteristic classes of a representation of a Lie groupoid. This is the first step to study characteristic classes of Lie groupoids endowed with a geometric structure, e.g., symplectic, symplectic-Nijenhuis, presymplectic, Pfaffian and contact groupoids. Our theory also allows us to give an alternative, simple and direct definition of the the existing notion of characteristic classes of a representation of a Lie algebroid (see [6, 7, 12]), and relate them with our definition of characteristic classes for Lie groupoids.

All the groupoids appearing in this paper are assumed to be Hausdorff.

Acknowledgments. The author would like to thank Marius Crainic, Ioan Mărcut and Eckhard Meinrenken for useful comments and suggestions. The author was partly supported by CNPq grant PQ1 304410/2020-9, FAPERJ grants JCNE 262012932021 and ARC 26211412201,

MARÍA AMELIA SALAZAR

and by the Serrapilheira Institute grant Serra-1912-3205. The author is grateful to Max Planck Institute for Mathematics in Bonn for its hospitality and financial support. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brazil (CAPES) – Finance code 001.

2. Relative cohomology of Lie groupoids

We will review the cochain complexes for Lie groupoids, and define the relative cohomology with respect to a Lie subgroupoid. We will give the main properties of these notions and exhibit equivalences between cochain complexes. In particular we show that the cohomology of a Lie groupoid relative to a proper subgroupoid computes its differentiable cohomology (see Theorem 2.5).

2.1. The simplicial manifold B_pG . Let $G \Rightarrow M$ be a Lie groupoid, with source and target maps denoted by $s, t: G \to M$ and unit map $u: M \to G$ (units are denoted by elements of M). The inverse of g is denoted by i(g) or as g^{-1} . Elements $g, h \in G$ are *composable* if s(g) = t(h); in this case their product is denoted as gh or as m(g, h). We denote by

$$B_p G = \{ (g_1, \dots, g_p) | \mathsf{s}(g_i) = \mathsf{t}(g_{i+1}), \ 0 < i < p \}$$

the space of *p*-arrows; by convention $B_0G = M$. Every *p*-arrow comes with p+1 base points (m_0, \ldots, m_p) , where $m_i = \mathsf{s}(g_i) = \mathsf{t}(g_{i+1})$. The collection of spaces B_pG defines a simplicial manifold $B_{\bullet}G$ called the *nerve* of the groupoid. The face map $\partial_i \colon B_pG \to B_{p-1}G$ drops the *i*-th base point:

(2)
$$\partial_i(g_1, \dots, g_p) = \begin{cases} (g_2, \dots, g_p), & i = 0, \\ (g_1, \dots, g_i g_{i+1}, \dots, g_p), & 0 < i < p, \\ (g_1, \dots, g_{p-1}), & i = p, \end{cases}$$

while the *i*-th degeneracy map $\epsilon_i \colon B_p G \to B_{p+1} G$ is defined by

(3)
$$\epsilon_i(g_1, \dots, g_p) = (g_1, \dots, g_i, m_i, g_{i+1}, \dots, g_p), \quad i = 0, \dots, p.$$

The manifolds B_pG also come equipped with p+1 commuting right G-actions

(4)
$$(g_1, \dots, g_p) \cdot a = \begin{cases} (a^{-1}g_1, g_2, \dots, g_p) & i = 0, \\ (g_1, \dots, g_i a, a^{-1}g_{i+1}, \dots, g_p) & 0 < i < p, \\ (g_1, \dots, g_{p-1}, g_p a) & i = p; \end{cases}$$

the *i*-th action has anchor map given by $(g_1, \ldots, g_p) \mapsto m_i$.

Define a simplicial fiber bundle

(5)
$$\kappa_p \colon E_p G = B_p G \times_M G \to B_p G.$$

The simplicial structure on $E_{\bullet}G$ comes from the identification $B_pG \times_M G \simeq B_p(G \ltimes G)$, where $G \ltimes G \rightrightarrows G$ is the action groupoid, with the *G*-action by left multiplication. For each p, κ_p is a principal *G*-bundle, with anchor map

$$\pi_p(g_1,\ldots,g_p;g)=\mathsf{s}(g),$$

and principal action

(6)
$$a \cdot (g_1, \dots, g_p; g) = (g_1, \dots, g_p; ga^{-1}).$$

Note that the face and degeneracy maps (2) and (3) for E_pG are *G*-equivariant with respect to this action. They are also principal bundle morphisms, making $\kappa \colon EG \to BG$ into a simplicial principal bundle. The actions (4) lift to p+1 commuting right *G*-actions on $E_pG = B_pG \times_M G$

(7)
$$(g_1, \dots, g_p; g) \cdot a = \begin{cases} (a^{-1}g_1, g_2, \dots, g_p; g) & i = 0, \\ (g_1, \dots, g_i a, a^{-1}g_{i+1}, \dots, g_p; g) & 0 < i < p, \\ (g_1, \dots, g_{p-1}, g_p a; a^{-1}g) & i = p; \end{cases}$$

the *i*-th action has anchor map $(g_1, \ldots, g_p; g) \mapsto m_i$.

2.2. The groupoid complex. The groupoid cochain complex $(C(G), \delta)$ has graded components

$$\mathsf{C}^p(G) = C^\infty(B_p G),$$

while the differential δ is given on *p*-cochains by $\delta = \sum_{i=0}^{p+1} (-1)^i \partial_i^*$. Its cohomology is called the *differentiable cohomology of G*.

For the simplicial manifold $E_{\bullet}G$ as in (5), we denote its cochain complex by $(\mathsf{C}(EG), \delta)$; note that the inclusion j: $\mathsf{C}(G) \to \mathsf{C}(EG)$ given in degree p by the pullback κ_p^* identifies $\mathsf{C}(G)$ with the G-invariant subcomplex of $\mathsf{C}(EG)$ with respect to the action dual to (6).

Consider now a wide Lie subgroupoid $K \subseteq G$ (i.e. the inclusion $K \hookrightarrow G$ is an injective immersion of Lie groupoids and K has the same base M). For each of the actions (4) we can consider the restriction to K-actions.

Definition 2.1. The Lie groupoid complex relative to K is the K-invariant subcomplex

 $\mathsf{C}(G)^K$.

That is, it consists of elements of $C^p(G)$ which are invariant w.r.t. p+1 commuting K-actions. We denote the resulting cohomology groups of $(C(G)^K, \delta)$ by $H^{\bullet}(G, K)$.

Remark 2.2. Sometimes it is useful to work with the *normalized K-invariant subcomplex*

 $\widetilde{\mathsf{C}}(G)^K$

consisting of functions $f \in C(G)^K$ with the property that $f(g_1, \ldots, g_p) = 0$ whenever $g_i \in K$ for some *i*.

Similarly, one defines the K-invariant subcomplex $C(EG)^K$ and its normalized version $\widetilde{C}(EG)^K$

$$\widetilde{\mathsf{C}}(EG)^K \subseteq \mathsf{C}(EG)^K \subseteq \mathsf{C}(EG)$$

with respect to the K-actions (7). A function $f \in \mathsf{C}(EG)^K$ is normalized if $f(g_1, \ldots, g_p; g) = 0$ whenever $g_i \in K$ for some $i = 1, \ldots p$. The map $j: \mathsf{C}(G)^K \to \mathsf{C}(EG)^K$ includes $\mathsf{C}(G)^K$ as the G-invariant subcomplex of $\mathsf{C}(EG)^K$ with respect to the action (6), and similarly for $j: \widetilde{\mathsf{C}}(G)^K \to \widetilde{\mathsf{C}}(EG)^K$.

MARÍA AMELIA SALAZAR

Remark 2.3. The usual cup product

$$(f_1 \cup f_2)(g_1, \dots, g_{p+q}) = f_1(g_1, \dots, g_p)f_2(g_{p+1}, \dots, g_{p+q})$$

defines a product structure $C^{p}(G) \times C^{q}(G) \to C^{p+q}(G)$, which passes to cohomology and turns $(C^{\bullet}(G)^{K}, \delta) \subseteq (C^{\bullet}(G), \delta)$ into a DG subalgebra and $H^{\bullet}(G, K)$ (and $H^{\bullet}(G)$) into a graded algebra.

Similarly, the normalized K-invariant subcomplex is a DG subalgebra and its cohomology is a graded algebra.

2.3. Equivalences between complexes. Let K be a wide Lie subgroupoid of G. The main result of this section is that the normalized Lie groupoid complex relative to K computes the same cohomology as the differentiable cohomology, whenever K is proper (see Theorem 2.5). In fact, in general the following holds:

Theorem 2.4. The inclusion $\iota \colon \widetilde{\mathsf{C}}(G)^K \to \mathsf{C}(G)^K$ is a homotopy equivalence.

In preparation for the proof of Theorem 2.4, recall that $(C(EG), \delta)$ is acyclic. To see this, consider the maps

 $h_p: E_pG \to E_{p+1}G, \ (g_1, \ldots, g_p; g) \mapsto (g_1, \ldots, g_p, g; m)$

where $m = \mathbf{s}(g)$. The map

(8)
$$h: \mathbf{C}^p(EG) \to \mathbf{C}^{p-1}(EG), \quad f \mapsto (-1)^p h_{p-1}^* f$$

satisfies $[\mathbf{h}, \delta] = 1 - \mathbf{i} \circ \mathbf{p}$, where $\mathbf{i} = \pi_0^* \colon C^\infty(M) \to \mathsf{C}^0(EG)$ and $\mathbf{p} \colon \mathsf{C}^0(EG) \to C^\infty(M)$ is the left inverse to \mathbf{i} given by pullback under the inclusion $\mathbf{u} \colon M \hookrightarrow E_0 G = G$. Similarly, $(\mathsf{C}(EG)^K, \delta)$ is acyclic with $\mathbf{h}, \mathbf{p}, \mathbf{i}$ restricting to maps for this complex.

Proof of Theorem 2.4. Denote by $\widetilde{\mathsf{C}}(G) \subseteq \mathsf{C}(G)$ the subcomplex consisting of cochains with the property that $f(g_1 \ldots, g_p) = 0$ whenever g_i is a unit for some *i*. It is well know that the inclusion $\iota \colon \widetilde{\mathsf{C}}(G) \to \mathsf{C}(G)$ is a homotopy equivalence, and one can exhibit a homotopy inverse $N \colon \mathsf{C}(G) \to \widetilde{\mathsf{C}}(G)$ (see e.g. [14]). Observe that $\widetilde{\mathsf{C}}(G)^K = \widetilde{\mathsf{C}}(G) \cap \mathsf{C}(G)^K$. We will then show that N restricts to a homotopy inverse $N^K \colon \mathsf{C}(G)^K \to \widetilde{\mathsf{C}}(G)^K$ of $\iota \colon \widetilde{\mathsf{C}}(G)^K \to \mathsf{C}(G)^K$.

For the definition of N (and that of the homotopy **n** below), we have an alternative description E_pG via the change of coordinates $(g_1, \ldots, g_p; g) \mapsto (g_1g_2 \cdots g_pg, g_2 \cdots g_pg, \ldots, g_pg, g)$ as

$$E_pG = \{(g_0, \dots, g_p) \in G^{p+1} \mid \mathsf{s}(g_0) = \dots = \mathsf{s}(g_p)\}.$$

The principal G action (6) becomes $a \cdot (g_0, \ldots, g_p) = (g_0 a^{-1}, \ldots, g_p a^{-1})$. The face maps $\partial_i \colon E_p G \to E_{p-1} G$ drop the *i*-th entry, while the degeneracy maps $\epsilon_i \colon E_p G \to E_{p+1} G$ repeat the *i*-th entry. In this form, the normalized subcomplex $\widetilde{C}(EG) \subseteq C(EG)$ consists of those functions on $E_p G$ with the property that $f(g_1 \ldots, g_p) = 0$ whenever $g_i = g_{i+1}$ for some $i = 0, \ldots, p-1$, while the K-invariant subcomplex $C(EG)^K$ consists of those functions with the property that $f(k_0 g_0, \ldots, k_p g_p) = f(g_0, \ldots, g_p)$ whenever $k_0, \ldots, k_p \in K$, and its normalized version $\widetilde{C}(EG)^K$ of function $f \in C^p(EG)^K$ such that $f(g_0, \ldots, g_p) = 0$ whenever $g_i g_{i+1}^{-1} \in K$ for some $i = 0, \ldots, p-1$. Setting

$$(\delta_{g_0}\otimes\cdots\otimes\delta_{g_p})(f):=f(g_0,\ldots,g_p),$$

as in [14] we define $N: \mathsf{C}^p(EG) \to \widetilde{\mathsf{C}}^p(EG)$ by

 $N(f)(g_0,\ldots,g_p) = \delta_{g_0} \otimes (\delta_{g_1} - \delta_{g_0}) \otimes \cdots \otimes (\delta_{g_n} - \delta_{g_{p-1}})(f)$

(it follows by a direct computation that $N(f) \subseteq \widetilde{\mathsf{C}}^p(G)$). Note that N is G-equivariant and it restricts to a map $N^K \colon \mathsf{C}(EG)^K \to \widetilde{\mathsf{C}}(EG)^K = \widetilde{\mathsf{C}}(EG) \cap \mathsf{C}(EG)^K$. In degree p = 0, $N = \mathrm{id}_{\mathsf{C}^0(EG)}$ so that $\mathsf{i} = \mathsf{i} \circ N$; a computation shows that N is a cochain map. It is also a computation to see that

$$N \circ \iota = \mathrm{id}_{\widetilde{\mathsf{C}}(EG)}$$

for the inclusion $\iota: \widetilde{\mathsf{C}}^p(EG) \to \mathsf{C}^p(EG)$. For $\mathsf{v} = \mathrm{id}_{\mathsf{C}(EG)} - \iota \circ N$, we will show (see computation below) that $\mathsf{v} = [\mathsf{n}, \delta]$ for some *G*-equivariant maps $\mathsf{n}: \mathsf{C}^p(EG) \to \mathsf{C}^{p-1}(EG)$ that restrict to the *K*-invariant subcomplex $\mathsf{C}(EG)^K$, thus concluding that $\iota: \widetilde{\mathsf{C}}(EG)^K \to \mathsf{C}(EG)^K$ is a homotopy equivalence. As all the maps involved are *G*-equivariant, it follows that ι restricts to the homotopy equivalence $\iota: \widetilde{\mathsf{C}}(G)^K \to \mathsf{C}(G)^K$ of the *G*-invariant subcomplexes.

First, we define $\mathbf{n}_1: \mathbf{C}^1(EG) \to \mathbf{C}^0(EG)$ as $\mathbf{n}_1 = 0$. As $\mathbf{v}_0 = 0$ then $\mathbf{v}_0 = \mathbf{n}_1 \circ \delta$. Let us argue recursively and suppose that we have constructed *G*-equivariant maps $\mathbf{n}_i: \mathbf{C}^i(EG) \to \mathbf{C}^{i-1}(EG)$ for $i = 1, \ldots, p-1$ with the property that

$$\mathsf{v}_{i-1} = \delta \circ \mathsf{n}_{i-1} + \mathsf{n}_i \circ \delta.$$

Let $\bar{\mathbf{v}}_{p-1} = \mathbf{v}_{p-1} - \delta \circ \mathbf{n}_{p-1}$, then $\bar{\mathbf{v}}_{n-1} \circ \delta = \mathbf{v}_{n-1}$ of

$$\mathbf{v}_{p-1} \circ \delta = \mathbf{v}_{p-1} \circ \delta - \delta \circ \mathbf{n}_{p-1} \circ \delta = \delta \circ \mathbf{v}_{p-2} - \delta \circ \mathbf{n}_{p-1} \circ \delta = 0,$$

hence $\bar{v}_{p-1}|_{\mathrm{Im}(\delta: C^{p-2}(EG) \to C^{p-1}(EG))} = 0$. As

$$\operatorname{Im}(\delta \colon \mathsf{C}^{p-2}(EG) \to \mathsf{C}^{p-1}(EG)) = \ker(\delta \colon \mathsf{C}^{p-1}(EG) \to \mathsf{C}^p(EG)),$$

 $\bar{\mathbf{v}}_{p-1}$ passes to the quotient $\mathbf{C}^{p-1}(EG)/\ker \delta$. Note that

$$\delta \colon \mathsf{C}^{p-1}(EG) / \ker \delta \to \mathsf{C}^p(EG)$$

is injective with left inverse $h: C^p(EG) \to C^{p-1}(EG)/\ker \delta$ (the homotopy operator (8) after passing to the quotient). Define $n_p: C^p(EG) \to C^{p-1}(EG)$ by

(9)
$$\mathsf{n}_p(f)(g_0,\ldots,g_{p-1}) = \bar{\mathsf{v}}_{p-1}(\mathsf{h}(g_0\cdot f))(\mathsf{t}(g_0),g_1g_0^{-1},\ldots,g_{p-1}g_0^{-1}).$$

The map n_p is G-equivariant and $n_p \circ \delta = \bar{v}_{p-1}$, thus $n_p \circ \delta = v_{p-1} - \delta \circ n_{p-1}$ as desired. \Box

Assume that $K \subseteq G$ is a proper Lie subgroupoid, in the sense that $(\mathbf{t}, \mathbf{s}) : K \to M \times M$ is a proper map. We will use properness of the Lie subgroupoid K of G to produce averaging operators, turning any cocycle on $\mathsf{C}(G)$ into a K-invariant cocycle on $\mathsf{C}^K(G)$. For this purpose recall [6, 10, 23] that a proper Lie groupoid admits *(left invariant) properly supported normalized* Haar system $\mu = {\mu_m}_{m \in M}$, i.e. a family of smooth measures on $\mathsf{t}^{-1}(m)$ such that:

• for any compactly supported function $f \in C^{\infty}(G)$, the formula

$$m \mapsto \int_{a \in \mathsf{t}^{-1}(m)} f(a) \ \mu_m(a)$$

defines a smooth function on M,

- μ is left invariant, i.e. $(L_q)_*\mu_m = \mu_{t(q)}$ for all $g \in s^{-1}(m)$,
- t restricts to a proper map $\operatorname{supp}(\mu) \to M$, and

• $\int_{\mathsf{t}^{-1}(m)} \mu_m = 1$ for all $m \in M$.

Choose such a Haar system μ on K and for each p denote by

(10)
$$\operatorname{Av}^{(i)} \colon C^{\infty}(B^{p}G) \to C^{\infty}(B^{p}G)$$

the induced averaging operator with respect to the *i*-th K-action (4). The operators $Av^{(i)}$ commute since the actions commute, and we denote by

(11)
$$Av = Av^{(0)} \circ \dots \circ Av^{(p)}$$

the total K^{p+1} -averaging operator.

Theorem 2.5. Suppose $K \subseteq G$ is a proper Lie subgroupoid. The averaging operator

Av: $\mathsf{C}^{\bullet}(G) \to \mathsf{C}^{\bullet}(G)^K$

defined by (11) is a homotopy equivalence with homotopy inverse given by the inclusion

 $\iota\colon \mathsf{C}^{\bullet}(G)^K\to\mathsf{C}^{\bullet}(G).$

Hence, the inclusion $\iota \colon \widetilde{\mathsf{C}}^{\bullet}(G)^K \to \mathsf{C}^{\bullet}(G)$ is a homotopy equivalence.

Proof. As in (10), denote by

$$\operatorname{Av}_{EG} \colon \mathsf{C}(EG) \to \mathsf{C}(EG)$$

the total averaging operator with respect to the K-actions (7). This averaging operator is a G-equivariant (w.r.t (6)) cochain map, restricting to (10) on the G-invariant subcomplex C(G). Note that Av_{EG} is a left inverse of the inclusion $\iota: C(EG)^K \to C(EG)$. We will see that $\iota \circ \operatorname{Av}_{EG}$ is homotopic to the identity with a G-equivariant homotopy $n: C^{\bullet}(EG) \to C^{\bullet-1}(EG)$, i.e. $v := \operatorname{id}_{C(EG)} - \iota \circ \operatorname{Av}_{EG} = [n, \delta]$, thus concluding that on the G-invariant complexes $\operatorname{Av}: C(G) \to C(G)^K$ is a homotopy equivalence with $\iota: C(G)^K \to C(G)$ a homotopy inverse.

The construction of \mathbf{n} is completely analogous to that of the proof of Theorem 2.4, with Av_{EG} playing the role of N. For the base case one defines $\mathbf{n}_1 \colon \mathsf{C}^1(EG) \to \mathsf{C}^0(EG)$ by

$$\mathsf{n}_1(f)(g) = \int_{a \in \mathsf{t}^{-1}(t(g)) \cap K} f(a^{-1};g) \ \mu_{\mathsf{t}(g)}(a).$$

For the inductive case one defines n_p as in (9).

To conclude that $\iota: \widetilde{\mathsf{C}}^{\bullet}(G)^K \to \mathsf{C}^{\bullet}(G)$ is a homotopy equivalence we use Theorem 2.4. \Box

3. Relative cohomology of Lie Algebroids

In this section we introduce the Chevalley-Eilenberg complex of a Lie algebroid relative to a subalgebroid. In the case where the algebroid comes from a Lie groupoid G, we also define the Chevalley-Eilenberg complex of a Lie algebroid relative to a subgroupoid of G. These notions extend the standard definition of the \mathfrak{k} -basic and K-invariant subcomplex, respectively, of the Chevalley-Eilenberg complex of a Lie algebra \mathfrak{g} , with $\mathfrak{k} \subseteq \mathfrak{g}$ a Lie algebra, and K a Lie subgroup of a Lie group G integrating \mathfrak{g} .

3.1. The Chevalley-Eilenberg complex and its geometry. Let $A \Rightarrow M$ be a Lie algebroid with anchor map $a: A \to TM$. Let $(C(A), d_{CE})$ be the Lie algebroid complex (or Chevalley-Eilenberg complex), with p-cochains

$$\mathsf{C}^p(A) = \Gamma(\wedge^p A^*),$$

and with the Chevalley-Eilenberg differential $\mathsf{d}_{CE} \colon \mathsf{C}^p(A) \to \mathsf{C}^{p+1}(A)$ defined by

$$(\mathsf{d}_{CE}\alpha)(\xi_1,\dots,\xi_{p+1}) = \sum_i (-1)^{i+1} \mathsf{a}(\xi_i) (\alpha(\xi_1,\dots,\hat{\xi}_i,\dots,\xi_{p+1})) + \sum_{i< j} (-1)^{i+j} \alpha([\xi_i,\xi_j],\xi_1,\dots,\hat{\xi}_i,\dots,\hat{\xi}_j,\dots,\xi_{p+1});$$

here $\alpha \in \Gamma(\wedge^p A^*)$ and $\xi_1, \ldots, \xi_{p+1} \in \Gamma(A)$. For $\xi \in \Gamma(A)$, we denote by

$$\iota_{\xi} \colon \mathsf{C}^{\bullet}(A) \to \mathsf{C}^{\bullet-1}(A), \ \mathcal{L}_{\xi} \colon \mathsf{C}^{\bullet}(A) \to \mathsf{C}^{\bullet}(A)$$

the operators given by contraction and the Lie derivative $\mathcal{L}_{\xi} = \mathsf{d}_{CE}\iota_{\xi} + \iota_{\xi}\mathsf{d}_{CE}$ respectively. On $\mathsf{C}^{0}(A) = C^{\infty}(M)$ we have that $\mathcal{L}_{\xi} = \mathcal{L}_{\mathsf{a}(\xi)}$.

Assume now that A is the Lie algebroid of $G \Rightarrow M$. Thus, A is the vector bundle whose sections are the left-invariant vector fields of G (tangent to the t-fibers); for $\xi \in \Gamma(A)$ we denote by ξ^L the corresponding left-invariant vector field. The anchor map $\mathbf{a}: A \to TM$ is characterized by the property that $\xi^L \sim_{\mathbf{s}} -\mathbf{a}(\xi)$. Denote by $(\Omega_{\mathcal{F}}(G), \mathbf{d}_{Rh})$ be the de Rham complex of t-foliated forms of G, and denote by $(\Omega_{\mathcal{F}}(G)^L, \mathbf{d}_{Rh})$ the subcomplex of left-invariant forms. For $\alpha \in C(A)$ we denote by α^L the left-invariant foliated form on G. This operation intertwines the Chevalley-Eilenberg and the de Rham differentials, and the contraction ι_{ξ} and Lie derivative \mathcal{L}_{ξ} operators of C(A) with the usual contraction of vector fields ι_{ξ^L} and Lie derivative of vector fields \mathcal{L}_{ξ^L} , respectively. Thus

(12)
$$(\mathsf{C}(A), \mathsf{d}_{CE}) \longleftrightarrow (\Omega_{\mathcal{F}}(G)^{L}, \mathsf{d}_{Rh})$$
$$\iota_{\xi} \colon \mathsf{C}(A) \to \mathsf{C}(A) \longleftrightarrow \iota_{\xi^{L}} \colon \Omega_{\mathcal{F}}(G)^{L} \to \Omega_{\mathcal{F}}(G)^{L}$$
$$\mathcal{L}_{\xi} \colon \mathsf{C}(A) \to \mathsf{C}(A) \longleftrightarrow \mathcal{L}_{\xi^{L}} \colon \Omega_{\mathcal{F}}(G)^{L} \to \Omega_{\mathcal{F}}(G)^{L}.$$

Actually the Lie derivatives arise as the infinitesimal generators of an action of Bis(G) – the group of bisections of G:

$$Bis(G) = \{b: M \to G \mid t \circ b = id, \ s \circ b \text{ a diffeomorphism} \}$$

with multiplication and inverse given by

$$(b \cdot a)(m) = b(m)a(\mathbf{s} \circ b(m)), \quad b^{-1}(m) = i(b((\mathbf{s} \circ b)^{-1}(m))), \text{ for } m \in M.$$

It acts on C(A) by

(13)
$$b \cdot \alpha := C_b^*(\alpha), \quad b \in \operatorname{Bis}(G), \ \alpha \in \mathsf{C}(A),$$

where C_b is the Lie groupoid diffeomorphism given by conjugation by b

(14)
$$C_b \colon G \to G, \ g \mapsto b(\mathsf{t}(g))^{-1}gb(\mathsf{s}(g)).$$

MARÍA AMELIA SALAZAR

Note that C_b^* on (13) is just the pullback of the Lie algebroid isomorphism $\text{Lie}(C_b) = TC_b|_A$ induced by C_b , and therefore

(15)
$$b \cdot (\mathsf{d}_{CE}\alpha) = \mathsf{d}_{CE}(b \cdot \alpha).$$

By differentiating the action (13) one obtains an operator $\nabla \colon \Gamma(A) \times \mathsf{C}^{\bullet}(A) \to \mathsf{C}^{\bullet}(A)$

(16)
$$\nabla_{\xi}(\alpha) = \frac{d}{d\epsilon} \mid_{\epsilon=0} \varphi_{\xi}^{\epsilon} \cdot \alpha ;$$

here φ_{ξ}^{ϵ} is the family of bisection defined by the flow of ξ^{L} as $\varphi_{\xi}^{\epsilon}(m) = \varphi_{\xi^{L}}^{\epsilon}(m), m \in M$. Note that by left-invariance of ξ^{L} ,

(17)
$$\varphi_{\xi^L}^{\epsilon}(g) = g\varphi_{\xi}^{\epsilon}(\mathbf{s}(g)).$$

Lemma 3.1. Let G be a Lie groupoid with Lie algebroid A. For $\alpha \in C^{\bullet}(A)$ and $\xi \in \Gamma(A)$

$$\mathcal{L}_{\xi}(\alpha) = \nabla_{\xi}(\alpha).$$

Proof. Denote by α^L the left-invariant foliated form induced by α . By property (17) and left invariance of α^L one finds that

$$(\varphi_{\xi}^{\epsilon} \cdot \alpha)|_m = (\varphi_{\xi^L}^{\epsilon})^* (\alpha^L)|_m, \text{ for } m \in M$$

Hence, by the usual formula using flows of vector fields,

$$\nabla_{\xi}(\alpha)(m) = (\mathcal{L}_{\xi^L} \alpha^L)(m), \text{ for } m \in M.$$

The result follows by (12).

3.2. The Lie algebroid complex relative to a subgroupoid. Assume that A is the Lie algebroid of G and that B is the Lie algebroid of a wide Lie subgroupoid $K \subseteq G$. Thus, $B \subseteq A$ is a Lie subalgebroid over the same base M. The B-horizontal subspace is defined as

$$\mathsf{C}(A)_{B-\mathrm{hor}} := \bigcap_{\eta \in \Gamma(B)} \ker \iota_{\eta},$$

which, in degree p, we can think of naturally as the space of sections of $\wedge^p(A/B)^*$. The action (13) when restricted to Bis(K) induces K-representations on $\wedge^{\bullet}(A/B)^*$. Explicitly, given $k \in K$ and $\alpha \in \wedge^{\bullet}(A/B)^*_{\mathfrak{s}(k)}$,

(18)
$$k \cdot \alpha = (b \cdot \alpha)_{\mathsf{t}(k)},$$

where $b \in Bis(K)$ is any bisection with the property that b(t(k)) = k.

Definition 3.2. The Lie algebroid complex relative to K is the K-basic subcomplex defined as

(19)
$$\mathsf{C}(A)_{K-\text{basic}} := \{ \alpha \in \mathsf{C}(A)_{B-\text{hor}} \mid k \cdot \alpha_{\mathsf{s}(k)} = \alpha_{\mathsf{t}(k)} \text{ for all } k \in K \}.$$

We denote the resulting cohomology groups of $(\mathsf{C}(A)_{K-\text{basic}}, \mathsf{d}_{CE})$ by $H^{\bullet}(A, K)$.

Observe that by Proposition 3.6, $C(A)_{K-\text{basic}}$ is a subcomplex of $(C(A), \mathsf{d}_{CE})$.

10

Г		٦	
L		I	
L		J	

3.3. The Lie algebroid complex relative to a subalgebroid. Consider now a Lie subalgebroid $B \subseteq A$ over the same base M. The *B*-horizontal subspace

$$\mathsf{C}(A)_{B-\mathrm{hor}} \subseteq \mathsf{C}(A)$$

is defined as in (19) (which we think of as the space of sections of $\wedge^{\bullet}(A/B)^*$). By the identity $[\mathcal{L}_{\eta}, \iota_{\xi}] = \iota_{[\eta,\xi]}$, one checks that the Lie derivative induces *B*-representations on $\wedge^{\bullet}(A/B)^*$ with flat connections¹

(20)
$$\mathcal{L}: \Gamma(B) \times \mathsf{C}^{\bullet}(A)_{B-\mathrm{hor}} \to \mathsf{C}^{\bullet}(A)_{B-\mathrm{hor}}, \quad (\eta, \alpha) \mapsto \mathcal{L}_{\eta}(\alpha).$$

By Lemma 3.1 if A and B are the Lie algebroids of G and of a wide Lie subgroupoid $K \subseteq G$ respectively, then (20) are the infinitesimal representations of the K-actions (18). The B-invariant subspace is analogously defined as the subcomplex

$$\mathsf{C}(A)_{B-\mathrm{inv}} := \cap_{\eta \in \Gamma(B)} \ker \mathcal{L}_{\eta}.$$

Definition 3.3. The *Lie algebroid complex relative to B* is the *B*-basic subcomplex defined as the intersection of the *B*-horizontal and the *B*-invariant subspaces

$$\mathsf{C}(A)_{B-\text{basic}} := \{ \alpha \in \mathsf{C}(A) \mid \iota_{\eta} \alpha = 0 \text{ and } \mathcal{L}_{\eta}(\alpha) = 0 \text{ for all } \eta \in \Gamma(B) \}.$$

We denote the resulting cohomology groups of $(\mathsf{C}(A)_{B-\text{basic}}, \mathsf{d}_{CE})$ by $H^{\bullet}(A, B)$.

Since $\mathcal{L}_{\eta} = \mathsf{d}_{CE}\iota_{\eta} + \iota_{\eta}\mathsf{d}_{CE}$, it follows that the *B*-invariant and the *B*-basic subspaces are subcomplexes of $(\mathsf{C}(A), \mathsf{d}_{CE})$.

Remark 3.4. The usual wedge product turns $(\mathsf{C}(A)_{K-\text{basic}}, \mathsf{d}_{CE}) \subseteq (\mathsf{C}(A), \mathsf{d}_{CE})$ into a DG subalgebra and $H^{\bullet}(A, K)$ (and $H^{\bullet}(A)$) into a graded algebra. Analogous result holds for $H^{\bullet}(A, B)$.

Remark 3.5. For Lie algebras, our definition for the Lie algebroid complex of \mathfrak{g} relative to a Lie subalgebra \mathfrak{k} coincides with the standard basic subcomplex of the Chevalley-Eilenberg complex $C(\mathfrak{g})$. More precisely, the Lie algebroid complex relative to \mathfrak{k} coincides with the \mathfrak{k} -basic subcomplex $C(\mathfrak{g})_{\mathfrak{k}-\text{basic}}$ of $C(\mathfrak{g})$ defined using the coadjoint action of \mathfrak{k} on $C(\mathfrak{g})$.

Similarly, when \mathfrak{g} is the Lie algebras of a Lie group G, and $K \subseteq G$ is a Lie subgroup, the *K*-basic subcomplex $C(\mathfrak{g})_{K-\text{basic}}$ coincides with the *K*-basic subcomplex $C(\mathfrak{g})_{K-\text{basic}}$ of $C(\mathfrak{g})$, now defined using the coadjoint action of K on $C(\mathfrak{g})$.

In the case where A and B are the Lie algebroids of the Lie groupoid G and a wide Lie subgroupoid $K \subseteq G$ respectively, the following result explains the relation between the two relative Lie algebroid complexes.

Proposition 3.6. Let K be a wide Lie subgroupoid of G and denote by B and A their Lie algebroids, respectively. Then $C(A)_{K-\text{basic}}$ is a subcomplex of $(C(A)_{B-\text{basic}}, d_{CE})$. Moreover, if K has connected t-fibers then equality holds.

Proof. By Lemma 3.1 and (15), the inclusion $C(A)_{K-\text{basic}} \subseteq C(A)_{B-\text{basic}}$ is a subcomplex.

To see that we have equality if the t-fibers are connected, we consider $\alpha \in C(A)_{B-\text{basic}}$ and define the fiber bundle map over M

$$e: K \to \wedge^{\bullet} (A/B)^*, \quad k \mapsto k \cdot \alpha_{\mathfrak{s}(k)}.$$

¹Note that for p = 1, \mathcal{L} coincides with the dual of the Bott connection $\Gamma(B) \times \Gamma(A/B) \to \Gamma(A/B)$, $(\eta, \xi \mod B) \to [\eta, \xi] \mod B$

This map intertwines t and the projection, keeping the base M fixed. Note that for any $m \in M \subseteq K$ and $\eta \in \Gamma(B)$,

$$d_m e(\eta) = \frac{d}{d\epsilon} \mid_{\epsilon=0} \varphi_{\eta}^{\epsilon}(m) \cdot (\alpha) = \mathcal{L}_{\eta}(\alpha)_m = 0.$$

For any other $k \in K$,

$$d_k e(\eta^L) = \frac{d}{d\epsilon} \mid_{\epsilon=0} \varphi_{\eta^L}^{\epsilon}(k) \cdot \alpha = \frac{d}{d\epsilon} \mid_{\epsilon=0} k \cdot \left(\varphi_{\eta}^{\epsilon}(\mathbf{s}(k)) \cdot \alpha\right) = k \cdot \left(\mathcal{L}_{\eta}(\alpha)_{\mathbf{s}(k)}\right) = 0,$$

since $\varphi_{\eta^L}^{\epsilon}(k) = k \cdot \varphi_{\eta}^{\epsilon}(\mathbf{s}(k))$. This implies that de = 0 on ker $(d\mathbf{t})$ and therefore by connectedness of the t-fibers, it is constant along the fibers, i.e. for $k \in K$,

$$k \cdot \alpha_{\mathsf{s}(k)} = \mathsf{t}(k) \cdot \alpha_{\mathsf{t}(k)} = \alpha_{\mathsf{t}(k)}.$$

4. The transitive case

In this section construct extension maps from the basic subcomplexes of the isotropy Lie groups and isotropy Lie algebras (see Remark 3.5), to the relative cochain complexes of transitive Lie groupoids and Lie algebroids respectively. The outcome is that in the transitive case, the relative cochain complexes of Lie groupoids and Lie algebroids are naturally isomorphic to the cochain basic subcomplexes of their isotropy Lie groups and Lie algebras respectively (see Theorems 4.1 and 4.3).

4.1. Lie groupoid cohomology relative to a transitive subgroupoid. Let G be a Lie groupoid over M and let $K \subseteq G$ be a wide Lie subgroupoid. Assume that K is *transitive*, in the sense that it has only one orbit: $t(s^{-1}(m)) = M$, for $m \in M$. Hence, G is transitive as well.

Fix now $z \in M$ and let $K_z := s^{-1}(z) \cap t^{-1}(z) \cap K \subseteq K$, $G_z := s^{-1}(z) \cap t^{-1}(z) \subseteq G$ be the isotropy groups. Let $f \in \mathsf{C}^p(G_z)^{K_z}$ and define

(21)
$$\operatorname{ext}(f)(g_1, \dots, g_p) \coloneqq f(k_0^{-1}g_1k_1, \dots, k_{p-1}^{-1}g_pk_p), \quad (g_1, \dots, g_p) \in B_pG;$$

here $k_0, \ldots, k_n \in K$ are so that $\mathbf{s}(k_i) = z$ and $\mathbf{t}(k_i) = m_i$ for all $i = 0, \ldots, p$. This is well defined as f is invariant w.r.t. p + 1 commuting K-actions (4). As a direct consequence of this definition we have:

Theorem 4.1. Suppose K is a transitive Lie subgroupoid of the Lie groupoid G. The extension operator

$$\operatorname{ext} \colon \mathsf{C}^{\bullet}(G_z)^{K_z} \to \mathsf{C}^{\bullet}(G)^K$$

defined by (21) is an isomorphism of complexes, which respects the cup product. Moreover, the restriction map

$$r \colon \mathsf{C}^{\bullet}(G)^K \to \mathsf{C}^{\bullet}(G_z)^{K_z}$$

is a cochain map inverse to ext, which also respects the cup product.

An analogous result holds for $\widetilde{\mathsf{C}}^{\bullet}(G_z)^{K_z}$ and $\widetilde{\mathsf{C}}^{\bullet}(G)^K$.

4.2. Lie algebroid cohomology relative to a transitive subgroupoid. Let A be the Lie algebroid of G, and B the Lie algebroid of a wide Lie subgroupoid $K \subseteq G$. Assume that K is transitive. In this case, the anchor map a of A, and its restriction to B, is point-wise surjective onto TM. Therefore, the kernels

$$\mathfrak{g} := \ker \mathsf{a} \subseteq A, \ \mathfrak{k} := B \cap \ker \mathsf{a} \subseteq B$$

are constant rank ideals of A and B, respectively, fitting into the following two exact sequences over M

$$0 \longrightarrow \mathfrak{g} \longrightarrow A \xrightarrow{\mathsf{a}} TM \longrightarrow 0 \quad \text{and} \quad 0 \longrightarrow \mathfrak{k} \longrightarrow B \xrightarrow{\mathsf{a}} TM \longrightarrow 0.$$

Hence, the inclusion $\mathfrak{g} \subseteq A$ gives the identification

 $A/B \simeq \mathfrak{g}/\mathfrak{k},$

and therefore the restriction map $(C(A), d_{CE}) \rightarrow (C(\mathfrak{g}), d_{CE})$ induces a natural identification of the *B*-horizontal subspace of C(A) with the \mathfrak{k} -horizontal subspace of $C(\mathfrak{g})$:

(22)
$$\mathsf{C}(A)_{B-\mathrm{hor}} = \mathsf{C}(\mathfrak{g})_{\mathfrak{k}-\mathrm{hor}}.$$

We also have a K-basic subcomplex of $C(\mathfrak{g})$ defined as follows. Given a bisection $b \in Bis(K)$, the differential of (14) -the conjugation map by b- sends \mathfrak{g} to \mathfrak{g} , and \mathfrak{k} to \mathfrak{k} ; therefore we have actions of K on $\wedge^{\bullet}(\mathfrak{g}/\mathfrak{k})^*$ analogous to the actions (18). We define the K-basic subcomplex of $C(\mathfrak{g})$ as

$$\mathsf{C}(\mathfrak{g})_{K-\text{basic}} = \{ \alpha \in \mathsf{C}(\mathfrak{g})_{\mathfrak{k}-\text{hor}} \mid k \cdot \alpha_{\mathsf{s}(g)} = \alpha_{\mathsf{t}(g)}, \ k \in K \}.$$

From the construction is clear that the natural identification (22) restricts to their K-basic subcomplexes:

(23)
$$C(A)_{K-\text{basic}} = C(\mathfrak{g})_{K-\text{basic}}.$$

Fix now $z \in M$. The Lie algebras of the isotropy groups K_z and G_z are \mathfrak{k}_z and \mathfrak{g}_z respectively. Given² $\alpha \in \mathsf{C}(\mathfrak{g}_z)_{K_z-\text{basic}}$, we define $\operatorname{ext}(\alpha) \in \mathsf{C}(\mathfrak{g})$ as

(24)
$$\operatorname{ext}(\alpha)_m := C_b^*(\alpha), \quad m \in M,$$

where $b: M \to K$ is any bisection with the property that s(b(m)) = z, and C_b is conjugation by b (see (14)).

Proposition 4.2. For ext: $C^{\bullet}(\mathfrak{g}_z)_{K_z-\text{basic}} \to C^{\bullet}(\mathfrak{g})$ as above, we have the following:

(a) ext is well defined, i.e., $ext(\alpha)_m$, $m \in M$ does not depend on the choice of $b \in Bis(K)$,

(b) the image of ext is contained in $C(\mathfrak{g})_{K-\text{basic}}$, and

(c) ext is a map of complexes, i.e., $\operatorname{ext} \circ \mathsf{d}_{CE} = \mathsf{d}_{CE} \circ \operatorname{ext}$.

Proof. In general, for $b, a \in Bis(G)$,

$$C_{b \cdot a} = C_a \circ C_b.$$

²By Remark 3.5, this is the K-basic subcomplex (w.r.t. coadjoint action of K) of the Chevalley-Eilenberg complex of the Lie algebra \mathfrak{g}_z .

MARÍA AMELIA SALAZAR

Using this identity we check (a) as follows: for $b, a \in Bis(K)$ with s(b(m)) = s(a(m)) = z, we have that $b^{-1} \cdot a \in Bis(K)$ is such that $b^{-1} \cdot a(z) \in K_z$. For $\alpha \in C^{\bullet}(\mathfrak{g}_z)_{K_z-\text{basic}}$,

$$C_b^*(\alpha) = C_b^*((b^{-1} \cdot a(z)) \cdot \alpha) = C_b^*(C_{b^{-1} \cdot a}^*(\alpha)) = C_b^*(C_{b^{-1}}^* \circ C_a^*(\alpha)) = C_a^*(\alpha)$$

where in the first equality we used that α is invariant under the coadjoint action of K_z .

For (b), note that as $\mathfrak{k} = \ker(\mathfrak{a} : B \to TM)$, then for any bisection b of K, the differential of C_b sends \mathfrak{k} to \mathfrak{k} . Hence the image of ext lies inside $\mathsf{C}(\mathfrak{g})_{\mathfrak{k}-\mathrm{hor}}$. Moreover, α is also K-invariant. Indeed, for any $a \in \mathrm{Bis}(K)$, let $m \in M$ and $x = \mathsf{s}(a(m))$; for $b \in \mathrm{Bis}(K)$ with $\mathsf{s}(b(m)) = z$, we have that $\mathsf{s}(a^{-1} \cdot b(x)) = z$, hence

$$a \cdot (\operatorname{ext}\alpha)_x = a \cdot \left(C^*_{a^{-1} \cdot b}(\alpha)\right) = C^*_a(C^*_{a^{-1}} \circ C^*_b(\alpha)) = C^*_b(\alpha) = \operatorname{ext}(\alpha)_m.$$

Item (c) follows by a straightforward computation by noting that $DC_b : \mathfrak{g} \to \mathfrak{g}$ is a Lie algebroid morphism and that both $\mathfrak{g}, \mathfrak{g}_z$ have trivial anchor.

By our previous Proposition 4.2, and keeping in mind the natural identification (23), we obtain the following:

Theorem 4.3. Suppose K is a transitive Lie subgroupoid of the Lie groupoid G. The extension operator

$$\operatorname{ext} \colon \mathsf{C}^{\bullet}(\mathfrak{g}_z)_{K_z - \operatorname{basic}} \to \mathsf{C}^{\bullet}(A)_{K - \operatorname{basic}}$$

defined by (24) is an isomorphism of complexes, which respects the wedge product. The restriction map

$$r \colon \mathsf{C}^{\bullet}(A)_{K-\text{basic}} \to \mathsf{C}^{\bullet}(\mathfrak{g}_z)_{K_z-\text{basic}}$$

is a cochain map inverse to ext, which also respects the wedge product.

5. VAN EST MAPS

In this section we study the van Est maps in the context of relative cohomology. We define tubular structures and a Cartan decomposition for a pair (G, K) consisting of a Lie groupoid G and Lie subgroupoid K, and make use of this notion to define van Est integration maps (see Theorem 5.12). We pay special attention to the cases where K is proper, and in addition K is transitive (see Theorem 5.17). We conclude with the example of $GL_n(\mathbb{C})$, where we apply our results.

5.1. Tubular structures and Cartan decomposition for (G, K). Let K be a closed embedded Lie subgroupoid of G, e.g. any proper Lie subgroupoid K. Recall from [20] that in this case, the space of left cosets G/K is a Hausdorff manifold and the projection $G \to G/K$ is a submersion. Denote by A and B the Lie algebroids of G and K respectively.

Denote by $\overline{t}: G/K \to M$ the surjective submersion induced by the target map. Note that $\overline{u}: M \to G/K$, the map induced by the unit map of G, is a canonical section of \overline{t} . Along this unit section $M \subseteq G/K$, the \overline{t} -fibers give a canonical decomposition of the tangent space $T(G/K)|_M = TM \oplus A/B$. In particular, the normal bundle of M in G/K is naturally identified with A/B. This motivates the following definition:

14

Definition 5.1. A tubular structure of (G, K) is diffeomorphism of bundles over M

$$\begin{array}{c} A/B \xrightarrow{\phi} G/K \\ q \downarrow \qquad \qquad \downarrow \overline{\mathfrak{t}} \\ M \xrightarrow{\operatorname{id}_M} M \end{array}$$

where $q: A/B \to M$ denotes the projection, such that $\bar{\mathfrak{t}} \circ \phi = q$ and $\phi(0_m) = \bar{\mathfrak{u}}(m)$ for $m \in M$ **Definition 5.2.** A *Cartan decomposition* of (G, K) is a *K*-equivariant diffeomorphism

 $\begin{array}{ccc} (25) & & A/B_q \times_{\mathfrak{t}} K \xrightarrow{\Psi} G \\ & q & & \downarrow_{\mathfrak{t}} \\ & & M \xrightarrow{\operatorname{id}_M} & M \end{array}$

of the form

(26) $\Psi(\xi,k) = \mathsf{m}(\psi(\xi),k),$

where $\psi : A/B \to G$ is a smooth map taking values on the isotropy groups, such that $\mathfrak{t} \circ \psi = q$ and taking M to K. Here K acts on the fiber product by the diagonal action (acting on A/Bby the action dual to (18) and by left multiplication on K) and on G by left multiplication.

Remark 5.3. A Cartan decomposition of (G, K) induces a tubular structure for G/K. Indeed, the map

(27)
$$\phi: A/B \to G/K, \quad \xi \mapsto \psi(\xi)K$$

is the required diffeomorphism.

Remark 5.4. In Definition 5.2 note that the commutativity of the diagram (25) is equivalent to the condition $\mathbf{t} \circ \psi = q$. Moreover, the reason to impose that ψ takes values on the isotropy groups is that in order to ensure that the multiplication $\mathsf{m}(\psi(\xi), k)$ is well-defined for $(\xi, k) \in A/B_q \times_{\mathbf{t}} K$, we need that ψ sends q-fibers to s-fibers so that $\mathsf{s}(\psi(\xi)) = q(\xi) = \mathsf{t}(k)$.

Furthermore, the fact that (27) is a bijective map imposes further restrictions for the pair (G, K). In particular, K is forced to have the same orbits as G and, denoting by $G(m, m') = s^{-1}(m') \cap t^{-1}(m)$, for $m, m' \in M$ -the arrows of G with source m' and target m (and similarly for K),

$$G(m, m') \cdot K(m', m) \subseteq \psi((A/B)_m).$$

The following example explains the relation that our Definitions 5.1 and 5.2 have with standard results in Lie theory. See e.g. [16, Theorem 6.31] and [22, Theorem 32.5].

Example 5.5. Let G be a Lie group with finitely many connected components and Lie algebra \mathfrak{g} , and let $K \subseteq G$ be a maximal compact subgroup with Lie algebra \mathfrak{k} . Then the diffeomorphism $G/K \simeq \mathfrak{g}/\mathfrak{k}$ is a tubular structure of (G, K).

Suppose that G is, moreover, a non-compact semisimple Lie group. Let θ be a Cartan involution of \mathfrak{g} . Then the eigenspaces \mathfrak{k} and \mathfrak{p} corresponding to the eigenvalues +1 and -1, respectively, form a Cartan pair of $\mathfrak{g}: \mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ and

(28)
$$[\mathfrak{k},\mathfrak{k}] \subseteq \mathfrak{k}, \ [\mathfrak{k},\mathfrak{p}] \subseteq \mathfrak{p}, \ \mathrm{and} \ [\mathfrak{p},\mathfrak{p}] \subseteq \mathfrak{k}.$$

Let K be the maximal compact subgroup with Lie algebra \mathfrak{k} . Then the global Cartan decomposition

(29)
$$\mathfrak{p} \times K \to G, \quad (\xi, k) \mapsto \mathsf{m}(\exp(\xi), k)$$

is a Cartan decomposition of the pair (G, K); here $\mathfrak{g}/\mathfrak{k}$ is canonically identified with \mathfrak{p} .

Example 5.6. A transitive Lie groupoids always admits proper subgroupoid with a tubular structure, under the assumption that the isotropy Lie groups have finitely many connected components. To see this we can assume that the groupoid is the Atiyah groupoid $\mathcal{G} := (P \times P)/G$ of a principal G-bundle $P \to M$, where G a Lie group with finitely many components, acting on $P \times P$ by the diagonal action.

Consider now the maximal compact subgroup K of G and choose a reduction of the structural group to K, i.e. a principal K-subbundle $Q \to M$. Such a Q always exists as this is equivalent to having a section of the bundle $P \times_G (G/K) \to M$ with contractible fibers G/K. Then, we have the isomorphism of principal G-bundles

$$Q \times_K G \xrightarrow{\simeq} P, \quad [p,g] \mapsto p \cdot g,$$

and therefore,

$$(P \times P)/G \simeq (Q \times Q) \times_{K \times K} (G \times G)/G \simeq (Q \times Q) \times_{K \times K} G$$
$$(Q \times Q)/K \simeq (Q \times Q) \times_{K \times K} (K \times K)/K \simeq (Q \times Q) \times_{K \times K} K,$$

where on the right hand side the right action of $K \times K$ on G (and K) is given by $(a, b) \cdot g = a^{-1}gb$. Hence, $\mathcal{K} := (Q \times Q)/K \subseteq (P \times P)/G = \mathcal{G}$ is a Lie subgroupoid which is proper as the fiber of $Q \to M$ is compact, and with the property that the isotropy group K of $(Q \times Q)/K$ is the maximal compact subgroup of the isotropy group G of $(P \times P)/G$.

Now, the space of left cosets \mathcal{G}/\mathcal{K} is diffeomorphic as a bundle over M to the space of left cosets $\overline{\mathcal{G}}/\overline{\mathcal{K}}$, where $\overline{\mathcal{G}}$ and $\overline{\mathcal{K}}$ are the Lie subgroupoids of \mathcal{G} and \mathcal{K} given by the union of the isotropy groups. Analogously, the quotient A/B of the Lie algebroids of \mathcal{G} and \mathcal{K} is isomorphic as a vector bundle to the quotient $\overline{A}/\overline{B}$ of the kernels of the anchors $\overline{A} = \ker(\mathbf{a}) \subseteq A$, $\overline{B} = \ker(\mathbf{a}) \subseteq B$. As \overline{A} and \overline{B} are locally trivial Lie algebra bundles with fibers $\overline{A}_m \simeq \mathfrak{g}$, $\overline{B}_m \simeq \mathfrak{k}$ – the Lie algebras of G and K, respectively – then the isomorphism $\mathfrak{g}/\mathfrak{k} \simeq G/K$ extends to a tubular structure

$$A/B \simeq A/B \rightarrow \mathcal{G}/\mathcal{K} \simeq \mathcal{G}/\mathcal{K}$$

Similarly a transitive Lie groupoid with non-compact semisimple isotropy Lie group admit a proper subgroupoid with a Cartan decomposition. In this case, the Cartan decomposition for the isotropy Lie groups (29) extends to a diffeomorphism $A/B_q \times_t \bar{\mathcal{K}} \xrightarrow{\bar{\Psi}} \bar{\mathcal{G}}$ that extends naturally to a Cartan decomposition

$$A/B_q \times_{\mathsf{t}} \mathcal{K} \xrightarrow{\Psi} \mathcal{G}.$$

5.2. Van Est differentiation map for relative complexes. The van Est differentiation map is a cochain map

$$VE_G \colon \mathsf{C}^{\bullet}(G) \to \mathsf{C}^{\bullet}(A).$$

(In appendix A we recall from [19] how the van Est differentiation map is obtained using the Perturbation Lemma.) Explicitly, the commuting left *G*-actions on B_pG obtained from (4) in the standard fashion have generating vector fields $\xi^{(i)} \in \mathfrak{X}(B_pG), \xi \in \Gamma(A), i = 0, \ldots, p$.

Theorem 5.7 ([27]). The map VE_G is given by the formula

(30)
$$VE_G(f)(\xi_1, \dots, \xi_p) = \sum_{s \in S_p} \operatorname{sign}(s) \ \mathcal{L}_{\xi_{s(1)}^{(1)}} \cdots \mathcal{L}_{\xi_{s(p)}^{(p)}}(f)|_M$$

for $f \in C^p(G)$ and $\xi_1, \ldots, \xi_p \in \Gamma(A)$. Here the sum is over the permutation group S_p and M is regarded as a submanifold of B_pG consisting of constant trivial p-arrows.

The previous result is also considered in [6] and the formula (30) is found in [17]. We obtain the following result:

Theorem 5.8. Let K be a wide Lie subgroupoid of G. Then VE_G restricts to a cochain map of the K-basic subcomplexes

$$VE_G \colon \mathsf{C}^{\bullet}(G)^K \to \mathsf{C}^{\bullet}(A)_{K-\text{basic}}$$

Moreover, when restricted to $\widetilde{\mathsf{C}}^{\bullet}(G)^K$, VE_G is an algebra map.

Of course, in Theorem 5.8 we are referring to the products of Remarks 2.3 and 3.4.

Proof. Let $f \in C^p(G)^K$. To see that $\iota_\eta(VE(f)) = 0$ for any $\eta \in \Gamma(B)$, note that the Lie derivatives on expression (30) commute as the *G*-actions are commutative. Hence $VE(f)(\xi_1, \ldots, \xi_{p-1}, \eta)$ is computed as the sum (with appropriate signs) of expressions of the form

$$\mathcal{L}_{\xi_{s(1)}^{(1)}}\cdots\mathcal{L}_{\xi_{s(p-1)}^{(p-1)}}\mathcal{L}_{\eta^{(i)}}(f)|_{M}.$$

But f is invariant w.r.t. the p+1 K-actions, and so $\mathcal{L}_{\eta^{(i)}}(f) = 0, i = 1, \dots, p$.

To see that VE(f) is K invariant, let $b \in Bis(K)$ and $m \in M$. Then

(31)

$$(b \cdot VE_G(f))(\xi_1, \dots, \xi_p)(m) = \sum_{s \in S_p} \operatorname{sign}(s) \mathcal{L}_{(C_{b,*}(\xi_{s(1)}))^{(1)}} \cdots \mathcal{L}_{(C_{b,*}(\xi_{s(p)}))^{(p)}}(f)|_{(C_b(m), \dots, C_b(m))}.$$

Now, the generator vector fields $\xi^{(i)} \in \mathfrak{X}(B_pG), \ \xi \in \Gamma(A)$ are of the form

$$\xi_{g_1,\dots,g_p}^{(i)} = \begin{cases} (\xi_{g_1}^R, 0_{g_2}, \dots, 0_{g_p}) & i = 0, \\ (0_{g_1},\dots, di(\xi)_{g_i}^L, \xi_{g_{i+1}}^R,\dots, 0_{g_p}) & 0 < i < p, \\ (0_{g_1},\dots, 0_{g_{p-1}}, di(\xi)_{g_p}^L) & i = p. \end{cases}$$

where $\xi^L, \xi^R \in \mathfrak{X}(G)$ denote the left and right invariant vector fields associated to ξ , respectively. From this description, it is a straightforward computation to see for any $b \in Bis(G)$, the pushforward of $\xi^{(i)}$ by the diffeomorphism $C_b^p: B_pG \to B_pG, \ C_b^p = C_b \times \cdots \times C_b$, where C_b is conjugation by b, is such that

$$C_{b,*}^{p}(\xi^{(i)}) = (C_{b,*}(\xi))^{(i)}$$

Since f is K-invariant, $(C_b^{p,-1})^*(f) = f$ for any $b \in Bis(G)$. Hence, we compute

$$\mathcal{L}_{(C_{b,*}(\xi_{s(1)}))^{(1)}}\cdots\mathcal{L}_{(C_{b,*}(\xi_{s(p)}))^{(p)}}(f) = \mathcal{L}_{(C_{b,*}(\xi_{s(1)}))^{(1)}}\cdots\mathcal{L}_{(C_{b,*}(\xi_{s(p)}))^{(p)}}((C_{b}^{p,-1})^{*}(f))$$
$$= \mathcal{L}_{C_{b,*}^{p}}(\xi_{s(1)}^{(1)})\cdots\mathcal{L}_{C_{b,*}^{p}}(\xi_{s(p)}^{(p)})((C_{b}^{p,-1})^{*}(f)) = (C_{b}^{p,-1})^{*}(\mathcal{L}_{\xi_{s(1)}^{(1)}}\cdots\mathcal{L}_{\xi_{s(p)}^{(p)}}(f)).$$

MARÍA AMELIA SALAZAR

In the last line we use recursively that $\mathcal{L}_{C_{b,*}^{p}(\xi^{(i)})}((C_{b}^{p,-1})^{*}(F)) = (C_{b}^{p,-1})^{*}(\mathcal{L}_{\xi^{(i)}}(F))$ for any function $F \in C^{\infty}(B_{p}G)$. Hence,

(32)
$$\mathcal{L}_{\left(C_{b,*}(\xi_{s(1)})\right)^{(1)}} \cdots \mathcal{L}_{\left(C_{b,*}(\xi_{s(p)})\right)^{(p)}}(f)|_{(C_{b}(m),\dots,C_{b}(m))} = \mathcal{L}_{\xi_{s(1)}^{(1)}} \cdots \mathcal{L}_{\xi_{s(p)}^{(p)}}(f)|_{(m,\dots,m)}$$

Putting together (31) and (32) we obtain that $VE(f) \in \mathsf{C}^p(A)^K$.

Finally, to see that when restricted to $\widetilde{\mathsf{C}}^{\bullet}(G)^K$, VE_G becomes an algebra map, recall the following. When considering the normalized subcomplex $\widetilde{\mathsf{C}}^{\bullet}(G) \subseteq \mathsf{C}^{\bullet}(G)$ of functions with the property that $f(g_1, \ldots, g_p) = 0$ whenever g_i is a unit for some i, the restriction $VE_G : \widetilde{\mathsf{C}}^{\bullet}(G) \to \mathsf{C}^{\bullet}(A)$ is an algebra map [6, 17]. As $\widetilde{\mathsf{C}}^{\bullet}(G)^K \subseteq \widetilde{\mathsf{C}}^{\bullet}(G)$ and $\mathsf{C}^{\bullet}(A)_{K-\text{basic}} \subseteq \mathsf{C}^{\bullet}(A)$ are subalgebras, then $VE_G|_{\widetilde{\mathsf{C}}^{\bullet}(G)^K}$ is also an algebra map.

5.3. Van Est differentiation maps in the proper case. Choose any left invariant left invariant normalized Haar system μ on the proper wide subgroupoid K and denote by $VE_{G/K}$ the composition

(33)
$$VE_{G/K} \colon \mathsf{C}^{\bullet}(G) \xrightarrow{\operatorname{Av}} \mathsf{C}^{\bullet}(G)^K \xrightarrow{VE_G} \mathsf{C}^{\bullet}(A)_{K-\text{basic}}$$

here Av is the homotopy equivalence given by the averaging operator (11) associated to μ .

The purpose of this section is to derive $VE_{G/K}$ using the Perturbation Lemma. We refer to Appendix A, where we summarize some of the results and techniques of [19] to obtain VE_G (see (30)) via the Perturbation Lemma.

Consider the van Est double complex $\mathsf{D}^{p,q}(G) = \Gamma(\pi_p^*(\wedge^q A^*))$, where $\pi_p \colon E_p G = B_p G \times_M G \to M$, $\pi_p(g_1, \ldots, g_p; g) = \mathsf{s}(g)$ – see (64) – that relates $\mathsf{C}^{\bullet}(G)$ with $\mathsf{C}^{\bullet}(A)$. For the relative version, the two complexes $\mathsf{C}^{\bullet}(G)$, $\mathsf{C}^{\bullet}(A)_{K-\text{basic}}$ are related by the K-basic subcomplex

(34)
$$(\mathsf{D}^{\bullet,\bullet}(G)_{K-\mathrm{basic}},\delta,\mathsf{d})$$

of $\mathsf{D}^{\bullet,\bullet}(G)$; the K-action is

(35)
$$(a \cdot \psi)(g_1, \dots, g_p; g) = a \cdot \psi(g_1, \dots, g_p; ga)$$

The double complex comes with horizontal and vertical augmentation maps i: $C^{\bullet}(A)_{K-\text{basic}} \rightarrow D^{0,\bullet}(G)_{K-\text{basic}}$ given by the pullback π_0^* and by j: $C^{\bullet}(G) \rightarrow D^{\bullet,0}(G)_{K-\text{basic}}$ given in degree p by the pullback map κ_p^* , where $\kappa_p \colon E_pG \rightarrow B_pG$ is the projection. The augmentation maps define cochain maps to the total complex

$$C^{\bullet}(G) \xrightarrow{j} \operatorname{Tot}^{\bullet}(\mathsf{D}(G)_{K-\text{basic}}) \xleftarrow{i} C^{\bullet}(A)_{K-\text{basic}}$$

The double complex also comes with a horizontal homotopy operator $h_K \colon D^{\bullet,\bullet}(G)_{K-\text{basic}} \to D^{\bullet-1,\bullet}(G)_{K-\text{basic}}$ given by

(36)
$$(\mathsf{h}_{K}\psi)(g_{1},\ldots,g_{p-1};g) = (-1)^{p} \int_{\mathsf{t}^{-1}(m)} \psi(g_{1},\ldots,g_{p-1},ga;a^{-1})\mu_{m}(a);$$

here m = s(g). Indeed, a direct calculation shows that $\delta h_K + h_K \delta = 1 - i \circ p_K$ where p_K vanishes on elements of bidegree (p,q) with p > 0, while

(37)
$$\mathbf{p}_{K}(\psi) = \int_{\mathbf{t}^{-1}(m)} \psi(a^{-1}) \mu_{m}(a),$$

for $\psi \in \Gamma(\pi_0^*(\wedge^q A^*)))_{K-\text{basic}}$.

Using the Perturbation Lemma A.1, we obtain a cochain map

$$\mathsf{p}_K \circ (1 + \mathsf{dh}_K)^{-1} \circ \mathsf{j} \colon \mathsf{C}^{\bullet}(G) \to \mathsf{C}^{\bullet}(A)_{K-\text{basic}}.$$

Remark 5.9. For K = M, one recovers the homotopy operator

$$(\mathsf{h}\psi)(g_1,\ldots,g_{p-1};g) = (-1)^p \psi(g_1,\ldots,g_{p-1},g;m).$$

and the map $\mathbf{p} = \mathbf{u}^*$, discussed the Appendix A. In particular, we recover the formula $VE_G = \mathbf{p} \circ (1 + \mathbf{dh})^{-1} \circ \mathbf{j}$ from [19].

The following form of $VE_{G/K}$ via the Perturbation Lemma will allow us to conclude in Theorem 5.12 that in the presence of tubular structures for (G, K), $VE_{G/K}$ becomes a homotopy equivalence.

Proposition 5.10. For any proper wide Lie subgroupoid $K \subseteq G$, the cochain map $p_K \circ (1 + dh_K)^{-1} \circ j$ equals

$$\mathsf{p} \circ (1 + \mathsf{dh})^{-1} \circ \mathsf{j} \circ \operatorname{Av}$$
.

In particular, $VE_{G/K} = \mathbf{p}_K \circ (1 + \mathbf{dh}_K)^{-1} \circ \mathbf{j}$.

In preparation for the proof of Proposition 5.10, in analogy with (10) we denote by

$$\operatorname{Av}_{\mathsf{D}(G)}^{(i)} \colon \mathsf{D}^{p,q}(G) \to \mathsf{D}^{p,q}(G)$$

the induced averaging operator with respect to the i-th K-action (7), and by

$$\operatorname{Av}_{\mathsf{D}(G)} = \operatorname{Av}_{\mathsf{D}(G)}^{(0)} \circ \cdots \circ \operatorname{Av}_{\mathsf{D}(G)}^{(p)}$$

the total K^{p+1} -averaging operator. It is immediate to see that

(38)
$$\mathbf{h}_{K}|_{\mathbf{D}^{p,q}(G)} = \mathbf{h} \circ \operatorname{Av}_{\mathbf{D}(G)}^{(p)}|_{\mathbf{D}^{p,q}(G)}, \quad \mathbf{p}_{K} = \mathbf{p} \circ \operatorname{Av}_{\mathbf{D}(G)}^{(0)},$$

(39)
$$\mathsf{h} \circ \operatorname{Av}_{\mathsf{D}(G)}^{(p-1)}|_{\mathsf{D}^{p,q}(G)} = \operatorname{Av}_{\mathsf{D}(G)}^{(p-1)} \circ \mathsf{h}|_{\mathsf{D}^{p,q}(G)}, \quad \operatorname{Av}_{\mathsf{D}(G)} \circ \mathsf{j} = \mathsf{j} \circ \operatorname{Av}.$$

Lemma 5.11. For all i, the averaging operators commute with the vertical differential:

$$\mathsf{d} \circ \operatorname{Av}_{\mathsf{D}(G)}^{(i)} = \operatorname{Av}_{\mathsf{D}(G)}^{(i)} \circ \mathsf{d}.$$

Proof. By (66) we have that $\mathbf{d} = -(1)^p \mathbf{d}_{Rh}$ is the foliated de Rham differential on elements of bidegree (p,q), under the isomorphism $\pi_p^* A \cong T_{\mathcal{F}}(E_p G)$. For a section $\xi \in \Gamma(A)$, the pullback $\pi_p^*(\xi) \in \Gamma(\pi_p^* A)$ is identified with $\hat{\xi} \in \Gamma(T_{\mathcal{F}} E_p)$ given by

$$\hat{\xi}_{(g_1,\ldots,g_p;g)} = (0_{g_1},\ldots,0_{g_p},\xi_g^L).$$

Hence, the flow of $\hat{\xi}$ is given by

$$\varphi_{\hat{\xi}}^{\epsilon}(g_1,\ldots,g_p;g) = \left(g_1,\ldots,g_p;g\varphi_{\xi}^{\epsilon}(\mathsf{s}(g))\right)$$

(cf. (17)). From this expression it is immediate to see that $\varphi_{\hat{\xi}}^{\epsilon}$ commutes with the p+1 *G*-actions (7) and therefore, for $\psi \in C^{\infty}(E_pG)$,

$$\mathsf{d}_{Rh}\left(\operatorname{Av}_{\mathsf{D}(G)}^{(i)}\psi\right)(\hat{\xi}) = \operatorname{Av}_{\mathsf{D}(G)}^{(i)}\left(\mathsf{d}_{Rh}\psi\right)(\hat{\xi})$$

(i = 0, ..., p). Since $\Gamma(T_{\mathcal{F}}E_p)$ is generated by sections $\hat{\xi}$, $\xi \in \Gamma(A)$ then $\mathsf{d}_{Rh} \circ \operatorname{Av}_{\mathsf{D}(G)}^{(i)} \psi = \operatorname{Av}_{\mathsf{D}(G)}^{(i)} \circ \mathsf{d}_{Rh} \psi$ for any $\psi \in C^{\infty}(E_pG)$. Hence, $\mathsf{d}_{Rh} \circ \operatorname{Av}_{\mathsf{D}(G)}^{(i)} \psi = \operatorname{Av}_{\mathsf{D}(G)}^{(i)} \circ \mathsf{d}_{Rh} \psi$ for $\psi \in \Gamma(\wedge^q T_{\mathcal{F}}^* E_p) \simeq \mathsf{D}^{p,q}(G)$ and any $q \ge 0$.

Proof of Proposition 5.10. At degree p, we compute $\mathbf{p}_K \circ (1 + \mathbf{dh}_K)^{-1} \circ \mathbf{j}$:

$$\begin{aligned} \mathsf{p}_{K} \circ (\mathsf{dh}_{K})^{p} \circ \mathsf{j} &= \mathsf{p} \circ \operatorname{Av}_{\mathsf{D}(G)}^{(0)} (\mathsf{dh}_{K})^{p} \circ \mathsf{j} \\ &= \mathsf{p} \circ \operatorname{Av}_{\mathsf{D}(G)}^{(0)} \circ (\mathsf{dh} \operatorname{Av}_{\mathsf{D}(G)}^{(1)} \circ \cdots \circ \mathsf{dh} \operatorname{Av}_{\mathsf{D}(G)}^{(p)}) \circ \mathsf{j} \\ &= \mathsf{p} \circ (\mathsf{d} \operatorname{Av}_{\mathsf{D}(G)}^{(0)} \mathsf{h} \circ \mathsf{d} \operatorname{Av}_{\mathsf{D}(G)}^{(1)} \mathsf{h} \circ \cdots \circ \mathsf{d} \operatorname{Av}_{\mathsf{D}(G)}^{(p-1)} \mathsf{h} \operatorname{Av}_{\mathsf{D}(G)}^{(p)}) \circ \mathsf{j} \\ &= \mathsf{p} \circ (\mathsf{dh} \operatorname{Av}_{\mathsf{D}(G)}^{(0)} \circ \mathsf{dh} \operatorname{Av}_{\mathsf{D}(G)}^{(1)} \circ \cdots \circ \mathsf{dh} \operatorname{Av}_{\mathsf{D}(G)}^{(p-1)} \operatorname{Av}_{\mathsf{D}(G)}^{(p)}) \circ \mathsf{j} \\ &= \dots \\ &= \mathsf{p} \circ (\mathsf{dh})^{p} \circ \operatorname{Av}_{\mathsf{D}(G)} \circ \mathsf{j} \\ &= \mathsf{p} \circ (\mathsf{dh})^{p} \circ \mathsf{j} \circ \operatorname{Av}. \end{aligned}$$

In the second equality we used (38), in the third equality we applied Lemma 5.11, in the fourth we used the left equation of (39), and in the last equality we used the right equation of (39). \Box

5.4. Van Est integration maps in the proper case. Let K be a proper wide Lie subgroupoid of G. We will use the Perturbation Lemma to obtain van Est integration maps (see Appendix A for background on the Perturbation Lemma). For the discussion, another interpretation of the K-basic (double) complex will be convenient (see the parallel in the case of Lie groups in [19]). In this case, we will make use of a tubular structure to obtain a vertical homotopy operator on the double complex and define the van Est integration map $R_{G/K}: C^{\bullet}(A)_{K-\text{basic}} \to C^{\bullet}(G).$

The manifold G/K comes equipped with a left G-action

$$a \cdot gK = agK$$

with anchor $\overline{t}: G/K \to M$. Let $(\Omega_{\mathcal{F}}(G/K), \mathsf{d}_{Rh})$ be the de Rham complex of \overline{t} -foliated forms. In analogy with (12), the restriction to $M \subseteq G$ takes such a form to a section of $(A/B)^*$, and induces an isomorphism of differential complexes,

(40)
$$(\Omega^{\bullet}_{\mathcal{F}}(G/K)^L, \mathsf{d}_{Rh}) \cong (\mathsf{C}^{\bullet}(A)_{K-\text{basic}}, \mathsf{d}_{CE}).$$

Similarly, elements in the double complex $\mathsf{D}^{p,q}(G)_{K-\text{basic}}$ as in (34) may be identified with functions

(41)
$$\beta \colon B_p G \to \Omega^q_{\mathcal{F}}(G/K)$$

such that $\beta(g_1, \ldots, g_p) \in \Omega^q(\bar{\mathfrak{t}}^{-1}(m))$ for $m = \mathfrak{s}(g_p)$, and smoothly depending on (g_1, \ldots, g_p) . In these terms, the two differentials are

(42)
$$(\mathsf{d}\beta)(g_1,\ldots,g_p) = (-1)^p \mathsf{d}_{Rh}\beta(g_1,\ldots,g_p),$$

where d_{Rh} is the de Rham differential, and

(43)
$$(\delta\beta)(g_1, \dots, g_{p+1}) = \beta(g_2, \dots, g_{p+1}) + \sum_{i=1}^p (-1)^i \beta(g_1, \dots, g_i g_{i+1}, \dots, g_{p+1})$$
$$+ (-1)^{p+1} L(g_{p+1})^* \beta(g_1, \dots, g_p)$$

where $L(a): \bar{t}^{-1}(s(a)) \to \bar{t}^{-1}(t(a))$ is the action of $a \in G$. (For p = 0, this is to be interpreted as $(\delta\beta)(g_1) = \beta - L(g_1)^*\beta$.) The horizontal augmentation map i is the inclusion of the invariant forms (40), while j is the inclusion of $C^{\infty}(B_pG)$ into the space of maps $\beta: B_pG \to \Omega^0_{\mathcal{F}}(G/K)$ such that $\beta(g_1, \ldots, g_p) \in C^{\infty}(\bar{t}^{-1}(m))$ is constant on $\bar{t}^{-1}(m)$, for any given (g_1, \ldots, g_p) with $m = s(g_p)$.

Suppose now that (G, K) admits a tubular structure. We transport the scalar multiplication in A/B to a retraction along \bar{t} -fibers

(44)
$$\lambda \colon [0,1] \times G/K \to G/K, \ (t,gK) \mapsto \lambda_t(gK).$$

Here $\lambda_0 = \bar{u} \circ \bar{t}$. The retraction determines a homotopy operator

$$T: \Omega^q_{\mathcal{F}}(G/K) \to \Omega^{q-1}_{\mathcal{F}}(G/K)$$

given by pullback under λ followed by integration over [0, 1]. This has the properties $T \circ T = 0$ and

$$T\beta|_K = 0$$

It defines a vertical homotopy operator k on the double complex, where

$$(\mathsf{k}\beta)(g_1,\ldots,g_p) = (-1)^p T(\beta(g_1,\ldots,g_p)).$$

That is, $[\mathbf{k}, \mathbf{d}] = 1 - \mathbf{j} \circ \mathbf{q}$ where, for β of bidegree (p, 0),

$$(\mathsf{q}\beta)(g_1,\ldots,g_p)=\beta(g_1,\ldots,g_p)(\mathsf{s}(g_p)K).$$

The properties of T imply that

$$\mathsf{k} \circ \mathsf{k} = 0, \ \mathsf{q} \circ \mathsf{k} = 0.$$

By the Perturbation Lemma A.1, we obtain a cochain map

(45)
$$R_{G/K} = \mathbf{q} \circ (1 + \delta \mathbf{k})^{-1} \circ \mathbf{i} \colon \mathsf{C}^{\bullet}(A)_{K-\text{basic}} \to \mathsf{C}^{\bullet}(G).$$

This formula together with Proposition 5.10 imply that the relative cohomology of the Lie algebroid computes the differentiable cohomology of the Lie groupoid, as stated precisely below:

Theorem 5.12. The van Est integration map $R_{G/K}$ is a homotopy equivalence, which is a right inverse to the van Est differentiation map $VE_{G/K}$

$$VE_{G/K} \circ R_{G/K} = \mathrm{id}_{\mathsf{C}(A)_{K-\mathrm{basic}}}$$

Moreover, if the tubular structure comes from a Cartan decomposition as in Remark 5.3, then $R_{G/K}$ takes values in $\widetilde{\mathsf{C}}^{\bullet}(G)^{K}$, is a right inverse to the restriction $VE_G: \widetilde{\mathsf{C}}^{\bullet}(G)^{K} \to \mathsf{C}^{\bullet}(A)_{K-\text{basic}}$, and induces an algebra isomorphism in cohomology.

MARÍA AMELIA SALAZAR

Remark 5.13. Following Example 5.5, let G be a Lie group with finitely many connected components and Lie algebra \mathfrak{g} , and let $K \subseteq G$ be a maximal compact subgroup. In this context, the van Est integration map (45)

$$R_{G/K} \colon \mathsf{C}^{\bullet}(\mathfrak{g})_{K-\text{basic}} \to \mathsf{C}^{\bullet}(G)$$

and the first part of Theorem 5.12 first appeared in [19, Theorem 3.6] (see also [3]), where also an explicit formula was given in terms of the tubular structure of (G, K). As a bonus from Theorem 5.12 we obtain that, if we are in the setting of Example 5.5 (e.g. if G is a non-compact semisimple Lie group) then $R_{G/K}$ takes values in the normalized K-invariant subcomplex $\tilde{C}^{\bullet}(G)^{K}$ and it becomes an algebra map when passing to cohomology.

To recall the formula for $R_{G/K}$, we denote by $\lambda_t = \lambda(t, \cdot) \colon G/K \to G/K$ the induced scalar multiplication by t, with λ as in (44). For $(g_1, \ldots, g_p) \in G^p$ and $(t_1, \ldots, t_p) \in [0, 1]^p$, let

(46)
$$\gamma_{t_1,\ldots,t_p}^{(p)}(g_1,\ldots,g_p) = \left(\lambda_{t_1} \circ L(g_1) \cdots \circ \lambda_{t_p} \circ L(g_p)\right)(eK).$$

For a fixed $(g_1, \ldots, g_p) \in G^p$, this defines a map $\gamma^{(p)}(g_1, \ldots, g_p) \colon [0, 1]^p \to G/K$. On the other hand,

(47)
$$C^{\bullet}(\mathfrak{g})_{K-\text{basic}} \simeq \Omega^{\bullet}(G/K)^L, \quad \alpha \mapsto \alpha_{G/K} := \alpha \circ \theta$$

where $\alpha_{G/K} \in \Omega^{\bullet}(G/K)^{L}$ is the *G*-invariant form obtained by precomposing α with the left invariant Maurer-Cartan form θ of *G* and then passing to the quotient G/K. The van Est integration map is then given by the formula

(48)
$$R_{G/K}(\alpha)(g_1\dots,g_p) = \int_{[0,1]^p} \gamma^{(p)}(g_1,\dots,g_p)^* \alpha_{G/K}.$$

Proof of Theorem 5.12. By Proposition 5.10, $VE_{G/K}$ is obtained from the Perturbation Lemma using the double complex $\mathsf{D}^{\bullet,\bullet}(G)_{K-\text{basic}}$ with horizontal homotopy h_K and projection $\mathsf{p}_K \colon \mathsf{D}^{0,\bullet}(G)_{K-\text{basic}} \to \mathsf{C}^{\bullet}(A)_{K-\text{basic}}$. Then by Lemma A.2 for the first part it suffices to show that $\mathsf{h}_K \circ \mathsf{k} = 0$ and $\mathsf{p}_K \circ \mathsf{k} = 0$. Given β as in (41),

$$(\mathsf{h}_K\beta)(g_1,\ldots,g_{p-1})(gK) = (-1)^p \int_{\mathsf{t}^{-1}(m)} L((ga)^{-1})^*\beta(g_1,\ldots,g_{p-1},ga) \ \mu_m(a);$$

hence $h\beta = 0$ when $\beta(g_1, \ldots, g_p)|_{s(g_p)K} = 0$ for all $(g_1, \ldots, g_p) \in B_pG$. In particular, this applies when β is in the range of T. This shows $h_{K} \circ k = 0$; the argument for $p_K \circ k = 0$ is similar.

To see that $R_{G/K}$ takes values in $C^{\bullet}(G)^{K}$ when having a Cartan decomposition, denote by $D^{p,q}(G,K)_{K-\text{basic}} \subseteq D(G)_{K-\text{basic}}^{p,q}$ the space of invariant elements β w.r.t. the p+1 K actions (7) (i.e. we restrict the G-actions to K), and so that $\beta(g_1,\ldots,g_p) = 0$ whenever $g_i \in K$ for some *i*. Note that q maps $D^{p,0}(G,K)_{K-\text{basic}}$ to $\widetilde{C}^p(G)^K$. We claim that $D^{\bullet,\bullet}(G,K)_{K-\text{basic}}$ is a subcomplex of $(D(G)_{K-\text{basic}}^{\bullet,\bullet}, \delta, \mathsf{d}_{CE})$, and that the rest of the maps k,i involved in the definition of $R_{G/K}$ send $D(G,K)_{K-\text{basic}}$ to $D(G,K)_{K-\text{basic}}$. Indeed, for the last p+1 action we have that

$$(a \cdot \beta)(g_1, \dots, g_p) = L(a^{-1})^* \beta(g_1, \dots, g_p a)$$

for $a \in K$, $t(a) = s(g_p)$. A direct computation then shows that $D(G, K)_{K-\text{basic}}$ is closed under δ and d_{CE} as in (43) and (42), respectively; similarly, i maps $C^{\bullet}(A)_{K-\text{basic}} = \Omega^q_{\mathcal{F}}(G/K)^L$ to

 $\mathsf{D}^{0,q}(G,K)_{K-\text{basic}}$ -the space of left K-invariant foliated forms of G/K. Moreover, the tubular structure (27) of G/K is K-equivariant hence

$$\lambda_t(a \cdot gK) = a \cdot \lambda_t(gK)$$

for $a \in K$, s(a) = t(g). This implies that

$$L(a^{-1})^*T(\beta(g_1,\ldots,g_pa)) = T(L(a^{-1})^*\beta(g_1,\ldots,g_pa)).$$

With this, a direct computation shows that $D(G, K)_{K-\text{basic}}$ is closed under k, hence implying that $R_{G/K}$ lands in $\tilde{C}^{\bullet}(G)^{K}$. To see that in this case $R_{G/K}$ is an algebra map in cohomology, note that in this case $VE_G|_{\tilde{C}^{\bullet}(G)^{K}} \circ R_{G/K} = \text{id}$. As $VE_G|_{\tilde{C}^{\bullet}(G)^{K}}$ is an algebra map by Theorem 5.8, we then obtain that when passing to cohomology $R_{G/K}$ is an isomorphism with inverse the algebra map $VE_G|_{\tilde{C}^{\bullet}(G)^{K}}$. This implies that in cohomology $R_{G/K}$ is an algebra map. \Box

5.5. The proper and transitive case. By Example 5.6 we know that any transitive Lie groupoid G satisfying a very mild assumption admits a proper transitive subgroupoid K such that the pair (G, K) has a tubular structure as in Definition 5.1 (e.g. when the isotropy Lie groups of G have finitely many connected components). Moreover in some cases the tubular structure comes from a Cartan decomposition as in Definition 5.2 (e.g. when the isotropy Lie groups of G are non-compact semisimple). In this section we put together some results from the proper and transitive cases, and follow the notation of Section 4.

Assume that K is a proper wide and transitive Lie subgroupoid of G and fix $z \in M$. Recall from [10, Appendix 10.4] that any measure μ_M on M determines a Haar system μ by the formula

$$\int_{\mathsf{t}^{-1}(m)} f(a)\mu_m(a) = \int_M \left(\int_{K_m} f(ka)\mathsf{d}k\right)\mu_M;$$

here dk is the normalized invariant Haar measure on the compact Lie group K_m and $a \in t^{-1}(m) \mapsto \int_{K_m} f(ka) dk$ is regarded as a function on $M = t^{-1}(m)/K_m$ as it is K_m -invariant. In particular we may take μ_M to be the delta-distribution at $z \in M$. The resulting properly supported normalized Haar system becomes

(49)
$$\int_{\mathsf{t}^{-1}(m)} f(a)\mu_m(a) = \int_{K_m} f(kh)\mathsf{d}k$$

where $h \in K$ is any arrow with s(h) = z and t(h) = m.

Lemma 5.14. Suppose K is a proper and transitive subgroupoid of G. The following diagram commutes

$$C^{p}(G) \xrightarrow{\operatorname{Av}} C^{p}(G)^{K}$$

$$\downarrow r$$

$$C^{p}(G_{z}) \xrightarrow{\operatorname{Av}} C^{p}(G_{z})^{K_{z}}$$

Here r denotes the restriction map, the bottom averaging operator is associated to the normalized invariant Haar measure dk on K_z and the upper averaging operator to the Haar system defined by (49). *Proof.* For $f \in C^{\infty}(B_pG)$ and $(g_1, \ldots, g_p) \in B_pG_z$, we have that $r \circ \operatorname{Av}^{(i)}(f)(g_1, \ldots, g_p) = \operatorname{Av}^{(i)}(f)(g_1, \ldots, g_p)$, where $\operatorname{Av}^{(i)}$ is the averaging operator w.r.t the *i*-th K-action induced by the Haar system (49). For 1 < i < p,

$$Av^{(i)}(f)(g_1, \dots, g_p) = \int_{t^{-1}(z) \cap K} f(g_1, \dots, g_i a, a^{-1}g_{i+1}, \dots, g_p) \mu_z(a)$$
$$= \int_{K_z} f(g_1, \dots, g_i(kh), (kh)^{-1}g_{i+1}, \dots, g_p) dk$$

with $h \in K$ any arrow with s(h) = t(h) = z. Letting h = z, the above equation becomes

$$\int_{K_z} f(g_1, \dots, g_i k, k^{-1} g_{i+1}, \dots, g_p) \mathsf{d}k = \operatorname{Av}^{(i)}(r(f))(g_1, \dots, g_p)$$

where now $\operatorname{Av}^{(i)}$ is the one induced by dk. For $i = 1, \ldots, p$, we again obtain that $r \circ \operatorname{Av}^{(i)}(f) = \operatorname{Av}^{(i)} \circ r(f)$. The conclusion now follows by consecutively applying the *i*-th averaging operators so as to obtain that $r \circ \operatorname{Av}(f) = \operatorname{Av} \circ r(f)$ (see (11)).

When dealing with pairs (G, K) that admit a tubular structure ϕ (or a Cartan decomposition), the restriction ϕ_z to their isotropies at z is also a tubular structure (respectively a Cartan decomposition) of (G_z, K_z) . These are the tubular structures considered in the following result:

Lemma 5.15. Suppose that K is a proper wide and transitive subgroupoid of G such that (G, K) has a Cartan decomposition. Let A be the Lie algebroid of G and \mathfrak{g}_z its isotropy Lie algebra at z. Then the following diagrams commute

$$C^{p}(A)_{K-\text{basic}} \xrightarrow{R_{G/K}} \widetilde{C}^{p}(G)^{K} \qquad C^{p}(A)_{K-\text{basic}} \xrightarrow{R_{G/K}} \widetilde{C}^{p}(G)^{K}$$

$$\downarrow r \qquad \qquad \uparrow \text{ext} \qquad \uparrow \text{ext}$$

$$C^{p}(\mathfrak{g}_{z})_{K_{z}-\text{basic}} \xrightarrow{R_{G_{z}/K_{z}}} \widetilde{C}^{p}(G_{z})^{K_{z}} \qquad C^{p}(\mathfrak{g}_{z})_{K_{z}-\text{basic}} \xrightarrow{R_{G_{z}/K_{z}}} \widetilde{C}^{p}(G_{z})^{K_{z}}$$

Proof. The left diagram clearly commutes. The right diagram commutes because r and ext are inverse to each other, and the commutativity of the left diagram.

Passing to the van Est differentiation maps, we have the following general result for transitive Lie subgroupoids, which proof is analogous to that of Lemma 5.15.

Corollary 5.16. Let K be a transitive, wide Lie subroupoid of G and let $z \in M$. Then the diagrams

commute (an analogous result holds for the normalized invariant subcomplexes). Here $G_z \subseteq G$, and $K_z \subseteq K$ are the isotropy Lie groups at z, $\mathfrak{g} = \ker(\mathfrak{a})$, and the upper and lower ext are given by equations (21) and (24), respectively. For the following result we consider the Haar system (49) on K, and the invariant Haar measure dk on K_x . We also consider the tubular structure of (G, K) given by the Cartan decomposition, and its induced tubular structure on (G_z, K_z) .

Theorem 5.17. Let K be a proper wide and transitive subgroupoid of G such that (G, K) admits a Cartan decomposition. Let A be the Lie algebroid of G and \mathfrak{g}_z its isotropy Lie algebra at z. Then the following is a commutative diagram of homotopy equivalences

In particular,

$$r \circ VE_G/K \colon \mathsf{C}^{\bullet}(G) \to \mathsf{C}^{\bullet}(\mathfrak{g}_z)_{K_z - \text{basic}} \text{ and } \operatorname{ext} \circ R_{G_z/K_x} \colon \mathsf{C}^{\bullet}(\mathfrak{g}_z)_{K_z - \text{basic}} \to \mathsf{C}^{\bullet}(G)$$

are homotopy equivalences with $(r \circ VE_G/K) \circ (\text{ext} \circ R_{G_z/K_x}) = \mathrm{id}_{\mathsf{C}^{\bullet}(g_z)_{K_z-\mathrm{basic}}}$

Proof. First, it is known that $r: C^{\bullet}(G) \to C^{\bullet}(G_z)$ is a homotopy equivalence for transitive groupoids (see e.g. [6]). That the rest of the maps involved in the diagram and R_{G_z/K_x} are homotopy equivalences is a consequence of Theorems 4.1, 4.3 and 5.12. The commutativity of the left diagram is Corollary 5.16

Also, by the commutativity of the diagram above, of the one on the right in Lemma 5.15, and Theorem 5.12,

$$(r \circ VE_G/K) \circ (\text{ext} \circ R_{G_z/K_x}) = (VE_{G_z/K_z} \circ r) \circ (\text{ext} \circ R_{G_z/K_x})$$
$$= VE_{G_z/K_z} \circ R_{G_z/K_x} = \text{id}_{\mathsf{C}^{\bullet}(g_z)_{K_z}-\text{basic}}.$$

To conclude the proof, note that $\operatorname{ext} \circ R_{G_z/K_x} \colon \mathsf{C}^{\bullet}(g_z)_{K_z-\operatorname{basic}} \to \mathsf{C}^{\bullet}(G)$ can be written as the composition

$$\mathsf{C}^{\bullet}(g_z)_{K_z-\text{basic}} \stackrel{R_{G_z/K_x}}{\longrightarrow} \widetilde{\mathsf{C}}^{\bullet}(G_z)^{K_z} \stackrel{\text{ext}}{\longrightarrow} \widetilde{\mathsf{C}}^{\bullet}(G)^K \stackrel{\iota}{\longrightarrow} \mathsf{C}^{\bullet}(G)^K,$$

where all the maps are homotopy equivalences (ι by Theorem 2.4, and ext by Theorem 4.1). \Box

5.6. Example: The polar decomposition and the cohomology of $GL_n(\mathbb{C})$. Let $GL_n = GL_n(\mathbb{C})$ and let \mathfrak{gl}_n be its Lie algebra. We will explicitly find generators of $H(GL_n)$ at the cochain level, described in terms of cochains in $\mathsf{C}^{\bullet}(\mathfrak{gl}_n)_{U(n)-\text{basic}}$, where $U(n) = \{B \in GL_n \mid BB^* = I\}$. We will also describe the van Est map $VE_{GL_n} \colon \mathsf{C}^{\bullet}(GL_n) \to \mathsf{C}^{\bullet}(\mathfrak{gl}_n)$ in terms of the cochain generators of the respective cohomologies.

In the first part we explicitly find the generators of $H(\mathfrak{gl}_n, U(n))$ at the cochain level. Then we follow the strategy of Remark 5.13 with $G = GL_n$ and K = U(n), to find the generators of $H(GL_n)$ as the image of the generators of $H(\mathfrak{gl}_n, U(n))$ via the Est integration map $R_{GL_n/U(n)} \colon \mathsf{C}^{\bullet}(\mathfrak{gl}_n)_{U(n)-\text{basic}} \to \mathsf{C}^{\bullet}(GL_n).$

First of all recall from [15] that

$$H(\mathfrak{gl}_n) = \wedge (u'_1, u'_3, \dots, u'_{2n-1}, b_1, b_3, \dots, b_{2n-1}),$$

where the u's and b's are given (at the cochain level) by $u_q = \operatorname{Re}(\Phi_{2q-1})$, $b_q = \operatorname{Im}(\Phi_{2q-1})$ for q odd and $u'_q = \operatorname{Im}(\Phi_{2q-1})$, $b_q = \operatorname{Re}(\Phi_{2q-1})$ for q even, with

$$\Phi_{2q-1}(A_1, \dots, A_{2q-1}) = \sum_{\gamma \in S_{2p-1}} \operatorname{sgn}(\gamma) \operatorname{tr}(A_{\gamma(1)} \cdots A_{\gamma(2q-1)})$$

being the Ad–invariant skew-symmetric multilinear maps on $\mathfrak{gl}_n.$ In turn,

$$H(\mathfrak{gl}_n, U(n)) = (\wedge^{\bullet} \mathfrak{p}^*)_{U(n)-\mathrm{inv}} = \wedge (u_1, u_3, \dots, u_{2n-1}),$$

where the u's are given (at the cochain level) by

(50)
$$u_{2q-1} = i^{q-1} \Phi_{2q-1} \in C^{2q-1}(\mathfrak{gl}_n)_{U(n)-\text{basic}}$$

Note that the inclusion map

$$\iota \colon H(\mathfrak{gl}_n, U(n)) \to H(\mathfrak{gl}_n)$$

sends u_{2q-1} to u'_{2q-1} and it is injective. As a consequence, we obtain the following result.

Corollary 5.18. The van Est map $VE_{GL_n} : H(GL_n) \to H(\mathfrak{gl}_n)$ is injective.

Proof. This is a direct consequence of the commutativity of the diagram

$$\widetilde{\mathsf{C}}^{\bullet}(GL_n)^{U(n)} \xrightarrow{\iota} \mathsf{C}^{\bullet}(GL_n)$$

$$VE_{GL_n} \downarrow \qquad \qquad \downarrow VE_{GL_n}$$

$$\mathsf{C}^{\bullet}(\mathfrak{gl}_n)_{U(n)-\text{basic}} \xrightarrow{\iota} \mathsf{C}^{\bullet}(\mathfrak{gl}_n)$$

and the fact the lower map is injective in cohomology, the upper map is a homotopy equivalence by Theorem 2.5 and the left map as well by Theorem 5.12. $\hfill \Box$

For the second step, we use the *polar decomposition* of GL_n (see e.g. [16]). It comes from the polar decomposition of \mathfrak{gl}_n as

$$\mathfrak{gl}_n=\mathfrak{u}(n)\oplus\mathfrak{p},\quad A=\frac{1}{2}(A-A^*)+\frac{1}{2}(A+A^*),\ A\in\mathfrak{gl}_n.$$

with $\mathfrak{u}(n) = \{A \in \mathfrak{gl}_n \mid A^* = -A\}$ -the Lie algebra of U(n)-, while $\mathfrak{p} = \{A \in \mathfrak{gl}_n \mid A = A^*\}$. Note that $\mathfrak{u}(n)$ and \mathfrak{p} correspond to the eigenspace 1 and -1, respectively, of the Cartan involution $\theta(X) = -X^*$. The polar decomposition of GL_n is then defined as the U(n)-equivariant diffeomorphism

$$\mathfrak{p} \times U(n) \to GL_n, \quad (X,U) \mapsto e^X U,$$

(note the similarity with (29) in the case of a non-compact semisimple Lie group G), thus it is a Cartan decomposition for the pair $(GL_n, U(n))$. Applying Theorem 5.12 we obtain the following description of $H(GL_n)$.

Theorem 5.19. The van Est integration map (45)

$$R_{GL_n/U(n)} \colon \mathsf{C}^{\bullet}(\mathfrak{gl}_n)_{U(n)-\text{basic}} \to \mathsf{C}^{\bullet}(GL_n)$$

is a homotopy equivalence and it induces an algebra isomorphism in cohomology. In particular,

$$H(GL_n) = \wedge (v_1, \dots, v_{2n-1})$$

where $v_{2p-1} := R_{GL_n/U(n)}(u_{2p-1})$ has degree 2p-1 and is a normalized U(n)-invariant cochain (in the sense of Remark 2.2).

Finally, with the previous result at hand, we obtain the following description of the degree one generator $v_1 \in C^1(GL_n)$:

Corollary 5.20. The first cohomology class $v_1 = R_{GL_n/U(n)}(u_1) \in H^1(GL_n)$ is explicitly given at the cochain level by

$$v_1(A) = \operatorname{tr}(X) \in \mathbb{R}, \quad A \in GL_{\mathfrak{s}}$$

where $A = e^X U$ ($X \in \mathfrak{p}$) is the polar decomposition of the invertible matrix A.

Proof. First, we identify the quotient $GL_n/U(n)$ with the space of positive-definite Hermitian matrices $P = \text{Im}(e_{|\mathfrak{p}}:\mathfrak{p}\to GL_n)$. The scalar multiplication of P is inherited by the diffeomorphism given by the restriction of the exponential $e:\mathfrak{p}\to P$:

(51)
$$\lambda_t \colon P \to P, \ e^X \mapsto e^{tX}$$

We apply $R_{GL_n/U(n)}$ as in (48) to $u_1 = \text{tr}|_{\mathfrak{p}}$. For this notice that under the identification $C^1(\mathfrak{gl}_n,\mathfrak{u}(n)) \simeq \Omega^1(P)^{GL_n}$ given in (47), the 1-form $\text{tr}_P = \text{tr}_{GL_n/U(n)}$ is written as the composition

$$\operatorname{tr}_P \colon TP \stackrel{\theta}{\longrightarrow} \mathfrak{gl}_n = \mathfrak{p} \oplus \mathfrak{u}(n) \stackrel{\operatorname{tr}|_{\mathfrak{p}}}{\longrightarrow} \mathbb{R}$$

(all diagonal entries of a Hermitian matrix are real), where θ is the left invariant MC-form restricted to TP. Now, the map $\gamma_A \colon [0,1] \to P$, $A \in GL_n$ appearing in $R_{GL_n/U(n)}$ is just

$$\gamma_A(t) = \lambda_t(AU(n)) = \lambda_t(e^X UU(n)) = \lambda_t(e^X U(n)) = e^{tX}$$

where $A = e^{X}U$ is the polar decomposition of A. We have that

$$\gamma_A^*(\operatorname{tr}_P)\left(\frac{d}{dt}\right) = \operatorname{tr}_P \circ \gamma_{A,*}\left(\frac{d}{dt}\right) = \operatorname{tr}_P\left(\frac{d}{dt}e^{tX}\right) = \operatorname{tr}_P(L_{e^{tX}}(X)) = \operatorname{tr}(X),$$

where the third equality follows by using that $t \mapsto e^t X \in GL_n$ is an integral curve of the left invariant vector field determined by $X \in \mathfrak{p} \subseteq \mathfrak{gl}_n$, and the last equality by left-invariance of tr_P . Thus

$$R_{GL_n/U(n)}(u_1)(A) = \int_{[0,1]} \gamma_A^*(\operatorname{tr}_P)\left(\frac{d}{dt}\right) = \int_{[0,1]} \operatorname{tr}(X) = \operatorname{tr}(X).$$

6. Characteristic classes

In this section we introduce the definition of characteristic classes of a representation of a Lie groupoid, and relate them with the existing notion of characteristic classes of a representation of a Lie algebroid (see [6, 7, 12]). This is the first step to study characteristic classes of Lie groupoids endowed with a geometric structure, e.g. symplectic, symplectic-Nijenhuis, presymplectic, Pfaffian and contact groupoids. We also hope that the approach using the groupoid will bring a new insight into existing characteristic classes of Poisson manifolds (the modular class), foliations (Bott's characteristic classes), and other type of geometries encoded by Lie algebroids.

Throughout we deal with cohomologies with real coefficients.

6.1. Background on representations. A representation of a Lie groupoid $G \Rightarrow M$ is a (left) G-action on a vector bundle $V \to M$ so that for any $g \in G$ with $\mathbf{s}(g) = m, \mathbf{t}(g) = m'$, the induced map $V_m \to V_m$, $e \mapsto ge$ is a linear isomorphism. Alternatively, a representation can be encoded as follows: associated to any vector bundle $V \to M$, we consider the gauge groupoid $GL(V) \Rightarrow M$ of linear isomorphism between the fibers

 $GL(V) = \{\phi \colon V_m \to V_{m'} \mid m, m' \in M \text{ and } \phi \text{ is a linear isomorphism}\}$

with source and target $s(\phi) = m, t(\phi) = m'$ and multiplication given by composition. A representation of a Lie groupoid $G \rightrightarrows M$ is equivalently described as a Lie groupoid morphism

$$\Phi \colon G \to GL(V)$$

over the identity map of M.

Passing to the infinitesimal picture, a representation of a Lie algebroid $A \to M$ on a vector bundle $V \to M$ is a connection-like operator $\nabla : \Gamma(A) \times \Gamma(V) \to \Gamma(V)$, satisfying the Leibniz rule w.r.t. the anchor map, and so that it is flat. In analogy with the groupoid case a representation can be alternatively described as a Lie algebroid morphism

$$\Psi \colon A \to \mathfrak{gl}(V)$$

over the identity map, where $\mathfrak{gl}(V)$ is the Lie algebroid of GL(V).

If A is the Lie algebroid of G, then any representation $\Phi: G \to GL(V)$ induces a representation of A by

(52) $\Psi := \operatorname{Lie}(\Phi) \colon A \to \mathfrak{gl}(V)$

with $\operatorname{Lie}(\Phi) = d\Phi|_A$.

The following result is used in the next section and may be interesting in its own right.

Corollary 6.1. The van Est map $VE_{GL(V)}: H(GL(V)) \to H(\mathfrak{gl}(V))$ is injective.

Proof. Note that the inclusion $\iota: H(\mathfrak{gl}(V)), U(h)) \to H(\mathfrak{gl}(V))$ is injective. This follows from the injectivity of the inclusion $\iota_z: H(\mathfrak{gl}(V_z)), U(h_z) \to H(\mathfrak{gl}(V_z))$ by Corollary 5.18, the fact that restriction $r: H(\mathfrak{gl}(V), U(h)) \to H(\mathfrak{gl}(V_z)), U(h_z))$ is an isomorphism, and the commutativity of the diagram

$$C(\mathfrak{gl}(V_z))_{U(h_z)-\text{basic}} \xrightarrow{\iota_z} C(\mathfrak{gl}(V_z))$$

$$\uparrow^r \qquad \uparrow^r$$

$$C(\mathfrak{gl}(V))_{U(h)-\text{basic}} \xrightarrow{\iota} C(\mathfrak{gl}(V)).$$

Now, in the left commutative square of (59), the left arrow and $\iota_{GL(V)}$ are homotopy equivalences. Hence, $VE_{GL(V)}: H(GL(V)) \to H(\mathfrak{gl}(V))$ is injective.

6.2. Characteristic classes for algebroids revisited. In this section we focus on an alternative, simple and direct definition of the characteristic classes of representations of Lie algebroids first defined in [6] (see also [7, 12]). Our approach relies on the notion of relative cohomology and its properties. Let A be a Lie algebroid over M, V a *complex* vector bundle over M of complex rank n, and a representation of A on V:

$$\Psi \colon A \to \mathfrak{gl}(V).$$

Fix a (fiberwise) Hermitian metric h of V and denote by $U(h) \subseteq GL(V)$ the transitive and proper Lie subgroupoid

$$U(h) = \{\phi \colon V_m \to V_{m'} \in GL(V) \mid \phi \colon (V_m, h_m) \to (V_{m'}, h_{m'}) \text{ is an isometry} \}$$

Fix a point $z \in M$ and an isometry

(53)
$$F: (\mathbb{C}^n, h_0) \to (V_z, h_z)$$

where h_0 is the standard Hermitian metric of \mathbb{C}^n . Note that conjugation by F

$$C_F \colon GL(V_z) \to GL_n, \ C_F(\phi) := F^{-1} \circ \phi \circ F$$

is a Lie group isomorphism sending the Lie subgroup $U(h_z)$ onto the unitary group U(n). Consider the following composition of cochain maps

(54)
$$M_{\Psi} \colon \mathsf{C}^{\bullet}(\mathfrak{gl}_n)_{U(n)-\text{basic}} \xrightarrow{C_F^*} \mathsf{C}^{\bullet}(\mathfrak{gl}(V_z))_{U(h_z)-\text{basic}} \xrightarrow{\operatorname{ext}} \mathsf{C}^{\bullet}(\mathfrak{gl}(V))_{U(h)-\text{basic}} \xrightarrow{\Psi^*} \mathsf{C}^{\bullet}(A)$$

Recall from Section 5.6 that

$$H(\mathfrak{gl}_n, U(n)) = \wedge (u_1, u_3, \dots, u_{2n-1})$$

where the u's are explicitly defined at the cochain level by trace operators (50).

Definition 6.2. The *characteristic classes* of the representation of the Lie algebroid A on the complex vector bundle V are defined as

$$u_{2p-1}(V) := [M_{\Psi}(u_{2p-1})] \in H^{2p-1}(A), \quad 1 \le p \le n,$$

where n is the rank of V. If V is a *real* vector bundle then they are defined as $u_{2p-1}(V) := u_{2p-1}(V_{\mathbb{C}})$, where $V_{\mathbb{C}}$ is the complexification of V.

Remark 6.3. This definition should be compared with the explicit approach in [6]. By inspection, definition 6.2 agrees with [6, formula (45)] in local coordinates.

Lemma 6.4. The characteristic classes of the representation Ψ do not depend on the choice of $z \in M$, F as in (53), or the Hermitian metric h on V.

To prove Lemma 6.4 we will need the following result:

Lemma 6.5. Fix $z \in M$ and F as in (53), then

• for any other point $m \in M$ and isometry $F' : (\mathbb{C}^n, h_0) \to (V_m, h_m)$,

$$\operatorname{ext}_z \circ C_F^* = \operatorname{ext}_m \circ C_{F'}^*$$

• for any other Hermitian metric h' of V, and any isometry $H: (V,h) \to (V,h')$ fixing M,

$$C_H^* \colon \mathsf{C}(\mathfrak{gl}(V)) \to \mathsf{C}(\mathfrak{gl}(V))$$

is homotopic to the identity.

Proof of Lemma 6.4. By the first item of Lemma 6.5 we already know that they do not depend on the point, nor the isometry (53), so fix $z \in M$ and the isometry F as in (53). Denote the inclusions by

 $\iota_h \colon \mathsf{C}(\mathfrak{gl}(V))_{U(h)-\mathrm{basic}} \to \mathsf{C}(\mathfrak{gl}(V)), \ \ \iota_{h'} \colon \mathsf{C}(\mathfrak{gl}(V))_{U(h')-\mathrm{basic}} \to \mathsf{C}(\mathfrak{gl}(V))$

Consider the diagram

$$C(\mathfrak{gl}(V_{z}))_{U(h_{z})-\text{basic}} \xrightarrow{\text{ext}} C(\mathfrak{gl}(V))_{U(h)-\text{basic}} \xrightarrow{\iota_{h}} C(\mathfrak{gl}(V))$$

$$C(\mathfrak{gl}_{n})_{U(n)-\text{basic}} \xrightarrow{C_{F}^{*}} C_{H_{z}}^{*} \xrightarrow{C_{H_{z}}^{*}} C_{H_{z}}^{*} \xrightarrow{C_{H}^{*}} C_{H}^{*} \xrightarrow{C_{H}^{*}} C(\mathfrak{gl}(V))_{U(h')-\text{basic}} \xrightarrow{\iota_{h'}} C(\mathfrak{gl}(V))$$

with $F' := H_z \circ F : (\mathbb{C}^n, h_0) \to (V_z, h'_z)$. Then the left triangle is clearly commutative, and also the right square is commutative. The middle square is also commutative as $C^*_{H_z} \circ r = r \circ C^*_H$ where r are the restriction maps, and the fact that r and ext are inverse to each other.

Denote by $u_{2p-1}(V,h)$ ($u_{2p-1}(V,h')$ respectively) the characteristic classes of Definition 6.2 defined by the composition (54) using the Hermitian metric h (h' respectively). We compute:

$$u_{2p-1}(V,h) = \Psi^*[\iota_h \circ \text{ext} \circ C_F^*(u_{2p-1})] = \Psi^*[C_H^* \circ \iota_h \circ \text{ext} \circ C_F^*(u_{2p-1})]$$
$$= \Psi^*[\iota_{h'} \circ \text{ext} \circ C_{F'}^*(u_{2p-1})] = [u_{2p-1}(V,h')]$$

where in the second equality we used the second item of Lemma 6.5

Proof of Lemma 6.5. For the first item, note that, since the restriction map

 $r_m \colon \mathsf{C}(\mathfrak{gl}(V))_{U(h)-\mathrm{basic}} \to \mathsf{C}(\mathfrak{gl}(V_m))_{U(h_m)-\mathrm{basic}}$

is inverse to the extension map ext_m , the equation to prove holds if and only if

$$C_{F'^{-1}}^* \circ r_m \circ \operatorname{ext}_z \circ C_F^* = \operatorname{id}_{\mathsf{C}^{\bullet}(\mathfrak{gl}_n)_{U(n)-\operatorname{basic}}}.$$

To check this last equation, note that we can write $r_m \circ \text{ext}_z$ as the pullback by conjugation

$$r_m \circ \operatorname{ext}_z = C^*_{b(m)},$$

where $b: M \to U(h)$ is any bisection with the property that s(b(m)) = z (i.e. $b(m): (V_z, h_z) \to (V_m, h_m)$ is an isometry). Therefore,

$$C^*_{F'^{-1}} \circ r_m \circ \text{ext}_z \circ C^*_F = C^*_{F'^{-1}} \circ C^*_{b(m)} \circ C^*_F = C^*_{F'^{-1} \circ b(m) \circ F}.$$

As $U := F'^{-1} \circ b(m) \circ F$ is the composition of isometries

$$(\mathbb{C}^n, h_0) \xrightarrow{F} (V_z, h_z) \xrightarrow{b(m)} (V_m, h_m) \xrightarrow{F'^{-1}} (\mathbb{C}^n, h_0),$$

 $C_U: GL_n \to GL_n$ is conjugation by the unitary map $U \in U(n), C_U^*: \mathfrak{gl}_n^* \to \mathfrak{gl}_n^*$ is the coadjoint action by U, and hence $C_U^*: \mathsf{C}(\mathfrak{gl}_n)_{U(n)-\text{basic}} \to \mathsf{C}(\mathfrak{gl}_n)_{U(n)-\text{basic}}$ is the identity map on the U(n)-basic subcomplex, as we desired.

30

For the second item, we will construct a homotopy h between the identity map and C_H^* as follows. Denote by $GL(V)_M$ the space of linear automorphisms of V fixing the base M. Let $t \mapsto H_t \in GL(V)_M$ be a smooth path connecting the identity and H:

$$H_t \colon V \xrightarrow{\simeq} V, \quad H_0 = \mathrm{id}_V, \quad H_1 = H.$$

Observe that $GL(V)_M \subseteq GL(V)$ is the Lie groupoid given by the union of the isotropy Lie groups of GL(V), and that $H_t \in \text{Bis}(GL(V)_M)$.

Infinitesimally, the Lie algebroid $\operatorname{Lie}(GL(V)_M) = \mathfrak{gl}(V)_M$ –the space of linear transformations from V into V fixing the base M– is a Lie subalgebroid of $\mathfrak{gl}(V)$ given by the kernel of the anchor map, and H_t induces a path of sections $t \mapsto X_t \in \mathfrak{gl}(V)_M$ defined by

$$\frac{d}{dt}H_t = X_t \circ H_t$$

We can then have the induced Lie derivatives $\mathcal{L}_{X_t} = [\mathsf{d}_{CE}, \iota_{X_t}]$ along $X_t, \mathcal{L}_{X_t} \colon \mathsf{C}^{\bullet}(\mathfrak{gl}(V)) \to \mathsf{C}^{\bullet}(\mathfrak{gl}(V))$, satisfying the property

(55)
$$\frac{d}{dt}C^*_{H_t}(\omega) = \mathcal{L}_{X_t}(C^*_{H_t}(\omega)), \quad \omega \in \mathsf{C}^{\bullet}(\mathfrak{gl}(V)).$$

Define $h: C^{\bullet}(\mathfrak{gl}(V)) \to C^{\bullet-1}(\mathfrak{gl}(V))$ by

$$\mathsf{h}(\omega) := \int_0^1 \iota_{X_t} \big(C^*_{H_t}(\omega) \big) \, dt.$$

We compute:

$$C_{H}^{*}(\omega) - \omega = \int_{0}^{1} \frac{d}{dt} C_{H_{t}}^{*}(\omega) dt = \int_{0}^{t} \mathcal{L}_{X_{t}} \left(C_{H_{t}}^{*}(\omega) \right) dt = \int_{0}^{1} \mathsf{d}_{CE} \iota_{X_{t}} \left(C_{H_{t}}^{*}(\omega) \right) + \iota_{X_{t}} \mathsf{d}_{CE} \left(C_{H_{t}}^{*}(\omega) \right) dt = \int_{0}^{1} \mathsf{d}_{CE} \iota_{X_{t}} \left(C_{H_{t}}^{*}(\omega) \right) + \iota_{X_{t}} \mathsf{d}_{CE} \left(C_{H_{t}}^{*}(\omega) \right) dt$$
$$= \mathsf{d}_{CE} \mathsf{h}(\omega) + \mathsf{hd}_{CE}(\omega),$$

where in the last equality we used that d_{CE} commutes with $C^*_{H_t}$ since $C_{H_t}: GL(V) \to GL(V)$ is a groupoid morphism for each t.

6.3. Characteristic classes for Lie groupoids. Let G be a Lie groupoid over M, V a *complex* vector bundle over M of complex rank n, and a representation of G on V:

$$\Phi \colon G \to GL(V).$$

Fix a Hermitian metric h of V, a point $z \in M$ and an isometry $F: (\mathbb{C}^n, h_0) \to (V_z, h_z)$. Using the same conventions of Section 6.2, consider the following composition of cochain maps

(56)
$$M_{\Phi}: \widetilde{\mathsf{C}}^{\bullet}(GL_n)^{U(n)} \xrightarrow{C_F^*} \widetilde{\mathsf{C}}^{\bullet}(GL(V_z))^{U(h_z)} \xrightarrow{\operatorname{ext}} \widetilde{\mathsf{C}}^{\bullet}(GL(V))^{U(h)} \xrightarrow{\Phi^*} \mathsf{C}^{\bullet}(G).$$

From Section 5.6 we know that

(57)
$$H(GL_n) = \wedge (v_1, \dots, v_{2n-1})$$

where $v_{2p-1} := R_{GL_n/U(n)}(u_{2p-1})$ is the cochain of $\widetilde{\mathsf{C}}^{\bullet}(GL_n)^{U(n)}$ given in Theorem 5.19.

Definition 6.6. The *characteristic classes* of the representation Φ of the Lie groupoid G on the complex vector bundle V are defined as

$$v_{2p-1}(V) := M_{\Phi}(v_{2p-1}) \in H^{2p-1}(G), \quad 1 \le p \le n,$$

where n is the rank of V. If V is a *real* representation then they are defined as $v_{2p-1}(V) := v_{2p-1}(V_{\mathbb{C}})$, where $V_{\mathbb{C}}$ is the complexification of V.

Lemma 6.7. The characteristic classes of the representation Φ do not depend on the choice of $z \in M$, F as in (53), nor the Hermitian metric h of V.

Before proving Lemma 6.7, let us point out the relation between the characteristic classes of a representation of G, and those of the induced representation on its algebroid:

Proposition 6.8. Let $\Psi: A \to \mathfrak{gl}(V)$ be the induced representation (52) of $\Phi: G \to GL(V)$ on the Lie algebroid A of G. Then the characteristic classes of Ψ are the image of the characteristic classes of Φ via the van Est map $VE_G: H(G) \to H(A)$:

$$VE_G(v_{2p-1}(V)) = u_{2p-1}(V), \quad 1 \le p \le n.$$

Proof. This follows by the fact that $VE_{GL_n}(v_{2p-1}) = u_{2p-1}$, $1 \leq p \leq n$ (as $VE_{GL_n} \circ R_{GL_n/U(n)} = \operatorname{id}_{\mathsf{C}^{\bullet}(\mathfrak{gl}_n)_{U(n)-\operatorname{basic}}}$) and the commutativity of the diagram

$$(58) \qquad \widetilde{\mathsf{C}}(GL_{n})^{U(n)} \xrightarrow{C_{F}^{*}} \widetilde{\mathsf{C}}(GL(V_{z}))^{U(h_{z})} \xrightarrow{\operatorname{ext}} \widetilde{\mathsf{C}}(GL(V))^{U(h)} \xrightarrow{\Phi^{*}} \mathsf{C}(G)$$

$$\downarrow^{VE_{GL_{n}}} \qquad \qquad \downarrow^{VE_{GL(V_{z})}} \qquad \qquad \downarrow^{VE_{GL(V)}} \qquad \qquad \downarrow^{VE_{GL(V)}} \qquad \qquad \downarrow^{VE_{G}}$$

$$\mathsf{C}(\mathfrak{gl}_{n})_{U(n)-\operatorname{basic}} \xrightarrow{C_{F}^{*}} \mathsf{C}(\mathfrak{gl}(V_{z}))_{U(h_{z})-\operatorname{basic}} \xrightarrow{\operatorname{ext}} \mathsf{C}(\mathfrak{gl}(V))_{U(h)-\operatorname{basic}} \xrightarrow{\Psi^{*}} \mathsf{C}(A)$$

where the middle square commutes by Corollary 5.18.

Proof of Lemma 6.7. The right square of the diagram (58) factors through the inclusions as

(59)
$$\widetilde{\mathsf{C}}(GL(V))^{U(h)} \xrightarrow{\iota_{GL(V)}} \mathsf{C}(GL(V)) \xrightarrow{\Phi^*} \mathsf{C}(G)$$

$$\downarrow^{VE_{GL(V)}} \qquad \qquad \downarrow^{VE_{GL(V)}} \qquad \qquad \downarrow^{VE_{GL(V)}} \qquad \qquad \downarrow^{VE_{GL(V)}} \qquad \qquad \downarrow^{VE_{GL(V)}} \mathsf{C}(\mathfrak{gl}(V))_{U(h)-\text{basic}} \xrightarrow{\iota} \mathsf{C}(\mathfrak{gl}(V)) \xrightarrow{\Psi^*} \mathsf{C}(A).$$

To see that the classes do not depend on the choices, recall that $VE_{GL(V)}: H(GL(V)) \rightarrow H(\mathfrak{gl}(V))$ is injective by Corollary 6.1. In this case, the result follows at once by this fact, (59) and Lemma 6.4.

The main properties of the characteristic classes are:

Proposition 6.9. For any representations V, W of G, the characteristic classes satisfy the following properties:

(a) $v_{2p-1}(V \oplus W) = v_{2p-1}(V) + v_{2p-1}(W).$ (b) $v_{2p-1}(V \otimes W) = v_{2p-1}(V) \operatorname{rank}(W) + \operatorname{rank}(V)v_{2p-1}(W).$ (c) $v_{2p-1}(V^*) = i^*v_{2p-1}(V), \text{ where } i: G \to G \text{ is the inverse map. In particular,}$ $v_1(V^*) = -v_1(V).$

(d) $v_{2p-1}(V) = 0$ if p is even and V is real.

Proof. Property (c) follows directly from the definition of the action on the dual V^* . The proofs of (a), (b) and (d) are similar, we will show (b) and (d).

For (b), consider the Lie subgroup $GL(V) \otimes GL(W) \subseteq GL(V \otimes W)$ consisting of the tensor product $\phi \otimes \psi \colon V_m \otimes W_m \to V_{m'} \otimes W_{m'}$ of linear isomorphisms $\phi \colon V_m \to V_{m'}, \psi \colon W_m \to W_{m'}$. Similarly, if h, h' are Hermitian metrics for V and W, respectively, the Lie subgroupoid $U(h) \otimes U(h') \subseteq GL(V) \otimes GL(W)$ is proper and transitive and is a subgroupoid of $U(h \otimes h')$.

If $\Phi_V \colon G \to GL(V)$ and $\Phi_W \colon G \to GL(W)$ denote the representations of G, then its induced representation on $V \otimes W$ factors as $\Phi_V \otimes \Phi_W \colon G \to GL(V) \otimes GL(W)$, and

commutes, where $\iota: GL(V) \otimes GL(W) \to GL(V \otimes W)$ is the inclusion. Denoting by $n = \operatorname{rank}(V), m = \operatorname{rank}(W)$, it is then a straightforward check to see that the composition

$$M_{\Phi_V^* \otimes \Phi_W^*} \colon \widetilde{\mathsf{C}}(GL_{nm})^{U(nm)} \to \mathsf{C}(G)$$

from (56) factors as

$$\widetilde{\mathsf{C}}(GL_{nm})^{U(nm)} \xrightarrow{\iota^*} \widetilde{\mathsf{C}}(GL_n \otimes GL_m)^{U(n) \otimes U(m)} \xrightarrow{C_F^* \otimes C_F^*} \widetilde{\mathsf{C}}(GL(V_z) \otimes GL(W)^{U(h_z) \otimes U(h'_z)} \xrightarrow{\operatorname{ext} \otimes \operatorname{ext}} \widetilde{\mathsf{C}}(GL(V) \otimes GL(W))^{U(h) \otimes U(h')} \xrightarrow{\Phi_V^* \otimes \Phi_W^*} \mathsf{C}(G)$$

It is then enough to see that for $v_{2p-1}(nm) \in \widetilde{\mathsf{C}}(GL_{nm})^{U(nm)}$ the generators as in (57)

$$\iota^*(v_{2p-1(nm)}) = v_{2p-1}(n)m + nv_{2p-1}(m),$$

with $v_{2p-1}(n) \in \widetilde{\mathsf{C}}(GL_n)^{U(n)}, v_{2p-1}(m) \in \widetilde{\mathsf{C}}(GL_m)^{U(m)}$ the generators, accordingly.

Indeed, we use the Kronecker product $A \otimes B \in GL_{nm}$ for matrices $A \in GL_n, B \in GL_m$ which represents the tensor product of the two invertible maps A, B. By general properties of the Kronecker product one can verify that if $[A] = e^X \mod U(n)$ and $[B] = e^Y \mod U(m)$, then $[A \otimes B] = e^X \otimes e^Y \mod U(nm)$ and $\lambda_t(e^X \otimes e^Y) = e^{tX} \otimes e^{tY}$ (λ_t as in (51)). Therefore for $A_1, \ldots, A_q \in GL_n, B_1, \ldots, B_q \in GL_m$

$$\gamma_{t_1,\dots,t_q}^{(q)}(A_1 \otimes B_1,\dots,A_q \otimes B_q) = \gamma_{t_1,\dots,t_q}^{(q)}(A_1,\dots,A_q) \otimes \gamma_{t_1,\dots,t_q}^{(q)}(B_1,\dots,B_q)$$

 γ as in (46), and by bilinearity of \otimes and the fact that

(60)
$$(A \otimes B)(C \otimes D) = AC \otimes BD,$$

it follows that

$$\theta\left(\frac{\partial}{\partial t_i}(\gamma_{t_1,\dots,t_q}^{(q)}(A_1\otimes B_1,\dots,A_q\otimes B_q))\right) = \\\theta\left(\frac{\partial}{\partial t_i}(\gamma_{t_1,\dots,t_q}^{(q)}(A_1,\dots,A_q))\right)\otimes I_m + I_n\otimes\theta\left(\frac{\partial}{\partial t_i}(\gamma_{t_1,\dots,t_q}^{(q)}(A_1,\dots,A_q))\right)$$

with θ the left invariant MC-form of GL_{nm} , GL_n and GL_m , respectively. Again, by property (60) and $\operatorname{tr}(A \otimes B) = \operatorname{tr}(A)\operatorname{tr}(B)$, and recalling that $u_q \in C^q(\mathfrak{gl}_{nm})_{U(nm)-\text{basic}}$, q = 2p - 1 are sums of traces as in (50), we further obtain that

$$\iota^{*}(v_{q}(nm))(A_{1} \otimes B_{1}, \dots, A_{q} \otimes B_{q}) = \int_{[0,1]^{q}} \gamma^{(q)}(A_{1} \otimes B_{1}, \dots, A_{q} \otimes B_{q})^{*}\iota^{*}(u_{q,GL_{nm}/Unm}) = \operatorname{tr}(I_{m}) \int_{[0,1]^{q}} \gamma^{(q)}(A_{1}, \dots, A_{q})^{*}u_{q,GL_{n}/Un} + \operatorname{tr}(I_{n}) \int_{[0,1]^{q}} \gamma^{(q)}(B_{1}, \dots, B_{q})^{*}u_{q,GL_{m}/Um} = mv_{q}(n)(A_{1}, \dots, A_{q}) + nv_{q}(m)(B_{1}, \dots, B_{q}).$$

For (d) notice that the obvious inclusion $\mathfrak{gl}_n(\mathbb{R})\subseteq\mathfrak{gl}_n(\mathbb{C})$ factors through the diagram

and that the restriction to $\mathfrak{gl}_n(\mathbb{R})$ of the generators u_{2q-1} of $H(\mathfrak{gl}_n(\mathbb{C}), U(n))$ vanish for q even (see [15]).

APPENDIX A. VAN EST MAPS USING THE PERTURBATION LEMMA

In this appendix we recall some of the constructions and results of [17] (see also [19]). Let $(D^{\bullet,\bullet}, \mathsf{d}, \delta)$ be a double complex, concentrated in non-negative degrees. Let $(X^{\bullet}, \mathsf{d})$ be a cochain complex. A morphism of double complexes

 $i\colon X\to D$

(where X[•] is regarded as a double complex concentrated in bidegrees $(0, \bullet)$) will be called a *horizontal augmentation map*. Passing to total complexes, i becomes a cochain map from X[•] to the cochain complex (Tot[•](D), d + δ). The Perturbation Lemma, due to Brown [2] and Gugenheim [13], allows us to turn a homotopy operator for the horizontal differential δ into a homotopy operator with respect to the total differential d + δ .

Lemma A.1 (Perturbation Lemma). Suppose $h: D \to D$ is a linear map of bidegree (-1, 0) such that

$$[\mathbf{h}, \delta] = 1 - \mathbf{i} \circ \mathbf{p}$$

for some degree 0 map $\mathbf{p} \colon \mathsf{D}^{0,\bullet} \to \mathsf{X}^{\bullet}$. Put $\mathbf{h}' = \mathbf{h}(1 + \mathbf{d}\mathbf{h})^{-1}$ and $\mathbf{p}' = \mathbf{p}(1 + \mathbf{d}\mathbf{h})^{-1}$. Then
$$[\mathbf{h}', \mathbf{d} + \delta] = 1 - \mathbf{i} \circ \mathbf{p}'.$$

Here, $[\cdot, \cdot]$ denotes the graded commutator, e.g. $[h, \delta] = h\delta + \delta h$.

We shall assume from now on that $p \circ i = id_X$, so that i is injective and $i \circ p$ is a projection onto the image of i. Then also $p' \circ i = id_X$, and $i \circ p'$ is again a projection. In other words, $i: X^{\bullet} \to \text{Tot}^{\bullet}(D)$ is a homotopy equivalence, with p' a homotopy inverse.

In our applications, there is another cochain complex $(\mathbf{Y}^{\bullet}, \delta)$, with a vertical augmentation map

$$j: Y \rightarrow D$$

(thus Y^{\bullet} is regarded as a double complex concentrated in bidegrees $(\bullet, 0)$). The horizontal homotopy h allows us to 'invert' the second cochain map in

$$Y^{\bullet} \stackrel{j}{\longrightarrow} \operatorname{Tot}^{\bullet}(\mathsf{D}) \stackrel{i}{\longleftarrow} X^{\bullet},$$

thereby producing a cochain map $\mathbf{p}' \circ \mathbf{j} = \mathbf{p} \circ (1 + \mathsf{dh})^{-1} \circ \mathbf{j} \colon \mathbf{Y}^{\bullet} \to \mathbf{X}^{\bullet}$. On elements of degree p, this is given by

(61)
$$(-1)^p \mathsf{p} \circ (\mathsf{dh})^p \circ \mathsf{j} \colon \mathsf{Y}^p \to \mathsf{X}^p,$$

Consider now the situation that the vertical differential has a homotopy operator

$$\mathsf{k} \colon \mathsf{D}^{p,q} \to \mathsf{D}^{p,q-1}, \ [\mathsf{d},\mathsf{k}] = 1 - \mathsf{j} \circ \mathsf{q},$$

where $q: D^{0,\bullet} \to Y^{\bullet}$ is a cochain map for d with $q \circ j = id_Y$. Then we can apply the Perturbation Lemma A.1 to this vertical homotopy, and we obtain a cochain map $q \circ (1 + \delta k)^{-1} \circ i: X^{\bullet} \to Y^{\bullet}$ given on degree p elements by

(62)
$$(-1)^p \mathbf{q} \circ (\delta \mathbf{k})^p \circ \mathbf{i} \colon \mathbf{X}^p \to \mathbf{Y}^p$$

The following result will be used to relate van Est 'integration' and 'differentiation' maps.

Lemma A.2. Suppose the homotopy operators h, k satisfy

$$h \circ k = 0, \quad p \circ k = 0.$$

Then (62) followed by (61) is the identity map of X^p .

For our application we start with a Lie groupoid G with Lie algebroid A. The two complexes X = C(A), Y = C(G) are related by a van Est double complex, due to Crainic [6]. Recall from the discussion around (5) that

$$\kappa \colon EG \to BG$$

is a simplicial principal bundle, where for each p, we denote by $\pi_p: E_p G \to M$ the anchor map of the G-action on E_pG . The van Est double complex

(64)
$$(\mathsf{D}^{\bullet,\bullet}(G), \delta, \mathsf{d})$$

is defined as follows.

• The bigraded summands of the double complex are

(65)
$$\mathsf{D}^{p,q}(G) = \Gamma(\pi_n^*(\wedge^q A^*)$$

• δ is the simplicial differential on sections of the simplicial vector bundle

$$\pi^*_{\bullet}(\wedge^q A^*) \to E_{\bullet}G.$$

• $d = (-1)^p d_{CE}$ on elements of bidegree (p,q), is the Chevalley-Eilenberg differential on

).

(66)
$$\mathsf{D}^{p,\bullet}(G) \cong \mathsf{C}^{\bullet}(\pi_p^*A)$$

Specifically, let \mathcal{F} be the foliation of E_pG given by the κ_p -fibers; thus $T_{\mathcal{F}}E_pG$ is the vertical bundle. The isomorphism $\pi_p^*A \cong T_{\mathcal{F}}E_pG$ defines an algebroid structure on π_p^*A .

Furthermore, the double complex comes with horizontal and vertical augmentation maps:

- i: C[•](A) → D^{0,•}(G) is given in degree q by the pullback π₀^{*} (using (66) for p = 0).
 j: C[•](G) → D^{•,0}(G) is given in degree p by the pullback map κ_p^{*} (using (5)).

MARÍA AMELIA SALAZAR

The augmentation maps define cochain maps to the total complex

$$C^{\bullet}(G) \xrightarrow{j} Tot^{\bullet}(D(G)) \xleftarrow{i} C^{\bullet}(A).$$

Differentiation. The van Est double complex has an horizontal homotopy. Consider the maps

$$h_p: E_pG \to E_{p+1}G, \ (g_1, \dots, g_p; g) \mapsto (g_1, \dots, g_p, g; m)$$

where $m = \mathbf{s}(g)$. Since $\pi_{p+1} \circ h_p = \pi_p$, these lift to fiberwise isomorphisms of vector bundles $\pi_p^*(\wedge^q A^*) \to \pi_{p+1}^*(\wedge^q A^*)$, defining a pullback map on sections. The map

$$h\colon \mathsf{D}^{p,q}(G)\to \mathsf{D}^{p-1,q}(G), \quad \psi\mapsto (-1)^p h_{p-1}^*\psi$$

satisfies $[\mathbf{h}, \delta] = 1 - \mathbf{i} \circ \mathbf{p}$, where $\mathbf{p} \colon \mathsf{D}^{0,\bullet}(G) \to \mathsf{C}^{\bullet}(A)$ is the left inverse to $\mathbf{i} = \pi_0^*$ given by pullback under the inclusion $u \colon M \hookrightarrow E_0 G = G$:

$$\mathsf{p} = \mathsf{u}^* \colon \Gamma(\pi_0^*(\wedge^q A^*)) \to \Gamma(\wedge^q A^*).$$

The Perturbation Lemma A.1 gives a new projection $\mathbf{p}' = \mathbf{p} \circ (1 + d\mathbf{h})^{-1}$, which is a cochain map for the total differential $\mathbf{d} + \delta$, with $\mathbf{p}' \circ \mathbf{i} = \mathbf{id}$. Thus, \mathbf{i} is a homotopy equivalence, with \mathbf{p}' a homotopy inverse. As a result from [19], we have the following:

Theorem A.3. The van Est differentiation map (30) is obtained as

$$VE_G = \mathbf{p} \circ (1 + \mathsf{dh})^{-1} \circ \mathbf{j} \colon \mathsf{C}^{\bullet}(G) \to \mathsf{C}^{\bullet}(A).$$

References

- J.-M. Bismut and J. Lott. Flat vector bundles, direct images and higher real analytic torsion. Journal of the American Mathematical Society, 8(2):291–363, 1995.
- [2] R. Brown. The twisted Eilenberg-Zilber theorem. In Simposio di Topologia (Messina, 1964), pages 33–37. Edizioni Oderisi, Gubbio, 1965.
- [3] A. Cabrera, I. Mărcuţ, and M. A. Salazar. On local integration of Lie brackets. J. Reine Angew. Math., 760:267–293, 2020.
- [4] E. Cartan. Sur les invariants intégraux de certains espaces homogènes clos et les propriétés topologiques de ces espaces. Ann. Soc. Polon. Math., 8:181–225, 1930.
- [5] C. Chevalley and S. Eilenberg. Cohomology theory of Lie groups and Lie algebras. Trans. Am. Math. Soc., 63:85–124, 1948.
- [6] M. Crainic. Differentiable and algebroid cohomology, van Est isomorphisms, and characteristic classes. Comment. Math. Helv., 78(4):681–721, 2003.
- [7] M. Crainic and R. L. Fernandes. Secondary characteristic classes of Lie algebroids. In Quantum field theory and noncommutative geometry, volume 662 of Lecture Notes in Phys., pages 157–176. Springer, Berlin, 2005.
- [8] M. Crainic and I. Mărcuţ. A normal form theorem around symplectic leaves. J. Differ. Geom., 92(3):417–461, 2012.
- [9] M. Crainic and I. Mărcuţ. Reeb-Thurston stability for symplectic foliations. Math. Ann., 363(1-2):217-235, 2015.
- [10] M. Crainic and J. a. N. Mestre. Measures on differentiable stacks. J. Noncommut. Geom., 13(4):1235–1294, 2019.
- [11] S. Evens, J.-H. Lu, and A. Weinstein. Transverse measures, the modular class and a cohomology pairing for Lie algebroids. Q. J. Math., Oxf. II. Ser., 50(200):417–436, 1999.
- [12] R. L. Fernandes. Lie algebroids, holonomy and characteristic classes. Adv. Math., 170(1):119–179, 2002.
- [13] V. K. A. M. Gugenheim. On the chain-complex of a fibration. Illinois J. Math., 16:398–414, 1972.
- [14] A. Guichardet. Cohomologie des groupes topologiques et des algèbres de Lie, volume 2 of Textes Mathématiques [Mathematical Texts]. CEDIC, Paris, 1980.

36

- [15] F. Kamber and P. Tondeur. Foliated bundles and characteristic classes. Lecture notes in mathematics. Springer, 1975.
- [16] A. W. Knapp. *Lie groups beyond an introduction*, volume 140 of *Prog. Math.* Boston, MA: Birkhäuser, 2nd ed. edition, 2002.
- [17] D. Li-Bland and E. Meinrenken. On the van Est homomorphism for Lie groupoids. Enseign. Math., 61(1-2):93–137, 2015.
- [18] I. Mărcut and F. Zeiser. The Poisson cohomology of $\mathfrak{sl}_2^*(\mathbb{R})$. J. Symplectic Geom., 21(3):603–652, 2023.
- [19] E. Meinrenken and M. A. Salazar. Van Est differentiation and integration. Math. Ann., 376(3-4):1395–1428, 2020.
- [20] I. Moerdijk and J. Mrčun. On the integrability of Lie subalgebroids. Adv. Math., 204(1):101–115, 2006.
- [21] J. D. Stasheff. Continuous cohomology of groups and classifying spaces. Bull. Am. Math. Soc., 84:513–530, 1978.
- [22] M. Stroppel. Locally compact groups. EMS Textb. Math. Zürich: European Mathematical Society Publishing House, 2006.
- [23] J. L. Tu. La conjecture de Novikov pour les feuilletages hyperboliques. K-Theory, 16(2):129–184, 1999.
- [24] W. T. van Est. Group cohomology and Lie algebra cohomology in Lie groups. I, II. Nederl. Akad. Wetensch. Proc. Ser. A. 56 = Indagationes Math., 15:484–492, 493–504, 1953.
- [25] W. T. van Est. On the algebraic cohomology concepts in Lie groups. I, II. Nederl. Akad. Wetensch. Proc. Ser. A. 58 = Indag. Math., 17:225–233, 286–294, 1955.
- [26] W. T. van Est. Une application d'une méthode de Cartan-Leray. Nederl. Akad. Wetensch. Proc. Ser. A. 58 = Indag. Math., 17:542–544, 1955.
- [27] A. Weinstein and P. Xu. Extensions of symplectic groupoids and quantization. Journal f
 ür die reine und angewandte Mathematik, 417:159–190, 1991.

Instituto de Matematica e Estatistica, Universidade Federal Fluminense (Brazil) Email address: mariasalazar@id.uff.br