

SPIN LINK HOMOLOGY

ELIJAH BODISH, BEN ELIAS, AND DAVID E. V. ROSE

ABSTRACT. We put a new spin on Khovanov–Rozansky homology. That is, we equip Λ^n -colored \mathfrak{sl}_{2n} Khovanov–Rozansky homology with an involution whose ± 1 -eigenspaces are link invariants. When $n = 1, 2, 3$ (and assuming technical conjectures for $n \geq 4$), we prove that this refined invariant categorifies the spin-colored \mathfrak{so}_{2n+1} quantum link polynomial. Along the way, we partially develop the theory of quantum \mathfrak{so}_{2n+1} webs and make contact with ι quantum groups.

CONTENTS

1. Introduction	1
2. Conventions	12
3. Background on quantum groups and link invariants	16
4. Type B Intertwiners	20
5. Background on categorified quantum groups	42
6. Involutions on (part of) the categorified quantum group	54
7. Background on decompositions and equivariant categories	71
8. An equivariant category from the categorified quantum group	81
9. Background on link homology	95
10. Spin link homology	104
Appendix A. Proof of Lemma 4.27	122
Appendix B. Relation to Wenzl’s approach using the q -Clifford algebra	127
Appendix C. More on equivariant categories	129
References	135

1. INTRODUCTION

For each finite-dimensional simple complex Lie algebra \mathfrak{g} , work of Reshetikhin–Turaev [58] defines a Laurent polynomial-valued invariant of a link $\mathcal{L} \subset S^3$ with components colored by finite-dimensional representations of the corresponding quantum group $U_q(\mathfrak{g})$. In the case that $\mathfrak{g} = \mathfrak{sl}_2$ and the link is colored by the defining representation, this invariant is the much-celebrated Jones polynomial [29]. In pioneering work [32], Khovanov showed that the Jones polynomial admits a *categorification* taking the form of a bigraded homological link invariant $H(\mathcal{L})$ from which the Jones polynomial can be recovered by taking the Euler characteristic. Subsequent works of Khovanov–Rozansky [36] (and others [46, 12, 49]) construct analogous *link homology theories* in the $\mathfrak{g} = \mathfrak{sl}_N$ case. Khovanov–Rozansky homology is defined in terms of explicit chain complexes, and a number of subsequent formulations make it even more accessible and computable (e.g. the cobordism and foam formulations given in [4, 53, 19]). These explicit descriptions underlie many applications of Khovanov(–Rozansky) homology in low-dimensional topology (e.g. [56, 51, 57]).

In his ICM address [33], Khovanov posed the “difficult open problem” of categorifying the “polynomial invariants of links associated to arbitrary \mathfrak{g} and their irreducible representations. The first solution

to this problem was obtained by Webster in [73]. Therein, he defines algebras that categorify tensor product representations of $U_q(\mathfrak{g})$, and constructs derived functors corresponding to tangles. These derived functors categorify the Reshetikhin–Turaev invariants of tangles, hence, in particular, the $U_q(\mathfrak{g})$ Reshetikhin–Turaev link invariants. However, Webster’s categorical invariants are famously difficult to compute: for example, we are unaware of anyone having computed the Webster homology of the unknot outside of type A . Lack of computability has severely hindered the development and application of these link homologies.

In the present paper we provide a new construction for link homology (not a priori related to Webster’s) in the case that $\mathfrak{g} = \mathfrak{so}_{2n+1}$ and the link is colored by the *spin representation*. Our construction is based on a new “folded categorical skew Howe duality,” a surprising connection between spin-colored \mathfrak{so}_{2n+1} link invariants and Λ^n -colored \mathfrak{sl}_{2n} link invariants that is only visible at the categorified level. Along the way, we study endomorphisms of tensor powers of the quantum spin representation in detail; this work thus constitutes the first steps towards solving Kuperberg’s “webs problem” in type B . We also make contact with quantum groups and their categorifications.

1.1. Our construction, in a nutshell. The starting point for our investigation is a meta principle pioneered by Lusztig [43]: that representation-theoretic structures in non-simply laced type should arise via the process of *folding* categorical structures in simply laced type along diagram automorphisms. This principle is well-known; nevertheless, until now, the implementation of this principle in the context of link homology has remained elusive. The example we study is the folding of the root system A_{2n-1} to the root system of type B_n .

Our construction was originally motivated by examination of the invariants of colored unknots. Consider first the case $n = 2$. The root system $B_2 (= C_2)$ can be viewed as folded from the root system A_3 , with (the highest weight of) the \mathfrak{sl}_4 representation $\Lambda^2 \mathbb{C}^4$ corresponding to (the highest weight of) the spin representation S of \mathfrak{so}_5 :

$$\circ \text{---} \circ \text{---} \circ \xrightarrow{\text{fold}} \circ \text{---} \circ \text{---} \circ.$$

The associated quantum invariants $P_{\mathfrak{g}}$ of unknots (colored by these representations) are

$$(1.1) \quad P_{\mathfrak{sl}_4} \left(\bigcirc^{\Lambda^2} \right) = q^4 + q^2 + 2 + q^{-2} + q^{-4} \quad \text{and} \quad P_{\mathfrak{so}_5} \left(\bigcirc^S \right) = q^4 + q^2 + q^{-2} + q^{-4}.$$

The starting observation is that the latter invariant is obtained from the former by removing the summand 2, and that categorification provides a rigorous means for doing so.

When N is understood, let Λ^k denote the exterior product $\Lambda^k \mathbb{C}^N$, or its quantum analogue. In the \mathfrak{sl}_N Khovanov–Rozansky theory, the invariant of a Λ^k -colored unknot is a degree-shifted version of the cohomology of the Grassmannian $\text{Gr}_k(\mathbb{C}^N)$. In the case of (1.1), the \mathfrak{sl}_4 homology of the Λ^2 -colored unknot is $\mathbf{q}^{-4} H^*(\text{Gr}_2(\mathbb{C}^4))$, which has a basis indexed by partitions λ with Young diagrams fitting inside a 2×2 box; here the power of \mathbf{q} indicates a shift of this graded vector space (down, by 4). Observe that the graded dimension of $\mathbf{q}^{-4} H^*(\text{Gr}_2(\mathbb{C}^4))$ is exactly $q^4 + q^2 + 2 + q^{-2} + q^{-4}$. On the other hand, there is a natural involution on $H^*(\text{Gr}_2(\mathbb{C}^4))$, given by taking the transpose partition (induced by the involution on $\text{Gr}_2(\mathbb{C}^4)$ that takes the perpendicular 2-plane). If we instead take the graded trace of this involution, we recover $q^4 + q^2 + q^{-2} + q^{-4}$.

This behavior persists for all $n \geq 2$. Specifically, the \mathfrak{sl}_{2n} Khovanov–Rozansky homology of the Λ^n -colored unknot is isomorphic to $\mathbf{q}^{-n^2} H^*(\text{Gr}_n(\mathbb{C}^{2n}))$ and this space admits an analogous involution τ with graded trace that is equal to the spin-colored \mathfrak{so}_{2n+1} unknot invariant. Precisely,

$$\text{Tr} \left(\tau \bigcirc^{\Lambda^n} \mathbf{q}^{-n^2} H^*(\text{Gr}_n(\mathbb{C}^{2n})) \right) = \prod_{i=1}^n (q^{2i-1} + q^{1-2i}) = P_{\mathfrak{so}_{2n+1}} \left(\bigcirc^S \right).$$

A consequence of the main results of this paper is that these observations may be extended to all links.

Theorem 1.1 (Corollary 10.33 and Remark 10.35). Let $\mathcal{L} \subset S^3$ be a link.

- For all $n \geq 1$, the Λ^n -colored \mathfrak{sl}_{2n} Khovanov–Rozansky homology $H_{\mathfrak{sl}_{2n}}(\mathcal{L}^{\Lambda^n})$ admits an involution τ that preserves the bidegree. The bigraded eigenspaces of τ are link invariants.
- Assume that $n = 1, 2, 3$, or that Conjectures 4.40, 4.42, and 10.36 hold. Then

$$(1.2) \quad \mathrm{Tr} \left(\tau \left(\bigcirc_{H_{\mathfrak{sl}_{2n}}(\mathcal{L}^{\Lambda^n})} \right) \right) = \tilde{P}_{\mathfrak{so}_{2n+1}}(\mathcal{L}^S),$$

i.e., the pair $(H_{\mathfrak{sl}_{2n}}(\mathcal{L}^{\Lambda^n}), \tau)$ categorifies an appropriate renormalization $\tilde{P}_{\mathfrak{so}_{2n+1}}(\mathcal{L}^S)$ of the spin-colored \mathfrak{so}_{2n+1} quantum link polynomials. \square

The two parts of this theorem represent two distinct goals of this paper. The first is to define a new categorical link invariant, and this we achieve for all $n \geq 1$. The second goal is to determine what link polynomial our invariant categorifies. We expect it to categorify the spin-colored link polynomial for all $n \geq 1$, but technical issues of a combinatorial nature (primarily in the decategorification) obstruct our ability to verify this for $n > 3$. With sufficient time, it should be possible to check the relevant conjectures for any fixed value of n .

Theorem 1.1 is a slight reformulation and decategorification of our main construction, which defines *spin link homology* $H_{\mathfrak{so}_{2n+1}}(\mathcal{L}^S)$, a link invariant, valued in bigraded super vector spaces, that refines Λ^n -colored \mathfrak{sl}_{2n} Khovanov–Rozansky homology. As will be clear from our construction, this invariant is as computable¹ as the Khovanov–Rozansky theory. We thus propose that $H_{\mathfrak{so}_{2n+1}}(\mathcal{L}^S)$ are the first readily computable link homologies associated to simple complex $\mathfrak{g} \neq \mathfrak{sl}_N$.

Before proceeding, a word of caution. As the picture is painted above, it seems as though the Dynkin diagram automorphism of A_{2n-1} has been directly transformed into an involution τ on type A link homology. This is not at all what we do! Our construction of τ arises through duality, which we now explain.

1.2. Our construction, in more depth. Suppose one wishes to compute the \mathfrak{sl}_N Reshetikhin–Turaev invariant of a tangle with components colored by fundamental representations. One can view this invariant as a morphism between an m -fold tensor product of the representations Λ^a (with $0 \leq a \leq N$), for some² natural number m . Pioneering results of Cautis, Kamnitzer, Licata, and Morrison [14, 15] establish a duality between the subcategory of $\mathbf{Rep}(U_q(\mathfrak{sl}_N))$ consisting of such m -fold tensor products, and the idempotent quantum group $\check{U}_q(\mathfrak{gl}_m)$. In order to state the relevant representation theory correctly, we now replace \mathfrak{sl}_N with \mathfrak{gl}_N ; there is little distinction³ between \mathfrak{gl}_N and \mathfrak{sl}_N link invariants.

A consequence of the *quantum skew Howe duality* proved in [15] is the existence of a full functor

$$(1.3) \quad \check{U}_q(\mathfrak{gl}_m) \xrightarrow{\mathrm{SH}} \mathbf{Rep}(U_q(\mathfrak{gl}_N)).$$

Here, one views the idempotent algebra $\check{U}_q(\mathfrak{gl}_m)$ as a category with one object for each \mathfrak{gl}_m weight $\mathbf{a} = (a_1, \dots, a_m)$, and $\mathrm{SH}(\mathbf{a}) = \Lambda^{\mathbf{a}} := \Lambda^{a_1} \otimes \dots \otimes \Lambda^{a_m}$. After factoring through a quotient $\check{U}_q^{\leq N}(\mathfrak{gl}_m)$, the functor induced by (1.3) is fully faithful. Consequently, the link polynomials $P_{\mathfrak{sl}_N}(\mathcal{L})$, which are defined using the braided monoidal structure on $\mathbf{Rep}(U_q(\mathfrak{gl}_N))$, can be described entirely in terms of $\check{U}_q^{\leq N}(\mathfrak{gl}_m)$. The same is true at the categorical level [11], with $H_{\mathfrak{sl}_N}(\mathcal{L})$ admitting a formulation in the bounded homotopy category of complexes over an analogous quotient $\check{\mathcal{U}}_q^{\leq N}(\mathfrak{gl}_m)$ of the categorified quantum group $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$. The categorified quantum group is reviewed in full detail in §5 and the

¹That said, we do not include computations beyond the unknot here. The paper is already quite long, and (our) PhD students need things to do.

²The number m could be higher than the number of boundary points of the tangle, as one may need to add additional copies Λ^0 and/or Λ^N of the trivial representation to be the source or target of “cup” and “cap” tangles.

³The modern view is that categorical constructions are most naturally associated with \mathfrak{gl}_N , so we denote them thusly; however, as in the literature, we continue to refer to the link invariants as \mathfrak{sl}_N (Khovanov–Rozansky) link homology.

analogous quotient is recalled in Definition 9.6 (where it is denoted $\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)$). In order to precisely state our results, we remind the reader that 1-morphisms of $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$ are generated by elements $\mathcal{E}_i^{(k)} \mathbb{1}_{\mathbf{a}}$ and $\mathcal{F}_i^{(k)} \mathbb{1}_{\mathbf{a}}$ that lift divided powers of the Chevalley generators of $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$.

Remark 1.2. A categorification of [15] is given in [53]. Therein, a 2-category of \mathfrak{gl}_N foams is constructed, which categorifies the image of (1.3). As in the decategorified case, this foam 2-category is in duality with $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$.

Now consider the case where $N = 2n$. As we show in this paper, upon passage through (categorical) skew Howe duality, the Dynkin automorphism of \mathfrak{gl}_{2n} manifests as an involution on $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$ which is akin to the Chevalley involution. Namely, it swaps the Chevalley generators $e_i \leftrightarrow f_i$ in $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$ and sends a weight \mathbf{a} to the weight $2\mathbf{n} - \mathbf{a} = (2n - a_1, \dots, 2n - a_m)$. Precisely, we prove the following result at the categorical level.

Theorem 1.3 (Theorems 6.5 and 6.16 and Corollaries 6.8, 6.25 and 10.4). For each $n \geq 1$, there is an order 4 automorphism⁴ τ_n of the categorified quantum group $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$ that swaps generating 1-morphisms $\mathcal{E}_i^{(k)} \mathbb{1}_{\mathbf{n}+\mathbf{a}} \leftrightarrow \mathcal{F}_i^{(k)} \mathbb{1}_{\mathbf{n}-\mathbf{a}}$. Further, τ_n restricts to an involution on the 2-subcategory $\check{\mathcal{J}}\mathcal{E}_q(\mathfrak{gl}_m) \subset \check{\mathcal{U}}_q(\mathfrak{gl}_m)$ generated by the 1-endomorphisms $\mathcal{F}_i^{(k)} \mathcal{E}_i^{(k)} \mathbb{1}_{\mathbf{a}}$ and $\mathcal{E}_i^{(k)} \mathcal{F}_i^{(k)} \mathbb{1}_{\mathbf{a}}$. This involution descends to the corresponding 2-subcategory $\check{\mathcal{J}}\mathcal{E}_q^{\leq 2n}(\mathfrak{gl}_m) \subset \check{\mathcal{U}}_q^{\leq 2n}(\mathfrak{gl}_m)$, where it extends the involution on $H^*(\mathrm{Gr}_n(\mathbb{C}^{2n}))$. \square

To explain the final sentence of Theorem 1.3, we point out that the endomorphism algebra of the identity 1-morphism $\mathbb{1}_{\mathbf{a}}$ of the weight \mathbf{a} in $\check{\mathcal{U}}_q^{\leq N}(\mathfrak{gl}_m)$ is isomorphic to $\otimes_{i=1}^m H^*(\mathrm{Gr}_{a_i}(\mathbb{C}^N))$, which is the \mathfrak{sl}_N Khovanov–Rozansky homology of the \mathbf{a} -colored unlink (up to degree shift). When $N = 2n$, our involution τ of $\check{\mathcal{J}}\mathcal{E}_q^{\leq 2n}(\mathfrak{gl}_m)$ restricts in weight $\mathbf{n} = (n, \dots, n)$ to give an involution of $\otimes_{i=1}^m H^*(\mathrm{Gr}_n(\mathbb{C}^{2n}))$, which agrees with (the m -fold tensor product of) the involution from §1.1. Note that the weight \mathbf{n} corresponds to labeling each component of the unlink with Λ^n , the fundamental representation which folds to the spin representation in type B .

Remark 1.4. We began this project by searching for an extension of the involution on $H^*(\mathrm{Gr}_n(\mathbb{C}^{2n}))$ to all of $\check{\mathcal{U}}_q^{\leq 2n}(\mathfrak{gl}_m)$. We were bemused to find that it extended not to an involution, but an automorphism of order 4. That its restriction to $\check{\mathcal{J}}\mathcal{E}_q^{\leq 2n}(\mathfrak{gl}_m)$ is an involution is crucial to our constructions below.

The setting for our link invariant is the monoidal subcategory $\mathcal{B}_m^n \subset \check{\mathcal{J}}\mathcal{E}_q^{\leq 2n}(\mathfrak{gl}_m)$ of 1-endomorphisms of the object \mathbf{n} , or, more precisely, the associated *equivariant category* $(\mathcal{B}_m^n)^\tau$ of this category with respect to the involution $\tau = \tau_n$. Objects of $(\mathcal{B}_m^n)^\tau$ are equivariant structures: pairs (X, φ_X) , where X is an object of \mathcal{B}_m^n and φ_X is an isomorphism $\varphi_X: X \xrightarrow{\cong} \tau(X)$ in \mathcal{B}_m^n satisfying $\tau(\varphi_X) \circ \varphi_X = \mathrm{id}_X$. Morphisms $f: (X, \varphi_X) \rightarrow (Y, \varphi_Y)$ are morphisms $f: X \rightarrow Y$ in \mathcal{B}_m^n satisfying $\varphi_Y \circ f = \tau(f) \circ \varphi_X$.

The equivariant category $(\mathcal{B}_m^n)^\tau$ itself also admits a $\mathbb{Z}/2$ -action, given by $\varphi_X \mapsto -\varphi_X$. In particular, the monoidal identity $\mathbb{1} = \mathbb{1}_{\mathbf{n}}$ of \mathcal{B}_m^n gives rise to two equivariant objects $(\mathbb{1}, \mathrm{id}_{\mathbb{1}})$ and $(\mathbb{1}, -\mathrm{id}_{\mathbb{1}})$ which are exchanged by this $\mathbb{Z}/2$ -action.

The Rickard complexes of [13, 14, 11] (which generalize the Rouquier complex [17, 62]) associate a bounded complex $C(\beta_{i_1}^\pm \cdots \beta_{i_r}^\pm) \in \mathcal{K}(\mathcal{B}_m^n)$ to each word in the generators $\{\beta_i\}_{i=1}^{m-1}$ of the m -strand braid group Br_m . We show that these complexes determine *equivariant Rickard complexes* $C^\tau(\beta_{i_1}^\pm \cdots \beta_{i_r}^\pm) \in \mathcal{K}((\mathcal{B}_m^n)^\tau)$. In the setting of $\mathcal{K}(\mathcal{B}_m^n)$, braid relations are categorified by canonical homotopy equivalences, and therefore the Rickard complexes canonically associate a complex $C(\beta) \in \mathcal{K}(\mathcal{B}_m^n)$

⁴Our automorphism τ_n is unrelated to other symmetries of $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$ appearing in the literature, despite its similar action on 1-morphisms.

to each $\beta \in \text{Br}_m$. We show that the same is true for the equivariant Rickard complexes, and then apply appropriate representable functors to obtain invariants of links.

Theorem 1.5 (Theorems 10.21 and 10.28). The equivariant Rickard complexes assign a complex $C^\tau(\beta) \in \mathcal{K}((\mathcal{B}_m^n)^\tau)$ to each $\beta \in \text{Br}_m$, which is well-defined up to canonical homotopy equivalence. Let $\text{sVect}_{\mathbb{K}}^{\mathbb{Z}}$ denote the category of \mathbb{Z} -graded super vector spaces and let $\mathcal{R}: (\mathcal{B}_m^n)^\tau \rightarrow \text{sVect}_{\mathbb{K}}^{\mathbb{Z}}$ be the representable functor with even and odd components

$$\mathcal{R}(x)_{\bar{0}} = \text{Hom}_{(\mathcal{B}_m^n)^\tau}((\mathbb{1}, \text{id}_{\mathbb{1}}), x), \quad \mathcal{R}(x)_{\bar{1}} = \text{Hom}_{(\mathcal{B}_m^n)^\tau}((\mathbb{1}, -\text{id}_{\mathbb{1}}), x).$$

Then, the homology $H_{\mathfrak{so}_{2n+1}}(\mathcal{L}_\beta)$ of the complex $\mathbf{q}^{-mn^2} \mathcal{R}(C^\tau(\beta))$ is an invariant of the link $\mathcal{L}_\beta \subset S^3$ arising as the braid closure of β . \square

Remark 1.6. There is no difference between super vector spaces and $\mathbb{Z}/2$ -graded vector spaces when only considered as monoidal categories. Given this, one would not typically use super vector spaces unless the braiding on this braided monoidal category (which does distinguish it from $\mathbb{Z}/2$ -graded vector spaces) were relevant. While it is not relevant in this paper, we use the language of super vector spaces rather than $\mathbb{Z}/2$ -graded vector spaces for two reasons. First, to avoid confusion, as many other gradings already appear in the paper. Second, because our results are conveniently stated in terms of super dimension (which, if one considers the relevant pivotal structure on super vector spaces, equals the dimension in this category).

1.3. Our construction, decategorified. For each $n \geq 1$, our construction produces a homological link invariant. Conditional upon the assumptions in Theorem 1.1, an appropriate normalization of the graded Euler characteristic of $H_{\mathfrak{so}_{2n+1}}(\mathcal{L}_\beta)$ equals the spin-colored \mathfrak{so}_{2n+1} link polynomial $\bar{P}_{\mathfrak{so}_{2n+1}}(\mathcal{L}_\beta^S) := P_{\mathfrak{so}_{2n+1}}(\mathfrak{m}\mathcal{L}_\beta^S)$ of the mirror link $\mathfrak{m}\mathcal{L}$.

Theorem 1.7 (Theorem 10.32 and Remark 10.35). Assume that $n = 1, 2, 3$ or that Conjectures 4.40, 4.42, and 10.36 hold. If $\beta \in \text{Br}_m$ and $\epsilon(\beta)$ denotes the braid exponent, then

$$\bar{P}_{\mathfrak{so}_{2n+1}}(\mathcal{L}_\beta^S) = (-1)^{n\epsilon(\beta) + m\binom{n+1}{2}} q^{\frac{1}{2}n\epsilon(\beta)} \sum_i (-1)^i \dim_{\mathbf{q}}(H_{\mathfrak{so}_{2n+1}}^i(\mathcal{L}_\beta^S)).$$

\square

This result and Theorem 1.1 are essentially reformulations of one another. Indeed, for an equivariant object (X, φ_X) in $(\mathcal{B}_m^n)^\tau$, the space of morphisms $\text{Hom}_{\mathcal{B}_m^n}(\mathbb{1}, X)$ admits an involutory action of τ , whose $+1$ -eigenspace is $\text{Hom}_{(\mathcal{B}_m^n)^\tau}((\mathbb{1}, \text{id}_{\mathbb{1}}), -)$ and whose -1 -eigenspace is $\text{Hom}_{(\mathcal{B}_m^n)^\tau}((\mathbb{1}, -\text{id}_{\mathbb{1}}), -)$. Meanwhile, the dimension of a super vector space $V = V_{\bar{0}} \oplus V_{\bar{1}}$ is $\dim(V) = \dim(V_{\bar{0}}) - \dim(V_{\bar{1}})$; if $V_{\bar{0}}$ and $V_{\bar{1}}$ arise as the ± 1 -eigenspaces of an involution τ on V , this also equals $\text{Tr}(\tau)$.

From this perspective, Theorem 1.5 can be repackaged as the statement that τ descends to an involution on the complex $\text{Hom}_{\mathcal{B}_m^n}(\mathbb{1}, C(\beta))$ that computes Λ^n -colored \mathfrak{sl}_{2n} Khovanov–Rozansky homology (and is compatible with braid/Markov moves). Theorem 1.7 then implies Theorem 1.1. We emphasize that the conjectural results on which those theorems rely for $n \geq 4$ are primarily related to open questions in the decategorified setting. Regardless, for all $n \geq 1$ we still obtain the involution τ , and hence our link invariants from Theorem 1.5.

1.4. Intertwiners for the spin representation. Theorem 1.7 relies on new results that we establish concerning the endomorphism algebras of tensor powers $S^{\otimes m}$ of the spin representation in $\text{Rep}(U_q(\mathfrak{so}_{2n+1}))$. We believe these results to be of independent interest. Although they do not depend on our categorified construction, they are informed by it.

For the moment, we forget the involution τ and focus on the (surjective) algebra homomorphism

$$(1.4) \quad 1_{\mathbf{n}} \dot{U}_q(\mathfrak{gl}_2) 1_{\mathbf{n}} \rightarrow \text{End}_{U_q(\mathfrak{gl}_{2n})}(\Lambda^n \otimes \Lambda^n)$$

arising from the $m = 2$ case of (1.3) restricted to endomorphisms of the object $\mathbf{n} = (n, n)$. For $k \geq 1$, let $\chi^{(k)} := f^{(k)} e^{(k)} \mathbb{1}_{\mathbf{n}} \in \dot{U}_q(\mathfrak{gl}_2)$ where $e^{(k)} := \frac{1}{[k]!} e^k$ and $f^{(k)} := \frac{1}{[k]!} f^k$ are the divided power elements. It is straightforward to compute from the relations of $\dot{U}_q(\mathfrak{gl}_2)$ (found in Definition 5.1) that

$$(1.5) \quad \chi^{(k)} \cdot \chi = [k][k+1]\chi^{(k)} + [k+1]^2\chi^{(k+1)}.$$

Consequently, we obtain endomorphisms (also) denoted by $\chi^{(k)} \in \text{End}_{U_q(\mathfrak{gl}_{2n})}(\Lambda^n \otimes \Lambda^n)$ that again satisfy (1.5). Moreover, [15] describes explicitly how the braiding $R_{n,n} \in \text{End}_{U_q(\mathfrak{gl}_{2n})}(\Lambda^n \otimes \Lambda^n)$ can be expressed as a linear combination of the elements $\{\chi^{(k)}\}_{0 \leq k \leq n}$.

The left-hand side of (1.4) is categorified by $\mathbb{1}_{\mathbf{n}} \dot{\mathcal{U}}_q(\mathfrak{gl}_2) \mathbb{1}_{\mathbf{n}}$, with indecomposable 1-morphisms $\mathbf{X}^{(k)} := \mathcal{F}^{(k)} \mathcal{E}^{(k)} \mathbb{1}_{\mathbf{n}}$ corresponding to $\chi^{(k)}$ in the Grothendieck ring. There is a direct sum decomposition

$$(1.6) \quad \mathbf{X}^{(k)} \otimes \mathbf{X} \cong [k][k+1]\mathbf{X}^{(k)} \oplus [k+1]^2\mathbf{X}^{(k+1)}$$

lifting (1.5), where here multiplication by a quantum integer corresponds to an appropriate direct sum of grading shifts of $\mathbf{X}^{(k)}$. For example, $[2]\mathbf{X} = \mathbf{q}^1\mathbf{X} \oplus \mathbf{q}^{-1}\mathbf{X}$. Note that e.g. $[k][k+1]\mathbf{X}^{(k)}$ can be viewed either as a direct sum of shifted copies of $\mathbf{X}^{(k)}$, or equivalently as a formal tensor product of $\mathbf{X}^{(k)}$ with a ‘‘multiplicity space’’ whose graded dimension equals $[k][k+1]$.

Now we reintroduce the involution τ . As we show, the $\mathbf{X}^{(k)}$ admit equivariant structures, and τ induces a $\mathbb{Z}/2$ -action on the multiplicity spaces in (1.6). In Corollary 8.10, we compute the trace of this action. Up to a sign, the traces of τ on the above multiplicity spaces of dimension $[k][k+1]$ and $[k+1]^2$ are given by Laurent polynomials that we denote ‘‘ $[k][k+1]$ ’’ and ‘‘ $[k+1][k+1]$ ’’. Paralleling (1.1), these latter quantities are obtained from their unfolded counterparts by replacing certain coefficients with integers congruent to them modulo 2. For example,

$$[3][4] = [6] + [4] + [2] = q^5 + 2q^3 + 3q + 3q^{-1} + 2q^{-3} + q^{-5}$$

while

$$‘‘[3][4]’’ = [6] - [4] + [2] = q^5 + q + q^{-1} + q^{-5}.$$

Please allow us to introduce this ‘‘incorrect’’ arithmetic on quantum numbers, which we sympathetically call the *devil’s arithmetic*, in §4 below.

Theorems 1.1 and 1.7 suggest the existence of elements $\mathbf{X}^{(k)} \in \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$ analogous to the elements $\chi^{(k)}$. Working solely at the decategorified level, we establish the following.

Theorem 1.8 (Theorems 4.2 and 4.39). There is a basis $\{\mathbf{X}^{(k)}\}_{k=0}^n \subset \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$ in which the braiding is given by

$$(1.7) \quad R_{S,S} = q^{\frac{n}{2}} \sum_{k=0}^n q^{-k} \mathbf{X}^{(k)}.$$

The structure coefficients for multiplication in this basis are determined by

$$(1.8) \quad \mathbf{X}^{(k)} \mathbf{X}^{(1)} = (-1)^k ‘‘[k][k+1]’’ \mathbf{X}^{(k)} + (-1)^k ‘‘[k+1][k+1]’’ \mathbf{X}^{(k+1)}.$$

Further, the elements $\mathbf{X}_i^{(k)} := \text{id}_S^{\otimes i-1} \otimes \mathbf{X}^{(k)} \otimes \text{id}_S^{\otimes m-i-1}$ for $1 \leq i \leq m-1$ and $0 \leq k \leq n$ generate $\text{End}_{U_q(\mathfrak{so}_{2n+1})}((S)^{\otimes m})$. \square

We propose the elements $\mathbf{X}^{(k)}$ as a new *canonical basis* for the endomorphism space $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$. Indeed, they arise as a twisted decategorification of the indecomposable objects $\mathbf{X}^{(k)} = \mathcal{F}^{(k)} \mathcal{E}^{(k)} \mathbb{1}_{\mathbf{n}}$ which correspond to canonical basis elements in $\dot{U}_q(\mathfrak{gl}_2)$.

In more detail, there is a decategorification procedure for categories such as $(\mathcal{B}_m^n)^\tau$, called the *weighted Grothendieck group* $K_0^\tau(-)$. We recall this construction (in the case of $\mathbb{Z}/2$ -actions) in §7.4. Loosely stated, while the structure coefficients in the ordinary Grothendieck group are dimensions of multiplicity spaces, the structure coefficients of the weighted Grothendieck group are the super

dimensions of those same multiplicity spaces. As we show, an appropriate renormalization of the classes $[\mathbf{X}^{(k)}]_\tau \in K_0^\tau((\mathcal{B}_2^n)^\tau)$ satisfy equation (1.8). Since all indecomposable 1-morphisms in \mathcal{B}_2^n are of the form $\mathbf{X}^{(k)}$ for $0 \leq k \leq n$, standard results on weighted Grothendieck groups give the following.

Theorem 1.9 (Corollary 10.10). For all $n \geq 1$, there is an isomorphism of $\mathbb{C}(q)$ -algebras

$$\mathbb{C}(q) \otimes_{\mathbb{Z}[q^\pm]} K_0^\tau((\mathcal{B}_2^n)^\tau) \xrightarrow{\cong} \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$$

that sends the class of (an appropriate equivariant structure on) $\mathbf{X}^{(k)}$ to $\mathbf{X}^{(k)}$. \square

Since this is the folded analogue of (1.4), we refer to this result as a *folded skew Howe duality*. More generally, we propose the following generalization.

Conjecture 1.10. For all $n \geq 1$ and $m \geq 2$, there is an isomorphism of $\mathbb{C}(q)$ -algebras

$$\mathbb{C}(q) \otimes_{\mathbb{Z}[q^\pm]} K_0^\tau((\mathcal{B}_m^n)^\tau) \xrightarrow{\cong} \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$$

that sends the class of (an appropriate equivariant structure on) $\mathcal{F}_i^{(k)} \mathcal{E}_i^{(k)} \mathbb{1}_n$ to $\mathbf{X}_i^{(k)}$.

Further evidence for Conjecture 1.10 is as follows. In Proposition 4.38, we prove that the elements \mathbf{X}_i and $\mathbf{X}_{i\pm 1}$ satisfy a ‘‘Reidemeister III’’-like relation (4.40) in $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$. In Theorem 8.22 and Corollary 10.10, we then prove that this relation is categorified by a direct sum decomposition in $(\mathcal{B}_m^n)^\tau$. Consequently, Conjecture 1.10 would follow from the following two conjectures:

- That the ‘‘Reidemeister III’’-like relation (4.40), together with (1.8), gives a presentation for the $\mathbb{C}(q)$ -algebra $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$.
- That the Grothendieck group $K_0^\tau((\mathcal{B}_m^n)^\tau)$ is generated as a ring by the classes of the $\mathbf{X}_i^{(k)}$, and the dimensions of $K_0^\tau((\mathcal{B}_m^n)^\tau)$ and $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$ are equal.

We discuss these conjectures further in the body of the paper.

Remark 1.11. Theorem 1.9 and Conjecture 1.10 can be viewed as a categorification of an instance of [26, Theorem 1.1]. There, working in the setting of a connected almost simple algebraic group G equipped with a Dynkin automorphism σ , Hong and Shen prove that the dimensions of invariant spaces for the algebraic group G_σ associated to (G, σ) via folding can be computed as the trace of the involution induced by σ on invariant spaces for G . In the $G = SL_{2n}$ case, folded skew Howe duality asserts that the invariant space itself can be recovered via folding by considering involutions on the categories that categorify these invariant spaces.

We emphasize that folded skew Howe duality is necessarily a product of categorification: it shows that the algebra $\mathbb{1}_n \check{U}_q(\mathfrak{gl}_m) \mathbb{1}_n$ has a ‘‘secret’’ relation with $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$ that is only visible by first considering the categorification $\mathbb{1}_n \check{\mathcal{U}}_q(\mathfrak{gl}_m) \mathbb{1}_n$, then passing to the equivariant (subquotient) category $(\mathcal{B}_m^n)^\tau$, and finally taking the weighted Grothendieck ring. This procedure remains invisible at the decategorified level: there is no obvious method to modify the relations of (1.6) into those of (1.8). Said another way, if one only knows the ordinary dimension $\dim V_0 + \dim V_1$ for a super vector space V , one can not deduce the value of the super dimension $\dim V_0 - \dim V_1$.

Nevertheless, algebras related to $K_0^\tau((\mathcal{B}_m^n)^\tau)$ have previously appeared in the literature.

1.5. Relationship to existing spin literature and further results. The endomorphism algebra $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$ has previously been studied by Wenzl [74] and Reshetikhin [59], and also by Deligne [18] and McNamara–Savage [50] in the classical ($q = 1$) setting. All of these works (save for Wenzl’s), use the diagrammatic language for monoidal categories to describe intertwiners between tensor products of S and the vector representation of (quantum) \mathfrak{so}_{2n+1} .

In order to establish the results in §1.4, we further develop the diagrammatic calculus for the spin representation of $U_q(\mathfrak{so}_{2n+1})$. Our advances in this direction are the expression (1.7) for the braiding

in terms of our canonical basis from Theorem 1.8, the “ $H = I$ ” relation given in Lemma 4.23 below, and the interpretation of numerous structure coefficients as arising from the devil’s arithmetic.

Reshetikhin notes that his relation [59, Equation 5.29], which we reestablish below in Corollary 4.36, can be used to define a q -analogue of the Clifford algebra. Later, in [74, Equation 3.9], Wenzl constructs a distinguished endomorphism $C \in \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$ using these q -Clifford algebras. Via calculations with this element, Wenzl goes on to establish a relation to an algebra defined twenty years earlier by Gavrilik–Klimyk⁵.

Definition 1.12 ([23]). Let $U'_q(\mathfrak{so}_m)$ be the $\mathbb{C}(q)$ -algebra generated by b_1, \dots, b_{m-1} subject to the relations

$$b_i b_j = b_j b_i \quad |i - j| > 1,$$

and

$$b_i^2 b_{i\pm 1} + b_{i\pm 1} b_i^2 = [2] b_i b_{i\pm 1} b_i + b_{i\pm 1}.$$

Wenzl’s calculations in [74, Section 4] and [74, Theorem 5.2] give the following.

Theorem 1.13 (Wenzl). There is a surjective algebra homomorphism

$$(1.9) \quad U'_{-q^2}(\mathfrak{so}_m) \rightarrow \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$$

such that $b_i \mapsto \text{id}_S^{\otimes i-1} \otimes C \otimes \text{id}_S^{\otimes m-i-1}$.

Remark 1.14. A consequence of Wenzl’s theorem is that the vector space $S^{\otimes m}$ decomposes into a direct sum of irreducible representations of $U'_{-q^2}(\mathfrak{so}_m)$. However, the irreducible $U'_{-q^2}(\mathfrak{so}_m)$ -representations that appear are not q - (or $-q^2$)-analogues of irreducible representations of \mathfrak{so}_m . Wenzl overcomes this issue by studying “non-classical” representations of $U'_{-q^2}(\mathfrak{so}_m)$ first introduced in [27]; see [74, Theorem 2.1].

In Appendix B, we explicitly relate our approach to Wenzl’s endomorphism C . Proposition B.10 implies that

$$(1.10) \quad \mathbf{X} = C - \frac{1}{[2]} \text{id}_{S \otimes S}.$$

The most novel feature of our work in this direction is the simple formula (1.7) for the braiding after passing from Wenzl’s C to our \mathbf{X} . To our knowledge, such a formula has not previously appeared in the literature on the quantum spin representation. Further, our element \mathbf{X} also (conjecturally) sheds light on the kernel of (1.9); see Conjecture 1.20.

Remark 1.15. With the computations of Lemma 4.37, we can reprove the existence of the homomorphism (1.9). Note that we do not reprove the surjectivity of (1.9), but rather use Wenzl’s result to establish surjectivity in Theorem 1.8.

1.6. **Categorifying $U'_{-q^2}(\mathfrak{so}_m)$.** Taken together, our results thus far suggest an approach to the categorification of the algebra $U'_{-q^2}(\mathfrak{so}_m)$ itself. Inspired by (1.10) we make the following definition.

Definition 1.16. Inside $U'_{-q^2}(\mathfrak{so}_2)$, let $b = b_1$. For $k \geq 0$, define elements $\{x^{(k)}\}_{k \geq 0} \subset U'_{-q^2}(\mathfrak{so}_2)$ recursively as follows:

$$(1.11) \quad \begin{aligned} x^{(0)} &:= 1, & x^{(1)} &= x := b - \frac{1}{[2]}, \\ x^{(k)} x &= (-1)^k [k+1][k] x^{(k)} + (-1)^k [k+1][k+1] x^{(k+1)}. \end{aligned}$$

Set $x_i := b_i - \frac{1}{[2]}$ in $U'_{-q^2}(\mathfrak{so}_m)$ and define $x_i^{(k)}$ analogously.

⁵Their goal was to define a q -analogue of \mathfrak{so}_m which, unlike the usual Drinfeld–Jimbo quantum \mathfrak{so}_m , was adapted to the chain of embeddings $\mathfrak{so}_{m-1} \subset \mathfrak{so}_m$.

It is easy to deduce that $\{x^{(k)}\}_{k \geq 0}$ is a basis for $U'_{-q^2}(\mathfrak{so}_2)$. Paralleling Theorem 1.9, we obtain a categorification of $U'_{-q^2}(\mathfrak{so}_2)$ in which indecomposables correspond to this basis.

Theorem 1.17 (Corollary 8.12). For all $n \geq 1$, consider the involution τ_n of $\mathbb{1}_n \check{\mathcal{U}}_q(\mathfrak{gl}_2) \mathbb{1}_n$ given by Theorem 1.3. There is an isomorphism of $\mathbb{C}(q)$ -algebras

$$U'_{-q^2}(\mathfrak{so}_2) \xrightarrow{\cong} \mathbb{C}(q) \otimes_{\mathbb{Z}[q^\pm]} K_0^{\tau_n}((\mathbb{1}_n \check{\mathcal{U}}_q(\mathfrak{gl}_2) \mathbb{1}_n)^{\tau_n}).$$

that is intertwined with the isomorphism in Theorem 1.9 by Wenzl's homomorphism (1.9). In other words, it sends $x^{(k)}$ to the class of (an appropriate equivariant structure on) $\mathbf{X}^{(k)} = \mathcal{F}^{(k)} \mathcal{E}^{(k)} \mathbb{1}_n$. \square

We direct the reader distressed by the appearance of seemingly many (one for each $n \geq 1$) categorifications of $U'_{-q^2}(\mathfrak{so}_2)$ to §6.4. The $m = 2$ case of the results there give equivalences between the $\mathbb{1}_n \check{\mathcal{U}}_q(\mathfrak{gl}_2) \mathbb{1}_n$ (for various n) that intertwine the involutions τ_n .

Continuing the parallel, we expect that Theorem 1.17 generalizes from $m = 2$ to all $m \geq 2$. Recall that $\check{\mathcal{F}}_q(\mathfrak{gl}_m)$ is the full subcategory of $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$ generated by the 1-endomorphisms $\mathcal{F}_i^{(k)} \mathcal{E}_i^{(k)}$ and $\mathcal{E}_i^{(k)} \mathcal{F}_i^{(k)}$. We let $\mathcal{B}_m := \mathbb{1}_n \check{\mathcal{F}}_q(\mathfrak{gl}_m) \mathbb{1}_n$, suppressing n from the notation (all these categories as n varies are identified in §6.4).

Conjecture 1.18. There is an isomorphism of $\mathbb{C}(q)$ -algebras

$$U'_{-q^2}(\mathfrak{so}_m) \xrightarrow{\cong} \mathbb{C}(q) \otimes_{\mathbb{Z}[q^\pm]} K_0^\tau((\mathcal{B}_m)^\tau).$$

that sends $x_i^{(k)}$ to the class of (an appropriate equivariant structure on) $\mathbf{X}_i^{(k)} = \mathcal{F}_i^{(k)} \mathcal{E}_i^{(k)} \mathbb{1}_n$.

We again discuss some evidence for this conjecture. The relations of Definition 1.12 can be transformed into the following statement.

Proposition 1.19. The elements $x_i^{(k)} \in U'_{-q^2}(\mathfrak{so}_m)$ satisfy

$$(1.12) \quad \begin{aligned} x_i x_j &= x_j x_i \quad |i - j| > 1, \\ x_i x_{i \pm 1} x_i &= x_i^{(2)} x_{i \pm 1} + x_{i \pm 1} x_i^{(2)} + [2] x_i^{(2)} + x_i. \end{aligned}$$

These relations (together with the definition of $x_i^{(2)}$) give a presentation for $U'_{-q^2}(\mathfrak{so}_m)$. \square

In this context, we think of (1.12) as a variant on the usual quantum group Serre relations. We refer to it as the *devil's Serre relation*. Replacing x_i with $\mathbf{X}_i \in \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$ transforms the devil's Serre relation into the ‘‘Reidemeister III’’-like relation (4.40) discussed above, so Conjecture 1.18 is again intertwined with Conjecture 1.10 by Wenzl's theorem. As additional evidence for Conjecture 1.18, Theorem 8.22 and Corollary 8.23 provide a lift of the relations (1.12) to isomorphisms in the τ_n -equivariant category of \mathcal{B}_m and establish the existence of the $\mathbb{C}(q)$ -algebra homomorphism appearing in Conjecture 1.18.

We make one further statement about the decategorified setting, inspired by categorification. Recall Wenzl's surjective homomorphism

$$U'_{-q^2}(\mathfrak{so}_m) \rightarrow \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m}).$$

We propose above that the left-hand side is categorified (in the equivariant sense) by \mathcal{B}_m , and the right-hand side by $(\mathcal{B}_m^n)^\tau$. The relationship between \mathcal{B}_m and $(\mathcal{B}_m^n)^\tau$ is the projection to $\mathbb{1}_n$ of the surjection

$$\check{\mathcal{U}}_q(\mathfrak{gl}_m) \rightarrow \check{\mathcal{U}}_q^{\leq 2n}(\mathfrak{gl}_m),$$

which categorifies the surjection from $\dot{U}_q(\mathfrak{gl}_m)$ to a particular Schur algebra. The object $\mathbf{X}_i^{(k)}$ is sent to the zero object in this quotient, for any $k > n$.

Analogously, Wenzl's homomorphism satisfies

$$x_i^{(k)} \mapsto \begin{cases} X_i^{(k)} & 1 \leq k \leq n, \\ 0 & k > n, \end{cases}$$

by e.g. Remark 4.26 below. Denote by $I^{>n} \subset U'_{-q^2}(\mathfrak{so}_m)$ the two sided ideal generated by the elements $x_i^{(n+1)}$ for $i = 1, \dots, m-1$. We propose that the ideal $I^{>n}$ is exactly the kernel of Wenzl's homomorphism.

Conjecture 1.20. The algebra homomorphism in Theorem 1.13 induces an isomorphism

$$U'_{-q^2}(\mathfrak{so}_m)^{\leq n} := U'_{-q^2}(\mathfrak{so}_m)/I^{>n} \xrightarrow{\cong} \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m}).$$

It is straightforward to verify this conjecture when $n = 1$, by comparing the algebras with the Temperley–Lieb algebra. One can also verify the case $n = 2$ by direct calculation, using type $B_2 = C_2$ webs [38, 7]. For example, compare (1.12) to [7, equation (5.56b)].

Remark 1.21. A consequence of the conjecture is that the algebra $U'_{-q^2}(\mathfrak{so}_m)^{\leq n}$ is a finite dimensional algebra. Assume that we knew that $U'_{-q^2}(\mathfrak{so}_m)^{\leq n}$ is finite dimensional. Since finite dimensional $U'_{-q^2}(\mathfrak{so}_m)$ -modules are completely reducible [27], it would follow that $U'_{-q^2}(\mathfrak{so}_m)^{\leq n}$ is a finite direct sum of algebras $\text{End}(L)$, where L is a simple module for $U'_{-q^2}(\mathfrak{so}_m)$ that is annihilated by $I^{>n}$. This is the folded skew Howe duality analogue of [15, Lemma 4.4.2]. We expect that one can use [74, Theorem 5.3(a)] to show that each such L is isomorphic to a direct summand of the $U'_{-q^2}(\mathfrak{so}_m)$ -module $S^{\otimes m}$, and that this exhausts all (isomorphism classes) of irreducible summands appearing in $S^{\otimes m}$. Then, Conjecture 1.20 would follow by mimicking the proof of [15, Theorem 4.4.1].

The takeaway is that the proof sketch in Remark 1.21 would prove Conjecture 1.20 once one can show that $U'_{-q^2}(\mathfrak{so}_m)^{\leq n}$ is finite dimensional. The resolution of our Conjecture 4.40 would imply finite-dimensionality (so Proposition 4.41 shows that we already know finite-dimensionality when $n = 3$).

1.7. Relationship to ι quantum groups. The algebra $U'_q(\mathfrak{so}_m)$ appearing in Theorem 1.13 is a special case of a so-called ι quantum group. The latter are algebras that arise in the theory of quantum symmetric pairs. We refer the reader to Wang's 2022 ICM address [70] for comprehensive details and references to the vast body of work on this subject, and recall only the immediately pertinent information here.

Quantum symmetric pairs are parametrized by Satake diagrams [70, Section 1.2], which determine a semisimple Lie algebra \mathfrak{g} and an involution θ of \mathfrak{g} . The pair $(\mathfrak{g}, \mathfrak{g}^\theta)$ consisting of the Lie algebra and its fixed-point subalgebra is a (classical) symmetric pair. Analogously, a *quantum symmetric pair* is a tuple $(U_q(\mathfrak{g}), U'_q(\mathfrak{g}^\theta))$ where $U'_q(\mathfrak{g}^\theta)$ is a (coideal) subalgebra of $U_q(\mathfrak{g})$, referred to as an ι quantum group.

Remark 1.22. In the quasi-split case⁶, the Satake diagram is determined by a pair (D, t) , where D is the Dynkin diagram associated to \mathfrak{g} , and t is a Dynkin diagram involution⁷, see e.g. [70, Example 1.1]. A quasi-split pair is called split if $t = \text{id}$ and, in this case, θ is the Chevalley involution which swaps generators e_i and f_i , and acts by -1 on the Cartan subalgebra \mathfrak{h} .

The algebra $U'_q(\mathfrak{so}_m)$ can be identified with the ι quantum group associated to the split symmetric pair $(\mathfrak{sl}_m, \mathfrak{so}_m)$; see [42, Remark 2.4]. In particular, there is an embedding $U'_q(\mathfrak{so}_m) \rightarrow U_q(\mathfrak{sl}_m)$ given by $b_i \mapsto f_i + q^{-1}e_i k_i^{-1}$, which induces the isomorphism $U'_q(\mathfrak{so}_m) \xrightarrow{\cong} U'_q(\mathfrak{so}_m)$. Henceforth, we identify $U'_q(\mathfrak{so}_m)$ with $U'_q(\mathfrak{so}_m)$.

⁶This means there are no “black dots” on the Satake diagram.

⁷In [70] this Dynkin involution is denoted by τ .

We now contrast our results with the future goals and recent results from the ι quantum group literature. The punchline is that our constructions do **not** agree. Hence, we are hopeful that the constructions in the present paper will shed new light on the theory of ι quantum groups.

Using based modules for $U_q(\mathfrak{g})$ and the quasi K -matrix [70, Theorem 3.1], it is possible to construct the so-called ι canonical basis in the restriction of a finite dimensional based $U_q(\mathfrak{g})$ -module to $U_q^\iota(\mathfrak{g}^\theta)$. Then, in analogy with Lusztig’s construction of canonical bases for quantum groups using tensor products of based modules, Bao and Wang [3], defined the ι canonical basis in the modified (i.e. idempotent) form of the ι quantum group $U_q^\iota(\mathfrak{g}^\theta)$.

Example 1.23. In the special case of the split quantum symmetric pair corresponding to (A_1, id) , the ι canonical basis is computed explicitly in [70, Example 3.3]. Here, there are two modified versions $\dot{U}_q^\iota(\mathfrak{so}_2)1_{\bar{0}}$ and $\dot{U}_q^\iota(\mathfrak{so}_2)1_{\bar{1}}$ of the ι quantum group $U_q^\iota(\mathfrak{so}_2)$, where the parity corresponds to the parity of weights in irreducible \mathfrak{sl}_2 representations.

Denoting the generator of $\dot{U}_q^\iota(\mathfrak{so}_2)1_\epsilon$ by b_ϵ , the ι canonical basis consists of the ι divided powers, which are given by

$$(1.13) \quad b_\epsilon^{(0)} = 1, \quad b_\epsilon^{(1)} = b_\epsilon, \quad \text{and} \quad b_\epsilon^{(k)}b_\epsilon = [k+1]b_\epsilon^{(k+1)} + \delta_{\bar{k}, \epsilon} [k]b_\epsilon^{(k-1)}$$

where $\bar{k} = k \bmod 2$.

From this example, one can see that the basis $b_\epsilon^{(k)}$ for $U_{-q^2}^\iota(\mathfrak{so}_2)$ does **not** agree with our basis $x^{(k)}$ (nor is our basis dependent on parity). We henceforth refer to our elements $\{x^{(k)}\}$ as the *devil’s divided powers*.

Remark 1.24. Bao and Wang’s ι canonical basis for $U_q^\iota(\mathfrak{g}^\theta)$ is characterized as the unique basis which is “asymptotically compatible” with the ι canonical basis of tensor products of highest weight $U_q(\mathfrak{g})$ modules [3, Theorem G]. The ι canonical basis of a based $U_q(\mathfrak{g})$ module is defined using the quasi K -matrix [3, Theorem E], which is an element of the larger ambient quantum group $U_q(\mathfrak{g})$ [3, Proposition C].

It would be very interesting to give an algebraic description of the devil’s divided power basis obtained from our approach to the categorification of the ι quantum group $U_{-q^2}^\iota(\mathfrak{so}_m)$. One might imagine defining an ι canonical basis for non-classical representations from Remark 1.14, then describing the ι canonical basis of $U_{-q^2}^\iota(\mathfrak{so}_m)$ as the elements which are compatible with the ι canonical basis in the non-classical representations. However, since non-classical representations are not restricted from $U_{-q^2}(\mathfrak{sl}_m)$, it is unclear how to mimic Bao and Wang’s construction.

Although Bao–Wang’s theory of ι canonical bases is developed for all Satake diagrams, until recently there was only one quantum symmetric pair admitting a categorification: the quasi-split quantum symmetric pair corresponding to (A_{2n}, t) , where t is the non-trivial Dynkin diagram automorphism [2]. In very recent work [9, 10], Brundan, Wang, and Webster found an approach to categorifying the based algebras in Example 1.23 using the nilBrauer category. Evidently, our equivariant categorification of the (A_1, id) case is not naïvely related to the nilBrauer category, as they categorify different bases of the same algebra. The resolution to the problem posed in Remark 1.24 may clarify whether there is a categorical connection with the nilBrauer category.

In another direction, recall that the Drinfeld–Jimbo quantum groups possess quantum Weyl group elements [28, Chapter 8] that generate an action of the corresponding braid group on finite-dimensional representations. Similarly, it is expected that there exists an ι quantum Weyl group corresponding to the relative Weyl group of the associated Satake diagram. There has been extensive progress defining the ι quantum analogue of the braid group action on ι quantum groups themselves [70, Section 7]. Further, there is a proposal for the ι quantum Weyl group elements in $U_q^\iota(\mathfrak{so}_2)$ given in [77, Section 16.3]. These elements would act on modules for the ι quantum group so that the ι quantum braid group action on the ι quantum group is “conjugation” by the ι quantum Weyl group elements.

In our setting, the decategorification of the equivariant Rickard complexes provide natural candidates for the ι quantum Weyl group elements. We propose the following.

Definition 1.25. The *devil's quantum Weyl group generators* in $U_{-q^2}^\iota(\mathfrak{so}_m)$ are the formal expressions

$${}^\iota T_i := \sum_{k \geq 0} q^{-k} \mathbf{x}_i^{(k)}$$

for $1 \leq i \leq m - 1$.

As a consequence of Theorem 1.8, the elements ${}^\iota T_i$ act invertibly in any representation of $U_{-q^2}^\iota(\mathfrak{so}_m)$ that contains $I^{>N}$ in its kernel for some $N > 0$. One can combine Theorem 1.8, equation (1.10), and Wenzl's Theorem 1.13 to yield the following.

Theorem 1.26 (Theorem 4.39). The surjective algebra homomorphism in Theorem 1.13 is such that

$${}^\iota T_i \mapsto q^{-n/2} \text{id}_S^{\otimes(i-1)} \otimes R_{S,S} \otimes \text{id}_S^{\otimes(m-i-1)}.$$

Remark 1.27. One can view Theorem 1.26 as a type B analogue of a well-known family of results in type A that relate the quantum Weyl group elements with the R -matrix via Howe duality; see [69, Theorem 6.5] and [15, Corollary 6.2.3].

We now outline the remainder of the paper. First, we point out that the majority of our results (e.g. all those explicitly stated thus far) are contained in:

- Section 4, where we study quantum \mathfrak{so}_{2n+1} representation theory,
- Section 6, where we introduce an involution on a 2-subcategory of categorified quantum \mathfrak{gl}_m ,
- Section 8, where we study the equivariant category for this involution, and
- Section 10, where we define and study our link homology.

Each of these sections is immediately preceded by a section developing pertinent background (on quantum groups in §3, on their categorifications in §5, on decompositions/equivariant categories in §7, and on type A link homology in §9). The reader comfortable with this background may focus on the even-numbered sections and backfill the background material as needed. (That said, some of our treatment of the background is novel, and these sections do contain some new concepts/results.) We begin in §2 with our categorical conventions.

Acknowledgements. We thank William Ballinger, Jon Brundan, Lily Gergle, Jiuzu Hong, Peter McNamara, Alistair Savage, and Haihan Wu for helpful discussions. We also thank our once (and hopefully future) collaborator Logan Tatham, as aspects of this work grew out of our previous collaboration with him.

Funding. E.B. was partially supported by the NSF MSPRF-2202897 and the University of Oregon's Lokey Doctoral Fellowship. B.E. was partially supported by NSF grants DMS-2201387 and DMS-2039316. D.R. was partially supported by NSF CAREER grant DMS-2144463 and Simons Collaboration Grant 523992.

2. CONVENTIONS

2.1. The base ring \mathbb{K} . In this paper we use constructions at two different categorical levels. Downstairs, we have structures at the categorical level of traditional representation theory, e.g. the quantum group $U_q(\mathfrak{g})$, its category of finite-dimensional representations $\mathbf{Rep}(U_q(\mathfrak{g}))$, and algebras of intertwiners (endomorphism spaces in $\mathbf{Rep}(U_q(\mathfrak{g}))$). Upstairs, we have structures arising in categorical representations theory, e.g. the categorified quantum group $\mathcal{U}_q(\mathfrak{gl}_m)$ and related subcategories and quotients.

Downstairs⁸, our choice of scalars will not be particularly significant, provided the variable q is generic. Typically, we will use $\mathbb{C}(q)$ or at times $\mathbb{C}(q^{\frac{1}{d}})$ for an appropriate integer d ; the essential feature here is that some constructions in the representation theory of $U_q(\mathfrak{g})$ require the invertibility of certain quantum integers and the existence of certain fractional powers of q . (E.g. in type B , we require $[2]^{-1}$ and $q^{\frac{1}{2}}$.)

On the other hand, there are more subtleties for our choice of base ring upstairs, which we henceforth always denote by \mathbb{K} . The categorified quantum group $\mathcal{U}_q(\mathfrak{gl}_m)$ will be a \mathbb{K} -linear category and our categorified link invariants will be \mathbb{K} -modules. The reader who so desires may set $\mathbb{K} = \mathbb{C}$ and ignore any further technical discussion. However, there is general interest both in developing structures in categorical representation theory and in defining link homologies integrally (in the present setting, $\mathbb{Z}[\frac{1}{2}]$ is more appropriate) or over other fields. This leads to a number of technicalities, due both to subtleties when dealing with equivariant categories and to the choice of \mathbb{K} in the existing literature.

We now mention some of these technicalities. The category $\mathcal{U}_q(\mathfrak{gl}_m)$, which plays a central role for us upstairs, can be defined for any commutative ring \mathbb{K} . However, some common assertions about $\mathcal{U}_q(\mathfrak{gl}_m)$ (e.g. that certain objects are indecomposable) assume that \mathbb{K} is an integral domain or is local. Additionally, most literature concerning the Grothendieck group of $\mathcal{U}_q(\mathfrak{gl}_m)$ assumes that \mathbb{K} is a field (however, note the exception [35]). Further, when dealing with equivariant categories associated with involutions, it will be essential that 2 is invertible in \mathbb{K} . Finally, the results in the literature that construct braid group actions in the context of categorical representation theory are generally only proved under the assumption that \mathbb{K} is a field (although it is folklore that they hold integrally).

Although the goal at the outset of this project was to define the titular spin link homology, along the way we established results in categorical representation theory that we believe are of independent interest. Given that our results speak to a number of different audiences, we have attempted to work (and to set up for future work) over as general of a commutative ring \mathbb{K} as possible. We have attempted (and hopefully succeeded!) to clearly indicate what is being assumed about \mathbb{K} in each section of the paper, and we often reemphasize what is being assumed of \mathbb{K} in most of our “major” results. In particular, much of our work concerning the categorified quantum group $\mathcal{U}_q(\mathfrak{gl}_m)$ is done over a general integral domain, as are our decategorification results when $m = 2$. When considering equivariant categories associated with involutions, we will additionally impose the condition that 2 is invertible (in order to diagonalize $\mathbb{Z}/2$ -representations). Finally, we will assume that \mathbb{K} is a field (in which $2 \neq 0$) for certain decategorification results/conjectures, when dealing with the categorified braiding, and ultimately when defining our categorified link invariant.

To the last point, although we work over a field when defining spin link homology, the construction can in principle be carried out over the ring $\mathbb{K} = \mathbb{Z}[\frac{1}{2}]$. This would require:

- (1) A proof that the (bounded) Rickard complexes appearing in Definition 9.11 are invertible over \mathbb{Z} ,
- (2) a proof that the Rickard complexes braid over \mathbb{Z} , and
- (3) a subsequent adaptation of the proofs of Propositions 10.25, 10.26, and 10.27 to the integral setting.

We note that the first two items above are generally accepted folklore in the link homology community, but (to our knowledge) a proof⁹ has not appeared in the literature.

Meanwhile, we can say the most about the Grothendieck group of various (2-)categories considered under additional assumptions on \mathbb{K} . Assuming \mathbb{K} is a field of characteristic zero, we can use results

⁸All the algebras downstairs can be defined over more general rings. The resulting algebras have interesting modular representation theory, as well as connection to 3-manifold invariants. However, the typical approach to categorification (realizing the algebra as the Grothendieck group of an additive/triangulated category) means that downstairs the algebras are always in characteristic zero and generic q .

⁹We encourage someone to fill this gap in the literature.

of Webster [71, 72] which state that $\mathcal{U}_q(\mathfrak{gl}_m)$ is mixed. Further, when \mathbb{K} is algebraically closed, we can classify the indecomposable objects in the equivariant category of a mixed category. It is under these assumptions that we can say the most about weighted Grothendieck groups, thus we expect the decategorification conjectures (for $m > 2$) in Sections 1.4 and 1.6 to be most-easily accessible under this assumption. Nevertheless, we are able to establish some decategorification results over more general \mathbb{K} .

2.2. Categorical conventions. We now record our conventions for various categorical constructions which take place “upstairs,” in the terminology of §2.1. We state these results for categories, but remark that all constructions in this section extend to 2-categories by applying them to the constituent Hom-categories. For us, “2-category” will always mean weak 2-category (also called bicategories in the literature), although some that appear will be strict. As mentioned above, the base ring for these constructions is \mathbb{K} .

2.2.1. (Graded) linear categories. A \mathbb{K} -linear category will mean a category enriched in \mathbb{K} -modules; such a category is additive if and only if it admits all finite coproducts, which are necessarily biproducts. We refer to biproducts as *direct sums*. We can always pass from a \mathbb{K} -linear category to an additive \mathbb{K} -linear category by formally adjoining finite direct sums. Even in a \mathbb{K} -linear category which is not additive we can discuss direct sums, which may or may not exist.

Analogously, a \mathbb{Z} -graded \mathbb{K} -linear category is a category enriched in the category of \mathbb{Z} -graded \mathbb{K} -modules (which itself has morphisms the *degree-zero* linear maps). Throughout, we will denote various grading shift functors (that are not of a homological nature) by powers of \mathbf{q} . For example, given a \mathbb{Z} -graded \mathbb{K} -module V and $k \in \mathbb{Z}$, $\mathbf{q}^k V$ denotes the \mathbb{Z} -graded \mathbb{K} -module which, in degree m , agrees with V in degree $m - k$. Given a finitely generated free graded \mathbb{K} -module (e.g. a finite-dimensional graded vector space), we let $\dim_{\mathbf{q}}(V) \in \mathbb{Z}_{\geq 0}[q^{\pm}]$ denote its graded dimension. Our conventions therefore imply that $\dim_{\mathbf{q}}(\mathbf{q}^k V) = q^k \dim_{\mathbf{q}}(V)$.

The following standard construction allows to pass from a \mathbb{Z} -graded \mathbb{K} -linear category to an additive \mathbb{Z} -graded \mathbb{K} -linear category that is equipped with a grading shift autoequivalence.

Definition 2.1. Let \mathcal{A} be a \mathbb{Z} -graded \mathbb{K} -linear category. The *\mathbb{Z} -additive closure* of \mathcal{A} is the category whose objects are formal expressions $\bigoplus_{i \in I} \mathbf{q}^{k_i} X_i$, where I is a finite set, $k_i \in \mathbb{Z}$, and X_i are objects of \mathcal{A} . Morphisms are given by matrices:

$$\mathrm{Hom}\left(\bigoplus_{i \in I} \mathbf{q}^{k_i} X_i, \bigoplus_{j \in J} \mathbf{q}^{\ell_j} Y_j\right) := \bigoplus_{i \in I, j \in J} \mathbf{q}^{\ell_j - k_i} \mathrm{Hom}(X_i, Y_j).$$

Next, for an additive category \mathcal{A} we let $K_0(\mathcal{A})$ denote its (split) Grothendieck group. This is the quotient of the free abelian group generated by isomorphism classes $[X]$ of objects $X \in \mathcal{A}$ by the relation $[X \oplus Y] = [X] + [Y]$. An additive category is *Krull–Schmidt* if each object can be decomposed as a finite sum of objects with local endomorphism rings (graded local, in the event that \mathcal{A} is graded). In such a category, $K_0(\mathcal{A})$ is free abelian with basis the classes of non-isomorphic indecomposable objects. If \mathcal{A} is \mathbb{Z} -graded and is equipped with a grading shift autoequivalence (e.g. if we’re in the setting of Definition 2.1), we can endow $K_0(\mathcal{A})$ with the structure of a $\mathbb{Z}[q^{\pm}]$ -module via the relation $q[X] := [\mathbf{q}X]$.

A pre-additive category \mathcal{A} is *Karoubian* if every idempotent endomorphism in \mathcal{A} splits. Any Krull–Schmidt category is Karoubian, and in categories enriched in (graded) finitely dimensional \mathbb{K} -vector spaces (over a field \mathbb{K}), Karoubian implies Krull–Schmidt; see e.g. [22, Theorem 11.53]. If \mathcal{A} is not necessarily Karoubian, we can pass to its *Karoubi envelope* $\mathrm{Kar}(\mathcal{A})$ wherein objects are idempotent endomorphisms. The category $\mathrm{Kar}(\mathcal{A})$ is always Karoubian, and is equivalent to \mathcal{A} in the event that \mathcal{A} is itself Karoubian.

2.2.2. *Categories of complexes.* Next, we establish our conventions for homological algebra. We emphasize at the outset that all complexes considered in this paper will be bounded.

Given a \mathbb{K} -linear category \mathcal{A} , we let $\mathcal{A}[\mathfrak{t}^\pm]$ denote the category of (cohomologically indexed) finite sequences in \mathcal{A} . Explicitly, objects are sequences $X = (X_i)_{i \in \mathbb{Z}}$ with $X_i = 0$ for all but finitely many $i \in \mathbb{Z}$, and morphisms are given by

$$\mathrm{Hom}_{\mathcal{A}[\mathfrak{t}^\pm]}(X, Y) := \bigoplus_{k \in \mathbb{Z}} \mathrm{Hom}_{\mathcal{A}[\mathfrak{t}^\pm]}^k(X, Y), \quad \mathrm{Hom}_{\mathcal{A}[\mathfrak{t}^\pm]}^k(X, Y) := \prod_{i \in \mathbb{Z}} \mathrm{Hom}_{\mathcal{A}}^k(X_i, Y_{i+k}).$$

This category is enriched in \mathbb{Z} -graded \mathbb{K} -modules. By slight abuse of notation, we will denote objects in $\mathcal{A}[\mathfrak{t}^\pm]$ by $\bigoplus_{i \in \mathbb{Z}} \mathfrak{t}^i X_i$. Here, in contrast to Definition 2.1, we view the grading (shift) \mathfrak{t} as having a homological nature.

Definition 2.2. Let \mathcal{A} be a \mathbb{K} -linear category. A bounded *chain complex* over \mathcal{A} is a pair (X, d_X) with $X \in \mathcal{A}[\mathfrak{t}^\pm]$ and $d_X \in \mathrm{End}_{\mathcal{A}[\mathfrak{t}^\pm]}^1(X)$ such that $d_X^2 = 0$.

The *dg category of bounded chain complexes* over \mathcal{A} is the category $\mathcal{C}(\mathcal{A})$ with objects bounded chain complexes (X, d_X) over \mathcal{A} and morphism spaces the complexes of \mathbb{K} -modules

$$(2.1) \quad \mathrm{Hom}_{\mathcal{C}(\mathcal{A})}((X, d_X), (Y, d_Y)) := \left(\bigoplus_{k \in \mathbb{Z}} \mathfrak{t}^k \mathrm{Hom}_{\mathcal{A}[\mathfrak{t}^\pm]}^k(X, Y), D \right)$$

where the differential D is defined by $D(f) = d_Y \circ f - (-1)^{|f|} f \circ d_X$.

One benefit of working with $\mathcal{C}(\mathcal{A})$ is that other familiar categories of chain complexes are easily recovered from it, so it provides a unified setting for studying homological algebra. The (usual) category $\mathrm{Ch}(\mathcal{A})$ of bounded chain complexes has the same objects as $\mathcal{C}(\mathcal{A})$ and morphisms

$$\mathrm{Hom}_{\mathrm{Ch}(\mathcal{A})}((X, d_X), (Y, d_Y)) := \ker(D: \mathrm{Hom}_{\mathcal{A}[\mathfrak{t}^\pm]}^0(X, Y) \rightarrow \mathrm{Hom}_{\mathcal{A}[\mathfrak{t}^\pm]}^1(X, Y)),$$

while the homotopy category $\mathcal{K}(\mathcal{A})$ of bounded chain complexes has the same objects as $\mathcal{C}(\mathcal{A})$ and morphisms given by zeroth homology:

$$\mathrm{Hom}_{\mathcal{K}(\mathcal{A})}((X, d_X), (Y, d_Y)) := H^0(\mathrm{Hom}_{\mathcal{C}(\mathcal{A})}((X, d_X), (Y, d_Y))).$$

It follows that $\mathrm{Ch}(\mathcal{A})$ is a (non-full) subcategory of $\mathcal{C}(\mathcal{A})$, while $\mathcal{K}(\mathcal{A})$ is a quotient of $\mathrm{Ch}(\mathcal{A})$.

Further unpacking the definitions, we see that morphisms in $\mathrm{Ch}(\mathcal{A})$ are *chain maps*: morphisms $f \in \mathrm{Hom}_{\mathcal{A}[\mathfrak{t}^\pm]}^0(X, Y)$ such that $d_Y \circ f = f \circ d_X$. Similarly, morphisms in $\mathcal{K}(\mathcal{A})$ are homotopy classes of chain maps: the quotient of the space of chain maps by those that are *null-homotopic*, i.e. those that can be written as $D(h) = d_Y \circ h + h \circ d_X$ for $h \in \mathrm{Hom}_{\mathcal{A}[\mathfrak{t}^\pm]}^{-1}(X, Y)$. We will denote the corresponding equivalence relation on chain maps by \sim .

A morphism $f \in \mathrm{Hom}_{\mathcal{C}(\mathcal{A})}^0((X, d_X), (Y, d_Y))$ in $\mathcal{C}(\mathcal{A})$ is called a *homotopy equivalence* provided there exists $g \in \mathrm{Hom}_{\mathcal{C}(\mathcal{A})}^0((Y, d_Y), (X, d_X))$ such that $\mathrm{id}_X \sim g \circ f$ and $\mathrm{id}_Y \sim f \circ g$. We will write $(X, d_X) \simeq (Y, d_Y)$ if there exists a homotopy equivalence between these complexes, and refer to such complexes as *homotopy equivalent*. Note that homotopy equivalent complexes are isomorphic in $\mathcal{K}(\mathcal{A})$. In the case that \mathcal{A} is abelian, we let $\mathcal{D}(\mathcal{A})$ denote the *bounded derived category* of \mathcal{A} , which is the localization of $\mathcal{K}(\mathcal{A})$ at the class of quasi-isomorphisms.

Finally, note that we can consider $\mathcal{C}(\mathcal{A})$ in the event that \mathcal{A} is itself \mathbb{Z} -graded. In this case, the Hom-spaces (2.1) are complexes of \mathbb{Z} -graded \mathbb{K} -modules, hence are $\mathbb{Z} \times \mathbb{Z}$ -graded, while the Hom-spaces in $\mathrm{Ch}(\mathcal{A})$ are only \mathbb{Z} -graded (by \mathfrak{q} -degree). We require homotopy equivalences and quasi-isomorphisms to have \mathfrak{q} -degree zero, so this \mathbb{Z} -grading is inherited by $\mathcal{K}(\mathcal{A})$ and $\mathcal{D}(\mathcal{A})$.

2.2.3. *Over a field.* When \mathbb{K} is a field, we denote the category of \mathbb{K} -vector spaces by $\text{Vect}_{\mathbb{K}}$. Given an abelian group Γ , we denote by $\text{Vect}_{\mathbb{K}}^{\Gamma}$ the category of Γ -graded vector spaces and degree-zero linear maps.

We let $\text{sVect}_{\mathbb{K}}$ denote the category of super vector spaces. The objects in this category are $\mathbb{Z}/2$ -graded vector spaces and morphisms are degree-zero linear maps. As \mathbb{K} -linear monoidal categories, $\text{sVect}_{\mathbb{K}}$ and $\text{Vect}_{\mathbb{K}}^{\mathbb{Z}/2}$ are indistinguishable, but $\text{sVect}_{\mathbb{K}}$ has a non-trivial symmetric monoidal structure. We refer to the categorical dimension¹⁰ computed in the symmetric monoidal category $\text{sVect}_{\mathbb{K}}$ as the *super dimension*. The latter is characterized by an even line having super dimension $+1$ and an odd line having super dimension -1 .

In $\text{sVect}_{\mathbb{K}}$, we will denote grading shift (in super degree) by \mathbf{s} , since we also consider Γ -graded super vector spaces $\text{sVect}_{\mathbb{K}}^{\Gamma}$ and wish to reserve \mathbf{q} to denote grading shift in that setting. The super dimension of a \mathbb{Z} -graded super vector space $V = V_0 \oplus V_1$ is equal to

$$(2.2) \quad \dim_{\mathbf{q}}(V) = \dim_{\mathbf{q}}(V_0) - \dim_{\mathbf{q}}(V_1).$$

Note that we therefore use $\dim_{\mathbf{q}}$ to denote both the graded dimension of a graded vector space and the graded super dimension of a graded super vector space. There is no cause for confusion, since we can view vector spaces as super vector spaces whose odd degree part is zero.

3. BACKGROUND ON QUANTUM GROUPS AND LINK INVARIANTS

3.1. **The quantum group.** We recall background on the quantum group $U_q(\mathfrak{g})$ associated to a finite-dimensional simple complex Lie algebra \mathfrak{g} , for the purpose of establishing conventions and notation. Given such a Lie algebra \mathfrak{g} , there is an associated root system $\Phi(\mathfrak{g})$ and we let

$$\{\alpha_i\}_{i \in I} \subset \Phi_+(\mathfrak{g}) \subset \Phi(\mathfrak{g})$$

denote the subsets of simple roots and positive roots, respectively. The simple roots are indexed by the finite set I of nodes in the corresponding Dynkin diagram, and the cardinality $|I|$ of this subset is the rank of \mathfrak{g} . If $W_{\mathfrak{g}}$ is the corresponding Weyl group (generated by reflections in the root hyperplanes), then we denote the standard $W_{\mathfrak{g}}$ -invariant inner product on $\alpha, \alpha' \in \mathbb{R}\Phi(\mathfrak{g})$ by (α, α') , which is uniquely determined by the condition that $(\alpha, \alpha) = 2$ when $\alpha \in \Phi(\mathfrak{g})$ is a root of minimal length (i.e. a short root). For $i, j \in I$, we will occasionally abbreviate by writing $i \cdot j := (\alpha_i, \alpha_j)$. The coroot associated to a root α is $\alpha^{\vee} := \frac{2}{(\alpha, \alpha)}\alpha$, therefore the Cartan matrix $(a_{ij})_{i, j \in I}$ is given by

$$a_{ij} := (\alpha_i^{\vee}, \alpha_j) = 2 \frac{(\alpha_i, \alpha_j)}{(\alpha_i, \alpha_i)} = 2 \frac{i \cdot j}{i \cdot i}.$$

Write $\rho, \rho^{\vee} \in \mathbb{R}\Phi(\mathfrak{g})$ to denote the elements such that $(\rho, \alpha_i^{\vee}) = 1$ and $(\rho^{\vee}, \alpha_i) = 1$, for all $i \in I$.

Let q be an indeterminate. Given $i \in I$ and $m \in \mathbb{Z}_{\geq 0}$, set

$$q_i := q^{\frac{(\alpha_i, \alpha_i)}{2}}, \quad [m]_i := [m]_{q_i} := \frac{q_i^m - q_i^{-m}}{q_i - q_i^{-1}}$$

so if α_i is a short simple root then $[m]_i$ agrees with the usual quantum integer $[m] := \frac{q^m - q^{-m}}{q - q^{-1}}$.

Definition 3.1. Let $U_q(\mathfrak{g})$ be the $\mathbb{C}(q)$ -algebra generated by elements $e_i, f_i, k_i^{\pm 1}$ for $i \in I$, subject to the following relations:

- (1) $k_i k_i^{-1} = 1 = k_i^{-1} k_i$,
- (2) $k_i k_j = k_j k_i$,
- (3) $k_i e_j = q^{(\alpha_i, \alpha_j)} e_j k_i$,

¹⁰Computed in a general rigid symmetric monoidal category as the scalar multiple of the identity in the following composition of the braiding and the evaluation/coevaluation: $\mathbb{1} \rightarrow X \otimes X^* \rightarrow X^* \otimes X \rightarrow \mathbb{1}$.

- (4) $k_i f_j = q^{-(\alpha_i, \alpha_j)} f_j k_i$,
- (5) $e_i f_i = f_i e_i + \frac{k_i - k_i^{-1}}{q_i - q_i^{-1}}$,
- (6) $e_i f_j = f_j e_i$ if $i \neq j$, and
- (7) the “quantum Serre relations” (of e.g. [28, 4.3 (R5) and (R6)]).

Define the divided powers by $e_i^{(a)} := \frac{e_i^a}{[a]_i!}$ and $f_i^{(a)} := \frac{f_i^a}{[a]_i!}$.

The $\mathbb{C}(q)$ -algebra $U_q(\mathfrak{g})$ is a Hopf algebra and, following the conventions in [16, 66], the structure maps are given on generators as follows:

- (1) $\Delta(e_i) = e_i \otimes k_i + 1 \otimes e_i$, $\Delta(f_i) = f_i \otimes 1 + k_i^{-1} \otimes f_i$, and $\Delta(k_i^\pm) = k_i^\pm \otimes k_i^\pm$.
- (2) $\mathbf{S}(e_i) = -e_i k_i^{-1}$, $\mathbf{S}(f_i) = -k_i f_i$, and $\mathbf{S}(k_i^\pm) = k_i^\mp$.
- (3) $\epsilon(e_i) = 0$, $\epsilon(f_i) = 0$, and $\epsilon(k_i) = 1$.

In the present paper, we will be interested in the following quantum groups. In both cases, we let $\{\epsilon_i\}_{i=1}^N$ be the standard basis for the vector space \mathbb{R}^N .

Example 3.2. Let $\mathfrak{g} = \mathfrak{sl}_N$, the simple Lie algebra of type A_{N-1} . In this case $I = \{1, \dots, N-1\}$ and $\Phi(\mathfrak{sl}_N)$ is the root system with simple roots:

$$\{\alpha_i = \epsilon_i - \epsilon_{i+1}\}_{i=1}^{N-1} \subset \mathbb{R}^N.$$

In this case, $\mathbb{R}\Phi(\mathfrak{sl}_N)$ is a hyperplane in \mathbb{R}^N and the inner product is given by restricting the standard inner product $(\epsilon_i, \epsilon_j) = \delta_{ij}$ on \mathbb{R}^N to this hyperplane.

Example 3.3. Let $\mathfrak{g} = \mathfrak{so}_{2n+1}$, the simple Lie algebra of type B_n . In this case $I = \{1, \dots, n\}$ and $\Phi(\mathfrak{so}_{2n+1})$ is the root system with simple roots:

$$\{\alpha_i = \epsilon_i - \epsilon_{i+1}\}_{i=1}^{n-1} \cup \{\alpha_n = \epsilon_n\} \subset \mathbb{R}^n.$$

In this case, $\mathbb{R}\Phi(\mathfrak{so}_{2n+1}) = \mathbb{R}^n$, and the inner product $(\epsilon_i, \epsilon_j) = 2\delta_{ij}$ is a multiple of the standard inner product on \mathbb{R}^n .

Later, we will also be interested in the quantum group $\dot{U}_q(\mathfrak{gl}_m)$ associated with the (non-simple) Lie algebra \mathfrak{gl}_m . See §5.1 for the relevant definition.

3.2. Representations of $U_q(\mathfrak{g})$. We let $\mathbf{Rep}(U_q(\mathfrak{g}))$ denote the category of (type 1) finite-dimensional representations of $U_q(\mathfrak{g})$. Every representation $V \in \mathbf{Rep}(U_q(\mathfrak{g}))$ admits a weight space decomposition $V \cong \bigoplus_{\mu \in X(\mathfrak{g})} V_\mu$ indexed by the weight lattice $X(\mathfrak{g})$. Recall that the latter consists of all $\mu \in \mathbb{R}\Phi(\mathfrak{g})$ so that $(\alpha^\vee, \mu) \in \mathbb{Z}$. When $v \in V_\mu$, we will write $\text{wt}(v) = \mu$. Further, $\mathbf{Rep}(U_q(\mathfrak{g}))$ is semisimple, with irreducibles $V(\lambda)$ indexed by dominant weights $\lambda \in X_+(\mathfrak{g})$.

Since $U_q(\mathfrak{g})$ is a Hopf algebra, $\mathbf{Rep}(U_q(\mathfrak{g}))$ is a rigid monoidal category. Moreover, this category is pivotal and braided. We now review the latter in depth, since it will play an important role in our considerations. For this, we must work with representations over the field $\mathbb{C}(q^{\frac{1}{d}})$, where d is the index¹¹ of the root lattice in the weight lattice, which we do for the duration.

Given $V, W \in \mathbf{Rep}(U_q(\mathfrak{g}))$, let $\text{flip}_{V,W}$ denote the $\mathbb{C}(q^{\frac{1}{d}})$ -linear map $V \otimes W \rightarrow W \otimes V$ that sends $v \otimes w \mapsto w \otimes v$. The braiding on $\mathbf{Rep}(U_q(\mathfrak{g}))$ is the invertible operator

$$(3.1) \quad R_{V,W} := \text{flip}_{V,W} \circ R: V \otimes W \rightarrow W \otimes V$$

where R is the *universal R-matrix*. We now recall the approach to the later taken in [37], following [30, 68, 66]. For this, we need the following ingredients:

¹¹This value for d always suffices. For certain \mathfrak{g} it is possible to use a smaller value for d ; see e.g. [41] for details. (There, this parameter is denoted D .)

- Let $q^{(\text{wt}(-), \text{wt}(-))}$ denote the operator that acts on weight vectors of the form $v \otimes w \in V_{\mu_1} \otimes W_{\mu_2}$ in a tensor product $V \otimes W$ of finite-dimensional representations by

$$(3.2) \quad q^{(\text{wt}(-), \text{wt}(-))}(v \otimes w) := q^{(\mu_1, \mu_2)} v \otimes w.$$

- Let T_i be the operator that acts on weight vectors $v \in V_\mu$ as

$$(3.3) \quad T_i(v) := \sum_{\substack{a, b \geq 0 \\ b-a = (\alpha_i^\vee, \mu)}} (-q_i)^b e_i^{(a)} f_i^{(b)}(v).$$

The element T_i in (3.3) is equal to Lusztig's quantum Weyl group element $T''_{i,+1}$, in the simplified form given e.g. in [11, Remark 2.1]. These elements satisfy the relations of the type \mathfrak{g} braid group $B_{\mathfrak{g}}$. We have that $\text{wt}(T_i(v)) = s_i(\text{wt}(v))$, where $s_i \in W_{\mathfrak{g}}$ is the corresponding generator for the Weyl group. Given any $w \in W_{\mathfrak{g}}$, we can choose a reduced expression $w = s_{i_1} \cdots s_{i_\ell}$ and set

$$T_w := T_{i_1} \cdots T_{i_\ell}$$

which is well-defined by Matsumoto's theorem [48]. Of particular importance is the operator T_{w_0} that is associated with the longest element $w_0 \in W_{\mathfrak{g}}$.

In [37, Theorem 3], Kirillov–Reshetikhin describe the universal R matrix in terms of the operator T_{w_0} . Their result, in the conventions¹² of [30, Theorem 7.1], is as follows.

Theorem 3.4. $R = q^{(\text{wt}(-), \text{wt}(-))} \circ (T_{w_0}^{-1} \otimes T_{w_0}^{-1}) \circ \Delta(T_{w_0})$

For our considerations in Section 4.2 below, we will use that the operator $(T_{w_0}^{-1} \otimes T_{w_0}^{-1}) \circ \Delta(T_{w_0})$ admits a certain description. For this, let $U_q(\mathfrak{g})^{\geq 0}$ and $U_q(\mathfrak{g})^{\leq 0}$ denote the $\mathbb{C}(q)$ -subalgebras of $U_q(\mathfrak{g})$ generated by $\{e_i, k_i^\pm\}_{i \in I}$ and $\{f_i, k_i^\pm\}_{i \in I}$, respectively. These algebras are graded, respectively by $\mathbb{Z}_{\geq 0} \Phi_+(\mathfrak{g})$ and $\mathbb{Z}_{\geq 0}(-\Phi_+(\mathfrak{g}))$, and for $\mu \in \mathbb{Z}_{\geq 0} \Phi_+(\mathfrak{g})$ we denote the homogeneous components as $U_q(\mathfrak{g})_{\mu}^{\geq 0}$ and $U_q(\mathfrak{g})_{-\mu}^{\leq 0}$. We let $U_q(\mathfrak{g})^{>0}$ and $U_q(\mathfrak{g})^{<0}$ denote the graded $\mathbb{C}(q)$ -subalgebras generated by the $\{e_i\}_{i \in I}$ and $\{f_i\}_{i \in I}$ alone.

Proposition 3.5. There exist elements $\{x_{i,\mu}^+\}_{i=1}^{d_\mu} \subset U_q(\mathfrak{g})^{>0}$ and $\{y_{i,\mu}^-\}_{i=1}^{d_\mu} \subset U_q(\mathfrak{g})^{<0}$ so that

$$(T_{w_0}^{-1} \otimes T_{w_0}^{-1}) \circ \Delta(T_{w_0}) = 1 \otimes 1 + \sum_{\substack{\mu \in \mathbb{Z}_{\geq 0} \Phi_+(\mathfrak{g}) \\ \mu \neq 0}} \sum_{i=1}^{d_\mu} x_{i,\mu}^+ \otimes y_{i,\mu}^-$$

Proof. As noted e.g. in [68, Lemma 3.3.13] the “standard” R -matrix appearing in Theorem 3.4 admits a description of the form

$$q^{(\text{wt}(-), \text{wt}(-))} \circ \left(1 \otimes 1 + \sum_{\substack{\mu \in \mathbb{Z}_{\geq 0} \Phi_+(\mathfrak{g}) \\ \mu \neq 0}} \sum_{i=1}^{d_\mu} x_{i,\mu}^+ \otimes y_{i,\mu}^- \right)$$

with $\{x_{i,\mu}^+\}_{i=1}^{d_\mu} \subset U_q(\mathfrak{g})^{>0}$ and $\{y_{i,\mu}^-\}_{i=1}^{d_\mu} \subset U_q(\mathfrak{g})^{<0}$. The result then follows from Theorem 3.4. \square

¹²In [30], the authors state that they work with Lusztig's $T''_{i,-1}$; however, it is clear from [30, Lemma 5.6] that they in fact work with $T''_{i,+1}$.

3.3. Quantum link polynomials. Our approach to link invariants in the present paper is through the classical Alexander and Markov Theorems. We let $\text{Br}_m = \langle \beta_1, \dots, \beta_{m-1} \mid \beta_i \beta_{i+1} \beta_i = \beta_{i+1} \beta_i \beta_{i+1} \rangle$ denote the (type A) braid group on m strands.

Theorem 3.6 ([1]). Let $\mathcal{L} \subset S^3$ be a link in the 3-sphere. There is a braid β such that the braid closure \mathcal{L}_β is isotopic to \mathcal{L} .

Theorem 3.7 ([47]). Given two braids $\beta \in \text{Br}_m$ and $\beta' \in \text{Br}_{m'}$, their closures \mathcal{L}_β and $\mathcal{L}_{\beta'}$ are isotopic if and only if β' can be obtained from β by a sequence of the following ‘‘Markov moves.’’

(MI) Conjugate β in Br_m .

(MII) Replace $\beta \in \text{Br}_m$ by $\beta_{m+1}^{\pm 1} \beta \in \text{Br}_{m+1}$ or vice-versa.

Namely, we will first find invariants of braids $\beta \mapsto T_\beta \in X$ taking values in a set or category X (possibly with additional structure). We then compose with a function or functor $X \xrightarrow{[-]} A$ such that $[T_{\beta\beta'}] = [T_{\beta'}\beta]$ and $[T_\beta] = [T_{\beta_{m+1}^{\pm 1}\beta}]$. It follows from Theorems 3.6 and 3.7 that $[T_\beta]$ is an invariant of the link \mathcal{L}_β . In practice, we will encounter $[T_\beta]$ that are conjugation invariant, but only satisfy invariance under the second Markov move (MII) up to a sign and a power of q . In this case $[T_\beta]$ will be an invariant of *framed* links, i.e. links endowed with a framing on their normal bundle. It is possible to introduce a renormalization factor to obtain an invariant of unframed links, but we prefer to work in the present setup. See Remarks 3.9 and 3.10 for details.

In this section, we recall the Reshetikhin–Turaev link invariants defined using the quantum group $U_q(\mathfrak{g})$ associated to a finite-dimensional simple complex Lie algebra \mathfrak{g} . Our exposition is adapted to links presented as braid closures; see [58] for the original construction, which treats all link (and more-generally tangle) diagrams.

Let $\beta = \beta_{i_1}^{\epsilon_1} \cdots \beta_{i_d}^{\epsilon_d} \in \text{Br}_m$ for $\epsilon_i = \pm 1$. Given an m -tuple $\vec{\lambda} = (\lambda_1, \dots, \lambda_m)$ of highest weights of finite-dimensional irreducible representations of \mathfrak{g} , this induces a coloring of β , where the strand meeting the k^{th} point at the bottom of β is colored by the weight λ_k . If the crossing $\beta_{i_s}^{\epsilon_s}$ is colored by the pair $(\lambda_{l_s}, \lambda_{r_s})$ at its bottom, then we set

$$R(\beta, \vec{\lambda}) := R_{\lambda_{l_1}, \lambda_{r_1}}^{\epsilon_1} \cdots R_{\lambda_{l_d}, \lambda_{r_d}}^{\epsilon_d}$$

where here $R_{\lambda, \mu}$ denotes the isomorphism $R_{V(\lambda), V(\mu)}: V(\lambda) \otimes V(\mu) \rightarrow V(\mu) \otimes V(\lambda)$ from (3.1), tensored with appropriate identity morphisms.

We say that β is $\vec{\lambda}$ -balanced if the strand meeting the k^{th} point at the top of β is also colored by λ_k . In this case, $R(\beta, \vec{\lambda})$ is a braid invariant valued in $\text{End}_{U_q(\mathfrak{g})}(V(\lambda_1) \otimes \cdots \otimes V(\lambda_d))$. Since $\mathbf{Rep}(U_q(\mathfrak{g}))$ is pivotal, composing with the quantum trace Tr_q (i.e. taking a closure in the associated graphical calculus) yields an invariant of braid conjugacy classes taking values in $\mathbb{C}(q^{\frac{1}{d}})$.

Theorem 3.8 ([58, Section 6.1]). The braid conjugacy invariant

$$(3.4) \quad P_{\mathfrak{g}}(\mathcal{L}_\beta^{\vec{\lambda}}) := \text{Tr}_q(R(\beta, \vec{\lambda}))$$

descends to give an invariant of the framed link \mathcal{L}_β .

A priori, $P_{\mathfrak{g}}(\mathcal{L}_\beta^{\vec{\lambda}})$ takes values in $\mathbb{C}(q^{\frac{1}{d}})$; however, it is possible to show¹³ that $P_{\mathfrak{g}}(\mathcal{L}_\beta^{\vec{\lambda}}) \in \mathbb{Z}[q^{\frac{1}{d}}, q^{-\frac{1}{d}}]$. We thus refer to $P_{\mathfrak{g}}(\mathcal{L}_\beta^{\vec{\lambda}})$ as the $\vec{\lambda}$ -colored $U_q(\mathfrak{g})$ link polynomial. When $\vec{\lambda} = (\lambda, \dots, \lambda)$, we will denote this invariant simply by $P_{\mathfrak{g}}(\mathcal{L}_\beta^\lambda)$

¹³An even stronger result in [41] shows that there exists $r(\mathcal{L}) \in \mathbb{Q}$ depending only on the linking matrix of \mathcal{L} such that $q^{r(\mathcal{L})} P_{\mathfrak{g}}(\mathcal{L}^{\vec{\lambda}}) \in \mathbb{Z}[q^{\pm 2}]$.

Remark 3.9. As defined in (3.4), $P_{\mathfrak{g}}(\mathcal{L}_{\beta}^{\vec{\lambda}})$ changes by a factor of $\nu_{\lambda} = (-1)^{(2\lambda, \rho^{\vee})} q^{\pm(\lambda, \lambda + 2\rho)}$ upon applying a positive/negative Reidemeister I move on a λ -colored component, i.e. graphically:

$$(3.5) \quad P_{\mathfrak{g}} \left(\text{positive crossing} \right) = (-1)^{(2\lambda, \rho^{\vee})} q^{(\lambda, \lambda + 2\rho)} P_{\mathfrak{g}} \left(\left| \lambda \right. \right) \quad \text{and} \quad P_{\mathfrak{g}} \left(\text{negative crossing} \right) = (-1)^{(2\lambda, \rho^{\vee})} q^{-(\lambda, \lambda + 2\rho)} P_{\mathfrak{g}} \left(\left| \lambda \right. \right).$$

It follows that two diagrams for a given link yield the same value of $P_{\mathfrak{g}}(\mathcal{L}_{\beta}^{\vec{\lambda}})$ if and only if they have the same writhe ($= \#\{\text{positive crossings}\} - \#\{\text{negative crossings}\}$).

Remark 3.10. Let $P(\beta)$ be a braid-conjugacy invariant which satisfies the \pm -Markov II moves only up to a factor of $c^{\pm 1}$ for some invertible scalar c . For example, $P(\beta) = P_{\mathfrak{g}}(\mathcal{L}_{\beta}^{\vec{\lambda}})$ satisfies this property with $c = (-1)^{(2\lambda, \rho^{\vee})} q^{(\lambda, \lambda + 2\rho)}$ by (3.5). It is well known that $P(\beta)$ then descends to an invariant of framed links. Since an appropriate citation has eluded us, we provide a quick sketch. Let, $\epsilon(\beta)$ denote the braid exponent, then $c^{-\epsilon(\beta)} P(\beta)$ is invariant under the Markov moves MI and MII, therefore is an invariant of unframed links. Since the braid exponent equals the writhe of the braid closure, two braid closures have the same value for $P(\beta)$ if they have the same writhe, i.e. if they have the same (blackboard) framing.

Remark 3.11. The link polynomials $P_{\mathfrak{g}}(\mathcal{L}_{\beta}^{\vec{\lambda}})$ are defined using the braiding on $\mathbf{Rep}(U_q(\mathfrak{g}))$. The latter is essentially unique up to conventions [31], but not literally unique since e.g. the inverse to a braiding is again a braiding. Since we wish to use Theorem 3.4 and Proposition 3.5 below, our conventions for the braiding/ R -matrix are taken from [30, 66] and they lead to behavior described in Remark 3.9. However, it is common in the link homology literature to work with link polynomials that correspond to the inverse choice of R -matrix. We denote these link polynomials by $\overline{P}_{\mathfrak{g}}(\mathcal{L}_{\beta}^{\vec{\lambda}})$ and observe that

$$(3.6) \quad \overline{P}_{\mathfrak{g}}(\mathcal{L}_{\beta}^{\vec{\lambda}})(q) = P_{\mathfrak{g}}(\mathcal{L}_{\beta}^{\vec{\lambda}})(q^{-1}) = P_{\mathfrak{g}}(\mathfrak{m}\mathcal{L}_{\beta}^{\vec{\lambda}})(q) = P_{\mathfrak{g}}(\mathcal{L}_{\mathfrak{m}\beta}^{\vec{\lambda}})(q)$$

where here $\mathfrak{m}\mathcal{L}$ and $\mathfrak{m}\beta$ denote the mirror link/braid (obtained by switching over/under information at all crossings). Consequently, we have

$$\overline{P}_{\mathfrak{g}} \left(\text{positive crossing} \right) = (-1)^{(2\lambda, \rho^{\vee})} q^{-(\lambda, \lambda + 2\rho)} \overline{P}_{\mathfrak{g}} \left(\left| \lambda \right. \right) \quad \text{and} \quad \overline{P}_{\mathfrak{g}} \left(\text{negative crossing} \right) = (-1)^{(2\lambda, \rho^{\vee})} q^{(\lambda, \lambda + 2\rho)} \overline{P}_{\mathfrak{g}} \left(\left| \lambda \right. \right).$$

In Section 4.5, we further study the invariant $P_{\mathfrak{g}}(\mathcal{L}_{\beta}^{\vec{\lambda}})$ for $\mathfrak{g} = \mathfrak{so}_{2n+1}$ and when each entry of $\vec{\lambda}$ corresponds to the spin representation. Later, in Section 9, we consider the $\mathfrak{g} = \mathfrak{sl}_N$ case of the invariant $P_{\mathfrak{g}}(\mathcal{L}_{\beta}^{\vec{\lambda}})$ in more detail, and review its categorification via \mathfrak{sl}_N Khovanov–Rozansky homology. This forms the foundation for our categorification of the invariant from §4.5.

4. TYPE B INTERTWINERS

In this section, we study representations of $U_q(\mathfrak{so}_{2n+1})$ in depth. In this case, there is a distinguished *spin representation* $S \in \mathbf{Rep}(U_q(\mathfrak{so}_{2n+1}))$, and our main result is the construction of a “canonical basis” for the endomorphism algebra $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$. Although we discuss this material before our categorification results that in appear in later sections, this presentation is somewhat ahistorical: the existence of this basis (and the corresponding structure coefficients for multiplication) was initially suggested by the techniques appearing in §8.

In order to state this result precisely, we need the following, which we affectionately refer to as the *devil’s product*.

Definition 4.1. Let $m \leq n \in \mathbb{Z}_{\geq 0}$ and set

$$“[m][n]” = “[n][m]” := \sum_{i=0}^{m-1} (-1)^i [n + m - 2i - 1].$$

In other words, the expression

$$“[m][n]” = [n + m - 1] - [n + m - 3] + [n + m - 5] - \cdots + (-1)^{m-1} [n - m + 1]$$

is obtained from the usual expression for the product of quantum integers

$$[m][n] = \sum_{i=0}^{m-1} [n + m - 2i - 1] = [n + m - 1] + [n + m - 3] + [n + m - 5] + \cdots + [n - m + 1]$$

by introducing alternating signs. It is essential that $m \leq n$ above; one does not obtain the same result by swapping m and n in the formula. Note that if $m = 0$, the summation is empty, so by convention “[0][n]” = 0 for all $n \in \mathbb{Z}_{\geq 0}$. See Appendix A for further discussion.

We now precisely record the main result of this section.

Theorem 4.2. Let $S \in \mathbf{Rep}(U_q(\mathfrak{so}_{2n+1}))$ be the spin representation. There is a basis

$$\{\mathbf{X}^{(i)}\}_{i=0}^n \subset \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$$

such that

$$\mathbf{X}^{(i)}\mathbf{X}^{(1)} = (-1)^i “[i + 1][i + 1]”\mathbf{X}^{(i+1)} + (-1)^i “[i][i + 1]”\mathbf{X}^{(i)}$$

and so that the braiding is given by

$$R_{S,S} = q^{\frac{n}{2}} \sum_{i=0}^n q^{-i} \mathbf{X}^{(i)}.$$

□

This result is established in Propositions 4.25 and 4.28. Later on, Theorems 8.11, 8.22, and 10.21 categorify this result.

4.1. Representations of $U_q(\mathfrak{so}_{2n+1})$. In this section, we review the fundamental representations $\{V(\varpi_1), \dots, V(\varpi_{n-1}), S\}$ of $U_q(\mathfrak{so}_{2n+1})$. In doing so, we will follow the conventions and notation of Example 3.3. Further, as mentioned in Section 3.2, we consider representations over the field $\mathbb{C}(q^{\frac{1}{2}})$.

For $1 \leq k \leq n-1$, the fundamental weights are $\varpi_k = \epsilon_1 + \cdots + \epsilon_k$ and the representations $V(\varpi_k)$ are quantized analogues of the exterior powers $\Lambda^k(\mathbb{C}^{2n+1})$, while $\varpi_n = \frac{1}{2}(\epsilon_1 + \cdots + \epsilon_n)$ and $S := V(\varpi_n)$ is the (quantum) *spin representation*. We begin by studying the latter in depth, since it plays a leading role in the present work.

For $i, j = \{1, \dots, n\}$, let

$$\|i, j\| := \begin{cases} -2 & \text{if } j = i \\ 2 & \text{if } j = i + 1 \\ 0 & \text{else} \end{cases}$$

and for $J \subset \{1, \dots, n\}$ denote $\|i, J\| = \sum_{j \in J} \|i, j\| \in \{-2, 0, 2\}$. Also, set

$$q^J = \prod_{j \in J} (-q)(-q^2)^{n-j} = \prod_{j \in J} (-1)^{n-j+1} q^{2(n-j)+1}$$

and $q^{-J} = (q^J)^{-1}$.

Definition 4.3. The *spin representation* of $U_q(\mathfrak{so}_{2n+1})$ is the $\mathbb{C}(q^{\frac{1}{2}})$ -vector space S with basis $\{x_J\}_{J \in \mathcal{P}}$ where \mathcal{P} is the power set of $\{1, 2, \dots, n\}$. For $i \in \{1, \dots, n-1\}$, the action of $U_q(\mathfrak{so}_{2n+1})$ on S is specified as follows:

- (1) $k_i^{\pm 1} x_J = q^{\pm \|i, J\|} x_J$,
- (2) $k_n^{\pm 1} x_J = q^{\pm (1 + \|n, J\|)} x_J$
- (3) $e_i x_J = x_{J \setminus \{i\} \cup \{i+1\}}$ if $i \in J$ and $i+1 \notin J$,
- (4) $e_n x_J = x_{J \setminus \{n\}}$ if $n \in J$,
- (5) $f_i x_J = x_{J \setminus \{i+1\} \cup \{i\}}$ if $i+1 \in J$ and $i \notin J$,
- (6) $f_n x_J = x_{J \cup \{n\}}$ if $n \notin J$,

and where all e_j and f_j for $j \in \{1, \dots, n\}$ act by zero otherwise.

Using the relations in Definition 3.1, it is straightforward to check that S is indeed an irreducible $U_q(\mathfrak{so}_{2n+1})$ -representation. Note that $x_\emptyset \in S$ is thus a highest weight vector with $\text{wt}(x_\emptyset) = \varpi_n = \frac{1}{2}(\epsilon_1 + \dots + \epsilon_n)$, and hence $S \cong V(\varpi_n)$.

Lemma 4.4. Let $x_J \in S$ and $i \in \{1, \dots, n\}$. Exactly one of the following holds:

- (1) $(\alpha_i^\vee, \text{wt}(x_J)) = 1$ and $T_i x_J = -q_i f_i x_J$,
- (2) $(\alpha_i^\vee, \text{wt}(x_J)) = -1$ and $T_i x_J = e_i x_J$, or
- (3) $(\alpha_i^\vee, \text{wt}(x_J)) = 0$ and $T_i x_J = x_J$.

Proof. Using Definition 4.3 and the formulae for the simple roots from Example 3.3, we compute that

$$(4.1) \quad \text{wt}(x_J) = \frac{1}{2} \left(\sum_{i \notin J} \epsilon_i - \sum_{i \in J} \epsilon_i \right).$$

It is then an easy consequence that $(\alpha_i^\vee, \text{wt}(x_J)) \in \{-1, 0, 1\}$ for all $\{i\}, J \subset \{1, \dots, n\}$, thus S decomposes as a sum of 1- and 2-dimensional irreducible representations upon restriction to the subalgebra $U_{q_i}(\mathfrak{sl}_2) \subset U_q(\mathfrak{so}_{2n+1})$ generated by e_i, f_i , and $k_i^{\pm 1}$. The result then follows from (3.3). \square

Using Lemma 4.4, we can compute the action of T_{w_0} on certain basis vectors $x_J \in S$, where as always $w_0 \in W_{\mathfrak{so}_{2n+1}}$ denotes the longest element. Recall that the type B_n Weyl group $W_{\mathfrak{so}_{2n+1}}$ is the *hyperoctahedral group* which, as always, admits a presentation as a Coxeter group

$$W_{\mathfrak{so}_{2n+1}} := \langle s_1, \dots, s_n \mid s_i^2 = e, s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1} \text{ if } i \leq n-2, s_{n-1} s_n s_{n-1} s_n = s_n s_{n-1} s_n s_{n-1} \rangle.$$

However, $W_{\mathfrak{so}_{2n+1}}$ can also be identified with the subgroup of $\text{Perm}(\{-n, \dots, -1, 1, \dots, n\}) \cong \mathfrak{S}_{2n}$ consisting of permutations that satisfy $\sigma(-i) = -\sigma(i)$ via the assignment:

$$s_i \mapsto \begin{cases} (i-n, i-1-n)(n-i, n-i+1) & \text{if } 1 \leq i \leq n-1 \\ (-1, 1) & \text{if } i = n. \end{cases}$$

(Here (k, ℓ) denotes the transposition that interchanges k and ℓ .) Under this inclusion, the longest element $w_0 \in W_{\mathfrak{so}_{2n+1}}$ is identified with the longest element of \mathfrak{S}_{2n} , the half-twist permutation.

Lemma 4.5.

$$T_{w_0}(x_{\{1, \dots, i\}}) = q^{\{i+1, \dots, n\}} x_{\{i+1, \dots, n\}}$$

Proof. Set

$$w_{\{1, \dots, i\}}^{\{1, \dots, n\}} := (s_n)(s_{n-1} s_n) \cdots (s_{i+1} \cdots s_{n-1} s_n), \quad w_{\{1, \dots, n\}}^{\{i+1, \dots, n\}} := (s_i \cdots s_n)(s_{i-1} \cdots s_n) \cdots (s_1 \cdots s_n),$$

and let $w_0^{1, \dots, n-1}$ denote the longest element of the parabolic subgroup $\langle s_1, \dots, s_{n-1} \rangle \subset W_{\mathfrak{so}_{2n+1}}$. For each $i \in \{1, \dots, n\}$ the longest element $w_0 \in W_{\mathfrak{so}_{2n+1}}$ can be written as

$$w_0 = w_{\{1, \dots, n\}}^{\{i+1, \dots, n\}} w_0^{1, \dots, n-1} w_{\{1, \dots, i\}}^{\{1, \dots, n\}}$$

which e.g. can be verified using the aforementioned inclusion $W_{\mathfrak{so}_{2n+1}} \hookrightarrow \mathfrak{S}_{2n}$.

Next, Definition 4.3 gives that

$$x_{\{1, \dots, n\}} = (f_n)(f_{n-1}f_n) \cdots (f_{i+2} \cdots f_{n-1}f_n)(f_{i+1} \cdots f_{n-1}f_n)x_{\{1, \dots, i\}}$$

and

$$x_{\{i+1, \dots, n\}} = (e_i \cdots e_{n-1}e_n) \cdots (e_2 \cdots e_{n-1}e_n)(e_1 \cdots e_{n-1}e_n)x_{\{1, \dots, n\}},$$

so repeated application of Lemma 4.4 implies that

$$T_{w_{\{1, \dots, i\}}}^{\{1, \dots, n\}}(x_{\{1, \dots, i\}}) = q^{\{i+1, \dots, n\}}x_{\{1, \dots, n\}}$$

and

$$T_{w_{\{1, \dots, n\}}}^{\{i+1, \dots, n\}}(x_{\{1, \dots, n\}}) = x_{\{i+1, \dots, n\}}.$$

Since $(\alpha_i^\vee, \text{wt}(x_{\{1, \dots, n\}})) = 0$ for $1 \leq i \leq n-1$, Lemma 4.4 also gives that

$$T_{w_0^{1, \dots, n-1}}(x_{\{1, \dots, n\}}) = x_{\{1, \dots, n\}}$$

and the result follows. \square

We next discuss the remaining fundamental representations.

Definition 4.6. For $i = 1, 2, \dots, n-1$, let $V_i := V(\varpi_i)$ denote the Weyl module for $U_q(\mathfrak{so}_{2n+1})$ with fixed highest weight vector $v_i^+ \in V_i$ of weight $\varpi_i = \epsilon_1 + \epsilon_2 + \cdots + \epsilon_i$.

We will let $v_i^- \in V_i$ denote the unique vector in the lowest weight space of V_i such that $v_i^+ = T_{w_0}(v_i^-)$. Note that $\text{wt}(v_i^-) = w_0(\varpi_i) = -\varpi_i$. Extending this notation, we will also write $V_0 := \mathbb{C}(q^{\frac{1}{2}})$ to denote the trivial $U_q(\mathfrak{so}_{2n+1})$ -module (with fixed highest weight vector $v_0^+ := 1 = v_0^-$) and denote the distinguished highest and lowest weight vectors of S by $v_n^+ := v_\emptyset$ and $v_n^- := v_{\{1, \dots, n\}}$.

The representation V_1 admits the following explicit description, which we will use below to study certain morphisms in $\mathbf{Rep}(U_q(\mathfrak{so}_{2n+1}))$.

Proposition 4.7. The representation V_1 has a basis $\{a_1, a_2, \dots, a_n, u, b_n, \dots, b_2, b_1\}$ such that

$$\text{wt}(a_i) = \epsilon_i, \quad \text{wt}(u) = 0, \quad \text{wt}(b_i) = -\epsilon_i.$$

With respect to this basis, the $U_q(\mathfrak{so}_{2n+1})$ action is given by

- (1) $f_i a_i = a_{i+1}$ and $f_i b_{i+1} = b_i$ for $1 \leq i \leq n-1$,
- (2) $e_i a_i = a_{i-1}$ and $e_i b_i = b_{i+1}$ for $1 \leq i \leq n-1$,
- (3) $f_n a_n = u$ and $f_n u = [2]b_n$,
- (4) $e_n b_n = u$ and $e_n u = [2]a_n$,
- (5) $k_i v = q^{(\alpha_i, \text{wt}(v))}x$, for $1 \leq i \leq n$,

and where all e_i and f_i for $1 \leq i \leq n$ act by zero otherwise.

Proof. An exercise using Definition 3.1. \square

If we identify $v_1^+ = a_1$, then a computation using (3.3) shows that $v_1^- = b_1$.

4.2. Trivalent vertices. We now begin our study of morphisms in $\mathbf{Rep}(U_q(\mathfrak{so}_{2n+1}))$. It is a standard fact that

$$(4.2) \quad S \otimes S \cong \mathbb{C}(q^{\frac{1}{2}}) \oplus \left(\bigoplus_{i=1}^{n-1} V(\varpi_i) \right) \oplus V(2\varpi_n) = \left(\bigoplus_{i=0}^{n-1} V_i \right) \oplus V(2\varpi_n)$$

and we first describe the inclusions of the submodules V_i for $0 \leq i \leq n-1$. Note: in this subsection we will break our conventions from §3.1 by using the symbol I to denote arbitrary subsets of $\{1, \dots, n\}$ (as opposed to letting it denote the set of Dynkin nodes, as is done in other sections of the paper). Given such $I \subset \{1, \dots, n\}$, we will denote its complement by $I^c = \{1, \dots, n\} \setminus I$.

Proposition 4.8. For each $0 \leq i \leq n-1$, there is a unique map $Y_i: V_i \rightarrow S \otimes S$ of $U_q(\mathfrak{so}_{2n+1})$ -modules such that

$$(4.3) \quad Y_i(v_i^+) := \sum_{\substack{I \cap J = \emptyset \\ I \cup J = \{i+1, \dots, n\}}} q^J x_I \otimes x_J.$$

Proof. Equation (4.1) implies that $Y_i(v_i^+)$ is a non-zero vector in the ϖ_i -weight space of $S \otimes S$, so it suffices to show that it is a highest weight vector, i.e. that $U_q(\mathfrak{so}_{2n+1})^{>0} \cdot Y_i(v_i^+) = 0$. Recall that e_ℓ acts on tensor products via $\Delta(e_\ell) = e_\ell \otimes k_\ell + 1 \otimes e_\ell$. For $\ell \in \{1, \dots, n-1\}$, we thus compute:

$$\begin{aligned} e_\ell \cdot Y_i(v_i^+) &= \sum_{\substack{I \cap J = \emptyset \\ I \cup J = \{i+1, \dots, n\}}} q^J e_\ell(x_I) \otimes k_\ell(x_J) + \sum_{\substack{I \cap J = \emptyset \\ I \cup J = \{i+1, \dots, n\}}} q^J x_I \otimes e_\ell(x_J) \\ &= \sum_{\substack{I \cap J = \emptyset \\ I \cup J = \{i+1, \dots, n\} \\ \ell \in I, \ell+1 \notin I}} q^J q^{\|\ell, J\|} x_{I \setminus \{\ell\} \cup \{\ell+1\}} \otimes x_J + \sum_{\substack{I \cap J = \emptyset \\ I \cup J = \{i+1, \dots, n\} \\ \ell \in J, \ell+1 \notin J}} q^J x_I \otimes x_{J \setminus \{\ell\} \cup \{\ell+1\}} \end{aligned}$$

which is zero when $\ell \leq i$, since both summations are empty. If $\ell \in \{i+1, \dots, n-1\}$, we then have

$$\sum_{\substack{I \cap J = \{\ell+1\} \\ I \cup J = \{i+1, \dots, n\} \setminus \{\ell\}}} \left(q^{J \cup \{\ell\} \setminus \{\ell+1\}} + q^J q^{\|\ell, J\|} \right) x_I \otimes x_J.$$

Since

$$q^{J \cup \{\ell\} \setminus \{\ell+1\}} = \frac{(-q)(-q^2)^{n-\ell}}{(-q)(-q^2)^{n-(\ell+1)}} q^J = -q^2 q^J = -q^J q^{\|\ell, J\|},$$

it follows that $e_\ell \cdot Y_i(v_i^+) = 0$ in this case as well. Similarly, one can compute that $e_n \cdot Y_i(v_i^+) = 0$. \square

We next compute the value of Y_i on our distinguished lowest-weight vectors $v_i^- \in V_i$.

Lemma 4.9. We have that

$$(4.4) \quad Y_i(v_i^-) = \sum_{\substack{I \cap J = \emptyset \\ I \cup J = \{i+1, \dots, n\}}} q^J x_{J^c} \otimes x_{I^c}.$$

Proof. Let $w_i^- \in S \otimes S$ denote the right-hand side of (4.4). We leave it as an exercise to verify that $\text{wt}(w_i^-) = -\varpi_i$ and that w_i^- is a lowest weight vector, i.e. that $U_q(\mathfrak{so}_{2n+1})^{<0} \cdot w_i^- = 0$. It then follows that $Y_i(v_i^-) = \chi \cdot w_i^-$ for some $\chi \in \mathbb{C}(q^{\frac{1}{2}})$. We will show that $\chi = 1$ by computing the value of $(T_{w_0}^{-1} \otimes T_{w_0}^{-1}) \circ \Delta(T_{w_0})$ on both $Y_i(v_i^-)$ and $\chi \cdot w_i^-$.

First, since $x_{\{1, \dots, n\}} \in S$ is a lowest weight vector, Proposition 3.5 gives that

$$(4.5) \quad ((T_{w_0}^{-1} \otimes T_{w_0}^{-1}) \circ \Delta(T_{w_0}))(w_i^-) = \chi \cdot x_{\{1, \dots, n\}} \otimes x_{\{1, \dots, i\}} + \sum_{\substack{I', J' \\ I' \neq \{1, \dots, n\}}} \xi_{I', J'} x_{I'} \otimes x_{J'},$$

for some $\xi_{I', J'} \in \mathbb{C}(q^{\frac{1}{2}})$. Next, we compute that

$$\begin{aligned} ((T_{w_0}^{-1} \otimes T_{w_0}^{-1}) \circ \Delta(T_{w_0}))(Y_i(v_i^-)) &= (T_{w_0}^{-1} \otimes T_{w_0}^{-1} \circ Y_i)(T_{w_0} v_i^-) \\ &= (T_{w_0}^{-1} \otimes T_{w_0}^{-1} \circ Y_i)(v_i^+) \\ &= \sum_{\substack{I \cap J = \emptyset \\ I \cup J = \{i+1, \dots, n\}}} q^J T_{w_0}^{-1}(x_I) \otimes T_{w_0}^{-1}(x_J). \end{aligned}$$

Now, Lemma 4.5 gives that $T_{w_0}^{-1}(x_\emptyset) = x_{\{1, \dots, n\}}$ and $T_{w_0}^{-1}(x_{\{i+1, \dots, n\}}) = q^{-\{i+1, \dots, n\}} x_{\{1, \dots, i\}}$, while $T_{w_0}^{-1}x_I \in \mathbb{C}(q^{\frac{1}{2}}) \cdot x_{I^c}$ for all I , so

$$(4.6) \quad \begin{aligned} ((T_{w_0}^{-1} \otimes T_{w_0}^{-1}) \circ \Delta(T_{w_0}))(\Upsilon_i(v_i^-)) &= q^{-\{i+1, \dots, n\}} q^{\{i+1, \dots, n\}} x_{\{1, \dots, n\}} \otimes x_{\{1, \dots, i\}} \\ &+ \sum_{\substack{I', J' \\ I' \neq \{1, \dots, n\}}} \zeta_{I', J'} x_{I'} \otimes x_{J'}, \end{aligned}$$

for some $\zeta_{I', J'} \in \mathbb{C}(q^{\frac{1}{2}})$. The result then follows by comparing the coefficients of $x_{\{1, \dots, n\}} \otimes x_{\{1, \dots, i\}}$ in (4.5) and (4.6). \square

Example 4.10. We record the action of Υ_1 on the basis for V_1 from Proposition 4.7. For $\ell \in \{1, \dots, n\}$, let $\sigma_\ell: \{1\}^c \rightarrow \{\ell\}^c$ be the unique order preserving bijection, e.g. if $n = 4$ then $\sigma_3(2) = 1, \sigma_3(3) = 2$, and $\sigma_3(4) = 4$. We then have

$$\begin{aligned} \Upsilon_1(a_\ell) &= \sum_{I \subset \{2, \dots, n\}} q^I x_{\sigma_\ell(\{2, \dots, n\} \setminus I)} \otimes x_{\sigma_\ell(I)}, & \Upsilon_1(b_\ell) &= \sum_{I \subset \{2, \dots, n\}} q^I x_{\sigma_\ell(\{2, \dots, n\} \setminus I) \cup \{\ell\}} \otimes x_{\sigma_\ell(I) \cup \{\ell\}} \\ \Upsilon_1(u) &= \sum_{I \subset \{2, \dots, n\}} q^I x_{\sigma_n(\{2, \dots, n\} \setminus I) \cup \{n\}} \otimes x_{\sigma_n(I)} + \sum_{I \subset \{2, \dots, n\}} q^{-1} \cdot q^I x_{\sigma_n(\{2, \dots, n\} \setminus I)} \otimes x_{\sigma_n(I) \cup \{n\}}. \end{aligned}$$

Next, we compute the composition of the braiding (3.1) with the morphisms Υ_i from Proposition 4.8. Below, we will use this result to give a formula for the braiding on $S \otimes S$ in terms of (morphisms built from) the morphisms Υ_i .

Lemma 4.11. For $0 \leq i \leq n-1$, we have

$$R_{S,S} \circ \Upsilon_i = q^{-\binom{n-2i}{2}} q^{-\{i+1, \dots, n\}} \Upsilon_i.$$

Proof. The decomposition (4.2) implies that

$$\dim(\text{Hom}_{U_q(\mathfrak{so}_{2n+1})}(V_i, S \otimes S)) = 1$$

so $R_{S,S} \circ \Upsilon_i$ is necessarily a scalar multiple of Υ_i . We identify this scalar by computing the value of each on the highest weight vector $v_i^+ \in V_i$. Proposition 4.8 gives that

$$\Upsilon_i(v_i^+) = x_{\{i+1, \dots, n\}} \otimes x_\emptyset + \sum_{\substack{I \cap J = \emptyset, J \neq \emptyset \\ I \cup J = \{i+1, \dots, n\}}} q^J x_I \otimes x_J$$

so Proposition 3.5 gives that

$$((T_{w_0}^{-1} \otimes T_{w_0}^{-1}) \circ \Delta(T_{w_0}))(\Upsilon_i(v_i^+)) = x_{\{i+1, \dots, n\}} \otimes x_\emptyset + \sum_{\substack{I', J' \\ J' \neq \emptyset}} \xi_{I', J'} x_{I'} \otimes x_{J'},$$

for some $\xi_{I', J'} \in \mathbb{C}(q^{\frac{1}{2}})$. Theorem 3.4 and equation (3.1) then give that the coefficient of $x_\emptyset \otimes x_{\{i+1, \dots, n\}}$ in $R_{S,S} \circ \Upsilon_i(v_i^+)$ is

$$q^{(\text{wt}(-), \text{wt}(-))}(x_{\{i+1, \dots, n\}} \otimes x_\emptyset) = q^{(\sum_{\ell=1}^n \epsilon_\ell/2, \sum_{\ell=1}^i \epsilon_\ell/2 - \sum_{\ell=i+1}^n \epsilon_\ell/2)} = q^{-\binom{n-2i}{2}}.$$

The result now follows from observing that the coefficient of $x_\emptyset \otimes x_{\{i+1, \dots, n\}}$ in $\Upsilon_i(v_i^+)$ is $q^{\{i+1, \dots, n\}}$. \square

Remark 4.12. Since V_0 is the monoidal identity, we will later draw the map Υ_0 as a cup, so that $R_{S,S} \circ \Upsilon_0$ is a curl. In this case we see that Lemma 4.11 recovers the (negative) twist coefficient for the braiding from Remark 3.9. In particular, we have

$$(4.7) \quad q^{-\frac{n}{2}} q^{-\{1, \dots, n\}} = (-1)^{\binom{n+1}{2}} q^{-\frac{n(2n+1)}{2}} = (-1)^{(2\varpi_n, \rho^\vee)} q^{-(\varpi_n, \varpi_n + 2\rho)},$$

where for the type B_n root system:

$$(4.8) \quad 2\rho = \sum_{i=1}^n (2(n-i) + 1)\epsilon_i, \quad \rho^\vee = \frac{1}{2} \sum_{i=1}^n (n-i+1)\epsilon_i, \quad \varpi_n = \frac{1}{2} \sum_{i=1}^n \epsilon_i.$$

Next, note that the fundamental $U_q(\mathfrak{so}_{2n+1})$ -representations $V_1, \dots, V_{n-1}, V_n = S$ are all self-dual, since, for $1 \leq i \leq n$, the highest weight of V_i^* is $-w_0(\varpi_i) = -(-\varpi_i) = \varpi_i$. We hence can consider the compositions

$$(4.9) \quad V_0 = \mathbb{C}(q^{\frac{1}{2}}) \rightarrow V_i \otimes V_i^* \xrightarrow{\cong} V_i \otimes V_i \quad \text{and} \quad V_i \otimes V_i \xrightarrow{\cong} V_i^* \otimes V_i \rightarrow \mathbb{C}(q^{\frac{1}{2}}) = V_0$$

of these isomorphisms with the canonical coevaluation and evaluation maps. We now aim to give an explicit description of these morphisms.

For each $1 \leq i \leq n$, fix a basis \mathbb{B}_i for V_i that contains our distinguished highest and lowest weight vectors v_i^+ and v_i^- . Given $v \in \mathbb{B}_i$, we let v^* denote the corresponding vector in the dual basis \mathbb{B}_i^* of V_i^* . The assignment $v_i^+ \mapsto (v_i^-)^*$ then determines an isomorphism $\varphi_i: V_i \xrightarrow{\cong} V_i^*$.

Example 4.13. The isomorphisms $\varphi_1: V_1 \rightarrow V_1^*$ and $\varphi_n: S \rightarrow S^*$ have the following explicit descriptions:

$$(4.10) \quad \varphi_1(a_i) = (-q^{-2})^{i-1} b_i^*, \quad \varphi_1(u) = (-q^{-2})^n [2]u^*, \quad \varphi_1(b_i) = -(-q^{-2})^{2n-i} a_i^*,$$

and

$$(4.11) \quad \varphi_n(x_I) = q^{-I} x_{I^c}^*.$$

We now describe the morphisms in (4.9).

Proposition 4.14. For each $1 \leq i \leq n$, there is a unique $U_q(\mathfrak{so}_{2n+1})$ -module homomorphism

$$\mathbf{U}_i: \mathbb{C}(q^{\frac{1}{2}}) \longrightarrow V_i \otimes V_i$$

such that

$$(4.12) \quad \mathbf{U}_i(1) = \sum_{v \in \mathbb{B}_i} v \otimes \varphi_i^{-1}(v^*),$$

and there is a unique $U_q(\mathfrak{so}_{2n+1})$ -module homomorphism

$$\mathbf{\Omega}^i: V_i \otimes V_i \longrightarrow \mathbb{C}(q^{\frac{1}{2}})$$

such that

$$\mathbf{\Omega}^i(v_i^+ \otimes v_i^-) = 1.$$

Consequently, these maps agree with those in (4.9).

Proof. Our choice of Hopf algebra structure $(U_q(\mathfrak{so}_{2n+1}); \Delta, \mathbf{S}, \epsilon)$ ensures that the canonical linear maps $\mathbf{coev}: \mathbb{C}(q^{\frac{1}{2}}) \longrightarrow V_i \otimes V_i^*$ and $\mathbf{ev}: V_i^* \otimes V_i \longrightarrow \mathbb{C}(q^{\frac{1}{2}})$ intertwine the action of $U_q(\mathfrak{so}_{2n+1})$. Thus, $\mathbf{U}_i := (\text{id} \otimes \varphi_i^{-1}) \circ \mathbf{coev}$ and $\mathbf{\Omega}^i := \mathbf{ev} \circ (\varphi_i \otimes \text{id})$ are $U_q(\mathfrak{so}_{2n+1})$ -module homomorphisms and are characterized by the indicated formulae. \square

We will write the morphisms in Proposition 4.14 using the graphical calculus for monoidal categories as follows:

$$\mathbf{U}_i = \begin{array}{c} i \\ \cup \\ i \end{array}, \quad \mathbf{\Omega}^i = \begin{array}{c} \cap \\ i \\ i \end{array}.$$

By construction, these morphism satisfy the relations:

$$(4.13) \quad \begin{array}{c} \text{cup} \\ \text{cap} \end{array} = \text{straight line} = \begin{array}{c} \text{cup} \\ \text{cap} \end{array}$$

Remark 4.15. The full monoidal subcategory of $U_q(\mathfrak{so}_{2n+1})$ -modules generated by the self-dual representations V_i for $i = 0, 1, \dots, n$ is a strict pivotal category, where we use the (non-standard) pivotal structure from [66]. With this choice of pivotal structure, all Frobenius-Schur indicators are $+1$, and therefore morphisms in the category can be described by an unoriented graphical calculus, see [65, 64]. It is possible to show that our choice of cup and cap morphisms (\mathbf{U}_i and $\mathbf{\Omega}^i$, respectively) agree with any other choice of cups and caps, up to simultaneous rescaling. See e.g. [7, Remark 5.2] for further details.

Example 4.16. By (4.11), we have

$$(4.14) \quad \mathbf{\Omega}^n(x_I \otimes x_J) = q^{-I} \delta_{J, I^c}, \quad \mathbf{U}_n(1) = \sum_{I \subset \{1, \dots, n\}} q^I x_{I^c} \otimes x_I,$$

and

$$(4.15) \quad n \bigcirc := \mathbf{\Omega}^n \circ \mathbf{U}_n(1) = \sum_{I \subset \{1, \dots, n\}} \frac{q^I}{q^{I^c}} = (-1)^{\binom{n+1}{2}} \prod_{i=1}^n (q^{2i-1} + q^{1-2i})$$

Note also that that $\mathbf{U}_n = \mathbf{Y}_0$.

Establishing the last equality in (4.15) is a straightforward exercise, as is the following generalization.

Lemma 4.17. If $d_i := \prod_{\ell=1}^i \frac{[4\ell - 2]}{[2\ell - 1]} = \prod_{\ell=1}^i (q^{2\ell-1} + q^{-2\ell+1})$, then

$$\sum_{\substack{I \cup J = \{i+1, \dots, n\} \\ I \cap J = \emptyset}} q^J q^{-I} = (-1)^{\binom{n-i+1}{2}} d_{n-i}.$$

□

We denote the morphisms $\mathbf{Y}_i: V_i \rightarrow S \otimes S$ from Proposition 4.8 in our graphical calculus as follows:

$$\begin{array}{c} S \quad S \\ \text{Y} \\ i \end{array} := \mathbf{Y}_i.$$

Here, and in the following, we use the color gray to graphically denote the spin representation. Composing these morphisms with (tensor products of) the cap and cup morphisms from Proposition 4.14 yields the following morphisms in $\mathbf{Rep}(U_q(\mathfrak{so}_{2n+1}))$.

Definition 4.18. For $0 \leq i \leq n-1$, let $\lambda^i: S \otimes S \rightarrow V_i$ be the $U_q(\mathfrak{so}_{2n+1})$ -module morphisms given in terms of the graphical calculus as follows:

$$(4.16) \quad \lambda^i = \begin{array}{c} \text{cup} \\ \text{cap} \end{array} := \begin{array}{c} \text{cup} \\ \text{cap} \end{array}.$$

Proposition 4.20. Set $\mathfrak{I}^{(i)} := Y_i \circ \lambda^i$, for $i = 0, 1, \dots, n-1$. For $i, j \in \{0, \dots, n-1\}$, we have that

$$(4.20) \quad \mathfrak{I}^{(i)} \mathfrak{I}^{(j)} = (-1)^{\binom{n-i+1}{2}} d_{n-i} \delta_{i,j} \cdot \mathfrak{I}^{(i)}.$$

If we further let $\mathfrak{I}^{(n)} := \text{id}_{S \otimes S}$, then $\{\mathfrak{I}^{(i)}\}_{i=0}^n$ is a basis for $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$.

Proof. Equation (4.20) follows from equation (4.18) and equation (4.19). The result then follows from the decomposition (4.2) and another application of Schur's lemma. \square

We conclude this section with a description of the braiding $R_{S,S} \in \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$ in terms of the basis from Proposition 4.20.

Proposition 4.21. For $1 \leq i \leq n$, set

$$(4.21) \quad b_{n-i} = \frac{q^{\frac{n}{2}}}{d_i} \left((q^{-2})^{\binom{i+1}{2}} - (-1)^{\binom{i+1}{2}} \right).$$

The braiding on $S \otimes S$ is given by

$$R_{S,S} = q^{\frac{n}{2}} \text{id}_{S \otimes S} + \sum_{i=1}^n b_{n-i} \mathfrak{I}^{(n-i)}.$$

Proof. By Proposition 4.20, there exist scalars $b_j \in \mathbb{C}(q^{\frac{1}{2}})$ for $0 \leq j \leq n$ so that

$$R_{S,S} = b_n \text{id}_{S \otimes S} + \sum_{i=1}^n b_{n-i} \mathfrak{I}^{(n-i)}.$$

Plugging the highest weight vector $x_\emptyset \otimes x_\emptyset \in S \otimes S$ into $R_{S,S}$ and using Proposition 3.5 gives that $b_n = q^{\frac{n}{2}}$. For the remaining coefficients, Lemmata 4.11 and 4.19 give

$$q^{\frac{n-2i}{2}} q^{-\{n-i+1, \dots, n\}} Y_{n-i} = R_{S,S} \circ Y_{n-i} = q^{\frac{n}{2}} Y_{n-i} + (-1)^{\binom{i+1}{2}} d_i b_{n-i} Y_{n-i},$$

thus

$$b_{n-i} = \frac{(-1)^{\binom{i+1}{2}} q^{\frac{n}{2}}}{d_i} \left(q^{-i} q^{-\{n-i+1, \dots, n\}} - 1 \right) = \frac{q^{\frac{n}{2}}}{d_i} \left((q^{-2})^{\binom{i+1}{2}} - (-1)^{\binom{i+1}{2}} \right). \quad \square$$

4.3. H- and X-morphisms. In this section, we consider two additional bases $\{H^{(i)}\}_{i=0}^n$ and $\{X^{(i)}\}_{i=0}^n$ for the algebra $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$. The first is obtained in a straightforward way: diagrammatically, it is the rotation of the basis from Proposition 4.20. The latter basis is a novel construction of this paper and is more subtle. Its construction was motivated by our categorified considerations appearing later, and is the ‘‘canonical basis’’ for this algebra, in a certain sense.

To begin, define $H^{(i)}$ using graphical calculus as follows:

$$(4.22) \quad H^{(i)} = \begin{array}{c} S \quad S \\ \diagdown \quad \diagup \\ \text{---} \\ \diagup \quad \diagdown \\ S \quad S \end{array} \quad := \quad \begin{array}{c} S \quad S \\ \diagdown \quad \diagup \\ \text{---} \\ \diagup \quad \diagdown \\ S \quad S \end{array}$$

We begin by computing the values of the ‘‘triangles’’:

$$\begin{array}{c} S \quad S \\ \diagdown \quad \diagup \\ \text{---} \\ \diagup \quad \diagdown \\ \text{---} \\ \text{---} \\ i \end{array} \quad := H^{(1)} \circ Y_i.$$

Lemma 4.22. For $0 \leq i \leq n-1$,

$$\mathbf{H}^{(1)} \circ \mathbf{Y}_i = (-1)^{n-i} \frac{[2(n-i)+1]}{[2]} \mathbf{Y}_i$$

Proof. Again, the decomposition 4.2 implies that there is some scalar χ so that $\mathbf{H}^{(1)} \circ \mathbf{Y}_i = \chi \mathbf{Y}_i$. Let

$$\pi_{\{i+1, \dots, n\} \otimes \emptyset} \in \text{End}_{\mathbb{C}(q^{1/2})}(S \otimes S)$$

denote the *linear operator* which projects to $x_{\{i+1, \dots, n\}} \otimes x_\emptyset$ with respect to the basis $\{x_I \otimes x_J\}_{I, J \subset \{1, \dots, n\}}$ of $S \otimes S$. (These are not $U_q(\mathfrak{so}_{2n+1})$ -module homomorphisms.) It follows from (4.3) that

$$\chi \cdot q^\emptyset \cdot x_{\{i+1, \dots, n\}} \otimes x_\emptyset = \pi_{\{i+1, \dots, n\} \otimes \emptyset} \circ \chi \mathbf{Y}_i(v_i^+) = \pi_{\{i+1, \dots, n\} \otimes \emptyset} \circ \mathbf{H}^{(1)} \circ \mathbf{Y}_i(v_i^+).$$

Combining (4.22) with (4.16) gives

$$\mathbf{H}^{(1)} \circ \mathbf{Y}_i = (\mathbf{\Omega}^n \otimes \text{id}_{S \otimes S} \otimes \mathbf{\Omega}^n) \circ (\text{id}_S \otimes \mathbf{Y}_1 \otimes \mathbf{Y}_1 \otimes \text{id}_S) \circ (\text{id}_S \otimes \mathbf{U}_1 \otimes \text{id}_S) \circ \mathbf{Y}_i,$$

so (4.3) and (4.12) imply that $\pi_{\{i+1, \dots, n\} \otimes \emptyset} \circ \mathbf{H}^{(1)} \circ \mathbf{Y}_i(v_i^+)$ is equal to

$$(4.23) \quad \sum_{\substack{I \subset \{i+1, \dots, n\} \\ v \in \{a_i, u, b_j\}}} q^I \pi_{\{i+1, \dots, n\} \otimes \emptyset} \circ (\mathbf{\Omega}^n \otimes \text{id}_{S \otimes S} \otimes \mathbf{\Omega}^n)(x_{\{i+1, \dots, n\} \setminus I} \otimes \mathbf{Y}_1(v) \otimes \mathbf{Y}_1(\varphi_1^{-1}(v^*)) \otimes x_I).$$

We now work to simplify this expression.

For $K \subset \{1, \dots, n\}$, similarly let $\pi_K \in \text{End}_{\mathbb{C}(q^{1/2})}(S)$ denote the linear operator projecting to x_K with respect to the basis $\{x_I\}_{I \subset \{1, \dots, n\}}$. Let $\ell \in \{1, \dots, n\}$ and $I \subset \{i+1, \dots, n\}$. Using Example 4.10, along with equation (4.14), we see that

$$\pi_\emptyset \circ (\text{id}_S \otimes \mathbf{\Omega}^n)(\mathbf{Y}_1(b_\ell) \otimes x_I) = 0,$$

and, unless $I = \{\ell\}$, we also have

$$\pi_\emptyset \circ (\text{id}_S \otimes \mathbf{\Omega}^n)(\mathbf{Y}_1(a_\ell) \otimes x_I) = 0.$$

Similarly, one finds that

$$\pi_\emptyset \circ (\text{id}_S \otimes \mathbf{\Omega}^n)(\mathbf{Y}_1(u) \otimes x_I) \neq 0$$

if and only if $I = \emptyset$ and

$$\pi_{\{i+1, \dots, n\}} \circ (\mathbf{\Omega}^n \otimes \text{id}_S)(x_I \otimes \mathbf{Y}_1(u)) \neq 0$$

if and only if $I = \{i+1, \dots, n\}$.

These computations, together with equation (4.10), imply that (4.23) simplifies to

$$\begin{aligned} & \sum_{\ell=i+1}^n (-1)^{\ell-1} (q^2)^{\ell-1} q^{\{\ell\}} \pi_{\{i+1, \dots, n\} \otimes \emptyset} \circ (\mathbf{\Omega}^n \otimes \text{id}_{S \otimes S} \otimes \mathbf{\Omega}^n)(x_{\{i+1, \dots, n\} - \{\ell\}} \otimes \mathbf{Y}_1(b_\ell) \otimes \mathbf{Y}_1(a_\ell) \otimes x_{\{\ell\}}) \\ & + \frac{(-1)^n (q^2)^n q^\emptyset}{[2]} \pi_{\{i+1, \dots, n\} \otimes \emptyset} \circ (\mathbf{\Omega}^n \otimes \text{id}_{S \otimes S} \otimes \mathbf{\Omega}^n)(x_{\{i+1, \dots, n\}} \otimes \mathbf{Y}_1(u) \otimes \mathbf{Y}_1(u) \otimes x_\emptyset). \end{aligned}$$

Finally, using our explicit formulae for Υ_1 and Ω^n , we conclude that

$$\begin{aligned}
\chi &= \sum_{\ell=i+1}^n (-1)^{\ell-1} (q^2)^{\ell-1} q^{\{\ell\}} q^{\{2,\dots,n\}} q^{\{i+2,\dots,n\}} q^{-\{i+1,\dots,\widehat{\ell},\dots,n\}} q^{-\{1,\dots,\widehat{\ell},\dots,n\}} \\
&\quad + (-1)^n \frac{(q^2)^n q^0 q^{-1} q^{\{i+2,\dots,n\}} q^{-1} q^{\{2,\dots,n\}} q^{-\{i+1,\dots,n\}} q^{-\{1,\dots,n\}}}{[2]} \\
&= \frac{q^{\{2,\dots,n\}} q^{\{i+2,\dots,n\}}}{q^{\{i+1,\dots,n\}} q^{\{1,\dots,n\}}} \left(\sum_{\ell=i+1}^n (-1)^{\ell-1} (q^2)^{\ell-1} q^{\{\ell\}} q^{\{\ell\}} q^{\{\ell\}} + (-1)^n \frac{(q^2)^n q^{-2}}{[2]} \right) \\
&= (-1)^n q^{-\{1,i+1\}} \left(\sum_{\ell=i+1}^n q^{6n-4\ell+1} + \frac{q^{2n-2}}{[2]} \right) \\
&= \frac{(-1)^{n+i}}{[2]} \left(\sum_{\ell=i+1}^n (q^{2n+2i-4\ell+4} + q^{2n+2i-4\ell+2}) + q^{-2n+2i} \right) \\
&= (-1)^{n-i} \frac{[2(n-i)+1]}{[2]}. \quad \square
\end{aligned}$$

Using this, we now express the morphism $H^{(1)}$ in terms of the basis from Proposition 4.20. Recall the devil's product from Definition 4.1, and the notation $\Xi^{(i)} := \Upsilon_i \circ \lambda^i$ and $\Xi^{(n)} = \text{id}_{S \otimes S}$ introduced in Proposition 4.20, both will be used extensively for the remainder of this section.

Lemma 4.23. We have that

$$H^{(1)} = \frac{1}{[2]} \text{id}_{S \otimes S} + \sum_{i=1}^n (-1)^{\binom{i}{2}} \frac{[i][i+1]}{d_i} \Xi^{(n-i)}.$$

Proof. By Proposition 4.20, there are scalars $\chi, \lambda_0, \dots, \lambda_{n-1} \in \mathbb{C}(q^{\frac{1}{2}})$ such that

$$(4.24) \quad H^{(1)} = \chi \text{id}_{S \otimes S} + \sum_{\ell=0}^{n-1} \lambda_\ell \Xi^{(\ell)}.$$

Since $\Xi^{(\ell)}(x_\emptyset \otimes x_\emptyset) = 0$ for $0 \leq \ell \leq n-1$, a similar argument to the one given in the proof of Lemma 4.22 implies that $\chi = 1/[2]$. Next, composing (4.24) with λ^{n-i} and applying Lemma 4.22, we obtain

$$(-1)^k \frac{[2(n-(n-i))+1]}{[2]} = \frac{1}{[2]} + \lambda_{n-i} (-1)^{\binom{i+1}{2}} d_i$$

and so

$$\lambda_{n-i} = (-1)^{\binom{i+1}{2}} (-1)^i \left(\frac{[2i+1] + (-1)^{i+1}}{[2]d_i} \right).$$

The result then follows from $m = i, n = i+1$ case of the identity

$$(4.25) \quad [2]^{[m][n]} = [n+m] + (-1)^{m-1} [n-m]$$

(which holds for $m \leq n$) and the identity

$$(-1)^{\binom{i+1}{2}} (-1)^i = (-1)^{\binom{i}{2}}. \quad \square$$

We now introduce the distinguished elements of $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$ appearing in Theorem 4.2.

Definition 4.24. Let

$$(4.26) \quad \mathbf{X} := \mathbf{X}^{(1)} := \mathbf{H}^{(1)} - \frac{1}{[2]} \mathbf{\Xi}^{(n)} = \mathbf{\Xi}^{(n-1)} + \sum_{\ell=2}^n (-1)^{\binom{\ell}{2}} \frac{“[\ell][\ell+1]”}{d_\ell} \mathbf{\Xi}^{(n-\ell)}.$$

More generally, let $\mathbf{X}^{(0)} := \text{id}_{S \otimes S}$ and for $0 \leq i \leq n-1$, set

$$(4.27) \quad \mathbf{X}^{(i+1)} := \frac{(-1)^i}{“[i+1]2”} \left(\mathbf{X}^{(i)} \mathbf{X} - (-1)^i “[i][i+1]” \mathbf{X}^{(i)} \right).$$

By design we have that

$$(4.28) \quad \mathbf{X}^{(i)} \mathbf{X} = (-1)^i “[i+1]2” \mathbf{X}^{(i+1)} + (-1)^i “[i][i+1]” \mathbf{X}^{(i)}.$$

Further, (4.27) implies that for $1 \leq i \leq n$,

$$(4.29) \quad \mathbf{X}^{(i)} = (-1)^{\binom{i}{2}} \frac{(\mathbf{X} + (-1)^i “[i-1][i]”)(\mathbf{X} + (-1)^{i-1} “[i-2][i-1]”)\cdots(\mathbf{X} + (-1)^0 “[0][1]”)}{“[i]2” \cdots “[1]2”}.$$

We now give an explicit unitriangular change of basis from $\{\mathbf{X}^{(i)}\}_{i=0}^n$ to $\{\mathbf{\Xi}^{(n-i)}\}_{i=0}^n$, generalizing (4.26).

Proposition 4.25. For $1 \leq i \leq n$,

$$(4.30) \quad \mathbf{X}^{(i)} = \sum_{\ell=i}^n n^{-i} \lambda_{n-\ell} \mathbf{\Xi}^{(n-\ell)}$$

where, for $i \leq \ell \leq n$,

$$(4.31) \quad n^{-i} \lambda_{n-\ell} := \frac{(-1)^{\binom{\ell-i+1}{2}}}{d_\ell} \cdot \prod_{t=1}^i \frac{“[\ell+1-t][\ell+t]”}{“[t]2”}.$$

Further, $n^{-i} \lambda_{n-i} = 1$ and thus $\{\mathbf{X}^{(i)}\}_{i=0}^n$ is a basis for $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$.

Proof. Note that $n^{-1} \lambda_{n-\ell} = (-1)^{\binom{\ell}{2}} \frac{“[\ell][\ell+1]”}{d_\ell}$ so $n^{-1} \lambda_{n-1} = 1$ and thus (4.26) implies the claim when $i = 1$. We now compute inductively that

$$\begin{aligned} \mathbf{X}^{(i+1)} &\stackrel{(4.27)}{=} \frac{(-1)^i}{“[i+1]2”} \left(\mathbf{X}^{(i)} \mathbf{X} - (-1)^i “[i][i+1]” \mathbf{X}^{(i)} \right) \\ &\stackrel{(4.30)}{=} \frac{(-1)^i}{“[i+1]2”} \left(\sum_{\ell=i}^n n^{-i} \lambda_{n-\ell} \mathbf{\Xi}^{(n-\ell)} \right) \left(\left(\sum_{\ell=1}^n (-1)^{\binom{\ell}{2}} \frac{“[\ell][\ell+1]”}{d_\ell} \mathbf{\Xi}^{(n-\ell)} \right) - (-1)^i “[i][i+1]” \right) \\ &\stackrel{(4.20)}{=} \frac{(-1)^i}{“[i+1]2”} \sum_{\ell=i}^n n^{-i} \lambda_{n-\ell} \left((-1)^\ell “[\ell][\ell+1]” - (-1)^i “[i][i+1]” \right) \mathbf{\Xi}^{(n-\ell)} \\ &= \frac{(-1)^i}{“[i+1]2”} \sum_{\ell=i+1}^n n^{-i} \lambda_{n-\ell} \left((-1)^\ell “[\ell][\ell+1]” - (-1)^i “[i][i+1]” \right) \mathbf{\Xi}^{(n-\ell)} \\ &= \frac{1}{“[i+1]2”} \sum_{s=1}^{n-i} n^{-i} \lambda_{n-i-s} \left((-1)^s “[i+s][i+s+1]” - “[i][i+1]” \right) \mathbf{\Xi}^{(n-i-s)} \\ &= \frac{1}{“[i+1]2”} \sum_{s=1}^{n-i} n^{-i} \lambda_{n-i-s} \left(\sum_{j=0}^{i+s-1} (-1)^{j-s} [2i-2(j-s)] - \sum_{j=0}^{i-1} (-1)^j [2i-2j] \right) \mathbf{\Xi}^{(n-i-s)} \\ &= \frac{1}{“[i+1]2”} \sum_{s=1}^{n-i} n^{-i} \lambda_{n-i-s} \left((-1)^s \sum_{j=0}^{s-1} (-1)^j [2i+s+1+s-2j-1] \right) \mathbf{\Xi}^{(n-i-s)} \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{[i+1]_2} \sum_{s=1}^{n-i} n^{-i} \lambda_{n-i-s} (-1)^s [s] [2i+s+1] \mathbf{I}^{(n-i-s)} \\
&= \sum_{\ell=i+1}^n n^{-i} \lambda_{n-\ell} (-1)^{\ell-i} \frac{[\ell-i][\ell+i+1]}{[i+1]_2} \mathbf{I}^{(n-\ell)} \\
&= \sum_{\ell=i+1}^n \frac{(-1)^{\binom{\ell-i+1}{2}}}{d_\ell} \cdot \prod_{t=1}^i \frac{[\ell+1-t][\ell+t]}{[t]_2} (-1)^{\ell-i} \frac{[\ell-i][\ell+i+1]}{[i+1]_2} \mathbf{I}^{(n-\ell)} \\
&= \sum_{\ell=i+1}^n \frac{(-1)^{\binom{\ell-(i+1)-1}{2}}}{d_\ell} \cdot \prod_{t=1}^{i+1} \frac{[\ell+1-t][\ell+t]}{[t]_2} \mathbf{I}^{(n-\ell)}
\end{aligned}$$

which establishes (4.30).

Next, multiplying both numerator and denominator of (4.31) by $[2]^i$ (in the $\ell = i$ case) and using (4.25), we compute

$$\begin{aligned}
n^{-i} \lambda_{n-i} &= \frac{1}{d_i} \prod_{t=1}^i \frac{[2i+1] + (-1)^{i-t} [2t-1]}{[2t]} \\
&= \frac{1}{d_i \prod_{t=1}^i [2t]} \prod_{\substack{t=1 \\ t \equiv i \pmod{2}}}^i ([2i+1] + [2t-1]) \prod_{\substack{t=1 \\ t \equiv i+1 \pmod{2}}}^i ([2i+1] - [2t-1]) \\
&= \frac{\left(\frac{[2i][2]}{[1]} \frac{[2i-2][6]}{[3]} \frac{[2i-4][10]}{[5]} \frac{[2i-6][14]}{[7]} \dots \right) \left(\frac{[4i-2][2]}{[2i-1]} \frac{[4i-6][4]}{[2i-3]} \frac{[4i-10][6]}{[2i-5]} \frac{[4i-14][8]}{[2i-7]} \dots \right)}{d_i \prod_{t=1}^i [2t]} \\
&= \frac{1}{d_i} \prod_{t=1}^i \frac{[4t-2]}{[2t-1]} = 1.
\end{aligned}$$

Finally, recall that Proposition 4.20 gives that $\{\mathbf{I}^{(i)}\}_{i=0}^n$ is a basis for $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$. Since there is a unitriangular matrix relating this basis and $\{\mathbf{X}^{(i)}\}_{i=0}^n$, the latter is also a basis. \square

Remark 4.26. In Definition 4.24, we defined $\mathbf{X}^{(i)}$ for $0 \leq i \leq n$; however, we could have used the same recursion to define $\mathbf{X}^{(i)}$ for all $i \geq 0$. In fact, nothing is gained, since for all $i > n$, we find that $\mathbf{X}^{(i)} = 0$. To see this, it suffices to show that $\mathbf{X}^{(n+1)} = 0$. For this, Proposition 4.25 gives that $\mathbf{X}^{(n)} = \mathbf{I}^{(0)}$ and thus

$$\mathbf{X}^{(n)} \mathbf{X} = \mathbf{I}^{(0)} \mathbf{X} \stackrel{(4.26), (4.18)}{=} (-1)^{\binom{n}{2}} (-1)^{\binom{n+1}{2}} [n][n+1] \mathbf{I}^{(0)} = (-1)^n [n][n+1] \mathbf{X}^{(n)}.$$

and therefore $\mathbf{X}^{(n+1)} := \mathbf{X}^{(n)} \mathbf{X} - (-1)^n [n][n+1] \mathbf{X}^{(n)} = 0$. In particular, this implies that when $i = n+1$, we can interpret the right-hand side of (4.29) as the minimal polynomial of \mathbf{X} .

We now arrive at the main result of this section, the description given in Theorem 4.2 of the braiding on $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$ in terms of the basis $\{\mathbf{X}^{(i)}\}_{i=0}^n$. (Note that we have already established the rest of this theorem.) For this, we first need a technical result whose proof we relegate to the appendix.

Lemma 4.27. Fix $\ell \geq 1$, and set $\rho_{\ell+1}^{(\ell)} := 1$. For $1 \leq t \leq \ell$ consider the recurrence relation

$$(4.32) \quad \rho_t^{(\ell)} := (-1)^{\binom{\ell+2-t}{2}} + q^{-1} \frac{[\ell+1-t][\ell+t]}{[t]_2} \rho_{t+1}^{(\ell)}.$$

Then $\rho_1^{(\ell)} = (q^{-2})^{\binom{\ell+1}{2}}$.

Proof. See Appendix A. \square

Proposition 4.28. The braiding $R_{S,S} \in \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$ is given in the basis $\{\mathbf{X}^{(i)}\}_{i=0}^n$ as:

$$(4.33) \quad R_{S,S} = q^{\frac{n}{2}} \sum_{i=0}^n q^{-i} \mathbf{X}^{(i)}.$$

Proof. Proposition 4.25 gives that

$$\begin{aligned} q^{\frac{n}{2}} \sum_{i=0}^n q^{-i} \mathbf{X}^{(i)} &= q^{\frac{n}{2}} \text{id}_{S \otimes S} + q^{\frac{n}{2}} \sum_{i=1}^n q^{-i} \sum_{\ell=i}^n q^{-\ell} \lambda_{n-\ell} \mathbf{I}^{(n-\ell)} \\ &= q^{\frac{n}{2}} \text{id}_{S \otimes S} + \sum_{\ell=1}^n q^{\frac{n}{2}} \left(\sum_{i=1}^{\ell} q^{-i} \lambda_{n-\ell} \right) \mathbf{I}^{(n-\ell)}. \end{aligned}$$

The result then follows from Proposition 4.21 and the following calculation employing Lemma 4.27:

$$\begin{aligned} q^{\frac{n}{2}} \sum_{i=1}^{\ell} q^{-i} \lambda_{n-\ell} &= \frac{q^{\frac{n}{2}}}{d_{\ell}} \sum_{i=1}^{\ell} q^{-i} (-1)^{\binom{\ell-i+1}{2}} \prod_{t=1}^i \frac{[\ell+1-t][\ell+t]}{[t]^2} \\ &= \frac{q^{\frac{n}{2}}}{d_{\ell}} q^{-1} \frac{[\ell][\ell+1]}{[1]^2} \rho_2^{(\ell)} \\ &= \frac{q^{\frac{n}{2}}}{d_{\ell}} \left(\rho_1^{(\ell)} - (-1)^{\binom{\ell+1}{2}} \right) \\ &= \frac{q^{\frac{n}{2}}}{d_{\ell}} \left((q^{-2})^{\binom{\ell+1}{2}} - (-1)^{\binom{\ell+1}{2}} \right) \stackrel{(4.21)}{=} b_{n-\ell}. \quad \square \end{aligned}$$

Remark 4.29. By Proposition 4.25, the inverse braiding $R_{S,S}^{-1} \in \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$ can also be expressed as a $\mathbb{C}(q^{\frac{1}{2}})$ -linear combination of the elements $\{\mathbf{X}^{(i)}\}_{i=0}^n$. It is possible to show (either by analogous arguments to those used thus far, or as a consequence of Theorem 10.21 below) that

$$(4.34) \quad R_{S,S}^{-1} = q^{-\frac{n}{2}} \sum_{i=0}^n q^i \mathbf{X}^{(i)}.$$

Note that (4.28) allows for any product of the $\{\mathbf{X}^{(i)}\}_{i=0}^n$ to be expanded as a linear combination of these elements. Since they are linearly independent, this implies that (4.28) in fact suffices to confirm the formula (4.34). We will use this latter fact below.

4.4. Further relations in $\mathbf{Rep}(U_q(\mathfrak{so}_{2n+1}))$. In this section, we establish relations between various morphisms in the subcategory of $\mathbf{Rep}(U_q(\mathfrak{so}_{2n+1}))$ monoidally generated by the fundamental representations S and $V_1 = V(\varpi_1)$. Our purpose is two-fold. First, these relations will be used later for our decategorification results. Second, this lays the groundwork for giving a generators-and-relations presentation for the fundamental subcategory of $\mathbf{Rep}(U_q(\mathfrak{so}_{2n+1}))$, akin to that obtained for \mathfrak{sl}_N in [15] and \mathfrak{sp}_{2n} in [7], which we will pursue in future work.

The relations established here will be most conveniently described using the graphical language for monoidal categories, and we will use our conventions for diagrams established above (e.g. gray strands correspond to S), but with one simplification: since we only consider V_k when $k = 1$, we will drop the label 1 from our previous diagrammatics and let unlabelled black strands correspond to V_1 . We also abbreviate

$$(4.35) \quad \mathbf{H} := \mathbf{H}^{(1)} = \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} =: \begin{array}{c} | \quad | \\ \hline | \quad | \end{array}.$$

Example 4.30. In our present conventions, the $i = 1$ case of Lemma 4.11 is the relation

$$\text{Curl} = q^{\frac{-n+2}{2}} q^{-\{2, \dots, n\}} \text{Cup}$$

which immediately implies that

$$\text{Curl} = q^{\frac{n-2}{2}} q^{\{2, \dots, n\}} \text{Cup}.$$

Example 4.31. Similarly, in our present color-coding, (4.15) reads as:

$$\bigcirc = (-1)^{\binom{n+1}{2}} \prod_{i=1}^n (q^{2i-1} + q^{1-2i})$$

Observe that the $i = \ell = n$ case of the equality $n^{-i} \lambda_{n-i} = 1$ established in Proposition 4.25 gives that this latter quantity equals

$$(-1)^{n+\binom{n}{2}} \prod_{t=1}^n \frac{[n+1-t][n+t]}{[t]^2}.$$

Lemma 4.32. The relations

$$\text{Cap} = (-1)^n \frac{[2n+1]}{[2]} \quad \text{and} \quad \text{Cup} = (-1)^{n-1} \frac{[2n-1]}{[2]} \text{Cup}.$$

hold in $\mathbf{Rep}(U_q(\mathfrak{so}_{2n+1}))$.

Proof. The first equality follows from Remark 4.14, Lemma 4.19, and [8, Equation 1.1a]. We leave it to the reader to fill in the details, using the outline of the proof in [8, Lemma 4.1]. The second equality is the $i = 1$ case of Lemma 4.22. \square

Several of the remaining proofs in this section will use the fact that “fork-slide” relations hold in this graphical language, e.g.

$$\text{Fork} = \text{Slide}.$$

Such relations are consequences of the naturality of the braiding.

Lemma 4.33. The relations

$$\text{Curl} = (-1)^n q^{-2n} \text{Cup} \quad \text{and} \quad \text{Curl} = (-1)^n q^{2n} \text{Cup}$$

hold in $\mathbf{Rep}(U_q(\mathfrak{so}_{2n+1}))$.

Proof. The first relation follows by applying a fork-slide, resolving a curl using the $i = 0$ case of Lemma 4.11, and then using Example 4.30. The second is an immediate consequence of the first. \square

Lemma 4.34. The morphisms

$$(4.36) \quad \text{Cup} \quad \text{and} \quad \text{Cup}$$

are a basis of $\text{Hom}_{U_q(\mathfrak{so}_{2n+1})}(V_1 \otimes S, S \otimes V_1)$. The vertical reflections of these diagrams give a basis for $\text{Hom}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes V_1, V_1 \otimes S)$.

Proof. Since $V_1 \otimes S \cong V(\varpi_1 + \varpi_n) \oplus S \cong S \otimes V_1$, it suffices to show linear independence. Suppose there were $x, y \in \mathbb{C}(q^{\frac{1}{2}})$ such that x times the first morphism in (4.36) plus y times the second morphism is zero. Using Lemma 4.32 to evaluate this relation in two different ways by composing respectively on the bottom and the side with trivalent vertices, we obtain the system of linear equations

$$(-1)^{n-1} \frac{[2n-1]}{[2]} x + (-1)^n \frac{[2n+1]}{[2]} y = 0 \quad \text{and} \quad (-1)^n \frac{[2n+1]}{[2]} x + (-1)^{n-1} \frac{[2n-1]}{[2]} y = 0.$$

This system has unique solution $x = 0 = y$. The argument for their vertical reflections is identical. \square

Proposition 4.35. The braiding $R_{S, V_1} \in \text{Hom}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes V_1, V_1 \otimes S)$ and its inverse R_{S, V_1}^{-1} are given by

$$\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} = q \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} + q^{-1} \begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array} \quad \text{and} \quad \begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array} = q^{-1} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} + q \begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array}$$

Proof. The second equality follows from the first using pivotality (e.g. by rotating all diagrams by 90°), so it suffices to establish the first. By Lemma 4.34, there exist $x, y \in \mathbb{C}(q^{\frac{1}{2}})$ such that the braiding on $S \otimes V_1$ is x times the first diagram in (4.36) plus y times the second. Attaching a trivalent vertex to this equality of morphisms on the bottom and right hand side respectively and applying Lemma 4.33 gives the equations

$$\begin{aligned} (-1)^n q^{-2n} &= (-1)^{n-1} \frac{[2n-1]}{[2]} x + (-1)^n \frac{[2n+1]}{[2]} y \\ (-1)^n q^{2n} &= (-1)^n \frac{[2n+1]}{[2]} x + (-1)^{n-1} \frac{[2n-1]}{[2]} y \end{aligned}$$

respectively. It is easy to verify that $x = q$ and $y = q^{-1}$ gives the unique solution. \square

It is convenient to rewrite Proposition 4.35 as:

$$(4.37) \quad \begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array} = -q^2 \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} + q \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \quad \text{and} \quad \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} = -q^{-2} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} + q^{-1} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array}.$$

Corollary 4.36. The relations

$$\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} = -q^{-2} \begin{array}{c} \text{H} \\ \text{H} \end{array} + (-1)^n q^{-2n-1} \begin{array}{c} \cup \\ \cup \end{array} \quad \text{and} \quad \begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array} = -q^2 \begin{array}{c} \text{H} \\ \text{H} \end{array} + (-1)^n q^{2n+1} \begin{array}{c} \cup \\ \cup \end{array}$$

hold in $\text{Hom}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S, V_1 \otimes V_1)$.

Proof. We prove the first equality, which implies the second. Applying a fork-slide, then Lemma 4.33, we find

$$\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} = (-1)^n q^{-2n} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array}.$$

We then compute

$$(-1)^n q^{-2n} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} = (-1)^n q^{-2n} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \stackrel{(4.37)}{=} (-1)^n q^{-2n} \left(-q^{-2} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} + q^{-1} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \right)$$

and the result follows from Lemma 4.33 and invertibility of the braiding. \square

Lemma 4.37. The relation

$$\begin{array}{|c|} \hline \text{---} \\ \hline \text{---} \\ \hline \text{---} \\ \hline \end{array} + \begin{array}{|c|} \hline \text{---} \\ \hline \text{---} \\ \hline \text{---} \\ \hline \end{array} = -[2]_{q^2} \begin{array}{|c|} \hline \text{---} \\ \hline \text{---} \\ \hline \text{---} \\ \hline \end{array} + \begin{array}{|c|} \hline \text{---} \\ \hline \text{---} \\ \hline \text{---} \\ \hline \end{array}$$

holds in $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes 3})$.

Proof. The relations (4.37) imply

$$(4.38) \quad \begin{array}{|c|} \hline \text{---} \\ \hline \text{---} \\ \hline \text{---} \\ \hline \end{array} = q \begin{array}{|c|} \hline \text{---} \\ \hline \text{---} \\ \hline \text{---} \\ \hline \end{array} + q^{-1} \begin{array}{|c|} \hline \text{---} \\ \hline \text{---} \\ \hline \text{---} \\ \hline \end{array}$$

which gives

$$\begin{array}{|c|} \hline \text{---} \\ \hline \text{---} \\ \hline \text{---} \\ \hline \end{array} + \begin{array}{|c|} \hline \text{---} \\ \hline \text{---} \\ \hline \text{---} \\ \hline \end{array} = -[2]_{q^2} \begin{array}{|c|} \hline \text{---} \\ \hline \text{---} \\ \hline \text{---} \\ \hline \end{array} + q^{-1} \begin{array}{|c|} \hline \text{---} \\ \hline \text{---} \\ \hline \text{---} \\ \hline \end{array} + q \begin{array}{|c|} \hline \text{---} \\ \hline \text{---} \\ \hline \text{---} \\ \hline \end{array}.$$

Applying a forkslide, Corollary 4.36, and Lemma 4.33, we simplify the last diagram as

$$\begin{array}{|c|} \hline \text{---} \\ \hline \text{---} \\ \hline \text{---} \\ \hline \end{array} = -q^{-2} \begin{array}{|c|} \hline \text{---} \\ \hline \text{---} \\ \hline \text{---} \\ \hline \end{array} + (-1)^n q^{-2n-1} (-1)^n q^{2n} \begin{array}{|c|} \hline \text{---} \\ \hline \text{---} \\ \hline \text{---} \\ \hline \end{array}$$

and the result follows. \square

It is natural to consider Lemma 4.37 in terms of the morphisms $\{\mathbf{X}^{(i)}\}_{i=0}^n$ from Definition 4.24. For $m \geq 3$ and $1 \leq r \leq m-1$, let

$$\mathbf{H}_r := \text{id}_{S^{\otimes r-1}} \otimes \mathbf{H} \otimes \text{id}_{S^{\otimes m-r-1}} \in \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$$

and

$$\mathbf{X}_r^{(i)} := \text{id}_{S^{\otimes r-1}} \otimes \mathbf{X}^{(i)} \otimes \text{id}_{S^{\otimes m-r-1}} \in \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m}).$$

As above, we abbreviate $\mathbf{X}_r := \mathbf{X}_r^{(1)}$ and by definition $\mathbf{X}_r^{(0)} = \text{id}_{S^{\otimes m}}$. When $|r-s| \geq 2$, these morphisms satisfy the far-commutativity relation:

$$\mathbf{H}_r \mathbf{H}_s = \mathbf{H}_s \mathbf{H}_r \quad \text{and} \quad \mathbf{X}_r^{(i)} \mathbf{X}_s^{(j)} = \mathbf{X}_s^{(j)} \mathbf{X}_r^{(i)}.$$

In this notation, Lemma 4.37 is the relation

$$(4.39) \quad \mathbf{H}_2 \mathbf{H}_1 \mathbf{H}_1 + \mathbf{H}_1 \mathbf{H}_1 \mathbf{H}_2 = -[2]_{q^2} \mathbf{H}_1 \mathbf{H}_2 \mathbf{H}_1 + \mathbf{H}_2$$

which we now express in terms of the \mathbf{X}_i .

Proposition 4.38 (devil's Serre relation). The relations

$$(4.40) \quad \mathbf{X}_i \mathbf{X}_{i\pm 1} \mathbf{X}_i = \mathbf{X}_i^{(2)} \mathbf{X}_{i\pm 1} + \mathbf{X}_{i\pm 1} \mathbf{X}_i^{(2)} + [2] \mathbf{X}_i^{(2)} + \mathbf{X}_i$$

hold in $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$ whenever $\min(i, i \pm 1) \geq 1$ and $\max(i, i \pm 1) \leq m$.

Proof. It suffices to consider the $m=3$ case, and we will establish the relation

$$\mathbf{X}_1 \mathbf{X}_2 \mathbf{X}_1 = \mathbf{X}_1^{(2)} \mathbf{X}_2 + \mathbf{X}_2 \mathbf{X}_1^{(2)} + [2] \mathbf{X}_1^{(2)} + \mathbf{X}_1$$

since the other relation is obtained by conjugating with appropriate braids (or by making a similar computation). Recall from Definition 4.24, that

$$\mathbf{X} = \mathbf{H} - \frac{1}{[2]} \quad \text{and} \quad \mathbf{X}^{(2)} = \frac{-1}{[2]_{q^2}} \mathbf{X}(\mathbf{X} + [2])$$

so

$$(4.41) \quad \mathbf{X}^{(2)} = \frac{-1}{[2]_{q^2}} \left(\mathbf{H} - \frac{1}{[2]} \right) \left(\mathbf{H} + \frac{[3]}{[2]} \right) \quad \text{and} \quad \frac{-1}{[2]_{q^2}} \mathbf{H}^2 = \mathbf{X}^{(2)} + \frac{1}{[2]} \mathbf{H} - \frac{[3]}{[2]^2 [2]_{q^2}}.$$

Directly computing, we find

$$\begin{aligned} \mathbf{X}_1 \mathbf{X}_2 \mathbf{X}_1 &= \mathbf{H}_1 \mathbf{H}_2 \mathbf{H}_1 - \frac{1}{[2]} (\mathbf{H}_1 \mathbf{H}_2 + \mathbf{H}_1^2 + \mathbf{H}_2 \mathbf{H}_1) + \frac{1}{[2]^2} (2\mathbf{H}_1 + \mathbf{H}_2) - \frac{1}{[2]^3} \\ &\stackrel{(4.39)}{=} \frac{-1}{[2]_{q^2}} (\mathbf{H}_2 \mathbf{H}_1^2 + \mathbf{H}_1^2 \mathbf{H}_2 - \mathbf{H}_2) - \frac{1}{[2]} (\mathbf{H}_1 \mathbf{H}_2 + \mathbf{H}_1^2 + \mathbf{H}_2 \mathbf{H}_1) + \frac{1}{[2]^2} (2\mathbf{H}_1 + \mathbf{H}_2) - \frac{1}{[2]^3}. \end{aligned}$$

Similarly,

$$\mathbf{X}_1^{(2)} \mathbf{X}_2 + \mathbf{X}_2 \mathbf{X}_1^{(2)} \stackrel{(4.41)}{=} \frac{-1}{[2]_{q^2}} \left(\left(\mathbf{H}_1 - \frac{1}{[2]} \right) \left(\mathbf{H}_1 + \frac{[3]}{[2]} \right) \left(\mathbf{H}_2 - \frac{1}{[2]} \right) + \left(\mathbf{H}_2 - \frac{1}{[2]} \right) \left(\mathbf{H}_1 - \frac{1}{[2]} \right) \left(\mathbf{H}_1 + \frac{[3]}{[2]} \right) \right)$$

which expands as

$$\frac{-1}{[2]_{q^2}} \left(\mathbf{H}_1^2 \mathbf{H}_2 + \mathbf{H}_2 \mathbf{H}_1^2 - \frac{2}{[2]} \mathbf{H}_1^2 + \frac{[2]_{q^2}}{[2]} \mathbf{H}_1 \mathbf{H}_2 + \frac{[2]_{q^2}}{[2]} \mathbf{H}_2 \mathbf{H}_1 + 2 \frac{[2]_{q^2}}{[2]^2} \mathbf{H}_1 - 2 \frac{[3]}{[2]^2} \mathbf{H}_2 + 2 \frac{[3]}{[2]^3} \right).$$

Thus, we see that $\mathbf{X}_1 \mathbf{X}_2 \mathbf{X}_1 - \mathbf{X}_1^{(2)} \mathbf{X}_2 - \mathbf{X}_2 \mathbf{X}_1^{(2)}$ equals

$$-\frac{[2]}{[2]_{q^2}} \mathbf{H}_1^2 + \frac{1}{[2][2]_{q^2}} \stackrel{(4.41)}{=} [2] \mathbf{X}_1^{(2)} + \mathbf{H}_1 - \frac{[3]}{[2][2]_{q^2}} + \frac{1}{[2][2]_{q^2}} = [2] \mathbf{X}_1^{(2)} + \mathbf{X}_1. \quad \square$$

We now establish the connection discussed above in Section 1.7 between U -quantum groups and our elements $\mathbf{X}^{(i)} \in \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$.

Theorem 4.39. The surjective $\mathbb{C}(q)$ -algebra homomorphism $U'_{-q^2}(\mathfrak{so}_m) \rightarrow \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$ from Wenzl's Theorem 1.13 is such that

$$\mathbf{x}_i^{(k)} \mapsto \mathbf{X}_i^{(k)} \quad \text{and} \quad \sum_{k=0}^n q^{-k} \mathbf{x}_i^{(k)} \mapsto q^{-\frac{n}{2}} \text{id}_S^{\otimes i-1} \otimes R_{S,S} \otimes \text{id}_S^{\otimes m-i-1}.$$

Proof. Recall from Definition 1.16 that $\mathbf{x}_i \in U'_{-q^2}(\mathfrak{so}_m)$ are defined by $\mathbf{x}_i := b_i - \frac{1}{[2]}$ where the b_i are the standard generators of $U'_{-q^2}(\mathfrak{so}_m)$ in Definition 1.12. Lemma 4.37 then implies that the assignment $\mathbf{x}_i \mapsto \mathbf{X}_i$ indeed defines an algebra homomorphism $U'_{-q^2}(\mathfrak{so}_m) \rightarrow \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$. Comparing Definition 1.16 and (4.27), we see that this sends $\mathbf{x}_i^{(k)} \mapsto \mathbf{X}_i^{(k)}$. In Proposition B.10, we show that our endomorphism $\mathbf{H} \in \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes 2})$ agrees with Wenzl's endomorphism C , hence this is precisely Wenzl's homomorphism from Theorem 1.13. The remaining claim follows from Proposition 4.28. \square

4.5. Spin link polynomials. We now characterize the spin-colored link polynomials $P_{\mathfrak{so}_{2n+1}}(\mathcal{L}_\beta^S)$ in terms of the elements $\mathbf{X}_r^{(i)}$. Indeed, by Proposition 4.28, the endomorphism $R(\beta, S) \in \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$ assigned to a braid $\beta \in \text{Br}_m$ can be written as a linear combination of $\{\mathbf{X}_r^{(i)}\}_{1 \leq r \leq m-1}$ which, for each r , satisfy (4.27), i.e.

$$(4.42) \quad \mathbf{X}_r^{(i+1)} := \frac{(-1)^i}{[i+1]_{q^2}^2} \left(\mathbf{X}_r^{(i)} \mathbf{X}_r - (-1)^{i \llbracket [i][i+1] \rrbracket} \mathbf{X}_r^{(i)} \right).$$

We now show that, modulo a pair of conjectures that we have only verified in low rank ($n = 1, 2, 3$), the link invariant $P_{\mathfrak{so}_{2n+1}}(\mathcal{L}_\beta^S)$ is characterized by the braiding formula (4.33), the relations (4.27) and (4.40), and a compatibility between the quantum traces on $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$ and $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m-1})$. We will use this characterization for our decategorification results in §10.6.

To begin, we conjecture that relations (4.27) and (4.40) suffice to establish more-general versions of Proposition 4.38.

Conjecture 4.40. Let A_m^n be a unital $\mathbb{C}(q)$ -algebra and let $\{\mathbf{X}_r^{(i)}\}_{\substack{1 \leq r \leq m-1 \\ 0 \leq i \leq n}} \subset A_m^n$ be a collection of elements satisfying $\mathbf{X}_r^{(0)} = 1$ and equations (4.27) and (4.40). Then, given $1 \leq a, b, c \leq n$, there is a relation of the form

$$\mathbf{X}_i^{(a)} \mathbf{X}_{i\pm 1}^{(b)} \mathbf{X}_i^{(c)} = \xi \cdot \mathbf{X}_{i\pm 1}^{(a')} \mathbf{X}_i^{(b')} \mathbf{X}_{i\pm 1}^{(c')} + \text{LOT}_{a,b,c}$$

in A_m^n , where $\xi \in \mathbb{C}(q)$, $1 \leq a', b', c' \leq n$, and $\text{LOT}_{a,b,c}$ is a linear combination of terms of the form $\mathbf{X}_i^{(k)} \mathbf{X}_{i\pm 1}^{(\ell)}$ and $\mathbf{X}_{i\pm 1}^{(\ell)} \mathbf{X}_i^{(k)}$.

Proposition 4.41. Conjecture 4.40 holds when $n = 1, 2, 3$.

Proof. This is immediate from (4.40) when $n = 1$, since in that case $\mathbf{X}_r^{(a)} = 0$ when $a \geq 2$. We leave it as a (sometimes challenging) exercise to confirm that the following relations in A_m^3 can be derived using only (4.27) and (4.40):

$$(4.43) \quad \mathbf{X}_i^{(a)} \mathbf{X}_{i\pm 1}^{(1)} \mathbf{X}_i^{(c)} \begin{cases} \mathbf{X}_i^{(2)} \mathbf{X}_{i\pm 1} \mathbf{X}_i = \mathbf{X}_{i\pm 1} \mathbf{X}_i^{(3)} - "[2]^2" \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1} - [2] \mathbf{X}_i^{(2)} \mathbf{X}_{i\pm 1} - "[2]^2" \mathbf{X}_i^{(2)} - "[2][3]" \mathbf{X}_i^{(3)} \\ \mathbf{X}_i^{(2)} \mathbf{X}_{i\pm 1} \mathbf{X}_i^{(2)} = "[2][3]" (\mathbf{X}_{i\pm 1} \mathbf{X}_i^{(3)} + \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1} + \mathbf{X}_i^{(2)}) - (q^6 - 2q^4 - 2q^{-4} + q^{-6}) \mathbf{X}_i^{(3)} \\ \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1} \mathbf{X}_i = "[2][3]" \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1} + "[3]^2" \mathbf{X}_i^{(3)} \\ \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1} \mathbf{X}_i^{(2)} = (q^4 + q^2 + q^{-2} + q^{-4}) \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1} + (q^7 - q^5 - q^{-5} + q^{-7}) \mathbf{X}_i^{(3)} \\ \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1} \mathbf{X}_i^{(3)} = -(q^8 + q^2 + q^{-2} + q^{-8}) \mathbf{X}_i^{(3)} \end{cases}$$

$$(4.44) \quad \mathbf{X}_i^{(a)} \mathbf{X}_{i\pm 1}^{(2)} \mathbf{X}_i^{(c)} \begin{cases} \mathbf{X}_i \mathbf{X}_{i\pm 1}^{(2)} \mathbf{X}_i = \mathbf{X}_{i\pm 1} \mathbf{X}_i^{(2)} \mathbf{X}_{i\pm 1} \\ \mathbf{X}_i^{(2)} \mathbf{X}_{i\pm 1}^{(2)} \mathbf{X}_i = \mathbf{X}_{i\pm 1} \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1} + \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1}^{(2)} + [2] \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1} + \mathbf{X}_i^{(2)} \mathbf{X}_{i\pm 1} \\ \mathbf{X}_i^{(2)} \mathbf{X}_{i\pm 1}^{(2)} \mathbf{X}_i^{(2)} = -[2] \mathbf{X}_{i\pm 1} \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1} + \mathbf{X}_i^{(2)} - "[2]^2" (\mathbf{X}_{i\pm 1} \mathbf{X}_i^{(3)} + \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1}) - "[2][3]" \mathbf{X}_i^{(3)} \\ \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1}^{(2)} \mathbf{X}_i = -"[2]^2" \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1} - [2] \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1}^{(2)} \\ \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1}^{(2)} \mathbf{X}_i^{(2)} = "[2][3]" \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1} + "[3]^2" \mathbf{X}_i^{(3)} \\ \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1}^{(2)} \mathbf{X}_i^{(3)} = -"[3][4]" \mathbf{X}_i^{(3)} \end{cases}$$


$$(4.45) \quad \mathbf{X}_i^{(a)} \mathbf{X}_{i\pm 1}^{(3)} \mathbf{X}_i^{(c)} \begin{cases} \mathbf{X}_i \mathbf{X}_{i\pm 1}^{(3)} \mathbf{X}_i = \mathbf{X}_{i\pm 1}^{(2)} \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1}^{(2)} \\ \mathbf{X}_i^{(2)} \mathbf{X}_{i\pm 1}^{(3)} \mathbf{X}_i = \mathbf{X}_{i\pm 1} \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1}^{(2)} \\ \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1}^{(3)} \mathbf{X}_i = \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1}^{(2)} \\ \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1}^{(3)} \mathbf{X}_i^{(2)} = \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1} \\ \mathbf{X}_i^{(3)} \mathbf{X}_{i\pm 1}^{(3)} \mathbf{X}_i^{(3)} = \mathbf{X}_i^{(3)} \end{cases}$$

as well as those obtained from these by reversing the ordering of every monomial. The relations for A_m^2 are obtained from these by setting $\mathbf{X}_r^{(3)} = 0$. \square

The next ingredient in our characterization of $P_{\mathfrak{so}_{2n+1}}(\mathcal{L}_\beta^S)$ is a relation between the quantum traces on $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$ and $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m-1})$. Recall that, in the present setting, the quantum traces are the $\mathbb{C}(q^{\frac{1}{2}})$ -linear maps

$$\text{Tr}_q : \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m}) \rightarrow \text{End}_{U_q(\mathfrak{so}_{2n+1})}(\mathbb{C}(q^{\frac{1}{2}})) \cong \mathbb{C}(q^{\frac{1}{2}})$$

that are most easily described on $W \in \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$ in the graphical calculus for $\mathbf{Rep}(U_q(\mathfrak{so}_{2n+1}))$ by:




Note that Tr_q is trace-like with respect to composition in $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$, i.e. $\text{Tr}_q(W_1 \circ W_2) = \text{Tr}_q(W_2 \circ W_1)$.

Conjecture 4.42. Let $\iota: \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m-1}) \rightarrow \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$ be the inclusion given by $\iota(W) := W \otimes \text{id}_S$. If $0 \leq k \leq n$ and $W \in \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m-1})$, then

$$(4.46) \quad \text{Tr}_q(X_{m-1}^{(k)} \circ \iota(W)) = (-1)^{n(k+1) + \binom{n-k}{2}} \prod_{t=1}^{n-k} \frac{[n+1-t][n+t]}{[t]^2} \text{Tr}_q(W).$$

Proposition 4.43. Conjecture 4.42 holds when $k = 0, n$. It holds for all $0 \leq k \leq n$ when $n = 1, 2, 3$.

Proof. Graphically, (4.46) is the equality




Since $X^{(n)} = \mathfrak{I}^{(0)} = \mathbf{U}_n \circ \mathbf{\Omega}^n$, this makes clear that the $k = n$ case follows from (4.13), provided we interpret the product on the right-hand side of (4.46) as equaling 1. Similarly, since $X^{(0)} = \text{id}_{S \otimes S}$, we see that this reduces the $k = 0$ case to the assertion

$$(-1)^{n + \binom{n}{2}} \prod_{t=1}^n \frac{[n+1-t][n+t]}{[t]^2} = \bigcirc$$

which holds by Example 4.31. The remaining low-rank cases ($n = 2, k = 1$ and $n = 3, k = 1, 2$) follow from direct computations using the graphical relations in $\mathbf{Rep}(U_q(\mathfrak{so}_{2n+1}))$ established in §4.4. \square

The astute reader will notice that the product in (4.46) appears in the $i = n$ case of (4.31). Indeed, Conjecture 4.42 is a consequence of Proposition 4.25 and the following more-elemental conjecture (which is obvious when $k = 0$ or n).

Conjecture 4.44. For $0 \leq k \leq n$, the following holds in $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$:



Returning to the task at hand, we finally arrive at our desired characterization of the spin link polynomials.

Theorem 4.45. Fix $n \geq 1$ and suppose that for $m \geq 1$ there are unital $\mathbb{C}(q^{\frac{1}{2}})$ -algebras A_m^n such that:

- A1. (each) A_m^n contain elements $\{\mathbf{X}_r^{(i)}\}_{\substack{1 \leq r \leq m-1 \\ 0 \leq i \leq n}}$ satisfying $\mathbf{X}_r^{(0)} = 1$ and equations (4.42) and (4.40),
- A2. there are $\mathbb{C}(q^{\frac{1}{2}})$ -linear maps $\iota: A_{m-1}^n \rightarrow A_m^n$ that send $\mathbf{X}_r^{(i)} \mapsto \mathbf{X}_r^{(i)}$ for $1 \leq r \leq m-2$ and trace-like $\mathbb{C}(q^{\frac{1}{2}})$ -linear maps $T_m^n: A_m^n \rightarrow \mathbb{C}(q^{\frac{1}{2}})$ that satisfy (4.46),
- A3. $A_1^n \cong \mathbb{C}(q^{\frac{1}{2}})$ and $T_1^n(1) = (-1)^{\binom{n+1}{2}} \prod_{i=1}^n (q^{2i-1} + q^{1-2i})$, and
- A4. the assignment $\beta_i \mapsto q^{\frac{n}{2}} \sum_{\ell=0}^n q^{-\ell} \mathbf{X}_i^{(\ell)}$ determines braid group representations $R_m^n: \text{Br}_m \rightarrow A_m^n$.

Assuming that

- (*) $n = 1, 2, 3$, or (more generally) that Conjectures 4.40 and 4.42 hold

we have that $T_m^n(R_m^n(\beta)) = P_{\mathfrak{so}_{2n+1}}(\mathcal{L}_\beta^S)$ for any braid β .

Proof. First observe that, under the assumptions (*), Definition 4.24 and Propositions 4.38, 4.43, and 4.28 imply that the remaining hypotheses A1.–A4. hold when $A_m^n = \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$ and $T_m^n = \text{Tr}_q$. Since $P_{\mathfrak{so}_{2n+1}}$ is defined in terms of Tr_q and $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$, the result follows once we show that the hypotheses suffice to compute $T_m^n(R_m^n(\beta))$.

Remark 4.29 implies that the braid group representation is necessarily given on the inverses of the Artin generators as

$$R_m^n(\beta_i^{-1}) = q^{-\frac{n}{2}} \sum_{\ell=0}^n q^\ell \mathbf{X}_i^{(\ell)}.$$

Given $\beta \in \text{Br}_m$, the element $R_m^n(\beta) \in A_m^n$ is thus a $\mathbb{C}(q^{\frac{1}{2}})$ -linear combination of words in the elements $\{\mathbf{X}_r^{(i)}\}_{\substack{1 \leq r \leq m-1 \\ 0 \leq i \leq n}}$, so we need only show that the hypotheses suffice to compute $T_m^n(W)$ for any such word $W \in A_m^n$. For this we argue inductively on m , with the base case $m = 1$ holding by Hypothesis A3.

Now, let $W = \mathbf{X}_{r_1}^{(i_1)} \cdots \mathbf{X}_{r_k}^{(i_k)}$ be a word in the elements $\{\mathbf{X}_r^{(i)}\}_{\substack{1 \leq r \leq m-1 \\ 0 \leq i \leq n}} \subset A_m^n$. With m fixed, we now argue, by induction on the length k of the word W , that we can compute $T_m^n(W)$. For the base case $k = 1$, we observe that a length-one word necessarily takes the form $W = \mathbf{X}_{m-1}^{(p)} \iota(W')$ with $0 \leq p \leq n$ and W' a word in $\{\mathbf{X}_r^{(i)}\}_{\substack{1 \leq r \leq m-2 \\ 0 \leq i \leq n}} \subset A_{m-1}^n$. Hypothesis A2. then gives that

$$T_m^n(W) = (-1)^{n(k+1) + \binom{n-k}{2}} \prod_{t=1}^{n-k} \frac{[n+1-t][n+t]^{n-t}}{[t]^{2n-t}} T_{m-1}^n(W')$$

which in turn can be computed using the inductive hypothesis (with respect to m). Now suppose that $k \geq 2$. Given $W = \mathbf{X}_{r_1}^{(i_1)} \cdots \mathbf{X}_{r_k}^{(i_k)}$, consider the corresponding word $s_{r_1} \cdots s_{r_k}$ in the standard Coxeter generators of the symmetric group \mathfrak{S}_m . By standard results on (reduced) expressions in the symmetric group [6, Theorem 3.3.1], it is possible to use a sequence of the moves

$$(4.47) \quad s_r s_r \rightarrow \emptyset \quad , \quad s_r s_{r \pm 1} s_r \leftrightarrow s_{r \pm 1} s_r s_{r \pm 1}$$

to pass from the word $s_{r_1} \cdots s_{r_k}$ to a (reduced) word in \mathfrak{S}_m wherein the generator s_{m-1} appears only once. Equation (4.42) implies that

$$\mathbf{X}_r^{(i)} \mathbf{X}_r^{(j)} \in \text{span}_{\mathbb{C}(q^{\frac{1}{2}})} \{\mathbf{X}_r^{(i)}\}_{i=0}^n$$

which, together with Proposition 4.41 and Conjecture 4.40, implies that (4.47) holds in A_m^n up to scalars, modulo words of shorter length. It follows that we can use (4.42) and 4.40 to obtain

$$W = \xi \cdot \iota(W') \mathbf{X}_{m-1}^{(p)} \iota(W'') + \sum_{j=1}^M \xi_j \cdot W_j$$

for $\xi, \xi_j \in \mathbb{C}(q^{\frac{1}{2}})$ and where W_j are words of length strictly less than k . By induction (on k), we can compute $T_m^n(W_j)$, while Hypothesis A2. allows us to compute

$$\begin{aligned} T_m^n(\iota(W')\mathcal{X}_{m-1}^{(p)}\iota(W'')) &= T_m^n(\mathcal{X}_{m-1}^{(p)}\iota(W'')\iota(W')) \\ &= (-1)^{n(k+1)+\binom{n-k}{2}} \prod_{t=1}^{n-k} \frac{[n+1-t][n+t]}{[t]^2} T_{m-1}^n(W''W') \end{aligned}$$

and the result follows (by induction on m). \square

Example 4.46. The Temperley–Lieb algebras at circle value $-(q+q^{-1})$ satisfy the hypotheses of Theorem 4.45 when $n = 1$. This recovers the known fact that the Jones polynomial equals the spin-colored $U_q(\mathfrak{so}_3)$ link invariant. (In fact, the Temperley–Lieb algebras are simply equal to $\text{End}_{U_q(\mathfrak{so}_3)}(S^{\otimes m})$.)

We will later use Theorem 4.45 to show¹⁴ that the link homology theories defined in Section 10 categorify the spin link polynomials $P_{\mathfrak{so}_{2n+1}}(\mathcal{L}_\beta^S)$.

Remark 4.47. According to Reshetikhin [60, Proposition 9.3], the relation from Corollary 4.36 (combined with the circle relations and the ribbon category relations) are sufficient to evaluate any closed braided graph, where each vertex is a grey-grey-black trivalent vertex. However, in the present paper, we do not categorify webs with edges colored by the vector representation. Thus, in order to connect the spin colored \mathfrak{so}_{2n+1} link polynomial with our categorification, we require Theorem 4.45.

5. BACKGROUND ON CATEGORIFIED QUANTUM GROUPS

In this section, we review the categorification of quantum groups in type A . We consider quantum \mathfrak{gl}_m , rather than \mathfrak{sl}_m , since it is the categorification of the former that is most relevant to our approach to link invariants.

5.1. The idempotent form of the quantum group. Categorified quantum \mathfrak{gl}_m does not actually categorify $U_q(\mathfrak{gl}_m)$; rather, it categorifies the following close relative which was first studied in [5, 43].

Definition 5.1. The *idempotent quantum group* $\dot{U}_q(\mathfrak{gl}_m)$ is the (non-unital) $\mathbb{C}(q)$ -algebra generated by mutually orthogonal idempotents $\mathbb{1}_{\mathbf{a}}$ for $\mathbf{a} \in \mathbb{Z}^m$ and elements $\mathbb{1}_{\mathbf{a}+\alpha_i}e_i\mathbb{1}_{\mathbf{a}}$ and $\mathbb{1}_{\mathbf{a}-\alpha_i}f_i\mathbb{1}_{\mathbf{a}}$ for $1 \leq i \leq m$. Set $e_i^{(r)} := \frac{1}{[r]!}e_i^r$ and $f_i^{(r)} := \frac{1}{[r]!}f_i^r$, then the relations are as follows:

- $e_i^{(r)}f_i^{(s)}\mathbb{1}_{\mathbf{a}} = \sum_t [\alpha_i^\vee(\mathbf{a})+r-s] f_i^{(s-t)}e_i^{(r-t)}\mathbb{1}_{\mathbf{a}}$,
- for $i \neq j$, $e_i^{(r)}f_j^{(s)}\mathbb{1}_{\mathbf{a}} = f_j^{(s)}e_i^{(r)}\mathbb{1}_{\mathbf{a}}$,
- for $|i-j| = 1$, $e_i e_j e_i \mathbb{1}_{\mathbf{a}} = (e_i^{(2)}e_j + e_j e_i^{(2)})\mathbb{1}_{\mathbf{a}}$ and $f_i f_j f_i \mathbb{1}_{\mathbf{a}} = (f_i^{(2)}f_j + f_j f_i^{(2)})\mathbb{1}_{\mathbf{a}}$,
- for $|i-j| > 1$, $e_i^{(r)}e_j^{(s)}\mathbb{1}_{\mathbf{a}} = e_j^{(s)}e_i^{(r)}\mathbb{1}_{\mathbf{a}}$ and $f_i^{(r)}f_j^{(s)}\mathbb{1}_{\mathbf{a}} = f_j^{(s)}f_i^{(r)}\mathbb{1}_{\mathbf{a}}$, and
- $e_i^{(s)}e_i^{(r)}\mathbb{1}_{\mathbf{a}} = [r+s]e_i^{(r+s)}\mathbb{1}_{\mathbf{a}}$ and $f_i^{(s)}f_i^{(r)}\mathbb{1}_{\mathbf{a}} = [r+s]f_i^{(r+s)}\mathbb{1}_{\mathbf{a}}$.

The *integral idempotent quantum group* ${}_{\mathbb{Z}}\dot{U}_q(\mathfrak{gl}_m)$ is the (non-unital) $\mathbb{Z}[q^\pm]$ -subalgebra of $\dot{U}_q(\mathfrak{gl}_m)$ generated by mutually orthogonal idempotents $\mathbb{1}_{\mathbf{a}}$ for $\mathbf{a} \in \mathbb{Z}^m$ and elements $\mathbb{1}_{\mathbf{a}+r\alpha_i}e_i^{(r)}\mathbb{1}_{\mathbf{a}}$ and $\mathbb{1}_{\mathbf{a}-\alpha_i}f_i^{(r)}\mathbb{1}_{\mathbf{a}}$ for $1 \leq i \leq m$ and $r \geq 1$.

Note that the (non-integral) idempotent quantum group is recovered from the integral version as $\dot{U}_q(\mathfrak{gl}_m) := \mathbb{C}(q) \otimes_{\mathbb{Z}[q^\pm]} {}_{\mathbb{Z}}\dot{U}_q(\mathfrak{gl}_m)$.

Remark 5.2. Since $\dot{U}_q(\mathfrak{gl}_m)$ is an algebra equipped with a system of mutually orthogonal idempotents indexed by $\mathbf{a} \in \mathbb{Z}^m$, we can consider it as a category wherein the objects are $\mathbf{a} \in \mathbb{Z}^m$ and $\text{Hom}_{\dot{U}_q(\mathfrak{gl}_m)}(\mathbf{a}, \mathbf{b})$ consists of elements of the form $\mathbb{1}_{\mathbf{b}}x\mathbb{1}_{\mathbf{a}}$.

¹⁴Assuming Conjectures 4.40, 4.42, and 10.36 when $n > 3$.

5.2. The categorified quantum group. For the rest of this chapter, let \mathbb{K} be an integral domain. We will explicitly assume \mathbb{K} is a field when we state results about the Grothendieck group.

We next define our version of the categorified quantum group $\mathcal{U}_q(\mathfrak{gl}_m)$. The original definition of $\mathcal{U}_q(\mathfrak{gl}_m)$ given in [44] by Mackaay–Stošić–Vaz (MSV) is a (signed) analogue of the Khovanov–Lauda–Rouquier [34, 63] categorified quantum group $\mathcal{U}_q(\mathfrak{sl}_m)$ wherein the \mathfrak{sl}_m weight lattice (the set of objects in $\mathcal{U}_q(\mathfrak{sl}_m)$) is replaced by the \mathfrak{gl}_m weight lattice. Our version is related, but not identical, to that used by MSV, and we now clarify the differences for the expert. Recall that bubbles are certain 2-morphisms living in the endomorphism algebra of an identity 1-morphism. The two differences are as follows:

- In MSV, bubbles may be identified with symmetric functions in an alphabet of variables. From the perspective of the \mathfrak{gl}_N foam 2-category in [53], which is related to $\mathcal{U}_q(\mathfrak{gl}_m)$ via categorical skew Howe duality, one should consider symmetric functions in m alphabets, and these bubbles should be viewed as symmetric functions in a *difference* of two adjacent alphabets. We extend scalars in the endomorphism algebras of all identity 1-morphisms accordingly. This modification has previously appeared (in a slightly different guise) in work of Webster [72].
- We work with a different orientation for the \mathfrak{sl}_m Dynkin diagram, and hence different scalars associated to degree-zero bubbles.

Remark 5.3. Experts may be familiar with Lauda’s paper [40], which parametrizes all choices of scalars one might use for the degree-zero bubbles. In the conventions thereof, for each \mathfrak{gl}_m weight $\mathbf{a} = (a_1, \dots, a_m) \in \mathbb{Z}^m$ we set

$$(5.1a) \quad c_{i,\mathbf{a}}^- = (-1)^{a_i}, \quad c_{i,\mathbf{a}}^+ = (-1)^{a_i-1}.$$

Via [40, Equations (2.1), (2.2)], this determines the parameters

$$(5.1b) \quad t_{i,i} = -1, \quad t_{i,i-1} = -1, \quad t_{i,i+1} = 1, \quad t_{i,k} = 1 \text{ otherwise.}$$

Note that (5.1a) are the values of the anti-clockwise and clockwise degree-zero bubbles, respectively.

Before proceeding, we note that neither of the aforementioned differences affect any salient properties of $\mathcal{U}_q(\mathfrak{gl}_m)$, and the following categorification result of Khovanov–Lauda still holds.

Theorem 5.4 ([34]). Suppose \mathbb{K} is a field. There is an isomorphism of $\mathbb{C}(q)$ -algebras

$$\mathbb{C}(q) \otimes_{\mathbb{Z}[q^\pm]} K_0(\text{Kar}(\mathcal{U}_q(\mathfrak{gl}_m))) \cong \dot{U}_q(\mathfrak{gl}_m).$$

□

We now give our definition of $\mathcal{U}_q(\mathfrak{gl}_m)$.

Definition 5.5. Let $m \geq 1$. The categorified quantum group $\mathcal{U}_q(\mathfrak{gl}_m)$ is the \mathbb{Z} -additive closure of the \mathbb{Z} -graded \mathbb{K} -linear 2-category given as follows.

- **Objects** are elements $\mathbf{a} \in \mathbb{Z}^m$.
- **1-morphisms** are generated by

$$\mathcal{E}_i \mathbb{1}_{\mathbf{a}} : \mathbf{a} \rightarrow \mathbf{a} + \alpha_i, \quad \mathcal{F}_i \mathbb{1}_{\mathbf{a}} : \mathbf{a} \rightarrow \mathbf{a} - \alpha_i,$$

for $1 \leq i \leq m-1$. Here $\alpha_i = (0, \dots, 1, -1, \dots, 0)$.

- **2-morphisms** are \mathbb{K} -linear combinations of “string diagrams” that are generated via horizontal and vertical composition by the generators:

$$(5.2) \quad \begin{array}{l} \begin{array}{c} \mathbf{a} + \alpha_i \uparrow \\ \bullet \\ \downarrow \\ i \end{array} \in \text{End}^2(\mathcal{E}_i \mathbb{1}_{\mathbf{a}}), \quad \begin{array}{c} \mathbf{a} + \alpha_i + \alpha_j \\ \diagup \quad \diagdown \\ i \quad j \\ \downarrow \end{array} \in \text{Hom}^{-i,j}(\mathcal{E}_i \mathcal{E}_j \mathbb{1}_{\mathbf{a}}, \mathcal{E}_j \mathcal{E}_i \mathbb{1}_{\mathbf{a}}), \\ \\ \begin{array}{c} \mathbf{a} \\ \curvearrowright \\ i \end{array} \in \text{Hom}^{1+\mathbf{a}_i-\mathbf{a}_{i+1}}(\mathcal{F}_i \mathcal{E}_i \mathbb{1}_{\mathbf{a}}, \mathbb{1}_{\mathbf{a}}), \quad \begin{array}{c} \mathbf{a} \\ \curvearrowleft \\ i \end{array} \in \text{Hom}^{1-\mathbf{a}_i+\mathbf{a}_{i+1}}(\mathcal{E}_i \mathcal{F}_i \mathbb{1}_{\mathbf{a}}, \mathbb{1}_{\mathbf{a}}), \\ \\ \begin{array}{c} i \\ \curvearrowright \\ \mathbf{a} \end{array} \in \text{Hom}^{1+\mathbf{a}_i-\mathbf{a}_{i+1}}(\mathbb{1}_{\mathbf{a}}, \mathcal{F}_i \mathcal{E}_i \mathbb{1}_{\mathbf{a}}), \quad \begin{array}{c} i \\ \curvearrowleft \\ \mathbf{a} \end{array} \in \text{Hom}^{1-\mathbf{a}_i+\mathbf{a}_{i+1}}(\mathbb{1}_{\mathbf{a}}, \mathcal{E}_i \mathcal{F}_i \mathbb{1}_{\mathbf{a}}) \end{array}$$

and for each $f \in \text{Sym}(\mathbb{X}_1 | \cdots | \mathbb{X}_m)$, a *new bubble* generator in each weight:

$$(5.3) \quad \boxed{f}^{\mathbf{a}} \in \text{End}^{2 \deg(f)}(\mathbb{1}_{\mathbf{a}}).$$

Here, the alphabets $\mathbb{X}_1, \dots, \mathbb{X}_m$ are viewed as formal. That is, for each $1 \leq i \leq m$, $\text{Sym}(\mathbb{X}_i)$ is the ring of symmetric functions, and $\text{Sym}(\mathbb{X}_1 | \cdots | \mathbb{X}_m) \cong \bigotimes_{i=1}^m \text{Sym}(\mathbb{X}_i)$.

These 2-morphisms are subject to the following local relations:

- (1) **Adjunction and cyclicity:** diagrams that are related by planar isotopy (rel. boundary) are equal. This is guaranteed by the following relations:
 - (a) The cap and cup morphisms in (5.2) are the units and counits for (graded) biadjunctions between $\mathcal{E}_i \mathbb{1}_{\mathbf{a}}$ and $\mathcal{F}_i \mathbb{1}_{\mathbf{a} + \alpha_i}$.
 - (b) The dot and crossing morphisms in (5.2) are cyclic with respect to this biadjoint structure, meaning that the two ways of building a dotted downward strand are equal, as are the two ways of building a downward crossing.
- (2) **Dot slide:**

$$(5.4) \quad \begin{array}{c} \begin{array}{c} \uparrow \quad \uparrow \\ \bullet \quad \bullet \\ \diagdown \quad \diagup \\ i \quad j \end{array} \mathbf{a} - \begin{array}{c} \uparrow \quad \uparrow \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ i \quad j \end{array} \mathbf{a} = \begin{array}{c} \uparrow \quad \uparrow \\ \bullet \quad \bullet \\ \diagdown \quad \diagup \\ i \quad j \end{array} \mathbf{a} - \begin{array}{c} \uparrow \quad \uparrow \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ i \quad j \end{array} \mathbf{a} = \begin{cases} \begin{array}{c} \uparrow \quad \uparrow \\ \uparrow \quad \uparrow \\ i \quad i \end{array} \mathbf{a} & \text{if } i = j \\ 0 & \text{else} \end{cases} \end{array}$$

- (3) **Quadratic KLR relation:**

$$(5.5) \quad \begin{array}{c} \begin{array}{c} \uparrow \quad \uparrow \\ \diagdown \quad \diagup \\ \uparrow \quad \uparrow \\ i \quad j \end{array} \mathbf{a} = \begin{cases} 0 & \text{if } i = j \\ (j - i) \left(\begin{array}{c} \uparrow \quad \uparrow \\ \bullet \quad \bullet \\ \uparrow \quad \uparrow \\ i \quad j \end{array} \mathbf{a} - \begin{array}{c} \uparrow \quad \uparrow \\ \uparrow \quad \uparrow \\ \bullet \quad \bullet \\ i \quad j \end{array} \mathbf{a} \right) & \text{if } i \cdot j = -1 \\ \begin{array}{c} \uparrow \quad \uparrow \\ \uparrow \quad \uparrow \\ i \quad j \end{array} \mathbf{a} & \text{if } i \cdot j = 0 \end{cases} \end{array}$$

(4) **Cubic KLR relation:**

$$(5.6) \quad \begin{array}{c} \begin{array}{ccc} \nearrow & \nearrow & \mathbf{a} \\ \searrow & \searrow & \\ i & j & k \end{array} - \begin{array}{ccc} \nearrow & \nearrow & \mathbf{a} \\ \searrow & \searrow & \\ i & j & k \end{array} = \begin{cases} \begin{array}{c} \uparrow \uparrow \uparrow \mathbf{a} \\ (j-i) \quad i \quad j \quad i \end{array} & \text{if } k = i \text{ and } i \cdot j = -1 \\ 0 & \text{else} \end{cases} \end{array}$$

(5) **New bubble relations:**

(a) **Algebra compatibility:**

$$\boxed{f} \boxed{g} \mathbf{a} = \boxed{fg} \mathbf{a}, \quad \boxed{f} \mathbf{a} + \boxed{g} \mathbf{a} = \boxed{f+g} \mathbf{a}$$

(b) **Bubbles to new bubbles:** Let us define spaded bubble notation:

$$(5.7) \quad \begin{array}{c} \begin{array}{c} \mathbf{a} \\ \circlearrowleft \\ \spadesuit+r \end{array} := \begin{array}{c} \mathbf{a} \\ \circlearrowleft \\ r-1-a_i+a_{i+1} \end{array} \in \text{End}^{2r}(\mathbb{1}_{\mathbf{a}}), \quad \begin{array}{c} \mathbf{a} \\ \circlearrowright \\ \spadesuit+r \end{array} := \begin{array}{c} \mathbf{a} \\ \circlearrowright \\ r-1+a_i-a_{i+1} \end{array} \in \text{End}^{2r}(\mathbb{1}_{\mathbf{a}}). \end{array}$$

When the number of dots is a non-negative integer, this picture is called a *real bubble*, and represents a genuine composition of a cup, a cap, and a number of dots. When the number of dots is a negative integer, both sides of (5.7) are formal symbols, often called *fake bubbles*. The following equation (5.8) serves two purposes: it gives a definition of fake bubbles as genuine morphisms, and it provides a relation between real bubbles and symmetric functions:

$$(5.8) \quad \begin{array}{c} \mathbf{a} \\ \circlearrowleft \\ \spadesuit+r \end{array} = (-1)^{a_i} \boxed{h_r(\mathbb{X}_{i+1} - \mathbb{X}_i)} \mathbf{a}, \quad \begin{array}{c} \mathbf{a} \\ \circlearrowright \\ \spadesuit+r \end{array} = (-1)^{a_i-1} \boxed{h_r(\mathbb{X}_i - \mathbb{X}_{i+1})} \mathbf{a}.$$

Thus when $r < 0$, a spaded bubble with decoration $\spadesuit+r$ is zero.

(c) **Dots to new bubbles:**

$$(5.9) \quad \begin{array}{c} \uparrow \\ \bullet \\ i \end{array} = \boxed{e_1(\mathbb{X}_i)} \begin{array}{c} \uparrow \\ i \end{array} - \begin{array}{c} \uparrow \\ i \end{array} \boxed{e_1(\mathbb{X}_i)} = \begin{array}{c} \uparrow \\ i \end{array} \boxed{e_1(\mathbb{X}_{i+1})} - \boxed{e_1(\mathbb{X}_{i+1})} \begin{array}{c} \uparrow \\ i \end{array}$$

(6) **Extended \mathfrak{sl}_2 relations:**

$$(5.10) \quad \begin{array}{c} \uparrow \downarrow \\ i \quad i \end{array} \mathbf{a} = \begin{array}{c} \nearrow \searrow \\ i \quad i \end{array} \mathbf{a} - \sum_{\substack{p+q+r= \\ a_i-a_{i+1}-1}} \begin{array}{c} \uparrow \\ i \\ \downarrow \\ p \end{array} \begin{array}{c} \uparrow \\ i \\ \downarrow \\ q \end{array} \begin{array}{c} \mathbf{a} \\ \circlearrowleft \\ \spadesuit+r \end{array}$$

$$(5.11) \quad \begin{array}{c} \downarrow \uparrow \\ i \quad i \end{array} \mathbf{a} = \begin{array}{c} \searrow \nearrow \\ i \quad i \end{array} \mathbf{a} - \sum_{\substack{p+q+r= \\ -a_i+a_{i+1}-1}} \begin{array}{c} \downarrow \\ i \\ \uparrow \\ p \end{array} \begin{array}{c} \downarrow \\ i \\ \uparrow \\ q \end{array} \begin{array}{c} \mathbf{a} \\ \circlearrowright \\ \spadesuit+r \end{array}$$

(By convention, the summations on the right-hand side are zero when this requires $p + q + r < 0$.)

(7) **Mixed \mathcal{E}, \mathcal{F} relations:** for $i \neq j$,

$$(5.12) \quad \begin{array}{c} \uparrow \\ i \end{array} \begin{array}{c} \downarrow \\ j \end{array} \mathbf{a} = \begin{array}{c} \uparrow \downarrow \\ i \quad j \end{array} \mathbf{a}, \quad \begin{array}{c} \downarrow \\ i \end{array} \begin{array}{c} \uparrow \\ j \end{array} \mathbf{a} = \begin{array}{c} \downarrow \uparrow \\ i \quad j \end{array} \mathbf{a}$$

Remark 5.6. We collect several easy consequences of the new bubble relations. Most important are the standard **Bubble relations**, which are signed versions of the relations originally appearing in [39].

(1) **Non-positive degree relations:**

$$\begin{array}{c} \mathbf{a} \\ \circlearrowleft \\ \spadesuit_{+r} \end{array} = \begin{cases} 0 & \text{if } r < 0 \\ (-1)^{a_i} & \text{if } r = 0 \end{cases}, \quad \begin{array}{c} \mathbf{a} \\ \circlearrowright \\ \spadesuit_{+r} \end{array} = \begin{cases} 0 & \text{if } r < 0 \\ (-1)^{a_i-1} & \text{if } r = 0 \end{cases}$$

Compare with Remark 5.3.

(2) **Infinite Grassmannian relation:** the equality

$$\left(\sum_{r=0}^{\infty} \begin{array}{c} \mathbf{a} \\ \circlearrowleft \\ \spadesuit_{+r} \end{array} t^r \right) \left(\sum_{r=0}^{\infty} \begin{array}{c} \mathbf{a} \\ \circlearrowright \\ \spadesuit_{+r} \end{array} t^r \right) = -1$$

holds in the formal power series ring $\text{End}(\mathbb{1}_{\mathbf{a}})[[t]]$.

Lastly, **symmetric generators slide:** let $f \in \text{Sym}(\mathbb{X}_1 | \cdots | \mathbb{X}_i + \mathbb{X}_{i+1} | \cdots | \mathbb{X}_m)$, then

$$(5.13) \quad \begin{array}{c} \uparrow \\ i \end{array} \begin{array}{c} \circlearrowleft \\ f \end{array} \mathbf{a} = \begin{array}{c} \circlearrowleft \\ f \end{array} \begin{array}{c} \uparrow \\ i \end{array} \mathbf{a}.$$

Remark 5.7. When $m = 1$, there are no generating 1-morphisms $\mathcal{E}_i \mathbb{1}_{\mathbf{a}}$ or $\mathcal{F}_i \mathbb{1}_{\mathbf{a}}$, so all 1-morphisms in $\mathcal{U}_q(\mathfrak{gl}_1)$ are (direct sums of shifts of) identity 1-morphisms $\mathbb{1}_{\mathbf{a}}$. Hence, $\text{End}(\mathbb{1}_{\mathbf{a}}) \cong \text{Sym}(\mathbb{X}_1)$ is the \mathbb{Z} -graded \mathbb{K} -algebra of new bubbles (5.3).

By convention, $\mathcal{U}_q(\mathfrak{gl}_0)$ is a 2-category with one object (the zero weight in a zero-dimensional lattice). Again all 1-morphisms are direct sums of shifts of the identity 1-morphism $\mathbb{1}_0$, but now $\text{End}(\mathbb{1}_0) = \mathbb{K}$.

Remark 5.8. Note that each strand in a string diagram in $\mathcal{U}_q(\mathfrak{gl}_m)$ carries a label $i \in \{1, \dots, m-1\}$, i.e. a label by a node of the \mathfrak{gl}_m Dynkin diagram. We will refer to these labels as *colors*, since later we will depict them using colored strands. See Convention 5.18.

Remark 5.9. The 2-category $\coprod_{m \geq 0} \mathcal{U}_q(\mathfrak{gl}_m)$ is monoidal, via the *external tensor product*

$$\boxtimes: \mathcal{U}_q(\mathfrak{gl}_{m_1}) \times \mathcal{U}_q(\mathfrak{gl}_{m_2}) \rightarrow \mathcal{U}_q(\mathfrak{gl}_{m_1+m_2})$$

which is given on objects by

$$((a_1, \dots, a_{m_1}), (b_1, \dots, b_{m_2})) \mapsto (a_1, \dots, a_{m_1}, b_1, \dots, b_{m_2})$$

and on 1- and 2-morphisms by “raising” the colors on the second factor by m_1 and then concatenating the 1- and 2-morphisms. For example, $\boxtimes: \mathcal{U}_q(\mathfrak{gl}_2) \times \mathcal{U}_q(\mathfrak{gl}_3) \rightarrow \mathcal{U}_q(\mathfrak{gl}_5)$ sends

$$\left(\mathcal{E}_1 \mathbb{1}_{(a,b)}, \mathcal{F}_1 \mathcal{E}_2 \mathbb{1}_{(c,d,e)} \right) \mapsto \mathcal{E}_1 \mathcal{F}_3 \mathcal{E}_4 \mathbb{1}_{(a,b,c,d,e)}, \quad \left(\begin{array}{c} \uparrow \\ 1 \end{array} \begin{array}{c} \circlearrowleft \\ (a,b) \end{array}, \begin{array}{c} \uparrow \downarrow \\ 1 \quad 2 \end{array} \begin{array}{c} \circlearrowright \\ (c,d,e) \end{array} \right) \mapsto \begin{array}{c} \uparrow \\ 1 \end{array} \begin{array}{c} \uparrow \downarrow \\ 3 \quad 4 \end{array} \begin{array}{c} \circlearrowright \\ (a,b,c,d,e) \end{array}.$$

5.3. Digression on parameters. The parameters governing the relations in quantum groups have evolved over time. We have chosen our sign conventions in the definition of $\mathcal{U}_q(\mathfrak{gl}_m)$ above carefully, to match signs appearing in foams and singular Soergel bimodules, for future ease of use. However, we will need to use relations derived by Lauda [39] and Khovanov–Lauda–Mackaay–Stošić (KLMS) [35], which were provided in a version of the 2-category defined with different sign conventions. In [40, Section 3.1], Lauda constructs a functor F that is an equivalence from one particular version of the categorified quantum group to any other version. By taking the source of F to be the version used in [39] and [35] and the target of F to be our version $\mathcal{U}_q(\mathfrak{gl}_m)$, we are able to translate relations in [35] into relations in $\mathcal{U}_q(\mathfrak{gl}_m)$.

Remark 5.10. In Remark 5.3 we discussed the parameters used for $\mathcal{U}_q(\mathfrak{gl}_m)$. To apply [40, Section 3.1], one also needs to know the parameters for the version used by [39] and [35]. In the language of [40, Section 3.1] this is $c_{i,\mathbf{a}} = 1$.

Lauda’s functor F rescales the cups and caps by certain scalars which depend in a complicated way on the ambient weight. Fortunately, when F is applied to a bubble, which has a cap paired with a cup in the same ambient weight, the complicated individual scaling factors multiply to an easy overall scaling factor. Indeed, an i -colored clockwise bubble in region \mathbf{a} is rescaled by $(-1)^{a_i-1}$. This is as it must be, since the degree zero clockwise bubble in [35] is equal to the scalar 1 (times an identity map), while the degree zero clockwise bubble in $\mathcal{U}_q(\mathfrak{gl}_m)$ is the scalar $(-1)^{a_i-1}$. Moreover, whenever any i -colored clockwise cup and clockwise cap appear in a region of the same weight \mathbf{a} , even if they are not closed into a bubble, they will contribute a scaling factor of $(-1)^{a_i-1}$. An example of such a pair appears in the RHS of (5.10).

Similarly, a pair of an i -colored counterclockwise cup and counterclockwise cap in weight \mathbf{a} will contribute $(-1)^{a_i}$ to the scaling factor. For example, the reader can confirm that the bubble slide relations from [39, Proposition 5.7] pick up a sign when translated to a relation in $\mathcal{U}_q(\mathfrak{gl}_m)$, giving

$$(5.14) \quad \begin{array}{c} \downarrow \\ \begin{array}{c} \textcirclearrowright \\ \blacklozenge_{+r} \end{array} \\ \downarrow \\ i \end{array} \quad \mathbf{a} = - \sum_{k=0}^2 (-1)^k \binom{2}{k} \begin{array}{c} \downarrow \\ \begin{array}{c} \textcirclearrowright \\ \blacklozenge_{+r-k} \end{array} \\ \downarrow \\ i \end{array} \quad \mathbf{a} \quad .$$

Meanwhile, the usual curl relations [39, Proposition 5.4] hold verbatim, e.g.

$$(5.15) \quad \begin{array}{c} \uparrow \\ \textcirclearrowleft \\ \downarrow \\ i \end{array} \quad \mathbf{a} = - \sum_{p+q=a_{i+1}-a_i} \begin{array}{c} \uparrow \\ \bullet_p \\ \downarrow \\ i \end{array} \quad \begin{array}{c} \textcirclearrowright \\ \blacklozenge_{+q} \end{array} \quad \mathbf{a} \quad .$$

While upwards-oriented crossings are not rescaled by F , sideways crossings do pick up a scaling factor, since they are defined as compositions of upward crossings with caps and cups. Again, the general scaling factor on a single sideways crossing is complicated, but the overall scaling factor on a pair of uni-colored sideways crossings (one facing right, one facing left) in the same region is -1 , independent of the weight \mathbf{a} . For example, this explains the coefficient of the first term on the RHS of (5.10), which differs from the relation in [39] by a sign.

5.4. Thick calculus. The 2-category $\mathcal{U}_q(\mathfrak{gl}_m)$ is not Karoubian, e.g. the idempotent 2-morphism

$$(5.16) \quad \begin{array}{c} \uparrow \quad \uparrow \\ \textcirclearrowleft \\ \downarrow \quad \downarrow \\ i \quad i \end{array} \quad \mathbf{a} \in \text{Hom}^0(\mathcal{E}_i \mathcal{E}_i \mathbb{1}_{\mathbf{a}})$$

does not have an image 1-morphism. In [35], KLMS construct a diagrammatic category equivalent to the Karoubi envelope of $\mathcal{U}_q(\mathfrak{gl}_2)$, which is commonly called the *thick calculus*, which we now recall. As above, we use a different sign convention from [35] in this paper; see Remark 5.14 which discusses the relevant rescaling.

Definition 5.11. Set $m = 2$ and let $\mathcal{E} := \mathcal{E}_1$ and $\mathcal{F} := \mathcal{F}_1$. The 2-category $\check{\mathcal{U}}_q(\mathfrak{gl}_2)$ is obtained from $\mathcal{U}_q(\mathfrak{gl}_2)$ by adjoining new *divided power* 1-morphisms

$$\mathcal{E}^{(k)} \mathbb{1}_{\mathbf{a}} : \mathbf{a} \rightarrow \mathbf{a} + k\alpha_1, \quad \mathcal{F}^{(k)} \mathbb{1}_{\mathbf{a}} : \mathbf{a} \rightarrow \mathbf{a} - k\alpha_1$$

(where $\mathcal{E}^{(1)} \mathbb{1}_{\mathbf{a}} = \mathcal{E} \mathbb{1}_{\mathbf{a}}$ and $\mathcal{F}^{(1)} \mathbb{1}_{\mathbf{a}} = \mathcal{F} \mathbb{1}_{\mathbf{a}}$), together with new *merge/split* 2-morphisms

$$(5.17) \quad \begin{array}{c} \begin{array}{c} k+\ell \\ \uparrow \\ \text{---} \text{---} \\ \downarrow \downarrow \\ k \quad \ell \\ \mathbf{a} \end{array} \in \text{Hom}^{-k\ell}(\mathcal{E}^{(k)} \mathcal{E}^{(\ell)} \mathbb{1}_{\mathbf{a}}, \mathcal{E}^{(k+\ell)} \mathbb{1}_{\mathbf{a}}), & \begin{array}{c} k+\ell \\ \downarrow \\ \text{---} \text{---} \\ \uparrow \uparrow \\ k \quad \ell \\ \mathbf{a} \end{array} \in \text{Hom}^{-k\ell}(\mathcal{F}^{(k)} \mathcal{F}^{(\ell)} \mathbb{1}_{\mathbf{a}}, \mathcal{F}^{(k+\ell)} \mathbb{1}_{\mathbf{a}}) \\ \\ \begin{array}{c} k \quad \ell \\ \uparrow \uparrow \\ \text{---} \text{---} \\ \downarrow \\ k+\ell \\ \mathbf{a} \end{array} \in \text{Hom}^{-k\ell}(\mathcal{E}^{(k+\ell)} \mathbb{1}_{\mathbf{a}}, \mathcal{E}^{(k)} \mathcal{E}^{(\ell)} \mathbb{1}_{\mathbf{a}}), & \begin{array}{c} k \quad \ell \\ \downarrow \downarrow \\ \text{---} \text{---} \\ \uparrow \\ k+\ell \\ \mathbf{a} \end{array} \in \text{Hom}^{-k\ell}(\mathcal{F}^{(k+\ell)} \mathbb{1}_{\mathbf{a}}, \mathcal{F}^{(k)} \mathcal{F}^{(\ell)} \mathbb{1}_{\mathbf{a}}). \end{array}$$

Here, the labels on the strands in the string diagram correspond to their *thickness*, which is the index k in $\mathcal{E}^{(k)}$, rather than their Dynkin label. (There is a unique Dynkin label when $m = 2$, thus no need to record it.) By convention, any (diagrammatically) “thin” strand has thickness 1; see (5.20) below.

These 2-morphisms satisfy the oriented (co)associativity relations:

$$(5.18) \quad \begin{array}{c} \begin{array}{c} k+\ell+m \\ \uparrow \\ \text{---} \text{---} \\ \downarrow \downarrow \\ k \quad \ell \quad m \\ \mathbf{a} \end{array} = \begin{array}{c} k+\ell+m \\ \uparrow \\ \text{---} \text{---} \\ \downarrow \downarrow \\ k \quad \ell \quad m \\ \mathbf{a} \end{array}, \quad \begin{array}{c} m \quad \ell \quad k \\ \downarrow \downarrow \downarrow \\ \text{---} \text{---} \\ \uparrow \\ k+\ell+m \\ \mathbf{a} \end{array} = \begin{array}{c} k \quad \ell \quad m \\ \downarrow \downarrow \downarrow \\ \text{---} \text{---} \\ \uparrow \\ k+\ell+m \\ \mathbf{a} \end{array}. \end{array}$$

As a consequence, there is a unique 2-morphism $\mathcal{E}^{(k)} \rightarrow \mathcal{E}^k$ built from splits called the *full split*, denoted

$$\begin{array}{c} \uparrow \dots \uparrow \\ \text{---} \text{---} \\ \downarrow \\ k \\ \mathbf{a} \end{array}.$$

Similarly, there is a unique 2-morphism $\mathcal{E}^k \rightarrow \mathcal{E}^{(k)}$ built from merges, called the *full merge*, that is denoted analogously. The full merge/split morphisms satisfy the following relations:

$$(5.19) \quad \begin{array}{c} \begin{array}{c} \uparrow \dots \uparrow \\ \text{---} \text{---} \\ \downarrow \\ k \\ \mathbf{a} \end{array} = \begin{array}{c} \uparrow \dots \uparrow \\ \boxed{\mathbf{HT}_k} \\ \downarrow \dots \downarrow \\ \mathbf{a} \end{array}, \quad \begin{array}{c} \text{---} \text{---} \\ \downarrow \downarrow \downarrow \\ \text{---} \text{---} \\ \uparrow \\ k \\ \mathbf{a} \end{array} = \begin{array}{c} \text{---} \text{---} \\ \boxed{\mathbf{HT}_k} \\ \downarrow \dots \downarrow \\ \mathbf{a} \end{array} \end{array}$$

and

$$(5.20) \quad \begin{array}{c} \begin{array}{c} k \\ \downarrow \\ \text{---} \text{---} \\ \uparrow \uparrow \\ k-1 \quad k-2 \quad \dots \\ \downarrow \\ k \\ \mathbf{a} \end{array} = \begin{array}{c} \uparrow \\ \mathbf{a} \end{array}, \quad \begin{array}{c} k \\ \uparrow \\ \text{---} \text{---} \\ \downarrow \downarrow \\ k-1 \quad k-2 \quad \dots \\ \downarrow \\ k \\ \mathbf{a} \end{array} = \begin{array}{c} \downarrow \\ \mathbf{a} \end{array} \end{array}$$

Here \mathbf{HT}_k denotes any string diagram depicting a reduced word for the half-twist permutation (i.e. the longest element of \mathfrak{S}_k).

- The assignment sending a Schur polynomial $\mathfrak{s}_\lambda(x_1, \dots, x_k)$ to the element in (5.27) determines a \mathbb{K} -algebra isomorphism $\mathbb{K}[x_1, \dots, x_k]^{\mathfrak{S}_k} \xrightarrow{\cong} \text{End}(\mathcal{E}^{(k)} \mathbb{1}_{\mathbf{a}})$. In particular,

$$(5.28) \quad \begin{array}{c} \uparrow \mathbf{a} \\ \textcircled{\mathfrak{s}_\lambda} \\ \textcircled{\mathfrak{s}_\mu} \\ \downarrow k \end{array} = \sum_{\nu \in P(k)} c'_{\lambda, \mu} \begin{array}{c} \uparrow \mathbf{a} \\ \textcircled{\mathfrak{s}_\nu} \\ \downarrow k \end{array},$$

where $c'_{\lambda, \mu}$ is a Littlewood-Richardson coefficient.

- Decorations migrate according to the coproduct for symmetric functions, e.g.

$$(5.29) \quad \begin{array}{c} k \quad \ell \\ \uparrow \quad \uparrow \\ \textcircled{\mathfrak{s}_\nu} \\ \downarrow k+\ell \end{array} \mathbf{a} = \sum c'_{\lambda, \mu} \begin{array}{c} k \quad \ell \\ \uparrow \quad \uparrow \\ \textcircled{\mathfrak{s}_\lambda} \quad \textcircled{\mathfrak{s}_\mu} \\ \downarrow k+\ell \end{array} \mathbf{a}$$

- For $\lambda \in P(k, \ell)$ and $\mu \in P(\ell, k)$,

$$(5.30) \quad \begin{array}{c} \uparrow \mathbf{a} \\ \textcircled{\mathfrak{s}_\lambda} \quad \textcircled{\mathfrak{s}_\mu} \\ \downarrow \end{array} = \begin{cases} (-1)^{|\widehat{\lambda}|} & \text{if } \mu = \widehat{\lambda} \\ 0 & \text{else.} \end{cases}$$

- There is an idempotent decomposition of the identity

$$(5.31) \quad \sum_{\lambda \in P(k, \ell)} (-1)^{|\widehat{\lambda}|} \begin{array}{c} k \quad \ell \\ \uparrow \quad \uparrow \\ \textcircled{\mathfrak{s}_\lambda} \\ \downarrow \textcircled{\mathfrak{s}_{\widehat{\lambda}}} \\ k \quad \ell \end{array} \mathbf{a} = \begin{array}{c} \uparrow \mathbf{a} \\ \uparrow \mathbf{a} \\ k \quad \ell \end{array}$$

Together with (5.30), this implies that $\mathcal{E}^{(k)} \mathcal{E}^{(\ell)} \mathbb{1}_{\mathbf{a}} \cong \bigoplus_{\lambda \in P(k, \ell)} \mathcal{E}^{(k+\ell)} \mathbb{1}_{\mathbf{a}}$ and that $\mathcal{F}^{(k)} \mathcal{F}^{(\ell)} \mathbb{1}_{\mathbf{a}} \cong \bigoplus_{\lambda \in P(k, \ell)} \mathcal{F}^{(k+\ell)} \mathbb{1}_{\mathbf{a}}$.

- For any $b_1, \dots, b_k \in \mathbb{Z}_{\geq 0}$ and any $0 \leq r < k$

$$(5.32) \quad \begin{array}{c} \uparrow \mathbf{a} \\ \textcircled{b_1} \quad \textcircled{b_r} \quad \textcircled{b_{r+1}} \quad \textcircled{b_k} \\ \downarrow k \end{array} = -1 \cdot \begin{array}{c} \uparrow \mathbf{a} \\ \textcircled{b_1} \quad \textcircled{b_{r+1}} \quad \textcircled{b_r} \quad \textcircled{b_k} \\ \downarrow k \end{array} .$$

- Generalizing (5.15), the thick curl relation:

$$\begin{aligned}
 (5.33) \quad \begin{array}{c} \uparrow \\ \text{thick curl} \\ \downarrow k \end{array} &= (-1)^{k^2} \sum_{\lambda, \mu} c_{\lambda, \mu}^{\nu - a_1 + a_2} \begin{array}{c} \uparrow \\ \text{bubble} \\ \downarrow k \end{array} \\
 &= (-1)^{k(k+a_1-1)} \sum_{\lambda, \mu} c_{\lambda, \mu}^{\nu - a_1 + a_2} \begin{array}{c} \uparrow \\ \text{bubble} \\ \downarrow k \end{array},
 \end{aligned}$$

holds. Here, the notation $\nu \pm r$ stands for the partition $(\nu_1 \pm r, \dots, \nu_k \pm r)$ and $\mathfrak{s}_\mu^\blacklozenge = \mathfrak{s}_{\mu-r}$ for r such that the resulting bubble has degree $2|\mu|$.


- There are decompositions into indecomposable 1-morphisms:


$$(5.34) \quad \mathcal{E}^{(k)} \mathcal{F}^{(\ell)} \mathbb{1}_{(a_1, a_2)} \cong \bigoplus_{j=0}^{\min(k, \ell)} \bigoplus_{[k-\ell+a_1-a_2, j]} \mathcal{F}^{(\ell-j)} \mathcal{E}^{(k-j)} \mathbb{1}_{(a_1, a_2)} \quad \text{if } k - \ell + a_1 - a_2 \geq 0$$

and

$$(5.35) \quad \mathcal{F}^{(\ell)} \mathcal{E}^{(k)} \mathbb{1}_{(a_1, a_2)} \cong \bigoplus_{j=0}^{\min(k, \ell)} \bigoplus_{[k-\ell-a_1+a_2, j]} \mathcal{E}^{(k-j)} \mathcal{F}^{(\ell-j)} \mathbb{1}_{(a_1, a_2)} \quad \text{if } \ell - k - a_1 + a_2 \geq 0$$

given via the *Stošić formulae*:

$$(5.36a) \quad \begin{array}{c} \uparrow \\ \text{thick curl} \\ \downarrow k \end{array} \begin{array}{c} \uparrow \\ \text{thick curl} \\ \downarrow \ell \end{array} \begin{array}{c} (a_1, a_2) \end{array} = \sum_{j=0}^{\min(k, \ell)} \sum_{\substack{\lambda, \mu, \nu \in P(j) \\ y \in P(j, k-j) \\ z \in P(j, \ell-j)}} (-1)^{\frac{j(j+1)}{2} + |y| + |z| + (k+\ell)j} c_{\lambda, \mu, \nu, y, z}^{N_j}$$


$$(5.36b) \quad \begin{array}{c} \downarrow \ell \\ \text{thick curl} \\ \uparrow k \end{array} \begin{array}{c} \downarrow \ell \\ \text{thick curl} \\ \uparrow k \end{array} \begin{array}{c} (a_1, a_2) \end{array} = \sum_{j=0}^{\min(k, \ell)} \sum_{\substack{\lambda, \mu, \nu \in P(j) \\ y \in P(j, k-j) \\ z \in P(j, \ell-j)}} (-1)^{\frac{j(j+1)}{2} + |y| + |z|} c_{\lambda, \mu, \nu, y, z}^{M_j}$$


Here, $N_j := (a_1 - a_2 + k - \ell - j)^j$, $M_j := (a_2 - a_1 + \ell - k - j)^j$, and $c_{\lambda, \mu, \nu, y, z}^\rho$ is an iterated Littlewood-Richardson coefficient, defined by the equality $\mathfrak{s}_\lambda \mathfrak{s}_\mu \mathfrak{s}_\nu \mathfrak{s}_y \mathfrak{s}_z = \sum c_{\lambda, \mu, \nu, y, z}^\rho \mathfrak{s}_\rho$. Observe that the signs in (5.36) differ from those in [35]; see Remark 5.14.

Moreover, the 1-morphisms

$$\begin{cases} \mathcal{F}^{(\ell)} \mathcal{E}^{(k)} \mathbb{1}_{(a_1, a_2)} & \text{if } \ell - k \leq a_1 - a_2 \\ \mathcal{E}^{(k)} \mathcal{F}^{(\ell)} \mathbb{1}_{(a_1, a_2)} & \text{if } k - \ell \leq a_2 - a_1 \end{cases}$$

(that appear on the right-hand side of (5.34) and (5.35)) constitute all of the indecomposable 1-morphisms in $\check{\mathcal{U}}_q(\mathfrak{gl}_2)$, up to grading shift. \square

Now we pass from $\mathcal{U}_q(\mathfrak{gl}_2)$ to the setting of $\mathcal{U}_q(\mathfrak{gl}_m)$ for general m . When $m \geq 3$, there is no combinatorial description of the Karoubi envelope of $\mathcal{U}_q(\mathfrak{gl}_m)$, and, indeed, the canonical basis for $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$, which the indecomposable 1-morphisms in $\text{Kar}(\mathcal{U}_q(\mathfrak{gl}_m))$ categorify when working in characteristic zero [71], is not even known explicitly. However, using the $m = 2$ case, it is possible to describe a partial idempotent completion $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$ of $\mathcal{U}_q(\mathfrak{gl}_m)$ that contains divided power morphisms

$$\mathcal{E}_i^{(k)} \mathbb{1}_{\mathbf{a}}: \mathbf{a} \rightarrow \mathbf{a} + k\alpha_i, \quad \mathcal{F}_i^{(k)} \mathbb{1}_{\mathbf{a}}: \mathbf{a} \rightarrow \mathbf{a} - k\alpha_i$$

for each $1 \leq i \leq m - 1$. See e.g. [53, Definition 2.2].

Definition 5.16. Let $m \geq 1$. The *thick categorified quantum group* $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$ is the \mathbb{Z} -additive closure of the 2-category given as follows.

- Objects are elements $\mathbf{a} \in \mathbb{Z}^m$.
- 1-morphisms are generated by

$$\mathcal{E}_i^{(k)} \mathbb{1}_{\mathbf{a}}: \mathbf{a} \rightarrow \mathbf{a} + k\alpha_i, \quad \mathcal{F}_i^{(k)} \mathbb{1}_{\mathbf{a}}: \mathbf{a} \rightarrow \mathbf{a} - k\alpha_i$$

for $1 \leq i \leq m - 1$, $k \in \mathbb{Z}_+$, and $\alpha_i = (0, \dots, 1, -1, \dots, 0)$.

- 2-morphisms are \mathbb{K} -linear combinations of “string diagrams” that are generated via horizontal and vertical composition by those in (5.2) and for each Dynkin node $i \in \{1, \dots, m - 1\}$ merge/split generators (5.17), modulo the relations in Definition 5.5 and the relations (5.19) and (5.20).

Proposition 5.17. The category $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$ is equivalent to the partial idempotent completion of $\mathcal{U}_q(\mathfrak{gl}_m)$ which adjoins all direct summands of $\mathcal{E}_i^k \mathbb{1}_{\mathbf{a}}$ and $\mathcal{F}_i^k \mathbb{1}_{\mathbf{a}}$ for $1 \leq i \leq m - 1$ and $k \geq 1$.

Proof (Sketch). The discussion in Remark 5.12 implies that one can present partial idempotent completions by adding one idempotent at a time, independently of other idempotents. Meanwhile, [35] proves that $\check{\mathcal{U}}_q(\mathfrak{gl}_2)$ is equivalent to the partial idempotent completion which adds all summands of $\mathcal{E}^k \mathbb{1}_{\mathbf{a}}$ and $\mathcal{F}^k \mathbb{1}_{\mathbf{a}}$. Thus using the construction of [35] independently for each $\mathcal{E}_i \mathbb{1}_{\mathbf{a}}$ and $\mathcal{F}_i \mathbb{1}_{\mathbf{a}}$, one has added all direct summands of $\mathcal{E}_i^k \mathbb{1}_{\mathbf{a}}$ and $\mathcal{F}_i^k \mathbb{1}_{\mathbf{a}}$. \square

As in Remark 5.9, we will view $\coprod_{m \geq 0} \check{\mathcal{U}}_q(\mathfrak{gl}_m)$ as a monoidal 2-category under the analogously defined external tensor product \boxtimes .

Convention 5.18. Note that the strands in the string diagrams for $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$ require both a Dynkin label $i \in \{1, \dots, m - 1\}$ and a thickness label $k \in \mathbb{Z}_+$, e.g. we have

$$\begin{array}{c} \begin{array}{c} i, k \quad i, \ell \\ \uparrow \quad \uparrow \\ \text{Y} \\ \downarrow \\ i, k + \ell \end{array} \mathbf{a} \in \text{Hom}^{-k\ell}(\mathcal{E}_i^{(k+\ell)} \mathbb{1}_{\mathbf{a}}, \mathcal{E}_i^{(k)} \mathcal{E}_i^{(\ell)} \mathbb{1}_{\mathbf{a}}) \end{array}$$

Instead of including both labels, we will denote the Dynkin label by coloring the relevant strands according to the following coloring of the \mathfrak{gl}_m Dynkin diagram:

$$\bullet \cdots \bullet \overset{\text{red}}{\bullet} \overset{\text{green}}{\bullet} \overset{\text{blue}}{\bullet} \cdots \bullet$$

$$1 \quad \quad i-1 \quad i \quad i+1 \quad m-1 \quad \cdot$$

To elaborate, we typically use **green** as our arbitrary index i in the Dynkin diagram. It is always assumed that **red** is one less than **green** and **blue** is one more than **green**. Meanwhile, **black** represents an arbitrary index (possibly equal to **red**, **green**, **blue**, or distant from these colors).

Example 5.19. Pairing Convention 5.18 with the conventions discussed in Remark 5.3, we have $t_{\bullet, \bullet} = -1$, $t_{\bullet, \bullet} = +1$, $t_{\bullet, \bullet} = -1$, etc.

Convention 5.20. Convention 5.13 explains how to define thick caps/cups and thick crossings for $\check{\mathcal{U}}_q(\mathfrak{gl}_2)$. We use the same recipe to define thick caps/cups and (unicolored) thick crossings for $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$. We similarly define thick crossings for different colors as

$$(5.37) \quad \begin{array}{c} \text{green} \\ \text{green} \\ \text{black} \\ \text{green} \end{array} \text{ crossing } \mathbf{a} := \begin{array}{c} \text{green} \\ \text{green} \\ \text{black} \\ \text{green} \end{array} \text{ bubble } \mathbf{a}$$

These thick crossings still satisfy fork-slide relations with merge/split morphisms as in Convention 5.13. For example,

$$(5.38) \quad \begin{array}{c} \text{green} \\ \text{green} \\ \text{black} \\ \text{green} \end{array} \text{ crossing } \mathbf{a} = \begin{array}{c} \text{green} \\ \text{green} \\ \text{black} \\ \text{green} \end{array} \text{ crossing } \mathbf{a}$$

which follows using (5.6), in the case where i and j are green and k is black.

6. INVOLUTIONS ON (PART OF) THE CATEGORIFIED QUANTUM GROUP

Recall from Section 1.2 that we seek an involution of the 2-category $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$, in order to carry out the equivariantization procedure sketched there (and defined concretely below in §7). For the sake of this paper, we work with the following definition, rather than the more-general notion in which the stated equality of functors is replaced by a natural isomorphism.

Definition 6.1. A (strict) involution of a \mathbb{K} -linear 2-category \mathcal{A} is a \mathbb{K} -linear 2-functor $\sigma: \mathcal{A} \rightarrow \mathcal{A}$ such that $\sigma \circ \sigma = \text{id}_{\mathcal{A}}$.

Now, fix $n \geq 1$ and $m \geq 1$. Let \mathbf{n} denote the \mathfrak{gl}_m weight (n, n, \dots, n) , thus $2\mathbf{n} = (2n, 2n, \dots, 2n)$. To begin, we seek an involution τ of $\mathcal{U}_q(\mathfrak{gl}_m)$ that

- is given on \mathfrak{gl}_m weights by sending $\mathbf{a} \mapsto 2\mathbf{n} - \mathbf{a}$,
- is determined on 1-morphisms by $\mathcal{E}_i \mathbb{1}_{\mathbf{a}} \mapsto \mathcal{F}_i \mathbb{1}_{2\mathbf{n} - \mathbf{a}}$, and
- acts on new bubble 2-morphisms by $\mathfrak{s}_{\lambda}(\mathbb{X}_i) \mapsto \mathfrak{s}_{\lambda^t}(\mathbb{X}_i)$.

The condition on objects and 1-morphisms guarantee that τ maps the indecomposable 1-morphism $\mathbf{X}_i := \mathcal{F}_i \mathcal{E}_i \mathbb{1}_{\mathbf{n}}$ to $\mathcal{E}_i \mathcal{F}_i \mathbb{1}_{\mathbf{n}} \cong \mathbf{X}_i$. As outlined in Section 1.4, an equivariant structure on \mathbf{X}_i gives a natural candidate for a 1-morphism categorifying the distinguished elements $\mathbf{X}_i \in \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S^{\otimes m})$. The condition on 2-morphisms is present in order for this involution to be compatible with the involution on $H^*(\text{Gr}_n(\mathbb{C}^{2n}))$ from Section 1.1.

As it turns out, these requirements (together with integrality assumptions) essentially determine τ on the generating morphisms of $\mathcal{U}_q(\mathfrak{gl}_m)$, up to signs. For example, taken together with equation (5.9), they imply that the dot endomorphism in $\text{End}^2(\mathcal{E}_i \mathbb{1}_{\mathbf{a}})$ must be sent to minus the dot endomorphism in $\text{End}^2(\mathcal{F}_i \mathbb{1}_{2\mathbf{n}-\mathbf{a}})$. In §6.1, we find conditions on the requisite signs that guarantee that the above recipe for τ is a well-defined automorphism of $\mathcal{U}_q(\mathfrak{gl}_m)$. Next, in §6.2 we investigate further conditions on these signs which would imply that τ is an involution. As it turns out, when $m \geq 3$ it is surprisingly difficult to satisfy these conditions! (See Theorem 6.11 for the precise statement.) Fortunately, as we show in §6.3, we *are* able to construct a symmetry τ of order 4 for all $m \geq 1$ which restricts to an involution on an appropriate 2-subcategory that suffices for our present considerations. To conclude this section, in §6.4 we discuss the dependence on our chosen value of n and in §6.5 we extend τ to thick calculus.

For the remainder of this section, \mathbb{K} is permitted to be any integral domain for which $2 \neq 0$.

6.1. A family of symmetries. We now define a family of autoequivalences of categorified \mathfrak{gl}_m , one for each¹⁵ $n \geq 1$. By slight abuse of notation, we will denote all of these autoequivalences by τ .

Remark 6.2. Each of these symmetries will send \mathcal{E}_i to \mathcal{F}_i , which is similar to the first automorphism of $\mathcal{U}_q(\mathfrak{sl}_m)$ defined by Khovanov–Lauda in [34, §3.3.2]. It would be easy to confuse our automorphism with theirs, as both act by rescaling the generating 2-morphisms and then reversing the orientation on string diagrams. However, these automorphisms are distinct: as discussed above, ours rescales the “dot” endomorphism by -1 and does not rescale the uni-colored crossing, while the opposite is true for the Khovanov–Lauda symmetry. If $2 = 0$ in \mathbb{K} (a case not permitted by our assumptions on \mathbb{K} above), then $1 = -1$ and the symmetry we define below will agree with Khovanov–Lauda’s automorphism.

Given the action on 2-morphisms, our symmetry is covariant for both 1-morphism and 2-morphism composition. Moving forward, we will say that τ is *covariant* to mean that it is covariant for 2-morphism composition, and *monoidal* to mean that it is covariant for 1-morphism composition.

Definition 6.3. Fix $n \geq 0$. For each $i \in \{1, \dots, m-1\}$, let $r_i, r'_i, l_i,$ and l'_i be functions from the \mathfrak{gl}_m weight lattice \mathbb{Z}^m to $\mathbb{Z}/2$. For each pair $i, j \in \{1, \dots, m-1\}$, let $v_{i,j}$ be a function from \mathbb{Z}^m to $\mathbb{Z}/2$. Associated to this data, define the following map τ on the generating data of $\mathcal{U}_q(\mathfrak{gl}_m)$. In the next theorem, we will state precise conditions on $r_i, r'_i, l_i, l'_i,$ and $v_{i,j}$ which imply that τ extends to a 2-functor $\tau: \mathcal{U}_q(\mathfrak{gl}_m) \rightarrow \mathcal{U}_q(\mathfrak{gl}_m)$.

- **Objects:** $\mathbf{n} + \mathbf{a} \xrightarrow{\tau} \mathbf{n} - \mathbf{a}$ (i.e. $\mathbf{a} \mapsto 2\mathbf{n} - \mathbf{a}$).
- **1-morphisms:** $\mathcal{E}_i \mathbb{1}_{\mathbf{n}+\mathbf{a}} \xrightarrow{\tau} \mathcal{F}_i \mathbb{1}_{\mathbf{n}-\mathbf{a}}$.
- **2-morphisms:**

$$(6.1) \quad \begin{array}{c} \begin{array}{ccc} \begin{array}{c} \uparrow \\ \bullet \\ \downarrow \end{array} \mathbf{n}+\mathbf{a} & \xrightarrow{\tau} - & \begin{array}{c} \downarrow \\ \bullet \\ \uparrow \end{array} \mathbf{n}-\mathbf{a} \\ \begin{array}{ccc} \boxed{e_r(\mathbb{X}_i)} \mathbf{n}+\mathbf{a} & \xrightarrow{\tau} & \boxed{h_r(\mathbb{X}_i)} \mathbf{n}-\mathbf{a} \\ \begin{array}{ccc} \begin{array}{c} \nearrow \\ \downarrow \\ \searrow \end{array} \mathbf{n}+\mathbf{a} & \xrightarrow{\tau} (-1)^{v_{\bullet\bullet}(\mathbf{a})} & \begin{array}{c} \searrow \\ \downarrow \\ \nearrow \end{array} \mathbf{n}-\mathbf{a} \\ \begin{array}{ccc} \begin{array}{c} \curvearrowright \end{array} \mathbf{n}+\mathbf{a} & \xrightarrow{\tau} (-1)^{r_{\bullet}(\mathbf{a})} & \begin{array}{c} \curvearrowleft \end{array} \mathbf{n}-\mathbf{a} \\ \begin{array}{ccc} \begin{array}{c} \curvearrowleft \end{array} \mathbf{n}+\mathbf{a} & \xrightarrow{\tau} (-1)^{r'_{\bullet}(\mathbf{a})} & \begin{array}{c} \curvearrowright \end{array} \mathbf{n}-\mathbf{a} \\ \begin{array}{ccc} \begin{array}{c} \curvearrowright \end{array} \mathbf{n}+\mathbf{a} & \xrightarrow{\tau} (-1)^{l_{\bullet}(\mathbf{a})} & \begin{array}{c} \curvearrowleft \end{array} \mathbf{n}-\mathbf{a} \\ \begin{array}{ccc} \begin{array}{c} \curvearrowleft \end{array} \mathbf{n}+\mathbf{a} & \xrightarrow{\tau} (-1)^{l'_{\bullet}(\mathbf{a})} & \begin{array}{c} \curvearrowright \end{array} \mathbf{n}-\mathbf{a} \end{array} \end{array} \end{array}$$

¹⁵Actually, in Definition 6.3, we do so for $n \geq 0$; we use the $n = 0$ case in §6.4. In fact, our definition works for all $n \in \mathbb{Z}$, but those for $n < 0$ are not of use for us.

Remark 6.4. We emphasize that τ depends on the fixed integer $n > 0$ appearing in \mathbf{n} , although our notation suppresses this. For further discussion regarding dependence on n , in some special cases, see Section 6.4.

Theorem 6.5. The map τ defined above extends to a unique (invertible, monoidal, covariant, graded, \mathbb{K} -linear) 2-functor $\tau: \mathcal{U}_q(\mathfrak{gl}_m) \rightarrow \mathcal{U}_q(\mathfrak{gl}_m)$ if and only if the following equalities hold (in $\mathbb{Z}/2$) for all \mathbf{a} in the weight lattice, and for all $i, j, k \in \{1, \dots, m-1\}$:

$$(6.2a) \quad r'_i(\mathbf{a}) \equiv r_i(\mathbf{a} + \alpha_i), \quad l'_i(\mathbf{a}) \equiv l_i(\mathbf{a} - \alpha_i).$$

$$(6.2b) \quad r_i(\mathbf{a}) + l_i(\mathbf{a} - \alpha_i) \equiv a_i - a_{i+1} = \langle \alpha_i^\vee, \mathbf{a} \rangle,$$

$$(6.2c) \quad v_{i,i}(\mathbf{a}) \equiv 0,$$

$$(6.2d) \quad v_{i,j}(\mathbf{a}) + v_{j,i}(\mathbf{a}) \equiv i \cdot j = \langle \alpha_i^\vee, \alpha_j \rangle$$

$$(6.2e) \quad v_{i,j}(\mathbf{a}) + v_{i,k}(\mathbf{a}) + v_{j,k}(\mathbf{a}) + v_{i,j}(\mathbf{a} + \alpha_k) + v_{i,k}(\mathbf{a} + \alpha_j) + v_{j,k}(\mathbf{a} + \alpha_i) \equiv 0.$$

Here, and for the remainder of this section, we use \equiv to denote equality modulo 2. Note that $i \cdot j$ in (6.2d) is odd when i and j are adjacent in the Dynkin diagram, and even otherwise.

In Remark 6.7 below we demonstrate that functions satisfying (6.2) do indeed exist, and in Theorem 6.16 we fix a choice (with additional desired properties) that we use for the duration. For easy reference, whenever (6.2) holds, the value of τ on non-generating morphisms is as follows:

$$(6.3) \quad \boxed{\mathfrak{s}_\lambda(\mathbb{X}_i)}^{\mathbf{n}+\mathbf{a}} \xrightarrow{\tau} \boxed{\mathfrak{s}_{\lambda^t}(\mathbb{X}_i)}^{\mathbf{n}-\mathbf{a}}$$

$$(6.4) \quad \begin{array}{c} \downarrow^{\mathbf{n}+\mathbf{a}} \\ \bullet \\ \uparrow^{\mathbf{n}-\mathbf{a}} \end{array} \xrightarrow{\tau} - \begin{array}{c} \uparrow^{\mathbf{n}-\mathbf{a}} \\ \bullet \\ \downarrow^{\mathbf{n}+\mathbf{a}} \end{array}$$

$$\begin{array}{c} \begin{array}{c} \nearrow^{\mathbf{n}+\mathbf{a}} \\ \searrow^{\mathbf{n}+\mathbf{a}} \end{array} \xrightarrow{\tau} (-1)^{v_{\bullet\bullet}(\mathbf{a}-\alpha_\bullet)+r_\bullet(\mathbf{a})+r_\bullet(\mathbf{a}+\alpha_\bullet)} \begin{array}{c} \searrow^{\mathbf{n}-\mathbf{a}} \\ \nearrow^{\mathbf{n}-\mathbf{a}} \end{array} \\ \begin{array}{c} \searrow^{\mathbf{n}+\mathbf{a}} \\ \nearrow^{\mathbf{n}+\mathbf{a}} \end{array} \xrightarrow{\tau} (-1)^{v_{\bullet\bullet}(\mathbf{a}-\alpha_\bullet)+l_\bullet(\mathbf{a}-\alpha_\bullet)+l_\bullet(\mathbf{a}-\alpha_\bullet+\alpha_\bullet)} \begin{array}{c} \nearrow^{\mathbf{n}-\mathbf{a}} \\ \searrow^{\mathbf{n}-\mathbf{a}} \end{array} \end{array}$$

and

$$(6.5) \quad \begin{array}{c} \searrow^{\mathbf{n}+\mathbf{a}} \\ \nearrow^{\mathbf{n}+\mathbf{a}} \end{array} \xrightarrow{\tau} (-1)^{v'_{\bullet\bullet}(\mathbf{a})} \begin{array}{c} \nearrow^{\mathbf{n}-\mathbf{a}} \\ \searrow^{\mathbf{n}-\mathbf{a}} \end{array},$$

where

$$(6.6) \quad v'_{i,j}(\mathbf{a}) \equiv v_{i,j}(\mathbf{a} - \alpha_i - \alpha_j) + r_i(\mathbf{a}) + r_j(\mathbf{a}) + r_i(\mathbf{a} - \alpha_j) + r_j(\mathbf{a} - \alpha_i).$$

Equation 6.3 follows from noting that the standard involution of symmetric functions, defined by sending elementary symmetric functions to complete symmetric functions as in equation (6.1), sends the Schur function associated to a partition to the Schur function associated to the transpose partition. It is routine to use equation (6.1), which defines τ on generators, along with the defining relations of $\mathcal{U}_q(\mathfrak{gl}_m)$, to verify equations (6.4) and (6.5)

Remark 6.6. We view (6.2a) (and (6.6)) as expressing the primed variables in terms of the unprimed ones, and the remaining equations as conditions on the unprimed variables. Note that (6.2c) also pairs with (6.6) to imply that $v'_{i,i}(\mathbf{a}) \equiv 0$.

The corresponding computation for counterclockwise bubbles is left to the reader. Given (6.2a), the two actions of τ again agree if and only if (6.2b) holds. To see this most effortlessly, one should study a bubble in a region labeled $\mathbf{n} + (\mathbf{a} - \alpha_i)$.

Lastly, recall that fake bubbles (where the number of dots in (5.7) is a negative number) are defined via (5.8) as formal symmetric functions, and τ acts on them accordingly. Since fake bubbles are not genuine compositions of a cap, cup, and dots, there is no requirement for τ to act on them “as though they were compositions”, e.g. by the sign $(-1)^{r_{\bullet}(\mathbf{a}) + l'_{\bullet}(\mathbf{a}) + c}$ (where c is a negative number) for fake clockwise green bubbles. However, fake bubbles do indeed satisfy this formula: (6.8) still holds when $c < 0$. The same computation as above serves as the proof, where now one uses (6.2b) rather than deriving (6.2b).

• **Cyclicity of crossings:**

Now consider the isotopy relation in $\mathcal{U}_q(\mathfrak{gl}_m)$ which states that both the left and right mates under adjunction of the upward crossing agree, i.e. that a downward crossing may be defined as either mate. For this purpose we temporarily assume that (6.5) is the definition of τ on the downward crossing, and check both compatibilities.

We compute:

$$(6.9) \quad \begin{array}{c} \text{downward crossing} \\ \text{with } \mathbf{n} + \mathbf{a} \text{ dots} \end{array} = \begin{array}{c} \text{downward crossing} \\ \text{with } \mathbf{n} + \mathbf{a} \text{ dots} \end{array} \xrightarrow{\tau} (-1)^{v_{\bullet\bullet}(\mathbf{a} - \alpha_{\bullet} - \alpha_{\bullet}) + r_{\bullet}(\mathbf{a}) + r_{\bullet}(\mathbf{a} - \alpha_{\bullet}) + r'_{\bullet}(\mathbf{a} - \alpha_{\bullet} - \alpha_{\bullet}) + r'_{\bullet}(\mathbf{a} - \alpha_{\bullet})} \begin{array}{c} \text{downward crossing} \\ \text{with } \mathbf{n} - \mathbf{a} \text{ dots} \end{array}$$

Thus, for consistency with (6.1), we need

$$(6.10) \quad v'_{\bullet\bullet}(\mathbf{a}) \equiv v_{\bullet\bullet}(\mathbf{a} - \alpha_{\bullet} - \alpha_{\bullet}) + r_{\bullet}(\mathbf{a}) + r_{\bullet}(\mathbf{a} - \alpha_{\bullet}) + r'_{\bullet}(\mathbf{a} - \alpha_{\bullet} - \alpha_{\bullet}) + r'_{\bullet}(\mathbf{a} - \alpha_{\bullet}).$$

Using (6.2a), one obtains (6.6), as desired. Note that when $\bullet = \bullet$, many terms cancel and (6.10) (or (6.6)) simply becomes $v'_{\bullet\bullet}(\mathbf{a}) \equiv v_{\bullet\bullet}(\mathbf{a} - 2\alpha_{\bullet})$.

For the other mate, we compute:

$$(6.11) \quad \begin{array}{c} \text{downward crossing} \\ \text{with } \mathbf{n} + \mathbf{a} \text{ dots} \end{array} = \begin{array}{c} \text{downward crossing} \\ \text{with } \mathbf{n} + \mathbf{a} \text{ dots} \end{array} \xrightarrow{\tau} (-1)^{v_{\bullet\bullet}(\mathbf{a} - \alpha_{\bullet} - \alpha_{\bullet}) + l'_{\bullet}(\mathbf{a}) + l'_{\bullet}(\mathbf{a} - \alpha_{\bullet}) + l_{\bullet}(\mathbf{a} - \alpha_{\bullet} - \alpha_{\bullet}) + l_{\bullet}(\mathbf{a} - \alpha_{\bullet})} \begin{array}{c} \text{downward crossing} \\ \text{with } \mathbf{n} - \mathbf{a} \text{ dots} \end{array},$$

which requires

$$v'_{\bullet\bullet}(\mathbf{a}) \equiv v_{\bullet\bullet}(\mathbf{a} - \alpha_{\bullet} - \alpha_{\bullet}) + l_{\bullet}(\mathbf{a} - \alpha_{\bullet} - \alpha_{\bullet}) + l_{\bullet}(\mathbf{a} - \alpha_{\bullet}) + l'_{\bullet}(\mathbf{a}) + l'_{\bullet}(\mathbf{a} - \alpha_{\bullet}).$$

Substituting (6.10) (which we now assume) for the left-hand side and applying (6.2a) gives the condition

$$r_{\bullet}(\mathbf{a}) + r_{\bullet}(\mathbf{a} - \alpha_{\bullet}) + r_{\bullet}(\mathbf{a} - \alpha_{\bullet}) + r_{\bullet}(\mathbf{a}) + l_{\bullet}(\mathbf{a} - \alpha_{\bullet} - \alpha_{\bullet}) + l_{\bullet}(\mathbf{a} - \alpha_{\bullet}) + l_{\bullet}(\mathbf{a} - \alpha_{\bullet}) + l_{\bullet}(\mathbf{a} - \alpha_{\bullet} - \alpha_{\bullet}) \equiv 0.$$

Using (6.2b) four times, this becomes

$$\langle \alpha_{\bullet}^{\vee}, \mathbf{a} \rangle + \langle \alpha_{\bullet}^{\vee}, \mathbf{a} - \alpha_{\bullet} \rangle + \langle \alpha_{\bullet}^{\vee}, \mathbf{a} - \alpha_{\bullet} \rangle + \langle \alpha_{\bullet}^{\vee}, \mathbf{a} \rangle \equiv 0$$

which holds by the symmetry $\langle \alpha_{\bullet}^{\vee}, \alpha_{\bullet} \rangle = \langle \alpha_{\bullet}^{\vee}, \alpha_{\bullet} \rangle$ of the type A Cartan matrix. Hence, (6.10) is the only requirement, so no additional relations are needed for cyclicity of crossings.

Before studying the remaining relations, let us record how τ acts on the rightward sideways crossing:

$$(6.12) \quad \begin{array}{c} \text{Rightward Sideways Crossing} \\ \text{with } n+a \text{ strands} \end{array} = \begin{array}{c} \text{Rightward Sideways Crossing} \\ \text{with } n+a \text{ strands} \end{array} \xrightarrow{\tau} (-1)^{v_{\bullet\bullet}(\mathbf{a}-\alpha_{\bullet})+r_{\bullet}(\mathbf{a})+r'_{\bullet}(\mathbf{a}-\alpha_{\bullet}+\alpha_{\bullet})} \begin{array}{c} \text{Rightward Sideways Crossing} \\ \text{with } n-a \text{ strands} \end{array} \\ \stackrel{(6.2a)}{=} (-1)^{v_{\bullet\bullet}(\mathbf{a}-\alpha_{\bullet})+r_{\bullet}(\mathbf{a})+r_{\bullet}(\mathbf{a}+\alpha_{\bullet})} \begin{array}{c} \text{Rightward Sideways Crossing} \\ \text{with } n-a \text{ strands} \end{array}$$

and the leftward sideways crossing:

$$(6.13) \quad \begin{array}{c} \text{Leftward Sideways Crossing} \\ \text{with } n+a \text{ strands} \end{array} = \begin{array}{c} \text{Leftward Sideways Crossing} \\ \text{with } n+a \text{ strands} \end{array} \xrightarrow{\tau} (-1)^{v_{\bullet\bullet}(\mathbf{a}-\alpha_{\bullet})+l'_{\bullet}(\mathbf{a})+l_{\bullet}(\mathbf{a}-\alpha_{\bullet}+\alpha_{\bullet})} \begin{array}{c} \text{Leftward Sideways Crossing} \\ \text{with } n+a \text{ strands} \end{array} \\ \stackrel{(6.2a)}{=} (-1)^{v_{\bullet\bullet}(\mathbf{a}-\alpha_{\bullet})+l_{\bullet}(\mathbf{a}-\alpha_{\bullet})+l_{\bullet}(\mathbf{a}-\alpha_{\bullet}+\alpha_{\bullet})} \begin{array}{c} \text{Leftward Sideways Crossing} \\ \text{with } n+a \text{ strands} \end{array}$$

• **Dot slide:**

Now consider the dot slide relation (5.4) when $i = j = \bullet$. We compute that

$$\left(\begin{array}{c} \text{Dot Slide} \\ \text{with } n+a \text{ strands} \end{array} - \begin{array}{c} \text{Dot Slide} \\ \text{with } n+a \text{ strands} \end{array} \right) \mapsto (-1)^{v_{\bullet\bullet}(\mathbf{a})+1} \left(\begin{array}{c} \text{Dot Slide} \\ \text{with } n-a \text{ strands} \end{array} - \begin{array}{c} \text{Dot Slide} \\ \text{with } n-a \text{ strands} \end{array} \right) \stackrel{(5.4)}{=} (-1)^{v_{\bullet\bullet}(\mathbf{a})} \begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} n-a \\ n-a \\ n-a \end{array}$$

and

$$\left(\begin{array}{c} \text{Dot Slide} \\ \text{with } n+a \text{ strands} \end{array} - \begin{array}{c} \text{Dot Slide} \\ \text{with } n+a \text{ strands} \end{array} \right) \mapsto (-1)^{v_{\bullet\bullet}(\mathbf{a})+1} \left(\begin{array}{c} \text{Dot Slide} \\ \text{with } n-a \text{ strands} \end{array} - \begin{array}{c} \text{Dot Slide} \\ \text{with } n-a \text{ strands} \end{array} \right) \stackrel{(5.4)}{=} (-1)^{v_{\bullet\bullet}(\mathbf{a})} \begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \end{array} \begin{array}{c} n-a \\ n-a \\ n-a \end{array}$$

However, the (far) right-hand side of (5.4) is sent by τ to the identity map of $\mathcal{F}_{\bullet}\mathcal{F}_{\bullet}\mathbb{1}_{n-a}$ with no sign. For this relation to be preserved by τ , we therefore need $v_{\bullet\bullet}(\mathbf{a}) \equiv 0$, which is (6.2c).

Finally, it is easy to confirm that the dot slide relation when $i \neq j$ induces no constraints, essentially because $-1 \cdot 0 = 0$.

• **Quadratic KLR:**

The $i = j$ case of the quadratic KLR relation (5.5) is obvious. In the case where $i \cdot j = 0$, the left-hand side picks up the sign $(-1)^{v_{i,j}(\mathbf{a})+v_{j,i}(\mathbf{a})}$ under τ , and the right-hand side picks up no sign. Therefore, this relation is preserved by τ if and only if $v_{i,j}(\mathbf{a}) \equiv v_{j,i}(\mathbf{a})$ when $i \cdot j = 0$, which gives one case of (6.2d).

Lastly, suppose that $i \cdot j = -1$. Under τ , the left-hand side picks up the sign $(-1)^{v_{i,j}(\mathbf{a})+v_{j,i}(\mathbf{a})}$ and the right-hand side picks up the sign (-1) from the dot. This relation is preserved by τ if and only if $v_{i,j}(\mathbf{a}) \equiv v_{j,i}(\mathbf{a}) + 1$ when $i \cdot j = -1$, which gives the other case of (6.2d). In more detail, suppose that $j = i + 1$ (the $j = i - 1$ case is similar). Following Convention 5.18, we compute

$$\left(\begin{array}{c} \text{Quadratic KLR} \\ \text{with } n+a \text{ strands} \end{array} = \begin{array}{c} \text{Quadratic KLR} \\ \text{with } n+a \text{ strands} \end{array} - \begin{array}{c} \text{Quadratic KLR} \\ \text{with } n+a \text{ strands} \end{array} \right) \xrightarrow{\tau} \left((-1)^{v_{\bullet\bullet}(\mathbf{a})+v_{\bullet\bullet}(\mathbf{a})} \begin{array}{c} \text{Quadratic KLR} \\ \text{with } n-a \text{ strands} \end{array} = - \begin{array}{c} \text{Quadratic KLR} \\ \text{with } n-a \text{ strands} \end{array} + \begin{array}{c} \text{Quadratic KLR} \\ \text{with } n-a \text{ strands} \end{array} \right).$$

We confirm this for (5.11), since the case of (5.10) is similar. In the first diagram on the right-hand side of (5.11), using (6.12) and (6.13), the two sideways crossings produce a total sign with exponent $2v_{i,i}(\mathbf{a} - \alpha_i) + r_i(\mathbf{a}) + r_i(\mathbf{a} + \alpha_i) + l_i(\mathbf{a}) + l_i(\mathbf{a} - \alpha_i)$. Applying (6.2b), the result is equivalent modulo 2 to $\langle \alpha_i^\vee, \mathbf{a} \rangle + \langle \alpha_i^\vee, \mathbf{a} + \alpha_i \rangle$. Since $\langle \alpha_i^\vee, \alpha_i \rangle = 2$, this exponent is even, as desired.

In each term of the summation on the right-hand side, the overall sign produced by τ is

$$(-1)^{p+q} (-1)^{r_i(\mathbf{a})+r'_i(\mathbf{a})+l_i(\mathbf{a})+l'_i(\mathbf{a})} (-1)^{a_{i+1}-a_i-1+r}.$$

Here we've e.g. used (6.8) with $c = a_{i+1} - a_i - 1 + r$. Since $p + q + r = a_{i+1} - a_i - 1$, the overall exponent is equivalent mod 2 to

$$\begin{aligned} r_i(\mathbf{a}) + r'_i(\mathbf{a}) + l_i(\mathbf{a}) + l'_i(\mathbf{a}) &\stackrel{(6.2a)}{\equiv} r_i(\mathbf{a}) + r_i(\mathbf{a} + \alpha_i) + l_i(\mathbf{a}) + l_i(\mathbf{a} - \alpha_i) \\ &\stackrel{(6.2b)}{\equiv} \langle \alpha_i^\vee, \mathbf{a} \rangle + \langle \alpha_i^\vee, \mathbf{a} + \alpha_i \rangle \equiv 2\langle \alpha_i^\vee, \mathbf{a} \rangle + \langle \alpha_i^\vee, \alpha_i \rangle \equiv 0 \end{aligned}$$

as desired.

- **Mixed \mathcal{E}, \mathcal{F} relations:**

Consider the first relation in (5.12), with ambient weight $\mathbf{n} + \mathbf{a}$ on the right. The identity map is sent by τ to an identity map, with no sign. Once again, using (6.12) and (6.13), the overall sign on the right-hand side has exponent

$$(6.15) \quad v_{j,i}(\mathbf{a} - \alpha_j) + r_j(\mathbf{a}) + r_j(\mathbf{a} + \alpha_i) + v_{i,j}(\mathbf{a} - \alpha_j) + l_j(\mathbf{a} - \alpha_j) + l_j(\mathbf{a} - \alpha_j + \alpha_i).$$

Applying (6.2b) and (6.2d), this is equivalent mod 2 to

$$\langle \alpha_j^\vee, \alpha_i \rangle + \langle \alpha_j^\vee, \mathbf{a} \rangle + \langle \alpha_j^\vee, \mathbf{a} + \alpha_i \rangle \equiv 0$$

as desired.

This concludes the proof of Theorem 6.5. \square

Remark 6.7. Let us discuss what it takes to define functions $r_i, l_i, r'_i, l'_i, v_{i,j}: \mathbb{Z}^m \rightarrow \mathbb{Z}/2$ for $i, j \in \{1, \dots, m-1\}$ that satisfy (6.2). Since (6.2) has no conditions relating $v_{i,j}$ to the remaining functions, we can discuss the choice of $v_{i,j}$ separately from the others. However, we emphasize that in subsequent sections, we impose further conditions on τ such that $v_{i,j}$ will cease to be decoupled from the remaining functions.

Observe that one can use (6.2a) to define r'_i and l'_i in terms of r_i and l_i and can then use (6.2b) to define l_i in terms of r_i . We thus can choose each r_i freely, and extrapolate the values of r'_i, l_i, l'_i . Similarly, (6.2c) and (6.2d) can be used to define $v_{i,j}$ for $i \geq j$ in terms of $v_{i,j}$ for $i < j$. Next, (6.2c) implies that (6.2d) holds when $i = j$, and together (6.2c) and (6.2d) imply that (6.2e) holds whenever $\{i, j, k\}$ are not distinct. Lastly, if (6.2e) holds for $\{i, j, k\}$ then it also holds for any permutation of $\{i, j, k\}$.

Thus, it suffices to find $v_{i,j}$ for $i < j$ that satisfy (6.2e). This can be accomplished in a number of ways, e.g. we can let each such $v_{i,j}: \mathbb{Z}^m \rightarrow \mathbb{Z}/2$ be constant.

6.2. Is it an involution? Remark 6.7 shows it is easy to find functions that determine a 2-automorphism τ as in Theorem 6.5. However, we desire more, namely that τ be an involution. Our next result shows that any choice gets us close.

Corollary 6.8. When the conditions (6.2) are satisfied (so that τ is well-defined), we have $\tau^4 = \text{id}_{\mathcal{U}_q(\mathfrak{gl}_m)}$.

Proof. It is clear that τ^2 acts as the identity on all objects and 1-morphisms. Moreover, τ^2 sends each diagram to itself, up to multiplication by some sign. Therefore, τ^4 is the identity on 2-morphisms. \square

Applying (6.16b), we derive the equivalent formula

$$(6.18) \quad r_i(\mathbf{a}) + r_j(\mathbf{a}) + r_i(\alpha_i + \alpha_j - \mathbf{a}) + r_j(\alpha_i + \alpha_j - \mathbf{a}) = \langle \alpha_i^\vee + \alpha_j^\vee, \mathbf{a} \rangle,$$

which bears a comforting similarity to (6.16b), but for the function $r_i + r_j$.

A second property one might desire is that the formula for τ acting on $\mathcal{U}_q(\mathfrak{gl}_m)$ is compatible with a formula for τ on $\mathcal{U}_q(\mathfrak{gl}_2)$, with regards to the standard inclusions of \mathfrak{gl}_2 into \mathfrak{gl}_m . For \mathfrak{gl}_2 , there is a single function $r: \mathbb{Z}^2 \rightarrow \mathbb{Z}/2$, and it satisfies

$$(6.19) \quad r(a, b) + r(1 - a, -1 - b) \equiv a + b$$

by (6.16b). We thus say that the family of functions $\{r_i\}$ is *defined locally* if there exists a function $r: \mathbb{Z}^2 \rightarrow \mathbb{Z}/2$ satisfying (6.19), for which

$$(6.20) \quad r_i(a_1, \dots, a_i, a_{i+1}, \dots, a_m) = r(a_i, a_{i+1}).$$

Theorem 6.11. If $m \geq 3$, then there is no solution to (6.2) and (6.16) for which $v_{i,j}$ is a constant function for all i, j , and $\{r_i\}$ is defined locally.

Proof. Following the discussion above, regardless of the value of $v_{i,j}$ for $i < j$, we need only define a function r satisfying (6.19) for which (6.18) holds. Without loss of generality, we can assume that $r(0, 0) = 0$, since adding a constant function to r will not affect whether r is a solution.

Assume that $m = 3$, $i = 1$, $j = 2$ and let $\mathbf{a} = (a, b, c)$; more generally, we can choose $j = i + 1$ and study three consecutive coordinates of a general \mathfrak{gl}_m weight for $m \geq 3$. We then see that (6.18) is equivalent to

$$r(a, b) + r(b, c) + r(1 - a, -b) + r(-b, -1 - c) \equiv a + c,$$

while (6.19) gives

$$r(1 - a, -b) + r(a, b - 1) \equiv a + b + 1 \quad \text{and} \quad r(-b, -1 - c) + r(b + 1, c) \equiv b + c + 1.$$

Thus

$$r(a, b) + r(b, c) + r(1 - a, -b) + r(-b, -1 - c) \equiv r(a, b) + r(a, b - 1) + r(b, c) + r(b + 1, c) + a + c$$

so we conclude that

$$(6.21) \quad r(a, b) + r(a, b - 1) \equiv r(b, c) + r(b + 1, c)$$

in $\mathbb{Z}/2$.

Let $\epsilon_{a,b} := r(a, b) + r(a, b - 1)$, then (6.21) shows that this element of $\mathbb{Z}/2$ is equal to $r(b, c) + r(b + 1, c)$ for any c , implying that $\epsilon_{a,b}$ is independent of a . We thus write it as ϵ_b , i.e.

$$r(a, b) + r(a, b - 1) \equiv \epsilon_b \quad \text{and} \quad r(b, c) + r(b + 1, c) \equiv \epsilon_b$$

for all $a, c \in \mathbb{Z}$.

Recalling our assumption that $r(0, 0) = 0$ we compute

$$r(0, -1) = \epsilon_0 \quad \text{and} \quad r(1, 0) = \epsilon_0,$$

but this contradicts (6.19), which requires that $r(1, 0) + r(0, -1) \equiv 1$. \square

It is still an open question, and a surprisingly thorny one, whether there is any (non-naïve) choice of functions for which τ is well-defined and an involution.

Remark 6.12. It is interesting to replace the function $v_{ij}(\mathbf{a})$ with

$$w_{ij}(\mathbf{a}) := v_{ij}(-\mathbf{a}) + r_i(\mathbf{a}) + r_j(\mathbf{a}).$$

After doing so, (6.16c) can be rephrased (using (6.16b)) as

$$(6.22) \quad w_{ij}(\mathbf{a}) + w_{ij}(\alpha_i + \alpha_j - \mathbf{a}) \equiv \langle \alpha_i^\vee + \alpha_j^\vee, \mathbf{a} \rangle,$$

which bears a comforting similarity to (6.16b). One can now construct functions q_r and q_w (depending on indices $i < j < k$) with the symmetry

$$(6.23) \quad q(\mathbf{a}) + q(\alpha_i + \alpha_j + \alpha_k - \mathbf{a}) \equiv 0$$

by the formulas

$$q_w(\mathbf{a}) := w_{ij}(\mathbf{a}) + w_{jk}(\mathbf{a}) + w_{ik}(\mathbf{a}) + w_{ij}(\alpha_i + \alpha_j + \alpha_k - \mathbf{a}) + w_{jk}(\alpha_i + \alpha_j + \alpha_k - \mathbf{a}) + w_{ik}(\alpha_i + \alpha_j + \alpha_k - \mathbf{a}),$$

and

$$q_r(\mathbf{a}) := r_i(\mathbf{a} - \alpha_k) + r_i(\mathbf{a} - \alpha_j) + r_j(\mathbf{a} - \alpha_i) + r_j(\mathbf{a} - \alpha_k) + r_k(\mathbf{a} - \alpha_i) + r_k(\mathbf{a} - \alpha_j).$$

The final equation (6.2e) can then be reformulated as

$$(6.24) \quad q_w \equiv q_r.$$

6.3. An involution on part of $\mathcal{U}_q(\mathfrak{gl}_m)$. Although we are unable to find $r_i, v_{i,j}: \mathbb{Z}^m \rightarrow \mathbb{Z}/2$ such that the 2-automorphism τ from Theorem 6.5 is an involution, we now show that it is possible to find such functions so that τ restricts to an involution on a certain full 2-subcategory of $\mathcal{U}_q(\mathfrak{gl}_m)$. Moreover, the functions we construct are such that $\{r_i\}$ is defined locally and $v_{i,j}$ is a constant function for all i, j ; c.f. Theorem 6.11.

Definition 6.13. Fix a \mathfrak{gl}_m weight $\mathbf{a} \in \mathbb{Z}^m$. Let $\mathcal{F}_q(\mathfrak{gl}_m)\mathbb{1}_{\mathbf{a}}$ denote the full 2-subcategory of $\mathcal{U}_q(\mathfrak{gl}_m)$ generated by the 1-endomorphisms $\mathcal{E}_i\mathcal{F}_i\mathbb{1}_{\mathbf{a}}$ and $\mathcal{F}_i\mathcal{E}_i\mathbb{1}_{\mathbf{a}}$, and their grading shifts. Let $\mathcal{F}_q(\mathfrak{gl}_m)$ be the full 2-subcategory containing all $\mathcal{F}_q(\mathfrak{gl}_m)\mathbb{1}_{\mathbf{a}}$ as \mathbf{a} varies.

Since $\mathcal{E}_i\mathcal{F}_i\mathbb{1}_{\mathbf{a}}$ and $\mathcal{F}_i\mathcal{E}_i\mathbb{1}_{\mathbf{a}}$ are 1-endomorphisms (mapping from \mathbf{a} to \mathbf{a}), $\mathcal{F}_q(\mathfrak{gl}_m)\mathbb{1}_{\mathbf{a}} = \mathbb{1}_{\mathbf{a}}\mathcal{F}_q(\mathfrak{gl}_m)\mathbb{1}_{\mathbf{a}}$ is a monoidal category. There is no interaction between $\mathcal{F}_q(\mathfrak{gl}_m)\mathbb{1}_{\mathbf{a}}$ and $\mathcal{F}_q(\mathfrak{gl}_m)\mathbb{1}_{\mathbf{a}'}$ for $\mathbf{a} \neq \mathbf{a}'$.

Theorem 6.14. If τ is well-defined, then it restricts to an involution on $\mathcal{F}_q(\mathfrak{gl}_m)$ if and only if $l_i(\mathbf{a}) \equiv r_i(-\mathbf{a})$ for all \mathbf{a} .

Proof. If τ is well-defined (i.e. (6.2) is satisfied), then τ^2 is the identity on new bubble generators and, since $v_{i,i} = 0$, it is also the identity on same-colored crossings. As observed in the proof of Lemma 6.9, the condition $l_i(\mathbf{a}) \equiv r_i(-\mathbf{a})$ is equivalent to the condition that τ^2 acts as the identity on cap/cup generators in weight $\mathbf{n} + \mathbf{a}$. This, in particular, establishes the “only if” assertion of the statement, since $\mathcal{F}_q(\mathfrak{gl}_m)$ contains all such 2-morphisms.

We thus assume that the condition $l_i(\mathbf{a}) \equiv r_i(-\mathbf{a})$ holds for all $\mathbf{a} \in \mathbb{Z}^m$. It remains to consider the action of τ^2 on crossings of differently colored strands. Here, τ^2 introduces rescalings by hard-to-control signs; the key observation is that we can track these contributions for 2-morphisms lying in $\mathcal{F}_q(\mathfrak{gl}_m)$.

To begin, consider the full subcategory $\mathcal{F}'_q(\mathfrak{gl}_m)$ of $\mathcal{F}_q(\mathfrak{gl}_m)$ generated by objects of the form $\mathcal{F}_i\mathcal{E}_i\mathbb{1}_{\mathbf{a}}$. We claim that if τ^2 is the identity on $\mathcal{F}'_q(\mathfrak{gl}_m)$, then it is the identity on all of $\mathcal{F}_q(\mathfrak{gl}_m)$. This is a consequence of (5.10), which describes the identity of $\mathcal{E}_i\mathcal{F}_i\mathbb{1}_{\mathbf{a}}$ as a sum of diagrams which factor through $\mathcal{F}_i\mathcal{E}_i\mathbb{1}_{\mathbf{a}}$ and $\mathbb{1}_{\mathbf{a}}$. More precisely, using (5.11), any morphism in $\mathcal{F}_q(\mathfrak{gl}_m)$ factors as a composition $f \circ g \circ h$ where g is a morphism in $\mathcal{F}'_q(\mathfrak{gl}_m)$ and f and h are built from cups, caps, dots, and same-colored crossings. Since f and h are fixed by τ^2 , we deduce the claim.

For the rest of the proof, we focus on $\mathcal{F}'_q(\mathfrak{gl}_m)\mathbb{1}_{\mathbf{a}}$ for some fixed weight \mathbf{a} . Let D be a diagram which represents a morphism in $\mathcal{F}'_q(\mathfrak{gl}_m)\mathbb{1}_{\mathbf{a}}$, and consider the diagram D' obtained from D by replacing all same-colored crossings with identity maps:

$$\begin{array}{c} \begin{array}{c} \uparrow \quad \uparrow \\ \diagdown \quad \diagup \\ \uparrow \quad \uparrow \end{array} \rightsquigarrow \begin{array}{c} \uparrow \quad \uparrow \\ | \quad | \\ \uparrow \quad \uparrow \end{array} \end{array}$$

The morphisms D and D' are completely unrelated; nonetheless, if τ^2 fixes D' then it fixes D . Hence, it suffices to show that τ^2 acts as the identity on diagrams with no same-colored crossings. Similarly, we can assume that our diagrams have no dots or new bubbles (τ^2 will act as the identity on any


For gRIII, it suffices to check the braid-like orientation




since the others can be obtained¹⁶ from this using gRII moves. Again, τ^2 is a 2-functor, so it preserves the relation (5.6). Thus, it must have the same sign on both sides of gRIII, which concludes the proof of the claim. \blacktriangle

Any two matching diagrams with the same underlying matching are related by the moves (6.27), hence have the same value of $T(\mathcal{M})$. We now pass to different matching diagram \mathcal{M}' having the same underlying matching as our given diagram \mathcal{M} , for which the value $T(\mathcal{M}')$ is easy to compute.

Pick the last color appearing in \mathcal{M} and use gRII and gRIII to “pull all instances of this color” to the far right. For example, if we start with (6.25) then this color is blue and the result is




for




For the remainder of this proof, we will use “blue” to mean this final color. In general, the subdiagram \mathcal{M}_{rg} containing the other colors will be an arbitrary colored matching diagram (without blue), and the subdiagram to the right of that can be an arbitrary matching diagram (with only blue).

The meeting of “blue” and “non-blue” strands occur only in a proscribed fashion. Consider the shading on \mathcal{M}' :



The only intersections of blue strands and non-blue strands occur when blue-shaded strips cross over non-blue strips:



Said another way, each region with blue in its shading is either pure blue, or blue and one other color, and in the latter case the region is a square. We call these *doubly-shaded squares*.

¹⁶See e.g. [52], which accomplishes the more difficult task of showing that only five oriented link Reidemeister moves suffice to obtain all others. Our task here is easier since we work with graphs, and have already established all versions of gRII.

The advantage of having done this manipulation is that each crossing in a doubly-shaded square is adjacent to a white-colored region, i.e. a region in weight \mathbf{a} . Hence, in each doubly-shaded square there are two crossings of the form:

$$(6.28) \quad \begin{array}{c} \text{red} \\ \diagup \quad \diagdown \\ \text{blue} \end{array}$$

and two crossings of the form:

$$(6.29) \quad \begin{array}{c} \text{blue} \\ \diagdown \quad \diagup \\ \text{red} \end{array}.$$

The signs induced by τ^2 on the two instances of (6.28) are the same, and therefore cancel out. The same goes for (6.29).

Thus τ^2 fixes \mathcal{M}' if and only if it fixes \mathcal{M}_{rg} , a colored matching with one fewer color. By induction on the number of colors, τ^2 fixes all diagrams representing morphisms in $\mathcal{FE}'_q(\mathfrak{gl}_m)$. \square

Remark 6.15. If we directly consider the original diagram \mathcal{M} from (6.25), there would be red-blue crossings in the green-shaded region. The action of τ^2 on this diagram can introduce different signs on these crossings than the ones in the white region, which makes it harder to argue that all signs cancel out. However, our proof shows that they must.

We now fix our preferred choice of involution on $\mathcal{FE}_q(\mathfrak{gl}_m)$, which we use henceforth.

Theorem 6.16. If

$$(6.30) \quad r_i(\mathbf{a}) := \begin{cases} a_i & \text{if } \langle \alpha_i^\vee, \mathbf{a} \rangle \equiv 1 \\ 0 & \text{else} \end{cases}, \quad l_i(\mathbf{a}) \equiv r_i(-\mathbf{a}), \quad v_{i,j}(\mathbf{a}) := \begin{cases} 1 & \text{if } j = i + 1 \\ 0 & \text{else} \end{cases}$$

then τ restricts to an involution on $\mathcal{FE}_q(\mathfrak{gl}_m)$.

Proof. Recall that $\langle \alpha_i^\vee, \mathbf{a} \rangle = a_i - a_{i+1}$. It is straightforward to check that these functions satisfy (6.2), where we let $r'_i(\mathbf{a})$ and $l'_i(\mathbf{a})$ be defined by (6.2a). Since $l_i(\mathbf{a}) \equiv r_i(-\mathbf{a})$, τ restricts to an involution on $\mathcal{FE}_q(\mathfrak{gl}_m)$ by Theorem 6.14. \square

6.4. Dependence on n . Although the notation obscures it, the assignments in Definition 6.3 depend on n , e.g. via the action of τ on objects. On the other hand, the formulae in Theorem 6.16 giving our preferred choice of τ do not depend on n . In this section, we clarify the dependence on n for our preferred τ .

Notation 6.17. For this section (and in various other places when we want to emphasize the pertinent value of n), we will write τ_n to denote the preferred τ which acts on objects by $\mathbf{a} \mapsto 2\mathbf{n} - \mathbf{a}$.

Note that τ_0 sends \mathbf{a} to $-\mathbf{a}$, while τ_n sends $\mathbf{n} + \mathbf{a}$ to $\mathbf{n} - \mathbf{a}$. Suppose there were an automorphism sh_n of $\mathcal{U}_q(\mathfrak{gl}_m)$ which mapped \mathbf{a} to $\mathbf{a} + \mathbf{n}$. Then, by conjugating τ_0 by sh_n , we obtain an automorphism which would act as

$$\mathbf{n} + \mathbf{a} \xrightarrow{\text{sh}_n^{-1}} \mathbf{a} \xrightarrow{\tau_0} -\mathbf{a} \xrightarrow{\text{sh}_n} \mathbf{n} - \mathbf{a}.$$

Ideally, this automorphism would be identified with τ_n .

We now set out to define automorphisms $\text{sh}_n: \mathcal{U}_q(\mathfrak{gl}_m) \rightarrow \mathcal{U}_q(\mathfrak{gl}_m)$ that send $\mathbf{a} \mapsto \mathbf{a} + \mathbf{n}$ and satisfy $\text{sh}_n \circ \tau_0 = \tau_n \circ \text{sh}_n$.

Proposition 6.18. The following assignments determine a 2-functor $\text{sh}_1: \mathcal{U}_q(\mathfrak{gl}_m) \rightarrow \mathcal{U}_q(\mathfrak{gl}_m)$.

- **Objects:** $\mathbf{a} \xrightarrow{\text{sh}_1} \mathbf{a} + \mathbf{1}$.
- **1-morphisms:** $\mathcal{E}_i \mathbb{1}_{\mathbf{a}} \xrightarrow{\text{sh}_1} \mathcal{E}_i \mathbb{1}_{\mathbf{a} + \mathbf{1}}$ and $\mathcal{F}_i \mathbb{1}_{\mathbf{a}} \xrightarrow{\text{sh}_1} \mathcal{F}_i \mathbb{1}_{\mathbf{a} + \mathbf{1}}$.

6.5. Extending to thick calculus. We now observe that our automorphism τ extends to the thick calculus of Section 5.4 and record its action there.

Proposition 6.22. Suppose that τ satisfies the conditions of Theorem 6.5. There is a unique extension of τ to $\tilde{\mathcal{U}}_q(\mathfrak{gl}_m)$ such that $\mathcal{E}_i^{(k)} \mathbb{1}_{n+\mathbf{a}} \mapsto \mathcal{F}_i^{(k)} \mathbb{1}_{n-\mathbf{a}}$ and

$$(6.33) \quad \begin{array}{c} k+\ell \\ \uparrow \\ k \quad \ell \\ \text{n+a} \end{array} \xrightarrow{\tau} \begin{array}{c} k+\ell \\ \downarrow \\ k \quad \ell \\ \text{n-a} \end{array} \xrightarrow{\tau} \begin{array}{c} k+\ell \\ \uparrow \\ k \quad \ell \\ \text{n+a} \end{array}, \quad \begin{array}{c} k \quad \ell \\ \uparrow \\ k+\ell \\ \text{n+a} \end{array} \xrightarrow{\tau} \begin{array}{c} k \quad \ell \\ \downarrow \\ k+\ell \\ \text{n-a} \end{array} \xrightarrow{\tau} \begin{array}{c} k \quad \ell \\ \uparrow \\ k+\ell \\ \text{n+a} \end{array}.$$

Moreover, under this extension

$$(6.34) \quad \begin{array}{c} \text{n+a} \\ \downarrow \\ k \end{array} \mapsto (-1)^{\sum_{s=0}^{k-1} r_{\bullet}(\mathbf{a}-s\alpha_{\bullet})} \begin{array}{c} \text{n-a} \\ \downarrow \\ k \end{array}, \quad \begin{array}{c} \text{n+a} \\ \uparrow \\ k \end{array} \mapsto (-1)^{\sum_{s=0}^{k-1} \ell_{\bullet}(\mathbf{a}+s\alpha_{\bullet})} \begin{array}{c} \text{n-a} \\ \uparrow \\ k \end{array},$$

$$\begin{array}{c} k \\ \downarrow \\ \text{n+a} \end{array} \mapsto (-1)^{\sum_{s=0}^{k-1} r'_{\bullet}(\mathbf{a}+s\alpha_{\bullet})} \begin{array}{c} k \\ \downarrow \\ \text{n-a} \end{array}, \quad \begin{array}{c} k \\ \uparrow \\ \text{n+a} \end{array} \mapsto (-1)^{\sum_{s=0}^{k-1} \ell'_{\bullet}(\mathbf{a}-s\alpha_{\bullet})} \begin{array}{c} k \\ \uparrow \\ \text{n-a} \end{array},$$

$$(6.35) \quad \begin{array}{c} \text{n+a} \\ \swarrow \searrow \\ k \quad \ell \end{array} \mapsto (-1)^{\sum_{\substack{0 \leq s \leq k-1 \\ 0 \leq t \leq \ell-1}} v_{\bullet}(\mathbf{a}+s\alpha_{\bullet}+t\alpha_{\bullet})} \begin{array}{c} \text{n-a} \\ \swarrow \searrow \\ k \quad \ell \end{array},$$

and

$$(6.36) \quad \begin{array}{c} \text{n+a} \\ \uparrow \\ \textcircled{s_{\lambda}} \\ \downarrow \\ k \end{array} \mapsto (-1)^{|\lambda|} \begin{array}{c} \text{n-a} \\ \downarrow \\ \textcircled{s_{\lambda}} \\ \uparrow \\ k \end{array}.$$

Proof. Since τ preserves the relations of $\mathcal{U}_q(\mathfrak{gl}_m)$, it follows from Definition 5.16 that the extension of τ to $\tilde{\mathcal{U}}_q(\mathfrak{gl}_m)$ is well-defined if and only if it preserves the relations in (5.19) and (5.20). The former follows by (6.2c), i.e. since τ introduces no signs on thin uni-colored crossings, regardless of ambient weight. The latter is a consequence of τ acting on dots by -1 (i.e. the first relation in (6.1)) and the formula:

$$(6.37) \quad \begin{array}{c} k \\ \uparrow \\ \textcircled{\bullet} \\ \downarrow \\ k \end{array} \begin{array}{c} \text{a} \\ \bullet \\ \bullet \\ \bullet \\ \bullet \end{array} = (-1)^{\binom{k}{2}} \begin{array}{c} k \\ \uparrow \\ \textcircled{\bullet} \\ \downarrow \\ k \end{array} \begin{array}{c} \text{a} \\ \bullet \\ \bullet \\ \bullet \\ \bullet \end{array}$$

which is a consequence of (5.32).

The action of τ on thick caps follows from applying τ , as defined in (6.1), to the right hand side of (5.23) then using (6.37). The action of τ on thick cups is analogous. The action of τ on thick crossings (with one or two colors) follows from applying τ to the right hand side of (5.24) or (5.37), then using (6.37) in conjunction with (5.25), in the case of one color crossings, and (5.38), in the case of two color crossings. Finally, the action of τ on $s_{\lambda} \in \text{End}(\mathcal{E}_i^{(k)} \mathbb{1}_{\mathbf{a}})$ comes from applying τ to the right hand side of (5.27). \square

Corollary 6.23. Our preferred τ from Theorem 6.16, acts on thick cap, cup, and upward crossing 2-morphisms as follows:

(6.38)

$$\begin{array}{c} \text{thick cap} \\ \xrightarrow{\tau} \begin{cases} \text{thick cap} & \text{if } \langle \alpha_{\bullet}^{\vee}, \mathbf{a} \rangle \equiv 0 \\ (-1)^{k \cdot a_{\bullet} - \binom{k}{2}} \text{thick cap} & \text{if } \langle \alpha_{\bullet}^{\vee}, \mathbf{a} \rangle \equiv 1 \end{cases}, \quad \begin{array}{c} \text{thick cup} \\ \xrightarrow{\tau} \begin{cases} \text{thick cup} & \text{if } \langle \alpha_{\bullet}^{\vee}, \mathbf{a} \rangle \equiv 0 \\ (-1)^{-k \cdot a_{\bullet} - \binom{k}{2}} \text{thick cup} & \text{if } \langle \alpha_{\bullet}^{\vee}, \mathbf{a} \rangle \equiv 1 \end{cases} \end{array} \end{array}$$

(6.39)

$$\begin{array}{c} \text{thick upward crossing} \\ \xrightarrow{\tau} \begin{cases} \text{thick upward crossing} & \text{if } \langle \alpha_{\bullet}^{\vee}, \mathbf{a} \rangle \equiv 0 \\ (-1)^{k \cdot a_{\bullet} + \binom{k+1}{2}} \text{thick upward crossing} & \text{if } \langle \alpha_{\bullet}^{\vee}, \mathbf{a} \rangle \equiv 1 \end{cases}, \quad \begin{array}{c} \text{thick downward crossing} \\ \xrightarrow{\tau} \begin{cases} \text{thick downward crossing} & \text{if } \langle \alpha_{\bullet}^{\vee}, \mathbf{a} \rangle \equiv 0 \\ (-1)^{-k \cdot a_{\bullet} + \binom{k+1}{2}} \text{thick downward crossing} & \text{if } \langle \alpha_{\bullet}^{\vee}, \mathbf{a} \rangle \equiv 1 \end{cases} \end{array} \end{array}$$

$$(6.40) \quad \begin{array}{c} \text{thick crossing} \\ \mapsto \begin{cases} (-1)^{k\ell} \text{thick crossing} & \text{if } \bullet = \bullet, \\ \text{thick crossing} & \text{otherwise.} \end{cases} \end{array}$$

(Recall our color Convention 5.18.) Consequently, τ^2 acts as the identity on all uni-colored thick diagrams. \square

Theorem 6.16 also extends to thick calculus, after introducing the relevant 2-categories. Note that for any weight \mathbf{a} , we have $\mathcal{E}_i^{(k)} \mathcal{F}_i^{(k)} \mathbb{1}_{\mathbf{a}} = \mathbb{1}_{\mathbf{a}} \mathcal{E}_i^{(k)} \mathcal{F}_i^{(k)} \mathbb{1}_{\mathbf{a}}$.


Definition 6.24. Fix a weight \mathbf{a} . Let $\check{\mathcal{F}}_q(\mathfrak{gl}_m) \mathbb{1}_{\mathbf{a}}$ denote the full 2-subcategory of $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$ generated by the 1-endomorphisms $\mathcal{E}_i^{(k)} \mathcal{F}_i^{(k)} \mathbb{1}_{\mathbf{a}}$ and $\mathcal{F}_i^{(k)} \mathcal{E}_i^{(k)} \mathbb{1}_{\mathbf{a}}$, and their grading shifts. Let $\check{\mathcal{E}}_q(\mathfrak{gl}_m)$ be the full 2-subcategory containing all $\check{\mathcal{F}}_q(\mathfrak{gl}_m) \mathbb{1}_{\mathbf{a}}$ as \mathbf{a} varies.

As before, each $\check{\mathcal{F}}_q(\mathfrak{gl}_m) \mathbb{1}_{\mathbf{a}}$ is a monoidal category and there is no interaction between $\check{\mathcal{F}}_q(\mathfrak{gl}_m) \mathbb{1}_{\mathbf{a}}$ and $\check{\mathcal{F}}_q(\mathfrak{gl}_m) \mathbb{1}_{\mathbf{a}'}$ for $\mathbf{a} \neq \mathbf{a}'$.

Corollary 6.25. Let τ be as in Theorem 6.16. The extension of τ to $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$ restricts to an involution on $\check{\mathcal{E}}_q(\mathfrak{gl}_m)$.

Proof. Fix $\mathbf{a} \in \mathbb{Z}^m$ and consider the monoidal category $\check{\mathcal{F}}_q(\mathfrak{gl}_m) \mathbb{1}_{\mathbf{a}}$, which is generated by $\mathcal{E}_i^{(k)} \mathcal{F}_i^{(k)} \mathbb{1}_{\mathbf{a}}$ and $\mathcal{F}_i^{(k)} \mathcal{E}_i^{(k)} \mathbb{1}_{\mathbf{a}}$. Note that if $a_i - a_{i+1} > 0$, then we can use the Stošić formula (5.36a) to write the identity morphism of $\mathcal{E}_i^{(k)} \mathcal{F}_i^{(k)} \mathbb{1}_{\mathbf{a}}$ as a linear combination of morphisms that factor through the objects $\mathcal{F}_i^{(\ell)} \mathcal{E}_i^{(\ell)} \mathbb{1}_{\mathbf{a}}$ for $0 \leq \ell \leq k$. (The objects being factored through cannot be rewritten in the same way.) Similarly, if $a_i - a_{i+1} < 0$ then we can use (5.36b) to write the identity morphism of $\mathcal{F}_i^{(k)} \mathcal{E}_i^{(k)} \mathbb{1}_{\mathbf{a}}$ as a linear combination of morphisms that factor through $\mathcal{E}_i^{(\ell)} \mathcal{F}_i^{(\ell)} \mathbb{1}_{\mathbf{a}}$ for $0 \leq \ell \leq k$. As observed in Corollary 6.23, τ^2 acts as the identity on all uni-colored diagrams, hence on tensor products of uni-colored diagrams. Since the diagrams in the Stošić formula are uni-colored, we see that τ^2 will equal the identity if and only if it acts as the identity the full subcategory generated by $\mathcal{E}_i^{(k)} \mathcal{F}_i^{(k)} \mathbb{1}_{\mathbf{a}}$ for $a_i - a_{i+1} \leq 0$ and $\mathcal{F}_i^{(k)} \mathcal{E}_i^{(k)} \mathbb{1}_{\mathbf{a}}$ for $a_i - a_{i+1} \geq 0$.

Next, we observe that when $a_i - a_{i+1} \leq 0$ the identity morphism of $\mathcal{E}_i^{(k)} \mathcal{F}_i^{(k)} \mathbb{1}_{\mathbf{a}}$ factors through the object $(\mathcal{E}_i \mathcal{F}_i \mathbb{1}_{\mathbf{a}})^{\otimes k}$. This follows by observing that, in this case, $\mathcal{E}_i^{(k)} \mathcal{F}_i^{(k)} \mathbb{1}_{\mathbf{a}}$ is a summand of $(\mathcal{E}_i \mathcal{F}_i \mathbb{1}_{\mathbf{a}})^{\otimes k}$ (which can be confirmed using Theorem 5.4 by expressing $(e_i f_i \mathbb{1}_{\mathbf{a}})^k \in {}_{\mathbb{Z}} \dot{U}_q(\mathfrak{gl}_m)$ in terms of the canonical basis for this \mathfrak{sl}_2 string). Alternatively, this can be done explicitly, e.g. when $a_i - a_{i+1} \leq 0$ and $k = 3$, we have



and similar formulae (for all $k \geq 2$) can be deduced from (5.10). Similarly, when $a_i - a_{i+1} \geq 0$ the identity morphism of $\mathcal{F}_i^{(k)} \mathcal{E}_i^{(k)} \mathbb{1}_{\mathbf{a}}$ factors through the object $(\mathcal{F}_i \mathcal{E}_i \mathbb{1}_{\mathbf{a}})^{\otimes k}$. Again, since τ^2 is the identity on tensor products of uni-colored diagrams, we see that τ^2 is the identity if and only if it acts as the identity on the full subcategory generated by $(\mathcal{E}_i \mathcal{F}_i \mathbb{1}_{\mathbf{a}})^{\otimes k}$ for $a_i - a_{i+1} \leq 0$ and $(\mathcal{F}_i \mathcal{E}_i \mathbb{1}_{\mathbf{a}})^{\otimes k}$ for $a_i - a_{i+1} \geq 0$. This holds by Theorem 6.16. \square

7. BACKGROUND ON DECOMPOSITIONS AND EQUIVARIANT CATEGORIES

There are two main goals in this section. First, we study direct sum decompositions of objects into indecomposables via composition pairings. Second, in Section 7.4, we recall the definition and elementary structure theory of equivariant categories. We combine these ideas to discuss the classification of indecomposable objects in equivariant categories. This material is well-known to experts, and mostly adapted from the overview given in [20].

Some basic results we will repeatedly quote in the sequel are the following.

- Lemma 7.25, which gives a basis for certain multiplicity spaces in the presence of known direct sum decompositions.
- Proposition 7.50, which states that one can decompose objects in equivariant categories by computing eigenspaces of a certain involution σ^* acting on multiplicity spaces.
- Lemma 7.52, which gives an efficient technique for computing the action of σ^* on multiplicity spaces.

Lemma 7.52 did not previously appear in [20].

We work with \mathbb{K} -linear categories over a commutative ring \mathbb{K} . Until §7.5, there are very few restrictions we need to place on the base ring \mathbb{K} for the results above to hold, so long as one studies objects whose (degree zero) morphism spaces are well-understood as \mathbb{K} -modules. However, in [20] and elsewhere in the literature, such results are typically proven under the assumption that \mathbb{K} is an algebraically closed field. Where it streamlines the exposition, we do assume that \mathbb{K} is an algebraically closed field. Supplementary material dealing with more general \mathbb{K} can be found in Appendix C.

7.1. Composition pairings and endopositive objects. We now discuss direct sum decompositions, and tools for studying them in various contexts. Throughout, we are motivated by the case of \mathbb{K} -linear categories when \mathbb{K} is a field, but we state results that hold over general base rings.

Definition 7.1 ([22, Definition 11.64]). Let \mathbb{K} be a commutative ring and let X and Y be objects in a graded additive \mathbb{K} -linear category \mathcal{A} . The *naïve composition pairing* of Y at X is the \mathbb{K} -bilinear pairing

$$\beta_{X,Y} : \text{Hom}(Y, X) \times \text{Hom}(X, Y) \longrightarrow \text{End}(X)$$

defined by $\beta_{X,Y}(f, g) = f \circ g$.

The naïve composition pairing is valued in $\text{End}(X)$ rather than in \mathbb{K} , but we will soon have cause to discuss \mathbb{K} -valued bilinear pairings as well.

Definition 7.2. Given sets of morphisms $\{p_1, \dots, p_d\} \in \text{Hom}(Y, X)$ and $\{\iota_1, \dots, \iota_d\} \in \text{Hom}(X, Y)$, we call them *dual sets* with respect to the naïve composition pairing $\beta_{X,Y}$ if

$$(7.1) \quad \beta_{X,Y}(p_j, \iota_\ell) = \delta_{j\ell} \text{id}_X.$$

Similarly, given \mathbb{K} -modules V and V' and a \mathbb{K} -bilinear pairing $\beta: V \times V' \rightarrow \mathbb{K}$, sets $\{p_1, \dots, p_d\} \in V$ and $\{\iota_1, \dots, \iota_d\} \in V'$ are called *dual sets* if

$$\beta(p_j, \iota_\ell) = \delta_{j\ell}.$$

In either context, the *naïve rank* of the pairing is the maximal integer d such that one can find dual sets of size d .

The following lemma is tautological, following from the definition of direct sums and summands.

Lemma 7.3. The naïve rank of the composition pairing is the multiplicity of X as a direct summand of Y .

Proof. Dual sets give (orthogonal) projection and inclusion maps for d copies of X appearing as summands within Y . \square

Remark 7.4. For a \mathbb{K} -valued bilinear pairing β , if one finds elements $\{p_1, \dots, p_d\}$ and $\{\iota_1, \dots, \iota_d\}$ for which the $d \times d$ matrix with entries $\beta(p_j, \iota_\ell)$ has invertible determinant in \mathbb{K} , the naïve rank of the pairing is at least d . This follows using standard techniques in linear algebra (e.g. Cramer's rule) to find \mathbb{K} -linear combinations $\{p'_1, \dots, p'_d\}$ of the original elements $\{p_j\}$ which are dual to $\{\iota_\ell\}$.

One drawback of the naïve composition pairing is that it is valued in the \mathbb{K} -algebra $\text{End}(X)$. When $\text{End}(X)$ is a (graded) local ring with (homogeneous) maximal ideal $J(X)$, then $\text{End}(X)/J(X)$ is a division algebra. In this case we can use techniques of linear algebra over division algebras.

Definition 7.5 ([22, Definition 11.70]). Let \mathbb{K} be a commutative ring and let X and Y be objects in a graded additive \mathbb{K} -linear category \mathcal{A} . Suppose that $\text{End}(X)$ is a (graded) local ring with (homogeneous) maximal ideal $J(X)$. The *local composition pairing*, which we also denote $\beta_{X,Y}$, is defined in the same way as the naïve composition pairing except that one takes the image of $f \circ g$ in the division algebra $\text{End}(X)/J(X)$.

We point out the following stronger result in this case.

Lemma 7.6 ([22, Corollary 11.71]). Let X have (graded) local endomorphism algebra. The (graded) rank¹⁷ of the local composition pairing $\beta_{X,Y}$ is equal to the (graded) multiplicity of X as a summand of Y . \square

Remark 7.7. This lemma says that we need not find true dual sets to find summands of Y , but need only find sets which are dual modulo the maximal ideal of $\text{End}(X)$. This is analogous to classical results involving idempotent lifting modulo the Jacobson radical [22, Proposition 11.69]. Those results require the category to be linear over a complete ring \mathbb{K} ; however, as shown in loc. cit., these assumptions can be omitted when the idempotent is factored as a composition of inclusion and projection maps.

Rather than working modulo the maximal ideal (or when this is not necessarily possible), we can instead focus on situations where the composition pairing is valued in \mathbb{K} rather than in $\text{End}(X)$. We thus introduce the following notion.

¹⁷One can make sense of the rank of a bilinear pairing valued in a division ring.

Definition 7.8. Let \mathbb{K} be a commutative ring and \mathcal{A} be a graded additive \mathbb{K} -linear category. An object $X \in \mathcal{A}$ will be called *endopositive* if the graded \mathbb{K} -algebra $\text{End}(X)$ is supported in degrees ≥ 0 and $\text{End}^0(X) = \mathbb{K} \cdot \text{id}_X$.

If \mathbb{K} is a field, then any endopositive object X is indecomposable in \mathcal{A} , and even in $\text{Kar}(\mathcal{A})$, since the endomorphism ring of X is graded local. In general, \mathbb{K} need not be a domain, or local, so $\text{End}(X)$ might have nontrivial idempotents. The reader should view endopositive objects as a stand-in for indecomposable objects when working in \mathbb{K} -linear categories over general commutative rings. In particular, endopositive objects become indecomposable after base change to a field.

For any endopositive object, computing composition pairings one degree at a time gives the following \mathbb{K} -valued pairing.

Definition 7.9. Let \mathcal{A} be a graded additive \mathbb{K} -linear category over a commutative ring \mathbb{K} and let X be an endopositive object. The (\mathbb{K} -valued) *graded composition pairing of Y at $\mathbf{q}^k X$* is the following restriction of the naïve composition pairing:

$$(7.2) \quad \beta_{X,Y}^k : \text{Hom}^{-k}(Y, X) \times \text{Hom}^k(X, Y) \longrightarrow \text{End}^0(X) = \mathbb{K} \cdot \text{id}_X.$$

Corollary 7.10. Let X be endopositive. The naïve rank of $\beta_{X,Y}^k$ is equal to the multiplicity of $\mathbf{q}^k X$ as a summand of Y . \square

Proof. This follows from Lemma 7.3. \square

Remark 7.11. If X is endopositive and \mathbb{K} is a field, then $\text{End}(X)$ is graded local, the graded Jacobson radical $J(X)$ is equal to $\text{End}^{>0}(X)$, and the inclusion of degree zero endomorphisms induces an isomorphism $\mathbb{K} \cdot \text{id}_X = \text{End}^0(X) \cong \text{End}(X)/J(X)$. In particular, $\beta_{X,Y}^k$ is a graded slice of the local composition pairing as in Definition 7.5.

Informally, one can view the naïve rank of $\beta_{X,Y}^k$ as the “number of independent inclusion maps” from $\mathbf{q}^k X$ to Y ; however, in general there is no canonical subspace of $\text{Hom}(X, Y)$ of this dimension spanned by such inclusion maps. We now define a canonical quotient of $\text{Hom}(X, Y)$ (having this dimension, when \mathbb{K} is a field), which can be viewed as the analogue of a multiplicity space.

Definition 7.12. Let the right and left kernels of $\beta_{X,Y}^k$ be the \mathbb{K} -linear subspaces

$$\begin{aligned} \text{Rker}(\beta_{X,Y}^k) &:= \{g \in \text{Hom}^k(X, Y) \mid \beta_{X,Y}^k(f, g) = 0 \text{ for all } f \in \text{Hom}^{-k}(Y, X)\} \\ \text{Lker}(\beta_{X,Y}^k) &:= \{f \in \text{Hom}^{-k}(Y, X) \mid \beta_{X,Y}^k(f, g) = 0 \text{ for all } g \in \text{Hom}^k(X, Y)\} \end{aligned}$$

and set

$$V^k(X, Y) := \text{Hom}^k(X, Y) / \text{Rker}(\beta_{X,Y}^k), \quad V^{-k}(Y, X) := \text{Hom}^{-k}(Y, X) / \text{Lker}(\beta_{X,Y}^k).$$

The induced pairing

$$(7.3) \quad \mathbf{L}_{X,Y}^k : V^{-k}(Y, X) \times V^k(X, Y) \rightarrow \mathbb{K}$$

is the *non-degenerate graded composition pairing*.

Remark 7.13. The notation $V^k(X, Y)$ and $V^{-k}(Y, X)$ does not treat X and Y symmetrically, so one must know from context that we consider the composition pairing of Y at X , and not vice versa. In practice, this distinction will be obvious, as X will be the object which is endopositive.

The following is clear.

Lemma 7.14. Let X be endopositive. The naïve rank of the graded composition pairing (of Y at $\mathbf{q}^k X$) equals the naïve rank of the non-degenerate graded composition pairing. If \mathbb{K} is a field, this rank is equal to $\dim V^k(X, Y)$. \square

We also make the following observations.

Lemma 7.15. Let X be an endopositive object in \mathcal{A} , and Y_1, Y_2 be arbitrary objects. The isomorphism $\text{Hom}(X, Y_1 \oplus Y_2) \cong \text{Hom}(X, Y_1) \oplus \text{Hom}(X, Y_2)$ induces an isomorphism $V^k(X, Y_1 \oplus Y_2) \cong V^k(X, Y_1) \oplus V^k(X, Y_2)$ for each $k \in \mathbb{Z}$. Similarly, $V^{-k}(Y_1 \oplus Y_2, X) \cong V^{-k}(Y_1, X) \oplus V^{-k}(Y_2, X)$.

Proof. It is straightforward to verify that the natural isomorphism of Hom-spaces induces an isomorphism

$$\text{Lker}(\beta_{X, Y_1 \oplus Y_2}^k) \cong \text{Lker}(\beta_{X, Y_1}^k) \oplus \text{Lker}(\beta_{X, Y_2}^k).$$

This implies the first statement, and the second is similar. \square

Notation 7.16. For endopositive X , let $V(X, Y) := \bigoplus_k V^k(X, Y)$ and $V(Y, X) := \bigoplus_k V^{-k}(Y, X)$.

Corollary 7.17. Let X be endopositive. If $Y \cong \bigoplus_{j=1}^d \mathbf{q}^{k_j} X$ for some shifts $k_j \in \mathbb{Z}$, then $V(Y, X)$ (resp. $V(X, Y)$) is a free graded \mathbb{K} -module with graded rank $\sum_{j=1}^d q^{k_j}$ (resp. $\sum_{j=1}^d q^{-k_j}$). If one chooses a decomposition as above with projection maps $\{p_j\}$ and inclusion maps $\{\iota_j\}$, then $\{p_j\} \subset \text{Hom}(Y, X)$ descends to a basis for $V(Y, X)$ and $\{\iota_j\} \subset \text{Hom}(X, Y)$ descends to a basis for $V(X, Y)$.

Proof. It is immediate to verify when $Y = X$ that $V(Y, X)$ and $V(X, Y)$ are free of rank 1 over \mathbb{K} , spanned by the identity map. The general result for $Y = \bigoplus_j \mathbf{q}^{k_j} X$ then follows from Lemma 7.15. Note that under the isomorphism

$$\text{Hom}(X, Y) \cong \text{Hom}(X, \bigoplus_j \mathbf{q}^{k_j} X) \cong \bigoplus_j \mathbf{q}^{k_j} \text{End}(X)$$

the identity maps of X on the right-hand side become the chosen inclusion maps on the left-hand side. \square

Example 7.18. When pondering Corollary 7.17, it is important to realize that the space $\text{Hom}(X, Y)$ is typically much larger than $V(X, Y)$. Suppose that X is endopositive and $Y \cong \mathbf{q}X \oplus \mathbf{q}^{-1}X$. Then, $\text{Hom}^{-1}(X, Y)$ is free over \mathbb{K} of rank 1 spanned by ι_{-1} and is isomorphic to $V^{-1}(X, Y)$. However, $\text{Hom}^1(X, Y)$ is spanned by ι_1 together with $\iota_{-1} \circ f$ for $f \in \text{End}^2(X)$. Since X is endopositive, any term of the form $\iota_{-1} \circ f$ lies in $\text{Rker}(\beta_{X, Y}^1)$. Thus, as guaranteed by Corollary 7.17, we see that $V^1(X, Y)$ is free of rank 1, spanned by the image of ι_1 .

This observation will be significant later when we consider equivariance: an automorphism of \mathcal{A} will induce an automorphism of $V^k(X, Y)$, but need not preserve the span elements in $\text{Hom}^k(X, Y)$ that descend to a basis. There are a number of tools for studying the action of an automorphism on multiplicity spaces. The first is the following, which allows us to work directly with $V(X, Y)$ by identify elements in the left and right kernels.

Lemma 7.19. If X is endopositive, then

$$\text{Rker}(X, -) := \bigoplus_k \text{Rker}(\beta_{X, -}^k) \quad \text{and} \quad \text{Lker}(-, X) := \bigoplus_k \text{Lker}(\beta_{X, -}^k)$$

are left and right ideals in \mathcal{A} , respectively. Further, any positive degree endomorphism in $\text{End}(X)$ is in $\text{Rker}(X, -) \cap \text{Lker}(-, X)$.

Proof. The statement about ideals is true more generally for any (endopositive) object X . To see that $\text{Rker}(X, -)$ is a left ideal, we must show that given $g \in \text{Rker}(X, Y) \subset \text{Hom}(X, Y)$ and any morphism $f: Y \rightarrow Z$, the composition $f \circ g$ is in $\text{Rker}(X, Z)$. Clearly, it suffices to consider homogenous g and f . Thus, the claim is that if $g \in \text{Rker}(\beta_{X, Y}^k)$ and $f \in \text{Hom}^\ell(Y, Z)$, then $f \circ g \in \text{Rker}(\beta_{X, Z}^{k+\ell})$, which is immediate from the definition of $\text{Rker}(\beta_{X, Y}^k)$. The argument for $\text{Lker}(-, X)$ is analogous. Finally, if $k > 0$ then $\text{Hom}^{-k}(X, X)$ is zero, which implies the final result. \square

If $(\mathcal{A}, \otimes, \mathbb{1})$ is a monoidal category, we can identify further elements in the kernel of $\beta_{X,Y}^k$ using the actions of the endomorphism algebra $\text{End}(\mathbb{1})$ on homomorphism spaces in \mathcal{A} via tensor product on the left and right.

Lemma 7.20. Let X be an endopositive object in a monoidal category \mathcal{A} . For $\ell > 0$ and any object Y , the left or right action by $\text{End}^\ell(\mathbb{1})$ sends $\text{Hom}^{k-\ell}(X, Y)$ to the right kernel of $\beta_{X,Y}^k$.

Proof. This follows immediately from Lemma 7.19, since

$$\text{id}_X \cdot \text{End}^\ell(\mathbb{1}) \subset \text{End}^\ell(X) \supset \text{End}^\ell(\mathbb{1}) \cdot \text{id}_X. \quad \square$$

Finally, we record a method for working with $V(X, Y)$ indirectly.

Remark 7.21. Let X be endopositive. Suppose we are given maps $\{\iota_1, \dots, \iota_d\} \subset \text{Hom}^k(X, Y)$ which are “candidate inclusions” and maps $\{p_1, \dots, p_d\} \subset \text{Hom}^{-k}(Y, X)$ which are “candidate projections.” If we compute the pairing matrix $\mathbf{L}_{X,Y}^k(p_j, \iota_\ell)$ and the determinant is invertible in \mathbb{K} , then these maps descend to linearly independent sets in $V^k(X, Y)$ and $V^{-k}(Y, X)$, respectively. If we know somehow that $\{\iota_1, \dots, \iota_d\}$ descend to a spanning set of $V^k(X, Y)$ (e.g. using Corollary 7.17, or because we work over a field and $d = \dim V^k(X, Y)$), then we obtain bases of $V^k(X, Y)$ and $V^{-k}(Y, X)$. In this case, an arbitrary morphism $f \in \text{Hom}^k(X, Y)$ need not be in the span of $\{\iota_1, \dots, \iota_d\}$, but its image in $V^k(X, Y)$ is in (the image of) this span. By non-degeneracy, $f = \sum_{j=1}^d a_j \iota_j$ in $V^k(X, Y)$ if and only if they have the same pairing against all p_j . Thus, one can compute the image of f in $V^k(X, Y)$ by computing all the pairings $\beta^k(p_j, f)$ and using elementary linear algebra.

7.2. Endopositive families of objects. We now focus on categories with distinguished collections of endopositive objects.

Definition 7.22. Let \mathbb{K} be a commutative ring and let \mathbb{B} be a set indexing a family of distinguished objects $\{X_b\}_{b \in \mathbb{B}}$ in a graded additive \mathbb{K} -linear category \mathcal{A} . We call $\{X_b\}_{b \in \mathbb{B}}$ an *endopositive family* of objects provided each X_b is endopositive and $\text{Hom}(X_b, X_{b'})$ is concentrated in strictly positive degrees when $b \neq b'$. We let $\mathcal{A}(\mathbb{B})$ denote the full subcategory of \mathcal{A} whose objects are isomorphic to direct sums of shifts of objects in the corresponding endopositive family.

Given an endopositive family, if $b \neq b'$, then X_b is not isomorphic to any grading shift of $X_{b'}$, as such an isomorphism would require a morphism of non-positive degree (either $X_b \rightarrow X_{b'}$ or $X_{b'} \rightarrow X_b$). Similarly, X_b is not isomorphic to a nonzero grading shift of itself. Note also that the condition of being an endopositive family is preserved by base change.

We informally summarize the condition that $\{X_b\}$ is an endopositive family with the equation:

$$(7.4) \quad \text{grdim}_{\mathbb{K}} \text{Hom}(X_b, X_{b'}) \in \delta_{b,b'} + q\mathbb{Z}_{\geq 0}[[q]].$$

This uses several abuses of notation! We have not assumed that $\text{Hom}(X_b, X_{b'})$ is a free module over \mathbb{K} , except in degree 0 when $b = b'$. Also, \mathbb{K} is not necessarily a field, yet for simplicity we choose to write $\text{grdim}_{\mathbb{K}}$ to denote the graded rank of a \mathbb{K} -module. Nonetheless, (7.4) encapsulates the idea of an endopositive family, and is mathematically accurate whenever \mathbb{K} is a field.

Example 7.23. For any commutative ring \mathbb{K} , the monoidal category $\mathbb{1}_{\mathbf{n}} \tilde{\mathcal{F}}_q(\mathfrak{gl}_2) \mathbb{1}_{\mathbf{n}}$ has an endopositive family $\{\mathcal{F}^{(k)} \mathcal{E}^{(k)} \mathbb{1}_{\mathbf{n}}\}_{k \geq 0}$. Note that $\mathcal{E}^{(k)} \mathcal{F}^{(k)} \mathbb{1}_{\mathbf{n}} \cong \mathcal{F}^{(k)} \mathcal{E}^{(k)} \mathbb{1}_{\mathbf{n}}$, by (5.34) and (5.35). See [35, Proposition 5.15] for a description of a (positively graded) basis for morphism spaces (the summary in [21, Example 2.21] may also be helpful). In this example, every object in $\mathbb{1}_{\mathbf{n}} \tilde{\mathcal{F}}_q(\mathfrak{gl}_2) \mathbb{1}_{\mathbf{n}}$ is a direct sum of shifts of distinguished objects.

Remark 7.24. Let \mathbb{K} be a field. As in Remark 7.11, the endopositive objects X_b are indecomposable with graded local endomorphism rings. Moreover, since every object in $\mathcal{A}(\mathbb{B})$ admits a decomposition

into objects with graded local endomorphism rings, one concludes that the category $\mathcal{A}(\mathbb{B})$ is Karoubian and graded Krull–Schmidt, see e.g. [22, Theorem 11.50]. The set \mathbb{B} therefore indexes the indecomposable objects of $\mathcal{A}(\mathbb{B})$, up to isomorphism and grading shift.

When \mathbb{K} is a field, Corollary 7.10 and Lemma 7.14 show that if $\{X_b\}_{b \in \mathbb{B}}$ is an endopositive family and $Y \in \mathcal{A}(\mathbb{B})$, then $Y \cong \bigoplus_{\substack{k \in \mathbb{Z} \\ b \in \mathbb{B}}} \mathbf{q}^k X_b^{\oplus d_{b,k}}$ for $d_{b,k} = \dim_{\mathbb{K}} V^k(X_b, Y)$. More generally, we have the following.

Lemma 7.25. Let $\{X_b\}_{b \in \mathbb{B}}$ be an endopositive family in \mathcal{A} , over any commutative ring \mathbb{K} , and let Y be an object of $\mathcal{A}(\mathbb{B})$. Fix any decomposition

$$Y \cong \bigoplus_{b \in \mathbb{B}} \bigoplus_{j=1}^{d_b} \mathbf{q}^{k_{b,j}} X_b,$$

i.e. fix inclusion maps $\{p_{b,j}\}_{j=1}^{d_b}$ and $\{\iota_{b,j}\}_{j=1}^{d_b}$ for the decomposition above. Then, for each $b \in \mathbb{B}$, the space $V(Y, X_b)$ is free over \mathbb{K} with graded dimension $\sum_{j=1}^{d_b} q^{k_{b,j}}$ and the elements $\{p_{b,j}\}$ descend to a basis. Similarly, $V(X_b, Y)$ is free with graded dimension $\sum_{j=1}^{d_b} q^{-k_{b,j}}$ and basis $\{\iota_{b,j}\}$.

Proof. This follows from Lemma 7.15, similar to the proof of Corollary 7.17, after noting that $V(X_b, X_{b'})$ is free over \mathbb{K} of graded rank $\delta_{b,b'}$, spanned by the identity map of X_b . \square

Finally, we record a supplement to Lemma 7.19 in the presence of an endopositive family. When paired with that result, this will guarantee that, when examining the graded composition pairing at X_b , any morphism which factors through $X_{b'}$ for $b' \neq b$ is automatically in the kernel.

Lemma 7.26. Let $\{X_b\}_{b \in \mathbb{B}}$ be an endopositive family. If $b \neq b' \in \mathbb{B}$, then any morphism in $\text{Hom}(X_b, X_{b'})$ is in $\text{Rker}(X_b, -) \cap \text{Lker}(-, X_{b'})$.

Proof. If $b \neq b'$, (7.4) implies that for each $k \in \mathbb{Z}$, at least one of $\text{Hom}^{-k}(X_{b'}, X_b)$ or $\text{Hom}^k(X_b, X_{b'})$ is identically zero. \square

7.3. Mixed categories. In the literature, endopositive families are typically studied in the context of mixed categories, which also possess a duality functor.

Definition 7.27 (c.f. [20, Definition 3.9]). Let \mathbb{K} be a commutative ring, and \mathcal{A} be a graded additive \mathbb{K} -linear category. We call \mathcal{A} a *positively graded category* if

- all morphism spaces in \mathcal{A} are free \mathbb{Z} -graded \mathbb{K} -modules of finite rank in each degree, and
- \mathcal{A} is equipped with an endopositive family $\{X_b\}_{b \in \mathbb{B}}$ such that $\mathcal{A} = \mathcal{A}(\mathbb{B})$.

Suppose further that \mathcal{A} is equipped with a \mathbb{K} -linear functor $\mathbb{D}: \mathcal{A} \rightarrow \mathcal{A}^{\text{op}}$ such that $\mathbb{D}^2 = \mathbb{1}_{\mathcal{A}}$ and $\mathbb{D}(\mathbf{q}^k X) = \mathbf{q}^{-k} \mathbb{D}(X)$. Then, we call such a (positively graded) \mathcal{A} *mixed* provided for each $b \in \mathbb{B}$, there is some $b^* \in \mathbb{B}$ such that $\mathbb{D}(X_b) \cong X_{b^*}$. If $b = b^*$ for all $b \in \mathbb{B}$, then we say \mathcal{A} is *self-dual mixed*.

As usual, we extend these notions to 2-categories via their Hom-categories.

Note that Remark 7.24 implies that a positively graded category over a field \mathbb{K} is Karoubian and graded Krull–Schmidt, and \mathbb{B} indexes the indecomposable objects up to isomorphism and grading shift.

Example 7.28. When \mathbb{K} is a field of characteristic zero, the 2-category $\text{Kar}(\mathcal{U}_q(\mathfrak{g}_m))$ is self-dual mixed, by [71].

Remark 7.29. With apologies to Webster, the definition of mixed in Definition 7.27 (which is adapted from [20, Definition 3.9]) differs from the definition of mixed in [71, Definitions 1.2 and 1.11]. The primary feature of both definitions is (7.4). Webster’s definition assumes the category is self-dual. When \mathbb{K} is a field, our definition coincides in the self-dual case with the special case of Webster’s where the orthodox and canonical bases coincide; see [71, Corollary 1.14].

We briefly note several advantages of a self-dual mixed category. The first is that it gives a reason to prefer X_b over $\mathbf{q}^k X_b$, as only when $k = 0$ is $\mathbf{q}^k X_b$ self-dual. The second is that any automorphism of the category which commutes with shifts and duality must preserve the self-dual distinguished objects X_b . The third (which we do not use in this paper) is that if X_b and Y are both self-dual objects, then one can identify $\text{Hom}(X_b, Y)$ with $\text{Hom}(Y, X_b)$ using \mathbb{D} . This transforms the composition pairing into a bilinear form on $\text{Hom}(X_b, Y)$.

7.4. Equivariant categories. In the theory of quantum groups and Hecke algebras, there are certain algebras with fixed bases such that the structure constants for multiplication have the form $P - Q$ for $P, Q \in \mathbb{Z}_{\geq 0}[q^{\pm}]$. As an immediately relevant example, recall (1.1) which stated that

$$(7.5) \quad P_{\text{st}_4} \left(\bigcirc^{\Lambda^2} \right) = [5] + [1] \quad \text{and} \quad P_{\text{so}_5} \left(\bigcirc^S \right) = [5] - [1].$$

The latter value appears as a structure constant in $\text{End}_{U_q(\mathfrak{so}_5)}(S \otimes S)$, when computing $\mathbf{X}^{(2)} \cdot \mathbf{X}^{(2)}$, so centralizer algebras for type B quantum groups are examples of such algebras.

Lusztig intuited that such structure constants may be realized by the trace of an involution acting on a graded vector space, making P (resp. Q) the graded dimension of the $+1$ (resp. -1) eigenspace. See e.g. [43, Remark 14.4.14]. We now briefly summarize the general exposition in [20] concerning how such algebras may be realized as the weighted Grothendieck rings of certain G -equivariant categories. We then give precise definitions only for the group $G = \mathbb{Z}/2$, since that is the case that concerns us in the present work. Throughout this section, \mathbb{K} can be an arbitrary commutative ring, although some of the definitions will be trivial when \mathbb{K} has characteristic two.

Let \mathcal{A} be a (graded) additive \mathbb{K} -linear category with a strict action of a group G . We can then form the G -equivariantization of \mathcal{A} , denoted \mathcal{A}^G . The objects of \mathcal{A}^G are pairs (X, φ) , where X is an object of \mathcal{A} such that $g \cdot X \cong X$ for all $g \in G$, and φ is a family of isomorphisms $\varphi_g: X \rightarrow g \cdot X$ that satisfy a natural compatibility constraint.

The category \mathcal{A}^G has a strict action of the dual group $G^* := \text{Hom}(G, \mathbb{K}^\times)$. The action of a homomorphism $\xi: G \rightarrow \mathbb{K}^\times$ will rescale φ_g by $\xi(g)$. That is,

$$\xi \cdot (X, \varphi) = (X, \xi \cdot \varphi), \quad (\xi \cdot \varphi)_g = \xi(g)\varphi_g.$$

Given an element $g \in G$, we can form the g -weighted Grothendieck group of \mathcal{A}^G , denoted $K_0^g(\mathcal{A}^G)$. This is the quotient of the usual (additive) Grothendieck group, base changed to \mathbb{K} (or any subring containing the images of all $\xi \in G^*$), by the relation $[\xi \cdot (X, \varphi)] = \xi(g)[(X, \varphi)]$ for all $\xi \in G^*$. For more details see [20, Section 3.1].

In the case that \mathcal{A} is monoidal and/or is equipped with a duality functor and the G -action preserves these structures, \mathcal{A}^G inherits these structures and the action of G^* respects them. In this way, $K_0^g(\mathcal{A}^G)$ inherits the structure of a ring and/or inherits the bar involution (the anti-linear action of duality on the Grothendieck group).

We now restrict our attention to $G = \mathbb{Z}/2$.

Definition 7.30 (c.f. [20, Definition 3.1]). A *strict action* of $\mathbb{Z}/2$ on an additive \mathbb{K} -linear category \mathcal{A} is an additive \mathbb{K} -linear autoequivalence $\sigma: \mathcal{A} \rightarrow \mathcal{A}$ and a natural isomorphism $\mathfrak{s}: \sigma \circ \sigma \xrightarrow{\cong} \mathbb{1}_{\mathcal{A}}$ such that¹⁸ $\mathfrak{s} \otimes \text{id}_\sigma = \text{id}_\sigma \otimes \mathfrak{s}$ as natural transformations $\sigma \circ \sigma \circ \sigma \rightarrow \sigma$.

Remark 7.31. From the group $\mathbb{Z}/2$ (here written multiplicatively), one obtains a monoidal category $\Omega_{\mathbb{Z}/2}$, often called the 2-groupoid of $\mathbb{Z}/2$. It has objects ± 1 , tensor product given by the group operation, and the only morphisms are identities. Equivalent to the definition above, a *strict action* of $\mathbb{Z}/2$ on \mathcal{A}

¹⁸Here, \otimes denotes horizontal composition of natural transformations; this is the tensor product on the monoidal category of endofunctors of \mathcal{A} .

is a strong¹⁹ (but not necessarily strict) monoidal functor $F: \Omega_{\mathbb{Z}/2} \rightarrow \text{End}(\mathcal{A})$ for which $F(+1) = \mathbb{1}_{\mathcal{A}}$ and $F(-1) =: \sigma$.

Remark 7.32. We call σ an *involution* if $\sigma \circ \sigma = \mathbb{1}_{\mathcal{A}}$ is an equality of functors. In this case, we can use $s = \text{id}_{\mathbb{1}_{\mathcal{A}}}$ to obtain a strict action of $\mathbb{Z}/2$ on \mathcal{A} .

From here on, we assume that σ is an involution and that $s = \text{id}_{\mathbb{1}_{\mathcal{A}}}$.

Remark 7.33. Consider an involution σ on a 2-category in the sense of Definition 6.1. If \mathbf{a} is an object fixed by σ , then σ induces a monoidal functor on the endomorphism category of \mathbf{a} . This functor will be an involution in the sense of Remark 7.32.

Definition 7.34 ([20, Definition 3.5]). Let \mathcal{A} be a category and let $\sigma: \mathcal{A} \rightarrow \mathcal{A}$ be an involution. An *equivariant object* is a pair $(X, \varphi_X: X \xrightarrow{\cong} \sigma X)$ such that $\sigma(\varphi_X) \circ \varphi_X = \text{id}_X$. A *morphism of equivariant objects* from (X, φ_X) to (Y, φ_Y) is a morphism $f \in \text{Hom}(X, Y)$ such that $\varphi_Y \circ f = \sigma(f) \circ \varphi_X$. Equivariant objects and morphisms of equivariant objects form the *equivariant category* \mathcal{A}^σ .

Applying σ to the equation $\sigma(\varphi_X) \circ \varphi_X = \text{id}_X$, we see that $\varphi_X \circ \sigma(\varphi_X) = \text{id}_{\sigma(X)}$. Thus, the condition determining an equivariant object can be rewritten simply as $\sigma(\varphi_X) = \varphi_X^{-1}$. We will refer to the choice of isomorphism $\varphi_X: X \rightarrow \sigma(X)$ satisfying this condition as an *equivariant structure* on X . We say that X is *equivariantizable* if some equivariant structure exists.

We now establish a perspective on the Hom-spaces in \mathcal{A}^σ that will prove useful.

Proposition 7.35. Let σ be an involution of a \mathbb{K} -linear category \mathcal{A} and let (X, φ_X) and (Y, φ_Y) be objects in \mathcal{A}^σ . For $f \in \text{Hom}(X, Y)$, the formula

$$(7.6) \quad \sigma * f = \sigma(\varphi_Y) \circ \sigma(f) \circ \varphi_X$$

determines a \mathbb{K} -linear action of the group $\langle \sigma * \rangle \cong \mathbb{Z}/2$ on $\text{Hom}(X, Y)$. This action respects composition, i.e. if further (Z, φ_Z) is an object in \mathcal{A}^σ and $g \in \text{Hom}(Y, Z)$, then

$$(7.7) \quad \sigma * (g \circ f) = (\sigma * g) \circ (\sigma * f).$$

Proof. We have that $\sigma(\varphi_X) = \varphi_X^{-1}$ and $\sigma(\varphi_Y) = \varphi_Y^{-1}$. We compute

$$\begin{aligned} \sigma * (\sigma * f) &= \sigma * (\sigma(\varphi_Y) \circ \sigma(f) \circ \varphi_X) = \sigma(\varphi_Y) \circ \sigma(\sigma(\varphi_Y) \circ \sigma(f) \circ \varphi_X) \circ \varphi_X \\ &= \varphi_Y^{-1} \circ \varphi_Y \circ f \circ \varphi_X^{-1} \circ \varphi_X = f, \end{aligned}$$

which shows that $\sigma *$ generates a $\mathbb{Z}/2$ -action. Since composition and σ are \mathbb{K} -linear, $\sigma * f$ is linear in f . We leave the reader to check compatibility with composition. \square

We **strongly emphasize** that the definition of $\sigma *$ on $\text{Hom}(X, Y)$ depends on the choice of isomorphisms φ_X and φ_Y , i.e. on the choice of equivariant objects (X, φ_X) and (Y, φ_Y) .

Corollary 7.36. Let (X, φ_X) and (Y, φ_Y) be equivariant objects and let $\text{Hom}(X, Y)^\sigma$ denote the $(\sigma *)$ -invariants in $\text{Hom}(X, Y)$. Then, $\text{Hom}_{\mathcal{A}^\sigma}((X, \varphi_X), (Y, \varphi_Y)) = \text{Hom}(X, Y)^\sigma$.

Proof. Since $\sigma(\varphi_Y) = \varphi_Y^{-1}$, we see that $f = \sigma * f$ if and only if $\varphi_Y \circ f = \sigma(f) \circ \varphi_X$. \square

Remark 7.37. Corollary 7.36 immediately shows that if (X, φ_X) is decomposable in \mathcal{A}^σ , then $X \in \mathcal{A}$ is decomposable. Indeed, any idempotents in $\text{End}_{\mathcal{A}^\sigma}((X, \varphi)) = \text{End}(X)^\sigma$ giving a direct sum decomposition in \mathcal{A}^σ also give such a decomposition in \mathcal{A} . The converse of this statement need not be true; see Proposition 7.46 below. Note also that if X is endopositive, then Corollary 7.36 implies that (X, φ_X) is endopositive.

¹⁹A monoidal functor always comes with data of coherence maps; a strong monoidal functor refers to when these maps are isomorphisms and a strict monoidal functor refers to when these isomorphisms are equalities.

Next, we review the weighted notion of Grothendieck group that can be applied to the category \mathcal{A}^σ . If (X, φ_X) is an object in \mathcal{A}^σ , then $\text{id}_X = \sigma(\varphi_X) \circ \varphi_X = \sigma(-\varphi_X) \circ (-\varphi_X)$. Thus, $(X, -\varphi_X)$ is an object of \mathcal{A}^σ as well. For $f \in \text{Hom}(X, Y)$, it is easy to verify that

$$f \in \text{Hom}_{\mathcal{A}^\sigma}((X, \varphi_X), (Y, \varphi_Y)) \iff f \in \text{Hom}_{\mathcal{A}^\sigma}((X, -\varphi_X), (Y, -\varphi_Y)).$$

This motivates the following.

Definition 7.38 ([20, Definition 3.6]). Let $\text{sgn}: \mathcal{A}^\sigma \rightarrow \mathcal{A}^\sigma$ be the involutive functor given on objects by $\text{sgn}((X, \varphi_X)) = (X, -\varphi_X)$ and on morphisms by $\text{sgn}(f) = f$.

In the context of the discussion at the start of this section, one should view sgn as the generator of the dual group $(\mathbb{Z}/2)^*$. We now introduce the weighted Grothendieck group of \mathcal{A}^σ as a quotient of the usual additive Grothendieck group $K_0(\mathcal{A}^\sigma)$. Classes in the latter are denoted $[(X, \varphi_X)]$.

Definition 7.39 ([20, Definition 3.7]). The σ -weighted Grothendieck group of \mathcal{A}^σ , denoted $K_0^\sigma(\mathcal{A}^\sigma)$, is the quotient of the Grothendieck group $K_0(\mathcal{A}^\sigma) \otimes_{\mathbb{Z}} \mathbb{Z}[\frac{1}{2}]$ by the relation

$$[(X, -\varphi_X)] \equiv -[(X, \varphi_X)].$$

For an object $(X, \varphi_X) \in \mathcal{A}^\sigma$, we continue to write $[(X, \varphi_X)]$ for its class in the ordinary Grothendieck group $K_0(\mathcal{A}^\sigma)$ and denote by $[(X, \varphi_X)]_\sigma$ the image of this class in $K_0^\sigma(\mathcal{A}^\sigma)$.

Remark 7.40. We change the base of the weighted Grothendieck group from \mathbb{Z} to $\mathbb{Z}[\frac{1}{2}]$ to eliminate 2-torsion; as a consequence, if $(X, \varphi_X) \cong (X, -\varphi_X)$ in \mathcal{A}^σ , then $[(X, \varphi_X)]_\sigma = 0$.

Finally, we precisely record properties of \mathcal{A} inherited by \mathcal{A}^σ under the assumption that σ is compatible with them. The proof is straightforward.

Proposition 7.41. Let \mathcal{A} be a \mathbb{K} -linear additive category with an involution σ .

- If \mathcal{A} is monoidal and σ is a monoidal functor, then $(X, \varphi_X) \otimes (Y, \varphi_Y) := (X \otimes Y, \varphi_X \otimes \varphi_Y)$ defines a monoidal structure on \mathcal{A}^σ . Tensor product of morphisms is given as in \mathcal{A} .
- If \mathcal{A} has a contravariant duality functor \mathbb{D} as in Definition 7.27 such that $\mathbb{D} \circ \sigma = \sigma \circ \mathbb{D}$, then $\mathbb{D}(X, \varphi) := (\mathbb{D}(X), \mathbb{D}(\sigma(\varphi)))$ defines a contravariant duality on \mathcal{A}^σ . The duality is given on morphisms as in \mathcal{A} . \square

7.5. Indecomposable objects in equivariant categories. Now, assume that \mathbb{K} is an integral domain. We aim to characterize the indecomposable objects in \mathcal{A}^σ . Observe that if $X \in \mathcal{A}$ is an equivariantizable object, then $X \cong \sigma(X)$. However, we warn the reader that, in general, the converse does not hold without additional assumptions. In particular, the theory greatly simplifies under Hypothesis 7.44 below, and simplifies further under the assumption that \mathbb{K} is an algebraically closed field of characteristic not equal to 2. The case when \mathbb{K} is not algebraically closed is discussed in Appendix C.

Notation 7.42. Let \mathbb{K} be an integral domain where 2 is invertible and let \mathcal{A} be a graded additive \mathbb{K} -linear category equipped with an involution σ which commutes with the shift autoequivalence \mathbf{q} . Let $\{X_b\}_{b \in \mathbb{B}}$ be an endopositive family which is preserved by σ , up to isomorphism. Define an action of σ on \mathbb{B} by $\sigma(X_b) \cong X_{\sigma(b)}$. Since $\sigma(X_{\sigma(b)}) \cong X_b$, there is a decomposition

$$\mathbb{B} = \mathbb{B}^{\text{fix}} \sqcup \mathbb{B}^{\text{free}}$$

where $\sigma(b) = b$ for all $b \in \mathbb{B}^{\text{fix}}$ and $\sigma(b) \neq b$ for all $b \in \mathbb{B}^{\text{free}}$.

In other words, \mathbb{B}^{fix} is the set of fixed points in \mathbb{B} under the σ -action and \mathbb{B}^{free} is the union of all orbits of size two.

Remark 7.43. If \mathcal{A} is self-dual mixed and σ commutes with \mathbb{D} , then σ preserves the set of self-dual indecomposable objects. Hence, it automatically preserves $\{X_b\}_{b \in \mathbb{B}}$, up to isomorphism.

For the remainder of this section, **we assume the following hypothesis.**

Hypothesis 7.44. Assume we are in the situation of Notation 7.42 and that X_b is equivariantizable for each $b \in \mathbb{B}^{\text{fix}}$.

When \mathbb{K} is an algebraically closed field with $2 \neq 0$, the second part of this hypothesis is superfluous, since it is straightforward to show that $X_b \cong \sigma(X_b)$ implies that X_b is equivariantizable. The results cited below from [20] are sketched there under the assumption that \mathbb{K} is an algebraically closed field where $2 \neq 0$. We prove them more generally under Hypothesis 7.44 in Appendix C.

Lemma 7.45 ([20, Proposition 3.10]). For all $b \in \mathbb{B}^{\text{fix}}$, there is an isomorphism $\varphi_b: X_b \xrightarrow{\cong} \sigma(X_b)$ such that $\sigma(\varphi_b) \circ \varphi_b = \text{id}_{X_b}$. For all orbits $\{b, \sigma(b)\} \subset \mathbb{B}^{\text{free}}$, there is an isomorphism

$$\psi_b: X_b \oplus X_{\sigma(b)} \xrightarrow{\cong} \sigma(X_b \oplus X_{\sigma(b)})$$

such that $\sigma(\psi_b) \circ \psi_b = \text{id}$. As equivariant objects, $(X_b, \varphi_b) \not\cong (X_b, -\varphi_b)$ for $b \in \mathbb{B}^{\text{fix}}$ while

$$(7.8) \quad (X_b \oplus X_{\sigma(b)}, \psi_b) \cong (X_b \oplus X_{\sigma(b)}, -\psi_b)$$

for $b \in \mathbb{B}^{\text{free}}$. □

Proposition 7.46 ([20, Proposition 3.10]). Fix isomorphisms φ_b as in Lemma 7.45 for all $b \in \mathbb{B}^{\text{fix}}$. If (X, φ_X) is an indecomposable object in \mathcal{A}^σ for which $X \in \mathcal{A}(\mathbb{B})$, then exactly one of the following holds.

- $(X, \varphi_X) \cong \mathbf{q}^k(X_b, \varphi_b)$ for some $b \in \mathbb{B}^{\text{fix}}$ and $k \in \mathbb{Z}$,
- $(X, \varphi_X) \cong \mathbf{q}^k(X_b, -\varphi_b)$ for some $b \in \mathbb{B}^{\text{fix}}$ and $k \in \mathbb{Z}$, or
- $(X, \varphi_X) \cong \mathbf{q}^k(X_b \oplus X_{\sigma(b)}, \psi_b) \cong \mathbf{q}^k(X_b \oplus X_{\sigma(b)}, -\psi_b)$ for some $b \in \mathbb{B}^{\text{free}}$ and $k \in \mathbb{Z}$. □

It is relatively straightforward to compute the morphism spaces between the self-dual indecomposable equivariant objects $(X_b, \pm\varphi_b)$ and $(X_b \oplus X_{\sigma(b)}, \psi_b)$.

Corollary 7.47. In \mathcal{A}^σ , the objects $\{(X_b, \varphi_b), (X_b, -\varphi_b)\}_{b \in \mathbb{B}^{\text{fix}}} \cup \{(X_b \oplus X_{\sigma(b)}, \psi_b)\}_{b \in \mathbb{B}^{\text{free}}}$ form an endopositive family. If \mathcal{A} is self-dual mixed, then \mathcal{A}^σ is self-dual mixed.

Remark 7.48. For general G , the equivariant category of a self-dual mixed category is mixed, but not necessarily self-dual. The equivariant category is self-dual mixed if and only if all representations of G are self dual, as is the case for $G = \mathbb{Z}/2$.

Notation 7.49. Using a slight abuse of notation, the full subcategory of \mathcal{A}^σ whose objects are direct sums of shifts of the endopositive family in Corollary 7.47 will be denoted $\mathcal{A}^\sigma(\mathbb{B})$.

Next, we discuss an effective algorithm to take an equivariant object (Y, φ_Y) in \mathcal{A}^σ (and not necessarily in $\mathcal{A}^\sigma(\mathbb{B})$) and find all indecomposable summands of the form in Corollary 7.47. Recall that if X and Y are equipped with equivariant structures, then $\text{Hom}(X, Y)$ admits an action of σ_* which depends on the choice of those structures. When $X = X_b$ for $b \in \mathbb{B}^{\text{fix}}$, we always assume that the choice of equivariant structure for this action is φ_b (rather than $-\varphi_b$), and we denote the corresponding action as σ_{b*} .

We leave it to the reader to verify that σ_{b*} preserves the kernels $\text{Rker}(\beta_{X_b, Y}^k)$ and $\text{Lker}(\beta_{X_b, Y}^k)$ of the graded composition pairing. Therefore, σ_{b*} descends to an action on $V^k(X_b, Y)$ and $V^{-k}(Y, X_b)$ for all $k \in \mathbb{Z}$. It is not hard to show that any element of the $+1$ -eigenspace of σ_{b*} is indeed a valid inclusion map making (X_b, φ_b) a direct summand of (Y, φ_Y) , while any element of the -1 -eigenspace gives a direct summand of the form $(X_b, -\varphi_b)$.

Proposition 7.50 ([20, Claim 3.16]). Let (Y, φ_Y) be an object in \mathcal{A}^σ .

- (1) For $b \in \mathbb{B}^{\text{free}}$, the multiplicity of $\mathbf{q}^k(X_b \oplus X_{\sigma(b)}, \psi_b)$ in (Y, φ_Y) is equal to the multiplicity of $\mathbf{q}^k X_b$ in Y .

- (2) For $b \in \mathbb{B}^{\text{fix}}$, the multiplicity of $\mathbf{q}^k(X_b, \varphi_b)$ in (Y, φ_Y) equals the dimension of the $+1$ -eigenspace of σ_{b*} in $V^k(X_b, Y)$.
- (3) For $b \in \mathbb{B}^{\text{fix}}$, the multiplicity of $\mathbf{q}^k(X_b, -\varphi_b)$ in (Y, φ_Y) equals the dimension of the -1 -eigenspace of σ_{b*} in $V^k(X_b, Y)$.

The above results allow for the computation of the weighted Grothendieck group of $\mathcal{A}^\sigma(\mathbb{B})$. Observe that (7.8) implies that $[(X_b \oplus X_{\sigma(b)}, \psi_b)]_\sigma = 0$ whenever $b \in \mathbb{B}^{\text{free}}$, since in this case

$$[(X_b \oplus X_{\sigma(b)}, \psi_b)]_\sigma = [(X_b \oplus X_{\sigma(b)}, -\psi_b)]_\sigma = -[(X_b \oplus X_{\sigma(b)}, \psi_b)]_\sigma.$$

Proposition 7.51 ([20, Prop. 3.17 and Cor. 3.19]). The σ -weighted Grothendieck group $K_0^\sigma(\mathcal{A}^\sigma(\mathbb{B}))$ has a basis

$$\{[(X_b, \varphi_b)]_\sigma\}_{b \in \mathbb{B}^{\text{fix}}}$$

in bijection with \mathbb{B}^{fix} . Moreover, if (Y, φ_Y) is any object in $\mathcal{A}^\sigma(\mathbb{B})$, then

$$(7.9) \quad [(Y, \varphi_Y)]_\sigma = \sum_{k \in \mathbb{Z}} \sum_{b \in \mathbb{B}^{\text{fix}}} \text{Tr}(\sigma_{b*}|_{V^k(X_b, Y)}) q^k [(X_b, \varphi_b)]_\sigma.$$

□

In light of (7.9), we conclude this section with remarks on the practicalities of computing the trace of σ_{b*} . It is often easy to compute σ_{b*} on $\text{Hom}^k(X_b, Y)$, but computing the action on $V^k(X_b, Y)$ can be difficult. In [20], the action of σ_{b*} on $V^k(X_b, Y)$ could be computed directly, since one could find a subset $\{\iota_1, \dots, \iota_d\} \subset \text{Hom}^k(X_b, Y)$ that is permuted by σ_{b*} and which descends to a basis of $V^k(X_b, Y)$. See e.g. [20, Sections 5.1, 5.4].

In our setting, finding such a “basis of inclusion maps” which is permuted by σ_{b*} is significantly more difficult. In some instances, we are able to overcome this difficulty by computing directly, e.g. the computations in §8.2. In others, e.g. in §8.3, we instead compute the action of σ_{b*} on $V^k(X_b, Y)$ using the techniques of Remark 7.21. The following result records this method explicitly.

Lemma 7.52. Suppose that $V^k(X_b, Y)$ is free over \mathbb{K} of rank d and choose $\{\iota_1, \dots, \iota_d\} \subset \text{Hom}^k(X_b, Y)$ and $\{p_1, \dots, p_d\} \subset \text{Hom}^{-k}(Y, X_b)$ which descend to bases of $V^k(X_b, Y)$ and $V^{-k}(Y, X_b)$ respectively. Assume also that the pairing matrix C between these bases is invertible over \mathbb{K} . If S is the pairing matrix between $\{\sigma_{b*}\iota_1, \dots, \sigma_{b*}\iota_d\}$ and $\{p_1, \dots, p_d\}$, then $C^{-1}S$ is the matrix encoding the action of σ_{b*} on $V^k(X_b, Y)$, in the basis induced by $\{\iota_1, \dots, \iota_d\}$.

Proof. Let $x \mapsto \bar{x}$ denote the quotient map $\text{Hom}^k(X_b, Y) \twoheadrightarrow V^k(X_b, Y)$, so $\{\bar{\iota}_1, \dots, \bar{\iota}_d\}$ is a basis for $V^k(X_b, Y)$. The action of σ_{b*} on $V^k(X_b, Y)$ is determined by the matrix Σ which describes how to write $\sigma_{b*}\bar{\iota}_r$ in terms of the basis $\{\bar{\iota}_1, \dots, \bar{\iota}_d\}$. It is easy to see that the pairing matrix S is equal to the composition of Σ and the pairing matrix C , i.e. $S = C \cdot \Sigma$. Thus, $\Sigma = C^{-1}S$. □

8. AN EQUIVARIANT CATEGORY FROM THE CATEGORIFIED QUANTUM GROUP

In this section we assume that \mathbb{K} is an integral domain and 2 is invertible in \mathbb{K} .

8.1. The $\mathcal{FE}\mathbb{1}_n$ equivariant category. We assume now, and for the duration, that we have chosen the functions in Theorem 6.16 and fixed $n \geq 1$. We thus have the involution $\tau: \tilde{\mathcal{FE}}_q(\mathfrak{gl}_m) \rightarrow \tilde{\mathcal{FE}}_q(\mathfrak{gl}_m)$ from Corollary 6.25. Since τ preserves the object $\mathbb{1}_n$, it preserves the endomorphism category of $\mathbb{1}_n$, which is the following monoidal category.

Definition 8.1. The monoidal category \mathcal{B}_m is $\mathbb{1}_n \tilde{\mathcal{FE}}_q(\mathfrak{gl}_m) \mathbb{1}_n$. Equivalently, it is the full 2-subcategory of $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$ generated by the 1-morphisms $\mathcal{E}_i^{(k)} \mathcal{F}_i^{(k)} \mathbb{1}_n$ and $\mathcal{F}_i^{(k)} \mathcal{E}_i^{(k)} \mathbb{1}_n$ for $i \in \{1, \dots, m-1\}$ and $k \in \mathbb{Z}_{\geq 0}$.

Note that both τ and \mathcal{B}_m depend on n , though, in light of the results from §6.4, the dependence is not significant. Hence, the absence of this dependence in our notation. We now aim to apply the techniques of §7.5 to study the corresponding equivariant category $(\mathcal{B}_m)^\tau$. Since the involution τ from Corollary 6.25 gives a monoidal involution of \mathcal{B}_m , Proposition 7.41 shows that $(\mathcal{B}_m)^\tau$ is monoidal.

Remark 8.2. Example 7.28 gives that $\text{Kar}(\mathcal{U}_q(\mathfrak{gl}_m))$ is self-dual mixed when \mathbb{K} is a field of characteristic zero, thus the same is true for the full 2-subcategory $\text{Kar}(\mathcal{B}_m)$. By Corollary 7.47, its τ -equivariant category is also self-dual mixed.

We begin by exploring equivariant objects. Note that

$$(8.1) \quad \begin{array}{c} \uparrow \downarrow \mathbf{n} \\ \downarrow \uparrow \mathbf{n} \\ \downarrow \downarrow \mathbf{n} \\ \uparrow \uparrow \mathbf{n} \\ \downarrow \downarrow \mathbf{n} \\ \uparrow \uparrow \mathbf{n} \end{array} = \begin{array}{c} \downarrow \downarrow \mathbf{n} \\ \uparrow \uparrow \mathbf{n} \\ \downarrow \downarrow \mathbf{n} \\ \uparrow \uparrow \mathbf{n} \\ \downarrow \downarrow \mathbf{n} \\ \uparrow \uparrow \mathbf{n} \end{array}, \quad \begin{array}{c} \downarrow \downarrow \mathbf{n} \\ \uparrow \uparrow \mathbf{n} \\ \downarrow \downarrow \mathbf{n} \\ \uparrow \uparrow \mathbf{n} \\ \downarrow \downarrow \mathbf{n} \\ \uparrow \uparrow \mathbf{n} \end{array} = \begin{array}{c} \downarrow \downarrow \mathbf{n} \\ \uparrow \uparrow \mathbf{n} \\ \downarrow \downarrow \mathbf{n} \\ \uparrow \uparrow \mathbf{n} \\ \downarrow \downarrow \mathbf{n} \\ \uparrow \uparrow \mathbf{n} \end{array}$$

which are special cases of the Stošić formulae (5.36) when $k = \ell$. Consequently

$$(8.2) \quad \mathcal{E}_i^{(k)} \mathcal{F}_i^{(k)} \mathbb{1}_{\mathbf{n}} \cong \mathcal{F}_i^{(k)} \mathcal{E}_i^{(k)} \mathbb{1}_{\mathbf{n}}.$$

Moreover, these 1-morphisms are indecomposable (see (5.34) and (5.35)).

Remark 8.3. More generally, for fixed i and over any commutative ring \mathbb{K} , the sets of objects $\{\mathcal{F}_i^{(k)} \mathcal{E}_i^{(k)} \mathbb{1}_{\mathbf{n}}\}_{k \geq 0}$ and $\{\mathcal{E}_i^{(k)} \mathcal{F}_i^{(k)} \mathbb{1}_{\mathbf{n}}\}_{k \geq 0}$ each form an endopositive family. This generalizes Example 7.23. To see this, note that the size of the morphism space e.g. between $\mathcal{F}_i^{(k)} \mathcal{E}_i^{(k)} \mathbb{1}_{\mathbf{n}}$ and $\mathcal{F}_i^{(l)} \mathcal{E}_i^{(l)} \mathbb{1}_{\mathbf{n}}$ depends on the ambient context. That they form an endopositive family within \mathcal{B}_2 was shown explicitly in [35, Proposition 5.15]. For $m > 2$, the morphism space in \mathcal{B}_m is larger than in \mathcal{B}_2 , since, at the very least, there are additional (positive degree) new bubble generators in $\text{End}_{\mathcal{B}_m}(\mathbb{1}_{\mathbf{n}}) \cong \text{Sym}(\mathbb{X}_1 | \cdots | \mathbb{X}_m)$. In fact, this is the only difference: as shown in [34, Proposition 3.11], if the color j does not appear on the boundary of a diagram, then, as a module over $\text{End}_{\mathcal{B}_m}(\mathbb{1}_{\mathbf{n}})$, there is a set of generators for the Hom-space that does not contain the color j .

Remark 8.4. Further, it is possible to describe a basis for the Hom-spaces in \mathcal{B}_m . There is a certain sesquilinear form on the quantum group $\dot{U}_q(\mathfrak{gl}_m)$ which gives an upper bound on the size of morphism spaces in $\mathcal{U}_q(\mathfrak{gl}_m)$; see [34, Corollary 3.14]. In the proof of [34, Proposition 3.11], Khovanov–Lauda give diagrammatic arguments to find a spanning set for morphism spaces. By construction, the size of this spanning set agrees with a corresponding value of the sesquilinear form [34, Proposition 3.12].

Khovanov–Lauda call the category *nondegenerate* [34, Definition 3.15] if their spanning set is actually a basis, in which case the sesquilinear form precisely controls the size of morphism spaces. We refer to this as the *Khovanov–Lauda Hom formula*. They use an action of the categorified quantum group on cohomology of partial flag varieties to prove nondegeneracy when \mathbb{K} is a field [34, Theorem 1.3].

Upon inspection, the diagrammatic spanning argument is valid over \mathbb{Z} . Since linearly independent elements in a \mathbb{Q} -vector space are necessarily linearly independent in the \mathbb{Z} -module that they generate, nondegeneracy over \mathbb{Z} (and hence over any commutative ring), follows from nondegeneracy over \mathbb{Q} .

The following objects and morphisms will be of fundamental importance moving forward.

Notation 8.5. Let $\mathbf{X}_i^{(k)} := \mathcal{F}_i^{(k)} \mathcal{E}_i^{(k)} \mathbb{1}_{\mathbf{n}}$, viewed as an object in the monoidal category \mathcal{B}_m , and denote

$$\text{LC}_k := \begin{array}{c} \downarrow \downarrow \mathbf{n} \\ \uparrow \uparrow \mathbf{n} \\ \downarrow \downarrow \mathbf{n} \\ \uparrow \uparrow \mathbf{n} \\ \downarrow \downarrow \mathbf{n} \\ \uparrow \uparrow \mathbf{n} \end{array}, \quad \text{RC}_k := \begin{array}{c} \downarrow \downarrow \mathbf{n} \\ \uparrow \uparrow \mathbf{n} \\ \downarrow \downarrow \mathbf{n} \\ \uparrow \uparrow \mathbf{n} \\ \downarrow \downarrow \mathbf{n} \\ \uparrow \uparrow \mathbf{n} \end{array}.$$

(Here, as usual, we follow Convention 5.18, i.e. LC_k and RC_k are i -colored.) We will also denote $\mathbf{X}_i^{(k)}$ by $\mathbf{X}_{\bullet}^{(k)}$ and abbreviate $\mathbf{X}_i := \mathbf{X}_i^{(1)}$.

Proposition 8.6. For each $k \geq 0$, the pair $(\mathbf{X}_{\bullet}^{(k)}, \mathbf{LC}_k)$ is an object in $(\mathcal{B}_m)^\tau$. The family $\{(\mathbf{X}_{\bullet}^{(k)}, \mathbf{LC}_k)\}_{k \geq 0}$ is endopositive.

Proof. We have $\mathbf{LC}_k \in \text{Hom}_{\mathcal{B}_m}(\mathbf{X}_{\bullet}^{(k)}, \tau(\mathbf{X}_{\bullet}^{(k)}))$ and a computation using equations (6.38), (6.39), and (6.40) shows that $\tau(\mathbf{LC}_k) = \mathbf{RC}_k$. We then compute

$$\tau(\mathbf{LC}_k) \circ \mathbf{LC}_k = \mathbf{RC}_k \circ \mathbf{LC}_k \stackrel{(8.1)}{=} \text{id}_{\mathbf{X}_{\bullet}^{(k)}}$$

so $(\mathbf{X}_{\bullet}^{(k)}, \mathbf{LC}_k)$ is an equivariant object, per Definition 7.34. The assertion that the family of all such objects is endopositive follows from Remarks 7.37 and 8.3. \square

Remark 8.7. The monoidal unit in \mathcal{B}_m is $\mathbb{1}_{\mathbf{n}}$ and the monoidal unit of $(\mathcal{B}_m)^\tau$ is the pair $(\mathbb{1}_{\mathbf{n}}, \text{id}_{\mathbb{1}_{\mathbf{n}}})$. In the notation above, this is the same as $(\mathbf{X}_{\bullet}^{(0)}, \mathbf{LC}_0)$ for any color i .

In the following sections, we study the monoidal structure of $(\mathcal{B}_m)^\tau$ and establish certain tensor product decompositions via the techniques in §7. We use the notation established there.

8.2. Divided power relation for \mathbf{X}_i . Our main goal in this subsection is to show that relation (4.28) holds for $(\mathbf{X}_{\bullet}^{(k)}, \mathbf{LC}_k)$ in the weighted Grothendieck group of $(\mathcal{B}_m)^\tau$.

Lemma 8.8. The sets

$$\left\{ \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \\ \text{Diagram 3} \\ \text{Diagram 4} \\ \text{Diagram 5} \end{array} \right\}_{\substack{0 \leq p \leq k \\ 0 \leq q \leq k-1}} \quad \text{and} \quad \left\{ \begin{array}{c} \text{Diagram 6} \\ \text{Diagram 7} \\ \text{Diagram 8} \\ \text{Diagram 9} \\ \text{Diagram 10} \end{array} \right\}_{0 \leq p, q \leq k}$$

descend to bases for $V(\mathbf{X}_{\bullet}^{(k)}, \mathbf{X}_{\bullet}^{(k)} \mathbf{X}_{\bullet})$ and $V(\mathbf{X}_{\bullet}^{(k+1)}, \mathbf{X}_{\bullet}^{(k)} \mathbf{X}_{\bullet})$, respectively.

Proof. Immediate from (5.31) and (5.36), together with Lemma 7.25. \square

We now consider the τ^* -action from Proposition 7.35 on $V(\mathbf{X}_{\bullet}^{(k)}, \mathbf{X}_{\bullet}^{(k)} \mathbf{X}_{\bullet})$ and $V(\mathbf{X}_{\bullet}^{(k+1)}, \mathbf{X}_{\bullet}^{(k)} \mathbf{X}_{\bullet})$, with respect to the isomorphisms $\mathbf{X}_{\bullet}^{(k)} \xrightarrow{\mathbf{LC}_k} \tau(\mathbf{X}_{\bullet}^{(k)})$ and $\mathbf{X}_{\bullet}^{(k)} \mathbf{X}_{\bullet} \xrightarrow{\mathbf{LC}_k \otimes \mathbf{LC}_1} \tau(\mathbf{X}_{\bullet}^{(k)} \mathbf{X}_{\bullet})$.

Proposition 8.9. Considered as elements in $V(\mathbf{X}_{\bullet}^{(k)}, \mathbf{X}_{\bullet}^{(k)} \mathbf{X}_{\bullet})$ and $V(\mathbf{X}_{\bullet}^{(k+1)}, \mathbf{X}_{\bullet}^{(k)} \mathbf{X}_{\bullet})$, we have that

$$\tau^* \left(\begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \\ \text{Diagram 3} \\ \text{Diagram 4} \\ \text{Diagram 5} \end{array} \right) = (-1)^{p+q+n+k+1} \sum_{y=0}^k (\delta_{y,q} - \delta_{y,k}) \left(\begin{array}{c} \text{Diagram 6} \\ \text{Diagram 7} \\ \text{Diagram 8} \\ \text{Diagram 9} \\ \text{Diagram 10} \end{array} \right)$$

and

$$\tau^* \left(\begin{array}{c} \text{Diagram 11} \\ \text{Diagram 12} \\ \text{Diagram 13} \\ \text{Diagram 14} \\ \text{Diagram 15} \end{array} \right) = (-1)^{p+q} \left(\begin{array}{c} \text{Diagram 16} \\ \text{Diagram 17} \\ \text{Diagram 18} \\ \text{Diagram 19} \\ \text{Diagram 20} \end{array} \right).$$

Proof. We compute, modulo terms that are in the kernel of the graded composition pairing. Each time we identify a term in the kernel, we depict it in gray and then drop it in subsequent steps.

$$\begin{aligned}
 (8.3) \quad \tau^* \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k \end{array} \right)^n &= (-1)^{p+q} \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k \end{array} \right)^n \stackrel{(5.4), (5.26)}{=} (-1)^{p+q} \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k \end{array} \right)^n - \sum_{\substack{y+z= \\ p+q-k}} \left(\begin{array}{c} \quad \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \right)^n \left(\begin{array}{c} \quad \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \right)^n \\
 &\stackrel{(8.1)}{=} (-1)^{p+q} \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k \end{array} \right)^n - \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \quad \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \right)^{p+q-k} - \sum_{y=1}^{p+q-k} \left(\begin{array}{c} \quad \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \right)^{p+q-k-y}.
 \end{aligned}$$

Here, we see that the gray terms are in the kernel of the graded composition pairing using Lemma 7.19, since they are compositions of positive degree endomorphisms of the endopositive object $\mathbf{X}_i^{(k)}$ with dotted cup maps (tensor the identity).

We next simplify the following subdiagram of the first term:

$$\begin{aligned}
 (8.4) \quad \left(\begin{array}{c} \quad \\ \downarrow \quad \downarrow \\ \quad \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \right)^{n+1, n-1} &\stackrel{(5.36a)}{=} \left(\begin{array}{c} \quad \\ \downarrow \quad \downarrow \\ \quad \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \right)^{n+1, n-1} + \sum_{\substack{r+s+t+u \\ =k}} (-1)^{r+k} \left(\begin{array}{c} \quad \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \right)^{n+1, n-1} \left(\begin{array}{c} \quad \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ \blacktriangle \quad \blacktriangle \end{array} \right)^{s+u} \\
 &\stackrel{(5.25)}{=} \left(\begin{array}{c} \quad \\ \downarrow \quad \downarrow \\ \quad \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \right)^{n+1, n-1} + \sum_{\substack{r+s+t+u \\ =k}} (-1)^{r+k} \left(\begin{array}{c} \quad \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \right)^{n+1, n-1} \left(\begin{array}{c} \quad \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ \blacktriangle \quad \blacktriangle \end{array} \right)^{s+u} \\
 &\stackrel{(5.30)}{=} \left(\begin{array}{c} \quad \\ \downarrow \quad \downarrow \\ \quad \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \right)^{n+1, n-1} + (-1)^k \left(\begin{array}{c} \quad \\ \downarrow \quad \downarrow \\ \quad \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \right)^{n+1, n-1} \left(\begin{array}{c} \quad \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ \blacktriangle \quad \blacktriangle \end{array} \right)^k \\
 &\stackrel{(5.26), (5.8)}{=} \left(\begin{array}{c} \quad \\ \downarrow \quad \downarrow \\ \quad \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \right)^{n+1, n-1} + (-1)^{n+k+1} \left(\begin{array}{c} \quad \\ \downarrow \quad \downarrow \\ \quad \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \right)^{n+1, n-1}
 \end{aligned}$$

Returning to the computation in (8.3), this gives

$$\tau^* \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k \end{array} \right)^n = (-1)^{p+q} \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k \end{array} \right)^n + (-1)^{n+k+1} \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k \end{array} \right)^n - \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \quad \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \right)^{p+q-k}$$

$$\begin{aligned}
& \stackrel{(5.18),(5.25),\underline{(5.26)},(8.1)}{=} (-1)^{p+q} \left(\begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} - \text{Diagram 3} \end{array} \right) \\
& \stackrel{(5.36a)}{=} (-1)^{p+q} \left(\begin{array}{c} \text{Diagram 2} - \text{Diagram 3} - \sum_{r+s+t+u=k} (-1)^{r+k} \text{Diagram 4} \end{array} \right) \\
& \stackrel{(5.31)}{=} (-1)^{p+q} \left(\begin{array}{c} \text{Diagram 2} - \sum_{j=0}^k (-1)^j \text{Diagram 5} - \sum_{r+s+t+u=k} (-1)^{r+k} \text{Diagram 4} \end{array} \right) \\
& \stackrel{(5.29)}{=} (-1)^{p+q} \left(\begin{array}{c} \text{Diagram 2} - \sum_{r+s+t+u=k} (-1)^{r+k} \sum_{y+z=r} (-1)^z \text{Diagram 6} \end{array} \right) \\
& = (-1)^{p+q+k} \left(\begin{array}{c} \text{Diagram 2} - \sum_{r+s+t+u=k} \text{Diagram 6} - \sum_{r+s+t+u=k} \sum_{y=1}^r (-1)^y \text{Diagram 7} \end{array} \right) \\
& \stackrel{(5.14)}{=} (-1)^{p+q+k} \left(\begin{array}{c} \text{Diagram 2} + \sum_{r+s+t+u=k} \sum_{j=0}^2 (-1)^j \binom{2}{j} \text{Diagram 8} \end{array} \right).
\end{aligned}$$

The last gray terms in this computation are again in the kernel of the graded composition pairing since they are compositions with positive degree endomorphisms of the endopositive object $\mathbf{X}_i^{(k)}$. The earlier gray terms are in the kernel by Lemma 7.26 and Remark 8.3, since they factor through $\mathbf{X}_i^{(\ell)}$ for $\ell \neq k$. We henceforth use both arguments tacitly when marking morphisms gray.

By Lemma 7.20, all terms in the final summation are in the kernel of the graded composition pairing unless $j = u$, thus we find

$$\begin{aligned}
\tau_* \left(\text{diagram} \right) &= (-1)^{p+q+k} \left((-1)^{n+1} \left(\text{diagram} \right) + \sum_{\substack{r+s+t \\ =k}} \left(\text{diagram} \right) - 2 \sum_{\substack{r+s+t \\ =k-1}} \left(\text{diagram} \right) \right. \\
&\quad \left. + \sum_{\substack{r+s+t \\ =k-2}} \left(\text{diagram} \right) \right) \\
&\stackrel{(5.8)}{=} (-1)^{p+q+k+n} \left(- \left(\text{diagram} \right) + \sum_{s=0}^k (k-s+1) \left(\text{diagram} \right) \right. \\
&\quad \left. - 2 \sum_{s=0}^{k-1} (k-s) \left(\text{diagram} \right) + \sum_{s=0}^{k-2} (k-s-1) \left(\text{diagram} \right) \right) \\
&= (-1)^{p+q+k+n} \left(- \left(\text{diagram} \right) + \left(\text{diagram} \right) + 2 \left(\text{diagram} \right) - 2 \left(\text{diagram} \right) \right) \\
&= (-1)^{p+q+k+n+1} \left(\left(\text{diagram} \right) - \left(\text{diagram} \right) \right).
\end{aligned}$$

Next, we compute

$$\begin{aligned}
 \tau^* \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k+1 \quad k+1 \end{array} \right)^n &= (-1)^{p+q} \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k+1 \quad k+1 \end{array} \right)^n \stackrel{(5.4),(5.26)}{=} (-1)^{p+q} \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k+1 \quad k+1 \end{array} \right)^n - \sum_{\substack{y+z= \\ p+q-k-1}} \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k+1 \quad k+1 \end{array} \right)^n \\
 &\stackrel{(8.4)}{=} (-1)^{p+q} \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k+1 \quad k+1 \end{array} \right)^n + (-1)^{n+k+2} \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k+1 \quad k+1 \end{array} \right)^n \\
 &\stackrel{(5.18),(5.25),(5.26)}{=} (-1)^{p+q} \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k+1 \quad k+1 \end{array} \right)^n + (-1)^{n+k} \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k+1 \quad k+1 \end{array} \right)^n.
 \end{aligned}
 \tag{8.5}$$

Using equation (10.2) below to simplify the last diagram in (8.5) gives

$$\tau^* \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k+1 \quad k+1 \end{array} \right)^n = (-1)^{p+q} \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k+1 \quad k+1 \end{array} \right)^n \stackrel{(5.26),(8.1)}{=} (-1)^{p+q} \left(\begin{array}{c} k \quad k \\ \downarrow \quad \downarrow \\ \bullet \quad \bullet \\ \downarrow \quad \downarrow \\ k+1 \quad k+1 \end{array} \right)^n. \quad \square$$

Corollary 8.10. We have

$$\mathrm{Tr}(\tau^*|_{V(\mathbf{x}_{\bullet}^{(k)}, \mathbf{x}_{\bullet}^{(k)}, \mathbf{x}_{\bullet})}) = (-1)^{n+k+1} \sum_{j=0}^{k-1} (-1)^j [2(k-j)] = (-1)^{n+k+1} "[k][k+1]"$$

and

$$\mathrm{Tr}(\tau^*|_{V(\mathbf{x}_{\bullet}^{(k+1)}, \mathbf{x}_{\bullet}^{(k)}, \mathbf{x}_{\bullet})}) = \sum_{j=0}^k (-1)^j [2(k-j)+1] = "[k+1]^2".$$

Proof. Using Proposition 8.9, we compute

$$\begin{aligned}
 \mathrm{Tr}(\tau^*|_{V(\mathbf{x}_{\bullet}^{(k)}, \mathbf{x}_{\bullet}^{(k)}, \mathbf{x}_{\bullet})}) &= \sum_{\substack{0 \leq p \leq k \\ 0 \leq q \leq k-1}} (-1)^{p+q+n+k+1} q^{2(p+q-k)+1} (\delta_{p,q} - \delta_{p,k}) \\
 &= (-1)^{n+k+1} q^{1-2k} \sum_{d=0}^{2k-1} (-1)^d q^{2d} \sum_{\substack{p+q=d \\ 0 \leq p \leq k \\ 0 \leq q \leq k-1}} (\delta_{p,q} - \delta_{p,k}).
 \end{aligned}$$

Now,

$$\sum_{\substack{p+q=d \\ 0 \leq p \leq k \\ 0 \leq q \leq k-1}} \delta_{p,q} = \begin{cases} 1 & d \text{ is even and } d \leq 2k-2 \\ 0 & \text{else} \end{cases} \quad \text{and} \quad \sum_{\substack{p+q=d \\ 0 \leq p \leq k \\ 0 \leq q \leq k-1}} \delta_{p,k} = \begin{cases} 1 & k \leq d \leq 2k-1 \\ 0 & \text{else} \end{cases},$$

so

$$\mathrm{Tr}(\tau_*|_{V(\mathbf{X}_{\bullet}^{(k)}, \mathbf{X}_{\bullet}^{(k)} \mathbf{x}_{\bullet})}) = (-1)^{n+k+1} q^{1-2k} \left(\sum_{r=0}^{k-1} q^{4r} - \sum_{s=k}^{2k-1} (-1)^s q^{2s} \right) = (-1)^{n+k+1} \sum_{j=0}^{k-1} (-1)^j [2(k-j)].$$

For the second statement, we have

$$\begin{aligned} \mathrm{Tr}(\tau_*|_{V(\mathbf{X}_{\bullet}^{(k+1)}, \mathbf{X}_{\bullet}^{(k)} \mathbf{x}_{\bullet})}) &= \sum_{0 \leq p, q \leq k} (-1)^{p+q} q^{2(p+q-k)} \delta_{p,q} = \sum_{0 \leq p \leq k} q^{4p-2k} \\ &= \sum_{j=0}^k (-1)^j [2(k-j) + 1]. \quad \square \end{aligned}$$

Theorem 8.11. The following equality holds in $K_0^\tau((\mathcal{B}_m)^\tau)$:

$$[(\mathbf{X}_{\bullet}^{(k)}, \mathbf{LC}_k)]_\tau \cdot [(\mathbf{X}_{\bullet}, \mathbf{LC}_1)]_\tau = (-1)^{n+k+1} [k][k+1] [(\mathbf{X}_{\bullet}^{(k)}, \mathbf{LC}_k)]_\tau + [k+1]^{2n} [(\mathbf{X}_{\bullet}^{(k+1)}, \mathbf{LC}_{k+1})]_\tau.$$

Proof. This follows from Proposition 7.51 and Corollary 8.10. \square

This result implies Theorem 1.17 from the introduction.

Corollary 8.12. Let \mathbb{K} be an integral domain in which 2 is invertible. The assignment $\mathbf{x}^{(k)} \mapsto [(\mathbf{X}^{(k)}, (-1)^{\binom{n+1}{2} + \binom{n-k+1}{2} + k} \mathbf{LC}_k)]_\tau$ determines an isomorphism

$$U'_{-q^2}(\mathfrak{so}_2) \xrightarrow{\cong} \mathbb{C}(q) \otimes_{\mathbb{Z}[q^\pm]} K_0^\tau((\mathcal{B}_2)^\tau).$$

Proof. Set $\mathbf{x}^{(k)} := [(\mathbf{X}^{(k)}, (-1)^{\binom{n+1}{2} + \binom{n-k+1}{2} + k} \mathbf{LC}_k)]_\tau$. Theorem 8.11 gives that

$$\begin{aligned} (-1)^{2\binom{n+1}{2} + \binom{n-k+1}{2} + \binom{n}{2} + k+1} \mathbf{x}^{(k)} \mathbf{x}^{(1)} &= (-1)^{n+2k+1 + \binom{n+1}{2} + \binom{n-k+1}{2}} [k][k+1] \mathbf{x}^{(k)} \\ &\quad + (-1)^{\binom{n+1}{2} + \binom{n-k}{2} + k+1} [k+1]^{2n} \mathbf{x}^{(k+1)}, \end{aligned}$$

i.e. that

$$\mathbf{x}^{(k)} \mathbf{x}^{(1)} = (-1)^k [k][k+1] \mathbf{x}^{(k)} + (-1)^k [k+1]^{2n} \mathbf{x}^{(k+1)}.$$

This is exactly the relation that defines $\mathbf{x}^{(k)} \in U'_{-q^2}(\mathfrak{so}_2)$ and $\{\mathbf{x}^{(k)}\}_{k \geq 0}$ are a $\mathbb{C}(q)$ -basis for $U'_{-q^2}(\mathfrak{so}_2)$, so $\mathbf{x}^{(k)} \mapsto \mathbf{x}^{(k)}$ defines a $\mathbb{C}(q)$ -algebra homomorphism $U'_{-q^2}(\mathfrak{so}_2) \rightarrow \mathbb{C}(q) \otimes_{\mathbb{Z}[q^\pm]} K_0^\tau((\mathcal{B}_2)^\tau)$. Since Hypothesis 7.44 holds for \mathcal{B}_m , Proposition 7.51 implies that this homomorphism is an isomorphism. \square

8.3. Devil's Serre relation for \mathbf{X}_i and \mathbf{X}_j . We now show that a categorification of the devil's Serre relation for $U'_{-q^2}(\mathfrak{so}_m)$ holds for \mathbf{X}_i and $\mathbf{X}_{i \pm 1}$. Precisely, we show that the relation (4.40) holds for the classes of $(\mathbf{X}_i, \mathbf{LC}_1)$ and $(\mathbf{X}_{i \pm 1}, \mathbf{LC}_1)$ in the weighted Grothendieck group of $(\mathcal{B}_m)^\tau$.

We begin by recording some isomorphisms in $\check{U}_q(\mathfrak{gl}_m)$. For each $1 \leq i \leq m-1$, equations (5.10), (5.11), and (5.36b) give the following isomorphisms with indicated projection maps:

$$(8.6) \quad \left(\begin{array}{c} \text{X} \\ \text{Y} \end{array} \begin{array}{c} \text{X} \\ \text{Y} \end{array} \right)^T : \mathcal{E}_i \mathcal{F}_i \mathbb{1}_{\mathbf{a}} \cong \mathcal{F}_i \mathcal{E}_i \mathbb{1}_{\mathbf{a}} \oplus \mathbb{1}_{\mathbf{a}}, \quad \text{if } \alpha_i^\vee(\mathbf{a}) = 1,$$

$$(8.7) \quad \left(\begin{array}{c} \text{X} \\ \text{Y} \end{array} \begin{array}{c} \text{X} \\ \text{Y} \end{array} \right)^T : \mathcal{F}_i \mathcal{E}_i \mathbb{1}_{\mathbf{a}} \cong \mathcal{E}_i \mathcal{F}_i \mathbb{1}_{\mathbf{a}} \oplus \mathbb{1}_{\mathbf{a}}, \quad \text{if } \alpha_i^\vee(\mathbf{a}) = -1,$$

$$(8.8) \quad \left(\begin{array}{c} \text{X} \\ \curvearrowright \\ \curvearrowleft \end{array} \right)^T : \mathcal{F}_i \mathcal{E}_i \mathbb{1}_{\mathbf{a}} \cong \mathcal{E}_i \mathcal{F}_i \mathbb{1}_{\mathbf{a}} \oplus [2] \mathbb{1}_{\mathbf{a}}, \quad \text{if } \alpha_i^\vee(\mathbf{a}) = -2,$$

and

$$(8.9) \quad \left(\begin{array}{c} \text{X} \\ \text{X} \\ \text{X} \end{array} \right)^T : \mathcal{F}_i^{(2)} \mathcal{E}_i^{(2)} \cong \mathcal{E}_i^{(2)} \mathcal{F}_i^{(2)} \oplus \mathcal{E}_i \mathcal{F}_i, \quad \text{if } \alpha_i^\vee(\mathbf{a}) = -1.$$

Further, for $j \neq i$ and all \mathbf{a} , we have

$$(8.10) \quad \begin{array}{c} \text{X} \\ \text{X} \end{array} : \mathcal{E}_i^{(k)} \mathcal{F}_j^{(l)} \mathbb{1}_{\mathbf{a}} \cong \mathcal{F}_j^{(l)} \mathcal{E}_i^{(k)} \mathbb{1}_{\mathbf{a}}.$$

which follows as a consequence of (5.12), (5.4), (5.37), and (5.38).

Convention 8.13. Recall our convention for colors in Convention 5.18 is that strands corresponding to the i th Dynkin node are **green**, while strands corresponding to the $i - 1$ and $i + 1$ Dynkin nodes are colored **red** and **blue**, respectively. In this subsection, we will color $j \in \{i \pm 1\}$ using **purple** — the color obtained from combining blue and red. Further, for the remainder of this subsection, we use the convention that all thick strands have thickness 2. (As always, thin strands are assumed to have thickness 1.)

Suppose that $j = i \pm 1$, then [67, Theorem 3] implies that, for all \mathbf{a} , there are isomorphisms with indicated projection maps:

$$(8.11) \quad \begin{array}{c} \left(\begin{array}{c} \text{X} \\ \text{X} \end{array} \right)^T \\ \left(\begin{array}{c} \text{X} \\ \text{X} \end{array} \right)^T \end{array} : \mathcal{E}_i \mathcal{E}_j \mathcal{E}_i \mathbb{1}_{\mathbf{a}} \cong \mathcal{E}_i^{(2)} \mathcal{E}_j \mathbb{1}_{\mathbf{a}} \oplus \mathcal{E}_j \mathcal{E}_i^{(2)} \mathbb{1}_{\mathbf{a}} \\ : \mathcal{F}_i \mathcal{F}_j \mathcal{F}_i \mathbb{1}_{\mathbf{a}} \cong \mathcal{F}_i^{(2)} \mathcal{F}_j \mathbb{1}_{\mathbf{a}} \oplus \mathcal{F}_j \mathcal{F}_i^{(2)} \mathbb{1}_{\mathbf{a}}.$$

Our aim is to decompose

$$(\mathbf{X}_i, \mathbf{LC}_1) \otimes (\mathbf{X}_j, \mathbf{LC}_1) \otimes (\mathbf{X}_i, \mathbf{LC}_1) = (\mathbf{X}_i \mathbf{X}_j \mathbf{X}_i, \mathbf{LC}_1 \otimes \mathbf{LC}_1 \otimes \mathbf{LC}_1)$$

in $\text{Kar}((\mathcal{B}_m)^\tau)$. The first step is to decompose $\mathbf{X}_i \mathbf{X}_j \mathbf{X}_i$ in $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$.

Lemma 8.14. Let $j = i \pm 1$. There is an isomorphism

$$(8.12) \quad \mathbf{X}_i \mathbf{X}_j \mathbf{X}_i \cong \mathbf{X}_i^{(2)} \mathbf{X}_j \oplus \mathbf{X}_j \mathbf{X}_i^{(2)} \oplus [2] \mathbf{X}_i^{(2)} \oplus \mathbf{X}_i \oplus A \oplus B$$

in $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$, where $A = A_1 \oplus A_2$ and $B = B_1 \oplus B_2$ for $A_1 := \mathcal{F}_j \mathcal{E}_i \mathcal{F}_i \mathcal{E}_j \mathbb{1}_{\mathbf{n}}$, $B_1 := \mathcal{F}_i^{(2)} \mathcal{E}_j \mathcal{F}_j \mathcal{E}_i^{(2)} \mathbb{1}_{\mathbf{n}}$, $\tau(A_1) =: A_2$, and $\tau(B_1) =: B_2$. Moreover, the set

$$\{\mathbf{X}_i^{(2)} \mathbf{X}_j, \mathbf{X}_j \mathbf{X}_i^{(2)}, \mathbf{X}_i^{(2)}, \mathbf{X}_i, A_1, A_2, B_1, B_2\}$$

of summands appearing in (8.12) is an endopositive family.

Proof. We have the following chain of isomorphisms:

$$\begin{aligned}
\mathcal{F}_i \mathcal{E}_i \mathcal{F}_j \mathcal{E}_j \mathcal{F}_i \mathcal{E}_i \mathbb{1}_{\mathbf{n}} &\stackrel{(8.10)}{\cong} \mathcal{F}_i \mathcal{F}_j \mathcal{E}_i \mathcal{F}_i \mathcal{E}_j \mathcal{E}_i \mathbb{1}_{\mathbf{n}} \\
&\stackrel{(8.6)}{\cong} \mathcal{F}_i \mathcal{F}_j \mathcal{F}_i \mathcal{E}_i \mathcal{E}_j \mathcal{E}_i \mathbb{1}_{\mathbf{n}} \oplus \mathcal{F}_i \mathcal{F}_j \mathcal{E}_j \mathcal{E}_i \mathbb{1}_{\mathbf{n}} \\
&\stackrel{(8.11), (8.7)}{\cong} \mathcal{F}_i^{(2)} \mathcal{F}_j \mathcal{E}_i^{(2)} \mathcal{E}_j \mathbb{1}_{\mathbf{n}} \oplus \mathcal{F}_i^{(2)} \mathcal{F}_j \mathcal{E}_j \mathcal{E}_i^{(2)} \mathbb{1}_{\mathbf{n}} \oplus \mathcal{F}_j \mathcal{F}_i^{(2)} \mathcal{E}_i^{(2)} \mathcal{E}_j \mathbb{1}_{\mathbf{n}} \oplus \mathcal{F}_j \mathcal{F}_i^{(2)} \mathcal{E}_j \mathcal{E}_i^{(2)} \mathbb{1}_{\mathbf{n}} \\
&\quad \oplus \mathcal{F}_i \mathcal{E}_j \mathcal{F}_j \mathcal{E}_i \mathbb{1}_{\mathbf{n}} \oplus \mathcal{F}_i \mathcal{E}_i \mathbb{1}_{\mathbf{n}} \\
&\stackrel{(8.8), (8.9), (8.10)}{\cong} \mathcal{F}_i^{(2)} \mathcal{E}_i^{(2)} \mathcal{F}_j \mathcal{E}_j \mathbb{1}_{\mathbf{n}} \oplus \mathcal{F}_i^{(2)} \mathcal{E}_j \mathcal{F}_j \mathcal{E}_i^{(2)} \mathbb{1}_{\mathbf{n}} \oplus [2] \mathcal{F}_i^{(2)} \mathcal{E}_i^{(2)} \mathbb{1}_{\mathbf{n}} \oplus \mathcal{F}_j \mathcal{E}_i^{(2)} \mathcal{F}_i^{(2)} \mathcal{E}_j \mathbb{1}_{\mathbf{n}} \\
&\quad \oplus \mathcal{F}_j \mathcal{E}_i \mathcal{F}_i \mathcal{E}_j \mathbb{1}_{\mathbf{n}} \oplus \mathcal{F}_j \mathcal{E}_j \mathcal{F}_i^{(2)} \mathcal{E}_i^{(2)} \mathbb{1}_{\mathbf{n}} \oplus \mathcal{E}_j \mathcal{F}_i \mathcal{E}_i \mathcal{F}_j \mathbb{1}_{\mathbf{n}} \oplus \mathcal{F}_i \mathcal{E}_i \mathbb{1}_{\mathbf{n}} \\
&\stackrel{(8.10)}{\cong} \mathcal{F}_i^{(2)} \mathcal{E}_i^{(2)} \mathcal{F}_j \mathcal{E}_j \mathbb{1}_{\mathbf{n}} \oplus \mathcal{F}_j \mathcal{E}_j \mathcal{F}_i^{(2)} \mathcal{E}_i^{(2)} \mathbb{1}_{\mathbf{n}} \oplus [2] \mathcal{F}_i^{(2)} \mathcal{E}_i^{(2)} \mathbb{1}_{\mathbf{n}} \oplus \mathcal{F}_i \mathcal{E}_i \mathbb{1}_{\mathbf{n}} \\
&\quad \oplus \mathcal{F}_j \mathcal{E}_i \mathcal{F}_i \mathcal{E}_j \mathbb{1}_{\mathbf{n}} \oplus \mathcal{E}_j \mathcal{F}_i \mathcal{E}_i \mathcal{F}_j \mathbb{1}_{\mathbf{n}} \oplus \mathcal{F}_i^{(2)} \mathcal{E}_j \mathcal{F}_j \mathcal{E}_i^{(2)} \mathbb{1}_{\mathbf{n}} \oplus \mathcal{F}_j \mathcal{E}_i^{(2)} \mathcal{F}_i^{(2)} \mathcal{E}_j \mathbb{1}_{\mathbf{n}}.
\end{aligned}$$

This yields (8.12), since $\tau(\mathcal{F}_j \mathcal{E}_i \mathcal{F}_i \mathcal{E}_j \mathbb{1}_{\mathbf{n}}) = \mathcal{E}_j \mathcal{F}_i \mathcal{E}_i \mathcal{F}_j \mathbb{1}_{\mathbf{n}}$ and $\tau(\mathcal{F}_i^{(2)} \mathcal{E}_j \mathcal{F}_j \mathcal{E}_i^{(2)} \mathbb{1}_{\mathbf{n}}) = \mathcal{F}_j \mathcal{E}_i^{(2)} \mathcal{F}_i^{(2)} \mathcal{E}_j \mathbb{1}_{\mathbf{n}}$.


That the summands appearing form an endopositive family is a straightforward but tedious verification using the Khovanov–Lauda Hom formula; see Remark 8.4. It can also be checked directly, as we now demonstrate for a pair of relevant Hom-spaces. (The others can be treated similarly.)

The proof of [34, Proposition 3.11] gives that the following diagrams:



possibly adorned with dots on strands and positive degree new-bubbles in the far right region, span $\text{End}_{\mathcal{U}_q(\mathfrak{gl}_m)}(\mathcal{F}_j \mathcal{E}_i \mathcal{F}_i \mathcal{E}_j \mathbb{1}_{\mathbf{n}})$. From (5.2), we compute that the degrees of these diagrams are 0, +4, +4, and +6 respectively. It follows that $\text{End}_{\mathcal{U}_q(\mathfrak{gl}_m)}(\mathcal{F}_j \mathcal{E}_i \mathcal{F}_i \mathcal{E}_j \mathbb{1}_{\mathbf{n}})$ is one-dimensional in degree zero and all other endomorphisms are in strictly positive degree, hence $\mathcal{F}_j \mathcal{E}_i \mathcal{F}_i \mathcal{E}_j \mathbb{1}_{\mathbf{n}}$ is an endopositive object.

The analogous diagrams for $\text{Hom}_{\mathcal{U}_q(\mathfrak{gl}_m)}(\mathcal{F}_j \mathcal{E}_i \mathcal{F}_i \mathcal{E}_j \mathbb{1}_{\mathbf{n}}, \mathcal{E}_j \mathcal{F}_i \mathcal{E}_i \mathcal{F}_j \mathbb{1}_{\mathbf{n}})$ are as follows.



Again, (5.2) gives that the degrees are +2, +4, +4, and +6. Thus, $\text{Hom}_{\mathcal{U}_q(\mathfrak{gl}_m)}(\mathcal{F}_j \mathcal{E}_i \mathcal{F}_i \mathcal{E}_j \mathbb{1}_{\mathbf{n}}, \mathcal{E}_j \mathcal{F}_i \mathcal{E}_i \mathcal{F}_j \mathbb{1}_{\mathbf{n}})$ can only be non-zero in strictly positive degrees. \square

We now proceed to compute the class of $(\mathbf{X}_i \mathbf{X}_j \mathbf{X}_i, \mathbf{LC}_1 \otimes \mathbf{LC}_1 \otimes \mathbf{LC}_1)$ in the weighted Grothendieck group, using the technique from Remark 7.21, as implemented in Lemma 7.52. This will be accomplished via a sequence of Lemmata which establish and study bases for the relevant factors in the domain of the non-degenerate graded composition pairing. For the rest of this subsection, all diagrams are now assumed to have the \mathfrak{gl}_m weight \mathbf{n} on the far right.

Notation 8.15. When applying the relation (5.5) for strands labelled by i and $j = i \pm 1$, we often deal with scalars of the form $\pm(i - j)$. For $j = i \pm 1$, we write

$$\epsilon_{ij} := j - i \quad \text{and} \quad \epsilon_{ji} := i - j.$$

We also write $\varphi_{ji} := (-1)^{a_j}$, when $\mathbf{a} = \mathbf{n} + \alpha_i$. By equation (5.8), this is the value of a counterclockwise j -colored degree-zero bubble in weight $\mathbf{n} + \alpha_i$. Note that $\epsilon_{ij} \varphi_{ji} = (-1)^{n-1}$ and $\epsilon_{ji} \varphi_{ji} = (-1)^n$ for $j = i \pm 1$.

Lemma 8.16. The maps

$$\text{pr} := \downarrow \begin{array}{c} \text{---} \\ \curvearrowright \\ \text{---} \end{array} \uparrow \quad \text{and} \quad \text{in} := \downarrow \begin{array}{c} \curvearrowleft \\ \text{---} \\ \uparrow \end{array} \uparrow$$

descend to bases for $V(\mathbf{X}_i \mathbf{X}_j \mathbf{X}_i, \mathbf{X}_i)$ and $V(\mathbf{X}_i, \mathbf{X}_i \mathbf{X}_j \mathbf{X}_i)$, respectively.

Proof. We use Lemma 7.25. Tracing through the isomorphisms in the proof of Lemma 8.14, we find the projection map $\mathbf{X}_i \mathbf{X}_j \mathbf{X}_i \rightarrow \mathbf{X}_i$ is

$$\downarrow \begin{array}{c} \text{---} \\ \curvearrowright \\ \text{---} \end{array} \uparrow \stackrel{(5.12)}{=} \text{pr}.$$

The statement for the map in can be obtained similarly, or follows from the computation of $\text{pr} \circ \text{in}$ in the proof of Lemma 8.17 below. \square

Lemma 8.17. The equality $\tau * \text{in} = \text{in}$ holds in $V(\mathbf{X}_i, \mathbf{X}_i \mathbf{X}_j \mathbf{X}_i)$.

Proof. First, we compute

$$\text{pr} \circ \text{in} = \downarrow \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \uparrow \stackrel{(5.12)}{=} \downarrow \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \uparrow \stackrel{(5.5)}{=} \epsilon_{ji} \cdot \downarrow \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \uparrow - \epsilon_{ji} \cdot \downarrow \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \uparrow.$$

The weight to the far right is \mathbf{n} , so the clockwise undotted i -colored bubble in weight $\mathbf{n} + \alpha_i$ has degree $2(1 - \alpha_i^\vee(\mathbf{n} + \alpha_i)) = -2$ and thus equals zero. Hence,

$$\text{pr} \circ \text{in} = -\epsilon_{ji} \cdot \downarrow \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \uparrow \stackrel{(5.7), (5.8)}{=} -\epsilon_{ji} \varphi_{ji} (-1)^n \text{id}_{\mathbf{X}_i} = -\text{id}_{\mathbf{X}_i}.$$

Next, we compute $\text{pr} \circ (\tau * \text{in})$. Since

$$\tau(\text{in}) = (-1)^{\ell'_i(\alpha_i) + r'_j(0)} \downarrow \begin{array}{c} \curvearrowleft \\ \text{---} \\ \uparrow \end{array} \uparrow \stackrel{(6.2), (6.30)}{=} \downarrow \begin{array}{c} \curvearrowleft \\ \text{---} \\ \uparrow \end{array} \uparrow$$

we have

$$\begin{aligned} \text{pr} \circ (\tau * \text{in}) &= \downarrow \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \uparrow \stackrel{(5.15), (5.7)}{=} - \sum_{\substack{p+q=0 \\ p \geq 0}} \downarrow \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \uparrow \stackrel{(5.8)}{=} -(-1)^{n-1} \downarrow \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \uparrow \\ &\stackrel{(5.12)}{=} (-1)^n \downarrow \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \uparrow \stackrel{(5.5)}{=} (-1)^n \epsilon_{ji} \downarrow \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \uparrow - (-1)^n \epsilon_{ji} \downarrow \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \uparrow \\ &\stackrel{(5.5), (5.4)}{=} (-1)^{n+1} \epsilon_{ji} \downarrow \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \uparrow \stackrel{(5.10), (5.8)}{=} (-1)^{n+1} \epsilon_{ji} \varphi_{ji} \downarrow \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \uparrow = -\text{id}_{\mathbf{X}_i}. \end{aligned}$$


It follows from Lemma 7.52 that $\tau * \text{in} = \text{in}$. \square

Lemma 8.18. The maps

$$\text{pr}_{-1} := \downarrow \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \uparrow \quad \text{and} \quad \text{pr}_{+1} := \downarrow \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \uparrow$$

descend to a basis for $V^{-1}(\mathbf{X}_i \mathbf{X}_j \mathbf{X}_i, \mathbf{X}_i^{(2)})$ and $V^{+1}(\mathbf{X}_i \mathbf{X}_j \mathbf{X}_i, \mathbf{X}^{(2)})$, respectively. Flipping each of pr_{-1} and pr_{+1} upside down and reversing orientation yields maps in_{-1} and in_{+1} that descend to a basis for $V^{-1}(\mathbf{X}_i^{(2)}, \mathbf{X}_i \mathbf{X}_j \mathbf{X}_i)$ and $V^{+1}(\mathbf{X}^{(2)}, \mathbf{X}_i \mathbf{X}_j \mathbf{X}_i)$, respectively.


Proof. We again use Lemma 7.25. The projections realized by the isomorphisms in the proof of Lemma 8.14 are



Applying (5.4) to the second diagram, and then using (5.6) and (5.5), we obtain the indicated maps. The argument for $\text{in}_{\pm 1}$ is similar, or follows from the computation that $\text{pr}_{\pm 1} \circ \text{in}_{\mp 1} = (-1)^n \text{id}_{\mathbf{X}^{(2)}}$ given in the proof of Lemma 8.19 below. \square

Lemma 8.19. The equalities $\tau * \text{in}_{\pm 1} = (-1)^n \text{in}_{\pm 1}$ hold in $V^{+1}(\mathbf{X}^{(2)}, \mathbf{X}_i \mathbf{X}_j \mathbf{X}_i)$.

Proof. We begin by computing




Note, for use below, that the same computation shows that $\text{pr}_{-1} \circ \text{in}_{+1} = (-1)^n \text{id}_{\mathbf{X}^{(2)}}$.

It follows from equations (6.1) and (6.33) and Theorem 6.16 that $\tau(\text{in}_{-1})$ is obtained from in_{-1} by reversing orientation and multiplying by $+1$. Now, we compute

$$\begin{aligned}
 \text{pr}_{+1} \circ (\tau * \text{in}_{-1}) &= \text{Diagram 1} \stackrel{(5.15)}{=} \text{Diagram 2} \stackrel{(5.8), (5.12), (5.4)}{=} (-1)^{n-1} \text{Diagram 3} \\
 &\stackrel{(5.5)}{=} (-1)^n \epsilon_{ji} \text{Diagram 4} - (-1)^n \epsilon_{ji} \text{Diagram 5} \stackrel{(5.25), (5.5), (5.4)}{=} (-1)^{n+1} \epsilon_{ji} \text{Diagram 6} \\
 &\stackrel{(5.25), (5.36b)}{=} (-1)^{n+1} \epsilon_{ji} \text{Diagram 7} \stackrel{(5.12), (5.4)}{=} (-1)^{n+1} \epsilon_{ji} \text{Diagram 8} \\
 &\stackrel{(5.5), (5.4)}{=} (-1)^{n+1} \epsilon_{ji}^2 \text{Diagram 9} \stackrel{(5.8)}{=} (-1)^{n+1} (-1)^n \text{Diagram 10} = \text{id}_{\mathbf{X}^{(2)}}.
 \end{aligned}$$

Lemma 7.52 then implies that $\tau * \text{in}_{-1} = (-1)^n \text{in}_{-1}$.

The argument for $\tau * \text{in}_{+1}$ is similar. We pair the equality $\text{pr}_{-1} \circ \text{in}_{+1} = (-1)^n \text{id}_{\mathbf{X}^{(2)}}$ established above with the computation of $\text{pr}_{-1} \circ (\tau * \text{in}_{+1})$ and apply Lemma 7.52. The computation of $\text{pr}_{-1} \circ (\tau * \text{in}_{+1})$ is exactly the same as for $\text{pr}_{+1} \circ (\tau * \text{in}_{-1})$ except that a -1 factor appears when computing $\tau(\text{in}_{+1})$ due to the presence of the dot, and we see



Regardless, we find that $\text{pr}_{-1} \circ (\tau * \text{in}_{+1}) = \text{id}_{\mathbf{X}^{(2)}}$ so $\tau * \text{in}_{+1} = (-1)^n \text{in}_{+1}$. □

Similar arguments to the proofs of Lemmata 8.18 and 8.16 give the following.

Lemma 8.20. The maps

$$\text{pr}_{\mathbf{X}_i^{(2)} \mathbf{X}_j} := \text{Diagram 1} \quad \text{and} \quad \text{pr}_{\mathbf{X}_j \mathbf{X}_i^{(2)}} := \text{Diagram 2}$$

descend to a bases for $V(\mathbf{X}_i \mathbf{X}_j \mathbf{X}_i, \mathbf{X}_i^{(2)} \mathbf{X}_j)$ and $V(\mathbf{X}_i \mathbf{X}_j \mathbf{X}_i, \mathbf{X}_j \mathbf{X}_i^{(2)})$. Flipping these diagrams upside down and reversing orientation yields maps, denoted $\text{in}_{\mathbf{X}_i^{(2)} \mathbf{X}_j}$ and $\text{in}_{\mathbf{X}_j \mathbf{X}_i^{(2)}}$, which descends to bases for $V(\mathbf{X}_i^{(2)} \mathbf{X}_j, \mathbf{X}_i \mathbf{X}_j \mathbf{X}_i)$ and $V(\mathbf{X}_j \mathbf{X}_i^{(2)}, \mathbf{X}_i \mathbf{X}_j \mathbf{X}_i)$. □

Lemma 8.21. The equalities

$$\tau * \text{in}_{\mathbf{X}_i^{(2)} \mathbf{X}_j} = -\text{in}_{\mathbf{X}_i^{(2)} \mathbf{X}_j} \quad \text{and} \quad \tau * \text{in}_{\mathbf{X}_j \mathbf{X}_i^{(2)}} = -\text{in}_{\mathbf{X}_j \mathbf{X}_i^{(2)}}$$

hold in $V(\mathbf{X}_i^{(2)} \mathbf{X}_j, \mathbf{X}_i \mathbf{X}_j \mathbf{X}_i)$ and $V(\mathbf{X}_j \mathbf{X}_i^{(2)}, \mathbf{X}_i \mathbf{X}_j \mathbf{X}_i)$, respectively.

Proof. First, we compute

$$\begin{aligned} \text{pr}_{\mathbf{X}_i^{(2)} \mathbf{X}_j} \circ \text{in}_{\mathbf{X}_i^{(2)} \mathbf{X}_j} &= \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \\ \text{Diagram 3} \\ \text{Diagram 4} \\ \text{Diagram 5} \\ \text{Diagram 6} \end{array} \\ &\stackrel{(5.12), (5.5), (5.4)}{=} \epsilon_{ij} \epsilon_{ji} \begin{array}{c} \text{Diagram 7} \\ \text{Diagram 8} \\ \text{Diagram 9} \\ \text{Diagram 10} \end{array} \\ &\stackrel{(5.4), (5.11)}{=} \begin{array}{c} \text{Diagram 11} \\ \text{Diagram 12} \\ \text{Diagram 13} \end{array} \stackrel{(5.20)}{=} \begin{array}{c} \text{Diagram 14} \\ \text{Diagram 15} \end{array}. \end{aligned}$$

In the penultimate step, we have omitted terms which are quickly seen to be zero using Lemma 7.26, as they contain endomorphisms of $\mathbf{X}_i^{(2)}$ that factor through \mathbf{X}_i .

Using equation (6.1), we see that $\tau(\text{in}_{\mathbf{X}_i^{(2)} \mathbf{X}_j})$ is obtained from $\text{in}_{\mathbf{X}_i^{(2)} \mathbf{X}_j}$ by reversing orientation and multiplying by

$$\begin{aligned} &(-1)^{v_{ij}(0) + v_{ji}(0) + r_j(\alpha_j) + r_j(\alpha_i + \alpha_j) + v_{ji}(0) + \ell_i(0) + \ell_i(\alpha_j) + v'_{ij}(\alpha_i + \alpha_j) + v_{ii}(0) + \ell_i(0) + \ell_i(\alpha_i)} \\ &\stackrel{(6.6)}{=} (-1)^{\ell_i(\alpha_j) + r_i(\alpha_i + \alpha_j) + r_i(\alpha_i) + \ell_i(\alpha_i)} \stackrel{(6.30)}{=} (-1)^{r_i(-\alpha_j) + r_i(\alpha_i + \alpha_j) + r_i(\alpha_i) + r_i(-\alpha_i)}. \end{aligned}$$

It follows from Theorem 6.16 that this is equal to

$$(-1)^{r_i(-\alpha_j) + r_i(\alpha_i + \alpha_j)} = \begin{cases} (-1)^{1+0} & \text{if } j = i - 1 \\ (-1)^{0+1} & \text{if } j = i + 1, \end{cases}$$

which, in either case, is equal to -1 . Thus,

$$\begin{aligned} -1 \cdot \text{pr}_{\mathbf{X}_i^{(2)} \mathbf{X}_j} \circ \tau * \text{in}_{\mathbf{X}_i^{(2)} \mathbf{X}_j} &= \begin{array}{c} \text{Diagram 16} \\ \text{Diagram 17} \\ \text{Diagram 18} \\ \text{Diagram 19} \\ \text{Diagram 20} \\ \text{Diagram 21} \\ \text{Diagram 22} \\ \text{Diagram 23} \\ \text{Diagram 24} \\ \text{Diagram 25} \\ \text{Diagram 26} \end{array} \\ &\stackrel{(5.6), (5.12)}{=} \begin{array}{c} \text{Diagram 27} \\ \text{Diagram 28} \\ \text{Diagram 29} \\ \text{Diagram 30} \end{array} \\ &\stackrel{(5.12), (5.6)}{=} \begin{array}{c} \text{Diagram 31} \\ \text{Diagram 32} \\ \text{Diagram 33} \\ \text{Diagram 34} \end{array} \stackrel{(5.5), (5.4)}{=} \begin{array}{c} \text{Diagram 35} \\ \text{Diagram 36} \end{array} \stackrel{(5.25), (5.20), (5.32), (5.11)}{=} \begin{array}{c} \text{Diagram 37} \\ \text{Diagram 38} \end{array}. \end{aligned}$$

It then follows from Lemma 7.52 that $\tau * \text{in}_{\mathbf{X}_i^{(2)} \mathbf{X}_j} = -\text{in}_{\mathbf{X}_i^{(2)} \mathbf{X}_j}$.

Similar computations (which we leave as an exercise) show that $\tau * \text{in}_{\mathbf{X}_j \mathbf{X}_i^{(2)}} = -\text{in}_{\mathbf{X}_j \mathbf{X}_i^{(2)}}$. \square

Theorem 8.22. The following equality holds in $K_0^\tau((\mathcal{B}_m)^\tau)$:

$$\begin{aligned} [(\mathbf{X}_i, \mathbf{LC}_1)]_\tau \cdot [(\mathbf{X}_{i\pm 1}, \mathbf{LC}_1)]_\tau \cdot [(\mathbf{X}_i, \mathbf{LC}_1)]_\tau &= [(\mathbf{X}_i, \mathbf{LC}_1)]_\tau + (-1)^n [2][(\mathbf{X}_i^{(2)}, \mathbf{LC}_2)]_\tau \\ &\quad - [(\mathbf{X}_i^{(2)}, \mathbf{LC}_2)]_\tau [(\mathbf{X}_{i\pm 1}, \mathbf{LC}_1)]_\tau - [(\mathbf{X}_{i\pm 1}, \mathbf{LC}_1)]_\tau [(\mathbf{X}_i^{(2)}, \mathbf{LC}_2)]_\tau. \end{aligned}$$

Proof. Using Proposition 7.51, the claim follows from Lemmata 8.14, 8.17, 8.19, and 8.21. \square

Combining Theorems 8.11 and 8.22, we arrive at the following result, which establishes the existence of the $\mathbb{C}(q)$ -algebra homomorphism appearing in Conjecture 1.18 from the introduction.

Corollary 8.23. The assignment $\mathbf{x}_i^{(k)} \mapsto [(\mathbf{X}_i^{(k)}, (-1)^{\binom{n+1}{2} + \binom{n-k+1}{2} + k} \mathbf{LC}_k)]_\tau$ determines a $\mathbb{C}(q)$ -algebra homomorphism $U_{-q^2}^l(\mathfrak{so}_2) \rightarrow K_0^\tau((\mathcal{B}_m)^\tau)$.

Proof. Set $\mathbf{x}_i^{(k)} := [(\mathbf{X}_i^{(k)}, (-1)^{\binom{n+1}{2} + \binom{n-k+1}{2} + k} \mathbf{LC}_k)]_\tau$ and $\mathbf{x}_i := \mathbf{x}_i^{(1)}$. In light of Corollary 8.12 and Proposition 1.19, it suffices to show that

$$(8.13) \quad \mathbf{x}_i \mathbf{x}_\ell = \mathbf{x}_\ell \mathbf{x}_i \quad \text{for } |i - \ell| > 1$$

and

$$(8.14) \quad \mathbf{x}_i \mathbf{x}_{i\pm 1} \mathbf{x}_i = \mathbf{x}_i^{(2)} \mathbf{x}_{i\pm 1} + \mathbf{x}_{i\pm 1} \mathbf{x}_i^{(2)} + [2]\mathbf{x}_i^{(2)} + \mathbf{x}_i.$$

Equation (8.13) follows from the isomorphism

$$\begin{array}{c} \text{X} \\ \text{X} \\ \text{X} \\ \text{X} \end{array} : [(\mathbf{X}_i, \mathbf{LC}_1)]_\tau \otimes [(\mathbf{X}_\ell, \mathbf{LC}_1)]_\tau \cong [(\mathbf{X}_\ell, \mathbf{LC}_1)]_\tau \otimes [(\mathbf{X}_i, \mathbf{LC}_1)]_\tau.$$

Finally, Theorem 8.22 gives that

$$\begin{aligned} (-1)^{\binom{n+1}{2} + \binom{n}{2} + 1} \mathbf{x}_i \mathbf{x}_{i\pm 1} \mathbf{x}_i &= (-1)^{\binom{n+1}{2} + \binom{n}{2} + 1} \mathbf{x}_i + (-1)^{n + \binom{n+1}{2} + \binom{n-1}{2} + 2} [2]\mathbf{x}_i^{(2)} \\ &\quad - (-1)^{\binom{n-1}{2} + 2 + \binom{n}{2} + 1} (\mathbf{x}_i^{(2)} \mathbf{x}_{i\pm 1} + \mathbf{x}_{i\pm 1} \mathbf{x}_i^{(2)}) \end{aligned}$$

which simplifies to (8.14). \square

9. BACKGROUND ON LINK HOMOLOGY

In this section, we review background material on link homology in type A . We also discuss certain representation-theoretic results which will allow us to deduce invariance results for our type B link homologies (defined below in Section 10) from their type A counterparts.

9.1. Type A link polynomials via Howe duality. In type A , work of Cautis–Kamnitzer–Licata [14] and Cautis–Kamnitzer–Morrison [15] shows that the quantum \mathfrak{sl}_N link polynomials $P_{\mathfrak{sl}_N}(\mathcal{L}_\beta^\lambda)$ can be computed using an auxiliary quantum group $U_q(\mathfrak{gl}_m)$ associated to the (non-simple) Lie algebra \mathfrak{gl}_m . This approach proceeds through (a quantization of) the Howe duality between \mathfrak{gl}_m and \mathfrak{gl}_N that we now briefly recall. Consider the vector space $\Lambda(\mathbb{C}^m \otimes \mathbb{C}^N)$, which admits actions of \mathfrak{gl}_m and \mathfrak{gl}_N that generate each others commutant. The weight space decomposition for the \mathfrak{gl}_m action (in degree k) is given by

$$(9.1) \quad \Lambda^k(\mathbb{C}^m \otimes \mathbb{C}^N) \cong \Lambda^k(\underbrace{\mathbb{C}^N \oplus \dots \oplus \mathbb{C}^N}_m) \cong \bigoplus_{\sum a_i = k} \Lambda^{a_1}(\mathbb{C}^N) \otimes \dots \otimes \Lambda^{a_m}(\mathbb{C}^N).$$

Most importantly for us, the (symmetric) braiding on the \mathfrak{sl}_N modules appearing as summands in the right-hand side of (9.1) admits a description in terms of the Weyl group action associated with the action of \mathfrak{gl}_m on the left-hand side. In the quantized setting, this remains true: the braiding on $U_q(\mathfrak{sl}_N)$ modules admits a description in terms of the quantum Weyl group action for $U_q(\mathfrak{gl}_m)$.

To formulate quantum skew Howe duality in the form most useful for our considerations, we use the idempotent form $\dot{U}_q(\mathfrak{gl}_m)$ of the quantum group from Section 5.1. Recall from Remark 5.2 that $\dot{U}_q(\mathfrak{gl}_m)$ is a category with objects $\mathbf{a} \in \mathbb{Z}^m$.

Theorem 9.1 ([14, Theorem 4.3], [15, Theorem 6.2.1]). There is a full functor

$$\mathrm{SH}_m^N: \dot{U}_q(\mathfrak{gl}_m) \rightarrow \mathbf{Rep}(U_q(\mathfrak{sl}_N))$$

given on objects by

$$(9.2) \quad \mathrm{SH}_m^N(\mathbf{a}) = \Lambda^{a_1}(\mathbb{C}^N) \otimes \cdots \otimes \Lambda^{a_m}(\mathbb{C}^N) =: \Lambda^{\mathbf{a}}(\mathbb{C}^N).$$

If we define the *renormalized quantum Weyl group* elements of $\dot{U}_q(\mathfrak{gl}_m)$ via

$$(9.3) \quad c_i^\pm \mathbb{1}_{\mathbf{a}} := \sum_{s \geq 0} (-q)^{\pm(s-a_{i+1})} f_i^{(\alpha_i^\vee(\mathbf{a})+s)} e_i^{(s)} \mathbb{1}_{\mathbf{a}},$$

then

$$(9.4) \quad \mathrm{SH}_m^N(c_i^\pm \mathbb{1}_{\mathbf{a}}) = (-q^{1/N})^{\mp a_i a_{i+1}} R_{\Lambda^{a_i}(\mathbb{C}^N), \Lambda^{a_{i+1}}(\mathbb{C}^N)}^{\mp 1}. \quad \square$$

By convention, $\Lambda^a(\mathbb{C}^N) = 0$ if $a < 0$ or $a > N$, so (9.3) sends $\mathbf{a} \in \mathbb{Z}^m$ to zero if any $a_i < 0$ or if any $a_i > N$. It follows that the functor SH_m^N factors through the following quotients of $\dot{U}_q(\mathfrak{gl}_m)$.

Definition 9.2. The *integral q -Schur algebra* ${}_{\mathbb{Z}}\dot{S}_q(\mathfrak{gl}_m)$ is the quotient of ${}_{\mathbb{Z}}\dot{U}_q(\mathfrak{gl}_m)$ by the ideal generated by all weight idempotents $\mathbb{1}_{\mathbf{a}}$ such that some $a_i < 0$. The *integral N -bounded quotient* of ${}_{\mathbb{Z}}\dot{S}_q(\mathfrak{gl}_m)$, denoted ${}_{\mathbb{Z}}\dot{S}_q^{\leq N}(\mathfrak{gl}_m)$, is the quotient of ${}_{\mathbb{Z}}\dot{S}_q(\mathfrak{gl}_m)$ by the ideal generated by the $\mathbb{1}_{\mathbf{a}}$ such that some $a_i > N$.

As in the case of the quantum group, there are non-integral versions of the above

$$\dot{S}_q(\mathfrak{gl}_m) := \mathbb{C}(q) \otimes_{\mathbb{Z}[q^\pm]} {}_{\mathbb{Z}}\dot{S}_q(\mathfrak{gl}_m), \quad \dot{S}_q^{\leq N}(\mathfrak{gl}_m) := \mathbb{C}(q) \otimes_{\mathbb{Z}[q^\pm]} {}_{\mathbb{Z}}\dot{S}_q^{\leq N}(\mathfrak{gl}_m)$$

which can also be defined as quotients of $\dot{U}_q(\mathfrak{gl}_m)$. The following now refines the first statement of Theorem 9.1.

Theorem 9.3 ([15, Theorem 4.4.1]). The functor $\dot{S}_q^{\leq N}(\mathfrak{gl}_m) \rightarrow \mathbf{Rep}(U_q(\mathfrak{sl}_N))$ is fully faithful. \square

Theorems 9.1 and 9.3 now provide a description of the $U_q(\mathfrak{sl}_N)$ link polynomial purely in terms of the N -bounded quotient of the q -Schur algebra.

Proposition 9.4. There is a unique $\mathbb{C}(q)$ -valued bilinear form $(-, -)_N$ on $\dot{S}_q^{\leq N}(\mathfrak{gl}_m)$ such that

- (1) $(\mathbb{1}_{\mathbf{b}} e_i x \mathbb{1}_{\mathbf{a}}, \mathbb{1}_{\mathbf{b}} y \mathbb{1}_{\mathbf{a}})_N = (\mathbb{1}_{\mathbf{b}-\alpha_i} x \mathbb{1}_{\mathbf{a}}, \mathbb{1}_{\mathbf{b}-\alpha_i} f_i y \mathbb{1}_{\mathbf{a}})_N$,
- (2) $(\mathbb{1}_{\mathbf{b}} f_i x \mathbb{1}_{\mathbf{a}}, \mathbb{1}_{\mathbf{b}} y \mathbb{1}_{\mathbf{a}})_N = (\mathbb{1}_{\mathbf{b}+\alpha_i} x \mathbb{1}_{\mathbf{a}}, \mathbb{1}_{\mathbf{b}+\alpha_i} e_i y \mathbb{1}_{\mathbf{a}})_N$,
- (3) $(\mathbb{1}_{\mathbf{b}} x e_i \mathbb{1}_{\mathbf{a}}, \mathbb{1}_{\mathbf{b}} y \mathbb{1}_{\mathbf{a}})_N = (\mathbb{1}_{\mathbf{b}} x \mathbb{1}_{\mathbf{a}+\alpha_i}, \mathbb{1}_{\mathbf{b}} y f_i \mathbb{1}_{\mathbf{a}+\alpha_i})_N$,
- (4) $(\mathbb{1}_{\mathbf{b}} x f_i \mathbb{1}_{\mathbf{a}}, \mathbb{1}_{\mathbf{b}} y \mathbb{1}_{\mathbf{a}})_N = (\mathbb{1}_{\mathbf{b}} x \mathbb{1}_{\mathbf{a}-\alpha_i}, \mathbb{1}_{\mathbf{b}} y e_i \mathbb{1}_{\mathbf{a}+\alpha_i})_N$,
- (5) $(\mathbb{1}_{\mathbf{b}} x \mathbb{1}_{\mathbf{a}}, \mathbb{1}_{\mathbf{b}'} y \mathbb{1}_{\mathbf{a}'})_N = 0$, unless $\mathbf{b} = \mathbf{b}'$ and $\mathbf{a} = \mathbf{a}'$, and
- (6) $(\mathbb{1}_{\mathbf{a}}, \mathbb{1}_{\mathbf{a}})_N = \prod_{i=1}^m \binom{N}{[a_i]}$.

Given $x \in \mathbb{1}_{\mathbf{a}} \dot{S}_q^{\leq N}(\mathfrak{gl}_m) \mathbb{1}_{\mathbf{a}}$, this bilinear form satisfies $\mathrm{Tr}_q(\mathrm{SH}_m^N(x)) = (\mathbb{1}_{\mathbf{a}}, x)_N$. Consequently, if $\beta = \beta_{i_1}^{\epsilon_1} \cdots \beta_{i_d}^{\epsilon_d} \in \mathrm{Br}_m$ is an \mathbf{a} -balanced braid, then

$$(9.5) \quad \overline{P}_{\mathfrak{sl}_N}(\mathcal{L}_\beta^{\mathbf{a}}) = (-q^{1/N})^{\epsilon(\beta, \mathbf{a})} (\mathbb{1}_{\mathbf{a}}, c_{i_1}^{\epsilon_1} \cdots c_{i_d}^{\epsilon_d} \mathbb{1}_{\mathbf{a}})_N.$$

Here, $\mathcal{L}_\beta^{\mathbf{a}}$ denotes the coloring of the braid closure \mathcal{L}_β determined by \mathbf{a} , and $\epsilon(\beta, \mathbf{a})$ is²⁰ the sum over the crossings in $\beta \mathbb{1}_{\mathbf{a}}$ of $\pm ab$, where a and b are the labels coloring the strands of the crossing and the sign is given by the sign of the crossing.

²⁰As the notation suggests, this can be viewed as a colored analogue of the exponent sum of β .

Proof. Properties (1) – (6) suffice to compute the value of such a bilinear form on $\dot{S}_q^{\leq N}(\mathfrak{gl}_m)$, e.g. using the triangular decomposition for $\dot{U}_q(\mathfrak{gl}_m)$ or the annular evaluation algorithm from [55, Theorems 1.2 and 3.2]. It thus suffices to show that such a form exists, and satisfies $\mathrm{Tr}_q(\mathrm{SH}_m^N(x)) = (\mathbb{1}_{\mathbf{a}}, x)_N$.

In fact, we can essentially use the latter as the definition. Consider the $\mathbb{C}(q)$ -linear anti-algebra involution of $\dot{U}_q(\mathfrak{gl}_m)$ given by $\mathbb{1}_{\mathbf{b}}e_i\mathbb{1}_{\mathbf{a}} \mapsto \mathbb{1}_{\mathbf{a}}f_i\mathbb{1}_{\mathbf{b}}$. This involution descends to $\dot{S}_q^{\leq N}(\mathfrak{gl}_m)$ and we denote the image of $x \in \dot{S}_q^{\leq N}(\mathfrak{gl}_m)$ under this involution by \bar{x} . For $x, y \in \dot{S}_q^{\leq N}(\mathfrak{gl}_m)$, set

$$(x, y)_N := \mathrm{Tr}_q(\mathrm{SH}_m^N(\bar{x}y))$$

which clearly satisfies (1) and (2). Properties (3) and (4) hold since Tr_q is trace-like, and (5) then follows from the mutual orthogonality of the weight idempotents $\mathbb{1}_{\mathbf{a}}$. Lastly, (6) holds since $\mathrm{Tr}_q(\mathrm{SH}_m^N(\mathbb{1}_{\mathbf{a}}))$ computes the quantum \mathfrak{sl}_N invariant of the \mathbf{a} -colored m -component unlink, which equals $\prod_{i=1}^m \left[\begin{matrix} N \\ a_i \end{matrix} \right]$. \square

Remark 9.5. Note that, in light of (9.4), it is the link polynomials $\overline{P}_{\mathfrak{sl}_N}(\mathcal{L})$ from Remark 3.11 that appear in (9.5).

9.2. Rickard complexes and colored \mathfrak{sl}_N link homology. We next discuss the categorification of the Howe duality approach to \mathfrak{sl}_N link polynomials from §9.1. The construction we present is entirely parallel to the decategorified story: first, one assigns to each braid $\beta \in \mathrm{Br}_m$ an invariant living in (the homotopy category of) a categorified analogue of the Schur algebra for \mathfrak{gl}_m . Passing to an N -bounded quotient and applying a trace-like functor yields a complex of graded vector spaces, whose homology is a link invariant which categorifies the \mathfrak{sl}_N link polynomial.

To begin in detail, we have categorical analogues of the q -Schur algebra and its N -bounded quotient.

Definition 9.6. Let $\mathcal{S}_q(\mathfrak{gl}_m)$ (respectively $\check{\mathcal{S}}_q(\mathfrak{gl}_m)$) be the quotient of $\mathcal{U}_q(\mathfrak{gl}_m)$ (respectively $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$) given by evaluating²¹ all formal alphabets \mathbb{X}_i in weight \mathbf{a} to alphabets of cardinality equal to a_i .

Let $\mathcal{S}_q^{\leq N}(\mathfrak{gl}_m)$ (respectively $\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)$) be the (further) quotient by the ideal $\mathcal{J}^{\leq N}$ generated by all new bubble morphisms $\boxed{\mathfrak{s}_\lambda(\mathbb{X}_i)}^{\mathbf{a}}$ wherein the partition λ does not fit inside an $a_i \times (N - a_i)$ box.

Remark 9.7. One might instead expect $\mathcal{S}_q^{\leq N}(\mathfrak{gl}_m)$ and $\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)$ to be the quotient by \mathfrak{gl}_m weights whose entries do not lie strictly between 0 and N . Such weights are indeed killed in $\mathcal{S}_q^{\leq N}(\mathfrak{gl}_m)$ and $\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)$ since $a \times (N - a)$ is not a box in this case (thus we quotient by $\mathrm{id}_{\mathbb{1}_{\mathbf{a}}} = \mathfrak{s}_\emptyset$). Our definition is a further quotient which ensures that Hom-spaces in $\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)$ have the correct size for the computation of (undeformed) \mathfrak{sl}_N link homology.

Remark 9.8. Let \mathbb{K} be a field of characteristic zero. As explained in [72], the 2-category $\mathrm{Kar}(\mathcal{S}_q(\mathfrak{gl}_m))$ is self-dual mixed; see Example 7.28. It follows from [71, Lemma 1.15] that $\mathrm{Kar}(\mathcal{S}_q^{\leq N}(\mathfrak{gl}_m))$ is self-dual mixed.

We call the 2-categories $\mathcal{S}_q(\mathfrak{gl}_m)$ and $\mathcal{S}_q^{\leq N}(\mathfrak{gl}_m)$, as well as their (partially) Karoubi completed variants, the *categorified Schur quotient* and the *categorified N -bounded Schur quotient*, respectively. The appropriateness of this terminology is given by the following.

Theorem 9.9. Let \mathbb{K} be a field. There are $\mathbb{Z}[q^\pm]$ -algebra isomorphisms

$$K_0(\mathrm{Kar}(\mathcal{S}_q(\mathfrak{gl}_m))) \cong_{\mathbb{Z}} \dot{S}_q(\mathfrak{gl}_m), \quad K_0(\mathrm{Kar}(\mathcal{S}_q^{\leq N}(\mathfrak{gl}_m))) \cong_{\mathbb{Z}} \dot{S}_q^{\leq N}(\mathfrak{gl}_m).$$

Proof. The statement for the (categorified) Schur algebra is the main result of [44]. The techniques therein can be used to establish the result for the N -bounded quotient. \square

²¹A consequence is that this kills all weights \mathbf{a} such that $a_i < 0$ for some $1 \leq i \leq m$, since, by convention, symmetric polynomials in alphabets of negative cardinality are the zero ring.

Before continuing, we make precise the final claim from Remark 9.7.

Proposition 9.10. Let \mathbb{K} be a commutative ring and let $\mathbb{1}_{\mathbf{a}}X\mathbb{1}_{\mathbf{a}}$ be a 1-endomorphism in $\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)$. Then, $\mathrm{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X)$ is a free \mathbb{K} -module and if $x \in \check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)$ denotes the class of X in $\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)$, then

$$(9.6) \quad \dim_{\mathbf{q}}(\mathbf{q}^{\sum_{i=1}^m a_i(a_i-N)} \mathrm{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X)) = (\mathbb{1}_{\mathbf{a}}, x)_N.$$

Proof. This is a consequence of the ‘‘annular evaluation algorithm’’ from [55, Theorems 1.2 and 3.2], which, in particular, shows that the same algorithm can be used to compute both sides of (9.6).

In some detail, first note that it suffices to establish the result when $\mathbb{K} = \mathbb{Z}$. Further, since

$$\mathrm{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, \mathbf{q}^k X) \cong \mathbf{q}^k \mathrm{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X)$$

and

$$\mathrm{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X_1 \oplus X_2) \cong \mathrm{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X_1) \oplus \mathrm{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X_2),$$

we can assume that X is given as a word in the 1-morphisms $\mathcal{E}_i^{(k)}$ and $\mathcal{F}_j^{(\ell)}$. By using the isomorphisms

$$\mathcal{E}_i^k \cong [k]! \mathcal{E}_i^{(k)} \quad \text{and} \quad \mathcal{F}_i^k \cong [k]! \mathcal{F}_i^{(k)},$$

the general case is a straightforward consequence of the case when X is a word in \mathcal{E}_i and \mathcal{F}_j , so we make this additional assumption.

The algorithm uses²² the following:

$$(9.7a) \quad \mathbf{q}^{\sum_{i=1}^m a_i(a_i-N)} \mathrm{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X_1 \mathbb{1}_{\mathbf{b}} X_2) \cong \mathbf{q}^{\sum_{i=1}^m b_i(b_i-N)} \mathrm{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{b}}, X_2 \mathbb{1}_{\mathbf{a}} X_1)$$

$$(9.7b) \quad \mathrm{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X_1 \mathcal{E}_i \mathcal{F}_i \mathbb{1}_{\mathbf{b}} X_2) \cong \mathrm{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X_1 \mathcal{F}_i \mathcal{E}_i \mathbb{1}_{\mathbf{b}} X_2) \\ \oplus [b_i - b_{i+1}] \mathrm{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X_1 X_2) \quad \text{if } b_i - b_{i+1} = \langle \alpha_i^\vee, \mathbf{b} \rangle \geq 0,$$

$$(9.7c) \quad \mathrm{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X_1 \mathcal{E}_i \mathcal{F}_i \mathbb{1}_{\mathbf{b}} X_2) \overset{\oplus}{\subset} \mathrm{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X_1 \mathcal{F}_i \mathcal{E}_i \mathbb{1}_{\mathbf{b}} X_2) \quad \text{if } b_i - b_{i+1} = \langle \alpha_i^\vee, \mathbf{b} \rangle < 0,$$

all of which hold integrally, to show that $\mathrm{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X)$ is isomorphic to a summand of a graded \mathbb{Z} -module of the form

$$\bigoplus_l \mathbf{q}^{k_l + \sum_{i=1}^m a_{l,i}(a_{l,i}-N)} \mathrm{End}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}_l}).$$

The algorithm proceeds by inducting on both the length of the word X , and the minimal weights (with respect to the standard partial order on the \mathfrak{gl}_m weight lattice) which appear in X . In doing so, it is crucial that we work in the (N -bounded) Schur quotient, which implies that $\mathbb{1}_{\mathbf{a}} \cong 0$ for sufficiently large \mathbf{a} .

Now, [34, Proposition 3.11], equation (5.8), and Definition 9.6 give a surjective ring homomorphism

$$(9.8) \quad H^*(\mathrm{Gr}_{a_{l,1}}(\mathbb{C}^N)) \otimes \cdots \otimes H^*(\mathrm{Gr}_{a_{l,m}}(\mathbb{C}^N)) \rightarrow \mathrm{End}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}_l}).$$

This map must be an isomorphism, since, after base change to a field, the dimension of the latter can be bounded below by $\prod_{j=1}^m q^{a_{l,j}(N-a_{l,j})} \begin{bmatrix} N \\ a_{l,j} \end{bmatrix}$ e.g. using the 2-representations of $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$ from [45] or [11] (which factor through $\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)$). Consequently, each $\mathbf{q}^{\sum_{i=1}^m a_{l,i}(a_{l,i}-N)} \mathrm{End}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}_l})$ is a free \mathbb{Z} -module (of graded rank $\prod_{i=1}^m \begin{bmatrix} N \\ a_{l,i} \end{bmatrix}$) so $\mathrm{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X)$ is a summand of a free graded \mathbb{Z} -module, thus is a free graded \mathbb{Z} -module.

²²The shift $\mathbf{q}^{\sum_{i=1}^m a_i(a_i-N)}$ here ensures that the assignment $X \mapsto \mathbf{q}^{\sum_{i=1}^m a_i(a_i-N)} \mathrm{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X)$ is indeed trace-like

Further, this procedure gives a recipe for computing the left-hand side of (9.6). Indeed, the equations (9.7) give

$$(9.9a) \quad \dim_{\mathbf{q}}(\mathbf{q}^{\sum_{i=1}^m a_i(a_i-N)} \text{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X_1 \mathbb{1}_{\mathbf{b}} X_2)) = \dim_{\mathbf{q}}(\mathbf{q}^{\sum_{i=1}^m b_i(b_i-N)} \text{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{b}}, X_2 \mathbb{1}_{\mathbf{a}} X_1))$$

$$(9.9b) \quad \dim_{\mathbf{q}}(\text{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X_1 \mathcal{E}_i \mathcal{F}_i \mathbb{1}_{\mathbf{b}} X_2)) = \text{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X_1 \mathcal{F}_i \mathcal{E}_i \mathbb{1}_{\mathbf{b}} X_2) \\ + [b_i - b_{i+1}] \dim_{\mathbf{q}}(\text{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X_1 X_2)) \quad \text{if } b_i - b_{i+1} = \langle \alpha_i^\vee, \mathbf{b} \rangle \geq 0,$$

$$(9.9c) \quad \dim_{\mathbf{q}}(\text{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X_1 \mathcal{E}_i \mathcal{F}_i \mathbb{1}_{\mathbf{b}} X_2)) = \dim_{\mathbf{q}}(\text{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X_1 \mathcal{F}_i \mathcal{E}_i \mathbb{1}_{\mathbf{b}} X_2)) \\ - [b_{i+1} - b_i] \dim_{\mathbf{q}}(\text{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X_1 X_2)) \quad \text{if } b_i - b_{i+1} = \langle \alpha_i^\vee, \mathbf{b} \rangle < 0,$$

which therefore can be used to write

$$\dim_{\mathbf{q}}(\mathbf{q}^{\sum_{i=1}^m a_i(a_i-N)} \text{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, X)) = \sum_{l=1}^{L_X} r_l q^{d_l} \cdot \dim_{\mathbf{q}}(\mathbf{q}^{\sum_{i=1}^m a_{l,i}(a_{l,i}-N)} \text{End}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}_l})).$$

However, the equations (9.9) hold with each instance of

$$\dim_{\mathbf{q}}(\mathbf{q}^{\sum_{i=1}^m a_i(a_i-N)} \text{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, -))$$

replaced by $(\mathbb{1}_{\mathbf{a}}, -)_N$, so the same recipe gives that

$$(\mathbb{1}_{\mathbf{a}}, x)_N = \sum_{l=1}^{L_X} r_l q^{d_l} (\mathbb{1}_{\mathbf{a}_l}, \mathbb{1}_{\mathbf{a}_l})_N.$$

Since

$$(\mathbb{1}_{\mathbf{a}_l}, \mathbb{1}_{\mathbf{a}_l})_N = \prod_{i=1}^m \begin{bmatrix} N \\ a_{l,i} \end{bmatrix} = \dim_{\mathbf{q}}(\mathbf{q}^{\sum_{i=1}^m a_{l,i}(a_{l,i}-N)} \text{End}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}_l})),$$

the result follows. \square

We next discuss the complexes categorifying the quantum Weyl group elements from Definition 9.3 defined in [13, 14, 11] by Cautis, Kamnitzer, and Licata (following pioneering work of Chuang and Rouquier [17, 62]). Adapted to our present setting of the categorified Schur quotient, these complexes are as follows.

Definition 9.11. For $a, b \geq 0$, the 2-strand Rickard complex is the chain complex

$$C_{a,b} := (\dots \xrightarrow{d} \mathbf{q}^{-k} \mathbf{t}^k \mathcal{F}^{(a-k)} \mathcal{E}^{(b-k)} \mathbb{1}_{a,b} \xrightarrow{d} \mathbf{q}^{-k-1} \mathbf{t}^{k+1} \mathcal{F}^{(a-k-1)} \mathcal{E}^{(b-k-1)} \mathbb{1}_{a,b} \xrightarrow{d} \dots) \\ = \left(\bigoplus_{k=0}^{\min(a,b)} \mathbf{q}^{-k} \mathbf{t}^k \mathcal{F}^{(a-k)} \mathcal{E}^{(b-k)} \mathbb{1}_{a,b}, d \right) \in \mathcal{C}(\check{\mathcal{S}}_q(\mathfrak{gl}_2))$$

with differential

$$(9.10) \quad d = \begin{array}{c} \begin{array}{ccc} a-k-1 & & b-k-1 \\ \downarrow & \leftarrow & \downarrow \\ & \leftarrow & \\ \downarrow & & \downarrow \\ a-k & & b-k \end{array} \\ (a, b) \end{array}.$$

More generally, for $1 \leq i \leq m-1$, the i^{th} Rickard complex is the chain complex

$$C(\beta_i) \mathbb{1}_{\mathbf{a}} := \mathbb{1}_{(a_1, \dots, a_{i-1})} \boxtimes C_{a_i, a_{i+1}} \boxtimes \mathbb{1}_{(a_{i+2}, \dots, a_m)} \in \mathcal{C}(\check{\mathcal{S}}_q(\mathfrak{gl}_m)).$$

The Rickard complex $C(\beta_i)\mathbb{1}_{\mathbf{a}}$ determines complexes in various quotients of $\mathcal{C}(\check{\mathcal{S}}_q(\mathfrak{gl}_2))$, (the homotopy category, the N -bounded quotient, etc.) which will we denote using the same notation. The following is a consequence of [13, 14, 19]; see the proof of [25, Proposition 2.25] for further details.

Theorem 9.12. Let \mathbb{K} be a field. The complex $C_{a,b}$ is invertible in $\mathcal{K}(\check{\mathcal{S}}_q(\mathfrak{gl}_2))$, with homotopy inverse the complex

$$\begin{aligned} C_{b,a}^\vee &:= (\dots \xrightarrow{d^\vee} \mathbf{q}^k \mathbf{t}^{-k} \mathcal{F}^{(b-k)} \mathcal{E}^{(a-k)} \mathbb{1}_{b,a} \xrightarrow{d^\vee} \mathbf{q}^{k-1} \mathbf{t}^{-k+1} \mathcal{F}^{(b-k+1)} \mathcal{E}^{(a-k+1)} \mathbb{1}_{b,a} \xrightarrow{d^\vee} \dots), \\ &= \left(\bigoplus_{k=0}^{\min(a,b)} \mathbf{q}^k \mathbf{t}^{-k} \mathcal{F}^{(b-k)} \mathcal{E}^{(a-k)} \mathbb{1}_{b,a}, d^\vee \right) \end{aligned}$$

i.e. $C_{b,a}^\vee \star C_{a,b} \simeq \mathbb{1}_{(a,b)} \simeq C_{b,a} \star C_{a,b}^\vee$. Here,

$$(9.11) \quad d^\vee = \begin{array}{c} \begin{array}{ccc} & b-k+1 & a-k+1 \\ & \downarrow & \uparrow \\ & \downarrow & \uparrow \\ & b-k & a-k \end{array} \\ (b, a) \end{array}.$$

Set $C(\beta_i^{-1})\mathbb{1}_{\mathbf{a}} := \mathbb{1}_{(a_1, \dots, a_{i-1})} \boxtimes C_{a_i, a_{i+1}}^\vee \boxtimes \mathbb{1}_{(a_{i+2}, \dots, a_m)}$. Given a braid word $\beta_{i_1}^{\epsilon_1} \dots \beta_{i_r}^{\epsilon_r}$, let

$$C(\beta_{i_1}^{\epsilon_1} \dots \beta_{i_r}^{\epsilon_r})\mathbb{1}_{\mathbf{a}} := C(\beta_{i_1}^{\epsilon_1}) \dots (\beta_{i_r}^{\epsilon_r})\mathbb{1}_{\mathbf{a}},$$

then these complexes satisfy the (colored) braid relations in $\mathcal{K}(\check{\mathcal{S}}_q(\mathfrak{gl}_m))$ up to canonical homotopy equivalence. \square

For each \mathfrak{gl}_m weight \mathbf{a} with $a_i \geq 0$ for $1 \leq i \leq m$, Theorem 9.12 canonically assigns a complex $C(\beta)\mathbb{1}_{\mathbf{a}} \in \mathcal{K}(\check{\mathcal{S}}_q(\mathfrak{gl}_m))$ to each braid $\beta \in \text{Br}_m$. For each $N \geq 1$, we therefore obtain complexes $C(\beta)\mathbb{1}_{\mathbf{a}} \in \mathcal{K}(\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m))$. (This complex is zero if $a_i \geq N$ for some $1 \leq i \leq m$.) Following [54, Section 6], we now recall a procedure that recovers the (colored) \mathfrak{sl}_N link homologies.

The functor

$$(9.12) \quad \text{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, -) : \mathbb{1}_{\mathbf{a}} \check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m) \mathbb{1}_{\mathbf{a}} \rightarrow \text{Vect}_{\mathbb{K}}^{\mathbb{Z}}$$

induces a functor on homotopy categories $\mathcal{K}(\mathbb{1}_{\mathbf{a}} \check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m) \mathbb{1}_{\mathbf{a}}) \rightarrow \mathcal{K}(\text{Vect}_{\mathbb{K}}^{\mathbb{Z}})$, that we denote similarly. Since $\mathcal{K}(\text{Vect}_{\mathbb{K}}^{\mathbb{Z}}) \cong \mathcal{D}(\text{Vect}_{\mathbb{K}}^{\mathbb{Z}}) \cong \text{Vect}_{\mathbb{K}}^{\mathbb{Z} \times \mathbb{Z}}$, the functor (9.12) assigns a bi-graded vector space to pairs $(\beta, \mathbf{a}) \in \text{Br}_m \times \mathbb{Z}_{\geq 0}^m$ such that β is \mathbf{a} -balanced, meaning $C(\beta)\mathbb{1}_{\mathbf{a}} = \mathbb{1}_{\mathbf{a}} C(\beta)\mathbb{1}_{\mathbf{a}}$. In down-to-earth terms, this is simply the homology of the complex $\text{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, \mathbb{1}_{\mathbf{a}} C(\beta)\mathbb{1}_{\mathbf{a}})$ of bi-graded vector spaces.

The following is a repackaging of results from [53, 54, 55] in the setup and conventions of the present paper; see also [11]. As above, if $(\beta, \mathbf{a}) \in \text{Br}_m \times \mathbb{Z}_{\geq 0}^m$ is a pair such that $C(\beta)\mathbb{1}_{\mathbf{a}} = \mathbb{1}_{\mathbf{a}} C(\beta)\mathbb{1}_{\mathbf{a}}$, then the corresponding colored braid closure is denoted by $\mathcal{L}_{\beta}^{\mathbf{a}}$.

Theorem 9.13. Let \mathbb{K} be a field. If $\mathbf{a} \in \mathbb{Z}_{\geq 0}^m$ and $\beta \in \text{Br}_m$ is a braid such that $C(\beta)\mathbb{1}_{\mathbf{a}} = \mathbb{1}_{\mathbf{a}} C(\beta)\mathbb{1}_{\mathbf{a}}$, then the complex

$$(9.13) \quad \llbracket \mathbb{1}_{\mathbf{a}} \beta \mathbb{1}_{\mathbf{a}} \rrbracket_N := \mathbf{q}^{\sum_{i=1}^m a_i (a_i - N)} \text{Hom}_{\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{a}}, \mathbb{1}_{\mathbf{a}} C(\beta)\mathbb{1}_{\mathbf{a}}) \in \mathcal{K}(\text{Vect}_{\mathbb{K}}^{\mathbb{Z}})$$

is an invariant of the framed colored link $\mathcal{L}_{\beta}^{\mathbf{a}}$. Denote the homology of the complex $\llbracket \mathbb{1}_{\mathbf{a}} \beta \mathbb{1}_{\mathbf{a}} \rrbracket_N$ by $H_{\mathfrak{sl}_N}(\mathcal{L}_{\beta}^{\mathbf{a}})$. Then, we have

$$(9.14) \quad \begin{aligned} \overline{P}_{\mathfrak{sl}_N}(\mathcal{L}_{\beta}^{\mathbf{a}}) &= (-1)^{\sum_{i=1}^m a_i (N - a_i)} (-q^{1/N})^{\epsilon(\beta, \mathbf{a})} \dim_{\mathbf{q}, \mathbf{t}} \left(H_{\mathfrak{sl}_N}^i(\mathcal{L}_{\beta}^{\mathbf{a}}) \right) \Big|_{t=-1} \\ &= (-1)^{\sum_{i=1}^m a_i (N - a_i)} (-q^{1/N})^{\epsilon(\beta, \mathbf{a})} \sum_i (-1)^i \dim_{\mathbf{q}} \left(H_{\mathfrak{sl}_N}^i(\mathcal{L}_{\beta}^{\mathbf{a}}) \right) \end{aligned}$$

i.e. $H_{\mathfrak{sl}_N}(\mathcal{L}_{\beta}^{\mathbf{a}})$ categorifies (a multiple of) the colored \mathfrak{sl}_N link polynomial. \square

Remark 9.14. It follows from [53, Theorem 4.12] and [54, Section 6] that the invariant $H_{\mathfrak{sl}_N}(\mathcal{L}_\beta^{\mathbf{a}})$ agrees with (colored) \mathfrak{sl}_N Khovanov–Rozansky link homology [36, 75, 76], up to a choice of conventions (for positive/negative crossings and grading shifts). Our conventions for the Rickard complexes in Definition 9.11 imply that if $\beta \in \text{Br}_m$, $\mathbf{a} = (a_1, \dots, a_m)$, and $\mathbf{a}' = (a_1, \dots, a_m, a_m)$, then

$$\llbracket \mathbb{1}_{\mathbf{a}'} \beta_m^\pm \mathbb{1}_{\mathbf{a}'} \rrbracket_N \simeq \mathbf{q}^{\mp a_m(N-a_m+1)} \mathbf{t}^{\pm a_m} \llbracket \mathbb{1}_{\mathbf{a}} \beta \mathbb{1}_{\mathbf{a}} \rrbracket_N.$$

In other words, a positive a -colored stabilization yields a shift of $\mathbf{q}^{-a(N-a+1)} \mathbf{t}^a$, while a negative a -colored stabilization yields the opposite shift of $\mathbf{q}^{a(N-a+1)} \mathbf{t}^{-a}$.

Remark 9.15. The precise decategorification result given in (9.14) follows from (9.4) and Remark 3.11. The astute reader will note that the first factor of ± 1 in (9.14) would not appear for the \mathfrak{sl}_N link polynomials defined in the conventions of [15]. They work with a different pivotal structure on $\text{Rep}(U_q(\mathfrak{sl}_N))$ than the one implicit in §3.1; in their conventions, the twist coefficients (3.5) lack the ± 1 factors.

The term $(-1)^{\sum_{i=1}^m a_i(N-a_i)}$ in (9.14) can be accounted for as follows. If we only multiply by $(-q^{1/N})^{\pm a_i a_{i+1}}$ for each \pm -crossing, the result is the \mathfrak{sl}_N polynomial of the mirror link, computed as in [15]. (The mirror since their \pm -crossing is our \mp -crossing.) However, we (implicitly) work with the pivotal structure studied in [66] while [15] work with the standard pivotal structure (which has positive circle values and no signs in the twist coefficients). To account for this discrepancy, we need to multiply by $(-1)^{a(N-a)}$ for each a -colored cap/cup pair in a link diagram. For closures of braids $\beta \in \text{Br}_m$, there are m such pairs.

9.3. An elaboration on Rickard canonicity. In order to define our type B link homology, we will need to establish an equivariant analogue of Theorem 9.12. Among other things, Theorem 9.12 states that when two braid words represent the two braids, then their Rickard complexes are *canonically* homotopy equivalent; we call this *Rickard canonicity*. In this section, we establish an effective method for understanding this canonical homotopy equivalence. We focus on the special case when $N = 2n$, which is the case relevant to our type B link homology.

Our method will be to study homotopy equivalences after applying a 2-functor from $\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_m)$ to the following simplified setting.

Definition 9.16. Let \mathcal{A}_n be the monoidal category of finitely generated free graded modules over $H^*(Gr_n(\mathbb{C}^{2n}))$. We treat this as a 2-category with one object, denoted (by abuse of notation) as \mathbf{n} .

We record some properties of this 2-category, which are mostly tautological. The identity 1-morphism $\mathbb{1}_{\mathbf{n}}$ in \mathcal{A}_n , i.e. the left regular module $H^*(Gr_n(\mathbb{C}^{2n}))$, is the only indecomposable 1-morphism in \mathcal{A}_n , up to grading shift and isomorphism. Left multiplication gives an identification $\text{End}_{\mathcal{A}_n}(\mathbb{1}_{\mathbf{n}}) = H^*(Gr_n(\mathbb{C}^{2n}))$, and the latter may be identified with the quotient of the ring $\text{Sym}(\mathbb{X})$ of symmetric functions in one alphabet \mathbb{X} by Schur polynomials indexed by partitions which do not fit in an $n \times n$ box.

Proposition 9.17. The following assignments determine a well-defined lax 2-functor

$$\Gamma: \text{Kar}(\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_m)) \rightarrow \mathcal{A}_n.$$

which is full on 2-morphisms. All objects of $\text{Kar}(\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_m))$ are sent to the unique object of \mathcal{A}_n . The identity 1-morphisms $\mathbb{1}_{\mathbf{a}}$ are sent to the zero module when $\mathbf{a} \neq \mathbf{n}$, and $\mathbb{1}_{\mathbf{n}} \mapsto \mathbb{1}_{\mathbf{n}}$. All other generating 1-morphisms \mathcal{E}_i and \mathcal{F}_i are sent to the zero module, and thus most generating 2-morphisms are sent to zero. The new bubbles $h_r(\mathbb{X}_i)$ are sent to $h_r(\mathbb{X})$ for all $r \geq 0$ and all $1 \leq i \leq m$.

Proof. Using the generators and relations for $\mathcal{U}_q(\mathfrak{gl}_m)$ in Definition 5.5, it is straightforward to check that the indicated assignments yield a lax 2-functor from $\mathcal{U}_q(\mathfrak{gl}_m)$ to \mathcal{A}_n . (This 2-functor is lax since when $\mathbf{a} \neq \mathbf{n}$, the structure map $H^*(Gr_n(\mathbb{C}^{2n})) = \mathbb{1}_{\Gamma(\mathbf{a})} \rightarrow \Gamma(\mathbb{1}_{\mathbf{a}}) = 0$ is not an isomorphism.)

To elaborate, both sides of most relations go immediately to zero, since they involve the 1-morphisms \mathcal{E}_i or \mathcal{F}_i . The only relation with any subtlety is (5.8). Recall that (5.8) is not a relation for fake bubbles (merely a notational convenience), while for real bubbles it is a relation equating a real bubble with a new bubble. It is crucial that all real bubbles in the region \mathbf{n} have strictly positive degree, so in this case the left-hand side of (5.8) is sent to zero, while the right-hand side $h_r(\mathbb{X}_i - \mathbb{X}_{i+1})$ is sent to $h_r(\mathbb{X} - \mathbb{X})$, which is zero when $r > 0$.

The 2-functor factors through $\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_m)$ by Definition 9.6 and is full on 2-morphisms because all $h_r(\mathbb{X})$ are in the image. Since projective modules for $H^*(Gr_n(\mathbb{C}^{2n}))$ are free, \mathcal{A}_n is Karoubian, and therefore there is an induced 2-functor $\text{Kar}(\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_m)) \rightarrow \mathcal{A}_n$. \square

Remark 9.18. All indecomposable 1-morphisms in $\text{Kar}(\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_m))$ except (shifts of) $\mathbb{1}_{\mathbf{n}}$ are sent by Γ to zero. In particular,

$$\Gamma(\mathcal{F}_i^{(k)} \mathcal{E}_i^{(k)} \mathbb{1}_{\mathbf{n}}) = 0, \quad \text{for } k > 0 \text{ and } i = 1, \dots, m-1.$$

Note that, by definition, the 1-morphism $\mathbb{1}_{\mathbf{n}}$ survives in the quotient.

In fact, for any weight \mathbf{a} with $a_i \neq a_{i+1}$ for some i , that $\mathbb{1}_{\mathbf{a}}$ is sent to zero is already implied by the fact that \mathcal{E}_i and \mathcal{F}_i are sent to zero. Indeed, in such weights there is at least one real bubble of degree zero, and consequently $\text{id}_{\mathbb{1}_{\mathbf{a}}} = h_0(\mathbb{X}_i - \mathbb{X}_{i+1})$ will be sent to zero.

We will use the symbol Γ to refer to several related 2-functors. We need not work with the full Karoubi envelope, but can restrict Γ to the partial idempotent completion $\check{\mathcal{S}}_q^{\leq 2n}(\mathfrak{gl}_m)$:

$$\Gamma: \check{\mathcal{S}}_q^{\leq 2n}(\mathfrak{gl}_m) \rightarrow \mathcal{A}_n.$$

Since Γ is additive, it induces a 2-functor between dg categories of chain complexes:

$$(9.15) \quad \Gamma: \mathcal{C}(\check{\mathcal{S}}_q^{\leq 2n}(\mathfrak{gl}_m)) \rightarrow \mathcal{C}(\mathcal{A}_n)$$

and hence also between homotopy categories. The targets of such functors provide a simplified setting to study Rickard complexes.

Lemma 9.19. The Rickard complexes $C(\beta_i) \mathbb{1}_{\mathbf{n}}$ and $C(\beta_i^{-1}) \mathbb{1}_{\mathbf{n}}$ are sent by Γ to the complexes $\mathbf{q}^{-n} \mathbf{t}^n \mathbb{1}_{\mathbf{n}}$ and $\mathbf{q}^n \mathbf{t}^{-n} \mathbb{1}_{\mathbf{n}}$ (with zero differential), respectively. More generally, if $\underline{\beta} = \beta_{i_1}^{\epsilon_1} \cdots \beta_{i_r}^{\epsilon_r}$ is a braid word and $\epsilon(\underline{\beta}) = \epsilon_1 + \cdots + \epsilon_r$ is the braid exponent, then the complex $C(\underline{\beta}) \mathbb{1}_{\mathbf{n}} = C(\beta_{i_1}^{\epsilon_1}) \mathbb{1}_{\mathbf{n}} \cdots C(\beta_{i_r}^{\epsilon_r}) \mathbb{1}_{\mathbf{n}}$ maps under (9.15) to the complex $(\mathbf{q}^{-n} \mathbf{t}^n)^{\epsilon(\underline{\beta})} \mathbb{1}_{\mathbf{n}}$ supported in a single degree.

Proof. The first statement is an obvious consequence of Remark 9.18 and the second follows from Γ being a 2-functor. \square

The following lemma is our effective method to understand morphisms between Rickard complexes in the homotopy category.

Lemma 9.20. Let \mathbb{K} be a field and let $\underline{\beta}$ and $\underline{\beta}'$ be two braid words for the same braid. The functor Γ from (9.15) induces an isomorphism

$$(9.16) \quad \text{Hom}_{\mathcal{K}(\check{\mathcal{S}}_q^{\leq 2n}(\mathfrak{gl}_m))}^0(C(\underline{\beta}) \mathbb{1}_{\mathbf{n}}, C(\underline{\beta}') \mathbb{1}_{\mathbf{n}}) \xrightarrow{\cong} \text{Hom}_{\mathcal{K}(\mathcal{A}_n)}^0(\Gamma(C(\underline{\beta}) \mathbb{1}_{\mathbf{n}}), \Gamma(C(\underline{\beta}') \mathbb{1}_{\mathbf{n}})) = \mathbb{K} \cdot \text{id}$$

(on the space of \mathbf{q} -degree zero morphisms). Further, a \mathbf{q} -degree zero chain map $f: C(\underline{\beta}) \mathbb{1}_{\mathbf{n}} \rightarrow C(\underline{\beta}') \mathbb{1}_{\mathbf{n}}$ in $\mathcal{C}(\check{\mathcal{S}}_q^{\leq 2n}(\mathfrak{gl}_m))$ will be a homotopy equivalence if and only if $\Gamma(f)$ is an isomorphism, and f will be nullhomotopic if and only if $\Gamma(f) = 0$.

Remark 9.21. Since \mathbb{K} is a field, it is immediate from the proof of Lemma 9.20 that f is a homotopy equivalence if and only if $\Gamma(f) \neq 0$. However, the following proof of the above lemma is designed to work over \mathbb{Z} , once invertibility of Rickard complexes is established over \mathbb{Z} .

Proof. Let $\underline{\beta}$ and $\underline{\beta}'$ be braid words for the same braid. Then, we necessarily have $\epsilon(\underline{\beta}) = \epsilon(\underline{\beta}')$, so Lemma 9.19 gives that $\Gamma(C(\underline{\beta})\mathbb{1}_{\mathbf{n}}) = \Gamma(C(\underline{\beta}')\mathbb{1}_{\mathbf{n}})$ is a complex supported in one degree, a shift of the free module $\mathbb{1}_{\mathbf{n}}$. Thus, the space

$$\mathrm{Hom}_{\mathcal{K}(\mathcal{A}_n)}^0(\Gamma(C(\underline{\beta})\mathbb{1}_{\mathbf{n}}), \Gamma(C(\underline{\beta}')\mathbb{1}_{\mathbf{n}}))$$

of \mathbf{q} -degree zero morphisms in the homotopy category consists solely of scalar multiples of the identity map. Since $\Gamma(C(\underline{\beta})\mathbb{1}_{\mathbf{n}})$ and $\Gamma(C(\underline{\beta}')\mathbb{1}_{\mathbf{n}})$ are one-term complexes supported in the same homological degree, the morphism space in the homotopy category agrees with the space

$$\mathrm{Hom}_{\mathrm{Ch}(\mathcal{A}_n)}^0(\Gamma(C(\underline{\beta})\mathbb{1}_{\mathbf{n}}), \Gamma(C(\underline{\beta}')\mathbb{1}_{\mathbf{n}}))$$

of \mathbf{q} -degree zero chain maps (which further agrees with the space of all \mathbf{q} -degree zero maps in $\mathcal{C}(\mathcal{A}_n)$). To summarize,

$$(9.17) \quad \mathrm{Hom}_{\mathcal{K}(\mathcal{A}_n)}^0(\Gamma(C(\underline{\beta})\mathbb{1}_{\mathbf{n}}), \Gamma(C(\underline{\beta}')\mathbb{1}_{\mathbf{n}})) = \mathbb{K} \cdot \mathrm{id} = \mathrm{Hom}_{\mathcal{C}(\mathcal{A}_n)}^{\mathbf{q}\text{-deg}=0}(\Gamma(C(\underline{\beta})\mathbb{1}_{\mathbf{n}}), \Gamma(C(\underline{\beta}')\mathbb{1}_{\mathbf{n}})),$$

i.e. elements in the latter are homotopy equivalences if and only if they are isomorphisms and are nullhomotopic if and only if they are zero.

By Theorem 9.12, $C(\underline{\beta})\mathbb{1}_{\mathbf{n}}$ and $C(\underline{\beta}')\mathbb{1}_{\mathbf{n}}$ are homotopy equivalent, and any choice of homotopy equivalence will induce an isomorphism

$$\mathrm{Hom}_{\mathcal{K}(\mathfrak{S}_{\bar{q}}^{\leq 2n}(\mathfrak{gl}_m))}^0(C(\underline{\beta})\mathbb{1}_{\mathbf{n}}, C(\underline{\beta}')\mathbb{1}_{\mathbf{n}}) \cong \mathrm{End}_{\mathcal{K}(\mathfrak{S}_{\bar{q}}^{\leq 2n}(\mathfrak{gl}_m))}^0(C(\underline{\beta})\mathbb{1}_{\mathbf{n}}).$$

Theorem 9.12 also gives that $C(\underline{\beta})$ is invertible, so the functor of tensoring with the inverse Rickard complex induces an isomorphism

$$\mathrm{End}_{\mathcal{K}(\mathfrak{S}_{\bar{q}}^{\leq 2n}(\mathfrak{gl}_m))}^0(C(\underline{\beta})\mathbb{1}_{\mathbf{n}}) \cong \mathrm{End}_{\mathcal{K}(\mathfrak{S}_{\bar{q}}^{\leq 2n}(\mathfrak{gl}_m))}^0(\mathbb{1}_{\mathbf{n}}) = \mathbb{K} \cdot \mathrm{id}.$$

The functor Γ sends $\mathrm{id}_{C(\underline{\beta})\mathbb{1}_{\mathbf{n}}} \in \mathrm{End}_{\mathcal{K}(\mathfrak{S}_{\bar{q}}^{\leq 2n}(\mathfrak{gl}_m))}^0(C(\underline{\beta})\mathbb{1}_{\mathbf{n}})$ to the identity map of $(\mathbf{q}^{-n}\mathbf{t}^n)^{\epsilon(\underline{\beta})}\mathbb{1}_{\mathbf{n}}$ in \mathcal{A}_n . Using the functoriality of Γ , this then implies that (9.16) is an isomorphism.

Since Γ is additive, it maps homotopy equivalences to homotopy equivalences and nullhomotopic maps to nullhomotopic maps. The final statement then follows from (9.17) and the isomorphism (9.16). \square

We use this lemma to pick distinguished canonical isomorphisms, thus clarifying Rickard canonicity.

Proposition 9.22. Let \mathbb{K} be a field and $\beta \in \mathrm{Br}_m$ be a braid. Given braid words $\underline{\beta}$ and $\underline{\beta}'$ for β , let $f_{\underline{\beta}', \underline{\beta}}: C(\underline{\beta})\mathbb{1}_{\mathbf{n}} \rightarrow C(\underline{\beta}')\mathbb{1}_{\mathbf{n}}$ be a homotopy equivalence for which $\Gamma(f_{\underline{\beta}', \underline{\beta}}) = \mathrm{id}_{(\mathbf{q}^{-n}\mathbf{t}^n)^{\epsilon(\underline{\beta})}\mathbb{1}_{\mathbf{n}}}$. If f is another homotopy equivalence satisfying this property, then $f \sim f_{\underline{\beta}', \underline{\beta}}$. Moreover, if $\underline{\beta}''$ is another braid word for β , then $f_{\underline{\beta}'', \underline{\beta}'} \circ f_{\underline{\beta}', \underline{\beta}} \sim f_{\underline{\beta}'', \underline{\beta}}$.

Proof. It follows from Lemma 9.20 that a chain map between such Rickard complexes is nullhomotopic if and only if it is zero after applying Γ . Thus, two chain maps between such Rickard complexes are homotopic if and only if they agree after one applies Γ . From this fact, the proposition is obvious. \square

Remark 9.23. We have focused here on the case $\mathbf{a} = \mathbf{n}$ for simplicity, and since this is the relevant setting for our spin link homology defined below. One might be able to make similar arguments for ‘‘Rickard canonicity’’ in arbitrary weight, by considering different quotient functors analogous to Γ .

10. SPIN LINK HOMOLOGY

At last, in this section we define our spin link homology theory. First, in §10.1 we introduce a monoidal category \mathcal{B}_m^n , defined as a quotient of \mathcal{B}_m , which inherits the involution τ . The equivariant category $(\mathcal{B}_m^n)^\tau$ will be the setting for type B_n spin-colored link homology. Then, in §10.2, we define complexes $C^\tau(\beta) \in \mathcal{C}((\mathcal{B}_m^n)^\tau)$ that determine a categorical braid group action. The latter fact is proved in §10.4 using a general result about lifting homotopy equivalences to equivariant categories that is established in §10.3. In §10.5 we apply an appropriate representable functor to $C^\tau(\beta)$ which yields a chain complex of super vector spaces whose homology is a bigraded super vector space valued braid invariant that is further invariant under the Markov moves, up to isomorphism. Hence, by Theorem 3.7, we obtain an invariant of links valued in isomorphism classes of bigraded super vector spaces. When $n = 1, 2, 3$, we prove this invariant categorifies the spin-colored quantum \mathfrak{so}_{2n+1} link polynomials²³, and we conjecture that this holds for all $n > 3$ as well. We therefore denote this new link homology theory by $H_{\mathfrak{so}_{2n+1}, \mathcal{S}}(\mathcal{L}_\beta)$.

Remark 10.1. Our construction bootstraps on the properties of the Rickard complexes from Theorem 9.12, namely that they are invertible and satisfy the braid relations. Since the proof of these results in the literature assumes that \mathbb{K} is a field, we will make this assumption when working with the Rickard complexes (essentially from Section 10.2 onward). Consequently, our link invariants will be defined over a field. However, once some folklore results are carefully established, it should be possible to refine our proofs to construct our link invariants over $\mathbb{K} = \mathbb{Z}[\frac{1}{2}]$, and thus over any commutative ring in which 2 is invertible.

10.1. $\mathcal{F}^{(k)}\mathcal{E}^{(k)}\mathbb{1}_n$ -generated Schur quotient. In §9.2, we saw that \mathfrak{sl}_N link homology may be formulated entirely in the setting of the N -bounded Schur quotient $\check{\mathcal{S}}_q^{\leq N}(\mathfrak{gl}_m)$ of the categorified quantum group $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$. We also saw in §6 that, in order to guarantee that τ is an involution, we must pass to $\check{\mathcal{F}}_q(\mathfrak{gl}_m)$, the full 2-subcategory of $\check{\mathcal{U}}_q(\mathfrak{gl}_m)$ generated by the 1-morphisms $\mathcal{F}_i^{(k)}\mathcal{E}_i^{(k)}\mathbb{1}_a$ and $\mathcal{E}_i^{(k)}\mathcal{F}_i^{(k)}\mathbb{1}_a$. Finally, to find a monoidal category categorifying the relations in the centralizer algebras studied in §4, we had to work in §8 with the equivariantization of the full 2-subcategory $\check{\mathcal{F}}_q(\mathfrak{gl}_m)\mathbb{1}_n$. To define our spin link homology, we will work in a setting that combines these desired features.

First, we verify that the 2-automorphism τ descends to an automorphism of the N -bounded Schur quotient from Definition 9.6. As discussed back in §1.1, and in light of the action of τ on weights, the relevant value here is $N = 2n$.

Lemma 10.2. If $\tau = \tau_n$ satisfies the conditions of Theorem 6.5, then it preserves the ideal $\mathcal{J}^{\leq 2n}$ from Definition 9.6. Consequently, it induces an automorphism of $\check{\mathcal{S}}_q^{\leq 2n}(\mathfrak{gl}_m)$.

Proof. Recall that $\mathcal{J}^{\leq N}$ is generated by all new bubble morphisms $\mathfrak{s}_\lambda(\mathbb{X}_i) \in \text{End}(\mathbb{1}_{n+\mathbf{a}})$ such that the partition λ does not fit inside an $(n + a_i) \times (n - a_i)$ box. (Note that here we consider ambient weight $n + \mathbf{a}$.) It follows from equation (6.3) that if $\mathfrak{s}_\lambda(\mathbb{X}_i) \in \text{End}(\mathbb{1}_{n+\mathbf{a}})$ is such that λ is not contained in an $(n + a_i) \times (n - a_i)$ box, then $\tau(\mathfrak{s}_\lambda(\mathbb{X}_i)) = \mathfrak{s}_{\lambda^t}(\mathbb{X}_i) \in \text{End}(\mathbb{1}_{n-\mathbf{a}})$ for λ^t not contained in an $(n - a_i) \times (n + a_i)$ box, as desired. \square

Recall that if τ satisfies the conditions of Theorem 6.16, then τ restricts to an involution on $\check{\mathcal{F}}_q(\mathfrak{gl}_m)$ by Corollary 6.25. We now define a Schur quotient category on which τ is an involution.

Definition 10.3. Let $\check{\mathcal{F}}_q^{\leq 2n}(\mathfrak{gl}_m)$ be the quotient of the 2-category $\check{\mathcal{F}}_q(\mathfrak{gl}_m)$ from Definition 6.24 by the intersection of $\check{\mathcal{F}}_q(\mathfrak{gl}_m)$ with the ideal $\mathcal{J}^{\leq 2n}$.

Corollary 10.4. The involution τ on $\check{\mathcal{F}}_q(\mathfrak{gl}_m)$ from Corollary 6.25 induces an involution τ on $\check{\mathcal{F}}_q^{\leq 2n}(\mathfrak{gl}_m)$. \square

²³Since $\mathfrak{sp}_4 \cong \mathfrak{so}_5$, we have thus also categorified the quantum \mathfrak{sp}_4 link invariant colored by the defining representation.

Finally, we arrive at our setting for type B_n spin link homology.

Definition 10.5. Let \mathcal{B}_m^n be the monoidal category $\mathbb{1}_n \check{\mathcal{F}}_q^{\leq 2n}(\mathfrak{gl}_m) \mathbb{1}_n$.

Corollary 10.6. The involution τ restricts from $\check{\mathcal{F}}_q^{\leq 2n}(\mathfrak{gl}_m)$ to a monoidal involution of \mathcal{B}_m^n .

Proof. In light of Corollary 10.4, it remains to note that $\tau(\mathbb{1}_n) = \mathbb{1}_n$. \square

Remark 10.7. The 2-category $\check{\mathcal{F}}_q^{\leq 2n}(\mathfrak{gl}_m)$ can equivalently be described as the full 2-subcategory of $\check{\mathcal{S}}_q^{\leq 2n}(\mathfrak{gl}_m)$ generated by all $\mathcal{F}_i^{(k)} \mathcal{E}_i^{(k)} \mathbb{1}_a$ and $\mathcal{E}_i^{(k)} \mathcal{F}_i^{(k)} \mathbb{1}_a$. Similarly, \mathcal{B}_m^n can be identified with the 2-subcategory of $\check{\mathcal{F}}_q^{\leq 2n}(\mathfrak{gl}_m)$ generated by the 1-morphisms $\mathbf{X}_i^{(k)} = \mathcal{F}_i^{(k)} \mathcal{E}_i^{(k)} \mathbb{1}_n$ and $\mathcal{E}_i^{(k)} \mathcal{F}_i^{(k)} \mathbb{1}_n$ for $i \in \{1, \dots, m-1\}$ and $k \in \mathbb{Z}_{\geq 0}$. Therefore, the following commutative diagram collects the various 2-categories introduced thus far.

$$\begin{array}{ccccc} \mathcal{B}_m & \hookrightarrow & \check{\mathcal{F}}_q(\mathfrak{gl}_m) & \hookrightarrow & \check{\mathcal{U}}_q(\mathfrak{gl}_m) \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{B}_m^n & \hookrightarrow & \check{\mathcal{F}}_q^{\leq 2n}(\mathfrak{gl}_m) & \hookrightarrow & \check{\mathcal{S}}_q^{\leq 2n}(\mathfrak{gl}_m) \end{array}$$

Here, all arrows are full and the horizontal arrows are faithful.

Remark 10.8. If \mathbb{K} is a field of characteristic zero, then \mathcal{B}_m^n is self-dual mixed. Further, if \mathbb{K} is also algebraically closed, then Hypothesis 7.44 holds and Corollary 7.47 implies that $(\mathcal{B}_m^n)^\tau$ is also self-dual mixed.

Before proceeding, we establish a decategorification result (Theorem 1.9) from §1.4.

Lemma 10.9. Let \mathbb{K} be an integral domain. The 1-morphisms $\mathbf{X}_\bullet^{(k)} \in \mathcal{B}_m^n$ are either indecomposable or zero, and they are non-zero if and only if $0 \leq k \leq n$.

Proof. The first statement follows from fullness of the functor $\mathcal{B}_m \rightarrow \mathcal{B}_m^n$. For the second statement, it suffices to argue that $\mathbf{X}_\bullet^{(k)}$ is non-zero in $\check{\mathcal{S}}_q^{\leq 2n}(\mathfrak{gl}_m)$ if and only if $0 \leq k \leq n$, which, in light of Theorem 9.9, can be easily checked in the Grothendieck ring. \square

Corollary 10.10. Let \mathbb{K} be an integral domain in which 2 is invertible. The direct sum decompositions from Theorems 8.11 and 8.22 hold in $(\mathcal{B}_m^n)^\tau$. Hence, in the $m = 2$ case, there is an isomorphism of $\mathbb{C}(q)$ -algebras

$$\mathbb{C}(q) \otimes_{\mathbb{Z}[q^\pm]} K_0^\tau((\mathcal{B}_2^n)^\tau) \xrightarrow{\cong} \text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$$

that sends $[(\mathbf{X}^{(k)}, (-1)^{\binom{n+1}{2} + \binom{n-k+1}{2} + k} \mathbf{LC}_k)]_\tau \mapsto \mathbf{X}^{(k)}$.

Proof. The first claim follows since the functor $\mathcal{B}_m \rightarrow \mathcal{B}_m^n$ intertwines the involution $\tau = \tau_n$ and sends $\mathbf{X}_i^{(k)} \in \mathcal{B}_m$ to $\mathbf{X}_i^{(k)} \in \mathcal{B}_m^n$. For the second, Lemma 10.9 implies that $\{\mathbf{X}^{(k)}\}_{0 \leq k \leq n}$ constitutes a complete set of indecomposable objects in \mathcal{B}_m^n , up to isomorphism and shift. Hence, Proposition 7.51 implies that the elements $[(\mathbf{X}^{(k)}, (-1)^{\binom{n+1}{2} + \binom{n-k+1}{2} + k} \mathbf{LC}_k)]_\tau$ for $0 \leq k \leq n$ give a basis for $K_0^\tau((\mathcal{B}_2^n)^\tau)$. Proposition 4.25 then implies that the indicated assignment is an isomorphism of $\mathbb{C}(q)$ -vector spaces. Finally, Corollary 8.12 and (4.28) imply that this map is a morphism of algebras. \square

10.2. Equivariant Rickard complexes. From here through the end of Section 10, we **assume that \mathbb{K} is a field in which $2 \neq 0$** . (We will occasionally reemphasize this hypothesis when stating some of our major results.)

We now promote the Rickard complexes from Definition 9.11 to the equivariant category $(\mathcal{B}_m^n)^\tau$. For $1 \leq \bullet \leq m-1$, recall that

$$C(\beta_\bullet)\mathbb{1}_n = (\dots \xrightarrow{d} \mathbf{q}^{-k}\mathbf{t}^k \mathcal{F}_\bullet^{(n-k)} \mathcal{E}_\bullet^{(n-k)} \mathbb{1}_n \xrightarrow{d} \mathbf{q}^{-k-1}\mathbf{t}^{k+1} \mathcal{F}_\bullet^{(n-k-1)} \mathcal{E}_\bullet^{(n-k-1)} \mathbb{1}_n \xrightarrow{d} \dots).$$

Since the chain objects in this complex are all of the form $\mathcal{F}_\bullet^{(\ell)} \mathcal{E}_\bullet^{(\ell)} \mathbb{1}_n$, this actually defines a complex

$$C(\beta_\bullet) := (\dots \xrightarrow{d} \mathbf{q}^{-k}\mathbf{t}^k \mathbf{X}_\bullet^{(n-k)} \xrightarrow{d} \mathbf{q}^{-k-1}\mathbf{t}^{k+1} \mathbf{X}_\bullet^{(n-k-1)} \xrightarrow{d} \dots) \in \mathcal{C}(\mathcal{B}_m^n).$$

Similarly, the inverse complex

$$C(\beta_\bullet^{-1}) := (\dots \xrightarrow{d^\vee} \mathbf{q}^k \mathbf{t}^{-k} \mathbf{X}_\bullet^{(n-k)} \xrightarrow{d^\vee} \mathbf{q}^{k-1} \mathbf{t}^{-k+1} \mathbf{X}_\bullet^{(n-k+1)} \xrightarrow{d^\vee} \dots)$$

lies in \mathcal{B}_m^n . Henceforth, we denote the components of the corresponding differentials appearing in (9.10) and (9.11) as follows:

$$d_k: \mathbf{q}\mathbf{X}_\bullet^{(k)} \rightarrow \mathbf{X}_\bullet^{(k-1)} \quad \text{and} \quad d_k^\vee: \mathbf{q}\mathbf{X}_\bullet^{(k)} \rightarrow \mathbf{X}_\bullet^{(k+1)}.$$

Lemma 10.11. The differentials in the Rickard complexes induce the following degree-zero morphisms in $(\mathcal{B}_m^n)^\tau$:

$$d_k: (\mathbf{q}\mathbf{X}_\bullet^{(k)}, \mathbf{LC}_k) \rightarrow (\mathbf{X}_\bullet^{(k-1)}, (-1)^{n-k+1} \mathbf{LC}_{k-1})$$

and

$$d_k^\vee: (\mathbf{q}\mathbf{X}_\bullet^{(k)}, (-1)^{n+k} \mathbf{LC}_k) \rightarrow (\mathbf{X}_\bullet^{(k+1)}, \mathbf{LC}_{k+1}).$$

Proof. Using the definitions of τ and d , we have

$$\tau(d_k) = \begin{array}{c} k-1 \quad k-1 \\ \uparrow \quad \uparrow \\ \text{---} \text{---} \text{---} \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \mathbf{n} \quad \text{and} \quad \tau(d_k^\vee) = \begin{array}{c} k+1 \quad k+1 \\ \uparrow \quad \uparrow \\ \text{---} \text{---} \text{---} \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \mathbf{n}.$$

It suffices to show that

$$(10.1) \quad \tau(d_k) \circ \mathbf{LC}_k = (-1)^{n-k+1} \mathbf{LC}_{k-1} \circ d_k \quad \text{and} \quad (-1)^{n+k} \tau(d_k^\vee) \circ \mathbf{LC}_k = \mathbf{LC}_{k+1} \circ d_k^\vee.$$

For the former, we compute

$$(10.2) \quad \begin{array}{c} k-1 \quad k-1 \\ \uparrow \quad \uparrow \\ \text{---} \text{---} \text{---} \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \mathbf{n} \stackrel{(5.26)}{=} \begin{array}{c} k-1 \quad k-1 \\ \uparrow \quad \uparrow \\ \text{---} \text{---} \text{---} \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \mathbf{n} \stackrel{(5.18)}{=} \begin{array}{c} k-1 \quad k-1 \\ \uparrow \quad \uparrow \\ \text{---} \text{---} \text{---} \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \mathbf{n} \stackrel{(5.26)}{=} \begin{array}{c} k-1 \quad k-1 \\ \uparrow \quad \uparrow \\ \text{---} \text{---} \text{---} \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \mathbf{n} \stackrel{(5.15)}{=} - \sum_{p+q=2k-2} \begin{array}{c} k-1 \quad k-1 \\ \uparrow \quad \uparrow \\ \text{---} \text{---} \text{---} \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \mathbf{n} \\ \stackrel{(5.8), (5.20), (5.32)}{=} (-1)^{n-k+1} \begin{array}{c} k-1 \quad k-1 \\ \uparrow \quad \uparrow \\ \text{---} \text{---} \text{---} \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \mathbf{n} \stackrel{(5.26)}{=} (-1)^{n-k+1} \begin{array}{c} k-1 \quad k-1 \\ \uparrow \quad \uparrow \\ \text{---} \text{---} \text{---} \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \mathbf{n} = (-1)^{n-k+1} \begin{array}{c} k-1 \quad k-1 \\ \uparrow \quad \uparrow \\ \text{---} \text{---} \text{---} \\ \downarrow \quad \downarrow \\ k \quad k \end{array} \mathbf{n}.$$

The second equality in (10.1) follows similarly. \square

Remark 10.12. The functor τ extends from \mathcal{B}_m^n to the category of complexes $\mathcal{C}(\mathcal{B}_m^n)$. Explicitly, given a complex $C := (\dots \xrightarrow{d_Y} \mathbf{t}^k Y_k \xrightarrow{d_Y} \mathbf{t}^{k+1} Y_{k+1} \xrightarrow{d_Y} \dots)$ over \mathcal{B}_m^n , we can apply τ to each term in the complex, obtaining the complex

$$(10.3) \quad \tau(C) := (\dots \xrightarrow{\tau(d_Y)} \mathbf{t}^k \tau(Y_k) \xrightarrow{\tau(d_Y)} \mathbf{t}^{k+1} \tau(Y_{k+1}) \xrightarrow{\tau(d_Y)} \dots).$$

If $f \in \text{Hom}_{\text{Ch}(\mathcal{B}_m^n)}(C, C')$ is a chain map, then so is $\tau(f): \tau(C) \rightarrow \tau(C')$.

Remark 10.13. Lemma 10.11 can be read as saying that the collection of maps $\{(-1)^{\binom{k+1}{2}} \mathbf{LC}_{n-k}\}_{0 \leq k \leq n}$ gives a chain map $C(\beta_\bullet) \rightarrow \tau(C(\beta_\bullet))$, i.e. a map in $\text{Ch}(\mathcal{B}_m^n)$. For this, the relevant computation is that

$$(10.4) \quad \binom{k+1}{2} + \binom{k+2}{2} = (k+1)^2 \equiv k+1 \pmod{2}$$

which, by (10.1), ensures that $\tau(d_{n-k}) \circ (-1)^{\binom{k+1}{2}} \mathbf{LC}_{n-k} = (-1)^{\binom{k+2}{2}} \mathbf{LC}_{n-k-1} \circ d_{n-k}$.

Definition 10.14. For $1 \leq \bullet \leq m-1$, the \bullet^{th} *equivariant Rickard complex* is the chain complex

$$(10.5) \quad C^\tau(\beta_\bullet) := \left(\bigoplus_{k=0}^n \mathbf{q}^{-k} \mathbf{t}^k(\mathbf{X}_\bullet^{(n-k)}), (-1)^{\binom{n+1}{2} + \binom{k+1}{2}} \mathbf{LC}_{n-k}, d \right) \in \mathcal{C}((\mathcal{B}_m^n)^\tau)$$

and the \bullet^{th} *equivariant inverse Rickard complex* is

$$(10.6) \quad C^\tau(\beta_\bullet^{-1}) := \left(\bigoplus_{k=0}^n \mathbf{q}^k \mathbf{t}^{-k}(\mathbf{X}_\bullet^{(n-k)}), (-1)^{\binom{n+1}{2} + \binom{k+1}{2}} \mathbf{LC}_{n-k}, d^\vee \right) \in \mathcal{C}((\mathcal{B}_m^n)^\tau).$$

For a braid word $\beta_{i_1}^{\epsilon_1} \dots \beta_{i_r}^{\epsilon_r}$, the *equivariant Rickard complex* is

$$C^\tau(\beta_{i_1}^{\epsilon_1} \dots \beta_{i_r}^{\epsilon_r}) := C^\tau(\beta_{i_1}^{\epsilon_1}) \dots C^\tau(\beta_{i_r}^{\epsilon_r}) \in \mathcal{C}((\mathcal{B}_m^n)^\tau).$$

Observe that (10.5) indeed determines a complex in $\mathcal{C}((\mathcal{B}_m^n)^\tau)$, since

$$d: (\mathbf{X}_\bullet^{(n-k)}, (-1)^{\binom{n+1}{2} + \binom{k+1}{2}} \mathbf{LC}_{n-k}) \rightarrow (\mathbf{X}_\bullet^{(n-k-1)}, (-1)^{\binom{n+1}{2} + \binom{k+2}{2}} \mathbf{LC}_{n-k-1})$$

is a morphism in $(\mathcal{B}_m^n)^\tau$ if and only if

$$d: (\mathbf{X}_\bullet^{(n-k)}, \mathbf{LC}_{n-k}) \rightarrow (\mathbf{X}_\bullet^{(n-k-1)}, (-1)^{\binom{k+1}{2} + \binom{k+2}{2}} \mathbf{LC}_{n-k-1})$$

is such a morphism. This holds by Lemma 10.11, using (10.4). Similarly, (10.6) gives a complex in $\mathcal{C}((\mathcal{B}_m^n)^\tau)$. Our conventions (e.g. the choice of global sign $(-1)^{\binom{n+1}{2}}$) are such that both complexes have the monoidal unit $(\mathbf{X}_\bullet^{(0)}, \mathbf{LC}_0) = (\mathbb{1}_n, \text{id}_{\mathbb{1}_n})$ appearing in the extremal homological degree.

Although we know that the complexes $C(\beta_{i_1}^{\epsilon_1} \dots \beta_{i_r}^{\epsilon_r})$ satisfy the braid relations in $\mathcal{K}(\mathcal{B}_m^n)$, at this point it is not obvious that the equivariant Rickard complexes satisfy the braid relations in $\mathcal{K}((\mathcal{B}_m^n)^\tau)$, or even that (10.5) and (10.6) are homotopy inverses. In the following subsection, we take a detour to prove a general result that will be used to establish these facts.

10.3. Lifting to equivariant homotopy equivalences. For this section, let \mathcal{A} be a \mathbb{K} -linear category equipped with an involution σ , as defined in Remark 7.32. Let $\mathbf{U}: \mathcal{A}^\sigma \rightarrow \mathcal{A}$ denote the “forgetful functor” which sends an equivariant object (X, φ) to the underlying object X and an equivariant morphism $f: (X, \varphi_X) \rightarrow (Y, \varphi_Y)$ to itself, now viewed as $f \in \text{Hom}(X, Y)$. This functor \mathbf{U} is faithful.

Recall from §2.2 that $\mathcal{C}(\mathcal{A})$ denotes the dg category of chain complexes over \mathcal{A} . In particular, the Hom-spaces in $\mathcal{C}(\mathcal{A})$ are themselves complexes. If $X = (\bigoplus_{i \in \mathbb{Z}} \mathbf{t}^i(X_i, \varphi_i), d_X)$ is a complex in $\mathcal{C}(\mathcal{A}^\sigma)$, then, applying the forgetful functor \mathbf{U} , we obtain a complex $\mathbf{U}(X) = (\bigoplus_{i \in \mathbb{Z}} \mathbf{t}^i X_i, d_X)$ in $\mathcal{C}(\mathcal{A})$.

Lemma 10.15. Let $X = (\bigoplus_{i \in \mathbb{Z}} \mathbf{t}^i(X_i, \varphi_i), d_X)$ and $Y = (\bigoplus_{i \in \mathbb{Z}} \mathbf{t}^i(Y_i, \psi_i), d_Y)$ be complexes in $\mathcal{C}(\mathcal{A}^\sigma)$. The map

$$\sigma_*: \mathrm{Hom}_{\mathcal{C}(\mathcal{A})}(\mathbf{U}(X), \mathbf{U}(Y)) \rightarrow \mathrm{Hom}_{\mathcal{C}(\mathcal{A})}(\mathbf{U}(X), \mathbf{U}(Y))$$

(given by applying σ_* degree-wise) is a chain map. That is, if D denotes the differential on the Hom complexes, then $D \circ \sigma_* = \sigma_* \circ D$.

Proof. Since $(\bigoplus_{i \in \mathbb{Z}} \mathbf{t}^i(X_i, \varphi_i), d_X)$ and $Y = (\bigoplus_{i \in \mathbb{Z}} \mathbf{t}^i(Y_i, \psi_i), d_Y)$ are complexes in $\mathcal{C}(\mathcal{A}^\sigma)$, their differentials are equivariant morphisms. Hence, Corollary 7.36 implies that both d_X and d_Y are fixed by σ_* . We thus compute

$$\begin{aligned} \sigma_*(D(f)) &= \sigma_*(d_Y \circ f) - (-1)^{|f|} \sigma_*(f \circ d_X) \\ &\stackrel{(7.7)}{=} (\sigma_* d_Y) \circ (\sigma_* f) - (-1)^{|f|} (\sigma_* f) \circ \sigma_*(d_X) \\ &= d_Y \circ (\sigma_* f) - (-1)^{|f|} (\sigma_* f) \circ d_X \\ &= D(\sigma_*(f)). \end{aligned} \quad \square$$

Next, given a morphism $f \in \mathrm{Hom}_{\mathcal{C}(\mathcal{A}^\sigma)}(X, Y)$, we can consider the corresponding morphism $\mathbf{U}(f) \in \mathrm{Hom}_{\mathcal{C}(\mathcal{A})}(\mathbf{U}(X), \mathbf{U}(Y))$. Note that if f is a chain map then so is $\mathbf{U}(f)$. Recall that the space of chain maps is denoted $\mathrm{Hom}_{\mathrm{Ch}(\mathcal{A})} \subset \mathrm{Hom}_{\mathcal{C}(\mathcal{A})}$.

Proposition 10.16. Let $X = (\bigoplus_{i \in \mathbb{Z}} \mathbf{t}^i(X_i, \varphi_i), d_X)$ and $Y = (\bigoplus_{i \in \mathbb{Z}} \mathbf{t}^i(Y_i, \psi_i), d_Y)$ be complexes in $\mathcal{C}(\mathcal{A}^\sigma)$ and let $f \in \mathrm{Hom}_{\mathrm{Ch}(\mathcal{A}^\sigma)}(X, Y)$. If $\mathbf{U}(f) \in \mathrm{Hom}_{\mathrm{Ch}(\mathcal{A})}(\mathbf{U}(X), \mathbf{U}(Y))$ is a homotopy equivalence, then f is a homotopy equivalence.

Proof. Suppose $\mathbf{U}(f)$ is a homotopy equivalence. Thus, there exists degree-zero $g \in \mathrm{Hom}_{\mathcal{C}(\mathcal{A})}(\mathbf{U}(Y), \mathbf{U}(X))$ such that $D(g) = 0$, and \mathbf{t} -degree -1 morphisms $h_X: \mathbf{U}(X) \rightarrow \mathbf{U}(X)$ and $h_Y: \mathbf{U}(Y) \rightarrow \mathbf{U}(Y)$ in $\mathcal{C}(\mathcal{A})$, such that

$$\mathrm{id}_X - g \circ f = D(h_X) \quad \text{and} \quad \mathrm{id}_Y - f \circ g = D(h_Y).$$

Set

$$G := \frac{1}{2}(g + \sigma_* g), \quad H_X := \frac{1}{2}(h_X + \sigma_* h_X), \quad H_Y := \frac{1}{2}(h_Y + \sigma_* h_Y)$$

and note that G , H_X , and H_Y are all morphisms in $\mathcal{C}(\mathcal{A}^\sigma)$. Using Lemma 10.15, we compute

$$D(G) = \frac{1}{2}D(g + \sigma_* g) = \frac{1}{2}(D(g) + D(\sigma_* g)) = \frac{1}{2}(D(g) + \sigma_*(D(g))) = 0$$

and that

$$\begin{aligned} \mathrm{id}_X - G \circ f &= \mathrm{id}_X - \frac{1}{2}(g + \sigma_* g) \circ f \\ &= \frac{1}{2}((\mathrm{id}_X - g \circ f) + (\mathrm{id}_X - (\sigma_* g) \circ f)) \\ &\stackrel{\text{Cor. 7.36}}{=} \frac{1}{2}((\mathrm{id}_X - g \circ f) + ((\sigma_* \mathrm{id}_X) - (\sigma_* g) \circ (\sigma_* f))) \\ &= \frac{1}{2}((\mathrm{id}_X - g \circ f) + \sigma_*(\mathrm{id}_X - g \circ f)) \\ &= \frac{1}{2}(D(h_X) + \sigma_*(D(h_X))) \\ &= \frac{1}{2}(D(h_X) + D(\sigma_* h_X)) \\ &= D\left(\frac{1}{2}(h_X + \sigma_* h_X)\right) \\ &= D(H_X). \end{aligned}$$

A similar computation shows that $\text{id}_Y - f \circ G = D(H_Y)$. Thus, f , G , H_X , and H_Y give the data of a homotopy equivalence in $\mathcal{C}(\mathcal{A}^\sigma)$. \square

10.4. Braid relations for equivariant Rickard complexes. We now prove the analogue of Theorem 9.12 for the equivariant Rickard complexes. In doing so, we will make use of the 2-functor $\Gamma: \text{Kar}(\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_m)) \rightarrow \mathcal{A}_n$ from §9.3. Since $\mathcal{B}_m^n \hookrightarrow \text{Kar}(\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_m))$, we can consider the 2-functor Γ restricted to \mathcal{B}_m^n , which we still denote by Γ . Observe that

$$\Gamma: \mathcal{B}_m^n \rightarrow \mathcal{A}_n$$

is a full and strongly monoidal, which follows from Proposition 9.17 and the observation that the monoidal unit $\mathbb{1}_n \in \mathcal{B}_m^n$ is, by definition, sent to the left regular representation of $H^*(Gr_n(\mathbb{C}^{2n}))$, which is the monoidal unit in \mathcal{A}_n . Recall that all indecomposable objects in \mathcal{B}_m^n other than (shifts of) $\mathbb{1}_n$ are sent to zero by Γ .

Given any (graded) commutative ring R , we can consider the monoidal category of finitely generated free (graded) R -modules \mathcal{A}_R . (The category \mathcal{A}_n is the special case when $R = H^*(Gr_n(\mathbb{C}^{2n}))$.) A ring automorphism of R determines an additive, monoidal endofunctor of \mathcal{A}_R , which is the identity on objects and is determined on morphisms by acting via the automorphism on $\text{End}_{\mathcal{A}_R}(R) \cong R$.

Definition 10.17. Define a monoidal involution of \mathcal{A}_n , which abusing notation we denote τ , to be the monoidal functor induced by the (graded) ring automorphism of $H^*(Gr_n(\mathbb{C}^{2n}))$ which acts on the basis of Schur polynomials by $\mathfrak{s}_\lambda(\mathbb{X}) \mapsto \mathfrak{s}_{\lambda^t}(\mathbb{X})$.

Lemma 10.18. We have the following equality of monoidal functors: $\Gamma \circ \tau = \tau \circ \Gamma$.

Proof. Immediate from the description of Γ in Proposition 9.17 and the description of τ in Definition 6.3. \square

Now, let $\underline{\beta}$ be a braid word and write the complex $C^\tau(\underline{\beta}) \in \mathcal{C}((\mathcal{B}_m^n)^\tau)$ as

$$(10.7) \quad C^\tau(\underline{\beta}) =: \left(\bigoplus_{j \in \mathbb{Z}} \mathfrak{t}^j (C_j(\underline{\beta}), \varphi_j(\underline{\beta})), d_{\underline{\beta}} \right).$$

It follows from Definition 10.14 that $C_j(\underline{\beta})$ is a direct sum of tensor products of shifts of objects of the form $\mathbf{X}_i^{(k)}$, and that $\varphi_j(\underline{\beta})$ is a (diagonal) matrix whose entries are tensor products of the morphisms $\pm \mathbf{LC}_k$. Further, Remark 10.13 implies that the collection of maps $\{\varphi_j(\underline{\beta})\}_{j \in \mathbb{Z}}$ gives a chain map $\varphi_{\underline{\beta}}: C(\underline{\beta}) \rightarrow \tau(C(\underline{\beta}))$ in $\mathcal{C}(\mathcal{B}_m^n)$.

Lemma 10.19. If $\underline{\beta}$ is a braid word, then $\Gamma(\mathbf{U}(\varphi_{\underline{\beta}})) = \text{id}_{(\mathfrak{q}^{-n} \mathfrak{t}^n)^{\epsilon(\underline{\beta})} \mathbb{1}_n}$.

Proof. Since Γ and \mathbf{U} are monoidal, it suffices to establish the result for $\underline{\beta} = \beta_i^{\pm 1}$. In this case, both $C(\beta_i^{\pm 1})$ and $\tau(C(\beta_i^{\pm 1}))$ are mapped by Γ to $\mathfrak{q}^{\mp n} \mathfrak{t}^{\pm n} \mathbf{X}_i^{(0)} = \mathfrak{q}^{\mp n} \mathfrak{t}^{\pm n} \mathbb{1}_n$, and the sole component of the chain map $\Gamma(\varphi_{\beta_i^{\pm 1}})$ is $(-1)^{\binom{n+1}{2} + \binom{n+1}{2}} \mathbf{LC}_0 = \text{id}_{\mathbb{1}_n}$. \square

Lemma 10.20. Let $\underline{\beta}$ and $\underline{\beta}'$ be two braid words for the same braid. If $f \in \text{Hom}_{\text{Ch}(\mathcal{B}_m^n)}(C(\underline{\beta}), C(\underline{\beta}'))$ is a chain map such that $\Gamma(f) = \text{id}_{(\mathfrak{q}^{-n} \mathfrak{t}^n)^{\epsilon(\underline{\beta})} \mathbb{1}_n}$, then the same is true for $\tau(f)$.

Proof. That $\tau(f)$ is a chain map follows from Remark 10.12. The result then follows from Lemma 10.18, since τ preserves the identity map of $\mathbb{1}_n$ both before and after applying Γ . \square

Pairing Lemmata 9.20, 10.19, and 10.20 with the results from §10.3, we now establish that the complexes $C^\tau(\beta_{i_1}^{\epsilon_1} \cdots \beta_{i_r}^{\epsilon_r})$ canonically satisfy the braid relations in $\mathcal{K}((\mathcal{B}_m^n)^\tau)$. Precisely, we have the following:

Theorem 10.21. Let \mathbb{K} be a field in which $2 \neq 0$. In $\mathcal{C}((\mathcal{B}_m^n)^\tau)$, there are homotopy equivalences

$$(10.8) \quad C^\tau(\beta_i\beta_{i+1}\beta_i) \simeq C^\tau(\beta_{i+1}\beta_i\beta_{i+1})$$

for $1 \leq i \leq m-2$ and

$$(10.9) \quad C^\tau(\beta_i)C^\tau(\beta_i^{-1}) \simeq (\mathbb{1}_n, \text{id}_{\mathbb{1}_n}) \simeq C^\tau(\beta_i^{-1})C^\tau(\beta_i)$$

for $1 \leq i \leq m-1$. These homotopy equivalences can be chosen so that their image under $\Gamma \circ \mathbf{U}$ is the identity map of the appropriate shift of $\mathbb{1}_n$. Consequently, given two braid words $\underline{\beta}$ and $\underline{\beta}'$ for the same braid, any two homotopy equivalences $C^\tau(\underline{\beta}) \simeq C^\tau(\underline{\beta}')$ that are given as compositions of (10.8) and (10.9) agree in $\mathcal{K}((\mathcal{B}_m^n)^\tau)$.

Proof. By Theorem 9.12, before accounting for equivariant structures, there are homotopy equivalences

$$f_i: C(\beta_i\beta_{i+1}\beta_i) \xrightarrow{\simeq} C(\beta_{i+1}\beta_i\beta_{i+1})$$

and

$$g_i: \mathbb{1}_n \xrightarrow{\simeq} C(\beta_i)C(\beta_i^{-1}), \quad g'_i: \mathbb{1}_n \xrightarrow{\simeq} C(\beta_i^{-1})C(\beta_i).$$

Moreover, using Lemma 9.20 we can choose f_i , g_i , and g'_i so that they each map to $\text{id}_{\mathbb{1}_n}$ under Γ .

Combining Lemma 10.19 and Lemma 10.20 with the definition of $\tau*$ from (7.6), we see that $\tau*f_i$, $\tau*g_i$, and $\tau*g'_i$ all also map to $\text{id}_{\mathbb{1}_n}$ under the quotient functor. It follows that the same is true for each of

$$f_i^\tau := \frac{1}{2}(f_i + \tau*f_i), \quad g_i^\tau := \frac{1}{2}(g_i + \tau*g_i), \quad g_i'^\tau := \frac{1}{2}(g'_i + \tau*g'_i).$$

Lemma 9.20 implies that each of f_i^τ , g_i^τ , and $g_i'^\tau$ are therefore homotopy equivalences in $\mathcal{C}(\mathcal{B}_m^n)$. Each of these maps is fixed by $\tau*$, so they give chain maps

$$(10.10) \quad f_i^\tau: C^\tau(\beta_i\beta_{i+1}\beta_i) \rightarrow C^\tau(\beta_{i+1}\beta_i\beta_{i+1}), \quad g_i^\tau: \mathbb{1}_n \rightarrow C^\tau(\beta_i)C^\tau(\beta_i^{-1}), \quad g_i'^\tau: \mathbb{1}_n \rightarrow C^\tau(\beta_i^{-1})C^\tau(\beta_i)$$

in $\mathcal{C}((\mathcal{B}_m^n)^\tau)$. Proposition 10.16 then shows that each of the maps in (10.10) is a homotopy equivalence.

The exact same argument as in the proof of Lemma 9.20 implies that

$$\text{Hom}_{\mathcal{K}((\mathcal{B}_m^n)^\tau)}(C^\tau(\underline{\beta}), C^\tau(\underline{\beta}')) \cong \mathbb{K}$$

when $\underline{\beta}$ and $\underline{\beta}'$ are words for the same braid. Thus one recovers an equivariant version of Lemma 9.20: if f is a chain map in this morphism space, and $\Gamma(\mathbf{U}(f)) = c \cdot \text{id}_{\mathbb{1}_n}$ for $c \in \mathbb{K}$, then either $c \neq 0$ and f is a homotopy equivalence, or $c = 0$ and f is nulhomotopic. Hence two chain maps f, g in this morphism space are homotopic if and only if they agree after applying $\Gamma \circ \mathbf{U}$. The final statement of the theorem is an immediate consequence. \square

Theorem 10.21 shows that we can canonically associate a complex over $(\mathcal{B}_m^n)^\tau$ to a braid β by choosing a braid word representative $\underline{\beta}$ and considering $C^\tau(\underline{\beta})$. We thus will slightly abuse notation and denote this complex as $C^\tau(\beta)$ moving forward.

10.5. Spin link homology. At last, we define our invariant. We proceed analogously to (9.13), and then establish that we indeed obtain an invariant of framed links $\mathcal{L} \subset S^3$. To simplify notation, we set

$$(\mathbb{1}_n, \pm) := (\mathbb{1}_n, \pm \text{id}_{\mathbb{1}_n}) \in (\mathcal{B}_m^n)^\tau.$$

For the duration, when we consider the $\tau*$ -action on Hom-spaces of the form $\text{Hom}_{\mathcal{B}_m^n}(\mathbb{1}_n, X)$, it will be assumed with respect to the equivariant structure $+\text{id}_{\mathbb{1}_n}$ on $\mathbb{1}_n$ (and whatever equivariant structure φ_X is currently being considered on X).

Recall from §2.2 that $\text{sVect}_{\mathbb{K}}^{\mathbb{Z}}$ denotes the category of \mathbb{Z} -graded super \mathbb{K} -vector spaces, and that we denote shift in super degree by \mathbf{s} . We will implicitly view $\text{Vect}_{\mathbb{K}}^{\mathbb{Z}} \subset \text{sVect}_{\mathbb{K}}^{\mathbb{Z}}$ as the full subcategory of super vector spaces concentrated in super degree zero.

Definition 10.22. Consider the additive functors $\mathcal{R}^+, \mathcal{R}^- : (\mathcal{B}_m^n)^\tau \rightarrow \text{Vect}_{\mathbb{K}}^{\mathbb{Z}}$ given by

$$(10.11) \quad \mathcal{R}^+(x) := \text{Hom}_{(\mathcal{B}_m^n)^\tau}((\mathbb{1}_n, +), x), \quad \mathcal{R}^-(x) := \text{Hom}_{(\mathcal{B}_m^n)^\tau}((\mathbb{1}_n, -), x)$$

and similarly denote the induced functors $\mathcal{R}^+, \mathcal{R}^- : \mathcal{K}((\mathcal{B}_m^n)^\tau) \rightarrow \mathcal{K}(\text{Vect}_{\mathbb{K}}^{\mathbb{Z}})$ given by applying (10.11) term-wise. Let

$$\{\beta\}_{2n+1, \mathcal{S}}^+ := \mathbf{q}^{-mn^2} \mathcal{R}^+(C^\tau(\beta)), \quad \{\beta\}_{2n+1, \mathcal{S}}^- := \mathbf{q}^{-mn^2} \mathcal{R}^-(C^\tau(\beta))$$

and set $\{\beta\}_{2n+1, \mathcal{S}} := \{\beta\}_{2n+1, \mathcal{S}}^+ \oplus \mathbf{s}\{\beta\}_{2n+1, \mathcal{S}}^- \in \mathcal{K}(\text{sVect}_{\mathbb{K}}^{\mathbb{Z}})$.

It is immediate from Theorem 10.21 that $\{\beta\}_{2n+1, \mathcal{S}} \in \mathcal{K}(\text{sVect}_{\mathbb{K}}^{\mathbb{Z}})$ is an invariant of braids $\beta \in \text{Br}_m$. We will establish that the assignment $\beta \mapsto \{\beta\}_{2n+1, \mathcal{S}}$ is further invariant under the Markov moves up to (appropriate) shift, and therefore is an invariant of the framed link \mathcal{L}_β given as the closure of β .

Our approach is as follows. Recall that the \mathfrak{sl}_{2n} link invariant $\llbracket \beta \mathbb{1}_n \rrbracket_{2n}$ comes from applying the functor $\text{Hom}(\mathbb{1}_n, -)$ to each term in the complex $C(\beta)$, yielding a complex whose chain group in degree j is $\text{Hom}_{\mathcal{B}_m^n}(\mathbb{1}_n, C_j(\beta))$. Using Corollary 7.36, one has

$$\text{Hom}_{(\mathcal{B}_m^n)^\tau}((\mathbb{1}_n, +), (C_j(\beta), \varphi_j(\beta))) = \text{Hom}_{\mathcal{B}_m^n}(\mathbb{1}_n, C_j(\beta))^\tau,$$

the subspace of invariants under the τ^* -action. Thus $\{\beta\}_{2n+1, \mathcal{S}}^+$ is a subcomplex of $\llbracket \beta \mathbb{1}_n \rrbracket_{2n}$, the subspace of τ^* -invariants. Similarly, $\{\beta\}_{2n+1, \mathcal{S}}^-$ is a subspace of $\llbracket \beta \mathbb{1}_n \rrbracket_{2n}$, the subcomplex of τ^* -anti-invariants (i.e. the isotypic component of the sign representation). Lemma 10.15 then implies that $\{\beta\}_{2n+1, \mathcal{S}}^\pm \subset \llbracket \beta \mathbb{1}_n \rrbracket_{2n}$ are in fact subcomplexes.

By Theorem 9.13, we know that $\llbracket \beta \mathbb{1}_n \rrbracket_{2n}$ satisfies the Markov moves, up to homotopy equivalence. We will observe that these homotopy equivalences restrict to $\{\beta\}_{2n+1, \mathcal{S}}^\pm$, and thus give Markov invariance for $\{\beta\}_{2n+1, \mathcal{S}}$.

We begin with a result concerning group actions on chain complexes of vector spaces.

Lemma 10.23. Let G be a finite group, and suppose $|G|$ is invertible in \mathbb{K} . Let $(X, d_X), (Y, d_Y) \in \mathcal{C}(\text{Vect}_{\mathbb{K}}^{\mathbb{Z}})$ be complexes of (graded) \mathbb{K} -vector spaces that admit a \mathbb{K} -linear action of G , i.e. the chain groups X_i and Y_i are G -representations and the differentials d_X and d_Y commute with the G -action. Suppose that $f : (X, d_X) \xrightarrow{\sim} (Y, d_Y)$ is a homotopy equivalence such that $f|_{X_i} : X_i \rightarrow Y_i$ is a morphism of G -representations. Then, f restricts to a homotopy equivalence between the G -invariant subcomplexes (X^G, d_X) and (Y^G, d_Y) of (X, d_X) and (Y, d_Y) . If f is an isomorphism of chain complexes in $\text{Ch}(\text{Vect}_{\mathbb{K}}^{\mathbb{Z}})$, then the restriction is as well.

Proof. Throughout, we let $Z = X$ or Y . First, observe that $(Z^G, d_Z) \subset (Z, d_Z)$ is indeed a subcomplex, where

$$Z_i^G := \{z \in Z_i \mid g \cdot z = z \ \forall g \in G\},$$

since $d_Z \circ (g \cdot) = (g \cdot) \circ d_Z$ for all $g \in G$. Since $f|_{X_i}$ is assumed to be a morphism of G -representations, $f|_{X^G} : (X^G, d_X) \rightarrow (Y^G, d_Y)$ is a chain map, so it remains to construct its homotopy inverse.

The argument is similar to the proof of Proposition 10.16. Let $k : (Y, d_Y) \rightarrow (X, d_X)$ be the homotopy inverse to f , so there exist h_X and h_Y such that

$$(10.12) \quad \text{id}_X - k \circ f = d_X \circ h_X + h_X \circ d_X \quad \text{and} \quad \text{id}_Y - f \circ k = d_Y \circ h_Y + h_Y \circ d_Y.$$

Set

$$k^G := \frac{1}{|G|} \sum_{g \in G} (g \cdot) \circ k, \quad h_X^G := \frac{1}{|G|} \sum_{g \in G} (g \cdot) \circ h_X, \quad h_Y^G := \frac{1}{|G|} \sum_{g \in G} (g \cdot) \circ h_Y$$

and note that $k^G \in \text{Hom}_{\text{Ch}(\text{Vect}_{\mathbb{K}}^{\mathbb{Z}})}((Y^G, d_Y), (X^G, d_X))$ and $h_Z^G \in \text{End}_{\text{e}(\text{Vect}_{\mathbb{K}}^{\mathbb{Z}})}((Z^G, d_Z))$. Averaging (10.12) over G (and using $f \circ (g \cdot) = (g \cdot) \circ f$ and $d_Z \circ (g \cdot) = (g \cdot) \circ d_Z$) gives

$$\left(\frac{1}{|G|} \sum_{g \in G} (g \cdot) \circ \text{id}_X \right) - k^G \circ f = d_X \circ h_X^G + h_X^G \circ d_X \quad \text{and} \quad \left(\frac{1}{|G|} \sum_{g \in G} (g \cdot) \circ \text{id}_Y \right) - f \circ k^G = d_Y \circ h_Y^G + h_Y^G \circ d_Y.$$

Since $\left(\frac{1}{|G|} \sum_{g \in G} (g \cdot) \circ \text{id}_Z \right) |_{Z^G} = \text{id}_{Z^G}$, this shows that $f|_{X^G}: (X^G, d_X) \rightarrow (Y^G, d_Y)$ is a homotopy equivalence.

For the final statement, note that the statement that f is an isomorphism is equivalent to being able to choose maps so that (10.12) is satisfied with $h_X = 0$ and $h_Y = 0$. It then follows that $h_X^G = 0$ and $h_Y^G = 0$, so $f|_{X^G}: (X^G, d_X) \rightarrow (Y^G, d_Y)$ is an isomorphism of chain complexes as well. \square

Remark 10.24. When $G = \mathbb{Z}/2$, Lemma 10.23 can also be applied to sign components (X^{sgn}, d_X) and (Y^{sgn}, d_Y) , rather than trivial components (X^G, d_X) and (Y^G, d_Y) . This is because, for an involutive linear map σ , the anti-invariants of σ agree with the invariants of $-\sigma$. More generally though, an analogous argument to the proof of Lemma 10.23 (using other central idempotents in the group algebra) establishes homotopy equivalences (and isomorphisms) between subcomplexes of isotypic components of (X, d_X) and (Y, d_Y) .

Proposition 10.25. The complex $\{\beta\}_{2n+1, \mathcal{S}}$ satisfies the first Markov move, up to isomorphism in $\text{Ch}(\text{sVect}_{\mathbb{K}}^{\mathbb{Z}})$. Explicitly, given braids $\beta, \beta' \in \text{Br}_m$, there is an isomorphism $\{\beta\beta'\}_{2n+1, \mathcal{S}} \cong \{\beta'\beta\}_{2n+1, \mathcal{S}}$ of chain complexes of super \mathbb{K} -vector spaces.

Proof. By Theorem 9.13, there is a homotopy equivalence of chain complexes

$$(10.13) \quad f: \llbracket \beta\beta' \rrbracket_{2n} \xrightarrow{\cong} \llbracket \beta'\beta \rrbracket_{2n}.$$

In fact, this is an isomorphism of chain complexes, that we now explicitly describe.

In the notation of (10.7), the term of the complex $\llbracket \beta\beta' \rrbracket_{2n}$ in homological degree k takes the form

$$\mathbf{q}^{-mn^2} \bigoplus_{i+j=k} \text{Hom}_{\mathcal{B}_m^n}(\mathbb{1}_{\mathbf{n}}, C_i(\beta)C_j(\beta')).$$

By definition, each of $C_i(\beta)$ and $C_j(\beta')$ can themselves be written as

$$C_i(\beta) = \bigoplus_{p=1}^{\ell_i} C_{i,p}, \quad C_j(\beta') = \bigoplus_{r=1}^{\ell'_j} C'_{j,r}$$

where each $C_{i,p}$ and $C'_{j,r}$ is a tensor product of objects of the form $\mathbf{X}_s^{(t)}$. The isomorphism (10.13) is then given summand-wise as follows:

$$(10.14) \quad \text{Hom}_{\mathcal{B}_m^n}(\mathbb{1}_{\mathbf{n}}, C_{i,p}C'_{j,r}) \xrightarrow{f} \text{Hom}_{\mathcal{B}_m^n}(\mathbb{1}_{\mathbf{n}}, C'_{j,r}C_{i,p})$$

Here, the sign $(-1)^{ij}$ ensures that this indeed gives a chain map, since the differential on the complex $C_i(\beta)C_j(\beta')$ follows the usual Koszul sign rule for a tensor product of complexes.

As noted above, $\{\beta\beta'\}_{2n+1,\mathcal{S}}^+$ and $\{\beta'\beta\}_{2n+1,\mathcal{S}}^+$ are precisely the $\mathbb{Z}/2$ -invariant subcomplexes for the τ^* -actions on $[[\beta\beta']_{2n}]$ and $[[\beta'\beta]_{2n}]$ (respectively), while $\{\beta\beta'\}_{2n+1,\mathcal{S}}^-$ and $\{\beta'\beta\}_{2n+1,\mathcal{S}}^-$ are the sign component subcomplexes. By Lemma 10.23 and Remark 10.24, it therefore suffices to show that the assignment (10.14) is a morphism of $\mathbb{Z}/2$ -representations, i.e. that $f \circ \tau^* = \tau^* \circ f$. For this, we compute

$$f \left(\tau^* \left(\begin{array}{c} \overbrace{C_{i,p}} \quad \overbrace{C'_{j,r}} \\ \uparrow \cdots \uparrow \quad \uparrow \cdots \uparrow \\ \xi \end{array} \right) \mathbf{n} \right) = f \left(\begin{array}{c} \overbrace{C_{i,p}} \quad \overbrace{C'_{j,r}} \\ \downarrow \cdots \downarrow \quad \downarrow \cdots \downarrow \\ \tau(\xi) \end{array} \right) \mathbf{n} = (-1)^{ij} \left(\begin{array}{c} \overbrace{C'_{j,r}} \quad \overbrace{C_{i,p}} \\ \uparrow \cdots \uparrow \\ \tau(\xi) \end{array} \right) \mathbf{n},$$

which agrees with the following calculation.

$$\tau^* f \left(\begin{array}{c} \overbrace{C_{i,p}} \quad \overbrace{C'_{j,r}} \\ \uparrow \cdots \uparrow \quad \uparrow \cdots \uparrow \\ \xi \end{array} \right) \mathbf{n} = (-1)^{ij} \tau^* \left(\begin{array}{c} \overbrace{C'_{j,r}} \quad \overbrace{C_{i,p}} \\ \uparrow \cdots \uparrow \\ \xi \end{array} \right) \mathbf{n} = (-1)^{ij} \left(\begin{array}{c} \overbrace{C'_{j,r}} \quad \overbrace{C_{i,p}} \\ \downarrow \cdots \downarrow \\ \tau(\xi) \end{array} \right) \mathbf{n}$$

The last equality is most easily seen by applying (6.38) and (6.39), which shows that, in the weights appearing in the computation, τ does not contribute any signs when acting on the caps/cups. \square

Next, we establish invariance of $\{\beta\}_{2n+1,\mathcal{S}}$ under the second Markov move, up to shift. In type A , Markov II invariance amounts to a homotopy equivalence

$$(10.15) \quad [[\beta \mathbb{1}_{\mathbf{n}}]_{2n}] \simeq \mathbf{q}^{\mp n(n+1)} \mathbf{t}^{\pm n} [[\beta_m^{\pm 1} \beta \mathbb{1}_{\mathbf{n}}]_{2n}]$$

where $\beta \in \text{Br}_m$ (and $\beta_m^{\pm 1} \beta \in \text{Br}_{m+1}$). Our approach, as in the proof of Proposition 10.25, is to prove that the chain map realizing (10.15) is a morphism of $\mathbb{Z}/2$ -representations for the relevant τ^* -actions. Unfortunately, the homotopy equivalence (10.15) is not explicitly given in the literature, but rather is typically proven indirectly; see e.g. [75, Lemma 14.7]. However, using results of Wu [75, Section 14.2] concerning the “topologically local” nature of the known Markov II invariance, we are able to explicitly write down²⁵ the relevant homotopy equivalence and check the $\mathbb{Z}/2$ -equivariance.

Let us explain the general outline in more detail. There are two essential ways in which $[[\beta \mathbb{1}_{\mathbf{n}}]_{2n}]$ a priori differs from $[[\beta_m^{\pm 1} \beta \mathbb{1}_{\mathbf{n}}]_{2n}]$. The first is obvious: the complex $C(\beta_m^{\pm 1} \beta)$ is obtained from $C(\beta)$ by tensoring with an extra factor $C(\beta_m^{\pm 1})$. The second is more subtle: the computation of Hom spaces takes place in different categories (associated to m and $m+1$, respectively). In particular, the endomorphism ring of $\mathbb{1}_{\mathbf{n}}$ in \mathcal{B}_{m+1}^n has an extra alphabet’s worth of symmetric functions $\text{Sym}(\mathbb{X}_{m+1})$, and therefore an extra tensor factor of $H^*(\text{Gr}_n(\mathbb{C}^{2n}))$, compared to the endomorphism ring in \mathcal{B}_m^n .

In [75, Lemma 14.8], Wu actually proves a stronger result than the homotopy equivalence (10.15). His results imply that²⁶ given any object $X \in \mathcal{B}_m^n$, there is a homotopy equivalence

$$(10.16) \quad [[\beta_m^{\pm 1} X]_{2n}] \xrightarrow{\simeq} \mathbf{q}^{\mp n(n+1)} \mathbf{t}^{\pm n} [[X]_{2n}].$$

²⁴Recall that $\beta_m^{\pm 1} \beta \in \text{Br}_{m+1}$ is obtained by sending $\beta \in \text{Br}_m$ to its image after applying the standard inclusion $\text{Br}_m \hookrightarrow \text{Br}_{m+1}$, and then multiplying with the new Artin generator or its inverse.

²⁵The explicit form of this homotopy equivalence may be of independent interest, even in the setting of type A link homologies.

²⁶This is the content, in our present setup, of the phrase “knotted MOY graphs” in Wu’s Lemma 14.8.

(Here, we slightly abuse notation in setting $\llbracket X \rrbracket_{2n} := \mathbf{q}^{-mn^2} \text{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_m)}(\mathbb{1}_n, X)$ for $X \in \mathcal{B}_m^n$.) Below, we explicitly construct a chain map for this homotopy equivalence. By construction, our chain map is natural in X , hence these chain maps assemble into the desired homotopy equivalence $\llbracket \beta_m^{\pm 1} C \rrbracket_{2n} \xrightarrow{\sim} \mathbf{q}^{\mp n(n+1)} \mathbf{t}^{\pm n} \llbracket C \rrbracket_{2n}$ for any complex $C \in \mathcal{C}(\mathcal{B}_m^n)$.

We now proceed with the argument in detail, which uses separate constructions of the chain map in (10.16) for β_m and β_m^{-1} . We emphasize that our arguments do not constitute a re-proof of Markov II invariance for \mathfrak{sl}_{2n} link homology, as we rely on [75] for the existence of the homotopy equivalence (10.16).

Proposition 10.26. The complex $\{\beta\}_{2n+1, \mathcal{S}}$ satisfies the positive second Markov move, up to homotopy equivalence and degree shift. Explicitly, given $\beta \in \text{Br}_m$, there is a homotopy equivalence $\mathbf{q}^{-n(n+1)} \mathbf{t}^n \{\beta\}_{2n+1, \mathcal{S}} \simeq \{\beta_m \beta\}_{2n+1, \mathcal{S}}$ of chain complexes of super \mathbb{K} -vector spaces.

Proof. As discussed above, given any object $X \in \mathcal{B}_m^n$, [75, Lemma 14.8] implies that there is a homotopy equivalence

$$(10.17) \quad f: \llbracket \beta_m X \rrbracket_{2n} \xrightarrow{\sim} \mathbf{q}^{-n(n+1)} \mathbf{t}^n \llbracket X \rrbracket_{2n}.$$

We will describe the homotopy equivalence f explicitly, in such a way that it is natural in X .

For this, consider

$$(10.18) \quad \begin{aligned} \llbracket \beta_m X \rrbracket_{2n} &= \mathbf{q}^{-(m+1)n^2} \left(\text{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_n, \mathcal{F}_m^{(n)} \mathcal{E}_m^{(n)} X) \rightarrow \dots \right. \\ &\quad \left. \dots \rightarrow \mathbf{q}^{1-n} \mathbf{t}^{n-1} \text{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_n, \mathcal{F}_m \mathcal{E}_m X) \xrightarrow{\delta} \mathbf{q}^{-n} \mathbf{t}^n \text{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_n, X) \right) \end{aligned}$$

and (as indicated) let δ denote the final differential in this complex. Given that

$$\mathbf{q}^{-n(n+1)} \mathbf{t}^n \llbracket X \rrbracket_{2n} = \mathbf{q}^{-mn^2 - n(n+1)} \mathbf{t}^n \text{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_m)}(\mathbb{1}_n, X)$$

is a chain complex of graded \mathbb{K} -vector spaces that is non-zero only in homological degree n , the homotopy equivalence (10.17) can be described by finding a surjective, degree-zero, \mathbb{K} -linear map

$$(10.19) \quad f: \text{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_n, X) \twoheadrightarrow \text{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_m)}(\mathbb{1}_n, X)$$

such that $\text{Im}(\delta) \subset \ker(f)$. Indeed, by (10.17) we know that

$$\mathbf{q}^{-n} \mathbf{t}^n \text{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_n, X) / \text{Im}(\delta) \cong \mathbf{q}^{-n} \mathbf{t}^n \text{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_m)}(\mathbb{1}_n, X)$$

thus such an f will necessarily be a quasi-isomorphism of bounded complexes of \mathbb{Z} -graded \mathbb{K} -vector spaces, hence a homotopy equivalence.

To construct f , we examine $\text{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_n, X)$. We know by [34, Proposition 3.11] and (5.8) that $\text{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_n, X)$ is spanned by morphisms of the form

$$(10.20) \quad \begin{array}{c} X \\ \downarrow \\ \boxed{\xi}^n \\ \uparrow \\ \boxed{p(\mathbb{X}_{m+1})} \end{array}$$

with $p \in \text{Sym}(\mathbb{X})$ and $\xi \in \text{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_m)}(\mathbb{1}_n, X)$. Here, any new bubbles in $\text{Sym}(\mathbb{X}_1 | \dots | \mathbb{X}_m)$ have been included as part of ξ , which is why only the last alphabet \mathbb{X}_{m+1} remains. In fact, we can winnow (10.20) down to a basis for $\text{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_n, X)$. The map

$$(10.21) \quad H^*(\text{Gr}_n(\mathbb{C}^{2n})) \otimes_{\mathbb{K}} \text{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_m)}(\mathbb{1}_n, X) \rightarrow \text{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_n, X)$$

sending $p(\mathbb{X}) \otimes \xi$ to the morphism in (10.20) is surjective, and Proposition 9.10 implies that

$$\begin{aligned} \dim_{\mathbf{q}}(\mathrm{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_{\mathbf{n}}, X)) &= q^{n^2} \binom{2n}{n} \dim_{\mathbf{q}}(\mathrm{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{n}}, X)) \\ &= \dim_{\mathbf{q}}\left(H^*(\mathrm{Gr}_n(\mathbb{C}^{2n})) \otimes_{\mathbb{K}} \mathrm{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{n}}, X)\right). \end{aligned}$$

Thus, (10.21) is an isomorphism, and we can consider a basis in which the $p(\mathbb{X})$ range amongst Schur polynomials associated to partitions contained in an $n \times n$ box.

Now, we define f by the formula

$$(10.22) \quad f \left(\begin{array}{c} X \\ \boxed{p(\mathbb{X}_{m+1})} \quad \downarrow \quad \mathbf{n} \\ \boxed{\xi} \end{array} \right) := \begin{array}{c} X \\ \boxed{p(\mathbb{X}_m)} \quad \downarrow \quad \mathbf{n} \\ \boxed{\xi} \end{array} \in \mathrm{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{n}}, X),$$

which identifies the alphabets \mathbb{X}_{m+1} and \mathbb{X}_m . By the above discussion, this is a well-defined, degree-zero, \mathbb{K} -linear map $\mathrm{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_{\mathbf{n}}, X) \rightarrow \mathrm{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{n}}, X)$ which is surjective and satisfies the desired relation $f \circ \tau_* = \tau_* \circ f$.

Next, we establish that $\mathrm{Im}(\delta) \subset \ker(f)$. For this, we first claim that $\mathrm{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_{\mathbf{n}}, \mathcal{F}_m \mathcal{E}_m X)$ is spanned by morphisms of the form

$$(10.23) \quad \begin{array}{c} X \\ \curvearrowright \quad \downarrow \quad \mathbf{n} \\ \bullet \quad \boxed{\eta} \end{array}$$

with $\eta \in \mathrm{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_{\mathbf{n}}, X)$ and $r \geq 0$. Here, we use **magenta** to depict the final Dynkin label m . The reason that morphisms of the form (10.23) constitute a spanning set is that the color **magenta** does not appear within the object X , so the basis in [34, Proposition 3.11] will separate **magenta** from the other colors.

The differential δ takes a morphism as in (10.23) and post-composes with an anti-clockwise **magenta** cap. Thus $\mathrm{Im}(\delta)$ is spanned by elements of the form

$$\begin{array}{c} X \\ \curvearrowright \quad \downarrow \quad \mathbf{n} \\ \bullet \quad \boxed{\eta} \end{array} \stackrel{(5.7)}{=} \begin{array}{c} X \\ \curvearrowright \quad \downarrow \quad \mathbf{n} \\ \blacklozenge_{+r+1} \quad \bullet \quad \boxed{\eta} \end{array} \stackrel{(5.8)}{=} (-1)^n \begin{array}{c} X \\ \boxed{h_{r+1}(\mathbb{X}_{m+1} - \mathbb{X}_m)} \quad \downarrow \quad \mathbf{n} \\ \boxed{\eta} \end{array}$$

with $\eta \in \mathrm{Hom}_{\mathcal{S}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_{\mathbf{n}}, X)$ and $r \geq 0$. We thus compute

$$f \left(\begin{array}{c} X \\ \boxed{h_{r+1}(\mathbb{X}_{m+1} - \mathbb{X}_m)} \quad \downarrow \quad \mathbf{n} \\ \boxed{\eta} \end{array} \right) = \begin{array}{c} X \\ \boxed{h_{r+1}(\mathbb{X}_m - \mathbb{X}_m)} \quad \downarrow \quad \mathbf{n} \\ \boxed{\eta} \end{array} f \left(\begin{array}{c} X \\ \downarrow \quad \mathbf{n} \\ \boxed{\eta} \end{array} \right) = 0.$$

Finally, we conclude as sketched above. The homotopy equivalence (10.17) is natural in X , so by applying it to every chain object we obtain a homotopy equivalence

$$f: \llbracket \beta_m C_j(\beta) \mathbb{1}_{\mathbf{n}} \rrbracket_{2n} \xrightarrow{(10.17)} \mathbf{q}^{-n(n+1)} \mathbf{t}^n \llbracket C_j(\beta) \mathbb{1}_{\mathbf{n}} \rrbracket_{2n}$$

that satisfies $f \circ \tau_* = \tau_* \circ f$. Thus, using Lemma 10.23 and Remark 10.24 as in the proof of Proposition 10.25, we deduce that $\mathbf{q}^{-n(n+1)} \mathbf{t}^n \{\beta\}_{2n+1, \mathcal{S}} \simeq \{\beta_m \beta\}_{2n+1, \mathcal{S}}$. \square

Proposition 10.27. The complex $\{\beta\}_{2n+1, \mathcal{S}}$ satisfies the negative second Markov move, up to homotopy equivalence and degree shift. Explicitly, given $\beta \in \mathrm{Br}_m$, there is a homotopy equivalence $\mathbf{q}^{n(n+1)} \mathbf{t}^{-n} \{\beta\}_{2n+1, \mathcal{S}} \simeq \{\beta_m^{-1} \beta\}_{2n+1, \mathcal{S}}$ of chain complexes of super \mathbb{K} -vector spaces.

Proof. The proof again follows that of Propositions 10.25 and 10.26. Namely, we explicitly describe the homotopy equivalence

$$(10.24) \quad f: \mathbf{q}^{n(n+1)}\mathbf{t}^{-n} \llbracket \beta \rrbracket_{2n} \xrightarrow{\cong} \llbracket \beta_m^{-1} \beta \rrbracket_{2n},$$

observe that $f \circ \tau_* = \tau_* \circ f$, and then deduce the result from Lemma 10.23 and Remark 10.24. Again, the chain map f in (10.24) is induced from a map defined for each $X \in \mathcal{B}_m^n$ in [75, Lemma 14.8] that is natural in X :

$$(10.25) \quad f: \mathbf{q}^{n(n+1)}\mathbf{t}^{-n} \llbracket X \rrbracket_{2n} \xrightarrow{\cong} \llbracket \beta_m^{-1} X \rrbracket_{2n}.$$

Now, we let δ denote the first differential in

$$(10.26) \quad \begin{aligned} \llbracket \beta_m^{-1} X \rrbracket_{2n} = \mathbf{q}^{-(m+1)n^2} \left(\mathbf{q}^n \mathbf{t}^{-n} \mathrm{Hom}_{\tilde{\mathfrak{S}}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_n, X) \xrightarrow{\delta} \mathbf{q}^{n-1} \mathbf{t}^{1-n} \mathrm{Hom}_{\tilde{\mathfrak{S}}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_n, \mathcal{F}_m \mathcal{E}_m X) \rightarrow \right. \\ \left. \cdots \rightarrow \mathrm{Hom}_{\tilde{\mathfrak{S}}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_n, \mathcal{F}_m^{(n)} \mathcal{E}_m^{(n)} X) \right). \end{aligned}$$

Since

$$\mathbf{q}^{n(n+1)}\mathbf{t}^{-n} \llbracket X \rrbracket_{2n} = \mathbf{q}^{-mn^2+n(n+1)}\mathbf{t}^{-n} \mathrm{Hom}_{\tilde{\mathfrak{S}}_q^{\leq 2n}(\mathfrak{gl}_m)}(\mathbb{1}_n, X)$$

is a chain complex of graded \mathbb{K} -vector spaces that is non-zero only in homological degree $-n$, the homotopy equivalence (10.25) can be described by finding an injective, degree-zero, \mathbb{K} -linear map

$$(10.27) \quad f: \mathbf{q}^{2n^2} \mathrm{Hom}_{\tilde{\mathfrak{S}}_q^{\leq 2n}(\mathfrak{gl}_m)}(\mathbb{1}_n, X) \hookrightarrow \mathrm{Hom}_{\tilde{\mathfrak{S}}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_n, X)$$

such that $\mathrm{Im}(f) \subset \ker(\delta)$. Indeed, by (10.25) we know that

$$\ker \left(\delta: \mathrm{Hom}_{\tilde{\mathfrak{S}}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_n, X) \rightarrow \mathbf{q}^{-1} \mathrm{Hom}_{\tilde{\mathfrak{S}}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_n, \mathcal{F}_m \mathcal{E}_m X) \right) \cong \mathbf{q}^{2n^2} \mathrm{Hom}_{\tilde{\mathfrak{S}}_q^{\leq 2n}(\mathfrak{gl}_m)}(\mathbb{1}_n, X)$$

so such a map must be a homotopy equivalence.

Let n^n denote the partition of n^2 whose Young diagram is an $n \times n$ box. Given $\xi \in \mathrm{Hom}_{\tilde{\mathfrak{S}}_q^{\leq 2n}(\mathfrak{gl}_m)}(\mathbb{1}_n, X)$, define f in (10.27) by

$$f \left(\begin{array}{c} X \\ \downarrow \\ \mathbb{1}_n \\ \downarrow \\ \xi \end{array} \right) := \boxed{\mathfrak{s}_{n^n}(\mathbb{X}_m + \mathbb{X}_{m+1})} \begin{array}{c} X \\ \downarrow \\ \mathbb{1}_n \\ \downarrow \\ \xi \end{array}.$$

Since the partition n^n is self-transpose, we have $f \circ \tau_* = \tau_* \circ f$ by (6.3).

The salient properties of f will follow from the expansion

$$(10.28) \quad \mathfrak{s}_{n^n}(\mathbb{X}_m + \mathbb{X}_{m+1}) = \sum_{\lambda \subseteq n^n} \mathfrak{s}_{\lambda^c}(\mathbb{X}_m) \mathfrak{s}_{\lambda}(\mathbb{X}_{m+1}).$$

Being the cohomology ring of a compact orientable manifold, $H^*(\mathrm{Gr}_n(\mathbb{C}^{2n}))$ has a \mathbb{K} -linear functional I which picks out the top cohomology class (corresponding to integration). Explicitly, this map I satisfies

$$I(\mathfrak{s}_{\lambda}) = \begin{cases} 1 & \text{if } \lambda = n^n \\ 0 & \text{else.} \end{cases}$$

Applying I to elements in $H^*(\mathrm{Gr}_n(\mathbb{C}^{2n})) \cong \mathrm{Sym}(\mathbb{X}_{m+1})/\sim$ will take an element in $H^*(\mathrm{Gr}_n(\mathbb{C}^{2n}))^{\otimes 2} \cong \mathrm{Sym}(\mathbb{X}_m | \mathbb{X}_{m+1})/\sim$ to an element of $H^*(\mathrm{Gr}_n(\mathbb{C}^{2n})) \cong \mathrm{Sym}(\mathbb{X}_m)/\sim$.

Applying this map to (the class of) $\mathfrak{s}_{n^n}(\mathbb{X}_m + \mathbb{X}_{m+1})$ will yield 1, by (10.28). Consequently, post-composing the isomorphism

$$\mathrm{Hom}_{\tilde{\mathfrak{S}}_q^{\leq 2n}(\mathfrak{gl}_{m+1})}(\mathbb{1}_n, X) \stackrel{(10.21)}{\cong} H^*(\mathrm{Gr}_n(\mathbb{C}^{2n})) \otimes_{\mathbb{K}} \mathrm{Hom}_{\tilde{\mathfrak{S}}_q^{\leq 2n}(\mathfrak{gl}_m)}(\mathbb{1}_n, X)$$

with the map $I \otimes \text{id}: H^*(\text{Gr}_n(\mathbb{C}^{2n})) \otimes_{\mathbb{K}} \text{Hom}_{\mathfrak{S}_q^{\leq 2n}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{n}}, X) \rightarrow \mathbf{q}^{2n^2} \text{Hom}_{\mathfrak{S}_q^{\leq 2n}(\mathfrak{gl}_m)}(\mathbb{1}_{\mathbf{n}}, X)$, one obtains a left inverse to f . Thus, f is injective, as desired.

The map δ in (10.26) is given by post-composition with an anti-clockwise magenta cup (we again use magenta to depict the final Dynkin label m). We compute that

$$(10.29) \quad \delta \circ f \left(\begin{array}{c} X \\ | \\ \xi \\ \square \\ \mathbf{n} \end{array} \right) = \begin{array}{c} \text{magenta cup} \\ \text{magenta box } \mathfrak{s}_{n^{\mathbf{n}}}(\mathbb{X}_m + \mathbb{X}_{m+1}) \\ \text{magenta cup} \\ \begin{array}{c} X \\ | \\ \xi \\ \square \\ \mathbf{n} \end{array} \end{array} \stackrel{(5.13)}{=} \begin{array}{c} \text{magenta cup} \\ \text{magenta box } \mathfrak{s}_{n^{\mathbf{n}}}(\mathbb{X}_m + \mathbb{X}_{m+1}) \\ \text{magenta cup} \\ \begin{array}{c} X \\ | \\ \xi \\ \square \\ \mathbf{n} \end{array} \end{array} .$$

In weight $\mathbf{n} + \alpha_m$, we have that $\mathfrak{s}_{\lambda}(\mathbb{X}_m) = 0$ if λ does not fit inside an $(n-1) \times (n+1)$ box and $\mathfrak{s}_{\mu}(\mathbb{X}_{m+1}) = 0$ if μ does not fit inside an $(n+1) \times (n-1)$ box. If λ fits inside both an $(n-1) \times (n+1)$ box and an $n \times n$ box, then its complement λ^c with respect to the $n \times n$ box does not fit inside an $(n+1) \times (n-1)$ box. Thus, (10.28) implies that the last morphism in (10.29) is zero, so $\text{Im}(f) \subset \ker(\delta)$, as desired. \square

Combining Theorem 10.21 and Propositions 10.25, 10.26, and 10.27 with Theorem 3.7, we arrive at our main result.

Theorem 10.28. The homology of the complex $\{\beta\}_{2n+1, \mathcal{S}}$ of \mathbb{Z} -graded super \mathbb{K} -vector spaces is an invariant of the framed link \mathcal{L}_{β} , up to isomorphism of \mathbb{Z}^2 -graded super \mathbb{K} -vector spaces.

Proof. Let $\epsilon(\beta)$ be the exponent sum of β , and consider the renormalized invariant

$$(10.30) \quad \mathbf{q}^{n(n+1)\epsilon(\beta)} \mathbf{t}^{-n\epsilon(\beta)} \{\beta\}_{2n+1, \mathcal{S}}$$

which is invariant under the second Markov move by Propositions 10.26 and 10.27. Since $\epsilon(\beta)$ is unchanged by the braid relations and the first Markov move, Theorem 10.21 and Propositions 10.25 imply that (10.30) is invariant under these moves. Hence, by Theorem 3.7, (10.30) is an invariant of the (unframed) link \mathcal{L}_{β} , up to homotopy equivalence. Since $\epsilon(\beta)$ equals the writhe of the link diagram for \mathcal{L}_{β} given by taking the braid closure, the non-renormalized invariant $\{\beta\}_{2n+1, \mathcal{S}}$ is an invariant of the framed link \mathcal{L}_{β} (endowed with the blackboard framing), up to homotopy equivalence. It follows that the homology of $\{\beta\}_{2n+1, \mathcal{S}}$, which is \mathbb{Z}^2 graded via \mathbf{t} - and \mathbf{q} -degree, is an invariant of the framed link \mathcal{L}_{β} . \square

Definition 10.29. Let $H_{\mathfrak{so}_{2n+1}}(\mathcal{L}_{\beta}^{\mathcal{S}}) \in \text{sVect}_{\mathbb{K}}^{\mathbb{Z} \times \mathbb{Z}}$ denote the homology of the complex $\{\beta\}_{2n+1, \mathcal{S}}$.

We will refer to these link invariants as *Type B spin link homology*, or simply as *spin link homology*, in light of our decategorification results given in the following section.

Remark 10.30. As graded super vector spaces form a semisimple category, the homology of the complex $\{\beta\}_{2n+1, \mathcal{S}}$ is uniquely determined by the isomorphism class of $\{\beta\}_{2n+1, \mathcal{S}}$ in the homotopy category $\mathcal{K}(\text{sVect}_{\mathbb{K}}^{\mathbb{Z}})$. One can view spin link homology as an invariant valued in $\mathcal{K}(\text{sVect}_{\mathbb{K}}^{\mathbb{Z}})$ rather than in the category $\text{sVect}_{\mathbb{K}}^{\mathbb{Z} \times \mathbb{Z}}$ of bigraded super vector spaces.

10.6. Decategorification. We conclude this section by establishing the results from §1.1 and §1.2 that relate our spin link homology to the spin-colored \mathfrak{so}_{2n+1} link polynomials discussed in Section 4.

To begin, recall that if \mathcal{A} is a \mathbb{K} -linear category equipped with an involution σ , then there is an action of the group $\langle \text{sgn} \rangle \cong \mathbb{Z}/2$ on \mathcal{A}^{σ} defined by $\text{sgn}(X, \phi) := (X, -\phi)$. Further, the σ -weighted Grothendieck group $K_0^{\sigma}(\mathcal{A}^{\sigma})$ is defined by imposing the relation $[\text{sgn}(X, \phi)] = -[(X, \phi)]$ on $K_0(\mathcal{A}^{\sigma})$.

Although the category $\mathrm{sVect}_{\mathbb{K}}^{\mathbb{Z}}$ does not arise in this way, it also carries a $\mathbb{Z}/2$ action, via the *parity-reversing involution*

$$\mathbf{s}: \mathrm{sVect}_{\mathbb{K}}^{\mathbb{Z}} \rightarrow \mathrm{sVect}_{\mathbb{K}}^{\mathbb{Z}}$$

which e.g. sends $\mathbb{K}^{m|n} \mapsto \mathbb{K}^{n|m}$. We can similarly impose the relation $[\mathbf{s}V] = -[V]$ on $K_0(\mathrm{sVect}_{\mathbb{K}}^{\mathbb{Z}})$, and in the quotient

$$K_0^{\mathbf{s}}(\mathrm{sVect}_{\mathbb{K}}^{\mathbb{Z}}) := K_0(\mathrm{sVect}_{\mathbb{K}}^{\mathbb{Z}})/([\mathbf{s}V] = -[V])$$

we find that $[\mathbb{K}^{0|n}]_{\mathbf{s}} = -[\mathbb{K}^{n|0}]_{\mathbf{s}}$. It follows that there is an isomorphism $K_0^{\mathbf{s}}(\mathrm{sVect}_{\mathbb{K}}^{\mathbb{Z}}) \cong \mathbb{Z}[q^{\pm}]$ given by $[V]_{\mathbf{s}} \mapsto \dim_{\mathbf{q}}(V)$. (Recall that dimension of a super vector space is computed as in (2.2), i.e. it always means super dimension.)

Since the functor $\mathbf{q}^{-mn^2}(\mathcal{R}^+ \oplus \mathbf{s}\mathcal{R}^-)$ intertwines sgn and \mathbf{s} , there is a unique map $K_0^{\sigma}((\mathcal{B}_m^n)^{\tau}) \rightarrow \mathbb{Z}[q^{\pm}]$ such that the following diagram commutes:

$$\begin{array}{ccc} \mathcal{K}((\mathcal{B}_m^n)^{\tau}) & \xrightarrow{\mathbf{q}^{-mn^2}(\mathcal{R}^+ \oplus \mathbf{s}\mathcal{R}^-)} & \mathcal{K}(\mathrm{sVect}_{\mathbb{K}}^{\mathbb{Z}}) \\ K_0^{\tau} \downarrow & & \downarrow K_0^{\mathbf{s}} \\ K_0^{\tau}((\mathcal{B}_m^n)^{\tau}) & \longrightarrow & \mathbb{Z}[q^{\pm}] \end{array}$$

This is essentially the framework for our decategorification results, as we now explain.

For an additive category \mathcal{A} , one can identify the triangulated Grothendieck group of the bounded homotopy category $\mathcal{K}(\mathcal{A})$ with the split Grothendieck group of \mathcal{A} , the map being given by taking the Euler characteristic (alternative sum of classes) of each complex. See e.g. [61] for a careful proof. Hence, given a complex $C = \bigoplus_{i \in \mathbb{Z}} \mathbf{t}^i C^i$ in $\mathcal{K}(\mathcal{A}^{\sigma})$, we can consider its class $\sum_{i \in \mathbb{Z}} (-1)^i [C^i] \in K_0(\mathcal{A}^{\sigma})$ which therefore determines a well-defined class $[C]_{\sigma} := \sum_{i \in \mathbb{Z}} (-1)^i [C^i]_{\sigma}$ in the quotient $K_0^{\sigma}(\mathcal{A}^{\sigma})$. Similarly, given $C = \bigoplus_{i \in \mathbb{Z}} \mathbf{t}^i C^i \in \mathcal{K}(\mathrm{sVect}_{\mathbb{K}}^{\mathbb{Z}})$, we can consider $[C]_{\mathbf{s}} = \sum_{i \in \mathbb{Z}} (-1)^i \dim_{\mathbf{q}}(C^i) \in \mathbb{Z}[q^{\pm}] \cong K_0^{\mathbf{s}}(\mathrm{sVect}_{\mathbb{K}}^{\mathbb{Z}})$.

In all, we arrive at the following diagram:

$$(10.31) \quad \begin{array}{ccc} \mathcal{K}((\mathcal{B}_m^n)^{\tau}) & \xrightarrow{\mathbf{q}^{-mn^2}(\mathcal{R}^+ \oplus \mathbf{s}\mathcal{R}^-)} & \mathcal{K}(\mathrm{sVect}_{\mathbb{K}}^{\mathbb{Z}}) & , & C^{\tau}(\beta) & \longmapsto & \{\beta\}_{2n+1, \mathbf{s}} \\ K_0^{\tau} \downarrow & & \downarrow K_0^{\mathbf{s}} & & \downarrow & & \downarrow \\ K_0^{\tau}((\mathcal{B}_m^n)^{\tau}) & \longrightarrow & \mathbb{Z}[q^{\pm}] & & [C^{\tau}(\beta)]_{\tau} & \longmapsto & [\{\beta\}_{2n+1, \mathbf{s}}]_{\tau} \end{array}$$

Notation 10.31. To emphasize dependence on τ , we write $[\{\beta\}_{2n+1, \mathbf{s}}]_{\tau}$ instead of $[\{\beta\}_{2n+1, \mathbf{s}}]_{\mathbf{s}}$ both in (10.31) and for the duration. Since the Euler characteristic of a complex in an abelian category always equals that of its homology, we have

$$[\{\beta\}_{2n+1, \mathbf{s}}]_{\tau} = \dim_{\mathbf{q}, \mathbf{t}}(H_{\mathbf{s}\mathfrak{so}_{2n+1}}(\mathcal{L}_{\beta}^{\mathbf{S}}))|_{t=-1} = \sum_i (-1)^i \dim_{\mathbf{q}}(H_{\mathbf{s}\mathfrak{so}_{2n+1}}^i(\mathcal{L}_{\beta}^{\mathbf{S}})).$$

Theorem 10.32. Assume that $n = 1, 2, 3$. If $\beta \in \mathrm{Br}_m$, then

$$(10.32) \quad \bar{P}_{\mathbf{s}\mathfrak{so}_{2n+1}}(\mathcal{L}_{\beta}^{\mathbf{S}}) = (-1)^{n\epsilon(\beta)+m\binom{n+1}{2}} q^{\frac{1}{2}n\epsilon(\beta)} \sum_i (-1)^i \dim_{\mathbf{q}}(H_{\mathbf{s}\mathfrak{so}_{2n+1}}^i(\mathcal{L}_{\beta}^{\mathbf{S}})).$$

Proof. Let $\mathbf{m}\beta$ denote the mirror braid, i.e. the braid obtained from $\beta = \beta_{i_1}^{\epsilon_1} \cdots \beta_{i_\ell}^{\epsilon_\ell}$ by replacing each $\beta_i^{\pm 1}$ with β_i^{\mp} . By (3.6), it suffices to show that

$$P_{\mathbf{s}\mathfrak{so}_{2n+1}}(\mathcal{L}_{\beta}^{\mathbf{S}}) = (-1)^{n\epsilon(\mathbf{m}\beta)+m\binom{n+1}{2}} q^{\frac{1}{2}n\epsilon(\mathbf{m}\beta)} [\{\mathbf{m}\beta\}_{2n+1, \mathbf{s}}]_{\tau}.$$

For this, we will use Theorem 4.45.

Let $A_m^n = \mathbb{C}(q^{\frac{1}{2}}) \otimes_{\mathbb{Z}[q^{\pm 1}]} K_0^\tau((\mathcal{B}_m^n)^\tau)$ and set $\mathbf{X}_i^{(k)} := [(\mathbf{X}_i^{(k)}, (-1)^{\binom{n+1}{2} + (n-k+1)k} \mathbf{LC}_k)]_\tau \in A_m^n$. Corollaries 8.12 and 8.23 then imply that Condition A1. of Theorem 4.45 holds. Further, Definition 10.14 implies that Condition A4. holds, since $(-q^{\frac{1}{2}})^{n\epsilon(\mathbf{m}\beta)} [C^\tau(\mathbf{m}\beta)]_\tau$ for $\beta = \beta_i$ evaluates to

$$\beta_i \mapsto (-q^{\frac{1}{2}})^{-n} \sum_{k=0}^n q^k (-1)^n \mathbf{X}_i^{(n-k)} = (-q^{\frac{1}{2}})^{-n} (-q)^n \sum_{\ell=0}^n q^{-\ell} \mathbf{X}_i^{(\ell)} = q^{\frac{n}{2}} \sum_{\ell=0}^n q^{-\ell} \mathbf{X}_i^{(\ell)}.$$

Let $T_m^n: A_m^n \rightarrow \mathbb{C}(q^{\frac{1}{2}})$ be given by $(-1)^{m\binom{n+1}{2}}$ times the bottom horizontal map in (10.31). For $(Y, \varphi_Y) \in (\mathcal{B}_m^n)^\tau$, we thus have

$$\begin{aligned} (10.33) \quad T_m^n([(Y, \varphi_Y)]_\tau) &= (-1)^{m\binom{n+1}{2}} q^{-mn^2} \left(\dim_{\mathbf{q}}(\mathrm{Hom}_{(\mathcal{B}_m^n)^\tau}((\mathbb{1}_n, +), (Y, \varphi_Y))) \right. \\ &\quad \left. - \dim_{\mathbf{q}}(\mathrm{Hom}_{(\mathcal{B}_m^n)^\tau}((\mathbb{1}_n, -), (Y, \varphi_Y))) \right) \\ &= (-1)^{m\binom{n+1}{2}} q^{-mn^2} \left(\dim_{\mathbf{q}}(\mathrm{Hom}_{\mathcal{B}_m^n}(\mathbb{1}_n, Y)^\tau) - \dim_{\mathbf{q}}(\mathrm{Hom}_{\mathcal{B}_m^n}(\mathbb{1}_n, Y)^{\mathrm{sgn}}) \right) \\ &= (-1)^{m\binom{n+1}{2}} q^{-mn^2} \mathrm{Tr}(\tau^*|_{\mathrm{Hom}_{\mathcal{B}_m^n}(\mathbb{1}_n, Y)}). \end{aligned}$$

In particular, (10.33) implies that T_m^n is trace-like, since (10.14) gives an isomorphism $\mathrm{Hom}_{\mathcal{B}_m^n}(\mathbb{1}_n, Y_1 Y_2) \cong \mathrm{Hom}_{\mathcal{B}_m^n}(\mathbb{1}_n, Y_2 Y_1)$ that intertwines the τ^* -action.

We now verify Condition A3. from Theorem 4.45. First, Remark 5.7 and Definitions 9.6 and 10.5 imply that \mathcal{B}_1^n is the additive, \mathbb{Z} -graded, \mathbb{K} -linear, monoidal category generated by the monoidal unit $\mathbb{1}_n$. This implies that $A_1^n \cong \mathbb{C}(q^{\frac{1}{2}})$. Further, the isomorphism in (9.8) gives that $\mathrm{End}_{\mathcal{B}_1^n}(\mathbb{1}_n) \cong H^*(\mathrm{Gr}_n(\mathbb{C}^{2n}))$, with basis consisting of the new bubble generators for Schur functions $\mathfrak{s}_\lambda(\mathbb{X}_1)$ with λ contained in an $n \times n$ box. By (6.3), we have that

$$(10.34) \quad T_1^n([(\mathbb{1}_n, +)]_\tau) = (-1)^{\binom{n+1}{2}} q^{-n^2} \mathrm{Tr}(\tau^*|_{\mathrm{End}_{\mathcal{B}_1^n}(\mathbb{1}_n)}) = (-1)^{\binom{n+1}{2}} q^{-n^2} \sum_{\substack{\lambda \subset n^n \\ \lambda = \lambda^t}} q^{2|\lambda|}$$

and a fun inductive exercise gives that

$$(10.35) \quad \sum_{\substack{\lambda \subset n^n \\ \lambda = \lambda^t}} q^{2|\lambda|} = q^{n^2} \prod_{i=1}^n (q^{2i-1} + q^{1-2i}).$$

so $T_1^n([(\mathbb{1}_n, +)]_\tau) = (-1)^{\binom{n+1}{2}} \prod_{i=1}^n (q^{2i-1} + q^{1-2i})$ as desired.

Finally, we confirm Condition A2. The requisite linear map $\iota: A_{m-1}^n \rightarrow A_m^n$ is induced from the functor $(\mathcal{B}_{m-1}^n)^\tau \rightarrow (\mathcal{B}_m^n)^\tau$, so it remains to establish (4.46). For this, (10.33) shows that we must relate the τ^* actions on $\mathrm{Hom}_{\mathcal{B}_{m-1}^n}(\mathbb{1}_n, Y)$ and $\mathrm{Hom}_{\mathcal{B}_m^n}(\mathbb{1}_n, \mathbf{X}_{m-1}^{(k)} Y)$ for $Y \in \mathcal{B}_{m-1}^n$ (for appropriate equivariant structures on $\mathbb{1}_n, Y$, and \mathbf{X}_{m-1}).

For this, consider the linear map

$$(10.36) \quad \mathrm{Hom}_{\mathcal{B}_2^n}(\mathbb{1}_n, \mathbf{X}^{(k)}) \otimes_{\mathbb{K}} \mathrm{Hom}_{\mathcal{B}_{m-1}^n}(\mathbb{1}_n, Y) \rightarrow \mathrm{Hom}_{\mathcal{B}_m^n}(\mathbb{1}_n, \mathbf{X}_{m-1}^{(k)} Y)$$

given by precomposing horizontal composition with the map

$$\mathrm{Hom}_{\mathcal{B}_2^n}(\mathbb{1}_n, \mathbf{X}^{(k)}) \hookrightarrow \mathrm{Hom}_{\mathcal{B}_m^n}(\mathbb{1}_n, \mathbf{X}_{m-1}^{(k)})$$

which includes into the last \mathfrak{gl}_2 -string in \mathcal{B}_m^n . (In lay terms, we color an element in $\mathrm{Hom}_{\mathcal{B}_2^n}(\mathbb{1}_n, \mathbf{X}^{(k)})$ the color of the final Dynkin node in \mathcal{B}_m^n , and place it next to an element of $\mathrm{Hom}_{\mathcal{B}_m^n}(\mathbb{1}_n, Y)$.) As in the proof of Proposition 10.26, [34, Proposition 3.11] and (5.8) imply that (10.36) is surjective. Further, it descends to a map

$$(10.37) \quad \mathrm{Hom}_{\mathcal{B}_2^n}(\mathbb{1}_n, \mathbf{X}^{(k)}) \otimes_{H^*(\mathrm{Gr}_n(\mathbb{C}^{2n}))} \mathrm{Hom}_{\mathcal{B}_{m-1}^n}(\mathbb{1}_n, Y) \rightarrow \mathrm{Hom}_{\mathcal{B}_m^n}(\mathbb{1}_n, \mathbf{X}_{m-1}^{(k)} Y)$$

where $H^*(\mathrm{Gr}_n(\mathbb{C}^{2n}))$ acts by new bubbles (in the alphabet \mathbb{X}_1 in the first factor and the alphabet \mathbb{X}_{m-1} in the second). Computing dimensions using Proposition 9.10, we see that (10.37) is an isomorphism. Further, it is straightforward to see that this isomorphism intertwines the $\tau^* \otimes \tau^*$ action on the domain with the τ^* action on the codomain (provided we compute it with respect to the equivariant structure on $\mathbf{X}_{m-1}^{(k)} Y$ given as the tensor product of the equivariant structures on $\mathbf{X}_{m-1}^{(k)}$ and Y).

We have thus reduced to verifying that

$$\begin{aligned} & (-1)^{2\binom{n+1}{2}} q^{-2n^2} \mathrm{Tr}(\tau^* |_{\mathrm{Hom}_{\mathcal{B}_2^n}(\mathbb{1}_n, \mathbf{X}^{(k)})}) \\ &= (-1)^{n(k+1) + \binom{n-k}{2}} \prod_{t=1}^{n-k} \frac{[n+1-t][n+t]}{[t]^2} (-1)^{\binom{n+1}{2}} q^{-n^2} \mathrm{Tr}(\tau^* |_{\mathrm{End}_{\mathcal{B}_1^n}(\mathbb{1}_n)}) \\ &\stackrel{(10.34), (10.35)}{=} (-1)^{n(k+1) + \binom{n-k}{2}} \prod_{t=1}^{n-k} \frac{[n+1-t][n+t]}{[t]^2} (-1)^{\binom{n+1}{2}} \prod_{i=1}^n (q^{2i-1} + q^{1-2i}) \end{aligned}$$

where here the τ^* action is taken with respect to the equivariant objects $(\mathbf{X}^{(k)}, (-1)^{\binom{n+1}{2} + \binom{n-k+1}{2} + k} \mathbf{LC}_k)$ and $(\mathbb{1}_n, +)$. Equivalently, we must show that

$$(10.38) \quad q^{-2n^2} \mathrm{Tr}(\tau^* |_{\mathrm{Hom}_{\mathcal{B}_2^n}(\mathbb{1}_n, \mathbf{X}^{(k)})}) = (-1)^{-nk} \prod_{t=1}^{n-k} \frac{[n+1-t][n+t]}{[t]^2} \prod_{i=1}^n (q^{2i-1} + q^{1-2i})$$

for the action with respect to the equivariant objects $(\mathbf{X}^{(k)}, +\mathbf{LC}_k)$ and $(\mathbb{1}_n, +)$.

When $k = 0$, equation (10.38) becomes

$$(10.39) \quad q^{-2n^2} \mathrm{Tr}(\tau^* |_{\mathrm{End}_{\mathcal{B}_2^n}(\mathbb{1}_n)}) = \prod_{t=1}^n \frac{[n+1-t][n+t]}{[t]^2} \prod_{i=1}^n (q^{2i-1} + q^{1-2i}).$$

By Example 4.31, the right-hand side is equal to $(\prod_{i=1}^n (q^{2i-1} + q^{1-2i}))^2$. Since $\mathrm{End}_{\mathcal{B}_2^n}(\mathbb{1}_n)$ has a basis consisting of (new bubbles for) products $\mathfrak{s}_\lambda(\mathbb{X}_1) \mathfrak{s}_\mu(\mathbb{X}_2)$ with λ, μ contained in an $n \times n$ box, (10.39) holds by (6.3) and (10.35).

When $k = n$, equation (10.38) becomes

$$(10.40) \quad \mathrm{Tr}(\tau^* |_{\mathrm{Hom}_{\mathcal{B}_2^n}(\mathbb{1}_n, \mathbf{X}^{(n)})}) = (-1)^{-n^2} q^{2n^2} \prod_{i=1}^n (q^{2i-1} + q^{1-2i})$$

which we again verify directly. Equation (5.9) and Proposition 9.10 imply that $\mathrm{Hom}_{\mathcal{B}_2^n}(\mathbb{1}_n, \mathbf{X}^{(n)})$ has basis

$$\left\{ \bigcup_{\mathfrak{s}_\lambda}^n \mathbb{1}_n \right\}_{\lambda \subset n^n}$$

so we compute

$$\tau^* \left(\bigcup_{\mathfrak{s}_\lambda}^n \mathbb{1}_n \right) \stackrel{(6.3)}{=} \bigcirc_{\mathfrak{s}_{\lambda^t}}^n \mathbb{1}_n \stackrel{(5.33)}{=} (-1)^{n^2} \bigcup_{\mathfrak{s}_{\lambda^t}}^n \mathbb{1}_n.$$

Since a thickness- n cup in \mathfrak{gl}_2 weight (n, n) has degree n^2 , equation (10.35) implies (10.40).

The remaining cases of (4.46) are when $k = 1$ (when $n = 2, 3$) and $k = 2$ (when $n = 3$). As for $k = 0, n$, it is possible to verify (10.38) directly in these cases; however, the computations are tedious so we use a trick instead. We necessarily have that $T_2^n(\mathbf{X}_i^{(k)}) = p_{k,n} T_1^n([\mathbb{1}_n, +])_\tau$ for some Laurent

polynomials $p_{k,n}$. Further, the Markov II invariance established in Propositions 10.26 and 10.27 implies presently that

$$(10.41) \quad (-1)^{n+\binom{n+1}{2}} q^{-n(n+1)} T_1^n([\mathbb{1}_n, +]_\tau) = (-1)^n \sum_{k=0}^n q^{-k} T_2^n(X_i^{(n-k)})$$

and

$$(10.42) \quad (-1)^{n+\binom{n+1}{2}} q^{n(n+1)} T_1^n([\mathbb{1}_n, +]_\tau) = (-1)^n \sum_{k=0}^n q^k T_2^n(X_i^{(n-k)}).$$

Our computations for $T_2^n(X_i^{(0)})$ and $T_2^n(X_i^{(n)})$ pair with these equations to determine the outstanding values of $T_2^n(X_i^{(k)})$.

In detail, when $n = 2$, equations (10.41) and (10.42) yield

$$\begin{aligned} -q^{-6} &= 1 + q^{-1} \cdot p_{1,2} - q^{-2}(q + q^{-1})(q^3 + q^{-3}) \\ -q^6 &= 1 + q \cdot p_{1,2} - q^2(q + q^{-1})(q^3 + q^{-3}) \end{aligned}$$

and it is straightforward to check that

$$p_{1,2} = q^3 + q^{-3} = (-1)^{2(1+1)+\binom{2-1}{2}} \prod_{t=1}^{2-1} \frac{“[2+1-t][2+t]”}{“[t]^2”}$$

solves both, as desired. When $n = 3$, equations (10.41) and (10.42) give the system

$$\begin{aligned} q^{-12} &= 1 + q^{-1} \cdot p_{2,3} + q^{-2} p_{1,3} + q^{-3}(q + q^{-1})(q^3 + q^{-3})(q^5 + q^{-5}) \\ q^{12} &= 1 + q \cdot p_{2,3} + q^2 p_{1,3} + q^3(q + q^{-1})(q^3 + q^{-3})(q^5 + q^{-5}) \end{aligned}$$

and again it is straightforward linear algebra to verify that

$$p_{2,3} = -(q^5 + q + q^{-1} + q^{-5}) = (-1)^{3(2+1)+\binom{3-2}{2}} \prod_{t=1}^{3-2} \frac{“[3+1-t][3+t]”}{“[t]^2”}$$

and

$$p_{1,3} = -(q^3 + q^{-3})(q^5 + q^{-5}) = (-1)^{3(1+1)+\binom{3-1}{2}} \prod_{t=1}^{3-1} \frac{“[3+1-t][3+t]”}{“[t]^2”}$$

are the unique solutions, as desired. \square

As initially stated in Theorem 1.1, we can reformulate Theorem 10.32 in terms of the involution τ acting on Λ^n -colored \mathfrak{sl}_{2n} link homology.

Corollary 10.33. Let $\mathcal{L} \subset S^3$ be a link.

- For all $n \geq 1$, the n -colored \mathfrak{sl}_{2n} Khovanov–Rozansky homology $H_{\mathfrak{sl}_{2n}}(\mathcal{L}^n)$ admits an involution τ that preserves the bidegree. The bigraded eigenspaces of τ are link invariants.
- If $\beta \in \text{Br}_m$ and $n = 1, 2, 3$, then

$$(10.43) \quad \text{Tr} \left(\tau \curvearrowright H_{\mathfrak{sl}_{2n}}(\mathcal{L}_\beta^n) \right) = (-1)^{n\epsilon(\beta)+m\binom{n+1}{2}} q^{\frac{1}{2}n\epsilon(\beta)} \overline{P}_{\mathfrak{so}_{2n+1}}(\mathcal{L}_\beta^S).$$

Proof. The arguments establishing Theorems 10.21 and 10.28 can be repackaged as saying that the τ_* -action on the chain groups of $[[\beta \mathbb{1}_n]]_{2n}$ descends to a well-defined involution on $H_{\mathfrak{sl}_{2n}}(\mathcal{L}^n)$. The ± 1 -eigenspaces of this involution on $H_{\mathfrak{sl}_{2n}}^i(\mathcal{L}^n)$ are exactly the even/odd parts of the super vector space $H_{\mathfrak{so}_{2n+1}}^i(\mathcal{L}_\beta^S)$, so the result follows from Theorem 10.32. \square

More generally, we propose that the equality stated in Theorem 10.32 (and therefore in Corollary 10.33) holds for all n .

Conjecture 10.34. Given $\beta \in \text{Br}_m$, equation (10.32) additionally holds for $n \geq 4$. That is, Type B spin link homology categorifies the spin-colored \mathfrak{so}_{2n+1} link polynomial for all $n \geq 1$.

The proof of Theorem 10.32 would suffice to prove Conjecture 10.34 once additional technical details are verified. Specifically, the proof of Theorem 10.32 uses Theorem 4.45, which holds unqualified only when $n = 1, 2, 3$. For $n \geq 4$, Theorem 4.45 is conditional on Conjectures 4.40 and 4.42. Additionally, in the proof of Theorem 10.32 we only checked Condition A2. of Theorem 4.45 when $k = 0$, $k = n$, and for $1 \leq k \leq n - 1$ when $n = 1, 2, 3$. (Our “trick” used for these latter cases does not work for $n \geq 4$.) After establishing Condition A2., the rest of the proof of Theorem 10.32 generalizes word-for-word.

Remark 10.35. Conjecture 10.34 is therefore an immediate consequence of Conjectures 4.40 and 4.42 and the following conjecture.

Conjecture 10.36. Equation (10.38) additionally holds for all $n \geq 4$ and all $1 \leq k \leq n - 1$.

APPENDIX A. PROOF OF LEMMA 4.27

The goal of this section is to establish the identity (4.32).

A.1. On the devil’s product. We begin with an informal discussion on the devil’s arithmetic, and the derivation of some useful identities. Recall from Definition 4.1 that the devil’s product is defined for $m \leq n \in \mathbb{Z}_{\geq 0}$ by

$$(A.1) \quad “[m][n]” = “[n][m]” := [n + m - 1] - [n + m - 3] + [n + m - 5] \mp \cdots + (-1)^{m-1}[n - m + 1].$$

Using the relation $[2][n] = [n + 1] + [n - 1]$ involving the usual product of quantum integers, this implies the relation

$$(A.2) \quad [2] \cdot “[m][n]” = [n + m] + (-1)^{m-1}[n - m]$$

Before presenting additional helpful identities, we warn the reader of two unintuitive aspects of devil’s arithmetic already present above: lack of symmetry and dependence on parity. When multiplying two quantum numbers, one can expand the product as a sum as follows.

$$[2][n] = [n + 1] + [n - 1], \quad [3][n] = [n + 2] + [n] + [n - 2], \quad [4][n] = [n + 3] + [n + 1] + [n - 1] + [n - 3], \quad \text{etc.}$$

Multiplication is commutative, which gives rise to two ways to expand any product, e.g.

$$(A.3) \quad [3][5] = [7] + [5] + [3], \quad [5][3] = [7] + [5] + [3] + [1] + [-1],$$

and $[3][5] = [5][3]$ because $[1] + [-1] = 0$. In general, the equality between these two expansions follows from a cancellation of positive and negative quantum integers.

In contrast, the devil’s product does not admit two equivalent signed expansions! Paralleling the example in (A.3), we have

$$“[3][5]” := [7] - [5] + [3], \quad \text{which does **not** equal } [7] - [5] + [3] - [1] + [-1].$$

When deriving formulas for “[m][n]”, one cannot treat m and n symmetrically, since (A.1) assumes that $m \leq n$. Equation (A.2) further illustrates the asymmetry between m and n , as well as the dependence on the parity of the smaller number m . An easy consequence of that relation is that the $q = 1$ specialization of the devil’s product is given by

$$“[m][n]”|_{q=1} = \begin{cases} n & \text{if } m \text{ is odd,} \\ m & \text{if } m \text{ is even.} \end{cases}$$

We now record some alternative formulae for the devil’s product.

Notation A.1. If $k \in \mathbb{Z}_{\geq 0}$, then we write $[2]_k := [2]_{q^k} := q^k + q^{-k}$.

Note that $[2] = [2]_1$ divides $[2]_k$ if and only if k is odd and that $[k]$ divides $[m]$ when k divides m . Also, we record the identity

$$(A.4) \quad \frac{[k \cdot \ell]}{[k]} = [\ell]_{q^k}.$$

Lemma A.2. If $m \leq n$ in $\mathbb{Z}_{\geq 0}$, then

$$[m][n] = \begin{cases} [2]_n [m/2]_{q^2} & \text{if } m \text{ is even,} \\ [2]_m [n/2]_{q^2} & \text{if } m \text{ is odd and } n \text{ is even,} \\ \binom{[2]_m}{[2]} [n] & \text{if } m \text{ and } n \text{ are odd.} \end{cases}$$

Proof. Expanding the quantum integers as $[k] = \frac{q^k - q^{-k}}{q - q^{-1}}$ on the right-hand side of (A.2), we obtain

$$[2][m][n] = \frac{(q^{n+m} - q^{-n-m}) + (-1)^{m-1}(q^{n-m} - q^{m-n})}{q - q^{-1}} = \frac{(q^n + (-1)^m q^{-n})(q^m + (-1)^{m-1} q^{-m})}{q - q^{-1}}.$$

This is clearly equal to $[2]_n [m]$ or $[2]_m [n]$ depending on the parity of m . Now divide by $[2]$, and possibly use the $k = 2$ case of (A.4). \square

Lemma A.3. We have

$$(A.5) \quad [n]^{[2]} = [n]_{q^2}.$$

Proof. An immediate consequence of (A.2) and the $k = 2$ case of (A.4). \square

A.2. The proof of Lemma 4.27. Recall that equation (4.32) defines

$$\rho_t^{(\ell)} := (-1)^{\binom{\ell+2-t}{2}} + q^{-1} \frac{[\ell + 1 - t][\ell + t]}{[[t]_2]} \rho_{t+1}^{(\ell)}.$$

recursively for $1 \leq t \leq \ell$ by declaring that $\rho_{\ell+1}^{(\ell)} := 1$. We aim to prove that $\rho_1^{(\ell)} = (q^{-2})^{\binom{\ell+1}{2}}$. To do so, we will unravel the recurrence to obtain a formula for $\rho_1^{(\ell)}$, which will depend on the value of ℓ modulo 4.

Example A.4. Consider the case when $\ell = 4n$, for $n \in \mathbb{Z}_{\geq 0}$. We find that

$$\begin{aligned} \rho_1^{(4n)} &= 1 + q^{-1} \frac{[2]_{q^{4n+1}} [2n]_{q^2}}{[1]_{q^2}} \rho_2^{(4n)} \\ &= 1 + q^{-1} \frac{[2]_{q^{4n+1}} [2n]_{q^2}}{[1]_{q^2}} - q^{-2} \frac{[2]_{q^{4n+1}} [2]_{q^{4n-1}} [2n]_{q^2} [2n+1]_{q^2}}{[1]_{q^2} [2]_{q^2}} \rho_3^{(4n)} \\ &= \dots \\ &= 1 - q^{-2} [2]_{q^{4n+1}} [2]_{q^{4n-1}} \begin{bmatrix} 2n+1 \\ 2 \end{bmatrix}_{q^2} + q^{-4} [2]_{q^{4n+1}} [2]_{q^{4n-1}} [2]_{q^{4n+3}} [2]_{q^{4n-3}} \begin{bmatrix} 2n+2 \\ 4 \end{bmatrix}_{q^2} \pm \dots \\ &\quad + q^{-1} [2]_{q^{4n+1}} \begin{bmatrix} 2n \\ 1 \end{bmatrix}_{q^2} - q^{-3} [2]_{q^{4n+1}} [2]_{q^{4n-1}} [2]_{q^{4n+3}} \begin{bmatrix} 2n+1 \\ 3 \end{bmatrix}_{q^2} \pm \dots \end{aligned}$$

where we have reordered the terms in the final equation. The assertion is that this sum simplifies to $(q^{-2})^{\binom{4n+1}{2}}$.

Remark A.5. Before continuing, we sketch a proof of the $q = 1$ specialization of the result, when $\ell = 8$. This suggests a useful heuristic for the general arguments below. By example A.4, the assertion is that

$$(A.6) \quad \begin{aligned} 1 &= 2^0 \binom{4}{0} - 2^2 \binom{5}{2} + 2^4 \binom{6}{4} - 2^6 \binom{7}{6} + 2^8 \binom{8}{8} \\ &\quad + 2^1 \binom{4}{1} - 2^3 \binom{5}{3} + 2^5 \binom{6}{5} - 2^7 \binom{7}{7}. \end{aligned}$$

Here, we have suggestively written 1 as $2^0 \binom{4}{0}$ for the first term on the right-hand side.

We can organize this sum by highlighting the relevant entries of Pascal's triangle with solid (for the first row in (A.6)) and dashed (for the second) circles, and labeling them by the appropriate coefficient $\pm 2^a$:

$$(A.7) \quad \begin{array}{cccccccc} & & & & & & & & 1 \\ & & & & & & & & 1 & 1 \\ & & & & & & & & 1 & 2 & 1 \\ & & & & & & & & 1 & 3 & 3 & 1 \\ & & & & & & & & +2^0 \textcircled{1} & +2^1 \textcircled{4} & 6 & 4 & 1 \\ & & & & & & & & 1 & 5 & -2^2 \textcircled{10} & -2^3 \textcircled{10} & 5 & 1 \\ & & & & & & & & 1 & 6 & 15 & 20 & +2^4 \textcircled{15} & +2^5 \textcircled{6} & 1 \\ & & & & & & & & 1 & 7 & 21 & 35 & 35 & 21 & -2^6 \textcircled{7} & -2^7 \textcircled{1} \\ & & & & & & & & 1 & 8 & 28 & 56 & 70 & 56 & 28 & 8 & +2^8 \textcircled{1} \end{array}$$

Using the defining property of Pascal's triangle, we can remove each solid circle labeled by $\pm 2^a$ and add this value to the coefficient for both spots diagonally above it (and ignoring spots outside of the triangle). This leaves the corresponding sum unchanged, and yields the schematic:

$$(A.8) \quad \begin{array}{cccccccc} & & & & & & & & 1 \\ & & & & & & & & 1 & 1 \\ & & & & & & & & 1 & 2 & 1 \\ & & & & & & & & +2^0 \textcircled{1} & 3 & 3 & 1 \\ & & & & & & & & 1 & -2^1 \textcircled{4} & -2^2 \textcircled{6} & 4 & 1 \\ & & & & & & & & 1 & 5 & 10 & +2^3 \textcircled{10} & +2^4 \textcircled{5} & 1 \\ & & & & & & & & 1 & 6 & 15 & 20 & 15 & -2^5 \textcircled{6} & -2^6 \textcircled{1} \\ & & & & & & & & 1 & 7 & 21 & 35 & 35 & 21 & 7 & +2^7 \textcircled{1} \\ & & & & & & & & 1 & 8 & 28 & 56 & 70 & 56 & 28 & 8 & 1 \end{array}$$

The corresponding sum is exactly the $q = 1$ specialization of $\rho_1^{(7)}$. Iterating this procedure, we eventually see that $\rho_1^{(8)}|_{q=1} = \dots = \rho_1^{(0)}|_{q=1} = 1$.

Trying to adapt this argument for generic q is not straightforward. There are well-known analogues of Pascal's identity, which we record here:

$$(A.9) \quad \begin{bmatrix} x \\ y \end{bmatrix}_q = q^y \begin{bmatrix} x-1 \\ y \end{bmatrix}_q + q^{y-x} \begin{bmatrix} x-1 \\ y-1 \end{bmatrix}_q$$

and

$$(A.9) \quad \begin{bmatrix} x \\ y \end{bmatrix}_q = q^{-y} \begin{bmatrix} x-1 \\ y \end{bmatrix}_q + q^{-y+x} \begin{bmatrix} x-1 \\ y-1 \end{bmatrix}_q.$$

However, the cancellation between solid and dashed circles in (A.7), which simply used that $2^{m+1} - 2^m = 2^m$, now requires a cancellation of Laurent polynomials which is not guaranteed by the corresponding $q = 1$ cancellation. One must keep careful track of the powers of q , in both the quantum binomial coefficients (choosing between (A.8) and (A.9) as appropriate) and the complicated “powers of 2” appearing e.g. in Example A.4.

Remark A.6. In general, the scalars $\rho_t^{(\ell)}$ are structural coefficients in the type B web algebra which describe how to rewrite the expression for the spin braiding $R_{S,S} = q^{\frac{\ell}{2}} \sum_{\ell \geq 0} q^{-\ell} \mathbf{X}^{(\ell)}$, in the $\mathbf{\Xi}^{(\ell)}$ basis; see the proof of Proposition 4.28. At $q = 1$, one can use similar techniques to analyze the integers $\rho_t^{(\ell)}$ for $t > 1$. However, for generic q and $t > 1$, the desired cancellations do not work! At present, we do not have a closed formula for $\rho_t^{(\ell)}$ for $t > 1$, and it seems to be a difficult problem.

Returning to the task at hand, we now introduce notation to help make the sums expressing $\rho_1^{(\ell)}$ more compact.

Definition A.7. Given $c, d \in \mathbb{Z}_{\geq 0}$, write $(2)_c^d$ to denote the product consisting of the first d terms of

$$[2]_{c+1}[2]_{c-1}[2]_{c+3}[2]_{c-3} \cdots$$

and write $\{2\}_c^d$ to denote the product consisting of the first d terms of

$$[2]_{c-1}[2]_{c+1}[2]_{c-3}[2]_{c+3} \cdots$$

For $a, b \in \mathbb{Z}_{\geq 0}$, and $\epsilon \in \{\pm 1\}$, we then set

$$A(a, b, \epsilon) = \sum_{i \geq 0} (-1)^i q^{2i\epsilon} (2)_a^{2i} \begin{bmatrix} b+i \\ 2i \end{bmatrix}_{q^2}, \quad B(a, b, \epsilon) = \sum_{i \geq 0} (-1)^i q^{2i\epsilon} (2)_a^{2i+1} \begin{bmatrix} b+i \\ 2i+1 \end{bmatrix}_{q^2}$$

and

$$A'(a, b, \epsilon) = \sum_{i \geq 0} (-1)^i q^{2i\epsilon} \{2\}_a^{2i} \begin{bmatrix} b+i \\ 2i \end{bmatrix}_{q^2}, \quad B'(a, b, \epsilon) = \sum_{i \geq 0} (-1)^i q^{2i\epsilon} \{2\}_a^{2i+1} \begin{bmatrix} b+i \\ 2i+1 \end{bmatrix}_{q^2}.$$

Observe that, for all a and ϵ ,

$$(A.10) \quad A(a, 0, \epsilon) = 1 = A'(a, 0, \epsilon) \quad \text{and} \quad B(a, 0, \epsilon) = 0 = B'(a, 0, \epsilon).$$

Lemma A.8. For $n \geq 0$,

$$\begin{aligned} \rho_1^{(4n)} &= A(4n, 2n, -1) + q^{-1} B(4n, 2n, -1), \\ \rho_1^{(4n+1)} &= -A'(4n+2, 2n, -1) + q^{-1} B'(4n+2, 2n+1, -1), \\ \rho_1^{(4n+2)} &= -A(4n+2, 2n+1, -1) - q^{-1} B(4n+2, 2n+1, -1), \end{aligned}$$

and

$$\rho_1^{(4n+3)} = A'(4n+4, 2n+1, -1) - q^{-1} B'(4n+4, 2n+2, -1).$$

Proof. Following Example A.4, expand $\rho_1^{(\ell)}$ as a pair of alternating sums. □

Lemma A.9. For $b \geq 1$, we have

$$(A.11) \quad A(a, b, -1) = A(a, b-1, 1) - q^{a-2b-1} B(a, b, -1) - q^{-a-2b+1} B(a, b, 1),$$

$$(A.12) \quad B(a, b, 1) = q^{-2} B(a, b-1, -1) + q^{a+2b-1} A(a, b-1, 1) + q^{-a+2b-3} A(a, b-1, -1),$$

$$(A.13) \quad A'(a, b, 1) = A'(a, b-1, -1) - q^{a+2b+1} B'(a, b, 1) - q^{-a+2b-1} B'(a, b, -1),$$

and

$$(A.14) \quad B'(a, b, -1) = q^2 B'(a, b-1, 1) + q^{a-2b+1} A'(a, b-1, -1) + q^{-a-2b+3} A'(a, b-1, 1).$$

Proof. We prove the first and third formula and leave the second and fourth to the reader. We have

$$\begin{aligned} A(a, b, -1) &\stackrel{(A.8)}{=} A(a, b-1, 1) + \sum_{j \geq 0} (-1)^j q^{-2b} [2]_{a-2j+1} (2)_a^{2j-1} \begin{bmatrix} b-1+j \\ 2j-1 \end{bmatrix}_{q^2} \\ &= A(a, b-1, 1) - \sum_{i \geq 0} (-1)^i q^{-2b} (q^{a-2i-1} + q^{-a+2i+1}) (2)_a^{2i+1} \begin{bmatrix} b+i \\ 2i+1 \end{bmatrix}_{q^2} \\ &= A(a, b-1, 1) - q^{a-2b-1} B(a, b, -1) - q^{-a-2b+1} B(a, b, +1) \end{aligned}$$

and

$$\begin{aligned} A'(a, b, 1) &\stackrel{(A.9)}{=} A'(a, b-1, -1) + \sum_{j \geq 0} (-1)^j q^{2b} [2]_{a+2j-1} \{2\}_a^{2j-1} \begin{bmatrix} b-1+j \\ 2j-1 \end{bmatrix}_{q^2} \\ &= A'(a, b-1, -1) - \sum_{i \geq 0} (-1)^i q^{2b} (q^{a+2i+1} + q^{-a-2i-1}) \{2\}_a^{2i+1} \begin{bmatrix} b+i \\ 2i+1 \end{bmatrix}_{q^2} \\ &= A'(a, b-1, -1) - q^{a+2b+1} B'(a, b, 1) - q^{-a+2b-1} B'(a, b, -1). \end{aligned}$$

□

Lemma A.10. Fix $n \geq 0$. For $0 \leq k \leq 2n$, we have

$$(A.15) \quad \rho_1^{(4n)} = (-q^{-8n-2})^k (A(4n, 2n-k, -1) + q^{2k-1} B(4n, 2n-k, -1))$$

and

$$(A.16) \quad \rho_1^{(4n+1)} = (-q^{-8n-2})^k (-A'(4n+2, 2n-k, -1) + q^{-2k-1} B'(4n+2, 2n+1-k, -1)).$$

For $0 \leq k \leq 2n+1$ we have

$$(A.17) \quad \rho_1^{(4n+2)} = (-q^{-8n-6})^k (-A(4n+2, 2n+1-k, -1) - q^{2k-1} B(4n+2, 2n+1-k, -1))$$

and

$$(A.18) \quad \rho_1^{(4n+3)} = (-q^{-8n-6})^k (A'(4n+4, 2n+1-k, -1) - q^{-2k-1} B'(4n+4, 2n+2-k, -1)).$$

Proof. Induction on k . The $k=0$ base case is Lemma A.8. The induction step, in the case of equation (A.15), is illustrated as follows:

$$\begin{aligned} \rho_1^{(4n)} &= (-q^{-8n-2})^k (A(4n, 2n-k, -1) + q^{2k-1} B(4n, 2n-k, -1)) \\ &\stackrel{(A.11)}{=} (-q^{-8n-2})^k \left(A(4n, 2n-k-1, 1) - q^{4n-2(2n-k)-1} B(4n, 2n-k, -1) \right. \\ &\quad \left. - q^{-4n-2(2n-k)+1} B(4n, 2n-k, 1) + q^{2k-1} B(4n, 2n-k, -1) \right) \\ &= (-q^{-8n-2})^k (A(4n, 2n-k-1, 1) - q^{-8n+2k+1} B(4n, 2n-k, 1)) \\ &\stackrel{(A.12)}{=} (-q^{-8n-2})^k \left(A(4n, 2n-k-1, 1) - q^{-8n+2k+1} (q^{-2} B(4n, 2n-k-1, -1) \right. \\ &\quad \left. + q^{4n+2(2n-k)-1} A(4n, 2n-k-1, 1) + q^{-4n+2(2n-k)-3} A(4n, 2n-k-1, -1)) \right) \\ &= (-q^{-8n-2})^k (-q^{-8n+2k+1-2} B(4n, 2n-k-1, -1) - q^{-8n-2} A(4n, 2n-k-1, -1)) \\ &= (-q^{-8n-2})^{k+1} (A(4n, 2n-(k+1), -1) + q^{2(k+1)-1} B(4n, 2n-(k+1), -1)). \end{aligned}$$

For equation (A.17), an identical argument applies. For Equations A.16 and A.18, the argument is similar, but it instead employs equation (A.14) and then equation (A.13). □

At last, we arrive at the main result of this appendix.

Proof of Lemma 4.27. Recall that we must establish the identity

$$\rho_1^{(\ell)} = (q^{-2})^{\binom{\ell+1}{2}}.$$

Taking $k = 2n$ in (A.15) and applying (A.10) gives

$$\rho_1^{(4n)} = (-q^{-8n-2})^{2n} = (q^{-2})^{(4n+1)(2n)} = (q^{-2})^{\binom{4n+1}{2}}.$$

A similar argument using the $k = 2n + 1$ case of equation (A.17) shows that $\rho_1^{(4n+2)} = (q^{-2})^{\binom{4n+3}{2}}$.

For the remaining cases, we also use that $B'(a, 1, \epsilon) = [2]_{a-1}$ for all a, ϵ . It then follows from (A.10) and the $k = 2n$ case of (A.16) that

$$\rho_1^{(4n+1)} = (-q^{-8n-2})^{2n} (-1 + q^{-4n-1}[2]_{4n+1}) = (q^{-2})^{(4n+1)(2n+1)} = (q^{-2})^{\binom{4n+2}{2}}.$$

Finally, the same argument using the $k = 2n + 1$ case of (A.18) gives $\rho_1^{(4n+3)} = (q^{-2})^{\binom{4n+4}{2}}$. \square

APPENDIX B. RELATION TO WENZL'S APPROACH USING THE q -CLIFFORD ALGEBRA

In [74], Wenzl constructs an endomorphism $C \in \text{End}_{\mathbb{C}(q)}(S \otimes S)$ which commutes with the action of $U_q(\mathfrak{so}_{2n+1})$, but does so using a non-standard coproduct. We review Wenzl's construction here and slightly modify his conventions in order to construct an endomorphism of $S \otimes S$ which commutes with the action of $U_q(\mathfrak{so}_{2n+1})$ via the coproduct in Definition 3.1. Our main result is that Wenzl's endomorphism agrees with the (diagrammatically defined) endomorphism

$$H^{(1)} = \begin{array}{c} S \quad S \\ \diagdown \quad / \\ \text{---} 1 \text{---} \\ / \quad \diagdown \\ S \quad S \end{array}$$

from (4.35) in Section 4.

Definition B.1. Let $\text{Cl}_q(2n)$ be $\mathbb{C}(q)$ algebra with generators ψ_i, ψ_i^* , and $\omega_i^{\pm 1}$, for $i = 1, \dots, n$, subject to the relations

$$\begin{aligned} \psi_i^2 = 0 = (\psi_i^*)^2, \quad \psi_i \psi_j = -\psi_j \psi_i, \quad \psi_i \psi_j^* = -\psi_j^* \psi_i, \quad \psi_i^* \psi_j^* = -\psi_j^* \psi_i^*, \\ \omega_i \omega_i^{-1} = 1 = \omega_i^{-1} \omega_i, \quad \omega_i \omega_j = \omega_j \omega_i, \\ \omega_i \psi_i \omega_i^{-1} = q^2 \psi_i, \quad \omega_i \psi_i^* \omega_i^{-1} = q^{-2} \psi_i^*, \quad \omega_i \psi_j \omega_i^{-1} = \psi_j, \quad \omega_i \psi_j^* \omega_i^{-1} = \psi_j^*, \\ \psi_i \psi_i^* + q^2 \psi_i^* \psi_i = \omega_i^{-1}, \quad \psi_i \psi_i^* + q^{-2} \psi_i^* \psi_i = \omega_i, \end{aligned}$$

for all $1 \leq i \neq j \leq n$, **and** further quotiented by the relations

$$(B.1) \quad \omega_i \psi_i = \psi_i \quad \text{and} \quad \psi_i^* \omega_i = \psi_i^* \quad 1 \leq i \leq n.$$

Remark B.2. The algebra $\text{Cl}_q(2n)$ is a version of the q -Clifford algebra. The usual definition of the q -Clifford algebra [24] does not impose the relation (B.1). However, it is essential in establishing that Wenzl's element C commutes with $U_q(\mathfrak{so}_{2n+1})$.

Multiplying the relation $\psi_i \psi_i^* + q^2 \psi_i^* \psi_i = \omega_i^{-1}$ on the left by ω_i , and using $\omega_i \psi_i = \psi_i$ and $\omega_i \psi_i^* \omega_i^{-1} = q^{-2} \psi_i^*$, we deduce that

$$(B.2) \quad \psi_i \psi_i^* + \psi_i^* \psi_i = 1, \quad 1 \leq i \leq n.$$

Definition B.3. The *volume element* in $\text{Cl}_q(2n)$ is defined as

$$f := (\psi_1 \psi_1^* - \psi_1^* \psi_1)(\psi_2 \psi_2^* - \psi_2^* \psi_2) \dots (\psi_n \psi_n^* - \psi_n^* \psi_n).$$

For $1 \leq i \leq n$, it satisfies identities:

$$\omega_i f = f \omega_i, \quad \psi_i f = -f \psi_i, \quad \psi_i^* f = -f \psi_i^*.$$

Lemma B.4. There is a homomorphism of algebras $U_q(\mathfrak{so}_{2n+1}) \rightarrow \text{Cl}_q(2n)$ defined for $1 \leq i \leq n-1$ by

$$E_i \mapsto \psi_i \psi_{i+1}^*, \quad F_i \mapsto \psi_{i+1} \psi_i^*, \quad K_i \mapsto \omega_i \omega_i^{-1}$$

and

$$E_n \mapsto \psi_n f, \quad F_n \mapsto f \psi_n^*, \quad K_n \mapsto q \omega_n.$$

Proof. An elementary generators and relations check; see [24, Theorem 3.2]. \square

Notation B.5. Given $I \subset \{1, \dots, n\}$ so that $I = \{i_1, \dots, i_d\}$ with $i_1 < \dots < i_d$, we write $\psi_I^* := \psi_{i_1}^* \cdots \psi_{i_d}^*$. Also, let $\epsilon_I := \prod_{i \in I} (-1)^{n-i+1}$.

Lemma B.6. Let \mathbf{I} be the left ideal $\text{Cl}_q(2n) \cdot \langle \psi_i \mid 1 \leq i \leq n \rangle$. There is an isomorphism of vector spaces

$$\text{Cl}_q(2n)/\mathbf{I} \xrightarrow{\cong} S$$

such that $\epsilon_I \psi_I^* \mapsto x_I$. Moreover, the induced action of $U_q(\mathfrak{so}_{2n+1})$ on S via the homomorphism in Lemma B.4 coincides with the action in Definition 4.3.

Proof. See [24, Sections 2.1 and 4.1]. \square

Definition B.7. Let $\Omega_k := \omega_1 \omega_2 \cdots \omega_k$ and define the following element in $\text{Cl}_q(2n) \otimes \text{Cl}_q(2n)$:

$$C := \frac{\Omega_n f \otimes \Omega_n^{-1} f}{[2]} + \sum_{k=1}^n (\Omega_{k-1} \otimes \Omega_{k-1}^{-1}) \cdot (\psi_k \otimes \psi_k^* + \psi_k^* \otimes \psi_k).$$

Since $\text{Cl}_q(2n)^{\otimes 2}$ acts on $S^{\otimes 2}$, we obtain an operator in $\text{End}_{\mathbb{C}(q)}(S \otimes S)$, (also) denoted C .

Lemma B.8. The operator C is in $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$.

Proof. It suffices to show that the elements

$$\Delta(E_i) \mapsto \psi_i \psi_{i+1}^* \otimes \omega_i \omega_{i+1}^{-1} + 1 \otimes \psi_i \psi_{i+1}^*, \quad \Delta(F_i) \mapsto \psi_{i+1} \psi_i^* \otimes 1 + \omega_i^{-1} \omega_{i+1} \otimes \psi_{i+1} \psi_i^*$$

for $1 \leq i \leq n-1$, as well as the elements

$$\Delta(E_n) \mapsto \psi_n f \otimes q \omega_n + 1 \otimes \psi_n f, \quad \Delta(F_n) \mapsto f \psi_n^* \otimes 1 + q^{-1} \omega_n^{-1} \otimes f \psi_n^*,$$

commute with C in $\text{Cl}_q(2n) \otimes \text{Cl}_q(2n)$. The calculation to verify this follows the rubric outlined in the proof of [74, Lemma 3.4]. \square

Lemma B.9.

$$C \circ Y_i = (-1)^{n-i} \frac{[2(n-i)+1]}{[2]} Y_i$$

Proof. We proceed similarly to the proof of Lemma 4.22. There is some scalar χ so that $C \circ Y_i = \chi Y_i$. Write $\pi_{\{i+1, \dots, n\} \otimes \emptyset}$ to denote the *linear* operator which projects, with respect to the basis $\{x_I \otimes x_J\}$, to $x_{\{i+1, \dots, n\}} \otimes x_\emptyset$. Then, $\chi \cdot q^\emptyset \cdot x_{\{i+1, \dots, n\}} \otimes x_\emptyset = \pi_{\{i+1, \dots, n\} \otimes \emptyset} \circ C \circ Y_i(v_i^+)$.

Using equation (4.3), it follows from weight considerations that

$$\begin{aligned}
\pi_{\{i+1, \dots, n\} \otimes \emptyset} \circ C \circ \Upsilon_i(v_i^+) &= \sum_{\substack{I \cap J = \emptyset \\ I \cup J = \{i+1, \dots, n\}}} q^J \pi_{\{i+1, \dots, n\} \otimes \emptyset} \circ C \cdot (x_I \otimes x_J) \\
&= \frac{q^\emptyset}{[2]} (\Omega_n f \otimes \Omega_n^{-1} f) \cdot x_{\{i+1, \dots, n\}} \otimes x_\emptyset \\
&\quad + \sum_{k=i+1}^n q^{\{k\}} (\Omega_{k-1} \otimes \Omega_{k-1}^{-1}) \circ (\psi_k^* \otimes \psi_k) \cdot x_{\{i+1, \dots, n\} \setminus \{k\}} \otimes x_{\{k\}} \\
&= \frac{\epsilon_{\{i+1, \dots, n\}} \epsilon_\emptyset (q^{-2})^{n-i} (-1)^{n-i}}{[2]} \psi_{\{i+1, \dots, n\}}^* \otimes 1 \\
&\quad + \sum_{k=i+1}^n q^{\{k\}} \epsilon_{\{i+1, \dots, n\} \setminus \{k\}} \epsilon_{\{k\}} (q^{-2})^{k-i-1} (-1)^{k-i-1} \psi_{\{i+1, \dots, n\}}^* \otimes 1 \\
&= \frac{(q^{-2})^{n-i} (-1)^{n-i}}{[2]} x_{\{i+1, \dots, n\}} \otimes x_\emptyset \\
&\quad + \sum_{k=i+1}^n (-q) (-q^2)^{n-k} (q^{-2})^{k-i-1} (-1)^{k-i-1} x_{\{i+1, \dots, n\}} \otimes x_\emptyset \\
&= \frac{(-1)^{n-i}}{[2]} \left(q^{-2(n-i)} + \sum_{k=i+1}^n (q + q^{-1}) q^{2n-4k+2i+3} \right) x_{\{i+1, \dots, n\}} \otimes x_\emptyset \\
&= (-1)^{(n-i)} \frac{[2(n-i)+1]}{[2]} x_{\{i+1, \dots, n\}} \otimes x_\emptyset.
\end{aligned}$$

Thus, $\chi = (-1)^{(n-i)} \frac{[2(n-i)+1]}{[2]}$. □

We now make precise the connection between Wenzl's C and the diagrammatically defined endomorphisms from equation (4.22).

Proposition B.10. In $\text{End}_{U_q(\mathfrak{so}_{2n+1})}(S \otimes S)$, we have $C = \mathbf{H}^{(1)}$.

Proof. Note that $C \cdot x_\emptyset \otimes x_\emptyset = \frac{1}{[2]} x_\emptyset \otimes x_\emptyset$. Thanks to Lemma B.9, the proof of Lemma 4.23 can be applied to C in place of $\mathbf{H}^{(1)}$, in which case we deduce that

$$C = \frac{1}{[2]} \text{id}_{S \otimes S} + \sum_{k=1}^n (-1)^{\binom{k}{2}} \frac{[k][k+1]}{d_k} \mathbf{I}^{(n-k)} = \mathbf{H}^{(1)}.$$

□

APPENDIX C. MORE ON EQUIVARIANT CATEGORIES

In this appendix we discuss the basic structure of equivariant \mathbb{K} -linear categories, with minimal assumptions on the commutative base ring \mathbb{K} . We assume throughout that \mathbb{K} is an integral domain and 2 is invertible in \mathbb{K} .

We have two goals. The first goal is to justify that most of the results of §7, including all the results we used when defining our categorical link invariant, still hold over \mathbb{K} . The second goal is to shed some light on the difficulties involved in computing weighted Grothendieck groups, as they pertain to our conjectures, e.g. the folded skew Howe duality of Conjecture 1.10. Even if these conjectures are proven for $\mathbb{K} = \mathbb{C}$, there are additional issues one would need to overcome to prove the result for other base rings \mathbb{K} , and we wish to point these out.

C.1. When things are nice. We assume we are in the situation of Notation 7.42 henceforth. Fix $b \in \mathbb{B}^{\text{fix}}$. The main technical issue is that the distinguished object X_b need not be equivariantizable.

Since X_b is assumed endopositive, there is a one-dimensional (by abuse of notation, this means “free of rank 1 over \mathbb{K} ”) space of degree zero maps $X_b \rightarrow \sigma(X_b)$. Suppose this space is spanned by $\varphi_b: X_b \rightarrow \sigma(X_b)$, which is necessarily invertible, as otherwise X_b and $\sigma(X_b)$ would not be isomorphic. Since σ is an involution (hence an automorphism), $\sigma(\varphi_b)$ is also invertible, with inverse $\sigma(\varphi_b)^{-1} = \sigma(\varphi_b^{-1})$. Since $\text{End}(X_b) = \mathbb{K} \cdot \text{id}$, we must have that

$$(C.1) \quad \sigma(\varphi_b) \circ \varphi_b = D \cdot \text{id}_{X_b}$$

for $D \in \mathbb{K}^\times$. Note that this implies that $\sigma(\varphi_b^{-1}) = D^{-1}\varphi_b$.

Now, in order to equip X_b with an equivariant structure, we need $c \in \mathbb{K}$ such that $\sigma(c \cdot \varphi) \circ (c \cdot \varphi) = \text{id}_{X_b}$. By (C.1), it follows that an equivariant structure exists if and only if there exists $c \in \mathbb{K}$ such that $c^2 = D^{-1}$. Equivalently, we see that X_b is equivariantizable if and only if D is a square. This is not always the case, although it is guaranteed if we assume \mathbb{K} is an algebraically closed field (hence, the comment after Hypothesis 7.44).

Notation C.1. Let $\sigma\mathbb{B}^{\text{fix}}$ denote the subset of $b \in \mathbb{B}^{\text{fix}}$ such that there exists $\varphi_b: X_b \rightarrow \sigma(X_b)$ so $(X_b, \varphi_b) \in \mathcal{A}^\sigma$.

Clearly, if \mathbb{K} has all square roots, then $\sigma\mathbb{B}^{\text{fix}} = \mathbb{B}^{\text{fix}}$. This equality might still hold over general \mathbb{K} , depending on the structure of the category in question, but it is not guaranteed. In the next section, we will study additional equivariant objects associated to $b \in \mathbb{B}^{\text{fix}} \setminus \sigma\mathbb{B}^{\text{fix}}$ in the event that this set is non-empty. Note that for $b \in \sigma\mathbb{B}^{\text{fix}}$, Corollary 7.36 implies that the equivariant object (X_b, φ_b) is endopositive, as it identifies its endomorphism algebra with a subalgebra of $\text{End}(X_b)$.

We now consider a general construction: the induction functor $\mathcal{A} \rightarrow \mathcal{A}^\sigma$. Given any object $Y \in \mathcal{A}$, define

$$(C.2) \quad \psi_Y: Y \oplus \sigma(Y) \rightarrow \sigma(Y) \oplus Y, \quad \psi_Y := \begin{pmatrix} 0 & \text{id} \\ \text{id} & 0 \end{pmatrix}.$$

It is straightforward to check that $\text{ind} Y := (Y \oplus \sigma(Y), \psi_Y)$ is then an equivariant object. It is straightforward to extend this assignment to morphisms, thus we obtain a functor

$$(C.3) \quad \text{ind}: \mathcal{A} \rightarrow \mathcal{A}^\sigma, \quad Y \mapsto (Y \oplus \sigma(Y), \psi_Y).$$

An equivalent construction, tailored to the setting of Notation 7.42, is as follows. Let $b \in \mathbb{B}$ (whether fixed or not) and choose an isomorphism $\varphi_b: X_{\sigma(b)} \rightarrow \sigma(X_b)$. We then can consider the following morphism in \mathcal{A} :

$$(C.4) \quad \psi_b: X_b \oplus X_{\sigma(b)} \rightarrow \sigma(X_b) \oplus \sigma(X_{\sigma(b)}), \quad \psi_b := \begin{pmatrix} 0 & \varphi_b \\ \sigma(\varphi_b^{-1}) & 0 \end{pmatrix}.$$

It is straightforward to verify that $(X_b \oplus X_{\sigma(b)}, \psi_b) \in \mathcal{A}^\sigma$, and that φ_b determines an isomorphism between this object and $\text{ind} X_b$.

Next, we compute the space of endomorphisms of $(X_b \oplus X_{\sigma(b)}, \psi_b)$ in the case that $b \in \mathbb{B}^{\text{free}}$. Since there are no non-zero degree zero maps between X_b and $X_{\sigma(b)}$, so any degree zero endomorphism of $X_b \oplus X_{\sigma(b)}$ is given by a diagonal matrix

$$\delta = \begin{pmatrix} c_1 \text{id}_{X_b} & 0 \\ 0 & c_2 \text{id}_{\sigma(X_b)} \end{pmatrix}, \quad c_1, c_2 \in \mathbb{K}.$$

One checks that $\psi_b \circ \delta = \sigma(\delta) \circ \psi_b$ if and only if $c_1 = c_2$, thus $\text{End}_{\mathcal{A}^\sigma}^0((X_b \oplus X_{\sigma(b)}, \psi_b)) = \mathbb{K} \cdot \text{id}$. Since X_b and $X_{\sigma(b)}$ are objects in an endopositive family, it follows that there are no endomorphisms of $(X_b \oplus X_{\sigma(b)}, \psi_b)$ of negative degree, hence this object is endopositive. We have thus described the objects appearing in Lemma 7.45, and it is easy to verify the remaining claims appearing therein.

Finally, we study multiplicity spaces and decompositions for the objects, thus proving Proposition 7.50 under the assumption of Hypothesis 7.44. As a byproduct, this also establishes Proposition 7.46.

Fix an equivariant object $(Y, \varphi_Y) \in \mathcal{A}^\sigma$. Suppose $b \in {}^\sigma\mathbb{B}^{\text{fix}}$, and equip X_b with an equivariant structure φ_b . We hence can consider the σ_b^* -action on $\text{Hom}^k(X_b, Y)$, which we may diagonalize, since 2 is assumed invertible.

Suppose we have an inclusion map $\iota \in \text{Hom}^k(X_b, Y)$ lying in the +1-eigenspace, which pairs against a projection map $p: Y \rightarrow X_b$ to give the identity map of X_b . By symmetrizing, we can also assume that p lies in the +1-eigenspace for the σ_b^* action on $\text{Hom}^{-k}(Y, X_b)$, so ι and p induce inclusion and projection maps between (X_b, φ_b) and (Y, φ_Y) . Similarly, inclusion/projection maps in the -1 -eigenspace induce inclusion and projection maps between $(X_b, -\varphi_b)$ and (Y, φ_Y) . Moreover, this argument continues to work for families of orthogonal inclusions and projections. Finally, note that σ_b^* gives a semisimple action of $\mathbb{Z}/2$ on both $\text{Hom}^k(X_b, Y)$ and the kernel of the graded composition pairing, so that an eigenbasis of $V^k(X_b, Y)$ can be lifted to eigenvectors in $\text{Hom}^k(X_b, Y)$.

A separate argument holds for $b \in \mathbb{B}^{\text{free}}$, which does not require the σ^* action (but does involve similar formulae). Fix an isomorphism $\varphi_b: X_{\sigma(b)} \rightarrow \sigma(X_b)$ as above and choose arbitrary maps $\iota \in \text{Hom}^k(X_b, Y)$ and $p \in \text{Hom}^{-k}(Y, X_b)$. We can consider the maps in \mathcal{A}^σ between $(X_b \oplus X_{\sigma(b)}, \psi_b)$ and (Y, φ_Y) given by the matrices

$$(C.5) \quad I(\iota) := \begin{pmatrix} \iota & \varphi_Y^{-1} \circ \sigma(\iota) \circ \varphi_b \end{pmatrix} \quad \text{and} \quad P(p) := \begin{pmatrix} p \\ \varphi_b^{-1} \circ \sigma(p) \circ \varphi_Y \end{pmatrix}.$$

We leave the reader to confirm that these are indeed equivariant maps. Composing, we obtain

$$(C.6) \quad P(p) \circ I(\iota) = \begin{pmatrix} p \circ \iota & p \circ \varphi_Y^{-1} \circ \sigma(\iota) \circ \varphi_b \\ \varphi_b^{-1} \circ \sigma(p) \circ \varphi_Y \circ \iota & \varphi_b^{-1} \circ \sigma(p \circ \iota) \circ \varphi_b \end{pmatrix},$$

which is an endomorphism of $(X_b \oplus X_{\sigma(b)}, \psi_b)$.

As observed above, since $b \in \mathbb{B}^{\text{free}}$, this matrix must be a multiple of the identity, so, in particular, the off-diagonal entries are zero. In fact, we can explicitly compute which multiple. We have $p \circ \iota = \beta_{X_b, Y}^k(p, \iota) \cdot \text{id}_{X_b}$, and so

$$\varphi_b^{-1} \circ \sigma(p \circ \iota) \circ \varphi_b = \beta_{X_b, Y}^k(p, \iota) \cdot \text{id}_{X_{\sigma(b)}}$$

is also the same multiple of the identity. Thus, the composition pairing of $I(\iota)$ and $P(p)$ agrees with the composition pairing of ι and p . Consequently, the multiplicity of X_b in Y exactly equals the multiplicity of $(X_b \oplus X_{\sigma(b)}, \psi_b)$ in (Y, φ_Y) .

Remark C.2. We can say more under a common situation in representation theory, that of the *top summand*.

Let $I \subset \mathbb{B}$. Given an object $Z \in \mathcal{A}(\mathbb{B})$, the condition that $[Z] \in \mathbb{Z}[q^\pm] \cdot I$ is equivalent to the condition that only (shifts of) objects $\{X_b\}_{b \in I}$ appear in the unique decomposition of Z into distinguished summands. The associated *thick ideal* $\mathcal{I} \subset \mathcal{A}(\mathbb{B})$ consists of the morphisms in $\mathcal{A}(\mathbb{B})$ that factor through an object Z such that $[Z] \in \mathbb{Z}[q^\pm] \cdot I$. If I is closed under the action of σ on \mathbb{B} , then \mathcal{I} is invariant under σ .

Now, let (Y, φ_Y) be an equivariant object in $\mathcal{A}^\sigma(\mathbb{B})$ and suppose that $Y = X \oplus Z$ where $X = X_c$ for $c \in \mathbb{B}^{\text{fix}} \setminus I$ and $[Z] \in \mathbb{Z}[q^\pm] \cdot I$. We think of X as the top summand of Y , and Z as consisting of “lower terms.” In this context, let us prove that X is equivariantizable.

Consider X as the image of an idempotent $e \in \text{End}(Y)$, and note that $\varphi_X := \sigma(e) \circ \varphi_Y \circ e$ is a morphism from X to $\sigma(X)$. We compute

$$\begin{aligned} \sigma(\varphi_X) \circ \varphi_X &= e \circ \sigma(\varphi_Y) \circ \sigma(e) \circ \varphi_Y \circ e \\ &= e \circ \sigma(\varphi_Y) \circ \sigma(\text{id}_Y + (e - \text{id}_Y)) \circ \varphi_Y \circ e \\ &= e \circ \sigma(\varphi_Y) \circ \varphi_Y \circ e + e \circ \sigma(\varphi_Y) \circ \sigma(e - \text{id}_Y) \circ \varphi_Y \circ e, \end{aligned}$$

so $\sigma(\varphi_X) \circ \varphi_X \in e + e\mathcal{I}e$. By endopositivity, the degree zero morphisms in $e\mathcal{I}e$ are zero, thus $\sigma(\varphi_X) \circ \varphi_X = e$. Since e is the identity map of X , φ_X is an equivariant structure.

This technique can be used to prove that many distinguished objects are equivariantizable. For example, this provides an alternative route to showing that all endopositive objects in \mathcal{B}_2^n are equivariantizable, via [21, Example 2.21].

C.2. When things are not nice. We continue to assume that we are in the setting of Notation 7.42. In this section we discuss indecomposable objects associated with $b \in \mathbb{B}^{\text{fix}} \setminus \sigma\mathbb{B}^{\text{fix}}$. We do not know where to find an exploration of this theme in the literature.

Fix $b \in \mathbb{B}^{\text{fix}}$, possibly in $\sigma\mathbb{B}^{\text{fix}}$. Once and for all, fix an isomorphism $\varphi_b: X_b \rightarrow \sigma(X_b)$ spanning $\text{Hom}(X_b, \sigma(X_b))$ and define $D \in \mathbb{K}^\times$ by the formula $\sigma(\varphi_b) \circ \varphi_b = D \cdot \text{id}$. Let $\mathbb{X}_b := (X_b \oplus X_b, \psi_b)$ be the equivariant object defined in (C.4). Note that

$$\psi_b = \begin{pmatrix} 0 & \varphi_b \\ \sigma(\varphi_b^{-1}) & 0 \end{pmatrix} = \begin{pmatrix} 0 & \varphi_b \\ D^{-1}\varphi_b & 0 \end{pmatrix}.$$

To begin, we compute the degree zero endomorphisms of \mathbb{X}_b . Since X_b is endopositive, a degree zero endomorphism of $X_b \oplus X_b$ in \mathcal{A} consists of a 2×2 matrix of scalars multiples of id_{X_b} . A computation shows that such a matrix M satisfies $\psi_b \circ M = \sigma(M) \circ \psi_b$ if and only if

$$M = \begin{pmatrix} \alpha & D \cdot \beta \\ \beta & \alpha \end{pmatrix} \cdot \text{id}_{X_b}$$

for some $\alpha, \beta \in \mathbb{K}$. Thus, $\text{End}_{\mathcal{A}^\sigma}^0(\mathbb{X}_b)$ is spanned over \mathbb{K} by the identity and by the map

$$\gamma := \begin{pmatrix} 0 & D \\ 1 & 0 \end{pmatrix} \cdot \text{id}_{X_b}.$$

Since $\gamma^2 = D \cdot \text{id}$, the ring $\text{End}_{\mathcal{A}^\sigma}^0(\mathbb{X}_b)$ is isomorphic to $\mathbb{K}' := \mathbb{K}[x]/(x^2 - D)$, where x acts by γ . Since X_b is endopositive in \mathcal{A} , there are no endomorphisms of \mathbb{X}_b of negative degree.

If D has a square root in \mathbb{K} , or equivalently (as discussed in §C.1 when $b \in \sigma\mathbb{B}^{\text{fix}}$), then it is easy to decompose \mathbb{X}_b as a direct sum of two indecomposable equivariant objects. Explicitly, we have

$$\mathbb{X}_b \cong (X_b, c\varphi_b) \oplus (X_b, -c\varphi_b)$$

where $\pm c$ are the two square roots of D^{-1} . Correspondingly, in this case $\mathbb{K}' = \mathbb{K}[x]/(x^2 - c^{-2}) \cong \mathbb{K} \times \mathbb{K}$.

We focus now on the case when D does not have a square root in \mathbb{K} . In this case $\text{End}_{\mathcal{A}^\sigma}^0(\mathbb{X}_b) \cong \mathbb{K}'$ is a domain and consequently the object \mathbb{X}_b is indecomposable.

Remark C.3. An immediate consequence of this computation is that \mathcal{A}^σ will **not** be positively graded over \mathbb{K} (in the sense of Definition 7.27) if $\sigma\mathbb{B}^{\text{fix}} \neq \mathbb{B}^{\text{fix}}$. Indeed, it has an indecomposable object whose endomorphism ring in degree zero is not $\mathbb{K} \cdot \text{id}$, but rather some degree two extension of \mathbb{K} . Thus, (7.4) fails. It is possible to generalize Definition 7.27 to allow for field extensions, but we will not pursue this matter here.

Let (Y, φ_Y) be an equivariant object. We aim to analyze summands of (Y, φ_Y) of the form \mathbb{X}_b analogously to our treatment in §C.1 of the summands associated to $b \in \mathbb{B}^{\text{free}}$. Fix $b \in \mathbb{B}^{\text{fix}} \setminus \sigma\mathbb{B}^{\text{fix}}$ and define operators $\sigma^* \iota$ on $\iota \in \text{Hom}(X_b, Y)$ and $p \in \text{Hom}(Y, X_b)$ as follows

$$(C.7) \quad \sigma^* \iota := \varphi_Y^{-1} \circ \sigma(\iota) \circ \varphi_b, \quad \sigma^* p := \varphi_b^{-1} \circ \sigma(p) \circ \varphi_Y.$$

Since $\sigma(\varphi_b) \circ \varphi_b = D \cdot \text{id}_{X_b}$ and $D \neq 1$, the pair (X_b, φ_b) is not an equivariant object, so, contrasting (7.6), these operators are not involutions! Instead, a computation analogous to the one in the proof of Proposition 7.35 shows that

$$(C.8) \quad \sigma^*(\sigma^* \iota) = D \cdot \iota, \quad \sigma^*(\sigma^* p) = D^{-1} \cdot p.$$

Hence, we can endow $\text{Hom}(X_b, Y)$ and $\text{Hom}(Y, X_b)$ with \mathbb{K}' -module structures by letting $x \in \mathbb{K}'$ act by σ^* on $\text{Hom}(X_b, Y)$ and $D \cdot \sigma^*$ on $\text{Hom}(Y, X_b)$.

Observe, however, that $\text{End}(X_b)$ is only a \mathbb{K} -module and the graded composition pairing is only \mathbb{K} -linear. Nevertheless, one has the following adjunction

$$(C.9) \quad \beta_{X_b, Y}^k(p, \sigma^* \iota) = D \cdot \beta_{X_b, Y}^k(\sigma^* p, \iota)$$

To see why, note that since $\sigma(\varphi_b) = D \cdot \varphi_b^{-1}$ and $\sigma(\varphi_Y^{-1}) = \varphi_Y$, we have that

$$\sigma(\beta_{X_b, Y}^k(p, \sigma^* \iota)) = \sigma(p \circ \varphi_Y^{-1} \circ \sigma(\iota) \circ \varphi_b) = \sigma(p) \circ \sigma(\varphi_Y^{-1}) \circ \iota \circ \sigma(\varphi_b)$$

is conjugate to

$$D \cdot \beta_{X_b, Y}^k(\sigma^* p, \iota) = D \cdot \varphi_b^{-1} \circ \sigma(p) \circ \varphi_Y \circ \iota = \sigma(\varphi_b) \circ \sigma(p) \circ \sigma(\varphi_Y^{-1}) \circ \iota.$$

Since both the left- and right-hand sides of (C.9) are \mathbb{K} -multiples of the identity map, \mathbb{K} -linearity of σ establishes (C.9).

Now consider the \mathbb{K} -modules $\text{Hom}_{\mathcal{A}^\sigma}^k(\mathbb{X}_b, (Y, \varphi_Y))$ and $\text{Hom}_{\mathcal{A}^\sigma}^{-k}((Y, \varphi_Y), \mathbb{X}_b)$ which become \mathbb{K}' -modules under pre- and post-composition with $\text{End}_{\mathcal{A}^\sigma}^0(\mathbb{X}_b)$. Adapting (C.5) to the present context, define \mathbb{K} -linear maps

$$(C.10) \quad I: \text{Hom}^k(X_b, Y) \rightarrow \text{Hom}_{\mathcal{A}^\sigma}^k(\mathbb{X}_b, (Y, \varphi_Y)), \quad P: \text{Hom}^{-k}(Y, X_b) \rightarrow \text{Hom}_{\mathcal{A}^\sigma}^{-k}((Y, \varphi_Y), \mathbb{X}_b)$$

by

$$I(\iota) := (\iota \quad \sigma^* \iota) \quad \text{and} \quad P(p) := \begin{pmatrix} p \\ \sigma^* p \end{pmatrix}.$$

Using (C.8), we compute

$$(C.11) \quad I(\iota) \circ \gamma = I(\sigma^* \iota), \quad \gamma \circ P(p) = P(D \cdot \sigma^* p),$$

so the maps in (C.10) are in fact \mathbb{K}' -module homomorphisms. The computation (C.6) is unchanged, but now the off-diagonal entries need not be zero. Instead, (C.9) implies that

$$(C.12) \quad P(p) \circ I(\iota) = \beta_{X_b, Y}^k(p, \iota) \cdot \text{id}_{\mathbb{X}_b} + D^{-1} \beta_{X_b, Y}^k(p, \sigma^* \iota) \cdot \gamma.$$

In particular, if one defines the *wacky composition pairing* by

$$(C.13) \quad \beta^k: \text{Hom}^{-k}(Y, X_b) \times \text{Hom}^k(X_b, Y) \rightarrow \mathbb{K}', \quad (p, \iota) \mapsto P(p) \circ I(\iota)$$

then (C.11) shows that this pairing is \mathbb{K}' -bilinear.

We claim the maps (C.10) send the kernels of the graded composition pairing to those of the wacky composition pairing. To see why, note that if $p \circ \iota = 0$ for all p , then it is straightforward to verify that $p \circ (\sigma^* \iota) = 0$ for all p . Equation (C.12) immediately implies that $P(p) \circ I(\iota) = 0$ for all p . A similar argument shows that if $p \circ \iota = 0$ for all ι , then $P(p) \circ I(\iota) = 0$ for all ι . Since the maps (C.10) respect the kernels of composition pairings, one can similarly define a *wacky non-degenerate composition pairing* between $V^{-k}(Y, X_b)$ and $V^k(X_b, Y)$.

If \mathbb{K} is a field, one can continue as follows. The spaces $V^{-k}(Y, X_b)$ and $V^k(X_b, Y)$ are free over \mathbb{K}' , and hence their dimension over \mathbb{K} is even. One can choose a \mathbb{K} basis $\{p_1, xp_1, p_2, xp_2, \dots, p_n, xp_n\}$ for $V^{-k}(Y, X_b)$ where the action of $x \in \mathbb{K}'$ is in rational canonical form. There is a dual basis, and (C.9) implies that x is in (inverse) rational canonical form as well: if ι_1 is dual to p_1 then $\sigma^* \iota_1$ is dual to $\sigma^* p_1$. The elements $\{p_1, \dots, p_n\}$ and $\{\iota_1, \dots, \iota_n\}$ form dual sets as bases over \mathbb{K}' , and one deduces the following result, which extends Proposition 7.50.

Proposition C.4. Suppose \mathbb{K} is a field and let (Y, φ_Y) be an object in \mathcal{A}^σ .

- (1) For $b \in \mathbb{B}^{\text{free}}$, the multiplicity of $\mathbf{q}^k(X_b \oplus X_{\sigma(b)}, \psi_b)$ in (Y, φ_Y) is equal to the multiplicity of $\mathbf{q}^k X_b$ in Y .

- (2) For $b \in \sigma\mathbb{B}^{\text{fix}}$, the multiplicity of $\mathbf{q}^k(X_b, \varphi_b)$ in (Y, φ_Y) equals the dimension of the $+1$ -eigenspace of σ_b^* in $V^k(X_b, Y)$.
- (3) For $b \in \sigma\mathbb{B}^{\text{fix}}$, the multiplicity of $\mathbf{q}^k(X_b, -\varphi_b)$ in (Y, φ_Y) equals the dimension of the -1 -eigenspace of σ_b^* in $V^k(X_b, Y)$.
- (4) For $b \in \mathbb{B}^{\text{fix}} \setminus \sigma\mathbb{B}^{\text{fix}}$, the multiplicity of $\mathbf{q}^k(X_b \oplus X_b, \psi_b)$ in (Y, φ_Y) is equal to half the multiplicity of $\mathbf{q}^k X_b$ in Y , which is equal to the dimension of $V^k(X_b, Y)$ as a \mathbb{K}' vector space.

Recall from Notation 7.49 that $\mathcal{A}^\sigma(\mathbb{B})$ denotes the full subcategory of \mathcal{A}^σ whose objects (Y, φ_Y) satisfy $Y \in \mathcal{A}(\mathbb{B})$. When working over a field, the following version of Proposition 7.46 classifies the indecomposable objects in this category without assuming Hypothesis 7.44.

Proposition C.5. Suppose \mathbb{K} is a field. If (X, φ_X) is an indecomposable object in $\mathcal{A}^\sigma(\mathbb{B})$, then exactly one of the following holds:

- $(X, \varphi_X) \cong \mathbf{q}^k(X_b, \varphi_b)$ for some $b \in \sigma\mathbb{B}^{\text{fix}}$ and $k \in \mathbb{Z}$,
- $(X, \varphi_X) \cong \mathbf{q}^k(X_b, -\varphi_b)$ for some $b \in \sigma\mathbb{B}^{\text{fix}}$ and $k \in \mathbb{Z}$,
- $(X, \varphi_X) \cong \mathbf{q}^k(X_b \oplus X_{\sigma(b)}, \psi_b) \cong \mathbf{q}^k(X_b \oplus X_{\sigma(b)}, -\psi_b)$ for some $b \in \mathbb{B}^{\text{free}}$ and $k \in \mathbb{Z}$, or
- $(X, \varphi_X) \cong \mathbf{q}^k(X_b \oplus X_b, \psi_b) \cong \mathbf{q}^k(X_b \oplus X_b, -\psi_b)$ for some $b \in \mathbb{B}^{\text{fix}} \setminus \sigma\mathbb{B}^{\text{fix}}$ and $k \in \mathbb{Z}$.

Moreover, $\mathcal{A}^\sigma(\mathbb{B})$ is graded Krull–Schmidt.

Proof. We will analyze the possible indecomposable summands of an equivariant object (Y, φ_Y) . Recall the functor $\text{ind}: \mathcal{A} \rightarrow \mathcal{A}^\sigma$ from (C.3) and observe that (Y, φ_Y) is necessarily a summand of $\text{ind} Y$. (Precisely, we have $\text{ind} Y \cong (Y, \varphi_Y) \oplus (Y, -\varphi_Y)$.) We thus consider the possible indecomposable summands of $\text{ind} Y$.

Since ind is additive, if $Y \in \mathcal{A}(\mathbb{B})$, then $\text{ind} Y$ is a direct sum of shifts of $\text{ind} X_b$ for various $b \in \mathbb{B}$. If $b \in \sigma\mathbb{B}^{\text{fix}}$, then (as above) $\text{ind} X_b \cong (X_b, \varphi_b) \oplus (X_b, -\varphi_b)$. Otherwise $\text{ind} X_b$ is either $(X_b \oplus X_{\sigma(b)}, \psi_b)$ for $b \in \mathbb{B}^{\text{free}}$ or $(X_b \oplus X_b, \psi_b)$ for $b \in \mathbb{B}^{\text{fix}} \setminus \sigma\mathbb{B}^{\text{fix}}$. In either of these latter cases, $\text{ind} X_b$ is indecomposable having degree zero endomorphism ring \mathbb{K} or \mathbb{K}' and no negative degree endomorphisms.

Regardless, we have a direct sum decomposition of $\text{ind} Y$ into indecomposable objects whose endomorphism rings are graded local. It follows (e.g. arguing as in [22, Theorem 11.50]) that (Y, φ_Y) must decompose into these indecomposable summands. This implies the graded Krull–Schmidt property, and further shows that if (Y, φ_Y) is itself indecomposable, it must be one of the stated objects. \square

This yields the analogue of Proposition 7.51 in the present setup.

Proposition C.6. Suppose \mathbb{K} is a field. The σ -weighted Grothendieck group $K_0^\sigma(\mathcal{A}^\sigma(\mathbb{B}))$ has basis

$$\{[(X_b, \varphi_b)]_\sigma\}_{b \in \sigma\mathbb{B}^{\text{fix}}}$$

in bijection with $\sigma\mathbb{B}^{\text{fix}}$. Moreover, if (Y, φ_Y) is any object in $\mathcal{A}^\sigma(\mathbb{B})$, then

$$(C.14) \quad [(Y, \varphi_Y)]_\sigma = \sum_{k \in \mathbb{Z}} \sum_{b \in \sigma\mathbb{B}^{\text{fix}}} \text{Tr}(\sigma_b^*|_{V^k(X_b, Y)}) q^k [(X_b, \varphi_b)]_\sigma.$$

\square

Proof. An immediate consequence of Propositions C.4 and C.5. The new indecomposable objects $(X_b \oplus X_b, \psi_b)$ are zero in the weighted Grothendieck group. \square

Remark C.7. When \mathbb{K} is an algebraically closed field, Proposition 7.51 shows that the weighted Grothendieck group has a basis in bijection with \mathbb{B}^{fix} , while, for other fields, Proposition C.6 shows that it has a basis in bijection with the subset $\sigma\mathbb{B}^{\text{fix}}$. Since the latter could be a proper subset, the size of the weighted Grothendieck group can depend on subtle features of \mathbb{K} .

When \mathbb{K} is not a field, we do not know how to imitate the arguments above. Even assuming that $V^k(X_b, Y)$ is free as a \mathbb{K} -module, we do not know how to deduce that $V^k(X_b, Y)$ is free as a \mathbb{K}' -module, or equivalently, that x should act in rational canonical form. As a consequence, we do not know how to deduce that the multiplicity of \mathbb{X}_b in (Y, φ_Y) is as large as expected.

Example C.8. Let $\mathbb{K} = \mathbb{Z}[\frac{1}{2}, \frac{1}{79}]$ and $D = 79$, so that $\mathbb{K}' = \mathbb{Z}[\frac{1}{2}, \frac{1}{79}, \sqrt{79}]$. The ideal $(3, \sqrt{79} - 1)$ inside \mathbb{K}' is not principal, so it is not a free \mathbb{K}' -module. It is, however, a free \mathbb{K} -module of rank 2. The question of extending a free module to a non-free module is related to the ideal class group; this is an example²⁷ where the ideal class group is cyclic of order 3.

Remark C.9. As in the proof of Proposition C.5, we know that every equivariant object (Y, φ_Y) is a direct summand of $\text{ind } Y$, which has a decomposition into various shifts of $\text{ind } X_b$ whenever $Y \in \mathcal{A}(\mathbb{B})$. If \mathbb{K} is local, and $\mathbb{K}[x]/(x^2 - D)$ is local for all D arising from summands X_b of Y , then we can prove Proposition C.5 with the same proof. Otherwise, we do not know how to rule out the possibility of (Y, φ_Y) having even more exotic direct summands than the \mathbb{X}_b .

C.3. Considerations for categorification. Let $\mathcal{A} = \mathcal{B}_m^n$. In the body of this paper, we proved directly that our (endopositive) objects $\mathbf{X}_i^{(k)}$ are fixed by τ and have equivariant structures. Tensor products of these objects also admit equivariant structures; however, it is not obvious that any direct summand of such a tensor product will be equivariantizable.

Question C.10. For a given base ring \mathbb{K} , is every endopositive object in $\text{Kar}(\mathcal{B}_m^n)$ either in a free orbit for τ , or fixed by τ (up to isomorphism) and equivariantizable?

If not, this would complicate the computation of the weighted Grothendieck group, which would depend on subtle properties of \mathbb{K} (not just the characteristic). When $m = 2$, we understand all indecomposable objects in \mathcal{B}_m^n : they all take the form X_b for $b \in \tau \mathbb{B}^{\text{fix}}$. For $m > 2$, there is no comprehensive understanding of the indecomposable objects in \mathcal{B}_m^n . That said, all the direct summands we computed in order to prove the devil's Serre relations in Proposition 8.22 are either in $\tau \mathbb{B}^{\text{fix}}$ or \mathbb{B}^{free} . We are hopeful that $\mathbb{B}^{\text{fix}} = \tau \mathbb{B}^{\text{fix}}$, but, lacking much hard evidence, we find it irresponsible to posit such a conjecture at this time.

REFERENCES

- [1] J. Alexander. A lemma on a system of knotted curves. *Proc. Nat. Acad. Sci. U.S.A.*, 9(3):93–95, 1923.
- [2] H. Bao, P. Shan, W. Wang, and B. Webster. Categorification of quantum symmetric pairs I. *Quantum Topol.*, 9(4):643–714, 2018.
- [3] Huanchen Bao and Weiqiang Wang. Canonical bases arising from quantum symmetric pairs. *Invent. Math.*, 213(3):1099–1177, 2018.
- [4] D. Bar-Natan. Khovanov's homology for tangles and cobordisms. *Geom. Topol.*, 9:1443–1499, 2005. arXiv:0410495.
- [5] A. Beilinson, G. Lusztig, and R. MacPherson. A geometric setting for the quantum deformation of GL_n . *Duke Math. J.*, 61(2):655–677, 1990.
- [6] Anders Bjorner and Brenti Francesco. *Combinatorics of Coxeter Groups*, volume 231 of *Graduate Texts in Mathematics*. Springer-Verlag Berlin Heidelberg, 1 edition, 2005.
- [7] E. Bodish, B. Elias, D. E. V. Rose, and L. Tatham. Type C webs. arXiv:2103.14997.
- [8] E. Bodish and H. Wu. Webs for the quantum orthogonal group. arXiv:2309.03623, 2023.
- [9] Jonathan Brundan, Weiqiang Wang, and Ben Webster. Nil-Brauer categorifies the split ι -quantum group of rank one, 2023. arXiv:2305.05877.
- [10] Jonathan Brundan, Weiqiang Wang, and Ben Webster. The nil-Brauer category, 2023. arXiv:2305.03876.
- [11] S. Cautis. Clasp technology to knot homology via the affine Grassmannian. *Math. Ann.*, 363(3-4):1053–1115, 2012. arXiv:1207.2074.
- [12] S. Cautis and J. Kamnitzer. Knot homology via derived categories of coherent sheaves. II. \mathfrak{sl}_m case. *Invent. Math.*, 174(1):165–232, 2008. arXiv:0710.3216.

²⁷Thanks to Nick Addington, Francis Dunn, Ellen Eischen, and Samantha Platt for finding and communicating this example.

- [13] S. Cautis and J. Kamnitzer. Braiding via geometric Lie algebra actions. *Compos. Math.*, 148(2):464–506, 2012. arXiv:1001.0619.
- [14] S. Cautis, J. Kamnitzer, and A. Licata. Categorical geometric skew Howe duality. *Invent. Math.*, 180(1):111–159, 2010. arXiv:0902.1795.
- [15] S. Cautis, J. Kamnitzer, and S. Morrison. Webs and quantum skew Howe duality. *Math. Ann.*, 360(1-2):351–390, 2014. arXiv:1210.6437.
- [16] V. Chari and A. Pressley. *A guide to quantum groups*. Cambridge Univ. Press, 1994.
- [17] J. Chuang and R. Rouquier. Derived equivalences for symmetric groups and sl_2 -categorification. *Ann. of Math.*, 167:245–298, 2008. arXiv:0407205.
- [18] Pierre Deligne. Letter to P. Etingof. 12/4/1996.
- [19] M. Ehrig, D. Tubbenhauer, and P. Wedrich. Functoriality of colored link homologies. *Proc. London Math. Soc.*, 2018.
- [20] B. Elias. Folding with Soergel bimodules. In *Categorification and higher representation theory*. Contemp. Math., 2017.
- [21] B. Elias and A. Lauda. Trace decategorification of the hecke category. *J. Algebra*, 2016.
- [22] B. Elias, S. Makisumi, U. Thiel, and G. Williamson. *Introduction to Soergel Bimodules*. RSME Springer Series. Springer Cham, 1 edition, 2020.
- [23] A.M. Gavriliuk and A. U. Klimyk. q -deformed orthogonal and pseudo-orthogonal algebras and their representations. *Letters in Mathematical Physics*, 21(3):215–220, 1991.
- [24] Hayashi, Takahiro. q -analogues of Clifford and Weyl algebras—spinor and oscillator representations of quantum enveloping algebras. *Comm. Math. Phys.*, 127(1):129–144, 1990.
- [25] M. Hogancamp, D. E. V. Rose, and P. Wedrich. A skein relation for singular Soergel bimodules. arXiv:2107.08117.
- [26] Jiuzu Hong and Linhui Shen. Tensor invariants, saturation problems, and Dynkin automorphisms. *Adv. Math.*, 285:629–657, 2015.
- [27] N.Z. Iorgov and A.U. Klimyk. Classification theorem on irreducible representations of the q -deformed algebra $U'_q(\mathfrak{so}_n)$. *Int. J. Math. Math. Sci.*, pages 225–262, 2005.
- [28] Jens Carsten Jantzen. *Lectures on Quantum Groups*, volume 6 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, first edition, 1996.
- [29] V. Jones. A polynomial invariant for knots via von Neumann algebras. *Bull. Amer. Math. Soc. (N.S.)*, 12(1):103–111, 1985.
- [30] J. Kamnitzer and P. Tingley. The crystal commutor and Drinfeld’s unitarized R -matrix. *J. Algebraic Combin.*, 29(3):315–335, 2009. arXiv:0707.2248.
- [31] S. M. Khoroshkin and V. N. Tolstoy. The uniqueness theorem for the universal R -matrix. *Lett. Math. Phys.*, 1992.
- [32] M. Khovanov. A categorification of the Jones polynomial. *Duke Math. J.*, 101(3):359–426, 2000. math.QA/9908171.
- [33] M. Khovanov. Link homology and categorification. *ICM Proceedings*, 2:989–999, 2006.
- [34] M. Khovanov and A. Lauda. A diagrammatic approach to categorification of quantum groups III. *Quantum Topology*, 1:1–92, 2010. arXiv:0807.3250.
- [35] M. Khovanov, A. Lauda, M. Mackaay, and M. Stošić. Extended graphical calculus for categorified quantum $sl(2)$. *Mem. Am. Math. Soc.*, 219, 2012. arXiv:1006.2866.
- [36] M. Khovanov and L. Rozansky. Matrix factorizations and link homology. *Fund. Math.*, 199(1):1–91, 2008. arXiv:0401268.
- [37] A. N. Kirillov and N. Reshetikhin. q -Weyl group and a multiplicative formula for universal R -matrices. *Comm. Math. Phys.*, 1990.
- [38] G. Kuperberg. Spiders for rank 2 Lie algebras. *Comm. Math. Phys.*, 180(1):109–151, 1996. arXiv:9712003.
- [39] A. D. Lauda. A categorification of quantum $sl(2)$. *Adv. Math.*, 225:3327–3424, 2008. arXiv:0803.3652.
- [40] A. D. Lauda. Parameters in categorified quantum groups. *Algebras and Representation Theory*, 2020. arXiv:1812.07654.
- [41] T. Le. Integrality and symmetry of quantum link invariants. *Duke Math. J.*, 2000.
- [42] G. Letzter. Subalgebras which appear in quantum Iwasawa decompositions. *Canad. J. Math.*, 49(6):1206–1223, 1997.
- [43] G. Lusztig. *Introduction to quantum groups*, volume 110 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 1993.
- [44] M. Mackaay, M. Stošić, and P. Vaz. A diagrammatic categorification of the q -Schur algebra. *Quantum Topol.*, 4(1):1–75, 2013. arXiv:1008.1348.
- [45] M. Mackaay and Y. Yonezawa. \mathfrak{sl}_N -Web categories. arXiv:1306.6242, 2013.
- [46] C. Manolescu. Link homology theories from symplectic geometry. *Adv. Math.*, 211(1):363–416, 2007. arXiv:0601629.
- [47] A. A. Markov. Über die freie Äquivalenz der geschlossen Zöpfe. *Recueil Math. Moscou*, 1:73–78, 1935.
- [48] H. Matsumoto. Generateurs et relations des groupes de Weyl generalises. *C. R. Acad. Sci Paris*, 258:3419–3422, 1964.

- [49] V. Mazorchuk and C. Stroppel. A combinatorial approach to functorial quantum \mathfrak{sl}_k knot invariants. *Amer. J. Math.*, 131(6):1679–1713, 2009. arXiv:0709.1971.
- [50] Peter J. McNamara and Alistair Savage. The spin Brauer category, 2023. arXiv:2312.11766.
- [51] Scott Morrison, Kevin Walker, and Paul Wedrich. Invariants of 4-manifolds from Khovanov-Rozansky link homology. *Geom. Topol.*, 26(8):3367–3420, 2022. arXiv:1907.12194.
- [52] M. Polyak. Minimal generating sets of Reidemeister moves. *Quantum Topol.*, 1, 2010. arXiv:0908.3127.
- [53] H. Queffelec and D. E. V. Rose. The \mathfrak{sl}_N foam 2-category: a combinatorial formulation of Khovanov-Rozansky homology via categorical skew Howe duality. *Adv. Math.*, 302:1251–1339, 2016. arXiv:1405.5920.
- [54] H. Queffelec and D. E. V. Rose. Sutured annular Khovanov-Rozansky homology. *Trans. Am. Math. Soc.*, 370:1285–1319, 2018. arXiv:1506.08188.
- [55] H. Queffelec, D. E. V. Rose, and A. Sartori. Annular evaluation and link homology. arXiv:1802.04131.
- [56] J. Rasmussen. Khovanov homology and the slice genus. *Invent. Math.*, 182(2):419–447, 2010. arXiv:0402131.
- [57] Q. Ren and M. Willis. Khovanov homology and exotic 4-manifolds. arXiv:2402.10452.
- [58] N. Yu. Reshetikhin and V. G. Turaev. Ribbon graphs and their invariants derived from quantum groups. *Comm. Math. Phys.*, 1:1–26, 1990.
- [59] Nicolai Reshetikhin. Invariants of tangles i. <https://math.berkeley.edu/reshtetik/Publications/QGInv-1-1987.pdf>, 1987.
- [60] Nicolai Reshetikhin. Invariants of tangles ii. <https://math.berkeley.edu/reshtetik/Publications/QGInv-2-1987.pdf>, 1987.
- [61] D. E. V. Rose. A note on the Grothendieck group of an additive category. *Vestn. Chelyab. Gos. Univ. Mat. Mekh. Inform.*, 2015.
- [62] R. Rouquier. Categorification of the braid groups. arXiv:0409593.
- [63] R. Rouquier. 2-Kac-Moody algebras. arXiv:0812.5023, 2008.
- [64] P. Selinger. Autonomous categories in which $A \cong A^*$. In *Proceedings of the 7th International Workshop on Quantum Physics and Logic*, pages 151–160, 2010.
- [65] P. Selinger. A survey of graphical languages for monoidal categories. In Bob Coecke, editor, *New structures for Physics*, volume 813 of *Lecture Notes in Physics*, pages 289–355. Springer, 2011.
- [66] N. Snyder and P. Tingley. The half-twist for $U_q(\mathfrak{g})$ representations. *Algebra & number theory*, 3(7):809–834, 2009.
- [67] M. Stošić. Indecomposable 1-morphisms of \dot{U}_3^+ and the canonical basis of $U_q^+(sl_3)$. arXiv:1105.4458, 2011.
- [68] P. Tingley. *Some results on the crystal commutator and $\widehat{\mathfrak{sl}}_N$ crystals*. PhD thesis, University of California, Berkeley, 2008.
- [69] Valerio Toledano Laredo. A Kohno-Drinfeld theorem for quantum Weyl groups. *Duke Math. J.*, 112(3):421–451, 2002.
- [70] W. Wang. Quantum symmetric pairs. arXiv:2112.10911.
- [71] B. Webster. Canonical bases and higher representation theory. *Compos. Math.*, 151(1):121–166, 2015. arXiv:1209.0051.
- [72] B. Webster. Comparison of canonical bases for Schur and universal enveloping algebras. *Transform. Groups*, 2017.
- [73] Ben Webster. Knot invariants and higher representation theory. *Mem. Amer. Math. Soc.*, 250(1191):v+141, 2017. arXiv:1309.3796.
- [74] H. Wenzl. Dualities for spin representations. arXiv:2005.11299.
- [75] H. Wu. A colored $\mathfrak{sl}(n)$ -homology for links in S^3 . *Diss. Math.*, 499, 2014. arXiv:0907.0695.
- [76] Y. Yonezawa. Quantum $(\mathfrak{sl}_N, \wedge V_n)$ link invariant and matrix factorizations. *Nagoya Math. J.*, 204:69–123, 2011. arXiv:0906.0220.
- [77] W. Zhang. *Relative braid group symmetries on quantum groups*. Phd thesis, University of Virginia, March 2023. Available at <https://doi.org/10.18130/hn08-4h31>.

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, BUILDING 2, CAMBRIDGE, MA 02139-4307, USA

Email address: ebodish@mit.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, FENTON HALL, EUGENE, OR 97403-1222, USA

Email address: belias@uoregon.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH CAROLINA, PHILLIPS HALL CB #3250, UNC-CH, CHAPEL HILL, NC 27599-3250, USA

Email address: davidrose@unc.edu