The Honeymoon Oberwolfach Problem: small cases

MARIE ROSE JERADE and MATEJA ŠAJNA*

University of Ottawa

July 2, 2024

Abstract

The Honeymoon Oberwolfach Problem HOP $(2m_1, 2m_2, \ldots, 2m_t)$ asks the following question. Given $n = m_1 + m_2 + \ldots + m_t$ newlywed couples at a conference and tround tables of sizes $2m_1, 2m_2, \ldots, 2m_t$, is it possible to arrange the 2n participants at these tables for 2n - 2 meals so that each participant sits next to their spouse at every meal, and sits next to every other participant exactly once? A solution to HOP $(2m_1, 2m_2, \ldots, 2m_t)$ is a decomposition of $K_{2n} + (2n - 3)I$, the complete graph K_{2n} with 2n - 3 additional copies of a fixed 1-factor I, into 2-factors, each consisting of disjoint I-alternating cycles of lengths $2m_1, 2m_2, \ldots, 2m_t$.

The Honeymoon Oberwolfach Problem was introduced in a 2019 paper by Lepine and Šajna. The authors conjectured that $HOP(2m_1, 2m_2, \ldots, 2m_t)$ has a solution whenever the obvious necessary conditions are satisfied, and proved the conjecture for several large cases, including the uniform cycle length case $m_1 = \ldots = m_t$, and the small cases with $n \leq 9$. In the present paper, we extend the latter result to all cases with $n \leq 20$ using a computer search.

Keywords: Honeymoon Oberwolfach Problem, 2-factorization, semi-uniform 1-factorization, HOP-colouring-orientation.

1 Introduction

The well-known Oberwolfach Problem, denoted $OP(m_1, \ldots, m_t)$, asks whether $n = m_1 + \ldots + m_t$ participants can be seated at t tables of sizes m_1, \ldots, m_t for several nights in a row so that each participant gets to sit next to every other participant exactly once. Thus, we are asking whether K_n , the complete graph on n vertices, admits a 2-factorization such that each 2-factor is a disjoint union of t cycles of lengths m_1, \ldots, m_t . The problem has been solved in many special cases — see [1, 2, 3, 5, 6, 7, 8, 9, 10, 12, 13] — but is in general still open.

In the 2019 paper [11], Lepine and the second author introduced a new variant of the Oberwolfach Problem, called the Honeymoon Oberwolfach Problem. This problem, denoted $HOP(m_1, \ldots, m_t)$, can be described as follows. We have $n = \frac{1}{2}(m_1 + \ldots + m_t)$ newlywed

^{*}Corresponding author. Email: msajna@uottawa.ca. Mailing address: Department of Mathematics and Statistics, University of Ottawa, Ottawa, ON, Canada.

couples attending a conference and t tables of sizes m_1, \ldots, m_t (where each $m_i \ge 3$). Is it possible to arrange the participants at these t round tables on 2n - 2 consecutive nights so that each couple sit together every night, and every participant sits next to every other participant exactly once?

In graph-theoretic terms we are asking whether $K_{2n} + (2n-3)I$, the multigraph obtained from the complete graph K_{2n} by adjoining 2n - 3 additional copies of a chosen 1-factor I, admits a decomposition into 2-factors, each a vertex-disjoint union of cycles of lengths m_1, \ldots, m_t , so that in each of these cycles, every other edge is a copy of an edge of I. A solution to $HOP(m_1, m_2, \ldots, m_t)$ is equivalent to a *semi-uniform 1-factorization of* K_{2n} of $type (m_1, m_2, \ldots, m_t)$; that is, a 1-factorization $\{F_1, F_2, \ldots, F_{2n-1}\}$ such that for all $i \neq 1$, the 2-factor $F_1 \cup F_i$ consists of disjoint cycles of lengths m_1, m_2, \ldots, m_t .

If $m_1 = \ldots = m_t = m$ and tm = 2n, then the symbol HOP (m_1, \ldots, m_t) is abbreviated as HOP(2n; m). Note that if HOP (m_1, \ldots, m_t) has a solution, then the m_i are all even and at least 4; these are the obvious necessary conditions.

In [11], the authors proposed the following conjecture.

Conjecture 1.1 [11] The obvious necessary conditions for $HOP(m_1, \ldots, m_t)$ to have a solution are also sufficient.

They also proved the conjecture in the following cases.

Theorem 1.2 [11] Let m and n be positive integers, $2 \le m \le n$. Then HOP(2n; 2m) has a solution if and only if $n \equiv 0 \pmod{m}$.

Theorem 1.3 [11] Let $2 \leq m_1 \leq \ldots \leq m_t$ be integers, and $n = m_1 + \ldots + m_t$. Then $HOP(2m_1, \ldots, 2m_t)$ has a solution in each of the following cases.

- (i) $m_i \equiv 0 \pmod{4}$ for all i.
- (ii) n is odd and $OP(m_1, \ldots, m_t)$ has a solution.
- (iii) n is odd and t = 2.
- (iv) n is odd, n < 40, and $m_1 \ge 3$.
- (v) $n \le 9$.

We remark that case (iv) of Theorem 1.3 is extended to $n \leq 60$ by the result of [12]. In this paper, we extend case (v) of Theorem 1.3, thus proving the following result.

Theorem 1.4 Let m_1, \ldots, m_t be integers with $m_i \ge 2$ for all i, and $n = m_1 + \ldots + m_t$ such that $n \le 20$. Then $HOP(2m_1, \ldots, 2m_t)$ has a solution.

To prove Theorem 1.4, we use the approach described in [11], combined with a computerassisted search. In Sections 2 and 3, we present the relevant terminology and tools from [11], in Section 4, we give the framework of the proof of Theorem 1.4, while in the appendices, we list the computational results supporting the proof.

2 Terminology

In this paper, graphs may contain parallel edges, but not loops. As usual, K_n and λK_n denote the complete graph and the λ -fold complete graph, respectively, on n vertices. For $m \geq 2$, the symbol C_m denotes the cycle of length m, or m-cycle.

Let G be a graph, and let H_1, \ldots, H_t be subgraphs of G. The collection $\{H_1, \ldots, H_t\}$ is called a *decomposition* of G if $\{E(H_1), \ldots, E(H_t)\}$ is a partition of E(G).

An *r*-factor in a graph G is an *r*-regular spanning subgraph of G, and an *r*-factorization of G is a decomposition of G into *r*-factors. A 2-factor of G consisting of disjoint cycles of lengths m_1, \ldots, m_t , respectively, is called a $(C_{m_1}, \ldots, C_{m_t})$ -factor of G, and a decomposition into $(C_{m_1}, \ldots, C_{m_t})$ -factors is called a $(C_{m_1}, \ldots, C_{m_t})$ -factorization of G.

For a positive integer n and $S \subseteq \mathbb{Z}_n^*$ such that S = -S, we define a *circulant* Circ(n; S) as the graph with vertex set $\{x_i : i \in \mathbb{Z}_n\}$ and edge set $\{x_i x_{i+d} : i \in \mathbb{Z}_n, d \in S\}$. An edge of the form $x_i x_{i+d}$ is said to be of *difference* d. Note that an edge of difference d is also of difference n - d, so we may assume that each difference is in $\{1, 2, \ldots, \lfloor \frac{n}{2} \rfloor\}$.

In this paper, the complete graph K_n will be viewed as the join of the circulant $\operatorname{Circ}(n-1;\mathbb{Z}_{n-1}^*)$ and the complete graph K_1 with vertex x_{∞} . Thus, $V(K_n) = \{x_i : i \in \mathbb{Z}_{n-1}\} \cup \{x_{\infty}\}$ and $E(K_n) = \{x_i x_j : i, j \in \mathbb{Z}_{n-1}, i \neq j\} \cup \{x_i x_{\infty} : i \in \mathbb{Z}_{n-1}\}$. If this is the case, then an edge of the form $x_i x_{\infty}$ will be called of *difference infinity*.

Let I be a chosen 1-factor in the graph K_{2n} . An edge of K_{2n} is said to be an *I-edge* if it belongs to E(I), and a *non-I-edge* otherwise. The symbol $K_{2n} + \lambda I$ denotes the graph K_{2n} with λ additional copies of each *I*-edge, for a total of $\lambda + 1$ copies of each *I*-edge. (Note that these additional copies of *I*-edges of K_{2n} are then also considered to be *I*-edges of $K_{2n} + \lambda I$.) A cycle C of $K_{2n} + \lambda I$, necessarily of even length, is said to be *I*-alternating if the *I*-edges and non-*I*-edges along C alternate. A 2-factor (or 2-factorization) of $K_{2n} + \lambda I$ is said to be *I*-alternating if each of its cycles is *I*-alternating.

Thus, a solution to the Honeymoon Oberwolfach Problem HOP (m_1, \ldots, m_t) is an *I*-alternating $(C_{m_1}, \ldots, C_{m_t})$ -factorization of $K_{2n} + (2n-3)I$ for $2n = m_1 + m_2 + \ldots + m_t$.

3 The Tools

As in [11], we use the symbol $4K_n^{\bullet}$ to denote the 4-fold complete graph with *n* vertices whose edges are coloured pink, blue, and black, and black edges are oriented so that each 4-set of parallel edges contains one pink edge, one blue edge, and two opposite black arcs.

Definition 3.1 [11] A 2-factorization \mathcal{D} of $4K_n^{\bullet}$ is said to be *HOP* if each cycle of \mathcal{D} satisfies the following condition:

(C) any two adjacent (that is, consecutive) edges satisfy one of the following:

- one is blue and the other pink; or
- both are black and directed in the same way;

- one is blue and the other black, directed towards the blue edge; or
- one is pink and the other black, directed away from the pink edge.

Theorem 3.2 [11] Let m_1, \ldots, m_t be integers greater than 1, and let $n = m_1 + \ldots + m_t$. Then $HOP(2m_1, \ldots, 2m_t)$ has a solution if and only if $4K_n^{\bullet}$ admits an $HOP(C_{m_1}, \ldots, C_{m_t})$ -factorization.

In the next proposition, the symbol $2K_n^{\circ}$ denotes the multigraph $2K_n$ whose edges are coloured pink and black so that each 2-set of parallel edges contains one pink edge and one black edge.

Proposition 3.3 [11] Assume *n* is even, and let the vertex set of $2K_n^{\circ}$ be $\{x_i : i \in \mathbb{Z}_{n-1}\} \cup \{x_{\infty}\}$. Let ρ be the permutation $\rho = (x_{\infty})(x_0 \ x_1 \ x_2 \ \dots \ x_{n-2})$, and let ρ_{\circ} denote the permutation on the edge set of $2K_n^{\circ}$ that is induced by ρ and that preserves the colour of the edges.

Suppose $2K_n^{\circ}$ admits a $(C_{m_1}, \ldots, C_{m_t})$ -factor F such that

(A1) each cycle in F of length at least 3 contains an even number of pink edges, and

(A2) F contains exactly one edge from each of the orbits of $\langle \rho_{\circ} \rangle$.

Then $4K_n^{\bullet}$ admits an HOP $(C_{m_1}, \ldots, C_{m_t})$ -factorization.

Proposition 3.4 [11] Assume *n* is even, and let the vertex set of $4K_n^{\bullet}$ be $\{x_i : i \in \mathbb{Z}_{n-1}\} \cup \{x_{\infty}\}$. Let ρ be the permutation $\rho = (x_{\infty})(x_0 \ x_1 \ x_2 \ \dots \ x_{n-2})$, and let ρ_{\bullet} denote the permutation on the edge set of $4K_n^{\bullet}$ that is induced by ρ and that preserves the colour (and orientation) of the edges.

Suppose $4K_n^{\bullet}$ admits edge-disjoint $(C_{m_1}, \ldots, C_{m_t})$ -factors F_1 and F_2 such that

(D1) each cycle in F_1 and F_2 satisfies Condition (C) in Definition 3.1, and

(D2) F_1 and F_2 jointly contain exactly one edge from each of the orbits of $\langle \rho_{\bullet} \rangle$.

Then $\mathcal{D} = \{\rho_{\bullet}^{i}(F_{1}), \rho_{\bullet}^{i}(F_{2}) : i \in \mathbb{Z}_{n-1}\}$ is an HOP $(C_{m_{1}}, \ldots, C_{m_{t}})$ -factorization of $4K_{n}^{\bullet}$.

Proposition 3.5 [11] Assume n is odd, and let the vertex set of $4K_n^{\bullet}$ be $\{x_i : i \in \mathbb{Z}_{n-1}\} \cup \{x_{\infty}\}$. Let ρ be the permutation $\rho = (x_{\infty})(x_0 \ x_1 \ x_2 \ \dots \ x_{n-2})$, and let ρ_{\bullet} denote the permutation on the edge set of $4K_n^{\bullet}$ that is induced by ρ and that preserves the colour (and orientation) of the edges.

Suppose $4K_n^{\bullet}$ admits pairwise edge-disjoint $(C_{m_1}, \ldots, C_{m_t})$ -factors F_1 , F_2 , and F_3 such that

(E1) each cycle in F_1 , F_2 , and F_3 satisfies Condition (C) in Definition 3.1;

(E2) each orbit of $\langle \rho_{\bullet} \rangle$ has edges either in $F_1 \cup F_2$ or in F_3 ;

(E3) if $e \in E(F_1 \cup F_2)$, then $\rho_{\bullet}^{\frac{n-1}{2}}(e) \in E(F_1 \cup F_2)$; and

(E4) $F_1 \cup F_2$ contains a pink and a blue edge of difference $\frac{n-1}{2}$.

Then $\mathcal{D} = \{\rho_{\bullet}^{i}(F_{1}), \rho_{\bullet}^{i}(F_{2}) : i = 0, 1, \dots, \frac{n-3}{2}\} \cup \{\rho_{\bullet}^{i}(F_{3}) : i \in \mathbb{Z}_{n-1}\} \text{ is an HOP } (C_{m_{1}}, \dots, C_{m_{t}}) - factorization of <math>4K_{n}^{\bullet}$.

The required 2-factors F from Proposition 3.3, F_1 and F_2 from Proposition 3.4, and F_1 , F_2 , and F_3 from Proposition 3.5 will are called the *starter 2-factors* (or *starters*) of the resulting 2-factorizations. Thus, we are referring to Propositions 3.3, 3.4, and 3.5 as the one-starter, two-starter, and three-starter approach, respectively.

4 Proof of Theorem 1.4

By the type of a $(C_{m_1}, \ldots, C_{m_t})$ -factor we mean the multiset $[m_1, \ldots, m_t]$.

PROOF. By Theorem 1.3(v), it suffices to consider $10 \le n \le 20$. For each value of n, we list all possible 2-factor types $[m_1, \ldots, m_t]$, and refer to the result that guarantees existence of a solution to HOP $(2m_1, \ldots, 2m_t)$. In cases where we are referring to Proposition 3.3, 3.4, or 3.5, appropriate starter 2-factors are given in the appendices.

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[2, 2, 2, 2, 2]	Theorem 1.2
[4, 2, 2, 2]	Proposition 3.3 (one starter)
[3, 3, 2, 2]	Proposition 3.4 (two starters)
[6, 2, 2]	Proposition 3.4 (two starters)
[5, 3, 2]	Proposition 3.3 (one starter)
[4, 4, 2]	Proposition 3.3 (one starter)
[4, 3, 3]	Proposition 3.4 (two starters)
[8, 2]	Proposition 3.3 (one starter)
[7,3]	Proposition 3.4 (two starters)
[6, 4]	Proposition 3.4 (two starters)
[5, 5]	Theorem 1.2
[10]	Theorem 1.2

CASE n = 10: see Appendix B.

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[3, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[5, 2, 2, 2]	Proposition 3.5 (three starters)
[4, 3, 2, 2]	Proposition 3.5 (three starters)
[3, 3, 3, 2]	Proposition 3.5 (three starters)
[7, 2, 2]	Proposition 3.5 (three starters)
[6, 3, 2]	Proposition 3.5 (three starters)
[5, 4, 2]	Proposition 3.5 (three starters)
[5, 3, 3]	Theorem 1.2
[4, 4, 3]	Theorem 1.2
[9, 2]	Theorem 1.2
[8,3]	Theorem 1.2
[7, 4]	Theorem 1.2
[6, 5]	Theorem 1.2
[11]	Theorem 1.2

CASE n = 11: see Appendix C.

CASE n = 12: see Appendix D.

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[2, 2, 2, 2, 2, 2]	Theorem 1.2
[4, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[3, 3, 2, 2, 2]	Proposition 3.4 (two starters)
[6, 2, 2, 2]	Proposition 3.4 (two starters)
[5, 3, 2, 2]	Proposition 3.3 (one starter)
[4, 4, 2, 2]	Proposition 3.3 (one starter)
[4, 3, 3, 2]	Proposition 3.4 (two starters)
[3, 3, 3, 3]	Theorem 1.2
[8, 2, 2]	Proposition 3.3 (one starter)
[7, 3, 2]	Proposition 3.4 (two starters)
[6, 4, 2]	Proposition 3.4 (two starters)
[5, 5, 2]	Proposition 3.4 (two starters)
[6, 3, 3]	Proposition 3.3 (one starter)
[5, 4, 3]	Proposition 3.3 (one starter)
[4, 4, 4]	Theorem 1.2
[10, 2]	Proposition 3.4 (two starters)
[9,3]	Proposition 3.3 (one starter)
[8, 4]	Theorem 1.2
[7, 5]	Proposition 3.3 (one starter)
[6, 6]	Theorem 1.2
[12]	Theorem 1.2

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[3, 2, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[5, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[4, 3, 2, 2, 2]	Proposition 3.5 (three starters)
[3, 3, 3, 2, 2]	Proposition 3.5 (three starters)
[7, 2, 2, 2]	Proposition 3.5 (three starters)
[6, 3, 2, 2]	Proposition 3.5 (three starters)
[5, 4, 2, 2]	Proposition 3.5 (three starters)
[5, 3, 3, 2]	Proposition 3.5 (three starters)
[4, 4, 3, 2]	Proposition 3.5 (three starters)
[4, 3, 3, 3]	Theorem 1.2
[9, 2, 2]	Proposition 3.5 (three starters)
[8, 3, 2]	Proposition 3.5 (three starters)
[7, 4, 2]	Proposition 3.5 (three starters)
[6, 5, 2]	Proposition 3.5 (three starters)
[7, 3, 3]	Theorem 1.2
[6, 4, 3]	Theorem 1.2
[5, 5, 3]	Theorem 1.2
[5, 4, 4]	Theorem 1.2
[11, 2]	Theorem 1.2
[10,3]	Theorem 1.2
[9, 4]	Theorem 1.2
[8,5]	Theorem 1.2
[7, 6]	Theorem 1.2
[13]	Theorem 1.2

CASE n = 13: see Appendix E.

CASE n = 14: see Appendix F.

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[2, 2, 2, 2, 2, 2, 2]	Theorem 1.2
[4, 2, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[3, 3, 2, 2, 2, 2]	Proposition 3.4 (two starters)
[6, 2, 2, 2, 2]	Proposition 3.4 (two starters)
[5, 3, 2, 2, 2]	Proposition 3.3 (one starter)
[4, 4, 2, 2, 2]	Proposition 3.3 (one starter)
[4, 3, 3, 2, 2]	Proposition 3.4 (two starters)
[3, 3, 3, 3, 2]	Proposition 3.3 (one starter)
[8, 2, 2, 2]	Proposition 3.3 (one starter)
[7, 3, 2, 2]	Proposition 3.4 (two starters)
[6, 4, 2, 2]	Proposition 3.4 (two starters)
[5, 5, 2, 2]	Proposition 3.4 (two starters)
[6, 3, 3, 2]	Proposition 3.3 (one starter)
[5, 4, 3, 2]	Proposition 3.3 (one starter)
[4, 4, 4, 2]	Proposition 3.3 (one starter)
[5, 3, 3, 3]	Proposition 3.4 (two starters)
[4, 4, 3, 3]	Proposition 3.4 (two starters)

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[10, 2, 2]	Proposition 3.4 (two starters)
[9, 3, 2]	Proposition 3.3 (one starter)
[8, 4, 2]	Proposition 3.3 (one starter)
[7, 5, 2]	Proposition 3.3 (one starter)
[6, 6, 2]	Proposition 3.3 (one starter)
[8, 3, 3]	Proposition 3.4 (two starters)
[7, 4, 3]	Proposition 3.4 (two starters)
[6, 5, 3]	Proposition 3.4 (two starters)
[6, 4, 4]	Proposition 3.4 (two starters)
[5, 5, 4]	Proposition 3.4 (two starters)
[12, 2]	Proposition 3.3 (one starter)
[11, 3]	Proposition 3.4 (two starters)
[10, 4]	Proposition 3.4 (two starters)
[9, 5]	Proposition 3.4 (two starters)
[8, 6]	Proposition 3.4 (two starters)
[7,7]	Theorem 1.2
[14]	Theorem 1.2

CASE n = 15: see Appendix G.

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[3, 2, 2, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[5, 2, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[4, 3, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[3, 3, 3, 2, 2, 2]	Proposition 3.5 (three starters)
[7, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[6, 3, 2, 2, 2]	Proposition 3.5 (three starters)
[5, 4, 2, 2, 2]	Proposition 3.5 (three starters)
[5, 3, 3, 2, 2]	Proposition 3.5 (three starters)
[4, 4, 3, 2, 2]	Proposition 3.5 (three starters)
[4, 3, 3, 3, 2]	Proposition 3.5 (three starters)
[3, 3, 3, 3, 3]	Theorem 1.2
[9, 2, 2, 2]	Proposition 3.5 (three starters)
[8, 3, 2, 2]	Proposition 3.5 (three starters)
[7, 4, 2, 2]	Proposition 3.5 (three starters)
[6, 5, 2, 2]	Proposition 3.5 (three starters)
[7, 3, 3, 2]	Proposition 3.5 (three starters)
[6, 4, 3, 2]	Proposition 3.5 (three starters)
[5, 5, 3, 2]	Proposition 3.5 (three starters)
[5, 4, 4, 2]	Proposition 3.5 (three starters)
[6, 3, 3, 3]	Theorem 1.2
[5, 4, 3, 3]	Theorem 1.2
[4, 4, 4, 3]	Theorem 1.2

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[11, 2, 2]	Proposition 3.5 (three starters)
[10, 3, 2]	Proposition 3.5 (three starters)
[9, 4, 2]	Proposition 3.5 (three starters)
[8, 5, 2]	Proposition 3.5 (three starters)
[7, 6, 2]	Proposition 3.5 (three starters)
[9, 3, 3]	Theorem 1.2
[8, 4, 3]	Theorem 1.2
[7, 5, 3]	Theorem 1.2
[6, 6, 3]	Theorem 1.2
[7, 4, 4]	Theorem 1.2
[6, 5, 4]	Theorem 1.2
[5, 5, 5]	Theorem 1.2
[13, 2]	Theorem 1.2
[12, 3]	Theorem 1.2
[11, 4]	Theorem 1.2
[10, 5]	Theorem 1.2
[9, 6]	Theorem 1.2
[8,7]	Theorem 1.2
[15]	Theorem 1.2

CASE n = 16: see Appendix H.

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[2, 2, 2, 2, 2, 2, 2, 2, 2]	Theorem 1.2
[4, 2, 2, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[3, 3, 2, 2, 2, 2, 2]	Proposition 3.4 (two starters)
[6, 2, 2, 2, 2, 2]	Proposition 3.4 (two starters)
[5, 3, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[4, 4, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[4, 3, 3, 2, 2, 2]	Proposition 3.4 (two starters)
[3, 3, 3, 3, 2, 2]	Proposition 3.3 (one starter)
[8, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[7, 3, 2, 2, 2]	Proposition 3.4 (two starters)
[6, 4, 2, 2, 2]	Proposition 3.4 (two starters)
[5, 5, 2, 2, 2]	Proposition 3.4 (two starters)
[6, 3, 3, 2, 2]	Proposition 3.3 (one starter)
[5, 4, 3, 2, 2]	Proposition 3.3 (one starter)
[4, 4, 4, 2, 2]	Proposition 3.3 (one starter)
[5, 3, 3, 3, 2]	Proposition 3.4 (two starters)
[4, 4, 3, 3, 2]	Proposition 3.4 (two starters)
[4, 3, 3, 3, 3]	Proposition 3.3 (one starter)
[10, 2, 2, 2]	Proposition 3.4 (two starters)
[9, 3, 2, 2]	Proposition 3.3 (one starter)
[8, 4, 2, 2]	Proposition 3.3 (one starter)
[7, 5, 2, 2]	Proposition 3.3 (one starter)
[6, 6, 2, 2]	Proposition 3.3 (one starter)

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[8, 3, 3, 2]	Proposition 3.4 (two starters)
[7, 4, 3, 2]	Proposition 3.4 (two starters)
[6, 5, 3, 2]	Proposition 3.4 (two starters)
[6, 4, 4, 2]	Proposition 3.4 (two starters)
[5, 5, 4, 2]	Proposition 3.4 (two starters)
[7, 3, 3, 3]	Proposition 3.3 (one starter)
[6, 4, 3, 3]	Proposition 3.3 (one starter)
[5, 5, 3, 3]	Proposition 3.3 (one starter)
[5, 4, 4, 3]	Proposition 3.3 (one starter)
[4, 4, 4, 4]	Theorem 1.2
[12, 2, 2]	Proposition 3.3 (one starter)
[11, 3, 2]	Proposition 3.4 (two starters)
[10, 4, 2]	Proposition 3.4 (two starters)
[9, 5, 2]	Proposition 3.4 (two starters)
[8, 6, 2]	Proposition 3.4 (two starters)
[7, 7, 2]	Proposition 3.4 (two starters)
[10, 3, 3]	Proposition 3.3 (one starter)
[9, 4, 3]	Proposition 3.3 (one starter)
[8, 5, 3]	Proposition 3.3 (one starter)
[7, 6, 3]	Proposition 3.3 (one starter)
[8, 4, 4]	Theorem 1.2
[7, 5, 4]	Proposition 3.3 (one starter)
[6, 6, 4]	Proposition 3.3 (one starter)
[6, 5, 5]	Proposition 3.3 (one starter)
[14, 2]	Proposition 3.4 (two starters)
[13, 3]	Proposition 3.3 (one starter)
[12, 4]	Theorem 1.2
[11, 5]	Proposition 3.3 (one starter)
[10, 6]	Proposition 3.3 (one starter)
[9,7]	Proposition 3.3 (one starter)
[8,8]	Theorem 1.2
[16]	Theorem 1.2

CASE n = 17: see Appendix I.

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[3, 2, 2, 2, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[5, 2, 2, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[4, 3, 2, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[3, 3, 3, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[7, 2, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[6, 3, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[5, 4, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[5, 3, 3, 2, 2, 2]	Proposition 3.5 (three starters)
[4, 4, 3, 2, 2, 2]	Proposition 3.5 (three starters)
[4, 3, 3, 3, 2, 2]	Proposition 3.5 (three starters)
[3, 3, 3, 3, 3, 3, 2]	Proposition 3.5 (three starters)

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[9, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[8, 3, 2, 2, 2]	Proposition 3.5 (three starters)
[7, 4, 2, 2, 2]	Proposition 3.5 (three starters)
[6, 5, 2, 2, 2]	Proposition 3.5 (three starters)
[7, 3, 3, 2, 2]	Proposition 3.5 (three starters)
[6, 4, 3, 2, 2]	Proposition 3.5 (three starters)
[5, 5, 3, 2, 2]	Proposition 3.5 (three starters)
[5, 4, 4, 2, 2]	Proposition 3.5 (three starters)
[6, 3, 3, 3, 2]	Proposition 3.5 (three starters)
[5, 4, 3, 3, 2]	Proposition 3.5 (three starters)
[4, 4, 4, 3, 2]	Proposition 3.5 (three starters)
[5, 3, 3, 3, 3]	Theorem 1.2
[4, 4, 3, 3, 3]	Theorem 1.2
[11, 2, 2, 2]	Proposition 3.5 (three starters)
[10, 3, 2, 2]	Proposition 3.5 (three starters)
[9, 4, 2, 2]	Proposition 3.5 (three starters)
$\begin{bmatrix} 0, 2, 2, -1 \end{bmatrix}$ [8 5 2 2]	Proposition 3.5 (three starters)
[7, 6, 2, 2]	Proposition 3.5 (three starters)
[1, 3, 2, 2] [9, 3, 3, 2]	Proposition 3.5 (three starters)
[8, 4, 3, 2]	Proposition 3.5 (three starters)
[0, 2, 3, 2]	Proposition 3.5 (three starters)
[6, 6, 3, 2]	Proposition 3.5 (three starters)
[7, 4, 4, 2]	Proposition 3.5 (three starters)
[6, 5, 4, 2]	Proposition 3.5 (three starters)
[5, 5, 5, 2]	Proposition 3.5 (three starters)
[8, 3, 3, 3]	Theorem 1.2
[7, 4, 3, 3]	Theorem 1.2
[6, 5, 3, 3]	Theorem 1.2
[6, 4, 4, 3]	Theorem 1.2
[5, 5, 4, 3]	Theorem 1.2
[5, 4, 4, 4]	Theorem 1.2
[13, 2, 2]	Proposition 3.5 (three starters)
[12, 3, 2]	Proposition 3.5 (three starters)
[11, 4, 2]	Proposition 3.5 (three starters)
[10, 5, 2]	Proposition 3.5 (three starters)
[9, 6, 2]	Proposition 3.5 (three starters)
[8, 7, 2]	Proposition 3.5 (three starters)
[11, 3, 3]	Theorem 1.2
[10, 4, 3]	Theorem 1.2
[9, 5, 3]	Theorem 1.2
[8, 6, 3]	Theorem 1.2
[7, 7, 3]	Theorem 1.2
[9, 4, 4]	Theorem 1.2
[8, 5, 4]	Theorem 1.2
[7, 6, 4]	Theorem 1.2
[7, 5, 5]	Theorem 1.2
[6, 6, 5]	Theorem 1.2

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[15, 2]	Theorem 1.2
[14, 3]	Theorem 1.2
[13, 4]	Theorem 1.2
[12, 5]	Theorem 1.2
[11, 6]	Theorem 1.2
[10, 7]	Theorem 1.2
[9, 8]	Theorem 1.2
[17]	Theorem 1.2

CASE n = 18: see Appendix J.

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[2, 2, 2, 2, 2, 2, 2, 2, 2]	Theorem 1.2
[4, 2, 2, 2, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[3, 3, 2, 2, 2, 2, 2, 2]	Proposition 3.4 (two starters)
[6, 2, 2, 2, 2, 2, 2]	Proposition 3.4 (two starters)
[5, 3, 2, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[4, 4, 2, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[4, 3, 3, 2, 2, 2, 2]	Proposition 3.4 (two starters)
[3, 3, 3, 3, 2, 2, 2]	Proposition 3.3 (one starter)
[8, 2, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[7, 3, 2, 2, 2, 2]	Proposition 3.4 (two starters)
[6, 4, 2, 2, 2, 2]	Proposition 3.4 (two starters)
[5, 5, 2, 2, 2, 2]	Proposition 3.4 (two starters)
[6, 3, 3, 2, 2, 2]	Proposition 3.3 (one starter)
[5, 4, 3, 2, 2, 2]	Proposition 3.3 (one starter)
[4, 4, 4, 2, 2, 2]	Proposition 3.3 (one starter)
[5, 3, 3, 3, 2, 2]	Proposition 3.4 (two starters)
[4, 4, 3, 3, 2, 2]	Proposition 3.4 (two starters)
[4, 3, 3, 3, 3, 2]	Proposition 3.3 (one starter)
[3, 3, 3, 3, 3, 3]	Theorem 1.2
[10, 2, 2, 2, 2]	Proposition 3.4 (two starters)
[9, 3, 2, 2, 2]	Proposition 3.3 (one starter)
[8, 4, 2, 2, 2]	Proposition 3.3 (one starter)
[7, 5, 2, 2, 2]	Proposition 3.3 (one starter)
[6, 6, 2, 2, 2]	Proposition 3.3 (one starter)
[8, 3, 3, 2, 2]	Proposition 3.4 (two starters)
[7, 4, 3, 2, 2]	Proposition 3.4 (two starters)
[6, 5, 3, 2, 2]	Proposition 3.4 (two starters)
[6, 4, 4, 2, 2]	Proposition 3.4 (two starters)
[5, 5, 4, 2, 2]	Proposition 3.4 (two starters)
[7, 3, 3, 3, 2]	Proposition 3.3 (one starter)
[6, 4, 3, 3, 2]	Proposition 3.3 (one starter)
[5, 5, 3, 3, 2]	Proposition 3.3 (one starter)
[5, 4, 4, 3, 2]	Proposition 3.3 (one starter)
[4, 4, 4, 4, 2]	Proposition 3.3 (one starter)

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[6, 3, 3, 3, 3]	Proposition 3.4 (two starters)
[5, 4, 3, 3, 3]	Proposition 3.4 (two starters)
[4, 4, 4, 3, 3]	Proposition 3.4 (two starters)
[12, 2, 2, 2]	Proposition 3.3 (one starter)
[11, 3, 2, 2]	Proposition 3.4 (two starters)
[10, 4, 2, 2]	Proposition 3.4 (two starters)
[9, 5, 2, 2]	Proposition 3.4 (two starters)
[8, 6, 2, 2]	Proposition 3.4 (two starters)
[7, 7, 2, 2]	Proposition 3.4 (two starters)
[10, 3, 3, 2]	Proposition 3.3 (one starter)
[9, 4, 3, 2]	Proposition 3.3 (one starter)
[8, 5, 3, 2]	Proposition 3.3 (one starter)
[7, 6, 3, 2]	Proposition 3.3 (one starter)
[8, 4, 4, 2]	Proposition 3.3 (one starter)
[7, 5, 4, 2]	Proposition 3.3 (one starter)
[6, 6, 4, 2]	Proposition 3.3 (one starter)
[6, 5, 5, 2]	Proposition 3.3 (one starter)
[9, 3, 3, 3]	Proposition 3.4 (two starters)
[8, 4, 3, 3]	Proposition 3.4 (two starters)
[7, 5, 3, 3]	Proposition 3.4 (two starters)
[6, 6, 3, 3]	Proposition 3.4 (two starters)
[7, 4, 4, 3]	Proposition 3.4 (two starters)
[6, 5, 4, 3]	Proposition 3.4 (two starters)
[5, 5, 5, 3]	Proposition 3.4 (two starters)
[6, 4, 4, 4]	Proposition 3.4 (two starters)
[5, 5, 4, 4]	Proposition 3.4 (two starters)
[14, 2, 2]	Proposition 3.4 (two starters)
[13, 3, 2]	Proposition 3.3 (one starter)
[12, 4, 2]	Proposition 3.3 (one starter)
[11, 5, 2]	Proposition 3.3 (one starter)
[10, 6, 2]	Proposition 3.3 (one starter)
[9, 7, 2]	Proposition 3.3 (one starter)
[8, 8, 2]	Proposition 3.3 (one starter)
[12, 3, 3]	Proposition 3.4 (two starters)
[11, 4, 3]	Proposition 3.4 (two starters)
[10, 5, 3]	Proposition 3.4 (two starters)
[9, 6, 3]	Proposition 3.4 (two starters)
[8, 7, 3]	Proposition 3.4 (two starters)
[10, 4, 4]	Proposition 3.4 (two starters)
[9, 5, 4]	Proposition 3.4 (two starters)
[8, 6, 4]	Proposition 3.4 (two starters)
[7, 7, 4]	Proposition 3.4 (two starters)
[8, 5, 5]	Proposition 3.4 (two starters)
[7, 6, 5]	Proposition 3.4 (two starters)
[6, 6, 6]	Theorem 1.2

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[16, 2]	Proposition 3.3 (one starter)
[15, 3]	Proposition 3.4 (two starters)
[14, 4]	Proposition 3.4 (two starters)
[13, 5]	Proposition 3.4 (two starters)
[12, 6]	Proposition 3.4 (two starters)
[11,7]	Proposition 3.4 (two starters)
[10, 8]	Proposition 3.4 (two starters)
[9, 9]	Theorem 1.2
[18]	Theorem 1.2

CASE n = 19: see Appendix K.

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[3, 2, 2, 2, 2, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[5, 2, 2, 2, 2, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[4, 3, 2, 2, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[3, 3, 3, 2, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[7, 2, 2, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[6, 3, 2, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[5, 4, 2, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[5, 3, 3, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[4, 4, 3, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[4, 3, 3, 3, 2, 2, 2]	Proposition 3.5 (three starters)
[3, 3, 3, 3, 3, 3, 2, 2]	Proposition 3.5 (three starters)
[9, 2, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[8, 3, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[7, 4, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[6, 5, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[7, 3, 3, 2, 2, 2]	Proposition 3.5 (three starters)
[6, 4, 3, 2, 2, 2]	Proposition 3.5 (three starters)
[5, 5, 3, 2, 2, 2]	Proposition 3.5 (three starters)
[5, 4, 4, 2, 2, 2]	Proposition 3.5 (three starters)
[6, 3, 3, 3, 2, 2]	Proposition 3.5 (three starters)
[5, 4, 3, 3, 2, 2]	Proposition 3.5 (three starters)
[4, 4, 4, 3, 2, 2]	Proposition 3.5 (three starters)
[5, 3, 3, 3, 3, 2]	Proposition 3.5 (three starters)
[4, 4, 3, 3, 3, 2]	Proposition 3.5 (three starters)
[4, 3, 3, 3, 3, 3]	Theorem 1.2
[11, 2, 2, 2, 2]	Proposition 3.5 (three starters)
[10, 3, 2, 2, 2]	Proposition 3.5 (three starters)
[9, 4, 2, 2, 2]	Proposition 3.5 (three starters)
[8, 5, 2, 2, 2]	Proposition 3.5 (three starters)
[7, 6, 2, 2, 2]	Proposition 3.5 (three starters)
[9, 3, 3, 2, 2]	Proposition 3.5 (three starters)
[8, 4, 3, 2, 2]	Proposition 3.5 (three starters)
[7, 5, 3, 2, 2]	Proposition 3.5 (three starters)

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[6, 6, 3, 2, 2]	Proposition 3.5 (three starters)
[7, 4, 4, 2, 2]	Proposition 3.5 (three starters)
[6, 5, 4, 2, 2]	Proposition 3.5 (three starters)
[5, 5, 5, 2, 2]	Proposition 3.5 (three starters)
[8, 3, 3, 3, 2]	Proposition 3.5 (three starters)
[7, 4, 3, 3, 2]	Proposition 3.5 (three starters)
[6, 5, 3, 3, 2]	Proposition 3.5 (three starters)
[6, 4, 4, 3, 2]	Proposition 3.5 (three starters)
[5, 5, 4, 3, 2]	Proposition 3.5 (three starters)
[5, 4, 4, 4, 2]	Proposition 3.5 (three starters)
[7, 3, 3, 3, 3]	Theorem 1.2
[6, 4, 3, 3, 3]	Theorem 1.2
[5, 5, 3, 3, 3]	Theorem 1.2
[5, 4, 4, 3, 3]	Theorem 1.2
[4, 4, 4, 4, 3]	Theorem 1.2
[13, 2, 2, 2]	Proposition 3.5 (three starters)
[12, 3, 2, 2]	Proposition 3.5 (three starters)
[11, 4, 2, 2]	Proposition 3.5 (three starters)
[10, 5, 2, 2]	Proposition 3.5 (three starters)
[9, 6, 2, 2]	Proposition 3.5 (three starters)
[8, 7, 2, 2]	Proposition 3.5 (three starters)
[11, 3, 3, 2]	Proposition 3.5 (three starters)
[10, 4, 3, 2]	Proposition 3.5 (three starters)
[9, 5, 3, 2]	Proposition 3.5 (three starters)
[8, 6, 3, 2]	Proposition 3.5 (three starters)
[7, 7, 3, 2]	Proposition 3.5 (three starters)
[9, 4, 4, 2]	Proposition 3.5 (three starters)
[8, 5, 4, 2]	Proposition 3.5 (three starters)
[7, 6, 4, 2]	Proposition 3.5 (three starters)
[7, 5, 5, 2]	Proposition 3.5 (three starters)
[6, 6, 5, 2]	Proposition 3.5 (three starters)
[10, 3, 3, 3]	Theorem 1.2
[9, 4, 3, 3]	Theorem 1.2
[8, 5, 3, 3]	Theorem 1.2
[7, 0, 3, 3]	Theorem 1.2
[8, 4, 4, 3]	Theorem 1.2
[7, 5, 4, 5]	Theorem 1.2
[0, 0, 4, 5]	Theorem 1.2
$\begin{bmatrix} 0, 0, 0, 0 \end{bmatrix}$	Theorem 1.2
$\begin{bmatrix} 1, 4, 4, 4 \end{bmatrix}$ [6 5 4 4]	Theorem 1.2
[0, 0, 4, 4] [5 5 5 4]	Theorem 1.2 Theorem 1.2
[0, 0, 0, 4] [15, 0, 9]	Proposition 3.5 (three starters)
[10, 2, 2] [14, 3, 2]	Proposition 3.5 (three starters)
[13, 4, 2]	$\frac{1}{2} = \frac{1}{2} + \frac{1}$
[12, 5, 2]	Proposition 3.5 (three starters)

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[11, 6, 2]	Proposition 3.5 (three starters)
[10, 7, 2]	Proposition 3.5 (three starters)
[9, 8, 2]	Proposition 3.5 (three starters)
[13, 3, 3]	Theorem 1.2
[12, 4, 3]	Theorem 1.2
[11, 5, 3]	Theorem 1.2
[10, 6, 3]	Theorem 1.2
[9, 7, 3]	Theorem 1.2
[8, 8, 3]	Theorem 1.2
[11, 4, 4]	Theorem 1.2
[10, 5, 4]	Theorem 1.2
[9, 6, 4]	Theorem 1.2
[8, 7, 4]	Theorem 1.2
[9, 5, 5]	Theorem 1.2
[8, 6, 5]	Theorem 1.2
[7, 7, 5]	Theorem 1.2
[7, 6, 6]	Theorem 1.2
[17, 2]	Theorem 1.2
[16, 3]	Theorem 1.2
[15, 4]	Theorem 1.2
[14, 5]	Theorem 1.2
[13, 6]	Theorem 1.2
[12,7]	Theorem 1.2
[11, 8]	Theorem 1.2
[10, 9]	Theorem 1.2
[19]	Theorem 1.2

CASE n = 20: see Appendix L.

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[2, 2, 2, 2, 2, 2, 2, 2, 2, 2]	Theorem 1.2
[4, 2, 2, 2, 2, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[3, 3, 2, 2, 2, 2, 2, 2, 2]	Proposition 3.4 (two starters)
[6, 2, 2, 2, 2, 2, 2, 2]	Proposition 3.4 (two starters)
[5, 3, 2, 2, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[4, 4, 2, 2, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[4, 3, 3, 2, 2, 2, 2, 2]	Proposition 3.4 (two starters)
[3, 3, 3, 3, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[8, 2, 2, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[7, 3, 2, 2, 2, 2, 2]	Proposition 3.4 (two starters)
[6, 4, 2, 2, 2, 2, 2]	Proposition 3.4 (two starters)
[5, 5, 2, 2, 2, 2, 2]	Proposition 3.4 (two starters)
[6, 3, 3, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[5, 4, 3, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[4, 4, 4, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[5, 3, 3, 3, 2, 2, 2]	Proposition 3.4 (two starters)

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[4, 4, 3, 3, 2, 2, 2]	Proposition 3.4 (two starters)
[4, 3, 3, 3, 3, 2, 2]	Proposition 3.3 (one starter)
[3, 3, 3, 3, 3, 3, 3, 2]	Proposition 3.4 (two starters)
[10, 2, 2, 2, 2, 2]	Proposition 3.4 (two starters)
[9, 3, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[8, 4, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[7, 5, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[6, 6, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[8, 3, 3, 2, 2, 2]	Proposition 3.4 (two starters)
[7, 4, 3, 2, 2, 2]	Proposition 3.4 (two starters)
[6, 5, 3, 2, 2, 2]	Proposition 3.4 (two starters)
[6, 4, 4, 2, 2, 2]	Proposition 3.4 (two starters)
[5, 5, 4, 2, 2, 2]	Proposition 3.4 (two starters)
[7, 3, 3, 3, 2, 2]	Proposition 3.3 (one starter)
[6, 4, 3, 3, 2, 2]	Proposition 3.3 (one starter)
[5, 5, 3, 3, 2, 2]	Proposition 3.3 (one starter)
[5, 4, 4, 3, 2, 2]	Proposition 3.3 (one starter)
[4, 4, 4, 4, 2, 2]	Proposition 3.3 (one starter)
[6, 3, 3, 3, 3, 2]	Proposition 3.4 (two starters)
[5, 4, 3, 3, 3, 2]	Proposition 3.4 (two starters)
[4, 4, 4, 3, 3, 2]	Proposition 3.4 (two starters)
[5, 3, 3, 3, 3, 3]	Proposition 3.3 (one starter)
[4, 4, 3, 3, 3, 3]	Proposition 3.3 (one starter)
[12, 2, 2, 2, 2]	Proposition 3.3 (one starter)
[11, 3, 2, 2, 2]	Proposition 3.4 (two starters)
[10, 4, 2, 2, 2]	Proposition 3.4 (two starters)
[9, 5, 2, 2, 2]	Proposition 3.4 (two starters)
[8, 6, 2, 2, 2]	Proposition 3.4 (two starters)
[7, 7, 2, 2, 2]	Proposition 3.4 (two starters)
[10, 3, 3, 2, 2]	Proposition 3.3 (one starter)
[9, 4, 3, 2, 2]	Proposition 3.3 (one starter)
[0, 0, 0, 2, 2] [7, 6, 2, 0, 0]	Proposition 3.3 (one starter)
[1, 0, 3, 2, 2]	Proposition 3.3 (one starter)
[0, 4, 4, 2, 2] [7 5 4 2 2]	Proposition 3.3 (one starter)
[1, 0, 4, 2, 2] [6 6 4 2 2]	Proposition 3.3 (one starter)
[0, 0, 4, 2, 2] [6, 5, 5, 2, 2]	Proposition 3.3 (one starter)
[0, 3, 3, 2, 2]	Proposition 3.4 (two starters)
[8, 4, 3, 3, 2]	Proposition 3.4 (two starters)
[7, 5, 3, 3, 2]	Proposition 3.4 (two starters)
[6, 6, 3, 3, 2]	Proposition 3.4 (two starters)
[7, 4, 4, 3, 2]	Proposition 3.4 (two starters)
[6, 5, 4, 3, 2]	Proposition 3.4 (two starters)
[5, 5, 5, 3, 2]	Proposition 3.4 (two starters)
[6, 4, 4, 4, 2]	Proposition 3.4 (two starters)
[5, 5, 4, 4, 2]	Proposition 3.4 (two starters)

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[8,3,3,3,3]	Proposition 3.3 (one starter)
[7, 4, 3, 3, 3]	Proposition 3.3 (one starter)
[6, 5, 3, 3, 3]	Proposition 3.3 (one starter)
[6, 4, 4, 3, 3]	Proposition 3.3 (one starter)
[5, 5, 4, 3, 3]	Proposition 3.3 (one starter)
[5, 4, 4, 4, 3]	Proposition 3.3 (one starter)
[4, 4, 4, 4, 4]	Theorem 1.2
[14, 2, 2, 2]	Proposition 3.4 (two starters)
[13, 3, 2, 2]	Proposition 3.3 (one starter)
[12, 4, 2, 2]	Proposition 3.3 (one starter)
[11, 5, 2, 2]	Proposition 3.3 (one starter)
[10, 6, 2, 2]	Proposition 3.3 (one starter)
[9, 7, 2, 2]	Proposition 3.3 (one starter)
[8, 8, 2, 2]	Proposition 3.3 (one starter)
[12, 3, 3, 2]	Proposition 3.4 (two starters)
[11, 4, 3, 2]	Proposition 3.4 (two starters)
[10, 5, 3, 2]	Proposition 3.4 (two starters)
[9, 6, 3, 2]	Proposition 3.4 (two starters)
[8,7,3,2]	Proposition 3.4 (two starters)
[10, 4, 4, 2]	Proposition 3.4 (two starters)
[9, 5, 4, 2]	Proposition 3.4 (two starters)
[8, 6, 4, 2]	Proposition 3.4 (two starters)
[7, 7, 4, 2]	Proposition 3.4 (two starters)
[8, 5, 5, 2]	Proposition 3.4 (two starters)
[7, 6, 5, 2]	Proposition 3.4 (two starters)
[6, 6, 6, 2]	Proposition 3.4 (two starters)
[11, 3, 3, 3]	Proposition 3.3 (one starter)
[10, 4, 3, 3]	Proposition 3.3 (one starter)
[9, 5, 3, 3]	Proposition 3.3 (one starter)
[8, 6, 3, 3]	Proposition 3.3 (one starter)
[7, 7, 3, 3]	Proposition 3.3 (one starter)
[9, 4, 4, 3]	Proposition 3.3 (one starter)
[8, 5, 4, 3]	Proposition 3.3 (one starter)
[7, 6, 4, 3]	Proposition 3.3 (one starter)
[7, 5, 5, 3]	Proposition 3.3 (one starter)
[6, 6, 5, 3]	Proposition 3.3 (one starter)
[8, 4, 4, 4]	Theorem 1.2
[7, 5, 4, 4]	Proposition 3.3 (one starter)
[6, 6, 4, 4]	Proposition 3.3 (one starter)
[6, 5, 5, 4]	Proposition 3.3 (one starter)
[5, 5, 5, 5]	Theorem 1.2
[16, 2, 2]	Proposition 3.3 (one starter)
[15, 3, 2]	Proposition 3.4 (two starters)
[14, 4, 2]	Proposition 3.4 (two starters)
[13, 5, 2]	Proposition 3.4 (two starters)
[12, 6, 2]	Proposition 3.4 (two starters)

2-factor type $[m_1, \ldots, m_t]$	$HOP(2m_1,\ldots,2m_t)$ has a solution by
[11, 7, 2]	Proposition 3.4 (two starters)
[10, 8, 2]	Proposition 3.4 (two starters)
[9, 9, 2]	Proposition 3.4 (two starters)
[14, 3, 3]	Proposition 3.3 (one starter)
[13, 4, 3]	Proposition 3.3 (one starter)
[12, 5, 3]	Proposition 3.3 (one starter)
[11, 6, 3]	Proposition 3.3 (one starter)
[10, 7, 3]	Proposition 3.3 (one starter)
[9, 8, 3]	Proposition 3.3 (one starter)
[12, 4, 4]	Theorem 1.2
[11, 5, 4]	Proposition 3.3 (one starter)
[10, 6, 4]	Proposition 3.3 (one starter)
[9, 7, 4]	Proposition 3.3 (one starter)
[8, 8, 4]	Theorem 1.2
[10, 5, 5]	Proposition 3.3 (one starter)
[9, 6, 5]	Proposition 3.3 (one starter)
[8, 7, 5]	Proposition 3.3 (one starter)
[8, 6, 6]	Proposition 3.3 (one starter)
[7, 7, 6]	Proposition 3.3 (one starter)
[18, 2]	Proposition 3.4 (two starters)
[17,3]	Proposition 3.3 (one starter)
[16, 4]	Theorem 1.2
[15, 5]	Proposition 3.3 (one starter)
[14, 6]	Proposition 3.3 (one starter)
[13,7]	Proposition 3.3 (one starter)
[12, 8]	Theorem 1.2
[11,9]	Proposition 3.3 (one starter)
[10, 10]	Theorem 1.2
[20]	Theorem 1.2

A Guide to computational results

In cases where Proposition 3.3, 3.4, or 3.5 is used to prove existence of a solution to $HOP(2m_1, \ldots, 2m_t)$, the remaining sections provide appropriate starter 2-factors.

- Where Proposition 3.3 is used: one starter, namely $F = \{C_0, C_1, \ldots, C_{t-1}\}$.
- Where Proposition 3.4 is used: two starters, namely $F_1 = \{C_0, C_1, \dots, C_{t-1}\}$ and $F_2 = \{C'_0, C'_1, \dots, C'_{t-1}\}.$
- Where Proposition 3.5 is used: two starters, namely $F_1 = \{C_0, C_1, \ldots, C_{t-1}\}$ and $F_3 = \{C'_0, C'_1, \ldots, C'_{t-1}\}$. Note that in all cases, F_1 contains a 2-cycle consisting of a pink and blue edge of difference $\frac{n-1}{2}$. The starter 2-factor F_2 (as denoted in Proposition 3.5) is obtained from F_1 by applying $\rho_{\bullet}^{\frac{n-1}{2}}$ and then replacing the 2-cycle containing a pink and

blue edge of difference $\frac{n-1}{2}$ with a directed 2-cycle (with the same vertices) containing black arcs of length $\frac{n-1}{2}$.

The vertices of $2K_n^{\circ}$ or $4K_n^{\bullet}$ are labelled $0, 1, \ldots, n-1$; label n-1 represents vertex x_{∞} . Edge orbits are given in the form [d, c] where $d \in \{1, 2, \ldots, \lfloor \frac{n-1}{2} \rfloor\} \cup \{n-1\}$ is the difference (with n-1 representing the infinity difference), and c is the colour-orientation: 0 for pink, 2 for blue, 1 for black (single starter cases) or black orientated forward (two- and three-starter cases), and -1 for black oriented backward. We consider an arc (u, u + d), for $u, d \in \mathbb{Z}_{n-1}$, oriented forward if $d < \frac{n-1}{2}$, and oriented backward if $d > \frac{n-1}{2}$. The case $d = \frac{n-1}{2}$ can occur only in the three-starter approach (that is, for odd n); however, in this case, black arcs of difference $\frac{n-1}{2}$ appear only in the starter 2-factor F_2 , which is not specified below.

Each t-cycle in a starter 2-factor is presented in the form

$$C = [v_0, [d_0, c_0], v_1, [d_1, c_1], v_2, \dots, v_{t-2}, [d_{t-2}, c_{t-2}], v_{t-1}, [d_{t-1}, c_{t-1}]],$$

where $v_0, v_1, \ldots, v_{t-1}$ are the consecutive vertices of the cycle, and $[d_i, c_i]$ represents the orbit of the edge $v_i v_{i+1}$, for $i \in \mathbb{Z}_t$, as explained above.

B Computational results for n = 10

• 2-factor type [4, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [6, [4, 1], 2, [4, 0], 7, [9, 1], 9, [9, 0]]\\ C_1 &= [5, [2, 0], 3, [2, 1]]\\ C_2 &= [4, [3, 0], 1, [3, 1]]\\ C_3 &= [0, [1, 0], 8, [1, 1]] \end{split}$$

• 2-factor type [3, 3, 2, 2]: two starters

$C_0 = [1, [9, 2], 9, [9, 0], 0, [1, 1]]$	$C'_0 = [9, [9, 1], 5, [1, -1], 6, [9, -1]]$
$C_1 = [8, [4, 1], 3, [4, 2], 7, [1, 0]]$	$C_1' = [4, [4, -1], 8, [4, 0], 3, [1, 2]]$
$C_2 = [6, [2, -1], 4, [2, 1]]$	$C_2' = [2, [2, 0], 0, [2, 2]]$
$C_3 = [5, [3, 0], 2, [3, 2]]$	$C_3' = [7, [3, -1], 1, [3, 1]]$

• 2-factor type [6, 2, 2]: two starters

$$C_{0} = [9, [9, 2], 3, [3, 0], 0, [2, -1], 7, [4, 1], 2, [2, 1], 4, [9, 1]]$$

$$C_{1} = [6, [4, 0], 1, [4, 2]]$$

$$C_{2} = [8, [3, -1], 5, [3, 1]]$$

$$C'_{0} = [5, [4, -1], 0, [9, -1], 9, [9, 0], 4, [1, 2], 3, [1, 0], 2, [3, 2]]$$

$$C'_{1} = [7, [1, -1], 6, [1, 1]]$$

$$C'_{2} = [1, [2, 0], 8, [2, 2]]$$
• 2-factor type [5, 3, 2]: one starter

 $C_0 = [3, [2, 0], 5, [1, 0], 4, [4, 0], 8, [2, 1], 6, [3, 0]]$ $C_1 = [1, [3, 1], 7, [4, 1], 2, [1, 1]]$ $C_2 = [0, [9, 0], 9, [9, 1]]$

- 2-factor type [4, 4, 2]: one starter $C_0 = [5, [2, 1], 7, [3, 0], 1, [1, 1], 0, [4, 0]]$ $C_1 = [3, [1, 0], 4, [2, 0], 2, [3, 1], 8, [4, 1]]$ $C_2 = [9, [9, 0], 6, [9, 1]]$
- 2-factor type [4,3,3]: two starters $C_0 = [0, [1,2], 1, [2,1], 8, [2,0], 6, [3,1]]$ $C_1 = [3, [9,1], 9, [9,-1], 7, [4,-1]]$ $C_2 = [5, [1,-1], 4, [2,2], 2, [3,0]]$
- $C'_{0} = [6, [2, -1], 4, [9, 2], 9, [9, 0], 0, [3, -1]]$ $C'_{1} = [8, [4, 1], 3, [4, 2], 7, [1, 0]]$ $C'_{2} = [1, [4, 0], 5, [3, 2], 2, [1, 1]]$
- 2-factor type [8, 2]: one starter $C_0 = [0, [3, 0], 3, [4, 0], 7, [9, 1], 9, [9, 0], 1, [4, 1], 6, [2, 1], 8, [3, 1], 2, [2, 0]]$ $C_1 = [4, [1, 0], 5, [1, 1]]$
- 2-factor type [7,3]: two starters $C_0 = [1, [4, -1], 5, [4, 0], 0, [9, 1], 9, [9, 2], 2, [3, 1], 8, [2, 0], 6, [4, 2]]$ $C_1 = [3, [4, 1], 7, [3, -1], 4, [1, -1]]$
 - $$\begin{split} C_0' &= [2, [2, -1], 0, [1, 1], 1, [3, 2], 7, [3, 0], 4, [1, 2], 5, [2, 1], 3, [1, 0]]\\ C_1' &= [8, [9, -1], 9, [9, 0], 6, [2, 2]] \end{split}$$
- 2-factor type [6, 4]: two starters
 - $C_0 = [0, [1, 0], 1, [4, 1], 5, [3, 2], 2, [3, 0], 8, [4, 2], 3, [3, 1]]$
 - $C_1 = [6, [9, -1], 9, [9, 0], 7, [3, -1], 4, [2, 2]]$
 - $$\begin{split} C_0' &= [0, [4, -1], 4, [1, 1], 3, [2, 1], 1, [2, 0], 8, [1, 2], 7, [2, -1]]\\ C_1' &= [2, [4, 0], 6, [1, -1], 5, [9, 2], 9, [9, 1]] \end{split}$$

C Computational results for n = 11

• 2-factor type [3, 2, 2, 2, 2]: three starters

$C_0 = [7, [10, 1], 10, [10, -1], 6, [1, 1]]$	$C_0' = [1, [1, -1], 2, [10, 0], 10, [10, 2]]$
$C_1 = [3, [1, 0], 4, [1, 2]]$	$C'_1 = [0, [5, 0], 5, [5, 2]]$
$C_2 = [5, [3, 0], 8, [3, 2]]$	$C_2' = [3, [4, -1], 7, [4, 1]]$
$C_3 = [9, [2, -1], 1, [2, 1]]$	$C'_3 = [4, [4, 0], 8, [4, 2]]$
$C_4 = [0, [2, 0], 2, [2, 2]]$	$C_4' = [9, [3, -1], 6, [3, 1]]$

• 2-factor type [5, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [9, [4, 0], 5, [1, -1], 4, [4, 2], 8, [1, 1], 7, [2, -1]] \\ C_1 &= [6, [3, -1], 3, [3, 1]] \\ C_2 &= [0, [2, 0], 2, [2, 2]] \\ C_3 &= [10, [10, -1], 1, [10, 1]] \\ C'_0 &= [8, [4, 1], 4, [2, 1], 2, [1, 0], 1, [4, -1], 7, [1, 2]] \\ C'_1 &= [0, [5, 0], 5, [5, 2]] \\ C'_2 &= [9, [3, 0], 6, [3, 2]] \\ C'_3 &= [3, [10, 0], 10, [10, 2]] \\ \bullet & 2\text{-factor type } [4, 3, 2, 2]: \text{ three starters} \end{split}$$

- $$\begin{split} C_0 &= [7, [2, 1], 5, [2, 0], 3, [4, -1], 9, [2, 2]] \\ C_1 &= [0, [2, -1], 8, [4, 2], 4, [4, 0]] \\ C_2 &= [1, [1, -1], 2, [1, 1]] \\ C_3 &= [10, [10, -1], 6, [10, 1]] \end{split}$$
- $C_1' = [1, [4, 1], 7, [1, 0], 8, [3, 2]]$ $C_2' = [0, [5, 0], 5, [5, 2]]$ $C_3' = [10, [10, 0], 4, [10, 2]]$

 $C'_0 = [6, [3, 0], 9, [3, 1], 2, [1, 2], 3, [3, -1]]$

- 2-factor type [3, 3, 3, 2]: three starters
 - $$\begin{split} C_0 &= [4, [2, -1], 6, [4, 0], 0, [4, 2]] \\ C_1 &= [5, [2, 2], 7, [1, 0], 8, [3, -1]] \\ C_2 &= [3, [2, 0], 1, [1, 2], 2, [1, -1]] \\ C_3 &= [10, [10, -1], 9, [10, 1]] \end{split} \qquad \begin{aligned} C_0' &= [8, [1, 1], 7, [3, 1], 4, [4, -1]] \\ C_1' &= [3, [2, 1], 1, [10, 0], 10, [10, 2]] \\ C_1' &= [3, [2, 1], 1, [10, 0], 10, [10, 2]] \\ C_2' &= [9, [3, 2], 6, [4, 1], 2, [3, 0]] \\ C_3' &= [0, [5, 0], 5, [5, 2]] \end{aligned}$$
- 2-factor type [7, 2, 2]: three starters

$$\begin{split} C_0 &= [4, [3, 0], 1, [1, 2], 0, [3, 1], 7, [2, -1], 9, [4, 0], 3, [2, 2], 5, [1, 1]] \\ C_1 &= [2, [4, -1], 8, [4, 1]] \\ C_2 &= [10, [10, 0], 6, [10, 2]] \\ C_0' &= [8, [2, 0], 6, [3, 2], 3, [1, 0], 2, [3, -1], 9, [2, 1], 1, [4, 2], 7, [1, -1] \\ C_1' &= [0, [5, 0], 5, [5, 2]] \\ C_2' &= [4, [10, -1], 10, [10, 1]] \end{split}$$

• 2-factor type [6, 3, 2]: three starters

$$\begin{split} C_0 &= [9, [4, -1], 5, [10, 1], 10, [10, 2], 7, [1, 1], 6, [3, 1], 3, [4, 0]] \\ C_1 &= [2, [2, 1], 4, [3, 2], 1, [1, 0]] \\ C_2 &= [8, [2, 0], 0, [2, 2]] \\ C'_0 &= [6, [2, -1], 4, [3, -1], 1, [1, 2], 2, [4, 1], 8, [1, -1], 9, [3, 0]] \\ C'_1 &= [7, [10, 0], 10, [10, -1], 3, [4, 2]] \\ C'_2 &= [0, [5, 0], 5, [5, 2]] \end{split}$$

• 2-factor type [5, 4, 2]: three starters

$$C_{0} = [0, [1, -1], 1, [3, 1], 8, [3, 0], 5, [1, 2], 6, [4, -1]] \quad C'_{0} = [9, [3, 2], 2, [10, 0], 10, [10, 2], 6, [2, 0], 8, [1, 1]]$$

$$C_{1} = [10, [10, -1], 7, [2, 2], 9, [4, 0], 3, [10, 1]] \quad C'_{1} = [4, [1, 0], 3, [4, 1], 7, [4, 2], 1, [3, -1]]$$

$$C_{2} = [2, [2, -1], 4, [2, 1]] \quad C'_{2} = [0, [5, 0], 5, [5, 2]]$$

D Computational results for n = 12

• 2-factor type [4, 2, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [6, [1, 0], 5, [3, 0], 2, [1, 1], 3, [3, 1]]\\ C_1 &= [11, [11, 0], 0, [11, 1]]\\ C_2 &= [1, [4, 0], 8, [4, 1]]\\ C_3 &= [7, [2, 0], 9, [2, 1]]\\ C_4 &= [10, [5, 0], 4, [5, 1]] \end{split}$$

• 2-factor type [3, 3, 2, 2, 2]: two starters

$$\begin{split} C_0 &= [3, [1, 1], 4, [11, 1], 11, [11, -1]] & C'_0 &= [6, [1, -1], 5, [5, 2], 0, [5, 0]] \\ C_1 &= [0, [5, -1], 5, [4, -1], 9, [2, -1]] & C'_1 &= [10, [5, 1], 4, [4, 1], 8, [2, 1]] \\ C_2 &= [6, [2, 0], 8, [2, 2]] & C'_2 &= [9, [3, 0], 1, [3, 2]] \\ C_3 &= [1, [1, 0], 2, [1, 2]] & C'_3 &= [11, [11, 0], 2, [11, 2]] \\ C_4 &= [10, [3, -1], 7, [3, 1]] & C'_4 &= [7, [4, 0], 3, [4, 2]] \end{split}$$

- 2-factor type [6, 2, 2, 2]: two starters $C_0 = [11, [11, 2], 4, [1, 0], 5, [5, 1], 10, [2, 2], 1, [4, 1], 8, [11, 0]]$ $C_1 = [3, [3, 0], 6, [3, 2]]$ $C_2 = [2, [5, 0], 7, [5, 2]]$ $C_3 = [9, [2, -1], 0, [2, 1]]$ $C'_0 = [10, [1, 2], 0, [5, -1], 5, [11, -1], 11, [11, 1], 1, [2, 0], 3, [4, -1]]$ $C'_1 = [9, [1, -1], 8, [1, 1]]$ $C'_2 = [6, [4, 0], 2, [4, 2]]$ $C'_3 = [7, [3, -1], 4, [3, 1]]$
- 2-factor type [5, 3, 2, 2]: one starter

$$\begin{split} C_0 &= [4, [2, 1], 2, [5, 1], 7, [2, 0], 9, [1, 1], 10, [5, 0]] \\ C_1 &= [6, [11, 1], 11, [11, 0], 5, [1, 0]] \\ C_2 &= [3, [3, 0], 0, [3, 1]] \\ C_3 &= [1, [4, 0], 8, [4, 1]] \end{split}$$

- 2-factor type [4, 4, 2, 2]: one starter
 - $$\begin{split} C_0 &= [11, [11, 0], 2, [3, 0], 5, [1, 1], 4, [11, 1]] \\ C_1 &= [0, [3, 1], 8, [2, 1], 10, [1, 0], 9, [2, 0]] \\ C_2 &= [3, [4, 0], 7, [4, 1]] \\ C_3 &= [1, [5, 0], 6, [5, 1]] \end{split}$$
- 2-factor type [4, 3, 3, 2]: two starters $C_0 = [2, [5, 0], 8, [1, 1], 9, [3, -1], 6, [4, 2]]$ $C_1 = [11, [11, 2], 7, [3, 1], 4, [11, 0]]$ $C_2 = [10, [4, -1], 3, [2, 0], 1, [2, 2]]$ $C_3 = [5, [5, -1], 0, [5, 1]]$
- $C'_{0} = [9, [4, 0], 2, [3, 2], 10, [2, 1], 8, [1, -1]]$ $C'_{1} = [3, [3, 0], 6, [5, 2], 1, [2, -1]]$ $C'_{2} = [7, [4, 1], 0, [11, 1], 11, [11, -1]]$ $C'_{3} = [5, [1, 0], 4, [1, 2]]$

- 2-factor type [8, 2, 2]: one starter $C_0 = [1, [1, 1], 0, [4, 1], 4, [2, 1], 6, [4, 0], 10, [3, 1], 7, [2, 0], 5, [3, 0], 2, [1, 0]]$ $C_1 = [8, [5, 0], 3, [5, 1]]$ $C_2 = [9, [11, 0], 11, [11, 1]]$
- 2-factor type [7, 3, 2]: two starters $C_0 = [5, [11, 1], 11, [11, -1], 3, [5, 2], 9, [1, 1], 8, [2, 1], 6, [5, 0], 1, [4, 1]]$ $C_1 = [4, [3, 2], 7, [5, 1], 2, [2, 0]]$ $C_2 = [0, [1, 0], 10, [1, 2]]$ $C'_0 = [3, [2, -1], 1, [4, 2], 5, [1, -1], 6, [4, -1], 10, [3, 0], 7, [11, 2], 11, [11, 0]]$ $C'_1 = [4, [5, -1], 9, [4, 0], 2, [2, 2]]$ $C'_2 = [0, [3, -1], 8, [3, 1]]$
- 2-factor type [6, 4, 2]: two starters
 - $$\begin{split} C_0 &= [0, [1, -1], 1, [3, -1], 4, [2, 1], 2, [5, 0], 7, [4, 2], 3, [3, 1]] \\ C_1 &= [9, [4, 0], 5, [5, 1], 10, [4, -1], 6, [3, 2]] \\ C_2 &= [8, [11, 0], 11, [11, 2]] \\ C'_0 &= [10, [1, 1], 0, [1, 2], 1, [11, -1], 11, [11, 1], 4, [5, -1], 9, [1, 0]] \\ C'_1 &= [3, [4, 1], 7, [5, 2], 2, [3, 0], 5, [2, -1]] \\ C'_2 &= [6, [2, 0], 8, [2, 2]] \end{split}$$
- 2-factor type [5, 5, 2]: two starters

$$\begin{split} C_0 &= [11, [11, 2], 9, [1, 0], 8, [4, 2], 1, [1, 1], 0, [11, 0]] \\ C_1 &= [5, [1, -1], 6, [4, 0], 10, [3, 2], 7, [4, 1], 3, [2, -1]] \\ C_2 &= [4, [2, 0], 2, [2, 2]] \\ C_0' &= [8, [4, -1], 1, [2, 1], 10, [3, -1], 2, [11, -1], 11, [11, 1]] \\ C_1' &= [4, [3, 1], 7, [1, 2], 6, [3, 0], 3, [5, 2], 9, [5, 0]] \\ C_2' &= [5, [5, -1], 0, [5, 1]] \end{split}$$

- 2-factor type [6,3,3]: one starter
 - $$\begin{split} C_0 &= [9, [3, 0], 1, [1, 1], 2, [1, 0], 3, [4, 1], 7, [3, 1], 4, [5, 1]] \\ C_1 &= [0, [5, 0], 5, [11, 0], 11, [11, 1]] \\ C_2 &= [6, [2, 0], 8, [2, 1], 10, [4, 0]] \end{split}$$
- 2-factor type [5, 4, 3]: one starter
 - $$\begin{split} C_0 &= [1, [2, 0], 3, [1, 1], 2, [4, 0], 9, [5, 1], 4, [3, 1]]\\ C_1 &= [10, [3, 0], 7, [1, 0], 8, [2, 1], 6, [4, 1]]\\ C_2 &= [11, [11, 0], 0, [5, 0], 5, [11, 1]] \end{split}$$

• 2-factor type [10, 2]: two starters

$$\begin{split} C_0 &= [2, [1, 0], 1, [2, 1], 3, [1, 2], 4, [4, 0], 8, [11, 1], 11, [11, -1], 9, [3, -1], 6, [1, 1], 7, [4, 1], 0, [2, 2]] \\ C_1 &= [5, [5, -1], 10, [5, 1]] \\ C_0' &= [9, [1, -1], 8, [5, 2], 3, [3, 0], 6, [4, 2], 2, [3, 1], 10, [2, -1], 1, [4, -1], 5, [2, 0], 7, [3, 2], 4, [5, 0]] \\ C_1' &= [0, [11, 0], 11, [11, 2]] \end{split}$$

$$\begin{split} C_0 &= [0, [3, 1], 8, [4, 0], 1, [4, 1], 5, [2, 1], 3, [3, 0], 6, [2, 0], 4, [5, 1], 9, [1, 1], 10, [1, 0]]\\ C_1 &= [2, [5, 0], 7, [11, 1], 11, [11, 0]] \end{split}$$

• 2-factor type [7,5]: one starter $C_0 = [10, [4, 0], 3, [1, 0], 2, [1, 1], 1, [5, 0], 6, [2, 1], 8, [3, 0], 5, [5, 1]]$ $C_1 = [9, [2, 0], 7, [3, 1], 4, [4, 1], 0, [11, 1], 11, [11, 0]]$

E Computational results for n = 13

• 2-factor type [3, 2, 2, 2, 2, 2]: three starters

$C_0 = [10, [5, 2], 3, [3, -1], 6, [4, 0]]$	$C'_0 = [1, [5, 0], 8, [4, 2], 4, [3, 1]]$
$C_1 = [0, [5, -1], 5, [5, 1]]$	$C_1' = [0, [6, 0], 6, [6, 2]]$
$C_2 = [12, [12, 0], 11, [12, 2]]$	$C_2' = [7, [12, -1], 12, [12, 1]]$
$C_3 = [7, [2, 0], 9, [2, 2]]$	$C_3^\prime = [10, [1, -1], 9, [1, 1]]$
$C_4 = [2, [1, 0], 1, [1, 2]]$	$C_4' = [3, [2, -1], 5, [2, 1]]$
$C_5 = [8, [4, -1], 4, [4, 1]]$	$C_5' = [11, [3, 0], 2, [3, 2]]$

• 2-factor type [5, 2, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [9, [5, 2], 2, [4, 1], 10, [2, 1], 8, [3, 1], 5, [4, 0]] \\ C_1 &= [1, [2, 0], 3, [2, 2]] \\ C_2 &= [0, [12, -1], 12, [12, 1]] \\ C_3 &= [7, [1, -1], 6, [1, 1]] \\ C_4 &= [4, [5, -1], 11, [5, 1]] \\ C_6 &= [1, [2, -1], 3, [4, -1], 7, [5, 0], 2, [4, 2], 10, [3, -1]] \\ C_1' &= [0, [6, 0], 6, [6, 2]] \\ C_2' &= [11, [3, 0], 8, [3, 2]] \\ C_3' &= [9, [12, 0], 12, [12, 2]] \\ C_4' &= [5, [1, 0], 4, [1, 2]] \end{split}$$

• 2-factor type [4, 3, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [10, [4, 0], 6, [5, 2], 11, [2, -1], 1, [3, 1]] \\ C_1 &= [7, [4, 2], 3, [5, 0], 8, [1, -1]] \\ C_2 &= [9, [12, -1], 12, [12, 1]] \\ C_3 &= [2, [2, 0], 4, [2, 2]] \\ C_4 &= [0, [5, -1], 5, [5, 1]] \end{split}$$

 $C'_{0} = [12, [12, 0], 10, [1, 1], 11, [3, -1], 8, [12, 2]]$ $C'_{1} = [3, [1, 2], 2, [1, 0], 1, [2, 1]]$ $C'_{2} = [0, [6, 0], 6, [6, 2]]$ $C'_{3} = [7, [3, 0], 4, [3, 2]]$ $C'_{4} = [9, [4, -1], 5, [4, 1]]$

• 2-factor type [3, 3, 3, 2, 2]: three starters

$$C_{0} = [8, [12, -1], 12, [12, 1], 7, [1, -1]] \qquad C_{0}' = [9, [4, 2], 1, [3, 1], 10, [1, 0]] \\ C_{1} = [11, [2, 0], 9, [1, 1], 10, [1, 2]] \qquad C_{1}' = [5, [2, -1], 3, [5, 1], 8, [3, -1]] \\ C_{2} = [2, [2, 2], 4, [4, 0], 0, [2, 1]] \qquad C_{2}' = [4, [12, 0], 12, [12, 2], 11, [5, -1]] \\ C_{3} = [6, [3, 0], 3, [3, 2]] \qquad C_{3}' = [0, [6, 0], 6, [6, 2]] \\ C_{4} = [1, [4, -1], 5, [4, 1]] \qquad C_{4}' = [2, [5, 0], 7, [5, 2]]$$

- 2-factor type [7, 2, 2, 2]: three starters $C_0 = [5, [3, 2], 2, [12, 0], 12, [12, 2], 4, [5, -1], 9, [3, 1], 6, [5, 1], 1, [4, 0]]$ $C_1 = [11, [4, -1], 3, [4, 1]]$ $C_2 = [0, [2, 0], 10, [2, 2]]$ $C_3 = [8, [1, 0], 7, [1, 2]]$ $C'_0 = [2, [1, 1], 1, [5, 0], 8, [4, 2], 4, [3, -1], 7, [3, 0], 10, [1, -1], 9, [5, 2]]$ $C'_1 = [0, [6, 0], 6, [6, 2]]$ $C'_2 = [11, [12, -1], 12, [12, 1]]$ $C'_3 = [3, [2, -1], 5, [2, 1]]$
- 2-factor type [6, 3, 2, 2]: three starters

 $C_{0} = [1, [5, 2], 8, [1, 1], 7, [2, 0], 9, [2, 2], 11, [12, 0], 12, [12, -1]]$ $C_{1} = [4, [1, -1], 3, [3, 2], 0, [4, 0]]$ $C_{2} = [5, [5, -1], 10, [5, 1]]$ $C_{3} = [2, [4, -1], 6, [4, 1]]$ $C'_{0} = [1, [12, 2], 12, [12, 1], 9, [2, 1], 7, [5, 0], 2, [4, 2], 10, [3, 0]]$ $C'_{1} = [3, [2, -1], 5, [1, 0], 4, [1, 2]]$ $C'_{2} = [0, [6, 0], 6, [6, 2]]$ $C'_{3} = [8, [3, -1], 11, [3, 1]]$

- 2-factor type [5, 4, 2, 2]: three starters
 - $C_{0} = [3, [1, -1], 4, [5, -1], 9, [2, 0], 7, [1, 2], 6, [3, 1]]$ $C_{1} = [1, [1, 1], 0, [2, -1], 2, [12, 0], 12, [12, 2]]$ $C_{2} = [5, [5, 0], 10, [5, 2]]$ $C_{3} = [8, [3, 0], 11, [3, 2]]$ $C'_{0} = [5, [3, -1], 2, [5, 1], 7, [4, 2], 3, [2, 1], 1, [4, 0]]$ $C'_{1} = [12, [12, -1], 11, [2, 2], 9, [1, 0], 10, [12, 1]]$ $C'_{2} = [0, [6, 0], 6, [6, 2]]$ $C'_{3} = [4, [4, -1], 8, [4, 1]]$
- 2-factor type [5, 3, 3, 2]: three starters

$$\begin{split} C_0 &= [12, [12, -1], 4, [3, 2], 1, [2, 1], 11, [2, 0], 9, [12, 1]] \\ C_1 &= [10, [2, 2], 8, [3, 0], 5, [5, 1]] \\ C_2 &= [6, [4, 1], 2, [1, -1], 3, [3, -1]] \\ C_3 &= [0, [5, 0], 7, [5, 2]] \end{split}$$

- $C'_0 = [5, [5, -1], 10, [12, 0], 12, [12, 2], 1, [4, 0], 9, [4, 2]]$ $C'_1 = [11, [3, 1], 8, [1, 1], 7, [4, -1]]$ $C'_{2} = [2, [2, -1], 4, [1, 0], 3, [1, 2]]$ $C'_{3} = [0, [6, 0], 6, [6, 2]]$
- 2-factor type [4, 4, 3, 2]: three starters
 - $C_0 = [8, [4, 1], 0, [1, -1], 11, [5, 2], 4, [4, 0]]$ $C_1 = \begin{bmatrix} 10, [12, -1], 12, [12, 1], 1, [4, -1], 5, [5, -1] \end{bmatrix} \qquad C_1' = \begin{bmatrix} 10, [12, 2], 12, [12, 0], 3, [2, 2], 1, [3, 0] \end{bmatrix}$ $C_2 = [3, [4, 2], 7, [5, 0], 2, [1, 1]]$ $C_3 = [6, [3, -1], 9, [3, 1]]$
- $C'_0 = [5, [1, 2], 4, [2, 0], 2, [5, 1], 7, [2, -1]]$ $C'_{2} = [11, [3, 2], 8, [1, 0], 9, [2, 1]]$ $C'_{3} = [0, [6, 0], 6, [6, 2]]$
- 2-factor type [9, 2, 2]: three starters
 - $C_0 = [0, [2, 2], 10, [5, -1], 3, [4, -1], 7, [5, 1], 2, [1, 0], 1, [4, 2], 5, [4, 0], 9, [12, 2], 12, [12, 0]]$ $C_1 = [4, [2, -1], 6, [2, 1]]$ $C_2 = [8, [3, 0], 11, [3, 2]]$ $C'_0 = [7, [2, 0], 9, [5, 2], 2, [3, -1], 5, [4, 1], 1, [5, 0], 8, [12, 1], 12, [12, -1], 11, [1, 2], 10, [3, 1]]$ $C'_1 = [0, [6, 0], 6, [6, 2]]$ $C'_{2} = [4, [1, -1], 3, [1, 1]]$
- 2-factor type [8, 3, 2]: three starters
 - $C_0 = [7, [5, 1], 2, [3, -1], 5, [4, -1], 9, [5, 0], 4, [4, 2], 0, [2, 1], 10, [4, 0], 6, [1, 2]]$
 - $C_1 = [8, [5, -1], 1, [2, 0], 11, [3, 2]]$
 - $C_2 = [12, [12, -1], 3, [12, 1]]$
 - $C_0' = [12, [12, 2], 5, [2, -1], 7, [4, 1], 3, [1, 1], 2, [1, 0], 1, [3, 1], 4, [5, 2], 9, [12, 0]]$
 - $C'_1 = [11, [3, 0], 8, [2, 2], 10, [1, -1]]$
 - $C'_{2} = [0, [6, 0], 6, [6, 2]]$
- 2-factor type [7, 4, 2]: three starters
 - $C_0 = [8, [1, 0], 9, [5, 2], 2, [2, 1], 0, [4, 0], 4, [3, -1], 1, [4, 1], 5, [3, 2]]$
 - $C_1 = [7, [4, -1], 3, [3, 1], 6, [12, 1], 12, [12, -1]]$
 - $C_2 = [11, [1, -1], 10, [1, 1]]$
 - $C'_0 = [8, [4, 2], 4, [3, 0], 7, [2, 2], 5, [5, -1], 10, [5, 0], 3, [12, 2], 12, [12, 0]]$
 - $C'_1 = [9, [5, 1], 2, [1, 2], 1, [2, 0], 11, [2, -1]]$
 - $C'_{2} = [0, [6, 0], 6, [6, 2]]$
- 2-factor type [6, 5, 2]: three starters
 - $C_0 = [5, [2, 1], 3, [12, 0], 12, [12, -1], 7, [5, 2], 2, [1, 1], 1, [4, -1]]$ $C_1 = [10, [4, 1], 6, [2, -1], 8, [4, 0], 0, [1, -1], 11, [1, 2]]$ $C_2 = [9, [5, -1], 4, [5, 1]]$ $C'_0 = [11, [12, 1], 12, [12, 2], 9, [5, 0], 4, [3, 2], 7, [1, 0], 8, [3, 1]]$ $C'_1 = [3, [2, 0], 5, [3, -1], 2, [4, 2], 10, [3, 0], 1, [2, 2]]$ $C'_{2} = [0, [6, 0], 6, [6, 2]]$

F Computational results for n = 14

• 2-factor type [4, 2, 2, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [6, [3, 0], 9, [13, 1], 13, [13, 0], 3, [3, 1]]\\ C_1 &= [0, [4, 1], 4, [4, 0]]\\ C_2 &= [12, [5, 1], 7, [5, 0]]\\ C_3 &= [10, [2, 0], 8, [2, 1]]\\ C_4 &= [1, [1, 0], 2, [1, 1]]\\ C_5 &= [5, [6, 1], 11, [6, 0]] \end{split}$$

• 2-factor type [3, 3, 2, 2, 2, 2]: two starters

$$\begin{aligned} C_0 &= [5, [3, 2], 2, [5, 0], 10, [5, -1]] \\ C_1 &= [12, [2, 0], 1, [6, 1], 7, [5, 2]] \\ C_2 &= [4, [2, -1], 6, [2, 1]] \\ C_3 &= [3, [6, 0], 9, [6, 2]] \\ C_4 &= [11, [3, -1], 8, [3, 1]] \\ C_5 &= [13, [13, -1], 0, [13, 1]] \end{aligned}$$

$$\begin{aligned} C_0' &= [8, [6, -1], 2, [13, 2], 13, [13, 0]] \\ C_1' &= [8, [6, -1], 2, [13, 2], 13, [13, 0]] \\ C_1' &= [8, [6, -1], 2, [13, 2], 13, [13, 0]] \\ C_1' &= [8, [6, -1], 2, [13, 2], 13, [13, 0]] \\ C_1' &= [8, [6, -1], 2, [13, 2], 13, [13, 0]] \\ C_1' &= [8, [6, -1], 2, [13, 2], 13, [13, 0]] \\ C_1' &= [8, [6, -1], 2, [13, 2], 13, [13, 0]] \\ C_1' &= [8, [6, -1], 2, [13, 2], 13, [13, 0]] \\ C_1' &= [8, [6, -1], 2, [13, 2], 13, [13, 0]] \\ C_1' &= [8, [6, -1], 2, [13, 2], 13, [13, 0]] \\ C_2' &= [7, [4, -1], 3, [4, 1]] \\ C_2' &= [7, [4, -1], 3, [4, 1]] \\ C_3' &= [11, [1, -1], 10, [1, 1]] \\ C_4' &= [9, [4, 0], 5, [4, 2]] \\ C_5' &= [12, [13, 0], 0, [1, 2]] \end{aligned}$$

• 2-factor type [6, 2, 2, 2, 2]: two starters

$$\begin{split} C_0 &= [6, [3, 2], 3, [13, -1], 13, [13, 1], 0, [5, 1], 8, [6, 0], 2, [4, 1]] \\ C_1 &= [11, [1, 0], 10, [1, 2]] \\ C_2 &= [5, [6, -1], 12, [6, 1]] \\ C_3 &= [1, [5, 0], 9, [5, 2]] \\ C_4 &= [4, [3, -1], 7, [3, 1]] \\ C_0' &= [5, [3, 0], 8, [5, -1], 3, [6, 2], 9, [2, 0], 11, [4, -1], 7, [2, 2]] \\ C_1' &= [6, [13, 0], 13, [13, 2]] \\ C_2' &= [10, [4, 0], 1, [4, 2]] \\ C_3' &= [12, [1, -1], 0, [1, 1]] \\ C_4' &= [4, [2, -1], 2, [2, 1]] \end{split}$$

• 2-factor type [5, 3, 2, 2, 2]: one starter

 $C_0 = [6, [2, 1], 8, [3, 0], 11, [13, 0], 13, [13, 1], 3, [3, 1]]$ $C_1 = [2, [2, 0], 0, [1, 0], 1, [1, 1]]$ $C_2 = [4, [6, 1], 10, [6, 0]]$ $C_3 = [5, [4, 0], 9, [4, 1]]$ $C_4 = [12, [5, 0], 7, [5, 1]]$

• 2-factor type [4, 4, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [5, [2, 1], 3, [5, 1], 8, [1, 0], 7, [2, 0]]\\ C_1 &= [12, [3, 1], 9, [5, 0], 1, [3, 0], 11, [1, 1]]\\ C_2 &= [13, [13, 0], 0, [13, 1]]\\ C_3 &= [4, [6, 0], 10, [6, 1]]\\ C_4 &= [2, [4, 1], 6, [4, 0]] \end{split}$$

• 2-factor type [4, 3, 3, 2, 2]: two starters

 $C_0 = [12, [4, 0], 3, [6, -1], 10, [4, 1], 1, [2, 2]]$ $C_1 = [4, [5, -1], 9, [2, 0], 11, [6, 2]]$ $C_2 = [2, [6, 0], 8, [13, 2], 13, [13, 1]]$ $C_3 = [0, [5, 0], 5, [5, 2]]$ $C_4 = [7, [1, -1], 6, [1, 1]]$

• 2-factor type [3, 3, 3, 3, 2]: one starter

 $C_{0} = [0, [5, 0], 8, [13, 1], 13, [13, 0]]$ $C_{1} = [11, [6, 1], 4, [5, 1], 12, [1, 1]]$ $C_{2} = [9, [6, 0], 2, [3, 0], 5, [4, 1]]$ $C_{3} = [6, [1, 0], 7, [3, 1], 10, [4, 0]]$ $C_{4} = [1, [2, 1], 3, [2, 0]]$

- $C'_{0} = [0, [4, -1], 9, [2, -1], 7, [3, 2], 10, [3, 0]]$ $C'_{1} = [4, [2, 1], 6, [5, 1], 11, [6, 1]]$ $C'_{2} = [3, [13, -1], 13, [13, 0], 12, [4, 2]]$ $C'_{3} = [1, [1, 0], 2, [1, 2]]$ $C'_{4} = [8, [3, -1], 5, [3, 1]]$
- [3, 2]: one starter , 13, [13, 0]] , 12, [1, 1]]
- 2-factor type [8, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [4, [1, 1], 5, [3, 1], 2, [6, 0], 8, [13, 0], 13, [13, 1], 12, [1, 0], 0, [3, 0], 10, [6, 1]]\\ C_1 &= [11, [2, 0], 9, [2, 1]]\\ C_2 &= [6, [5, 1], 1, [5, 0]]\\ C_3 &= [7, [4, 1], 3, [4, 0]] \end{split}$$

• 2-factor type [7, 3, 2, 2]: two starters $C_0 = [4, [2, -1], 6, [13, -1], 13, [13, 0], 8, [6, 2], 2, [6, 0], 9, [6, -1], 3, [1, 2]]$ $C_1 = [12, [2, 1], 10, [4, 0], 1, [2, 2]]$ $C_2 = [7, [4, -1], 11, [4, 1]]$

$$C_3 = [5, [5, -1], 0, [5, 1]]$$

- $C'_0 = [5, [1, 0], 6, [4, 2], 2, [6, 1], 9, [2, 0], 7, [3, 2], 10, [3, 0], 0, [5, 2]]$ $C'_1 = [8, [13, 1], 13, [13, 2], 3, [5, 0]]$
- $C'_2 = [11, [1, -1], 12, [1, 1]]$
- $C_3^\prime = [1, [3, -1], 4, [3, 1]]$
- 2-factor type [6, 4, 2, 2]: two starters

$$\begin{split} C_0 &= [2, [1, 2], 1, [3, 0], 11, [3, -1], 8, [13, 2], 13, [13, 0], 12, [3, 1]] \\ C_1 &= [0, [3, 2], 10, [6, 1], 4, [5, -1], 9, [4, 0]] \\ C_2 &= [7, [4, -1], 3, [4, 1]] \\ C_3 &= [5, [1, -1], 6, [1, 1]] \\ C'_0 &= [1, [6, -1], 8, [5, 2], 3, [6, 0], 10, [6, 2], 4, [5, 0], 12, [2, 1]] \\ C'_1 &= [11, [5, 1], 6, [1, 0], 5, [4, 2], 9, [2, -1]] \\ C'_2 &= [13, [13, -1], 7, [13, 1]] \\ C'_3 &= [2, [2, 0], 0, [2, 2]] \end{split}$$

• 2-factor type [5, 5, 2, 2]: two starters

$$\begin{split} C_0 &= [12, [1, 1], 0, [4, 1], 4, [5, 2], 9, [1, 0], 10, [2, 1]] \\ C_1 &= [1, [4, -1], 5, [2, -1], 7, [4, 0], 11, [13, 1], 13, [13, 2]] \\ C_2 &= [2, [6, 0], 8, [6, 2]] \\ C_3 &= [3, [3, -1], 6, [3, 1]] \\ C_0' &= [13, [13, 0], 4, [1, 2], 5, [3, 0], 8, [4, 2], 12, [13, -1]] \\ C_1' &= [11, [1, -1], 10, [3, 2], 7, [5, 0], 2, [2, 2], 0, [2, 0]] \\ C_2' &= [6, [5, -1], 1, [5, 1]] \\ C_3' &= [9, [6, -1], 3, [6, 1]] \end{split}$$

- 2-factor type [6, 3, 3, 2]: one starter $C_0 = [8, [6, 1], 1, [5, 1], 9, [13, 1], 13, [13, 0], 11, [6, 0], 5, [3, 1]]$ $C_1 = [4, [5, 0], 12, [4, 1], 3, [1, 0]]$ $C_2 = [7, [1, 1], 6, [4, 0], 10, [3, 0]]$ $C_3 = [0, [2, 0], 2, [2, 1]]$
- 2-factor type [5, 4, 3, 2]: one starter $C_0 = [6, [4, 1], 10, [5, 0], 5, [6, 1], 12, [5, 1], 7, [1, 0]]$ $C_1 = [1, [13, 1], 13, [13, 0], 4, [1, 1], 3, [2, 0]]$ $C_2 = [2, [6, 0], 9, [4, 0], 0, [2, 1]]$ $C_3 = [8, [3, 0], 11, [3, 1]]$
- 2-factor type [4, 4, 4, 2]: one starter
 - $C_0 = [4, [4, 0], 0, [13, 0], 13, [13, 1], 6, [2, 1]]$ $C_1 = [8, [3, 1], 11, [3, 0], 1, [1, 1], 2, [6, 0]]$ $C_2 = [9, [1, 0], 10, [6, 1], 3, [2, 0], 5, [4, 1]]$ $C_3 = [12, [5, 0], 7, [5, 1]]$
- 2-factor type [5, 3, 3, 3]: two starters $C_0 = [7, [2, 0], 9, [3, -1], 6, [6, 1], 12, [4, 1], 3, [4, 2]]$ $C_1 = [2, [2, 2], 0, [3, 0], 10, [5, 1]]$ $C_2 = [4, [4, 0], 8, [3, 2], 5, [1, 1]]$ $C_3 = [13, [13, 0], 11, [3, 1], 1, [13, 2]]$ $C'_0 = [0, [2, -1], 2, [4, -1], 6, [5, 0], 1, [6, 2], 7, [6, -1]]$ $C'_1 = [8, [2, 1], 10, [1, 2], 9, [1, 0]]$ $C'_2 = [5, [6, 0], 12, [5, 2], 4, [1, -1]]$ $C'_3 = [3, [5, -1], 11, [13, 1], 13, [13, -1]]$
- 2-factor type [4, 4, 3, 3]: two starters

 $C_0 = [11, [1, 2], 10, [6, -1], 3, [4, 1], 12, [1, 0]]$ $C_1 = [5, [13, 0], 13, [13, 2], 9, [1, 1], 8, [3, 1]]$ $C_2 = [4, [3, -1], 1, [5, 2], 6, [2, 0]]$ $C_3 = [7, [6, 0], 0, [2, 2], 2, [5, -1]]$
$$\begin{split} C_0' &= [13, [13, -1], 1, [5, 1], 6, [2, 1], 8, [13, 1]] \\ C_1' &= [9, [6, 2], 3, [1, -1], 4, [4, 0], 0, [4, -1]] \\ C_2' &= [11, [4, 2], 2, [3, 0], 5, [6, 1]] \\ C_3' &= [7, [3, 2], 10, [2, -1], 12, [5, 0]] \end{split}$$

• 2-factor type [10, 2, 2]: two starters

$$\begin{split} C_0 &= [5, [3, 2], 2, [6, 1], 9, [3, 0], 12, [2, 2], 1, [6, -1], 7, [1, 0], 6, [4, 2], 10, [3, -1], 0, [4, 0], 4, [1, 1]] \\ C_1 &= [11, [5, 0], 3, [5, 2]] \\ C_2 &= [13, [13, 0], 8, [13, 2]] \\ C_0' &= [1, [2, 0], 3, [5, -1], 11, [6, 2], 5, [6, 0], 12, [5, 1], 4, [3, 1], 7, [1, -1], 6, [13, 1], 13, [13, -1], 2, [1, 2]] \\ C_1' &= [9, [4, -1], 0, [4, 1]] \\ C_2' &= [10, [2, -1], 8, [2, 1]] \end{split}$$

- 2-factor type [9, 3, 2]: one starter
 - $$\begin{split} C_0 &= [7, [6, 0], 0, [2, 1], 2, [6, 1], 9, [3, 0], 12, [2, 0], 10, [4, 1], 1, [4, 0], 5, [3, 1], 8, [1, 1]] \\ C_1 &= [3, [13, 1], 13, [13, 0], 4, [1, 0]] \\ C_2 &= [6, [5, 1], 11, [5, 0]] \end{split}$$
- 2-factor type [8, 4, 2]: one starter $C_0 = [3, [13, 0], 13, [13, 1], 7, [3, 1], 10, [6, 1], 4, [5, 0], 12, [1, 1], 0, [2, 0], 2, [1, 0]]$ $C_1 = [6, [5, 1], 11, [3, 0], 1, [6, 0], 8, [2, 1]]$ $C_2 = [5, [4, 0], 9, [4, 1]]$
- 2-factor type [7, 5, 2]: one starter $C_0 = [8, [6, 1], 2, [6, 0], 9, [4, 1], 5, [1, 1], 4, [3, 1], 7, [3, 0], 10, [2, 1]]$ $C_1 = [1, [13, 1], 13, [13, 0], 0, [1, 0], 12, [4, 0], 3, [2, 0]]$ $C_2 = [11, [5, 1], 6, [5, 0]]$
- 2-factor type [6, 6, 2]: one starter $C_0 = [0, [6, 1], 6, [3, 0], 9, [6, 0], 3, [5, 0], 8, [13, 0], 13, [13, 1]]$ $C_1 = [4, [3, 1], 7, [2, 0], 5, [4, 1], 1, [4, 0], 10, [5, 1], 2, [2, 1]]$ $C_2 = [12, [1, 1], 11, [1, 0]]$
- 2-factor type [8,3,3]: two starters $C_0 = [8, [6,-1], 1, [4,-1], 5, [6,1], 12, [5,-1], 4, [6,0], 10, [1,2], 9, [2,-1], 11, [3,1]]$ $C_1 = [0, [13,0], 13, [13,-1], 6, [6,2]]$ $C_2 = [3, [1,1], 2, [5,0], 7, [4,2]]$ $C'_0 = [2, [5,2], 7, [3,0], 4, [2,2], 6, [1,0], 5, [5,1], 10, [3,2], 0, [2,1], 11, [4,0]]$ $C'_1 = [3, [13,2], 13, [13,1], 1, [2,0]]$ $C'_2 = [12, [3,-1], 9, [1,-1], 8, [4,1]]$ • 2-factor type [7,4,3]: two starters
 - $$\begin{split} C_0 &= [6, [1, 2], 5, [6, 1], 12, [13, -1], 13, [13, 0], 9, [1, -1], 8, [4, 2], 4, [2, 0]] \\ C_1 &= [1, [4, 1], 10, [5, 0], 2, [4, -1], 11, [3, 2]] \\ C_2 &= [3, [4, 0], 7, [6, 2], 0, [3, -1]] \\ C_0' &= [10, [5, 2], 2, [2, 1], 0, [1, 0], 12, [13, 1], 13, [13, 2], 11, [5, -1], 3, [6, 0]] \\ C_1' &= [7, [6, -1], 1, [3, 1], 4, [5, 1], 9, [2, -1]] \\ C_2' &= [8, [3, 0], 5, [1, 1], 6, [2, 2]] \end{split}$$

- 2-factor type [6, 5, 3]: two starters
 - $$\begin{split} C_0 &= [6, [3, 1], 3, [2, 0], 5, [1, 2], 4, [6, 0], 11, [13, 2], 13, [13, 1]] \\ C_1 &= [7, [5, 2], 12, [4, 1], 8, [6, -1], 1, [4, 0], 10, [3, -1]] \\ C_2 &= [0, [4, -1], 9, [6, 1], 2, [2, -1]] \\ C_0' &= [2, [2, 2], 0, [5, 1], 8, [1, -1], 9, [5, 0], 1, [4, 2], 5, [3, 0]] \\ C_1' &= [12, [1, 1], 11, [5, -1], 3, [1, 0], 4, [2, 1], 6, [6, 2]] \\ C_2' &= [10, [3, 2], 7, [13, 0], 13, [13, -1]] \end{split}$$
- 2-factor type [6, 4, 4]: two starters
 - $$\begin{split} C_0 &= [13, [13, 0], 1, [1, 2], 2, [5, 0], 7, [4, -1], 3, [2, 1], 5, [13, 2]] \\ C_1 &= [0, [6, 2], 6, [3, 0], 9, [2, 2], 11, [2, 0]] \\ C_2 &= [8, [4, 2], 12, [5, -1], 4, [6, 0], 10, [2, -1]] \\ C_0' &= [2, [3, 1], 12, [4, 1], 8, [1, 1], 7, [13, -1], 13, [13, 1], 1, [1, -1]] \\ C_1' &= [4, [6, -1], 10, [3, -1], 0, [4, 0], 9, [5, 2]] \\ C_2' &= [6, [1, 0], 5, [6, 1], 11, [5, 1], 3, [3, 2]] \end{split}$$
- 2-factor type [5, 5, 4]: two starters
 - $$\begin{split} C_0 &= [13, [13, 2], 10, [6, 0], 4, [3, 1], 7, [4, 1], 11, [13, 1]] \\ C_1 &= [1, [5, 0], 9, [4, 2], 5, [5, 1], 0, [1, 0], 12, [2, 2]] \\ C_2 &= [8, [2, 1], 6, [4, 0], 2, [1, 1], 3, [5, 2]] \\ C'_0 &= [12, [3, 0], 9, [5, -1], 4, [1, 2], 3, [3, -1], 6, [6, -1]] \\ C'_1 &= [2, [4, -1], 11, [6, 2], 5, [2, -1], 7, [13, -1], 13, [13, 0]] \end{split}$$
 - $C_2' = [10, [2, 0], 8, [6, 1], 1, [1, -1], 0, [3, 2]]$
- 2-factor type [12, 2]: one starter
 - $C_{0} = [0, [13, 1], 13, [13, 0], 9, [1, 1], 10, [2, 0], 12, [1, 0], 11, [5, 1], 3, [3, 1], 6, [5, 0], 1, [4, 1], 5, [2, 1], 7, [3, 0], 4, [4, 0]]$
 - $C_1 = [8, [6, 0], 2, [6, 1]]$
- 2-factor type [11, 3]: two starters
 - $$\begin{split} C_0 &= [7, [4, 2], 3, [5, 0], 8, [2, 2], 10, [6, 0], 4, [1, 2], 5, [6, 1], 12, [3, 0], 2, [2, -1], 0, [1, 1], 1, [5, 2], 9, [2, 0]] \\ C_1 &= [6, [5, 1], 11, [13, 2], 13, [13, 0]] \end{split}$$
 - $$\begin{split} C_0' &= [9, [2, 1], 7, [4, -1], 11, [1, -1], 12, [6, -1], 5, [3, 1], 2, [1, 0], 1, [3, 2], 4, [13, -1], 13, [13, 1], 8, [5, -1], \\ &0, [4, 1]] \end{split}$$
 - $C_1' = \left[10, [6, 2], 3, [3, -1], 6, [4, 0]\right]$
- 2-factor type [10,4]: two starters

 $C_{0} = [0, [4, 1], 4, [5, 1], 9, [6, 1], 2, [5, -1], 10, [2, 2], 8, [13, 0], 13, [13, -1], 11, [5, 2], 3, [4, 0], 12, [1, 1]]$ $C_{1} = [7, [2, 0], 5, [1, 2], 6, [5, 0], 1, [6, 2]]$ $C'_{0} = [1, [2, -1], 3, [3, 0], 6, [4, -1], 2, [3, -1], 12, [4, 2], 8, [1, 0], 9, [2, 1], 11, [6, -1], 5, [1, -1], 4, [3, 2]]$

 $C_1' = \begin{bmatrix} 13, [13, 1], 0, [6, 0], 7, [3, 1], 10, [13, 2] \end{bmatrix}$

• 2-factor type [9,5]: two starters

$$\begin{split} C_0 &= [6, [1, 1], 5, [2, 0], 3, [6, 1], 9, [13, 1], 13, [13, -1], 7, [3, 2], 4, [6, 0], 10, [2, 1], 12, [6, 2]] \\ C_1 &= [1, [1, 0], 2, [4, -1], 11, [3, -1], 8, [5, 1], 0, [1, 2]] \\ C_0' &= [2, [6, -1], 9, [5, 2], 4, [3, 0], 7, [1, -1], 6, [2, 2], 8, [5, 0], 0, [3, 1], 3, [5, -1], 11, [4, 1]] \\ C_1' &= [13, [13, 0], 12, [2, -1], 10, [4, 2], 1, [4, 0], 5, [13, 2]] \end{split}$$

• 2-factor type [8,6]: two starters

$$\begin{split} C_0 &= [13, [13, -1], 10, [5, 2], 5, [3, -1], 8, [4, 0], 12, [1, 1], 0, [3, 1], 3, [2, -1], 1, [13, 1]] \\ C_1 &= [9, [3, 2], 6, [4, 1], 2, [5, -1], 7, [3, 0], 4, [6, 2], 11, [2, 0]] \\ C_0' &= [1, [1, -1], 0, [2, 1], 2, [2, 2], 4, [5, 0], 9, [6, -1], 3, [4, 2], 7, [1, 0], 8, [6, 1]] \end{split}$$

 $C_0 = [1, [1, -1], 0, [2, 1], 2, [2, 2], 4, [0, 0], 9, [0, -1], 9, [4, 2], 7, [1, 0], 8, [0, -1], 0, [13, 0]]$ $C_1' = [13, [13, 2], 5, [6, 0], 12, [1, 2], 11, [5, 1], 6, [4, -1], 10, [13, 0]]$

G Computational results for n = 15

• 2-factor type [3, 2, 2, 2, 2, 2, 2]: three starters

 $\begin{array}{ll} C_0 = [9, [5, 0], 0, [6, 2], 6, [3, -1]] & C_0' = [4, [5, 2], 9, [6, 0], 1, [3, 1]] \\ C_1 = [11, [3, 0], 8, [3, 2]] & C_1' = [6, [5, -1], 11, [5, 1]] \\ C_2 = [14, [14, -1], 10, [14, 1]] & C_2' = [3, [1, 0], 2, [1, 2]] \\ C_3 = [12, [1, -1], 13, [1, 1]] & C_3' = [8, [14, 0], 14, [14, 2]] \\ C_4 = [4, [2, 0], 2, [2, 2]] & C_4' = [12, [2, -1], 10, [2, 1]] \\ C_5 = [1, [4, -1], 5, [4, 1]] & C_5' = [13, [6, -1], 5, [6, 1]] \\ C_6 = [7, [4, 0], 3, [4, 2]] & C_6' = [0, [7, 0], 7, [7, 2]] \end{array}$

• 2-factor type [5, 2, 2, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [11, [6, 2], 3, [6, -1], 9, [6, 0], 1, [1, 2], 0, [3, 0]] \\ C_1 &= [2, [14, -1], 14, [14, 1]] \\ C_2 &= [13, [5, -1], 4, [5, 1]] \\ C_3 &= [8, [4, 0], 12, [4, 2]] \\ C_4 &= [7, [2, 0], 5, [2, 2]] \\ C_5 &= [6, [4, -1], 10, [4, 1]] \\ C_0' &= [6, [3, 2], 9, [6, 1], 3, [14, 0], 14, [14, 2], 5, [1, 0]] \\ C_1' &= [12, [1, -1], 13, [1, 1]] \\ C_2' &= [2, [5, 0], 11, [5, 2]] \\ C_3' &= [10, [2, -1], 8, [2, 1]] \\ C_4' &= [4, [3, -1], 1, [3, 1]] \\ C_5' &= [0, [7, 0], 7, [7, 2]] \end{split}$$

• 2-factor type [4, 3, 2, 2, 2, 2]: three starters

$$C_{0} = [10, [5, -1], 5, [4, -1], 1, [6, -1], 9, [1, 1]]$$

$$C_{1} = [12, [5, 0], 3, [14, 1], 14, [14, 2]]$$

$$C_{2} = [2, [2, -1], 4, [2, 1]]$$

$$C_{3} = [6, [2, 0], 8, [2, 2]]$$

$$C_{4} = [11, [3, 0], 0, [3, 2]]$$

$$C_{5} = [7, [6, 0], 13, [6, 2]]$$

- 2-factor type [3, 3, 3, 2, 2, 2]: three starters
 - $$\begin{split} C_0 &= [3, [6, -1], 9, [5, 1], 4, [1, 1]] \\ C_1 &= [2, [4, 0], 12, [2, 2], 0, [2, -1]] \\ C_2 &= [1, [14, 1], 14, [14, -1], 11, [4, 1]] \\ C_3 &= [6, [1, 0], 7, [1, 2]] \\ C_4 &= [10, [3, 0], 13, [3, 2]] \\ C_5 &= [8, [3, -1], 5, [3, 1]] \end{split}$$
- $$\begin{split} C_0' &= [1, [4, 1], 5, [5, 1], 10, [1, -1], 9, [6, 1]] \\ C_1' &= [14, [14, -1], 11, [5, 2], 2, [14, 0]] \\ C_2' &= [4, [4, 0], 8, [4, 2]] \\ C_3' &= [6, [3, -1], 3, [3, 1]] \\ C_4' &= [12, [1, 0], 13, [1, 2]] \\ C_5' &= [0, [7, 0], 7, [7, 2]] \end{split}$$
- $$\begin{split} C_0' &= [1, [4, 2], 11, [2, 0], 13, [2, 1]] \\ C_1' &= [2, [4, -1], 6, [14, 0], 14, [14, 2]] \\ C_2' &= [8, [5, -1], 3, [6, 1], 9, [1, -1]] \\ C_3' &= [0, [7, 0], 7, [7, 2]] \\ C_4' &= [12, [6, 0], 4, [6, 2]] \\ C_5' &= [5, [5, 0], 10, [5, 2]] \end{split}$$
- 2-factor type [7, 2, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [8, [3, -1], 11, [2, 0], 13, [6, 1], 5, [3, 2], 2, [4, 0], 6, [6, -1], 0, [6, 2]] \\ C_1 &= [3, [4, -1], 7, [4, 1]] \\ C_2 &= [1, [14, -1], 14, [14, 1]] \\ C_3 &= [12, [2, -1], 10, [2, 1]] \\ C_4 &= [4, [5, 0], 9, [5, 2]] \\ C_6' &= [2, [4, 2], 12, [1, 1], 11, [6, 0], 5, [3, 1], 8, [2, 2], 6, [3, 0], 3, [1, -1]] \\ C_1' &= [0, [7, 0], 7, [7, 2]] \\ C_2' &= [10, [1, 0], 9, [1, 2]] \\ C_3' &= [1, [14, 0], 14, [14, 2]] \\ C_4' &= [4, [5, -1], 13, [5, 1]] \end{split}$$

• 2-factor type [6, 3, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [1, [4, -1], 5, [1, 0], 6, [6, 2], 12, [5, 1], 7, [14, -1], 14, [14, 1]] \\ C_1 &= [11, [5, -1], 2, [3, 1], 13, [2, 1]] \\ C_2 &= [8, [4, 0], 4, [4, 2]] \\ C_3 &= [9, [1, -1], 10, [1, 1]] \\ C_4 &= [0, [3, 0], 3, [3, 2]] \\ C'_0 &= [14, [14, 0], 10, [5, 2], 1, [2, -1], 3, [6, 0], 9, [3, -1], 6, [14, 2]] \\ C'_1 &= [8, [4, 1], 12, [1, 2], 13, [5, 0]] \\ C'_2 &= [0, [7, 0], 7, [7, 2]] \\ C'_3 &= [2, [2, 0], 4, [2, 2]] \\ C'_4 &= [11, [6, -1], 5, [6, 1]] \end{split}$$

• 2-factor type [5, 4, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [2, [4, 0], 6, [3, 2], 3, [2, 1], 1, [3, 0], 4, [2, 2]] \\ C_1 &= [0, [1, 2], 13, [4, 1], 9, [1, 0], 8, [6, 1]] \\ C_2 &= [12, [14, 0], 14, [14, 2]] \\ C_3 &= [11, [6, 0], 5, [6, 2]] \\ C_4 &= [10, [3, -1], 7, [3, 1]] \\ C'_0 &= [5, [2, 0], 3, [2, -1], 1, [1, 1], 2, [6, -1], 10, [5, 2]] \\ C'_1 &= [4, [4, 2], 8, [4, -1], 12, [1, -1], 13, [5, 0]] \\ C'_2 &= [0, [7, 0], 7, [7, 2]] \\ C'_3 &= [14, [14, -1], 9, [14, 1]] \\ C'_4 &= [6, [5, -1], 11, [5, 1]] \\ 2\text{-factor type } [5, 3, 3, 2, 2]: \text{ three starters} \end{split}$$

- $$\begin{split} C_0 &= [12, [5, 0], 7, [6, 1], 13, [2, 2], 11, [3, 0], 8, [4, 2]] \\ C_1 &= [6, [3, 1], 3, [2, 0], 5, [1, 2]] \\ C_2 &= [14, [14, -1], 0, [4, -1], 10, [14, 1]] \\ C_3 &= [1, [6, 0], 9, [6, 2]] \\ C_4 &= [4, [2, -1], 2, [2, 1]] \\ C_0' &= [2, [1, 1], 1, [3, -1], 4, [6, -1], 10, [14, 0], 14, [14, 2]] \\ C_1' &= [13, [4, 1], 3, [5, 2], 12, [1, 0]] \\ C_2' &= [8, [1, -1], 9, [4, 0], 5, [3, 2]] \\ C_3' &= [0, [7, 0], 7, [7, 2]] \\ C_4' &= [11, [5, -1], 6, [5, 1]] \end{split}$$
- 2-factor type [4, 4, 3, 2, 2]: three starters

$$\begin{split} C_0 &= [7, [14, 2], 14, [14, 0], 2, [1, -1], 1, [6, 1]] \\ C_1 &= [0, [4, 0], 4, [4, 1], 8, [5, 1], 13, [1, 2]] \\ C_2 &= [5, [6, 0], 11, [2, -1], 9, [4, 2]] \\ C_3 &= [10, [2, 0], 12, [2, 2]] \\ C_4 &= [6, [3, 0], 3, [3, 2]] \end{split}$$

- $$\begin{split} C_0' &= [9, [6, -1], 1, [1, 0], 2, [5, 2], 11, [2, 1]] \\ C_1' &= [4, [6, 2], 12, [5, -1], 3, [5, 0], 8, [4, -1]] \\ C_2' &= [6, [1, 1], 5, [14, -1], 14, [14, 1]] \\ C_3' &= [0, [7, 0], 7, [7, 2]] \\ C_4' &= [13, [3, -1], 10, [3, 1]] \end{split}$$
- 2-factor type [4, 3, 3, 3, 2]: three starters $C_0 = [12, [1, 1], 11, [3, -1], 0, [5, 0], 9, [3, 2]]$ $C_1 = [2, [3, 0], 5, [4, 2], 1, [1, -1]]$ $C_2 = [6, [3, 1], 3, [5, -1], 8, [2, 1]]$ $C_3 = [10, [6, -1], 4, [14, 1], 14, [14, -1]]$ $C_4 = [7, [6, 0], 13, [6, 2]]$

$$\begin{split} C_0' &= [6, [5, 2], 11, [1, 0], 12, [14, 2], 14, [14, 0]] \\ C_1' &= [1, [2, 0], 13, [4, 1], 3, [2, 2]] \\ C_2' &= [10, [1, 2], 9, [4, 0], 5, [5, 1]] \\ C_3' &= [2, [6, 1], 8, [4, -1], 4, [2, -1]] \\ C_4' &= [0, [7, 0], 7, [7, 2]] \end{split}$$

• 2-factor type [9, 2, 2, 2]: three starters

• 2-factor type [8, 3, 2, 2]: three starters

$$\begin{split} C_0 &= [14, [14, 2], 4, [6, 0], 12, [2, -1], 10, [3, 1], 13, [4, -1], 9, [6, 2], 3, [4, 0], 7, [4, 2], 11, [14, 0]] \\ C_1 &= [5, [3, 0], 8, [3, 2]] \\ C_2 &= [6, [5, -1], 1, [5, 1]] \\ C_3 &= [0, [2, 0], 2, [2, 2]] \\ C_0' &= [9, [3, -1], 12, [1, 0], 11, [5, 2], 6, [14, -1], 14, [14, 1], 5, [4, 1], 1, [2, 1], 13, [5, 0], 8, [1, 2]] \\ C_1' &= [0, [7, 0], 7, [7, 2]] \\ C_2' &= [4, [1, -1], 3, [1, 1]] \\ C_3' &= [10, [6, -1], 2, [6, 1]] \end{split}$$

- $C_{0} = [10, [5, 2], 5, [2, -1], 7, [1, 0], 6, [2, 2], 4, [3, 0], 1, [1, 2], 2, [4, 1], 12, [2, 0]]$ $C_{1} = [11, [6, -1], 3, [5, 0], 8, [3, 2]]$ $C_{2} = [9, [4, 0], 13, [4, 2]]$ $C_{3} = [0, [14, 0], 14, [14, 2]]$ $C'_{0} = [2, [1, -1], 3, [6, 0], 11, [2, 1], 13, [5, -1], 8, [3, -1], 5, [1, 1], 6, [6, 2], 12, [4, -1]]$ $C'_{1} = [10, [6, 1], 4, [3, 1], 1, [5, 1]]$ $C'_{2} = [0, [7, 0], 7, [7, 2]]$ $C'_{3} = [14, [14, -1], 9, [14, 1]]$
- 2-factor type [7, 4, 2, 2]: three starters

$$\begin{split} C_0 &= [5, [4, 1], 1, [2, 1], 13, [5, 1], 8, [1, 0], 7, [1, -1], 6, [3, 1], 9, [4, 2]] \\ C_1 &= [10, [4, 0], 0, [2, 2], 2, [5, 0], 11, [1, 2]] \\ C_2 &= [14, [14, -1], 3, [14, 1]] \\ C_3 &= [4, [6, -1], 12, [6, 1]] \\ C'_0 &= [3, [1, 1], 2, [6, 0], 10, [2, -1], 8, [4, -1], 4, [5, 2], 13, [2, 0], 11, [6, 2]] \\ C'_1 &= [6, [3, 0], 9, [3, 2], 12, [3, -1], 1, [5, -1]] \\ C'_2 &= [0, [7, 0], 7, [7, 2]] \\ C'_3 &= [14, [14, 0], 5, [14, 2]] \end{split}$$

• 2-factor type [6, 5, 2, 2]: three starters

$$\begin{split} C_0 &= [14, [14, 2], 2, [3, 0], 5, [6, 1], 11, [5, -1], 6, [2, 2], 8, [14, 0]] \\ C_1 &= [3, [2, 1], 1, [5, 0], 10, [4, 2], 0, [4, 0], 4, [1, 2]] \\ C_2 &= [13, [6, 0], 7, [6, 2]] \\ C_3 &= [12, [3, -1], 9, [3, 1]] \\ C_0' &= [3, [3, 2], 6, [2, 0], 8, [4, -1], 4, [14, 1], 14, [14, -1], 12, [5, 1]] \\ C_1' &= [10, [5, 2], 5, [6, -1], 11, [2, -1], 13, [4, 1], 9, [1, 0]] \\ C_2' &= [0, [7, 0], 7, [7, 2]] \\ C_3' &= [1, [1, -1], 2, [1, 1]] \end{split}$$
• 2-factor type [7, 3, 3, 2]: three starters

$$\begin{split} C_0 &= [0, [4, 0], 10, [6, 2], 4, [6, 1], 12, [5, 0], 3, [14, 2], 14, [14, 1], 13, [1, -1]] \\ C_1 &= [5, [6, 0], 11, [2, 2], 9, [4, 1]] \\ C_2 &= [2, [6, -1], 8, [1, 0], 7, [5, 2]] \\ C_3 &= [1, [5, -1], 6, [5, 1]] \\ C'_0 &= [6, [2, -1], 8, [3, -1], 11, [14, 0], 14, [14, -1], 9, [4, 2], 5, [3, 1], 2, [4, -1]] \\ C'_1 &= [3, [1, 2], 4, [3, 0], 1, [2, 1]] \\ C'_2 &= [10, [3, 2], 13, [1, 1], 12, [2, 0]] \\ C'_3 &= [0, [7, 0], 7, [7, 2]] \\ 2\text{-factor type } [6, 4, 3, 2]: \text{ three starters} \end{split}$$

$$\begin{split} C_0 &= [9, [14, 1], 14, [14, -1], 10, [3, 1], 13, [2, 2], 11, [4, -1], 1, [6, 0]] \\ C_1 &= [2, [2, -1], 4, [1, -1], 5, [2, 0], 3, [1, 2]] \\ C_2 &= [7, [1, 0], 8, [4, 2], 12, [5, 1]] \\ C_3 &= [6, [6, -1], 0, [6, 1]] \\ C_0' &= [1, [6, 2], 9, [5, 0], 4, [5, -1], 13, [3, 2], 2, [3, -1], 5, [4, 0]] \\ C_1' &= [3, [3, 0], 6, [2, 1], 8, [4, 1], 12, [5, 2]] \\ C_2' &= [10, [14, 0], 14, [14, 2], 11, [1, 1]] \\ C_3' &= [0, [7, 0], 7, [7, 2]] \end{split}$$

• 2-factor type [5, 5, 3, 2]: three starters

٠

$$C_{0} = [1, [6, -1], 7, [5, 0], 2, [2, -1], 0, [3, 1], 3, [2, 2]]$$

$$C_{1} = [5, [1, -1], 6, [4, -1], 10, [2, 1], 8, [1, 0], 9, [4, 2]]$$

$$C_{2} = [4, [5, -1], 13, [1, 2], 12, [6, 0]]$$

$$C_{3} = [14, [14, -1], 11, [14, 1]]$$

$$C'_{0} = [4, [5, 2], 13, [14, 0], 14, [14, 2], 9, [1, 1], 8, [4, 0]]$$

$$C'_{1} = [11, [6, 1], 5, [3, 0], 2, [4, 1], 6, [3, -1], 3, [6, 2]]$$

$$C'_{2} = [10, [2, 0], 12, [3, 2], 1, [5, 1]]$$

$$C'_{3} = [0, [7, 0], 7, [7, 2]]$$

• 2-factor type [5,4,4,2]: three starters

$$\begin{split} C_0 &= [1, [1, 2], 2, [4, 0], 12, [5, 2], 7, [14, -1], 14, [14, 0]] \\ C_1 &= [11, [3, 2], 8, [2, 0], 10, [5, -1], 5, [6, 1]] \\ C_2 &= [13, [4, 2], 3, [6, -1], 9, [5, 0], 0, [1, -1]] \\ C_3 &= [6, [2, -1], 4, [2, 1]] \\ C'_0 &= [10, [4, 1], 6, [5, 1], 1, [3, -1], 4, [6, 0], 12, [2, 2]] \\ C'_1 &= [13, [6, 2], 5, [3, 1], 2, [1, 0], 3, [4, -1]] \\ C'_2 &= [8, [3, 0], 11, [14, 2], 14, [14, 1], 9, [1, 1]] \\ C'_3 &= [0, [7, 0], 7, [7, 2]] \end{split}$$

• 2-factor type [11, 2, 2]: three starters

$$\begin{split} C_0 &= [3, [3, 0], 6, [1, 1], 7, [2, 2], 9, [4, 0], 13, [2, 1], 1, [4, 2], 5, [5, 0], 10, [6, -1], 4, [2, -1], 2, [4, -1], 12, [5, 2]] \\ C_1 &= [8, [14, -1], 14, [14, 1]] \\ C_2 &= [11, [3, -1], 0, [3, 1]] \\ C'_0 &= [6, [5, 1], 1, [1, -1], 2, [6, 0], 8, [3, 2], 5, [1, 0], 4, [6, 2], 12, [5, -1], 3, [6, 1], 11, [2, 0], 9, [1, 2], 10, [4, 1]] \\ C'_1 &= [0, [7, 0], 7, [7, 2]] \\ C'_2 &= [14, [14, 0], 13, [14, 2]] \end{split}$$

• 2-factor type [10, 3, 2]: three starters

$$\begin{split} C_0 &= [3, [3, 1], 0, [1, 1], 13, [14, 0], 14, [14, 2], 8, [4, 1], 4, [5, 0], 9, [3, -1], 6, [5, 2], 1, [4, -1], 5, [2, 1]] \\ C_1 &= [11, [1, -1], 12, [2, 0], 10, [1, 2]] \\ C_2 &= [2, [5, -1], 7, [5, 1]] \\ C'_0 &= [14, [14, 1], 12, [3, 0], 1, [2, 2], 3, [6, 1], 11, [6, 0], 5, [3, 2], 2, [6, -1], 8, [1, 0], 9, [4, 2], 13, [14, -1]] \\ C'_1 &= [10, [6, 2], 4, [2, -1], 6, [4, 0]] \\ C'_2 &= [0, [7, 0], 7, [7, 2]] \end{split}$$

• 2-factor type [9, 4, 2]: three starters

$$\begin{split} C_0 &= [14, [14, 1], 0, [1, 0], 1, [2, 1], 3, [2, 2], 5, [2, 0], 7, [6, 1], 13, [1, 2], 12, [1, 1], 11, [14, -1]] \\ C_1 &= [8, [6, 2], 2, [6, 0], 10, [6, -1], 4, [4, 1]] \\ C_2 &= [9, [3, 0], 6, [3, 2]] \\ C_0' &= [8, [5, -1], 3, [3, 1], 6, [5, 1], 11, [2, -1], 9, [4, 2], 13, [14, 0], 14, [14, 2], 2, [4, 0], 12, [4, -1]] \\ C_1' &= [4, [3, -1], 1, [5, 2], 10, [5, 0], 5, [1, -1]] \\ \end{split}$$

- $C'_2 = [0, [7, 0], 7, [7, 2]]$
- 2-factor type [8, 5, 2]: three starters

$$\begin{split} C_0 &= [14, [14, 1], 7, [5, 1], 2, [3, 0], 5, [6, 1], 11, [3, -1], 8, [2, 2], 10, [2, -1], 12, [14, -1]] \\ C_1 &= [4, [5, -1], 13, [4, 1], 3, [6, 2], 9, [3, 1], 6, [2, 0]] \\ C_2 &= [1, [1, -1], 0, [1, 1]] \\ C_0' &= [12, [3, 2], 1, [6, 0], 9, [1, 2], 10, [5, 0], 5, [6, -1], 13, [4, 2], 3, [1, 0], 2, [4, -1]] \\ C_1' &= [6, [5, 2], 11, [14, 0], 14, [14, 2], 8, [4, 0], 4, [2, 1]] \\ C_2' &= [0, [7, 0], 7, [7, 2] \end{split}$$

• 2-factor type [7, 6, 2]: three starters

$$\begin{split} C_0 &= [8, [1, 1], 9, [5, 2], 4, [5, 1], 13, [1, 0], 0, [6, 2], 6, [14, -1], 14, [14, 0]] \\ C_1 &= [5, [4, 2], 1, [4, 1], 11, [5, -1], 2, [1, -1], 3, [4, -1], 7, [2, 0]] \\ C_2 &= [10, [2, -1], 12, [2, 1]] \\ C_0' &= [1, [6, 0], 9, [14, 1], 14, [14, 2], 3, [5, 0], 8, [3, 1], 11, [1, 2], 12, [3, -1]] \\ C_1' &= [4, [2, 2], 6, [4, 0], 2, [3, 2], 5, [6, 1], 13, [3, 0], 10, [6, -1]] \\ C_2' &= [0, [7, 0], 7, [7, 2]] \end{split}$$

H Computational results for n = 16

• 2-factor type [4, 2, 2, 2, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [5, [6, 0], 11, [7, 1], 4, [6, 1], 13, [7, 0]] \\ C_1 &= [15, [15, 0], 3, [15, 1]] \\ C_2 &= [8, [1, 0], 9, [1, 1]] \\ C_3 &= [6, [4, 0], 10, [4, 1]] \\ C_4 &= [0, [3, 0], 12, [3, 1]] \\ C_5 &= [2, [5, 0], 7, [5, 1]] \\ C_6 &= [1, [2, 0], 14, [2, 1]] \end{split}$$

- 2-factor type [3, 3, 2, 2, 2, 2, 2]: two starters
 - $$\begin{split} C_0 &= [6, [2, 0], 8, [2, 2], 10, [4, 1]] \\ C_1 &= [13, [4, -1], 2, [6, 1], 11, [2, -1]] \\ C_2 &= [14, [1, -1], 0, [1, 1]] \\ C_3 &= [7, [3, -1], 4, [3, 1]] \\ C_4 &= [3, [15, -1], 15, [15, 1]] \\ C_5 &= [12, [7, 0], 5, [7, 2]] \\ C_6 &= [1, [7, -1], 9, [7, 1]] \end{split}$$

$$\begin{split} C_0' &= [0, [15, 2], 15, [15, 0], 13, [2, 1]] \\ C_1' &= [11, [3, 0], 8, [3, 2], 5, [6, -1]] \\ C_2' &= [6, [1, 0], 7, [1, 2]] \\ C_3' &= [1, [6, 0], 10, [6, 2]] \\ C_4' &= [3, [4, 0], 14, [4, 2]] \\ C_5' &= [9, [5, -1], 4, [5, 1]] \\ C_6' &= [2, [5, 0], 12, [5, 2]] \end{split}$$

- 2-factor type [6, 2, 2, 2, 2, 2]: two starters
 - $$\begin{split} C_0 &= [1, [7, 1], 9, [1, 0], 8, [4, 1], 12, [2, 2], 14, [4, -1], 3, [2, 1]] \\ C_1 &= [0, [7, 0], 7, [7, 2]] \\ C_2 &= [6, [4, 0], 10, [4, 2]] \\ C_3 &= [2, [15, -1], 15, [15, 1]] \\ C_4 &= [11, [6, 0], 5, [6, 2]] \\ C_5 &= [4, [6, -1], 13, [6, 1]] \\ C'_0 &= [14, [7, -1], 6, [2, -1], 8, [2, 0], 10, [1, 2], 9, [5, 0], 4, [5, 2]] \\ C'_1 &= [7, [15, 0], 15, [15, 2]] \\ C'_2 &= [12, [1, -1], 13, [1, 1]] \\ C'_3 &= [3, [3, -1], 0, [3, 1]] \\ C'_4 &= [11, [5, -1], 1, [5, 1]] \\ C'_5 &= [2, [3, 0], 5, [3, 2]] \end{split}$$
- 2-factor type [5, 3, 2, 2, 2, 2]: one starter

 $C_{0} = [4, [2, 1], 2, [4, 1], 13, [2, 0], 11, [4, 0], 7, [3, 1]]$ $C_{1} = [3, [3, 0], 0, [15, 1], 15, [15, 0]]$ $C_{2} = [1, [5, 0], 6, [5, 1]]$ $C_{3} = [8, [6, 0], 14, [6, 1]]$ $C_{4} = [5, [7, 0], 12, [7, 1]]$ $C_{5} = [10, [1, 0], 9, [1, 1]]$

• 2-factor type [4, 4, 2, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [2, [5, 0], 12, [7, 0], 5, [1, 1], 6, [4, 1]] \\ C_1 &= [3, [5, 1], 8, [7, 1], 0, [1, 0], 14, [4, 0]] \\ C_2 &= [9, [2, 0], 7, [2, 1]] \\ C_3 &= [15, [15, 0], 11, [15, 1]] \\ C_4 &= [13, [3, 0], 1, [3, 1]] \\ C_5 &= [10, [6, 0], 4, [6, 1]] \end{split}$$

• 2-factor type [4, 3, 3, 2, 2, 2]: two starters

$$\begin{split} C_0 &= [12, [5, 1], 2, [7, 1], 9, [6, 2], 0, [3, 0]] \\ C_1 &= [15, [15, -1], 6, [1, 2], 5, [15, 0]] \\ C_2 &= [1, [3, 1], 13, [6, 0], 4, [3, 2]] \\ C_3 &= [8, [5, 0], 3, [5, 2]] \\ C_4 &= [14, [7, 0], 7, [7, 2]] \\ C_5 &= [10, [1, -1], 11, [1, 1]] \end{split}$$

• 2-factor type [3, 3, 3, 3, 2, 2]: one starter

$$\begin{split} C_0 &= [0, [6, 0], 9, [15, 0], 15, [15, 1]] \\ C_1 &= [12, [6, 1], 3, [2, 0], 5, [7, 0]] \\ C_2 &= [8, [3, 0], 11, [2, 1], 13, [5, 0]] \\ C_3 &= [14, [5, 1], 4, [3, 1], 7, [7, 1]] \\ C_4 &= [1, [1, 0], 2, [1, 1]] \\ C_5 &= [6, [4, 0], 10, [4, 1]] \end{split}$$

• 2-factor type [8, 2, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [13, [4, 1], 2, [7, 0], 9, [3, 1], 12, [3, 0], 0, [4, 0], 11, [15, 1], 15, [15, 0], 6, [7, 1]] \\ C_1 &= [5, [2, 0], 3, [2, 1]] \\ C_2 &= [4, [5, 0], 14, [5, 1]] \\ C_3 &= [8, [1, 0], 7, [1, 1]] \\ C_4 &= [1, [6, 0], 10, [6, 1]] \end{split}$$

• 2-factor type [7, 3, 2, 2, 2]: two starters

$$\begin{split} C_0 &= [15, [15, 2], 6, [4, -1], 10, [5, 1], 5, [4, 0], 1, [6, 2], 7, [2, -1], 9, [15, 0]] \\ C_1 &= [2, [6, 0], 8, [4, 2], 12, [5, -1]] \\ C_2 &= [4, [6, -1], 13, [6, 1]] \\ C_3 &= [14, [3, 0], 11, [3, 2]] \\ C_4 &= [0, [3, -1], 3, [3, 1]] \\ C'_0 &= [3, [1, 1], 4, [1, 2], 5, [4, 1], 1, [1, 0], 0, [2, 1], 2, [7, 2], 10, [7, 0]] \\ C'_1 &= [15, [15, 1], 8, [1, -1], 9, [15, -1]] \\ C'_2 &= [14, [7, -1], 6, [7, 1]] \\ C'_3 &= [11, [2, 0], 13, [2, 2]] \\ C'_4 &= [12, [5, 0], 7, [5, 2]] \end{split}$$

$$\begin{split} C_0' &= [0, [5, -1], 10, [4, 2], 6, [7, -1], 13, [2, 0]] \\ C_1' &= [9, [15, 2], 15, [15, 1], 5, [4, 0]] \\ C_2' &= [1, [2, 2], 3, [1, 0], 4, [3, -1]] \\ C_3' &= [7, [4, -1], 11, [4, 1]] \\ C_4' &= [12, [2, -1], 14, [2, 1]] \\ C_5' &= [8, [6, -1], 2, [6, 1]] \end{split}$$

• 2-factor type [6, 4, 2, 2, 2]: two starters

$$C_{0} = [0, [2, 0], 13, [1, 1], 14, [3, -1], 11, [1, -1], 10, [2, -1], 8, [7, 2]]$$

$$C_{1} = [4, [2, 1], 6, [5, 2], 1, [4, -1], 5, [1, 0]]$$

$$C_{2} = [9, [7, -1], 2, [7, 1]]$$

$$C_{3} = [7, [5, -1], 12, [5, 1]]$$

$$C_{4} = [15, [15, 0], 3, [15, 2]]$$

$$C_{0}' = [6, [5, 0], 11, [1, 2], 12, [4, 1], 8, [7, 0], 0, [3, 1], 3, [3, 2]]$$

$$C_{1}' = [4, [15, 1], 15, [15, -1], 9, [2, 2], 7, [3, 0]]$$

$$C_{2}' = [5, [6, 0], 14, [6, 2]]$$

$$C_{3}' = [10, [6, -1], 1, [6, 1]]$$

$$C_{4}' = [2, [4, 0], 13, [4, 2]]$$
2-factor type [5, 5, 2, 2, 2]: two starters

$$C_{0} = [10, [2, 1], 8, [2, 0], 6, [7, -1], 14, [2, 2], 1, [6, 1]]$$

$$\begin{split} C_1 &= [4, [1, 1], 5, [5, 2], 0, [4, 0], 11, [15, 2], 15, [15, 0]] \\ C_2 &= [9, [6, 0], 3, [6, 2]] \\ C_3 &= [7, [5, -1], 2, [5, 1]] \\ C_4 &= [13, [1, 0], 12, [1, 2]] \\ C_0' &= [3, [2, -1], 5, [15, -1], 15, [15, 1], 9, [6, -1], 0, [3, -1]] \\ C_1' &= [2, [4, 2], 13, [5, 0], 8, [3, 1], 11, [1, -1], 10, [7, 1]] \\ C_2' &= [6, [7, 0], 14, [7, 2]] \\ C_3' &= [12, [4, -1], 1, [4, 1]] \\ C_4' &= [4, [3, 0], 7, [3, 2]] \end{split}$$

• 2-factor type [6, 3, 3, 2, 2]: one starter

•

$$\begin{split} C_0 &= [1, [6, 1], 10, [15, 0], 15, [15, 1], 5, [6, 0], 11, [7, 1], 3, [2, 1]] \\ C_1 &= [4, [4, 1], 0, [7, 0], 8, [4, 0]] \\ C_2 &= [12, [1, 0], 13, [1, 1], 14, [2, 0]] \\ C_3 &= [7, [5, 0], 2, [5, 1]] \\ C_4 &= [6, [3, 0], 9, [3, 1]] \end{split}$$

• 2-factor type
$$[5, 4, 3, 2, 2]$$
: one starter
 $C_0 = [9, [6, 1], 3, [4, 0], 7, [15, 0], 15, [15, 1], 8, [1, 1]]$
 $C_1 = [5, [1, 0], 6, [5, 0], 11, [4, 1], 0, [5, 1]]$
 $C_2 = [1, [3, 0], 13, [6, 0], 4, [3, 1]]$
 $C_3 = [12, [2, 0], 14, [2, 1]]$
 $C_4 = [2, [7, 0], 10, [7, 1]]$

- 2-factor type [4, 4, 4, 2, 2]: one starter $C_0 = [3, [7, 1], 11, [2, 0], 9, [5, 1], 14, [4, 0]]$ $C_1 = [6, [2, 1], 4, [4, 1], 0, [3, 0], 12, [6, 0]]$
 - $C_2 = [7, [6, 1], 13, [7, 0], 5, [5, 0], 10, [3, 1]]$ $C_3 = [2, [1, 0], 1, [1, 1]]$
 - $C_4 = [8, [15, 0], 15, [15, 1]]$
- 2-factor type [5, 3, 3, 3, 2]: two starters

$$\begin{split} C_0 &= [15, [15, 1], 3, [2, 1], 1, [3, -1], 4, [1, -1], 5, [15, -1]] \\ C_1 &= [2, [5, 2], 7, [1, 0], 8, [6, -1]] \\ C_2 &= [14, [1, 1], 0, [6, 2], 9, [5, 0]] \\ C_3 &= [10, [2, -1], 12, [6, 1], 6, [4, -1]] \\ C_4 &= [11, [2, 0], 13, [2, 2]] \\ C'_0 &= [1, [4, 0], 12, [5, -1], 7, [4, 1], 11, [3, 2], 8, [7, 1]] \\ C'_1 &= [0, [6, 0], 6, [7, -1], 14, [1, 2]] \\ C'_2 &= [3, [7, 2], 10, [3, 0], 13, [5, 1]] \\ C'_3 &= [5, [4, 2], 9, [7, 0], 2, [3, 1]] \\ C'_4 &= [4, [15, 0], 15, [15, 2]] \end{split}$$

• 2-factor type [4, 4, 3, 3, 2]: two starters

$$\begin{split} C_0 &= [10, [3, 0], 13, [6, 1], 4, [1, 1], 5, [5, 2]] \\ C_1 &= [3, [5, -1], 8, [15, 0], 15, [15, -1], 0, [3, 2]] \\ C_2 &= [9, [7, 1], 2, [6, 0], 11, [2, 2]] \\ C_3 &= [7, [1, -1], 6, [7, 2], 14, [7, 0]] \\ C_4 &= [12, [4, 0], 1, [4, 2]] \end{split}$$

- $\begin{array}{l} C_0' = [15, [15, 1], 7, [6, -1], 13, [5, 0], 8, [15, 2]] \\ 2]] & C_1' = [10, [4, 1], 14, [5, 1], 4, [7, -1], 12, [2, -1]] \\ C_2' = [0, [2, 1], 2, [1, 2], 1, [1, 0]] \\ C_3' = [11, [6, 2], 5, [4, -1], 9, [2, 0]] \\ C_4' = [3, [3, -1], 6, [3, 1]] \end{array}$
- 2-factor type [4, 3, 3, 3, 3]: one starter $C_0 = [9, [15, 0], 15, [15, 1], 12, [5, 1], 7, [2, 0]]$ $C_1 = [13, [3, 0], 1, [1, 1], 2, [4, 0]]$ $C_2 = [0, [7, 0], 8, [2, 1], 6, [6, 0]]$ $C_3 = [5, [1, 0], 4, [6, 1], 10, [5, 0]]$ $C_4 = [11, [3, 1], 14, [4, 1], 3, [7, 1]]$
- 2-factor type [10, 2, 2, 2]: two starters $C_0 = [8, [5, 1], 13, [2, 2], 0, [3, -1], 3, [7, 1], 11, [6, -1], 2, [2, 0], 4, [5, -1], 14, [15, 2], 15, [15, 0], 12, [4, -1]]$ $C_1 = [6, [1, -1], 5, [1, 1]]$ $C_2 = [9, [7, 0], 1, [7, 2]]$ $C_3 = [10, [3, 0], 7, [3, 2]]$ $C'_0 = [3, [3, 1], 6, [4, 2], 2, [5, 0], 12, [4, 1], 1, [15, 1], 15, [15, -1], 0, [5, 2], 5, [6, 1], 14, [4, 0], 10, [7, -1]]$ $C'_1 = [11, [2, -1], 9, [2, 1]]$ $C'_2 = [4, [6, 0], 13, [6, 2]]$ $C'_3 = [7, [1, 0], 8, [1, 2]]$

• 2-factor type [9, 3, 2, 2]: one starter

$$\begin{split} C_0 &= [7, [7, 1], 14, [1, 0], 13, [3, 0], 1, [7, 0], 8, [4, 0], 4, [15, 0], 15, [15, 1], 11, [2, 0], 9, [2, 1]] \\ C_1 &= [5, [1, 1], 6, [4, 1], 2, [3, 1]] \\ C_2 &= [3, [6, 0], 12, [6, 1]] \\ C_3 &= [0, [5, 0], 10, [5, 1]] \end{split}$$

• 2-factor type [8, 4, 2, 2]: one starter

$$\begin{split} C_0 &= [12, [6, 0], 6, [4, 1], 2, [7, 1], 10, [6, 1], 1, [1, 1], 0, [15, 1], 15, [15, 0], 9, [3, 1]] \\ C_1 &= [4, [1, 0], 3, [4, 0], 14, [3, 0], 11, [7, 0]] \\ C_2 &= [7, [2, 0], 5, [2, 1]] \\ C_3 &= [13, [5, 0], 8, [5, 1]] \end{split}$$

• 2-factor type [7, 5, 2, 2]: one starter

$$\begin{split} C_0 &= [14, [7, 1], 7, [5, 0], 12, [15, 0], 15, [15, 1], 13, [2, 0], 0, [6, 1], 6, [7, 0]] \\ C_1 &= [10, [6, 0], 4, [3, 0], 1, [5, 1], 11, [3, 1], 8, [2, 1]] \\ C_2 &= [3, [1, 0], 2, [1, 1]] \\ C_3 &= [9, [4, 0], 5, [4, 1]] \end{split}$$

• 2-factor type [6, 6, 2, 2]: one starter

$$\begin{split} C_0 &= [0, [3, 1], 3, [2, 1], 5, [3, 0], 8, [2, 0], 10, [6, 1], 1, [1, 1]]\\ C_1 &= [7, [15, 1], 15, [15, 0], 11, [7, 1], 4, [7, 0], 12, [1, 0], 13, [6, 0]]\\ C_2 &= [6, [4, 0], 2, [4, 1]]\\ C_3 &= [14, [5, 0], 9, [5, 1]] \end{split}$$

• 2-factor type [8, 3, 3, 2]: two starters

$$\begin{split} C_0 &= [0, [4, 2], 4, [7, -1], 11, [3, 0], 8, [4, 1], 12, [15, 2], 15, [15, 1], 14, [2, 0], 1, [1, -1]] \\ C_1 &= [10, [7, 0], 2, [1, 1], 3, [7, 2]] \\ C_2 &= [13, [7, 1], 5, [2, 1], 7, [6, 1]] \\ C_3 &= [6, [3, -1], 9, [3, 1]] \\ C'_0 &= [7, [4, 0], 11, [3, 2], 14, [5, 0], 4, [1, 2], 3, [6, -1], 9, [6, 0], 0, [2, -1], 13, [6, 2]] \\ C'_1 &= [2, [1, 0], 1, [4, -1], 12, [5, 2]] \\ C'_2 &= [6, [15, -1], 15, [15, 0], 8, [2, 2]] \\ C'_3 &= [5, [5, -1], 10, [5, 1]] \end{split}$$

• 2-factor type [7, 4, 3, 2]: two starters

$$\begin{split} C_0 &= [15, [15, -1], 14, [7, 2], 7, [1, 1], 6, [6, 1], 0, [7, 0], 8, [6, 2], 2, [15, 0]] \\ C_1 &= [4, [7, -1], 11, [2, 0], 13, [4, -1], 9, [5, 2]] \\ C_2 &= [1, [6, -1], 10, [2, 2], 12, [4, 0]] \\ C_3 &= [5, [2, -1], 3, [2, 1]] \\ C'_0 &= [0, [4, 2], 11, [1, 0], 10, [1, 2], 9, [5, -1], 14, [3, 0], 2, [3, 2], 5, [5, 0]] \\ C'_1 &= [13, [5, 1], 8, [4, 1], 4, [7, 1], 12, [1, -1]] \\ C'_2 &= [7, [6, 0], 1, [15, 1], 15, [15, 2]] \\ C'_3 &= [6, [3, -1], 3, [3, 1]] \end{split}$$

• 2-factor type [6, 5, 3, 2]: two starters

$$\begin{split} C_0 &= [10, [5, -1], 5, [3, -1], 2, [2, -1], 0, [3, 1], 3, [6, 2], 9, [1, 0]] \\ C_1 &= [11, [3, 0], 8, [7, -1], 1, [5, 1], 6, [7, 2], 13, [2, 1]] \\ C_2 &= [7, [3, 2], 4, [7, 1], 12, [5, 0]] \\ C_3 &= [15, [15, -1], 14, [15, 1]] \\ C'_0 &= [4, [6, -1], 13, [1, 2], 12, [4, -1], 1, [4, 0], 5, [5, 2], 10, [6, 0]] \\ C'_1 &= [0, [6, 1], 9, [7, 0], 2, [1, 1], 3, [4, 2], 14, [1, -1]] \\ C'_2 &= [7, [4, 1], 11, [15, 2], 15, [15, 0]] \\ C'_3 &= [8, [2, 0], 6, [2, 2]] \\ \bullet \ 2\text{-factor type } [6, 4, 4, 2]: \text{ two starters} \end{split}$$

- $$\begin{split} C_0 &= [9, [2, 0], 7, [7, 1], 14, [1, 2], 0, [15, -1], 15, [15, 1], 13, [4, 1]] \\ C_1 &= [12, [1, 1], 11, [7, 0], 4, [1, -1], 3, [6, 2]] \\ C_2 &= [2, [6, 0], 8, [3, -1], 5, [5, 2], 10, [7, -1]] \\ C_3 &= [1, [5, -1], 6, [5, 1]] \\ C_0' &= [4, [6, -1], 13, [2, 1], 0, [6, 1], 6, [4, 2], 2, [1, 0], 1, [3, 1]] \\ C_1' &= [10, [2, 2], 12, [5, 0], 7, [7, 2], 14, [4, 0]] \\ C_2' &= [9, [4, -1], 5, [3, 2], 8, [3, 0], 11, [2, -1]] \\ C_3' &= [15, [15, 0], 3, [15, 2]] \end{split}$$
- 2-factor type [5, 5, 4, 2]: two starters

$$\begin{split} C_0 &= [8, [5, 0], 13, [7, -1], 6, [6, 2], 12, [1, 1], 11, [3, 1]] \\ C_1 &= [15, [15, 1], 7, [6, 0], 1, [1, 2], 2, [3, -1], 5, [15, -1]] \\ C_2 &= [4, [4, -1], 0, [5, 2], 10, [7, 1], 3, [1, 0]] \\ C_3 &= [14, [5, -1], 9, [5, 1]] \\ C_0' &= [12, [1, -1], 11, [15, 2], 15, [15, 0], 5, [3, 2], 8, [4, 0]] \\ C_1' &= [1, [3, 0], 4, [6, 1], 10, [7, 2], 3, [4, 1], 14, [2, -1]] \\ C_2' &= [6, [7, 0], 13, [4, 2], 2, [2, 1], 0, [6, -1]] \\ C_3' &= [7, [2, 0], 9, [2, 2]] \end{split}$$

• 2-factor type [7, 3, 3, 3]: one starter

$$\begin{split} C_0 &= [12, [7, 0], 5, [2, 1], 7, [15, 1], 15, [15, 0], 14, [1, 1], 0, [6, 0], 9, [3, 0]] \\ C_1 &= [13, [4, 1], 2, [1, 0], 3, [5, 0]] \\ C_2 &= [8, [3, 1], 11, [5, 1], 1, [7, 1]] \\ C_3 &= [4, [6, 1], 10, [4, 0], 6, [2, 0]] \end{split}$$

• 2-factor type [6, 4, 3, 3]: one starter

 $C_0 = [11, [4, 0], 0, [5, 1], 10, [7, 1], 2, [15, 1], 15, [15, 0], 14, [3, 1]]$ $C_1 = [7, [2, 0], 5, [7, 0], 12, [4, 1], 1, [6, 1]]$ $C_2 = [9, [1, 0], 8, [2, 1], 6, [3, 0]]$ $C_3 = [4, [1, 1], 3, [5, 0], 13, [6, 0]]$ • 2-factor type [5, 5, 3, 3]: one starter

 $C_0 = [1, [15, 0], 15, [15, 1], 12, [7, 0], 5, [6, 1], 14, [2, 1]]$ $C_1 = [10, [4, 1], 6, [6, 0], 0, [3, 1], 3, [7, 1], 11, [1, 0]]$ $C_2 = [2, [2, 0], 4, [3, 0], 7, [5, 1]]$ $C_3 = [8, [1, 1], 9, [4, 0], 13, [5, 0]]$

• 2-factor type [5, 4, 4, 3]: one starter

$$\begin{split} C_0 &= [12, [6, 1], 3, [6, 0], 9, [4, 0], 5, [7, 1], 13, [1, 1]] \\ C_1 &= [0, [15, 1], 15, [15, 0], 2, [5, 1], 7, [7, 0]] \\ C_2 &= [8, [2, 1], 6, [4, 1], 10, [1, 0], 11, [3, 0]] \\ C_3 &= [14, [2, 0], 1, [3, 1], 4, [5, 0]] \end{split}$$

• 2-factor type [12, 2, 2]: one starter

$$\begin{split} C_0 &= [15, [15, 1], 14, [6, 1], 5, [5, 0], 0, [2, 1], 13, [7, 1], 6, [5, 1], 11, [7, 0], 4, [2, 0], 2, [1, 1], 3, [6, 0], 9, [1, 0], \\ &\quad 8, [15, 0]] \\ C_1 &= [7, [3, 0], 10, [3, 1]] \\ C_2 &= [1, [4, 0], 12, [4, 1]] \end{split}$$

• 2-factor type [11, 3, 2]: two starters

 $C_{0} = [6, [1, 2], 5, [5, 0], 0, [3, 2], 3, [4, 0], 14, [2, -1], 12, [5, 2], 7, [6, 0], 13, [2, 2], 11, [15, 0], 15, [15, 2], 8, [2, 0]]$ $C_{1} = [2, [7, 1], 10, [1, 0], 9, [7, 2]]$ $C_{2} = [4, [3, -1], 1, [3, 1]]$

$$\begin{split} C_0' &= [5, [6, 1], 11, [4, 1], 0, [7, -1], 8, [1, -1], 7, [2, 1], 9, [4, 2], 13, [6, -1], 4, [7, 0], 12, [6, 2], 3, [1, 1], 2, [3, 0]]\\ C_1' &= [14, [15, -1], 15, [15, 1], 10, [4, -1]]\\ C_2' &= [6, [5, -1], 1, [5, 1]] \end{split}$$

• 2-factor type [10, 4, 2]: two starters

$$\begin{split} C_0 &= [14, [7, 2], 6, [4, 0], 2, [4, -1], 13, [2, 2], 0, [5, -1], 5, [2, 0], 7, [5, 1], 12, [7, 1], 4, [3, -1], 1, [2, -1]] \\ C_1 &= [3, [5, 2], 8, [15, -1], 15, [15, 1], 11, [7, 0]] \\ C_2 &= [10, [1, 0], 9, [1, 2]] \\ C_0' &= [0, [4, 1], 4, [6, 2], 13, [5, 0], 3, [6, -1], 12, [4, 2], 1, [1, -1], 2, [7, -1], 9, [2, 1], 7, [1, 1], 6, [6, 0]] \\ C_1' &= [14, [6, 1], 5, [3, 1], 8, [3, 2], 11, [3, 0]] \\ C_2' &= [10, [15, 0], 15, [15, 2]] \end{split}$$

• 2-factor type [9, 5, 2]: two starters

$$\begin{split} C_0 &= [6, [3, 0], 3, [4, 2], 14, [5, 0], 4, [7, 2], 12, [4, -1], 1, [1, 0], 2, [15, 2], 15, [15, 1], 7, [1, 1]] \\ C_1 &= [0, [7, 1], 8, [3, 1], 5, [7, 0], 13, [3, -1], 10, [5, 2]] \\ C_2 &= [11, [2, -1], 9, [2, 1]] \\ C'_0 &= [1, [2, 0], 14, [3, 2], 2, [6, 0], 11, [4, 1], 0, [6, 2], 6, [15, -1], 15, [15, 0], 9, [6, -1], 3, [2, 2]] \\ C'_1 &= [13, [6, 1], 7, [1, -1], 8, [4, 0], 4, [7, -1], 12, [1, 2]] \\ C'_2 &= [5, [5, -1], 10, [5, 1]] \end{split}$$

• 2-factor type [8, 6, 2]: two starters

$$\begin{split} C_0 &= [6, [2, 2], 8, [1, 0], 9, [7, 2], 1, [3, 1], 13, [5, 0], 3, [7, -1], 11, [1, 2], 10, [4, 0]] \\ C_1 &= [15, [15, 2], 12, [3, 0], 0, [2, 1], 2, [5, 2], 7, [2, 0], 5, [15, 1]] \\ C_2 &= [4, [5, -1], 14, [5, 1]] \\ C'_0 &= [0, [7, 0], 7, [4, 2], 11, [1, 1], 10, [7, 1], 3, [3, -1], 6, [2, -1], 8, [1, -1], 9, [6, -1]] \\ C'_1 &= [2, [15, 0], 15, [15, -1], 12, [4, 1], 1, [3, 2], 4, [6, 1], 13, [4, -1]] \\ C'_2 &= [5, [6, 0], 14, [6, 2]] \end{split}$$

• 2-factor type [7,7,2]: two starters

$$\begin{split} C_0 &= [10, [1, -1], 9, [4, -1], 5, [2, 2], 3, [4, 0], 14, [15, 2], 15, [15, 0], 1, [6, -1]] \\ C_1 &= [0, [6, 0], 6, [4, 2], 2, [4, 1], 13, [1, 1], 12, [5, 0], 7, [1, 2], 8, [7, -1]] \\ C_2 &= [4, [7, 0], 11, [7, 2]] \\ C_0' &= [2, [5, 2], 12, [5, 1], 7, [2, 0], 9, [6, 2], 3, [3, 0], 0, [5, -1], 10, [7, 1]] \\ C_1' &= [13, [1, 0], 14, [6, 1], 5, [3, 2], 8, [3, -1], 11, [15, -1], 15, [15, 1], 1, [3, 1]] \\ C_2' &= [4, [2, -1], 6, [2, 1]] \end{split}$$

• 2-factor type [10, 3, 3]: one starter

 $C_0 = [1, [4, 0], 5, [1, 1], 6, [4, 1], 10, [7, 1], 2, [6, 1], 11, [7, 0], 4, [15, 1], 15, [15, 0], 0, [3, 0], 3, [2, 1]]$ $C_1 = [9, [3, 1], 12, [5, 0], 7, [2, 0]]$ $C_2 = [13, [5, 1], 8, [6, 0], 14, [1, 0]]$

• 2-factor type [9, 4, 3]: one starter

$$\begin{split} C_0 &= [9, [3, 1], 6, [6, 1], 12, [2, 0], 14, [1, 0], 0, [4, 1], 11, [7, 1], 4, [4, 0], 8, [15, 1], 15, [15, 0]] \\ C_1 &= [1, [1, 1], 2, [5, 1], 7, [3, 0], 10, [6, 0]] \\ C_2 &= [3, [5, 0], 13, [7, 0], 5, [2, 1]] \end{split}$$

• 2-factor type [8, 5, 3]: one starter

$$\begin{split} C_0 &= [1, [7, 1], 8, [5, 0], 3, [15, 0], 15, [15, 1], 10, [1, 1], 9, [6, 1], 0, [2, 1], 13, [3, 1]] \\ C_1 &= [5, [3, 0], 2, [4, 0], 6, [1, 0], 7, [4, 1], 11, [6, 0]] \\ C_2 &= [14, [5, 1], 4, [7, 0], 12, [2, 0]] \end{split}$$

• 2-factor type [7, 6, 3]: one starter

$$\begin{split} C_0 &= [14, [2, 0], 12, [15, 0], 15, [15, 1], 8, [1, 0], 7, [3, 1], 4, [6, 0], 13, [1, 1]] \\ C_1 &= [10, [5, 1], 0, [2, 1], 2, [7, 0], 9, [4, 0], 5, [4, 1], 1, [6, 1]] \\ C_2 &= [6, [5, 0], 11, [7, 1], 3, [3, 0]] \end{split}$$

• 2-factor type [7, 5, 4]: one starter

$$\begin{split} C_0 &= [10, [3, 1], 13, [15, 0], 15, [15, 1], 5, [7, 0], 12, [6, 1], 3, [3, 0], 6, [4, 0]]\\ C_1 &= [0, [1, 0], 14, [2, 0], 1, [5, 1], 11, [4, 1], 7, [7, 1]]\\ C_2 &= [4, [2, 1], 2, [6, 0], 8, [1, 1], 9, [5, 0]] \end{split}$$

• 2-factor type [6, 6, 4]: one starter

$$\begin{split} C_0 &= [4, [7, 1], 12, [3, 1], 0, [6, 1], 6, [2, 0], 8, [1, 0], 9, [5, 1]] \\ C_1 &= [15, [15, 1], 10, [5, 0], 5, [6, 0], 14, [7, 0], 7, [4, 1], 11, [15, 0]] \\ C_2 &= [13, [4, 0], 2, [1, 1], 3, [2, 1], 1, [3, 0]] \end{split}$$

- 2-factor type [6, 5, 5]: one starter
 - $$\begin{split} C_0 &= [2, [15, 1], 15, [15, 0], 3, [2, 0], 1, [5, 0], 6, [3, 0], 9, [7, 1]]\\ C_1 &= [8, [1, 0], 7, [7, 0], 14, [4, 1], 10, [6, 0], 4, [4, 0]]\\ C_2 &= [5, [6, 1], 11, [2, 1], 13, [1, 1], 12, [3, 1], 0, [5, 1]] \end{split}$$
- 2-factor type [14, 2]: two starters

$$\begin{split} C_0 &= [10, [4, 0], 14, [2, -1], 12, [3, 2], 9, [2, 1], 7, [2, 0], 5, [7, -1], 13, [15, 1], 15, [15, 2], 11, [6, 0], 2, [6, 1], 8, \\ &\quad [7, 2], 1, [1, 1], 0, [4, -1], 4, [6, -1]] \\ C_1 &= [3, [3, -1], 6, [3, 1]] \\ C_0' &= [0, [2, 2], 2, [4, 1], 13, [15, -1], 15, [15, 0], 9, [6, 2], 3, [1, -1], 4, [7, 0], 12, [5, -1], 7, [1, 2], 8, [3, 0], 5, \end{split}$$

- [5,1], 10, [4,2], 6, [7,1], 14, [1,0]]
- $C_1' = [1, [5, 0], 11, [5, 2]]$
- 2-factor type [13, 3]: one starter $C_0 = [8, [7, 0], 1, [2, 0], 14, [4, 1], 10, [1, 0], 11, [5, 0], 6, [3, 1], 9, [4, 0], 5, [3, 0], 2, [2, 1], 0, [15, 1], 15, [15, 0], 7, [6, 0], 13, [5, 1]]$ $C_1 = [3, [1, 1], 4, [7, 1], 12, [6, 1]]$
- 2-factor type [11,5]: one starter

$$\begin{split} C_0 &= [12, [1, 0], 13, [4, 0], 9, [6, 1], 3, [7, 0], 10, [6, 0], 1, [5, 1], 6, [1, 1], 5, [3, 1], 8, [15, 1], 15, [15, 0], 14, [2, 0]] \\ C_1 &= [2, [5, 0], 7, [3, 0], 4, [7, 1], 11, [4, 1], 0, [2, 1]] \end{split}$$

• 2-factor type [10, 6]: one starter

$$\begin{split} C_0 &= [9, [7, 1], 1, [1, 1], 2, [15, 0], 15, [15, 1], 6, [2, 1], 8, [3, 0], 11, [4, 1], 0, [5, 0], 5, [2, 0], 3, [6, 1]] \\ C_1 &= [13, [6, 0], 4, [7, 0], 12, [5, 1], 7, [3, 1], 10, [4, 0], 14, [1, 0]] \end{split}$$

• 2-factor type [9,7]: one starter

$$\begin{split} C_0 &= [9, [6, 1], 0, [3, 1], 3, [15, 0], 15, [15, 1], 5, [1, 0], 4, [7, 1], 11, [4, 0], 7, [7, 0], 14, [5, 1]]\\ C_1 &= [6, [5, 0], 1, [3, 0], 13, [1, 1], 12, [2, 1], 10, [2, 0], 8, [6, 0], 2, [4, 1]] \end{split}$$

I Computational results for n = 17

• 2-factor type [3, 2, 2, 2, 2, 2, 2, 2]: three starters

$C'_0 = [10, [7, 0], 3, [1, 1], 4, [6, 2]]$
$C_1' = [13, [2, -1], 15, [2, 1]]$
$C_2' = [12, [1, 0], 11, [1, 2]]$
$C_3' = [14, [3, -1], 1, [3, 1]]$
$C_4' = [16, [16, -1], 5, [16, 1]]$
$C_5' = [7, [5, -1], 2, [5, 1]]$
$C_6' = [6, [3, 0], 9, [3, 2]]$
$C_7' = [0, [8, 0], 8, [8, 2]]$

• 2-factor type [5, 2, 2, 2, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [7, [6, -1], 13, [3, 0], 10, [5, 2], 15, [1, 0], 14, [7, 2]] \\ C_1 &= [3, [1, -1], 4, [1, 1]] \\ C_2 &= [12, [6, 0], 6, [6, 2]] \\ C_3 &= [2, [7, -1], 9, [7, 1]] \\ C_4 &= [1, [4, -1], 5, [4, 1]] \\ C_5 &= [0, [5, -1], 11, [5, 1]] \\ C_6 &= [16, [16, -1], 8, [16, 1]] \\ C'_0 &= [12, [3, 2], 15, [7, 0], 6, [1, 2], 7, [6, 1], 1, [5, 0]] \\ C'_1 &= [0, [8, 0], 8, [8, 2]] \\ C'_2 &= [3, [2, -1], 5, [2, 1]] \\ C'_3 &= [13, [3, -1], 10, [3, 1]] \\ C'_4 &= [16, [16, 0], 4, [16, 2]] \\ C'_5 &= [14, [4, 0], 2, [4, 2]] \\ C'_6 &= [9, [2, 0], 11, [2, 2]] \end{split}$$

• 2-factor type [4, 3, 2, 2, 2, 2, 2]: three starters

$$C_{0} = [9, [5, 0], 14, [1, 2], 15, [3, 0], 2, [7, 2]]$$

$$C_{1} = [12, [16, 0], 16, [16, 2], 6, [6, -1]]$$

$$C_{2} = [0, [5, -1], 11, [5, 1]]$$

$$C_{3} = [5, [3, -1], 8, [3, 1]]$$

$$C_{4} = [4, [6, 0], 10, [6, 2]]$$

$$C_{5} = [13, [4, -1], 1, [4, 1]]$$

$$C_{6} = [7, [4, 0], 3, [4, 2]]$$

$$C_{1} = [7, [2, 3, 2], 5, [7, 0], 14, [16, 1], 16, [16, -1]]$$

$$C_{1}' = [7, [1, 0], 6, [6, 1], 12, [5, 2]]$$

$$C_{2}' = [0, [8, 0], 8, [8, 2]]$$

$$C_{2}' = [0, [8, 0], 8, [8, 2]]$$

$$C_{3}' = [9, [1, -1], 10, [1, 1]]$$

$$C_{4}' = [3, [2, 0], 1, [2, 2]]$$

$$C_{5}' = [4, [7, -1], 11, [7, 1]]$$

$$C_{6}' = [15, [2, -1], 13, [2, 1]]$$

• 2-factor type [3, 3, 3, 2, 2, 2, 2]: three starters

$C_0' = [10, [16, -1], 16, [16, 1], 9, [1, -1]]$
$C_1' = [11, [2, 0], 13, [7, -1], 6, [5, 2]]$
$C_2' = [12, [5, 1], 7, [4, 0], 3, [7, 2]]$
$C'_3 = [0, [8, 0], 8, [8, 2]]$
$C'_4 = [1, [1, 0], 2, [1, 2]]$
$C'_5 = [5, [6, 0], 15, [6, 2]]$
$C_6' = [4, [6, -1], 14, [6, 1]]$

• 2-factor type [7, 2, 2, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [9, [1, 0], 8, [1, -1], 7, [5, -1], 2, [1, 2], 1, [3, -1], 4, [7, 0], 11, [2, 2]] \\ C_1 &= [15, [7, -1], 6, [7, 1]] \\ C_2 &= [16, [16, -1], 14, [16, 1]] \\ C_3 &= [5, [2, -1], 3, [2, 1]] \\ C_4 &= [10, [3, 0], 13, [3, 2]] \\ C_5 &= [12, [4, -1], 0, [4, 1]] \end{split}$$

- $$\begin{split} C_0 &= [6, [2, 0], 4, [5, 1], 9, [5, 2], 14, [3, 1], 11, [1, 1], 10, [5, 0], 15, [7, 2]] \\ C_1 &= [0, [8, 0], 8, [8, 2]] \\ C_2 &= [2, [6, 0], 12, [6, 2]] \\ C_3 &= [16, [16, 0], 7, [16, 2]] \\ C_4 &= [5, [4, 0], 1, [4, 2]] \\ C_5 &= [13, [6, -1], 3, [6, 1]] \end{split}$$
- 2-factor type [6, 3, 2, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [1, [3, 0], 4, [3, 2], 7, [7, 0], 14, [6, -1], 8, [5, 2], 13, [4, -1]] \\ C_1 &= [2, [2, 0], 0, [6, 2], 6, [4, 1]] \\ C_2 &= [10, [5, -1], 5, [5, 1]] \\ C_3 &= [11, [16, 0], 16, [16, 2]] \\ C_4 &= [15, [4, 0], 3, [4, 2]] \\ C_5 &= [9, [3, -1], 12, [3, 1]] \\ C'_0 &= [7, [2, 1], 5, [5, 0], 10, [2, 2], 12, [6, 0], 6, [7, 2], 13, [6, 1]] \\ C'_1 &= [2, [1, 2], 1, [2, -1], 3, [1, 0]] \\ C'_2 &= [0, [8, 0], 8, [8, 2]] \\ C'_3 &= [9, [16, -1], 16, [16, 1]] \\ C'_4 &= [4, [7, -1], 11, [7, 1]] \\ C'_5 &= [15, [1, -1], 14, [1, 1]] \\ \bullet & 2\text{-factor type } [5, 4, 2, 2, 2, 2]: \text{ three starters} \\ C_0 &= [14, [1, 2], 13, [4, 0], 1, [5, 2], 12, [3, 1], 9, [5, 0]] \\ C_1 &= [10, [6, 2], 4, [16, -1], 16, [16, 1], 3, [7, 0]] \\ C_2 &= [5, [6, -1], 11, [6, 1]] \\ C_3 &= [8, [2, -1], 6, [2, 1]] \\ C_4 &= [15, [3, 0], 2, [3, 2]] \\ C_5 &= [0, [7, -1], 7, [7, 1]] \\ C'_0 &= [7, [5, -1], 12, [16, 0], 16, [16, 2], 9, [6, 0], 3, [4, 2]] \\ C'_1 &= [11, [3, -1], 14, [1, 0], 13, [5, 1], 2, [7, 2]] \\ C'_2 &= [0, [8, 0], 8, [8, 2]] \\ C'_3 &= [10, [4, -1], 6, [4, 1]] \\ C'_4 &= [15, [2, 0], 1, [2, 2]] \\ C'_5 &= [5, [1, -1], 4, [1, 1]] \\ \bullet & 2\text{-factor type } [5, 3, 3, 2, 2, 2]: \text{ three starters} \end{aligned}$$

 $C_{0} = [7, [2, 0], 5, [6, 1], 11, [2, 2], 13, [1, -1], 14, [7, 1]]$ $C_{1} = [1, [1, 2], 2, [1, 0], 3, [2, -1]]$ $C_{2} = [10, [4, 2], 6, [2, 1], 4, [6, 0]]$ $C_{3} = [0, [4, -1], 12, [4, 1]]$ $C_{4} = [15, [7, 0], 8, [7, 2]]$ $C_{5} = [16, [16, 0], 9, [16, 2]]$

- $$\begin{split} C_0' &= [16, [16, -1], 11, [3, 2], 14, [6, -1], 4, [5, 0], 9, [16, 1]] \\ C_1' &= [5, [4, 0], 1, [5, 2], 6, [1, 1]] \\ C_2' &= [13, [6, 2], 3, [7, -1], 10, [3, 0]] \\ C_3' &= [0, [8, 0], 8, [8, 2]] \\ C_4' &= [2, [5, -1], 7, [5, 1]] \\ C_5' &= [12, [3, -1], 15, [3, 1]] \end{split}$$
- 2-factor type [4, 4, 3, 2, 2, 2]: three starters
 - $$\begin{split} C_0 &= [1, [3, 0], 14, [1, 2], 15, [6, 0], 5, [4, 2]] \\ C_1 &= [6, [6, -1], 12, [1, 0], 13, [6, 1], 3, [3, 2]] \\ C_2 &= [7, [5, -1], 2, [7, 2], 9, [2, 0]] \\ C_3 &= [11, [1, -1], 10, [1, 1]] \\ C_4 &= [4, [4, -1], 0, [4, 1]] \\ C_5 &= [8, [16, -1], 16, [16, 1]] \end{split}$$
- 2-factor type [4, 3, 3, 3, 2, 2]: three starters
 - $$\begin{split} C_0 &= [8, [1, -1], 9, [5, 1], 4, [6, 0], 14, [6, 2]] \\ C_1 &= [5, [5, -1], 0, [2, 1], 2, [3, 1]] \\ C_2 &= [7, [4, 0], 3, [7, 1], 10, [3, 2]] \\ C_3 &= [6, [16, 0], 16, [16, -1], 13, [7, 2]] \\ C_4 &= [1, [2, 0], 15, [2, 2]] \\ C_5 &= [12, [1, 0], 11, [1, 2]] \end{split}$$
- 2-factor type [3, 3, 3, 3, 3, 2]: three starters
 - $C_{0} = [4, [4, 2], 8, [5, 1], 3, [1, 0]]$ $C_{1} = [9, [1, 1], 10, [3, 2], 13, [4, 0]]$ $C_{2} = [5, [4, -1], 1, [2, -1], 15, [6, 1]]$ $C_{3} = [6, [6, 2], 12, [5, 0], 7, [1, -1]]$ $C_{4} = [2, [16, 2], 16, [16, 1], 11, [7, 0]]$ $C_{5} = [14, [2, 0], 0, [2, 2]]$

- $$\begin{split} C_0' &= [7, [2, -1], 9, [3, 1], 6, [2, 1], 4, [3, -1]] \\ C_1' &= [2, [7, -1], 11, [6, 2], 1, [4, 0], 13, [5, 1]] \\ C_2' &= [12, [2, 2], 14, [7, 0], 5, [7, 1]] \\ C_3' &= [0, [8, 0], 8, [8, 2]] \\ C_4' &= [16, [16, 0], 3, [16, 2]] \\ C_5' &= [10, [5, 0], 15, [5, 2]] \end{split}$$
- $$\begin{split} C_0' &= [4, [1, 1], 3, [4, 1], 15, [2, -1], 1, [3, -1]] \\ C_1' &= [10, [4, 2], 6, [7, -1], 13, [3, 0]] \\ C_2' &= [14, [5, 2], 9, [7, 0], 2, [4, -1]] \\ C_3' &= [16, [16, 2], 7, [5, 0], 12, [16, 1]] \\ C_4' &= [0, [8, 0], 8, [8, 2]] \\ C_5' &= [11, [6, -1], 5, [6, 1]] \end{split}$$
- $C'_{0} = [10, [1, 2], 11, [7, 1], 4, [6, 0]]$ $C'_{1} = [9, [3, 0], 6, [7, 2], 13, [4, 1]]$ $C'_{2} = [7, [5, 2], 2, [16, -1], 16, [16, 0]]$ $C'_{3} = [5, [7, -1], 12, [3, -1], 15, [6, -1]]$ $C'_{4} = [3, [2, 1], 1, [3, 1], 14, [5, -1]]$ $C'_{5} = [0, [8, 0], 8, [8, 2]]$
- 2-factor type [9, 2, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [3, [5, 2], 8, [7, 1], 1, [3, 0], 14, [4, -1], 10, [3, 2], 13, [3, -1], 0, [16, -1], 16, [16, 1], 2, [1, 0]] \\ C_1 &= [12, [6, 0], 6, [6, 2]] \\ C_2 &= [7, [2, 0], 9, [2, 2]] \\ C_3 &= [15, [4, 0], 11, [4, 2]] \\ C_4 &= [5, [1, -1], 4, [1, 1]] \\ C_6 &= [10, [7, -1], 1, [16, 0], 16, [16, 2], 13, [4, 1], 9, [3, 1], 6, [7, 0], 15, [1, 2], 14, [5, 0], 3, [7, 2]] \\ C_1' &= [0, [8, 0], 8, [8, 2]] \\ C_2' &= [7, [5, -1], 12, [5, 1]] \\ C_3' &= [4, [2, -1], 2, [2, 1]] \\ C_4' &= [5, [6, -1], 11, [6, 1]] \end{split}$$

• 2-factor type [8, 3, 2, 2, 2]: three starters

$$\begin{array}{l} C_0 = [14, [4, 0], 10, [7, 2], 3, [6, 0], 13, [5, 1], 2, [2, 1], 4, [4, 2], 0, [7, 1], 9, [5, -1]] \\ C_1 = [5, [3, -1], 8, [3, 0], 11, [6, 2]] \\ C_2 = [16, [16, -1], 12, [16, 1]] \\ C_3 = [15, [2, 0], 1, [2, 2]] \\ C_4 = [6, [1, -1], 7, [1, 1]] \\ \hline \\ C_0' = [12, [2, -1], 10, [3, 1], 13, [5, 2], 2, [7, 0], 11, [3, 2], 14, [1, 0], 15, [4, 1], 3, [7, -1]] \\ C_1' = [4, [1, 2], 5, [4, -1], 9, [5, 0]] \\ C_2' = [0, [8, 0], 8, [8, 2]] \\ C_3' = [1, [6, -1], 7, [6, 1]] \\ C_4' = [16, [16, 0], 6, [16, 2]] \\ 2 \text{-factor type } [7, 4, 2, 2, 2]: \text{ three starters} \\ \hline \\ C_0 = [2, [1, 0], 3, [6, 2], 13, [1, -1], 14, [16, -1], 16, [16, 1], 12, [4, 0], 0, [2, 2]] \\ \hline \\ C_1 = [7, [1, 1], 6, [2, -1], 8, [7, 1], 1, [6, -1]] \\ C_2 = [10, [5, -1], 15, [5, 1]] \\ C_3 = [11, [7, 0], 4, [7, 2]] \\ C_4 = [9, [4, -1], 5, [4, 1]] \\ \hline \\ C_0' = [13, [6, 0], 3, [2, 1], 5, [1, 2], 4, [7, -1], 11, [5, 0], 6, [4, 2], 10, [3, -1]] \\ \hline \\ C_1' = [2, [5, 2], 7, [6, 1], 1, [2, 0], 15, [3, 1]] \\ \hline \\ C_2' = [0, [8, 0], 8, [8, 2]] \\ \hline \\ C_3' = [16, [16, 0], 14, [16, 2]] \\ \hline \\ C_4' = [12, [3, 0], 9, [3, 2]] \\ \hline \end{array}$$

• 2-factor type [6, 5, 2, 2, 2]: three starters

٠

٠

$$\begin{split} C_0 &= [3, [4, 2], 15, [7, 0], 8, [3, 2], 5, [6, 0], 11, [1, -1], 10, [7, -1]] \\ C_1 &= [2, [1, 2], 1, [3, -1], 4, [2, -1], 6, [16, 0], 16, [16, -1]] \\ C_2 &= [12, [5, -1], 7, [5, 1]] \\ C_3 &= [13, [4, -1], 9, [4, 1]] \\ C_4 &= [14, [2, 0], 0, [2, 2]] \\ C'_0 &= [9, [16, 1], 16, [16, 2], 11, [4, 0], 7, [6, 2], 13, [7, 1], 6, [3, 0]] \\ C'_1 &= [3, [1, 1], 2, [3, 1], 15, [1, 0], 14, [7, 2], 5, [2, 1]] \\ C'_2 &= [0, [8, 0], 8, [8, 2]] \\ C'_3 &= [10, [6, -1], 4, [6, 1]] \\ C'_4 &= [1, [5, 0], 12, [5, 2]] \\ 2\text{-factor type } [7, 3, 3, 2, 2]: \text{ three starters} \end{split}$$

$$\begin{split} C_0 &= [15, [7, 1], 8, [3, -1], 11, [1, 0], 10, [2, 1], 12, [4, 1], 0, [4, 2], 4, [5, 1]] \\ C_1 &= [1, [4, -1], 13, [1, 2], 14, [3, 0]] \\ C_2 &= [2, [4, 0], 6, [3, 1], 9, [7, 2]] \\ C_3 &= [5, [2, 0], 3, [2, 2]] \\ C_4 &= [16, [16, -1], 7, [16, 1]] \end{split}$$

- $$\begin{split} C_0' &= [1, [16, 2], 16, [16, 0], 5, [6, -1], 15, [3, 2], 12, [5, 0], 7, [5, -1], 2, [1, -1]] \\ C_1' &= [6, [5, 2], 11, [2, -1], 13, [7, 0]] \\ C_2' &= [3, [6, 1], 9, [1, 1], 10, [7, -1]] \\ C_3' &= [0, [8, 0], 8, [8, 2]] \\ C_4' &= [4, [6, 0], 14, [6, 2]] \end{split}$$
- 2-factor type [6, 4, 3, 2, 2]: three starters

$$\begin{split} C_0 &= [13, [3, 2], 10, [1, 0], 9, [2, 2], 11, [1, -1], 12, [5, -1], 1, [4, 0]] \\ C_1 &= [15, [4, 2], 3, [5, 0], 8, [2, -1], 6, [7, -1]] \\ C_2 &= [16, [16, 0], 2, [3, 1], 5, [16, 2]] \\ C_3 &= [7, [7, 0], 0, [7, 2]] \\ C_4 &= [14, [6, -1], 4, [6, 1]] \\ C_0' &= [9, [4, 1], 13, [1, 1], 14, [4, -1], 10, [5, 2], 15, [3, 0], 2, [7, 1]] \\ C_1' &= [5, [2, 1], 3, [3, -1], 6, [2, 0], 4, [1, 2]] \\ C_2' &= [16, [16, -1], 7, [5, 1], 12, [16, 1]] \\ C_3' &= [0, [8, 0], 8, [8, 2]] \\ C_4' &= [11, [6, 0], 1, [6, 2]] \end{split}$$

- 2-factor type [5, 5, 3, 2, 2]: three starters
 - $$\begin{split} C_0 &= [3, [4, -1], 15, [4, 2], 11, [16, -1], 16, [16, 1], 2, [1, 0]] \\ C_1 &= [6, [2, -1], 8, [7, 1], 1, [7, 0], 10, [3, 1], 13, [7, 2]] \\ C_2 &= [12, [5, 1], 7, [2, 0], 9, [3, 2]] \\ C_3 &= [5, [5, 0], 0, [5, 2]] \\ C_4 &= [4, [6, -1], 14, [6, 1]] \\ C'_0 &= [13, [2, 1], 11, [6, 0], 5, [4, 1], 9, [1, 2], 10, [3, -1]] \\ C'_1 &= [14, [4, 0], 2, [6, 2], 12, [7, -1], 3, [16, 0], 16, [16, 2]] \\ C'_2 &= [15, [5, -1], 4, [3, 0], 1, [2, 2]] \\ C'_3 &= [0, [8, 0], 8, [8, 2]] \\ C'_4 &= [6, [1, -1], 7, [1, 1]] \end{split}$$
- 2-factor type [5, 4, 4, 2, 2]: three starters

$$\begin{split} C_0 &= [2, [1, 0], 1, [7, 2], 10, [1, 1], 9, [3, 1], 6, [4, 1]] \\ C_1 &= [5, [6, 1], 15, [2, 1], 13, [6, 0], 7, [2, 2]] \\ C_2 &= [14, [6, 2], 8, [5, 0], 3, [16, 2], 16, [16, 0]] \\ C_3 &= [12, [4, 0], 0, [4, 2]] \\ C_4 &= [11, [7, -1], 4, [7, 1]] \\ C'_0 &= [15, [6, -1], 5, [2, 0], 7, [1, 2], 6, [5, -1], 11, [4, -1]] \\ C'_1 &= [9, [3, -1], 12, [7, 0], 3, [5, 2], 14, [5, 1]] \\ C'_2 &= [2, [1, -1], 1, [16, 1], 16, [16, -1], 4, [2, -1]] \\ C'_3 &= [0, [8, 0], 8, [8, 2]] \\ C'_4 &= [10, [3, 0], 13, [3, 2]] \end{split}$$

• 2-factor type [6, 3, 3, 3, 2]: three starters

$$\begin{split} C_0 &= [11, [6, 2], 5, [7, -1], 12, [3, 0], 15, [1, 2], 14, [5, 1], 9, [2, 0]] \\ C_1 &= [7, [3, -1], 10, [4, 1], 6, [1, -1]] \\ C_2 &= [4, [16, 2], 16, [16, 0], 13, [7, 1]] \\ C_3 &= [3, [3, 2], 0, [2, -1], 2, [1, 0]] \\ C_4 &= [8, [7, 0], 1, [7, 2]] \\ \hline \\ C_0' &= [2, [3, 1], 15, [6, -1], 5, [1, 1], 4, [6, 0], 10, [4, -1], 6, [4, 2]] \\ C_1' &= [14, [16, 1], 16, [16, -1], 3, [5, -1]] \\ C_2' &= [9, [2, 1], 11, [2, 2], 13, [4, 0]] \\ C_3' &= [12, [5, 0], 1, [6, 1], 7, [5, 2]] \\ C_4' &= [0, [8, 0], 8, [8, 2]] \\ \hline \\ \bullet \ 2\text{-factor type } [5, 4, 3, 3, 2]: \text{ three starters} \\ C_0 &= [9, [1, 0], 8, [5, 1], 13, [2, 1], 15, [5, -1], 10, [1, 2]] \\ C_1 &= [0, [6, -1], 6, [5, 0], 11, [3, 1], 14, [2, 2]] \\ C_2 &= [2, [1, -1], 3, [7, 0], 12, [6, 2]] \\ C_3 &= [16, [16, 0], 4, [3, 2], 7, [16, -1]] \\ C_4 &= [5, [4, 0], 1, [4, 2]] \\ \hline \\ C_0' &= [13, [4, 1], 1, [5, 2], 12, [2, 0], 10, [7, 2], 3, [6, 0]] \\ C_1' &= [16, [16, 1], 7, [2, -1], 9, [3, 0], 6, [16, 2]] \\ C_2' &= [11, [4, -1], 15, [3, -1], 2, [7, 1]] \\ C_3' &= [14, [7, -1], 5, [1, 1], 4, [6, 1]] \\ C_4' &= [0, [8, 0], 8, [8, 2]] \\ \bullet \ 2\text{-factor type } [4, 4, 4, 3, 2]: \text{ three starters} \end{aligned}$$

$$\begin{split} C_0 &= [5, [5, -1], 0, [2, 2], 2, [7, 1], 11, [6, 0]] \\ C_1 &= [9, [5, 2], 14, [1, 0], 13, [3, -1], 10, [1, -1]] \\ C_2 &= [15, [7, 2], 8, [7, 0], 1, [3, 2], 4, [5, 0]] \\ C_3 &= [3, [7, -1], 12, [6, 2], 6, [3, 0]] \\ C_4 &= [16, [16, 0], 7, [16, 2]] \end{split} \quad \begin{aligned} C_0' &= [13, [1, 1], 12, [5, 1], 7, [4, 1], 3, [6, 1]] \\ C_1' &= [11, [1, 2], 10, [4, 0], 6, [3, 1], 9, [2, 1]] \\ C_2' &= [5, [4, -1], 1, [2, -1], 15, [16, 1], 16, [16, -1]] \\ C_3' &= [4, [2, 0], 2, [4, 2], 14, [6, -1]] \\ C_4' &= [0, [8, 0], 8, [8, 2]] \end{split}$$

• 2-factor type [11, 2, 2, 2]: three starters $C_0 = [8, [1, 0], 7, [6, 2], 13, [6, 0], 3, [4, -1], 15, [5, 2], 10, [2, 0], 12, [5, 1], 1, [1, 2], 0, [2, 1], 14, [3, 0], 11, [3, 2]]$ $C_1 = [6, [16, 0], 16, [16, 2]]$ $C_2 = [4, [1, -1], 5, [1, 1]]$ $C_3 = [2, [7, 0], 9, [7, 2]]$ $C'_0 = [10, [5, 0], 15, [3, -1], 12, [7, 1], 3, [3, 1], 6, [2, 2], 4, [7, -1], 11, [2, -1], 13, [5, -1], 2, [4, 1], 14, [16, -1], 16, [16, 1]]$ $C'_1 = [0, [8, 0], 8, [8, 2]]$ $C'_2 = [9, [4, 0], 5, [4, 2]]$ $C'_3 = [7, [6, -1], 1, [6, 1]]$ • 2-factor type [10, 3, 2, 2]: three starters

 $C_0 = [10, [1, 1], 11, [5, 2], 0, [4, 0], 4, [5, 1], 9, [3, 2], 12, [6, 0], 2, [3, 1], 5, [1, 2], 6, [1, -1], 7, [3, 0]]$ $C_1 = [8, [5, 0], 3, [5, -1], 14, [6, 2]]$ $C_2 = [15, [2, -1], 1, [2, 1]]$ $C_3 = [13, [16, -1], 16, [16, 1]]$ $C_0' = [13, [7, -1], 4, [1, 0], 3, [7, 2], 12, [6, 1], 6, [4, 1], 2, [3, -1], 5, [7, 1], 14, [7, 0], 7, [2, 2], 9, [4, -1]]$ $C'_1 = [11, [4, 2], 15, [2, 0], 1, [6, -1]]$ $C'_{2} = [0, [8, 0], 8, [8, 2]]$ $C'_{3} = [10, [16, 0], 16, [16, 2]]$ • 2-factor type [9, 4, 2, 2]: three starters $C_0 = [10, [4, 1], 14, [1, 2], 13, [7, 1], 6, [7, 0], 15, [6, -1], 9, [6, 2], 3, [4, 0], 7, [7, 2], 0, [6, 0]]$ $C_1 = [4, [1, 0], 5, [3, 2], 2, [16, -1], 16, [16, 1]]$ $C_2 = [8, [3, -1], 11, [3, 1]]$ $C_3 = [1, [5, -1], 12, [5, 1]]$ $C_0' = [15, [6, 1], 9, [4, -1], 13, [1, 1], 12, [2, 0], 14, [7, -1], 7, [4, 2], 11, [5, 0], 6, [2, -1], 4, [5, 2]]$ $C'_1 = [5, [3, 0], 2, [1, -1], 1, [2, 1], 3, [2, 2]]$ $C'_{2} = [0, [8, 0], 8, [8, 2]]$ $C'_{3} = [10, [16, 0], 16, [16, 2]]$ • 2-factor type [8, 5, 2, 2]: three starters $C_0 = [4, [1, 1], 3, [3, 0], 0, [5, -1], 11, [16, 2], 16, [16, 1], 8, [2, 0], 6, [7, 1], 13, [7, 2]]$ $C_1 = [10, [5, 1], 15, [1, 2], 14, [7, -1], 5, [4, 1], 1, [7, 0]]$ $C_2 = [2, [5, 0], 7, [5, 2]]$ $C_3 = [12, [3, -1], 9, [3, 1]]$

- $$\begin{split} C_0' &= [11, [6, -1], 1, [2, 1], 15, [4, -1], 3, [16, -1], 16, [16, 0], 2, [4, 2], 14, [1, 0], 13, [2, 2]] \\ C_1' &= [7, [3, 2], 10, [6, 1], 4, [1, -1], 5, [4, 0], 9, [2, -1]] \\ C_2' &= [0, [8, 0], 8, [8, 2]] \\ C_3' &= [12, [6, 0], 6, [6, 2]] \end{split}$$
- 2-factor type [7, 6, 2, 2]: three starters

$$\begin{split} C_0 &= [6, [5, 1], 1, [16, 0], 16, [16, 2], 11, [4, 0], 15, [6, 1], 5, [2, 2], 3, [3, -1]] \\ C_1 &= [12, [2, 1], 10, [4, -1], 14, [5, 0], 9, [7, 2], 0, [2, 0], 2, [6, 2]] \\ C_2 &= [4, [7, -1], 13, [7, 1]] \\ C_3 &= [8, [1, 0], 7, [1, 2]] \\ C'_0 &= [10, [3, 0], 7, [3, 2], 4, [5, -1], 9, [3, 1], 6, [7, 0], 15, [1, -1], 14, [4, 2]] \\ C'_1 &= [13, [4, 1], 1, [5, 2], 12, [1, 1], 11, [6, 0], 5, [2, -1], 3, [6, -1]] \\ C'_2 &= [0, [8, 0], 8, [8, 2]] \\ C'_3 &= [2, [16, -1], 16, [16, 1]] \end{split}$$

• 2-factor type [9, 3, 3, 2]: three starters

 $C_0 = [13, [5, 2], 8, [2, -1], 10, [4, -1], 14, [7, 1], 7, [2, 1], 5, [6, 0], 15, [1, 1], 0, [16, 1], 16, [16, -1]]$ $C_1 = [2, [6, -1], 12, [1, 2], 11, [7, 0]]$ $C_2 = [3, [3, -1], 6, [5, 0], 1, [2, 2]]$ $C_3 = [9, [5, -1], 4, [5, 1]]$ $C_0' = [12, [6, 1], 6, [7, -1], 13, [4, 0], 9, [7, 2], 2, [3, 0], 5, [1, -1], 4, [16, 2], 16, [16, 0], 15, [3, 2]]$ $C'_1 = [14, [3, 1], 11, [1, 0], 10, [4, 2]]$ $C'_{2} = [7, [4, 1], 3, [2, 0], 1, [6, 2]]$ $C'_3 = [0, [8, 0], 8, [8, 2]]$ • 2-factor type [8, 4, 3, 2]: three starters $C_0 = [1, [7, 0], 10, [5, 2], 5, [1, 0], 4, [6, -1], 14, [3, 2], 11, [2, -1], 13, [5, -1], 2, [1, 1]]$ $C_1 = [6, [7, -1], 15, [1, 2], 0, [7, 1], 9, [3, 0]]$ $C_2 = [8, [16, 2], 16, [16, 1], 12, [4, 0]]$ $C_3 = [3, [4, -1], 7, [4, 1]]$ $C_0' = [5, [7, 2], 12, [6, 0], 6, [4, 2], 2, [2, 0], 4, [1, -1], 3, [6, 2], 13, [16, 0], 16, [16, -1]]$ $C'_1 = [1, [2, 1], 15, [5, 1], 10, [3, 1], 7, [6, 1]]$ $C'_{2} = [14, [3, -1], 11, [2, 2], 9, [5, 0]]$ $C'_{3} = [0, [8, 0], 8, [8, 2]]$

• 2-factor type [7, 5, 3, 2]: three starters

$$\begin{split} C_0 &= [10, [3, 1], 7, [1, 1], 6, [7, -1], 13, [4, -1], 1, [1, 0], 0, [1, 2], 15, [5, 1]] \\ C_1 &= [3, [6, 1], 9, [5, 2], 4, [4, 0], 8, [6, 2], 14, [5, 0]] \\ C_2 &= [11, [1, -1], 12, [6, -1], 2, [7, 1]] \\ C_3 &= [16, [16, 0], 5, [16, 2]] \\ C'_0 &= [14, [3, 2], 1, [16, -1], 16, [16, 1], 3, [3, 0], 6, [4, 1], 10, [2, 2], 12, [2, 0]] \\ C'_1 &= [9, [5, -1], 4, [7, 2], 13, [2, -1], 15, [3, -1], 2, [7, 0]] \\ C'_2 &= [11, [4, 2], 7, [2, 1], 5, [6, 0]] \\ C'_3 &= [0, [8, 0], 8, [8, 2]] \end{split}$$

• 2-factor type [6, 6, 3, 2]: three starters

$$\begin{split} C_0 &= [2, [7, -1], 11, [6, 1], 1, [16, 2], 16, [16, 0], 0, [1, -1], 15, [3, 1]] \\ C_1 &= [5, [3, -1], 8, [6, -1], 14, [5, 0], 9, [6, 2], 3, [4, 0], 7, [2, 2]] \\ C_2 &= [10, [2, 0], 12, [1, 1], 13, [3, 2]] \\ C_3 &= [4, [2, -1], 6, [2, 1]] \\ C_0' &= [11, [5, 1], 6, [1, 0], 7, [4, 2], 3, [6, 0], 13, [5, 2], 2, [7, 1]] \\ C_1' &= [10, [7, 0], 1, [16, 1], 16, [16, -1], 4, [5, -1], 15, [1, 2], 14, [4, 1]] \\ C_2' &= [12, [7, 2], 5, [4, -1], 9, [3, 0]] \\ C_3' &= [0, [8, 0], 8, [8, 2]] \end{split}$$

• 2-factor type [7, 4, 4, 2]: three starters

$$\begin{split} C_0 &= [6, [1, 0], 7, [4, 2], 3, [1, -1], 4, [6, 0], 14, [1, 2], 13, [4, 1], 9, [3, 1]] \\ C_1 &= [11, [1, 1], 12, [3, 2], 15, [2, 0], 1, [6, -1]] \\ C_2 &= [8, [3, -1], 5, [5, 1], 10, [16, 1], 16, [16, -1]] \\ C_3 &= [2, [2, -1], 0, [2, 1]] \\ C'_0 &= [5, [7, -1], 12, [5, 0], 7, [2, 2], 9, [5, -1], 14, [4, -1], 2, [7, 0], 11, [6, 2]] \\ C'_1 &= [16, [16, 2], 1, [3, 0], 4, [5, 2], 15, [16, 0]] \\ C'_2 &= [3, [6, 1], 13, [7, 1], 6, [4, 0], 10, [7, 2]] \\ C'_3 &= [0, [8, 0], 8, [8, 2]] \end{split}$$

• 2-factor type [6, 5, 4, 2]: three starters

$$\begin{split} C_0 &= [7, [4, -1], 11, [5, -1], 0, [1, 0], 15, [6, 1], 5, [1, 1], 6, [1, 2]] \\ C_1 &= [14, [1, -1], 13, [16, 2], 16, [16, 1], 10, [7, 1], 3, [5, 0]] \\ C_2 &= [1, [5, 2], 12, [4, 1], 8, [4, 0], 4, [3, -1]] \\ C_3 &= [9, [7, 0], 2, [7, 2]] \\ C'_0 &= [5, [4, 2], 1, [16, 0], 16, [16, -1], 12, [6, -1], 6, [3, 2], 3, [2, 0]] \\ C'_1 &= [2, [7, -1], 9, [2, -1], 11, [3, 0], 14, [6, 2], 4, [2, 1]] \\ C'_2 &= [13, [2, 2], 15, [5, 1], 10, [3, 1], 7, [6, 0]] \\ C'_3 &= [0, [8, 0], 8, [8, 2]] \end{split}$$

• 2-factor type [5, 5, 5, 2]: three starters

$$\begin{split} C_0 &= [13, [7, 1], 4, [5, -1], 15, [3, 1], 2, [4, -1], 14, [1, -1]] \\ C_1 &= [3, [16, 0], 16, [16, 2], 7, [2, 1], 5, [4, 0], 1, [2, 2]] \\ C_2 &= [8, [4, 2], 12, [1, 0], 11, [1, 2], 10, [4, 1], 6, [2, 0]] \\ C_3 &= [0, [7, 0], 9, [7, 2]] \\ C'_0 &= [13, [5, 1], 2, [3, 2], 15, [6, 1], 9, [6, 0], 3, [6, -1]] \\ C'_1 &= [6, [16, 1], 16, [16, -1], 4, [1, 1], 5, [6, 2], 11, [5, 0]] \\ C'_2 &= [7, [5, 2], 12, [2, -1], 14, [3, 0], 1, [7, -1], 10, [3, -1]] \\ C'_3 &= [0, [8, 0], 8, [8, 2]] \end{split}$$

• 2-factor type [13, 2, 2]: three starters

$$\begin{split} C_0 &= [7, [2, -1], 9, [4, 1], 5, [1, -1], 6, [6, -1], 12, [6, 0], 2, [3, 2], 15, [7, 0], 8, [7, -1], 1, [16, 1], 16, [16, -1], \\ &\quad 14, [4, -1], 10, [3, 1], 13, [6, 2]] \\ C_1 &= [3, [1, 0], 4, [1, 2]] \\ C_2 &= [0, [5, -1], 11, [5, 1]] \\ C'_0 &= [1, [3, -1], 4, [16, 0], 16, [16, 2], 5, [2, 1], 3, [7, 1], 12, [3, 0], 15, [5, 2], 10, [4, 0], 6, [4, 2], 2, [5, 0], 13, \\ &\quad [1, 1], 14, [7, 2], 7, [6, 1]] \\ C'_1 &= [0, [8, 0], 8, [8, 2]] \\ C'_2 &= [9, [2, 0], 11, [2, 2]] \end{split}$$

• 2-factor type [12, 3, 2]: three starters

$$\begin{split} &C_0 = [4, [7, -1], 13, [1, 1], 14, [2, 1], 0, [4, 2], 12, [4, 1], 8, [3, 0], 11, [6, -1], 5, [4, -1], 1, [2, 2], 3, [3, -1], 6, \\ & [7, 0], 15, [5, 1]] \\ &C_1 = [7, [16, -1], 16, [16, 1], 2, [5, -1]] \\ &C_2 = [10, [1, 0], 9, [1, 2]] \\ &C_0' = [4, [2, 0], 2, [5, 2], 13, [7, 1], 6, [5, 0], 1, [7, 2], 10, [16, 0], 16, [16, 2], 5, [6, 1], 15, [6, 0], 9, [6, 2], 3, [4, 0], \\ & 7, [3, 2]] \\ &C_1' = [12, [1, -1], 11, [3, 1], 14, [2, -1]] \\ &C_2' = [0, [8, 0], 8, [8, 2]] \\ &2\text{-factor type } [11, 4, 2]\text{: three starters} \\ &C_0 = [4, [3, 1], 1, [7, -1], 8, [7, 0], 15, [1, 2], 14, [2, 0], 12, [7, 2], 5, [5, -1], 10, [1, 0], 11, [2, 2], 9, [7, 1], 2, \\ & [2, -1]] \\ &C_1 = [6, [6, -1], 0, [3, -1], 13, [16, 1], 16, [16, -1]] \\ &C_2 = [7, [4, 0], 3, [4, 2]] \\ &C_0' = [14, [1, -1], 13, [3, 2], 10, [5, 1], 5, [1, 1], 4, [2, 1], 2, [4, -1], 6, [3, 0], 3, [6, 2], 9, [6, 0], 15, [16, 2], 16, \\ & [16, 0]] \\ &C_1' = [11, [4, 1], 7, [5, 0], 12, [5, 2], 1, [6, 1]] \\ &C_2' = [0, [8, 0], 8, [8, 2]] \end{aligned}$$

• 2-factor type [10, 5, 2]: three starters

$$\begin{split} C_0 &= [14, [3, 1], 11, [4, 1], 7, [16, 0], 16, [16, -1], 3, [5, 2], 8, [2, -1], 10, [6, 1], 4, [5, 0], 15, [6, 2], 5, [7, 1]] \\ C_1 &= [6, [7, 2], 13, [5, -1], 2, [7, -1], 9, [3, 0], 12, [6, -1]] \\ C_2 &= [0, [1, 0], 1, [1, 2]] \\ C_0' &= [12, [7, 0], 3, [4, 2], 7, [6, 0], 1, [3, 2], 4, [2, 0], 6, [4, -1], 2, [3, -1], 15, [2, 2], 13, [1, -1], 14, [2, 1]] \\ C_1' &= [9, [16, 1], 16, [16, 2], 11, [1, 1], 10, [5, 1], 5, [4, 0]] \\ C_2' &= [0, [8, 0], 8, [8, 2]] \end{split}$$

• 2-factor type [9, 6, 2]: three starters

$$\begin{split} C_0 &= [12, [1, 2], 13, [6, 0], 7, [5, 2], 2, [2, 1], 0, [4, -1], 4, [2, -1], 6, [16, 0], 16, [16, -1], 3, [7, -1]] \\ C_1 &= [10, [5, 0], 15, [1, -1], 14, [6, 2], 8, [3, 0], 11, [6, 1], 1, [7, 2]] \\ C_2 &= [5, [4, 0], 9, [4, 2]] \\ C_0' &= [15, [2, 0], 13, [4, 1], 1, [1, 1], 2, [16, 1], 16, [16, 2], 11, [5, 1], 6, [3, 1], 3, [7, 0], 12, [3, 2]] \\ C_1' &= [4, [5, -1], 9, [1, 0], 10, [3, -1], 7, [2, 2], 5, [7, 1], 14, [6, -1]] \\ C_2' &= [0, [8, 0], 8, [8, 2]] \end{split}$$

• 2-factor type [8,7,2]: three starters

$$\begin{split} C_0 &= [3, [4, -1], 7, [3, 1], 4, [7, 1], 13, [4, 0], 1, [2, -1], 15, [3, 2], 2, [2, 1], 0, [3, -1]] \\ C_1 &= [8, [3, 0], 5, [5, 1], 10, [4, 2], 14, [5, 0], 9, [16, 1], 16, [16, -1], 6, [2, 2]] \\ C_2 &= [12, [1, -1], 11, [1, 1]] \end{split}$$

$$\begin{split} C_0' &= [5, [7, -1], 14, [5, 2], 9, [2, 0], 11, [6, 2], 1, [16, 0], 16, [16, 2], 10, [6, 1], 4, [1, 0]] \\ C_1' &= [2, [5, -1], 13, [6, -1], 7, [1, 2], 6, [7, 0], 15, [4, 1], 3, [7, 2], 12, [6, 0]] \\ C_2' &= [0, [8, 0], 8, [8, 2]] \end{split}$$

J Computational results for n = 18

- 2-factor type [4, 2, 2, 2, 2, 2, 2, 2]: one starter
 - $$\begin{split} C_0 &= [2, [4, 0], 15, [5, 1], 3, [4, 1], 7, [5, 0]] \\ C_1 &= [1, [8, 0], 9, [8, 1]] \\ C_2 &= [10, [7, 0], 0, [7, 1]] \\ C_3 &= [11, [3, 0], 8, [3, 1]] \\ C_4 &= [12, [6, 0], 6, [6, 1]] \\ C_5 &= [5, [1, 0], 4, [1, 1]] \\ C_6 &= [17, [17, 0], 13, [17, 1]] \\ C_7 &= [16, [2, 0], 14, [2, 1]] \end{split}$$
- 2-factor type [3, 3, 2, 2, 2, 2, 2, 2]: two starters
 - $C_0 = [17, [17, 1], 11, [4, 1], 7, [17, -1]]$ $C'_0 = [8, [7, 0], 15, [5, 2], 3, [5, -1]]$ $C'_1 = [9, [2, 2], 11, [2, 0], 13, [4, -1]]$ $C_1 = [13, [5, 1], 8, [5, 0], 3, [7, 2]]$ $C_2 = [5, [1, 0], 4, [1, 2]]$ $C'_{2} = [2, [3, 0], 5, [3, 2]]$ $C'_3 = [1, [3, -1], 4, [3, 1]]$ $C_3 = [15, [1, -1], 16, [1, 1]]$ $C_4 = [2, [7, -1], 9, [7, 1]]$ $C'_4 = [7, [8, -1], 16, [8, 1]]$ $C_5 = [0, [6, 0], 6, [6, 2]]$ $C'_{5} = [10, [2, -1], 12, [2, 1]]$ $C_6 = [1, [6, -1], 12, [6, 1]]$ $C'_6 = [0, [17, 0], 17, [17, 2]]$ $C_7 = [10, [4, 0], 14, [4, 2]]$ $C'_7 = [14, [8, 0], 6, [8, 2]]$
- $C_{7} = [10, [4, 0], 14, [4, 2]]$ $C_{7}' = [14, [8, -2]]$ $C_{7}' = [14, [8, -2]]$ $C_{0} = [10, [17, -1], 17, [17, 1], 1, [6, 0], 7, [1, -1]], 6, [8, 2], 14, [4, 1]]$ $C_{1} = [2, [1, 0], 3, [1, 2]]$ $C_{2} = [5, [3, 0], 8, [3, 2]]$ $C_{3} = [15, [2, -1], 13, [2, 1]]$ $C_{4} = [9, [8, -1], 0, [8, 1]]$ $C_{5} = [12, [4, 0], 16, [4, 2]]$ $C_{6} = [11, [7, 0], 4, [7, 2]]$ $C_{0}' = [0, [1, 1], 16, [6, -1], 5, [4, -1], 9, [6, 1], 3, [8, 0], 11, [6, 2]]$ $C_{1}' = [10, [3, -1], 13, [3, 1]]$ $C_{2}' = [4, [17, 0], 17, [17, 2]]$ $C_{4}' = [1, [5, 0], 6, [5, 2]]$

 $C'_{5} = [14, [2, 0], 12, [2, 2]]$ $C'_{6} = [7, [5, -1], 2, [5, 1]]$ • 2-factor type [5, 3, 2, 2, 2, 2, 2]: one starter

 $C_0 = [15, [6, 1], 4, [17, 1], 17, [17, 0], 9, [7, 0], 2, [4, 1]]$ $C_1 = [14, [7, 1], 7, [4, 0], 3, [6, 0]]$ $C_2 = [6, [2, 0], 8, [2, 1]]$ $C_3 = [13, [3, 0], 16, [3, 1]]$ $C_4 = [5, [5, 0], 0, [5, 1]]$ $C_5 = [12, [1, 0], 11, [1, 1]]$ $C_6 = [1, [8, 0], 10, [8, 1]]$

• 2-factor type [4, 4, 2, 2, 2, 2, 2]: one starter

 $C_0 = [14, [8, 1], 5, [5, 1], 0, [6, 1], 11, [3, 1]]$ $C_1 = [16, [8, 0], 7, [6, 0], 1, [3, 0], 4, [5, 0]]$ $C_2 = [6, [17, 0], 17, [17, 1]]$ $C_3 = [13, [4, 0], 9, [4, 1]]$ $C_4 = [8, [7, 0], 15, [7, 1]]$ $C_5 = [2, [1, 0], 3, [1, 1]]$ $C_6 = [12, [2, 0], 10, [2, 1]]$

- 2-factor type [4, 3, 3, 2, 2, 2, 2]: two starters
 - $C_0 = [1, [17, 2], 17, [17, 0], 4, [8, -1], 13, [5, 1]]$ $C_1 = [12, [2, 0], 10, [7, 2], 3, [8, 1]]$ $C_2 = [6, [1, 0], 5, [4, 2], 9, [3, 1]]$ $C_3 = [2, [6, 0], 8, [6, 2]]$ $C_4 = [0, [3, 0], 14, [3, 2]]$ $C_5 = [16, [1, -1], 15, [1, 1]]$ $C_6 = [11, [4, -1], 7, [4, 1]]$
- 2-factor type [3, 3, 3, 3, 2, 2, 2]: one starter

 $C_0 = [13, [7, 1], 3, [1, 0], 2, [6, 0]]$ $C_1 = [0, [1, 1], 16, [2, 1], 14, [3, 1]]$ $C_2 = [4, [6, 1], 10, [8, 0], 1, [3, 0]]$ $C_3 = [6, [8, 1], 15, [7, 0], 8, [2, 0]]$ $C_4 = [9, [4, 0], 5, [4, 1]]$ $C_5 = [12, [5, 0], 7, [5, 1]]$ $C_6 = [11, [17, 0], 17, [17, 1]]$

• 2-factor type [8, 2, 2, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [7, [4, 1], 11, [4, 0], 15, [3, 0], 12, [2, 1], 14, [2, 0], 16, [3, 1], 2, [6, 1], 13, [6, 0]] \\ C_1 &= [17, [17, 0], 1, [17, 1]] \\ C_2 &= [6, [1, 0], 5, [1, 1]] \\ C_3 &= [4, [5, 0], 9, [5, 1]] \\ C_4 &= [3, [7, 0], 10, [7, 1]] \\ C_5 &= [0, [8, 0], 8, [8, 1]] \end{split}$$

$$\begin{split} C_0' &= [8, [3, -1], 5, [1, 2], 4, [8, 0], 13, [5, -1]] \\ C_1' &= [11, [8, 2], 3, [2, 1], 1, [7, 0]] \\ C_2' &= [14, [4, 0], 10, [2, 2], 12, [2, -1]] \\ C_3' &= [9, [7, -1], 16, [7, 1]] \\ C_4' &= [2, [5, 0], 7, [5, 2]] \\ C_5' &= [15, [17, -1], 17, [17, 1]] \\ C_6' &= [0, [6, -1], 6, [6, 1]] \end{split}$$

[o] [o]

• 2-factor type [7, 3, 2, 2, 2, 2]: two starters

$$\begin{split} &C_0 = [10, [8, 1], 1, [4, -1], 14, [17, 1], 17, [17, 2], 15, [8, 0], 6, [5, 1], 11, [1, -1]] \\ &C_1 = [12, [1, 1], 13, [7, 1], 3, [8, -1]] \\ &C_2 = [7, [3, -1], 4, [3, 1]] \\ &C_3 = [5, [5, 0], 0, [5, 2]] \\ &C_4 = [16, [7, 0], 9, [7, 2]] \\ &C_5 = [2, [6, 0], 8, [6, 2]] \\ &C'_0 = [11, [1, 0], 12, [3, 2], 9, [5, -1], 14, [17, 0], 17, [17, -1], 13, [8, 2], 4, [7, -1]] \\ &C'_1 = [1, [1, 2], 2, [4, 1], 15, [3, 0]] \\ &C'_2 = [6, [6, -1], 0, [6, 1]] \\ &C'_3 = [8, [2, 0], 10, [2, 2]] \\ &C'_4 = [7, [2, -1], 5, [2, 1]] \\ &C'_5 = [3, [4, 0], 16, [4, 2]] \\ e 2 \text{-factor type } [6, 4, 2, 2, 2, 2]: \text{ two starters} \\ &C_0 = [0, [5, 0], 12, [3, 2], 15, [6, 1], 9, [5, 1], 4, [2, 0], 6, [6, 2]] \\ &C_1 = [5, [5, 2], 10, [6, 0], 16, [5, -1], 11, [6, -1]] \\ &C_2 = [8, [17, 0], 17, [17, 2]] \\ &C_3 = [1, (1, 0], 2, [1, 2]] \\ &C_4 = [3, [4, -1], 7, [4, 1]] \\ &C_5 = [1, (1, 0), 2, [1, 2]] \\ &C'_0 = [9, [8, -1], 1, [2, 2], 16, [8, 0], 7, [17, 1], 17, [17, -1], 12, [3, -1]] \\ &C'_1 = [5, [8, 1], 14, [3, 0], 0, [8, 2], 8, [3, 1]] \\ &C'_2 = [4, [2, -1], 2, [2, 1]] \\ &C'_3 = [13, [7, 0], 6, [7, 2]] \\ &C'_4 = [11, [4, 0], 15, [4, 2]] \\ &C'_5 = [10, [7, -1], 3, [7, 1]] \\ e 2 \text{-factor type } [5, 5, 2, 2, 2, 2]: \text{ two starters} \\ &C_0 = [6, [4, -1], 10, [3, 1], 7, [2, 0], 9, [4, 2], 5, [1, -1]] \\ &C_1 = [11, [7, 0], 4, [5, 2], 16, [4, 0], 3, [1, 2], 2, [8, 1]] \\ &C_2 = [13, [5, -1], 1, [5, 1]] \\ &C_3 = [12, [2, -1], 14, [2, 1]] \\ &C_4 = [15, [7, -1], 8, [7, 1]] \\ &C_5 = [17, [17, -1], 0, [17, 1]] \\ &C'_0 = [8, [7, 2], 1, [3, -1], 4, [8, -1], 12, [5, 0], 7, [1, 1]] \\ &C'_1 = [14, [8, 2], 6, [8, 0], 15, [2, 2], 0, [4, 1], 13, [1, 0]] \\ &C'_3 = [11, [6, 0], 5, [6, 2]] \\ &C'_4 = [10, [17, 0], 7, [1, 7, 2]] \\ &C'_5 = [3, [6, -1], 9, [6, 1]] \\ \end{array}$$

• 2-factor type [6, 3, 3, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [6, [4, 0], 10, [7, 1], 3, [2, 1], 5, [5, 1], 0, [2, 0], 15, [8, 1]] \\ C_1 &= [12, [4, 1], 16, [5, 0], 4, [8, 0]] \\ C_2 &= [9, [7, 0], 2, [17, 1], 17, [17, 0]] \\ C_3 &= [13, [1, 0], 14, [1, 1]] \\ C_4 &= [7, [6, 0], 1, [6, 1]] \\ C_5 &= [11, [3, 0], 8, [3, 1]] \end{split}$$

• 2-factor type [5, 4, 3, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [11, [1, 0], 12, [7, 1], 5, [5, 0], 10, [3, 1], 7, [4, 1]] \\ C_1 &= [13, [3, 0], 16, [2, 1], 1, [1, 1], 0, [4, 0]] \\ C_2 &= [14, [5, 1], 2, [2, 0], 4, [7, 0]] \\ C_3 &= [3, [6, 0], 9, [6, 1]] \\ C_4 &= [15, [8, 0], 6, [8, 1]] \\ C_5 &= [8, [17, 0], 17, [17, 1]] \end{split}$$

• 2-factor type [4, 4, 4, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [6, [8, 0], 14, [8, 1], 5, [5, 1], 0, [6, 0]] \\ C_1 &= [4, [5, 0], 16, [2, 0], 1, [2, 1], 3, [1, 1]] \\ C_2 &= [15, [6, 1], 9, [1, 0], 8, [17, 1], 17, [17, 0]] \\ C_3 &= [2, [7, 0], 12, [7, 1]] \\ C_4 &= [7, [4, 0], 11, [4, 1]] \\ C_5 &= [10, [3, 0], 13, [3, 1]] \end{split}$$

• 2-factor type [5, 3, 3, 3, 2, 2]: two starters

$$\begin{split} C_0 &= [10, [1, 0], 11, [17, 2], 17, [17, 0], 2, [1, 2], 3, [7, -1]] \\ C_1 &= [9, [3, 2], 6, [6, 0], 12, [3, -1]] \\ C_2 &= [4, [8, 1], 13, [4, 0], 0, [4, 2]] \\ C_3 &= [5, [4, 1], 1, [3, 0], 15, [7, 2]] \\ C_4 &= [14, [6, -1], 8, [6, 1]] \\ C_5 &= [7, [8, 0], 16, [8, 2]] \\ C'_0 &= [11, [2, 1], 13, [7, 1], 3, [4, -1], 16, [8, -1], 8, [3, 1]] \\ C'_1 &= [1, [6, 2], 12, [5, 0], 0, [1, 1]] \\ C'_2 &= [6, [1, -1], 5, [17, 1], 17, [17, -1]] \\ C'_3 &= [2, [5, 2], 14, [7, 0], 4, [2, -1]] \\ C'_4 &= [10, [5, -1], 15, [5, 1]] \\ C'_5 &= [9, [2, 0], 7, [2, 2]] \end{split}$$

• 2-factor type [4, 4, 3, 3, 2, 2]: two starters

$$\begin{split} C_0 &= [10, [8, -1], 1, [2, 0], 3, [4, 2], 7, [3, -1]] \\ C_1 &= [6, [8, 0], 14, [6, 2], 8, [17, -1], 17, [17, 1]] \\ C_2 &= [4, [4, -1], 0, [8, 2], 9, [5, 0]] \\ C_3 &= [5, [7, -1], 15, [4, 1], 2, [3, 1]] \\ C_4 &= [16, [5, -1], 11, [5, 1]] \\ C_5 &= [13, [1, -1], 12, [1, 1]] \end{split}$$

• 2-factor type [4, 3, 3, 3, 3, 3]: one starter

$$\begin{split} C_0 &= [8, [4, 0], 4, [8, 0], 12, [7, 1], 5, [3, 1]] \\ C_1 &= [11, [2, 0], 9, [5, 1], 14, [3, 0]] \\ C_2 &= [10, [6, 1], 16, [2, 1], 1, [8, 1]] \\ C_3 &= [15, [4, 1], 2, [1, 0], 3, [5, 0]] \\ C_4 &= [6, [1, 1], 7, [7, 0], 0, [6, 0]] \\ C_5 &= [13, [17, 0], 17, [17, 1]] \end{split}$$

- $C'_{0} = [11, [3, 2], 8, [6, -1], 14, [7, 1], 7, [4, 0]]$ $C'_{1} = [12, [6, 0], 6, [2, -1], 4, [6, 1], 10, [2, 2]]$ $C'_{2} = [15, [3, 0], 1, [2, 1], 3, [5, 2]]$ $C'_{3} = [17, [17, 2], 13, [8, 1], 5, [17, 0]]$ $C'_{4} = [2, [7, 0], 9, [7, 2]]$
 - $C'_5 = [16, [1, 0], 0, [1, 2]]$

- 2-factor type [10, 2, 2, 2, 2]: two starters $C_0 = [7, [2, -1], 5, [3, 1], 8, [6, 1], 14, [4, 2], 10, [4, 0], 6, [6, 2], 0, [17, 0], 17, [17, 2], 2, [1, 0], 3, [4, 1]]$ $C_1 = [4, [8, -1], 12, [8, 1]]$ $C_2 = [13, [5, 0], 1, [5, 2]]$ $C_3 = [15, [1, -1], 16, [1, 1]]$ $C_4 = [9, [2, 0], 11, [2, 2]]$ $C'_0 = [1, [6, 0], 12, [1, 2], 13, [4, -1], 0, [17, -1], 17, [17, 1], 8, [8, 0], 16, [6, -1], 10, [3, -1], 7, [2, 1],$ 9, [8, 2]] $C'_1 = [4, [7, -1], 11, [7, 1]]$ $C'_2 = [14, [5, -1], 2, [5, 1]]$ $C'_3 = [3, [3, 0], 6, [3, 2]]$ $C'_4 = [5, [7, 0], 15, [7, 2]]$
- 2-factor type [9, 3, 2, 2, 2]: one starter
 - $$\begin{split} C_0 &= [13, [2, 1], 15, [5, 1], 10, [3, 0], 7, [4, 0], 3, [2, 0], 5, [17, 1], 17, [17, 0], 0, [8, 1], 9, [4, 1]] \\ C_1 &= [11, [5, 0], 16, [8, 0], 8, [3, 1]] \\ C_2 &= [4, [7, 0], 14, [7, 1]] \\ C_3 &= [12, [6, 0], 6, [6, 1]] \\ C_4 &= [2, [1, 0], 1, [1, 1]] \end{split}$$
- 2-factor type [8, 4, 2, 2, 2]: one starter
 - $$\begin{split} C_0 &= [11, [2, 1], 13, [5, 0], 8, [17, 0], 17, [17, 1], 4, [7, 1], 14, [2, 0], 16, [3, 0], 2, [8, 1]] \\ C_1 &= [3, [5, 1], 15, [8, 0], 7, [3, 1], 10, [7, 0]] \\ C_2 &= [12, [6, 0], 6, [6, 1]] \\ C_3 &= [5, [4, 0], 9, [4, 1]] \\ C_4 &= [0, [1, 0], 1, [1, 1]] \end{split}$$

• 2-factor type [7, 5, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [12, [6, 0], 1, [6, 1], 7, [7, 0], 0, [2, 1], 2, [8, 0], 10, [5, 0], 5, [7, 1]] \\ C_1 &= [14, [1, 1], 15, [5, 1], 3, [1, 0], 4, [2, 0], 6, [8, 1]] \\ C_2 &= [8, [3, 0], 11, [3, 1]] \\ C_3 &= [16, [17, 0], 17, [17, 1]] \\ C_4 &= [13, [4, 0], 9, [4, 1]] \end{split}$$

• 2-factor type [6, 6, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [0, [3, 1], 14, [2, 1], 16, [8, 0], 8, [2, 0], 10, [6, 1], 4, [4, 1]]\\ C_1 &= [2, [8, 1], 11, [4, 0], 15, [6, 0], 9, [3, 0], 12, [5, 1], 7, [5, 0]]\\ C_2 &= [1, [17, 0], 17, [17, 1]]\\ C_3 &= [5, [1, 0], 6, [1, 1]]\\ C_4 &= [3, [7, 0], 13, [7, 1]] \end{split}$$

• 2-factor type [8, 3, 3, 2, 2]: two starters

$$\begin{split} C_0 &= [8, [8, 1], 0, [6, 0], 11, [2, -1], 9, [7, 2], 2, [3, -1], 5, [4, 1], 1, [8, 0], 10, [2, 2]] \\ C_1 &= [3, [4, 2], 7, [7, 0], 14, [6, 1]] \\ C_2 &= [13, [17, -1], 17, [17, 1], 6, [7, -1]] \\ C_3 &= [12, [3, 0], 15, [3, 2]] \\ C_4 &= [4, [5, -1], 16, [5, 1]] \\ C_6' &= [17, [17, 0], 6, [4, -1], 2, [2, 1], 4, [8, -1], 13, [8, 2], 5, [4, 0], 9, [6, -1], 3, [17, 2]] \\ C_1' &= [0, [3, 1], 14, [2, 0], 16, [1, 2]] \\ C_2' &= [11, [7, 1], 1, [6, 2], 12, [1, 0]] \\ C_3' &= [15, [5, 0], 10, [5, 2]] \\ C_4' &= [7, [1, -1], 8, [1, 1]] \end{split}$$

• 2-factor type [7, 4, 3, 2, 2]: two starters

$$\begin{split} C_0 &= [8, [4, 0], 12, [5, -1], 7, [3, -1], 4, [5, 2], 9, [1, -1], 10, [6, -1], 16, [8, 1]] \\ C_1 &= [13, [7, -1], 3, [2, -1], 5, [1, 0], 6, [7, 2]] \\ C_2 &= [2, [8, -1], 11, [7, 1], 1, [1, 1]] \\ C_3 &= [15, [2, 0], 0, [2, 2]] \\ C_4 &= [17, [17, -1], 14, [17, 1]] \\ C'_0 &= [8, [1, 2], 9, [3, 1], 6, [17, 0], 17, [17, 2], 5, [5, 1], 0, [5, 0], 12, [4, -1]] \\ C'_1 &= [15, [2, 1], 13, [3, 0], 16, [3, 2], 2, [4, 1]] \\ C'_2 &= [4, [7, 0], 14, [4, 2], 10, [6, 1]] \\ C'_4 &= [3, [8, 0], 11, [8, 2]] \end{split}$$

• 2-factor type [6, 5, 3, 2, 2]: two starters

$$C_{0} = [16, [7, 1], 9, [1, 0], 10, [5, 2], 5, [5, 0], 0, [4, 2], 13, [3, -1]]$$

$$C_{1} = [2, [2, 2], 4, [6, 0], 15, [17, 2], 17, [17, 0], 6, [4, -1]]$$

$$C_{2} = [7, [7, -1], 14, [3, 1], 11, [4, 1]]$$

$$C_{3} = [1, [7, 0], 8, [7, 2]]$$

$$C_{4} = [12, [8, -1], 3, [8, 1]]$$

$$C_{0}' = [4, [2, -1], 6, [3, 0], 3, [5, 1], 8, [3, 2], 11, [4, 0], 15, [6, 2]]$$

$$C_{1}' = [7, [5, -1], 12, [1, -1], 13, [17, -1], 17, [17, 1], 9, [2, 1]]$$

$$C_{2}' = [0, [2, 0], 2, [1, 2], 1, [1, 1]]$$

$$C_{3}' = [14, [8, 0], 5, [8, 2]]$$

$$C_{4}' = [10, [6, -1], 16, [6, 1]]$$
2-factor type $[6, 4, 4, 2, 2]$: two starters

$$C_{0} = [0, [7, 0], 7, [17, 2], 17, [17, 0], 11, [1, 1], 12, [4, 1], 16, [1, 2]]$$

$$C_{1} = [4, [5, 2], 9, [7, 1], 2, [6, 0], 8, [4, -1]]$$

$$C_2 = [14, [1, 0], 15, [7, 2], 5, [8, -1], 13, [1, -1]]$$

$$C_3 = [1, [2, -1], 3, [2, 1]]$$

$$C_4 = [10, [4, 0], 6, [4, 2]]$$

$$C_0' = [4, [6, 2], 15, [5, -1], 3, [6, 1], 14, [2, 0], 12, [3, 2], 9, [5, 0]]$$

- $C_1^\prime = [6, [17, -1], 17, [17, 1], 11, [3, 0], 8, [2, 2]]$
- $C_2^\prime = [7, [7, -1], 0, [5, 1], 5, [8, 1], 13, [6, -1]]$
- $C_3^\prime = [1, [8, 0], 10, [8, 2]]$

$$C'_4 = [2, [3, -1], 16, [3, 1]]$$

• 2-factor type [5, 5, 4, 2, 2]: two starters

$$\begin{split} C_0 &= [5, [1, 0], 4, [3, -1], 1, [3, 2], 15, [7, 1], 8, [3, 1]] \\ C_1 &= [17, [17, -1], 14, [1, -1], 13, [6, 2], 2, [2, 0], 0, [17, 1]] \\ C_2 &= [7, [8, 2], 16, [5, 0], 11, [5, -1], 6, [1, 1]] \\ C_3 &= [9, [6, -1], 3, [6, 1]] \\ C_4 &= [10, [2, -1], 12, [2, 1]] \\ C'_0 &= [14, [5, 1], 9, [3, 0], 12, [7, 2], 2, [6, 0], 13, [1, 2]] \\ C'_1 &= [16, [5, 2], 11, [8, 0], 3, [2, 2], 1, [7, 0], 8, [8, 1]] \\ C'_2 &= [5, [7, -1], 15, [8, -1], 7, [17, 2], 17, [17, 0]] \\ C'_3 &= [0, [4, 0], 4, [4, 2]] \\ C'_4 &= [10, [4, -1], 6, [4, 1]] \end{split}$$

• 2-factor type [7, 3, 3, 3, 2]: one starter

$$\begin{split} C_0 &= [13, [2, 1], 11, [2, 0], 9, [17, 1], 17, [17, 0], 8, [6, 0], 2, [4, 1], 6, [7, 0]] \\ C_1 &= [0, [3, 1], 14, [4, 0], 1, [1, 0]] \\ C_2 &= [12, [8, 1], 4, [1, 1], 5, [7, 1]] \\ C_3 &= [10, [3, 0], 7, [8, 0], 16, [6, 1]] \\ C_4 &= [15, [5, 0], 3, [5, 1]] \end{split}$$

• 2-factor type [6, 4, 3, 3, 2]: one starter

$$\begin{split} C_0 &= [14, [1, 1], 13, [4, 1], 9, [3, 0], 6, [17, 1], 17, [17, 0], 4, [7, 1]] \\ C_1 &= [3, [5, 0], 8, [2, 0], 10, [6, 0], 16, [4, 0]] \\ C_2 &= [5, [3, 1], 2, [2, 1], 0, [5, 1]] \\ C_3 &= [11, [7, 0], 1, [6, 1], 12, [1, 0]] \\ C_4 &= [7, [8, 0], 15, [8, 1]] \end{split}$$

• 2-factor type [5, 5, 3, 3, 2]: one starter

$$\begin{split} C_0 &= [17, [17, 1], 4, [7, 1], 11, [3, 1], 8, [7, 0], 15, [17, 0]] \\ C_1 &= [9, [2, 0], 7, [2, 1], 5, [1, 0], 6, [8, 1], 14, [5, 1]] \\ C_2 &= [1, [1, 1], 0, [5, 0], 12, [6, 0]] \\ C_3 &= [2, [6, 1], 13, [3, 0], 10, [8, 0]] \\ C_4 &= [3, [4, 0], 16, [4, 1]] \end{split}$$

• 2-factor type [5, 4, 4, 3, 2]: one starter

$$\begin{split} C_0 &= [11, [4, 1], 15, [6, 0], 4, [17, 1], 17, [17, 0], 1, [7, 1]] \\ C_1 &= [5, [6, 1], 16, [8, 1], 8, [5, 0], 13, [8, 0]] \\ C_2 &= [2, [4, 0], 6, [3, 0], 3, [3, 1], 0, [2, 1]] \\ C_3 &= [14, [2, 0], 12, [5, 1], 7, [7, 0]] \\ C_4 &= [9, [1, 0], 10, [1, 1]] \end{split}$$

• 2-factor type [4, 4, 4, 4, 2]: one starter

$$\begin{split} C_0 &= [16, [3, 1], 13, [6, 0], 7, [17, 0], 17, [17, 1]] \\ C_1 &= [9, [5, 0], 14, [8, 0], 6, [5, 1], 11, [2, 1]] \\ C_2 &= [10, [7, 1], 0, [8, 1], 8, [4, 0], 12, [2, 0]] \\ C_3 &= [5, [4, 1], 1, [3, 0], 4, [6, 1], 15, [7, 0]] \\ C_4 &= [3, [1, 0], 2, [1, 1]] \end{split}$$

- 2-factor type [6, 3, 3, 3, 3]: two starters
- $$\begin{split} C_0 &= [12, [7, -1], 2, [8, -1], 10, [4, -1], 14, [3, 0], 0, [4, 1], 4, [8, 2]] \\ C_1 &= [7, [2, 2], 5, [2, 1], 3, [4, 0]] \\ C_2 &= [16, [2, 0], 1, [5, -1], 13, [3, 2]] \\ C_3 &= [8, [3, -1], 11, [17, -1], 17, [17, 1]] \\ C_4 &= [15, [8, 0], 6, [3, 1], 9, [6, 2]] \\ C_0' &= [1, [17, 0], 17, [17, 2], 10, [6, 1], 4, [2, -1], 6, [7, 0], 13, [5, 2]] \\ C_1' &= [9, [5, 1], 14, [6, -1], 8, [1, 1]] \\ C_2' &= [16, [8, 1], 7, [7, 2], 0, [1, 0]] \\ C_3' &= [15, [5, 0], 3, [1, -1], 2, [4, 2]] \\ C_4' &= [5, [6, 0], 11, [1, 2], 12, [7, 1]] \end{split}$$

• 2-factor type [5, 4, 3, 3, 3]: two starters

$$\begin{split} C_0 &= [5, [8, 1], 14, [6, -1], 3, [5, -1], 8, [5, 0], 13, [8, 2]] \\ C_1 &= [12, [1, -1], 11, [5, 1], 16, [6, 2], 10, [2, 0]] \\ C_2 &= [0, [1, 1], 1, [3, -1], 15, [2, 1]] \\ C_3 &= [17, [17, 2], 4, [3, 0], 7, [17, 1]] \\ C_4 &= [9, [7, -1], 2, [4, 1], 6, [3, 1]] \\ C'_0 &= [5, [4, 2], 9, [17, -1], 17, [17, 0], 13, [2, 2], 11, [6, 0]] \\ C'_1 &= [12, [5, 2], 0, [2, -1], 2, [1, 0], 3, [8, -1]] \\ C'_2 &= [7, [8, 0], 15, [1, 2], 14, [7, 1]] \\ C'_3 &= [8, [7, 0], 1, [3, 2], 4, [4, -1]] \\ C'_4 &= [16, [7, 2], 6, [4, 0], 10, [6, 1]] \end{split}$$

• 2-factor type [4, 4, 4, 3, 3]: two starters

 $C_2 = [15, [7, -1], 5, [3, 0], 8, [2, 1], 10, [5, 2]]$ $C'_2 = [1, [1, 2], 0, [3, 1], 14, [6, 1], 8, [7, 0]]$ $C_3 = [1, [4, 0], 14, [17, 2], 17, [17, 1]]$ $C_4 = [0, [2, 2], 2, [4, -1], 6, [6, 0]]$

- $C_0 = [9, [7, 1], 16, [8, 1], 7, [3, -1], 4, [5, 1]] \qquad C'_0 = [4, [5, -1], 9, [17, -1], 17, [17, 0], 11, [7, 2]]$ $C_1 = [12, [8, 0], 3, [8, 2], 11, [2, -1], 13, [1, 1]] \qquad C'_1 = [10, [3, 2], 13, [1, 0], 12, [6, -1], 6, [4, 1]]$ $C'_{3} = [5, [6, 2], 16, [8, -1], 7, [2, 0]]$ $C'_4 = [15, [5, 0], 3, [1, -1], 2, [4, 2]]$
- 2-factor type [12, 2, 2, 2]: one starter

 $C_0 = [15, [5, 0], 3, [3, 1], 6, [5, 1], 11, [1, 0], 10, [17, 0], 17, [17, 1], 14, [3, 0], 0, [1, 1], 16, [6, 0], 5, [2, 0], 7, [17, 1], 14, [1, 0], 10, [17, 0], 17, [17, 1], 14, [13, 0], 0, [1, 1], 16, [10, 0], 5, [12, 0], 7, [10, 0], 10, [10, 0], 1$ [6,1], 13, [2,1]] $C_1 = [12, [7, 0], 2, [7, 1]]$ $C_2 = [1, [8, 0], 9, [8, 1]]$ $C_3 = [8, [4, 0], 4, [4, 1]]$

• 2-factor type [11, 3, 2, 2]: two starters

 $C_0 = [12, [6, 1], 1, [5, -1], 13, [7, 2], 3, [4, -1], 7, [2, 1], 5, [4, 0], 9, [5, 2], 4, [7, 0], 14, [6, 2], 8, [6, 0], 2, [7, -1]]$ $C_1 = [16, [17, 2], 17, [17, 0], 0, [1, -1]]$ $C_2 = [6, [8, -1], 15, [8, 1]]$ $C_3 = [10, [1, 0], 11, [1, 2]]$ $C_0' = [8, [2, -1], 10, [5, 0], 15, [3, 2], 1, [1, 1], 0, [8, 0], 9, [4, 1], 13, [17, 1], 17, [17, -1], 16, [7, 1], 6, [4, 2], 2, 10, [10, 10],$ [6, -1]]

- $C'_1 = [4, [8, 2], 12, [5, 1], 7, [3, 0]]$
- $C'_{2} = [3, [2, 0], 5, [2, 2]]$
- $C'_{3} = [11, [3, -1], 14, [3, 1]]$
- 2-factor type [10, 4, 2, 2]: two starters
 - $C_0 = [6, [3, 1], 9, [2, -1], 7, [17, 1], 17, [17, 2], 0, [6, 0], 11, [8, 1], 2, [6, 2], 8, [7, -1], 15, [5, 1], 10, [4, 0]]$ $C_1 = [5, [1, 2], 4, [5, 0], 16, [2, 1], 1, [4, 1]]$ $C_2 = [14, [1, -1], 13, [1, 1]]$ $C_3 = [3, [8, 0], 12, [8, 2]]$

$$\begin{split} C_0' &= [7, [6, -1], 13, [5, -1], 1, [7, 1], 11, [3, -1], 14, [6, 1], 8, [4, -1], 12, [17, -1], 17, [17, 0], 6, [7, 2], 16, \\ & [8, -1]] \\ C_1' &= [5, [4, 2], 9, [1, 0], 10, [5, 2], 15, [7, 0]] \\ C_2' &= [2, [2, 0], 4, [2, 2]] \\ C_3' &= [0, [3, 0], 3, [3, 2]] \end{split}$$

• 2-factor type [9, 5, 2, 2]: two starters

$$\begin{split} C_0 &= [13, [17, 2], 17, [17, 0], 12, [3, -1], 9, [7, 1], 16, [8, 2], 8, [6, 1], 2, [5, -1], 7, [1, 1], 6, [7, 0]] \\ C_1 &= [1, [3, 0], 4, [6, -1], 15, [4, -1], 11, [1, 2], 10, [8, -1]] \\ C_2 &= [14, [6, 0], 3, [6, 2]] \\ C_3 &= [5, [5, 0], 0, [5, 2]] \\ C'_0 &= [11, [4, 0], 7, [7, 2], 14, [1, 0], 13, [4, 2], 9, [8, 0], 0, [8, 1], 8, [4, 1], 12, [2, 2], 10, [1, -1]] \\ C'_1 &= [15, [3, 1], 1, [5, 1], 6, [3, 2], 3, [2, 0], 5, [7, -1]] \\ C'_2 &= [17, [17, -1], 16, [17, 1]] \\ C'_3 &= [4, [2, -1], 2, [2, 1]] \end{split}$$

- 2-factor type [8, 6, 2, 2]: two starters
 - $$\begin{split} C_0 &= [12, [7, 1], 2, [4, 2], 6, [7, 0], 16, [2, -1], 14, [1, -1], 13, [4, -1], 9, [5, -1], 4, [8, 1]] \\ C_1 &= [15, [2, 2], 0, [5, 0], 5, [2, 1], 7, [6, -1], 1, [8, -1], 10, [5, 1]] \\ C_2 &= [8, [3, 0], 11, [3, 2]] \\ C_3 &= [17, [17, 0], 3, [17, 2]] \\ C'_0 &= [5, [7, -1], 15, [8, 2], 7, [2, 0], 9, [7, 2], 2, [1, 0], 3, [5, 2], 8, [4, 0], 4, [1, 1]] \\ C'_1 &= [0, [1, 2], 1, [8, 0], 10, [6, 2], 16, [4, 1], 12, [6, 1], 6, [6, 0]] \\ C'_2 &= [11, [3, -1], 14, [3, 1]] \\ C'_3 &= [13, [17, -1], 17, [17, 1]] \end{split}$$
- 2-factor type [7, 7, 2, 2]: two starters

$$\begin{split} C_0 &= [11, [5, 2], 16, [4, -1], 3, [5, 0], 15, [8, -1], 7, [6, 2], 1, [8, 0], 10, [1, 1]] \\ C_1 &= [17, [17, 2], 4, [1, -1], 5, [3, 1], 2, [7, 1], 12, [4, 0], 8, [1, 2], 9, [17, 0]] \\ C_2 &= [6, [7, 0], 13, [7, 2]] \\ C_3 &= [0, [3, 0], 14, [3, 2]] \\ C_0' &= [9, [6, 1], 3, [4, 1], 16, [1, 0], 15, [8, 2], 7, [2, 0], 5, [8, 1], 13, [4, 2]] \\ C_1' &= [1, [3, -1], 4, [17, -1], 17, [17, 1], 11, [6, 0], 0, [7, -1], 10, [2, 2], 12, [6, -1]] \\ C_2' &= [14, [5, -1], 2, [5, 1]] \\ C_3' &= [6, [2, -1], 8, [2, 1]] \end{split}$$

• 2-factor type [10, 3, 3, 2]: one starter

$$\begin{split} C_0 &= [8, [7, 1], 1, [8, 1], 10, [4, 1], 6, [3, 0], 3, [1, 0], 2, [17, 1], 17, [17, 0], 13, [2, 0], 15, [6, 1], 9, [1, 1]] \\ C_1 &= [7, [4, 0], 11, [7, 0], 4, [3, 1]] \\ C_2 &= [16, [6, 0], 5, [8, 0], 14, [2, 1]] \\ C_3 &= [12, [5, 0], 0, [5, 1]] \end{split}$$

• 2-factor type [9, 4, 3, 2]: one starter

$$\begin{split} C_0 &= [5, [8, 1], 14, [2, 1], 16, [7, 0], 9, [17, 0], 17, [17, 1], 2, [8, 0], 10, [4, 0], 6, [5, 1], 11, [6, 1]] \\ C_1 &= [7, [6, 0], 1, [7, 1], 8, [4, 1], 4, [3, 0]] \\ C_2 &= [3, [3, 1], 0, [2, 0], 15, [5, 0]] \\ C_3 &= [13, [1, 0], 12, [1, 1]] \end{split}$$

• 2-factor type [8, 5, 3, 2]: one starter

$$\begin{split} C_0 &= [1, [7, 0], 8, [4, 0], 12, [4, 1], 16, [5, 0], 4, [1, 1], 3, [6, 1], 9, [17, 1], 17, [17, 0]] \\ C_1 &= [13, [2, 0], 15, [1, 0], 14, [3, 0], 0, [6, 0], 6, [7, 1]] \\ C_2 &= [10, [5, 1], 5, [2, 1], 7, [3, 1]] \\ C_3 &= [11, [8, 0], 2, [8, 1]] \end{split}$$

• 2-factor type [7, 6, 3, 2]: one starter

 $C_0 = [1, [5, 0], 6, [5, 1], 11, [7, 1], 4, [4, 0], 8, [1, 1], 9, [8, 0], 0, [1, 0]]$ $C_1 = [13, [3, 1], 10, [17, 1], 17, [17, 0], 5, [2, 1], 7, [8, 1], 15, [2, 0]]$ $C_2 = [16, [4, 1], 12, [7, 0], 2, [3, 0]]$ $C_3 = [3, [6, 0], 14, [6, 1]]$

• 2-factor type [8, 4, 4, 2]: one starter

$$\begin{split} C_0 &= [4, [8, 0], 12, [6, 1], 1, [7, 0], 8, [17, 0], 17, [17, 1], 16, [5, 1], 11, [4, 1], 15, [6, 0]] \\ C_1 &= [3, [3, 1], 6, [1, 0], 7, [3, 0], 10, [7, 1]] \\ C_2 &= [14, [5, 0], 9, [4, 0], 5, [8, 1], 13, [1, 1]] \\ C_3 &= [2, [2, 0], 0, [2, 1]] \end{split}$$

• 2-factor type [7, 5, 4, 2]: one starter

$$\begin{split} C_0 &= [2, [4, 0], 6, [7, 1], 16, [6, 1], 10, [2, 1], 12, [8, 0], 4, [3, 1], 1, [1, 1]] \\ C_1 &= [5, [7, 0], 15, [17, 0], 17, [17, 1], 13, [2, 0], 11, [6, 0]] \\ C_2 &= [8, [1, 0], 7, [4, 1], 3, [3, 0], 0, [8, 1]] \\ C_3 &= [14, [5, 0], 9, [5, 1]] \end{split}$$

• 2-factor type [6, 6, 4, 2]: one starter

$$\begin{split} C_0 &= [12, [2, 1], 14, [3, 1], 11, [7, 1], 4, [1, 1], 3, [7, 0], 13, [1, 0]]\\ C_1 &= [0, [5, 1], 5, [4, 1], 1, [6, 1], 7, [17, 0], 17, [17, 1], 2, [2, 0]]\\ C_2 &= [6, [3, 0], 9, [6, 0], 15, [5, 0], 10, [4, 0]]\\ C_3 &= [16, [8, 0], 8, [8, 1]] \end{split}$$

• 2-factor type [6, 5, 5, 2]: one starter

$$\begin{split} C_0 &= [10, [17, 0], 17, [17, 1], 7, [4, 1], 3, [8, 1], 11, [2, 0], 9, [1, 1]]\\ C_1 &= [8, [4, 0], 4, [8, 0], 12, [6, 0], 1, [5, 0], 6, [2, 1]]\\ C_2 &= [14, [5, 1], 2, [6, 1], 13, [3, 0], 16, [1, 0], 0, [3, 1]]\\ C_3 &= [5, [7, 0], 15, [7, 1]] \end{split}$$

• 2-factor type [9, 3, 3, 3]: two starters

 $C_0 = [10, [3, 2], 7, [6, 0], 13, [8, -1], 5, [1, 1], 6, [8, 2], 15, [3, 0], 1, [1, 2], 2, [1, -1], 3, [7, 0]]$ $C_1 = [16, [17, 0], 17, [17, 2], 11, [5, -1]]$ $C_2 = [9, [5, 0], 14, [7, 1], 4, [5, 2]]$ $C_3 = [0, [5, 1], 12, [4, 1], 8, [8, 1]]$ $C'_0 = [7, [7, -1], 0, [2, 1], 2, [3, -1], 16, [4, 2], 3, [2, 0], 1, [7, 2], 11, [1, 0], 12, [17, 1], 17, [17, -1]]$ $C'_1 = [4, [2, 2], 6, [4, 0], 10, [6, -1]]$ $C'_{2} = [9, [6, 1], 15, [2, -1], 13, [4, -1]]$ $C'_3 = [5, [8, 0], 14, [6, 2], 8, [3, 1]]$ • 2-factor type [8, 4, 3, 3]: two starters $C_0 = [12, [2, 1], 14, [5, 2], 2, [6, 0], 13, [8, 2], 5, [1, -1], 6, [1, 0], 7, [1, 2], 8, [4, 0]]$ $C_1 = [9, [17, 2], 17, [17, 0], 16, [4, 1], 3, [6, 1]]$ $C_2 = [4, [6, -1], 10, [7, -1], 0, [4, -1]]$ $C_3 = [11, [4, 2], 15, [3, -1], 1, [7, 0]]$ $C'_0 = [11, [2, 2], 9, [5, 0], 14, [2, -1], 12, [7, 1], 2, [8, 1], 10, [7, 2], 0, [8, 0], 8, [3, 1]]$ $C'_{1} = [7, [6, 2], 13, [5, -1], 1, [3, 0], 15, [8, -1]]$ $C'_{2} = [3, [3, 2], 6, [1, 1], 5, [2, 0]]$ $C'_{3} = [16, [17, -1], 17, [17, 1], 4, [5, 1]]$ • 2-factor type [7, 5, 3, 3]: two starters $C_0 = [15, [6, -1], 4, [4, 0], 8, [6, 2], 2, [6, 1], 13, [7, -1], 3, [8, 1], 12, [3, -1]]$ $C_1 = [11, [1, 2], 10, [4, 1], 6, [7, 0], 16, [17, 2], 17, [17, 0]]$ $C_2 = [5, [2, 2], 7, [2, 0], 9, [4, -1]]$ $C_3 = [14, [3, 0], 0, [1, 1], 1, [4, 2]]$ $C'_0 = [16, [6, 0], 5, [5, 2], 0, [8, 0], 9, [7, 2], 2, [7, 1], 12, [5, 0], 7, [8, 2]]$ $C'_1 = [17, [17, 1], 11, [5, 1], 6, [8, -1], 14, [1, -1], 15, [17, -1]]$ $C'_{2} = [8, [5, -1], 13, [3, 1], 10, [2, 1]]$

- $C'_3 = [4, [1, 0], 3, [2, -1], 1, [3, 2]]$
- 2-factor type [6, 6, 3, 3]: two starters

$$\begin{split} C_0 &= [10, [8, -1], 1, [2, 0], 16, [7, -1], 9, [17, 2], 17, [17, 1], 12, [2, 1]] \\ C_1 &= [14, [7, 2], 7, [7, 0], 0, [8, 2], 8, [5, -1], 13, [6, -1], 2, [5, 0]] \\ C_2 &= [15, [4, 0], 11, [6, 2], 5, [7, 1]] \\ C_3 &= [4, [1, -1], 3, [3, 1], 6, [2, -1]] \\ C_0' &= [9, [3, 0], 12, [4, -1], 8, [2, 2], 6, [5, 1], 1, [1, 1], 0, [8, 1]] \\ C_1' &= [11, [4, 1], 15, [1, 2], 14, [17, 0], 17, [17, -1], 7, [4, 2], 3, [8, 0]] \\ C_2' &= [13, [3, 2], 16, [3, -1], 2, [6, 0]] \\ C_3' &= [4, [6, 1], 10, [5, 2], 5, [1, 0]] \end{split}$$

• 2-factor type [7, 4, 4, 3]: two starters

$$\begin{split} C_0 &= [0, [5, 0], 12, [1, -1], 11, [5, 2], 6, [7, 0], 13, [3, 2], 10, [1, 0], 9, [8, 2]] \\ C_1 &= [4, [3, 1], 1, [17, 0], 17, [17, -1], 3, [1, 2]] \\ C_2 &= [2, [4, 1], 15, [1, 1], 14, [7, 1], 7, [5, 1]] \\ C_3 &= [16, [6, 2], 5, [3, 0], 8, [8, 1]] \\ C'_0 &= [11, [4, 0], 7, [2, 1], 9, [17, 2], 17, [17, 1], 10, [4, -1], 14, [6, -1], 3, [8, -1]] \\ C'_1 &= [8, [7, 2], 1, [3, -1], 4, [2, 0], 2, [6, 1]] \\ C'_2 &= [15, [2, -1], 0, [5, -1], 5, [8, 0], 13, [2, 2]] \\ C'_3 &= [12, [4, 2], 16, [7, -1], 6, [6, 0]] \end{split}$$
2-factor type [6, 5, 4, 3]: two starters

- $$\begin{split} C_0 &= [7, [7, 2], 14, [2, 0], 16, [4, 2], 12, [8, 1], 4, [17, -1], 17, [17, 0]]\\ C_1 &= [15, [5, 0], 10, [1, 2], 11, [7, 0], 1, [8, 2], 9, [6, -1]]\\ C_2 &= [6, [6, 1], 0, [5, -1], 5, [8, -1], 13, [7, 1]]\\ C_3 &= [2, [1, 0], 3, [5, 1], 8, [6, 2]]\\ C_0' &= [7, [2, -1], 9, [3, -1], 12, [8, 0], 4, [1, 1], 5, [2, 2], 3, [4, -1]]\\ C_1' &= [13, [3, 0], 16, [1, -1], 15, [2, 1], 0, [7, -1], 10, [3, 2]]\\ C_2' &= [8, [6, 0], 2, [4, 1], 6, [5, 2], 11, [3, 1]]\\ C_3' &= [17, [17, 2], 14, [4, 0], 1, [17, 1]] \end{split}$$
- 2-factor type [5, 5, 5, 3]: two starters

 $C_{0} = [3, [7, 0], 10, [1, 1], 11, [2, 2], 9, [5, 0], 4, [1, 2]]$ $C_{1} = [15, [17, 0], 17, [17, 2], 7, [6, 1], 1, [4, 1], 14, [1, -1]]$ $C_{2} = [16, [3, 2], 13, [7, 1], 6, [6, -1], 12, [7, -1], 2, [3, 0]]$ $C_{3} = [0, [5, 1], 5, [3, 1], 8, [8, -1]]$ $C'_{0} = [9, [2, 1], 7, [1, 0], 6, [6, 2], 0, [17, -1], 17, [17, 1]]$ $C'_{1} = [2, [3, -1], 16, [5, 2], 11, [6, 0], 5, [7, 2], 15, [4, 0]]$ $C'_{2} = [12, [8, 1], 3, [2, -1], 1, [8, 2], 10, [4, -1], 14, [2, 0]]$

- $C'_3 = [4, [4, 2], 8, [5, -1], 13, [8, 0]]$
- 2-factor type [6, 4, 4, 4]: two starters

$$\begin{split} C_0 &= [13, [2, -1], 15, [7, 0], 8, [3, 2], 11, [5, 0], 16, [5, 2], 4, [8, 1]] \\ C_1 &= [6, [6, 1], 12, [7, 2], 5, [3, 0], 2, [4, 1]] \\ C_2 &= [7, [17, -1], 17, [17, 1], 3, [2, 1], 1, [6, -1]] \\ C_3 &= [14, [3, 1], 0, [7, -1], 10, [1, -1], 9, [5, 1]] \\ C_0' &= [6, [7, 1], 13, [3, -1], 10, [1, 2], 11, [1, 0], 12, [17, 2], 17, [17, 0]] \\ C_1' &= [16, [2, 0], 14, [6, 2], 3, [4, 0], 7, [8, 2]] \\ C_2' &= [9, [8, 0], 1, [4, 2], 5, [1, 1], 4, [5, -1]] \\ C_3' &= [8, [8, -1], 0, [2, 2], 15, [4, -1], 2, [6, 0]] \end{split}$$

• 2-factor type [5, 5, 4, 4]: two starters

$$\begin{split} C_0 &= [7, [3, 1], 4, [8, -1], 12, [8, 0], 3, [5, 2], 15, [8, 1]] \\ C_1 &= [17, [17, 0], 14, [6, 2], 8, [6, 0], 2, [3, 2], 5, [17, -1]] \\ C_2 &= [16, [5, 0], 11, [2, 2], 9, [1, -1], 10, [6, -1]] \\ C_3 &= [13, [4, 0], 0, [1, 1], 1, [5, 1], 6, [7, 2]] \\ C_0' &= [10, [7, 0], 0, [5, -1], 12, [8, 2], 4, [2, -1], 6, [4, -1]] \\ C_1' &= [15, [6, 1], 9, [2, 0], 7, [4, 1], 11, [17, 1], 17, [17, 2]] \\ C_2' &= [3, [4, 2], 16, [2, 1], 14, [1, 0], 13, [7, 1]] \\ C_3' &= [2, [1, 2], 1, [7, -1], 8, [3, 0], 5, [3, -1]] \end{split}$$

- 2-factor type [14, 2, 2]: two starters
 - $$\begin{split} C_0 = [6, [5, 0], 1, [7, 2], 8, [17, -1], 17, [17, 0], 16, [1, 2], 0, [2, 1], 15, [6, 1], 9, [4, -1], 13, [1, 1], 12, [8, -1], 3, \\ [2, -1], 5, [5, -1], 10, [1, -1], 11, [5, 1]] \end{split}$$
 - $C_1 = [2, [2, 0], 4, [2, 2]]$
 - $C_2 = [14, [7, -1], 7, [7, 1]]$
 - $$\begin{split} C_0' = [15, [17, 1], 17, [17, 2], 16, [4, 0], 3, [8, 1], 11, [5, 2], 6, [6, -1], 12, [7, 0], 5, [3, 2], 2, [6, 0], 8, [4, 2], 4, \\ [3, 0], 7, [6, 2], 1, [4, 1], 14, [1, 0]] \end{split}$$

$$C'_1 = [9, [8, 0], 0, [8, 2]]$$

- $C_2^\prime = [10, [3, -1], 13, [3, 1]]$
- 2-factor type [13, 3, 2]: one starter
 - $$\begin{split} C_0 &= [1, [17, 0], 17, [17, 1], 7, [1, 1], 8, [3, 0], 11, [8, 0], 2, [2, 1], 4, [5, 0], 16, [7, 1], 6, [4, 1], 10, [7, 0], 0, [5, 1], \\ &\quad 5, [4, 0], 9, [8, 1]] \\ C_0 &= [12, [1, 0], 12, [2, 0], 15, [2, 1]] \end{split}$$
 - $C_1 = [12, [1, 0], 13, [2, 0], 15, [3, 1]]$
 - $C_2 = [14, [6, 0], 3, [6, 1]]$
- 2-factor type [12, 4, 2]: one starter
 - $$\begin{split} C_0 &= [4, [8, 1], 13, [4, 0], 9, [7, 0], 16, [8, 0], 7, [7, 1], 14, [3, 0], 11, [4, 1], 15, [17, 1], 17, [17, 0], 3, [1, 0], 2, \\ &\quad [3, 1], 5, [1, 1]] \\ C_1 &= [0, [5, 0], 12, [6, 1], 1, [5, 1], 6, [6, 0]] \\ C_2 &= [8, [2, 0], 10, [2, 1]] \end{split}$$
- 2-factor type [11, 5, 2]: one starter

$$\begin{split} C_0 &= [17, [17, 1], 11, [3, 0], 8, [7, 1], 15, [6, 0], 4, [5, 1], 16, [1, 1], 0, [1, 0], 1, [4, 0], 5, [5, 0], 10, [3, 1], 13, [17, 0]] \\ C_1 &= [12, [7, 0], 2, [4, 1], 6, [8, 1], 14, [6, 1], 3, [8, 0]] \\ C_2 &= [7, [2, 0], 9, [2, 1]] \end{split}$$

• 2-factor type [10, 6, 2]: one starter

$$\begin{split} C_0 &= [10, [7, 0], 0, [8, 0], 8, [5, 1], 13, [1, 1], 12, [6, 0], 6, [1, 0], 7, [8, 1], 16, [2, 0], 1, [4, 1], 14, [4, 0]] \\ C_1 &= [11, [7, 1], 4, [17, 1], 17, [17, 0], 3, [5, 0], 15, [6, 1], 9, [2, 1]] \\ C_2 &= [2, [3, 0], 5, [3, 1]] \end{split}$$

• 2-factor type [9, 7, 2]: one starter

$$\begin{split} C_0 &= [9, [3, 1], 12, [4, 0], 8, [3, 0], 5, [6, 0], 11, [5, 1], 16, [8, 0], 7, [8, 1], 15, [5, 0], 10, [1, 0]] \\ C_1 &= [1, [1, 1], 2, [2, 0], 0, [6, 1], 6, [2, 1], 4, [17, 1], 17, [17, 0], 14, [4, 1]] \\ C_2 &= [3, [7, 0], 13, [7, 1]] \end{split}$$

- 2-factor type [8, 8, 2]: one starter $C_0 = [4, [7, 0], 14, [3, 1], 0, [4, 0], 13, [1, 0], 12, [7, 1], 5, [3, 0], 2, [4, 1], 6, [2, 1]]$ $C_1 = [7, [6, 0], 1, [8, 0], 10, [17, 0], 17, [17, 1], 11, [2, 0], 9, [6, 1], 15, [1, 1], 16, [8, 1]]$ $C_2 = [3, [5, 0], 8, [5, 1]]$
- 2-factor type [12, 3, 3]: two starters
 - $$\begin{split} C_0 &= [12, [1, -1], 11, [6, 1], 0, [7, 1], 7, [17, 2], 17, [17, 1], 14, [6, -1], 3, [6, 0], 9, [1, 2], 8, [3, 1], 5, [7, 0], 15, \\ &= [6, 2], 4, [8, 0]] \\ C_1 &= [13, [5, 0], 1, [2, -1], 16, [3, 2]] \\ C_2 &= [6, [4, -1], 2, [8, 2], 10, [4, 0]] \\ C_0' &= [0, [8, -1], 8, [1, 0], 7, [7, 2], 14, [2, 0], 12, [4, 1], 16, [3, -1], 13, [8, 1], 4, [5, 2], 9, [3, 0], 6, [5, -1], 1, \\ &= [1, 1], 2, [2, 2]] \\ C_1' &= [17, [17, 0], 11, [4, 2], 15, [17, -1]] \\ C_2' &= [10, [5, 1], 5, [2, 1], 3, [7, -1]] \\ 2 \text{ fractor type } [11, 4, 2], \text{ type startore} \end{split}$$
- 2-factor type [11, 4, 3]: two starters
 - $C_{0} = \begin{bmatrix} 16, [2, 1], 14, [17, -1], 17, [17, 1], 13, [3, 0], 10, [8, 1], 1, [6, 2], 12, [7, 0], 2, [3, 2], 5, [6, 0], 11, [2, -1], \\9, [7, 2] \end{bmatrix}$
 - $C_1 = [0, [7, -1], 7, [4, 0], 3, [5, 1], 8, [8, 2]]$
 - $C_2 = [6, [2, 2], 4, [6, 1], 15, [8, 0]]$
 - $$\begin{split} C_0' = [4, [7, 1], 11, [5, -1], 6, [4, 1], 10, [1, -1], 9, [3, 1], 12, [4, 2], 16, [3, -1], 2, [17, 0], 17, [17, 2], 14, [6, -1], \\ 3, [1, 0]] \end{split}$$
 - $C_1' = [13, [2, 0], 15, [8, -1], 7, [1, 1], 8, [5, 2]]$
 - $C'_2 = [0, [5, 0], 5, [4, -1], 1, [1, 2]]$
- 2-factor type [10, 5, 3]: two starters

$$\begin{split} C_0 &= [5, [17, -1], 17, [17, 1], 8, [5, 0], 3, [8, -1], 12, [5, -1], 7, [2, 1], 9, [4, 1], 13, [7, 2], 6, [6, 0], 0, [5, 2]] \\ C_1 &= [10, [6, -1], 4, [7, 1], 11, [8, 2], 2, [1, 1], 1, [8, 0]] \\ C_2 &= [14, [2, 0], 16, [1, -1], 15, [1, 2]] \\ C'_0 &= [10, [3, 2], 13, [17, 0], 17, [17, 2], 4, [5, 1], 16, [2, -1], 1, [3, 0], 15, [3, -1], 12, [7, -1], 5, [4, 2], 9, [1, 0]] \\ C'_1 &= [0, [3, 1], 14, [7, 0], 7, [4, -1], 3, [8, 1], 11, [6, 2]] \\ C'_2 &= [2, [4, 0], 6, [2, 2], 8, [6, 1]] \end{split}$$
• 2-factor type [9, 6, 3]: two starters

$$\begin{split} C_0 &= [16, [1, 2], 0, [6, 0], 6, [1, 1], 7, [2, 1], 9, [4, 1], 13, [7, 1], 3, [8, 1], 11, [7, 2], 4, [5, 0]] \\ C_1 &= [1, [3, -1], 15, [4, 2], 2, [7, 0], 12, [2, 2], 14, [8, -1], 5, [4, 0]] \\ C_2 &= [10, [2, -1], 8, [17, 2], 17, [17, 0]] \\ C'_0 &= [2, [6, 2], 8, [5, 1], 3, [3, 1], 0, [7, -1], 7, [4, -1], 11, [1, 0], 12, [6, -1], 6, [17, 1], 17, [17, -1]] \\ C'_1 &= [14, [2, 0], 16, [1, -1], 15, [6, 1], 4, [3, 2], 1, [8, 0], 9, [5, 2]] \\ C'_2 &= [10, [3, 0], 13, [8, 2], 5, [5, -1]] \end{split}$$

• 2-factor type [8,7,3]: two starters

$$\begin{split} C_0 &= [6, [7, 2], 16, [8, -1], 7, [6, -1], 13, [17, 0], 17, [17, -1], 5, [5, 2], 0, [3, 0], 3, [3, 1]] \\ C_1 &= [4, [4, 2], 8, [7, -1], 15, [1, 1], 14, [5, 0], 2, [8, 1], 10, [1, 2], 11, [7, 0]] \\ C_2 &= [1, [6, 0], 12, [3, -1], 9, [8, 2]] \\ C_0' &= [16, [1, -1], 0, [1, 0], 1, [6, 1], 7, [7, 1], 14, [3, 2], 11, [8, 0], 2, [2, 2], 4, [5, 1]] \\ C_1' &= [10, [4, 0], 6, [2, 1], 8, [4, 1], 12, [17, 2], 17, [17, 1], 3, [2, -1], 5, [5, -1]] \\ C_2' &= [9, [4, -1], 13, [2, 0], 15, [6, 2]] \end{split}$$

• 2-factor type [10, 4, 4]: two starters

$$\begin{split} C_0 &= [3, [7, 2], 13, [3, 1], 10, [4, 0], 14, [8, 1], 5, [1, -1], 4, [2, 1], 6, [5, 2], 1, [1, 0], 2, [8, 2], 11, [8, 0]] \\ C_1 &= [12, [4, -1], 8, [1, 2], 7, [7, 0], 0, [5, -1]] \\ C_2 &= [15, [1, 1], 16, [17, 1], 17, [17, -1], 9, [6, 1]] \\ C'_0 &= [8, [4, 2], 12, [17, 0], 17, [17, 2], 16, [5, 0], 4, [6, 2], 15, [4, 1], 11, [7, -1], 1, [5, 1], 13, [3, 0], 10, \\ &\qquad [2, -1]] \\ C'_1 &= [3, [6, 0], 14, [8, -1], 6, [6, -1], 0, [3, 2]] \\ C'_2 &= [2, [3, -1], 5, [2, 0], 7, [2, 2], 9, [7, 1]] \end{split}$$

• 2-factor type [9, 5, 4]: two starters

$$\begin{split} C_0 &= [1, [1, 0], 2, [4, 2], 15, [1, -1], 16, [7, -1], 6, [2, 0], 4, [1, 2], 5, [5, -1], 10, [2, -1], 12, [6, -1]] \\ C_1 &= [17, [17, 1], 9, [4, -1], 13, [7, 0], 3, [5, 2], 8, [17, -1]] \\ C_2 &= [11, [4, 0], 7, [7, 1], 14, [3, 2], 0, [6, 1]] \\ C_0' &= [8, [8, 1], 0, [8, 0], 9, [17, 2], 17, [17, 0], 16, [6, 2], 5, [3, 0], 2, [7, 2], 12, [1, 1], 11, [3, 1]] \\ C_1' &= [6, [2, 1], 4, [3, -1], 7, [4, 1], 3, [6, 0], 14, [8, 2]] \\ \end{split}$$

- $C'_2 = [1, [8, -1], 10, [5, 1], 15, [2, 2], 13, [5, 0]]$
- 2-factor type [8, 6, 4]: two starters

$$\begin{split} C_0 &= [11, [7, -1], 1, [5, 1], 13, [2, 0], 15, [4, 2], 2, [6, 0], 8, [1, 1], 9, [2, 2], 7, [4, -1]] \\ C_1 &= [3, [4, 0], 16, [7, 1], 6, [1, 2], 5, [5, 0], 0, [3, 2], 14, [6, -1]] \\ C_2 &= [12, [8, 1], 4, [17, 0], 17, [17, 2], 10, [2, -1]] \\ C'_0 &= [1, [3, 0], 15, [4, 1], 2, [5, 2], 7, [1, 0], 8, [3, 1], 11, [2, 1], 13, [3, -1], 10, [8, 2]] \\ C'_1 &= [16, [7, 0], 9, [5, -1], 4, [1, -1], 3, [8, -1], 12, [7, 2], 5, [6, 1]] \\ C'_2 &= [0, [6, 2], 6, [8, 0], 14, [17, 1], 17, [17, -1]] \end{split}$$

• 2-factor type [7,7,4]: two starters

$$\begin{split} C_0 &= [3, [7, -1], 10, [4, 0], 14, [8, 1], 5, [6, 2], 16, [4, 1], 12, [3, -1], 15, [5, -1]] \\ C_1 &= [13, [7, 0], 6, [17, 2], 17, [17, 0], 11, [7, 2], 4, [2, 1], 2, [1, 1], 1, [5, 1]] \\ C_2 &= [7, [2, 0], 9, [8, 2], 0, [8, 0], 8, [1, 2]] \\ C_0' &= [1, [1, 0], 0, [3, 2], 14, [6, 0], 8, [5, 2], 13, [3, 0], 10, [2, 2], 12, [6, -1]] \\ C_1' &= [4, [17, 1], 17, [17, -1], 16, [3, 1], 2, [8, -1], 11, [4, -1], 7, [1, -1], 6, [2, -1]] \\ C_2' &= [9, [4, 2], 5, [7, 1], 15, [5, 0], 3, [6, 1]] \end{split}$$

• 2-factor type [8, 5, 5]: two starters

$$\begin{split} C_0 &= [16, [6, -1], 10, [5, -1], 5, [8, 1], 13, [8, 2], 4, [2, 0], 6, [5, 2], 1, [6, 0], 12, [4, 1]] \\ C_1 &= [3, [8, -1], 11, [3, -1], 14, [3, 0], 0, [7, 1], 7, [4, 2]] \\ C_2 &= [15, [4, 0], 2, [7, 2], 9, [17, -1], 17, [17, 1], 8, [7, -1]] \\ C_0' &= [16, [1, 0], 15, [5, 1], 3, [3, 2], 6, [6, 1], 0, [1, -1], 1, [4, -1], 5, [8, 0], 14, [2, 2]] \\ C_1' &= [10, [6, 2], 4, [2, 1], 2, [7, 0], 9, [2, -1], 7, [3, 1]] \\ C_2' &= [17, [17, 2], 8, [5, 0], 13, [1, 2], 12, [1, 1], 11, [17, 0]] \end{split}$$

• 2-factor type [7, 6, 5]: two starters

$$\begin{split} C_0 &= [14, [1, 1], 15, [4, -1], 11, [3, -1], 8, [8, -1], 0, [1, -1], 16, [2, 2], 1, [4, 0]] \\ C_1 &= [13, [7, 1], 6, [2, 0], 4, [1, 2], 5, [5, 0], 10, [3, 2], 7, [6, -1]] \\ C_2 &= [9, [7, -1], 2, [17, 1], 17, [17, 2], 3, [8, 1], 12, [3, 0]] \\ C_0' &= [15, [17, 0], 17, [17, -1], 16, [7, 2], 6, [8, 0], 14, [2, -1], 12, [6, 2], 1, [3, 1]] \\ C_1' &= [9, [2, 1], 11, [8, 2], 3, [1, 0], 4, [6, 1], 10, [5, -1], 5, [4, 1]] \\ C_2' &= [13, [5, 1], 8, [6, 0], 2, [5, 2], 7, [7, 0], 0, [4, 2]] \end{split}$$

- 2-factor type [16, 2]: one starter
 - $$\begin{split} C_0 &= [11, [5, 0], 16, [4, 1], 12, [2, 0], 10, [5, 1], 5, [1, 1], 4, [3, 1], 7, [4, 0], 3, [6, 1], 14, [17, 0], 17, [17, 1], 2, \\ &\quad [1, 0], 1, [8, 0], 9, [6, 0], 15, [2, 1], 0, [8, 1], 8, [3, 0]] \\ C_1 &= [6, [7, 0], 13, [7, 1]] \end{split}$$
- 2-factor type [15, 3]: two starters

$$\begin{split} C_0 &= [15, [8, 1], 6, [17, 1], 17, [17, -1], 16, [8, 2], 8, [1, 0], 7, [5, 2], 2, [2, -1], 4, [3, 1], 1, [5, 0], 13, [4, -1], 9, \\ &\quad [3, 2], 12, [7, 0], 5, [8, -1], 14, [6, 1], 3, [5, -1]] \\ C_1 &= [11, [1, 2], 10, [7, -1], 0, [6, 0]] \end{split}$$

- $$\begin{split} C_0' &= [0, [5, 1], 5, [4, 2], 9, [3, 0], 6, [2, 1], 8, [17, 2], 17, [17, 0], 13, [6, -1], 7, [6, 2], 1, [4, 0], 14, [3, -1], 11, \\ &\quad [1, -1], 10, [7, 2], 3, [4, 1], 16, [1, 1], 15, [2, 0]] \\ C_1' &= [4, [8, 0], 12, [7, 1], 2, [2, 2]] \end{split}$$
- 2-factor type [14, 4]: two starters
 - $C_0 = [7, [6, 1], 1, [3, 0], 15, [2, 1], 0, [8, 2], 9, [1, 0], 10, [4, 2], 6, [2, 0], 8, [3, 1], 11, [6, 2], 5, [17, 0], 17, [17, 2], 16, [3, -1], 2, [1, -1], 3, [4, -1]]$
 - $C_1 = [13, [1, 2], 14, [7, -1], 4, [8, 0], 12, [1, 1]]$

$$\begin{split} C_0' &= [9, [3, 2], 12, [8, -1], 3, [5, 1], 15, [17, -1], 17, [17, 1], 2, [6, -1], 8, [8, 1], 0, [5, 0], 5, [2, 2], 7, [6, 0], 13, \\ & [7, 2], 6, [4, 0], 10, [4, 1], 14, [5, -1]] \\ C_0' &= [4, [5, 0], 14, [5, -1]] \\ C_0' &$$

- $C'_1 = [4, [7, 0], 11, [7, 1], 1, [2, -1], 16, [5, 2]]$
- 2-factor type [13, 5]: two starters
 - $C_{0} = [8, [2, 1], 6, [4, 1], 2, [2, 0], 4, [6, -1], 15, [4, 2], 11, [5, 0], 16, [6, 1], 5, [2, 2], 3, [7, -1], 10, [7, 0], 0, [4, -1], 13, [6, 2], 7, [1, -1]]$
 - $C_1 = [9, [3, 1], 12, [17, 2], 17, [17, 1], 1, [4, 0], 14, [5, -1]]$
 - $$\begin{split} C_0' = & [11, [7, 1], 1, [8, 2], 9, [17, 0], 17, [17, -1], 13, [5, 2], 8, [8, -1], 16, [3, 0], 2, [2, -1], 0, [3, 2], 14, [8, 0], 5, \\ & [7, 2], 15, [5, 1], 10, [1, 0]] \end{split}$$
 - $C_1' = [4, [3, -1], 7, [1, 1], 6, [6, 0], 12, [8, 1], 3, [1, 2]]$
- 2-factor type [12, 6]: two starters
 - $C_{0} = [5, [8, -1], 13, [1, 0], 14, [4, 2], 10, [17, 0], 17, [17, 2], 0, [3, 0], 3, [8, 2], 11, [7, 1], 4, [3, 1], 1, [2, 0], 16, [1, -1], 15, [7, 2]]$
 - $C_1 = [9, [7, -1], 2, [5, 2], 7, [5, 0], 12, [6, -1], 6, [2, 1], 8, [1, 1]]$
 - $C_0' = \begin{bmatrix} 1, [17, 1], 17, [17, -1], 0, [3, 2], 14, [2, -1], 16, [7, 0], 9, [1, 2], 10, [5, 1], 5, [8, 0], 13, [8, 1], 4, [2, 2], 6, \\ \begin{bmatrix} 6, 0 \end{bmatrix}, 12, \begin{bmatrix} 6, 1 \end{bmatrix} \end{bmatrix}$
 - $C_1' = [3, [5, -1], 15, [4, 1], 2, [6, 2], 8, [3, -1], 11, [4, 0], 7, [4, -1]]$
- 2-factor type [11,7]: two starters
 - $$\begin{split} C_0 = [4, [17, 1], 17, [17, 2], 15, [3, -1], 1, [4, 0], 14, [7, 2], 7, [4, -1], 11, [8, 1], 3, [6, 0], 9, [7, -1], 2, [3, 2], 5, \\ [1, 0]] \end{split}$$
 - $C_1 = \begin{bmatrix} 16, [7,0], 6, [6,2], 0, [5,1], 12, [2,0], 10, [3,1], 13, [5,2], 8, [8,-1] \end{bmatrix}$
 - $$\begin{split} C_0' = [5, [1, 2], 6, [2, 1], 4, [5, 0], 9, [1, -1], 8, [8, 2], 0, [2, -1], 2, [17, -1], 17, [17, 0], 10, [6, 1], 16, [5, -1], 11, \\ [6, -1]] \end{split}$$
 - $C_1' = \begin{bmatrix} 13, [2,2], 15, [3,0], 1, [4,2], 14, [7,1], 7, [4,1], 3, [8,0], 12, [1,1] \end{bmatrix}$
- 2-factor type [10, 8]: two starters
 - $C_0 = [3, [5, 1], 8, [7, 2], 15, [6, 1], 9, [7, -1], 16, [2, 0], 1, [1, -1], 0, [4, 2], 4, [3, -1], 7, [1, 1], 6, [3, 0]]$
 - $C_1 = [11, [6, -1], 5, [8, 2], 13, [17, 0], 17, [17, 2], 2, [5, 0], 14, [2, -1], 12, [2, 2], 10, [1, 0]]$
 - $$\begin{split} C_0' = [7, [2, 1], 9, [17, 1], 17, [17, -1], 6, [8, -1], 15, [4, -1], 11, [5, 2], 16, [6, 0], 5, [5, -1], 0, [3, 2], \\ 3, [4, 0]] \end{split}$$
 - $C_1' = [14, [7, 0], 4, [8, 1], 12, [1, 2], 13, [3, 1], 10, [8, 0], 2, [6, 2], 8, [7, 1], 1, [4, 1]]$

K Computational results for n = 19

• 2-factor type [3, 2, 2, 2, 2, 2, 2, 2, 2]: three starters

 $C_0 = [12, [1, 0], 11, [4, 2], 15, [3, 1]]$ $C'_0 = [6, [3, -1], 3, [1, 2], 2, [4, 0]]$ $C'_1 = [12, [8, -1], 4, [8, 1]]$ $C_1 = [5, [4, -1], 1, [4, 1]]$ $C'_{2} = [16, [1, -1], 15, [1, 1]]$ $C_2 = [3, [8, 0], 13, [8, 2]]$ $C_3 = [7, [7, 0], 0, [7, 2]]$ $C'_3 = [17, [7, -1], 10, [7, 1]]$ $C'_4 = [14, [3, 0], 11, [3, 2]]$ $C_4 = [4, [5, -1], 17, [5, 1]]$ $C'_5 = [5, [2, -1], 7, [2, 1]]$ $C_5 = [8, [2, 0], 6, [2, 2]]$ $C_6 = [2, [18, -1], 18, [18, 1]]$ $C'_6 = [13, [6, -1], 1, [6, 1]]$ $C'_7 = [18, [18, 0], 8, [18, 2]]$ $C_7 = [10, [6, 0], 16, [6, 2]]$ $C'_8 = [0, [9, 0], 9, [9, 2]]$ $C_8 = [14, [5, 0], 9, [5, 2]]$

• 2-factor type [5, 2, 2, 2, 2, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [7, [5, 2], 2, [8, -1], 10, [18, -1], 18, [18, 1], 6, [1, 0]] \\ C_1 &= [4, [6, -1], 16, [6, 1]] \\ C_2 &= [15, [8, 0], 5, [8, 2]] \\ C_3 &= [0, [5, -1], 13, [5, 1]] \\ C_4 &= [11, [3, 0], 8, [3, 2]] \\ C_5 &= [17, [2, 0], 1, [2, 2]] \\ C_6 &= [12, [3, -1], 9, [3, 1]] \\ C_7 &= [14, [7, -1], 3, [7, 1]] \\ C_7 &= [14, [7, -1], 3, [7, 1]] \\ C_0' &= [7, [7, 2], 14, [7, 0], 3, [1, 2], 2, [8, 1], 12, [5, 0]] \\ C_1' &= [0, [9, 0], 9, [9, 2]] \\ C_2' &= [4, [6, 0], 10, [6, 2]] \\ C_3' &= [17, [1, -1], 16, [1, 1]] \\ C_4' &= [13, [18, 0], 18, [18, 2]] \\ C_5' &= [8, [2, -1], 6, [2, 1]] \\ C_6' &= [11, [4, 0], 15, [4, 2]] \\ C_7' &= [5, [4, -1], 1, [4, 1]] \end{split}$$

• 2-factor type [4, 3, 2, 2, 2, 2, 2, 2]: three starters

$$\begin{array}{ll} C_0 = [10, [2, -1], 12, [6, 1], 6, [2, 0], 4, [6, 2]] & C_0' = [3, [2, 2], 1, [7, -1], 8, [6, 0], 2, [1, 1]] \\ C_1 = [0, [1, 2], 17, [1, 0], 16, [2, 1]] & C_1' = [11, [6, -1], 17, [7, 1], 10, [1, -1]] \\ C_2 = [14, [3, -1], 11, [3, 1]] & C_2' = [0, [9, 0], 9, [9, 2]] \\ C_3 = [13, [5, 0], 8, [5, 2]] & C_3' = [13, [8, -1], 5, [8, 1]] \\ C_4 = [9, [18, -1], 18, [18, 1]] & C_4' = [18, [18, 0], 14, [18, 2]] \\ C_5 = [2, [3, 0], 5, [3, 2]] & C_5' = [6, [8, 0], 16, [8, 2]] \\ C_6 = [3, [4, -1], 7, [4, 1]] & C_6' = [12, [5, -1], 7, [5, 1]] \\ C_7 = [1, [4, 0], 15, [4, 2]] & C_7' = [15, [7, 0], 4, [7, 2]] \end{array}$$

• 2-factor type [3, 3, 3, 2, 2, 2, 2, 2]: three starters

$C_0 = [17, [3, -1], 2, [1, 1], 1, [2, 1]]$	$C_0' = [1, [7, 1], 12, [1, -1], 13, [6, -1]]$
$C_1 = [4, [4, -1], 0, [3, 2], 3, [1, 0]]$	$C_1' = [15, [1, 2], 16, [6, 1], 10, [5, 0]]$
$C_2 = [13, [7, -1], 6, [3, 1], 9, [4, 1]]$	$C_2^\prime = [2, [2, -1], 4, [3, 0], 7, [5, 2]]$
$C_3 = [10, [2, 0], 12, [2, 2]]$	$C_3' = [0, [9, 0], 9, [9, 2]]$
$C_4 = [18, [18, 0], 5, [18, 2]]$	$C_4' = [17, [6, 0], 11, [6, 2]]$
$C_5 = [11, [4, 0], 15, [4, 2]]$	$C_5' = [5, [18, -1], 18, [18, 1]]$
$C_6 = [16, [8, 0], 8, [8, 2]]$	$C_6' = [3, [5, -1], 8, [5, 1]]$
$C_7 = [14, [7, 0], 7, [7, 2]]$	$C_7' = [6, [8, -1], 14, [8, 1]]$

• 2-factor type [7, 2, 2, 2, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [9, [1, 2], 8, [5, -1], 13, [1, 0], 14, [5, 1], 1, [3, 2], 4, [5, 0], 17, [8, -1]] \\ C_1 &= [5, [2, -1], 3, [2, 1]] \\ C_2 &= [6, [1, -1], 7, [1, 1]] \\ C_3 &= [18, [18, 0], 11, [18, 2]] \\ C_4 &= [16, [2, 0], 0, [2, 2]] \\ C_5 &= [15, [3, -1], 12, [3, 1]] \\ C_6 &= [10, [8, 0], 2, [8, 2]] \\ C'_0 &= [16, [3, 0], 13, [8, 1], 3, [18, 1], 18, [18, -1], 11, [4, -1], 7, [5, 2], 2, [4, 1]] \\ C'_1 &= [0, [9, 0], 9, [9, 2]] \\ C'_2 &= [17, [7, 0], 10, [7, 2]] \\ C'_3 &= [6, [6, -1], 12, [6, 1]] \\ C'_4 &= [5, [4, 0], 1, [4, 2]] \\ C'_5 &= [8, [6, 0], 14, [6, 2]] \\ C'_6 &= [15, [7, -1], 4, [7, 1]] \end{split}$$

• 2-factor type [6, 3, 2, 2, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [13, [2, -1], 11, [3, -1], 8, [18, 2], 18, [18, 0], 0, [1, 1], 1, [6, -1]] \\ C_1 &= [7, [7, 1], 14, [3, 1], 17, [8, 1]] \\ C_2 &= [2, [1, 0], 3, [1, 2]] \\ C_3 &= [16, [4, -1], 12, [4, 1]] \\ C_4 &= [15, [6, 0], 9, [6, 2]] \\ C_5 &= [4, [2, 0], 6, [2, 2]] \\ C_6 &= [5, [5, -1], 10, [5, 1]] \\ C'_0 &= [10, [7, -1], 17, [4, 0], 3, [1, -1], 2, [4, 2], 6, [18, -1], 18, [18, 1]] \\ C'_1 &= [7, [6, 1], 13, [2, 1], 15, [8, -1]] \\ C'_2 &= [0, [9, 0], 9, [9, 2]] \\ C'_3 &= [5, [7, 0], 16, [7, 2]] \\ C'_4 &= [12, [8, 0], 4, [8, 2]] \\ C'_5 &= [8, [3, 0], 11, [3, 2]] \\ C'_6 &= [1, [5, 0], 14, [5, 2]] \end{split}$$

$$\begin{split} C_0 &= [6, [6, 2], 12, [6, 0], 0, [1, 2], 1, [6, -1], 7, [1, 0]] \\ C_1 &= [15, [7, 1], 8, [18, 0], 18, [18, -1], 4, [7, 2]] \\ C_2 &= [11, [1, -1], 10, [1, 1]] \\ C_3 &= [16, [5, 0], 3, [5, 2]] \\ C_4 &= [14, [3, -1], 17, [3, 1]] \\ C_5 &= [9, [4, -1], 13, [4, 1]] \\ C_6 &= [5, [3, 0], 2, [3, 2]] \\ C'_0 &= [17, [7, -1], 10, [8, 2], 2, [7, 0], 13, [2, -1], 11, [6, 1]] \\ C'_1 &= [14, [18, 1], 18, [18, 2], 4, [8, 0], 12, [2, 1]] \\ C'_2 &= [0, [9, 0], 9, [9, 2]] \\ C'_3 &= [8, [2, 0], 6, [2, 2]] \\ C'_4 &= [1, [4, 0], 5, [4, 2]] \\ C'_5 &= [7, [8, -1], 15, [8, 1]] \\ C'_6 &= [3, [5, -1], 16, [5, 1]] \end{split}$$

• 2-factor type [5, 4, 2, 2, 2, 2, 2]: three starters

- 2-factor type [5, 3, 3, 2, 2, 2, 2]: three starters
 - $$\begin{split} C_0 &= [12, [5, 2], 17, [6, -1], 5, [1, 0], 6, [3, 2], 9, [3, 0]] \\ C_1 &= [15, [2, 0], 13, [18, 2], 18, [18, 1]] \\ C_2 &= [0, [4, -1], 14, [6, 1], 2, [2, -1]] \\ C_3 &= [16, [5, -1], 3, [5, 1]] \\ C_4 &= [1, [7, 0], 8, [7, 2]] \\ C_5 &= [4, [7, -1], 11, [7, 1]] \\ C_6 &= [10, [3, -1], 7, [3, 1]] \\ C'_0 &= [11, [5, 0], 6, [6, 2], 12, [4, 0], 8, [2, 1], 10, [1, 2]] \\ C'_1 &= [2, [4, 1], 16, [6, 0], 4, [2, 2]] \\ C'_2 &= [1, [4, 2], 5, [18, 0], 18, [18, -1]] \\ C'_3 &= [0, [9, 0], 9, [9, 2]] \\ C'_4 &= [17, [8, -1], 7, [8, 1]] \\ C'_5 &= [14, [1, -1], 15, [1, 1]] \\ \end{split}$$
 - $C_6' = [3, [8, 0], 13, [8, 2]]$
- 2-factor type [4, 4, 3, 2, 2, 2, 2]: three starters
 - $$\begin{split} C_0 &= [6, [1, 0], 5, [1, -1], 4, [8, -1], 14, [8, 2]] \\ C_1 &= [12, [1, 2], 13, [3, -1], 16, [2, -1], 0, [6, 0]] \\ C_2 &= [1, [8, 0], 11, [6, 1], 17, [2, 2]] \\ C_3 &= [3, [7, -1], 10, [7, 1]] \\ C_4 &= [9, [18, 0], 18, [18, 2]] \\ C_5 &= [2, [5, 0], 7, [5, 2]] \\ C_6 &= [8, [7, 0], 15, [7, 2]] \end{split}$$
- $C'_{0} = [1, [2, 1], 17, [4, 1], 13, [18, -1], 18, [18, 1]]$ $C'_{1} = [16, [2, 0], 14, [8, 1], 4, [6, 2], 10, [6, -1]]$ $C'_{2} = [2, [1, 1], 3, [3, 1], 6, [4, -1]]$ $C'_{3} = [0, [9, 0], 9, [9, 2]]$ $C'_{4} = [5, [3, 0], 8, [3, 2]]$ $C'_{5} = [12, [5, -1], 7, [5, 1]]$ $C'_{6} = [15, [4, 0], 11, [4, 2]$

• 2-factor type [4, 3, 3, 3, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [9, [7, 0], 16, [4, 2], 12, [5, 1], 7, [2, -1]] \\ C_1 &= [10, [8, 2], 0, [3, 1], 15, [5, 0]] \\ C_2 &= [1, [2, 2], 3, [1, 1], 2, [1, 0]] \\ C_3 &= [18, [18, 1], 13, [4, 0], 17, [18, 2]] \\ C_4 &= [6, [8, -1], 14, [8, 1]] \\ C_5 &= [11, [7, -1], 4, [7, 1]] \\ C_6 &= [5, [3, 0], 8, [3, 2]] \end{split}$$

• 2-factor type [3, 3, 3, 3, 3, 2, 2]: three starters

$$\begin{split} C_0 &= [10, [5, 1], 5, [3, 1], 2, [8, -1]]\\ C_1 &= [11, [7, 1], 4, [4, -1], 8, [3, -1]]\\ C_2 &= [0, [7, 0], 7, [6, 2], 1, [1, 1]]\\ C_3 &= [15, [1, 2], 14, [18, 0], 18, [18, -1]]\\ C_4 &= [13, [4, 1], 9, [8, 0], 17, [4, 2]]\\ C_5 &= [3, [5, 0], 16, [5, 2]]\\ C_6 &= [6, [6, -1], 12, [6, 1]] \end{split}$$

$$\begin{split} C_0' &= [14, [1, 2], 15, [2, 0], 17, [2, 1], 1, [5, -1]] \\ C_1' &= [10, [18, 0], 18, [18, -1], 4, [6, 2]] \\ C_2' &= [5, [7, 2], 12, [6, 0], 6, [1, -1]] \\ C_3' &= [11, [8, 0], 3, [5, 2], 8, [3, -1]] \\ C_4' &= [0, [9, 0], 9, [9, 2]] \\ C_5' &= [16, [4, -1], 2, [4, 1]] \\ C_6' &= [7, [6, -1], 13, [6, 1]] \end{split}$$

$$\begin{split} C_0' &= [8, [4, 0], 12, [7, 2], 1, [7, -1]] \\ C_1' &= [10, [1, -1], 11, [2, 0], 13, [3, 2]] \\ C_2' &= [15, [1, 0], 14, [18, 2], 18, [18, 1]] \\ C_3' &= [3, [5, -1], 16, [8, 2], 6, [3, 0]] \\ C_4' &= [5, [2, 2], 7, [8, 1], 17, [6, 0]] \\ C_5' &= [0, [9, 0], 9, [9, 2]] \\ C_6' &= [4, [2, -1], 2, [2, 1]] \end{split}$$

• 2-factor type [9, 2, 2, 2, 2, 2]: three starters $C_0 = [1, [6, -1], 7, [7, 0], 0, [2, 2], 16, [5, 0], 11, [5, -1], 6, [3, 2], 9, [7, 1], 2, [6, 0], 8, [7, 2]]$ $C_1 = [10, [4, 0], 14, [4, 2]]$ $C_2 = [4, [1, -1], 5, [1, 1]]$ $C_3 = [3, [8, 0], 13, [8, 2]]$ $C_4 = [12, [3, -1], 15, [3, 1]]$ $C_5 = [18, [18, -1], 17, [18, 1]]$ $C_6 = [11, [3, 0], 14, [18, 2], 18, [18, 0], 6, [5, 2], 1, [6, 1], 13, [7, -1], 2, [2, 0], 4, [6, 2], 16, [5, 1]]$ $C_1' = [0, [9, 0], 9, [9, 2]]$ $C_2' = [8, [1, 0], 7, [1, 2]]$ $C_3' = [5, [8, -1], 15, [8, 1]]$ $C_4' = [10, [2, -1], 12, [2, 1]]$ $C_5' = [3, [4, -1], 17, [4, 1]]$

• 2-factor type [8, 3, 2, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [10, [1, 1], 11, [3, 2], 14, [4, 0], 0, [1, 2], 1, [3, -1], 4, [1, 0], 3, [4, 2], 17, [7, 0]] \\ C_1 &= [9, [3, 1], 12, [4, 1], 16, [7, -1]] \\ C_2 &= [8, [2, 0], 6, [2, 2]] \\ C_3 &= [5, [8, 0], 13, [8, 2]] \\ C_4 &= [7, [8, -1], 15, [8, 1]] \\ C_5 &= [18, [18, -1], 2, [18, 1]] \end{split}$$

- $$\begin{split} C_0' &= [8, [2, 1], 6, [3, 0], 3, [4, -1], 17, [1, -1], 16, [5, 2], 11, [7, 1], 4, [18, 0], 18, [18, 2]] \\ C_1' &= [5, [2, -1], 7, [5, 0], 12, [7, 2]] \\ C_2' &= [0, [9, 0], 9, [9, 2]] \\ C_3' &= [10, [5, -1], 15, [5, 1]] \\ C_4' &= [2, [6, 0], 14, [6, 2]] \\ C_5' &= [1, [6, -1], 13, [6, 1]] \end{split}$$
- 2-factor type [7, 4, 2, 2, 2, 2]: three starters

.

$$\begin{split} &C_0 = [9, [2, -1], 7, [7, 2], 14, [18, 0], 18, [18, -1], 17, [1, 1], 0, [2, 1], 2, [7, 1]] \\ &C_1 = [12, [7, 0], 1, [8, 2], 11, [6, 1], 5, [7, -1]] \\ &C_2 = [15, [2, 0], 13, [2, 2]] \\ &C_3 = [6, [8, -1], 16, [8, 1]] \\ &C_4 = [4, [6, 0], 10, [6, 2]] \\ &C_5 = [3, [5, -1], 8, [5, 1]] \\ &C_0' = [2, [1, 0], 3, [1, 2], 4, [1, -1], 5, [8, 0], 15, [4, -1], 11, [18, 1], 18, [18, 2]] \\ &C_1' = [12, [4, 1], 8, [5, 0], 13, [6, -1], 7, [5, 2]] \\ &C_2' = [0, [9, 0], 9, [9, 2]] \\ &C_3' = [10, [4, 0], 6, [4, 2]] \\ &C_4' = [14, [3, -1], 17, [3, 1]] \\ &C_5' = [16, [3, 0], 1, [3, 2]] \\ &2\text{-factor type } [6, 5, 2, 2, 2, 2]: \text{ three starters} \\ &C_0 = [11, [5, 0], 16, [2, 2], 0, [4, 1], 14, [3, -1], 17, [7, 0], 6, [5, 2]] \\ &C_1 = [1, [8, 1], 9, [2, -1], 7, [3, 2], 10, [2, 0], 8, [7, -1]] \\ &C_2 = [18, [18, -1], 2, [18, 1]] \\ &C_3 = [4, [1, 0], 3, [1, 2]] \\ &C_4 = [15, [8, 0], 5, [8, 2]] \\ &C_5 = [12, [1, -1], 13, [1, 1]] \\ &C_0' = [7, [5, 1], 12, [18, 2], 18, [18, 0], 10, [3, 1], 13, [2, 1], 15, [8, -1]] \\ &C_1' = [3, [4, -1], 17, [7, 2], 6, [5, -1], 11, [3, 0], 14, [7, 1]] \\ &C_2' = [0, [9, 0], 9, [9, 2]] \\ &C_3' = [16, [6, -1], 4, [6, 1]] \\ &C_4' = [1, [4, 0], 5, [4, 2]] \\ &C_5' = [8, [6, 0], 2, [6, 2]] \\ \end{aligned}$$

• 2-factor type [7, 3, 3, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [3, [8, 2], 11, [3, 0], 8, [18, 1], 18, [18, 2], 0, [8, 0], 10, [8, -1], 2, [1, 1]] \\ C_1 &= [1, [3, -1], 16, [1, 2], 15, [4, 0]] \\ C_2 &= [7, [6, 1], 13, [4, 1], 17, [8, 1]] \\ C_3 &= [5, [7, -1], 12, [7, 1]] \\ C_4 &= [14, [5, 0], 9, [5, 2]] \\ C_5 &= [6, [2, -1], 4, [2, 1]] \end{split}$$

- $$\begin{split} C_0' &= [15, [3, 2], 12, [1, 0], 11, [1, -1], 10, [6, -1], 4, [3, 1], 7, [6, 2], 13, [2, 0]] \\ C_1' &= [14, [6, 0], 2, [4, -1], 16, [2, 2]] \\ C_2' &= [18, [18, -1], 5, [4, 2], 1, [18, 0]] \\ C_3' &= [0, [9, 0], 9, [9, 2]] \\ C_4' &= [8, [5, -1], 3, [5, 1]] \\ C_5' &= [17, [7, 0], 6, [7, 2]] \end{split}$$
- 2-factor type [6, 4, 3, 2, 2, 2]: three starters

.

$$\begin{split} C_0 &= [13, [7, 0], 6, [1, 1], 7, [5, -1], 2, [1, 2], 3, [4, 0], 17, [4, 2]] \\ C_1 &= [4, [8, -1], 14, [2, 2], 12, [2, 0], 10, [6, -1]] \\ C_2 &= [11, [5, 2], 16, [7, -1], 5, [6, 0]] \\ C_3 &= [18, [18, 0], 8, [18, 2]] \\ C_4 &= [0, [3, 0], 15, [3, 2]] \\ C_5 &= [1, [8, 0], 9, [8, 2]] \\ C_6 &= [11, [3, 1], 8, [1, 0], 7, [8, 1], 15, [18, 1], 18, [18, -1], 5, [6, 2]] \\ C_1' &= [17, [3, -1], 14, [7, 2], 3, [1, -1], 4, [5, 0]] \\ C_2' &= [6, [7, 1], 13, [6, 1], 1, [5, 1]] \\ C_3' &= [0, [9, 0], 9, [9, 2]] \\ C_4' &= [2, [4, -1], 16, [4, 1]] \\ C_5' &= [12, [2, -1], 10, [2, 1]] \\ 2\text{-factor type } [5, 5, 3, 2, 2, 2]: \text{ three starters} \\ C_0 &= [12, [8, -1], 2, [2, 1], 0, [7, -1], 7, [3, -1], 10, [2, -1]] \\ C_1 &= [13, [7, 2], 6, [2, 0], 8, [8, 1], 16, [1, 2], 17, [4, 0]] \\ C_2 &= [3, [7, 1], 14, [5, 0], 1, [2, 2]] \\ C_3 &= [5, [6, 0], 11, [6, 2]] \\ C_4 &= [4, [5, -1], 9, [5, 1]] \\ C_5 &= [18, [18, 0], 15, [18, 2]] \\ C_0' &= [12, [7, 0], 1, [4, 2], 15, [1, 0], 14, [3, 2], 11, [1, -1]] \\ C_1' &= [2, [1, 1], 3, [18, 1], 18, [18, -1], 7, [8, 2], 17, [3, 0]] \\ C_2' &= [8, [5, 2], 13, [8, 0], 5, [3, 1]] \\ C_3' &= [0, [9, 0], 9, [9, 2]] \\ C_4' &= [6, [4, -1], 10, [4, 1]] \\ C_5' &= [16, [6, -1], 4, [6, 1]] \\ \end{split}$$

• 2-factor type [5, 4, 4, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [15, [18, 1], 18, [18, -1], 8, [1, 2], 7, [5, 0], 2, [5, -1]] \\ C_1 &= [14, [5, 2], 9, [4, 1], 5, [1, 0], 6, [8, 1]] \\ C_2 &= [1, [2, -1], 3, [8, 0], 13, [3, 1], 16, [3, 2]] \\ C_3 &= [4, [4, 0], 0, [4, 2]] \\ C_4 &= [10, [7, 0], 17, [7, 2]] \\ C_5 &= [11, [1, -1], 12, [1, 1]] \end{split}$$

- $$\begin{split} C_0' &= [7, [4, -1], 3, [7, 1], 10, [8, -1], 2, [18, 2], 18, [18, 0]] \\ C_1' &= [15, [3, 0], 12, [7, -1], 5, [6, 2], 17, [2, 1]] \\ C_2' &= [4, [3, -1], 1, [5, 1], 6, [8, 2], 16, [6, 0]] \\ C_3' &= [0, [9, 0], 9, [9, 2]] \\ C_4' &= [14, [6, -1], 8, [6, 1]] \\ C_5' &= [13, [2, 0], 11, [2, 2]] \end{split}$$
- 2-factor type [6, 3, 3, 3, 2, 2]: three starters

$$\begin{split} C_0 &= [1, [7, 1], 12, [3, 0], 9, [7, 2], 16, [1, 0], 17, [2, -1], 15, [4, 2]] \\ C_1 &= [13, [1, -1], 14, [4, 1], 10, [3, -1]] \\ C_2 &= [7, [3, 2], 4, [18, 0], 18, [18, -1]] \\ C_3 &= [0, [6, 2], 6, [1, 1], 5, [5, 0]] \\ C_4 &= [2, [6, -1], 8, [6, 1]] \\ C_5 &= [11, [8, -1], 3, [8, 1]] \\ C_0' &= [10, [5, 2], 15, [4, 0], 11, [5, 1], 16, [4, -1], 12, [8, 2], 4, [6, 0]] \\ C_1' &= [7, [7, -1], 14, [8, 0], 6, [1, 2]] \\ C_2' &= [8, [5, -1], 3, [2, 1], 5, [3, 1]] \\ C_3' &= [13, [18, 1], 18, [18, 2], 2, [7, 0]] \\ C_4' &= [0, [9, 0], 9, [9, 2]] \\ C_5' &= [1, [2, 0], 17, [2, 2]] \\ 2\text{-factor type } [5, 4, 3, 3, 2, 2]: \text{ three starters} \\ C_0 &= [17, [8, 2], 7, [7, 0], 0, [4, 2], 14, [4, 0], 10, [7, 1]] \\ C_1 &= [9, [4, -1], 5, [18, 1], 18, [18, 2], 12, [3, 0]] \\ C_2 &= [15, [4, 1], 11, [8, 1], 3, [6, 1]] \\ C_3 &= [1, [7, -1], 8, [2, 1], 6, [5, 1]] \\ C_4 &= [16, [3, -1], 13, [3, 1]] \\ C_5 &= [2, [2, 0], 4, [2, 2]] \\ C_0' &= [14, [18, -1], 18, [18, 0], 17, [1, 2], 16, [5, 0], 11, [3, 2]] \\ C_1' &= [2, [1, 0], 3, [6, -1], 15, [8, -1], 7, [5, 2]] \\ C_2' &= [1, [7, 2], 8, [5, -1], 13, [6, 0]] \\ C_3' &= [10, [2, -1], 12, [8, 0], 4, [6, 2]] \\ C_4' &= [0, [9, 0], 9, [9, 2]] \\ C_5' &= [6, [1, -1], 5, [1, 1]] \\ 2\text{-factor type } [4, 4, 4, 3, 2, 2]: \text{ three starters} \\ \end{split}$$

$$\begin{split} C_0 &= [8, [2, 0], 10, [5, 1], 15, [8, 2], 5, [3, -1]] \\ C_1 &= [6, [18, 0], 18, [18, -1], 1, [3, 2], 16, [8, -1]] \\ C_2 &= [3, [1, -1], 2, [7, 1], 9, [2, 2], 11, [8, 0]] \\ C_3 &= [4, [8, 1], 14, [3, 0], 17, [5, 2]] \\ C_4 &= [7, [7, 0], 0, [7, 2]] \\ C_5 &= [12, [1, 0], 13, [1, 2]] \end{split}$$

$$\begin{split} C_0' &= [3, [5, -1], 16, [2, -1], 14, [1, 1], 15, [6, 1]] \\ C_1' &= [2, [5, 0], 7, [6, -1], 1, [18, 2], 18, [18, 1]] \\ C_2' &= [11, [7, -1], 4, [4, 2], 8, [3, 1], 5, [6, 0]] \\ C_3' &= [6, [6, 2], 12, [2, 1], 10, [4, 0]] \\ C_4' &= [0, [9, 0], 9, [9, 2]] \\ C_5' &= [13, [4, -1], 17, [4, 1]] \end{split}$$

• 2-factor type [5, 3, 3, 3, 3, 2]: three starters

$$\begin{split} C_0 &= [9, [5, 2], 14, [4, 0], 0, [3, 2], 15, [5, -1], 2, [7, 0]] \\ C_1 &= [4, [1, 1], 5, [2, 2], 7, [3, 0]] \\ C_2 &= [6, [3, 1], 3, [4, 1], 17, [7, -1]] \\ C_3 &= [12, [4, -1], 8, [5, 1], 13, [1, -1]] \\ C_4 &= [18, [18, 0], 10, [6, 1], 16, [18, 2]] \\ C_5 &= [1, [8, 0], 11, [8, 2]] \\ C_5 &= [1, [8, 0], 11, [8, 2]] \\ C_6' &= [8, [3, -1], 11, [1, 0], 12, [1, 2], 13, [2, -1], 15, [7, 1]] \\ C_1' &= [6, [4, 2], 2, [6, 0], 14, [8, -1]] \\ C_2' &= [10, [6, -1], 4, [18, 1], 18, [18, -1]] \\ C_3' &= [1, [6, 2], 7, [8, 1], 17, [2, 0]] \\ C_4' &= [3, [2, 1], 5, [7, 2], 16, [5, 0]] \\ C_5' &= [0, [9, 0], 9, [9, 2]] \end{split}$$

• 2-factor type [4, 4, 3, 3, 3, 2]: three starters

$$\begin{split} C_0 &= [11, [2, 0], 13, [1, -1], 12, [2, 2], 14, [3, 1]] & C'_0 &= [10, [7, 2], 3, [6, 0], 15, [5, 1], 2, [8, 1]] \\ C_1 &= [10, [7, 1], 3, [4, 0], 7, [2, 1], 9, [1, 2]] & C'_1 &= [14, [3, -1], 11, [18, 1], 18, [18, -1], 1, [5, -1]] \\ C_2 &= [4, [2, -1], 2, [18, 2], 18, [18, 0]] & C'_2 &= [6, [6, 1], 12, [4, 2], 16, [8, 0]] \\ C_3 &= [15, [7, -1], 8, [8, 2], 0, [3, 0]] & C'_3 &= [17, [8, -1], 7, [6, -1], 13, [4, -1]] \\ C_4 &= [16, [1, 1], 17, [6, 2], 5, [7, 0]] & C'_4 &= [8, [4, 1], 4, [1, 0], 5, [3, 2]] \\ C_5 &= [6, [5, 0], 1, [5, 2]] & C'_5 &= [0, [9, 0], 9, [9, 2]] \end{split}$$

• 2-factor type [11, 2, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [8, [4, 0], 12, [2, 1], 14, [1, 2], 13, [7, 0], 6, [7, 2], 17, [1, -1], 0, [3, 0], 15, [5, 2], 2, [1, 0], 3, [5, -1], 16, \\ & [8, 2]] \\ C_1 &= [1, [8, -1], 9, [8, 1]] \\ C_2 &= [18, [18, -1], 11, [18, 1]] \\ C_3 &= [4, [18, -1], 12, [18, 1]] \\ C_4 &= [4, [18, -1], 12, [18, 1]] \\ C_5 &= [4, [18, -1], 12, [18, 1]] \\ C_6 &= [4, [18, -1], 12, [18, 1]] \\ C_7 &= [4, [18, -1], 12, [18, 1]] \\ C_8 &= [4, [18, -1], 12, [18, 1]] \\ C_8 &= [4, [18, -1], 12, [18, 1]] \\ C_9 &= [4, [18, -1], 12, [18, -1], 12, [18, -1]] \\ C_9 &= [4, [18, -1], 12, [18, -1], 12, [18, -1]] \\ C_9 &= [4, [18, -1], 12, [18, -1], 12, [18, -1]] \\ C_9 &= [4, [18, -1], 12, [18, -1]] \\ C_9 &= [4, [18, -1], 12, [18, -1]] \\ C_9 &= [4, [18, -1], 12, [18, -1]] \\ C_9 &= [4, [18, -1], 12, [18, -1]] \\ C_9 &= [4, [18, -1], 12, [18, -1]] \\ C_9 &= [4, [18, -1], 12, [18, -1]] \\ C_9 &= [4, [18$$

- $C_3 = [4, [6, 0], 10, [6, 2]]$
- $C_4 = [7, [2, 0], 5, [2, 2]]$
- $$\begin{split} C_0' = [17, [3, 2], 14, [1, 1], 13, [8, 0], 5, [6, 1], 11, [5, 1], 16, [4, 2], 2, [5, 0], 7, [6, -1], 1, [7, -1], 12, [2, -1], 10, \\ [7, 1]] \end{split}$$
- $C_1' = [0, [9, 0], 9, [9, 2]]$
- $C_2' = [15, [18, 0], 18, [18, 2]]$
- $C'_3 = [8, [4, -1], 4, [4, 1]]$
- $C'_4 = [6, [3, -1], 3, [3, 1]]$
- 2-factor type [10, 3, 2, 2, 2]: three starters
 - $C_0 = [2, [4, 2], 16, [3, -1], 1, [18, 0], 18, [18, 2], 13, [6, 1], 7, [4, 0], 3, [3, 2], 6, [8, 0], 14, [6, 2], 8, [6, 0]]$
 - $C_1 = [11, [7, 1], 0, [6, -1], 12, [1, -1]]$
 - $C_2 = [15, [8, -1], 5, [8, 1]]$
 - $C_3 = [4, [5, 0], 9, [5, 2]]$
 - $C_4 = [10, [7, 0], 17, [7, 2]]$

$$\begin{split} C_0' &= [14, [2, 0], 16, [1, 1], 17, [4, 1], 3, [3, 1], 6, [5, -1], 1, [7, -1], 12, [8, 2], 4, [4, -1], 8, [18, -1], 18, [18, 1]] \\ C_1' &= [2, [3, 0], 5, [2, 2], 7, [5, 1]] \\ C_2' &= [0, [9, 0], 9, [9, 2]] \\ C_3' &= [13, [2, -1], 15, [2, 1]] \\ C_4' &= [11, [1, 0], 10, [1, 2]] \end{split}$$

• 2-factor type [9, 4, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [13, [3, 2], 16, [5, 0], 11, [4, -1], 7, [3, 1], 10, [5, 2], 15, [4, 0], 1, [5, -1], 14, [6, 2], 2, [7, 0]] \\ C_1 &= [17, [5, 1], 12, [3, 0], 9, [18, 2], 18, [18, 1]] \\ C_2 &= [0, [8, -1], 8, [8, 1]] \\ C_3 &= [6, [1, -1], 5, [1, 1]] \\ C_4 &= [4, [1, 0], 3, [1, 2]] \\ C_6' &= [11, [6, 0], 5, [2, 2], 3, [2, 0], 1, [4, 2], 15, [18, 0], 18, [18, -1], 7, [7, 1], 14, [8, 2], 4, [7, -1]] \\ C_1' &= [17, [7, 2], 6, [8, 0], 16, [3, -1], 13, [4, 1]] \\ C_2' &= [0, [9, 0], 9, [9, 2]] \\ C_3' &= [8, [6, -1], 2, [6, 1]] \\ C_4' &= [10, [2, -1], 12, [2, 1]] \end{split}$$

• 2-factor type [8, 5, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [7, [3, 0], 4, [4, -1], 0, [2, -1], 16, [3, 2], 1, [5, 0], 14, [6, 2], 8, [1, 0], 9, [2, 2]] \\ C_1 &= [6, [18, -1], 18, [18, 0], 15, [2, 1], 17, [3, 1], 2, [4, 2]] \\ C_2 &= [12, [1, -1], 13, [1, 1]] \\ C_3 &= [5, [5, -1], 10, [5, 1]] \\ C_4 &= [3, [8, -1], 11, [8, 1]] \\ C'_0 &= [11, [6, 0], 17, [1, 2], 16, [2, 0], 14, [18, 2], 18, [18, 1], 7, [6, -1], 13, [8, 0], 3, [8, 2]] \\ C'_1 &= [2, [6, 1], 8, [3, -1], 5, [5, 2], 10, [4, 1], 6, [4, 0]] \\ C'_2 &= [0, [9, 0], 9, [9, 2]] \\ C'_3 &= [1, [7, 0], 12, [7, 2]] \\ C'_4 &= [4, [7, -1], 15, [7, 1]] \end{split}$$

• 2-factor type [7, 6, 2, 2, 2]: three starters

$$\begin{split} C_0 &= [6, [7, -1], 13, [5, 0], 0, [1, 2], 1, [2, 0], 17, [3, 2], 2, [3, -1], 5, [1, -1]] \\ C_1 &= [12, [3, 0], 9, [5, 2], 14, [2, -1], 16, [6, 0], 10, [2, 2], 8, [4, -1]] \\ C_2 &= [4, [18, 0], 18, [18, 2]] \\ C_3 &= [3, [8, -1], 11, [8, 1]] \\ C_4 &= [7, [8, 0], 15, [8, 2]] \\ C'_0 &= [11, [7, 1], 4, [6, 1], 16, [1, 0], 15, [7, 2], 8, [1, 1], 7, [6, -1], 13, [2, 1]] \\ C'_1 &= [2, [18, -1], 18, [18, 1], 10, [4, 1], 6, [3, 1], 3, [7, 0], 14, [6, 2]] \\ C'_2 &= [0, [9, 0], 9, [9, 2]] \\ C'_3 &= [5, [4, 0], 1, [4, 2]] \\ C'_4 &= [12, [5, -1], 17, [5, 1]] \end{split}$$

• 2-factor type [9, 3, 3, 2, 2]: three starters

$$\begin{split} &C_0 = [2, [2, -1], 4, [3, -1], 7, [6, 0], 1, [8, 2], 9, [3, 0], 6, [6, -1], 0, [2, 2], 16, [5, 0], 3, [1, 2]] \\ &C_1 = [10, [2, 0], 12, [2, 1], 14, [4, 2]] \\ &C_2 = [8, [3, 1], 11, [18, 2], 18, [18, 0]] \\ &C_3 = [15, [8, -1], 5, [8, 1]] \\ &C_4 = [17, [4, -1], 13, [4, 1]] \\ &C_0' = [14, [18, 1], 18, [18, -1], 1, [3, 2], 4, [7, 1], 15, [5, 1], 10, [1, 0], 11, [6, 2], 17, [7, -1], 6, [8, 0]] \\ &C_1' = [16, [5, 2], 3, [1, 1], 2, [4, 0]] \\ &C_2' = [13, [5, -1], 8, [1, -1], 7, [6, 1]] \\ &C_3' = [0, [9, 0], 9, [9, 2]] \\ &C_4' = [12, [7, 0], 5, [7, 2]] \\ &2\text{-factor type } [8, 4, 3, 2, 2]: \text{ three starters} \\ &C_0 = [17, [1, 1], 0, [7, -1], 11, [3, 1], 14, [6, 1], 2, [4, 1], 6, [4, 2], 10, [2, 0], 12, [5, 1]] \\ &C_1 = [4, [4, 0], 8, [18, 1], 18, [18, -1], 3, [1, 2]] \\ &C_2 = [13, [6, 0], 1, [6, 2], 7, [6, -1]] \\ &C_3 = [9, [7, 0], 16, [7, 2]] \\ &C_4 = [15, [8, -1], 5, [8, 1]] \\ &C_0' = [16, [1, -1], 17, [4, -1], 3, [18, 0], 18, [18, 2], 4, [1, 0], 5, [7, 1], 12, [2, 1], 14, [2, 2]] \\ \end{split}$$

$$C'_1 = [6, [5, 2], 1, [8, 0], 11, [3, -1], 8, [2, -1]]$$

 $C_1 = [0, [5, 2], 1, [8, 0], 11, [5, -1], c$ $C_2 = [7, [8, 2], 15, [5, -1], 2, [5, 0]]$

 $C_3^\prime = [0, [9, 0], 9, [9, 2]]$

$$C'_4 = [13, [3, 0], 10, [3, 2]]$$

• 2-factor type [7, 5, 3, 2, 2]: three starters

$$\begin{split} C_0 &= [17, [4, 2], 3, [6, 1], 15, [3, 0], 0, [6, 2], 6, [1, 0], 7, [2, 2], 5, [6, 0]] \\ C_1 &= [4, [18, 1], 18, [18, -1], 10, [1, 2], 9, [7, 1], 2, [2, 0]] \\ C_2 &= [11, [3, -1], 14, [5, 0], 1, [8, 2]] \\ C_3 &= [12, [4, -1], 16, [4, 1]] \\ C_4 &= [13, [5, -1], 8, [5, 1]] \\ C_6' &= [4, [6, -1], 16, [1, -1], 15, [7, -1], 8, [5, 2], 13, [1, 1], 12, [8, 0], 2, [2, 1]] \\ C_1' &= [7, [4, 0], 11, [8, 1], 1, [2, -1], 17, [7, 2], 10, [3, 1]] \\ C_2' &= [3, [3, 2], 6, [8, -1], 14, [7, 0]] \\ C_3' &= [0, [9, 0], 9, [9, 2]] \\ C_4' &= [5, [18, 0], 18, [18, 2]] \end{split}$$

• 2-factor type [6, 6, 3, 2, 2]: three starters

$$\begin{split} C_0 &= [9, [4, 0], 5, [6, 1], 11, [7, 2], 4, [2, 1], 2, [5, 0], 7, [2, 2]] \\ C_1 &= [16, [1, 0], 17, [2, -1], 15, [7, -1], 8, [8, -1], 0, [6, 2], 12, [4, -1]] \\ C_2 &= [18, [18, -1], 10, [3, 2], 13, [18, 0]] \\ C_3 &= [1, [5, -1], 14, [5, 1]] \\ C_4 &= [6, [3, -1], 3, [3, 1]] \end{split}$$

- $$\begin{split} C_0' &= [8, [18, 1], 18, [18, 2], 1, [8, 1], 11, [7, 0], 4, [1, 2], 5, [3, 0]] \\ C_1' &= [3, [7, 1], 14, [6, 0], 2, [5, 2], 15, [2, 0], 17, [8, 2], 7, [4, 1]] \\ C_2' &= [16, [8, 0], 6, [4, 2], 10, [6, -1]] \\ C_3' &= [0, [9, 0], 9, [9, 2]] \\ C_4' &= [13, [1, -1], 12, [1, 1]] \end{split}$$
- 2-factor type [7, 4, 4, 2, 2]: three starters

$$\begin{split} C_0 &= [0, [3, 1], 3, [5, 2], 8, [3, -1], 11, [4, 0], 7, [5, 1], 12, [2, 1], 14, [4, 1]] \\ C_1 &= [17, [2, 2], 15, [4, -1], 1, [1, -1], 2, [3, 0]] \\ C_2 &= [5, [7, 2], 16, [6, 0], 4, [18, 2], 18, [18, 0]] \\ C_3 &= [9, [1, 0], 10, [1, 2]] \\ C_4 &= [13, [7, -1], 6, [7, 1]] \\ C'_0 &= [14, [3, 2], 17, [6, -1], 5, [8, 0], 13, [6, 2], 7, [5, 0], 2, [8, 2], 12, [2, 0]] \\ C'_1 &= [11, [4, 2], 15, [7, 0], 4, [6, 1], 10, [1, 1]] \\ C'_2 &= [18, [18, -1], 8, [5, -1], 3, [2, -1], 1, [18, 1]] \\ C'_3 &= [0, [9, 0], 9, [9, 2]] \\ C'_4 &= [6, [8, -1], 16, [8, 1]] \end{split}$$

• 2-factor type [6, 5, 4, 2, 2]: three starters

$$\begin{split} C_0 &= [8, [1, 1], 7, [3, -1], 10, [8, 0], 0, [3, 1], 3, [2, -1], 1, [7, 2]] \\ C_1 &= [9, [7, 0], 2, [7, -1], 13, [3, 2], 16, [5, 0], 11, [2, 2]] \\ C_2 &= [6, [18, 0], 18, [18, 2], 15, [3, 0], 12, [6, 2]] \\ C_3 &= [5, [6, -1], 17, [6, 1]] \\ C_4 &= [14, [8, -1], 4, [8, 1]] \\ C'_0 &= [15, [7, 1], 4, [5, 2], 17, [1, 0], 16, [4, 2], 12, [1, -1], 13, [2, 0]] \\ C'_1 &= [3, [5, -1], 8, [6, 0], 2, [18, 1], 18, [18, -1], 11, [8, 2]] \\ C'_2 &= [7, [2, 1], 5, [4, 0], 1, [5, 1], 6, [1, 2]] \\ C'_3 &= [0, [9, 0], 9, [9, 2]] \\ C'_4 &= [14, [4, -1], 10, [4, 1]] \end{split}$$

• 2-factor type [5, 5, 5, 2, 2]: three starters

$$\begin{split} C_0 &= [3, [4, -1], 17, [8, 1], 7, [5, 1], 12, [8, 2], 2, [1, 0]] \\ C_1 &= [6, [5, -1], 1, [3, 2], 4, [8, 0], 14, [18, 1], 18, [18, -1]] \\ C_2 &= [10, [2, 1], 8, [8, -1], 16, [7, 0], 5, [6, 2], 11, [1, 1]] \\ C_3 &= [9, [6, -1], 15, [6, 1]] \\ C_4 &= [0, [5, 0], 13, [5, 2]] \\ C'_0 &= [15, [18, 0], 18, [18, 2], 17, [4, 0], 3, [2, -1], 1, [4, 2]] \\ C'_1 &= [16, [3, 1], 13, [7, -1], 2, [6, 0], 8, [3, -1], 5, [7, 2]] \\ C'_2 &= [11, [1, 2], 10, [4, 1], 6, [1, -1], 7, [3, 0], 4, [7, 1]] \\ C'_3 &= [0, [9, 0], 9, [9, 2]] \\ C'_4 &= [12, [2, 0], 14, [2, 2]] \end{split}$$

• 2-factor type [8, 3, 3, 3, 2]: three starters

$$\begin{split} C_0 &= [14, [6, 1], 8, [5, 0], 13, [7, 2], 6, [3, 0], 9, [2, -1], 7, [3, 2], 10, [1, -1], 11, [3, -1]] \\ C_1 &= [0, [6, -1], 12, [5, 2], 17, [1, 0]] \\ C_2 &= [3, [1, 1], 4, [18, 1], 18, [18, -1]] \\ C_3 &= [1, [1, 2], 2, [4, 0], 16, [3, 1]] \\ C_4 &= [15, [8, 0], 5, [8, 2]] \\ C_0' &= [5, [7, 0], 12, [5, -1], 7, [18, 2], 18, [18, 0], 6, [8, -1], 16, [4, 2], 2, [7, 1], 13, [8, 1]] \\ C_1' &= [10, [6, 2], 4, [4, -1], 8, [2, 0]] \\ C_2' &= [15, [4, 1], 11, [6, 0], 17, [2, 2]] \\ C_3' &= [1, [2, 1], 3, [7, -1], 14, [5, 1]] \\ C_4' &= [0, [9, 0], 9, [9, 2]] \\ \bullet & 2\text{-factor type } [7, 4, 3, 3, 2]: \text{ three starters} \\ C_0 &= [2, [4, 1], 16, [2, 0], 0, [6, -1], 12, [3, -1], 9, [4, -1], 5, [3, 1], 8, [6, 2]] \\ C_1 &= [3, [4, 0], 17, [4, 2], 13, [2, 1], 11, [8, 1]] \\ C_2 &= [6, [5, 2], 1, [5, 0], 14, [8, -1]] \\ C_3 &= [7, [8, 2], 15, [7, -1], 4, [3, 0]] \\ C_4 &= [18, [18, 0], 10, [18, 2]] \\ C_0' &= [13, [6, 0], 1, [7, 1], 8, [6, 1], 14, [7, 2], 3, [1, -1], 4, [7, 0], 11, [2, 2]] \\ C_1' &= [12, [5, 1], 7, [1, 1], 6, [18, -1], 18, [18, 1]] \\ C_2' &= [17, [1, 0], 16, [1, 2], 15, [2, -1]] \\ C_3' &= [2, [8, 0], 10, [5, -1], 5, [3, 2]] \\ C_4' &= [0, [9, 0], 9, [9, 2]] \\ \bullet & 2\text{-factor type } [6, 5, 3, 3, 2]: \text{ three starters} \\ C_0 &= [3, [3, 0], 0, [5, 2], 5, [6, 0], 17, [2, -1], 15, [8, -1], 7, [4, 2]] \\ C_1' &= [13, [7, 0], 6, [5, -1], 1, [3, 2], 16, [2, 1], 14, [1, 1]] \\ \end{split}$$

$$\begin{aligned} C_0 &= [3, [3, 0], 0, [3, 2], 3, [0, 0], 17, [2, -1], 13, [8, -1], 7, [4, 2]] \\ C_1 &= [13, [7, 0], 6, [5, -1], 1, [3, 2], 16, [2, 1], 14, [1, 1]] \\ C_2 &= [12, [3, -1], 9, [1, 2], 8, [4, 0]] \\ C_3 &= [10, [8, 2], 2, [18, 0], 18, [18, -1]] \\ C_4 &= [4, [7, -1], 11, [7, 1]] \\ C_6 &= [15, [8, 1], 5, [6, 2], 11, [6, -1], 17, [3, 1], 14, [4, 1], 10, [5, 0]] \\ C_1' &= [4, [2, 0], 6, [2, 2], 8, [8, 0], 16, [18, 2], 18, [18, 1]] \\ C_2' &= [7, [5, 1], 2, [1, -1], 3, [4, -1]] \\ C_3' &= [13, [1, 0], 12, [7, 2], 1, [6, 1]] \\ C_4' &= [0, [9, 0], 9, [9, 2]] \end{aligned}$$

• 2-factor type [6, 4, 4, 3, 2]: three starters

$$\begin{split} C_0 &= [1, [7, 1], 8, [4, 1], 12, [2, 2], 10, [8, 0], 2, [3, -1], 17, [2, 1]] \\ C_1 &= [14, [7, 2], 7, [6, -1], 13, [18, -1], 18, [18, 0]] \\ C_2 &= [0, [3, 1], 3, [6, 2], 15, [1, -1], 16, [2, 0]] \\ C_3 &= [5, [6, 0], 11, [5, -1], 6, [1, 2]] \\ C_4 &= [4, [5, 0], 9, [5, 2]] \end{split}$$

- $$\begin{split} C_0' &= [3, [8, -1], 11, [6, 1], 5, [1, 0], 6, [8, 2], 16, [8, 1], 8, [5, 1]] \\ C_1' &= [2, [2, -1], 4, [3, 0], 7, [18, 1], 18, [18, 2]] \\ C_2' &= [13, [1, 1], 14, [4, 2], 10, [7, -1], 17, [4, 0]] \\ C_3' &= [12, [7, 0], 1, [4, -1], 15, [3, 2]] \\ C_4' &= [0, [9, 0], 9, [9, 2]] \end{split}$$
- 2-factor type [5, 5, 4, 3, 2]: three starters

$$\begin{split} C_0 &= [12, [2, 1], 10, [8, 0], 0, [4, -1], 14, [5, 2], 1, [7, 1]] \\ C_1 &= [2, [7, 0], 9, [18, 1], 18, [18, -1], 4, [1, 2], 3, [1, 1]] \\ C_2 &= [5, [7, 2], 16, [1, 0], 17, [6, 2], 11, [6, 0]] \\ C_3 &= [7, [8, 1], 15, [2, -1], 13, [6, -1]] \\ C_4 &= [6, [2, 0], 8, [2, 2]] \\ C_0' &= [3, [8, -1], 13, [8, 2], 5, [4, 0], 1, [3, -1], 16, [5, 1]] \\ C_1' &= [15, [3, 0], 12, [1, -1], 11, [3, 2], 14, [6, 1], 8, [7, -1]] \\ C_2' &= [7, [5, 0], 2, [4, 1], 6, [4, 2], 10, [3, 1]] \\ C_3' &= [17, [18, 2], 18, [18, 0], 4, [5, -1]] \\ C_4' &= [0, [9, 0], 9, [9, 2]] \end{split}$$

- 2-factor type [5, 4, 4, 4, 2]: three starters
 - $$\begin{split} C_0 &= [6, [6, 2], 12, [3, 0], 15, [1, 1], 16, [18, 1], 18, [18, -1]] \\ C_1 &= [8, [3, 1], 5, [6, 0], 17, [4, 2], 13, [5, 1]] \\ C_2 &= [7, [7, 1], 14, [4, 1], 0, [3, 2], 3, [4, 0]] \\ C_3 &= [10, [1, 2], 9, [8, 1], 1, [8, 0], 11, [1, -1]] \\ C_4 &= [2, [2, 0], 4, [2, 2]] \\ C_0' &= [5, [5, -1], 10, [7, 0], 3, [2, -1], 1, [7, -1], 12, [7, 2]] \\ C_1' &= [14, [3, -1], 17, [5, 0], 4, [2, 1], 6, [8, 2]] \\ C_2' &= [15, [5, 2], 2, [6, -1], 8, [8, -1], 16, [1, 0]] \\ C_3' &= [18, [18, 2], 13, [6, 1], 7, [4, -1], 11, [18, 0]] \\ C_4' &= [0, [9, 0], 9, [9, 2]] \end{split}$$
- 2-factor type [13, 2, 2, 2]: three starters
 - $$\begin{split} C_0 &= [6, [8, 2], 14, [3, 0], 11, [4, 2], 15, [7, 0], 8, [1, 2], 9, [8, 1], 1, [1, 0], 0, [3, 2], 3, [2, 0], 5, [7, 2], 12, [2, 1], 10, \\ &= [8, 0], 2, [4, 1]] \\ C_1 &= [18, [18, 0], 16, [18, 2]] \\ C_2 &= [4, [5, 0], 17, [5, 2]] \\ C_3 &= [7, [6, -1], 13, [6, 1]] \\ C_0' &= [16, [4, 0], 12, [1, -1], 11, [5, -1], 6, [2, 2], 4, [7, 1], 15, [4, -1], 1, [7, -1], 8, [5, 1], 3, [2, -1], 5, [8, -1], \\ &= 13, [18, -1], 18, [18, 1], 17, [1, 1]] \\ C_1' &= [0, [9, 0], 9, [9, 2]] \\ C_2' &= [2, [6, 0], 14, [6, 2]] \\ C_3' &= [7, [3, -1], 10, [3, 1]] \end{split}$$

• 2-factor type [12, 3, 2, 2]: three starters

$$\begin{split} &C_0 = [10, [3, 2], 7, [2, 0], 9, [4, 2], 5, [6, 1], 17, [3, 1], 14, [6, 0], 8, [5, 2], 13, [2, -1], 15, [1, 0], 16, [4, 1], 2, [2, 2], \\ &0, [8, 0]] \\ &C_1 = [12, [6, 2], 6, [5, 1], 1, [7, 0]] \\ &C_2 = [4, [18, 0], 18, [18, 2]] \\ &C_3 = [11, [8, -1], 3, [8, 1]] \\ &C_0' = [5, [8, 2], 15, [2, 1], 13, [3, 0], 10, [1, 2], 11, [6, -1], 17, [3, -1], 2, [5, 0], 7, [1, -1], 6, [5, -1], 1, [7, 2], 8, \\ &[4, 0], 4, [1, 1]] \\ &C_1' = [12, [18, 1], 18, [18, -1], 16, [4, -1]] \\ &C_2' = [0, [9, 0], 9, [9, 2]] \\ &C_3' = [3, [7, -1], 14, [7, 1]] \\ &2\text{-factor type } [11, 4, 2, 2]: \text{ three starters} \\ &C_0 = [1, [1, 0], 0, [8, -1], 10, [4, 2], 14, [1, -1], 15, [4, 0], 11, [5, 2], 6, [2, 0], 4, [6, 2], 16, [4, 1], 12, [5, 1], 7, \\ &[6, 1]] \\ &C_1 = [9, [4, -1], 13, [7, -1], 2, [3, 1], 17, [8, 1]] \\ &C_2 = [5, [3, 0], 8, [3, 2]] \\ &C_3 = [18, [18, 0], 3, [18, 2]] \\ &C_0' = [1, [1, 2], 2, [2, -1], 4, [5, 0], 17, [2, 2], 15, [7, 1], 8, [6, 0], 14, [3, -1], 11, [6, -1], 5, [7, 2], 16, [8, 0], 6, \\ &[5, -1]] \\ &C_1' = [12, [2, 1], 10, [7, 0], 3, [8, 2], 13, [1, 1]] \\ &C_2' = [0, [9, 0], 9, [9, 2]] \end{aligned}$$

- $C'_3 = [18, [18, -1], 7, [18, 1]]$
- 2-factor type [10, 5, 2, 2]: three starters

$$\begin{split} C_0 &= [13, [4, 2], 9, [8, -1], 17, [4, -1], 3, [7, 0], 14, [6, 2], 2, [2, 0], 4, [3, 1], 7, [7, -1], 0, [8, 1], 8, [5, 1]] \\ C_1 &= [18, [18, 2], 11, [5, 0], 6, [5, 2], 1, [7, 1], 12, [18, 0]] \\ C_2 &= [10, [6, -1], 16, [6, 1]] \\ C_3 &= [15, [8, 0], 5, [8, 2]] \\ C_0' &= [4, [7, 2], 15, [4, 1], 11, [2, -1], 13, [6, 0], 7, [18, 1], 18, [18, -1], 10, [2, 2], 12, [4, 0], 8, [5, -1], 3, [1, 1]] \\ C_1' &= [1, [3, 2], 16, [2, 1], 14, [3, -1], 17, [3, 0], 2, [1, -1]] \\ C_2' &= [0, [9, 0], 9, [9, 2]] \\ C_3' &= [5, [1, 0], 6, [1, 2]] \end{split}$$

• 2-factor type [9, 6, 2, 2]: three starters

$$\begin{split} C_0 &= [0, [5, 1], 5, [7, 2], 16, [3, -1], 1, [18, -1], 18, [18, 1], 7, [6, -1], 13, [4, 0], 17, [5, -1], 12, [6, 1]] \\ C_1 &= [2, [5, 0], 15, [1, 2], 14, [3, 1], 11, [2, 0], 9, [1, 1], 10, [8, 2]] \\ C_2 &= [6, [3, 0], 3, [3, 2]] \\ C_3 &= [8, [4, -1], 4, [4, 1]] \end{split}$$

- $$\begin{split} C_0' &= [14, [7, -1], 7, [6, 2], 13, [7, 0], 6, [2, -1], 4, [2, 2], 2, [8, 0], 12, [7, 1], 1, [4, 2], 15, [1, 0]] \\ C_1' &= [10, [2, 1], 8, [18, 0], 18, [18, 2], 16, [1, -1], 17, [6, 0], 5, [5, 2]] \\ C_2' &= [0, [9, 0], 9, [9, 2]] \\ C_3' &= [11, [8, -1], 3, [8, 1]] \end{split}$$
- 2-factor type [8,7,2,2]: three starters
 - $$\begin{split} C_0 &= [3, [3, 1], 0, [5, 0], 13, [18, 1], 18, [18, -1], 14, [2, 2], 12, [8, 1], 4, [6, 0], 16, [5, 2]] \\ C_1 &= [6, [5, -1], 11, [1, 0], 10, [7, 1], 17, [8, 2], 9, [2, 0], 7, [1, 2], 8, [2, 1]] \\ C_2 &= [1, [4, 0], 15, [4, 2]] \\ C_3 &= [2, [3, 0], 5, [3, 2]] \\ C_0' &= [4, [1, -1], 5, [4, 1], 1, [6, -1], 7, [3, -1], 10, [2, -1], 12, [4, -1], 16, [1, 1], 15, [7, -1]] \\ C_1' &= [3, [5, 1], 8, [6, 1], 14, [8, -1], 6, [18, 2], 18, [18, 0], 17, [6, 2], 11, [8, 0]] \\ C_2' &= [0, [9, 0], 9, [9, 2]] \end{split}$$
 - $C'_3 = [13, [7, 0], 2, [7, 2]]$
- 2-factor type [11, 3, 3, 2]: three starters
 - $$\begin{split} C_0 &= [15, [8, -1], 7, [6, 2], 13, [3, 0], 10, [8, 2], 2, [18, 0], 18, [18, -1], 9, [3, 1], 12, [5, 1], 17, [5, 2], 4, [7, -1], \\ &\quad 11, [4, 0]] \end{split}$$
 - $C_1 = [8, [6, -1], 14, [2, -1], 16, [8, 1]]$
 - $C_2 = [0, [3, -1], 3, [2, 0], 1, [1, 2]]$
 - $C_3 = [6, [1, -1], 5, [1, 1]]$
 - $C_0' = [12, [2, 2], 10, [1, 0], 11, [5, -1], 6, [7, 2], 13, [5, 0], 8, [18, 1], 18, [18, 2], 3, [4, -1], 7, [8, 0], 17, [6, 1], 5, [7, 1]]$
 - $C_1' = [2, [4, 2], 16, [2, 1], 14, [6, 0]]$
 - $C_2' = [4, [3, 2], 1, [4, 1], 15, [7, 0]]$
 - $C_3' = \left[0, [9, 0], 9, [9, 2]\right]$
- 2-factor type [10, 4, 3, 2]: three starters

$$\begin{split} C_0 &= [6, [4, -1], 10, [3, 0], 7, [8, 1], 15, [4, 2], 1, [3, 1], 16, [5, 0], 11, [2, 2], 13, [1, 0], 14, [18, 2], 18, [18, 1]] \\ C_1 &= [5, [7, 1], 12, [6, 2], 0, [1, 1], 17, [6, 0]] \\ C_2 &= [4, [4, 0], 8, [5, -1], 3, [1, 2]] \\ C_3 &= [2, [7, 0], 9, [7, 2]] \\ C_0' &= [5, [8, -1], 15, [6, 1], 3, [3, 2], 6, [8, 0], 14, [2, 1], 16, [6, -1], 10, [2, -1], 8, [5, 1], 13, [1, -1], 12, [7, -1]] \\ C_1' &= [1, [8, 2], 11, [4, 1], 7, [18, 0], 18, [18, -1]] \\ C_2' &= [17, [3, -1], 2, [2, 0], 4, [5, 2]] \\ C_3' &= [0, [9, 0], 9, [9, 2]] \end{split}$$

• 2-factor type [9, 5, 3, 2]: three starters

$$\begin{split} C_0 &= [10, [5, 1], 15, [1, -1], 14, [6, 2], 8, [4, 0], 4, [4, 2], 0, [3, 0], 3, [8, 2], 13, [1, 1], 12, [2, 0]] \\ C_1 &= [9, [2, -1], 7, [1, 2], 6, [8, 0], 16, [3, 1], 1, [8, 1]] \\ C_2 &= [2, [3, 2], 5, [18, -1], 18, [18, 0]] \\ C_3 &= [11, [6, -1], 17, [6, 1]] \end{split}$$

$$\begin{split} C_0' &= [1, [18, 2], 18, [18, 1], 16, [1, 0], 15, [5, -1], 10, [2, 2], 12, [4, 1], 8, [3, -1], 11, [6, 0], 5, [4, -1]] \\ C_1' &= [7, [5, 2], 2, [7, 1], 13, [8, -1], 3, [7, 0], 14, [7, -1]] \\ C_2' &= [4, [2, 1], 6, [7, 2], 17, [5, 0]] \\ C_3' &= [0, [9, 0], 9, [9, 2]] \end{split}$$

• 2-factor type [8, 6, 3, 2]: three starters

$$\begin{split} C_0 &= [7, [18, 2], 18, [18, 0], 12, [7, 1], 1, [1, 2], 2, [6, 0], 14, [1, 1], 15, [3, 2], 0, [7, 0]] \\ C_1 &= [10, [7, 2], 3, [5, 1], 16, [6, -1], 4, [5, 0], 17, [8, 2], 9, [1, 0]] \\ C_2 &= [6, [2, 1], 8, [3, 1], 11, [5, -1]] \\ C_3 &= [13, [8, -1], 5, [8, 1]] \\ C_0' &= [13, [6, 2], 7, [4, 0], 3, [2, 2], 5, [18, -1], 18, [18, 1], 4, [8, 0], 12, [4, 2], 16, [3, 0]] \\ C_1' &= [11, [1, -1], 10, [4, 1], 14, [6, 1], 2, [3, -1], 17, [2, -1], 15, [4, -1]] \\ C_2' &= [1, [7, -1], 8, [2, 0], 6, [5, 2]] \\ C_3' &= [0, [9, 0], 9, [9, 2]] \end{split}$$

- 2-factor type [7,7,3,2]: three starters
 - $$\begin{split} C_0 &= [12, [4, 2], 16, [4, -1], 2, [18, 0], 18, [18, 2], 9, [3, 1], 6, [1, 0], 7, [5, 1]] \\ C_1 &= [17, [6, 0], 11, [6, -1], 5, [2, -1], 3, [2, 2], 1, [1, 1], 0, [4, 0], 4, [5, 2]] \\ C_2 &= [10, [3, 0], 13, [1, 2], 14, [4, 1]] \\ C_3 &= [15, [7, 0], 8, [7, 2]] \\ C'_0 &= [15, [2, 0], 13, [7, 1], 2, [1, -1], 1, [3, -1], 16, [8, 2], 8, [18, -1], 18, [18, 1]] \\ C'_1 &= [12, [6, 2], 6, [8, 0], 14, [8, 1], 4, [5, -1], 17, [7, -1], 10, [3, 2], 7, [5, 0]] \\ C'_2 &= [3, [8, -1], 11, [6, 1], 5, [2, 1]] \\ C'_3 &= [0, [9, 0], 9, [9, 2]] \end{split}$$
- 2-factor type [9, 4, 4, 2]: three starters

$$\begin{split} C_0 &= [7, [2, 2], 9, [3, -1], 12, [1, -1], 13, [1, 0], 14, [4, 2], 0, [3, 0], 15, [5, 2], 2, [1, 1], 1, [6, 0]] \\ C_1 &= [4, [18, -1], 18, [18, 1], 5, [6, -1], 11, [7, 1]] \\ C_2 &= [8, [2, 1], 10, [7, -1], 3, [5, -1], 16, [8, -1]] \\ C_3 &= [17, [7, 0], 6, [7, 2]] \\ C_0' &= [4, [6, 2], 16, [8, 0], 6, [3, 2], 3, [4, 0], 17, [4, -1], 13, [2, -1], 11, [1, 2], 10, [8, 1], 2, [2, 0]] \\ C_1' &= [12, [18, 2], 18, [18, 0], 15, [8, 2], 7, [5, 0]] \\ C_2' &= [5, [3, 1], 8, [6, 1], 14, [5, 1], 1, [4, 1]] \\ C_3' &= [0, [9, 0], 9, [9, 2]] \end{split}$$

- 2-factor type [8, 5, 4, 2]: three starters
 - $$\begin{split} C_0 &= [3, [1, -1], 2, [5, 1], 7, [18, 2], 18, [18, 0], 8, [6, 2], 14, [3, 1], 11, [1, 0], 10, [7, -1]] \\ C_1 &= [4, [6, -1], 16, [4, 2], 12, [6, 0], 0, [1, 2], 1, [3, 0]] \\ C_2 &= [5, [8, 1], 15, [6, 1], 9, [4, 0], 13, [8, 2]] \\ C_3 &= [17, [7, 0], 6, [7, 2]] \end{split}$$

- $$\begin{split} C_0' &= [7, [5, -1], 2, [3, -1], 17, [2, 2], 15, [4, -1], 1, [7, 1], 12, [8, 0], 4, [1, 1], 5, [2, 1]] \\ C_1' &= [10, [3, 2], 13, [8, -1], 3, [5, 0], 8, [2, -1], 6, [4, 1]] \\ C_2' &= [16, [2, 0], 14, [18, 1], 18, [18, -1], 11, [5, 2]] \\ C_3' &= [0, [9, 0], 9, [9, 2]] \end{split}$$
- 2-factor type [7, 6, 4, 2]: three starters

$$\begin{split} C_0 &= [2, [1, 1], 1, [2, 0], 17, [7, -1], 10, [2, 2], 8, [6, 0], 14, [8, 2], 4, [2, 1]] \\ C_1 &= [13, [5, 1], 0, [18, 1], 18, [18, 2], 15, [3, 0], 12, [7, 2], 5, [8, 0]] \\ C_2 &= [3, [5, 0], 16, [8, 1], 6, [3, 1], 9, [6, 2]] \\ C_3 &= [7, [4, -1], 11, [4, 1]] \\ C_0' &= [18, [18, 0], 10, [5, 2], 15, [1, -1], 16, [6, -1], 4, [1, 0], 3, [4, 2], 17, [18, -1]] \\ C_1' &= [13, [2, -1], 11, [1, 2], 12, [4, 0], 8, [3, -1], 5, [3, 2], 2, [7, 0]] \\ C_2' &= [7, [6, 1], 1, [5, -1], 6, [8, -1], 14, [7, 1]] \\ C_3' &= [0, [9, 0], 9, [9, 2]] \end{split}$$

• 2-factor type [7, 5, 5, 2]: three starters

$$\begin{split} C_0 &= [0, [1, 1], 17, [6, 1], 11, [5, -1], 16, [6, 0], 10, [4, -1], 6, [7, 1], 13, [5, 2]] \\ C_1 &= [9, [1, 2], 8, [6, -1], 14, [7, 0], 7, [3, 2], 4, [5, 0]] \\ C_2 &= [2, [8, 1], 12, [7, -1], 1, [4, 0], 15, [18, 1], 18, [18, 2]] \\ C_3 &= [3, [2, -1], 5, [2, 1]] \\ C_0' &= [18, [18, -1], 15, [7, 2], 4, [3, 1], 1, [2, 0], 3, [4, 2], 17, [5, 1], 12, [18, 0]] \\ C_1' &= [11, [4, 1], 7, [3, 0], 10, [8, -1], 2, [6, 2], 8, [3, -1]] \\ C_2' &= [14, [1, 0], 13, [8, 2], 5, [1, -1], 6, [8, 0], 16, [2, 2]] \\ C_3' &= [0, [9, 0], 9, [9, 2]] \end{split}$$

• 2-factor type [6, 6, 5, 2]: three starters

$$\begin{split} C_0 &= [3, [6, 1], 15, [1, 0], 14, [2, -1], 12, [5, -1], 7, [6, 2], 13, [8, -1]] \\ C_1 &= [10, [1, -1], 9, [8, 2], 1, [2, 1], 17, [6, 0], 5, [3, -1], 2, [8, 1]] \\ C_2 &= [8, [4, -1], 4, [2, 2], 6, [18, -1], 18, [18, 1], 16, [8, 0]] \\ C_3 &= [0, [7, 0], 11, [7, 2]] \\ C_0' &= [4, [2, 0], 6, [3, 2], 3, [5, 0], 8, [7, -1], 1, [4, 2], 5, [1, 1]] \\ C_1' &= [2, [7, 1], 13, [3, 1], 10, [6, -1], 16, [4, 0], 12, [5, 2], 7, [5, 1]] \\ C_2' &= [17, [3, 0], 14, [1, 2], 15, [4, 1], 11, [18, 0], 18, [18, 2]] \\ C_3' &= [0, [9, 0], 9, [9, 2]] \end{split}$$

- 2-factor type [15, 2, 2]: three starters
 - $$\begin{split} C_0 &= [11, [1, 1], 12, [7, 2], 5, [3, 0], 2, [4, 2], 6, [18, 0], 18, [18, 2], 13, [6, 1], 7, [2, 0], 9, [6, 2], 15, [1, -1], 16, \\ &\qquad [2, 1], 14, [4, 0], 10, [7, -1], 3, [4, -1], 17, [6, -1]] \end{split}$$

$$C_1 = [0, [8, -1], 8, [8, 1]]$$
$$C_2 = [1, [3, -1], 4, [3, 1]]$$

- $$\begin{split} C_0' &= [1, [6, 0], 13, [8, 2], 5, [2, -1], 7, [5, 0], 12, [2, 2], 14, [4, 1], 10, [7, 0], 17, [1, 2], 16, [8, 0], 8, [5, 2], 3, [1, 0], \\ &2, [18, 1], 18, [18, -1], 15, [7, 1], 4, [3, 2]] \\ C_1' &= [0, [9, 0], 9, [9, 2]] \\ C_2' &= [6, [5, -1], 11, [5, 1]] \end{split}$$
- 2-factor type [14, 3, 2]: three starters
 - $$\begin{split} C_0 &= [17, [3, 2], 14, [6, 0], 2, [6, 1], 8, [3, -1], 5, [4, 1], 9, [4, 2], 13, [1, 1], 12, [4, 0], 16, [5, -1], 11, [7, 2], 4, \\ &\quad [1, 0], 3, [2, -1], 1, [6, 2], 7, [8, 0]] \\ C_1 &= [0, [8, -1], 10, [4, -1], 6, [6, -1]] \\ C_2 &= [18, [18, 0], 15, [18, 2]] \\ C_0' &= [10, [3, 0], 7, [2, 2], 5, [1, -1], 6, [2, 0], 4, [8, 1], 12, [2, 1], 14, [1, 2], 13, [7, -1], 2, [3, 1], 17, [18, -1], 18, \\ &\quad [18, 1], 1, [7, 0], 8, [7, 1], 15, [5, 2]] \\ C_1' &= [3, [8, 2], 11, [5, 0], 16, [5, 1]] \\ C_2' &= [0, [9, 0], 9, [9, 2]] \end{split}$$
- 2-factor type [13, 4, 2]: three starters
 - $$\begin{split} C_0 &= [3, [1, 2], 2, [7, 1], 13, [5, 1], 8, [2, 1], 6, [5, 0], 1, [2, -1], 17, [5, 2], 4, [7, -1], 11, [6, 1], 5, [7, 0], 12, [4, 2], \\ &\quad 16, [18, -1], 18, [18, 0]] \\ C_1 &= [0, [8, 0], 10, [4, 1], 14, [1, 1], 15, [3, 2]] \\ C_2 &= [7, [2, 0], 9, [2, 2]] \\ C_0' &= [16, [8, 2], 8, [6, -1], 14, [1, -1], 15, [3, 0], 12, [5, -1], 7, [6, 2], 13, [8, 1], 5, [3, 1], 2, [4, 0], 6, [18, 1], 18, \\ &\quad [18, 2], 1, [3, -1], 4, [6, 0]] \\ C_1' &= [11, [8, -1], 3, [4, -1], 17, [7, 2], 10, [1, 0]] \\ C_2' &= [0, [9, 0], 9, [9, 2]] \end{split}$$
- 2-factor type [12, 5, 2]: three starters
 - $$\begin{split} C_0 &= [7, [1, 1], 6, [7, 1], 17, [2, 0], 15, [4, -1], 11, [1, 2], 12, [6, 0], 0, [8, -1], 10, [6, 1], 16, [8, 2], 8, [3, 0], 5, \\ &\quad [18, 2], 18, [18, 1]] \\ C_1 &= [13, [8, 1], 3, [1, -1], 2, [6, -1], 14, [5, -1], 9, [4, 1]] \\ C_2 &= [1, [3, -1], 4, [3, 1]] \\ C_0' &= [7, [5, 0], 2, [5, 2], 15, [4, 0], 11, [2, 2], 13, [8, 0], 3, [4, 2], 17, [1, 0], 16, [6, 2], 10, [5, 1], 5, [7, 0], 12, [2, 1], 0 \end{split}$$
 - 14, [7, 2]] $C'_1 = [18, [18, 0], 8, [7, -1], 1, [3, 2], 4, [2, -1], 6, [18, -1]]$
 - $C'_{2} = [0, [9, 0], 9, [9, 2]]$
- 2-factor type [11, 6, 2]: three starters
 - $$\begin{split} C_0 = [7, [6, 0], 13, [7, 1], 2, [2, -1], 0, [3, 2], 15, [18, 0], 18, [18, -1], 4, [5, 2], 9, [1, -1], 10, [4, 0], 6, [3, -1], 3, \\ [4, 2]] \end{split}$$
 - $C_1 = [8, [3, 1], 11, [5, 1], 16, [7, 2], 5, [7, 0], 12, [2, 1], 14, [6, -1]]$
 - $C_2 = [17, [2, 0], 1, [2, 2]]$

$$\begin{split} C_0' &= [8, [8, 0], 16, [6, 1], 4, [7, -1], 15, [6, 2], 3, [3, 0], 6, [4, -1], 2, [8, -1], 12, [1, 1], 13, [4, 1], 17, [8, 1], 7, \\ & [1, 2]] \\ C_1' &= [18, [18, 1], 14, [5, 0], 1, [8, 2], 11, [1, 0], 10, [5, -1], 5, [18, 2]] \\ C_2' &= [0, [9, 0], 9, [9, 2]] \end{split}$$

• 2-factor type [10, 7, 2]: three starters

$$\begin{split} C_0 &= [17, [8, 2], 9, [6, 1], 3, [2, 0], 5, [18, 2], 18, [18, 1], 7, [5, 1], 2, [6, 0], 14, [2, 2], 12, [7, 0], 1, [2, -1]] \\ C_1 &= [11, [1, -1], 10, [3, 1], 13, [7, 2], 6, [2, 1], 4, [4, 0], 0, [3, -1], 15, [4, -1]] \\ C_2 &= [8, [8, -1], 16, [8, 1]] \\ C_0' &= [18, [18, 0], 14, [5, 2], 1, [7, 1], 12, [4, 1], 8, [5, 0], 13, [7, -1], 6, [1, 2], 5, [3, 0], 2, [4, 2], 16, [18, -1]] \\ C_1' &= [11, [6, 2], 17, [8, 0], 7, [3, 2], 4, [1, 0], 3, [6, -1], 15, [5, -1], 10, [1, 1]] \\ C_2' &= [0, [9, 0], 9, [9, 2]] \end{split}$$

• 2-factor type [9, 8, 2]: three starters

$$\begin{split} C_0 &= [18, [18, 2], 15, [7, 0], 8, [4, 1], 12, [2, -1], 10, [3, 1], 13, [6, 2], 1, [5, -1], 6, [6, 1], 0, [18, 0]] \\ C_1 &= [4, [1, 1], 3, [5, 0], 16, [7, -1], 9, [4, -1], 5, [2, 1], 7, [7, 1], 14, [3, -1], 11, [7, 2]] \\ C_2 &= [2, [3, 0], 17, [3, 2]] \\ C'_0 &= [18, [18, -1], 10, [5, 2], 5, [1, -1], 6, [2, 0], 4, [8, 2], 14, [1, 0], 15, [2, 2], 13, [6, 0], 1, [18, 1]] \\ C'_1 &= [8, [6, -1], 2, [1, 2], 3, [8, 0], 11, [4, 2], 7, [8, 1], 17, [5, 1], 12, [4, 0], 16, [8, -1]] \\ C'_2 &= [0, [9, 0], 9, [9, 2]] \end{split}$$

L Computational results for n = 20

• 2-factor type [4, 2, 2, 2, 2, 2, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [4, [8, 0], 12, [1, 1], 11, [8, 1], 3, [1, 0]] \\ C_1 &= [6, [19, 0], 19, [19, 1]] \\ C_2 &= [0, [5, 0], 5, [5, 1]] \\ C_3 &= [15, [2, 0], 13, [2, 1]] \\ C_4 &= [18, [3, 0], 2, [3, 1]] \\ C_5 &= [1, [6, 0], 7, [6, 1]] \\ C_6 &= [8, [9, 0], 17, [9, 1]] \\ C_7 &= [9, [7, 0], 16, [7, 1]] \\ C_8 &= [14, [4, 0], 10, [4, 1]] \end{split}$$

• 2-factor type [3, 3, 2, 2, 2, 2, 2, 2, 2]: two starters

$C_0 = [2, [1, -1], 1, [19, 1], 19, [19, -1]]$	$C_0' = [17, [7, 2], 10, [5, 0], 5, [7, -1]]$
$C_1 = [9, [7, 0], 16, [7, 1], 4, [5, 2]]$	$C_1' = [0, [1, 1], 1, [19, 2], 19, [19, 0]]$
$C_2 = [14, [9, -1], 5, [9, 1]]$	$C_2' = [15, [6, -1], 2, [6, 1]]$
$C_3 = [8, [1, 0], 7, [1, 2]]$	$C_3' = [16, [8, 0], 8, [8, 2]]$
$C_4 = [18, [4, 0], 3, [4, 2]]$	$C_4' = [18, [6, 0], 12, [6, 2]]$
$C_5 = [10, [3, 0], 13, [3, 2]]$	$C_5' = [4, [9, 0], 13, [9, 2]]$
$C_6 = [17, [2, -1], 0, [2, 1]]$	$C_6' = [11, [2, 0], 9, [2, 2]]$
$C_7 = [12, [3, -1], 15, [3, 1]]$	$C_7' = [7, [4, -1], 3, [4, 1]]$
$C_8 = [11, [5, -1], 6, [5, 1]]$	$C_8' = [14, [8, -1], 6, [8, 1]]$

• 2-factor type [6, 2, 2, 2, 2, 2, 2, 2]: two starters

$$\begin{split} C_0 &= [10, [8, 2], 18, [9, 0], 9, [19, 1], 19, [19, -1], 2, [4, 2], 6, [4, 0]] \\ C_1 &= [12, [1, -1], 13, [1, 1]] \\ C_2 &= [7, [9, -1], 17, [9, 1]] \\ C_3 &= [1, [4, -1], 16, [4, 1]] \\ C_4 &= [8, [7, -1], 15, [7, 1]] \\ C_5 &= [14, [3, -1], 11, [3, 1]] \\ C_6 &= [5, [1, 0], 4, [1, 2]] \\ C_7 &= [3, [3, 0], 0, [3, 2]] \\ C_0' &= [1, [7, 0], 13, [9, 2], 4, [19, 0], 19, [19, 2], 16, [8, 0], 8, [7, 2]] \\ C_1' &= [3, [5, 0], 17, [5, 2]] \\ C_2' &= [0, [5, -1], 14, [5, 1]] \\ C_3' &= [5, [2, 0], 7, [2, 2]] \\ C_4' &= [6, [6, -1], 12, [6, 1]] \\ C_5' &= [11, [2, -1], 9, [2, 1]] \\ C_6' &= [15, [6, 0], 2, [6, 2]] \\ C_7' &= [10, [8, -1], 18, [8, 1]] \end{split}$$

• 2-factor type [5, 3, 2, 2, 2, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [6, [3, 0], 3, [4, 1], 18, [9, 1], 8, [3, 1], 11, [5, 0]] \\ C_1 &= [10, [4, 0], 14, [5, 1], 0, [9, 0]] \\ C_2 &= [16, [1, 0], 17, [1, 1]] \\ C_3 &= [12, [8, 0], 4, [8, 1]] \\ C_4 &= [9, [19, 0], 19, [19, 1]] \\ C_5 &= [15, [6, 0], 2, [6, 1]] \\ C_6 &= [1, [7, 0], 13, [7, 1]] \\ C_7 &= [7, [2, 0], 5, [2, 1]] \end{split}$$

• 2-factor type [4, 4, 2, 2, 2, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [15, [3, 0], 12, [8, 0], 4, [2, 1], 2, [6, 1]] \\ C_1 &= [3, [6, 0], 16, [2, 0], 14, [8, 1], 6, [3, 1]] \\ C_2 &= [11, [1, 0], 10, [1, 1]] \\ C_3 &= [19, [19, 0], 0, [19, 1]] \\ C_4 &= [8, [7, 0], 1, [7, 1]] \\ C_5 &= [18, [5, 0], 13, [5, 1]] \\ C_6 &= [17, [9, 0], 7, [9, 1]] \\ C_7 &= [5, [4, 0], 9, [4, 1]] \end{split}$$

• 2-factor type [4, 3, 3, 2, 2, 2, 2, 2]: two starters

$$\begin{split} C_0 &= [16, [3, 1], 13, [9, 0], 3, [7, -1], 15, [1, 2]] \\ C_1 &= [18, [7, 1], 11, [9, -1], 1, [2, 1]] \\ C_2 &= [4, [1, 0], 5, [7, 2], 12, [8, 1]] \\ C_3 &= [19, [19, 0], 10, [19, 2]] \\ C_4 &= [14, [6, 0], 8, [6, 2]] \\ C_5 &= [2, [5, -1], 7, [5, 1]] \\ C_6 &= [0, [6, -1], 6, [6, 1]] \\ C_7 &= [17, [8, 0], 9, [8, 2]] \end{split}$$

• 2-factor type [3, 3, 3, 3, 2, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [2, [6, 0], 8, [2, 0], 6, [4, 1]] \\ C_1 &= [3, [9, 0], 12, [2, 1], 10, [7, 0]] \\ C_2 &= [17, [7, 1], 5, [4, 0], 1, [3, 0]] \\ C_3 &= [4, [9, 1], 13, [6, 1], 7, [3, 1]] \\ C_4 &= [18, [19, 0], 19, [19, 1]] \\ C_5 &= [16, [1, 0], 15, [1, 1]] \\ C_6 &= [9, [5, 0], 14, [5, 1]] \\ C_7 &= [11, [8, 0], 0, [8, 1]] \end{split}$$

• 2-factor type [8, 2, 2, 2, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [18, [4, 1], 14, [9, 1], 4, [4, 0], 8, [2, 1], 6, [9, 0], 16, [1, 0], 17, [2, 0], 0, [1, 1]] \\ C_1 &= [15, [7, 0], 3, [7, 1]] \\ C_2 &= [10, [3, 0], 13, [3, 1]] \\ C_3 &= [5, [6, 0], 11, [6, 1]] \\ C_4 &= [19, [19, 0], 12, [19, 1]] \\ C_5 &= [2, [5, 0], 7, [5, 1]] \\ C_6 &= [9, [8, 0], 1, [8, 1]] \end{split}$$

$$\begin{split} C_0' &= [2, [9, 2], 11, [8, -1], 0, [5, 0], 5, [3, -1]] \\ C_1' &= [7, [9, 1], 17, [7, 0], 10, [3, 2]] \\ C_2' &= [4, [3, 0], 1, [2, -1], 18, [5, 2]] \\ C_3' &= [8, [2, 0], 6, [2, 2]] \\ C_4' &= [13, [4, -1], 9, [4, 1]] \\ C_5' &= [3, [19, -1], 19, [19, 1]] \\ C_6' &= [14, [1, -1], 15, [1, 1]] \\ C_7' &= [16, [4, 0], 12, [4, 2]] \end{split}$$

• 2-factor type [7, 3, 2, 2, 2, 2, 2]: two starters

$$\begin{split} C_0 &= [10, [2, 1], 12, [9, 2], 3, [5, -1], 8, [7, -1], 15, [1, 0], 14, [7, 1], 2, [8, 1]] \\ C_1 &= [17, [1, -1], 16, [9, 1], 6, [8, -1]] \\ C_2 &= [18, [2, 0], 1, [2, 2]] \\ C_3 &= [5, [19, 0], 19, [19, 2]] \\ C_4 &= [0, [6, 0], 13, [6, 2]] \\ C_5 &= [11, [4, -1], 7, [4, 1]] \\ C_6 &= [9, [5, 0], 4, [5, 2]] \\ C'_0 &= [19, [19, 1], 5, [9, 0], 15, [5, 1], 1, [2, -1], 18, [8, 2], 7, [1, 1], 6, [19, -1]] \\ C'_1 &= [4, [8, 0], 12, [9, -1], 3, [1, 2]] \\ C'_2 &= [0, [3, -1], 16, [3, 1]] \\ C'_3 &= [9, [4, 0], 13, [4, 2]] \\ C'_4 &= [10, [7, 0], 17, [7, 2]] \\ C'_5 &= [11, [3, 0], 14, [3, 2]] \\ C'_6 &= [2, [6, -1], 8, [6, 1]] \end{split}$$

• 2-factor type [6, 4, 2, 2, 2, 2, 2]: two starters

$$\begin{split} C_0 &= [1, [9, 0], 10, [4, -1], 6, [1, 2], 7, [4, 0], 3, [1, 1], 4, [3, 2]] \\ C_1 &= [11, [2, 0], 9, [7, 1], 16, [1, -1], 15, [4, 2]] \\ C_2 &= [5, [5, 0], 0, [5, 2]] \\ C_3 &= [13, [8, -1], 2, [8, 1]] \\ C_4 &= [19, [19, -1], 14, [19, 1]] \\ C_5 &= [8, [9, -1], 17, [9, 1]] \\ C_6 &= [12, [6, -1], 18, [6, 1]] \\ C'_0 &= [1, [1, 0], 2, [7, 2], 9, [4, 1], 5, [6, 0], 18, [7, -1], 11, [9, 2]] \\ C'_1 &= [10, [6, 2], 16, [3, 0], 0, [2, 2], 17, [7, 0]] \\ C'_2 &= [7, [8, 0], 15, [8, 2]] \\ C'_3 &= [6, [3, -1], 3, [3, 1]] \\ C'_4 &= [19, [19, 0], 4, [19, 2]] \\ C'_5 &= [12, [2, -1], 14, [2, 1]] \\ C'_6 &= [13, [5, -1], 8, [5, 1]] \end{split}$$

- 2-factor type [5, 5, 2, 2, 2, 2, 2]: two starters
 - $C_{0} = [5, [1, 2], 4, [5, -1], 9, [9, -1], 18, [7, 0], 6, [1, -1]]$ $C_{1} = [12, [1, 1], 13, [2, -1], 11, [8, -1], 3, [7, 2], 10, [2, 0]]$ $C_{2} = [2, [4, 0], 17, [4, 2]]$ $C_{3} = [0, [8, 0], 8, [8, 2]]$ $C_{4} = [14, [7, -1], 7, [7, 1]]$ $C_{5} = [1, [5, 0], 15, [5, 2]]$ $C_{6} = [19, [19, -1], 16, [19, 1]]$

- $$\begin{split} C_0' &= [16, [3, 1], 0, [2, 1], 2, [9, 1], 11, [3, -1], 8, [8, 1]] \\ C_1' &= [15, [1, 0], 14, [2, 2], 12, [5, 1], 7, [19, 0], 19, [19, 2]] \\ C_2' &= [9, [4, -1], 13, [4, 1]] \\ C_3' &= [3, [3, 0], 6, [3, 2]] \\ C_4' &= [1, [9, 0], 10, [9, 2]] \\ C_5' &= [18, [6, -1], 5, [6, 1]] \\ C_6' &= [4, [6, 0], 17, [6, 0]] \end{split}$$
- $C'_6 = [4, [6, 0], 17, [6, 2]]$
- 2-factor type [6, 3, 3, 2, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [6, [9, 1], 16, [4, 0], 12, [4, 1], 8, [9, 0], 18, [8, 0], 7, [1, 0]] \\ C_1 &= [10, [8, 1], 2, [7, 1], 9, [1, 1]] \\ C_2 &= [11, [6, 0], 17, [7, 0], 5, [6, 1]] \\ C_3 &= [4, [3, 0], 1, [3, 1]] \\ C_4 &= [3, [19, 0], 19, [19, 1]] \\ C_5 &= [0, [5, 0], 14, [5, 1]] \end{split}$$

$$C_6 = [15, [2, 0], 13, [2, 1]]$$

• 2-factor type [5, 4, 3, 2, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [9, [19, 0], 19, [19, 1], 1, [5, 0], 15, [9, 1], 5, [4, 1]] \\ C_1 &= [11, [4, 0], 7, [5, 1], 2, [6, 0], 8, [3, 1]] \\ C_2 &= [6, [6, 1], 12, [9, 0], 3, [3, 0]] \\ C_3 &= [17, [2, 0], 0, [2, 1]] \\ C_4 &= [18, [8, 0], 10, [8, 1]] \\ C_5 &= [16, [7, 0], 4, [7, 1]] \\ C_6 &= [13, [1, 0], 14, [1, 1]] \end{split}$$

• 2-factor type [4, 4, 4, 2, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [8, [8, 0], 0, [6, 0], 6, [7, 0], 13, [5, 0]] \\ C_1 &= [5, [2, 1], 7, [8, 1], 18, [7, 1], 11, [6, 1]] \\ C_2 &= [10, [19, 1], 19, [19, 0], 17, [2, 0], 15, [5, 1]] \\ C_3 &= [9, [3, 0], 12, [3, 1]] \\ C_4 &= [14, [9, 0], 4, [9, 1]] \\ C_5 &= [2, [1, 0], 3, [1, 1]] \\ C_6 &= [1, [4, 0], 16, [4, 1]] \end{split}$$

• 2-factor type [5, 3, 3, 3, 2, 2, 2]: two starters

$$\begin{split} C_0 &= [4, [9, 1], 14, [1, 0], 15, [3, 2], 12, [6, 1], 6, [2, 1]] \\ C_1 &= [8, [8, 2], 0, [9, 0], 10, [2, -1]] \\ C_2 &= [16, [6, -1], 3, [19, 0], 19, [19, 2]] \\ C_3 &= [9, [8, -1], 1, [9, 2], 11, [2, 0]] \\ C_4 &= [17, [4, -1], 13, [4, 1]] \\ C_5 &= [18, [6, 0], 5, [6, 2]] \\ C_6 &= [7, [5, -1], 2, [5, 1]] \end{split}$$

- $$\begin{split} C_0' &= [11, [9, -1], 2, [3, -1], 18, [7, 1], 6, [19, 1], 19, [19, -1]] \\ C_1' &= [17, [5, 2], 3, [8, 1], 14, [3, 0]] \\ C_2' &= [4, [1, 2], 5, [7, -1], 12, [8, 0]] \\ C_3' &= [10, [5, 0], 15, [2, 2], 13, [3, 1]] \\ C_4' &= [1, [4, 0], 16, [4, 2]] \\ C_5' &= [0, [7, 0], 7, [7, 2]] \\ C_6' &= [9, [1, -1], 8, [1, 1]] \end{split}$$
- 2-factor type [4, 4, 3, 3, 2, 2, 2]: two starters

$$\begin{split} C_0 &= [16, [7, 2], 9, [8, 0], 17, [2, 2], 15, [1, 0]] \\ C_1 &= [7, [4, 1], 3, [1, -1], 4, [9, -1], 13, [6, 1]] \\ C_2 &= [5, [9, 1], 14, [19, 2], 19, [19, 0]] \\ C_3 &= [1, [9, 2], 10, [9, 0], 0, [1, 1]] \\ C_4 &= [11, [5, 0], 6, [5, 2]] \\ C_5 &= [18, [3, -1], 2, [3, 1]] \\ C_6 &= [12, [4, 0], 8, [4, 2]] \end{split}$$

- 2-factor type [4, 3, 3, 3, 3, 2, 2]: one starter $C_0 = [7, [19, 0], 19, [19, 1], 6, [2, 1], 8, [1, 0]]$ $C_1 = [12, [3, 1], 9, [9, 1], 18, [6, 1]]$ $C_2 = [16, [5, 0], 2, [6, 0], 15, [1, 1]]$ $C_3 = [4, [3, 0], 1, [7, 1], 13, [9, 0]]$ $C_4 = [17, [7, 0], 5, [2, 0], 3, [5, 1]]$ $C_5 = [10, [4, 0], 14, [4, 1]]$ $C_6 = [0, [8, 0], 11, [8, 1]]$
- 2-factor type [3, 3, 3, 3, 3, 3, 3, 2]: two starters

$C_0 = [7, [5, 1], 12, [19, 2], 19, [19, 0]]$	$C_0' = [4, [3, 1], 7, [7, 2], 14, [9, 0]]$
$C_1 = [4, [2, 2], 2, [7, 0], 14, [9, 1]]$	$C_1' = [1, [5, 2], 15, [1, 0], 16, [4, 1]]$
$C_2 = [0, [3, 2], 16, [3, 0], 13, [6, 1]]$	$C_2' = [18, [4, -1], 3, [9, -1], 12, [6, -1]]$
$C_3 = [5, [5, -1], 10, [5, 0], 15, [9, 2]]$	$C_3^\prime = [13, [4, 0], 9, [7, -1], 2, [8, 2]]$
$C_4 = [1, [2, -1], 18, [4, 2], 3, [2, 0]]$	$C_4' = [5, [1, 2], 6, [8, 0], 17, [7, 1]]$
$C_5 = [8, [2, 1], 6, [3, -1], 9, [1, 1]]$	$C_5^\prime = [10, [1, -1], 11, [19, -1], 19, [19, 1]]$
$C_6 = [11, [6, 0], 17, [6, 2]]$	$C_6' = [0, [8, -1], 8, [8, 1]]$

• 2-factor type [10, 2, 2, 2, 2, 2]: two starters $C_0 = [1, [4, 1], 5, [2, 1], 7, [1, 1], 8, [3, 1], 11, [8, 1], 0, [3, 2], 3, [5, 0], 17, [8, 2], 6, [3, 0], 9, [8, -1]]$ $C_1 = [13, [1, 0], 14, [1, 2]]$ $C_2 = [19, [19, -1], 16, [19, 1]]$ $C_3 = [18, [6, -1], 12, [6, 1]]$ $C_4 = [15, [5, -1], 10, [5, 1]]$ $C_5 = [4, [2, 0], 2, [2, 2]]$

$$C'_{0} = [18, [1, 2], 0, [6, 0], 13, [8, -1], 5, [6, -1]]$$

$$C'_{1} = [3, [19, -1], 19, [19, 1], 8, [3, 0], 11, [8, 2]]$$

$$C'_{2} = [17, [4, -1], 2, [7, 0], 14, [3, 2]]$$

$$C'_{3} = [15, [8, 1], 7, [2, 0], 9, [6, 2]]$$

$$C'_{4} = [4, [7, -1], 16, [7, 1]]$$

$$C'_{5} = [1, [5, -1], 6, [5, 1]]$$

$$C_6' = [10, [2, -1], 12, [2, 1]]$$

$$\begin{split} C_0' &= [8, [8, 0], 16, [6, 2], 10, [3, -1], 13, [6, 0], 0, [19, 2], 19, [19, 0], 6, [5, 2], 1, [1, -1], 2, [2, -1], 4, [4, -1]] \\ C_1' &= [14, [7, -1], 7, [7, 1]] \\ C_2' &= [12, [9, 0], 3, [9, 2]] \\ C_3' &= [17, [7, 0], 5, [7, 2]] \\ C_4' &= [9, [9, -1], 18, [9, 1]] \\ C_5' &= [15, [4, 0], 11, [4, 2]] \end{split}$$

• 2-factor type [9, 3, 2, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [1, [2, 1], 18, [1, 0], 17, [19, 1], 19, [19, 0], 15, [8, 0], 4, [1, 1], 5, [9, 1], 14, [2, 0], 12, [8, 1]] \\ C_1 &= [2, [5, 0], 16, [9, 0], 7, [5, 1]] \\ C_2 &= [8, [3, 0], 11, [3, 1]] \\ C_3 &= [9, [4, 0], 13, [4, 1]] \\ C_4 &= [3, [7, 0], 10, [7, 1]] \\ C_5 &= [0, [6, 0], 6, [6, 1]] \end{split}$$

• 2-factor type [8, 4, 2, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [7, [19, 1], 19, [19, 0], 14, [6, 1], 8, [2, 1], 6, [4, 1], 10, [7, 0], 17, [4, 0], 13, [6, 0]] \\ C_1 &= [18, [7, 1], 11, [2, 0], 9, [5, 1], 4, [5, 0]] \\ C_2 &= [16, [3, 0], 0, [3, 1]] \\ C_3 &= [1, [8, 0], 12, [8, 1]] \\ C_4 &= [15, [9, 0], 5, [9, 1]] \\ C_5 &= [3, [1, 0], 2, [1, 1]] \end{split}$$

• 2-factor type [7, 5, 2, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [13, [1, 0], 12, [2, 0], 10, [5, 0], 15, [1, 1], 16, [4, 0], 1, [2, 1], 3, [9, 1]] \\ C_1 &= [14, [5, 1], 9, [9, 0], 18, [7, 1], 6, [4, 1], 2, [7, 0]] \\ C_2 &= [7, [19, 0], 19, [19, 1]] \\ C_3 &= [4, [6, 0], 17, [6, 1]] \\ C_4 &= [0, [8, 0], 11, [8, 1]] \\ C_5 &= [8, [3, 0], 5, [3, 1]] \end{split}$$

• 2-factor type [6, 6, 2, 2, 2, 2]: one starter

$$\begin{split} C_0 &= [17, [2, 1], 0, [1, 1], 1, [4, 1], 5, [4, 0], 9, [5, 1], 14, [3, 0]] \\ C_1 &= [3, [7, 1], 10, [2, 0], 8, [3, 1], 11, [5, 0], 16, [7, 0], 4, [1, 0]] \\ C_2 &= [13, [8, 0], 2, [8, 1]] \\ C_3 &= [15, [9, 0], 6, [9, 1]] \\ C_4 &= [7, [19, 0], 19, [19, 1]] \\ C_5 &= [12, [6, 0], 18, [6, 1]] \end{split}$$

- 2-factor type [8, 3, 3, 2, 2, 2]: two starters $C_0 = [17, [6, 1], 11, [6, 0], 5, [5, 1], 10, [3, 2], 13, [7, 0], 1, [3, 1], 4, [4, -1], 0, [2, 2]]$ $C_1 = [9, [1, -1], 8, [4, 1], 12, [3, -1]]$ $C_2 = [16, [2, 0], 18, [8, 2], 7, [9, -1]]$
 - $C_{3} = [6, [9, 0], 15, [9, 2]]$ $C_{4} = [14, [7, -1], 2, [7, 1]]$ $C_{5} = [3, [19, 0], 19, [19, 2]]$ $C'_{0} = [2, [3, 0], 18, [1, 2], 17, [8, 0], 9, [4, 2], 5, [1, 1], 4, [2, -1], 6, [5, -1], 11, [9, 1]]$ $C'_{1} = [1, [7, 2], 13, [6, -1], 0, [1, 0]]$ $C'_{2} = [10, [6, 2], 16, [2, 1], 14, [4, 0]]$ $C'_{3} = [3, [5, 0], 8, [5, 2]]$ $C'_{4} = [12, [19, -1], 19, [19, 1]]$ $C'_{5} = [15, [8, -1], 7, [8, 1]]$
- 2-factor type [7, 4, 3, 2, 2, 2]: two starters $C_0 = [5, [2, 1], 3, [9, 1], 13, [7, 0], 6, [3, 1], 9, [6, 1], 15, [4, -1], 11, [6, 2]]$ $C_1 = [0, [5, 0], 14, [19, 2], 19, [19, 0], 16, [3, 2]]$ $C_2 = [4, [8, 0], 12, [4, 2], 8, [4, 1]]$ $C_3 = [2, [1, 0], 1, [1, 2]]$ $C_4 = [10, [7, -1], 17, [7, 1]]$ $C_5 = [18, [8, -1], 7, [8, 1]]$ $C'_0 = [5, [6, -1], 11, [9, -1], 1, [2, -1], 3, [4, 0], 18, [8, 2], 7, [5, 1], 2, [3, -1]]$

$$\begin{split} C_0 &= [0, [0, -1], 12, [0, -1], 12, [0, -1], 0, [1, 0], 0, [0, 0], 10, [0, 2], 1, [0, 1], 2, [0, -1], 2, [0, -1], 2, [0, -1], 12, [0, -1], 13, [0, 0]] \\ C_2 &= [15, [6, 0], 9, [5, 2], 14, [1, -1]] \\ C_3' &= [6, [2, 0], 8, [2, 2]] \\ C_4' &= [10, [19, -1], 19, [19, 1]] \\ C_5' &= [13, [9, 0], 4, [9, 2]] \end{split}$$

• 2-factor type [6, 5, 3, 2, 2, 2]: two starters

$$\begin{split} C_0 &= [13, [9, -1], 4, [2, -1], 2, [7, 2], 9, [1, 1], 8, [2, 0], 10, [3, 1]] \\ C_1 &= [14, [2, 1], 16, [4, 2], 1, [9, 1], 11, [8, 1], 3, [8, 0]] \\ C_2 &= [6, [7, -1], 18, [6, 2], 5, [1, 0]] \\ C_3 &= [19, [19, 0], 0, [19, 2]] \\ C_4 &= [7, [9, 0], 17, [9, 2]] \\ C_5 &= [15, [3, 0], 12, [3, 2]] \\ C_6 &= [1, [1, 2], 0, [4, 0], 4, [8, 2], 12, [4, 1], 8, [6, 0], 2, [1, -1]] \\ C_1' &= [6, [7, 0], 13, [8, -1], 5, [2, 2], 7, [19, -1], 19, [19, 1]] \\ C_2' &= [11, [4, -1], 15, [3, -1], 18, [7, 1]] \\ C_3' &= [16, [6, -1], 10, [6, 1]] \\ C_4' &= [14, [5, 0], 9, [5, 2]] \\ C_5' &= [3, [5, -1], 17, [5, 1]] \end{split}$$

• 2-factor type [6, 4, 4, 2, 2, 2]: two starters

$$\begin{split} C_0 &= [16, [7, 1], 4, [1, 2], 3, [19, 0], 19, [19, -1], 14, [2, -1], 12, [4, 1]] \\ C_1 &= [18, [7, 0], 6, [2, 1], 8, [7, 2], 15, [3, -1]] \\ C_2 &= [9, [8, 0], 17, [3, 2], 1, [1, -1], 2, [7, -1]] \\ C_3 &= [13, [6, 0], 0, [6, 2]] \\ C_4 &= [11, [4, 0], 7, [4, 2]] \\ C_5 &= [5, [5, 0], 10, [5, 2]] \\ C'_0 &= [1, [3, 1], 17, [1, 0], 16, [4, -1], 12, [19, 1], 19, [19, 2], 2, [1, 1]] \\ C'_1 &= [11, [3, 0], 14, [6, -1], 8, [5, -1], 3, [8, 2]] \\ C'_2 &= [0, [9, 2], 9, [5, 1], 4, [9, 0], 13, [6, 1]] \\ C'_3 &= [7, [2, 0], 5, [2, 2]] \\ C'_4 &= [15, [9, -1], 6, [9, 1]] \\ C'_5 &= [10, [8, -1], 18, [8, 1]] \end{split}$$
e 2-factor type [5, 5, 4, 2, 2, 2]: two starters \\ C_0 &= [7, [3, 1], 10, [19, 1], 19, [19, -1], 18, [3, 2], 15, [8, 0]] \\ C_1 &= [17, [3, 0], 14, [9, -1], 5, [1, -1], 4, [2, -1], 2, [4, 2]] \\ C_2 &= [0, [3, -1], 3, [9, 0], 12, [4, -1], 8, [8, 2]] \\ C_3 &= [1, [5, 0], 6, [5, 2]] \\ C_4 &= [13, [2, 0], 11, [2, 2]] \\ C_5 &= [9, [7, -1], 16, [7, 1]] \end{aligned}

$$\begin{split} C_0' &= [3, [1, 1], 2, [6, -1], 8, [6, 0], 14, [9, 1], 4, [1, 2]] \\ C_1' &= [18, [2, 1], 1, [6, 2], 7, [4, 0], 11, [19, 2], 19, [19, 0]] \\ C_2' &= [9, [6, 1], 15, [4, 1], 0, [9, 2], 10, [1, 0]] \\ C_3' &= [12, [5, -1], 17, [5, 1]] \\ C_4' &= [13, [7, 0], 6, [7, 2]] \\ C_5' &= [16, [8, -1], 5, [8, 1]] \end{split}$$

• 2-factor type [7, 3, 3, 3, 2, 2]: one starter

$$\begin{split} C_0 &= [13, [6, 0], 7, [1, 0], 6, [7, 1], 18, [2, 0], 16, [6, 1], 10, [1, 1], 9, [4, 0]] \\ C_1 &= [5, [9, 1], 14, [7, 0], 2, [3, 0]] \\ C_2 &= [8, [4, 1], 12, [9, 0], 3, [5, 0]] \\ C_3 &= [17, [3, 1], 1, [5, 1], 15, [2, 1]] \\ C_4 &= [11, [8, 0], 0, [8, 1]] \\ C_5 &= [4, [19, 0], 19, [19, 1]] \end{split}$$

• 2-factor type [6, 4, 3, 3, 2, 2]: one starter

$$\begin{split} C_0 &= [16, [5, 0], 2, [8, 0], 13, [6, 0], 0, [9, 0], 9, [1, 1], 8, [8, 1]] \\ C_1 &= [15, [9, 1], 6, [2, 1], 4, [5, 1], 18, [3, 1]] \\ C_2 &= [7, [4, 0], 3, [2, 0], 1, [6, 1]] \\ C_3 &= [14, [3, 0], 11, [1, 0], 10, [4, 1]] \\ C_4 &= [12, [7, 0], 5, [7, 1]] \\ C_5 &= [17, [19, 0], 19, [19, 1]] \end{split}$$

• 2-factor type [5, 5, 3, 3, 2, 2]: one starter

$$\begin{split} C_0 &= [15, [6, 0], 2, [3, 1], 18, [7, 1], 11, [4, 0], 7, [8, 1]]\\ C_1 &= [13, [3, 0], 16, [4, 1], 1, [19, 0], 19, [19, 1], 4, [9, 1]]\\ C_2 &= [0, [6, 1], 6, [8, 0], 17, [2, 0]]\\ C_3 &= [5, [9, 0], 14, [2, 1], 12, [7, 0]]\\ C_4 &= [10, [1, 0], 9, [1, 1]]\\ C_5 &= [3, [5, 0], 8, [5, 1]] \end{split}$$

• 2-factor type [5, 4, 4, 3, 2, 2]: one starter

$$\begin{split} C_0 &= [9, [8, 0], 1, [1, 0], 0, [6, 1], 6, [8, 1], 14, [5, 1]] \\ C_1 &= [18, [6, 0], 12, [3, 1], 15, [1, 1], 16, [2, 0]] \\ C_2 &= [5, [3, 0], 2, [9, 0], 11, [4, 1], 7, [2, 1]] \\ C_3 &= [8, [9, 1], 17, [4, 0], 13, [5, 0]] \\ C_4 &= [4, [19, 0], 19, [19, 1]] \\ C_5 &= [10, [7, 0], 3, [7, 1]] \end{split}$$

• 2-factor type [4, 4, 4, 4, 2, 2]: one starter

$$\begin{split} C_0 &= [5, [8, 0], 16, [6, 1], 10, [7, 1], 17, [7, 0]] \\ C_1 &= [12, [6, 0], 18, [8, 1], 7, [5, 1], 2, [9, 0]] \\ C_2 &= [9, [5, 0], 4, [1, 1], 3, [9, 1], 13, [4, 0]] \\ C_3 &= [19, [19, 1], 1, [1, 0], 0, [4, 1], 15, [19, 0]] \\ C_4 &= [11, [3, 0], 14, [3, 1]] \\ C_5 &= [6, [2, 0], 8, [2, 1]] \end{split}$$

• 2-factor type [6, 3, 3, 3, 3, 2]: two starters

$$\begin{split} C_0 &= [4, [2, 1], 2, [8, 1], 13, [1, 1], 12, [6, -1], 18, [7, 0], 11, [7, 2]] \\ C_1 &= [16, [8, -1], 5, [9, -1], 14, [2, -1]] \\ C_2 &= [8, [2, 0], 6, [5, 2], 1, [7, -1]] \\ C_3 &= [7, [4, -1], 3, [7, 1], 10, [3, -1]] \\ C_4 &= [15, [4, 0], 0, [9, 1], 9, [6, 2]] \\ C_5 &= [17, [19, -1], 19, [19, 1]] \end{split}$$

- $C'_0 = [16, [8, 0], 8, [5, 1], 13, [9, 2], 4, [6, 0], 10, [4, 1], 14, [2, 2]]$ $C'_1 = [19, [19, 2], 3, [3, 1], 0, [19, 0]]$ $C'_{2} = [1, [8, 2], 9, [3, 0], 6, [5, -1]]$ $C'_3 = [15, [6, 1], 2, [3, 2], 5, [9, 0]]$ $C'_4 = [11, [4, 2], 7, [5, 0], 12, [1, -1]]$ $C'_{5} = [18, [1, 0], 17, [1, 2]]$
- 2-factor type [5, 4, 3, 3, 3, 2]: two starters

$$\begin{split} C_0 &= [11, [4, 2], 7, [1, 0], 8, [6, 2], 14, [7, -1], 2, [9, 0]] \\ C_1 &= [16, [19, 0], 19, [19, 2], 10, [2, -1], 12, [4, -1]] \\ C_2 &= [15, [9, 1], 6, [6, 1], 0, [4, 1]] \\ C_3 &= [17, [6, -1], 4, [5, 1], 18, [1, 1]] \\ C_4 &= [13, [7, 1], 1, [8, 2], 9, [4, 0]] \\ C_5 &= [3, [2, 0], 5, [2, 2]] \\ C_0' &= [18, [19, 1], 19, [19, -1], 8, [3, -1], 5, [9, -1], 15, [3, 1]] \\ C_1' &= [16, [7, 0], 9, [5, 2], 4, [3, 0], 7, [9, 2]] \\ C_2' &= [14, [2, 1], 12, [5, 0], 17, [3, 2]] \\ C_3' &= [1, [1, 2], 0, [6, 0], 6, [5, -1]] \\ C_4' &= [3, [7, 2], 10, [1, -1], 11, [8, 0]] \\ C_5' &= [13, [8, -1], 2, [8, 1]] \end{split}$$

• 2-factor type [4, 4, 4, 3, 3, 2]: two starters

 $C_0 = [16, [3, 0], 13, [8, 2], 2, [1, 0], 3, [6, 2]]$ $C_1 = [4, [9, 2], 14, [3, 1], 11, [2, 1], 9, [5, 0]]$ $C_2 = [7, [9, 0], 17, [5, -1], 12, [19, 2], 19, [19, 1]]$ $C'_2 = [4, [19, -1], 19, [19, 0], 3, [5, 1], 8, [4, 2]]$ $C_3 = [10, [5, 2], 15, [7, 1], 8, [2, 0]]$ $C_4 = [0, [6, 1], 6, [7, -1], 18, [1, 1]]$ $C_5 = [5, [4, -1], 1, [4, 1]]$

- $C'_0 = [12, [1, -1], 13, [6, 0], 7, [2, -1], 5, [7, 2]]$ $C'_1 = [2, [6, -1], 15, [1, 2], 14, [3, -1], 17, [4, 0]]$ $C'_{3} = [1, [2, 2], 18, [8, 0], 10, [9, -1]]$ $C'_4 = [6, [3, 2], 9, [7, 0], 16, [9, 1]]$ $C'_{5} = [0, [8, -1], 11, [8, 1]]$
- 2-factor type [5, 3, 3, 3, 3, 3]: one starter $C_0 = [1, [3, 0], 4, [6, 0], 10, [7, 0], 3, [9, 1], 12, [8, 0]]$

 $C_1 = [5, [7, 1], 17, [8, 1], 9, [4, 1]]$ $C_2 = [14, [19, 0], 19, [19, 1], 16, [2, 0]]$ $C_3 = [18, [3, 1], 2, [2, 1], 0, [1, 1]]$ $C_4 = [15, [9, 0], 6, [5, 1], 11, [4, 0]]$ $C_5 = [8, [5, 0], 13, [6, 1], 7, [1, 0]]$

• 2-factor type [4, 4, 3, 3, 3, 3]: one starter

$$\begin{split} C_0 &= [7, [2, 1], 5, [9, 1], 15, [5, 0], 10, [3, 0]]\\ C_1 &= [3, [1, 0], 4, [9, 0], 14, [5, 1], 9, [6, 1]]\\ C_2 &= [12, [7, 1], 0, [1, 1], 1, [8, 1]]\\ C_3 &= [2, [3, 1], 18, [7, 0], 6, [4, 0]]\\ C_4 &= [17, [4, 1], 13, [2, 0], 11, [6, 0]]\\ C_5 &= [8, [8, 0], 16, [19, 0], 19, [19, 1]] \end{split}$$

- 2-factor type [12, 2, 2, 2, 2]: one starter
 - $$\begin{split} C_0 &= [15, [8, 0], 7, [19, 1], 19, [19, 0], 4, [5, 0], 9, [4, 0], 13, [8, 1], 2, [3, 0], 18, [4, 1], 3, [3, 1], 6, [5, 1], 1, [7, 1], \\ &\quad 8, [7, 0]] \\ C_1 &= [11, [1, 0], 12, [1, 1]] \\ C_2 &= [5, [9, 0], 14, [9, 1]] \end{split}$$
 - $C_3 = [0, [2, 0], 17, [2, 1]]$
 - $C_4 = [10, [6, 0], 16, [6, 1]]$
- 2-factor type [11, 3, 2, 2, 2]: two starters
 - $C_{0} = \begin{bmatrix} 18, [5, 1], 13, [3, -1], 16, [8, 0], 5, [19, 2], 19, [19, 1], 11, [6, 0], 17, [5, 2], 3, [7, 1], 15, [7, 0], 8, [2, 1], 10, \\ \begin{bmatrix} 8, 2 \end{bmatrix} \end{bmatrix}$
 - $C_1 = [14, [5, -1], 9, [9, 2], 0, [5, 0]]$
 - $C_2 = [7, [6, -1], 1, [6, 1]]$
 - $C_3 = [2, [9, -1], 12, [9, 1]]$
 - $C_4 = [6, [2, 0], 4, [2, 2]]$
 - $$\begin{split} C_0' &= [0, [3, 1], 16, [9, 0], 6, [4, 2], 10, [8, 1], 2, [4, 0], 17, [7, 2], 5, [19, 0], 19, [19, -1], 11, [2, -1], 9, [8, -1], 1, \\ & [1, 2]] \\ C_1' &= [15, [7, -1], 8, [6, 2], 14, [1, 0]] \end{split}$$
 - $C_1 = [13, [7, -1], 3, [0, 2], 14, [1], C_2' = [3, [4, -1], 18, [4, 1]]$ $C_3' = [7, [3, 0], 4, [3, 2]]$
 - $C_4' = [13, [1, -1], 12, [1, 1]]$
- 2-factor type [10, 4, 2, 2, 2]: two starters

$$\begin{split} C_0 &= [18, [7, 0], 11, [8, 2], 0, [5, 1], 14, [7, -1], 2, [5, 0], 16, [8, 1], 5, [1, 2], 4, [9, -1], 13, [19, -1], 19, [19, 1]] \\ C_1 &= [10, [9, 2], 1, [8, 0], 12, [4, 2], 8, [2, 0]] \\ C_2 &= [17, [2, -1], 15, [2, 1]] \\ C_3 &= [9, [6, -1], 3, [6, 1]] \\ C_4 &= [6, [1, -1], 7, [1, 1]] \\ C_6' &= [6, [6, 0], 12, [2, 2], 10, [9, 0], 0, [6, 2], 13, [9, 1], 4, [19, 0], 19, [19, 2], 14, [7, 1], 7, [4, 0], 11, [5, 2]] \\ C_1' &= [16, [8, -1], 8, [5, -1], 3, [7, 2], 15, [1, 0]] \\ C_2' &= [17, [3, -1], 1, [3, 1]] \\ C_3' &= [2, [3, 0], 18, [3, 2]] \\ C_4' &= [9, [4, -1], 5, [4, 1]] \end{split}$$

• 2-factor type [9, 5, 2, 2, 2]: two starters

 $C_0 = [4, [1, 2], 3, [5, -1], 8, [9, 0], 17, [9, 2], 7, [6, 0], 13, [6, 1], 0, [7, 2], 12, [9, -1], 2, [2, 0]]$ $C_1 = [10, [5, 1], 5, [8, 1], 16, [1, 0], 15, [6, 2], 9, [1, -1]]$ $C_2 = [14, [4, -1], 18, [4, 1]]$ $C_3 = [1, [19, -1], 19, [19, 1]]$ $C_4 = [6, [5, 0], 11, [5, 2]]$ $C_0' = [3, [4, 0], 18, [19, 2], 19, [19, 0], 5, [9, 1], 14, [3, 2], 17, [8, 0], 6, [6, -1], 0, [1, 1], 1, [2, 2]]$ $C'_{1} = [8, [8, -1], 16, [7, 0], 4, [8, 2], 15, [3, 0], 12, [4, 2]]$ $C'_{2} = [13, [2, -1], 11, [2, 1]]$ $C'_{3} = [2, [7, -1], 9, [7, 1]]$ $C'_{4} = [7, [3, -1], 10, [3, 1]]$ • 2-factor type [8, 6, 2, 2, 2]: two starters $C_0 = [7, [4, 0], 11, [6, -1], 5, [7, 1], 12, [4, 2], 16, [4, -1], 1, [6, 0], 14, [7, 2], 2, [5, -1]]$ $C_1 = [4, [19, 0], 19, [19, 2], 6, [6, 1], 0, [8, 0], 8, [9, 1], 17, [6, 2]]$ $C_2 = [15, [3, -1], 18, [3, 1]]$ $C_3 = [9, [1, 0], 10, [1, 2]]$ $C_4 = [13, [9, 0], 3, [9, 2]]$ $C'_0 = [7, [19, 1], 19, [19, -1], 13, [7, -1], 6, [2, 1], 8, [3, 2], 11, [7, 0], 4, [8, 2], 12, [5, 0]]$ $C'_1 = [0, [5, 1], 5, [9, -1], 15, [5, 2], 1, [3, 0], 17, [4, 1], 2, [2, -1]]$

- $C'_2 = [3, [8, -1], 14, [8, 1]]$
- $C'_3 = [18, [2, 0], 16, [2, 2]]$
- $C'_4 = [10, [1, -1], 9, [1, 1]]$
- 2-factor type [7, 7, 2, 2, 2]: two starters

$$\begin{split} C_0 &= [13, [7, 2], 1, [8, 0], 9, [2, 1], 11, [19, 1], 19, [19, 2], 16, [8, 1], 8, [5, 0]] \\ C_1 &= [6, [4, -1], 10, [3, 0], 7, [4, 2], 3, [1, 0], 4, [8, 2], 15, [2, -1], 17, [8, -1]] \\ C_2 &= [14, [5, -1], 0, [5, 1]] \\ C_3 &= [18, [3, -1], 2, [3, 1]] \\ C_4 &= [5, [7, -1], 12, [7, 1]] \\ C_0' &= [13, [1, -1], 14, [4, 0], 10, [5, 2], 15, [7, 0], 3, [4, 1], 7, [9, 1], 16, [3, 2]] \\ C_1' &= [6, [2, 2], 4, [19, -1], 19, [19, 0], 1, [1, 2], 0, [9, -1], 9, [1, 1], 8, [2, 0]] \\ C_2' &= [18, [6, 0], 5, [6, 2]] \\ C_3' &= [17, [6, -1], 11, [6, 1]] \\ C_4' &= [12, [9, 0], 2, [9, 2]] \end{split}$$

• 2-factor type [10, 3, 3, 2, 2]: one starter

$$\begin{split} C_0 &= [6, [5, 1], 1, [2, 1], 18, [8, 1], 10, [2, 0], 8, [1, 1], 9, [6, 0], 3, [7, 1], 15, [9, 0], 5, [5, 0], 0, [6, 1]] \\ C_1 &= [13, [8, 0], 2, [9, 1], 12, [1, 0]] \\ C_2 &= [4, [19, 1], 19, [19, 0], 16, [7, 0]] \\ C_3 &= [11, [4, 0], 7, [4, 1]] \\ C_4 &= [17, [3, 0], 14, [3, 1]] \end{split}$$

• 2-factor type [9, 4, 3, 2, 2]: one starter

$$\begin{split} C_0 &= [19, [19, 1], 5, [7, 0], 17, [7, 1], 10, [3, 1], 13, [2, 1], 11, [1, 1], 12, [6, 0], 18, [2, 0], 16, [19, 0]] \\ C_1 &= [8, [4, 0], 4, [3, 0], 7, [4, 1], 3, [5, 1]] \\ C_2 &= [15, [5, 0], 1, [1, 0], 2, [6, 1]] \\ C_3 &= [9, [9, 0], 0, [9, 1]] \\ C_4 &= [6, [8, 0], 14, [8, 1]] \end{split}$$

• 2-factor type [8, 5, 3, 2, 2]: one starter

$$\begin{split} C_0 &= [17, [1, 0], 18, [6, 0], 5, [4, 1], 9, [7, 1], 16, [3, 1], 13, [6, 1], 7, [5, 0], 2, [4, 0]] \\ C_1 &= [0, [8, 1], 11, [1, 1], 10, [7, 0], 3, [5, 1], 8, [8, 0]] \\ C_2 &= [1, [19, 1], 19, [19, 0], 4, [3, 0]] \\ C_3 &= [6, [9, 0], 15, [9, 1]] \\ C_4 &= [14, [2, 0], 12, [2, 1]] \end{split}$$

• 2-factor type [7, 6, 3, 2, 2]: one starter

$$\begin{split} C_0 &= [12, [6, 0], 18, [8, 0], 7, [9, 0], 16, [7, 0], 4, [4, 1], 8, [9, 1], 17, [5, 1]] \\ C_1 &= [15, [1, 0], 14, [4, 0], 10, [5, 0], 5, [6, 1], 11, [2, 0], 13, [2, 1]] \\ C_2 &= [9, [7, 1], 2, [1, 1], 1, [8, 1]] \\ C_3 &= [6, [19, 0], 19, [19, 1]] \\ C_4 &= [0, [3, 0], 3, [3, 1]] \end{split}$$

• 2-factor type [8, 4, 4, 2, 2]: one starter

 $C_0 = [3, [2, 0], 1, [1, 0], 0, [9, 1], 9, [9, 0], 18, [5, 1], 13, [1, 1], 12, [4, 1], 8, [5, 0]]$

- $C_1 = [6, [2, 1], 4, [8, 1], 15, [6, 0], 2, [4, 0]]$
- $C_2 = [10, [7, 1], 17, [7, 0], 5, [8, 0], 16, [6, 1]]$
- $C_3 = [7, [19, 0], 19, [19, 1]]$
- $C_4 = [11, [3, 0], 14, [3, 1]]$
- 2-factor type [7, 5, 4, 2, 2]: one starter
 - $C_0 = [8, [3, 1], 11, [19, 0], 19, [19, 1], 4, [5, 1], 9, [7, 0], 2, [4, 1], 17, [9, 1]]$
 - $C_1 = [7, [3, 0], 10, [9, 0], 1, [1, 1], 0, [4, 0], 15, [8, 0]]$
 - $C_2 = [6, [1, 0], 5, [8, 1], 13, [5, 0], 18, [7, 1]]$
 - $C_3 = [3, [6, 0], 16, [6, 1]]$
 - $C_4 = [14, [2, 0], 12, [2, 1]]$
- 2-factor type [6, 6, 4, 2, 2]: one starter

$$\begin{split} C_0 &= [5, [5, 0], 0, [9, 1], 10, [6, 0], 16, [8, 0], 8, [9, 0], 18, [6, 1]]\\ C_1 &= [15, [19, 1], 19, [19, 0], 9, [5, 1], 4, [3, 1], 7, [4, 1], 11, [4, 0]]\\ C_2 &= [6, [7, 0], 13, [7, 1], 1, [3, 0], 17, [8, 1]]\\ C_3 &= [3, [1, 0], 2, [1, 1]]\\ C_4 &= [12, [2, 0], 14, [2, 1]] \end{split}$$

- 2-factor type [6, 5, 5, 2, 2]: one starter
 - $$\begin{split} C_0 &= [10, [8, 0], 2, [2, 1], 0, [6, 1], 6, [8, 1], 14, [1, 1], 15, [5, 0]]\\ C_1 &= [9, [4, 1], 5, [3, 1], 8, [3, 0], 11, [7, 0], 4, [5, 1]]\\ C_2 &= [13, [1, 0], 12, [6, 0], 18, [4, 0], 3, [2, 0], 1, [7, 1]]\\ C_3 &= [7, [9, 0], 17, [9, 1]]\\ C_4 &= [16, [19, 0], 19, [19, 1]] \end{split}$$
- 2-factor type [9, 3, 3, 3, 2]: two starters $C_0 = [19, [19, 2], 18, [8, -1], 7, [6, -1], 13, [3, 1], 10, [9, 0], 0, [5, 2], 5, [1, 1], 4, [4, 0], 8, [19, 1]]$ $C_1 = [1, [5, 0], 6, [9, -1], 16, [4, 2]]$ $C_2 = [15, [6, 1], 9, [8, 0], 17, [2, 2]]$ $C_3 = [11, [9, 1], 2, [1, 0], 3, [8, 2]]$ $C_4 = [12, [2, -1], 14, [2, 1]]$ $C'_0 = [4, [19, -1], 19, [19, 0], 16, [7, -1], 9, [3, -1], 6, [4, -1], 2, [1, -1], 1, [9, 2], 10, [7, 0], 17, [6, 2]]$ $C'_1 = [7, [5, -1], 12, [2, 0], 14, [7, 2]]$ $C'_2 = [15, [4, 1], 11, [8, 1], 3, [7, 1]]$ $C'_4 = [8, [3, 0], 5, [3, 2]]$
- 2-factor type [8, 4, 3, 3, 2]: two starters
 - $$\begin{split} C_0 &= [7, [1, 0], 6, [5, 1], 11, [8, 1], 0, [5, 2], 14, [1, 1], 13, [19, 0], 19, [19, -1], 15, [8, 2]] \\ C_1 &= [8, [9, 1], 18, [5, 0], 4, [2, 2], 2, [6, -1]] \\ C_2 &= [12, [4, 0], 16, [7, -1], 9, [3, 2]] \\ C_3 &= [17, [3, 0], 1, [4, 2], 5, [7, 1]] \\ C_4 &= [10, [7, 0], 3, [7, 2]] \\ C'_0 &= [10, [1, -1], 11, [5, -1], 16, [9, 0], 7, [1, 2], 6, [4, 1], 2, [8, 0], 13, [9, -1], 4, [6, 2]] \\ C'_1 &= [15, [19, 2], 19, [19, 1], 0, [2, 0], 17, [2, -1]] \\ C'_2 &= [9, [6, 0], 3, [4, -1], 18, [9, 2]] \\ C'_3 &= [12, [2, 1], 14, [6, 1], 1, [8, -1]] \\ C'_4 &= [8, [3, -1], 5, [3, 1]] \end{split}$$
- 2-factor type [7, 5, 3, 3, 2]: two starters

$$\begin{split} C_0 &= [3, [1, 0], 4, [4, 1], 8, [7, 2], 15, [9, 1], 6, [8, -1], 14, [1, 1], 13, [9, -1]] \\ C_1 &= [19, [19, 0], 2, [8, 1], 10, [1, -1], 9, [2, 2], 11, [19, -1]] \\ C_2 &= [7, [2, 1], 5, [6, 0], 18, [8, 2]] \\ C_3 &= [0, [3, 2], 16, [4, 0], 12, [7, 1]] \\ C_4 &= [1, [3, -1], 17, [3, 1]] \\ C_6 &= [17, [2, -1], 15, [1, 2], 14, [7, -1], 2, [3, 0], 18, [6, 1], 5, [5, 2], 10, [7, 0]] \\ C_1' &= [9, [6, -1], 3, [9, 2], 12, [5, 0], 7, [4, 2], 11, [2, 0]] \\ C_2' &= [13, [6, 2], 0, [4, -1], 4, [9, 0]] \\ C_3' &= [16, [8, 0], 8, [19, 2], 19, [19, 1]] \\ C_4' &= [6, [5, -1], 1, [5, 1]] \end{split}$$
• 2-factor type [6, 6, 3, 3, 2]: two starters

$$\begin{split} C_0 &= [4, [9, -1], 14, [8, 2], 3, [4, 1], 18, [7, 0], 11, [2, 2], 9, [5, 0]] \\ C_1 &= [13, [7, 1], 1, [7, 2], 8, [4, -1], 12, [2, 0], 10, [5, 2], 5, [8, 0]] \\ C_2 &= [2, [6, 2], 15, [4, 0], 0, [2, 1]] \\ C_3 &= [17, [1, -1], 16, [19, 2], 19, [19, 0]] \\ C_4 &= [6, [1, 0], 7, [1, 2]] \\ C_0' &= [10, [4, 2], 6, [5, 1], 1, [3, 0], 17, [6, 1], 4, [5, -1], 18, [8, -1]] \\ C_1' &= [9, [7, -1], 2, [19, 1], 19, [19, -1], 15, [1, 1], 16, [3, 1], 0, [9, 1]] \\ C_2' &= [7, [6, -1], 13, [8, 1], 5, [2, -1]] \\ C_3' &= [14, [3, -1], 11, [3, 2], 8, [6, 0]] \\ C_4' &= [3, [9, 0], 12, [9, 2]] \\ 2\text{-factor type } [7, 4, 4, 3, 2]: \text{ two starters} \\ C_0 &= [16, [1, 0], 17, [7, -1], 10, [19, 1], 19, [19, -1], 4, [9, 1], 13, [6, 2], 0, [3, 1]] \\ C_1 &= [14, [7, 1], 7, [2, 1], 5, [6, -1], 11, [3, -1]] \\ C_2 &= [8, [4, 1], 12, [9, 2], 2, [1, -1], 3, [5, 0]] \\ C_3 &= [6, [9, -1], 15, [6, 0], 9, [3, 2]] \\ C_4 &= [1, [2, 0], 18, [2, 2]] \\ C_0' &= [17, [1, 2], 18, [4, -1], 3, [6, 1], 16, [8, 0], 8, [8, 2], 0, [9, 0], 9, [8, 1]] \\ C_1' &= [6, [8, -1], 14, [1, 1], 13, [7, 0], 1, [5, 2]] \\ \end{split}$$

- $C'_{2} = [12, [7, 2], 5, [5, -1], 10, [3, 0], 7, [5, 1]]$
- $C_3' = [19, [19, 0], 4, [2, -1], 2, [19, 2]]$
- $C_4^\prime = [15, [4, 0], 11, [4, 2]]$
- 2-factor type [6, 5, 4, 3, 2]: two starters

$$\begin{split} C_0 &= [6, [4, 1], 10, [19, 2], 19, [19, 0], 13, [6, 2], 7, [7, 1], 0, [6, 0]] \\ C_1 &= [5, [4, -1], 1, [3, 1], 4, [5, 2], 18, [3, -1], 2, [3, 0]] \\ C_2 &= [12, [3, 2], 15, [7, 0], 3, [8, 2], 11, [1, 0]] \\ C_3 &= [14, [5, 1], 9, [7, -1], 16, [2, 1]] \\ C_4 &= [8, [9, -1], 17, [9, 1]] \\ C_6 &= [1, [1, 2], 0, [4, 0], 4, [1, 1], 5, [2, -1], 3, [7, 2], 15, [5, 0]] \\ C_1' &= [18, [5, -1], 13, [8, 1], 2, [6, 1], 8, [1, -1], 7, [8, -1]] \\ C_2' &= [12, [2, 0], 14, [4, 2], 10, [19, -1], 19, [19, 1]] \\ C_3' &= [17, [8, 0], 9, [2, 2], 11, [6, -1]] \\ C_4' &= [6, [9, 0], 16, [9, 2]] \end{split}$$

• 2-factor type [5, 5, 5, 3, 2]: two starters

$$\begin{split} C_0 &= [5, [1, -1], 6, [4, 1], 2, [5, 0], 7, [2, 1], 9, [4, 2]] \\ C_1 &= [10, [7, -1], 17, [1, 1], 16, [2, 0], 14, [5, 1], 0, [9, 2]] \\ C_2 &= [13, [5, -1], 8, [9, -1], 18, [19, 1], 19, [19, 2], 4, [9, 0]] \\ C_3 &= [1, [8, 1], 12, [3, 0], 15, [5, 2]] \\ C_4 &= [11, [8, 0], 3, [8, 2]] \end{split}$$

$$\begin{split} C_0' &= [14, [19, 0], 19, [19, -1], 10, [1, 2], 11, [6, -1], 17, [3, 1]] \\ C_1' &= [3, [4, 0], 18, [3, 2], 2, [2, -1], 4, [1, 0], 5, [2, 2]] \\ C_2' &= [1, [8, -1], 9, [4, -1], 13, [3, -1], 16, [9, 1], 7, [6, 1]] \\ C_3' &= [12, [6, 0], 6, [6, 2], 0, [7, 1]] \\ C_4' &= [15, [7, 0], 8, [7, 2]] \end{split}$$

• 2-factor type [6, 4, 4, 4, 2]: two starters

$$\begin{split} C_0 &= [15, [19, 0], 19, [19, 2], 12, [8, 0], 4, [3, 2], 1, [1, 0], 2, [6, 2]] \\ C_1 &= [14, [5, -1], 0, [2, 1], 17, [9, -1], 7, [7, -1]] \\ C_2 &= [6, [5, 2], 11, [2, -1], 13, [3, -1], 16, [9, 0]] \\ C_3 &= [18, [6, 1], 5, [4, 2], 9, [6, 0], 3, [4, -1]] \\ C_4 &= [10, [2, 0], 8, [2, 2]] \\ C_0' &= [13, [8, 2], 5, [4, 0], 9, [8, -1], 1, [7, 2], 8, [9, 1], 18, [5, 0]] \\ C_1' &= [17, [6, -1], 4, [7, 1], 16, [19, -1], 19, [19, 1]] \\ C_2' &= [10, [9, 2], 0, [3, 0], 3, [3, 1], 6, [4, 1]] \\ C_3' &= [14, [7, 0], 2, [5, 1], 7, [8, 1], 15, [1, 2]] \\ C_4' &= [11, [1, -1], 12, [1, 1]] \end{split}$$

- 2-factor type [5, 5, 4, 4, 2]: two starters
 - $$\begin{split} C_0 &= [0, [8, 1], 8, [1, 1], 9, [4, -1], 5, [4, 2], 1, [1, 0]] \\ C_1 &= [4, [7, 0], 11, [5, 2], 6, [1, -1], 7, [19, -1], 19, [19, 1]] \\ C_2 &= [10, [4, 0], 14, [1, 2], 13, [8, 0], 2, [8, 2]] \\ C_3 &= [18, [2, 1], 16, [4, 1], 12, [3, 0], 15, [3, 2]] \\ C_4 &= [3, [5, -1], 17, [5, 1]] \\ C_0' &= [4, [7, 2], 11, [19, 0], 19, [19, 2], 12, [6, 0], 6, [2, -1]] \\ C_1' &= [16, [5, 0], 2, [2, 2], 0, [8, -1], 8, [2, 0], 10, [6, 2]] \\ C_2' &= [18, [3, -1], 15, [7, 1], 3, [6, 1], 9, [9, 1]] \\ C_3' &= [7, [9, -1], 17, [3, 1], 1, [7, -1], 13, [6, -1]] \\ C_4' &= [5, [9, 0], 14, [9, 2]] \end{split}$$
- 2-factor type [8, 3, 3, 3, 3]: one starter

$$\begin{split} C_0 &= [13, [5, 0], 8, [8, 0], 0, [7, 1], 7, [8, 1], 15, [1, 0], 14, [3, 0], 11, [6, 0], 17, [4, 0]] \\ C_1 &= [3, [19, 0], 19, [19, 1], 1, [2, 0]] \\ C_2 &= [9, [1, 1], 10, [5, 1], 5, [4, 1]] \\ C_3 &= [4, [2, 1], 6, [9, 0], 16, [7, 0]] \\ C_4 &= [18, [6, 1], 12, [9, 1], 2, [3, 1]] \end{split}$$

• 2-factor type [7, 4, 3, 3, 3]: one starter

$$\begin{split} C_0 &= [12, [4, 1], 8, [8, 0], 16, [5, 0], 2, [1, 1], 3, [1, 0], 4, [19, 0], 19, [19, 1]] \\ C_1 &= [18, [9, 0], 9, [9, 1], 0, [6, 1], 6, [7, 0]] \\ C_2 &= [7, [7, 1], 14, [4, 0], 10, [3, 0]] \\ C_3 &= [15, [5, 1], 1, [3, 1], 17, [2, 1]] \\ C_4 &= [5, [6, 0], 11, [2, 0], 13, [8, 1]] \end{split}$$

• 2-factor type [6, 5, 3, 3, 3]: one starter

$$\begin{split} C_0 &= [12, [7, 1], 0, [6, 0], 13, [9, 1], 4, [3, 0], 1, [5, 1], 6, [6, 1]] \\ C_1 &= [14, [4, 0], 18, [3, 1], 15, [4, 1], 11, [8, 1], 3, [8, 0]] \\ C_2 &= [7, [2, 1], 5, [7, 0], 17, [9, 0]] \\ C_3 &= [2, [19, 1], 19, [19, 0], 16, [5, 0]] \\ C_4 &= [10, [2, 0], 8, [1, 1], 9, [1, 0]] \end{split}$$

• 2-factor type [6, 4, 4, 3, 3]: one starter

$$\begin{split} C_0 &= [16, [7, 0], 9, [8, 0], 1, [2, 1], 18, [5, 0], 13, [2, 0], 15, [1, 1]] \\ C_1 &= [4, [6, 1], 10, [5, 1], 5, [3, 0], 8, [4, 0]] \\ C_2 &= [6, [4, 1], 2, [19, 0], 19, [19, 1], 0, [6, 0]] \\ C_3 &= [17, [3, 1], 14, [7, 1], 7, [9, 1]] \\ C_4 &= [12, [1, 0], 11, [8, 1], 3, [9, 0]] \end{split}$$

- 2-factor type [5, 5, 4, 3, 3]: one starter
 - $C_0 = [0, [2, 1], 17, [19, 1], 19, [19, 0], 9, [4, 0], 13, [6, 1]]$
 - $C_1 = [16, [6, 0], 3, [4, 1], 7, [3, 0], 10, [8, 1], 2, [5, 1]]$
 - $C_2 = [8, [7, 1], 1, [5, 0], 6, [7, 0], 18, [9, 1]]$
 - $C_3 = [11, [3, 1], 14, [2, 0], 12, [1, 0]]$
 - $C_4 = [15, [9, 0], 5, [1, 1], 4, [8, 0]]$
- 2-factor type [5, 4, 4, 4, 3]: one starter
 - $C_0 = [11, [9, 1], 1, [1, 1], 2, [6, 1], 8, [7, 1], 15, [4, 1]]$
 - $C_1 = [14, [2, 1], 12, [8, 1], 4, [3, 0], 7, [7, 0]]$
 - $C_2 = [16, [2, 0], 18, [4, 0], 3, [5, 0], 17, [1, 0]]$
 - $C_3 = [10, [19, 0], 19, [19, 1], 13, [8, 0], 5, [5, 1]]$
 - $C_4 = [9, [3, 1], 6, [6, 0], 0, [9, 0]]$
- 2-factor type [14, 2, 2, 2]: two starters

$$\begin{split} C_0 &= [7, [2, 2], 5, [19, 0], 19, [19, 2], 12, [9, 0], 3, [4, 2], 18, [1, 0], 17, [3, 2], 14, [5, -1], 0, [6, -1], 6, [2, 0], 4, \\ &\quad [4, 1], 8, [8, 1], 16, [1, 2], 15, [8, 0]] \\ C_1 &= [10, [3, -1], 13, [3, 1]] \end{split}$$

- $C_2 = [11, [2, -1], 9, [2, 1]]$
- $C_3 = [1, [1, -1], 2, [1, 1]]$

- $$\begin{split} C_0' &= [18, [4, -1], 14, [19, 1], 19, [19, -1], 12, [5, 2], 7, [4, 0], 11, [6, 1], 17, [9, 2], 8, [8, -1], 16, [7, 0], 4, [5, 1], \\ &9, [7, 2], 2, [3, 0], 5, [8, 2], 13, [5, 0]] \\ C_1' &= [6, [6, 0], 0, [6, 2]] \\ C_2' &= [15, [7, -1], 3, [7, 1]] \\ C_3' &= [10, [9, -1], 1, [9, 1]] \end{split}$$
- 2-factor type [13, 3, 2, 2]: one starter
 - $$\begin{split} C_0 &= [11, [3, 0], 14, [1, 1], 13, [5, 1], 18, [4, 1], 3, [3, 1], 0, [7, 0], 7, [8, 1], 15, [5, 0], 1, [4, 0], 5, [7, 1], 12, [9, 0], \\ &\quad 2, [19, 1], 19, [19, 0]] \\ C_1 &= [9, [1, 0], 8, [9, 1], 17, [8, 0]] \end{split}$$
 - $C_2 = [10, [6, 0], 16, [6, 1]]$
 - $C_3 = [4, [2, 0], 6, [2, 1]]$
- 2-factor type [12, 4, 2, 2]: one starter
 - $$\begin{split} C_0 = [10, [2, 0], 12, [3, 1], 9, [6, 0], 3, [5, 0], 8, [2, 1], 6, [4, 0], 2, [5, 1], 16, [8, 0], 5, [19, 1], 19, [19, 0], 11, [6, 1], \\ 17, [7, 1]] \end{split}$$
 - $C_1 = [14, [7, 0], 7, [8, 1], 15, [3, 0], 18, [4, 1]]$
 - $C_2 = [4, [9, 0], 13, [9, 1]]$
 - $C_3 = [1, [1, 0], 0, [1, 1]]$
- 2-factor type [11, 5, 2, 2]: one starter

$$\begin{split} C_0 &= [11, [9, 1], 2, [7, 1], 14, [8, 1], 6, [7, 0], 18, [3, 0], 15, [9, 0], 5, [5, 0], 10, [2, 1], 12, [3, 1], 9, [4, 0], 13, [2, 0]] \\ C_1 &= [17, [6, 0], 4, [4, 1], 8, [8, 0], 16, [6, 1], 3, [5, 1]] \\ C_2 &= [1, [1, 0], 0, [1, 1]] \\ C_3 &= [7, [19, 0], 19, [19, 1]] \end{split}$$

• 2-factor type [10, 6, 2, 2]: one starter

$$\begin{split} C_0 &= [4, [8, 0], 12, [9, 0], 3, [8, 1], 14, [7, 0], 2, [4, 0], 17, [1, 0], 16, [4, 1], 1, [2, 1], 18, [19, 0], 19, [19, 1]] \\ C_1 &= [9, [6, 0], 15, [2, 0], 13, [6, 1], 7, [7, 1], 0, [9, 1], 10, [1, 1]] \\ C_2 &= [8, [3, 0], 5, [3, 1]] \\ C_3 &= [6, [5, 0], 11, [5, 1]] \end{split}$$

• 2-factor type [9, 7, 2, 2]: one starter

$$\begin{split} C_0 &= [0, [9, 1], 10, [5, 0], 5, [6, 1], 11, [9, 0], 1, [3, 0], 4, [5, 1], 9, [7, 0], 2, [3, 1], 18, [1, 1]]\\ C_1 &= [7, [1, 0], 6, [19, 0], 19, [19, 1], 3, [6, 0], 16, [8, 0], 8, [7, 1], 15, [8, 1]]\\ C_2 &= [13, [4, 0], 17, [4, 1]] \end{split}$$

- $C_3 = [14, [2, 0], 12, [2, 1]]$
- 2-factor type [8, 8, 2, 2]: one starter
 - $C_0 = [11, [8, 1], 3, [2, 0], 1, [6, 1], 7, [5, 0], 12, [6, 0], 6, [7, 0], 13, [2, 1], 15, [4, 1]]$
 - $C_1 = [18, [19, 0], 19, [19, 1], 0, [3, 0], 16, [7, 1], 9, [5, 1], 14, [4, 0], 10, [8, 0], 2, [3, 1]]$
 - $C_2 = [8, [9, 0], 17, [9, 1]]$
 - $C_3 = [4, [1, 0], 5, [1, 1]]$

• 2-factor type [12, 3, 3, 2]: two starters

$$\begin{split} &C_0 = [7, [5, 2], 12, [2, 0], 10, [4, 2], 14, [1, 0], 15, [8, 1], 4, [7, -1], 16, [3, 2], 13, [7, 0], 6, [4, -1], 2, [6, 2], 8, \\ & [9, 0], 17, [9, 1]] \\ &C_1 = [1, [4, 0], 5, [6, -1], 18, [2, 2]] \\ &C_2 = [9, [2, -1], 11, [8, 0], 0, [9, 2]] \\ &C_3 = [19, [19, 0], 3, [19, 2]] \\ &C_0' = [16, [4, 1], 1, [5, -1], 15, [7, 2], 8, [9, -1], 17, [5, 0], 3, [1, 2], 4, [1, -1], 5, [6, 0], 11, [8, 2], 0, [1, 1], 18, \\ & [5, 1], 13, [3, 0]] \\ &C_1' = [12, [6, 1], 6, [8, -1], 14, [2, 1]] \\ &C_2' = [19, [19, -1], 2, [7, 1], 9, [19, 1]] \\ &C_3' = [7, [3, -1], 10, [3, 1]] \\ &2\text{-factor type } [11, 4, 3, 2]: \text{ two starters} \\ &C_0 = [1, [1, 2], 2, [3, 0], 5, [1, -1], 4, [2, 2], 6, [5, -1], 11, [3, 1], 8, [9, 0], 17, [19, 1], 19, [19, 2], 18, [5, 1], 13, \\ & [7, 0]] \\ &C_1 = [10, [2, -1], 12, [7, -1], 0, [5, 0], 14, [4, 2]] \\ &C_2 = [9, [7, 2], 16, [6, -1], 3, [6, 0]] \\ &C_3 = [15, [8, -1], 7, [8, 1]] \\ &C_0' = [2, [5, 2], 7, [1, 0], 8, [4, -1], 4, [7, 1], 11, [8, 2], 0, [6, 1], 13, [3, -1], 16, [2, 0], 14, [1, 1], 15, [2, 1], 17, \\ & [4, 1]] \\ &C_1' = [6, [4, 0], 10, [9, 2], 1, [8, 0], 9, [3, 2]] \\ &C_2' = [18, [6, 2], 5, [19, 0], 19, [19, -1]] \\ \end{split}$$

- $C_3' = [12, [9, -1], 3, [9, 1]]$
- 2-factor type [10, 5, 3, 2]: two starters

$$\begin{split} C_0 &= [9, [5, -1], 14, [4, 0], 18, [1, -1], 17, [8, 1], 6, [2, 1], 8, [7, 2], 1, [2, 0], 3, [3, 2], 0, [3, 1], 16, [7, 1]] \\ C_1 &= [4, [3, 0], 7, [5, 2], 12, [3, -1], 15, [5, 0], 10, [6, 2]] \\ C_2 &= [2, [8, -1], 13, [19, 1], 19, [19, -1]] \\ C_3 &= [5, [6, -1], 11, [6, 1]] \\ C'_0 &= [2, [4, 1], 17, [8, 0], 6, [1, 2], 5, [9, 0], 15, [4, 2], 11, [7, 0], 4, [9, -1], 14, [2, 2], 16, [9, 1], 7, [5, 1]] \\ C'_1 &= [3, [4, -1], 18, [9, 2], 8, [2, -1], 10, [1, 1], 9, [6, 0]] \\ C'_2 &= [0, [1, 0], 1, [8, 2], 12, [7, -1]] \\ C'_3 &= [13, [19, 0], 19, [19, 2]] \end{split}$$
• 2-factor type [9, 6, 3, 2]: two starters

$$\begin{split} C_0 &= [13, [9, 0], 4, [1, 1], 5, [5, 2], 0, [7, 1], 12, [8, -1], 1, [2, 1], 18, [4, 0], 3, [7, 2], 10, [3, -1]] \\ C_1 &= [7, [1, 0], 8, [8, 1], 16, [9, 1], 6, [5, 1], 11, [19, 2], 19, [19, 1]] \\ C_2 &= [14, [5, 0], 9, [6, 1], 15, [1, 2]] \\ C_3 &= [17, [4, -1], 2, [4, 1]] \end{split}$$

- $$\begin{split} C_0' &= [14, [5, -1], 9, [6, 2], 15, [9, -1], 5, [7, 0], 12, [4, 2], 16, [3, 0], 0, [6, -1], 13, [9, 2], 3, [8, 0]] \\ C_1' &= [8, [6, 0], 2, [8, 2], 10, [7, -1], 17, [19, 0], 19, [19, -1], 11, [3, 2]] \\ C_2' &= [7, [3, 1], 4, [2, -1], 6, [1, -1]] \\ C_3' &= [18, [2, 0], 1, [2, 2]] \end{split}$$
- 2-factor type [8, 7, 3, 2]: two starters

$$\begin{split} C_0 &= [19, [19, 0], 17, [4, -1], 13, [8, 2], 5, [9, 0], 15, [5, 1], 1, [2, -1], 18, [1, 1], 0, [19, 2]] \\ C_1 &= [11, [1, 0], 10, [3, 2], 7, [5, -1], 12, [8, 0], 4, [5, 2], 9, [6, 1], 3, [8, -1]] \\ C_2 &= [8, [6, 0], 14, [2, 2], 16, [8, 1]] \\ C_3 &= [2, [4, 0], 6, [4, 2]] \\ C'_0 &= [10, [6, 2], 16, [6, -1], 3, [7, 0], 15, [7, 2], 8, [9, -1], 17, [1, -1], 18, [5, 0], 13, [3, -1]] \\ C'_1 &= [2, [4, 1], 6, [1, 2], 7, [2, 1], 5, [7, -1], 12, [3, 1], 9, [9, 1], 0, [2, 0]] \\ C'_2 &= [14, [19, -1], 19, [19, 1]] \end{split}$$

• 2-factor type [10, 4, 4, 2]: two starters

$$\begin{split} C_0 &= [3, [6, 0], 9, [4, -1], 5, [3, 2], 2, [8, 0], 10, [2, -1], 8, [9, 1], 17, [5, -1], 12, [8, 2], 1, [5, 0], 15, [7, 2]] \\ C_1 &= [13, [2, 0], 11, [4, 2], 7, [7, 0], 14, [1, 2]] \\ C_2 &= [6, [19, 2], 19, [19, 0], 18, [5, 1], 4, [2, 1]] \\ C_3 &= [0, [3, -1], 16, [3, 1]] \\ C_0' &= [15, [3, 0], 12, [8, -1], 4, [6, 2], 10, [4, 0], 14, [7, -1], 7, [1, -1], 6, [19, 1], 19, [19, -1], 16, [8, 1], 5, [9, 2]] \\ C_1' &= [11, [9, -1], 1, [1, 0], 2, [7, 1], 9, [2, 2]] \\ C_2' &= [17, [1, 1], 18, [4, 1], 3, [5, 2], 8, [9, 0]] \\ C_3' &= [13, [6, -1], 0, [6, 1]] \end{split}$$

• 2-factor type [9, 5, 4, 2]: two starters

$$\begin{split} C_0 &= [13, [9, 0], 4, [7, -1], 16, [8, -1], 8, [3, 2], 11, [1, -1], 12, [3, -1], 15, [1, 1], 14, [6, -1], 1, [7, 1]] \\ C_1 &= [3, [3, 0], 6, [8, 2], 17, [4, 0], 2, [3, 1], 5, [2, 2]] \\ C_2 &= [18, [9, -1], 9, [2, -1], 7, [7, 2], 0, [1, 0]] \\ C_3 &= [10, [19, -1], 19, [19, 1]] \\ C_6 &= [15, [9, 1], 5, [4, -1], 1, [5, 1], 6, [6, 2], 0, [6, 0], 13, [1, 2], 14, [2, 1], 12, [8, 1], 4, [8, 0]] \\ C_1' &= [16, [5, 2], 11, [7, 0], 18, [9, 2], 8, [2, 0], 10, [6, 1]] \\ C_2' &= [17, [5, 0], 3, [4, 1], 7, [5, -1], 2, [4, 2]] \\ C_3' &= [19, [19, 0], 9, [19, 2]] \end{split}$$

- 2-factor type [8, 6, 4, 2]: two starters
 - $\begin{aligned} C_0 &= [0, [1, 2], 1, [3, 1], 17, [7, 0], 5, [8, 2], 16, [19, 0], 19, [19, 2], 13, [2, 0], 15, [4, 1]] \\ C_1 &= [18, [7, -1], 11, [1, -1], 10, [8, -1], 2, [4, 2], 6, [3, -1], 9, [9, 0]] \\ C_2 &= [12, [9, -1], 3, [5, 2], 8, [6, 0], 14, [2, -1]] \\ C_3 &= [7, [3, 0], 4, [3, 2]] \end{aligned}$

- $$\begin{split} C_0' &= [13, [7, 2], 6, [19, -1], 19, [19, 1], 16, [2, 1], 14, [5, 0], 0, [8, 1], 8, [4, -1], 4, [9, 1]] \\ C_1' &= [2, [9, 2], 11, [1, 0], 10, [5, 1], 15, [6, 2], 9, [8, 0], 1, [1, 1]] \\ C_2' &= [3, [4, 0], 7, [2, 2], 5, [7, 1], 17, [5, -1]] \\ C_3' &= [12, [6, -1], 18, [6, 1]] \end{split}$$
- 2-factor type [7, 7, 4, 2]: two starters

$$\begin{split} C_0 &= [17, [5, -1], 3, [3, 0], 6, [9, 1], 15, [6, 2], 2, [19, -1], 19, [19, 0], 8, [9, 2]] \\ C_1 &= [16, [5, 1], 11, [4, 1], 7, [7, 0], 0, [9, -1], 10, [3, 2], 13, [5, 0], 18, [2, 2]] \\ C_2 &= [1, [8, 2], 12, [8, 1], 4, [9, 0], 14, [6, 1]] \\ C_3 &= [5, [4, 0], 9, [4, 2]] \\ C_0' &= [11, [8, -1], 3, [7, 2], 15, [6, 0], 9, [2, -1], 7, [6, -1], 1, [3, 1], 4, [7, 1]] \\ C_1' &= [12, [7, -1], 5, [1, 1], 6, [4, -1], 2, [3, -1], 18, [5, 2], 13, [1, -1], 14, [2, 0]] \\ C_2' &= [19, [19, 1], 0, [8, 0], 8, [2, 1], 10, [19, 2]] \\ C_3' &= [17, [1, 0], 16, [1, 2]] \end{split}$$

• 2-factor type [8, 5, 5, 2]: two starters

$$\begin{split} C_0 &= [17, [3, -1], 14, [5, 2], 9, [7, 1], 2, [6, -1], 8, [4, 0], 4, [3, 1], 7, [8, 1], 15, [2, 1]] \\ C_1 &= [13, [9, -1], 3, [6, 0], 16, [8, 2], 5, [19, -1], 19, [19, 1]] \\ C_2 &= [10, [9, 1], 0, [1, 1], 1, [8, -1], 12, [1, 2], 11, [1, 0]] \\ C_3 &= [18, [7, 0], 6, [7, 2]] \\ C_0' &= [0, [5, -1], 5, [7, -1], 12, [5, 0], 17, [9, 2], 7, [8, 0], 15, [4, 2], 11, [5, 1], 6, [6, 1]] \\ C_1' &= [8, [9, 0], 18, [2, 2], 1, [3, 0], 4, [2, -1], 2, [6, 2]] \\ C_2' &= [16, [2, 0], 14, [4, -1], 10, [1, -1], 9, [4, 1], 13, [3, 2]] \\ C_3' &= [19, [19, 0], 3, [19, 2]] \end{split}$$

• 2-factor type [7, 6, 5, 2]: two starters

$$\begin{split} C_0 &= [19, [19, -1], 18, [7, 1], 6, [5, -1], 1, [3, 2], 4, [5, 0], 9, [3, 1], 12, [19, 1]] \\ C_1 &= [10, [6, -1], 16, [6, 0], 3, [3, -1], 0, [2, 2], 17, [4, 0], 2, [8, 2]] \\ C_2 &= [5, [2, 1], 7, [1, 2], 8, [3, 0], 11, [4, 2], 15, [9, 0]] \\ C_3 &= [14, [1, -1], 13, [1, 1]] \\ C_0' &= [6, [6, 1], 12, [5, 2], 7, [19, 0], 19, [19, 2], 15, [4, -1], 0, [8, -1], 8, [2, 0]] \\ C_1' &= [2, [9, 1], 11, [2, -1], 9, [5, 1], 14, [8, 1], 3, [9, 2], 13, [8, 0]] \\ C_2' &= [5, [6, 2], 18, [1, 0], 17, [7, -1], 10, [9, -1], 1, [4, 1]] \\ C_3' &= [4, [7, 0], 16, [7, 2]] \end{split}$$

- 2-factor type [6, 6, 6, 2]: two starters
 - $C_{0} = [8, [4, 2], 4, [7, 0], 16, [4, -1], 12, [1, 2], 13, [9, -1], 3, [5, 0]]$ $C_{1} = [19, [19, -1], 5, [7, 2], 17, [7, 1], 10, [8, 0], 18, [7, -1], 11, [19, 1]]$ $C_{2} = [15, [4, 1], 0, [5, -1], 14, [8, 2], 6, [1, 0], 7, [5, 2], 2, [6, 0]]$ $C_{3} = [1, [8, -1], 9, [8, 1]]$

$$\begin{split} C_0' &= [8, [6, 1], 2, [2, 1], 0, [6, -1], 6, [1, -1], 7, [2, -1], 9, [1, 1]] \\ C_1' &= [1, [9, 2], 11, [3, -1], 14, [9, 0], 5, [5, 1], 10, [6, 2], 4, [3, 0]] \\ C_2' &= [3, [4, 0], 18, [2, 2], 16, [3, 1], 13, [2, 0], 15, [3, 2], 12, [9, 1]] \\ C_3' &= [19, [19, 0], 17, [19, 2]] \end{split}$$

• 2-factor type [11, 3, 3, 3]: one starter

$$\begin{split} C_0 &= [19, [19, 1], 7, [9, 1], 16, [8, 1], 8, [7, 0], 15, [6, 1], 9, [3, 0], 12, [1, 0], 13, [7, 1], 1, [2, 1], 18, [1, 1], 0, [19, 0]] \\ C_1 &= [11, [8, 0], 3, [3, 1], 6, [5, 0]] \\ C_2 &= [4, [6, 0], 17, [4, 1], 2, [2, 0]] \\ C_3 &= [5, [9, 0], 14, [4, 0], 10, [5, 1]] \end{split}$$

• 2-factor type [10, 4, 3, 3]: one starter

$$\begin{split} C_0 &= [7, [2, 1], 5, [4, 1], 9, [6, 1], 15, [1, 0], 14, [4, 0], 10, [2, 0], 12, [8, 0], 4, [1, 1], 3, [5, 1], 17, [9, 1]] \\ C_1 &= [19, [19, 1], 0, [6, 0], 6, [7, 1], 18, [19, 0]] \\ C_2 &= [8, [7, 0], 1, [9, 0], 11, [3, 1]] \\ C_3 &= [2, [5, 0], 16, [3, 0], 13, [8, 1]] \end{split}$$

• 2-factor type [9, 5, 3, 3]: one starter

$$\begin{split} C_0 &= [2, [6, 1], 8, [7, 0], 1, [6, 0], 14, [1, 1], 15, [3, 1], 18, [2, 0], 16, [7, 1], 4, [2, 1], 6, [4, 0]] \\ C_1 &= [5, [5, 1], 10, [9, 0], 0, [3, 0], 3, [9, 1], 13, [8, 1]] \\ C_2 &= [11, [4, 1], 7, [5, 0], 12, [1, 0]] \\ C_3 &= [19, [19, 0], 17, [8, 0], 9, [19, 1]] \end{split}$$

• 2-factor type [8, 6, 3, 3]: one starter

$$\begin{split} C_0 &= [2, [9, 0], 11, [3, 0], 8, [8, 0], 0, [2, 1], 17, [7, 1], 5, [9, 1], 14, [6, 0], 1, [1, 1]] \\ C_1 &= [6, [4, 0], 10, [3, 1], 13, [2, 0], 15, [8, 1], 7, [5, 1], 12, [6, 1]] \\ C_2 &= [9, [7, 0], 16, [19, 1], 19, [19, 0]] \\ C_3 &= [4, [5, 0], 18, [4, 1], 3, [1, 0]] \end{split}$$

- 2-factor type [7, 7, 3, 3]: one starter
 - $$\begin{split} C_0 &= [17, [1, 0], 16, [3, 0], 0, [9, 1], 9, [4, 1], 13, [6, 0], 7, [4, 0], 11, [6, 1]] \\ C_1 &= [5, [2, 0], 3, [8, 1], 14, [7, 0], 2, [9, 0], 12, [19, 0], 19, [19, 1], 4, [1, 1]] \\ C_2 &= [15, [3, 1], 18, [8, 0], 10, [5, 0]] \\ C_3 &= [6, [2, 1], 8, [7, 1], 1, [5, 1]] \end{split}$$
- 2-factor type [9, 4, 4, 3]: one starter
 - $C_0 = [11, [9, 0], 2, [6, 1], 8, [7, 1], 1, [8, 1], 12, [3, 0], 9, [5, 0], 14, [8, 0], 6, [1, 1], 7, [4, 1]]$
 - $C_1 = [0, [9, 1], 10, [19, 1], 19, [19, 0], 15, [4, 0]]$
 - $C_2 = [5, [7, 0], 17, [6, 0], 4, [1, 0], 3, [2, 0]]$
 - $C_3 = [13, [5, 1], 18, [2, 1], 16, [3, 1]]$

• 2-factor type [8, 5, 4, 3]: one starter

$$\begin{split} C_0 &= [15, [9, 1], 5, [8, 1], 16, [9, 0], 7, [1, 0], 6, [6, 1], 0, [5, 0], 14, [6, 0], 8, [7, 1]] \\ C_1 &= [17, [4, 0], 13, [19, 0], 19, [19, 1], 2, [8, 0], 10, [7, 0]] \\ C_2 &= [1, [2, 1], 3, [4, 1], 18, [5, 1], 4, [3, 1]] \\ C_3 &= [12, [1, 1], 11, [2, 0], 9, [3, 0]] \end{split}$$

• 2-factor type [7, 6, 4, 3]: one starter

$$\begin{split} C_0 &= [0, [5, 0], 14, [3, 1], 11, [5, 1], 6, [6, 1], 12, [3, 0], 9, [4, 0], 13, [6, 0]] \\ C_1 &= [18, [9, 1], 8, [7, 1], 1, [4, 1], 16, [1, 0], 17, [7, 0], 10, [8, 1]] \\ C_2 &= [4, [1, 1], 3, [19, 1], 19, [19, 0], 2, [2, 0]] \\ C_3 &= [15, [9, 0], 5, [2, 1], 7, [8, 0]] \end{split}$$

• 2-factor type [7, 5, 5, 3]: one starter

$$\begin{split} C_0 &= [15, [5, 0], 10, [19, 1], 19, [19, 0], 4, [5, 1], 9, [7, 1], 16, [3, 1], 13, [2, 1]] \\ C_1 &= [18, [6, 0], 5, [4, 1], 1, [1, 0], 0, [2, 0], 2, [3, 0]] \\ C_2 &= [12, [4, 0], 8, [9, 0], 17, [6, 1], 11, [8, 1], 3, [9, 1]] \\ C_3 &= [7, [7, 0], 14, [8, 0], 6, [1, 1]] \end{split}$$

• 2-factor type [6, 6, 5, 3]: one starter

$$\begin{split} C_0 &= [14, [8, 0], 3, [6, 1], 16, [3, 0], 13, [19, 0], 19, [19, 1], 10, [4, 0]] \\ C_1 &= [4, [7, 0], 11, [2, 0], 9, [2, 1], 7, [9, 1], 17, [5, 1], 12, [8, 1]] \\ C_2 &= [18, [7, 1], 6, [1, 1], 5, [3, 1], 2, [6, 0], 8, [9, 0]] \\ C_3 &= [0, [4, 1], 15, [5, 0], 1, [1, 0]] \end{split}$$

• 2-factor type [7, 5, 4, 4]: one starter

$$\begin{split} C_0 &= [19, [19, 1], 6, [5, 0], 1, [4, 0], 5, [3, 1], 8, [4, 1], 12, [2, 0], 10, [19, 0]] \\ C_1 &= [16, [3, 0], 13, [9, 0], 3, [1, 1], 2, [6, 0], 15, [1, 0]] \\ C_2 &= [17, [6, 1], 4, [7, 0], 11, [8, 0], 0, [2, 1]] \\ C_3 &= [14, [7, 1], 7, [8, 1], 18, [9, 1], 9, [5, 1]] \end{split}$$

• 2-factor type [6, 6, 4, 4]: one starter

$$\begin{split} C_0 &= [10, [19, 0], 19, [19, 1], 7, [8, 0], 18, [9, 0], 8, [1, 1], 9, [1, 0]] \\ C_1 &= [11, [8, 1], 0, [4, 1], 15, [2, 1], 13, [9, 1], 4, [7, 0], 16, [5, 0]] \\ C_2 &= [1, [2, 0], 3, [5, 1], 17, [3, 1], 14, [6, 0]] \\ C_3 &= [12, [7, 1], 5, [3, 0], 2, [4, 0], 6, [6, 1]] \end{split}$$

• 2-factor type [6, 5, 5, 4]: one starter

$$\begin{split} C_0 &= [19, [19, 0], 18, [1, 0], 0, [1, 1], 1, [4, 1], 5, [8, 1], 13, [19, 1]] \\ C_1 &= [14, [4, 0], 10, [2, 0], 12, [3, 1], 15, [9, 0], 6, [8, 0]] \\ C_2 &= [3, [5, 0], 8, [6, 1], 2, [9, 1], 11, [6, 0], 17, [5, 1]] \\ C_3 &= [9, [2, 1], 7, [3, 0], 4, [7, 1], 16, [7, 0]] \end{split}$$

- 2-factor type [16, 2, 2]: one starter
 - $$\begin{split} C_0 &= [3, [1, 1], 2, [5, 0], 16, [2, 1], 14, [9, 0], 5, [2, 0], 7, [6, 0], 13, [7, 1], 1, [8, 1], 12, [6, 1], 18, [9, 1], 9, [3, 1], 6, \\ &\quad [8, 0], 17, [7, 0], 10, [1, 0], 11, [3, 0], 8, [5, 1]] \\ C_1 &= [19, [19, 0], 4, [19, 1]] \\ C_2 &= [15, [4, 0], 0, [4, 1]] \end{split}$$
- 2-factor type [15, 3, 2]: two starters
 - $$\begin{split} C_0 &= [15, [2, 2], 17, [6, 0], 11, [2, -1], 9, [9, 1], 18, [5, 1], 4, [1, -1], 3, [5, 2], 8, [8, 1], 0, [5, -1], 5, [5, 0], 10, \\ &\quad [6, 1], 16, [3, -1], 13, [7, -1], 6, [1, 2], 7, [8, 0]] \\ C_1 &= [12, [2, 0], 14, [7, 1], 2, [9, 2]] \\ C_2 &= [19, [19, -1], 1, [19, 1]] \end{split}$$
 - $C_0' = \begin{bmatrix} 15, [3,2], 18, [8,-1], 7, [2,1], 5, [4,0], 1, [19,2], 19, [19,0], 12, [9,-1], 3, [7,2], 10, [1,1], 9, [7,0], 16, \\ [3,1], 0, [8,2], 11, [3,0], 8, [6,2], 14, [1,0] \end{bmatrix}$
 - $C'_1 = [13, [9, 0], 4, [6, -1], 17, [4, 2]]$
 - $C'_2 = [6, [4, -1], 2, [4, 1]]$
- 2-factor type [14, 4, 2]: two starters
 - $C_{0} = [7, [9, 1], 17, [6, 0], 4, [2, -1], 2, [1, 1], 3, [3, -1], 0, [3, 2], 16, [19, -1], 19, [19, 1], 5, [3, 0], 8, [7, 1], 15, [1, 2], 14, [4, 0], 10, [8, 1], 18, [8, 2]]$
 - $C_1 = [1, [8, 0], 9, [4, 2], 13, [2, 1], 11, [9, -1]]$
 - $C_2 = [6, [6, -1], 12, [6, 1]]$
 - $$\begin{split} C_0' &= [1, [8, -1], 12, [4, -1], 8, [9, 2], 18, [7, -1], 6, [5, 0], 11, [5, 2], 16, [9, 0], 7, [2, 2], 9, [4, 1], 5, [3, 1], 2, \\ & [7, 0], 14, [1, -1], 13, [6, 2], 0, [1, 0]] \\ C_0' &= [2, [5, 1], 15, [2, 0], 15, [5, -1], 10, [5, 0]] \end{split}$$
 - $C'_1 = [3, [5, 1], 17, [2, 0], 15, [5, -1], 10, [7, 2]]$
 - $C_2' = [19, [19, 0], 4, [19, 2]]$
- 2-factor type [13, 5, 2]: two starters
 - $C_0 = [13, [7, -1], 1, [2, -1], 3, [3, 0], 0, [6, 1], 6, [6, 2], 12, [1, 0], 11, [1, 2], 10, [5, 0], 15, [2, 1], 17, [7, 2], 5, [3, 1], 2, [2, 0], 4, [9, 2]]$
 - $C_1 = [19, [19, 0], 7, [8, -1], 18, [4, -1], 14, [5, -1], 9, [19, 2]]$
 - $C_2 = [8, [8, 0], 16, [8, 2]]$
 - $$\begin{split} C_0' &= [18, [7, 0], 11, [2, 2], 13, [8, 1], 5, [9, -1], 14, [4, 1], 10, [7, 1], 3, [9, 0], 12, [3, 2], 9, [9, 1], 0, [6, 0], 6, \\ &[1, 1], 7, [3, -1], 4, [5, 2]] \end{split}$$
 - $C_1' = [15, [5, 1], 1, [4, 2], 16, [1, -1], 17, [4, 0], 2, [6, -1]]$
 - $C'_2 = [19, [19, -1], 8, [19, 1]]$
- 2-factor type [12, 6, 2]: two starters
 - $$\begin{split} C_0 &= [18, [9, 1], 8, [3, -1], 5, [4, 2], 1, [9, 0], 10, [2, 2], 12, [9, -1], 2, [2, 0], 0, [9, 2], 9, [3, 0], 6, [2, -1], 4, \\ & [7, -1], 16, [2, 1]] \\ C_1 &= [19, [19, 1], 3, [5, 1], 17, [4, 1], 13, [6, 1], 7, [4, -1], 11, [19, -1]] \end{split}$$
 - $C_2 = [15, [1, 0], 14, [1, 2]]$

$$\begin{split} C_0' &= [18, [7, 1], 6, [3, 2], 3, [6, 0], 16, [6, -1], 10, [3, 1], 13, [19, 2], 19, [19, 0], 11, [8, 1], 0, [5, -1], 14, [6, 2], \\ &\quad 8, [1, 1], 7, [8, 0]] \\ C_1' &= [15, [8, -1], 4, [5, 0], 9, [8, 2], 17, [4, 0], 2, [1, -1], 1, [5, 2]] \\ C_2' &= [12, [7, 0], 5, [7, 2]] \end{split}$$

• 2-factor type [11, 7, 2]: two starters

$$\begin{split} C_0 &= [0, [1, 0], 18, [2, 2], 16, [4, 0], 1, [5, 2], 6, [3, -1], 9, [4, -1], 13, [9, -1], 3, [2, -1], 5, [7, -1], 12, [8, 1], \\ &\quad 4, [4, 1]] \\ C_1 &= [8, [19, -1], 19, [19, 0], 7, [7, 2], 14, [1, -1], 15, [5, 0], 10, [8, -1], 2, [6, 2]] \\ C_2 &= [11, [6, -1], 17, [6, 1]] \\ C_0' &= [2, [8, 2], 13, [8, 0], 5, [5, 1], 10, [1, 1], 11, [19, 2], 19, [19, 1], 14, [6, 0], 8, [1, 2], 9, [2, 1], 7, [7, 1], 0, [2, 0]] \\ C_1' &= [3, [4, 2], 18, [5, -1], 4, [7, 0], 16, [9, 1], 6, [9, 2], 15, [3, 1], 12, [9, 0]] \\ C_2' &= [1, [3, 0], 17, [3, 2]] \end{split}$$

• 2-factor type [10, 8, 2]: two starters

$$\begin{split} C_0 &= [9, [6, 1], 15, [5, 1], 1, [9, -1], 11, [7, 1], 18, [19, 1], 19, [19, -1], 2, [8, 2], 10, [3, 0], 7, [1, 2], 8, [1, 0]] \\ C_1 &= [5, [9, 2], 14, [8, 0], 3, [3, -1], 0, [7, 2], 12, [6, 0], 6, [2, -1], 4, [7, -1], 16, [8, 1]] \\ C_2 &= [17, [4, 0], 13, [4, 2]] \\ C_0' &= [12, [2, 1], 14, [3, 2], 11, [1, 1], 10, [5, -1], 15, [1, -1], 16, [8, -1], 5, [3, 1], 2, [4, -1], 6, [7, 0], 18, [6, -1]] \end{split}$$

- $C_{0} = [12, [2, 1], 14, [3, 2], 11, [1, 1], 10, [3, -1], 13, [1, -1], 10, [8, -1], 5, [3, 1], 2, [4, -1], 6, [7, 0], 18, [0, -1], C_{1}' = [17, [6, 2], 4, [9, 0], 13, [5, 2], 8, [5, 0], 3, [4, 1], 7, [2, 2], 9, [9, 1], 0, [2, 0]]$ $C_{2}' = [1, [19, 0], 19, [19, 2]]$
- 2-factor type [9,9,2]: two starters

$$\begin{split} C_0 &= [7, [4, 2], 11, [4, 0], 15, [7, 2], 8, [8, 0], 0, [2, 2], 17, [6, 0], 4, [6, 1], 10, [4, 1], 14, [7, -1]] \\ C_1 &= [16, [19, 2], 19, [19, 0], 5, [1, 2], 6, [5, 1], 1, [1, 0], 2, [3, -1], 18, [6, -1], 12, [3, 2], 9, [7, 0]] \\ C_2 &= [13, [9, 0], 3, [9, 2]] \end{split}$$

$$\begin{split} C_0' &= [18, [4, -1], 3, [5, 0], 17, [8, -1], 9, [3, 1], 12, [8, 2], 4, [7, 1], 16, [5, -1], 2, [2, 1], 0, [1, 1]] \\ C_1' &= [1, [6, 2], 7, [1, -1], 8, [2, 0], 6, [8, 1], 14, [9, -1], 5, [5, 2], 10, [3, 0], 13, [2, -1], 11, [9, 1]] \\ C_2' &= [19, [19, -1], 15, [19, 1]] \end{split}$$

- 2-factor type [14, 3, 3]: one starter
 - $$\begin{split} C_0 = [12, [6, 1], 18, [5, 0], 13, [7, 1], 1, [7, 0], 8, [5, 1], 3, [19, 0], 19, [19, 1], 5, [3, 0], 2, [2, 1], 4, [6, 0], 10, [4, 0], \\ 14, [3, 1], 17, [1, 1], 16, [4, 1]] \end{split}$$
 - $C_1 = [7, [1, 0], 6, [9, 0], 15, [8, 1]]$
 - $C_2 = [9, [2, 0], 11, [8, 0], 0, [9, 1]]$
- 2-factor type [13, 4, 3]: one starter
 - $$\begin{split} C_0 &= [13, [1, 0], 12, [6, 1], 18, [8, 0], 10, [1, 1], 11, [4, 0], 15, [7, 1], 3, [2, 1], 5, [3, 0], 2, [7, 0], 9, [19, 1], 19, \\ &\quad [19, 0], 4, [4, 1], 8, [5, 1]] \end{split}$$
 - $C_1 = [7, [9, 0], 16, [3, 1], 0, [2, 0], 17, [9, 1]]$
 - $C_2 = [1, [6, 0], 14, [8, 1], 6, [5, 0]]$

- 2-factor type [12, 5, 3]: one starter
 - $$\begin{split} C_0 &= [5, [8, 1], 13, [1, 0], 12, [4, 1], 16, [6, 1], 3, [7, 0], 15, [9, 1], 6, [4, 0], 10, [3, 1], 7, [7, 1], 14, [5, 0], 9, [2, 0], \\ &\quad 11, [6, 0]] \\ C_1 &= [19, [19, 0], 1, [1, 1], 2, [3, 0], 18, [5, 1], 4, [19, 1]] \\ C_2 &= [17, [2, 1], 0, [8, 0], 8, [9, 0]] \end{split}$$
- 2-factor type [11, 6, 3]: one starter $C_0 = [3, [6, 1], 16, [8, 1], 8, [5, 1], 13, [3, 0], 10, [4, 1], 14, [4, 0], 18, [2, 0], 1, [1, 0], 0, [5, 0], 5, [7, 1], 12, [9, 0]]$ $C_1 = [2, [9, 1], 11, [7, 0], 4, [3, 1], 7, [1, 1], 6, [19, 0], 19, [19, 1]]$ $C_2 = [9, [8, 0], 17, [2, 1], 15, [6, 0]]$
- 2-factor type [10, 7, 3]: one starter
 - $\begin{aligned} C_0 &= [8, [7, 0], 15, [9, 1], 6, [1, 1], 7, [6, 1], 13, [4, 1], 9, [3, 0], 12, [19, 1], 19, [19, 0], 2, [5, 0], 16, [8, 1]] \\ C_1 &= [18, [7, 1], 11, [8, 0], 3, [2, 0], 1, [3, 1], 4, [6, 0], 17, [2, 1], 0, [1, 0]] \\ C_2 &= [14, [4, 0], 10, [5, 1], 5, [9, 0]] \end{aligned}$
- 2-factor type [9, 8, 3]: one starter
 - $$\begin{split} C_0 &= [14, [4, 1], 10, [6, 1], 4, [1, 0], 5, [9, 0], 15, [2, 0], 17, [4, 0], 13, [7, 0], 6, [3, 0], 3, [8, 1]] \\ C_1 &= [19, [19, 0], 16, [5, 0], 11, [3, 1], 8, [7, 1], 1, [1, 1], 2, [2, 1], 0, [9, 1], 9, [19, 1]] \\ C_2 &= [7, [5, 1], 12, [6, 0], 18, [8, 0]] \end{split}$$
- 2-factor type [11, 5, 4]: one starter

$$\begin{split} C_0 &= [15, [3, 1], 12, [9, 0], 3, [1, 1], 4, [4, 0], 0, [7, 1], 7, [5, 1], 2, [4, 1], 17, [7, 0], 10, [9, 1], 1, [19, 0], 19, [19, 1]] \\ C_1 &= [13, [2, 1], 11, [5, 0], 16, [2, 0], 18, [6, 0], 5, [8, 0]] \\ C_2 &= [6, [3, 0], 9, [1, 0], 8, [6, 1], 14, [8, 1]] \end{split}$$

• 2-factor type [10, 6, 4]: one starter

$$\begin{split} C_0 &= [5, [1, 0], 4, [2, 0], 6, [8, 1], 17, [2, 1], 0, [3, 1], 3, [4, 1], 18, [8, 0], 10, [5, 0], 15, [6, 0], 9, [4, 0]] \\ C_1 &= [19, [19, 1], 14, [7, 1], 7, [5, 1], 12, [1, 1], 13, [3, 0], 16, [19, 0]] \\ C_2 &= [1, [9, 1], 11, [9, 0], 2, [6, 1], 8, [7, 0]] \end{split}$$

• 2-factor type [9,7,4]: one starter

$$\begin{split} C_0 &= [16, [5, 0], 11, [2, 0], 9, [6, 0], 15, [19, 0], 19, [19, 1], 10, [9, 0], 1, [4, 1], 5, [7, 0], 17, [1, 1]] \\ C_1 &= [4, [3, 1], 7, [1, 0], 8, [6, 1], 2, [2, 1], 0, [3, 0], 3, [9, 1], 12, [8, 1]] \\ C_2 &= [18, [5, 1], 13, [7, 1], 6, [8, 0], 14, [4, 0]] \end{split}$$

• 2-factor type [10, 5, 5]: one starter

$$\begin{split} C_0 &= [13, [3, 0], 16, [2, 0], 14, [6, 1], 8, [6, 0], 2, [4, 1], 6, [5, 0], 1, [2, 1], 18, [5, 1], 4, [1, 0], 5, [8, 0]] \\ C_1 &= [7, [4, 0], 11, [1, 1], 12, [3, 1], 9, [8, 1], 17, [9, 0]] \\ C_2 &= [19, [19, 1], 15, [7, 1], 3, [7, 0], 10, [9, 1], 0, [19, 0]] \end{split}$$

• 2-factor type [9, 6, 5]: one starter

$$\begin{split} C_0 &= [10, [7, 0], 17, [4, 0], 13, [8, 0], 5, [2, 1], 3, [4, 1], 18, [5, 0], 4, [9, 0], 14, [3, 1], 11, [1, 0]] \\ C_1 &= [15, [5, 1], 1, [6, 1], 7, [2, 0], 9, [3, 0], 6, [9, 1], 16, [1, 1]] \\ C_2 &= [12, [19, 0], 19, [19, 1], 2, [6, 0], 8, [8, 1], 0, [7, 1]] \end{split}$$

• 2-factor type [8, 7, 5]: one starter

$$\begin{split} C_0 &= [16, [5, 1], 11, [1, 0], 10, [5, 0], 15, [6, 1], 9, [1, 1], 8, [4, 1], 12, [9, 0], 3, [6, 0]] \\ C_1 &= [18, [19, 0], 19, [19, 1], 6, [8, 1], 14, [3, 0], 17, [2, 0], 0, [7, 1], 7, [8, 0]] \\ C_2 &= [2, [3, 1], 5, [4, 0], 1, [7, 0], 13, [9, 1], 4, [2, 1]] \end{split}$$

• 2-factor type [8, 6, 6]: one starter

$$\begin{split} C_0 &= [15, [2, 1], 13, [8, 0], 5, [1, 1], 4, [3, 0], 7, [6, 1], 1, [5, 0], 6, [19, 0], 19, [19, 1]] \\ C_1 &= [8, [6, 0], 14, [3, 1], 17, [1, 0], 18, [4, 0], 3, [7, 1], 10, [2, 0]] \\ C_2 &= [12, [4, 1], 16, [5, 1], 11, [8, 1], 0, [9, 1], 9, [7, 0], 2, [9, 0]] \end{split}$$

• 2-factor type [7, 7, 6]: one starter

$$\begin{split} C_0 &= [5, [5, 0], 10, [6, 0], 16, [2, 1], 14, [8, 1], 6, [6, 1], 0, [2, 0], 17, [7, 0]]\\ C_1 &= [18, [5, 1], 4, [3, 1], 1, [8, 0], 9, [3, 0], 12, [1, 0], 13, [9, 0], 3, [4, 1]]\\ C_2 &= [19, [19, 0], 7, [1, 1], 8, [7, 1], 15, [4, 0], 11, [9, 1], 2, [19, 1]] \end{split}$$

• 2-factor type [18, 2]: two starters

$$\begin{split} C_0 &= [11, [1, 0], 10, [7, 2], 3, [19, -1], 19, [19, 1], 2, [2, -1], 4, [4, 1], 0, [6, 0], 6, [6, 2], 12, [1, -1], 13, [7, 0], 1, \\ &= [8, 2], 9, [1, 1], 8, [6, -1], 14, [4, -1], 18, [8, -1], 7, [9, 0], 16, [1, 2], 17, [6, 1]] \\ C_1 &= [15, [9, -1], 5, [9, 1]] \end{split}$$

- $$\begin{split} C_0' &= [13, [5, 0], 8, [7, -1], 1, [3, 1], 4, [9, 2], 14, [5, -1], 0, [3, 0], 3, [2, 2], 5, [8, 0], 16, [2, 1], 18, [8, 1], 7, [4, 2], \\ &\quad 11, [5, 1], 6, [3, -1], 9, [7, 1], 2, [4, 0], 17, [5, 2], 12, [2, 0], 10, [3, 2]] \\ C_1' &= [19, [19, 0], 15, [19, 2]] \end{split}$$
- 2-factor type [17, 3]: one starter

$$\begin{split} C_0 &= [3, [1, 0], 4, [3, 1], 7, [8, 0], 18, [6, 0], 5, [3, 0], 8, [6, 1], 2, [8, 1], 10, [7, 0], 17, [19, 0], 19, [19, 1], 1, [4, 0], \\ &\quad 16, [1, 1], 15, [4, 1], 0, [9, 1], 9, [5, 0], 14, [2, 0], 12, [9, 0]] \\ C_1 &= [6, [5, 1], 11, [2, 1], 13, [7, 1]] \end{split}$$

- 2-factor type [15,5]: one starter
 - $$\begin{split} C_0 &= [2, [5, 1], 16, [4, 1], 12, [19, 1], 19, [19, 0], 3, [3, 1], 0, [4, 0], 4, [1, 1], 5, [5, 0], 10, [1, 0], 11, [2, 0], 13, [7, 0], \\ &\quad 6, [2, 1], 8, [9, 1], 18, [3, 0], 15, [6, 0]] \end{split}$$
 - $C_1 = [17, [8, 0], 9, [8, 1], 1, [6, 1], 14, [7, 1], 7, [9, 0]]$
- 2-factor type [14, 6]: one starter
 - $C_0 = [9, [7, 0], 2, [8, 0], 13, [2, 1], 15, [8, 1], 7, [4, 0], 3, [19, 0], 19, [19, 1], 10, [9, 0], 0, [1, 1], 1, [7, 1], 8, [3, 1], 11, [5, 1], 16, [9, 1], 6, [3, 0]]$
 - $C_1 = [14, [4, 1], 18, [6, 1], 5, [1, 0], 4, [6, 0], 17, [5, 0], 12, [2, 0]]$
- 2-factor type [13, 7]: one starter
 - $$\begin{split} C_0 &= [18, [4, 0], 3, [5, 1], 8, [1, 1], 7, [3, 1], 10, [6, 1], 4, [7, 1], 16, [19, 1], 19, [19, 0], 13, [2, 1], 15, [5, 0], 1, [8, 0], \\ &9, [9, 0], 0, [1, 0]] \end{split}$$
 - $C_1 = [14, [2, 0], 12, [7, 0], 5, [6, 0], 11, [9, 1], 2, [4, 1], 6, [8, 1], 17, [3, 0]]$

• 2-factor type [11,9]: one starter

$$\begin{split} C_0 &= [11, [7, 0], 4, [19, 0], 19, [19, 1], 5, [3, 0], 2, [4, 0], 6, [6, 1], 0, [9, 1], 9, [9, 0], 18, [8, 0], 10, [3, 1], 13, [2, 1]] \\ C_1 &= [16, [1, 1], 15, [7, 1], 8, [1, 0], 7, [4, 1], 3, [5, 0], 17, [5, 1], 12, [8, 1], 1, [6, 0], 14, [2, 0]] \end{split}$$

Acknowledgements

The second author gratefully acknowledges support by the Natural Sciences and Engineering Research Council of Canada (NSERC).

References

- P. Adams, D. Bryant, Two-factorisations of complete graphs of orders fifteen and seventeen, Australas. J. Combin. 35 (2006), 113–118.
- B. Alspach, R. Häggkvist, Some observations on the Oberwolfach problem, J. Graph Theory 9 (1985), 177–187.
- [3] B. Alspach, P. J. Schellenberg, D. R. Stinson, D. Wagner, The Oberwolfach problem and factors of uniform odd length cycles, J. Combin. Theory Ser. A 52 (1989), 20–43.
- [4] R. D. Baker, R. M. Wilson, Nearly Kirkman triple systems, Utilitas Math. 11 (1977), 289–296.
- [5] D. Bryant, P. Danziger, On bipartite 2-factorizations of $K_n I$ and the Oberwolfach problem, J. Graph Theory 68 (2011), 22–37.
- [6] A. Deza, F. Franek, W. Hua, M. Meszka, A. Rosa, Solutions to the Oberwolfach problem for orders 18 to 40, J. Combin. Math. Combin. Comput. 74 (2010), 95–102.
- [7] F. Franek, J. Holub, A. Rosa, Two-factorizations of small complete graphs II: The case of 13 vertices, J. Combin. Math. Combin. Comput. 51 (2004), 89–94.
- [8] F. Franek, A. Rosa, Two-factorizations of small complete graphs, J. Statist. Plann. Inference 86 (2000), 435–442.
- [9] S. Glock, F. Joos, J. Kim, D. Kühn, D. Osthus, Resolution of the Oberwolfach problem, J. Eur. Math. Soc. 23 (2021), 2511–2547.
- [10] D. G. Hoffman, P. J. Schellenberg, The existence of C_k -factorizations of $K_{2n} F$, Discrete Math. 97 (1991), 243–250.
- [11] D. Lepine, M. Šajna, On the Honeymoon Oberwolfach Problem, J. of Combin. Des. 27 (2019), 420–447.
- [12] F. Salassa, G. Dragotto, T. Traetta, M. Buratti, F. Della Croce, Merging combinatorial design and optimization: the Oberwolfach problem, Australas. J. Combin. 79 (2021), 141–166.
- [13] T. Traetta, A complete solution to the two-table Oberwolfach problems, J. Combin. Theory Ser. A 120 (2013), 984–997.