

A RANK-TWO CASE OF LOCAL-GLOBAL COMPATIBILITY FOR $l = p$

YUJI YANG

ABSTRACT. We prove the classical $l = p$ local-global compatibility conjecture for certain regular algebraic cuspidal automorphic representations of weight 0 for GL_2 over CM fields. Using an automorphy lifting theorem, we show that if the automorphic side comes from a twist of Steinberg at $v \mid l$, then the Galois side has nontrivial monodromy at v . Based on this observation, we will give a definition of the Fontaine–Mazur \mathcal{L} -invariants attached to certain automorphic representations.

CONTENTS

1. Introduction	1
2. Some Important Results	3
3. Local-Global Compatibility	6
4. An application	7
References	8

1. INTRODUCTION

The goal of this work is to prove a new case of local-global compatibility for automorphic Galois representations over CM fields at $l = p$. Let F be a CM field and let $G_F = \mathrm{Gal}(\overline{F}/F)$ denote the absolute Galois group. Let π be a regular algebraic cuspidal automorphic representation of $\mathrm{GL}_n(\mathbb{A}_F)$. Fix a prime l and an isomorphism $\iota : \overline{\mathbb{Q}}_l \cong \mathbb{C}$. Conjecturally there exists a continuous semisimple Galois representation $r_\iota(\pi) : G_F \rightarrow \mathrm{GL}_n(\overline{\mathbb{Q}}_l)$ that is potentially semistable (which is known to be equivalent to de Rham) above l , such that we have an isomorphism of Weil–Deligne representations

$$\mathrm{WD}(r_\iota(\pi)|_{G_{F_v}})^{\mathrm{F-ss}} \otimes_\iota \mathbb{C} \cong \mathrm{rec}_{F_v}(\pi_v \otimes |\det|^{\frac{1-n}{2}})$$

for any finite place v in F . Here on the automorphic side π_v is the local component of $\pi = \otimes'_v \pi_v$ at v , and rec_{F_v} is the local Langlands correspondence for F_v normalized as in [HT01]; on the Galois side $G_{F_v} = \mathrm{Gal}(\overline{F}_v/F_v)$ is the decomposition group at v , and WD is a functor taking a continuous semisimple Galois representation to a Weil–Deligne representation by Grothendieck’s l -adic monodromy theorem when $v \nmid l$ and by a construction of Fontaine [Fon94] when $v \mid l$ and $r_\iota(\pi)|_{G_{F_v}}$ is potentially semistable. We may then study the above isomorphism for all finite places not dividing l and for all places above l separately. The former is called the $l \neq p$ case and the latter is called the $l = p$ case.

From now on we will focus on the $l = p$ case. If π is polarizable, the $l = p$ local-global compatibility has been established by work of many people [BLGGT12, BLGGT14a, Car14], etc. For non-polarizable π , the representations $r_\iota(\pi)$ have been constructed by [HLTT16] and [Sch15] using different methods. Recently, A’Campo proved that under certain conditions the

representation $r_l(\pi)$ is de Rham with specified Hodge–Tate weights [A’C24, Theorem 4.3.3], and Hevesi proved that under the same conditions the $l = p$ local-global compatibility holds up to semisimplification [Hev23, Theorem 1.1]. This means that to prove the $l = p$ local-global compatibility, we only need to compare the monodromy operators of the two Weil–Deligne representations. It turns out when π is of rank 2 and weight 0, we can solve the problem most of the time.

Theorem 1.1. *Let F be a CM field and let π be a regular algebraic cuspidal automorphic representation of $\mathrm{GL}_2(\mathbb{A}_F)$ of weight 0. There is a set of rational primes l of Dirichlet density one such that for any isomorphism $\iota: \overline{\mathbb{Q}}_l \cong \mathbb{C}$ and any finite place $v \mid l$ in F , we have*

$$\mathrm{WD}(r_l(\pi)|_{G_{F_v}})^{\mathrm{F}\text{-ss}} \otimes_{\iota} \mathbb{C} \cong \mathrm{rec}_{F_v}(\pi_v \otimes |\det|^{-1/2}).$$

Remark 1.2. For simplicity, here we present a compromised version of our main theorem. For the full version with more technical conditions, see Theorem 3.1.

Method of proof. By [Hev23, Theorem 1.1], it suffices to show that when π_v is special (i.e. a twist of Steinberg), the Galois representation $r_l(\pi)|_{G_{F_v}}$ has nontrivial monodromy. Assume otherwise. Combined with [A’C24, Theorem 4.3.3], after a base change we may assume that $r_l(\pi)|_{G_{F_v}}$ is crystalline. Next we apply a potential automorphy theorem to find an automorphic representation π_1 that is unramified above l such that there is a congruence between mod l Galois representations $\bar{r}_l(\pi) \cong \bar{r}_l(\pi_1)$ when restricted to the absolute Galois group of some finite extension of F . Then we are able to apply an automorphy lifting theorem to find an automorphic representation π_2 that is unramified above l such that $r_l(\pi) \cong r_l(\pi_2)$ when restricted to the absolute Galois group of some field extension of F . From there we deduce a contradiction with genericity.

The idea to approach the local-global compatibility conjecture using automorphy lifting theorems is inspired by Luu [Luu15]. This method has been used in [AN20] and [Yan21] to prove cases of the $l \neq p$ local-global compatibility, where the isomorphism is known up to semisimplification by the main result of [Var24]. However, in the $l = p$ case there is extra subtlety coming from the construction of the WD functor. We need to use the fact that this construction is independent of field extensions in some sense. See the next chapter for a review with more details.

Our method is confined to rank two and weight 0. The proof of our potential automorphy theorem [AN20, Theorem 3.9] uses a moduli space of Hilbert–Blumenthal abelian varieties which realizes a fixed mod l Galois representation, as in Taylor’s original work on the subject [Tay02]. This method only works for GL_2 . On the other hand, our automorphy lifting theorem [ACC⁺23, Theorem 6.1.1] (of Fontaine–Laffaille version) does not allow a change of weight, and our potential automorphy theorem produces an automorphic representation of weight 0 as an input of the automorphy lifting theorem. Thus we are also confined to weight 0.

When dealing with the $l \neq p$ local-global compatibility, [AN20] also has the restriction to rank two and weight 0. By using the ordinary automorphy lifting theorem [ACC⁺23, Theorem 6.1.2], [Yan21] removes the weight 0 condition at the cost of imposing an ordinarity assumption on the automorphic representation at the prime l (the residue characteristic of the coefficients). However, this can not be translated to the $l = p$ local-global compatibility problem, since the ordinary automorphy lifting theorem does not guarantee that the lift π_2 is unramified above l , which is essential for our proof.

To go beyond rank two, we need to upgrade our potential automorphy theorems. When π is polarizable such theorems are known [HSBT10, BLGHT11, BLGGT14b] by using Dwork

family of motives. In the non-polarizable case, Qian proved potential automorphy theorems under general settings [Qia23, Theorem 1.1] using Dwork families. Recently in [Mat23], based on [BLGGT14b, Qia23], Matsumoto proved many versions of potential automorphy theorems, and through a careful investigation of the Jordan canonical form of the Weil–Deligne representations, he generalized the $l \neq p$ local-global compatibility results [AN20, Yan21] to GL_n . If this method can be translated to the $l = p$ case, then we believe analogous versions of our main theorems can be proved in higher rank.

Notations. Throughout F will be a CM field. If v is a place of F , F_v is the completion of F at v . Let \mathbb{A}_F be the ring of adèles of F . Fix \overline{F} an algebraic closure of F . We write $G_F = \text{Gal}(\overline{F}/F)$ for the absolute Galois group, and similarly $G_{F_v} = \text{Gal}(\overline{F}_v/F_v)$.

For a non-archimedean local field K , let I_K be the inertia subgroup and let W_K be the Weil group.

We write ϵ_l for the l -adic cyclotomic character. We normalize our Hodge–Tate weights such that ϵ_l has Hodge–Tate weight -1 . Let B_{dR} , B_{st} , B_{cris} be Fontaine’s rings of de Rham, semistable, and crystalline periods, respectively. We take K and E to be finite extensions of \mathbb{Q}_l , with E sufficiently large to contain the images of all embeddings $\tau : K \hookrightarrow \overline{\mathbb{Q}_l}$. Let K_0 be the maximal unramified extension of \mathbb{Q}_l inside K . If $\rho : G_K \rightarrow \text{GL}(V) \cong \text{GL}_d(E)$ is a de Rham representation, we write $D_{\text{dR}}(\rho) = D_{\text{dR}}(V) = (B_{\text{dR}} \otimes_{\mathbb{Q}_l} V)^{G_K}$, which is a $K \otimes_{\mathbb{Q}_l} E$ -module. We define D_{st} and D_{cris} analogously but they are both $K_0 \otimes_{\mathbb{Q}_l} E$ -modules. We write $\text{HT}_{\tau}(\rho)$ for the multiset of τ -labelled Hodge–Tate weights of ρ , which are the degrees i at which $\text{gr}^i(D_{\text{dR}}(V) \otimes_{K_0 \otimes_{\mathbb{Q}_l} E, \tau} E) \neq 0$, counted with multiplicities equal to the E -dimensions of the corresponding graded pieces.

Let π be a regular algebraic cuspidal automorphic representation of $\text{GL}_2(\mathbb{A}_F)$. We say that π has weight 0 if it has the same infinitesimal character as the trivial (algebraic) representation of $\text{Res}_{F/\mathbb{Q}} \text{GL}_2$. For any finite place v in F , π_v is the local component of $\pi = \otimes'_v \pi_v$ at v , and rec_{F_v} is the local Langlands correspondence for F_v of [HT01].

For a Weil–Deligne representation (r, N) , its Frobenius semisimplification is $(r, N)^{\text{F-ss}} = (r^{\text{ss}}, N)$, and its semisimplification is $(r, N)^{\text{ss}} = (r^{\text{ss}}, 0)$.

Acknowledgements. I would like to thank Patrick Allen for bringing my attention to this project and answering my questions in the early stage with great patience. I would like to thank Liang Xiao and Yiwen Ding for their helpful suggestions on an early draft of this paper. I would like to thank Shanxiao Huang and Yiqin He for helpful discussions on p -adic Hodge theory and the \mathcal{L} -invariants.

2. SOME IMPORTANT RESULTS

We first recall Fontaine’s construction of the WD functor following [Fon94, All16]. Let K and E be finite extensions of \mathbb{Q}_l with E sufficiently large.

For fixed finite extension L/K , we write $G_{L/K} = \text{Gal}(L/K)$ and make the following definition.

Definition 2.1. A $(\varphi, N, G_{L/K})$ -module is a finite free $L_0 \otimes_{\mathbb{Q}_l} E$ -module together with operators φ and N , and a $\text{Gal}(L/K)$ -action, satisfying the following:

- φ is E -linear and L_0 -semilinear (this means $\varphi(ax) = \sigma(a)\varphi(x)$ for any $a \in L_0$ and $x \in D$, where $\sigma \in \text{Gal}(L_0/\mathbb{Q}_l)$ is the absolute arithmetic Frobenius);
- N is $L_0 \otimes_{\mathbb{Q}_l} E$ -linear;

- $N\varphi = l\varphi N$;
- the $\text{Gal}(L/K)$ -action is E -linear and L_0 -semilinear and commutes with φ and N .

We first associate a Weil–Deligne representation (r_D, N_D) to a $(\varphi, N, G_{L/K})$ -module D . Let $N_D = N$. Extend the action of $\text{Gal}(L/K)$ to W_K by letting I_L act trivially. For $w \in W_K$, let $\alpha(w) \in \mathbb{Z}$ be such that the image of w in W_K/I_K is $\sigma^{-\alpha(w)}$, where σ is the image of the absolute arithmetic Frobenius. We then define an $L_0 \otimes_{\mathbb{Q}_l} E$ -linear action r_D of W_K on D by $r_D(w) = w\varphi^{\alpha(w)}$. Since $L_0 \otimes_{\mathbb{Q}_l} E \cong \prod_{\tau: L_0 \rightarrow E} E$, we have a decomposition $D = \prod_{\tau: L_0 \rightarrow E} D_\tau$ where $D_\tau = D \otimes_{L_0 \otimes_{\mathbb{Q}_l} E, \tau \otimes 1} E$, and an induced Weil–Deligne representation (r_τ, N_τ) of dimension $[L_0 : \mathbb{Q}_l]$ over E . By [BM02, 2.2.1], the isomorphism class of (r_τ, N_τ) is independent of τ , so we may denote any element in the isomorphism class by $\text{WD}(D)$.

Now let $\rho : G_K \rightarrow \text{GL}_d(E)$ be a continuous potentially semistable representation. Choose L/K such that $\rho|_{G_L}$ is semistable. There is a standard way to associate it with a $(\varphi, N, G_{L/K})$ -module

$$D = D_{\text{st}, L}(\rho) = (B_{\text{st}} \otimes_{\mathbb{Q}_l} V_\rho)^{G_L}.$$

Still by [BM02, §2.2.1], the isomorphism class of $\text{WD}(D_{\text{st}, L}(\rho))$ is independent of the choice of L . Hence we may set $\text{WD}(\rho) = \text{WD}(D_{\text{st}, L}(\rho))$.

Remark 2.2. Unlike in the $l \neq p$ case, in the $l = p$ case the WD functor is not faithful: it forgets the information from the filtration.

Next we explain some important results used in our proof. The first is an automorphy lifting theorem [AN20, Theorem 2.1], which is essentially a special case of [ACC⁺23, Theorem 6.1.1].

Theorem 2.3. *Let F be a CM field and let $l \geq 5$ be a prime that is unramified in F . Let $\rho : G_F \rightarrow \text{GL}_2(\overline{\mathbb{Q}_l})$ be a continuous representation satisfying the following conditions:*

- (1) ρ is unramified almost everywhere.
- (2) For each place $v \mid l$ of F , the representation $\rho|_{G_{F_v}}$ is crystalline with labelled Hodge–Tate weights all equal to $\{0, 1\}$.
- (3) $\bar{\rho}$ is decomposed generic, $\bar{\rho}|_{G_{F(\zeta_l)}}$ is absolutely irreducible with enormous image. There exists $\sigma \in G_F - G_{F(\zeta_l)}$ such that $\bar{\rho}(\sigma)$ is a scalar.
- (4) There exists a regular algebraic cuspidal automorphic representation π of $\text{GL}_2(\mathbb{A}_F)$ of weight 0 satisfying
 - (a) $\bar{\rho} \cong \bar{r}_l(\pi)$.
 - (b) For each place $v \mid l$ of F , π_v is unramified.

Then ρ is automorphic: there exists a cuspidal automorphic representation Π of $\text{GL}_2(\mathbb{A}_F)$ of weight 0 such that $\rho \cong r_l(\Pi)$. Moreover, if v is a finite place of F and either $v \mid l$ or both ρ and π are unramified at v , then Π_v is unramified.

We give the definitions of decomposed generic and enormous image in the technical condition 3. Take a continuous mod l representation $\bar{\rho} : G_F \rightarrow \text{GL}_2(\overline{\mathbb{F}_l})$.

Definition 2.4. Let ad^0 denote the space of trace zero matrices in $M_{2 \times 2}(\overline{\mathbb{F}_l})$ with the adjoint $\text{GL}_2(\overline{\mathbb{F}_l})$ -action. We say that $\bar{\rho}$ has *enormous image* if $H = \bar{\rho}(G_F)$ is absolutely irreducible and satisfies the following:

- (1) H has no nontrivial l -power order quotient.
- (2) $H^0(H, \text{ad}^0) = H^1(H, \text{ad}^0) = 0$.
- (3) For any simple $\overline{\mathbb{F}_l}[H]$ -submodule $W \subseteq \text{ad}^0$, there is a regular semisimple $h \in H$ such that $W^{h^l} \neq 0$.

- Definition 2.5.** (1) We say that a prime $p \neq l$ is *decomposed generic* for $\bar{\rho}$ if p splits completely in F , and for any place $v \mid p$ in F , $\bar{\rho}$ is unramified at v with the quotient of the two eigenvalues of $\bar{\rho}(\text{Frob}_v)$ not equal to $1, p, p^{-1}$. Here Frob_v is the geometric Frobenius element at v .
- (2) We say that $\bar{\rho}$ is *decomposed generic* if there exists $p \neq l$ that is decomposed generic for $\bar{\rho}$.

Next is a potential automorphy theorem [AN20, Theorem 3.9].

Theorem 2.6. *Suppose that F is a CM field, l is an odd prime that is unramified in F , and k/\mathbb{F}_l finite. Let \mathcal{O} be a discrete valuation ring finite over $W(k)$ with residue field k . Let $\bar{\rho} : G_F \rightarrow \text{GL}_2(k)$ be a continuous absolutely irreducible representation such that*

- $\det(\bar{\rho}) = \bar{\epsilon}_l^{-1}$;
- for each $v \mid l$, $\bar{\rho}|_{G_{F_v}}$ admits a crystalline lift $\rho_v : G_{F_v} \rightarrow \text{GL}_2(\mathcal{O})$ with all labelled Hodge–Tate weights equal to $\{0, 1\}$.

Suppose moreover that F^{avoid}/F is a finite extension. Then we can find

- a finite CM extension F_1/F that is linearly disjoint from F^{avoid} over F with l unramified in F_1 ;
- a regular algebraic cuspidal automorphic representation π for $\text{GL}_2(\mathbb{A}_{F_1})$ of weight 0, unramified at places above l ;
- an isomorphism $\iota : \overline{\mathbb{Q}}_l \xrightarrow{\sim} \mathbb{C}$

such that (composing $\bar{\rho}$ with some embedding $k \hookrightarrow \overline{\mathbb{F}}_l$)

$$\bar{\tau}_\iota(\pi) \cong \bar{\rho}|_{G_{F_1}}.$$

If $\bar{v}_0 \nmid l$ is a finite place of F^+ , then we can moreover find F_1 and π as above with π unramified above \bar{v}_0 .

Finally we state two remarkable recent results. The first is [A’C24, Theorem 4.3.3]. The original theorem is proved for π of rank n and arbitrary weight, but for simplicity we only state the special case when π is of rank 2 and weight 0.

Theorem 2.7. *Let F be a CM field, $\iota : \overline{\mathbb{Q}}_l \rightarrow \mathbb{C}$ an isomorphism and π a regular algebraic cuspidal automorphic representation of $\text{GL}_2(\mathbb{A}_F)$ of weight 0. If $\bar{\tau}_\iota(\pi)$ is absolutely irreducible and decomposed generic, then for every place $v \mid l$ of F the representation $r_\iota(\pi)|_{G_{F_v}}$ is potentially semistable with τ -labelled Hodge–Tate weights $\text{HT}_\tau = \{0, 1\}$ for each embedding $\tau : F \hookrightarrow \overline{\mathbb{Q}}_l$. If π_v and π_{v^c} are unramified, then $r_\iota(\pi)|_{G_{F_v}}$ is crystalline.*

The second is the up-to-monodromy $l = p$ local-global compatibility [Hev23, Theorem 1.1, Remark 1.2].

Theorem 2.8. *Let F be a CM field, $\iota : \overline{\mathbb{Q}}_l \rightarrow \mathbb{C}$ an isomorphism and π a regular algebraic cuspidal automorphic representation of $\text{GL}_n(\mathbb{A}_F)$. If $\bar{\tau}_\iota(\pi)$ is absolutely irreducible and decomposed generic, then for every place $v \mid l$ of F ,*

$$\text{WD}(r_\iota(\pi)|_{G_{F_v}})^{\text{ss}} \otimes_\iota \mathbb{C} \cong \text{rec}_{F_v}(\pi_v \otimes |\det|^{\frac{1-n}{2}})^{\text{ss}}.$$

We also modify the statement for simplicity. The original theorem proves that under certain partial order, the monodromy of the Galois side is “less than” the monodromy of the automorphic side ([Hev23, Remark 1.2]). In particular, if the Galois side has nontrivial monodromy, so does the automorphic side. It remains to prove the converse.

3. LOCAL-GLOBAL COMPATIBILITY

Now we are ready to prove our main theorem.

Theorem 3.1. *Let F be a CM field and let π be a regular algebraic cuspidal automorphic representation of $\mathrm{GL}_2(\mathbb{A}_F)$ of weight 0. Fix an isomorphism $\iota : \overline{\mathbb{Q}}_l \cong \mathbb{C}$. Suppose that*

- (1) $l \geq 5$ is a prime that is unramified in F ,
- (2) $\overline{r}_l(\pi)$ is decomposed generic, $\overline{r}_l(\pi)(G_{F(\zeta_l)})$ is enormous, and there is a $\sigma \in G_F - G_{F(\zeta_l)}$ such that $\overline{r}_l(\pi)(\sigma)$ is a scalar.

Then for any finite place $v \mid l$ in F , we have

$$\mathrm{WD}(r_l(\pi)|_{G_{F_v}})^{\mathrm{F}\text{-ss}} \otimes_l \mathbb{C} \cong \mathrm{rec}_{F_v}(\pi_v \otimes |\det|^{-1/2}).$$

Proof. Fix a prime $p \neq l$ for which $\overline{r}_l(\pi)$ is decomposed generic. By Theorem 2.7 we know that $r_l(\pi)|_{G_{F_v}}$ is potentially semistable, so it makes sense to consider the associated Weil–Deligne representation. By Theorem 2.8, it suffices to show that if $v \mid l$ is a finite place at which π is special (i.e., π_v is a twist of Steinberg), then $\mathrm{WD}(r_l(\pi)|_{G_{F_v}})$ has nontrivial monodromy. Let N be the monodromy operator. To show that N is nontrivial it suffices to do so after restricting to a finite extension. In particular, we may go to a solvable base change that is disjoint from $\overline{F}^{\ker(\overline{r}_l(\pi))}$ in which l is unramified and p is totally split, and assume that

- π_v is an unramified twist of Steinberg,
- $r_l(\pi)|_{G_{F_v}}$ is semistable,
- $\overline{r}_l(\pi)$ is unramified at v and v^c .

Now assume that $N = 0$, and therefore $r_l(\pi)|_{G_{F_v}}$ is crystalline. We apply Theorem 2.6 with $\rho = r_l(\pi)$ and F^{avoid} equal to the Galois closure of $\overline{F}^{\ker(\overline{r}_l(\pi))}(\zeta_l)/\mathbb{Q}$, to get a CM Galois extension F_1/F linearly disjoint from F^{avoid} over F with l unramified in F_1 , and a regular algebraic cuspidal automorphic representation π_1 for $\mathrm{GL}_2(\mathbb{A}_{F_1})$ unramified at places above l and of weight 0, such that $\overline{r}_l(\pi)|_{G_{F_1}} \cong \overline{r}_l(\pi_1)$.

We wish to apply the automorphy lifting theorem Theorem 2.3. To do so, we check the following assumptions:

- (0) The prime l is unramified in F_1 . This is guaranteed by Theorem 2.6.
- (1) $r_l(\pi)|_{G_{F_1}}$ is unramified almost everywhere. We know that $r_l(\pi)$ is unramified almost everywhere by the main result of [HLT16] and so is the restriction.
- (2) For each place $v_1 \mid l$ of F_1 , the representation $r_l(\pi)|_{G_{F_1, v_1}}$ is crystalline with labelled Hodge–Tate weights all equal to $\{0, 1\}$. Since l is unramified in both F and F_1 and hence $(F_v)_0 = F_v$, $(F_{1, v_1})_0 = F_{1, v_1}$, writing V for the \mathbb{Q}_l -vector space underlying $r_l(\pi)|_{G_{F_v}}$, we can use Galois descent to compute the dimensions

$$\begin{aligned} \dim_{\mathbb{Q}_l} V &= \dim_{F_v} (B_{\mathrm{cris}} \otimes_{\mathbb{Q}_l} V)^{G_{F_v}} \\ &= \dim_{F_v} ((B_{\mathrm{cris}} \otimes_{\mathbb{Q}_l} V)^{G_{F_1, v_1}})^{\mathrm{Gal}(F_1, v_1/F_v)} \\ &= \dim_{F_1, v_1} (B_{\mathrm{cris}} \otimes_{\mathbb{Q}_l} V)^{G_{F_1, v_1}}. \end{aligned}$$

Thus the restriction $r_l(\pi)|_{G_{F_1, v_1}}$ must also be crystalline.

- (3) $\overline{r}_l(\pi)|_{G_{F_1}}$ is decomposed generic (by a similar argument using Chebotarev density to choose a prime at which $\overline{r}_l(\pi)|_{G_{F_1}}$ is decomposed generic, as in the proof of [AN20, Theorem 4.1]). $\overline{r}_l(\pi)|_{G_{F_1(\zeta_l)}}$ is absolutely irreducible (encoded in the definition of enormous image) with enormous image (by our choice of F^{avoid}). There exists $\sigma \in G_{F_1} - G_{F_1(\zeta_l)}$ such that $\overline{r}_l(\pi)(\sigma)$ is a scalar (still by our choice of F^{avoid}).

- (4) There exists a regular algebraic cuspidal automorphic representation π_1 of $\mathrm{GL}_2(\mathbb{A}_{F_1})$ of weight 0 such that $\bar{r}_l(\pi_1) \cong \bar{r}_l(\pi)|_{G_{F_1}}$. For each place $v_1 \mid l$ of F_1 , π_{1,v_1} is unramified. This is by Theorem 2.6.

The automorphy lifting theorem gives a cuspidal automorphic representation Π of $\mathrm{GL}_2(\mathbb{A}_{F_1})$ of weight 0 such that $r_l(\Pi) \cong r_l(\pi)|_{G_{F_1}}$ and Π_{v_1} is unramified at all $v_1 \mid l$ in F_1 . Now

$$\mathrm{rec}_{F_1, v_1}(\Pi_{v_1} \otimes |\det|^{-1/2}) \cong \mathrm{WD}(r_l(\Pi)|_{G_{F_1, v_1}})^{\mathrm{ss}} \otimes_l \mathbb{C} \cong \mathrm{WD}(r_l(\pi)|_{G_{F_1, v_1}})^{\mathrm{ss}} \otimes_l \mathbb{C}.$$

Under the local Langlands correspondence for F_{1, v_1} , the last object corresponds to a subquotient of an induction of two characters whose quotient equal to $|\cdot|^{\pm 1}$, which can not be generic. This contradicts the fact that Π_{v_1} is generic. \square

Proof of Theorem 1.1. It follows directly from Theorem 3.1 and [AN20, Lemma 2.9]. \square

4. AN APPLICATION

In [Maz94, §9], Mazur defined a collection of invariants attached to the isomorphism classes of certain filtered (φ, N) -modules called the two-dimensional monodromy modules. We first recall the definitions. Let D be a filtered (φ, N) -module, which is a K_0 -vector space with actions of φ and N and an exhaustive and separated decreasing filtration $\{\mathrm{Fil}^i\}_i$ on $D_K := D \otimes_{K_0} K$, satisfying the following:

- (1) φ is K_0 -semilinear;
- (2) N is K_0 -linear;
- (3) $N\varphi = l\varphi N$.

Note that D has a basis of φ -eigenvectors, and N takes φ -eigenvectors to φ -eigenvectors or 0.

Definition 4.1. A *(two-dimensional) monodromy module* D is a two-dimensional filtered (φ, N) -module with the following properties:

- (1) N is nonzero;
- (2) There is $j_0 \in \mathbb{Z}$ for which $\mathrm{Fil}^{j_0} D_K$ is 1-dimensional;
- (3) $N(D) \otimes_{K_0} K \neq \mathrm{Fil}^{j_0} D_K$.

Definition 4.2. Let D be a two-dimensional monodromy module. The *Fontaine–Mazur \mathcal{L} -invariant* $\mathcal{L}(D)$ is the unique element of K such that $x - \mathcal{L}(D) \cdot Nx$ spans $\mathrm{Fil}^{j_0} D_K$, where x is a φ -eigenvector with $Nx \neq 0$.

Let f be a classical normalized eigenform of $\Gamma_0(N)$ of even weight $k \geq 2$ (and Nebentypus character χ , for some integer N), and $\rho_f : G_{\mathbb{Q}} \rightarrow \mathrm{GL}_2(E)$ its associated l -adic Galois representation for a suitable E/\mathbb{Q}_l . If $\rho_{f, l} := \rho_f|_{G_{\mathbb{Q}_l}}$ is semistable non-crystalline, it is automatically non-critical, which means that $D_{\mathrm{st}}(\rho_{f, l})$ is a monodromy module, and therefore we may make sense of the Fontaine–Mazur \mathcal{L} -invariant $\mathcal{L}(D_{\mathrm{st}}(\rho_{f, l}))$ attached to f . Note that this \mathcal{L} -invariant only depends on $\rho_{f, l}$, so in particular it only depends on f and l .

Now we return to the settings of this paper and slightly generalize the above idea. Let F be a CM field and let l be a rational prime that is inert in F . Fix an isomorphism $\iota : \overline{\mathbb{Q}_l} \cong \mathbb{C}$. Let π be a regular algebraic cuspidal automorphic representation of $\mathrm{GL}_2(\mathbb{A}_F)$ of weight 0. Assume that π_v is special for the finite place v in F lying above l . The proof of Theorem 3.1 shows that $r_l(\pi)|_{G_{F_v}}$ is (potentially) semistable non-crystalline, and is non-critical since $G_{F_v} \cong G_{\mathbb{Q}_l}$. Hence we can make sense of the Fontaine–Mazur \mathcal{L} -invariant $\mathcal{L}(D_{\mathrm{st}}(r_l(\pi)|_{G_{F_v}}))$ attached to π . Similar to the classical modular form case, the \mathcal{L} -invariant only depends on $r_l(\pi)|_{G_{F_v}}$, so in particular it only depends on π and l . More generally, if we only assume that l is unramified in

F , then we need to assume that $r_\iota(\pi)|_{G_{F_v}}$ is non-critical to make $D_{\text{st}}(r_\iota(\pi)|_{G_{F_v}})$ a monodromy module.

Once we have the definition, it is a natural question to ask if it coincides with other versions of \mathcal{L} -invariants other than the one due to Fontaine–Mazur. In the classical modular form case the question is extensively studied and the answer is yes (for instance see the introduction of [BDI10]), but in general the question is widely open. We hope to see the question being answered with the emergence of generalizations of more versions of \mathcal{L} -invariants.

REFERENCES

- [A'C24] Lambert A'Campo, *Rigidity of automorphic Galois representations over CM fields*, Int. Math. Res. Not. IMRN (2024), no. 6, 4541–4623. MR 4721650
- [ACC⁺23] Patrick B. Allen, Frank Calegari, Ana Caraiani, Toby Gee, David Helm, Bao V. Le Hung, James Newton, Peter Scholze, Richard Taylor, and Jack A. Thorne, *Potential automorphy over CM fields*, Ann. of Math. (2) **197** (2023), no. 3, 897–1113. MR 4564261
- [All16] Patrick B. Allen, *Deformations of polarized automorphic Galois representations and adjoint Selmer groups*, Duke Math. J. **165** (2016), no. 13, 2407–2460. MR 3546966
- [AN20] Patrick B. Allen and James Newton, *Monodromy for some rank two Galois representations over CM fields*, Doc. Math. **25** (2020), 2487–2506. MR 4213130
- [BDI10] Massimo Bertolini, Henri Darmon, and Adrian Iovita, *Families of automorphic forms on definite quaternion algebras and Teitelbaum's conjecture*, Astérisque (2010), no. 331, 29–64. MR 2667886
- [BLGGT12] Thomas Barnet-Lamb, Toby Gee, David Geraghty, and Richard Taylor, *Local-global compatibility for $l = p$, I*, Ann. Fac. Sci. Toulouse Math. (6) **21** (2012), no. 1, 57–92. MR 2954105
- [BLGGT14a] ———, *Local-global compatibility for $l = p$, II*, Ann. Sci. Éc. Norm. Supér. (4) **47** (2014), no. 1, 165–179. MR 3205603
- [BLGGT14b] ———, *Potential automorphy and change of weight*, Ann. of Math. (2) **179** (2014), no. 2, 501–609. MR 3152941
- [BLGHT11] Tom Barnet-Lamb, David Geraghty, Michael Harris, and Richard Taylor, *A family of Calabi-Yau varieties and potential automorphy II*, Publ. Res. Inst. Math. Sci. **47** (2011), no. 1, 29–98. MR 2827723
- [BM02] Christophe Breuil and Ariane Mézard, *Multiplicités modulaires et représentations de $\text{GL}_2(\mathbf{Z}_p)$ et de $\text{Gal}(\overline{\mathbf{Q}}_p/\mathbf{Q}_p)$ en $l = p$* , Duke Math. J. **115** (2002), no. 2, 205–310, With an appendix by Guy Henniart. MR 1944572
- [Car14] Ana Caraiani, *Monodromy and local-global compatibility for $l = p$* , Algebra Number Theory **8** (2014), no. 7, 1597–1646. MR 3272276
- [Fon94] Jean-Marc Fontaine, *Représentations l -adiques potentiellement semi-stables*, Astérisque (1994), no. 223, 321–347, Périodes p -adiques (Bures-sur-Yvette, 1988). MR 1293977
- [Hev23] Bence Hevesi, *Ordinary parts and local-global compatibility at $l = p$* , arXiv e-prints (2023), arXiv:2311.13514.
- [HLTT16] Michael Harris, Kai-Wen Lan, Richard Taylor, and Jack Thorne, *On the rigid cohomology of certain Shimura varieties*, Res. Math. Sci. **3** (2016), Paper No. 37, 308. MR 3565594
- [HSBT10] Michael Harris, Nick Shepherd-Barron, and Richard Taylor, *A family of Calabi-Yau varieties and potential automorphy*, Ann. of Math. (2) **171** (2010), no. 2, 779–813. MR 2630056
- [HT01] Michael Harris and Richard Taylor, *The geometry and cohomology of some simple Shimura varieties*, Annals of Mathematics Studies, vol. 151, Princeton University Press, Princeton, NJ, 2001, With an appendix by Vladimir G. Berkovich. MR 1876802
- [Luu15] Martin Luu, *Deformation theory and local-global compatibility of Langlands correspondences*, Mem. Amer. Math. Soc. **238** (2015), no. 1123, vii+101. MR 3402383
- [Mat23] Kojiro Matsumoto, *On the potential automorphy and the local-global compatibility for the monodromy operators at $p \neq l$ over CM fields*, arXiv e-prints (2023), arXiv:2312.01551.
- [Maz94] B. Mazur, *On monodromy invariants occurring in global arithmetic, and Fontaine's theory, p -adic monodromy and the Birch and Swinnerton-Dyer conjecture* (Boston, MA, 1991), Contemp. Math., vol. 165, Amer. Math. Soc., Providence, RI, 1994, pp. 1–20. MR 1279599
- [Qia23] Lie Qian, *Potential automorphy for GL_n* , Invent. Math. **231** (2023), no. 3, 1239–1275. MR 4549090

- [Sch15] Peter Scholze, *On torsion in the cohomology of locally symmetric varieties*, Ann. of Math. (2) **182** (2015), no. 3, 945–1066. MR 3418533
- [Tay02] Richard Taylor, *Remarks on a conjecture of Fontaine and Mazur*, J. Inst. Math. Jussieu **1** (2002), no. 1, 125–143. MR 1954941
- [Var24] Ila Varma, *Local-global compatibility for regular algebraic cuspidal automorphic representations when $l \neq p$* , Forum Math. Sigma **12** (2024), Paper No. e21, 32. MR 4706792
- [Yan21] Yuji Yang, *An Ordinary Rank-Two Case of Local-Global Compatibility for Automorphic Representations of Arbitrary Weight Over CM Fields*, arXiv e-prints (2021), arXiv:2111.00318.

BEIJING INTERNATIONAL CENTER FOR MATHEMATICAL RESEARCH, PEKING UNIVERSITY
Email address: yujiy@pku.edu.cn