

LOG-HÖLDER REGULARITY OF STATIONARY MEASURES

GRIGORII MONAKOV

ABSTRACT. We consider Lipschitz and Hölder continuous random dynamical systems defined by a distribution with a finite logarithmic moment. We prove that under suitable non-degeneracy conditions every stationary measure must be log-Hölder continuous.

1. INTRODUCTION

Hölder continuity is an important concept that appears naturally in various fields, which include but are not limited to: stochastic processes (see, for example, [MP]), dynamical systems (see [BP, BrK, BV, BBS] and references therein), and spectral theory (e.g. see [CS, DG, GS, HV, Mu, V]).

Recently Hölder continuity was established for stationary measures of random dynamical systems under very general assumptions:

Theorem 1.1 (Gorodetski, Kleptsyn, M., [GKM]). *Let M be a closed Riemannian manifold. Suppose that μ is a probability distribution on $\text{Diff}^1(M)$ such that $\int \|f\|_{\text{Diff}^1}^\gamma d\mu(f) < \infty$ for some $\gamma > 0$. Suppose also that there is no probability measure m on the manifold M invariant under every map $f \in \text{supp}(\mu)$. Then every stationary measure of a random dynamical system defined by the distribution μ is Hölder continuous.*

This theorem gives rise to a series of natural questions, that are known to be important in that area. Namely, one can ask

- What do we get if we impose a weaker condition on the tails of distribution μ ? For instance, if we only assume that $\int (\log \|f\|_{\text{Diff}^1})^\gamma d\mu(f) < \infty$.
- Is it possible to weaken the regularity assumption on the dynamical system? For example, what if we only assume that homeomorphisms $f \in \text{supp}(\mu)$ are Hölder continuous and not differentiable?

The aim of present paper is to address these questions. Our main results are the following two theorems (for rigorous formulations and further discussion see Section 2):

Theorem 1.2. *Let μ be a probability measure on $\text{Lip}(M)$ – space of bi-Lipschitz homeomorphisms of M , satisfying the following assumptions:*

- Lipschitz constant has finite α -logarithmic moment with respect to μ .
- There is no measure m on M , such that $f_*m = m$ for μ -a.e. f .

Then every μ -stationary measure is $\alpha/2$ -log-Hölder.

Theorem 1.3. *Let μ be a probability measure on $\text{Hol}(M)$ – space of bi-Hölder homeomorphisms of M , satisfying the following assumptions:*

Date: July 2, 2024.

- Hölder constant has a finite logarithmic moment with respect to μ .
- There is no measure m on M , such that $f_*m = m$ for μ -a.e. f .

Then every μ -stationary measure is log-Hölder.

Stationary measures of random dynamical systems are analogues of invariant measures of deterministic maps, and their properties are of crucial importance for many results in random dynamics, see [A, BH, BQ, BL, Fu1, Fu2, Kif1, Kif2, LQ, M] and references therein.

Various results regarding regularity of stationary measures are known. A well studied special case is action of random projective maps on a projective space.

Let A_1, A_2, \dots be i.i.d. random matrices from $SL(k, \mathbb{R})$, distributed with respect to measure μ . Every $A \in SL(k, \mathbb{R})$ induces a projective map $f_A : \mathbb{RP}^{k-1} \rightarrow \mathbb{RP}^{k-1}$. Denote by S_μ the closed semigroup in $SL(k, \mathbb{R})$ generated by matrices from $\text{supp}(\mu)$. Under certain irreducibility assumptions (which we won't define rigorously since they are beyond the scope of our paper) on S_μ Guivarc'h was able to prove the following regularity result for the stationary measure:

Theorem 1.4 (Guivarc'h, [Gu]). *Suppose that, in the setting above, S_μ is strongly irreducible and proximal, and*

$$\mathbb{E}_\mu \|A_1\|^\gamma = \int_{SL(k, \mathbb{R})} \|A\|^\gamma d\mu(A) < \infty$$

for some $\gamma > 0$. Then the corresponding random dynamical system on \mathbb{RP}^{k-1} has unique stationary measure ν and ν is Hölder continuous.

A regularity estimate similar to the one in Theorem 1.4 is one of the key ingredients in the proof of the following Central Limit Theorem for matrix products:

Theorem 1.5 (Le Page, [L], Guivarc'h, Raugi, [GR], Gol'dsheid, Margulis, [GM]). *Let $\{A_k, k \geq 1\}$ be independent and identically distributed random matrices in $SL(k, \mathbb{R})$, distributed with respect to measure μ . Assume that S_μ is proximal and strongly irreducible and*

$$(1) \quad \mathbb{E}_\mu \|A_1\|^\gamma = \int_{SL(k, \mathbb{R})} \|A\|^\gamma d\mu(A) < \infty$$

for some $\gamma > 0$. Then there exists a $a > 0$ such that the random variables

$$\frac{\log \|A_n \dots A_1\| - n\lambda_F}{\sqrt{n}},$$

where $\lambda_F > 0$ is the Lyapunov exponent, converge in distribution to $\mathcal{N}(0, a^2)$.

It is known that assumption (1), which is usually referred to as *finite exponential moment*, is not optimal for the Central Limit Theorem. Recently Benuist and Quint were able to improve it in the following way:

Theorem 1.6 (Benuist, Quint, [BQ16]). *Let $\{A_k, k \geq 1\}$ be independent and identically distributed random matrices in $SL(k, \mathbb{R})$, distributed with respect to measure μ . Assume that S_μ is proximal and strongly irreducible and*

$$(2) \quad \mathbb{E}_\mu [(\log \|A_1\|)^2] = \int_{SL(k, \mathbb{R})} (\log \|A\|)^2 d\mu(A) < \infty.$$

Then there exists $a > 0$ such that the random variables

$$\frac{\log \|A_n \dots A_1\| - n\lambda_F}{\sqrt{n}},$$

where $\lambda_F > 0$ is the Lyapunov exponent, converge in distribution to $\mathcal{N}(0, a^2)$.

Condition (2), which is usually referred to as *finite second moment* is known to be optimal. In order to establish this more general result (as well as other limit theorems in the same setting) the authors proved a form of log-Hölder regularity of the stationary measure (see [BQ16, Proposition 4.5] for details) under *finite second moment* assumption. This result illustrates the importance of log-Hölder continuity for proving limit theorems with optimal assumptions.

Log-Hölder regular measures appears naturally in other problems as well. For example, in [CS] Craig and Simon proved that integrated density of states for discrete Schrödinger operators with ergodic potentials is log-Hölder. Curiously, the assumption they found sufficient was a finite logarithmic moment of the function that generates the potential, which is very similar to assumptions (2) and (5). Regularity of integrated density of states of discrete Schrödinger operators was extensively studied by many authors, see [B, BK, CK, DG, GK, GS, HV, MS, Mu, ST, V] and references therein.

2. MAIN RESULTS

Let M be a smooth closed Riemannian manifold of dimension k and μ be a Borel probability measure on $\text{Homeo}(M)$, the set of continuous homeomorphisms of M . Consider the corresponding random dynamical system, given by the compositions

$$T_n := f_n \circ \dots \circ f_1,$$

where $f_i \in \text{Homeo}(M)$ are chosen randomly and independently, with respect to the distribution μ .

If an initial point $x \in M$ is distributed with respect to a probability measure ν , one can consider the distribution $\mu * \nu$ of its random image $f(x)$. In other words, $\mu * \nu$ is the μ -averaged push-forward image of the measure ν :

$$\mu * \nu := \int_{\text{Homeo}(M)} (f_* \nu) \, d\mu(f).$$

The measure ν is called μ -stationary if $\mu * \nu = \nu$. Let \mathcal{M} denote the space of all Borel probability measures on M equipped with the weak-* topology.

Definition 2.1. A measure $\nu \in \mathcal{M}$ is (C, α) -Hölder if

$$\forall x \in X \quad \forall r > 0 \quad \nu(B_r(x)) \leq Cr^\alpha.$$

A measure $\nu \in \mathcal{M}$ is (C, α) -log-Hölder if

$$\forall x \in X \quad \forall r > 0 \quad \nu(B_r(x)) \leq C |\log(r)|^{-\alpha}.$$

Let $\text{Hol}(M)$ denote the space of all bi-Hölder continuous homeomorphisms of M and $\text{Lip}(M)$ denote the space of all bi-Lipschitz continuous homeomorphisms of M . We define Hölder and Lipschitz constants by the following formulae:

Definition 2.2. For $f \in \text{Hol}(M)$ define

$$\gamma(f) = \min \left(\sup \left\{ \gamma > 0 : \sup_{x,y \in M} \frac{d(f(x), f(y))}{d(x,y)^\gamma} < \infty \right\}, \right. \\ \left. \sup \left\{ \gamma > 0 : \sup_{x,y \in M} \frac{d(x,y)}{d(f(x), f(y))^\gamma} < \infty \right\} \right)$$

and

$$L(f) = \max \left(\sup_{x,y \in M} \frac{d(f(x), f(y))}{d(x,y)^{\frac{\gamma(f)}{2}}}, \sup_{x,y \in M} \frac{d(x,y)}{d(f(x), f(y))^{\frac{\gamma(f)}{2}}} \right).$$

For $f \in \text{Lip}(M)$ define

$$\mathfrak{L}(f) = \max \left(\sup_{x,y \in M} \frac{d(f(x), f(y))}{d(x,y)}, \sup_{x,y \in M} \frac{d(x,y)}{d(f(x), f(y))} \right).$$

Now we are ready to formulate our main results.

Theorem 2.3. *Let μ be a probability measure on $\text{Hol}(M)$, satisfying the following assumptions:*

- *There exists $\beta > 0$ such that*

$$(3) \quad \int_{\text{Hol}(M)} \left(\frac{1 + \log(L(f))}{\gamma(f)} \right)^\beta d\mu(f) < \infty.$$

- **(no invariant measure)** *There is no measure $m \in \mathcal{M}$, such that $f_*m = m$ for μ -a.e. f .*

Then there exist $\alpha > 0$ and C such that every μ -stationary probability measure ν on M is (α, C) -log-Hölder.

Moreover, we prove the following regularity estimate for a nonstationary random dynamical systems after finitely many iterations:

Theorem 2.4. *Let K be a compact set (with respect to weak-* topology) in the space of Borel probability measures on $\text{Homeo}(M)$, satisfying the following assumptions:*

- *For every $\mu \in K$ we have $\text{supp}(\mu) \in \text{Hol}(M)$.*
- *There exists $\beta > 0$ and C_0 such that for every $\mu \in K$*

$$\int_{\text{Hol}(M)} \left(\frac{1 + \log(L(f))}{\gamma(f)} \right)^\beta d\mu(f) < C_0.$$

- **(no deterministic image)** *For every $\mu \in K$ there are no measures $m_1, m_2 \in \mathcal{M}$, such that $f_*m_1 = m_2$ for μ -a.e. f .*

Then there exist $\alpha > 0$, C and $\kappa < 1$ such that for every initial measure ν_0 , every sequence of measures $\mu_1, \mu_2, \dots \in K$, every number of iterations $n \in \mathbb{N}$, and every $x \in M$ one has:

$$\text{if } r > \kappa^n \text{ then } [\mu_n * \mu_{n-1} * \dots * \mu_1 * \nu_0](B_r(x)) < C |\log(r)|^{-\alpha}.$$

We would like to mention the following connection between the **no invariant measure** and the **no deterministic image** assumptions (for proof see [GKM]):

Proposition 2.5. *Let μ be a probability measure on $\text{Homeo}(M)$ that satisfies **no invariant measure** condition (there is no common invariant measure for all $f \in \text{supp } \mu$). Then there exists $k \in \mathbb{N}$ such that μ^{*k} satisfies **no deterministic images** condition (i.e. there are no probability measures ν, ν' on M such that $f_*\nu = \nu'$ with $f = f_k \circ f_{k-1} \circ \dots \circ f_1$ for $\mu \times \mu \times \dots \times \mu$ -almost all $(f_1, f_2, \dots, f_k) \in (\text{Homeo}(M))^k$).*

Applying Theorem 2.4 to a one-point set $K = \{\mu\}$ and taking into account Proposition 2.5 we obtain the following

Corollary 2.6. *Assume that μ satisfies the assumptions of Theorem 2.3. Then there exist $\alpha > 0$, C and $\kappa < 1$ such that for every initial measure ν_0 , every number of iterations $n \in \mathbb{N}$, and every $x \in M$ one has:*

$$(4) \quad \text{if } r > \kappa^n \text{ then } [\mu^{*n} * \nu_0](B_r(x)) < C |\log(r)|^{-\alpha}.$$

Remark 2.7. It is also easy to see that Theorem 2.3 follows from Corollary 2.6, hence all we need to prove is Theorem 2.4.

Using similar techniques we are also able to provide more refined estimates for the case of bi-Lipschitz homeomorphisms.

Theorem 2.8. *Let μ be a probability measure on $\text{Lip}(M)$, satisfying the following assumptions:*

- *There exists $\alpha > 0$ such that*

$$(5) \quad \int_{\text{Lip}(M)} (\log(\mathfrak{L}(f)))^\alpha d\mu(f) < \infty.$$

- **(no invariant measure)** *There is no measure $m \in \mathcal{M}$, such that $f_*m = m$ for μ -a.e. f .*

Then there exists C , such that every stationary measure $\nu \in \mathcal{M}$ is $(C, \alpha/2)$ -log-Hölder.

Similar to the case of Hölder continuous homeomorphisms, a nonstationary version of Theorem 2.8 is available.

Theorem 2.9. *Let K be a compact set (with respect to weak- $*$ topology) in the space of Borel probability measures on $\text{Homeo}(M)$, satisfying the following assumptions:*

- *For every $\mu \in K$ we have $\text{supp}(\mu) \subset \text{Lip}(M)$.*
- *There exists $\alpha > 0$ and C_0 such that for every $\mu \in K$*

$$\int_{\text{Lip}(M)} (\log(\mathfrak{L}(f)))^\alpha d\mu(f) < C_0.$$

- **(no deterministic image)** *For every $\mu \in K$ there are no measures $m_1, m_2 \in \mathcal{M}$, such that $f_*m_1 = m_2$ for μ -a.e. f .*

Then there exist C and $\kappa < 1$ such that for every initial measure ν_0 , every sequence of measures $\mu_1, \mu_2, \dots \in K$, every number of iterations $n \in \mathbb{N}$, and every $x \in M$ one has:

$$\text{if } r > \kappa^n \text{ then } [\mu_n * \mu_{n-1} * \dots * \mu_1 * \nu_0](B_r(x)) < C |\log(r)|^{-\frac{\alpha}{2}}.$$

The following corollary also holds:

Corollary 2.10. *Assume that μ satisfies the assumptions of Theorem 2.8. Then there exist C and $\kappa < 1$ such that for every initial measure ν_0 , every number of iterations $n \in \mathbb{N}$, and every $x \in M$ one has:*

$$\text{if } r > \kappa^n \text{ then } [\mu^{*n} * \nu_0](B_r(x)) < C |\log(r)|^{-\frac{\alpha}{2}}.$$

3. HÖLDER CONTINUOUS HOMEOMORPHISMS

In order to describe regularity of a measure $\nu \in \mathcal{M}$ we will study the following integral:

$$\iint_{M^2} |\log(d(x, z))|^\alpha d\nu(x) d\nu(z)$$

for $\alpha > 0$. To apply the machinery introduced in [GKM] we need to find a function φ_α such that its convolution with itself has a singularity at the origin equivalent to $|\log(r)|^\alpha$.

Definition 3.1. Define

$$\varphi_\alpha(x) = \begin{cases} \frac{(-\log(x))^{\frac{\alpha-1}{2}}}{x^{\frac{k}{2}}}, & \text{if } 0 < x < \frac{1}{e}, \\ 0, & \text{otherwise,} \end{cases}$$

and for $r > 0$

$$\begin{aligned} U_\alpha(r) &= \int_{\mathbb{R}^k} \varphi_\alpha(|\bar{x}|) \varphi_\alpha(|\bar{r} - \bar{x}|) d\bar{x} = \\ &= \int_{\{|\bar{x}| < \frac{1}{e}\} \cap \{|\bar{r} - \bar{x}| < \frac{1}{e}\}} \varphi_\alpha(|\bar{x}|) \varphi_\alpha(|\bar{r} - \bar{x}|) d\bar{x}, \end{aligned}$$

where $\bar{x} \in \mathbb{R}^k$ and $\bar{r} = (r, 0, \dots, 0) \in \mathbb{R}^k$.

The function $U_\alpha(r)$ has the same singularity at the origin as $|\log(r)|^\alpha$ (we will formalize that statement in Proposition 3.4), but that is still not good enough for us, because the integral

$$\iint_{M^2} U_\alpha(d(x, z)) d\nu(x) d\nu(z)$$

can be infinite. To fix that problem we define the following family of cut-offs:

Definition 3.2. Define

$$\varphi_{\alpha, \varepsilon}(x) = \begin{cases} \varphi_\alpha(x), & \text{if } x \geq \varepsilon; \\ \varphi_\alpha(\varepsilon), & \text{if } 0 < x < \varepsilon \end{cases}$$

and

$$\begin{aligned} U_{\alpha, \varepsilon}(r) &= \int_{\mathbb{R}^k} \varphi_{\alpha, \varepsilon}(|\bar{x}|) \varphi_{\alpha, \varepsilon}(|\bar{r} - \bar{x}|) d\bar{x} = \\ &= \int_{\{|\bar{x}| < \frac{1}{e}\} \cap \{|\bar{r} - \bar{x}| < \frac{1}{e}\}} \varphi_{\alpha, \varepsilon}(|\bar{x}|) \varphi_{\alpha, \varepsilon}(|\bar{r} - \bar{x}|) d\bar{x}, \end{aligned}$$

where $\bar{x} \in \mathbb{R}^k$ and $\bar{r} = (r, 0, \dots, 0) \in \mathbb{R}^k$.

An important symmetry that the function $U_{\alpha, \varepsilon}$ possesses is described in the following

Remark 3.3. Note that the definition of $U_{\alpha, \varepsilon}$ can be reformulated as follows: for any two points $x, z \in \mathbb{R}^k$ consider the integral

$$(6) \quad I_{\alpha, \varepsilon}(x, z) := \int_{\mathbb{R}^k} \varphi_{\alpha, \varepsilon}(|y - x|) \varphi_{\alpha, \varepsilon}(|y - z|) d\text{Leb}(y).$$

Due to the spherical symmetry, this integral depends only on the distance $r = |x - z|$ between these two points:

$$I_{\alpha,\varepsilon}(x, z) = U_{\alpha,\varepsilon}(|x - z|)$$

for some function $U_{\alpha,\varepsilon}$ of $r = |x - z|$. We can take this as a definition of $U_{\alpha,\varepsilon}(r)$.

Let us denote by $c_{\alpha,k}$ the following constant:

$$c_{\alpha,k} = \frac{k\omega_k}{\alpha},$$

where ω_k is the volume of a k -dimensional unit ball. We will collect some useful properties of the function $U_{\alpha,\varepsilon}$ in the following

Proposition 3.4. *For every dimension $k \geq 1$ the following holds:*

(I) *For every $\alpha > 0$ and $\varepsilon > 0$ the function $U_{\alpha,\varepsilon}$ is non-increasing.*

(II) *For every $\alpha_0 > 0$ there exist $r_0 > 0$ and $\varepsilon_0 > 0$, such that for every $0 < \alpha < \alpha_0$, every $0 < \varepsilon < \varepsilon_0$, and every $0 < r < r_0$ we have:*

$$(7) \quad U_{\alpha,\varepsilon}(r) < 2c_{\alpha,k}(-\log(r))^\alpha.$$

(III) *For every $\alpha_0 > 0$ there exist $r_0 > 0$ and $\varepsilon_0 > 0$, such that for every $0 < \alpha < \alpha_0$, every $0 < \varepsilon < \varepsilon_0$, and every $\varepsilon < r < r_0$ we have:*

$$(8) \quad U_{\alpha,\varepsilon}(r) > \frac{c_{\alpha,k}}{2}(-\log(r))^\alpha.$$

(IV) *For every $\alpha_0 > 0$ and every $\delta > 0$ there exist $\varepsilon_0 > 0$ and $r_0 > 0$ such that for every $0 < \alpha < \alpha_0$, every $0 < \varepsilon < \varepsilon_0$, and every $0 < r_1 \leq r_2 \leq r_0$ we have*

$$(9) \quad U_{\alpha,\varepsilon}(r_1) \leq (1 + \delta) \left(\frac{\log(r_1)}{\log(r_2)} \right)^\alpha U_{\alpha,\varepsilon}(r_2).$$

We will postpone the proof of Proposition 3.4 until Appendix A. We are ready to define the energy that will be convenient to work with and will allow us to capture the log-Hölder regularity of a measure $\nu \in \mathcal{M}$.

Definition 3.5.

$$\mathcal{E}_{\alpha,\varepsilon}(\nu) = \iint_{M^2} U_{\alpha,\varepsilon}(d(x, z)) d\nu(x) d\nu(z).$$

Let us point out a few useful properties of $\mathcal{E}_{\alpha,\varepsilon}$:

Lemma 3.6. *For every $\alpha > 0$, every $\nu \in \mathcal{M}$ and every $0 < \varepsilon < 1$ we have the following upper bound:*

$$\mathcal{E}_{\alpha,\varepsilon}(\nu) \leq \frac{\omega_k(-\log(\varepsilon))^{\alpha-1}}{e^k \varepsilon^k}.$$

Proof. A straightforward computation shows that

$$\mathcal{E}_{\alpha,\varepsilon}(\nu) \leq U_{\alpha,\varepsilon}(0) \leq \frac{\omega_k}{e^k} \varphi_\alpha^2(\varepsilon) = \frac{\omega_k(-\log(\varepsilon))^{\alpha-1}}{e^k \varepsilon^k}.$$

□

Lemma 3.7. *For every $\alpha > 0$ there exists $C_{\alpha,k} < \infty$ and $\varepsilon_0 > 0$, such that for every $0 < \varepsilon < \varepsilon_0$, every $\nu \in \mathcal{M}$, every $0 < \varepsilon < r < \frac{1}{e}$ and every ball $B_r(x) \subset M$ the following holds:*

$$(10) \quad \nu(B_r(x)) \leq C_{\alpha,k} \sqrt{\mathcal{E}_{\alpha,\varepsilon}(\nu)} (-\log(r))^{-\frac{\alpha}{2}}.$$

Proof. Let us choose ε_0 such that Part **(III)** of Proposition 3.4 is applicable. Then for small enough r applying Markov inequality to the definition of $\mathcal{E}_{\alpha,\varepsilon}$ we obtain:

$$\mathcal{E}_{\alpha,\varepsilon}(\nu) \geq U_{\alpha,\varepsilon}(2r) \cdot \nu(B_r(x))^2 \geq \frac{C_{\alpha,k}}{2} (-\log(2r))^\alpha \cdot \nu(B_r(x))^2.$$

Choosing $C_{\alpha,k}$ big enough we can ensure that (10) holds for all r such that $\varepsilon < r < 1/e$. \square

The following series of statements describes the change of the energy $\mathcal{E}_{\alpha,\varepsilon}(\nu)$ after applying a Hölder continuous homeomorphism.

Proposition 3.8. *For every $\delta > 0$ there exists $\varepsilon_0 > 0$, such that for every $0 < \alpha < 1$ there exist $A = A(\alpha, \delta)$, such that for every $0 < \varepsilon < \varepsilon_0$, every $f \in \text{Hol}(M)$ and every $\nu \in \mathcal{M}$ the following holds:*

$$(11) \quad \mathcal{E}_{\alpha,\varepsilon}(f_*\nu) \leq \left(\frac{2(1 + \log(L(f)))}{\gamma(f)} \right)^\alpha ((1 + \delta)\mathcal{E}_{\alpha,\varepsilon}(\nu) + A).$$

Proof. According to the Part **(IV)** of Proposition 3.4 we can choose $\varepsilon_0 > 0$ and $r_0 > 0$, such that for every $0 < \alpha < 1$, every $0 < \varepsilon < \varepsilon_0$ and every $0 < r_1 \leq r_2 \leq r_0$ we have

$$U_{\alpha,\varepsilon}(r_1) \leq (1 + \delta) \left(\frac{\log(r_1)}{\log(r_2)} \right)^\alpha U_{\alpha,\varepsilon}(r_2).$$

Let us split the integral from the definition of energy the $\mathcal{E}_{\alpha,\varepsilon}(f_*\nu)$ in the following way:

$$\begin{aligned} \mathcal{E}_{\alpha,\varepsilon}(f_*\nu) &= \iint_{M \times M} U_{\alpha,\varepsilon}(d(x, z)) \, df_*\nu(x) \, df_*\nu(z) = \\ &= \iint_{M \times M} U_{\alpha,\varepsilon}(d(f(x), f(z))) \, d\nu(x) \, d\nu(z) = \\ &= \iint_{d(x,z) < r_0} U_{\alpha,\varepsilon}(d(f(x), f(z))) \, d\nu(x) \, d\nu(z) + \\ &\quad + \iint_{d(x,z) \geq r_0} U_{\alpha,\varepsilon}(d(f(x), f(z))) \, d\nu(x) \, d\nu(z). \end{aligned}$$

To estimate the first summand we take $r_2 = d(x, z)$ and $r_1 = \min(d(f(x), f(z)), r_2)$ and we deduce that

$$\begin{aligned} U_{\alpha,\varepsilon}(d(f(x), f(z))) &\leq U_{\alpha,\varepsilon}(r_1) \leq (1 + \delta) \left(\frac{\log(r_1)}{\log(r_2)} \right)^\alpha U_{\alpha,\varepsilon}(r_2) \leq \\ &\leq (1 + \delta) \left(\frac{\log(d(f(x), f(z)))}{\log(d(x, z))} \right)^\alpha U_{\alpha,\varepsilon}(d(x, z)) \end{aligned}$$

for any $x, z \in M$, such that $d(x, z) \leq r_0$. By definition of $\gamma(f)$ and $L(f)$ we know that

$$d(f(x), f(z)) \geq \left(\frac{d(x, z)}{L(f)} \right)^{\frac{2}{\gamma(f)}}$$

for any $x, z \in M$. After substituting that into previous inequality we arrive to

$$\begin{aligned} U_{\alpha,\varepsilon}(d(f(x), f(z))) &\leq \\ &\leq (1 + \delta) \left(\frac{2(\log(d(x, z)) - \log(L(f)))}{\gamma(f) \log(d(x, z))} \right)^\alpha U_{\alpha,\varepsilon}(d(x, z)) \leq \\ &\leq (1 + \delta) \left(\frac{2(1 + \log(L(f)))}{\gamma(f)} \right)^\alpha U_{\alpha,\varepsilon}(d(x, z)). \end{aligned}$$

For the first summand the last inequality gives us the following:

$$\begin{aligned} &\iint_{d(x,z) < r_0} U_{\alpha,\varepsilon}(d(f(x), f(z))) d\nu(x) d\nu(z) \leq \\ &\leq (1 + \delta) \iint_{M \times M} \left(\frac{2(1 + \log(L(f)))}{\gamma(f)} \right)^\alpha U_{\alpha,\varepsilon}(d(x, z)) d\nu(x) d\nu(z) \leq \\ &\leq (1 + \delta) \left(\frac{2(1 + \log(L(f)))}{\gamma(f)} \right)^\alpha \mathcal{E}_{\alpha,\varepsilon}(\nu). \end{aligned}$$

Let us take r_0 and ε_0 small enough to ensure that the conclusion of Part **(II)** of Proposition 3.4 also holds. That allows us to estimate the second summand as follows:

$$\begin{aligned} &\iint_{d(x,z) \geq r_0} U_{\alpha,\varepsilon}(d(f(x), f(z))) d\nu(x) d\nu(z) \leq \\ &\leq \iint_{M \times M} U_{\alpha,\varepsilon} \left(\left(\frac{r_0}{L(f)} \right)^{\frac{2}{\gamma(f)}} \right) d\nu(x) d\nu(z) \leq \\ &\leq 2c_{\alpha,k} \left(-\log \left(\left(\frac{r_0}{L(f)} \right)^{\frac{2}{\gamma(f)}} \right) \right)^\alpha \leq \\ &\leq 2c_{\alpha,k} (-\log(r_0))^\alpha \left(\frac{2(1 + \log(L(f)))}{\gamma(f)} \right)^\alpha. \end{aligned}$$

Taking $A = 2c_{\alpha,k} (-\log(r_0))^\alpha$ finishes the proof. \square

An immediate corollary can be formulated as follows:

Corollary 3.9. *For every $\delta > 0$ there exists $\varepsilon_0 > 0$, such that for every $0 < \alpha < 1$ there exists $C < \infty$, such that for every $0 < \varepsilon < \varepsilon_0$, every $f \in \text{Hol}(M)$ and every $\nu \in \mathcal{M}$, such that $\mathcal{E}_{\alpha,\varepsilon}(\nu) > C$ the following holds:*

$$\begin{aligned} (12) \quad &\left(\frac{2(1 + \log(L(f)))}{\gamma(f)} \right)^{-\alpha} \frac{1}{1 + \delta} \leq \\ &\leq \frac{\mathcal{E}_{\alpha,\varepsilon}(f_*\nu)}{\mathcal{E}_{\alpha,\varepsilon}(\nu)} \leq \\ &\leq \left(\frac{2(1 + \log(L(f)))}{\gamma(f)} \right)^\alpha (1 + \delta). \end{aligned}$$

Another corollary of this statement that would be convenient to use is

Corollary 3.10. *There exists $\varepsilon_0 > 0$, such that for every $0 < \alpha < 1$ there exists $C < \infty$, such that for every $0 < \varepsilon < \varepsilon_0$, every $f \in \text{Hol}(M)$ and every $\nu \in \mathcal{M}$ the following holds:*

$$(13) \quad \mathcal{E}_{\alpha,\varepsilon}(f_*\nu) \leq \max \left(2 \left(\frac{2(1 + \log(L(f)))}{\gamma(f)} \right)^\alpha \mathcal{E}_{\alpha,\varepsilon}(\nu), C \right).$$

Proof. Applying Corollary 3.9 to $\delta = \frac{1}{2}$, $\tilde{\nu} := f_*\nu$ and $\tilde{f} = f^{-1}$ we conclude that there exists $C < \infty$, such that if $\mathcal{E}_{\alpha,\varepsilon}(\tilde{\nu}) = \mathcal{E}_{\alpha,\varepsilon}(f_*\nu) > C$ then

$$\mathcal{E}_{\alpha,\varepsilon}(f_*^{-1}f_*\nu) \geq \frac{1}{1 + \frac{1}{2}} \left(\frac{2(1 + \log(L(f^{-1})))}{\gamma(f^{-1})} \right)^{-\alpha} \mathcal{E}_{\alpha,\varepsilon}(f_*\nu),$$

and the inequality (13) follows. \square

In what follows it will be convenient to use the following notation:

Definition 3.11. We say that two positive numbers, A and A' , are (δ, C) -close, if

$$A < (1 + \delta)A' + C \quad \text{and} \quad A' < (1 + \delta)A + C;$$

this can be equivalently rewritten as

$$\frac{1}{1 + \delta}A - \frac{1}{1 + \delta}C < A' < (1 + \delta)A + C.$$

We denote it $A \approx_{(\delta,C)} A'$.

Starting from this point the proof follows the ideas of the one in [GKM], but since some technical details are different we repeat it completely. We start by

Definition 3.12. For $\nu \in \mathcal{M}$ define

$$(14) \quad \rho_{\alpha,\varepsilon}[\nu](y) = \int_M \varphi_{\alpha,\varepsilon}(d(x, y)) d\nu(x);$$

and

$$\Theta_{\alpha,\varepsilon}[\nu] = \rho_{\alpha,\varepsilon}^2[\nu](y) d\text{Leb}(y).$$

The motivation behind formula (14) is that $\rho_{\alpha,\varepsilon}[\nu]$ can be viewed as a vector in $L^2(M, \text{Leb})$ (which is crucial for the proof of Proposition 3.27) that captures local regularity of the measure ν . We will also need

Definition 3.13.

$$(15) \quad \begin{aligned} \tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu) &= \Theta_{\alpha,\varepsilon}[\nu](M); \\ \theta_{\alpha,\varepsilon}[\nu] &= \frac{\Theta_{\alpha,\varepsilon}[\nu]}{\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu)}. \end{aligned}$$

The energy $\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu)$ plays an important role in our proof, since it mimics the behaviour of $\mathcal{E}_{\alpha,\varepsilon}(\nu)$ and at the same time can be represented as a square of the norm of an $L^2(M, \text{Leb})$ vector $\rho_{\alpha,\varepsilon}$.

Lemma 3.14.

$$\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu) = \iint_{M \times M} K_{\alpha,\varepsilon}(x, z) d\nu(x) d\nu(z),$$

where

$$(16) \quad K_{\alpha,\varepsilon}(x, z) := \int_M \varphi_{\alpha,\varepsilon}(d(x, y)) \varphi_{\alpha,\varepsilon}(d(z, y)) d\text{Leb}(y).$$

Proof. It suffices to substitute (14) that defines $\rho_{\alpha,\varepsilon}^2[\nu](y)$, into (15), obtaining a triple integral

$$\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu) = \iiint_{M \times M \times M} \varphi_{\alpha,\varepsilon}(d(x,y)) \varphi_{\alpha,\varepsilon}(d(z,y)) \, d\nu(x) \, d\nu(z) \, d\text{Leb}(y),$$

and then change the order of integration. \square

The following proposition claims that $\mathcal{E}_{\alpha,\varepsilon}$ and $\tilde{\mathcal{E}}_{\alpha,\varepsilon}$ are comparable for measures with high energy.

Proposition 3.15. *For every $\alpha > 0$ and every $\delta > 0$ there exists $C > 0$ such that for every $\varepsilon > 0$ if $\mathcal{E}_{\alpha,\varepsilon}(\nu) > C$ or $\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu) > C$ then one has*

$$\frac{\mathcal{E}_{\alpha,\varepsilon}(\nu)}{\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu)} \in (1 - \delta, 1 + \delta).$$

Remark 3.16. We can reformulate that statement using notations introduced above: for every $\alpha > 0$ and every $\delta > 0$ there exists $C < \infty$, such that for every $\varepsilon > 0$ and every $\nu \in \mathcal{M}$ one has

$$\mathcal{E}_{\alpha,\varepsilon}(\nu) \approx_{(\delta,C)} \tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu).$$

Joining Proposition 3.15 and Corollary 3.9 we get

Corollary 3.17. *For every $\delta > 0$ there exists $\varepsilon_0 > 0$, such that for every $R < \infty$ there exists $\alpha_0 > 0$, such that for every $0 < \alpha < \alpha_0$ there exists $C > 0$ such that for every $f \in \text{Hol}(M)$ with $L(f) < R$ and $\gamma(f)^{-1} < R$, every $0 < \varepsilon < \varepsilon_0$ and every ν , such that $\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu) > C$ or $\mathcal{E}_{\alpha,\varepsilon}(\nu) > C$ one has*

$$(17) \quad \frac{\tilde{\mathcal{E}}_{\alpha,\varepsilon}(f_*\nu)}{\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu)} \in (1 - \delta, 1 + \delta).$$

To prove Proposition 3.15 we will need the following:

Lemma 3.18. *For every $\alpha > 0$, the interaction potentials $K_{\alpha,\varepsilon}(x, z)$ and $U_{\alpha,\varepsilon}(d(x, z))$ are comparable in the following sense:*

$$(18) \quad \forall \delta > 0 \exists C : \quad \forall \varepsilon > 0, \forall x, z \in M$$

$$K_{\alpha,\varepsilon}(x, z) \approx_{(\delta,C)} U_{\alpha,\varepsilon}(d(x, z)), \text{ i.e.}$$

$$\frac{1}{1 + \delta} (U_{\alpha,\varepsilon}(d(x, z)) - C) < K_{\alpha,\varepsilon}(x, z) < (1 + \delta) U_{\alpha,\varepsilon}(d(x, z)) + C.$$

Let us postpone the proof of Lemma 3.18 for a moment and deduce Proposition 3.15.

Proof of Proposition 3.15. It suffices to integrate (18) w.r.t. $\nu \times \nu(x, z)$. As constant C does not depend on these points nor on the measure ν , one gets

$$\frac{1}{1 + \delta} (\mathcal{E}_{\alpha,\varepsilon}(\nu) - C) < \tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu) < (1 + \delta) \mathcal{E}_{\alpha,\varepsilon}(\nu) + C$$

with the same constant C . As it was noticed in Remark 3.16, this is an equivalent form of Proposition 3.15. \square

Proof of Lemma 3.18. Let $\alpha > 0$ be given. Take any $r_0 > 0$ and divide the integral (16) defining $K_{\alpha,\varepsilon}(x, z)$ into two parts, depending on whether the distance $d(x, y)$ exceeds r_0 :

$$K_{\alpha,\varepsilon}(x, z) = \int_{B_{r_0}(x)} \varphi_{\alpha,\varepsilon}(d(x, y))\varphi_{\alpha,\varepsilon}(d(z, y)) \, d\text{Leb}(y) + \\ + \int_{M \setminus B_{r_0}(x)} \varphi_{\alpha,\varepsilon}(d(x, y))\varphi_{\alpha,\varepsilon}(d(z, y)) \, d\text{Leb}(y).$$

Denote the first and the second summands as $K_{\alpha,\varepsilon}^{(r_0)}(x, z)$ and $\overline{K}_{\alpha,\varepsilon}^{(r_0)}(x, z)$. Note that the second summand is uniformly bounded: indeed, the factor $\varphi_{\alpha,\varepsilon}(d(x, y))$ doesn't exceed a constant $\varphi_{\alpha,\varepsilon}(r_0) \leq \varphi_\alpha(r_0)$, while the second factor $\varphi_{\alpha,\varepsilon}(d(y, z))$ is a function with the integral on M that is bounded uniformly in $z \in M$. The latter uniform bound can be seen by again decomposing the integral in two:

$$(19) \quad \int_M \varphi_{\alpha,\varepsilon}(d(y, z)) \, d\text{Leb}(y) = \int_{B_{r_0}(z)} \varphi_{\alpha,\varepsilon}(d(y, z)) \, d\text{Leb}(y) + \\ + \int_{M \setminus B_{r_0}(z)} \varphi_{\alpha,\varepsilon}(d(y, z)) \, d\text{Leb}(y).$$

The second summand in (19) does not exceed $\text{vol}(M) \cdot \varphi_\alpha(r_0)$. The first one can be estimated uniformly in $z \in M$ by passage to the geodesic coordinates centred at z , comparing it to the same integral in a ball in \mathbb{R}^k : the Jacobian of the change of variables is (for r_0 smaller than the injectivity radius in M) uniformly bounded, and the integral of $\frac{(-\log(r))^{\frac{\alpha-1}{2}}}{r^{\frac{k}{2}}}$ on $B_{r_0}(0) \subset \mathbb{R}^k$ converges.

We thus have

$$(20) \quad \overline{K}_{\alpha,\varepsilon}^{(r_0)}(x, z) \leq C_\alpha^K(r_0)$$

for some constant $C_\alpha^K(r_0)$.

Now, let us transform the first integral, $K_{\alpha,\varepsilon}^{(r_0)}(x, z)$. For sufficiently small $r_0 > 0$ (smaller than the injectivity radius at every point) one can take the geodesic coordinates at any point $x \in M$ in a ball of radius r_0 . Thus, we can take a point $x' \in \mathbb{R}^k$, let $\psi : B_{r_0}(x') \rightarrow B_{r_0}(x)$ be geodesic coordinates, and denote $z' := \psi^{-1}(z)$.

Fix any $\delta_1 > 0$. Due to the compactness of M , for a sufficiently small r_0 the Jacobian and the bi-Lipschitz constants of ψ are δ_1 -close to 1:

$$\mathfrak{L}(\psi) < 1 + \delta_1, \quad \text{Jac}(\psi) \in \left(\frac{1}{1 + \delta_1}, 1 + \delta_1 \right).$$

Making a change of variables $y = \psi(y')$ in the integral for $K_{\alpha,\varepsilon}^{(r_0)}(x, z)$, we get

$$K_{\alpha,\varepsilon}^{(r_0)}(x, z) = \int_{B_{r_0}(x)} \varphi_{\alpha,\varepsilon}(d(x, y))\varphi_{\alpha,\varepsilon}(d(z, y)) \, d\text{Leb}_M(y) = \\ = \int_{B_{r_0}(x')} \varphi_{\alpha,\varepsilon}(d(\psi(x'), \psi(y'))) \varphi_{\alpha,\varepsilon}(d(\psi(z'), \psi(y'))) \, d\text{Leb}_M(\psi(y'));$$

all the three quotients

$$\frac{\varphi_{\alpha,\varepsilon}(d(\psi(x'), \psi(y')))}{\varphi_{\alpha,\varepsilon}(|x' - y'|)}, \quad \frac{\varphi_{\alpha,\varepsilon}(d(\psi(z'), \psi(y')))}{\varphi_{\alpha,\varepsilon}(|z' - y'|)}, \quad \frac{d\text{Leb}_M}{d\psi_*\text{Leb}_{\mathbb{R}^k}} \Big|_y$$

are close to 1: the first two by a factor $(1 + \delta_1)^{\frac{k+\alpha}{2}}$ since

$$1 \leq \frac{\varphi_{\alpha,\varepsilon}(r_1)}{\varphi_{\alpha,\varepsilon}(r_2)} \leq \left(\frac{\log(r_1)}{\log(r_2)}\right)^{\frac{\alpha-1}{2}} \left(\frac{r_2}{r_1}\right)^{\frac{k}{2}} < \left(\frac{r_2}{r_1}\right)^{\frac{k+\alpha}{2}} \quad \text{for any } 0 < r_1 \leq r_2 < \frac{1}{e},$$

and the last one by the factor $(1 + \delta_1)$. Thus, $K_{\alpha,\varepsilon}^{(r_0)}(x, z)$ differs from

$$(21) \quad I_{\alpha,\varepsilon}^{(r_0)}(x', z') := \int_{B_{r_0}(x')} \varphi_{\alpha,\varepsilon}(|x' - y'|) \varphi_{\alpha,\varepsilon}(|z' - y'|) \, d\text{Leb}_{\mathbb{R}^k}(y')$$

by the factor at most $(1 + \delta_1)^{k+\alpha+1}$. Now, $I_{\alpha,\varepsilon}^{(r_0)}(x', z')$ is also a part of $I_{\alpha,\varepsilon}(x', z')$ (define by (6)), that differs from it by

$$\bar{I}_{\alpha,\varepsilon}^{(r_0)}(x', z') := \int_{\mathbb{R}^k \setminus B_{r_0}(x')} \varphi_{\alpha,\varepsilon}(|x' - y'|) \varphi_{\alpha,\varepsilon}(|z' - y'|) \, d\text{Leb}_{\mathbb{R}^k}(y'),$$

that is bounded uniformly in ε, x', z' (for $|x' - z'| < \frac{r_0}{2}$ both functions outside of $B_{r_0}(x')$ are bounded and have compact supports, for $|x' - z'| \geq \frac{r_0}{2}$ we have $\bar{I}_{\alpha,\varepsilon}^{(r_0)}(x', z') \leq U_{\alpha,\varepsilon}(\frac{r_0}{2})$). Thus, first choosing δ_1 so that $(1 + \delta_1)^{k+\alpha+1} < 1 + \delta$ and then accordingly choosing r_0 , we obtain the desired estimate (18). \square

In what follows it will be convenient to use Wasserstein metric in the space of probability measures on M . Let us recall its definition, as well as definition of the total variation distance between measures:

Definition 3.19. Let ν_1, ν_2 be two probability measures on a measure space (M, \mathcal{B}) . Then the *Wasserstein distance* between them is defined as

$$W(\nu_1, \nu_2) = \inf_{\gamma} \iint_{M \times M} d(x, y) \, d\gamma(x, y),$$

where the infimum is taken over all probability measures γ on $(M \times M, \mathcal{B} \times \mathcal{B})$ with the marginals (projections on the x and y coordinates) $P_x(\gamma) = \nu_1$ and $P_y(\gamma) = \nu_2$.

Definition 3.20. Let ν_1, ν_2 be two probability measures on a measure space (M, \mathcal{B}) . Then the *total variation distance* between them is

$$\text{TV}(\nu_1, \nu_2) = \sup_{B \in \mathcal{B}} |\nu_1(B) - \nu_2(B)|.$$

Also, we will need the following statement that can be found, for example, in [Vi, Theorem 6.15]: the Wasserstein metric is bounded from above by the diameter times the total variation distance.

Lemma 3.21. *For every two probability measures ν_1, ν_2 on a manifold M we have*

$$W(\nu_1, \nu_2) \leq \text{diam}(M) \cdot \text{TV}(\nu_1, \nu_2).$$

Equipped with these notions we would like to prove the following

Proposition 3.22. *For every $\delta > 0$ and every $R < \infty$ there exists $\alpha_0 > 0$, such that for every $0 < \alpha < \alpha_0$ there exists $C > 0$, such that for every $\varepsilon > 0$, every $f \in \text{Hol}(M)$, such that $\gamma(f)^{-1}, L(f) < R$ and every $\nu \in \mathcal{M}$ such that $\mathcal{E}_{\alpha,\varepsilon}(\nu) > C$ or $\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu) > C$ one has*

$$(22) \quad W(f_* \theta_{\alpha,\varepsilon}[\nu], \theta_{\alpha,\varepsilon}[f_* \nu]) < \delta.$$

Proof. For given $\alpha, \varepsilon > 0$ and measure ν , consider a (non-probability) measure $m_{\alpha, \varepsilon}(\nu)$ on $M \times M \times M$, given by

$$m_{\alpha, \varepsilon}(\nu) = \varphi_{\alpha, \varepsilon}(d(x, y))\varphi_{\alpha, \varepsilon}(d(z, y)) d\nu(x) d\text{Leb}(y) d\nu(z).$$

Denote by π_1, π_2 the projections of $M \times M \times M$ on the first and second coordinates respectively, and by $\pi_{1,3}$ the projection on $M \times M$ corresponding to the first and third coordinates. Then directly from definition

$$(\pi_2)_* m_{\alpha, \varepsilon}(\nu) = \Theta_{\alpha, \varepsilon}(\nu);$$

also, define

$$(23) \quad \widehat{\Theta}_{\alpha, \varepsilon}(\nu) := (\pi_{1,3})_* m_{\alpha, \varepsilon}(\nu) = K_{\alpha, \varepsilon}(x, z) \nu(dx) \nu(dz);$$

the second equality is due to (16). Finally, consider the measure

$$(24) \quad \Theta'_{\alpha, \varepsilon}(\nu) := (\pi_1)_* m_{\alpha, \varepsilon}(\nu)$$

as well as the normalizations of these measures,

$$(25) \quad \widehat{\theta}_{\alpha, \varepsilon}(\nu) := \frac{1}{\widetilde{\mathcal{E}}_{\alpha, \varepsilon}(\nu)} \widehat{\Theta}_{\alpha, \varepsilon}(\nu), \quad \theta'_{\alpha, \varepsilon}(\nu) := \frac{1}{\widetilde{\mathcal{E}}_{\alpha, \varepsilon}(\nu)} \Theta'_{\alpha, \varepsilon}(\nu).$$

For a high-energy measure ν most of the measure $m_{\alpha, \varepsilon}(\nu)$ is concentrated near the diagonal, and hence the projections on the first and on the second coordinates are close to each other. The following lemma formalizes this argument:

Lemma 3.23. *For every δ_1, α there exists C' such that for every $\varepsilon > 0$ and ν with $\widetilde{\mathcal{E}}_{\alpha, \varepsilon}(\nu) > C'$ one has*

$$W(\theta_{\alpha, \varepsilon}(\nu), \theta'_{\alpha, \varepsilon}(\nu)) < \delta_1.$$

Proof. Take $r_0 := \frac{\delta_1}{2}$ and let

$$A_{r_0} := \{(x, y, z) \in M^3 \mid d(x, y) < r_0\}, \quad \overline{A}_{r_0} := M^3 \setminus A_{r_0}.$$

From the proof of Lemma 3.18, we have

$$m_{\alpha, \varepsilon}(\nu)(\overline{A}_{r_0}) = \int \overline{K}_{\alpha, \varepsilon}^{(r_0)}(x, z) d\nu(x) d\nu(z) \leq C_{\alpha}^K(r_0),$$

where the second inequality is due to (20). Thus, the non-normalized measure $m_{\alpha, \varepsilon}(\nu)(\overline{A}_{r_0})$ does not exceed a constant $C_{\alpha}^K(r_0)$.

Now, we can couple the normalized measures $\theta_{\alpha, \varepsilon}(\nu)$ and $\theta'_{\alpha, \varepsilon}(\nu)$ using the projection of $\frac{1}{\widetilde{\mathcal{E}}_{\alpha, \varepsilon}(\nu)} m_{\alpha, \varepsilon}(\nu)$ on the first two coordinates; this coupling leads to the upper bound for the Wasserstein distance:

$$(26) \quad W(\theta_{\alpha, \varepsilon}(\nu), \theta'_{\alpha, \varepsilon}(\nu)) \leq r_0 \cdot \frac{m_{\alpha, \varepsilon}(\nu)(A_{r_0})}{\widetilde{\mathcal{E}}_{\alpha, \varepsilon}(\nu)} + \text{diam}(M) \cdot \frac{m_{\alpha, \varepsilon}(\nu)(\overline{A}_{r_0})}{\widetilde{\mathcal{E}}_{\alpha, \varepsilon}(\nu)} \leq r_0 + \frac{C_{\alpha}^K(r_0)}{\widetilde{\mathcal{E}}_{\alpha, \varepsilon}(\nu)},$$

where the first summand corresponds to the points with $d(x, y) < r_0$, and the second to the points with $d(x, y) \geq r_0$. As we chose $r_0 = \frac{\delta_1}{2}$, it suffices to require that

$$\widetilde{\mathcal{E}}_{\alpha, \varepsilon}(\nu) > \frac{2C_{\alpha}^K(r_0)}{\delta_1} =: C'$$

to ensure that the total Wasserstein distance does not exceed δ_1 . \square

On the other hand, for the measures $\theta'_{\alpha,\varepsilon}(\nu)$ the analogue of Proposition 3.22 can be established directly, and one can even estimate the total variations distance:

Lemma 3.24. *For every $\delta_2 > 0$ and every $R > 0$ there exists $\alpha_2 > 0$ such that for every $0 < \alpha < \alpha_2$ there exists $C > 0$ such that for every $\varepsilon > 0$, every $f \in \text{Hol}(M)$ with $\gamma(f)^{-1}, L(f) < R$ and every ν such that $\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu) > C$ or $\mathcal{E}_{\alpha,\varepsilon}(\nu) > C$ one has*

$$(27) \quad \text{TV}(f_*\theta'_{\alpha,\varepsilon}(\nu), \theta'_{\alpha,\varepsilon}(f_*\nu)) < \delta_2,$$

and, hence,

$$(28) \quad W(f_*\theta'_{\alpha,\varepsilon}(\nu), \theta'_{\alpha,\varepsilon}(f_*\nu)) < \delta_2 \cdot \text{diam}(M).$$

Proof. We will first use Lemma 3.18 together with Part **(IV)** of Proposition 3.4 to compare $K_{\alpha,\varepsilon}(x, z)$ with $K_{\alpha,\varepsilon}(f(x), f(z))$. Fix α_0 and δ_3 so small that

$$(1 + \delta_3)^2 \cdot [2R(1 + \log(R))]^{\alpha_0} < 1 + \frac{\delta_2}{10}.$$

Then from Lemma 3.18 we know that there exists $C > 0$ such that for any x, z, ε we have

$$(29) \quad K_{\alpha,\varepsilon}(x, z) \approx_{(\delta_3, C)} U_{\alpha,\varepsilon}(x, z), \quad K_{\alpha,\varepsilon}(f(x), f(z)) \approx_{(\delta_3, C)} U_{\alpha,\varepsilon}(f(x), f(z)).$$

Without loss of generality we can assume that $d(x, z) < d(f(x), f(z))$. Then by Part **(IV)** of Proposition 3.4 we can choose a particular $r_0 = r_0(\alpha_0, \delta_3)$, such that if $d(f(x), f(z)) < r_0$ then

$$(30) \quad \frac{U_{\alpha,\varepsilon}(d(x, z))}{U_{\alpha,\varepsilon}(d(f(x), f(z)))} < (1 + \delta_3) \left(\frac{\log(d(x, z))}{\log(d(f(x), f(z)))} \right)^\alpha.$$

Notice that inequality (30) only holds for $\varepsilon < \varepsilon_0(\alpha_0, \delta_3)$. That can be guaranteed by taking C big enough, because, according to Lemma 3.6, the energy $\mathcal{E}_{\varepsilon,\alpha}(\nu)$ for $\varepsilon \geq \varepsilon_0$ has an explicit upper bound.

Using the fact that $L(f), \gamma(f)^{-1} < R$ we conclude that

$$\begin{aligned} \left(\frac{\log(d(x, z))}{\log(d(f(x), f(z)))} \right)^\alpha &\leq \\ &\leq \left(\frac{\frac{2}{\gamma(f)} \log [d(f(x), f(z))L(f)^{-1}]}{\log(d(f(x), f(z)))} \right)^\alpha < \\ &< [2R(1 + \log(R))]^{\alpha_0} \end{aligned}$$

and

$$(31) \quad U_{\alpha,\varepsilon}(d(x, z)) < (1 + \delta_3) [2R(1 + \log(R))]^{\alpha_0} U_{\alpha,\varepsilon}(d(f(x), f(z))).$$

On the other hand, if $d(f(x), f(z)) > r_0$ then

$$d(x, z) \geq \left(\frac{r_0}{L(f)} \right)^{\frac{2}{\gamma(f)}} > \left(\frac{r_0}{R} \right)^{\frac{2}{R}}$$

and hence

$$(32) \quad U_{\alpha,\varepsilon}(d(x, z)) < U_{\alpha_0} \left(\left(\frac{r_0}{R} \right)^{\frac{2}{R}} \right).$$

Combining estimates (29), (31) and (32) we conclude that

$$(33) \quad K_{\alpha,\varepsilon}(x, z) \approx_{(\frac{\delta_2}{10}, C'')} K_{\alpha,\varepsilon}(f(x), f(z))$$

for some explicit constant C'' .

Now, applying f_*^{-1} does not change the total variations distance, so instead of (27) we can show the equivalent statement

$$(34) \quad \text{TV}(\theta'_{\alpha,\varepsilon}(\nu), f_*^{-1}\theta'_{\alpha,\varepsilon}(f_*\nu)) < \delta_2.$$

The measures here can be obtained as projections of measures on $M \times M$:

$$\begin{aligned} \hat{\theta}_{\alpha,\varepsilon}(\nu) &= (\pi_1)_* \hat{\theta}_{\alpha,\varepsilon}(\nu), \\ f_*^{-1}\hat{\theta}_{\alpha,\varepsilon}(f_*\nu) &= (\pi_1)_* \left(f_*^{-1}\hat{\theta}_{\alpha,\varepsilon}(f_*\nu) \right) \end{aligned}$$

where π_1 is the projection on the first coordinate. To obtain the desired (34), we will actually show that for sufficiently high energy $\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu)$ even before projection one has

$$(35) \quad \text{TV}(\hat{\theta}_{\alpha,\varepsilon}(\nu), f_*^{-1}\hat{\theta}_{\alpha,\varepsilon}(f_*\nu)) < \delta_2.$$

To do so, note that both these measures are absolutely continuous with respect to $\nu \times \nu$:

$$\begin{aligned} \hat{\theta}_{\alpha,\varepsilon}(\nu) &= \frac{1}{\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu)} K_{\alpha,\varepsilon}(x, z) \nu(dx) \nu(dz), \\ f_*^{-1}\hat{\theta}_{\alpha,\varepsilon}(f_*\nu) &= \frac{1}{\tilde{\mathcal{E}}_{\alpha,\varepsilon}(f_*\nu)} K_{\alpha,\varepsilon}(f(x), f(z)) \nu(dx) \nu(dz). \end{aligned}$$

Now, (33) implies that once $K_{\alpha,\varepsilon}(x, z) > C_2 := \frac{20C''}{\delta_2}$, the $K_{\alpha,\varepsilon}$ parts of these densities are close to each other:

$$\frac{1}{1 + \frac{\delta_2}{6}} < \frac{K_{\alpha,\varepsilon}(f(x), f(z))}{K_{\alpha,\varepsilon}(x, z)} < 1 + \frac{\delta_2}{6}.$$

On the other hand, due to Corollary 3.17, once the energy $\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu)$ is sufficiently large, the normalization constants are also close to each other:

$$\frac{1}{1 + \frac{\delta_2}{6}} \tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu) < \tilde{\mathcal{E}}_{\alpha,\varepsilon}(f_*\nu) < \left(1 + \frac{\delta_2}{6}\right) \tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu).$$

Multiplying the two inequalities, we get that on the set

$$A := \{(x, z) \mid K_{\alpha,\varepsilon}(x, z) > C_2\}$$

the quotient of densities w.r.t. $\nu \times \nu$ of the two measures is in the interval

$$\left(\frac{1}{\left(1 + \frac{\delta_2}{6}\right)^2}, \left(1 + \frac{\delta_2}{6}\right)^2 \right).$$

Finally, the $\hat{\theta}_{\alpha,\varepsilon}(\nu)$ -measure of its complement does not exceed $\frac{C_2}{\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu)}$. Hence, if the energy $\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu)$ is sufficiently high to make sure that $\frac{C_2}{\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu)} < \frac{\delta_2}{6}$, the part that the normalized measures have in common is at least

$$\frac{1 - \frac{\delta_2}{6}}{\left(1 + \frac{\delta_2}{6}\right)^2} > 1 - \frac{\delta_2}{2},$$

and hence the total variation (35) indeed does not exceed δ_2 .

An application of Lemma 3.21 concludes the proof of the upper bound (28) for the Wasserstein distance. \square

Lemma 3.23 and Lemma 3.24 together imply Proposition 3.22. Indeed, we have:

$$\begin{aligned} W(f_*\theta_{\alpha,\varepsilon}(\nu), \theta_{\alpha,\varepsilon}(f_*\nu)) &\leq W(f_*\theta_{\alpha,\varepsilon}(\nu), f_*\theta'_{\alpha,\varepsilon}(\nu)) + \\ &\quad + W(f_*\theta'_{\alpha,\varepsilon}(\nu), \theta'_{\alpha,\varepsilon}(f_*\nu)) + W(\theta'_{\alpha,\varepsilon}(f_*\nu), \theta_{\alpha,\varepsilon}(f_*\nu)) \end{aligned}$$

The first and the third summands can be estimated directly using Lemma 3.23. It suffices to note that since $L(f), \gamma(f)^{-1} < R$ we have control over the increase of the Wasserstein distance after application of f . Indeed, a simple application of Jensen's inequality gives us

$$W(f_*\theta_{\alpha,\varepsilon}(\nu), f_*\theta'_{\alpha,\varepsilon}(\nu)) < R \cdot W(\theta_{\alpha,\varepsilon}(\nu), \theta'_{\alpha,\varepsilon}(\nu))^{\frac{2}{R}}.$$

For the last summand, to apply Lemma, we note that an upper bound on $L(f)$ and $\gamma(f)^{-1}$ implies that $f_*\nu$ is of sufficiently high energy provided that ν is of high energy.

Finally, the second summand is estimated directly by Lemma 3.24. \square

Proposition 3.25. *If a probability measure μ on $\text{Hol}(M)$ satisfies assumption (3) then for every $\delta > 0$ there exists $\alpha_0 > 0$ such that for every $0 < \alpha < \alpha_0$ there exists $C > 0$ such that for every $\varepsilon > 0$ and every measure $\nu \in \mathcal{M}$ such that $\mathcal{E}_{\alpha,\varepsilon}(\nu) > C$ the following formula holds:*

$$(36) \quad \frac{\mathbb{E}_\mu \left[\tilde{\mathcal{E}}_{\alpha,\varepsilon}(f_*\nu) \right]}{\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu)} \in (1 - \delta, 1 + \delta).$$

Proof. Let us fix some $\delta_E > 0$ and pick $\tilde{\delta} > 0$, such that $(1 + \tilde{\delta})^2 < 1 + \delta_E$. It follows directly from condition (3) and Hölder inequality that there exists $\alpha_0 > 0$, such that for every $0 < \alpha < \alpha_0$ one has

$$(37) \quad \int_{\text{Hol}(M)} \left(\frac{2(1 + \log(L(f)))}{\gamma(f)} \right)^\alpha d\mu(f) < 1 + \tilde{\delta}.$$

According to Corollary 3.9 we can choose C such that inequality (12) holds for $\tilde{\delta}$. Let $f \in \text{Hol}(M)$ be a random homeomorphism distributed with respect to the measure μ . Then integrating inequality (12) with respect to the measure μ and combining it with inequality (37) we get:

$$(38) \quad \frac{1}{1 + \delta_E} < \frac{1}{(1 + \tilde{\delta})^2} \leq \frac{\mathbb{E}_\mu [\mathcal{E}_{\alpha,\varepsilon}(f_*\nu)]}{\mathcal{E}_{\alpha,\varepsilon}(\nu)} \leq (1 + \tilde{\delta})^2 < 1 + \delta_E.$$

Let us now return back to the original (36). Namely, let $\delta > 0$ be given; choose and fix $\delta_E > 0$ such that $(1 + \delta_E)^3 < 1 + \delta$, and let α_0 be chosen w.r.t. δ_E so that (38) holds.

By Proposition 3.15, there exists $C_E > 0$ such that for any measure ν' on M and any $\varepsilon > 0$ one has

$$(39) \quad \frac{1}{1 + \delta_E} \mathcal{E}_{\alpha,\varepsilon}(\nu') - \frac{C_E}{1 + \delta_E} < \tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu') < (1 + \delta_E) \mathcal{E}_{\alpha,\varepsilon}(\nu') + C_E$$

Applying this for $\nu' = f_*\nu$ and taking the expectation w.r.t. μ provides

$$\begin{aligned} \frac{1}{1 + \delta_E} \mathbb{E}_\mu [\mathcal{E}_{\alpha,\varepsilon}(f_*\nu)] - \frac{C_E}{1 + \delta_E} &< \mathbb{E}_\mu \left[\tilde{\mathcal{E}}_{\alpha,\varepsilon}(f_*\nu) \right] \\ &< (1 + \delta_E) \mathbb{E}_\mu [\mathcal{E}_{\alpha,\varepsilon}(f_*\nu)] + C_E; \end{aligned}$$

using (38), we thus get

$$\frac{1}{(1 + \delta_E)^2} \mathcal{E}_{\alpha, \varepsilon}(\nu) - \frac{C_E}{1 + \delta_E} < \mathbb{E}_\mu \left[\tilde{\mathcal{E}}_{\alpha, \varepsilon}(f_* \nu) \right] < (1 + \delta_E)^2 \mathcal{E}_{\alpha, \varepsilon}(\nu) + C_E.$$

Finally, using (39) with $\nu' = \nu$ to estimate $\mathcal{E}_{\alpha, \varepsilon}$ via $\tilde{\mathcal{E}}_{\alpha, \varepsilon}$, we get

$$\begin{aligned} \frac{1}{(1 + \delta_E)^2} \cdot \frac{1}{(1 + \delta_E)} \left(\tilde{\mathcal{E}}_{\alpha, \varepsilon}(\nu) - C_E \right) - \frac{C_E}{1 + \delta_E} &< \mathbb{E}_\mu \left[\tilde{\mathcal{E}}_{\alpha, \varepsilon}(f_* \nu) \right] \\ &< (1 + \delta_E)^2 \cdot \left((1 + \delta_E) \tilde{\mathcal{E}}_{\alpha, \varepsilon}(\nu) + C_E \right) + C_E; \end{aligned}$$

as $(1 + \delta_E)^3 < 1 + \delta$, we obtain the desired

$$(1 + \delta)(\tilde{\mathcal{E}}_{\alpha, \varepsilon}(\nu) - C') < \mathbb{E}_\mu \left[\tilde{\mathcal{E}}_{\alpha, \varepsilon}(f_* \nu) \right] < (1 + \delta) \tilde{\mathcal{E}}_{\alpha, \varepsilon}(\nu) + C'$$

for some constant C' . As $\delta > 0$ was arbitrary (36) follows. \square

Remark 3.26. Notice that if instead of one measure μ on $\text{Hol}(M)$ we consider a compact K of such measures, such that there exists $\beta > 0$ and C_0 such that for every $\mu \in K$

$$\int_{\text{Hol}(M)} \left(\frac{1 + \log(L(f))}{\gamma(f)} \right)^\beta d\mu(f) < C_0$$

then constants α_0 and C in Proposition 3.25 can be chosen uniformly for all measures in K .

Proposition 3.27. *Under the assumptions of Theorem 2.4 there exists $\alpha > 0$, $\lambda < 1$, and $C < \infty$, such that for every $\varepsilon > 0$, every $\nu \in \mathcal{M}$ and every $\mu \in K$ if $\mathcal{E}_{\alpha, \varepsilon}(\nu) > C$ then*

$$(40) \quad \tilde{\mathcal{E}}_{\alpha, \varepsilon}(\mu * \nu) < \lambda \tilde{\mathcal{E}}_{\alpha, \varepsilon}(\nu).$$

Proof. Fix some small number $\delta > 0$, take $\lambda = 1 - \delta$, and assume the contrary: for any $\alpha > 0$ and $C < \infty$ there exists a measure $\mu \in K$, an $\varepsilon > 0$ and a probability measure ν with energy $\tilde{\mathcal{E}}_{\alpha, \varepsilon}(\nu) > C$, such that

$$(41) \quad \tilde{\mathcal{E}}_{\alpha, \varepsilon}(\mu * \nu) \geq (1 - \delta) \tilde{\mathcal{E}}_{\alpha, \varepsilon}(\nu).$$

Recall the following standard statement:

Lemma 3.28. *Assume that in some Hilbert space \mathcal{H} a probability measure is given; in other words, one is given a random variable v taking values in this space. Assume that this measure has finite second moment, and let $\bar{v} := \mathbb{E}v$ be its expectation. Then*

$$(42) \quad \mathbb{E}\langle v - \bar{v}, v - \bar{v} \rangle = \mathbb{E}\langle v, v \rangle - \langle \bar{v}, \bar{v} \rangle$$

Let us apply this statement to $L^2(M, \text{Leb})$. Namely, for a given measure ν its averaged image $\mu * \nu$ is the expectation of $f_* \nu$, where the homeomorphism f of M is taken randomly w.r.t. the measure μ . Hence, the same applies for the density $\rho_{\alpha, \varepsilon}[\nu]$, given by (14) (see Definition 3.12):

$$(43) \quad \rho_{\alpha, \varepsilon}[\mu * \nu] = \mathbb{E}_\mu \rho_{\alpha, \varepsilon}[f_* \nu].$$

Substituting this into (42) (with $v = \rho_{\alpha,\varepsilon}[f_*\nu]$), and taking into account the definition (15), we get

$$(44) \quad \mathbb{E}_\mu \left[\int_M (\rho_{\alpha,\varepsilon}[f_*\nu](y) - \rho_{\alpha,\varepsilon}[\mu * \nu](y))^2 d\text{Leb}(y) \right] = \\ = \mathbb{E}_\mu \left[\tilde{\mathcal{E}}_{\alpha,\varepsilon}(f_*\nu) \right] - \tilde{\mathcal{E}}_{\alpha,\varepsilon}(\mu * \nu);$$

in particular,

$$(45) \quad \tilde{\mathcal{E}}_{\alpha,\varepsilon}(\mu * \nu) \leq \mathbb{E}_\mu \left[\tilde{\mathcal{E}}_{\alpha,\varepsilon}(f_*\nu) \right].$$

By Proposition 3.25 we can find sufficiently small $\alpha > 0$ and sufficiently large $C < \infty$ (uniformly in μ), such that for any $\varepsilon > 0$ and any measure ν on M with $\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu) > C$ the inequality (36) holds, and thus (taking only the upper bound)

$$(46) \quad \mathbb{E}_\mu \left[\tilde{\mathcal{E}}_{\alpha,\varepsilon}(f_*\nu) \right] \leq (1 + \delta) \tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu).$$

Denote by ν a probability measure with $\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu) > C$, for which the inequality (41) holds. Substituting (46) and (41) in the right hand side of (44), we get

$$(47) \quad \mathbb{E}_\mu \left[\int_M (\rho_{\alpha,\varepsilon}(f_*\nu)(y) - \rho_{\alpha,\varepsilon}(\mu * \nu)(y))^2 d\text{Leb}(y) \right] < 2\delta \tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu).$$

and thus

$$(48) \quad \mathbb{E}_\mu \left[\int_M \frac{(\rho_{\alpha,\varepsilon}(f_*\nu)(y) - \rho_{\alpha,\varepsilon}(\mu * \nu)(y))^2}{\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu)} d\text{Leb}(y) \right] < 2\delta.$$

The final step is to show that measures

$$\theta_{\alpha,\varepsilon}(f_*\nu) = \frac{\rho_{\alpha,\varepsilon}(f_*\nu)(y)^2}{\tilde{\mathcal{E}}_{\alpha,\varepsilon}(f_*\nu)} d\text{Leb}(y) \quad \text{and} \quad \theta_{\alpha,\varepsilon}(\mu * \nu) = \frac{\rho_{\alpha,\varepsilon}(\mu * \nu)(y)^2}{\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\mu * \nu)} d\text{Leb}(y)$$

are close with high probability. And indeed, we have the following statement, estimating (even) the total variations distance:

Lemma 3.29. *Under the assumptions above*

$$(49) \quad \mathbb{P}_\mu \left[\text{TV}(\theta_{\alpha,\varepsilon}(f_*\nu), \theta_{\alpha,\varepsilon}(\mu * \nu)) > 10\sqrt[8]{\delta} \right] < 4\sqrt{\delta},$$

and hence

$$(50) \quad \mathbb{P}_\mu \left[W(\theta_{\alpha,\varepsilon}(f_*\nu), \theta_{\alpha,\varepsilon}(\mu * \nu)) > 10\text{diam}(M)\sqrt[8]{\delta} \right] < 4\sqrt{\delta}.$$

Proof of Lemma 3.29. We start by working with the non-normalized measures $\Theta_{\alpha,\varepsilon}(f_*\nu)$ and $\Theta_{\alpha,\varepsilon}(\mu * \nu)$. To simplify the notation, denote

$$g_f(y) := \rho_{\alpha,\varepsilon}[f_*\nu](y), \quad \bar{g}(y) := \rho_{\alpha,\varepsilon}[\mu * \nu](y).$$

Therefore, we have

$$\Theta_{\alpha,\varepsilon}[f_*\nu] = (g_f(y))^2 d\text{Leb}(y) \quad \text{and} \quad \Theta_{\alpha,\varepsilon}[\mu * \nu] = (\bar{g}(y))^2 d\text{Leb}(y).$$

From (41) we have

$$\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu) \leq \frac{1}{1 - \delta} \tilde{\mathcal{E}}_{\alpha,\varepsilon}(\mu * \nu).$$

Joining with (47), we have

$$(51) \quad \mathbb{E}_f \left[\int_M |g_f(y) - \bar{g}(y)|^2 d\text{Leb}(y) \right] \leq \frac{2\delta}{1-\delta} \tilde{\mathcal{E}}_{\alpha,\varepsilon}(\mu * \nu).$$

By Markov inequality, with the probability at least $1 - \frac{2\sqrt{\delta}}{1-\delta}$, one has

$$(52) \quad \int_M |g_f(y) - \bar{g}(y)|^2 d\text{Leb}(y) \leq \sqrt{\delta} \tilde{\mathcal{E}}_{\alpha,\varepsilon}(\mu * \nu) \\ = \sqrt{\delta} \Theta_{\alpha,\varepsilon}[\mu * \nu](M) = \sqrt{\delta} \int_M \bar{g}^2(y) d\text{Leb}(y).$$

Now, for any f for which (52) holds, consider the set

$$X_f := \{y \in M : |g_f(y) - \bar{g}(y)| \leq \sqrt[8]{\delta} \cdot \bar{g}(y)\}.$$

Again from Markov inequality type argument, one has

$$(53) \quad \Theta_{\alpha,\varepsilon}[\mu * \nu](X_f) \geq \left(1 - \sqrt[4]{\delta}\right) \Theta_{\alpha,\varepsilon}[\mu * \nu](M).$$

Indeed, on the complement $M \setminus X_f$ one has $|g_f(y) - \bar{g}(y)|^2 > \sqrt[4]{\delta} \cdot \bar{g}^2(y)$; integrating, one gets

$$\int_{M \setminus X_f} |g_f(y) - \bar{g}(y)|^2 d\text{Leb}(y) > \sqrt[4]{\delta} \cdot \Theta_{\alpha,\varepsilon}[\mu * \nu](M \setminus X_f).$$

Thus, if (53) did not hold, it would imply

$$\Theta_{\alpha,\varepsilon}[\mu * \nu](M \setminus X_f) > \sqrt[4]{\delta} \cdot \Theta_{\alpha,\varepsilon}[\mu * \nu](M),$$

hence providing a contradiction with (52).

Now, (53) implies that the non-normalized measures $\Theta_{\alpha,\varepsilon}[f_*\nu]$ and $\Theta_{\alpha,\varepsilon}[\mu * \nu]$ share a common part

$$\Xi_f := (1 - \sqrt[8]{\delta})^2 \bar{g}^2(y) \cdot \mathbf{1}_{X_f}(y) d\text{Leb}(y)$$

of measure at least

$$(54) \quad \Xi_f(M) \geq (1 - \sqrt[8]{\delta})^2 (1 - \sqrt[4]{\delta}) \cdot \Theta_{\alpha,\varepsilon}[\mu * \nu](M).$$

Finally, note that the normalization constants are also close with high probability, namely, for f such that the inequality (52) holds. Indeed, the inequality (52) can be rewritten as

$$\|g_f - \bar{g}\|_{L_2(M)}^2 \leq \sqrt{\delta} \cdot \|\bar{g}\|_{L_2(M)}^2;$$

hence, for any f for which it holds, the triangle inequality implies

$$(55) \quad \Theta_{\alpha,\varepsilon}[f_*\nu](M) = \|g_f\|_{L_2(M)}^2 \leq (1 + \sqrt[4]{\delta})^2 \Theta_{\alpha,\varepsilon}[\mu * \nu](M).$$

Now, the normalized measures $\theta_{\alpha,\varepsilon}[f_*\nu]$ and $\theta_{\alpha,\varepsilon}[\mu * \nu]$ share the common part $\frac{1}{C_f} \Xi_f$, where

$$C_f = \max(\Theta_{\alpha,\varepsilon}[f_*\nu](M), \Theta_{\alpha,\varepsilon}[\mu * \nu](M))$$

is the maximum of two normalization constants. Using (54) and (55), we see that this part has measure at least

$$\frac{1}{C_f} \Xi_f(M) \geq \frac{(1 - \sqrt[8]{\delta})^2 (1 - \sqrt[4]{\delta})}{(1 + \sqrt[4]{\delta})^2} \geq 1 - 10 \sqrt[8]{\delta}.$$

Hence, for any f satisfying (52) the total variation distance between the normalized measures does not exceed

$$\mathrm{TV}(\theta_{\alpha,\varepsilon}(f_*\nu), \theta_{\alpha,\varepsilon}(\mu * \nu)) \leq 10\sqrt[8]{\delta}.$$

As (52) holds with the probability at least $1 - \frac{2\sqrt{\delta}}{1-\delta} > 1 - 4\sqrt{\delta}$, this establishes conclusion (49). Finally, this immediately implies conclusion (50) due to Lemma 3.21. \square

We are now ready to conclude the proof of Proposition 3.27. Namely, the conclusion (50) states that with high probability the measure $\theta_{\alpha,\varepsilon}(f_*\nu)$ is close to the deterministic one, $\theta_{\alpha,\varepsilon}(\mu * \nu)$. The next step is to show that with high probability the first measure is close to f_* -image of a given measure, $\theta_{\alpha,\varepsilon}(\nu)$.

It easily follows from inequality (3) that we can choose $R < \infty$ (again, uniformly in μ), such that

$$\mathbb{P}_\mu [L(f) < R \text{ and } \gamma(f)^{-1} < R] > 1 - \sqrt[4]{\delta}$$

By Proposition 3.22 we can choose C big enough, so that for any $f \in \mathrm{Hol}(M)$, such that $L(f), \gamma(f)^{-1} < R$ inequality (22) holds, and using triangle inequality we arrive to

$$(56) \quad \mathbb{P}_\mu \left[W(f_*\theta_{\alpha,\varepsilon}(\nu), \theta_{\alpha,\varepsilon}(\mu * \nu)) > 10\mathrm{diam}(M)\sqrt[8]{\delta} + \delta \right] < 5\sqrt[4]{\delta}.$$

Now consider the sequence $\delta_n = \frac{1}{n}$ and denote by μ_n and ν_n the corresponding measures and corresponding parameters by α_n, ε_n , for which (41) holds. Then the measures $\theta_{\alpha_n, \varepsilon_n}(\nu_n)$ have a weakly convergent subsequence. Furthermore, we can take another subsequence, such that the measures $\theta_{\alpha_{n_k}, \varepsilon_{n_k}}(\mu_{n_k} * \nu_{n_k})$ also converge. Passing to a subsequence one more time we can guarantee (due to compactness of K) that

$$\lim_{k \rightarrow \infty} \mu_{n_k} = \mu$$

for some $\mu \in K$. It remains to notice that due to inequality (56) the limit measure

$$m = \lim_{k \rightarrow \infty} \theta_{\alpha_{n_k}, \varepsilon_{n_k}}(\nu_{n_k})$$

has an almost surely constant image under the action of $f \in \mathrm{supp}(\mu)$, that is equal to

$$\tilde{m} = \lim_{k \rightarrow \infty} \theta_{\alpha_{n_k}, \varepsilon_{n_k}}(\mu_{n_k} * \nu_{n_k}) = \lim_{k \rightarrow \infty} \theta_{\alpha_{n_k}, \varepsilon_{n_k}}(\mu * \nu_{n_k}),$$

which contradicts our assumption. \square

Finally, we note that the conclusion of Proposition 3.27 can be modified to include all possible measures ν (without assuming the energy higher than C). To do so, we need to adjust the upper bound (40) in order to include low-energy measures ν :

Lemma 3.30. *Assume that the measure μ satisfies the condition (3) with some β , and that $\alpha < \beta$ and C are given. Then there exists C' such that for any $\varepsilon > 0$ and any measure ν on M with $\mathcal{E}_{\alpha,\varepsilon}(\nu) \leq C$ one has*

$$\mathcal{E}_{\alpha,\varepsilon}(\mu * \nu) \leq C'.$$

Corollary 3.31. *In the assumptions of Proposition 3.27 one can conclude that there exist $\alpha > 0, \tilde{C} < \infty, \lambda < 1$, such that for every $\mu \in K$, every $\varepsilon > 0$ and every measure ν on M*

$$(57) \quad \mathcal{E}_{\alpha,\varepsilon}(\mu * \nu) < \max(\lambda\mathcal{E}_{\alpha,\varepsilon}(\nu), \tilde{C}).$$

Proof of Lemma 3.30. Condition (3) implies the finiteness of the expectation

$$(58) \quad C_I := \mathbb{E}_\mu \left(\frac{2(1 + \log(L(f)))}{\gamma(f)} \right)^\alpha < \infty.$$

Now, (45) implies that for every measure ν we have

$$\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\mu * \nu) \leq \mathbb{E}_\mu \tilde{\mathcal{E}}_{\alpha,\varepsilon}(f_*\nu).$$

At the same time Proposition 3.15 for $\delta = \frac{1}{2}$ implies that for some constant C_1 one has for any measure ν'

$$\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu') \leq \max(2\mathcal{E}_{\alpha,\varepsilon}(\nu'), C_1).$$

Applying this for $\nu' = f_*\nu$ and joining it with Corollary 3.10, we get

$$\begin{aligned} \tilde{\mathcal{E}}_{\alpha,\varepsilon}(f_*\nu) &\leq \max(2\mathcal{E}_{\alpha,\varepsilon}(f_*\nu), C_1) \leq \\ &\leq 2 \max \left(2 \left(\frac{2(1 + \log(L(f)))}{\gamma(f)} \right)^\alpha \mathcal{E}_{\alpha,\varepsilon}(\nu), C_2 \right) + C_1, \end{aligned}$$

where C_2 is equal to C from Corollary (3.10). Taking the expectation w.r.t. μ and using (58), we finally get a uniform bound

$$\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\mu * \nu) \leq 4C_I C + 2C_2 + C_1 =: C'. \quad \square$$

Proof of Corollary 3.31. In the proof of Proposition 3.27 we can take α arbitrarily small, so we can assume that $\alpha < \beta$. Applying Proposition 3.15 to both sides of (40), where we take δ sufficiently small so that $\frac{1-\delta}{1+\delta} > \lambda$, allows to conclude that there exists some constant C_3 such that for all ε and ν

$$\mathcal{E}_{\alpha,\varepsilon}(\mu * \nu) < \lambda' \mathcal{E}_{\alpha,\varepsilon}(\nu) \quad \text{if } \mathcal{E}_{\alpha,\varepsilon}(\nu) > C_3,$$

where $\lambda' = \frac{1+\delta}{1-\delta} \lambda < 1$. Meanwhile, Lemma 3.30 implies that there exists some C_4 such that

$$\mathcal{E}_{\alpha,\varepsilon}(\mu * \nu) < C_4 \quad \text{if } \mathcal{E}_{\alpha,\varepsilon}(\nu) \leq C_3.$$

Taking $\tilde{C} := C_4$, we get the desired (57). \square

Now we are ready to complete the proof of Theorem 2.4 (and hence of Theorem 2.3).

Proof of Theorem 2.4. Let α, \tilde{C} be as in Corollary 3.31: for any $\mu \in K$, any $\varepsilon > 0$ and for any measure ν one has

$$\mathcal{E}_{\alpha,\varepsilon}(\mu * \nu) < \max(\lambda \mathcal{E}_{\alpha,\varepsilon}(\nu), \tilde{C});$$

applying this n times, we get

$$\mathcal{E}_{\alpha,\varepsilon}(\mu^n * \nu) < \max(\lambda^n \mathcal{E}_{\alpha,\varepsilon}(\nu), \tilde{C}).$$

Recall that Lemma 3.6 gives a uniform upper bound

$$\mathcal{E}_{\alpha,\varepsilon}(\nu) \leq \frac{\omega_k (-\log(\varepsilon))^{\alpha-1}}{e^k \varepsilon^k},$$

(and, without loss of generality, we can assume that $\alpha < 1$) choose $\varepsilon := \lambda^{\frac{2}{k}}$, then we have

$$\mathcal{E}_{\alpha,\varepsilon}(\mu^n * \nu) < \max \left(\frac{\omega_k}{e^k}, \tilde{C} \right).$$

Now, applying Lemma 3.7 for any $r > \varepsilon = \kappa^n$, where $\kappa := \lambda^{\frac{1}{k}}$, we deduce that

$$(\mu^n * \nu)(B_r(x)) < C_{\alpha,k} \sqrt{\max\left(\frac{\omega_k}{e^k}, \tilde{C}\right)} \cdot (-\log(r))^{-\frac{\alpha}{2}},$$

which completes the proof of Theorem 2.4. \square

4. LIPSCHITZ HOMEOMORPHISMS

In this section we will outline the proof of Theorem 2.9. Since it utilizes the same method as the proof of Theorem 2.4 we will give modified versions of key propositions avoiding technical details. In this section we will work with a fixed $\alpha > 0$.

First, we will need an analog of Proposition 3.8 to have an estimate for $\mathcal{E}_{\alpha,\varepsilon}(f_*\nu)$ for a Lipschitz continuous f . It can be formulated as follows:

Proposition 4.1. *For every $\delta > 0$ there exists $\varepsilon_0 > 0$ and $B < \infty$, such that every $0 < \varepsilon < \varepsilon_0$, every $f \in \text{Lip}(M)$ and every $\nu \in \mathcal{M}$ the following holds:*

$$(59) \quad \mathcal{E}_{\alpha,\varepsilon}(f_*\nu) \leq (1 + \delta \log(\mathfrak{L}(f)))^\alpha (\mathcal{E}_{\alpha,\varepsilon}(\nu) + B).$$

In this case useful corollaries can be formulated as follows:

Corollary 4.2. *For every $\delta > 0$ there exists $C < \infty$ such that for every $\varepsilon > 0$, every $f \in \text{Lip}(M)$ and every $\nu \in \mathcal{M}$, such that $\mathcal{E}_{\alpha,\varepsilon}(\nu) > C$ the following holds:*

$$(60) \quad \frac{1}{1 + \delta} (1 + \delta \log(\mathfrak{L}(f)))^{-\alpha} < \frac{\mathcal{E}_{\alpha,\varepsilon}(f_*\nu)}{\mathcal{E}_{\alpha,\varepsilon}(\nu)} < (1 + \delta) (1 + \delta \log(\mathfrak{L}(f)))^\alpha.$$

Corollary 4.3. *There exists $C < \infty$ such that for every $\varepsilon > 0$, every $f \in \text{Lip}(M)$ and every $\nu \in \mathcal{M}$ the following holds:*

$$\mathcal{E}_{\alpha,\varepsilon}(f_*\nu) < \max(2(1 + \log(\mathfrak{L}(f)))^\alpha \mathcal{E}_{\alpha,\varepsilon}(\nu), C).$$

Notice, that Proposition 3.15 does not depend on the regularity of f , hence it remains unchanged. That, once again, allows us to establish an estimate for $\tilde{\mathcal{E}}_{\alpha,\varepsilon}(f_*\nu)$ (analog of Corollary 3.17):

Corollary 4.4. *For every $\delta > 0$ and every $R < \infty$ there exists $C > 0$ such that for every $f \in \text{Lip}(M)$ with $\mathfrak{L}(f) < R$, every $\varepsilon > 0$ and every ν such that $\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu) > C$ or $\mathcal{E}_{\alpha,\varepsilon}(\nu) > C$ one has*

$$\frac{\tilde{\mathcal{E}}_{\alpha,\varepsilon}(f_*\nu)}{\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu)} \in (1 - \delta, 1 + \delta).$$

Next on the list is the analog of Proposition 3.22, which is formulated below.

Proposition 4.5. *For any $\delta > 0$, any $R < \infty$ and any $\alpha \in (0, 1]$ there exists $C > 0$ such that for any $\varepsilon > 0$, any $f \in \text{Lip}(M)$, such that $\mathfrak{L}(f) < R$ and any $\nu \in \mathcal{M}$ such that $\mathcal{E}_{\alpha,\varepsilon}(\nu) > C$ or $\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu) > C$ one has*

$$W(f_*\theta_{\alpha,\varepsilon}[\nu], \theta_{\alpha,\varepsilon}[f_*\nu]) < \delta.$$

Once again, the proof repeats the one of Proposition 3.22 up to minor computational details. Finally, the uniform estimate (5) allows us to establish the following analog of Proposition 3.25:

Proposition 4.6. *If a probability measure μ on $\text{Lip}(M)$ satisfies assumption (5) then for every $\delta > 0$ there exists $C > 0$ such that for every $\varepsilon > 0$ and every measure $\nu \in \mathcal{M}$ such that $\mathcal{E}_{\alpha,\varepsilon}(\nu) > C$ the following formula holds:*

$$\frac{\mathbb{E}_\mu \left[\tilde{\mathcal{E}}_{\alpha,\varepsilon}(f_*\nu) \right]}{\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu)} \in (1 - \delta, 1 + \delta).$$

Utilizing these modified statements we are able to repeat the proof of the contracting property for $\tilde{\mathcal{E}}_{\alpha,\varepsilon}$ (analog of Proposition 3.27):

Proposition 4.7. *Under the assumptions of Theorem 2.8 there exists $\lambda < 1$ and $C < \infty$, such that for every $\mu \in K$, every $\varepsilon > 0$ and every $\nu \in \mathcal{M}$ if $\mathcal{E}_{\alpha,\varepsilon}(\nu) > C$ then*

$$\tilde{\mathcal{E}}_{\alpha,\varepsilon}(\mu * \nu) < \lambda \tilde{\mathcal{E}}_{\alpha,\varepsilon}(\nu).$$

From here Theorem 2.9 (and hence Theorem 2.8) follows.

APPENDIX A. TECHNICAL LEMMATA

In this section we prove establish some properties of functions φ_α and $\varphi_{\alpha,\varepsilon}$ and then prove Proposition 3.4.

A straightforward computation allows us to prove the following

Lemma A.1. *For any $0 < r < \frac{1}{e}$*

$$\int_{\bar{x} \in \mathbb{R}^k, r < |\bar{x}| < \frac{1}{e}} \varphi_\alpha(|\bar{x}|)^2 d\bar{x} = c_{\alpha,k} ((-\log(r))^\alpha - 1)$$

Proof.

$$\begin{aligned} \int_{\bar{x} \in \mathbb{R}^k, r < |\bar{x}| < \frac{1}{e}} \varphi_\alpha(|\bar{x}|)^2 d\bar{x} &= \\ &= \int_{r < y < \frac{1}{e}} \int_{\theta \in \mathbb{S}^{k-1}} \frac{(-\log(y))^{\alpha-1}}{y^k} J(\theta) d\theta y^{k-1} dy = \\ &= k\omega_k \int_{r < y < \frac{1}{e}} \frac{(-\log(y))^{\alpha-1}}{y} dy = \\ &= c_{\alpha,k} ((-\log(r))^\alpha - 1), \end{aligned}$$

where $J(\theta)y^{k-1}$ is a Jacobian of a k -dimensional spherical coordinates transformation, ω_k is a volume of a k -dimensional unit ball and

$$c_{\alpha,k} = \frac{k\omega_k}{\alpha}.$$

□

Let us define a function

$$V_{\alpha,\varepsilon}(r) = \begin{cases} c_{\alpha,k} (-\log(r))^\alpha, & \text{for } \varepsilon < r < 1/e; \\ c_{\alpha,k} (-\log(\varepsilon))^\alpha, & \text{for } r \leq \varepsilon. \end{cases}$$

Our next goal is to prove

Proposition A.2. *For every $\alpha_0 > 0$ and every $\delta > 0$ there exists $\varepsilon_0 > 0$ and $r_0 > 0$, such that for every $0 < \alpha < \alpha_0$, for every $0 < \varepsilon < \varepsilon_0$, and every $0 \leq r < r_0$ we have*

$$\frac{1}{1 + \delta} < \frac{U_{\alpha, \varepsilon}(r)}{V_{\alpha, \varepsilon}(r)} < 1 + \delta.$$

We will start with two useful lemmata.

Lemma A.3. *For every $\alpha > 0$ we have*

$$\frac{2 \int_{x_1 > 2r} \varphi_\alpha^2(|\bar{x}|) d\bar{x}}{c_{\alpha, k} (-\log(r))^\alpha} \xrightarrow{r \rightarrow 0} 1, \quad \text{where } \bar{x} = (x_1, x_2, \dots, x_k) \in \mathbb{R}^k.$$

Moreover, if we fix $\alpha_0 > 0$ then the convergence is uniform with respect to $\alpha \in (0, \alpha_0)$.

Proof. We start with the following upper bound for the integral in question:

$$2 \int_{x_1 > 2r} \varphi_\alpha^2(|\bar{x}|) d\bar{x} < \int_{|\bar{x}| > 2r} \varphi_\alpha^2(|\bar{x}|) d\bar{x} = c_{\alpha, k} ((-\log(2r))^\alpha - 1).$$

To estimate the integral from below we first represent it as follows:

$$\begin{aligned} 2 \int_{x_1 > 2r} \varphi_\alpha^2(|x|) dx &= \int_{|x| > 2r} \varphi_\alpha^2(|x|) dx - \int_{|x| > 2r, |x_1| < 2r} \varphi_\alpha^2(|x|) dx = \\ &= c_{\alpha, k} ((-\log(2r))^\alpha - 1) - \int_{|x| > 2r, |x_1| < 2r} \varphi_\alpha^2(|x|) dx. \end{aligned}$$

It remains to show that

$$(61) \quad \frac{\int_{|\bar{x}| > 2r, |x_1| < 2r} \varphi_\alpha^2(|x|) dx}{c_{\alpha, k} (-\log(r))^\alpha} \xrightarrow{r \rightarrow 0} 0$$

uniformly in $\alpha \in (0, \alpha_0)$. First of all, notice that

$$\int_{2r < |\bar{x}| < 4r} \varphi_\alpha^2(|\bar{x}|) d\bar{x} = c_{\alpha, k} ((-\log(2r))^\alpha - (-\log(4r))^\alpha).$$

For any $x > y > 1$ and $\alpha > 0$ we have

$$x^\alpha - y^\alpha < \alpha(x^{\alpha-1} + y^{\alpha-1})(x - y),$$

which implies

$$\begin{aligned} \int_{2r < |\bar{x}| < 4r} \varphi_\alpha^2(|\bar{x}|) d\bar{x} &< \\ &< c_{\alpha, k} \cdot \alpha \left((-\log(2r))^{\alpha-1} + (-\log(4r))^{\alpha-1} \right) (\log(4r) - \log(2r)), \end{aligned}$$

hence it is enough to prove that

$$\frac{\int_{|\bar{x}| > 4r, |x_1| < 2r} \varphi_\alpha^2(|x|) dx}{c_{\alpha, k} (-\log(r))^\alpha} \xrightarrow{r \rightarrow 0} 0$$

uniformly for $\alpha \in (0, \alpha_0)$. We proceed by performing polar change of coordinates. For $\bar{x} = (x_1, x_2, \dots, x_k) \in \mathbb{R}^k$ we write:

$$\begin{cases} x_1 &= t \cos(\theta_{k-1}); \\ x_2 &= t \cos(\theta_{k-2}) \sin(\theta_{k-1}); \\ &\vdots \\ x_{k-2} &= t \cos(\theta_2) \sin(\theta_3) \dots \sin(\theta_{k-1}); \\ x_{k-1} &= t \cos(\theta_1) \sin(\theta_2) \dots \sin(\theta_{k-1}); \\ x_k &= t \sin(\theta_1) \sin(\theta_2) \dots \sin(\theta_{k-1}), \end{cases}$$

where $t = |\bar{x}|$ and $\theta_1 \in [0, 2\pi)$, $\theta_j \in [0, \pi)$ for $2 \leq j < k$. In polar coordinates the domain $\{\bar{x} : |\bar{x}| > 4r, |x_1| < 2r\}$ becomes $\{\bar{x} : t > 4r, -2r/t < \cos(\theta_{k-1}) < 2r/t\}$. The Jacobian is equal to

$$J = (-1)^{k-1} t^{k-1} \prod_{j=2}^{k-1} \sin^{j-1}(\theta_j),$$

hence for some $C_k > 0$ that depends only on the dimension k we have:

$$\begin{aligned} \int_{|x| > 4r, |x_1| < 2r} \varphi_\alpha^2(|x|) dx &= \\ &= C_k \left| \int_{4r}^{1/e} \int_{\arccos(-2r/t)}^{\arccos(2r/t)} \sin^{k-2}(\theta_{k-1}) \frac{(-\log(t))^{\alpha-1}}{t^k} t^{k-1} d\theta_{k-1} dt \right| < \\ &< C_k \int_{4r}^{1/e} \int_{\arccos(-2r/t)}^{\arccos(2r/t)} \frac{(-\log(t))^{\alpha-1}}{t} d\theta_{k-1} dt. \end{aligned}$$

Since $2r/t < 1/2$ we have

$$\arccos(2r/t) - \arccos(-2r/t) < 100 \frac{r}{t}.$$

Hence

$$\begin{aligned} \int_{|x| > 4r, |x_1| < 2r} \varphi_\alpha^2(|x|) dx &< 100 C_k r \int_{4r}^{1/e} \frac{(-\log(t))^{\alpha-1}}{t^2} dt \leq \\ &\leq 100 C_k r (-\log(4r))^{\alpha-1} \int_{4r}^{1/e} \frac{1}{t^2} dt \leq 100 C_k (-\log(4r))^{\alpha-1}. \end{aligned}$$

Dividing the last quantity by $(-\log(r))^\alpha$ we finish the proof of equality (61) and the lemma. \square

Lemma A.4. *For every $\alpha > 0$ we have*

$$\frac{\int_{|x_1| \leq 2r} \varphi_\alpha(|\bar{x}|) \varphi_\alpha(|\bar{r} - \bar{x}|) d\bar{x}}{(-\log(r))^\alpha} \rightarrow_{r \rightarrow 0} 0.$$

Moreover, if we fix $\alpha_0 > 0$ then the convergence is uniform with respect to $\alpha \in (0, \alpha_0)$.

Proof. First of all, notice that

$$\int_{|x_1| \leq 2r} \varphi_\alpha(|\bar{x}|) \varphi_\alpha(|\bar{r} - \bar{x}|) d\bar{x} < 8 \int_{0 < x_1 < r/2} \varphi_\alpha(|\bar{x}|) \varphi_\alpha(|\bar{r} - \bar{x}|) d\bar{x}.$$

Let us split the integral on the right in two parts:

$$\begin{aligned} & \int_{0 < x_1 < r/2} \varphi_\alpha(|\bar{x}|) \varphi_\alpha(|\bar{r} - \bar{x}|) d\bar{x} = \\ & = \int_{0 < x_1 < r/2, |\bar{x}| > r/2} \varphi_\alpha(|\bar{x}|) \varphi_\alpha(|\bar{r} - \bar{x}|) d\bar{x} + \int_{|\bar{x}| < r/2} \varphi_\alpha(|\bar{x}|) \varphi_\alpha(|\bar{r} - \bar{x}|) d\bar{x}. \end{aligned}$$

The first summand can be estimated by

$$\int_{0 < x_1 < r/2, |\bar{x}| > r/2} \varphi_\alpha^2(|\bar{x}|) d\bar{x},$$

which goes to zero uniformly for $\alpha \in (0, \alpha_0)$ after division by $(-\log(r))^\alpha$ according to equation (61). As for the second summand, we write

$$\begin{aligned} (62) \quad & \int_{|\bar{x}| < r/2} \varphi_\alpha(|\bar{x}|) \varphi_\alpha(|\bar{r} - \bar{x}|) d\bar{x} < \varphi_\alpha(r/2) \int_{|\bar{x}| < r/2} \varphi_\alpha(|\bar{x}|) d\bar{x} = \\ & = k\omega_k \frac{(-\log(r/2))^{\frac{\alpha-1}{2}}}{(r/2)^{k/2}} \int_0^{r/2} \frac{(-\log(t))^{\frac{\alpha-1}{2}}}{t^{k/2}} t^{k-1} dt. \end{aligned}$$

We will focus on the last integral. If $\alpha \leq 1$ then

$$\int_0^{r/2} (-\log(t))^{\frac{\alpha-1}{2}} t^{k/2-1} dt \leq \int_0^{r/2} t^{k/2-1} dt = \frac{2}{k} \left(\frac{r}{2}\right)^{k/2}.$$

Combining that with formula (62) we get

$$\frac{\int_{|\bar{x}| < r/2} \varphi_\alpha(|\bar{x}|) \varphi_\alpha(|\bar{r} - \bar{x}|) d\bar{x}}{(-\log(r))^\alpha} < 2\omega_k \frac{(-\log(r/2))^{\frac{\alpha-1}{2}}}{(-\log(r))^\alpha}$$

and uniform convergence follows.

If $\alpha > 1$ after taking the last integral in formula (62) by parts we obtain:

$$\begin{aligned} (63) \quad & \int_0^{r/2} (-\log(t))^{\frac{\alpha-1}{2}} t^{k/2-1} dt = \\ & = \frac{2(-\log(t))^{\frac{\alpha-1}{2}} t^{k/2}}{k} \Big|_0^{r/2} + \frac{\alpha-1}{k} \int_0^{r/2} (-\log(t))^{\frac{\alpha-3}{2}} t^{k/2-1} dt. \end{aligned}$$

Since r goes to zero we can assume that $-\log(r/2) > 2(\alpha_0 - 1)/k$. In that case we can write

$$\begin{aligned} & \frac{\alpha-1}{k} \int_0^{r/2} (-\log(t))^{\frac{\alpha-3}{2}} t^{k/2-1} dt = \\ & = \frac{\alpha-1}{k} \int_0^{r/2} (-\log(t))^{\frac{\alpha-3}{2}} (-\log(t)) (-\log(t))^{-1} t^{k/2-1} dt < \\ & < \frac{\alpha-1}{k} (-\log(r/2))^{-1} \int_0^{r/2} (-\log(t))^{\frac{\alpha-1}{2}} t^{k/2-1} dt < \\ & < \frac{1}{2} \int_0^{r/2} (-\log(t))^{\frac{\alpha-1}{2}} t^{k/2-1} dt. \end{aligned}$$

Together with formula (63) the last inequality implies

$$\int_0^{r/2} (-\log(t))^{\frac{\alpha-1}{2}} t^{k/2-1} dt < \frac{4}{k} \left(-\log\left(\frac{r}{2}\right)\right)^{\frac{\alpha-1}{2}} \left(\frac{r}{2}\right)^{k/2}$$

and the uniform convergence follows. \square

We are ready to finish

Proof of Proposition A.2. Let us fix $\alpha_0 > 0$ and $\delta > 0$. We start by considering the case $r \geq \varepsilon$. Then, for the upper bound we write

$$U_{\alpha,\varepsilon}(r) < U_{\alpha}(r) < 2 \int_{|x_1| > 2r} \varphi_{\alpha}(|\bar{x}|) \varphi_{\alpha}(|\bar{r} - \bar{x}|) d\bar{x} + \int_{|x_1| \leq 2r} \varphi_{\alpha}(|\bar{x}|) \varphi_{\alpha}(|\bar{r} - \bar{x}|) d\bar{x}.$$

According to Lemma A.4 for r small enough (with the smallness being uniform in $\alpha \in (0, \alpha_0)$) we can guarantee that

$$\frac{\int_{|x_1| \leq 2r} \varphi_{\alpha}(|\bar{x}|) \varphi_{\alpha}(|\bar{r} - \bar{x}|) d\bar{x}}{c_{\alpha,k} (-\log(r))^{\alpha}} < \delta.$$

As for the first summand, we estimate it (again, for r small enough with the smallness being uniform in $\alpha \in (0, \alpha_0)$) using Lemma A.3 as follows:

$$2 \int_{|x_1| > 2r} \varphi_{\alpha}(|\bar{x}|) \varphi_{\alpha}(|\bar{r} - \bar{x}|) d\bar{x} < 2 \int_{|x_1| > r} \varphi_{\alpha}^2(|\bar{x}|) d\bar{x} < (1 + \delta) c_{\alpha,k} (-\log(r))^{\alpha}.$$

Combining these inequalities we get (for $r > \varepsilon$ and r small enough)

$$\frac{U_{\alpha,\varepsilon}(r)}{V_{\alpha,\varepsilon}(r)} < 1 + 2\delta.$$

To estimate this quotient from below notice that

$$\begin{aligned} U_{\alpha,\varepsilon}(r) &> 2 \int_{|x_1| > r+\varepsilon} \varphi_{\alpha}(|\bar{x}|) \varphi_{\alpha}(|\bar{r} - \bar{x}|) d\bar{x} \geq \\ &\geq 2 \int_{|x_1| > 2r} \varphi_{\alpha}(|\bar{x}|) \varphi_{\alpha}(|\bar{r} - \bar{x}|) d\bar{x} > 2 \int_{|x_1| > 2r} \varphi_{\alpha}^2(|\bar{x}|) d\bar{x}. \end{aligned}$$

Using Lemma A.3 once again we conclude that for r small enough (with the smallness being uniform in $\alpha \in (0, \alpha_0)$) we have

$$\frac{2 \int_{|x_1| > 2r} \varphi_{\alpha}^2(|\bar{x}|) d\bar{x}}{c_{\alpha,k} (-\log(r))^{\alpha}} > \frac{1}{1 + \delta},$$

which finishes the proof in the case $r \geq \varepsilon$.

Let us now assume that $r < \varepsilon$. In that case the upper bound can be established as follows:

$$U_{\alpha,\varepsilon}(r) < U_{\alpha,\varepsilon}(0) = \int_{|\bar{x}| < \varepsilon} \frac{(-\log(\varepsilon))^{\alpha-1}}{\varepsilon^k} d\bar{x} + \int_{|\bar{x}| > \varepsilon} \frac{(-\log(|\bar{x}|))^{\alpha-1}}{|\bar{x}|^k} d\bar{x}.$$

Applying Lemma A.1 to the second summand and computing the first we arrive to

$$U_{\alpha,\varepsilon}(r) < \omega_k (-\log(\varepsilon))^{\alpha-1} + c_{\alpha,k} ((-\log(\varepsilon))^{\alpha} - 1),$$

which guarantees that for ε small enough (with the smallness once again being uniform in $\alpha \in (0, \alpha_0)$) and $r < \varepsilon$ we have

$$\frac{U_{\alpha,\varepsilon}(r)}{V_{\alpha,\varepsilon}(r)} < 1 + \delta.$$

To finish the proof of Proposition A.2 it remains to notice that for $r < \varepsilon$

$$U_{\alpha,\varepsilon}(r) > U_{\alpha,\varepsilon}(\varepsilon).$$

As we saw in the first part of the proof, for ε small enough (with the smallness being uniform in $\alpha \in (0, \alpha_0)$) we have

$$\frac{U_{\alpha,\varepsilon}(\varepsilon)}{V_{\alpha,\varepsilon}(\varepsilon)} > \frac{1}{1+\delta}$$

and the lower bound follows. \square

Finally, using Proposition A.2 we are able to prove Proposition 3.4

- Proof of Proposition 3.4.* **(I)** First, notice that the function $\varphi_{\alpha,\varepsilon}$ is non-increasing. Using that fact Part **(I)** follows from the definition of $U_{\alpha,\varepsilon}$, for details see [GKM, Lemma 6.6].
- (II)** Choosing $\delta = 1$ in Proposition A.2 we obtain inequality (7).
- (III)** Choosing $\delta = 1$ in Proposition A.2 we obtain inequality (8).
- (IV)** Thanks to Proposition A.2 it is enough to prove inequality (9) for $V_{\alpha,\varepsilon}(r_1)$ and $V_{\alpha,\varepsilon}(r_2)$, which easily follows from the definition. \square

ACKNOWLEDGEMENTS

The author is grateful to A. Gorodetski and V. Kleptsyn for fruitful discussions. The author was supported in part by NSF grant DMS-2247966 (PI: A. Gorodetski).

REFERENCES

- [A] L. Arnold, Random dynamical systems. *Dynamical systems (Montecatini Terme, 1994)*, pp. 1–43, Lecture Notes in Math., **1609**, Springer, Berlin, 1995.
- [BBS] V. Baladi, M. Benedicks, D. Schnellmann, Whitney-Hölder continuity of the SRB measure for transversal families of smooth unimodal maps. *Invent. Math.*, **201** (2015), no.3, 773–844.
- [BV] L. Barreira, C. Valls, Hölder Grobman-Hartman linearization. *Discrete Contin. Dyn. Syst.*, **18** (2007), no.1, 187–197.
- [BQ] Y. Benoist, J. Quint, Random walks on reductive groups. *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics* **62**, Springer, Cham, 2016, xi+323 pp.
- [BQ16] Y. Benoist, J. Quint, Central Limit Theorem for Linear Groups, *The Annals of Probability*, **44** (2016), pp. 1308–1340.
- [BL] P. Bougerol and J. Lacroix, *Products of Random Matrices with Applications to Schrödinger Operators*, Birkhauser, Boston, 1985.
- [B] J. Bourgain, On the Furstenberg measure and density of states for the Anderson-Bernoulli model at small disorder, *J. Anal. Math.* **117** (2012), pp. 273–295.
- [BK] J. Bourgain, A. Klein, *Bounds on the density of states for Schrödinger operators*, *Invent. Math.* **194** (2013), no.1, pp. 41–72.
- [BrK] M. Brin, Yu. Kifer, Dynamics of Markov chains and stable manifolds for random diffeomorphisms, *Ergodic Theory and Dyn. Syst.*, **7** (1987), pp. 351–374.
- [BP] M. Brin, Ya. Pesin, Partially hyperbolic dynamical systems, *Izv.Acad.Nauk SSSR ser. mat.* **1** (1974), pp. 170–212.
- [BH] A. Brown, F. Hertz, Measure rigidity for random dynamics on surfaces and related skew products, *J. Amer. Math. Soc.* **30** (2017), pp. 1055–1132.
- [CK] M. Campanino, A. Klein, A supersymmetric transfer matrix and differentiability of the density of states in the one-dimensional Anderson model. *Comm. Math. Phys.* **104** (1986), no.2, pp. 227–241.
- [CS] W. Craig, B. Simon, Log Hölder continuity of the integrated density of states for stochastic Jacobi matrices, *Comm. Math. Phys.* **90** (1983), no.2, 207–218.

- [DG] D. Damanik, A. Gorodetski, Hölder continuity of the integrated density of states for the Fibonacci Hamiltonian. *Comm. Math. Phys.* **323** (2013), no.2, pp. 497–515.
- [Fu1] A. Furman, Random walks on groups and random transformations, *Handbook of dynamical systems*, vol. 1A, pp. 931–1014, North-Holland, Amsterdam, 2002.
- [Fu2] A. Furman, What is ... a stationary measure? *Notices Amer. Math. Soc.* **58** (2011), no. 9, pp. 1276–1277.
- [GK] Z. Gan, H. Krüger, Optimality of log Hölder continuity of the integrated density of states. *Math. Nachr.*, **284** (2011), no.14-15, 1919–1923.
- [GM] I. Gol'dsheid, G. Margulis, Lyapunov exponents of a product of random matrices. *Uspekhi Mat. Nauk*, **44** (1989), no.5, pp. 13–60.
- [GS] M. Goldstein, W. Schlag, Hölder continuity of the integrated density of states for quasi-periodic Schrödinger equations and averages of shifts of subharmonic functions. *Ann. of Math.* **2**, 154 (2001), no.1, 155–203.
- [GKM] A. Gorodetski, V. Kleptsyn, G. Monakov, Hölder regularity of stationary measures, arXiv:2209.12342
- [Gu] Y. Guivarc'h, Produits de matrices aleatoires et applications aux proprietes geometriques des sous-groupes du groupe lineaire, *Ergodic Theory Dynam. Systems* **10** (1990), pp. 483–512.
- [GR] Y. Guivarc'h, A. Raugi, Products of random matrices: convergence theorems, *Random matrices and their applications (Brunswick, Maine, 1984)*, pp. 31–54, *Contemp. Math.*, 50, Amer. Math. Soc., Providence, RI, 1986.
- [HV] E. Hart, B. Virág, Hölder Continuity of the Integrated Density of States in the One-Dimensional Anderson Model. *Commun. Math. Phys.* **355** (2017), pp. 839–863.
- [Kif1] Yu. Kifer, Random dynamics and its applications, *Proceedings of the International Congress of Mathematicians*, Vol. II (Berlin, 1998). Doc. Math. 1998, Extra Vol. II, pp. 809–818.
- [Kif2] Yu. Kifer, Ergodic theory of random transformations, *Progress in Probability and Statistics*, **10**, Birkhäuser Boston, Inc., Boston, MA, 1986. x+210 pp.
- [L] E. Le Page, Théorèmes limites pour les produits de matrices aléatoires, in: *Probability Measures on Groups*, H. Heyer, ed., Springer-Verlag, New York, 1982.
- [LQ] P. Liu, M. Qian, Smooth ergodic theory of random dynamical systems, *Lecture Notes in Mathematics*, **1606**, Springer-Verlag, Berlin, 1995, xii+221 pp.
- [M] D. Malicet, Random Walks on $\text{Homeo}(S^1)$, *Commun. Math. Phys.* **356** (2017), pp. 1083–1116.
- [MS] P. March, A. Sznitman, Some connections between excursion theory and the discrete Schrödinger equation with random potentials. *Probab. Theory Related Fields* **75** (1987), no.1, pp. 11–53.
- [MP] P. Mörters, Yu. Peres, Brownian motion. *Camb. Ser. Stat. Probab. Math.*, **30**, Cambridge University Press, Cambridge, 2010. xii+403 pp.
- [Mu] P. Munger, Frequency dependence of Hölder continuity for quasiperiodic Schrödinger operators. *J. Fractal Geom.* **6** (2019), no.1, pp. 53–65.
- [ST] B. Simon, M. Taylor, Harmonic analysis on $\text{SL}(2, \mathbb{R})$ and smoothness of the density of states in the one-dimensional Anderson model. *Comm. Math. Phys.* **101** (1985), no.1, pp. 1–19.
- [V] I. Veselić, Existence and regularity properties of the integrated density of states of random Schrödinger operators. *Lecture Notes in Math.*, **1917** Springer-Verlag, Berlin, 2008. x+142 pp.
- [Vi] C. Villani, Optimal Transport. Old and New, Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], **338**, Springer-Verlag, Berlin, 2009, xxii + 973 pp.

GRIGORII MONAKOV, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, IRVINE, CA 92697, USA

Email address: gmonakov@uci.edu