# CIRCULAR SYMMETRY-BREAKING AND TOPOLOGICAL NOETHER CURRENTS

### J MORAVA

ABSTRACT. We propose a toy model for the algebraic topology of bubbling as circular symmetry-breaking, in terms of Noether currents and cobordism of manifolds with circle actions free along boundaries. This leads to an interpretation of Planck's radiation law  $[9](\S2.4),[13]$  as expressing the loss of a Noether symmetry when a bubble

$$|[S^2//\mathbb{T}]| \to \mathrm{pt}$$

(analogous to blowing up or down in projective geometry) collapses; cf the von Kármán street of sparks left when a candle flame wisps out [15] .

**1** In these notes 'a space X with circle action' will usually mean the geometric realization  $|X/\mathbb{T}| := |[X/\mathbb{T}]|$  of the topological transformation groupoid defined by an action of the circle group  $\mathbb{T}$  on the space X. The collapse map

$$[X/\mathbb{T}] \to [\mathrm{pt}/\mathbb{T}]$$

thus defines a ring homomorphism

$$H^*_{\mathbb{T}}(\mathrm{pt}) \cong \mathbb{Z}[c] \to H^*_{\mathbb{T}}(X)$$

and therefore a natural characteristic class in the equivariant cohomology  $[3](\S 8)$  of  $[X/\mathbb{T}]$ , which I suggest calling, the (topological) Noether charge of the space (*ie* topological groupoid)  $[X/\mathbb{T}]$ , *ie* with circle action <sup>1</sup>. The universal example  $E\mathbb{T}$  then has charge  $c \in H^2(B\mathbb{T})$ .

Since Gauss, the Chern class c is understood as a measure of curvature. On the other hand, the integral Federer-de Rham current  $b \in H_2(B\mathbb{T},\mathbb{Z})$ is (using Quillen's conventions) the class  $[s^{-1}(0)]$  of the zero-locus of the canonical line bundle over  $\mathbb{C}P^{\infty}$ ; under cap product it represents a Lefschetz hyperplane intersection operator, a  $\delta$ -function supported by the codimension two cycle  $\mathbb{C}P^{n-1} \subset \mathbb{C}P^n$ ,  $n \to \infty$ .

Kronecker duality pairs these by integration, but both terms make sense as elements of the (torsion-free) Swan - Tate cohomology  $t_{\mathbb{T}}H\mathbb{Z} \cong \mathbb{Z}[c^{\pm 1}]$ , where they can be compared over  $\mathbb{Q}$ .

<sup>&</sup>lt;sup>1</sup>If  $(M, \omega) \ni X$  is a symplectic manifold w.c.a. there is a theory of moment maps  $M \to \text{Lie}_{\mathbb{T}} \dots$ 

#### J MORAVA

**2** The Swan-Tate cohomology  $t_{\mathbb{G}}E$  (of a complex-oriented equivariant  $E_{\infty}$  ring-spectrum E) is defined for any compact Lie group G, but when G is  $\mathbb{T}$ , as below, things simplify considerably, for  $t_{\mathbb{T}}H$  is then the periodic cyclic homology of the singular cochain algebra [1, 7, 11]. This account is concerned mostly with the structure of (the coefficient ring of)  $t_{\mathbb{T}}H\mathbb{Z}$ , but because MU = E is the (torsion-free) universal example, it is convenient to start there (or skip to §2.3).

**2.1** If [X] is a compact cx-oriented 2*d*-dimensional smooth manifold with circle action, perhaps with boundary, such that the restriction of the circle action to the boundary is free, *ie* making the boundary quotient a 2(d-1) manifold endowed with a canonical circle bundle, it seems reasonable to call  $[X/\mathbb{T}]$  a manifold with meromorphic circle action Then

$$t^*_{\mathbb{T}}MU^* \ni [X/\mathbb{T}] \mapsto \partial_{\mathbb{T}}[X/\mathbb{T}] := [(\partial X)/\mathbb{T} \to B\mathbb{T}] \in MU_{*-2}B\mathbb{T} ,$$

defines the boundary operator in the diagram<sup>2</sup>

where  $b \in \operatorname{Hom}_{FG}(F_E, \mathbb{G}_m)$ , *ie* satisfying

$$b(T) = 1 + \sum_{k>1} b_k T^k$$
,  $b(T_0 +_E T_1) = b(T_0) \cdot b(T_1)$ ,

is the Ravenel-Wilson-Katz Cartier generating character for the formal group  $F_E$  defined by the cx orientation of E.

**2.2** The product of two such manifolds, *ie* with circle actions free along the boundary, is a manifold with  $\mathbb{T} \times \mathbb{T}$  structure; restricting along the diagonal  $\mathbb{T} \to \mathbb{T} \times \mathbb{T}$  defines a manifold  $[(X \times Y)/\mathbb{T}]$  with circle action, again free along the boundary

$$\partial(X_0 \times X_1) = X_0 \times \partial X_1 \cup_{\partial X_0 \times \partial X_1} \partial X_0 \times X_1 ,$$

defining a ring structure on  $t^*_{\mathbb{T}}MU$ . One wonders about Massey products.

## **2.3 Examples** $E = H\mathbb{Z} \& K$

If G is a finite group,  $t_G^*K \cong R_{\mathbb{C}}(G) \otimes \mathbb{Q}$ , whereas  $t_{\mathbb{T}}^*K \cong (1-q)^{-1}K_{\mathbb{T}}$ ; where  $K_{\mathbb{T}} \cong R_{\mathbb{C}}(\mathbb{T})$  is Atiyah - Segal T-equivariant K-theory (of a point), and  $q: \mathbb{T} \subset \mathbb{C}^{\times}$  sends  $q \to e^{i\theta}$ .

Recall from [3](Prop 3.4) that localization kills (c)-torsion, so

 $t_{\mathbb{T}}H^*[M/\mathbb{T}] \cong H\mathbb{Z}^*(\operatorname{Fix}_{\mathbb{T}}M) \otimes t_{\mathbb{T}}H\mathbb{Z}$ 

 $<sup>^{2}</sup>$ A preliminary account of such Hopf algebra biextensions can be found in the appendix to 11 of the 1959/60 Seminaire Cartan-Moore published in an adjacent timeline, but see more recently [4]

*ie* with the grading scrambled by weights coming from normal bundles [3](§7); see [5] for  $t_{\mathbb{Z}_2}H\mathbb{F}_2$ . Re SU(2), recall  $BSU(2) \cong \mathbb{H}P^{\infty}$  is not an *H*-space.

**2.4** When  $E = H\mathbb{Z}$  is Eilenberg - Mac Lane cohomology, we now have an exact sequence

0

$$0 \longrightarrow H^*B\mathbb{T} = \mathbb{Z}[c] \longrightarrow t_{\mathbb{T}}H\mathbb{Z} = \mathbb{Z}[c^{\pm 1}] \xrightarrow{\partial_{\mathbb{T}}} H_{*-2}B\mathbb{T} = \Gamma[b_*] \longrightarrow$$

$$0 \longrightarrow c^3 \longrightarrow c^3 \longrightarrow c^3 \longrightarrow 0$$

$$0 \longrightarrow c^2 \longrightarrow c^2 \longrightarrow c^2 \longrightarrow 0$$

$$0 \longrightarrow c \longrightarrow c \longrightarrow c \longrightarrow c \longrightarrow 0$$

$$0 \longrightarrow 1 \longrightarrow 1 \longrightarrow c^{-1} \longrightarrow 0$$

$$0 \longrightarrow 0 \longrightarrow c^{-1} \longrightarrow 1$$

$$0 \longrightarrow 0 \longrightarrow c^{-2} \longrightarrow b_1$$

$$0 \longrightarrow 0 \longrightarrow c^{-3} \longrightarrow b_2 \dots$$

in which  $\partial^0_{\mathbb{T}} : (\{0\} \in \mathbb{C}) \to (\mathbb{C} - \{0\}) = \mathbb{C}^{\times} \in \text{free } \mathbb{T}\text{-spaces.}$  The boundary operator thus defines a Rota-Baxter structure: invertible in negative degree and zero in positive degree.

Here  $\Gamma_*[b] = \mathbb{Z}[b_i \mid b_i b_k = (i, k) b_{I+k}]$  is the free divided power algebra on one generator; thus  $b_k \mapsto b^k/k! \in \Gamma_*(b) \otimes \mathbb{Q} \cong \mathbb{Q}[b]$ . The grading forces this sequence to split, as

$$t_{\mathbb{T}}^*H = \mathbb{Z}[c] \oplus c^{-1}\mathbb{Z}[c^{-1}]$$

with  $\partial_{\mathbb{T}} c^{-k} = b_{k-1}$ , defining a representation of a Heisenberg group [8](§3).

It is useful to think of  $c^{-1}$  as the two-disk  $D^2$  with its usual circle action; then

 $\partial_{\mathbb{T}} c^0 = 0, \partial_{\mathbb{T}} c^{-1} = 1, \partial_{\mathbb{T}} c^{-2} = [S^3/\mathbb{T} \cong S^2 \to \mathbb{C}P^{\infty} \cong B\mathbb{T}] = b \in MU_2B\mathbb{T}$ , making  $[D^2/\mathbb{T}]$  to  $[S^2/\mathbb{T}]$  much as open are to closed strings.

# §3 Some applications

**3.1** The degrading functor  $V_* \mapsto V_* \otimes \mathbb{Z}((T))|_0 := V||T|| \in \mathbb{Z}((T))$  – Mod is defined on  $\mathbb{Z}$  - graded modules by

$$\operatorname{Hom}^{d}(V,W) \ni \phi \mapsto [\Sigma_{i>-\infty} v_{i}T^{i} \to T^{d}\Sigma_{i>-\infty} \phi(v_{i})T^{i}], \ v_{i} \in V_{i}$$

so we have from  $[5](\S I)$  that

$$b = -T^{-1}\log(1 - c^{-1}T), \ c = -b^{-1}B^{-}(-bT) \in (t_{\mathbb{T}}H\mathbb{Q})||T||$$

with Bernoulli operator  $B^{-}(D) = \frac{D}{e^{D}-1}$ , because

$$(1 - c^{-1}T)^{-1} = \exp(bT)$$

(and more generally in the Lazard ring.) Note that  $B_k/k = h_k$  are the cumulants of the uniform distribution.

These relations interpret b as a de Rham current corresponding to the Noether charge c, *ie* as something like its supporting cycle; but they make sense only after rationalizing. In particular, b and c have opposed natural variances. In this formalism T is a kind of place-holder for a scaling action of Adams' operations in  $\mathbb{N}^{\times}$ , interpreted as an asymptotic expansion [10].

**3.2** The topological Mellin transform identifies Spec  $t_{\mathbb{T}}H\mathbb{Z}$  as a localization  $\mathbb{G}_m = \operatorname{Spec} \mathbb{Z}[c^{\pm 1}]$  of the ungraded cohomology  $H^*B\mathbb{T} \cong H^*\mathbb{C}P^\infty$ :

**Proposition** The differential form [9] (§2.1 Proposition)

 $d(bT) = -d\log(1 - c^{-1}T) \in \Omega^1_{\mathbb{Z}}(t_{\mathbb{T}}H\mathbb{Z})$ 

maps by  $c^{-1}T \rightarrow e^{\varepsilon}$  to the tempered distribution-valued one-form

$$d(bT) = \frac{d(c^{-1}T)}{1 - c^{-1}T} = \frac{e^{\varepsilon}d\varepsilon}{1 - e^{\varepsilon}} = \operatorname{li}_0(\varepsilon) \cdot d\varepsilon = d\operatorname{li}_1(\varepsilon) \in \mathcal{S}'(\mathbb{R}) \cdot d\varepsilon \cong \Omega^1_{\operatorname{temp}}(\mathbb{R})$$

where

$$li_0(x) = -(1 - e^{-x})^{-1}, \ li_1(x) = -\log|1 - e^x|$$

**3.3** We interpret db as Haar measure, translation-invariant with respect to  $+_{\mathbb{G}_m}$ , pulled back along the exponential map to the Lie algebra of  $\mathbb{G}_m$ , defining a moment map  $b : \mathbb{G}_m \to \text{Lie}_{\mathbb{T}}$  (as in [3](§7)) for the circle. We can regard b as defining a transcendental extension of  $\mathbb{C}(c)$ .

The resulting linear operator [14]

$$\pi_!^{\mathbb{G}_m}: \Omega^1_{\mathbb{Z}}(t_{\mathbb{T}}H\mathbb{Z}) \to \mathcal{S}'(\mathbb{R}) \cdot d\varepsilon$$

sends one-forms to de Rham Harvey-Lawson one-currents on the  $\varepsilon$ -line, defining a pairing

$$\mathcal{S}(\mathbb{R}) \times \mathbb{Z}[c^{\pm 1}] \cdot dc \to \mathbb{R} \cdot d\varepsilon$$

in which  $\varepsilon = -\log c$ ,  $d\varepsilon = -c^{-1}dc$ , suggesting  $\varepsilon$  as a measure of topological entropy or information loss.

# 4 Afterthoughts

4.1 Convolutions of tempered distributions are not necessarily tempered, but the holomorphic family

$$\gamma^s_+ := \frac{x_+^{s-1}}{\Gamma(s)} \in \mathcal{S}'(\mathbb{R})\{s\}$$

of Gel'fand-Ŝilov tempered distributions  $(x \in \mathbb{R})$  represents the group of fractional differentiation operations  $\partial^{-s}$  on functions such as  $B^{-}(x) \sim (e^{x} - 1)^{-1} = \lim_{i \to 0} (x)$ .

Prop 3 of [6] asserts that the function

$$li_{s}(x) = \gamma_{+}^{s} * li_{0} = Li_{s}(e^{x}) = 1 + \sum_{n \ge 1} \frac{e^{nx}}{n^{s}} \in C^{\infty}(\mathbb{R}_{<0})\{s\}, \ li_{s}|_{x=0} \mapsto \zeta(s)$$

(as defined on the negative real line) extends to a holomorphic family  $s \mapsto li_s \equiv -\pi \cot \pi s \cdot \gamma^s_+ \in \mathcal{S}'(\mathbb{R})\{s\}$  of tempered distributions on the whole real line. Consequently [9](§2.2)

The divided moments

$$\gamma^s_+(\varepsilon) * db = (\gamma^s_+ * dli_1)(\varepsilon)$$

are tempered distributions.

**4.2** Note that the distribution-valued 'measure'  $\gamma^1_+(\varepsilon) * db(\varepsilon)$  on  $\mathbb{R}$  has no 'value' at x = 0, though the Taylor series of  $\lim_{s \to \infty} (x)$  at s = 1 is

$$(1-s)^{-1} + \log \left| \frac{x}{e^x - 1} \right| (1-s)^0 + \sum_{k \ge 1} t_k \cdot (1-s)^k \dots$$

with tempered distributions  $t_*$ .

Nevertheless, these generalized moments may be useful in statistical mechanics. The Stefan/Boltzmann asymptotic expansion  $[9](\S2.4),[12]$ 

$$T^{-4}\boldsymbol{\rho} \sim 12h\kappa^4 \cdot \gamma_+^4 * dli_1$$

(for the radiance<sup>3</sup>  $\rho$  of an incandescent body in terms of its temperature) is how we understand the colors of the stars.

 $<sup>{}^3\</sup>kappa = kh^{-1} \sim 20.836 \times 10^9$  Hertz per Kelvin

#### J MORAVA

#### References

- A Adem, R L Cohen, W G Dwyer, Generalized Tate homology, homotopy fixed points and the transfer. Algebraic topology (Evanston 1988) 1–13, Contemp Math 96, AMS 1989
- M Atiyah, The moment map in symplectic geometry. Global Riemannian geometry (Durham, 1983) 43 — 51, Ellis Horwood Ser. Math. Appl., Horwood, Chichester, 1984.
- 3. \_\_\_\_, R Bott, The moment map and equivariant cohomology. Topology 23 (1984) 1  $_{\mbox{28}}$
- 4. Ching-Li Chai, Tate-linear structures on formal varieties, https://www2.math.upenn.edu/~chai/papers\_pdf/TL\_survey\_v2.pdf §3.3
- 5. W Dwyer, C Wilkerson, Smith theory revisited, Ann Math 127 (1988) 191 198
- 6. Charles L Epstein, J Morava, Tempering the polylogarithm, https://arxiv.org/abs/math/0611240
- 7. E Getzler, J D S Jones, S Petrack , Differential forms on loop spaces and the cyclic bar complex, Topology 30, 339 371 (1991)
- J Morava, Heisenberg groups in algebraic topology, in Topology, Geometry and Quantum Field Theory, 235 - 246, LMS Lecture Notes 308, Cambridge 2004 https://arxiv.org/abs/math/0305250 §3
- 9. ——-Periods for topological circle actions, https://arxiv.org/abs/2301.05772
- 10. —, Swan-Tate cohomology of equivariant circle actions, https://arxiv.org/abs/2403.19714
- 11. D Quillen, Algebra cochains and cyclic cohomology. IHES Publ. Math. 68 (1988) 139
   174 http://www.numdam.org/item/PMIHES\_1988\_68\_139\_0.pdf
- 12. https://en.wikipedia.org/wiki/Stefan%E2%80%93Boltzmann\_law
- 13. https://en.wikipedia.org/wiki/Planck%27s\_law
- 14. https://en.wikipedia.org/wiki/Sokhotski%E2%80%93Plemelj\_theorem
- 15. https://en.wikipedia.org/wiki/Gaston\_Bachelard

Department of Mathematics, The Johns Hopkins University, Baltimore, Maryland

Email address: jmorava1@jhu.edu

 $\mathbf{6}$