

SOME REMARKS ON REAL MINIMAL NILPOTENT ORBITS AND SYMMETRIC PAIRS

TAKAYUKI OKUDA

ABSTRACT. For a non-compact simple Lie algebra \mathfrak{g} over \mathbb{R} , we denote by $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}}$ the unique complex nilpotent orbit in $\mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C}$ containing all minimal real nilpotent orbits in \mathfrak{g} . In this paper, we give a complete classification of symmetric pairs $(\mathfrak{g}, \mathfrak{h})$ such that $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}} \cap \mathfrak{g}^d = \emptyset$, where \mathfrak{g}^d denotes the dual Lie algebra of $(\mathfrak{g}, \mathfrak{h})$. Furthermore, for symmetric pairs (G, H) with real simple Lie group G , we apply our classification to theorems given by T. Kobayashi [J. Lie Theory (2023)], and study bounded multiplicity properties of restrictions on H of infinite-dimensional irreducible G -representations with minimum Gelfand–Kirillov dimension.

1. INTRODUCTION

Let \mathfrak{g} be a non-compact simple Lie algebra over \mathbb{R} . We write $\mathcal{N}(\mathfrak{g})$ for the nilpotent cone in \mathfrak{g} , and $\mathcal{N}(\mathfrak{g})/G$ for the set of real nilpotent (adjoint) orbits in \mathfrak{g} . A non-zero real nilpotent orbit \mathcal{O} in \mathfrak{g} with minimum dimension is called *minimal*. Let us put $\mathcal{N}_{\min}(\mathfrak{g}) \subset \mathcal{N}(\mathfrak{g})$ the union of all minimal real nilpotent orbits in \mathfrak{g} . The complexification of \mathfrak{g} is denoted by $\mathfrak{g}_{\mathbb{C}} := \mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C} = \mathfrak{g} + \sqrt{-1}\mathfrak{g}$. Then as proved in [12, Theorems 1.1 and 1.3], there exists uniquely a complex nilpotent (adjoint) orbit $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}}$ in $\mathfrak{g}_{\mathbb{C}}$ containing $\mathcal{N}_{\min}(\mathfrak{g})$. We denote by $m(\mathfrak{g})$ half of the complex dimension of $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}}$. Note that the orbit $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}}$ is minimal as a complex nilpotent orbit in most cases but not for some real simple \mathfrak{g} (see Section 2 for more details).

Let us also fix an involutive automorphism σ on \mathfrak{g} , and put $\mathfrak{h} := \{X \in \mathfrak{g} \mid \sigma(X) = X\}$ and $\mathfrak{q} := \{X \in \mathfrak{g} \mid \sigma(X) = -X\}$. Then $(\mathfrak{g}, \mathfrak{h})$ is a symmetric pair. Take a Cartan involution θ on \mathfrak{g} commuting with σ , and write $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ for the Cartan decomposition. Then as in [13], the dual Lie algebra \mathfrak{g}^d of $(\mathfrak{g}, \mathfrak{h})$ is defined by

$$\mathfrak{g}^d := (\mathfrak{k} \cap \mathfrak{h}) + \sqrt{-1}(\mathfrak{k} \cap \mathfrak{q}) + \sqrt{-1}(\mathfrak{p} \cap \mathfrak{h}) + (\mathfrak{p} \cap \mathfrak{q}).$$

Note that \mathfrak{g}^d is a real form of $\mathfrak{g}_{\mathbb{C}}$, and up to inner-automorphisms on $\mathfrak{g}_{\mathbb{C}}$, the real form \mathfrak{g}^d of $\mathfrak{g}_{\mathbb{C}}$ depends only on (\mathfrak{g}, σ) , but not on θ .

2020 *Mathematics Subject Classification.* Primary 22E46; Secondary 22E45, 53C35, 32M15, 53C15, 57S30.

The third author is supported by JSPS Grants-in-Aid for Scientific Research JP20K03589, JP20K14310, JP22H0112 and JP24K06714.

In this paper, we address the following question:

Question 1.1. *Whether $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}}$ meets \mathfrak{g}^d or not?*

We find that the orbit $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}}$ meets \mathfrak{g}^d in most cases. More precisely, our main result gives a complete classification of symmetric pairs $(\mathfrak{g}, \mathfrak{h})$ with $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}} \cap \mathfrak{g}^d = \emptyset$ (see Theorems 3.2 and 3.4 in Section 3 for the classification).

Our work is motivated by branching problems of “small” infinite-dimensional representations of simple Lie groups as below: let G be a non-compact connected simple Lie group with Lie algebra \mathfrak{g} . The set of irreducible objects in the category of smooth admissible representations of G of finite length with moderate growth is denoted by $\text{Irr}(G)$. Then for each infinite-dimensional irreducible object $\Pi \in \text{Irr}(G)$, the Gelfand-Kirillov dimension $\text{DIM}(\Pi)$ of Π satisfies

$$m(\mathfrak{g}) \leq \text{DIM}(\Pi).$$

By combining our classification with T. Kobayashi’s results in [8, 9], we have the following theorem, mentioned in [9, Remark 5.7]:

Theorem 1.2. *Let G be a non-compact connected simple Lie group, and $\Pi \in \text{Irr}(G)$ with $m(\mathfrak{g}) = \text{DIM}(\Pi)$. Then for any symmetric pair (G, H) , the restriction $\Pi|_H$ has the bounded multiplicity property*

$$\sup_{\pi \in \text{Irr}(H)} [\Pi|_H : \pi] < \infty,$$

where $[\Pi|_H : \pi]$ denotes the multiplicity of π in $\Pi|_H$ given by $\dim_{\mathbb{C}} \text{Hom}_H(\Pi|_H, \pi)$.

Question 1.1 is also related to the following topological problem: let G/H^a be the symmetric space corresponding to the associated symmetric pair of (G, H) , in the sense of [2], and let $SL_2(\mathbb{R})$ or $PSL_2(\mathbb{R})$ be a subgroup of G corresponding to a real minimal nilpotent orbit by the Jacobson–Morozov theorem (see Section 5 for the details). In Theorem 5.1, we show that $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}} \cap \mathfrak{g}^d = \emptyset$ if and only if the action of the subgroup on G/H^a is proper.

2. PRELIMINARIES FOR MINIMAL REAL NILPOTENT ORBITS

Let \mathfrak{g} be a non-compact simple Lie algebra over \mathbb{R} . In this section, we recall some facts and give some observations for minimal real nilpotent orbits in \mathfrak{g} .

As in Section 1, we denote by $\mathcal{N}(\mathfrak{g})$ and $\mathcal{N}(\mathfrak{g})/G$ the nilpotent cone and the set of nilpotent (adjoint) orbits in \mathfrak{g} . With respect to the closure ordering, $\mathcal{N}(\mathfrak{g})/G$ is a partially ordered set, and the zero orbit is the minimum in $\mathcal{N}(\mathfrak{g})/G$. A non-zero real nilpotent orbit \mathcal{O} in \mathfrak{g} is called minimal if its dimension is minimum, or equivalently, it is minimal in the partially ordered set $(\mathcal{N}(\mathfrak{g})/G) \setminus \{0\}$ (see [12, Theorem 1.3]). We write $\mathcal{N}_{\min}(\mathfrak{g})$ for the union of all minimal real nilpotent orbits in \mathfrak{g} .

For the number of minimal real nilpotent orbits, the following is known:

Fact 2.1 (See e.g. [12]). *Let $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ be a Cartan decomposition of \mathfrak{g} .*

- (i) *If \mathfrak{g} is absolutely-simple and $(\mathfrak{g}, \mathfrak{k})$ is of Hermitian-type, then \mathfrak{g} admits two minimal real nilpotent orbits $\mathcal{O}_{\min,1}$ and $\mathcal{O}_{\min,2}$ with $\mathcal{O}_{\min,2} = -\mathcal{O}_{\min,1}$.*
- (ii) *If \mathfrak{g} is absolutely-simple but $(\mathfrak{g}, \mathfrak{k})$ is not of Hermitian-type, then the minimal real nilpotent orbit \mathcal{O}_{\min} in \mathfrak{g} is unique.*
- (iii) *If \mathfrak{g} is not absolutely-simple, then the minimal real nilpotent orbit \mathcal{O}_{\min} in \mathfrak{g} is unique. Furthermore, in this situation, \mathfrak{g} admits two complex structures $\{\pm J\}$, and \mathcal{O}_{\min} can be considered as the minimal complex nilpotent orbit in the complex simple Lie algebra $(\mathfrak{g}, \pm J)$.*

We denote by $\mathfrak{g}_{\mathbb{C}} = \mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C} = \mathfrak{g} + \sqrt{-1}\mathfrak{g}$ the complexification of \mathfrak{g} . Then there uniquely exists a complex nilpotent adjoint orbit $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}}$ in $\mathfrak{g}_{\mathbb{C}}$ with $\mathcal{N}_{\min}(\mathfrak{g}) \subset \mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}}$ (see [12, Theorems 1.1 and 1.3]). We define $m(\mathfrak{g}) \in \mathbb{Z}_{\geq 1}$ as half of the complex dimension of $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}}$. Note that for each real minimal nilpotent orbit \mathcal{O}_{\min} in \mathfrak{g} , we have $m(\mathfrak{g}) = \frac{1}{2} \dim_{\mathbb{R}} \mathcal{O}_{\min}$.

We also recall that for each complex simple Lie algebra $\mathfrak{l}_{\mathbb{C}}$, the minimal complex nilpotent (adjoint) orbit $\mathcal{O}_{\min}^{\mathbb{C}}$ in $\mathfrak{l}_{\mathbb{C}}$ is defined as the unique non-zero complex nilpotent orbit in $\mathfrak{l}_{\mathbb{C}}$ which is contained in the closure of any non-zero complex nilpotent orbit in $\mathfrak{l}_{\mathbb{C}}$. Note that $\dim_{\mathbb{C}} \mathcal{O}_{\min}^{\mathbb{C}} \leq \dim_{\mathbb{C}} \mathcal{O}^{\mathbb{C}}$ holds for any non-zero complex nilpotent orbit $\mathcal{O}^{\mathbb{C}}$ and the equality holds only if $\mathcal{O}^{\mathbb{C}} = \mathcal{O}_{\min}^{\mathbb{C}}$. Throughout this paper, we define $n(\mathfrak{l}_{\mathbb{C}})$ as half of the complex dimension of $\mathcal{O}_{\min}^{\mathbb{C}}$. Table 1 gives the formula of $n(\mathfrak{l}_{\mathbb{C}})$ (see e.g. [4, Lemma 4.3.5 and Chapter 8.4]). Furthermore, for a complex semisimple Lie algebra $\mathfrak{l}'_{\mathbb{C}}$, we also put $n(\mathfrak{l}'_{\mathbb{C}}) := \sum_i n(\mathfrak{l}^i_{\mathbb{C}})$, where $\mathfrak{l}'_{\mathbb{C}} = \bigoplus_i \mathfrak{l}^i_{\mathbb{C}}$ denotes the complex simple ideal decomposition of $\mathfrak{l}'_{\mathbb{C}}$.

TABLE 1. List of $n(\mathfrak{l}_{\mathbb{C}})$ for complex simple $\mathfrak{l}_{\mathbb{C}}$

$\mathfrak{l}_{\mathbb{C}}$	$\mathfrak{sl}_n(\mathbb{C})$	$\mathfrak{so}_n(\mathbb{C})$	$\mathfrak{sp}_n(\mathbb{C})$	$\mathfrak{e}_6^{\mathbb{C}}$	$\mathfrak{e}_7^{\mathbb{C}}$	$\mathfrak{e}_8^{\mathbb{C}}$	$\mathfrak{f}_4^{\mathbb{C}}$	$\mathfrak{g}_2^{\mathbb{C}}$
$n(\mathfrak{l}_{\mathbb{C}})$	$n - 1$	$n - 3$	n	11	17	29	8	3

If our real Lie algebra \mathfrak{g} is absolutely-simple, then its complexification $\mathfrak{g}_{\mathbb{C}}$ is complex simple and the closure of $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}}$ contains the minimal complex nilpotent orbit $\mathcal{O}_{\min}^{\mathbb{C}}$ in $\mathfrak{g}_{\mathbb{C}}$, and thus $n(\mathfrak{g}_{\mathbb{C}}) \leq m(\mathfrak{g})$. Furthermore, if \mathfrak{g} is not absolutely simple, for the complex structures $\{\pm J\}$ on \mathfrak{g} , we have complex simple Lie algebras (\mathfrak{g}, J) and $(\mathfrak{g}, -J)$. The complexification $\mathfrak{g}_{\mathbb{C}}$ of \mathfrak{g} can be identified with $(\mathfrak{g}, J) \oplus (\mathfrak{g}, -J)$ by

$$\mathfrak{g}_{\mathbb{C}} \rightarrow (\mathfrak{g}, J) \oplus (\mathfrak{g}, -J), \quad X + \sqrt{-1}Y \mapsto (X + JY, X - JY).$$

Let us denote by \mathcal{O}_{\min} the unique minimal nilpotent orbit in \mathfrak{g} . Then $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}}$ can be identified with $\mathcal{O}_{\min} \times \mathcal{O}_{\min}$. In particular, we obtain

$$n(\mathfrak{g}_{\mathbb{C}}) = n((\mathfrak{g}, J)) + n((\mathfrak{g}, -J)) = m(\mathfrak{g}).$$

The situation $n(\mathfrak{g}_{\mathbb{C}}) < m(\mathfrak{g})$ occurs only if \mathfrak{g} is absolutely-simple, and classified as below:

Fact 2.2 (See also [3], [8, Proposition 30], [10, Corollary 5.9], [12, Proposition 4.1]). *The following conditions on a non-compact absolutely-simple Lie algebra \mathfrak{g} are equivalent:*

- (i) $n(\mathfrak{g}_{\mathbb{C}}) < m(\mathfrak{g})$.
- (ii) $\mathcal{O}_{\min}^{\mathbb{C}} \cap \mathfrak{g} = \emptyset$.
- (iii) $\mathcal{O}_{\min}^{\mathbb{C}} \neq \mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}}$.
- (iv) \mathfrak{g} is isomorphic to one of the Lie algebras in Table 2.

Furthermore, if \mathfrak{g} satisfies the equivalent conditions above, then the symmetric pair $(\mathfrak{g}, \mathfrak{k})$ is not of Hermitian-type, and \mathfrak{g} has a unique minimal real nilpotent orbit.

TABLE 2. List of \mathfrak{g} with $n(\mathfrak{g}_{\mathbb{C}}) < m(\mathfrak{g})$

\mathfrak{g}	$m(\mathfrak{g})$	$(\mathfrak{g}_{\mathbb{C}}, \mathfrak{k}_{\mathbb{C}})$	$n(\mathfrak{g}_{\mathbb{C}})$
\mathfrak{su}_{2n}^* ($n \geq 2$)	$4n - 4$	$(\mathfrak{sl}_{2n}(\mathbb{C}), \mathfrak{sp}_n(\mathbb{C}))$	$2n - 1$
$\mathfrak{so}_{n-1,1}$ ($n \geq 5$)	$n - 2$	$(\mathfrak{so}_n(\mathbb{C}), \mathfrak{so}_{n-1}(\mathbb{C}))$	$n - 3$
$\mathfrak{sp}_{m,n}$ ($m, n \geq 1$)	$2(m + n) - 1$	$(\mathfrak{sp}_{m+n}(\mathbb{C}), \mathfrak{sp}_m(\mathbb{C}) \oplus \mathfrak{sp}_n(\mathbb{C}))$	$m + n$
$\mathfrak{e}_{6(-26)}$	16	$(\mathfrak{e}_6^{\mathbb{C}}, \mathfrak{f}_4^{\mathbb{C}})$	11
$\mathfrak{f}_{4(-20)}$	11	$(\mathfrak{f}_4^{\mathbb{C}}, \mathfrak{so}_9(\mathbb{C}))$	8

The following observation will be applied in the next section:

Proposition 2.3. *Suppose that $n(\mathfrak{g}_{\mathbb{C}}) < m(\mathfrak{g})$. Let \mathfrak{g}' be a non-compact real form of $\mathfrak{g}_{\mathbb{C}}$. Then $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}} \cap \mathfrak{g}' = \emptyset$ if and only if $(\mathfrak{g}, \mathfrak{g}')$ is isomorphic to $(\mathfrak{su}_{2n}^*, \mathfrak{su}_{2n-1,1})$ ($n \geq 2$) or $(\mathfrak{so}_{2n-1,1}, \mathfrak{so}_{2n}^*)$ ($n \geq 4$).*

Proof of Proposition 2.3. As stated in [12, Section 4.1] or [11, Proposition 7.8 and Theorem 7.10], a complex nilpotent orbit $\mathcal{O}^{\mathbb{C}}$ in $\mathfrak{g}_{\mathbb{C}}$ meets a fixed real form \mathfrak{g}' if and only if the weighted Dynkin diagram of $\mathcal{O}^{\mathbb{C}}$ satisfies the matching condition to the Satake diagram $S_{\mathfrak{g}'}$ of \mathfrak{g}' . For the cases where $n(\mathfrak{g}_{\mathbb{C}}) < m(\mathfrak{g})$, or equivalently, $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}} \neq \mathcal{O}_{\min}^{\mathbb{C}}$, the list of weighted Dynkin diagrams of $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}}$ can be found in [12, Table 1]. By comparing it with the list of Satake diagrams of absolutely simple Lie algebras (see e.g. [1] or [5, Chapter X]), one can easily obtain the claim in Proposition 2.3. \square

3. CLASSIFICATIONS

In this section, we classify symmetric pairs $(\mathfrak{g}, \mathfrak{h})$ with $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}} \cap \mathfrak{g}^d = \emptyset$.

3.1. In the case where \mathfrak{g} is absolutely-simple. Let \mathfrak{g} be a non-compact absolutely simple Lie algebra over \mathbb{R} , and $(\mathfrak{g}, \mathfrak{h})$ a symmetric pair.

One of the main results of this paper is given below:

Theorem 3.1. *Suppose that $n(\mathfrak{g}_{\mathbb{C}}) < m(\mathfrak{g})$. Then $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}} \cap \mathfrak{g}^d \neq \emptyset$.*

Proof of Theorem 3.1. Let $(\mathfrak{g}, \mathfrak{h})$ be a symmetric pair with absolutely simple \mathfrak{g} . By Proposition 2.3, we only need to check that the pair $(\mathfrak{g}, \mathfrak{g}^d)$ is not isomorphic to $(\mathfrak{su}_{2n}^*, \mathfrak{su}_{2n-1,1})$ ($n \geq 2$) nor $(\mathfrak{so}_{2n-1,1}, \mathfrak{so}_{2n}^*)$ ($n \geq 4$). By the tables in [13, Section 1], one can see the following:

- If $\mathfrak{g} \simeq \mathfrak{su}_{2n}^*$ ($n \geq 2$), then \mathfrak{g}^d is isomorphic to \mathfrak{su}_{2n}^* , $\mathfrak{su}_{2n-2j,2j}$, $\mathfrak{sl}_{2n}(\mathbb{R})$ or $\mathfrak{su}_{n,n}$.
- If $\mathfrak{g}^d \simeq \mathfrak{so}_{2n}^*$ ($n \geq 4$), then \mathfrak{g} is isomorphic to \mathfrak{so}_{2n}^* , $\mathfrak{so}_{n,n}$ or $\mathfrak{so}_{2n-2j,2j}$.

These complete the proof. \square

By combining Theorem 3.1 with Fact 2.2 and tables of dual pairs of symmetric pairs in [13], we have the following classification result:

Theorem 3.2. *Let $(\mathfrak{g}, \mathfrak{h})$ be a symmetric pair such that \mathfrak{g} is absolutely-simple. Then the five conditions on $(\mathfrak{g}, \mathfrak{h})$ below are equivalent:*

- (i) $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}} \cap \mathfrak{g}^d = \emptyset$.
- (ii) $m(\mathfrak{g}) = n(\mathfrak{g}_{\mathbb{C}}) < m(\mathfrak{g}^d)$.
- (iii) $\mathcal{O}_{\min}^{\mathbb{C}} = \mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}}$ but $\mathcal{O}_{\min}^{\mathbb{C}} \neq \mathcal{O}_{\min, \mathfrak{g}^d}^{\mathbb{C}}$.
- (iv) The simple Lie algebra \mathfrak{g}^d can be found in Table 2 but \mathfrak{g} can not.
- (v) The symmetric pair $(\mathfrak{g}, \mathfrak{h})$ is isomorphic to one of the symmetric pairs in Table 3.

TABLE 3. List of $(\mathfrak{g}, \mathfrak{h})$ with absolutely simple \mathfrak{g} such that $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}} \cap \mathfrak{g}^d = \emptyset$

$(\mathfrak{g}, \mathfrak{h})$	\mathfrak{g}^d	$\mathfrak{h}^a = \mathfrak{g} \cap \mathfrak{g}^d$
$(\mathfrak{sl}_{2n}(\mathbb{R}), \mathfrak{sp}_n(\mathbb{R}))$ ($n \geq 2$)	\mathfrak{su}_{2n}^*	$\mathfrak{sl}_n(\mathbb{C}) \oplus \mathfrak{so}_2$
$(\mathfrak{su}_{2n-2j,2j}, \mathfrak{sp}_{n-j,j})$ ($n \geq 2$)	\mathfrak{su}_{2n}^*	$\mathfrak{sp}_{n-j,j}$
$(\mathfrak{su}_{n,n}, \mathfrak{sp}_n(\mathbb{R}))$ ($n \geq 2$)	\mathfrak{su}_{2n}^*	\mathfrak{so}_{2n}^*
$(\mathfrak{so}_{m,n}, \mathfrak{so}_{m-1,n})$ ($m, n \geq 2$ with $(m, n) \neq (2, 2)$)	$\mathfrak{so}_{m+n-1,1}$	$\mathfrak{so}_{1,n} \oplus \mathfrak{so}_{m-1}$
$(\mathfrak{sp}_n(\mathbb{R}), \mathfrak{sp}_{n-j}(\mathbb{R}) \oplus \mathfrak{sp}_j(\mathbb{R}))$ ($n \geq 2, n-1 \geq j \geq 1$)	$\mathfrak{sp}_{n-j,j}$	$\mathfrak{su}_{n-j,j} \oplus \mathfrak{so}_2$
$(\mathfrak{sp}_{2n}(\mathbb{R}), \mathfrak{sp}_n(\mathbb{C}))$ ($n \geq 1$)	$\mathfrak{sp}_{n,n}$	$\mathfrak{sp}_n(\mathbb{C})$
$(\mathfrak{e}_{6(6)}, \mathfrak{f}_{4(4)})$	$\mathfrak{e}_{6(-26)}$	$\mathfrak{su}_6^* \oplus \mathfrak{su}_2$
$(\mathfrak{e}_{6(2)}, \mathfrak{f}_{4(4)})$	$\mathfrak{e}_{6(-26)}$	$\mathfrak{sp}_{3,1}$
$(\mathfrak{e}_{6(-14)}, \mathfrak{f}_{4(-20)})$	$\mathfrak{e}_{6(-26)}$	$\mathfrak{f}_{4(-20)}$
$(\mathfrak{f}_{4(4)}, \mathfrak{so}_{5,4})$	$\mathfrak{f}_{4(-20)}$	$\mathfrak{sp}_{2,1} \oplus \mathfrak{su}_2$

Remark 3.3. *In Table 3, the Riemannian symmetric pair $(\mathfrak{g}, \mathfrak{k})$ is of Hermitian type if and only if \mathfrak{g} is isomorphic to $\mathfrak{su}_{2n-2j, 2j}$, $\mathfrak{su}_{n, n}$, $\mathfrak{so}_{m, n}$ with $\min\{m, n\} = 2$, $\mathfrak{sp}_n(\mathbb{R})$, $\mathfrak{sp}_{2n}(\mathbb{R})$ or $\mathfrak{e}_{6(-14)}$.*

3.2. In the case where \mathfrak{g} is not absolutely simple. Let \mathfrak{g} be a non-compact simple Lie algebra over \mathbb{R} and $(\mathfrak{g}, \mathfrak{h})$ a symmetric pair. In this subsection, we assume that \mathfrak{g} is not absolutely simple. Then \mathfrak{g} admits complex structures $\{\pm J\}$ (see Section 2). For the involution σ on \mathfrak{g} defining \mathfrak{h} , either one of the following two situations occurs:

- $\sigma \circ J = (-J) \circ \sigma$.
- $\sigma \circ J = J \circ \sigma$.

We fix a Cartan involution θ on \mathfrak{g} commuting with σ . Then $\theta \circ J = (-J) \circ \theta$, and the Cartan decomposition of \mathfrak{g} corresponding to θ can be written as $\mathfrak{g} = \mathfrak{u} + J\mathfrak{u}$ by a compact real form \mathfrak{u} of $(\mathfrak{g}, \pm J)$.

Let us consider the case where $\sigma \circ J = (-J) \circ \sigma$, that is, σ is anti- \mathbb{C} -linear on $(\mathfrak{g}, \pm J)$. Then the subalgebra \mathfrak{h} is a real form of $(\mathfrak{g}, \pm J)$, and the real form \mathfrak{g}^d of $\mathfrak{g}_{\mathbb{C}} = (\mathfrak{g}, J) \oplus (\mathfrak{g}, -J)$ can be written as

$$\{(X, \theta\sigma X) \in (\mathfrak{g}, J) \oplus (\mathfrak{g}, -J) \mid X \in \mathfrak{g}\}.$$

In particular, the simple Lie algebra \mathfrak{g}^d over \mathbb{R} is isomorphic to \mathfrak{g} .

Next, we shall suppose that $\sigma \circ J = J \circ \sigma$, that is, σ is \mathbb{C} -linear on $(\mathfrak{g}, \pm J)$. Then \mathfrak{h} is $\pm J$ -stable. In this situation, $\mathfrak{u} \cap \mathfrak{h}$ gives a real form of $(\mathfrak{h}, \pm J)$. We define the real form

$$\mathfrak{g}_0^d := (\mathfrak{u} \cap \mathfrak{h}) + J(\mathfrak{u} \cap \mathfrak{q})$$

of $(\mathfrak{g}, \pm J)$. Then the real form \mathfrak{g}^d of $\mathfrak{g}_{\mathbb{C}} = (\mathfrak{g}, J) \oplus (\mathfrak{g}, -J)$ can be identified with $\mathfrak{g}_0^d \oplus \mathfrak{g}_0^d$, and \mathfrak{h} can be understood as the complexification of the maximal compact subalgebra $\mathfrak{u} \cap \mathfrak{h}$ of \mathfrak{g}_0^d .

The theorem below gives a classification of $(\mathfrak{g}, \mathfrak{h})$ with $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}} \cap \mathfrak{g}^d = \emptyset$ in the setting above:

Theorem 3.4. *Let $(\mathfrak{g}, \mathfrak{h})$ be a symmetric pair with simple \mathfrak{g} admitting two complex structures $\{\pm J\}$.*

- (1) *Suppose that $\sigma \circ J = (-J) \circ \sigma$. Then $m(\mathfrak{g}) = n(\mathfrak{g}_{\mathbb{C}}) = m(\mathfrak{g}^d)$, and $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}} \cap \mathfrak{g}^d \neq \emptyset$.*
- (2) *Suppose that $\sigma \circ J = J \circ \sigma$. Then the following six conditions on $(\mathfrak{g}, \mathfrak{h})$ are equivalent:*
 - (i) $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}} \cap \mathfrak{g}^d = \emptyset$.
 - (ii) $m(\mathfrak{g}) = n(\mathfrak{g}_{\mathbb{C}}) < m(\mathfrak{g}^d)$ where we put $m(\mathfrak{g}^d) = 2m(\mathfrak{g}_0^d)$.
 - (iii) $\mathcal{O}_{\min} \cap \mathfrak{g}_0^d = \emptyset$.
 - (iv) $n((\mathfrak{g}, \pm J)) < m(\mathfrak{g}_0^d)$.
 - (v) \mathfrak{g}_0^d is isomorphic to one of the Lie algebras in Table 2.
 - (vi) The complex symmetric pair $(\mathfrak{g}, \mathfrak{h})$ is isomorphic to one of the symmetric pairs in Table 4.

TABLE 4. List of $(\mathfrak{g}, \mathfrak{h})$ with non-absolutely-simple \mathfrak{g} such that $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}} \cap \mathfrak{g}^d = \emptyset$

$(\mathfrak{g}, \mathfrak{h})$	$\mathfrak{g}_0^d = \mathfrak{h}^a$
$(\mathfrak{sl}_{2n}(\mathbb{C}), \mathfrak{sp}_n(\mathbb{C}))$ ($n \geq 2$)	\mathfrak{su}_{2n}^*
$(\mathfrak{so}_n(\mathbb{C}), \mathfrak{so}_{n-1}(\mathbb{C}))$ ($n \geq 5$)	$\mathfrak{so}_{n-1,1}$
$(\mathfrak{sp}_{m+n}(\mathbb{C}), \mathfrak{sp}_m(\mathbb{C}) \oplus \mathfrak{sp}_n(\mathbb{C}))$ ($n+m \geq 2, n, m \geq 1$).	$\mathfrak{sp}_{m,n}$
$(\mathfrak{e}_6^{\mathbb{C}}, \mathfrak{f}_4^{\mathbb{C}})$	$\mathfrak{e}_{6(-26)}$
$(\mathfrak{f}_4^{\mathbb{C}}, \mathfrak{so}_9(\mathbb{C}))$	$\mathfrak{f}_{4(-20)}$

Proof of Theorem 3.4. As we mentioned in Section 2, under the identification of $\mathfrak{g}_{\mathbb{C}}$ with $(\mathfrak{g}, J) \oplus (\mathfrak{g}, -J)$, the orbit $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}}$ coincides with $\mathcal{O}_{\min} \times \mathcal{O}_{\min}$, and $n(\mathfrak{g}_{\mathbb{C}}) = m(\mathfrak{g})$.

First, let us suppose that $\sigma \circ J = (-J) \circ \sigma$. Then the real form \mathfrak{g}^d of $\mathfrak{g}_{\mathbb{C}}$ is simple but not absolutely-simple since \mathfrak{g}^d is isomorphic to \mathfrak{g} . Thus by the arguments in Section 2 again, we have $n(\mathfrak{g}_{\mathbb{C}}) = m(\mathfrak{g}^d)$ and $\mathcal{O}_{\min, \mathfrak{g}^d}^{\mathbb{C}} = \mathcal{O}_{\min} \times \mathcal{O}_{\min}$. Thus we have $m(\mathfrak{g}) = n(\mathfrak{g}_{\mathbb{C}}) = m(\mathfrak{g}^d)$, and $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}} \cap \mathfrak{g}^d \neq \emptyset$. This completes the proof of the claim (1).

Next, suppose that $\sigma \circ J = J \circ \sigma$. The equivalence among (2i), (2iii), (2iv) and (2v) comes from $\mathfrak{g}^d = \mathfrak{g}_0^d \oplus \mathfrak{g}_0^d$, $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}} = \mathcal{O}_{\min} \times \mathcal{O}_{\min}$ and Fact 2.2. The equivalence (2iv) \Leftrightarrow (2ii) is followed by $n(\mathfrak{g}_{\mathbb{C}}) = n(\mathfrak{g}, J) + n(\mathfrak{g}, -J)$, $n(\mathfrak{g}, J) = n(\mathfrak{g}, -J)$ and $m(\mathfrak{g}^d) = 2m(\mathfrak{g}_0^d)$. Finally, we obtain the equivalence (2v) \Leftrightarrow (2vi) by the observation that $(\mathfrak{g}, \pm J)$ is isomorphic to the complexification of \mathfrak{g}_0^d , and \mathfrak{h} is the complexification of the maximal compact subalgebra $\mathfrak{u} \cap \mathfrak{g}_0^d$ of \mathfrak{g}_0^d in $(\mathfrak{g}, \pm J)$. The proof of the claim (2) has been completed. \square

Remark 3.5. For each symmetric pair $(\mathfrak{g}, \mathfrak{h})$, the dual Lie algebra of the complexified symmetric pair $(\mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C}, \mathfrak{h} \otimes_{\mathbb{R}} \mathbb{C})$ can be written as $\mathfrak{g}^d \oplus \mathfrak{g}^d$. Therefore, if $(\mathfrak{g}, \mathfrak{h})$ satisfies the conditions in Theorem 3.2, then $(\mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C}, \mathfrak{h} \otimes_{\mathbb{R}} \mathbb{C})$ satisfies the conditions in Theorem 3.4. Note that the converse claim does not hold. In fact, suppose \mathfrak{g} is absolutely simple and both $m(\mathfrak{g})$ and $m(\mathfrak{g}^d)$ are greater than $n(\mathfrak{g}_{\mathbb{C}})$, then $(\mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C}, \mathfrak{h} \otimes_{\mathbb{R}} \mathbb{C})$ satisfies the conditions in Theorem 3.4 but the conditions in Theorem 3.2 do not hold for $(\mathfrak{g}, \mathfrak{h})$. The symmetric pair $(\mathfrak{g}, \mathfrak{h}) = (\mathfrak{su}_{2n}^*, \mathfrak{sp}_{2n-j,j})$ with $\mathfrak{g}^d = \mathfrak{su}_{2n}^*$ is one of examples.

4. BOUNDED MULTIPLICITY BRANCHING FOR SYMMETRIC PAIRS

In this section, as an application of Theorem 3.1, we give a proof of Theorem 1.2 stated in Section 1.

Let G be a non-compact simple Lie group with Lie algebra \mathfrak{g} . For $\Pi \in \text{Irr}(G)$ and a reductive subgroup H of G , as in [8, 9], we say that the restriction $\Pi|_H$ has the bounded multiplicity property if

$$\sup_{\pi \in \text{Irr}(H)} [\Pi|_H : \pi] < \infty.$$

T. Kobayashi [9] gives the following theorem for bounded multiplicity properties of restrictions of irreducible G -representation with “small” Gelfand–Kirillov dimensions for symmetric pairs:

Fact 4.1 ([8, Theorem 3], [9, Theorems 5.5 (1) and 6.1]). *Let $\Pi \in \text{Irr}(G)$.*

- (1) *Suppose that $n(\mathfrak{g}_{\mathbb{C}}) = \text{DIM}(\Pi)$. Then for any symmetric pair (G, H) , the restriction $\Pi|_H$ has the bounded multiplicity property.*
- (2) *Suppose that $m(\mathfrak{g}) = \text{DIM}(\Pi)$. Take a symmetric pair (G, H) such that the corresponding symmetric pair $(\mathfrak{g}, \mathfrak{h})$ of Lie algebras satisfies the condition “ $\sigma\mu = -\mu$ ” (see below for the details). Then the restriction $\Pi|_H$ has the bounded multiplicity property.*

Let us explain the condition “ $\sigma\mu = -\mu$ ” for a symmetric pair $(\mathfrak{g}, \mathfrak{h})$ with simple \mathfrak{g} . As in Section 1, we denote by σ the involution on \mathfrak{g} defining \mathfrak{h} , and put $\mathfrak{q} := \{X \in \mathfrak{g} \mid \sigma(X) = -X\}$. Fix a Cartan involution θ on \mathfrak{g} commuting with σ , and write $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ for the Cartan decomposition. We take a maximal abelian subspace $\mathfrak{a}^{-\sigma}$ of $\mathfrak{p} \cap \mathfrak{q}$, and extend it to a maximal abelian subspace \mathfrak{a} of \mathfrak{p} . Note that \mathfrak{a} is σ -stable. The dual of the vector space \mathfrak{a} is denoted by \mathfrak{a}^* , and we use the same symbol σ for the involution on \mathfrak{a}^* induced by σ on \mathfrak{a} . Then the root system $\Sigma(\mathfrak{g}, \mathfrak{a})$ is known to be σ -stable. Let us also fix a lexicographic ordering on $\mathfrak{a}^{-\sigma}$, and extend it to that on \mathfrak{a} . The positive system of the root system $\Sigma(\mathfrak{g}, \mathfrak{a})$ corresponding to the fixed ordering is denoted by $\Sigma^+(\mathfrak{g}, \mathfrak{a})$. We write μ for the highest element in $\Sigma^+(\mathfrak{g}, \mathfrak{a})$. Then the vectors $\sigma\mu$ and $-\mu$ in \mathfrak{a}^* are both defined. Note that the condition “ $\sigma\mu = -\mu$ ” depends only on $(\mathfrak{g}, \mathfrak{h})$.

Theorem 1.2 can be obtained as a corollary to Kobayashi’s theorem (Fact 4.1), Theorem 3.1 and the following proposition:

Proposition 4.2. *Let $(\mathfrak{g}, \mathfrak{h})$ be a symmetric pair with simple \mathfrak{g} . Then the condition “ $\sigma\mu = -\mu$ ” holds if and only if $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}} \cap \mathfrak{g}^d \neq \emptyset$.*

Proof of Proposition 4.2. Let us denote by the coroot $\mu^{\vee} \in \mathfrak{a}$ of the highest root $\mu \in \Sigma(\mathfrak{g}, \mathfrak{a})$. We denote by $\mathcal{O}_{\mu^{\vee}}$ the real adjoint orbit in \mathfrak{g} through μ^{\vee} , and by $\mathcal{O}_{\mu^{\vee}}^{\mathbb{C}}$ the complex adjoint orbit in $\mathfrak{g}_{\mathbb{C}}$ through μ^{\vee} . We shall prove that the following six conditions on $(\mathfrak{g}, \mathfrak{h})$ are equivalent:

- (i) $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}}$ meets \mathfrak{g}^d .
- (ii) $\mathcal{O}_{\mu^{\vee}}^{\mathbb{C}}$ meets \mathfrak{g}^d .
- (iii) $\mathcal{O}_{\mu^{\vee}}$ meets $\mathfrak{g} \cap \mathfrak{g}^d = \mathfrak{h}^{\mathfrak{a}} = (\mathfrak{k} \cap \mathfrak{h}) + (\mathfrak{p} \cap \mathfrak{q})$.
- (iv) $\mathcal{O}_{\mu^{\vee}}$ meets $\mathfrak{a}^{-\sigma} \cap \mathfrak{a}_+$ at one point, where \mathfrak{a}_+ denotes the closed Weyl chamber corresponding to the positive system $\Sigma^+(\mathfrak{g}, \mathfrak{a})$.
- (v) $\mu^{\vee} \in \mathfrak{a}^{-\sigma}$.
- (vi) $\sigma\mu = -\mu$.

Let us fix any non-zero root vector $X \in \mathfrak{g}_{\mu}$. Then one can find $Y \in \mathfrak{g}_{-\mu}$ such that (μ^{\vee}, X, Y) is an \mathfrak{sl}_2 -triple, and $X \in \mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}}$ by [12, Theorem

1.1]. Thus we obtain the equivalence (i) \Leftrightarrow (ii) by [11, Lemma 4.7]. Furthermore, by [11, Proposition 4.6], we have the equivalence (ii) \Leftrightarrow (iii). The equivalence (iii) \Leftrightarrow (iv) comes from [11, Lemma 4.9 and Fact 5.1]. Since μ is the highest root of $\Sigma^+(\mathfrak{g}, \mathfrak{a})$, we have $\mu^\vee \in \mathfrak{a}_+$, and hence (iv) \Leftrightarrow (v). The equivalence (v) \Leftrightarrow (vi) is easy. \square

Let us also give a remark for almost irreducible branching laws as below. Let (G, H) be a symmetric pair with a non-compact connected absolutely simple Lie group G , and denote by $(\mathfrak{g}, \mathfrak{h})$ the corresponding pair of Lie algebras. Suppose G has a minimal representation $\Pi_{\min} \in \text{Irr}(G)$ (see [9, Lemma 5.8] and [14] for the construction of minimal representations in suitable situations). Then $n(\mathfrak{g}_{\mathbb{C}}) = \text{DIM}(\Pi_{\min})$, and consequently, $n(\mathfrak{g}_{\mathbb{C}}) = m(\mathfrak{g})$. Consider the restriction $\Pi_{\min}|_H$ of Π_{\min} to H . Since $\text{DIM}(\Pi_{\min}) = n(\mathfrak{g}_{\mathbb{C}})$, by Fact 4.1 (1), the restriction $\Pi_{\min}|_H$ exhibits the bounded multiplicity property (refer to [8, Theorem 7]).

In such a situation, as mentioned by Kobayashi [9, Remark 6.4], the following theorem holds as an application of [8, Theorem 10]:

Theorem 4.3. *In the setting above, suppose that $(\mathfrak{g}, \mathfrak{h})$ can be found in Table 3, or equivalently “ $\sigma\mu \neq -\mu$ ”. Then the restriction $\Pi_{\min}|_H$ is almost irreducible, that is, the H -module $\Pi_{\min}|_H$ remains irreducible or decomposes into a finite sum of irreducible representations of H .*

Proof. Let us state Condition (\star) on $(\mathfrak{g}, \mathfrak{h})$ as below:

Condition (\star) : There exists a Riemannian symmetric pair $(\mathfrak{g}_0, \mathfrak{k}_0)$ such that $n(\mathfrak{g}_{\mathbb{C}}) < m(\mathfrak{g}_0)$ and $(\mathfrak{g}_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}}) := (\mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C}, \mathfrak{h} \otimes_{\mathbb{R}} \mathbb{C})$ is isomorphic to $(\mathfrak{g}_0 \otimes_{\mathbb{R}} \mathbb{C}, \mathfrak{k}_0 \otimes_{\mathbb{R}} \mathbb{C})$.

Kobayashi [8, Theorem 10] proved that under the condition (\star) , the restriction $\Pi_{\min}|_H$ is almost irreducible.

We note that $(\mathfrak{g}^d, \mathfrak{k}(\mathfrak{g}^d))$ is the unique Riemannian symmetric pair whose complexification is isomorphic to $(\mathfrak{g}_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$. Thus Condition (\star) is nothing but $n(\mathfrak{g}_{\mathbb{C}}) < m(\mathfrak{g}^d)$. Thus the claim of Theorem 4.3 follows directly from [8, Theorem 10], Theorem 3.2 and Proposition 4.2. \square

Remark 4.4. *See also [7] for other triples (G, H, π) with almost irreducible branching laws.*

5. PROPER $SL_2(\mathbb{R})$ -ACTIONS ASSOCIATED TO REAL MINIMAL NILPOTENT ORBITS

We also give another application of our results in Section 3 for proper $SL_2(\mathbb{R})$ -actions on symmetric spaces (cf. [11]).

Let $(\mathfrak{g}, \mathfrak{h})$ be a symmetric pair with a simple Lie algebra \mathfrak{g} . We denote by $(\mathfrak{g}, \mathfrak{h}^a)$ the associated pair of $(\mathfrak{g}, \mathfrak{h})$ in the sense of [13]. Note that $\mathfrak{g} \cap \mathfrak{g}^d = \mathfrak{h}^a$. Let (G, H^a) be a symmetric pair of Lie groups whose symmetric pair of Lie algebras is isomorphic to $(\mathfrak{g}, \mathfrak{h}^a)$. We also suppose that G is linear and connected.

For each Lie algebra homomorphism, $\rho : \mathfrak{sl}_2(\mathbb{R}) \rightarrow \mathfrak{g}$, we denote by $\mathcal{O}_{\rho, \text{nilp}}^{\mathbb{R}}$ and $\mathcal{O}_{\rho, \text{nilp}}^{\mathbb{C}}$ for the real nilpotent adjoint orbit and the complex nilpotent adjoint orbit through the nilpotent element

$$\rho \left(\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right) \in \mathfrak{g}$$

in \mathfrak{g} and in $\mathfrak{g}_{\mathbb{C}} := \mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C}$, respectively. Note that by the Jacobson–Morozov theorem and Kostant’s theorem for real nilpotent orbits in \mathfrak{g} (see [4, Chapter 9] for the details), the correspondence $\rho \mapsto \mathcal{O}_{\rho, \text{nilp}}^{\mathbb{R}}$ defines a bijection from the set of all $\text{Int}(\mathfrak{g})$ -conjugacy classes of $\rho : \mathfrak{sl}_2(\mathbb{R}) \rightarrow \mathfrak{g}$ onto the set of all real nilpotent orbits in \mathfrak{g} , where $\text{Int}(\mathfrak{g})$ denotes the group of all inner-automorphisms on \mathfrak{g} .

Let us use the same symbol ρ for the Lie group homomorphism $\rho : SL_2(\mathbb{R}) \rightarrow G$ corresponding to $\rho : \mathfrak{sl}_2(\mathbb{R}) \rightarrow \mathfrak{g}$ (see [11, Lemma 5.4]). Then by [11, Theorem 10.1], which is a corollary to the properness criterion by T. Kobayashi [6, Theorem 4.1], the $SL_2(\mathbb{R})$ -action on the symmetric space G/H^a via $\rho : SL_2(\mathbb{R}) \rightarrow G$ is proper if and only if $\mathcal{O}_{\rho, \text{nilp}}^{\mathbb{C}} \cap \mathfrak{g}^d = \emptyset$.

We shall consider a Lie algebra homomorphism $\rho_{\min} : \mathfrak{sl}_2(\mathbb{R}) \rightarrow \mathfrak{g}$ such that $\mathcal{O}_{\rho_{\min}, \text{nilp}}^{\mathbb{R}}$ is one of the real minimal nilpotent orbit in \mathfrak{g} . Then $\mathcal{O}_{\rho_{\min}, \text{nilp}}^{\mathbb{C}} = \mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}}$ and the hyperbolic element

$$\rho_{\min} \left(\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \right) \in \mathfrak{g}$$

is conjugate to the coroot $\mu^{\vee} \in \mathfrak{a}$ of the highest element μ of $\Sigma^+(\mathfrak{g}, \mathfrak{a})$ defined in Section 4.

Then by Theorems 3.2 and 3.4, we have the theorem below:

Theorem 5.1. *The following three conditions on $(\mathfrak{g}, \mathfrak{h})$ are equivalent:*

- (i) *The $SL_2(\mathbb{R})$ -action on G/H^a via ρ_{\min} is proper.*
- (ii) $\mathcal{O}_{\min, \mathfrak{g}}^{\mathbb{C}} \cap \mathfrak{g}^d = \emptyset$.
- (iii) *The symmetric pair $(\mathfrak{g}, \mathfrak{h})$ is isomorphic to one of the symmetric pairs in Tables 3 or 4.*

ACKNOWLEDGEMENTS.

The author would like to give heartfelt thanks to Toshiyuki Kobayashi for his encouragement to write this paper. I am also indebted to Koichi Tojo and Ryusei Tsukada for many helpful comments. I also would like to express my gratitude to the anonymous referees for their meticulous review of the manuscript. The author is supported by JSPS Grants-in-Aid for Scientific Research JP20K03589, JP20K14310, JP22H01124, and JP24K06714.

REFERENCES

- [1] Shôrô Araki, *On root systems and an infinitesimal classification of irreducible symmetric spaces*, J. Math. Osaka City Univ. **13** (1962), 1–34. MR 153782
- [2] Marcel Berger, *Les espaces symétriques noncompacts*, Ann. Sci. École Norm. Sup. (3) **74** (1957), 85–177. MR 0104763
- [3] Ranee Brylinski, *Geometric quantization of real minimal nilpotent orbits*, vol. 9, 1998, Symplectic geometry, pp. 5–58. MR 1636300
- [4] David H. Collingwood and William M. McGovern, *Nilpotent orbits in semisimple Lie algebras*, Van Nostrand Reinhold Mathematics Series, Van Nostrand Reinhold Co., New York, 1993. MR 1251060
- [5] Sigurdur Helgason, *Differential geometry, Lie groups, and symmetric spaces*, Graduate Studies in Mathematics, vol. 34, American Mathematical Society, Providence, RI, 2001, Corrected reprint of the 1978 original. MR 1834454
- [6] Toshiyuki Kobayashi, *Proper action on a homogeneous space of reductive type*, Math. Ann. **285** (1989), no. 2, 249–263. MR 1016093
- [7] ———, *Branching problems of Zuckerman derived functor modules*, Representation theory and mathematical physics, Contemp. Math., vol. 557, Amer. Math. Soc., Providence, RI, 2011, pp. 23–40. MR 2848919
- [8] ———, *Multiplicity in restricting minimal representations*, Lie theory and its applications in physics, Springer Proc. Math. Stat., vol. 396, Springer, Singapore, [2022] ©2022, pp. 3–20. MR 4607949
- [9] ———, *Bounded multiplicity branching for symmetric pairs*, J. Lie Theory **33** (2023), no. 1, 305–328. MR 4567759
- [10] Toshiyuki Kobayashi and Yoshiki Oshima, *Classification of symmetric pairs with discretely decomposable restrictions of (\mathfrak{g}, K) -modules*, J. Reine Angew. Math. **703** (2015), 201–223. MR 3353547
- [11] Takayuki Okuda, *Classification of semisimple symmetric spaces with proper $SL(2, \mathbb{R})$ -actions*, J. Differential Geom. **94** (2013), no. 2, 301–342. MR 3080484
- [12] ———, *Smallest complex nilpotent orbits with real points*, J. Lie Theory **25** (2015), no. 2, 507–533. MR 3346070
- [13] Toshio Ôshima and Jirô Sekiguchi, *The restricted root system of a semisimple symmetric pair*, Group representations and systems of differential equations (Tokyo, 1982), Adv. Stud. Pure Math., vol. 4, North-Holland, Amsterdam, 1984, pp. 433–497. MR 810638
- [14] Hiroyoshi Tamori, *Classification of minimal representations of real simple Lie groups*, Math. Z. **292** (2019), no. 1-2, 387–402. MR 3968907

(T. Okuda) GRADUATE SCHOOL OF ADVANCED SCIENCE AND ENGINEERING,
HIROSHIMA UNIVERSITY, 1-3-1 KAGAMIYAMA, HIGASHI-HIROSHIMA CITY, HIROSHIMA, 739-8526, JAPAN.

Email address: okudatak@hiroshima-u.ac.jp