ON CANONICAL NUMBER SYSTEMS AND MONOGENITY OF PURE NUMBER FIELDS

HAMID BEN YAKKOU AND BRAHIM BOUDINE

ABSTRACT. Let m be a rational integer $(m \neq 0, \pm 1)$ and consider a pure number field $K = \mathbb{Q}(\sqrt[n]{m})$ with $n \geq 3$. Most papers discussing the monogenity of pure number fields focus only on the case where m is square-free. In this paper, based on a classical theorem of Ore on the prime ideal decomposition [27, 29], we study the monogenity of K with m is not necessarily square-free. As an application, several examples of the application of monogenic fields to CNS (Canonical Number System) are shown. In this way, our results extend some parts of previously established results [3, 5, 23].

1. INTRODUCTION

Consider a monic irreducible polynomial $F(x) \in \mathbb{Z}[x]$ of degree $n, K := \mathbb{Q}(\alpha)$ a number field with α a roof of F(x) and \mathbb{Z}_K the ring of integers of K. A primitive element of \mathbb{Z}_K is an element θ verifying $K = \mathbb{Q}(\theta)$. Let $\widehat{\mathbb{Z}}_K$ be the set of all primitive elements of \mathbb{Z}_K . We say that K is monogenic if there exists $\theta \in \widehat{\mathbb{Z}}_K$ such that $(1, \theta, \theta^2, ..., \theta^{n-1})$ is a basis of the \mathbb{Z} -module \mathbb{Z}_K . In this case, we say that θ is a generator of a power integral basis of \mathbb{Z}_K (shortly, GPIB of \mathbb{Z}_K).

The monogenity of number fields is among the most active classical problems in algebraic number theory (cf. [1, 5, 8, 6, 10, 11, 17, 18, 24, 31, 32]). By Hensel's discovery in 1894, the monogenity of K is encoded by the solvability of an index form equation

$$I(x_2, ..., x_n) = \pm 1 \text{ in } x_2, ..., x_n \in \mathbb{Z}$$
(1.1)

associated with an integral basis $\{1, \omega_2, ..., \omega_n\}$ of K (see [8] and [28, 2.2, Section 6, p. 64]).

In [17, 18, 20], Győry provided general algorithms to decide whether K is monogenic or not and to determine all power integral bases in \mathbb{Z}_K and gave effective bounds for the solutions of (1.1) and gave algorithms for solving index form equations. He also studied in [19, 21] the monogenity of relative extensions. For detailed surveys on discriminant and index form theory and its applications, including related Diophantine equations and monogenity of number fields, see the book [8] by Evertse and Győry.

For quartic number fields see [7] by Davis and Spearman and [13] by Gaál, Pethő and Pohst. In [11], Gaál and Győry described an algorithm to solve index form equations in quintic number fields and made very extensive computations for power integral bases in totally real quintic number fields with Galois group S_5 . In [6], Bilu, Gaál and Győry studied the monogenity of sextic number fields. In [32], Pethő and Ziegler gave an efficient

²⁰²⁰ Mathematics Subject Classification. 11A63, 11R04, 11R16, 11R21, 11Y40.

Key words and phrases. Canonical number system, Newton polygon, Theorem of Ore, prime ideal factorization, power integral basis, monogenity.

criterion to decide whether the maximal order of a biquadratic field has a unit power integral basis or not. For multiquadratic number fields, see [31] by Pethő and Pohst. For a survey on monogenity with a focus on efficient algorithms for several classes of number fields, see the books [8, 10]. For another bibliography of monogenity, see Narkiewicz's book [28].

Recently, the interest in the study of monogenity of pure number fields $\mathbb{Q}(\sqrt[6]{m})$ increased clearly. Despite the lack of a standard classification, there has been some progress in this direction. There are several works that consider the case when m is a square-free rational integer: we remain for example that Gassert [14] investigated when $\mathbb{Z}_K = \mathbb{Z}[\alpha]$, Ahmad et al. [1, 2] explored the case n = 6, Hameed and Nakahara [22] studied the case n = 8, Gaál and Remete studied the cases $3 \leq n \leq 9$. In [3, 4, 5], Ben Yakkou et al. studied several classes of pure number fields with variables exponents.

In the present paper, we study the monogenity of any pure number field $K = \mathbb{Q}(\sqrt[n]{m})$ without the condition that m is square-free. In this way, our results extend some part of several of previously published works.

2. Main results

Let p be a prime. Throughout this paper, \mathbb{F}_p denotes the finite field with p elements. For $m \in \mathbb{Z}$, $\nu_p(m)$ stands for the p-adic valuation of m and PD(m) stands for the set of prime divisors of m. For two positive rational integers d and u, we denote by $N_p(d)$ the number of monic irreducible polynomials of degree d in $\mathbb{F}_p[x]$ and by $N_p(d, u, m)$ the number of monic irreducible factors of degree d of the polynomial $x^u - \overline{m}$ in $\mathbb{F}_p[x]$.

Now, let us give our results. We start by the following theorem which gives infinite families of non-monogenic pure number fields; that is \mathbb{Z}_K has no power integral basis.

Theorem 2.1. Let $F(x) = x^n - m \in \mathbb{Z}[x]$ be an irreducible polynomial, $K := \mathbb{Q}(\alpha)$ a pure number field with α a root of F(x). Assume that $n = u \cdot p^r$, where p is an odd prime not dividing $u \cdot m$ and let $\nu = \nu_p(m^{p-1}-1)$. If $\min\{r+1,\nu\}N(d,u,m) > N_p(d)$ for some positive integer d, then K is not monogenic.

Remark 2.2. Note that the condition *m* is square free is not required for the above theorem.

Remark 2.3. The above theorem implies [5, Theorem 2.4] and [3, Theorem 2.2], where the special cases $n = p^r$ and $n = 2^5 \cdot 5^s$ with m is square-free are previously and respectively studied.

As a consequence of the above result, the following three corollaries provide infinite parametric families of non-monogenic pure number fields with variable exponents.

Corollary 2.4. For $F(x) = x^{5^r \cdot 7^s} - m$. If any one of the following conditions holds:

- (1) $r \ge 1, s \ge 7$ and $m^6 \equiv 1 \pmod{7^8}$.
- (2) r > 5, s > 1 and $m^4 \equiv 1 \pmod{5^6}$,

then K is not monogenic.

Corollary 2.5. For $F(x) = x^{3^r \cdot 11^s} - m$. If any one of the following conditions holds:

- (1) $r \ge 1, s \ge 11$ and $m^{10} \pmod{11^{12}}$.
- (2) $r \ge 2, s \ge 1 \text{ and } m^2 \equiv 1 \pmod{27}$,

then K is not monogenic.

Corollary 2.6. For $F(x) = x^{5^r \cdot 11^s} - m$. If any one of the following conditions holds:

- (1) $r \ge 1, s \ge 2, m \equiv -1 \pmod{11}$ and $m^{10} \equiv 1 \pmod{1331}$.
- (2) $r \ge 6, s \ge 1 \text{ and } m^4 \equiv 1 \pmod{5^6}$,

then K is not monogenic.

Recall that a monic polynomial $F(x) \in \mathbb{Z}[x]$ is called monogenic if it is irreducible over \mathbb{Q} and $\mathbb{Z}_K = \mathbb{Z}[\alpha]$, where $K = \mathbb{Q}(\alpha)$ with $F(\alpha) = 0$. But, it may happen that the field K is monogenic despite its defining polynomial F(x) is not monogenic (see e.g [24] by Jones). In the following result, we identify new infinite parametric families of monogenic pure number fields defined by non-monogenic binomials.

Theorem 2.7. Let $F(x) = x^n - a^u \in \mathbb{Z}[x]$ such that $u \ge 2$, gcd(u, n) = 1, $PD(n) \subseteq PD(a)$ and a is square-free. Then F(x) is irreducible over \mathbb{Q} . Let $K := \mathbb{Q}(\alpha)$ a pure number field with α a root of F(x) and \mathbb{Z}_K the ring of integers of K. Then, we have the following:

- (1) $\mathbb{Z}[\alpha] \neq \mathbb{Z}_K$.
- (2) There exist $\theta \in \widehat{\mathbb{Z}}_K$ such that $\mathbb{Z}_K = \mathbb{Z}[\theta]$; K monogenic.

Remark 2.8.

- (1) Note that in above theorem a^u is not square-free.
- (2) Recall that in [14], Gassert gave a necessary and sufficient condition so that $\mathbb{Z}[\alpha] = \mathbb{Z}_K$ when u = 1. In the above theorem, we have $u \ge 2$, α is not a GPIB of \mathbb{Z}_K but \mathbb{Z}_K admits a GPIB θ different to α .

3. Preliminaries

In what follows, $K = \mathbb{Q}(\alpha)$ is a number field with ring of integers \mathcal{O}_K with $\alpha \in \mathbb{Z}_K$ a root of a monic irreducible polynomial $F(x) \in \mathbb{Z}[x]$ of degree n. Recall that for any θ in $\widehat{\mathbb{Z}}_K, \mathbb{Z}[\theta]$ is a sub- \mathbb{Z} -module of \mathbb{Z}_K with rank n. Thus, the index of $\mathbb{Z}[\theta]$ in \mathbb{Z}_K is finite. This index is denoted by $(\mathbb{Z}_K : \mathbb{Z}[\theta])$. So, K is monogenic if $(\mathbb{Z}_K : \mathbb{Z}[\theta]) = 1$ for some primitive element θ of K. But, if K does not have any such θ , then K is not monogenic.

The greatest common divisor of the indices of all integral primitive elements of K is called the index of K and is denoted by i(K), that is

$$i(K) = \gcd\{(\mathbb{Z}_K : \mathbb{Z}[\theta]) \mid \theta \in \mathbb{Z}_K\}.$$

A prime p dividing i(K) is called a prime common index divisor of K. If K is monogenic, then i(K) = 1. Therefore a field having a prime common index divisor is not monogenic.

For a positive integer d, we denote $L_p(d)$ the number of distinct prime ideals of \mathbb{Z}_K , that are factors of $p\mathbb{Z}_K$ with residue degree d and $N_p(d)$ denote the number of monic and irreducible polynomial having degree d in $\mathbb{Z}/p\mathbb{Z}[x]$. Then, we have the following lemma which provides a sufficient condition for a prime p to be a common index divisor of a number field:

Lemma 3.1. [28, Theorems 4.33 and 4.34] With the above notation, if $L_p(d) > N_p(d)$ for some positive integer d, then p is a common index divisor of K. In particular, K is not monogenic.

To prove our result, we shall use the index theorem of Ore which gives $\nu_p((\mathbb{Z}_K : \mathbb{Z}[\theta]))$ and the factorization of $p\mathbb{Z}_K$. This theorem allows us to apply the above lemma (for non-monogenity) as well as to test if $\nu_p((\mathbb{Z}_K : \mathbb{Z}[\theta])) = 1$ or not (for monogenity). Ore's Theorem is based on Newton's polygon techniques. So, let us briefly describe the related notions and the use of this technique. For more details, we refer to [29] by Ore, [27] by Montes and Nart, and to [15, 16] by Guàrdia, Montes and Nart.

Let p be a prime, ν_p the p-adic valuation For any polynomial $F(x) := \sum_{i=0}^n a_i x^i \in \mathbb{Z}_p[x]$, the p-adic valuation of F(x) is $\nu_p(F(x)) = \min\{\nu_p(a_i) \mid 0 \le i \le n\}$. Let $\phi(x)$ be a monic polynomial in $\mathbb{Z}[x]$, where $\overline{\phi(x)}$ is irreducible in $\mathbb{F}_p[x]$. Let

$$F(x) = a_l(x)\phi(x)^l + a_{l-1}(x)\phi(x)^{l-1} + \dots + a_1(x)\phi(x) + a_0(x)$$

be the ϕ -adic development of F(x): $\deg(a_j(x)) < \deg(\phi(x))$ for each $j \in \{0, 1, ..., l\}$. We consider the set $\mathfrak{S}_{\{F,\phi\}} = \{(j, \nu_p(a_j(x))) \mid a_j(x) \neq 0 \text{ and } 0 \leq j \leq l\}$. The lower boundary convex envelope of $\mathfrak{S}_{\{F,\phi\}}$ is called the ϕ -Newton polygon of F(x), denoted by $N_{\phi}(F)$. It is made up of some sides S_k with different slopes $\lambda_k \in \mathbb{Q}$, then we write $N_{\phi}(F) =$ $S_1 + S_2 + ... + S_g$, where g is the number of its sides with different slopes. In particular, the ϕ -principal Newton polygon of F(x), denoted by $N_{\phi}^+(F)$, is the polygon formed by the sides of $N_{\phi}(F)$ which have negative slopes. Every side S of $N_{\phi}^+(F)$ joins two points: 1) the beginning point $(s, \nu_p(a_s(x)))$, and 2) the final point $(r, \nu_p(a_r(x)))$, where $0 \leq s < r \leq n$. In this case, the length of S is l(S) := r - s, its height is $h(S) := \nu_p(a_s(x)) - \nu_p(a_r(x))$, its slope is $-\lambda = -\frac{h(S)}{l(S)} < 0$, its degree is $d(S) := \gcd(l(S), h(S))$, and its ramification index is $e(S) = \frac{l(S)}{d(S)}$. Let d = d(S) and e = e(S). Then, the residual polynomial of F(x)associated to S is

$$R_{\lambda}(F)(y) = c_{s+de}y^d + c_{s+(d-1)e}y^{d-1} + \dots + c_{s+e}y + c_s \in \mathbb{F}_{\phi}[y] := (\mathbb{F}_p[x]/\langle \overline{\phi(x)} \rangle)[y],$$

where

$$c_i = \begin{cases} 0, & \text{if } (i, v_p(a_i(x))) \text{ lies strictly above } N_{\phi}^+(F), \\ \frac{a_i(x)}{p^{v_p(a_i(x))}} & \text{mod } (p, \phi(x)), & \text{if } (i, v_p(a_i(x))) \text{ lies on } N_{\phi}^+(F). \end{cases}$$

The polynomial F(x) is called ϕ -regular, if for each side S_{λ} of $N_{\phi}^+(F)$, $R_{\lambda}(F)(y)$ is separable in $\mathbb{F}_{\phi}[y]$. Further, F(x) is called *p*-regular, if it is ϕ -regular, for every monic polynomial $\phi(x) \in \mathbb{Z}[x]$ such that $\overline{\phi(x)}$ is a monic irreducible factor of $\overline{F(x)}$ in $\mathbb{F}_p[x]$.

Now, we have all the necessary tools to give Theorem of Ore (see [16, 27, 29]).

Theorem 3.2 (Theorem of Ore). With the above notations, suppose that

- $\overline{F(x)} = \prod_{i=1}^{t} \overline{\phi_i(x)}^{l_i}$ is the factorization of $\overline{F(x)}$ into a product of powers of distinct monic irreducible polynomials in $\mathbb{F}_p[x]$,
- $N^+_{\phi_i}(F) = S_{i1} + S_{i2} + ... + S_{ir_i}$ and λ_{ij} is the slope of S_{ij} , for each $i \in \{1, 2, ..., t\}$ and $j \in \{1, 2, ..., r_i\}$,

• $R_{\lambda_{ij}}(F)(y) = \prod_{s=1}^{s_{ij}} \psi_{ijs}(y)^{n_{ijs}}$ is the factorization of $R_{\lambda_{ij}}(F)(y)$ into a product of powers of distinct irreducible polynomials in $\mathbb{F}_{\phi_i}[x]$, for each $i \in \{1, 2, ..., t\}$ and $j \in \{1, 2, ..., r_i\}$.

Then, we have

and

- (1) $\nu_p((\mathbb{Z}_K : \mathbb{Z}[\alpha])) \ge \sum_{i=1}^t ind_{\phi_i}(F)$. The equality holds if F(x) is p-regular. (2) If F(x) is p-regular $(n_{ijs} = 1)$, then
 - $(x) \quad b \in p \quad regulation \quad (reg) \quad (reg) \quad (reg) \quad t \quad r_s$

$$p\mathbb{Z}_K = \prod_{i=1}^r \prod_{j=1}^{r_i} \prod_{s=1}^{r_{j}} \mathfrak{p}_{ijs}^{e_{ij}},$$

where e_{ij} is the ramification index of the side S_{ij} , and \mathfrak{p}_{ijs} is a prime ideal of \mathbb{Z}_K its residue degree over p is $f_{ijs} = deg(\phi_i) \times deg(\psi_{ijs})$.

Example 3.3. Consider the quartic pure field $K = \mathbb{Q}(\sqrt[4]{17})$. In this case, we have $\alpha = \sqrt[4]{17}$ and $F(x) = x^4 - 17$. Let p = 2. Reducing modulo 2, we get $\overline{F(x)} = \overline{\phi_1(x)}$ in $\mathbb{F}_2[x]$. The ϕ_1 -adic development of F(x) is

$$F(x) = -16 + 4\phi_1(x) + 6\phi_1(x)^2 + 4\phi_1(x)^3 + \phi_1(x)^4.$$

Here, we have $\nu_0 = \nu_2(-16) = 4$, $\nu_1 = \nu_2(4) = 2$, $\nu_2 = \nu_2(6) = 1$, $\nu_3 = \nu_2(4) = 2$, and $\nu_4 = \nu_2(1) = 0$. It follows that $N^+_{\phi_1}(F) = S_{11} + S_{12} + S_{13}$ has three sides of degree 1 each joining the points (0, 4), (1, 2), (2, 1) and (4, 0) (see FIGURE 1). Moreover, for j = 1, 2, 3, the residual polynomial $R_{\lambda_{1j}}(F)(y)$ has degree 1. So, it is separable. Thus, F(x) is 2-regular. By Theorem 3.2, we have

 $2\mathbb{Z}_K = \mathfrak{p}_{11} \cdot \mathfrak{p}_{12} \cdot \mathfrak{p}_{13}^2$ with the same residue degree $f(\mathfrak{p}_{1j}/2) = 1$ for j = 1, 2, 3,

$$\nu_2((\mathbb{Z}_K : \mathbb{Z}[\alpha])) = \operatorname{ind}_{\phi_1}(F) = 3.$$

In this case, by Lemma 3.1, the prime 2 divides i(K). Hence, K is not monogenic.




FIGURE 1. $N_{\phi_1}^+(F)$ with respect to ν_2 .

4. PROOFS OF THE MAIN RESULTS

In this section, we prove our results. To prove Theorem 2.1, we will use Lemma 4.1 below. Let p be a prime. Assume that p divides n and does not divide m, and set $n = u \cdot p^r$. Let $\phi(x) \in \mathbb{Z}[x]$ be a monic irreducible polynomial. Then $\overline{\phi(x)}$ is a monic irreducible factor of $\overline{F(x)}$ in $\mathbb{F}_p[x]$ if and only if $\overline{\phi(x)}$ is a monic irreducible factor of $\overline{x^u - m}$ in $\mathbb{F}_p[x]$. Let

- $x^u \underline{m} = \phi(x)U(x) + pT(x)$, where U(x) and $T(x) \in \mathbb{Z}[x]$ two polynomials such that $\overline{\phi(x)}$ does not divide $\overline{U(x)T(x)}$.
- $H(x) = m^{p^r-1}T(x) + \frac{1}{p^{r+1}} \sum_{j=0}^{p^r-2} {p^r \choose j} m^j (pT(x))^{p^r-j}.$
- Let V(x) and R(x) be the quotient and the remainder upon the Euclidean division of H(x) by $\phi(x)$, respectively.
- $A_0(x) = p^{r+1}R(x) + m^{p^r} m$, and $\nu_0 = \nu_p(A_0(x))$.

In [4], Ben Yakkou and Didi proved the following result:

Lemma 4.1. Let F(x), p, and $\phi(x)$ be as in above. Then $N_{\phi}^+(F)$ is the Newton polygon joining the points $\{(0, \nu_0)\} \cup \{(p^j, k - j) \mid 0 \le j \le r\}$ in the euclidean plane.

We recall also that if a prime integer p does not divide a non-zero rational integer m, then for every positive integer k, $\nu = \nu_p(m^{p^u} - m) = \nu_p(m^{p^u-1} - 1) = \nu_p(m^{p-1} - 1)$ (see [3]).

Proof of Theorem 2.1. With the assumptions of Theorem 2.1, let $\phi(x)$ be a monic irreducible factor of F(x) modulo p of degree d. We distinguish two cases:

- If $\nu \geq r+1$, then $\nu_0 \geq \min\{\nu, r+1\} \geq r+1$. It follows by Lemma 4.1 that the principal Newton polygon $N_{\phi}^+(F) = S_1 + \cdot + S_{r+1}$ has r+1 sides of degree 1 each. More precisely, $N_{\phi}^+(F)$ is the lower convex hull of the points $(0, \nu_0)$, $(1, r), (p, r-1), \ldots$, and $(p^r, 0)$ (see FIGURE 2). Thus, for $j = 1, \ldots, r+1$, the residual polynomial $R_{\lambda_j}(F)(y)$ is separable in $\mathbb{F}_{\phi}[y]$ since it is of degree 1. By Ore's Theorem (Theorem 3.2), the factor $\phi(x)$ provides r+1 distinct prime ideals of \mathbb{Z}_K of residue degree equals $\deg(R_{\lambda_j}(F)(y)) \times \deg(\phi(x)) = 1 \times d = d$ lying above $p\mathbb{Z}_K$. Since the number of monic irreducible factors of $\overline{F(x)}$ equals $N_p(d, u, m)$, there exist $(r+1) \times N_p(d, u, m)$ distinct prime ideals of \mathbb{Z}_K which divide $p\mathbb{Z}_K$.
- If $r \ge \nu$, then $\nu_0 \ge \min\{\nu, r+1\} = \nu$. Thus, by Lemma 4.1, $N_{\phi}^+(F) = S_1 + \cdot + S_{\nu}$ has ν sides of degree 1 each. Explicitly, $N_{\phi}^+(F)$ is the lower convex hull of the points $(0,\nu)$, $(p^{r-\nu+1},\nu-1)$, $(p^{r-\nu+2},\nu-2)$, ... and $(p^r,0)$. Thus, $R_{\lambda_j}(F)(y)$, $j=1,\ldots,r+1$ is separable. Thus, for $k=1,\ldots,N_p(d,u,m)$, F(x) is ϕ_k -regular, for every factor $\phi_k(x)$ of F(x) of degree d. The ramification indices of the sides of $N_{\phi}^+(F)$ are $e_1 = e(S_1) = p^{r-\nu+1}$, $e_j = e(S_j) = (p-1) \cdot p^{r-\nu+j-1}$, for $j=2,\ldots,\nu$. By Theorem 3.2, we get

$$p\mathbb{Z}_K = \prod_{k=1}^{N_p(d,s,b)} \prod_{j=1}^{\nu} \mathfrak{p}_{kj}^{ej} \mathfrak{a},$$

where **a** is a proper ideal of \mathbb{Z}_K , and \mathfrak{p}_{kj} is a prime ideal of \mathbb{Z}_K with residue degree $f(\mathfrak{p}_j/p) = d$, for $k = 1, \ldots, N_p(d, s, b)$ and $j = 1, \ldots, \nu$. So there are at least $\nu \times N_p(d, s, b)$ prime ideals of residue degree d lying above $p\mathbb{Z}_K$. By Lemma 3.1, if $\nu \times N_p(d, s, b) > N_p(d)$, then p divides i(K), and so K is not monogenic.

The proofs of Corollaries 2.4, 2.6, 2.5 are similar. Since they are obtained by a direct application of Theorem 2.1, we propose only the proof of Corollary 2.4.




FIGURE 2. $N_{\phi}^+(F)$ when $\nu \ge r+1$.

Proof of Corollary 2.4.

- (1) $r \geq 1, s \geq 7$, and $m^6 \equiv 1 \pmod{7^8}$. Here, we have $F(x) = x^{5^r \cdot 7^s} m \equiv (x^{5^r} m)^{7^s} \pmod{7} \equiv (\phi_1(x)V(x))^{7^s} \pmod{7}$, with $\phi_1(x) = x b_m$ and $V(b_m) \not\equiv 0 \pmod{p}$ (since $x^{5^r} m$ is square free in $\mathbb{F}_7[x]$, because 7 does not divide 5^r). Precisely, modulo 7, we have $b_1 = 1, b_2 = 4, b_3 = 5, b_4 = 2, b_5 = 3, b_6 = 6$. So, ϕ_1 is an irreducible factor of $x^{2^r} m$ modulo 7. According to the notations of Theorem 2.1, for d = 1, we have $N_7(1, 2^r, m) = 1$. Notice also that $N_7(1) = 1$. By applying Theorem 2.1, K is not monogenic if the condition $\min\{s + 1, \nu_7(m^6 1)\} \times 1 > 7$ holds, which is equivalent to $s \geq 7$, and $m^6 \equiv 1 \pmod{7^8}$.
- (2) $r \ge 5$, $s \ge 1$, and $m^4 \equiv 1 \pmod{5^6}$. In this case, for p = 5 and d = 1, we have $N_5(1, 7^s, m) = 1$. Then, similarly to the above case, by changing the role of r and s, and applying Theorem 2.1, we obtain the desired result.

Now, let us prove Theorem 2.7.

Proof of Theorem 2.7. Let p be a prime dividing a. Then $F(x) \equiv \phi_1(x)^n \pmod{p}$, where $\phi_1(x) = x$. Note that $\nu_p(a^u) = u$, since a is square-free. As gcd(u, n) = 1, $N_{\phi_1}^+(F) = S_1$ has only one side of degree 1 joining the points (0, u) and (n, 0). Thus, $deg(S_1) = gcd(u, n) = 1$, and so the degree of the residual polynomial $R_{\lambda_1}(F)(y)$ equals to 1. By Theorem of the residual polynomial, F(x) is irreducible in $\mathbb{Q}_p(x)$. So, it is irreducible in $\mathbb{Q}(x)$.

(1) Since deg $(R_{\lambda_1}(F)(y)) = 1$, the polynomial $R_{\lambda_1}(F)(y)$ is separable in $\mathbb{F}_{\phi_1}[y]$. Thus, F(x) is ϕ_1 -regular. So, it is *p*-regular. By using Theorem 3.2(2), we get

$$\nu_p(\mathbb{Z}_K : \mathbb{Z}[\alpha]) \ge ind_{\phi_1}(F) = \frac{(n-1)(u-1)}{2} \ge 2.$$

It follows that $\mathbb{Z}[\alpha] \subsetneq \mathbb{Z}_K$.

(2) Let $(t,s) \in (\mathbb{N}^*)^2$ be the positive solution of the Diophantine equation ux - ny = 1, and $\theta = \frac{\alpha^t}{a^s}$. We have

$$\theta^n = \left(\frac{\alpha^t}{a^s}\right)^n = \frac{(\alpha^n)^t}{a^{ns}} = \frac{a^{ut}}{a^{ns}} = a^{ut-ns} = a$$

Thus, $G(\theta) = 0$, where $G(x) = x^n - a \in \mathbb{Z}[x]$. Since *m* is square free, F(x) is irreducible over \mathbb{Q} (it is a *q*-Eisenstein type polynomial with respect to any prime *q* dividing *a*). Then, we consider the field $K' = \mathbb{Q}(\theta)$. Obviously $K' \subseteq K$. On the

other hand, $\alpha^t = a^s \theta \in K'$. It follows that

$$\alpha = \alpha^1 = \alpha^{ut-ns} = \frac{\alpha^{ut}}{\alpha^{ns}} = \frac{(\alpha^t)^u}{(\alpha^n)^s} = \frac{(\alpha^t)^u}{a^s} \in K'.$$

So $K = \mathbb{Q}(\alpha) = \mathbb{Q}(\theta)$. By [28, Propositions 2.9 and 2.13], one has

$$\Delta(G) = n^n \cdot a^{n-1} = \left(\mathbb{Z}_K : \mathbb{Z}[\theta]\right)^2 \cdot D_K,\tag{4.1}$$

where D_K the field discriminant of K. Let q be a prime. As $PD(n) \subseteq PD(a)$, by the above relation, if q divides the index $(\mathbb{Z}_K : \mathbb{Z}[\theta])$, then q divides a. So to show $\mathbb{Z}_K = \mathbb{Z}[\theta]$ we need to prove that $q \nmid (\mathbb{Z}_K : \mathbb{Z}[\theta])$, for every $q \in PD(a)$. Let $q \in PD(a)$. We have $G(x) \equiv \phi_1(x)^n \pmod{q}$, where $\phi_1(x) = x$. As a is square free, $\nu_q(a) = 1$. It follows that $N_{\phi_1}^+(F) = S_1$ with respect to q, has only one side of degree 1 joining the points (0, 1) and (n, 0). So, by applying Ore's Theorem with the polynomial G(x), we see that $\nu_q((\mathbb{Z}_K : \mathbb{Z}[\theta])) = 0$. This completes the proof.

5. Application to the existence of Canonical number system

Let $N_{K/\mathbb{Q}}$ be the norm function over K. The pair (θ, N_{θ}) , where $N_{\theta} = \{0, 1, \ldots, |N_{K/\mathbb{Q}}(\theta)| - 1\}$ is called a canonical number system (CNS) in \mathbb{Z}_K for an integer θ if each element of \mathbb{Z}_K can be represented uniquely as $a_0 + a_1\theta + \cdots + a_l\theta^l$, where $\pm a_i \in N_{\theta}$ for every $i = 0, \ldots, l$ and $a_l \neq 0$. We call θ the base of the number system, N_{θ} the set of digits, and the minimal polynomial F(x) of θ is called the CNS polynomial. For more details and some applications, we refer to [9, 30]. The most important connection between CNS and monogenity of is in the following result (see [25] by Kovács, and [26] by Kovács and Pethő).

Theorem 5.1. Let K is a number field of degree $n \ge 3$, $\theta \in \mathbb{Z}_K$, and $F(x) = x^n + a_{n-1} + \cdots + a_0$ such that $1 \le a_{n-1} \le \cdots \le a_0$, $a_0 \ge 2$, and $|N_{K/\mathbb{Q}}(\theta)| > 2$. Then

$$(\theta, N_{\theta})$$
 is a CNS in $\mathbb{Z}_K \Leftrightarrow \mathbb{Z}_K = \mathbb{Z}[\theta].$

Using the above theorem, Hameed, Nakahara, and Husnine identify in Theorems 2.1 and 3.1 of [23] CNS polynomials of type $x^{2^r} - m$ and $x^n - m$ with the condition m is square free. By the above theorem and our results, we deduce that:

- Under the hypotheses of Theorem 2.7, \mathbb{Z}_K admits a CNS, and such system is explicitly given (we combine Theorem 5.1 and Proof of Theorem 2.7). In this direction, our results extend those given in [23].
- Under the hypotheses of Theorem 2.1, \mathbb{Z}_K has no CNS.

We propose the following examples.

Example 5.2. Let $F(x) = x^6 - m^5$ with $(m \neq 0, \pm 1)$, *m* square free, 6 divides *m*, and $K = \mathbb{Q}(\sqrt[6]{m^5})$ (here $\alpha = \sqrt[6]{m^5}$). According to the notations of Proof of Theorem 2.7, t = 5, s = 4, and $\theta = \frac{(\sqrt[6]{m^5})^5}{m^4}$. By Theorem 2.7, θ generates a power integral basis of \mathbb{Z}_K . Note that the minimal polynomial of θ is $G(x) = x^6 - m$. By [28, Propositions 2.9 and 2.13], we have

$$6^{6} \cdot m^{5} = \Delta(G) = D_{K/\mathbb{Q}}(1, \theta, \dots, \theta^{5}) = (-1)^{\frac{6\times5}{2}} N_{K/\mathbb{Q}}(G'(\theta)) = -N_{K/\mathbb{Q}}(6\theta^{5}) = -6^{6} (N_{K/\mathbb{Q}}(\theta))^{5}$$

Example 5.3. Let $F(x) = x^{2^r \cdot 3^k} - 6^w$ with $w \ge 5$, and $w \equiv 1, 5 \pmod{6}$. Let t, s, and θ be as in Proof of Theorem 2.7, then (θ, N_{θ}) is CNS \mathbb{Z}_K , where the set of digits is $N_{\theta} = \{1, \ldots, 6\}$.

DECLARATIONS

Conflict of interest: There is no conflict of interest related to this paper. The authors have freely chosen this journal without any consideration.

DATA AVAILABILITY

Not applicable.

AUTHORS CONTRIBUTIONS

The authors are equally contributed to this paper. All authors read and improved the final manuscript.

References

- S. Ahmad, T, Nakahara and S. M. Husnine, Power integral bases for certain pure sextic fields, Int. J. Number Theory, 10(2014), 2257–2265.
- [2] S. Ahmad, T. Nakahara and A. Hameed, On certain pure sextic fields related to a problem of Hasse, Int. J. Algebra Comput., 26(2016), 577–583.
- [3] H. Ben Yakkou, A. Chillali and L. El Fadil, On Power integral bases for certain pure number fields defined by $x^{2^r \cdot 5^s} m$, Comm. Algebra, 49 (2021), 2916–2926.
- [4] H. Ben Yakkou and J. Didi, On monogenity of certain pure number fields of degrees 2^r · 3^k · 7^s, Math. Bohem., (2023). Doi: 10.21136/MB.2023.0071-22.
- [5] H. Ben Yakkou and L. El Fadil, On monogenity of certain pure number fields defined by x^{p^r} m, Int. J. Number theory, 17 (2021), 2235-2242.
- [6] Y. Bilu, I. Gaál and K. Győry, Index form equations in sextic fields: a hard computation, Acta. Arith., 115(2004), 85–96.
- [7] C. T. Davis and B. K. Spearman, The index of quartic field defined by a trinomial x⁴ + ax + b, J. Algebra Appl., 17 (2018), 1850197, 18 pp.
- [8] J. H. Evertse and K. Győry, Discriminant Equations in Diophantine Number Theory, Cambridge Univ. Press (2017).
- [9] J. H. Evertse and K. Győry, A. Pethő, et al. Number systems over general orders. Acta Math. Hungar., 159(2019), 187–205.
- [10] I. Gaál, Diophantine Equations and Power Integral Bases, Theory and algorithm, 2nd edn., Birkhäuser, (Boston, 2019).
- [11] I. Gaál and K. Győry, Index form equations in quintic fields, Acta. Arith., 89(1999), 379-396.
- [12] I. Gaál and L. Remete, Power integral bases and monogenity of pure fields, J. Number Theory, 173(2017), 129–146.
- [13] I. Gaál, A. Pethő and M. Pohst, On the resolution of index form equations in quartic number fields, J. Symbolic Comput., 16(1993), 563–584.
- [14] T. A. Gassert, A note on the monogenity of power maps, Albanian J. Math., 11(2017), 3-12.

- [15] J. Guàrdia, J. Montes and E. Nart, Higher Newton polygons in the computation of discriminants and prime ideal decomposition in number fields, J. Théor. Nombres Bordeaux, 23 (2011), 667–669.
- [16] J. Guàrdia, J. Montes and E. Nart, Newton polygons of higher order in algebraic number theory, Trans. Amer. Math. Soc., 364 (2012), 361–416.
- [17] K.Győry, Sur les polynômes à coefficients entiers et de discriminant donné, Acta. Arith., 23(1973), 419–426.
- [18] K.Győry, Sur les polynômes à coefficients entiers et de discriminant donne III, Publ. Math. Debrecen, 23(1976), 141–165.
- [19] K.Győry, On polynomials with integer coefficients and given discriminant, IV, Publ. Math. Debrecen, 25(1978), 155–167.
- [20] K. Győry, Corps de nombres algébriques d'anneau d'entiers monogène, In: "Séminaire Delange-Pisot-Poitou", 20e année: 1978/1979. Théorie des nombres, Fasc. 2 (French), Secrétariat Math., Paris, 1980, pp. Exp. No. 26, 7.
- [21] K. Győry, On discriminants and indices of integers of an algebraic number field, J. Reine Angew. Math., 324(1981), 114–126.
- [22] A. Hameed and T. Nakahara, Integral bases and relative monogenity of pure octic fields, Bull. Math. Soc. Sci. Math. Roumaine, 58(2015), 419–433.
- [23] A. Hameed, T. Nakahara, S. M. Husnine and S. Ahmad, On existence of canonical number system in certain classes of pure algebraic number fields, J. Prime Res. Math., 7 (2011) 19–24.
- [24] L. Jones, Infinite families of non-monogenic trinomials, Acta Sci. Math., (2021), 95–105.
- [25] B. Kovács, Canonical number systems in algebraic number fields, Acta. Math. Acad. Sci. Hungar., 37(1981), 405–407.
- [26] B. Kovács and A. Pethő, Number systems in integral domains, especially in orders of algebraic number fields, Acta Sci. Math. (Szeged) 55 (1991) 287–299.
- [27] J. Montes and E. Nart, On a theorem of Ore, J. Algebra 146(1992), 318-334.
- [28] W. Narkiewicz, Elementary and Analytic Theory of Algebraic Numbers, 3rd edn., Springer Monographs in Mathematics (Springer-Verlag, Berlin, 2004).
- [29] Ø. Ore, Newtonsche Polygone in der Theorie der algebraischen Körper, Math. Ann., 99(1928), 84–117.
- [30] A. Pethő, On a polynomial transformation and its application to the construction of a public key cryptosystem, in Computational Number Theory, eds. A. Pethő, M. Pohst, H. C. Williams and H. G. Zimmer (de Gruyter, Berlin, 1991), pp. 31–44.
- [31] A. Pethő and M. Pohst, On the indices of multiquadratic number fields, Acta. Arith., 153(2012), 393–414.
- [32] A. Pethő, V. Ziegler, On biquadratic fields that admit unit power integral basis, Acta Math. Hungar., 133(2011), 221–241.

FACULTY OF SCIENCES DHAR EL MAHRAZ, P.O. BOX 1874 ATLAS-FES, SIDI MOHAMED BEN ABDELLAH UNIVERSITY, MOROCCO

Email address: beyakouhamid@gmail.com, brahimboudine.bb@gmail.com