

AN INTEGRAL REPRESENTATION OF $\mathcal{R}(s)$ DUE TO GABCKE.

J. ARIAS DE REYNA

ABSTRACT. Gabcke proved a new integral expression for the auxiliary Riemann function

$$\mathcal{R}(s) = 2^{s/2} \pi^{s/2} e^{\pi i(s-1)/4} \int_{-\frac{1}{2} \searrow \frac{1}{2}} \frac{e^{-\pi i u^2/2 + \pi i u}}{2i \cos \pi u} U\left(s - \frac{1}{2}, \sqrt{2\pi} e^{\pi i/4} u\right) du,$$

where $U(\nu, z)$ is the usual parabolic cylinder function.

We give a new, shorter proof, which avoids the use of the Mordell integral. And we write it in the form

$$(1) \quad \mathcal{R}(s) = -2^s \pi^{s/2} e^{\pi i s/4} \int_{-\infty}^{\infty} \frac{e^{-\pi x^2} H_{-s}(x\sqrt{\pi})}{1 + e^{-2\pi \omega x}} dx.$$

where $H_\nu(z)$ is the generalized Hermite polynomial.

1. INTRODUCTION

The auxiliary function of Riemann is defined by the integral

$$\mathcal{R}(s) = \int_{0 \swarrow 1} \frac{x^{-s} e^{\pi i x^2}}{e^{\pi i x} - e^{-\pi i x}} dx.$$

The position of the zeros of this function is connected with the zeros of the Riemann zeta function [2]. In a paper by Gabcke [3], posted on arXiv and not published, he proved an integral expression for $\mathcal{R}(s)$ using the parabolic cylinder function $U(\nu, z)$ as defined in [1]. We prefer to give it in terms of the Hermite functions [4]. We show that

$$(2) \quad \mathcal{R}(s) = -2^s \pi^{s/2} e^{\pi i s/4} \int_{-\infty}^{\infty} \frac{e^{-\pi x^2} H_{-s}(x\sqrt{\pi})}{1 + e^{-2\pi \omega x}} dx, \quad \omega = e^{\pi i/4}.$$

We give a different proof of that of Gabcke, which is simpler and does not use the Mordell integral. Furthermore, our proof presents some integrals that may be of independent interest.

2. SOMME LEMMAS

Lemma 1. For $z = x + iy$ with $0 < x - y < 1$, we have

$$\int_L \frac{e^{-z\zeta}}{1 + e^{-\zeta}} d\zeta = \frac{\pi}{\sin \pi z},$$

where L is a line through 0 in direction $e^{\pi i/4}$.

Proof. This is an exercise in complex analysis. Let $\omega = e^{\pi i/4}$, then $\zeta = \omega u$ and

$$\int_L \frac{e^{-z\zeta}}{1 + e^{-\zeta}} d\zeta = \omega \int_{-\infty}^{\infty} \frac{e^{-\omega z u}}{1 + e^{-\omega u}} du.$$

2020 *Mathematics Subject Classification.* Primary 11M06; Secondary 30D99.

Key words and phrases. función zeta, representation integral.

Since $\omega z = 2^{-1/2}(1+i)(x+iy)$ our hypothesis implies that $0 < \alpha := \operatorname{Re}(\omega z) < 2^{-1/2}$ the integrand decays as $e^{-(\alpha-2^{-1/2})u}$ for $u \rightarrow -\infty$, and as $e^{-\alpha u}$ when $u \rightarrow +\infty$. Therefore, the integral is well defined and absolutely convergent. It is clear that the same happens to the integral along $L+w$ for any complex number w . We have

$$\int_L \frac{e^{-z\zeta}}{1+e^{-\zeta}} d\zeta - \int_{L+2\pi i} \frac{e^{-z\zeta}}{1+e^{-\zeta}} d\zeta = 2\pi i \operatorname{Res}_{\zeta=\pi i} \frac{e^{-z\zeta}}{1+e^{-\zeta}} = 2\pi i e^{-\pi iz}.$$

On the other hand, by the periodicity of e^z we have

$$\int_{L+2\pi i} \frac{e^{-z\zeta}}{1+e^{-\zeta}} d\zeta = \int_L \frac{e^{-z(\zeta+2\pi i)}}{1+e^{-\zeta}} d\zeta = e^{-2\pi iz} \int_L \frac{e^{-z\zeta}}{1+e^{-\zeta}} d\zeta.$$

It follows that

$$\int_L \frac{e^{-z\zeta}}{1+e^{-\zeta}} d\zeta = 2\pi i \frac{e^{-\pi iz}}{1-e^{-2\pi iz}} = \frac{\pi}{\sin \pi z}. \quad \square$$

The parabolic cylinder functions are solutions of the differential equation

$$\frac{d^2 w}{dz^2} = \left(\frac{z^2}{4} + a \right) w.$$

The principal solution $U(a, z)$ is defined by the condition $U(a, z) \sim z^{-a-\frac{1}{2}} e^{-z^2/4}$ it is an entire function in z and in a . There are two other alternative notation, Whittaker's $D_\nu(z)$ and Hermite functions $H_\nu(z)$. They are related by

$$D_\nu(z) = U(-\nu - 1/2, z) = 2^{-\nu/2} e^{-z^2/4} H_\nu(z/\sqrt{2}).$$

Therefore, they are equivalent. We will use the notation of Hermite functions that is more familiar to a general mathematician. It is true that the Hermite functions are the most unusual in the specific bibliography. One exception is the book by Lebedev [4]. For ν a nonnegative integer the functions $H_\nu(z)$ are the usual Hermite polynomials. Since they are entire functions, we may define them by its power series expansion [4, eq. (10.4.3)] (for ν not a nonnegative integer)

$$H_\nu(z) = \frac{1}{2\Gamma(-\nu)} \sum_{n=0}^{\infty} \Gamma\left(\frac{n-\nu}{2}\right) \frac{(-2z)^n}{n!}.$$

In Lebedev [4, eq. (10.5.2)] we find the following integral representation

$$(3) \quad H_\nu(z) = \frac{1}{\Gamma(-\nu)} \int_0^\infty e^{-t^2-2tz} t^{-\nu-1} dt, \quad \operatorname{Re} \nu < 0.$$

Proposition 2. *For any complex numbers s and z we have*

$$(4) \quad \int_{0\uparrow} x^s e^{x^2/4-xz} dx = 2i\sqrt{\pi} e^{-z^2} H_s(z),$$

where we integrate along any line of direction $e^{i\theta}$ with $\pi/4 < \theta < 3\pi/4$ that cuts the positive real axis.

Proof. An application of Cauchy's Theorem proves that the integral do not depend on the line of integration in the conditions of the proposition. We take a vertical line in the proof. First, we prove that the integral is an entire function of z and of s . Consider the line of integration $x = 1 + iu$ with $u \in \mathbf{R}$. Assume that $|s| \leq R$ and $|z| \leq R$ for some positive real number $R > 1$. We have

$$|x^s| \leq |\exp((\sigma + it)(\log|1+iu| + i \arctan u))| \leq |1+iu|^\sigma e^{\pi|t|} \leq e^{\pi R}(1+|u|)^R.$$

$$x^2/4 - zx = \frac{1}{4}(1 + iu)^2 - z - izu = \frac{1}{4} - \frac{1}{4}u^2 + \frac{1}{2}iu - z - izu,$$

so that

$$|\exp(x^2/4 - zx)| \leq e^{-u^2/4 + R + R|u| + \frac{1}{4}}.$$

Therefore,

$$|x^{-s} e^{x^2/4 - zx}| \ll_R (1 + |u|)^R e^{-u^2/4 + R|u|}.$$

Hence, the integrand is bounded by an absolutely integrable function of u . Since the integrand is holomorphic in s and z , it follows that the integral defines an entire function of z and s .

When $\sigma > -1$, applying Cauchy's Theorem, we may move the line of integration to $x = iu$, since the singularity at $x = 0$ is integrable, and the bound we have proved slightly modified still applies. We find that our integral is equal in this case to

$$ie^{\pi is/2} \int_0^\infty u^s e^{-u^2/4 - izu} du + ie^{-\pi is/2} \int_0^\infty u^s e^{-u^2/4 + izu} du.$$

By (3) this is equal to

$$ie^{\pi is/2} 2^{1+s} \Gamma(1+s) H_{-s-1}(iz) + ie^{-\pi is/2} 2^{1+s} \Gamma(1+s) H_{-s-1}(-iz).$$

Equivalently

$$i2^{s+1} \Gamma(1+s) [e^{\pi is/2} H_{-s-1}(iz) + e^{-\pi is/2} H_{-s-1}(-iz)].$$

In Lebedev [4, eq. (10.3.4)] it is proved that

$$H_\nu(z) = \frac{2^\nu \Gamma(\nu+1)}{\sqrt{\pi}} e^{z^2} [e^{\pi i \nu/2} H_{-\nu-1}(iz) + e^{-\pi i \nu/2} H_{-\nu-1}(-iz)].$$

Hence, our integral

$$\int_{0\uparrow} x^s e^{x^2/4 - zx} dx = 2i\sqrt{\pi} e^{-z^2} H_s(z),$$

as we wanted to prove. This has been proved only for $\sigma > -1$ but is then true for any s and z by analyticity. \square

3. MAIN THEOREM

Theorem 3. *For any complex number s we have*

$$(5) \quad \mathcal{R}(s) = -2^s \pi^{s/2} e^{\pi is/4} \int_{-\infty}^\infty \frac{e^{-\pi x^2} H_{-s}(x\sqrt{\pi})}{1 + e^{-2\pi \omega x}} dx.$$

Proof. By definition

$$\mathcal{R}(s) = \int_{0\swarrow 1} \frac{x^{-s} e^{\pi i x^2}}{e^{\pi i x} - e^{-\pi i x}} dx.$$

Take $x = \frac{1}{2} - \omega u$ as the line of integration. Any x in this line satisfies the conditions of lemma 1. Therefore,

$$\mathcal{R}(s) = \frac{1}{2\pi i} \int_{0\swarrow 1} x^{-s} e^{\pi i x^2} \left(\int_L \frac{e^{-x\zeta}}{1 + e^{-\zeta}} d\zeta \right) dx,$$

where L is the line $\zeta = \omega v$ with $v \in \mathbf{R}$. The function

$$-i \frac{x^{-s} e^{\pi i x^2 - x\zeta}}{1 + e^{-\zeta}}, \quad \zeta = \omega v, \quad x = \frac{1}{2} - \omega u,$$

is absolutely integrable in $(u, v) \in \mathbf{R}^2$. In fact, we have for $|s| \leq R$

$$\begin{aligned} |x^{-s}| &= |\exp(-(\sigma + it)(\log|\frac{1}{2} - \omega u| + i \arg(\frac{1}{2} - \omega u)))| \\ &\ll_R \max(2^{\frac{3R}{2}}, (1 + |u|)^R) e^{\pi R} \ll_R (1 + |u|)^R. \end{aligned}$$

$$\pi i x^2 - x\zeta = \pi i (\frac{1}{2} - \omega u)^2 - (\frac{1}{2} - \omega u)\omega v = \frac{1}{4}\pi i - \pi u^2 - \pi i \omega u - \frac{1}{2}\omega v + iuv,$$

so that

$$|\exp(\pi i x^2 - x\zeta)| = \exp(-\pi u^2 + \pi u/\sqrt{2} - v/2\sqrt{2}).$$

$1 + e^{-\zeta} = 1 + e^{-\omega v}$ is a continuous function that does not vanish for $v \in \mathbf{R}$ and

$$|1 + e^{-\zeta}| = |1 + e^{-v/\sqrt{2} - iv/\sqrt{2}}| \gg \max(e^{-v/\sqrt{2}} - 1, 1).$$

Hence,

$$\left| -i \frac{x^{-s} e^{\pi i x^2 - x\zeta}}{1 + e^{-\zeta}} \right| \ll_R (1 + |u|)^R \exp(-\pi u^2 + \pi u/\sqrt{2}) \cdot \frac{e^{-v/2\sqrt{2}}}{\max(e^{-v/\sqrt{2}} - 1, 1)},$$

is bounded by the product of an integrable function of u by an integrable function of v .

Hence, we may interchange the order of integration so that

$$\mathcal{R}(s) = \frac{1}{2\pi i} \int_L \left(\int_{0 \swarrow 1} x^{-s} e^{\pi i x^2 - x\zeta} dx \right) \frac{d\zeta}{1 + e^{-\zeta}}.$$

In the inner integral change variable putting $\xi = 2\sqrt{\pi}\omega x$, when x runs through the line $0 \swarrow 1$ the variable ξ runs through $\sqrt{\pi}\omega - 2i\sqrt{\pi}\omega$ that is a vertical line $0 \downarrow$ so that by Proposition 2

$$\begin{aligned} \int_{0 \swarrow 1} x^{-s} e^{\pi i x^2 - x\zeta} dx &= -\frac{2^s \pi^{s/2} e^{\pi i s/4}}{2\sqrt{\pi}\omega} \int_{0 \uparrow} \xi^{-s} e^{\xi^2/4 - \xi\zeta/2\sqrt{\pi}\omega} d\xi \\ &= -\omega 2^s \pi^{s/2} e^{\pi i s/4} e^{-\frac{\zeta^2}{4\pi i}} H_{-s} \left(\frac{\zeta}{2\sqrt{\pi}\omega} \right). \end{aligned}$$

Hence,

$$\mathcal{R}(s) = -\frac{\omega}{2\pi i} 2^s \pi^{s/2} e^{\pi i s/4} \int_L \frac{e^{-\frac{\zeta^2}{4\pi i}} H_{-s} \left(\frac{\zeta}{2\sqrt{\pi}\omega} \right)}{1 + e^{-\zeta}} d\zeta.$$

Finally, take the parametrization $\zeta = 2\pi\omega v$ of the integration line to get

$$\mathcal{R}(s) = -2^s \pi^{s/2} e^{\pi i s/4} \int_{-\infty}^{\infty} \frac{e^{-\pi v^2} H_{-s}(\sqrt{\pi}v)}{1 + e^{-2\pi\omega v}} dv. \quad \square$$

REFERENCES

- [1] M. ABRAMOWITZ, I. A. STEGUN, *Handbook of Mathematical Functions with Formulas, Graphs and Mathematical Tables*, National Bureau of Standards Appl. Series, vol 55, U. S. Government Printing Office, Washington, DC, 1964.
- [2] J. ARIAS DE REYNA, *Riemann's auxiliary function: Basic Results*, [arXiv:2406.02403](https://arxiv.org/abs/2406.02403).
- [3] W. GABCKE, *A Parabolic Cylinder Function in the Riemann-Siegel Integral Formula*, [arXiv:1512.01186v1](https://arxiv.org/abs/1512.01186v1), 6pp. (2015).
- [4] N. N. LEBEDEV, *Special functions and their applications*, Revised ed., translated from the Russian and ed. by Richard A. Silverman. Dover Publ. Inc., New York 1972.




FIGURE 1. x-ray of $H_{-\frac{1}{2}-10i}(z\sqrt{\pi})e^{-\pi z^2}(1+e^{-2\pi\omega z})^{-1}$ in the square $(-6,6)^2$




FIGURE 2. x-ray of $H_{-\frac{1}{2}-20i}(z\sqrt{\pi})e^{-\pi z^2}(1+e^{-2\pi\omega z})^{-1}$ in the square $(-6,6)^2$