A generalized version of Holmstedt's formula for the K-functional

Irshaad Ahmed¹, Alberto Fiorenza² and Amiran Gogatishvili³

¹Department of Mathematics, Sukkur IBA University, Sukkur, Pakistan.

irshaad.ahmed@iba-suk.edu.pk

²Università di Napoli Federico II, Dipartimento di Architettura, via Monteoliveto, 3, 80134 - Napoli, Italy and Consiglio Nazionale delle Ricerche, Istituto per le Applicazioni del Calcolo "Mauro Picone", Sezione di Napoli, via Pietro Castellino, 111, 80131 - Napoli, Italy.

fiorenza@unina.it

³Institute of Mathematics of the Czech Academy of Sciences - Žitná, 115 67 Prague 1, Czech Republic. gogatish@math.cas.cz

Abstract

Let (A_0, A_1) be a compatible couple of quasi-normed spaces, and let Φ_0 and Φ_1 be two general parameters of K-interpolation method. We compute K-functional for the couple $((A_0, A_1)_{\Phi_0}, (A_0, A_1)_{\Phi_1})$ in terms of K-functional for the couple (A_0, A_1) .

Key words: K-functional, K-interpolation spaces, Holmstedt's formula

MSC 2020: 46B70

1 Introduction

Let (A_0, A_1) be a compatible couple of quasi-normed spaces. For each $f \in A_0 + A_1$ and t > 0, the Peetre's K-functional is defined by

$$K(t, f) = K(t, f; A_0, A_1)$$

= $\inf\{\|f_0\|_{A_0} + t\|f_1\|_{A_1} : f_0 \in A_0, f_1 \in A_1, f = f_0 + f_1\}.$

Let Φ be a quasi-normed space of Lebesgue measurable functions defined on $(0, \infty)$ with monotone quasi-norm, that is, $|g| \leq |h|$ implies $||g||_{\Phi} \leq ||h||_{\Phi}$. Assume that $t \mapsto \min(1, t) \in \Phi$. The general K-interpolation $\bar{A}_{\Phi} = (A_0, A_1)_{\Phi}$ is formed of those $f \in A_0 + A_1$ for which the quasi-norm

$$||f||_{\bar{A}_{\Phi}} = ||K(t,f)||_{\Phi}$$

is finite; see [7]. We call Φ as general parameter of K-interpolation method.

Let $(\theta, q) \in ([0, 1] \times [1, \infty]) \setminus (\{0, 1\} \times [1, \infty])$. In particular, let Φ be given by

$$\|g\|_{\Phi} = \left(\int_0^\infty [t^{-\theta}|g(t)|]^q \frac{dt}{t}\right)^{1/q}$$

with the usual modification when $q = \infty$. Then we put $\bar{A}_{\Phi} = \bar{A}_{\theta,q}$, and $\bar{A}_{\theta,q}$ is the classical scale of K-interpolation spaces (see [5, 6, 24]). In 1970, Tord Holmstedt computed, up to equivalence of constants, K-functional for the couple $(\bar{A}_{\theta_0,q_0}, \bar{A}_{\theta_1,q_1})$ in terms of K-functional for the couple (A_0, A_1) . The precise formula is as follows. Let $0 < q_0, q_1 \le \infty$ and $0 < \theta_0 < \theta_1 < 1$. Then for all $f \in A_0 + A_1$ and for all t > 0, we have

$$\begin{split} K(t^{\theta_1-\theta_0}, f; \bar{A}_{\theta_0, q_0}, \bar{A}_{\theta_1, q_1}) &\approx & \|u^{-\theta_0-1/q_0} K(u, f)\|_{q_0, (0, t)} \\ &+ t^{\theta_1-\theta_0} \|u^{-\theta_1-1/q_1} K(u, f)\|_{q_1, (t, \infty)}, \end{split}$$

(see [22, Theorem 2.1]). Since then several authors have obtained generalizations or variants of Holmstedt's formula (see, for instance, [1–4, 8–12, 14–21, 23]).

Let Φ_0 and Φ_1 be two general parameters of K-interpolation method. In this paper, we compute the K-functional for the couple $(\bar{A}_{\Phi_0}, \bar{A}_{\Phi_1})$ in terms of K-functional for the couple (A_0, A_1) under certain appropriate separation conditions imposed on Φ_0 and Φ_1 . Our computation provides a general frame work to derive Holmstedt-type estimates. In particular, all those Holmstedt type formulae (except the one in [3, Theorem 3.5]) contained in the papers [2–4, 8–10, 12, 14–21, 23], which express the K-functional of the couple $(\bar{A}_{\Phi_0}, \bar{A}_{\Phi_1})$, for special cases of Φ_0 and Φ_1 , in terms of that of (A_0, A_1) , can be derived immediately by our general computation.

2 A notation and slowly varying functions

2.1 Notation

Let f and g be two positive functions defined on $(0, \infty)$. We write

$$f(t) \lesssim g(t), t > 0$$

if there exists a constant c > 0 such that

$$f(t) \le cg(t), \ t > 0.$$

We simply write

if

$$f(t) \leq g(t), t > 0$$
, and $g(t) \leq f(t), t > 0$.

 $f(t) \approx g(t), \ t > 0,$

2.2 Slowly varying functions

Let $b: (0, \infty) \to (0, \infty)$ be a Lebesgue measurable function. Following [20], we say b is slowly varying on $(0, \infty)$ if for every $\varepsilon > 0$, there are positive functions g_{ε} and $g_{-\varepsilon}$ on $(0, \infty)$ such that g_{ε} is non-decreasing and $g_{-\varepsilon}$ is non-increasing, and we have

$$t^{\varepsilon}b(t) \approx g_{\varepsilon}(t)$$
 and $t^{-\varepsilon}b(t) \approx g_{-\varepsilon}(t)$ for all $t \in (0, \infty)$.

We denote the class of all slowly varying functions by SV. Let $\mathbb{A} = (\alpha_0, \alpha_\infty) \in \mathbb{R}^2$. Define

$$\ell^{\mathbb{A}}(t) = \begin{cases} (1 - \ln t)^{\alpha_0}, & 0 < t \le 1\\ (1 + \ln t)^{\alpha_\infty}, & t > 1, \end{cases}$$

Then $\ell^{\mathbb{A}} \in SV$. In addition, the class SV contains compositions of appropriate log-functions, $\exp|\log t|^{\alpha}$ with $\alpha \in (0, 1)$, etc.

We collect in next Proposition some elementary properties of slowly varying functions, which we will need in Section 4. The proofs of these assertions can be carried out as in [20, Lemma 2.1] or [13, Proposition 3.4.33].

Proposition 2.1 Given $b, b_1, b_2 \in SV$, the following assertions hold:

- (i) $b_1b_2 \in SV$ and $b^r \in SV$ for each $r \in \mathbb{R}$.
- (ii) If $\alpha > 0$, then

$$\int_0^t s^{\alpha} b(s) \frac{ds}{s} \approx t^{\alpha} b(t), \quad t > 0.$$

(iii) If $\alpha > 0$, then

$$\int_{t}^{\infty} s^{-\alpha} b(s) \frac{ds}{s} \approx t^{-\alpha} b(t), \quad t > 0.$$

(iv) Assume that

$$\int_0^1 b(t) \frac{dt}{t} < \infty,$$

 $and \ set$

$$B(t) = \int_0^t b(s) \frac{ds}{s}, \quad t > 0.$$

Then $B \in SV$, and $b(t) \leq B(t), t > 0$.

(v) Assume that

$$\int_1^\infty b(t) \frac{dt}{t} < \infty,$$

 $and \ set$

$$\tilde{B}(t) = \int_{t}^{\infty} b(s) \frac{ds}{s}, \quad t > 0.$$

Then $\tilde{B} \in SV$, and $b(t) \leq \tilde{B}(t), t > 0$.

3 Main result

Let Φ_0 and Φ_1 be two general parameters of K-interpolation method. In order to formulate our results we need the following conditions on a positive weight ρ defined on $(0, \infty)$.

(C₁)
$$\frac{t \|\chi_{(t,\infty)}\|_{\Phi_0}}{\|\min(u,t)\|_{\Phi_1}} \lesssim \rho(t) \lesssim \frac{\|\min(u,t)\|_{\Phi_0}}{\|u\chi_{(0,t)}(u)\|_{\Phi_1}}, \quad t > 0$$

(C₂)
$$\|\chi_{(0,t)}(u)u\|\min(s,u)\|_{\Phi_1}^{-1}\|_{\Phi_0} \lesssim \rho(t), t > 0$$

(C₃)
$$\|\chi_{(t,\infty)}(u)u\| \min(s,u)\|_{\Phi_0}^{-1}\|_{\Phi_1} \lesssim \frac{1}{\rho(t)}, t > 0.$$

(C₄) $t\|\chi_{(t,\infty)}\|_{\Phi_0} \lesssim \|u\chi_{(0,t)}(u)\|_{\Phi_0} + t\rho(t)\|\chi_{(t,\infty)}\|_{\Phi_1}, t > 0.$

Lemma 3.1 [4, Theorem 4 (Case 1)] Let (A_0, A_1) be a compatible couple of quasi-normed spaces, and let Φ_0 and Φ_1 be two general parameters of K-interpolation method. Assume that a weight ρ satisfies the conditions (C_1) , (C_2) and (C_3) . Then, for all $f \in A_0 + A_1$ and for all t > 0, the following two-sided estimate

$$K(\rho(t), f; \bar{A}_{\Phi_0}, \bar{A}_{\Phi_1}) \approx \|\chi_{(0,t)}(u)K(u, f)\|_{\Phi_0} + \rho(t)\|\chi_{(t,\infty)}(u)K(u, f)\|_{\Phi_1} + \|\chi_{(t,\infty)}\|_{\Phi_0}K(t, f) + \rho(t)\|u\chi_{(0,t)}(u)\|_{\Phi_1}\frac{K(t, f)}{t},$$
(3.1)

holds.

The main result of this paper is following.

Theorem 3.2 Let (A_0, A_1) be a compatible couple of quasi-normed spaces, and let Φ_0 and Φ_1 be two general parameters of K-interpolation method. Put

$$\rho(t) = \frac{\|\min(u, t)\|_{\Phi_0}}{\|\min(u, t)\|_{\Phi_1}}, \ t > 0.$$

(i) Assume that ρ satisfies the conditions (C₂) and (C₃). Then, for all $f \in A_0 + A_1$ and for all t > 0, the following two-sided estimate

$$K(\rho(t), f; \bar{A}_{\Phi_0}, \bar{A}_{\Phi_1}) \approx \|\chi_{(0,t)}(u)K(u, f)\|_{\Phi_0} + \rho(t)\|\chi_{(t,\infty)}(u)K(u, f)\|_{\Phi_1} + \|\chi_{(t,\infty)}\|_{\Phi_0}K(t, f),$$
(3.2)

holds.

(ii) Assume that ρ satisfies the conditions (C₂), (C₃) and (C₄). Then, for all $f \in A_0 + A_1$ and for all t > 0, the following two-sided estimate

$$K(\rho(t), f; \bar{A}_{\Phi_0}, \bar{A}_{\Phi_1}) \approx \|\chi_{(0,t)}(u)K(u, f)\|_{\Phi_0} + \rho(t)\|\chi_{(t,\infty)}(u)K(u, f)\|_{\Phi_1},$$
(3.3)

holds.

Proof. First we derive the formula (3.2) using Lemma 3.1. Since (for j = 0, 1)

$$\|\min(u,t)\|_{\Phi_j} \approx \|u\chi_{(0,t)}(u)\|_{\Phi_j} + t\|\chi_{(t,\infty)}\|_{\Phi_j}, \ t > 0,$$

the condition (C_1) is satisfied trivially by our choice of ρ . Next note that

$$\rho(t) \| u\chi_{(0,t)}(u) \|_{\Phi_1} \frac{K(t,f)}{t} \lesssim \| \chi_{(t,\infty)} \|_{\Phi_0} K(t,f) + \| u\chi_{(0,t)}(u) \|_{\Phi_0} \frac{K(t,f)}{t}.$$
(3.4)

Since $t \mapsto K(t, f)/t$ is non-increasing (see [5, Proposition 1.2]), we have

$$\|\chi_{(0,t)}(u)K(u,f)\|_{\Phi_0} \ge \frac{K(t,f)}{t} \|u\chi_{(0,t)}(u)\|_{\Phi_0},$$

therefore, (3.4) reduces to

$$\rho(t) \| u\chi_{(0,t)}(u) \|_{\Phi_1} \frac{K(t,f)}{t} \lesssim \| \chi_{(t,\infty)} \|_{\Phi_0} K(t,f) + \| \chi_{(0,t)}(u) K(u,f) \|_{\Phi_0}.$$
(3.5)

Finally, in view of (3.5), the estimate (3.2) follows from (3.1).

Next we derive (3.3). Since $t \mapsto K(t, f)$ is non-decreasing (see [5, Proposition 1.2]) and $t \mapsto K(t, f)/t$ is non-increasing, we have

$$\|\chi_{(0,t)}(u)K(u,f)\|_{\Phi_0} + \rho(t)\|\chi_{(t,\infty)}(u)K(u,f)\|_{\Phi_1} \ge \frac{K(t,f)}{t}\|u\chi_{(0,t)}(u)\|_{\Phi_0} + \rho(t)K(t,f)\|\chi_{(t,\infty)}\|_{\Phi_1},$$

combining above inequality with the condition (C_4) , we arrive at

$$\|\chi_{(t,\infty)}\|_{\Phi_0}K(t,f) \lesssim \|\chi_{(0,t)}(u)K(u,f)\|_{\Phi_0} + \rho(t)\|\chi_{(t,\infty)}(u)K(u,f)\|_{\Phi_1}, \ t > 0,$$

now inserting the above estimate in (3.2) yields (3.3). The proof is complete.

Remark 3.3 The essential difference between the general computation in our main result and that of [4, Theorem 4] is that the weight ρ is explicitly given in our main result.

4 Scope of the main result

As mentioned in the Introduction, almost all those Holmstedt-type estimates, contained in the papers [2–4,8–10,12,14–21,23], which express the couple $(\bar{A}_{\Phi_0}, \bar{A}_{\Phi_1})$, for special cases of Φ_0 and Φ_1 , in terms of (A_0, A_1) , can be derived immediately by our general computation. In this section, we illustrate this in some instances.

(i) Let $0 < q_0, q_1 \le \infty, 0 \le \theta_0, \theta_1 \le 1$, and let b_j (j = 0, 1) be a slowly varying function. Define

$$\|g\|_{\Phi_{\theta_j,q_j;b_j}} = \left(\int_0^\infty \left[t^{-\theta_j} b_j(t) |g(t)|\right]^{q_j} \frac{dt}{t}\right)^{1/q_j},$$

with the usual modification when $q_j = \infty$ (j = 0, 1). First consider the case $0 < \theta_0 < \theta_1 < 1$. Thanks to Proposition 2.1, it is easy to compute that

$$\|\min(u,t)\|_{\Phi_{\theta_j,q_j;b_j}} \approx t^{1-\theta_j}b_j(t), \ t>0.$$

The conditions (C_2) , (C_3) and (C_4) are also met again thanks to Proposition 2.1. As a consequence of Theorem 3.2 (ii), we recover an extension of Holmstedt's formula contained in [20, Theorem 3.1 (a)]. When $b_j \equiv 1$, we get back the classical Holmstedt's formula.

Next we consider the case when $\theta_0 = 0$ and $\theta_1 = 1$. Again thanks to Proposition 2.1, the conditions (C_2) and (C_3) are met, and also we have the following computations

$$\|\min(u,t)\|_{\Phi_{0,q_0;b_0}} \approx t \left(\int_t^\infty b_0^{q_0}(s) \frac{ds}{s}\right)^{1/q_0}, \ t > 0,$$

and

$$\|\min(u,t)\|_{\Phi_{1,q_1;b_1}} \approx \left(\int_0^t b_1^{q_1}(s)\frac{ds}{s}\right)^{1/q_1}, \ t > 0.$$

Let us put $\bar{A}_{j,q_j;b_j} = \bar{A}_{\Phi_{j,q_j;b_j}}$ (j = 0, 1). As a consequence of Theorem 3.2 (i), for all $f \in A_0 + A_1$ and t > 0, we get the following formula

$$\begin{split} K\left(\rho(t), f; \bar{A}_{0,q_0:b_0}, \bar{A}_{1,q_1:b_1}\right) &\approx & \left(\int_0^t b_0^{q_0}(s) K^{q_0}(s, f) \frac{ds}{s}\right)^{1/q_0} + \rho(t) \left(\int_t^\infty b_1^{q_1}(s) K^{q_1}(s, f) \frac{ds}{s}\right)^{1/q_1} \\ &+ \left(\int_t^\infty b_0^{q_0}(s) \frac{ds}{s}\right)^{1/q_0} K(t, f), \end{split}$$

where

$$\rho(t) = \frac{t \left(\int_t^\infty b_0^{q_0}(s) \frac{ds}{s}\right)^{1/q_0}}{\left(\int_0^t b_1^{q_1}(s) \frac{ds}{s}\right)^{1/q_1}}, \ t > 0.$$

This formula also follows from [4, Example 5], however computations in [4, Example 5] are tedious as compared to our straightforward computations.

Next we consider the case when $\theta_0 = \theta_1 = 0$ with $q_0 \neq q_1$. This time (thanks to Proposition 2.1) we have (for j = 0, 1)

$$\|\min(u,t)\|_{\Phi_{j,q_j;b_j}} \approx t \left(\int_t^\infty b_j^{q_j}(s) \frac{ds}{s}\right)^{1/q_j}, \ t > 0.$$

The condition (C_4) is met while the conditions (C_2) and (C_3) turn into

$$\left(\int_0^t \frac{b_0^{q_0}(x)}{\left(\int_x^\infty b_1^{q_1}(s)\frac{ds}{s}\right)^{q_0/q_1}}\frac{dx}{x}\right)^{1/q_0} \lesssim \frac{\left(\int_t^\infty b_0^{q_0}(s)\frac{ds}{s}\right)^{1/q_0}}{\left(\int_t^\infty b_1^{q_1}(s)\frac{ds}{s}\right)^{1/q_1}}, \ t > 0,$$

and

$$\left(\int_{t}^{\infty} \frac{b_{1}^{q_{1}}(x)}{\left(\int_{x}^{\infty} b_{0}^{q_{0}}(s)\frac{ds}{s}\right)^{q_{1}/q_{0}}}\frac{dx}{x}\right)^{1/q_{1}} \lesssim \frac{\left(\int_{t}^{\infty} b_{1}^{q_{1}}(s)\frac{ds}{s}\right)^{1/q_{1}}}{\left(\int_{t}^{\infty} b_{0}^{q_{0}}(s)\frac{ds}{s}\right)^{1/q_{0}}}, \quad t > 0.$$

As a consequence of Theorem 3.2 (ii), we recover the Holmstedt-type estimate contained in [3, Theorem 3.2]. It is worthy of mention that it has been established in [3, Theorem 3.2] that a sufficient condition for the previous two estimates to be held is that the following function $(6^{\infty} + q_0) + d_{\infty})^{1/q_0(1+\epsilon)}$

$$t \mapsto \frac{\left(\int_t^\infty b_0^{q_0}(s)\frac{ds}{s}\right)^{1/q_0(1+\epsilon)}}{\left(\int_t^\infty b_1^{q_1}(s)\frac{ds}{s}\right)^{1/q_1}}$$

is equivalent to a nondecreasing function for some $\epsilon > 0$.

(ii) Next we turn to the papers [17–19]. Let $0 \le \theta \le 1$, and let a, b be slowly varying functions. Let E, F be rearrangement invariant spaces (see [5]), and let \tilde{E} and \tilde{F} be the corresponding rearrangement invariant spaces with respect to homogeneous measure dt/t (see [15, p. 135]). Define

$$\|g\|_{\Phi_{\theta,b,E}} = \left\|t^{-\theta}b(t)g(t)\right\|_{\tilde{E}}$$

$$\|g\|_{\Phi_{\mathcal{L};\theta,b,E,a,F}} = \left\|b(t) \left\|s^{-\theta}a(s)g(s)\right\|_{\tilde{F}(0,t)}\right\|_{\tilde{E}}.$$

and

$$\|g\|_{\Phi_{\mathcal{R};\theta,b,E,a,F}} = \left\|b(t) \left\|s^{-\theta}a(s)g(s)\right\|_{\tilde{F}(t,\infty)}\right\|_{\tilde{E}},$$

Using the estimates in [15, Lemma 2.3], we can easily make the following straightforward computations. For $0 < \theta < 1$, we have

$$\|\min(u,t)\|_{\Phi_{\theta,q,E}} \approx t^{1-\theta}b(t), \ t > 0,$$

$$\|\min(u,t)\|_{\Phi_{\mathcal{L};\theta,b,E,a,F}} \approx t^{1-\theta}a(t)\|b\|_{\tilde{E}(t,\infty)}, \ t > 0$$

and

and

$$\|\min(u,t)\|_{\Phi_{\mathcal{R};\theta,b,E,a,F}} \approx t^{1-\theta}a(t)\|b\|_{\tilde{E}(0,t)}, \ t>0.$$

And also we have

$$\|\min(u,t)\|_{\Phi_{0,q,E}} \approx t \|b\|_{\tilde{E}(t,\infty)}, \quad t > 0,$$
$$\|\min(u,t)\|_{\Phi_{1,q,E}} \approx \|b\|_{\tilde{E}(0,t)}, \quad t > 0.$$

Now using the above computations and estimates in [15, Lemma 2.3], we can immediately derive, from Theorem 3.2, all the Holmstedt-type estimates contained in [17, Theorem 3.1], [17, Theorem 3.2], [17, Theorem 3.4], [18, Theorem 3.1], [18, Theorem 3.2], [19, Theorem 3.1], [19, Theorem 3.2], [19, Theorem 3.3] and [19, Theorem 3.4]. It is worthy of mention that authors of above mentioned papers have to carry out a lot of computations in each separate case. While, on the other hand, our main result provides a general framework where all these separate cases can be treated simultaneously by means of handful computations.

Acknowledgements. The research of A. Gogatishvili was partially supported by the Czech Academy of Sciences (RVO 67985840), by Czech Science Foundation (GAČR), grant no: 23-04720S, and by Shota Rustaveli National Science Foundation of Georgia (SRNSFG), grant no: FR21-12353.

References

- I. Ahmed, D. E. Edmunds, W. D. Evans and G. E. Karadzhov, Reiteration theorems for the K-interpolation method in limiting cases, Math. Nachr. 284 (2011), 421-442.
- [2] I. Ahmed and T. Ejaz, Reiteration theorems for two-parameter limiting real interpolation methods, J. Funct. Spaces vol. 2016, Article ID 8214643, 9 pages, 2016. https://doi.org/10.1155/2016/8214643
- [3] I. Ahmed, A. Fiorenza and A. Gogatishvili, Holmstedt's formula for the K-functional: the limit case $\theta_0 = \theta_1$, Math. Nachr. **296** (2023), 5474-5492. https://doi.org/10.1002/mana.202200440
- [4] I. Ahmed, G. E. Karadzhov and A. Raza, General Holmstedt's formulae for the K-functional, J. Funct. Spaces vol. 2017, Article ID 4958073, 9 pages, 2017. https://doi.org/10.1155/2017/4958073
- [5] C. Bennett and R. Sharpley, Interpolation of Operators, Academic Press, Boston, 1988.
- [6] J. Bergh and J. Löfström, Interpolation Spaces. An Introduction, Springer, Berlin, 1976.
- [7] Y. A. Brudnyi and N. Y. Krugljak, Interpolation Functors and Interpolation Spaces, Volume I, North-Holland, Amsterdam, 1991.
- [8] R. Ya. Doktorski, Reiteration relations of the real interpolation method, Soviet Math. Dokl. 44 (1992), 665-669.
- [9] R. Ya. Doktorski, Reiteration formulae for the real interpolation method including L or R limiting spaces, J. Funct. Spaces vol. 2020, Article ID 6685993, 15 pages, 2020. https://doi.org/10.1155/2020/6685993
- [10] R. Ya. Doktorski, Some reiteration theorems for *R*, *L*, *RR*, *RL*, *LR*, and *LL* limiting interpolation spaces, *J. Funct. Spaces* Vol. 2021, Article ID 8513304, 31 pages, 2021. https://doi.org/10.1155/2021/8513304
- [11] R. Ya. Doktorski, P. Fernández-Martínez and T. Signes, Reiteration theorem for *R* and *L*-spaces with the same parameter, J. Math. Anal. Appl. 508 (2022): 125846.
- [12] R. Ya. Doktorski, P. Fernández-Martínez and T. Signes, The K-functional and reiteration theorems for left and right spaces, Part I, J. Math. Anal. Appl. 531 (2024): 127882.
- [13] D. E. Edmunds and W. D. Evans, Hardy operators, Function Spaces and Embeddings, Springer, Berlin, 2004.
- [14] W. D. Evans, B. Opic and L. Pick, Real interpolation with logarithmic functors, J. Inequal. Appl. 7 (2002), 187-269.
- [15] P. Fernández-Martínez and T. Signes, Real interpolation with symmetric spaces and slowly varying functions, Q. J. Math. 63 (2012), 133-164.
- [16] P. Fernández-Martínez and T. Signes, Limit cases of reiteration theorems, Math. Nachr. 288 (2015), 25-47.

- [17] P. Fernández-Martínez and T. Signes, General reiteration theorems for *R* and *L* classes: Case of left *R*-spaces and right *L*-spaces, *J. Math. Anal. Appl.* **494** (2021) 124649.
- [18] P. Fernández-Martínez and T. Signes, General reiteration theorems for *L* and *R* classes: Case of right *R*-spaces and left *L*-spaces, *Mediterr. J. Math.* 19, 193 (2022).
- [19] P. Fernández-Martínez and T. Signes, General reiteration theorems for R and L classes: mixed interpolation of R and L-spaces, Positivity 26, 47 (2022). https://doi.org/10.1007/s11117-022-00888-z
- [20] A. Gogatishvili, B. Opic and W. Trebels, Limiting reiteration for real interpolation with slowly varying functions, *Math. Nachr.* 278 (2005), 86-107.
- [21] H. P. Heinig, Interpolation of quasi-normed spaces involving weights, CMS Conference Proceedings, 1 (1981), 245-267.
- [22] T. Holmstedt, Interpolation of quasi-normed spaces, Math. Scand. 26 (1970), 177-199.
- [23] L. E. Persson, Interpolation with a function parameter, Math. Scand. 59 (1986), 199-222.
- [24] H. Triebel, Interpolation Theory, Function Spaces, Differential Operators, North-Holland, Amsterdam, 1978.